# A UNIQUE CARTAN SUBALGEBRA RESULT FOR BERNOULLI ACTIONS OF WEAKLY AMENABLE GROUPS

CHANGYING DING

ABSTRACT. We show that if  $\Gamma \curvearrowright (X^{\Gamma}, \mu^{\Gamma})$  is a Bernoulli action of an i.c.c. nonamenable group  $\Gamma$  which is weakly amenable with Cowling-Haagerup constant 1, and  $\Lambda \curvearrowright (Y, \nu)$  is a free ergodic p.m.p. algebraic action of a group  $\Lambda$ , then the isomorphism  $L^{\infty}(X^{\Gamma}) \rtimes \Gamma \cong L^{\infty}(Y) \rtimes \Lambda$  implies that  $L^{\infty}(X^{\Gamma})$  and  $L^{\infty}(Y)$  are unitarily conjugate. This is obtained by showing a new rigidity result of non properly proximal groups and combining it with a rigidity result of properly proximal groups from [BIP21].

#### 1. INTRODUCTION

The group measure space construction associates to every probability measure preserving (p.m.p.) action  $\Gamma \curvearrowright (X, \mu)$  of a countable group  $\Gamma$ , a finite von Neumann algebra  $L^{\infty}(X) \rtimes \Gamma$  [MvN43]. When the action is free and ergodic,  $L^{\infty}(X) \rtimes \Gamma$  is a II<sub>1</sub> factor and  $L^{\infty}(X)$  is a Cartan subalgebra. With the discovery of Popa's deformation/rigidity theory [Pop06a, Pop06b, Pop06c], spectacular progress has been made in the classification and structural results of II<sub>1</sub> factors (see surveys [Pop07, Vae10, Ioa18]), and specifically, group measure space II<sub>1</sub> factors arising from Bernoulli actions have been shown to possess extreme rigidity (see e.g. [Pop06b, Pop06c, Pop08, Ioa11, IPV13]). In fact, a conjecture of Popa states that if  $\Gamma \curvearrowright (X^{\Gamma}, \mu^{\Gamma})$  is a Bernoulli action of a nonamenable group  $\Gamma$ , then  $L^{\infty}(X^{\Gamma}) \rtimes \Gamma$  has a unique Cartan subalgebra, up to unitary conjugacy [Ioa18, Problem III]. Such unique Cartan subalgebra results play a crucial role in the classification of group measure space II<sub>1</sub> factors to the classification of the orbit equivalence relations of the corresponding group actions [Sin55].

Although this conjecture of Popa remains open in its full generality, significant progress has been made towards it during the last 15 years. To mention a few breakthrough results, Ioana showed that  $L^{\infty}(X^{\Gamma}) \rtimes \Gamma$  has a unique group measure space Cartan subalgebra when  $\Gamma$  has Property (T) [Ioa11]. Subsequently, Ioana, Popa and Vaes showed the same conclusion holds if  $\Gamma$  is a product group [IPV13]. In another direction, Popa and Vaes proved that  $L^{\infty}(Y) \rtimes \Gamma$  has a unique Cartan subalgebra for any free ergodic p.m.p. action  $\Gamma \curvearrowright Y$ , when  $\Gamma$  is weakly amenable and satisfies that either  $\Gamma$  has positive first  $\ell^2$ -Betti number [PV14a] or  $\Gamma$  is biexact [PV14b]. Building upon [PV14a], Ioana showed that the same conclusion holds if  $\Gamma$  is a free product group [Ioa15]. More recently, Boutonnet, Ioana and Peterson generalized [PV14b] to groups that are weakly amenable and properly proximal [BIP21].

In view of [PV14a, PV14b], to make further progress towards this conjecture, one may first consider the question of showing  $L^{\infty}(X^{\Gamma}) \rtimes \Gamma$  has a unique group measure space Cartan subalgebra for weakly amenable  $\Gamma$ , as suggested in [Bou14, Chapter V]. Moreover, in light of [BIP21], it suffices to consider this question for groups that are non properly proximal. Our main theorem is towards this direction.

#### CHANGYING DING

**Theorem 1.1.** Let  $\Gamma$  be a countable discrete nonamenable i.c.c. group,  $(X_0, \mu_0)$  a diffuse standard probability space and  $\Gamma \curvearrowright (X_0^{\Gamma}, \mu_0^{\Gamma}) =: (X, \mu)$  a Bernoulli action. Suppose  $L^{\infty}(X) \rtimes \Gamma \cong L\Lambda$ for some countable discrete group  $\Lambda$ . If  $\Gamma$  is non properly proximal and weakly amenable with Cowling-Haagerup constant 1, then  $\Lambda \cong \Sigma \rtimes \Gamma$  for some infinite abelian group  $\Sigma$  such that  $\Gamma \curvearrowright \Sigma$ by automorphisms, and  $\Gamma \curvearrowright X$ ,  $\Gamma \curvearrowright \widehat{\Sigma}$  are conjugate p.m.p. actions.

Piecing Theorem 1.1 and [BIP21] together, we obtain the following result that does not involve proper proximality. Recall that an algebraic action  $\Lambda \curvearrowright Y$  is a homomorphism from  $\Lambda$  to Aut(Y), with Y a compact metrizable abelian group.

**Theorem 1.2.** Let  $\Gamma$  be a countable discrete *i.c.c.* nonamenable group,  $(X_0, \mu_0)$  a diffuse standard probability space and  $\Gamma \curvearrowright^{\sigma}(X_0^{\Gamma}, \mu_0^{\Gamma}) =: (X, \mu)$  the Bernoulli action. Suppose  $\Gamma$  is weakly amenable with Cowling-Haagerup constant 1. For any group  $\Lambda$  and any free ergodic p.m.p. algebraic action  $\Lambda \curvearrowright Y$ , if  $L^{\infty}(X) \rtimes \Gamma \cong L^{\infty}(Y) \rtimes \Lambda$ , then  $L^{\infty}(X)$  and  $L^{\infty}(Y)$  are unitarily conjugate.

As a side note, we point out that [Bou14, Question V.1.2] can be answered by piecing together the properly proximal case and non properly proximal case, similar to Theorem 1.2, only using existing results in the literature.

**Theorem 1.3.** Let  $\Gamma$  be a nonamenable i.c.c. weakly amenable group and  $\Gamma \curvearrowright X$  a free ergodic p.m.p. action. Let  $\Lambda$  be a nonamenable group and  $\Lambda \curvearrowright [0,1]^{\Lambda}$  a Bernoulli action. If  $L^{\infty}(X) \rtimes \Gamma \cong L^{\infty}([0,1]^{\Lambda}) \rtimes \Lambda$ , then  $L^{\infty}(X)$  and  $L^{\infty}([0,1]^{\Lambda})$  are unitarily conjugate.

Indeed, when  $\Gamma$  is properly proximal, this is a special case of [BIP21, Theorem 1.5]. When  $\Gamma$  is non properly proximal, [Din24, Theorem 1.3] shows that  $\Gamma \curvearrowright X$  and  $\Lambda \curvearrowright [0,1]^{\Lambda}$  are actually conjugate. In both cases, we conclude that  $L^{\infty}(X)$  and  $L^{\infty}([0,1]^{\Lambda})$  are unitarily conjugate.

Turning back to Theorem 1.1, we remark that its proof follows the strategy of [IPV13] closely and exploits the rigidity of non properly proximal groups in the setting of Bernoulli actions by building upon results in [Din24]. We also note that Theorem 1.1 holds when  $\Gamma$  has Property (T) or is a product group as well by [IPV13].

Some concrete examples beyond [IPV13] are covered by Theorem 1.1. Groups considered in [TD20, Example 0.9] are of the form  $(\bigoplus_S \mathbb{Z}_2) \rtimes \mathbb{F}_n$ , where  $n \ge 2$  and S is an infinite set on which  $\mathbb{F}_n$  acts on amenably. They are inner amenable [TD20] and thus non properly proximal [BIP21], and  $\Lambda_{\rm cb}((\bigoplus_S \mathbb{Z}_2) \rtimes \mathbb{F}_n) = \Lambda_{\rm cb}(\mathbb{F}_n) = 1$  by [OP10, Proposition 3.2, Corollary 3.3]. However, [IPV13] does not apply to these since the centralizer of any infinite subgroup of these groups is amenable [TD20].

Before outlining the proof, let us make some remarks on the assumptions of Theorem 1.2. In addition to assuming weak amenability of  $\Gamma$ , we also assume  $\Lambda_{cb}(\Gamma) = 1$  and  $\Lambda \curvearrowright Y$  is an algebraic action. The action  $\Lambda \curvearrowright Y$  being algebraic allows us to realize  $L^{\infty}(Y) \rtimes \Lambda$  as a group von Neumann algebra  $L(\widehat{Y} \rtimes \Lambda)$ , for which we may use the comultiplication map arising from a group instead of a group measure space [PV10]. Using this comultiplication map enables us to analyze the relation between proper proximality and the comultiplication map in Section 5. The reason to assume  $\Lambda_{cb}(\Gamma) = 1$  is present in Section 3. Roughly speaking, in the relative setting, the complete metric approximation property plays a role that is similar to local reflexivity. This assumption is also present in [Iso20, Proposition 7.3] to avoid the same technical difficulty.

Let us finish the introduction by outlining the proof of Theorem 1.1 informally. It is built on the recently developed notions of proper proximality and biexactness for von Neumann algebras [DKEP23, DP23], and Popa's deformation/rigidity theory, in particular, the breakthrough work [Ioa11, IPV13]. In view of [IPV13], to prove Theorem 1.1, it suffices to show  $\Delta(L\Gamma)$  can be unitarily conjugate into  $L\Gamma \otimes L\Gamma$ , where  $\Delta : M := L\Lambda \to L\Lambda \otimes L\Lambda$  is the comultiplication map first considered in [PV10]. First, we use the fact that  $M \otimes M$  is biexact relative to  $\{M \otimes L\Gamma, L\Gamma \otimes M\}$  [BO08, DP23] to conclude that  $\Delta(L\Gamma) \subset M \otimes M$  is properly proximal relative to  $\{M \otimes L\Gamma, L\Gamma \otimes M\}$  [DP23]. In Proposition 4.2, we play this, together with malnormality of  $\Gamma < \mathbb{Z} \wr \Gamma$ , against the non properly proximality of  $\Gamma$  to conclude  $\Delta(L\Gamma)$  can be unitarily conjugate into  $M \otimes L\Gamma$ . Next, although  $M \otimes L\Gamma$  has no relative biexactness, using the machinery from [DP23], we may still obtain certain proper proximality for  $\Delta(L\Gamma) \subset M \otimes L\Gamma$  by Proposition 3.1. This allows us to exploit the non proper proximality of  $\Gamma$  again in Proposition 4.2 and conclude that  $\Delta(L\Gamma)$  may be unitarily conjugate into  $L\Gamma \otimes L\Gamma$ .

Acknowledgment. I would like to thank Daniel Drimbe, Adrian Ioana, Jesse Peterson, Sorin Popa and Stefaan Vaes for their useful comments. I am also very grateful to Jesse Peterson and Sorin Popa for their encouragement.

#### 2. Preliminaries

2.1. The small-at-infinity boundary and boundary pieces. In this section we recall the notion of the small-at-infinity boundary for von Neumann algebras developed in [DKEP23, DP23], which is a noncommutative analogue of the corresponding boundary for groups introduced by Ozawa [Oza04, BO08].

Let M be a finite von Neumann algebra. An M-boundary piece  $\mathbb{X}$  is a hereditary C\*-subalgebra  $\mathbb{X} \subset \mathbb{B}(L^2M)$  such that  $M \cap M(\mathbb{X}) \subset M$  and  $JMJ \cap M(\mathbb{X}) \subset JMJ$  are weakly dense, and  $\mathbb{X} \neq \{0\}$ , where  $M(\mathbb{X})$  denotes the multiplier algebra of  $\mathbb{X}$ . For convenience, we will always assume  $\mathbb{X} \neq \{0\}$ . Given an M-boundary piece  $\mathbb{X}$ , define  $\mathbb{K}^L_{\mathbb{X}}(M) \subset \mathbb{B}(L^2M)$  to be the  $\|\cdot\|_{\infty,2}$  closure of  $\mathbb{B}(L^2M)\mathbb{X}$ , where  $\|T\|_{\infty,2} = \sup_{\hat{a}\in(M)_1} \|T\hat{a}\|$  and  $(M)_1 = \{a \in M \mid \|a\| \leq 1\}$ . Set  $\mathbb{K}_{\mathbb{X}}(M) = \mathbb{K}^L_{\mathbb{X}}(M)^* \cap \mathbb{K}^L_{\mathbb{X}}(M)$ , then  $\mathbb{K}_{\mathbb{X}}(M)$  is a C\*-subalgebra that contains M and JMJ in its multiplier algebra. Put  $\mathbb{K}^{\infty,1}_{\mathbb{X}}(M) = \overline{\mathbb{K}_{\mathbb{X}}(M)^{\|\cdot\|_{\infty,1}} \subset \mathbb{B}(L^2M)$ , where  $\|T\|_{\infty,1} = \sup_{a,b\in(M)_1} \langle T\hat{a}, \hat{b} \rangle$ , and the small-at-infinity boundary for M relative to  $\mathbb{X}$  is given by

$$\mathbb{S}_{\mathbb{X}}(M) = \{ T \in \mathbb{B}(L^2M) \mid [T, x] \in \mathbb{K}_{\mathbb{X}}^{\infty, 1}(M), \text{ for any } x \in M' \}.$$

When  $\mathbb{X} = \mathbb{K}(L^2 M)$ , we omit  $\mathbb{X}$  in the above notations.

The following instance of boundary pieces is extensively used in this paper. Let M be a finite von Neumann algebra and  $\{P_i\}_{i=1}^n$  a family of von Neumann subalgebras. Recall from [DKEP23] that the M-boundary piece  $\mathbb{X}$  associated with  $\{P_i\}_{i=1}^n$  is the hereditary C\*-subalgebra of  $\mathbb{B}(L^2M)$ generated by  $\{xJyJe_{P_i} \mid i = 1, \ldots, n, x, y \in M\}$ . If  $[e_{P_i}, e_{P_j}] = 0$  for  $i, j = 1, \ldots, n$ , we have  $\mathbb{X}$ coincides with the hereditary C\*-subalgebra generated by  $\{xJyJ(\bigvee_{i=1}^n e_{P_i}) \mid x, y \in M\}$  [DKE22, Lemma 3.2]. When the family only contains one von Neumann subalgebra  $P \subset M$ , we usually denote the M-boundary piece by  $\mathbb{X}_P$ .

2.2. Biexactness and proper proximality. With the notion of the small-at-infinity boundary in hand, we recall proper proximality and biexactness for von Neumann algebras, which were introduced in [DKEP23] and [DP23], respectively. These are generalizations of their corresponding notions for groups introduced in [BIP21] and [Oza04, BO08], respectively.

Let M be a von Neumann algebra and  $\mathbb{X}$  an M-boundary piece. Given a von Neumann subalgebra  $N \subset pMp$  with a nonzero projection  $p \in M$ , we say N is not properly proximal relative to  $\mathbb{X}$  in M if there exists an N-central state  $\varphi : p\mathbb{S}_{\mathbb{X}}(M)p \to \mathbb{C}$  such that  $\varphi_{|pMp}$  is normal. Equivalently, N is not properly proximal relative to  $\mathbb{X}$  in M if there exists some nonzero projection  $z \in \mathcal{Z}(N)$  and an Nz-bimodular u.c.p. map  $\phi : z\mathbb{S}_{\mathbb{X}}(M)z \to Nz$  such that  $\phi_{|zMz}$  is normal. When  $\mathbb{X}$  is the

*M*-boundary piece associated with a family of von Neumann subalgebras  $\{P_i\}_{i=1}^n$  of *M*, we say N is not properly proximal relative to  $\{P_i\}_{i=1}^n$  in M.

We say M is biexact relative to X if there exist nets of u.c.p. maps  $\phi_i : M \to \mathbb{M}_{n(i)}(\mathbb{C})$  and  $\psi_i : \mathbb{M}_{n(i)}(\mathbb{C}) \to \mathbb{S}_{\mathbb{X}}(M)$  such that  $\psi_i \circ \phi_i(x) \to x$  in the *M*-topology of  $\mathbb{S}_{\mathbb{X}}(M)$ . Here, by an equivalent characterization [DP23, Lemma 3.4], we say a net  $\{x_i\} \subset \mathbb{S}_{\mathbb{X}}(M)$  converging to 0 in the M-topology if there exists a net of projections  $p_i \in M$  such that  $p_i \to 1$  strongly and  $\|p_i x_i p_i\| \to 0$ . When X is the M-boundary piece associated with a family of von Neumann subalgebras  $\{P_i\}_{i=1}^n$  of M, we say M is biexact relative to  $\{P_i\}_{i=1}^n$ .

These notions coincide with the corresponding notions of groups if we consider group von Neumann algebras: a discrete group  $\Gamma$  is properly proximal (resp. biexact) relative to a family of subgroups  $\{\Lambda_i\}_{i=1}^n$  if and only if  $L\Gamma$  is properly proximal (resp. biexact) relative to  $\{L\Lambda_i\}_{i=1}^n$ [DKEP23, DP23].

2.3. Normal biduals. A bidual characterization of proper proximality will be crucial to our arguments. In this section, we briefly recall necessary notions around normal biduals from [DKEP23, DP23].

Let M be a finite von Neumann algebra and  $A \subset \mathbb{B}(L^2M)$  a C\*-subalgebra such that M and JMJ are contained in its multiplier algebra. Denote by  $A_J^{\sharp}$  the set of functionals  $\varphi \in A^*$  such that for any  $T \in A$ ,

$$M \times M \ni (a, b) \mapsto \varphi(aTb) \in \mathbb{C}, \ JMJ \times JMJ \ni (a, b) \mapsto \varphi(aTb) \in \mathbb{C}$$

are both separately normal.

The normal bidual of A, denoted by  $A_J^{\sharp*}$ , may be identified with a corner of  $A^{**}$  and be viewed as a von Neumann algebra. Denote by  $p_{nor} \in M(A)^{**}$  the support projection of both identity representations of M and JMJ. Equivalently,  $p_{nor}$  is the support projection of states in  $M(A)^*$  that are normal when restricted to M and JMJ. Then  $M(A)_J^{\sharp*}$  may be identified with  $p_{\text{nor}}M(A)^{**}p_{\text{nor}}$ and  $A_J^{\sharp*} = q_A p_{\text{nor}} M(A)^{**} p_{\text{nor}}$ , where  $q_A \in M(A)^{**}$  is the identity of  $A^{**}$  in  $M(A)^{**}$ .

Throughout the paper, we reserve the notation  $p_{\rm nor}$  for the above projection and set  $\iota_{\rm nor}$ :  $\mathbb{B}(L^2M) \ni T \mapsto p_{\mathrm{nor}}\pi_u(T)p_{\mathrm{nor}} \in \mathbb{B}(L^2M)_J^{\sharp*}$ , where  $\pi_u: \mathbb{B}(L^2M) \to \mathbb{B}(L^2M)^{**}$  is the universal representation. Notice that  $\iota_{nor}$  is no longer a \*-homomorphism, but  $\iota_{nor}$  restricts to normal representations on M and JMJ.

We consider the following bidual version of the small-at-infinity boundary for a von Neumann algebra M:

$$\tilde{\mathbb{S}}(M) = |\{T \in \mathbb{B}(L^2 M)_J^{\sharp*} \mid [T, x] \in \mathbb{K}(M)_J^{\sharp*}, \text{ for all } x \in J M J\},\$$

 $\mathbb{S}(M) = |\{I \in \mathbb{B}(L^{-}M)_{J}^{*} | [I, x] \in \mathbb{K}(M)_{J}^{*}, \text{ for all } x \}$ where we view JMJ as in  $\mathbb{B}(L^{2}M)_{J}^{\sharp*}$  through the representation  $\iota_{\text{nor}}$ .

By [DKEP23, Lemma 8.5], we have that if a countable discrete group  $\Gamma$  is non properly proximal, then there exists a  $L\Gamma$ -central state  $\varphi : \tilde{\mathbb{S}}(L\Gamma) \to \mathbb{C}$  such that  $\varphi_{|L\Gamma} = \tau$ .

Next we collect a few lemmas for Section 4.

**Lemma 2.1.** e Let  $\Gamma$  be a countable discrete group with a family of subgroups  $\{\Sigma_i\}_{i=1}^n$ . Denote by  $M = L\Gamma$ ,  $\mathbb{Y}$  the M-boundary piece associated with  $\{L\Sigma_i\}_{i=1}^n$ ,  $P_F$  the orthogonal projection from  $\ell^2 \Gamma$  onto  $\overline{\operatorname{sp}\{\delta_g \mid g \in F(\bigcup_{i=1}^n \Sigma_i)F\}}$  for a finite subset  $F \subset \Gamma$ , and  $q_{\mathbb{Y}}$  the unit in  $(\mathbb{K}_{\mathbb{Y}}(M)_J^{\sharp})^*$ . Then  $q_{\mathbb{Y}} = \lim_{F} \iota_{\mathrm{nor}}(P_F)$ , where the limit is over finite subsets of  $\Gamma$  and is taken in  $(\mathbb{B}(L^2 M)_J^{\sharp})^*$ with the weak<sup>\*</sup> topology.

Proof. Denote by  $\mathbb{Y}_0$  the hereditary C\*-subalgebra generated by  $\{xJyJ(\vee_{i=1}^n e_{L\Sigma_i}) \mid x, y \in C_r^*\Gamma\}$ . One checks  $\{P_F \mid F \subset \Gamma \text{ finite}\}$  forms an approximate unit for  $\mathbb{Y}_0$ . Since  $\iota_{\operatorname{nor}}(\mathbb{Y}_0) \subset (\mathbb{K}_{\mathbb{Y}}(M)_J^{\sharp})^*$  is weak\* dense [DKE22, Lemma 3.7], one has  $\lim_F \iota_{\operatorname{nor}}(P_F) = q_{\mathbb{Y}}$ .

**Lemma 2.2.** Let  $\Gamma$  be a countable discrete group with a subgroup  $\Lambda < \Gamma$  and a family of subgroups  $\{\Sigma_i\}_{i=1}^n$ . Denote by  $M = L\Gamma$  and  $\mathbb{Y}$  the M-boundary piece associated with  $\{L\Sigma_i\}_{i=1}^n$ . Set  $q_{\mathbb{Y}}$  be the unit in  $(\mathbb{K}_{\mathbb{Y}}(M)_J^{\sharp})^*$  and P the orthogonal projection from  $\ell^2\Gamma$  to  $\overline{\operatorname{sp}}\{\delta_t \mid t \in \bigcup_{j=1}^d t_j\Lambda s_j\}$  for a some  $t_j, s_j \in \Gamma$ . Then  $[q_{\mathbb{Y}}, \iota_{\operatorname{nor}}(P)] = 0$ .

Proof. For a subset  $S \subset \Gamma$ , denote by  $P_S$  the orthogonal projection onto  $\overline{\operatorname{sp}\{\delta_t \mid t \in S\}}$ . Note that for any subgroup  $\Gamma_0$ , the projection  $\pi_u(P_{\Gamma_0})$  commutes with  $p_{\operatorname{nor}}$  [Din24, Lemma 3.4]. Since  $P_{t\Gamma_0 s} = \operatorname{Ad}(\lambda_t^* \rho_s)(P_{\Gamma_0})$ , we have  $\pi_u(P_{t\Gamma_0 s})$  commutes with  $p_{\operatorname{nor}}$ . Moreover, for two commuting projections  $Q_1, Q_2 \in \mathbb{B}(L^2 M)^{**}$  such that  $[Q_1, p_{\operatorname{nor}}] = [Q_2, p_{\operatorname{nor}}] = 0$ , one has  $Q_1 \vee Q_2 = Q_1 + Q_2 - Q_1 Q_2$  commutes with  $p_{\operatorname{nor}}$ .

By Lemma 2.1 one has  $q_{\mathbb{Y}} = \lim_{F} \iota_{\operatorname{nor}}(P_F)$ , where  $F \subset \Gamma$  is a finite set and  $P_F \in \mathbb{B}(\ell^2\Gamma)$  is the orthogonal projection onto  $\operatorname{sp}\{\delta_t \mid t \in F(\cup_{i=1}^n \Sigma_i)F\}$ . From the above discussion, we see that  $P_F$  and P are in the multiplicative domain of  $\iota_{\operatorname{nor}}$ . Since  $[P_F, P] = 0$ , we have  $[\iota_{\operatorname{nor}}(P_F), \iota_{\operatorname{nor}}(P)] = 0$  and hence  $[q_{\mathbb{Y}}, \iota_{\operatorname{nor}}(P)] = 0$ .

Given a group  $\Gamma$  with a family of subgroups  $\mathcal{G}$ , recall that a set  $S \subset \Gamma$  is small relative to  $\mathcal{G}$  if there exists some  $n \in \mathbb{N}$ ,  $\{s_i, t_i\}_{i=1}^n \subset \Gamma$  and  $\{\Sigma_i\}_{i=1}^n \subset \mathcal{G}$  such that  $S \subset \bigcup_{i=1}^n s_i \Sigma_i t_i$ , see e.g. [BO08, Chapter 15]. We say a subgroup  $\Lambda < \Gamma$  is almost malnormal relative to  $\mathcal{G}$  if  $s\Lambda s^{-1} \cap \Lambda$ is small relative to  $\mathcal{G}$  for any  $s \in \Gamma \setminus \Lambda$ . One checks that if  $\mathcal{G}$  only contains normal subgroups, then  $\Lambda < \Gamma$  being almost malnormal relative to  $\mathcal{G}$  is equivalent to  $s\Lambda t \cap \Lambda$  is small relative to  $\mathcal{G}$ for any  $s, t \in \Gamma$  with at least one of s, t is not in  $\Lambda$ .

**Lemma 2.3.** Let  $\Gamma$  be a countable discrete group with a subgroup  $\Lambda < \Gamma$  and a family of subgroups  $\{\Sigma_i\}_{i=1}^n$ . Set  $M = L\Gamma$  and denote by  $\mathbb{X}$  the M-boundary piece associated with  $L\Lambda$ , and by  $\mathbb{Y}$  the M-boundary piece associated with  $\{L\Sigma_i\}_{i=1}^n$ .

Take  $\{t_k\}_{k\in\mathbb{N}} \subset \Gamma$  a transversal for  $\Gamma/\Lambda$  and put  $p_{k,\ell} = P_{t_k\Lambda t_\ell} = \operatorname{Ad}(\lambda_{t_k}\rho_{t_\ell})(e_{L\Lambda}) \in \mathbb{B}(L^2M)$ , where  $P_{t_k\Lambda t_\ell}$  denotes the orthogonal projection from  $\ell^2\Gamma$  to  $\operatorname{sp}\{\delta_g \mid g \in t_k\Lambda t_\ell\}$ .

Suppose  $\Lambda$  is almost malnormal relative to  $\{\Sigma_i\}_{i=1}^n$  and each  $\Sigma_i$  is normal. Then  $\{q_{\mathbb{Y}}^{\perp}\iota_{\operatorname{nor}}(p_{k,\ell})\}_{k,\ell\in\mathbb{N}}$  is a family of pairwise orthogonal projections and  $\sum_{k,\ell\in\mathbb{N}}q_{\mathbb{Y}}^{\perp}\iota_{\operatorname{nor}}(p_{k,\ell}) = q_{\mathbb{Y}}^{\perp}q_{\mathbb{X}}$ , where  $q_{\mathbb{X}}$  and  $q_{\mathbb{Y}}$  are identities of  $(\mathbb{K}_{\mathbb{X}}(M)_J^{\sharp})^*$  and  $(\mathbb{K}_{\mathbb{Y}}(M)_J^{\sharp})^*$ , respectively.

Proof. First note that  $p_{k,\ell}p_{k',\ell'} = P_{t_k\Lambda t_\ell\cap t_{k'}\Lambda t_{\ell'}} \in \mathbb{K}_{\mathbb{Y}}(M)$  if  $k \neq k'$  or  $\ell \neq \ell'$ , since  $t_{k'}^{-1}t_k\Lambda t_\ell t_{\ell'}^{-1}\cap \Lambda$  is small relative to  $\{\Sigma_i\}_{i=1}^n$ . As  $\{p_{k,\ell}\}$  is in the multiplicative domain of  $\iota_{\mathrm{nor}}$  and  $[q_{\mathbb{Y}}, \iota_{\mathrm{nor}}(p_{k,\ell})] = 0$  by Lemma 2.2, we have  $q_{\mathbb{Y}}^{\perp}\iota_{\mathrm{nor}}(p_{k,\ell})q_{\mathbb{Y}}^{\perp}\iota_{\mathrm{nor}}(p_{k',\ell'}) = \delta_{k,k'}\delta_{\ell,\ell'}q_{\mathbb{Y}}^{\perp}\iota_{\mathrm{nor}}(p_{k,\ell})$ .

To see  $\sum_{k,\ell\in\mathbb{N}} q_{\mathbb{Y}}^{\perp} \iota_{\operatorname{nor}}(p_{k,\ell}) = q_{\mathbb{Y}}^{\perp} q_{\mathbb{X}}$ , we first notice that  $\{p_n := \bigvee_{k,\ell\leq n} p_{k,\ell}\}_{n\in\mathbb{N}}$  is an approximate unit for the hereditary C\*-subalgebra generated by  $\{xJyJe_{L\Lambda} \mid x, y \in C_r^*\Gamma\}$ . Thus by the same argument as in Lemma 2.1, we have  $q_{\mathbb{X}} = \lim_{n} \iota_{\operatorname{nor}}(p_n)$  and hence

$$q_{\mathbb{Y}}^{\perp}q_{\mathbb{X}} = \lim_{n} q_{\mathbb{Y}}^{\perp}\iota_{\operatorname{nor}}(\vee_{k,\ell \leq n}p_{k,\ell}) = \sum_{k,\ell \in \mathbb{N}} q_{\mathbb{Y}}^{\perp}\iota_{\operatorname{nor}}(p_{k,\ell}).$$

#### 2.4. Popa's intertwining-by-bimodules.

**Theorem 2.4** ([Pop06b]). Let  $(M, \tau)$  be a tracial von Neumann algebra and  $P \subset pMp, Q \subset M$ be von Neumann subalgebras. Then the following are equivalent:

- (1) There exist projections  $p_0 \in P, q_0 \in Q, a *-homomorphism \theta : p_0 P p_0 \rightarrow q_0 Q q_0$  and a non-zero partial isometry  $v \in q_0 M p_0$  such that  $\theta(x)v = vx$ , for all  $x \in p_0 P p_0$ .
- (2) There is no sequence  $u_n \in \mathcal{U}(P)$  satisfying  $||E_Q(x^*u_ny)||_2 \to 0$ , for all  $x, y \in pMp$ .

If one of these equivalent conditions holds, we write  $P \prec_M Q$ .

2.5. Relative amenability. Given a finite von Neumann algebra M with a von Neumann subalgebra  $Q \subset M$ . Recall from [OP10] that a von Neumann subalgebra  $P \subset pMp$  with  $p \in \mathcal{P}(M)$  is amenable relative to Q in M if there exists a u.c.p. map  $\phi: p\langle M, e_Q \rangle p \to P$  such that  $\phi|_{pMp} = E_P$ , where  $E_P: pMp \to P$  is the normal conditional expectation. Following [IPV13], we say P is strongly nonamenable relative to Q in M if for any nonzero projection  $p' \in P' \cap pMp$ , we have Pp' is not amenable relative to Q in M.

The following is an abstraction of [Ioa15, Corollary 2.12] (also a straightforward relativization of [Din24, Lemma 4.1]), and thus we omit the proof.

**Lemma 2.5.** Let  $(M, \tau)$  and N be tracial von Neumann algebras and  $Q \subset M$  a von Neumann subalgebra. Suppose there exist another tracial von Neumann algebra  $(M, \tilde{\tau})$  such that  $M \subset M$ and  $\tilde{\tau}_{|M} = \tau$ , and a net of trace preserving automorphisms  $\{\alpha_t\}_{t \in \mathbb{R}} \subset \operatorname{Aut}(M)$  such that  $\alpha_{t|Q} \in \mathbb{R}$ Aut(Q), and such that  $\alpha_{t|M} \to \mathrm{id}_M$  in the point- $\|\cdot\|_2$  topology, as  $t \to 0$ . Set  $\tilde{\alpha}_t = \alpha_t \otimes \mathrm{id}_N \in$  $\operatorname{Aut}(M \otimes N).$ 

If a von Neumann subalgebra  $P \subset p(M \otimes N)p$  is amenable relative to  $Q \otimes N$  in  $M \otimes N$ , where  $p \in \mathcal{P}(M \otimes N)$  is some nonzero projection, then for any  $0 < \delta \leq 1$ , one of the following is true.

- (1) There exists  $t_{\delta} > 0$  such that  $\inf_{u \in \mathcal{U}(P)} \|E_{M \otimes \mathcal{Q}}(\tilde{\alpha}_{t_{\delta}}(u))\|_{2} > (1-\delta) \|p\|_{2}$ .
- (2) There exists a net  $\{\eta_k\} \subset \mathcal{K}^{\perp}$ , where  $\mathcal{K}$  is the closure of  $(M \otimes N)e_{Q \otimes N}(\tilde{M} \otimes N)$  inside  $L^2(\langle \tilde{M} \otimes N, e_{Q \otimes N} \rangle)$ , such that  $\|x\eta_k - \eta_k x\|_2 \to 0$  for all  $x \in P$ ,  $\limsup_k \|y\eta_k\|_2 \le 2\|y\|_2$ for all  $y \in p(M \otimes N)p$  and  $\limsup_k \|p\eta_k\|_2 > 0$ .

Specializing to the situation of Bernoulli actions, we obtain the following result.

**Proposition 2.6.** Let  $\Gamma$  be a nonamenable group and denote by  $M = L^{\infty}(X^{\Gamma}) \rtimes \Gamma$  the von Neumann algebra associated with its Bernoulli action. Let N be a finite von Neumann algebra and  $P \subset p(M \otimes N)p$  a von Neumann subalgebra with some nonzero projection  $p \in M \otimes N$ . Suppose P is amenable relative to  $L\Gamma \overline{\otimes} N$ , and P has no direct summand that is amenable relative to  $1 \otimes N$ , then P is rigid relative to deformation deformation associated with the Bernoulli action.

Moreover, if  $\Gamma$  is i.c.c. and N is a factor, then one of the following is true.

- (1) There exists some partial isometry  $v \in M \overline{\otimes} N$  such that  $v^*(\mathcal{N}_{p(M \overline{\otimes} N)p}(P)'')v \subset L\Gamma \overline{\otimes} N$ and  $vv^* = p$ . (2)  $\mathcal{N}_{p(M\overline{\otimes}N)p}(P)'' \prec_{M\overline{\otimes}N} L^{\infty}(X^{\Gamma})\overline{\otimes}N$ .

*Proof.* Denote by  $\tilde{M}$  and  $\{\alpha_t\} \subset \operatorname{Aut}(\tilde{M})$  from [Pop06b]. It suffices to show that (2) of Lemma 2.5 does not occur by Popa's transversality lemma [Pop08, Lemma 2.1]. In fact, note that as  $M \otimes N$ bimodules we have

$$L^{2}(\langle \tilde{M} \,\overline{\otimes}\, N, e_{L\Gamma \,\overline{\otimes}\, N} \rangle) \ominus \mathcal{K} = (L^{2}(\tilde{M} \ominus M) \otimes L^{2}N) \otimes_{L\Gamma \,\overline{\otimes}\, N} (L^{2}\tilde{M} \otimes L^{2}N) \prec L^{2}M \otimes L^{2}N \otimes L^{2}M,$$

since  $L^2(\tilde{M} \ominus M)$  is a weakly coarse *M*-bimodule. If (2) were the case, then one would have almost *P*-central vectors in  $L^2M \otimes L^2N \otimes L^2M$ , which implies that *P* has a direct summand is amenable relative to  $1 \otimes N$  in  $M \otimes N$ .

The moreover part follows from the same argument as in the beginning of the proof of [IPV13, Corollary 4.3] by noticing that  $P \not\prec_{M \overline{\otimes} N} 1 \otimes N$  since P has no direct summand amenable relative to  $1 \otimes N$  in  $M \overline{\otimes} N$  [Ioa15, Remark 2.2].

2.6. Weak amenability and W\*CMAP. Recall from [CH89] that a discrete group  $\Gamma$  is weakly amenable if there exists a net of finitely supported functions  $\varphi_i : \Gamma \to \mathbb{C}$  such that  $\varphi_i \to 1$ pointless,  $m_{\varphi_i}(u_g) = \varphi_i(g)u_g$  extends to a c.b. map on  $L\Gamma$  and there exists some C > 0 satisfying  $\sup_i ||m_{\varphi_i}||_{cb} \leq C$ . The Cowling-Haagerup constant  $\Lambda_{cb}(\Gamma)$  is the infimum of all C for which such a net  $\varphi_i$  exists. Similarly, a von Neumann algebra M has the W\* complete metric approximation property (W\*CMAP) if there exists a net of normal finite rank c.c.  $\theta_i : M \to M$  such that  $\theta_i \to id_M$  in the point-weak\* topology. Given a discrete group  $\Gamma$ , one has  $L\Gamma$  has W\*CMAP if and only if  $\Lambda_{cb}(\Gamma) = 1$ .

#### 3. BIEXACTNESS AND A DICHOTOMY OF SUBALGEBRAS

In this section, we obtain a dichotomy of von Neumann subalgebras of  $M \otimes N$ , where M is assumed to be biexact relative to some von Neumann subalgebra. This can be seen as a relative version of [DKEP23, Theorem 7.1] and [Din24, Proposition 2.3], where  $N = \mathbb{C}$ . Since the arguments we employ here are somewhat C\*-algebraic, we in addition assume N has W\*CMAP. This extra technical condition is used in the same way as in [Iso20, Proposition 7.3], and is similar to how exactness is used in [Oza04]. Specifically, we prove the following.

**Proposition 3.1.** Let M, N be finite von Neumann algebras and  $Q \subset M$  a von Neumann subalgebra. Suppose M is biexact relative to Q and N has  $W^*CMAP$ . Then for any von Neumann subalgebra  $P \subset p(M \otimes N)p$  for some nonzero projection  $p \in M \otimes N$ , we have P is either properly proximal relative to  $Q \otimes N$  in  $M \otimes N$ , or there exists some nonzero projection  $z \in \mathcal{Z}(P)$  such that Pz is amenable relative to N in  $M \otimes N$ .

We first consider the situation that N is without W\*CMAP.

**Lemma 3.2.** Let M, N be finite von Neumann algebras and  $Q \subset M$  a von Neumann subalgebra. Suppose M is biexact relative to Q. Then for any von Neumann subalgebra  $P \subset p(M \otimes N)p$  with  $0 \neq p \in \mathcal{P}(M \otimes N)$  that is not properly proximal relative to  $Q \otimes N$  in  $M \otimes N$ , there exist a nonzero central projection  $z \in \mathcal{Z}(P)$  and a u.c.p. map  $\phi \in \mathbb{B}(L^2M) \otimes N \to Pz$  such that  $\phi_{|M \otimes_{\min} N} = \psi$ , where  $\psi : M \otimes N \to Pz$  is a normal u.c.p. map with  $\psi_{|Pz} = \text{id}$ .

*Proof.* We first check that  $\mathbb{S}_{\mathbb{X}_Q}(M) \otimes_{\min} N \subset \mathbb{S}_{\mathbb{X}_{Q\overline{\otimes}N}}(M\overline{\otimes}N)$ , for which it suffices to verify that  $\mathbb{K}_{\mathbb{X}_Q}(M)^{\infty,1} \otimes_{\mathrm{alg}} N \subset \mathbb{K}_{\mathbb{X}_{Q\overline{\otimes}N}}^{\infty,1}(M\overline{\otimes}N)$ . Indeed, for any  $a, b, c, d \in M, T \in \mathbb{B}(L^2M)$  and  $x \in N$ , note that

 $aJbJe_{Q}Te_{Q}cJdJ\otimes x=(a\otimes 1)J(b\otimes 1)Je_{Q\,\overline{\otimes}\,N}(T\otimes 1)e_{Q\,\overline{\otimes}\,N}(c\otimes x)J(d\otimes 1)J\in \mathbb{K}_{\mathbb{X}_{Q\,\overline{\otimes}\,N}}(M\,\overline{\otimes}\,N).$ 

Since  $\mathbb{B}(L^2M) \ni T \mapsto T \otimes 1 \in \mathbb{B}(L^2M \otimes L^2N)$  is continuous from both *M*-topology to  $M \otimes N$ -topology and *JMJ*-topology to  $J(M \otimes N)J$ -topology, we have  $\mathbb{K}^{\infty,1}_{\mathbb{X}_Q}(M) \otimes_{\mathrm{alg}} N \subset \mathbb{K}^{\infty,1}_{\mathbb{X}_Q \otimes N}(M \otimes N)$ .

Denote by  $\theta_i : \mathbb{B}(L^2M) \to \mathbb{S}_{\mathbb{X}_Q}(M)$  a net of normal u.c.p. maps such that  $\theta_{i|M} \to \mathrm{id}_M$  in the point *M*-topology, which is given by the assumption that *M* is biexact relative to *Q*. Consider

 $\tilde{\theta_i} := \theta_i \otimes \operatorname{id}_N : \mathbb{B}(L^2 M) \,\overline{\otimes}\, N \to \mathbb{S}_{\mathbb{X}_Q}(M) \otimes_{\min} N \subset \mathbb{S}_{\mathbb{X}_Q \,\overline{\otimes}\, N}(M \,\overline{\otimes}\, N).$ 

Since  $P \subset p(M \otimes N)p$  is not properly proximal relative to  $Q \otimes N$ , there exists a nonzero central projection  $z \in P$  and a P-bimodular u.c.p. map  $\psi : \mathbb{S}_{\mathbb{X}_{Q\overline{\otimes}N}}(M\overline{\otimes}N) \to Pz$  such that  $\psi_{|M\overline{\otimes}N}$  is normal. Denote by  $\phi$  a weak<sup>\*</sup> limit point of  $\psi \circ \tilde{\theta}_i : \mathbb{B}(L^2 M) \otimes N \to Pz$ .

Now we show  $\phi_{|M\otimes_{\min}N} = \psi$ . First note that for any  $x \in M \otimes_{\min} N$ , we have  $\tilde{\theta}_i(x) \to x$  in the  $M \otimes N$ -topology. To see this, one checks that  $(\theta_i(a) - a) \otimes b \to 0$  in the  $M \otimes N$ -topology for any  $a \in M$  and  $b \in N$ . Moreover,  $\psi$  is continuous from the weak  $M \otimes N$ -topology to the weak operator topology on Pz as  $\psi$  is u.c.p. and  $\psi_{|M \otimes N}$  is normal. Altogether, for any  $x \in M \otimes_{\min} N$ and any  $\omega \in (Pz)_*$ , we have

$$\lim_{i} \langle \psi(\tilde{\theta}_{i}(x)), \omega \rangle = \lim_{i} \langle \tilde{\theta}_{i}(x), \omega \circ \psi \rangle = \langle x, \omega \circ \psi \rangle,$$
  
i.e.,  $\phi(x) = \lim_{i} \psi(\tilde{\theta}_{i}(x)) = \psi(x).$ 

Proof of Proposition 3.1. Since N has  $W^*CMAP$ , there exists a net of finite rank normal c.c. maps  $\phi_n : N \to N$  such that  $\phi_n \to \mathrm{id}_N$  in point-weak<sup>\*</sup> and

$$\tilde{\phi}_n := \mathrm{id} \otimes \phi_n : \mathbb{B}(L^2 M) \overline{\otimes} N \to \mathbb{B}(L^2 M) \otimes_{\min} N.$$

Consider  $\phi \circ \tilde{\phi}_n : \mathbb{B}(L^2 M) \overline{\otimes} N \to Pz$ , where  $\phi : \mathbb{B}(L^2 M) \overline{\otimes} N \to Pz$  is from Lemma 3.2, denote by  $\Phi$  a point weak<sup>\*</sup> limit of  $\phi \circ \tilde{\phi}_n$ . For any  $x \in M \otimes N$  and  $\varphi \in (Pz)_*$ , since  $\tilde{\phi}_n$  maps  $M \otimes N$  to  $M \otimes_{\min} N$  and  $\phi_n \to \operatorname{id}_{M \overline{\otimes} N}$  in the point weak<sup>\*</sup> topology, we have

$$\langle \Phi(x), \varphi \rangle = \lim_{n} \langle \phi(\tilde{\phi}_n(x)), \varphi \rangle = \lim_{n} \langle \tilde{\phi}_n(x), \varphi \circ \psi \rangle = \varphi(\psi(x)),$$

i.e.,  $\Phi_{|M \otimes N} = \psi$ . Therefore  $\Phi : \mathbb{B}(L^2 M) \otimes N \to Pz$  is a conditional expectation with  $\Phi_{|M \otimes N}$ normal, which shows Pz is amenable relative to N in  $M \otimes N$  by [OP10].  $\square$ 

Let M be a finite von Neumann algebra and  $N \subset M$  a von Neumann subalgebra. Recall that M is solid relative to N if for any von Neumann subalgebra  $P \subset M$  such that  $P \not\prec_M N$ , one has  $P' \cap M$  is amenable relative to N in M. The following recovers [Iso20, Proposition 7.3].

**Corollary 3.3.** Let M and N be finite von Neumann algebras. If M is biexact relative to a von Neumann subalgebra  $Q \subset M$  and N has  $W^*CMAP$ , then  $M \otimes N$  is solid relative to  $Q \otimes N$ .

*Proof.* Suppose  $P \subset M \overline{\otimes} N$  is a von Neumann subalgebra such that  $P \not\prec_{M \overline{\otimes} N} Q \overline{\otimes} N$ , from which we obtain a sequence of unitary  $\{u_n\} \subset \mathcal{U}(P)$  with  $||E_{Q\overline{\otimes}N}(xu_ny)||_2 \to 0$  for any  $x, y \in M\overline{\otimes}N$ . As in [DKEP23], a point weak<sup>\*</sup> limit point  $\phi := \lim_{n \to 0} \operatorname{Ad}(u_n) : \mathbb{S}_{\mathbb{X}_{Q\overline{\otimes}N}}(M\overline{\otimes}N) \to P' \cap (M\overline{\otimes}N)$ gives a conditional expectation. Then the proof of Proposition 3.1 shows that  $P' \cap (M \otimes N)$  is amenable relative to N in  $M \otimes N$  and hence amenable relative to  $Q \otimes N$  in  $M \otimes N$ .  $\square$ 

#### 4. Concentration of states on the boundary and relative amenability

In this section, we show a relative version of the main technical result in [Din24] and [DKE22]. The following lemma shows that under certain conditions, one may relate the basic construction and the small-at-infinity boundary.

**Lemma 4.1.** Let  $\Gamma$  be a countable discrete group with subgroups  $\Lambda$  and  $\Sigma_i$ ,  $1 \leq i \leq n$ . Suppose  $\Lambda < \Gamma$  is almost malnormal relative to the family  $\{\Sigma_i\}_{i=1}^n$  and each  $\Sigma_i$  is normal. Denote by  $\mathbb{X}$  the  $L\Gamma$ -boundary piece associated with  $L\Lambda$  and  $\mathbb{Y}$  the  $L\Gamma$ -boundary piece associated with  $\{L\Sigma_i\}_{i=1}^n$ . Then there exists a u.c.p. map  $\phi : \langle L\Gamma, e_{L\Lambda} \rangle \to q_{\mathbb{Y}}^{\perp} \tilde{\mathbb{S}}_{\mathbb{Y}}(L\Gamma) q_{\mathbb{Y}}^{\perp}$  such that  $\phi(x) = q_{\mathbb{Y}}^{\perp} q_{\mathbb{X}} \iota_{\mathrm{nor}}(x)$  for any  $x \in L\Gamma$ , where  $q_{\mathbb{X}}$  and  $q_{\mathbb{Y}}$  denote the identity of  $\mathbb{K}_{\mathbb{X}}(L\Gamma)_{I}^{\sharp*}$  and  $\mathbb{K}_{\mathbb{Y}}(L\Gamma)_{I}^{\sharp*}$ , respectively.

*Proof.* We follow the proof in [Din24] closely. Denote by  $\{t_k\}_{k\geq 1} \subset \Gamma$  a transversal of  $\Gamma/\Lambda$ . Set  $M = L\Gamma$ ,  $N = L\Lambda$  and  $u_k = \lambda_{t_k} \in \mathcal{U}(M)$ .

For each  $n \ge 1$ , consider the c.p. map  $\psi_n : \langle M, e_N \rangle \to \langle M, e_N \rangle$  given by

$$\psi_n(x) = (\sum_{k \le n} u_k e_N u_k^*) x (\sum_{j \le n} u_j e_N u_j^*).$$

Notice that the image of  $\psi_n$  lies in the \*-algebra  $A_0 := \sup\{u_k a e_N u_i^* \mid a \in N, j, k \ge 1\}$ .

By [Din24, Lemma 3.4], we have  $\{\iota_{nor}(Ju_k Je_N Ju_k^*J)\}_{k\geq 1} \subset \mathbb{B}(L^2 M)_J^{\sharp*}$  is a family of pairwise orthogonal projections and let  $e = \sum_{k\geq 1} \iota_{nor}(Ju_k Je_N Ju_k^*J)$ . Set  $\pi: A_0 \to q_{\mathbb{Y}}^{\perp} \mathbb{B}(L^2 M)_J^{\sharp*} q_{\mathbb{Y}}^{\perp}$  to be the linear map satisfying  $\pi(u_k ae_N u_j^*) = q_{\mathbb{Y}}^{\perp} \iota_{nor}(u_k a) e\iota_{nor}(u_j^*) q_{\mathbb{Y}}^{\perp}$ .

We first show that  $\pi$  is a \*-homomorphism. Indeed, we claim that for any  $x \in M$ , we have

(1) 
$$q_{\mathbb{Y}}^{\perp} e \iota_{\mathrm{nor}}(x) e q_{\mathbb{Y}}^{\perp} = q_{\mathbb{Y}}^{\perp} \iota_{\mathrm{nor}}(E_N(x)) e q_{\mathbb{Y}}^{\perp}.$$

By normality of  $\iota_{nor}$  and  $E_N$ , it suffices to check this for  $x \in \mathbb{C}\Gamma$ . Compute

$$\begin{aligned} q_{\mathbb{Y}}^{\perp} e\iota_{\mathrm{nor}}(x) eq_{\mathbb{Y}}^{\perp} &= \mathrm{Ad}(q_{\mathbb{Y}}^{\perp}) (\sum_{k,j} \iota_{\mathrm{nor}}((Ju_{k}Je_{N}Ju_{k}^{*}J)x(Ju_{j}Je_{N}Ju_{j}^{*}J)) \\ &= \mathrm{Ad}(q_{\mathbb{Y}}^{\perp})(\iota_{\mathrm{nor}}(E_{N}(x))e + \mathrm{Ad}(q_{\mathbb{Y}}^{\perp})(\sum_{k\neq j} \iota_{\mathrm{nor}}((Ju_{k}Je_{N}Ju_{k}^{*}J)x(Ju_{j}Je_{N}Ju_{j}^{*}J))). \end{aligned}$$

Note that for  $x = \lambda_t \in \mathbb{C}\Gamma$ , one has  $(Ju_k Je_N Ju_k^* J)x(Ju_j Je_N Ju_j^* J) = \lambda_t P_{t^{-1}\Lambda t_k \cap \Lambda t_j}$  which is in  $\mathbb{K}_{\mathbb{Y}}(M)$  if  $k \neq j$  since  $\Lambda$  is almost malnormal relative to  $\{\Sigma_i\}_{i=1}^n$  and hence the (1) follows. Here, for a subset  $S \subset \Gamma$ ,  $P_S$  denotes the orthogonal projection from  $\ell^2\Gamma$  to  $\overline{sp}\{\delta_g \in g \in S\}$ .

Moreover, we have that  $[e, q_{\mathbb{Y}}] = 0$ . To see this, recall from Lemma 2.1 that  $q_{\mathbb{Y}} = \lim_{F} \iota_{nor}(e_F)$ , where  $\{e_F\} \in \mathbb{K}_{\mathbb{Y}}(M)$  is an approximate unit of the form that  $e_F$  is the orthogonal projection on to  $\operatorname{sp}\{\delta_g \mid g \in F(\bigcup_{i=1}^n \Sigma_i)F\}$  for any finite subset  $F \subset \Gamma$ . Since  $Ju_k Je_N Ju_k^* J$  is the projection on to  $\operatorname{sp}\{\delta_g \mid g \in \Lambda t_k\}$ , we have  $[\iota_{nor}(Ju_k Je_N Ju_k^* J), q_{\mathbb{Y}}] = 0$  by Lemma 2.2 and hence  $[e, q_{\mathbb{Y}}] = 0$  as well. Together with (1), it then follows that  $\pi$  is a homomorphism.

We then show  $\pi$  is a contraction and thus  $\pi$  extends to the C\*-algebra  $A := \overline{A_0}^{\|\cdot\|}$ . To this end, it suffices to show that for any  $\sum_{i=1}^{d} u_{k_i} a_i e_N u_{j_i}^* \in A_0$  with  $a_i \in N$  and  $k_i, j_i \geq 1$ , and unit vectors  $\xi, \eta \in \mathcal{H}$  we have  $|\langle \pi(\sum_{i=1}^{d} u_{k_i} a_i e_N u_{j_i}^*) \xi, \eta \rangle| \leq ||\sum_{i=1}^{d} u_{k_i} a_i e_N u_{j_i}^*||$ , where  $\mathcal{H}$  is the Hilbert space on which  $\mathbb{B}(L^2 M)_I^{\sharp*}$  is represented.

For each  $k \in \mathbb{N}$ , set  $Q_k = q_{\mathbb{Y}}^{\perp} \sum_{i=1}^{d} \iota_{\operatorname{nor}}(P_{t_{j_i}\Lambda t_k^{-1}})$  and  $R_k = q_{\mathbb{Y}}^{\perp} \sum_{i=1}^{d} \iota_{\operatorname{nor}}(P_{t_{k_i}\Lambda t_k^{-1}})$ . Note that  $\{Q_k\}$  and  $\{R_k\}$  are two families of pairwise orthogonal projections as for  $k \neq r$  we have  $Q_kQ_r = q_{\mathbb{Y}}^{\perp} \sum_{i,j=1}^{d} \iota_{\operatorname{nor}}(P_{t_{j_i}\Lambda t_k^{-1}\cap t_{j_j}\Lambda t_r^{-1}}) = 0$  and similarly  $R_kR_r = 0$ . Moreover, observe that

$$\sum_{i=1}^{d} \iota_{\text{nor}}(e_{N}u_{j_{i}}^{*}Ju_{k}^{*}J)Q_{k} = q_{\mathbb{Y}}^{\perp} \sum_{i=1}^{d} \iota_{\text{nor}}(e_{N}u_{j_{i}}^{*}Ju_{k}J), \ \sum_{i=1}^{d} \iota_{\text{nor}}(e_{N}u_{k_{i}}^{*}Ju_{k}^{*}J)R_{k} = q_{\mathbb{Y}}^{\perp} \sum_{i=1}^{d} \iota_{\text{nor}}(e_{N}u_{k_{i}}^{*}Ju_{k}^{*}J)$$

Thus we compute

$$\begin{aligned} |\langle \pi (\sum_{i=1}^{d} u_{k_{i}} a_{i} e_{N} u_{j_{i}}^{*}) \xi, \eta \rangle | &\leq \sum_{k \geq 0} |\sum_{i=1}^{d} \langle q_{\mathbb{Y}}^{\perp} \iota_{\operatorname{nor}}(e_{N} u_{j_{i}}^{*} J u_{k}^{*} J) \xi, q_{\mathbb{Y}}^{\perp} \iota_{\operatorname{nor}}(J u_{k} J u_{k_{i}} e_{N} a_{i})^{*} \eta \rangle | \\ &= \sum_{k \geq 0} |\sum_{i=1}^{d} \langle q_{\mathbb{Y}}^{\perp} \iota_{\operatorname{nor}}(e_{N} u_{j_{i}}^{*} J u_{k}^{*} J) Q_{k} \xi, q_{\mathbb{Y}}^{\perp} \iota_{\operatorname{nor}}(J u_{k} J u_{k_{i}} e_{N} a_{i})^{*} R_{k} \eta \rangle | \\ &\leq \sum_{k \geq 0} ||\iota_{\operatorname{nor}}(J u_{k} J (\sum_{i=1}^{d} u_{k_{i}} a_{i} e_{N} u_{j_{i}}^{*}) J u_{k}^{*} J)|||Q_{k} \xi|||R_{k} \eta|| \\ &\leq ||\sum_{i=1}^{d} u_{k_{i}} a_{i} e_{N} u_{j_{i}}^{*}||. \end{aligned}$$

Next we show  $[\pi(A_0), \iota_{nor}(x)] = 0$  for any  $x \in JMJ$ , and thus

$$\psi_n \circ \pi : \langle M, e_N \rangle \to q_{\mathbb{Y}}^{\perp} \tilde{\mathbb{S}}_{\mathbb{Y}}(M) q_{\mathbb{Y}}^{\perp}.$$

For this, it suffices to check that  $\iota_{nor}(Ju_s^*J)e\iota_{nor}(Ju_sJ) = e$  for any  $s \in \Gamma$ . Indeed, observe that  $e = \sum_{k\geq 1} \iota_{nor}(Ju_kJe_NJu_k^*J)$  is independent of the choice of traversals of  $\Gamma/\Lambda$  and  $\Gamma = \bigcup_{k\in\mathbb{N}} t_k\Lambda = \bigcup_{k\in\mathbb{N}} st_k\Lambda$ .

Finally, consider the sub-unital c.p. map  $\phi_n := \psi_n \circ \pi : \langle M, e_N \rangle \to q_{\mathbb{Y}}^{\perp} \tilde{\mathbb{S}}_{\mathbb{Y}}(M) q_{\mathbb{Y}}^{\perp}$ , and denote by  $\phi$  a weak<sup>\*</sup> limit point of  $\phi_n$ . We claim that  $\phi(x) = q_{\mathbb{Y}}^{\perp} q_{\mathbb{X}} \iota_{\operatorname{nor}}(x)$  for any  $x \in M$ .

In fact, for any  $x \in M$ , we have

$$\begin{split} \phi(x) &= \lim_{n \to \infty} \pi \Big( \sum_{0 \le k, \ell \le n} (u_k E_N(u_k^* x u_\ell) e_N u_\ell^*) \Big) = q_{\mathbb{Y}}^\perp \lim_{n \to \infty} \sum_{0 \le k, \ell \le n} \iota_{\operatorname{nor}}(u_k E_N(u_k^* x u_\ell)) e_{\iota_{\operatorname{nor}}}(u_\ell^*) \\ &= q_{\mathbb{Y}}^\perp \lim_{n \to \infty} \sum_{0 \le k, \ell \le n} \big( \iota_{\operatorname{nor}}(u_k) e_{\iota_{\operatorname{nor}}}(u_k^*) \big) \iota_{\operatorname{nor}}(x) \big( \iota_{\operatorname{nor}}(u_\ell) e_{\iota_{\operatorname{nor}}}(u_\ell^*) \big), \end{split}$$

where the last equation follows from (1). Finally, note that by Lemma 2.3,  $\{p_k\}_{k\geq 0}$  is a family of pairwise orthogonal projections, where

$$p_k := q_{\mathbb{Y}}^{\perp} \iota_{\operatorname{nor}}(u_k) e \iota_{\operatorname{nor}}(u_k^*) = q_{\mathbb{Y}}^{\perp} \sum_{r \ge 0} \iota_{\operatorname{nor}}(J u_r J u_k e_N u_k^* J u_r^* J),$$

and  $\sum_{k\geq 0} p_k = \sum_{k,r\geq 0} q_{\mathbb{Y}}^{\perp} \iota_{\text{nor}}(Ju_r Ju_k e_B u_k^* Ju_r^* J) = q_{\mathbb{Y}}^{\perp} q_{\mathbb{X}}$ . Therefore, we conclude that  $\phi(x) = q_{\mathbb{Y}}^{\perp} q_{\mathbb{X}} \iota_{\text{nor}}(x)$ , as desired.

**Proposition 4.2.** Let  $\Gamma$  be a countable discrete group with two families of subgroups  $\{\Lambda_i\}_{i=1}^n$  and  $\{\Sigma_i\}_{i=1}^n$  such that each  $\Lambda_i$  is almost malnormal relative to  $\{\Sigma_j\}_{j=1}^n$  and and each  $\Sigma_i$  is normal. Set  $M = L\Gamma$  and denote by  $\mathbb{X}$  and  $\mathbb{Y}$  the M-boundary pieces associated with  $\{L\Lambda_i\}_{i=1}^n$  and  $\{L\Sigma_i\}_{i=1}^n$ , respectively.

Suppose  $N \subset pMp$ , for some nonzero  $p \in \mathcal{P}(M)$ , is a von Neumann subalgebra that has no amenable direct summand, such that N is properly proximal relative to  $\{L\Lambda_i\}_{i=1}^n$  in M.

If there exists an N-central state  $\varphi : \tilde{\mathbb{S}}_{\mathbb{Y}}(M) \to \mathbb{C}$  such that  $\varphi_{|pMp}$  is a faithful normal state, then there exists a partition of unity  $\{p_i\}_{i=1}^n \subset \mathcal{P}(\mathcal{Z}(N' \cap pMp))$  such that  $Np_i$  is amenable relative to  $L\Lambda_i$  in M for each  $1 \leq i \leq n$ . *Proof.* Denote by  $q_{\mathbb{X}}$  and  $q_{\mathbb{Y}}$  the identities of  $\mathbb{K}_{\mathbb{X}}(M)_{J}^{\sharp*}$  and  $\mathbb{K}_{\mathbb{Y}}(M)_{J}^{\sharp*}$ , respectively.

Note that  $\varphi(q_{\mathbb{Y}}) = 0$ . Indeed, if this is not the case, we may then consider the *M*-bimodular u.c.p. map

$$\mathbb{B}(L^2M) \ni T \mapsto q_{\mathbb{Y}}\iota_{\mathrm{nor}}(T)q_{\mathbb{Y}} \in q_{\mathbb{Y}}\mathbb{B}(L^2M)_I^{\mathfrak{p}*}q_{\mathbb{Y}}.$$

Since  $\mathbb{K}_{\mathbb{Y}}(M) \subset \mathbb{B}(L^2M)$  is a hereditary C\*-algebra with M contained in its multiplier algebra, we see that  $q_{\mathbb{Y}} \mathbb{B}(L^2M)_J^{\sharp*} q_{\mathbb{Y}} = \mathbb{K}_{\mathbb{Y}}(M)_J^{\sharp*}$  and  $[q_{\mathbb{Y}}, M] = 0$ . The non-vanishing of  $\varphi$  on  $q_{\mathbb{Y}}$  then yields an N-central state on  $\mathbb{B}(L^2M)$  which entails that N has an amenable direct summand.

On the other hand, notice that  $q_{\mathbb{X}} \in \tilde{\mathbb{S}}_{\mathbb{Y}}(M)$  as  $[q_{\mathbb{X}}, JMJ] = 0$ , and we have  $\varphi(q_{\mathbb{X}}^{\perp}) = 0$  since N is properly proximal relative to X. In fact, it is clear that

$$q_{\mathbb{X}}^{\perp} \tilde{\mathbb{S}}_{\mathbb{X}}(M) q_{\mathbb{X}}^{\perp} \subset \mathbb{B}(L^2 M)_J^{\sharp *} \cap (JMJ)' \subset \tilde{\mathbb{S}}_{\mathbb{Y}}(M)$$

and hence restricting  $\varphi$  to  $q_{\mathbb{X}}^{\perp} \tilde{\mathbb{S}}_{\mathbb{X}}(M) q_{\mathbb{X}}^{\perp}$  gives us an *N*-central state on  $\tilde{\mathbb{S}}_{\mathbb{X}}(M)$ . As  $\iota_{\text{nor}}(p\mathbb{S}_{\mathbb{X}}(M)p) \subset \tilde{\mathbb{S}}_{\mathbb{X}}(M)$ , we yields an *N*-central state on  $p\mathbb{S}_{\mathbb{X}}(M)p$  that restricts to a normal state on pMp, contradicting the assumption that *N* is properly proximal relative to  $\mathbb{X}$  in *M*.

From the above discussion, we conclude that  $\varphi(q_{\mathbb{Y}}^{\perp}q_{\mathbb{X}}) = 1$ . Furthermore, if we denote by  $\mathbb{X}_i$  the boundary piece associated with  $L\Lambda_i$  and by  $q_i$  the identity of  $\mathbb{K}_{\mathbb{X}_i}(M)_J^{\sharp*}$ , we then have  $\varphi(q_{\mathbb{Y}}^{\perp}q_iq_{\mathbb{Y}}^{\perp}) > 0$  for some  $1 \leq i \leq n$ . This is a consequence of the facts that  $q_{\mathbb{X}} = \bigvee_{i=1}^n q_i$  and  $\{q_i\}_{i=1}^n$  are pairwise commuting by [DKE22, Lemma 3.10].

By Lemma 4.1, we have an *M*-bimodular map  $\phi : \langle M, e_{L\Lambda_i} \rangle \to q_{\mathbb{Y}}^{\perp} \tilde{\mathbb{S}}_{\mathbb{Y}}(M) q_{\mathbb{Y}}^{\perp}$  for each *i*. Composing with  $\psi_i := \varphi(\cdot)/\varphi(q_{\mathbb{Y}}^{\perp}q_iq_{\mathbb{Y}}^{\perp})$  yields an *N*-central state which is normal on *M* and hence we may find some projection  $p_i \in \mathcal{Z}(N' \cap pMp)$  such that  $Np_i$  is amenable relative to  $L\Lambda_i$  in *M*, where  $p_i$  is the support of  $\psi_i$  on  $\mathcal{Z}(N' \cap pMp)$ .

Lastly, we claim that  $\forall p_i = p$ . Indeed, observe that  $\varphi(q_i p_i^{\perp}) = 0$  and  $[p_i q_i, p_j q_j] = 0$  and hence  $\varphi(\forall p_i) \geq \varphi(\forall p_i q_i) = \varphi(\forall q_i) = \varphi(p)$ . Since  $\varphi_{|pMp|}$  is a faithful normal state, it follows that  $\forall p_i = p$ .

#### 5. Some properties of the comultiplication map

The comultiplication map  $\Delta$  was first considered in [PV10] and has been an indispensable tool to obtain superrigidity results (e.g. [Ioa11, IPV13]). Since our proof of Theorem 1.1 follows the strategy of [IPV13] closely, we collect some properties of the comultiplication map in relation to proper proximality in this section.

Recall that if a von Neumann algebra M is isomorphic to  $L\Lambda$  for some group  $\Lambda$ , then the associated comultiplication is given by

$$\Delta: M \ni u_t \to u_t \otimes u_t \in M \overline{\otimes} M,$$

for  $t \in \Lambda$ , where  $u_t \in L\Lambda$  is the canonical unitary corresponding to  $t \in \Lambda$ .

Observe that the comultiplication extends to an isometry  $V : L^2M \ni \delta_t \mapsto \delta_t \otimes \delta_t \in L^2M \otimes L^2M$ , and we denote by  $\phi = \operatorname{Ad}(V) : \mathbb{B}(L^2M \otimes L^2M) \to \mathbb{B}(L^2M)$  the corresponding u.c.p. map.

**Lemma 5.1.** Let  $M \cong L\Lambda$ ,  $\Delta$  and  $\phi$  be as above. Then the following are true.

- (1) We have  $\phi(x) = (\Delta^{-1} \circ E_{\Delta(M)})(x)$  and  $\phi(JxJ) = J(\Delta^{-1} \circ E_{\Delta(M)})(x)J$  for any  $x \in M \otimes M$ , where  $E_{\Delta(M)} : M \otimes M \to \Delta(M)$  is the normal conditional expectation.
- (2) Both  $\phi(\mathbb{K}(L^2M) \otimes_{\min} \mathbb{B}(L^2M))$  and  $\phi(\mathbb{B}(L^2M) \otimes_{\min} \mathbb{K}(L^2M))$  are in  $\mathbb{K}(L^2M)$ .
- (3) The map  $\phi$  is continuous in  $\|\cdot\|_{\infty,1}$  and  $\|\cdot\|_{\infty,2}$ .

*Proof.* (1) is straightforward to check.

(2) Let  $\{\xi_n\} \subset (L^2 M)_1$  be an sequence that converges to 0 weakly and write each  $\xi_n = \sum_{t \in \Lambda} \alpha_{n,t} \delta_t$ . Since  $\xi_n \to 0$  weakly, for any finite subset  $F \subset \Lambda$ , we have  $\lim_{n\to\infty} ||P_F \xi_n|| = \lim_{n\to\infty} (\sum_{t\in F} |\alpha_{n,t}|^2)^{1/2} = 0$ , where  $P_F \in \mathbb{K}(\ell^2 \Lambda)$  is the finite rank projection corresponding to F.

Then for any contraction  $T \in \mathbb{B}(L^2M)$ , we have

$$\|(P_F \otimes T)V\xi_n\| = \|(1 \otimes T)(\sum_{t \in F} \alpha_{n,t}\delta_t \otimes \delta_t)\| \le (\sum_{t \in F} |\alpha_{n,t}|^2)^{1/2} \to 0$$

as  $n \to \infty$ . It then follows that  $\phi$  maps  $\mathbb{K}(L^2M) \otimes_{\min} \mathbb{B}(L^2M)$  and  $\mathbb{B}(L^2M) \otimes_{\min} \mathbb{K}(L^2M)$  to  $\mathbb{K}(L^2M)$ .

(3) For any  $T \in \mathbb{B}(L^2 M \otimes M)$ , note that

$$\|\phi(T)\|_{\infty,1} = \sup_{a,b \in (M)_1} \langle TV\hat{a}, V\hat{b} \rangle = \sup_{a,b \in (M)_1} \langle T\widehat{\Delta(a)}, \widehat{\Delta(b)} \rangle \le \|T\|_{\infty,1}.$$

The proof for  $\|\cdot\|_{\infty,2}$  is similar.

**Lemma 5.2.** Let  $\Lambda$  be a countable discrete group and  $M = L\Lambda$ . Denote by  $\Delta : L\Lambda \to L\Lambda \overline{\otimes} L\Lambda$ the comultiplication map and  $\mathbb{X}$  the  $M \overline{\otimes} M$ -boundary piece associated with  $\{M \otimes 1, 1 \otimes M\}$ . Then we have  $\phi : \mathbb{K}^{\infty,1}_{\mathbb{X}}(M \overline{\otimes} M) \to \mathbb{K}^{\infty,1}(M)$ , where  $\phi$  is the u.c.p. map defined above.

*Proof.* Set  $\tilde{\Lambda} = (\Lambda \times \{e\}) \cup (\{e\} \times \Lambda) \subset \Lambda \times \Lambda$  and denote by  $P_S : \ell^2(\Lambda \times \Lambda) \to \overline{\operatorname{sp}\{\delta_t \mid t \in S\}}$  the orthogonal projection for any subset  $S \subset \Lambda \times \Lambda$ .

By Lemma 5.1, (2), we have  $\phi(P_{F\tilde{\Lambda}F}) \in \mathbb{K}(L^2M)$  for any finite subset  $F \subset \Lambda$ . Denote by  $\mathbb{X}_0$  the hereditary C\*-subalgebra generated by  $\{xJyJP_{\tilde{\Lambda}} \mid x, y \in C_r^*(\Lambda \times \Lambda)\}$ . By the proof of [Din24, Lemma 3.5], we have  $\mathbb{X}_0 \subset \mathbb{K}^{\infty,1}_{\mathbb{X}}(M \overline{\otimes} M)$  is dense in  $\|\cdot\|_{\infty,1}$ .

One checks that  $\{P_{F\tilde{\Lambda}F}\}_F$  forms an approximate unit in  $\mathbb{X}_0$  and hence  $\phi(\mathbb{X}_0) \subset \mathbb{K}(L^2M)$ . Moreover, since  $\phi$  is continuous in  $\|\cdot\|_{\infty,1}$  and  $\mathbb{X}_0 \subset \mathbb{K}^{\infty,1}_{\mathbb{X}}(M \otimes M)$  is dense in  $\|\cdot\|_{\infty,1}$  by Lemma 5.1, (3), we conclude  $\phi(\mathbb{K}^{\infty,1}_{\mathbb{X}}(M \otimes M)) \subset \mathbb{K}^{\infty,1}(M)$ .

Given a finite von Neumann algebra M with a von Neumann subalgebra N, we denote by  $E_N^M$ :  $M \to N$  the normal conditional expectation.

**Corollary 5.3.** Let  $\Lambda$  be a countable discrete group and  $N \subset L\Lambda =: M$  a von Neumann subalgebra. Denote by  $\Delta : M \to M \overline{\otimes} M$  the comultiplication map and  $\mathbb{X}$  the  $M \overline{\otimes} M$ -boundary piece associated with  $\{M \otimes 1, 1 \otimes M\}$ . Then there exists a u.c.p. map  $\phi : \tilde{\mathbb{S}}_{\mathbb{X}}(M \overline{\otimes} M) \to \tilde{\mathbb{S}}(N)$  such that  $\phi(x) = E_N^M \circ \Delta^{-1} \circ E_{\Delta(M)}^{M \overline{\otimes} M}(x)$  for all  $x \in M \overline{\otimes} M$ .

In particular, if there exists an N-central state  $\varphi : \mathbb{S}(N) \to \mathbb{C}$  with  $\varphi_{|N} = \tau_N$ , then  $\psi := \varphi \circ \phi : \widetilde{\mathbb{S}}_{\mathbb{X}}(M \boxtimes M) \to \mathbb{C}$  is  $\Delta(N)$ -central with  $\psi(x) = \tau(x)$  for any  $x \in M \boxtimes M$ , where  $\tau$  is a trace on  $M \boxtimes M$ .

Proof. Let  $\phi : \mathbb{B}(L^2M \otimes L^2M) \to \mathbb{B}(L^2M)$  be the u.c.p. map as above. By Lemma 5.1, (1), we have  $\phi^*$  maps  $\mathbb{B}(L^2M)_J^{\sharp}$  to  $\mathbb{B}(L^2(M \overline{\otimes} M))_J^{\sharp}$  and hence by taking the bidual of  $\phi$  we obtain  $\tilde{\phi} : (\mathbb{B}(L^2(M \overline{\otimes} M))_J^{\sharp})^* \to (\mathbb{B}(L^2M)_J^{\sharp})^*$ , which satisfies  $\tilde{\phi}(\iota_{\mathrm{nor}}(x)) = \iota_{\mathrm{nor}}(\phi(x))$  for any  $x \in M \overline{\otimes} M$  and  $J(M \overline{\otimes} M)J$ . Moreover, from Lemma 5.2, we have  $\tilde{\phi} : \mathbb{K}_{\mathbb{X}}(M \overline{\otimes} M) \to \mathbb{K}^{\infty,1}(M) \subset (\mathbb{K}(M)_J^{\sharp})^*$ . By continuity of  $\tilde{\phi}$ , it follows that  $\tilde{\phi} : (\mathbb{K}_{\mathbb{X}}(M \overline{\otimes} M)_J^{\sharp})^* \to (\mathbb{K}(M)_J^{\sharp})^*$ . Thus we have  $\tilde{\phi} :$ 

 $\tilde{\mathbb{S}}_{\mathbb{X}}(M \overline{\otimes} M) \to \tilde{\mathbb{S}}(M)$  with  $\tilde{\phi}(x) = (\Delta^{-1} \circ E_{\Delta(M)}^{M \overline{\otimes} M})(x)$  for any  $x \in M \overline{\otimes} M$ . Combining  $\tilde{\phi}$  with [Din24, Lemma 3.2] yields a u.c.p. map  $\psi : \tilde{\mathbb{S}}_{\mathbb{X}}(M \overline{\otimes} M) \to \tilde{\mathbb{S}}(N)$  with the desired property.  $\Box$ 

**Lemma 5.4.** Let  $\Lambda$  be an i.c.c. countable discrete group,  $M = L\Lambda$  and  $N, P \subset M$  von Neumann subalgebras with  $\mathcal{Z}(N) = \mathbb{C}$ . Denote by  $\Delta : M \to M \otimes M$  the comultiplication map given by  $M = L\Lambda$ .

Suppose there exist some nonzero projection  $p \in \Delta(N)' \cap (M \otimes M)$  and some partial isometry  $v \in M \otimes M$  such that  $v^*(\Delta(N)p)v \subset P \otimes M$  and  $vv^* = p$ .

Then there exists a c.c.p. map  $\phi : \mathbb{B}(L^2(P \otimes M)) \to \mathbb{B}(L^2N)$  such that

- (1) For any  $x \in P \overline{\otimes} M$  we have  $\phi(x) = (\Delta^{-1} \circ E_{\Delta(N)}^{M \overline{\otimes} M})(pvxv^*p).$
- (2) For any  $y \in N$ , we have  $\phi(J(v^*\Delta(y)pv)J) = \tau(p)JyJ$ , where  $\tau$  is the trace on  $M \otimes M$ .
- (3) Denoting by  $\mathbb{X}_{P\otimes 1}$  the  $P \overline{\otimes} M$  boundary piece associated with  $\{P \otimes 1\}$ , we have  $\phi$  maps  $\mathbb{K}^{\infty,1}_{\mathbb{X}_{P\otimes 1}}(P \overline{\otimes} M)$  to  $\mathbb{K}^{\infty,1}(N)$ .

Proof. Setting  $q = v^* v$ , we have pv = vq and hence  $v^*(\Delta(N)p)v = (v^*\Delta(N)v)q$  and  $[q, v^*\Delta(N)v] = 0$ . We may extend v to a unitary  $u \in M \otimes M$ , which satisfies uq = vq and pu = pv, and it follows that  $v^*(\Delta(N)p)v = (u^*\Delta(N)u)q$  with  $[q, u^*\Delta(N)u] = 0$ .

Put  $N_1 = u^* \Delta(N) u$  and  $\tau$  to be the canonical trace on  $M \otimes M$ .

Since N is a factor, the map  $V_0 : L^2(N_1, \tau) \ni \hat{x} \to q\hat{x} \in L^2(N_1q, \tau)$  has norm  $\tau(q)^{1/2}$  and  $\operatorname{Ad}(V_0) : \mathbb{B}(L^2(N_1q, \tau)) \to \mathbb{B}(L^2(N_1, \tau))$  satisfies  $\operatorname{Ad}(V_0)(qx) = \tau(q)x$  and  $\operatorname{Ad}(V_0)(JqxJ) = \tau(q)JxJ$  for any  $x \in N_1$ .

Consider

$$V: L^2N \xrightarrow{\Delta} L^2(\Delta(N), \tau) \xrightarrow{u^*Ju^*J} L^2(N_1, \tau) \xrightarrow{V_0} L^2(N_1q, \tau) \xrightarrow{\iota} L^2(P \overline{\otimes} M, \tau),$$

where  $\iota: L^2(N_1q, \tau) \to L^2(P \overline{\otimes} M, \tau)$  is the inclusion map, and set

$$\phi := \operatorname{Ad}(V) : \mathbb{B}(L^2(P \otimes M, \tau)) \to \mathbb{B}(L^2 N).$$

For any  $a \in N$ , one checks that  $V\hat{a} = u^*\Delta(a)u\hat{q}$  and thus  $\phi(x) = \Delta^{-1} \circ E_{\Delta(N)}(uqxqu^*)$  for  $x \in P \otimes M$ , and  $\phi(J(u^*\Delta(y)uq)J) = \tau(p)JyJ$  for any  $y \in N$ . In particular,  $\tau(p)^{-1}\phi$  is a u.c.p. map.

Now we show  $\phi(\mathbb{K}_{\mathbb{X}_{P\overline{\otimes}1}}^{\infty,1}(P\overline{\otimes}M)) \subset \mathbb{K}^{\infty,1}(N)$ . Indeed, from Lemma 5.1, (2) and (3), we see that if  $a_i \in (N)_1$  is a sequence converging to 0 weakly, then  $\|K\Delta(\hat{a}_i)\| \to 0$  for any  $K \in \mathbb{K}_{\mathbb{X}_{M\otimes 1}}(M\overline{\otimes}M)$ . It follows that

$$||Ku^*\Delta(a_i)u\hat{q}|| = ||Ku^*Jqu^*J\Delta(\hat{a}_i)|| \to 0,$$

as  $i \to \infty$  for any  $K \in \mathbb{K}_{\mathbb{X}_{M\otimes 1}}(M \overline{\otimes} M)$ , since  $\mathbb{K}_{\mathbb{X}_{M\otimes 1}}(M \overline{\otimes} M)$  is invariant under pre and post composition with  $M \overline{\otimes} M$  and  $J(M \overline{\otimes} M)J$ . Since  $\operatorname{Ad}(e_{P \overline{\otimes} M}) : \mathbb{K}_{\mathbb{X}_{M\otimes 1}}^{\infty,1}(M \overline{\otimes} M) \to \mathbb{K}_{\mathbb{X}_{P\otimes 1}}^{\infty,1}(P \overline{\otimes} M)$ is onto, we have

$$\phi(\mathbb{K}^{\infty,1}_{\mathbb{X}_{P\otimes 1}}(P\overline{\otimes}M)) \subset \mathbb{K}^{\infty,1}(N).$$

**Corollary 5.5.** Let  $\Lambda$  be an i.c.c. countable discrete group,  $P \subset M := L\Lambda$  a von Neumann subalgebra and  $N \subset M$  a subfactor. Denote by  $\Delta : M \to M \otimes M$  the comultiplication map.

Suppose there exist a nonzero projection  $p \in \Delta(N)' \cap (M \otimes M)$  and a partial isometry  $v \in M \otimes M$ such that  $v^*(\Delta(N)p)v \subset M \otimes P$  and  $vv^* = p$ . Then there exists a c.c.p. map  $\phi : \tilde{\mathbb{S}}_{\mathbb{X}_{P\otimes 1}}(P \otimes M) \to \tilde{\mathbb{S}}(N)$  such that  $\phi(\iota_{nor}(x)) = (\iota_{nor} \circ \Delta^{-1} \circ E^{M \otimes M}_{\Delta(N)})(pvxv^*p)$  for any  $x \in P \otimes M$ .

#### CHANGYING DING

In particular, if there exists an N-central state  $\varphi : \mathbb{S}(N) \to \mathbb{C}$  with  $\varphi_{|N} = \tau$ , then  $\psi := \varphi \circ \phi : \mathbb{S}_{\mathbb{X}_{P \otimes 1}}(P \otimes M) \to \mathbb{C}$  is  $v^*(\Delta(N)p)v$ -central and  $\psi(x) = \tau(qxq)$  for  $x \in P \otimes M$ , where  $q = v^*v$ .

*Proof.* This follows from Lemma 5.4, similar to the argument in Corollary 5.3.

#### 6. Proof of main theorems

Now we are ready to demonstrate rigidity of non properly proximal groups in the context of Bernoulli actions.

**Proposition 6.1.** Let  $\Gamma$  be a countable discrete group that is i.c.c. and nonamenable. Suppose  $\Gamma$  is non properly proximal with  $\Lambda_{cb}(\Gamma) = 1$ , and  $L(\mathbb{Z} \wr \Gamma) \cong L\Lambda$  for some group  $\Lambda$ . Then there exists a unitary  $u \in L\Lambda$  such that  $u^*\Delta(L\Gamma)u \subset L\Gamma \otimes L\Gamma$ , where  $\Delta : L\Lambda \to L\Lambda \otimes L\Lambda$  is the comultiplication map.

*Proof.* Set  $M = L(\mathbb{Z} \wr \Gamma)$  and  $N = \Delta(L\Gamma)$ . Since  $(\mathbb{Z} \wr \Gamma) \times (\mathbb{Z} \wr \Gamma)$  is biexact relative to  $\{\Gamma \times (\mathbb{Z} \wr \Gamma), (\mathbb{Z} \wr \Gamma) \times \Gamma\}$  [BO08, Chapter 15], and N has no amenable direct summand, we have N is properly proximal relative to  $\{L\Gamma \otimes M, M \otimes L\Gamma\}$  in  $M \otimes M$  as well by [DP23].

Furthermore, as  $\Gamma$  is non properly proximal, we have a  $L\Gamma$ -central state  $\varphi_0 : \mathbb{S}(L\Gamma) \to \mathbb{C}$  such that  $\varphi_{|L\Gamma} = \tau$ , which yields an N-central state  $\varphi : \tilde{\mathbb{S}}_{\mathbb{X}}(M \otimes M) \to \mathbb{C}$  with  $\varphi_{|M \otimes M} = \tau$  by Corollary 5.3, where  $\mathbb{X}$  denotes the  $M \otimes M$ -boundary piece associated with  $\{M \otimes 1, 1 \otimes M\}$ .

Observe that  $\Gamma < \mathbb{Z} \wr \Gamma$  is almost malnormal and thus  $\Gamma \times (\mathbb{Z}^{\Gamma} \rtimes \Gamma)$  (resp.  $(\mathbb{Z}^{\Gamma} \rtimes \Gamma) \times \Gamma$ ) is almost malnormal relative to  $\{e\} \times (\mathbb{Z}^{\Gamma} \rtimes \Gamma)$  (resp.  $(\mathbb{Z}^{\Gamma} \rtimes \Gamma) \times \{e\}$ ) in  $(\mathbb{Z} \wr \Gamma) \times (\mathbb{Z} \wr \Gamma)$ . Now we are in the situation which Proposition 4.2 applies to, with  $\Lambda_1 = \Gamma \times (\mathbb{Z} \wr \Gamma) \times (\mathbb{Z} \wr \Gamma)$ . Now and  $\Sigma_1 = \{e\} \times (\mathbb{Z} \wr \Gamma), \Sigma_2 = (\mathbb{Z} \wr \Gamma) \times \{e\}$ , and it follows that there exists a partition of unity  $p_1, p_2 \in \mathcal{Z}(N' \cap (M \otimes M))$  such that  $Np_1$  is amenable relative to  $L\Gamma \otimes M$  in  $M \otimes M$ , and  $Np_2$  is amenable relative to  $M \otimes L\Gamma$  in  $M \otimes M$ .

Since  $\Gamma$  is nonamenable, we have N is strongly nonamenable relative to  $1 \otimes M$  and  $M \otimes 1$ ,  $N \not\prec_{M \overline{\otimes} M} L(\oplus_{\Gamma} \mathbb{Z}) \overline{\otimes} M$  and  $N \not\prec_{M \overline{\otimes} M} M \overline{\otimes} L(\oplus_{\Gamma} \mathbb{Z})$  by [IPV13, Proposition 7.2]. It then follows from Proposition 2.6 that we may find partial isometries  $v_1, v_2 \in M \overline{\otimes} M$  such that  $v_1^*(Np_1)v_1 \subset M \overline{\otimes} L\Gamma$  with  $v_1v_1^* = p_1$ , and  $v_2^*(Np_2)v_2 \subset L\Gamma \overline{\otimes} M$  with  $v_2v_2^* = p_2$ .

We then consider  $N_1 := v_1^*(Np_1)v_1$ . One checks that since N is strongly nonamenable relative to  $1 \otimes M$  in  $M \otimes M$ , we have  $N_1$  is also strongly nonamenable relative to  $1 \otimes L\Gamma$  in  $M \otimes L\Gamma$ . Thus by Proposition 3.1, one has  $N_1$  is properly proximal relative to  $L\Gamma \otimes L\Gamma$ .

By Corollary 5.5, we have  $\psi : \tilde{\mathbb{S}}_{\mathbb{X}_{1\otimes L\Gamma}}(M \otimes L\Gamma) \to \mathbb{C}$  is an  $N_1$ -central state with  $\psi(q_1 x q_1) = \tau(p_1)^{-1}\tau(q_1 x q_1)$  for  $x \in M \otimes L\Gamma$ , where  $q_1 = v_1^* v_1$ . We may then apply Proposition 4.2, which yields that  $N_1$  is amenable relative to  $L\Gamma \otimes L\Gamma$  in  $M \otimes L\Gamma$ .

As  $N_1 \not\prec_{M \overline{\otimes} L\Gamma} L(\oplus_{\Gamma} \mathbb{Z}) \overline{\otimes} L\Gamma$ , by invoking Proposition 2.6 one more time, we obtain a partial isometry  $w_1 \in M \overline{\otimes} L\Gamma$  such that  $w_1^* N_1 w_1 \subset L\Gamma \overline{\otimes} L\Gamma$  and  $w_1 w_1^* = q_1$ .

Similarly, we obtain a partial isometry  $w_2 \in L\Gamma \otimes M$  such that  $w_2^*(v_2^*Np_2v_2)w_2 \subset L\Gamma \otimes L\Gamma$  and  $w_2w_2^* = v_2v_2^*$ .

Finally, observe that  $\operatorname{Ad}(v_2w_2)(p_2) \in L\Gamma \otimes L\Gamma$  and  $1 - \operatorname{Ad}(v_1w_1)(p_1) \in L\Gamma \otimes L\Gamma$  are equivalent as  $p_1 + p_2 = 1$ . Thus we may find a partial isometry  $w'_2 \in L\Gamma \otimes \Gamma$  that implements their equivalence. Set  $u = v_1w_1 + v_2w_2w'_2$  and one verifies that u is a unitary and  $u^*Nu \subset L\Gamma \otimes L\Gamma$ .  $\Box$ 

*Proof of Theorem 1.1.* This follows directly from the proof of [IPV13, Theorem 8.2], as its step 1 is established in Proposition 6.1.  $\Box$ 

The following elementary lemma may be derived directly from [Ioa11, Theorem 6.2]. We nevertheless give a straightforward proof.

**Lemma 6.2.** Let H be an abelian group and  $\Gamma$  be an i.c.c. group that acts on H by automorphisms. Suppose  $\Gamma \curvearrowright \widehat{H}$  is conjugate to the Bernoulli action  $\Gamma \curvearrowright [0,1]^{\Gamma}$ . Then for any nontrivial normal abelian subgroup  $K \triangleleft H \rtimes \Gamma$ , we have K < H.

*Proof.* First we show that  $K \cap H$  is nontrivial. Indeed, from  $K \cap H = \{e\}$  one sees that K is in the centralizer of H in  $H \rtimes \Gamma$ . As  $LH \subset L(H \rtimes \Gamma)$  is maximal abelian, we have  $LK \subset LH$ , contradicting  $K \cap H = \{e\}$ .

We denote by  $E_S$  the conditional expectation from  $\overline{\otimes}_{\Gamma} L^{\infty}([0,1])$  to  $\overline{\otimes}_{S} L^{\infty}([0,1])$  for a subset  $S \subset \Gamma$ . For any function  $f \in \overline{\otimes}_{\Gamma} L^{\infty}([0,1])$  and any  $\varepsilon > 0$ , one may consider

$$F_{f,\varepsilon} := \{ t \in \Gamma \mid \|P_t(f) - f\|_2 > \varepsilon \},\$$

where  $P_t = E_{\Gamma \setminus \{t\}}$ . We show that for a non-scalar f, we may find some  $\varepsilon$  such that  $F_{f,\varepsilon}$  is a nonempty finite set.

If  $F_{f,\varepsilon}$  was empty for any  $\varepsilon > 0$ , then one would have  $P_t(f) = f$  for any  $t \in \Gamma$  and hence  $(\prod_{t \in S} P_t)(f) = f$  for any finite  $S \subset \Gamma$ . For any  $\varepsilon > 0$ , take  $S \subset \Gamma$  a finite set such that  $||E_S(f) - f||_2 < \varepsilon$ . Then  $||\tau(f) - f||_2 = ||(\prod_{t \in S} P_s)(E_S(f) - f)||_2 < \varepsilon$  and hence  $f = \tau(f)$ .

To see  $F_{f,\varepsilon}$  is finite for any  $\varepsilon > 0$ , we may find some finite subset  $S \subset \Gamma$  such that  $||f - E_S(f)||_2 < \varepsilon/2$ . Notice that for any  $t \notin S$ , we have  $P_t(E_S(f)) = E_S(f)$  and hence  $P_t(f) - f = P_t(f - E_S(f)) + E_S(f)) - f$ . As  $P_t$  is  $|| \cdot ||_2$ -continuous, we have  $||P_t(f) - f||_2 \le 2||f - E_S(f))||_2 < \varepsilon$  for  $t \notin S$ .

Since  $K \cap H$  is nontrivial, we may take some nontrivial  $g \in K \cap H$  and view  $\lambda_g \in LH \cong L^{\infty}([0,1])^{\Gamma}$ as a function that is not a scalar as  $\tau_{L(H \rtimes \Gamma)}(\lambda_g) = 0$ . Thus we may find some  $\varepsilon > 0$  such that  $F_g := F_{\lambda_g,\varepsilon} \subset \Gamma$  is a finite nonempty set. Note that if  $s \in \Gamma$  fixes g, then  $sF_g = F_g$  as  $\pi_t(\sigma_s(f)) = \pi_{s^{-1}t}(f)$  for any  $t \in \Gamma$ , where we denote by  $\sigma$  the Bernoulli action  $\Gamma \curvearrowright [0,1]^{\Gamma}$ .

For any  $hs \in K$  with  $h \in H$  and  $s \in \Gamma$ , since K is abelian and normal, we have  $[hs, tgt^{-1}] = e$ for any  $t \in \Gamma$ . In particular, this implies  $t^{-1}stF_g = F_g$  for any  $t \in \Gamma$ , which in turn shows that  $\{t^{-1}st \mid t \in \Gamma\}$  is finite, as  $F_g \subset \Gamma$  is a nonempty finite set. As  $\Gamma$  is i.c.c., we conclude s must be trivial and hence K < H.

Proof of Theorem 1.2. Suppose  $\Gamma$  is in addition properly proximal, this follows from [BIP21, Theorem 1.5].

If  $\Gamma$  is non properly proximal, then by Theorem 1.1 one obtains the conclusion of [IPV13, Theorem 8.2]: there exists a group isomorphism  $\delta : \widehat{Y} \rtimes \Lambda \to H \rtimes \Gamma$  for some abelian group H and  $\Gamma \curvearrowright^{\alpha} H$  by automorphisms, a \*-isomorphism  $\theta : L(H) \to L(\oplus_{\Gamma} \mathbb{Z})$  satisfying  $\theta \circ \alpha_g = \sigma_g \circ \theta$  for all  $g \in \Gamma$ , a character  $\eta : \mathbb{Z}\wr\Gamma \to \mathbb{C}$  and a unitary  $w \in L(\mathbb{Z}\wr\Gamma)$  such that the isomorphism  $\pi : L(\widehat{Y} \rtimes \Lambda) = L(\mathbb{Z}\wr\Gamma)$  is given by  $\pi = \operatorname{Ad}(w) \circ \pi_{\eta} \circ \pi_{\theta} \circ \pi_{\delta}$ , with

$$\pi_{\delta}: L(Y \rtimes \Lambda) \ni \lambda_g \to \lambda_{\delta(g)} \in L(H \rtimes \Gamma),$$
$$\pi_{\theta}: L(H) \rtimes \Gamma \ni au_g \mapsto \theta(a)u_g \in L(\oplus_{\Gamma}\mathbb{Z}) \rtimes \Gamma,$$
$$\pi_{\eta}: L(\mathbb{Z} \wr \Gamma) \ni \lambda_g \to \eta(g)\lambda_g \in L(\mathbb{Z} \wr \Gamma).$$

By Lemma 6.2, we have  $\delta(\widehat{Y}) < H$  and hence  $\operatorname{Ad}(w^*) \circ \pi(L^{\infty}(Y)) = L(\bigoplus_{\Gamma} \mathbb{Z}).$ 

## CHANGYING DING

### References

[BIP21]	Rémi Boutonnet, Adrian Ioana, and Jesse Peterson, Properly proximal groups and their von Neumann algebras. Ann. Sci. Éc. Norm. Supér. (4) 54 (2021), no. 2, 445–482
[BO08]	Nathanial P. Brown and Narutaka Ozawa, C <sup>*</sup> -algebras and finite-dimensional approximations, Graduate Studies in Mathematics, vol. 88, American Mathematical Society, Providence, PL 2008
[Bou14]	Rémi Boutonnet, Several rigidity features of von Neumann algebras, Theses, Ecole normale supérieure de lyon - ENS LYON June 2014
[CH89]	Michael Cowling and Uffe Haagerup, Completely bounded multipliers of the Fourier algebra of a simple Lie group of real rank one Invent Math <b>96</b> (1989) no 3 507–549 MB 996553
[Din24]	Changying Ding, First $\ell^2$ -Betti numbers and proper proximality, Adv. Math. <b>438</b> (2024), 109467. MR 4686743
[DKE22]	Changying Ding and Srivatsav Kunnawalkam Elayavalli, On the structure of relatively biexact group von neumann algebras, arXiv:2211.05298, 2022.
[DKEP23]	Changying Ding, Srivatsav Kunnawalkam Elayavalli, and Jesse Peterson, Properly proximal von Neu- mann algebras, Duke Math. J. <b>172</b> (2023), no. 15, 2821–2894. MR 4675043
[DP23]	Changying Ding and Jesse Peterson <i>Bieract von neumann algebras</i> arXiv 2309 10161 2023
[Ioa11]	Adrian Ioana, $W^*$ -superrigidity for Bernoulli actions of property (T) groups, J. Amer. Math. Soc. 24 (2011), no. 4, 1175–1226. MR 2813341
[Ioa15]	, Cartan subalgebras of amalgamated free product II <sub>1</sub> factors, Ann. Sci. Éc. Norm. Supér. (4) 48 (2015), no. 1, 71–130, With an appendix by Ioana and Stefaan Vaes. MR 3335839
[Ioa18]	, <i>Rigidity for von neumann algebras</i> , Proceedings of the International Congress of Mathemati- cians. Volume II, 2018, pp. 1635–1668.
[IPV13]	Adrian Ioana, Sorin Popa, and Stefaan Vaes, A class of superrigid group von Neumann algebras, Ann. of Math. (2) <b>178</b> (2013), no. 1, 231–286.
[Iso20]	Yusuke Isono, On fundamental groups of tensor product II <sub>1</sub> factors, J. Inst. Math. Jussieu <b>19</b> (2020), no. 4, 1121–1139. MR 4120805
[MvN43]	F. I. Murray and I. yon Neumann. On rings of operators, IV App. of Math. (2) 44 (1943) 716-808
[OP10]	Narutaka Ozawa and Sorin Popa, On a class of $II_1$ factors with at most one Cartan subalgebra, Ann. of Math. (2) <b>172</b> (2010), no. 1, 713–749.
$\left[O_{22}04\right]$	Narutaka Orawa Solid von Neumann algebras Acta Math <b>192</b> (2004) no 1 111–117
[Pop06a]	Sorin Popa, some rigidity results for non-commutative Bernoulli shifts, J. Funct. Anal. 230 (2006), no. 2, 273–328. MB 2186215
[Pop06b]	$\frac{100}{2}$ , strong rigidity of II <sub>1</sub> factors arising from malleable actions of w-rigid groups. I, Invent. Math. <b>165</b> (2006), no. 2, 369–408.
[Pop06c]	, strong rigidity of II <sub>1</sub> factors arising from malleable actions of w-rigid groups. II, Invent. Math. <b>165</b> (2006), no. 2, 409–451.
[Pop07]	, Deformation and rigidity for group actions and von Neumann algebras, International Congress of Mathematicians. Vol. I, Eur. Math. Soc., Zürich, 2007, pp. 445–477.
[Pop08]	, On the superrigidity of malleable actions with spectral gap, J. Amer. Math. Soc. 21 (2008), no. 4, 981–1000.
[PV10]	Sorin Popa and Stefaan Vaes, Group measure space decomposition of II <sub>1</sub> factors and $W^*$ -superrigidity, Invent. Math. <b>182</b> (2010), no. 2, 371–417. MR 2729271
[PV14a]	, Unique Cartan decomposition for $II_1$ factors arising from arbitrary actions of free groups, Acta Math. <b>212</b> (2014), no. 1, 141–198.
[PV14b]	, Unique Cartan decomposition for II <sub>1</sub> factors arising from arbitrary actions of hyperbolic groups, J. Reine Angew. Math. <b>694</b> (2014), 215–239.
[Sin55]	I. M. Singer, Automorphisms of finite factors, Amer. J. Math. 77 (1955), 117–133.
[TD20]	Robin D. Tucker-Drob, <i>Invariant means and the structure of inner amenable groups</i> , Duke Math. J. <b>169</b> (2020), no. 13, 2571–2628. MR 4142752
[Vae10]	Stefaan Vaes, <i>Rigidity for von Neumann algebras and their invariants</i> , Proceedings of the International Congress of Mathematicians. Volume III, Hindustan Book Agency, New Delhi, 2010, pp. 1624–1650.

DEPARTMENT OF MATHEMATICS, UCLA, LOS ANGELES, CA 90095, USA

 $Email \ address: \ {\tt cding@math.ucla.edu}$