

Minimax Optimal Goodness-of-Fit Testing with Kernel Stein Discrepancy

Omar Hagrass¹, Bharath Sriperumbudur¹, and Krishnakumar Balasubramanian²

¹Department of Statistics, Pennsylvania State University

²Department of Statistics, University of California, Davis

¹{oih3, bks18}@psu.edu

²kbala@ucdavis.edu

Abstract

We explore the minimax optimality of goodness-of-fit tests on general domains using the kernelized Stein discrepancy (KSD). The KSD framework offers a flexible approach for goodness-of-fit testing, avoiding strong distributional assumptions, accommodating diverse data structures beyond Euclidean spaces, and relying only on partial knowledge of the reference distribution, while maintaining computational efficiency. We establish a general framework and an operator-theoretic representation of the KSD, encompassing many existing KSD tests in the literature, which vary depending on the domain. We reveal the characteristics and limitations of KSD and demonstrate its non-optimality under a certain alternative space, defined over general domains when considering χ^2 -divergence as the separation metric. To address this issue of non-optimality, we propose a modified, minimax optimal test by incorporating a spectral regularizer, thereby overcoming the shortcomings of standard KSD tests. Our results are established under a weak moment condition on the Stein kernel, which relaxes the bounded kernel assumption required by prior work in the analysis of kernel-based hypothesis testing. Additionally, we introduce an adaptive test capable of achieving minimax optimality up to a logarithmic factor by adapting to unknown parameters. Through numerical experiments, we illustrate the superior performance of our proposed tests across various domains compared to their unregularized counterparts.

MSC 2010 subject classification: Primary: 62G10; Secondary: 65J20, 65J22, 46E22, 47A52.

Keywords and phrases: Goodness-of-fit test, maximum mean discrepancy, kernel Stein discrepancy, reproducing kernel Hilbert space, covariance operator, U-statistics, Bernstein's inequality, minimax separation, adaptivity, wild bootstrap, spectral regularization

1 Introduction

Given $\mathbb{X}_n := (X_i)_{i=1}^n \stackrel{i.i.d.}{\sim} P$, where P is defined on a measurable space \mathcal{X} , a goodness-of-fit test involves evaluating the hypotheses

$$H_0 : P = P_0 \quad \text{vs.} \quad H_1 : P \neq P_0,$$

where P_0 denotes a fixed known distribution. See, for example, Lehmann and Romano (2006); Ingster and Suslina (2003) for a textbook-level treatment of the subject. From a methodological perspective, several different approaches exist for constructing goodness-of-fit tests, a few of which

we discuss next. Approaches based on the concept of nearest-neighbor statistics have been explored when \mathcal{X} is a d -dimensional Euclidean space (Bickel and Breiman, 1983; Schilling, 1983a,b), or a manifold (Ebner et al., 2018). Other tests for the case of manifolds include works by Mardia et al. (1984), Jupp (2008) and Boente et al. (2014), to name a few, which are invariably limited to the case of P_0 being uniform or P_0 being the von Mises–Fisher distribution on the sphere. Yet another approach for goodness-of-fit testing involves using the innovation process (Khmaladze, 1988, 1993, 2016), which, to the best of our knowledge, is limited to Euclidean spaces. Techniques based on optimal transport have been used by Hallin et al. (2021) for goodness-of-fit testing in the Euclidean setting, although such approaches could potentially be extended for certain non-Euclidean spaces with additional computational hardships. Goodness-of-fit tests based on variations of χ^2 tests have been analyzed, for example, in Khmaladze (2013) and Balakrishnan and Wasserman (2019) for discrete distributions.

Despite several decades of work on this classical problem, existing tests suffer from at least one of the following drawbacks:

- (i) They rely on strong distributional assumptions, e.g., P_0 is Gaussian or belongs to some classical parametric family.
- (ii) They cannot handle many non-Euclidean data, such as graphs, strings, functions, etc., which is prevalent in many modern applications such as social networks, bioinformatics, etc.
- (iii) They require the complete knowledge of P_0 and cannot handle distributions that are known only up to a normalization constant, even if \mathcal{X} is Euclidean, e.g., a restricted Boltzmann machine (Fischer and Igel, 2012).
- (iv) They are not in general efficiently computable and may need additional samples from P_0 to make them computable even if the complete knowledge of P_0 is available.

In recent years, a non-parametric testing framework based on the concept of reproducing kernel Hilbert space (RKHS) embedding of probability distributions (Muandet et al., 2017; Smola et al., 2007; Sriperumbudur et al., 2010) has gained significant traction, in an attempt to address the issues mentioned above. This approach entails embedding a probability measure P into an RKHS (Aronszajn, 1950), denoted as \mathcal{H}_K , through its corresponding mean element, expressed as:

$$\mu_P = \int_{\mathcal{X}} K(\cdot, x) dP(x) \in \mathcal{H}_K,$$

where $K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ represents the unique reproducing kernel (r.k.) associated with \mathcal{H}_K . By employing this embedding, a pseudo-metric can be defined on the space of probability measures, referred to as the *maximum mean discrepancy* (MMD) (Gretton et al., 2012, 2006; Sriperumbudur et al., 2010; Sriperumbudur, 2016; Simon-Gabriel and Schölkopf, 2018), defined as

$$D_{\text{MMD},K}(P, Q) = \|\mu_P - \mu_Q\|_{\mathcal{H}_K} = \sup \left\{ \int_{\mathcal{X}} f(x) d(P - Q)(x) : f \in \mathcal{H}_K, \|f\|_{\mathcal{H}_K} \leq 1 \right\},$$

where in an alternate interpretation, the discrepancy between P and Q is measured w.r.t. test functions, f whose RKHS norm is bounded by one. Thus, with a fixed P_0 , a consistent goodness-of-fit test can be constructed using the following estimator of $D_{\text{MMD},K}^2$ as a test statistic, i.e.,

$$\hat{D}_{\text{MMD},K}^2(P, P_0) := \frac{1}{n(n-1)} \sum_{i \neq j} \langle K(\cdot, X_i) - \mu_0, K(\cdot, X_j) - \mu_0 \rangle_{\mathcal{H}_K}$$

$$= \frac{1}{n(n-1)} \sum_{i \neq j} K(X_i, X_j) - \frac{2}{n} \sum_{i=1}^n \mu_0(X_i) + \|\mu_0\|_{\mathcal{H}_K}^2, \quad (1)$$

and using the $1 - \alpha$ quantile of the asymptotic null distribution of $\hat{D}_{\text{MMD},K}^2(P, P_0)$ as the critical level (Balasubramanian et al., 2021, Theorem 1), while assuming $\mu_0 := \mu_{P_0}$ and $\|\mu_0\|_{\mathcal{H}_K}^2$ are computable in closed form. Note that the above test addresses issues (i) and (ii) since (1) does not make any assumptions about P_0 and is well-defined for any $(X_i)_i \subset \mathcal{X}$ as long as K can be defined on \mathcal{X} , which alleviates the requirement of \mathcal{X} being Euclidean, and hence the strength of this framework. While this provides a convincing solution to issues (i) and (ii) raised above, it still suffers from issues in (iii) and (iv) since $\hat{D}_{\text{MMD},K}^2(P, P_0)$ is computable only if μ_0 and $\|\mu_0\|_{\mathcal{H}_K}^2$ are computable in closed form, which in turn is possible only for a few (K, P_0) pairs—e.g., both K and P_0 are Gaussian—, and definitely not possible if P_0 is not fully specified.

When $\mathcal{X} = \mathbb{R}^d$, recently, Chwialkowski et al. (2016) and Liu et al. (2016) proposed a goodness-of-fit test based on kernel Stein discrepancy (KSD), which is defined w.r.t. a suitable class of test functions $\{\mathcal{S}_{P_0} f : f \in \mathcal{H}_K^d\}$ that satisfy $\int_{\mathcal{X}} (\mathcal{S}_{P_0} f)(x) dP_0(x) = 0$ for all $f \in \mathcal{H}_K^d =: \mathcal{H}_K \times \dots \times \mathcal{H}_K$, so that

$$\sup_{f \in \mathcal{H}_K^d} \int_{\mathcal{X}} (\mathcal{S}_{P_0} f)(x) d(P - P_0)(x) = \sup_{f \in \mathcal{H}_K^d} \int_{\mathcal{X}} (\mathcal{S}_{P_0} f)(x) dP(x) =: D_{\text{KSD}}(P, P_0), \quad (2)$$

where $\mathcal{S}_{P_0} f := \langle \nabla \log p_0, f \rangle_2 + \nabla \cdot f$ is called the *Stein operator* with p_0 being the density of P_0 w.r.t. the Lebesgue measure, ∇ is the gradient operator, and $\nabla \cdot f$ is the divergence of f . Using the reproducing property of K , it follows from (2) that

$$\begin{aligned} D_{\text{KSD}}(P, P_0) &= \left\| \int_{\mathbb{R}^d} (\nabla \log p_0(x) K(\cdot, x) + \nabla K(\cdot, x)) dP(x) \right\|_{\mathcal{H}_K^d} = \left\| \int_{\mathbb{R}^d} K_0(\cdot, x) dP(x) \right\|_{\mathcal{H}_{K_0}} \\ &= D_{\text{MMD},K_0}(P, P_0), \end{aligned} \quad (3)$$

where $K_0(x, y) = \langle \nabla \log p_0(x) K(\cdot, x) + \nabla K(\cdot, x), \nabla \log p_0(y) K(\cdot, y) + \nabla K(\cdot, y) \rangle_{\mathcal{H}_K^d}$ is a reproducing kernel, called the *Stein kernel* with \mathcal{H}_{K_0} (Stein RKHS) as the corresponding RKHS. Therefore D_{KSD} defined w.r.t. K is the same as D_{MMD,K_0} defined w.r.t. the Stein kernel, K_0 , which means the test statistic in (1) is now computable w.r.t. K_0 since the corresponding mean element w.r.t. P_0 is zero and K_0 depends on P_0 only through $\nabla \log p_0$, i.e., it is sufficient to know P_0 only up to normalization. Therefore, the KSD test addresses the issues raised in (i), (iii) and (iv) but caters only to Euclidean data and therefore suffers from (ii). This issue of (ii) can be resolved by replacing \mathcal{S}_{P_0} with the corresponding Stein operator defined on functions, graphs, discrete spaces, manifolds, point processes, etc. See Section 1.1 for more details.

In light of the discussion so far, it should be clear that KSD provides a powerful framework for goodness-of-fit testing that is devoid of the issues (i)–(iv) unlike many existing tests. The central goal of this paper is to hence investigate and develop minimax optimal goodness-of-fit tests based on KSD.

1.1 General formulation of KSD with examples

We now introduce a general framework of kernel Stein discrepancy that subsumes KSD defined in (2) for $\mathcal{X} = \mathbb{R}^d$. Let \mathcal{H} be a Hilbert space on \mathcal{X} and ψ_{P_0} be defined on $\mathcal{X} \times \mathcal{X}$ such that

$\psi_{P_0}(\cdot, x) \in \mathcal{H}$, $\forall x \in \mathcal{X}$ and $\mathbb{E}_{P_0} \psi_{P_0}(y, X) = 0$, $\forall y \in \mathcal{X}$. Define the Stein operator \mathcal{S}_{P_0} acting on $f \in \mathcal{H}$ as

$$(\mathcal{S}_{P_0} f)(x) := \langle \psi_{P_0}(\cdot, x), f \rangle_{\mathcal{H}}.$$

Thus, it is clear that $\mathbb{E}_{P_0}(\mathcal{S}_{P_0} f)(X) = \langle \mathbb{E}_{P_0} \psi_{P_0}(\cdot, X), f \rangle_{\mathcal{H}} = 0$ if $\mathbb{E}_{P_0} \|\psi_{P_0}(\cdot, X)\|_{\mathcal{H}} < \infty$. The kernel Stein discrepancy is defined as

$$\begin{aligned} D_{\text{KSD}}(P, P_0) &:= D_{\text{KSD}}(P, \mathcal{S}_{P_0}, \mathcal{H}) := \sup_{f \in \mathcal{H}: \|f\|_{\mathcal{H}} \leq 1} |\mathbb{E}_P(\mathcal{S}_{P_0} f)(X) - \mathbb{E}_{P_0}(\mathcal{S}_{P_0} f)(X)| \\ &= \|\mathbb{E}_P \psi_{P_0}(\cdot, X)\|_{\mathcal{H}} = \sqrt{\mathbb{E}_{P \times P} K_0(X, Y)}, \end{aligned} \quad (4)$$

where

$$K_0(x, y) := \langle \psi_{P_0}(\cdot, x), \psi_{P_0}(\cdot, y) \rangle_{\mathcal{H}} = \langle K_0(\cdot, x), K_0(\cdot, y) \rangle_{\mathcal{H}_{K_0}} \quad (5)$$

is the *Stein kernel* and \mathcal{H}_{K_0} is the (Stein) RKHS associated with K_0 . In the following, we provide some examples that specialize this general framework.

Example 1 (KSD as MMD). *By choosing \mathcal{H} to be the RKHS \mathcal{H}_K associated with the kernel $K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ and defining $\psi_{P_0}(\cdot, x) = K(\cdot, x) - \mathbb{E}_{P_0} K(\cdot, X)$ yields MMD (Gretton et al., 2012, 2006), i.e., $D_{\text{KSD}}(P, \mathcal{S}_{P_0}, \mathcal{H}_K) = D_{\text{MMD}, K}(P, P_0)$ with*

$$K_0(x, y) = K(x, y) - \mathbb{E}_{P_0} K(x, X) - \mathbb{E}_{P_0} K(X, y) + \mathbb{E}_{P_0 \times P_0} K(X, Y).$$

Example 2 (KSD on $\mathcal{X} = \mathbb{R}^d$). *Suppose $\mathcal{X} = \mathbb{R}^d$ and P_0 have a smooth density p_0 w.r.t. the Lebesgue measure. By defining $\psi_{P_0}(\cdot, x) = K(\cdot, x) \nabla_x \log p_0(x) + \nabla_x K(\cdot, x) \in \mathcal{H}_K^d =: \mathcal{H}$ yields the KSD (Chwialkowski et al., 2016; Liu et al., 2016) defined in (3) with*

$$\begin{aligned} K_0(x, y) &= \nabla_x \log p_0(x)^\top \nabla_y \log p_0(y) K(x, y) + \nabla_y \log p_0(y)^\top \nabla_x K(x, y) \\ &\quad + \nabla_x \log p_0(x)^\top \nabla_y K(x, y) + \nabla_x \nabla_y K(x, y), \end{aligned}$$

which is the same K_0 defined below (3).

Example 3 (KSD on a Riemannian manifold). *Suppose $\mathcal{X} = (\mathcal{M}, g)$, where \mathcal{M} is a smooth d -dimensional Riemannian manifold (with either empty boundary or the kernel K defined on $\mathcal{X} \times \mathcal{X}$ vanishes on the boundary) with Riemannian metric g and P_0 has a smooth density p_0 . The choice of $\psi_{P_0}(\cdot, x) = K(\cdot, x) \nabla_{\theta_x} \log(p_0(x) J(\theta)) + \nabla_{\theta_x} K(\cdot, x) \in \mathcal{H}_K^d$, $x \in \mathcal{M}$, yields the manifold kernel Stein discrepancy (mKSD⁽¹⁾) (Xu and Matsuda, 2021) which also covers the directional Stein discrepancy (dKSD) (Xu and Matsuda, 2020) as a special case when \mathcal{M} is the unit hypersphere \mathbb{S}^{d-1} . Here, $\theta = (\theta^1, \dots, \theta^d)$ is a coordinate system on \mathcal{M} that covers \mathcal{M} almost everywhere, $J = \sqrt{\det g}$ is the volume element, and θ_x is the coordinate representation of $x \in \mathcal{M}$. Other variations of mKSD discussed in (Xu and Matsuda, 2021) are covered by choosing appropriate ψ_{P_0} . Note that this setting covers P_0 defined on a torus, sphere (e.g., uniform, von Mises-Fisher, Fisher-Bingham distribution), rotation group (e.g., Fisher distribution), etc.*

Example 4 (KSD on Hilbert spaces). *For a Gibbs measure, P_0 defined on a separable Hilbert space, \mathcal{X} , i.e., $\frac{dP_0}{dN_C}$ exists on \mathcal{X} with $\frac{dP_0}{dN_C} \propto \exp(-U)$, where N_C is a Gaussian measure with mean zero and covariance operator C , the Stein operator on \mathcal{X} is defined through $\psi_{P_0}(\cdot, x) = \text{Tr}[CD^2 k(\cdot, x)] - \langle Dk(\cdot, x), x + CDU(x) \rangle_{\mathcal{X}}$, $x \in \mathcal{X}$, where D denotes the Fréchet differential (Wynne et al., 2023, Definition 3.2). Examples of distributions that can be tested in this setting include Gaussian processes on a separable Hilbert space.*

Similar Stein operators have been defined on exponential random graph models (Xu and Reintert, 2021, 2022), point processes (Yang et al., 2019), Lie groups (Qu et al., 2023), discrete distributions (Yang et al., 2018), latent variable models (Kanagawa et al., 2023), and censored data (Fernandez et al., 2020). To avoid technical notations that are required to define these operators, we do not introduce them here, however, the proposed framework subsumes all these non-Euclidean scenarios involving graph data, point process data, discrete data, censored data, and functional data.

1.2 Minimax framework

Since the goal of our work is to investigate the minimax optimality of the KSD test, we first introduce the minimax framework for hypothesis testing pioneered by Burnashev (1979) and Ingster (1987, 1993), which is essential to understand our contributions.

Let $\phi(\mathbb{X}_n)$ be any test that rejects H_0 when $\phi = 1$ and fails to reject H_0 when $\phi = 0$. Denote the class of all such asymptotic α -level tests to be Φ_α . Let \mathcal{C} be a set of probability measures on \mathcal{X} . The Type-II error of a test $\phi \in \Phi_\alpha$ w.r.t. \mathcal{P}_Δ is defined as $R_\Delta(\phi) = \sup_{P \in \mathcal{P}_\Delta} \mathbb{E}_{P^n}(1 - \phi)$, where

$$\mathcal{P}_\Delta := \{P \in \mathcal{C} : \rho^2(P, P_0) \geq \Delta\},$$

is the class of Δ -separated alternatives in probability metric ρ , with Δ being referred to as the *separation boundary* or *contiguity radius*. Of course, the interest is in letting $\Delta \rightarrow 0$ as $n \rightarrow \infty$ (i.e., shrinking alternatives) and analyzing R_Δ for a given test, ϕ , i.e., whether $R_\Delta(\phi) \rightarrow 0$. In the asymptotic setting, the *minimax separation* or *critical radius* Δ^* is the fastest possible order at which $\Delta \rightarrow 0$ such that $\liminf_{n \rightarrow \infty} \inf_{\phi \in \Phi_\alpha} R_\Delta(\phi) \rightarrow 0$, i.e., for any Δ such that $\Delta/\Delta^* \rightarrow \infty$, there is no test $\phi \in \Phi_\alpha$ that is consistent over \mathcal{P}_Δ . A test is *asymptotically minimax optimal* if it is consistent over \mathcal{P}_Δ with $\Delta \asymp \Delta^*$.

In this work, we consider the class of Δ -separated alternatives in χ^2 -divergence as

$$\mathcal{P} := \mathcal{P}_{\theta, \Delta} := \left\{ P : \frac{dP}{dP_0} - 1 \in \text{Ran}(\Upsilon_{P_0}^\theta), \chi^2(P, P_0) = \left\| \frac{dP}{dP_0} - 1 \right\|_{L^2(P_0)}^2 \geq \Delta \right\}, \theta > 0,$$

where $\Upsilon_{P_0} : L^2(P_0) \rightarrow L^2(P_0)$, $f \mapsto \int_{\mathcal{X}} K_0(\cdot, x)f(x) dP_0(x)$ is the integral operator induced by K_0 , and investigate the optimality of the KSD test w.r.t. \mathcal{P} . The condition $dP/dP_0 - 1 \in \text{Ran}(\Upsilon_{P_0}^\theta)$ captures the smoothness of $dP/dP_0 - 1$ and the degree of smoothness is controlled by θ , i.e., the larger θ is, more smooth is $dP/dP_0 - 1$. $\text{Ran}(\Upsilon_{P_0}^\theta)$ —range space of $\Upsilon_{P_0}^\theta$ —for $\theta \in (0, \frac{1}{2}]$ can be interpreted as an interpolation space obtained by the real interpolation of \mathcal{H}_{K_0} (corresponds to $\theta = \frac{1}{2}$) and $L^2(P_0)$ (corresponds to $\theta = 0$) at scale θ (Steinwart and Scovel, 2012, Theorem 4.6).

We would like to highlight that unlike the classical smoothness classes such as Sobolev or Hölder which are defined for functions defined on $\mathcal{X} = \mathbb{R}^d$, the smoothness class we considered above is defined through the spectrum of the operator Υ_{P_0} that takes into account the interaction between K_0 and P_0 , and applies to domains that are not necessarily Euclidean. Such a smoothness condition is standard in kernel-based methods like kernel regression (Caponnetto and Vito, 2007) and has also been used in two-sample and goodness-of-fit tests (Balasubramanian et al., 2021; Hagrass et al., 2022, 2023). Moreover, the alternative class \mathcal{P} also appears in (Balasubramanian et al., 2021; Hagrass et al., 2023), which investigates the minimax optimality of the MMD test w.r.t. \mathcal{P} (where K_0 corresponds to the kernel in Example 1 in Section 1.1).

1.3 Contributions

Having established the importance of KSD as a discrepancy measure for goodness-of-fit testing, in this work, we investigate the minimax optimality of the goodness-of-fit test based on the test statistic in (1) where K is chosen to be K_0 defined in (5). To the best of our knowledge, only consistency of such a test has been established for $\mathcal{X} = \mathbb{R}^d$ with ψ_{P_0} as defined in Example 2 of Section 1.1 (Chwialkowski et al., 2016; Liu et al., 2016). To this end, the following are the main contributions of our work.

(i) **Non-optimality of KSD test.** First, in Section 3, we provide an operator theoretic representation of D_{KSD} (defined in (4)) by showing that

$$D_{\text{KSD}}^2(P, P_0) = \langle \Upsilon_{P_0} u, u \rangle_{L^2(P_0)} = \sum_{i \geq 1} \lambda_i \langle u, \tilde{\phi}_i \rangle_{L^2(P_0)}^2,$$

where $(\lambda_i, \phi_i)_i$ are the eigenvalues and eigenfunctions of Υ_{P_0} , and $u := \frac{dP}{dP_0} - 1$. This representation will be critical in the study of the optimality of the KSD test. Next, we introduce the KSD test by using the test statistic in (1) with K being replaced by K_0 , and computing the critical level by i.i.d. weighted bootstrap (Dehling and Mikosch, 1994)—also called wild bootstrap. Under the smoothness condition, $u = \frac{dP}{dP_0} - 1 \in \text{Ran}(\Upsilon_{P_0}^\theta)$, $\theta > 0$, we show in Theorem 1 that the KSD test has a separation radius $\Delta_n = o(n^{-2\theta/(2\theta+1)})$ as $n \rightarrow \infty$, irrespective of the decay rate of the eigenvalues of Υ_{P_0} —the eigenvalue decay rate determines the smoothness of functions in \mathcal{H}_{K_0} . On the other hand, we show in Theorem 2 that the minimax separation radius for \mathcal{P} is $\Delta_n = o(n^{-4\theta\beta/(4\theta\beta+1)})$ as $n \rightarrow \infty$ if $\lambda_i \asymp i^{-\beta}$, $\beta > 1$ (i.e., polynomial decay of eigenvalues) and $\Delta_n = o(\sqrt{\log n}/n)$ as $n \rightarrow \infty$ if $\lambda_i \asymp e^{-i}$ (i.e., exponential decay of eigenvalues). These results together establish the non-optimality of the KSD test w.r.t. \mathcal{P} .

(ii) **Minimax optimal regularized KSD test.** In Section 4, we construct a modified version of the KSD test and show it to be minimax optimal w.r.t. \mathcal{P} . This modification is based on the following observations. Note that D_{KSD}^2 defined above downweights the Fourier coefficients of u , i.e., $(\langle u, \tilde{\phi}_i \rangle_{L^2(P_0)})_i$ by the eigenvalues of Υ_{P_0} which decay to zero as $i \rightarrow \infty$. This means that if the differences in P and P_0 are hidden in the high frequency (i.e., large i) coefficients of u , then these differences may not be sufficiently captured by the empirical version of $D_{\text{KSD}}^2(P, P_0)$, which is the test statistic of the KSD test. Hence the non-optimality of the KSD test can be attributed to this limitation of D_{KSD}^2 . To address this issue, in Section 4, we modify $D_{\text{KSD}}^2(P, P_0)$ as $D_\lambda^2(P, P_0) := \sum_i \lambda_i g_\lambda(\lambda_i) \langle u, \tilde{\phi}_i \rangle_{L^2(P_0)}^2$ where $g_\lambda : (0, \infty) \rightarrow (0, \infty)$ is the spectral regularizer satisfying $x g_\lambda(x) \asymp 1$, i.e., $g_\lambda(x)$ is an approximation to x^{-1} as $\lambda \rightarrow 0$ (an example of such a regularizer is the Tikhonov regularizer, $g_\lambda(x) = (x + \lambda)^{-1}$). This property of the spectral regularizer ensures that $D_\lambda^2(P, P_0) \rightarrow \sum_i \langle u, \tilde{\phi}_i \rangle_{L^2(P_0)}^2$ as $\lambda \rightarrow 0$, thereby removing the above mentioned limitation of KSD. A similar observation has been made w.r.t. the limitation of MMD and was addressed through spectral regularization in MMD-based two-sample (Hagrass et al., 2022) and goodness-of-fit (Balasubramanian et al., 2021; Hagrass et al., 2023) tests.

In Section 4, we first show the spectral regularized version of KSD, i.e., $D_\lambda^2(P, P_0)$ to be equal to $\|g^{1/2}(\mathcal{A}_{P_0})\Psi_P\|_{\mathcal{H}_{K_0}}^2$ where $\Psi_P := \int K_0(\cdot, x) dP(x)$ and $\mathcal{A}_{P_0} := \int K_0(\cdot, x) \otimes_{\mathcal{H}_{K_0}} K_0(\cdot, x) dP_0(x)$, which are the mean element of P and covariance operator of P_0 w.r.t. K_0 . If an empirical version of $D_\lambda^2(P, P_0)$ is used as a test statistic, then it still suffers from (iv) since $g^{1/2}(\mathcal{A}_{P_0})$ is in general computable for only some (K_0, P_0) pairs and may require samples from P_0 to estimate \mathcal{A}_{P_0} , which in turn defeats the purpose of using KSD over MMD. Therefore, we consider a modification to D_λ^2 as $\tilde{D}_\lambda^2 := \|g_\lambda^{1/2}(\mathcal{A}_P)\Psi_P\|_{\mathcal{H}_{K_0}}^2$ which can be empirically estimated based on the given samples

(X_1, \dots, X_n) and does not require either additional samples from P_0 or the complete knowledge of P_0 , but only requires information about P_0 up to a normalization constant. The intuition for the modification is that if P is close to P_0 , then the above modification is a good approximation to D_λ and if P is farther from P_0 , then the testing problem is anyway easy enough that the effect of regularization is not significant. Using a U -statistic estimator of \tilde{D}_λ as the test statistic, we propose a goodness-of-fit test using wild bootstrap and establish its minimax optimality w.r.t. \mathcal{P} in Corollaries 6 and 7 (also see Corollaries B.1, B.2 and Theorem 5). This contribution therefore overcomes all the shortcomings raised in (i)–(iv).

(iii) Unbounded Stein kernel. The key technical contribution in (i) and (ii) is the main results of the paper (Theorems 1, 2, 5 and Corollaries 6, 7—also Corollaries B.1, B.2) are obtained without assuming the Stein kernel K_0 to be bounded. While the results related to MMD are obtained for bounded kernels (Li and Yuan, 2019; Balasubramanian et al., 2021; Hagrass et al., 2023), such an assumption on K_0 is quite severe, since, for example, in Example 2, the boundedness of K_0 imposes that p_0 (the density of P_0) cannot have tails thinner than $e^{-\|x\|_1}$. Therefore, the results of this paper are obtained under a weak moment condition on the Stein kernel K_0 . We would like to highlight that this requirement of K_0 being unbounded causes serious technical challenges and our proof techniques significantly differ, for example, from that used in Hagrass et al. (2023) for the analysis of the regularized MMD test.

(iv) Adaptive test. Note that the regularized test presented in Section 4 depends on the regularization parameter $\lambda > 0$ which has to be chosen appropriately. The optimal choice of λ that yields a minimax optimal test depends on the unknown parameters θ and β (in the case of polynomial decay of eigenvalues). Therefore, it is important to construct a minimax optimal test that does not require information about θ and β . To this end, by aggregating tests constructed for different λ , in Section 4.1, we construct an adaptive test (precisely, a union test) that is minimax optimal w.r.t. \mathcal{P} up to a log log factor.

(v) Interpretation for \mathcal{P} . Since the class of alternatives, \mathcal{P} as defined in Section 1.2 is abstract, to develop a deeper understanding of this class, in Section 5, we provide concrete examples to elaborate the range space condition. Specifically, through the provided examples, we highlight the differences in the alternative classes when considering MMD-based tests that have access to full information about P_0 and KSD-based tests that only have partial information about P_0 .

(vi) Numerical experiments. In Section 6, we provide numerical experiments on \mathbb{R}^d , \mathbb{S}^d (sphere), and a separable Hilbert space, comparing the performance of KSD and regularized KSD tests. We show that the regularized KSD tests have superior empirical performance over the KSD test in all these scenarios.

The proofs of the results are provided in Section 8 and the technical results are collected in appendices.

2 Definitions & Notation

For a topological space \mathcal{X} , $L^r(\mathcal{X}, \mu)$ denotes the Banach space of r -power ($r \geq 1$) μ -integrable function, where μ is a finite non-negative Borel measure on \mathcal{X} . For $f \in L^r(\mathcal{X}, \mu) =: L^r(\mu)$, $\|f\|_{L^r(\mu)} := (\int_{\mathcal{X}} |f|^r d\mu)^{1/r}$ denotes the L^r -norm of f . $\mu^n := \mu \times \dots \times \mu$ is the n -fold product measure. \mathcal{H}_K denotes a reproducing kernel Hilbert space with a reproducing kernel $K : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$. $[f]_{\sim}$ denotes the equivalence class associated with $f \in L^r(\mathcal{X}, \mu)$, that is the collection of functions $g \in L^r(\mathcal{X}, \mu)$ such that $\|f - g\|_{L^r(\mu)} = 0$. For two measures P and Q , $P \ll Q$ denotes

that P is dominated by Q which means, if $Q(A) = 0$ for some measurable set A , then $P(A) = 0$.

Let H_1 and H_2 be abstract Hilbert spaces. $\mathcal{L}(H_1, H_2)$ denotes the space of bounded linear operators from H_1 to H_2 . For $S \in \mathcal{L}(H_1, H_2)$, S^* denotes the adjoint of S . $S \in \mathcal{L}(H) := \mathcal{L}(H, H)$ is called self-adjoint if $S^* = S$. For $S \in \mathcal{L}(H)$, $\text{Tr}(S)$, $\|S\|_{\mathcal{L}^2(H)}$, and $\|S\|_{\mathcal{L}^\infty(H)}$ denote the trace, Hilbert-Schmidt and operator norms of S , respectively. For $x, y \in H$, $x \otimes_H y$ is an element of the tensor product space of $H \otimes H$ which can also be seen as an operator from $H \rightarrow H$ as $(x \otimes_H y)z = x\langle y, z \rangle_H$ for any $z \in H$.

For constants a and b , $a \lesssim b$ (resp. $a \gtrsim b$) denotes that there exists a positive constant c (resp. c') such that $a \leq cb$ (resp. $a \geq c'b$). $a \asymp b$ denotes that there exists positive constants c and c' such that $cb \leq a \leq c'b$. We denote $[\ell]$ for $\{1, \dots, \ell\}$.

3 Non-optimality of kernel Stein discrepancy

We begin by examining the operating characteristics of the kernel Stein discrepancy (KSD) based goodness-of-fit tests, by viewing KSD from an operator perspective. This examination will assist in demonstrating the non-optimality of such tests and in introducing the concept of spectral regularization. Throughout the paper, we maintain the following assumption.

Assumption A₀. $(\mathcal{X}, \mathcal{B})$ is a second countable (i.e., completely separable) space endowed with Borel σ -algebra \mathcal{B} . (\mathcal{H}_{K_0}, K_0) is an RKHS of real-valued functions on \mathcal{X} with a continuous reproducing kernel K_0 satisfying $\int_{\mathcal{X}} K_0(x, x) dP_0(x) < \infty$.

From (4), we have

$$\begin{aligned} D_{\text{KSD}}^2(P, P_0) &= \mathbb{E}_{P \times P}(K_0(X, Y)) = \mathbb{E}_{P \times P} \langle K_0(\cdot, X), K_0(\cdot, Y) \rangle_{\mathcal{H}_{K_0}} \\ &\stackrel{(*)}{=} \langle \mathbb{E}_P K_0(\cdot, X) - \mathbb{E}_{P_0} K_0(\cdot, X), \mathbb{E}_P K_0(\cdot, Y) - \mathbb{E}_{P_0} K_0(\cdot, Y) \rangle_{\mathcal{H}_{K_0}} \\ &= \langle \mathbb{E}_{P_0} (K_0(\cdot, X)u(X)), \mathbb{E}_{P_0} (K_0(\cdot, Y)u(Y)) \rangle_{\mathcal{H}_{K_0}}, \end{aligned}$$

where \mathcal{H}_{K_0} represents the RKHS associated with the Stein kernel K_0 , which is a symmetric positive definite kernel due to its definition by the inner product $\langle \psi_{P_0}(\cdot, x), \psi_{P_0}(\cdot, y) \rangle_{\mathcal{H}}$. Here, $u := \frac{dP}{dP_0} - 1$, requiring $P_0 \gg P$ (i.e., P is absolutely continuous w.r.t. P_0) so that the Radon-Nikodym derivative dP/dP_0 is well defined. In the above computation, $(*)$ is derived by noticing that $\mathbb{E}_{P_0} K_0(x, Y) = \langle \psi_{P_0}(\cdot, x), \mathbb{E}_{P_0} \psi_{P_0}(\cdot, Y) \rangle_{\mathcal{H}} = 0$ for all $x \in \mathcal{X}$.

By defining the operator $\Upsilon_{P_0} : L^2(P_0) \rightarrow L^2(P_0)$, $f \mapsto \int K_0(\cdot, x)f(x) dP_0(x)$, it can verified that $\Upsilon_{P_0} = \mathfrak{J}\mathfrak{J}^*$, where $\mathfrak{J} : \mathcal{H}_{K_0} \rightarrow L^2(P_0)$, $f \mapsto [f]_{\sim}$ and $\mathfrak{J}^* : L^2(P_0) \rightarrow \mathcal{H}_{K_0}$, $f \mapsto \int K_0(\cdot, x)f(x) dP_0(x)$. Under **A₀**, it follows that Υ_{P_0} is a positive self-adjoint trace class operator since $\text{Tr}(\Upsilon_{P_0}) = \text{Tr}(\mathfrak{J}\mathfrak{J}^*) = \text{Tr}(\mathfrak{J}^*\mathfrak{J}) = \text{Tr}(\int K_0(\cdot, x) \otimes_{\mathcal{H}_{K_0}} K_0(\cdot, x) dP_0(x)) = \int K_0(x, x) dP_0(x) < \infty$. Thus, the spectral theorem (Reed and Simon, 1980, Theorems VI.16, VI.17) yields that

$$\Upsilon_{P_0} = \sum_{i \in I} \lambda_i \tilde{\phi}_i \otimes_{L^2(P_0)} \tilde{\phi}_i,$$

where $(\lambda_i)_i \subset \mathbb{R}^+$ are the eigenvalues and $(\tilde{\phi}_i)_i$ are the orthonormal system of eigenfunctions of Υ_{P_0} that span $\overline{\text{Ran}(\Upsilon_{P_0})}$ with the index set I being either countable in which case $\lambda_i \rightarrow 0$ or finite. Since $\tilde{\phi}_i$ represents an equivalence class in $L^2(P_0)$, by defining $\phi_i := \frac{\tilde{\mathfrak{J}}^* \tilde{\phi}_i}{\lambda_i}$, it is clear that

$\mathfrak{J}\phi_i = [\phi_i]_{\sim} = \tilde{\phi}_i$ and $\phi_i \in \mathcal{H}_{K_0}$. Throughout the paper, ϕ_i refers to this definition. Consequently,

$$D_{\text{KSD}}^2(P, P_0) = \langle \mathfrak{J}^*u, \mathfrak{J}^*u \rangle_{\mathcal{H}_{K_0}} = \langle \Upsilon_{P_0}u, u \rangle_{L^2(P_0)} = \sum_{i \geq 1} \lambda_i \langle u, \tilde{\phi}_i \rangle_{L^2(P_0)}^2, \quad (6)$$

which shows that KSD is a weighted sum of the squared Fourier components of u with the weights being the eigenvalues of the operator Υ_{P_0} . This representation is similar to that of MMD presented in Hagrass et al. (2023, Section 3), which shows that the squared MMD has a similar representation with Υ_{P_0} replaced by the operator $\mathcal{T} : L^2(P_0) \rightarrow L^2(P_0)$, $f \mapsto \int \bar{K}(\cdot, x)f(x) dP_0(x)$, where $\bar{K}(x, y) = \langle K(\cdot, x) - \mathbb{E}_{P_0}K(\cdot, X), K(\cdot, y) - \mathbb{E}_{P_0}K(\cdot, Y) \rangle_{\mathcal{H}_K}$. Therefore, KSD can be viewed as MMD defined through the Stein kernel K_0 . Hence, the goodness-of-fit test based on $D_{\text{KSD}}^2(P, P_0)$ naturally inherits the non-optimality of the MMD-based test (Hagrass et al., 2023, Theorem 1) if K_0 is assumed to be bounded. However, for the reasons aforementioned in (iii) of Section 1.3, it is important to relax the boundedness assumption on K_0 . Therefore, our key contribution in the KSD non-optimality result presented in Theorem 1 is to do away with the boundedness assumption and replace it with a moment condition on P_0 (see also Remark 2). To this end, we define the following class of alternatives,

$$\mathcal{P} := \mathcal{P}_{\theta, \Delta} := \left\{ P \in \mathcal{S} : \frac{dP}{dP_0} - 1 \in \text{Ran}(\Upsilon_{P_0}^\theta), \chi^2(P, P_0) = \left\| \frac{dP}{dP_0} - 1 \right\|_{L^2(P_0)}^2 \geq \Delta \right\},$$

for $\theta > 0$, where

$$\mathcal{S} := \{P : \mathbb{E}_P[K_0(X, X)]^r \leq cr! \kappa^r, \forall r > 2, \text{ and } \mathbb{E}_P[K_0(X, X)] < \infty\},$$

for some $c, \kappa > 0$. Observe that if K_0 is bounded then both the conditions in \mathcal{S} are satisfied for any distribution P . Thus, when $\psi_{P_0}(\cdot, x) = K(\cdot, x) - \mathbb{E}_{P_0}K(\cdot, X)$ and K is bounded, we recover the alternative space in Hagrass et al. (2023). Define the U -statistic estimator of (4) as

$$S := \hat{D}_{\text{KSD}}^2 = \frac{1}{n(n-1)} \sum_{i \neq j} K_0(X_i, X_j),$$

which can be used as a test statistic in a goodness-of-fit test (Liu et al., 2016; Schrab et al., 2022). We consider a testing threshold based on a bootstrap approach (Arcones and Giné, 1992; Huskova and Janssen, 1993; Dehling and Mikosch, 1994), specifically the i.i.d. weighted bootstrap (Dehling and Mikosch, 1994)—also called *wild bootstrap*—as described below. Let

$$S_\varepsilon := \frac{1}{n(n-1)} \sum_{i \neq j} \varepsilon_i \varepsilon_j K_0(X_i, X_j),$$

where $\varepsilon \in \{-1, 1\}^n$ is a random vector of n i.i.d Rademacher random variables. Then it can be shown that under H_0 , the conditional distribution of S_ε given the samples \mathbb{X}_n , converges to the same distribution as S (Dehling and Mikosch, 1994, Theorem 3.1) if $\mathbb{E}_{P_0}K_0^2(X, Y) < \infty$ —this condition is satisfied in our case since $\mathbb{E}_{P_0}K_0^2(X, Y) \leq [\mathbb{E}_{P_0}K_0(X, X)]^2 < \infty$. Hence, the $(1 - \alpha)$ quantile of S_ε defined as

$$q_{1-\alpha} := \inf \{q \in \mathbb{R} : P_\varepsilon\{S_\varepsilon \leq q | \mathbb{X}_n\} \geq 1 - \alpha\},$$

can be used as a test threshold.

Remark 1. In practice, the calculation of the exact quantile $q_{1-\alpha}$ is computationally expensive. Thus, a Monte Carlo approximation is often used to approximate the exact quantile using B independent realizations of S_ε defined as

$$q_{1-\alpha}^B := \inf \left\{ q \in \mathbb{R} : \frac{1}{B} \sum_{k=1}^B \mathbb{I}(S_{\varepsilon_k} \leq q) \geq 1 - \alpha \right\},$$

where $(\varepsilon_k)_{k=1}^B$ are B randomly selected Rademacher vectors out of 2^n total possibilities. This can be justified using the Dvoretzky-Kiefer-Wolfowitz inequality (see Dvoretzky et al. 1956, Massart 1990) to show that the approximation error between $q_{1-\alpha}$ and $q_{1-\alpha}^B$ gets arbitrarily small for sufficiently large B . Hence, all the results involving $q_{1-\alpha}$ can be extended to $q_{1-\alpha}^B$ for sufficiently large B .

Before presenting the results related to the KSD test, we introduce the following assumption.

Assumption A₁. The pair (P_0, K_0) satisfies $\mathbb{E}_{P_0}[K_0(X, X)]^r \leq cr! \kappa^r$, $\forall r > 2$, for some $c, \kappa > 0$.

Remark 2. Assumption **A₁** is weaker than assuming K_0 to be bounded. As aforementioned in (iii) of Section 1.3, for example, if K_0 is chosen as in Example 2 with K being a bounded translation-invariant kernel on \mathbb{R}^d , then it imposes strong assumptions on P_0 , i.e., p_0 cannot decay faster than $e^{-\|x\|_1}$. On the other hand, Assumption **A₁** imposes the condition $\int |\nabla \log p_0|^{2r} p_0(x) dx \leq cr! \kappa^2$ which of course is satisfied by p_0 that decays faster than $e^{-\|x\|_1}$.

The following result (proved in Section 8.1) provides the separation boundary of the wild bootstrap test based on \hat{D}_{KSD}^2 .

Theorem 1 (Separation boundary of wild bootstrap KSD test). Suppose **A₀** holds and $n^{\frac{2\theta}{2\theta+1}} \Delta_n \rightarrow 0$ as $n \rightarrow \infty$. Then for any decay rate of $(\lambda_i)_i$, if **A₁** holds, we have

$$\liminf_{n \rightarrow \infty} \inf_{\mathcal{P}} P_{H_1} \{ \hat{D}_{\text{KSD}}^2 \geq q_{1-\alpha} \} < 1,$$

for any $\theta > 1$. Furthermore if $\sup_i \|\phi_i\|_\infty < \infty$, then

$$\liminf_{n \rightarrow \infty} \inf_{\mathcal{P}} P_{H_1} \{ \hat{D}_{\text{KSD}}^2 \geq q_{1-\alpha} \} < 1,$$

for any $\theta > 0$.

Remark 3. (i) Note that Theorem 1 remains valid even if the testing threshold $q_{1-\alpha}$ is replaced with any threshold that converges in probability to the $(1-\alpha)$ quantile of the asymptotic distribution of \hat{D}_{KSD}^2 .

(ii) The uniform boundedness assumption $\sup_i \|\phi_i\|_\infty < \infty$ implies that the kernel K_0 is bounded, which follows from Mercer's theorem (see Steinwart and Scovel 2012, Lemma 2.6). But this condition does not hold in general. For example, it holds in any d for Gaussian kernels on \mathbb{R}^d while it does not hold for $d \geq 3$ for Gaussian kernel defined on \mathbb{S}^{d-1} . See (Minh et al., 2006, Theorem 5) and (Hagrass et al., 2022, Remark 6) for details. However, we present the result with this assumption in Theorem 1 to show its similarity to that of MMD (see Haggras et al. 2023, Theorem 1) and to understand its impact on the behavior of the test in contrast to that of **A₁**.

Next, we will show in Theorem 2 (proved in Section 8.2) that the above separation boundary is not optimal (i.e., a better rate can be achieved). We refer the reader to Section 1.2 for the notation used in the following result.

Theorem 2 (Minimax separation). *If $\lambda_i \asymp i^{-\beta}$, $\beta > 1$, and*

$$n^{\frac{4\theta\beta}{4\theta\beta+1}} \Delta_n \rightarrow 0,$$

then $\liminf_{n \rightarrow \infty} \inf_{\phi \in \Phi_\alpha} R_{\Delta_n}(\phi) > 0$, provided one of the following holds: (a) $\theta \geq 1$ under \mathbf{A}_1 , and $\mathbb{E}_{P_0}[K_0(x, Y)]^2 < \infty$; $\forall x \in \mathcal{X}$. (b) $\theta \geq \frac{1}{4\beta}$ under $\sup_i \|\phi_i\|_\infty < \infty$.

Suppose $\lambda_i \asymp e^{-\tau i}$, $\tau > 0$, $\theta > 0$ and

$$\frac{n}{\sqrt{\log n}} \Delta_n \rightarrow 0.$$

Then $\liminf_{n \rightarrow \infty} \inf_{\phi \in \Phi_\alpha} R_{\Delta_n}(\phi) > 0$.

Remark 4. (i) *The combination of the above result with the attained separation boundary of our proposed test (see Section 4), as demonstrated in Corollaries 6 and 7, reveals the non-optimality of the test based on \hat{D}_{KSD} . This is evident since $\inf_{\beta > 1} \frac{4\theta\beta}{4\theta\beta+1} = \frac{4\theta}{4\theta+1} > \frac{2\theta}{2\theta+1}$ and $1 > \frac{2\theta}{2\theta+1}$ for any $\theta > 0$.*

(ii) *If we refrain from assuming a specific decay rate of the eigenvalues λ_i and instead work with $\sum_i \lambda_i < \infty$ (which holds since Υ_{P_0} is a trace class operator), then the general condition on the separation boundary becomes $n^{\frac{4\theta}{4\theta+1}} \Delta_n \rightarrow 0$.*

4 Optimal KSD-based tests via spectral regularization

In the preceding section, we demonstrated the non-optimality of the KSD-based goodness-of-fit test. This deficiency can be ascribed to the representation of KSD in (6) as a weighted sum of the squared Fourier components of u , which suggests that the KSD could overlook discrepancies between P and P_0 occurring at higher frequency components as $\lambda_i \rightarrow 0$. This inherent limitation mirrors the reasons behind the non-optimality of the MMD, as discussed in Hagrass et al. (2023). To address this issue, we use the idea of spectral regularization as considered in Balasubramanian et al. (2021) and Hagrass et al. (2023, 2022) by using a regularization function designed to avoid neglecting the higher frequency components. For that matter, let

$$D_\lambda^2(P, P_0) := \langle \Upsilon_{P_0} g_\lambda(\Upsilon_{P_0}) u, u \rangle_{L^2(P_0)} = \sum_{i \geq 1} \lambda_i g_\lambda(\lambda_i) \langle u, \tilde{\phi}_i \rangle_{L^2(P_0)}^2, \quad (7)$$

where $g_\lambda : (0, \infty) \rightarrow (0, \infty)$, $\lambda > 0$ is a *spectral regularizer* (satisfies $\lim_{\lambda \rightarrow 0} x g_\lambda(x) \asymp 1$, i.e., $g_\lambda(x)$ is an approximation to x^{-1} with the approximation being controlled by λ), defined as

$$g_\lambda(\mathcal{C}) := \sum_{i \geq 1} g_\lambda(\tau_i) (\alpha_i \otimes_H \alpha_i) + g_\lambda(0) \left(\mathbf{I} - \sum_{i \geq 1} \alpha_i \otimes_H \alpha_i \right),$$

with \mathcal{C} being any compact, self-adjoint operator defined on a separable Hilbert space, H and $(\tau_i, \alpha_i)_i$ are the eigenvalues and eigenfunctions of \mathcal{C} . A common choice for regularization is the *Tikhonov regularizer*, $g_\lambda(x) = \frac{1}{x+\lambda}$, where $\lambda > 0$ is regularization parameter. By letting $\lambda \rightarrow 0$, $\frac{\lambda_i}{\lambda_i+\lambda} \rightarrow 1$, hence uniformly weighting the frequency components. Another choice we will use is by setting $g_\lambda(x) = \max\{1, (x+\lambda)^{-1}\}$ —we refer to this regularizer as *TikMax regularizer*—which only reweighs frequency components corresponding to $\lambda_i < 1 - \lambda$ and keeps the same weights corresponding to lower frequency components when $\lambda_i > 1 - \lambda$.

The results we present below hold for any spectral regularization function that satisfies the following sufficient conditions:

$$\begin{aligned}
(E_1) \quad & \sup_{x \in \Gamma} |xg_\lambda(x)| \leq C_1, & (E_2) \quad & \sup_{x \in \Gamma} |\lambda g_\lambda(x)| \leq C_2, \\
(E_3) \quad & \sup_{\{x \in \Gamma: xg_\lambda(x) < B_3\}} |B_3 - xg_\lambda(x)|x^{2\varphi} \leq C_3\lambda^{2\varphi}, & (E_4) \quad & \inf_{x \in \Gamma} g_\lambda(x)(x + \lambda) \geq C_4,
\end{aligned}$$

where $\Gamma := [0, \kappa]$, $\varphi \in (0, \xi]$, the constant ξ is called the *qualification* of g_λ , and $(C_i)_{i=1}^4$ and B_3 are finite positive constants (all independent of $\lambda > 0$). These are common conditions on the spectral regularizer employed in the inverse problem literature (see, for instance, Engl et al., 1996) and has also been employed in kernel ridge regression (Bauer et al., 2007) and kernel testing (Hagrass et al., 2023, 2022). The key condition is (E_3) which provides a quantitative estimate on the amplified version of the approximation error in approximating x^{-1} by $g_\lambda(x)$, and the qualification ξ imposes a bound on the amplification factor. It is easy to verify that both Tikhonov and TikMax regularizers have $\xi = \infty$.

The regularized KSD in (7) can be written in a form that is more suitable for estimation as

$$\begin{aligned}
D_\lambda^2(P, P_0) &= \langle \Upsilon_{P_0} g_\lambda(\Upsilon_{P_0})u, u \rangle_{L^2(P_0)} = \langle \mathfrak{J}\mathfrak{J}^* g_\lambda(\mathfrak{J}\mathfrak{J}^*)u, u \rangle_{L^2(P_0)} = \langle g_\lambda(\mathcal{A}_{P_0})\Psi_P, \Psi_P \rangle_{\mathcal{H}_{K_0}} \\
&= \left\| g_\lambda^{1/2}(\mathcal{A}_{P_0})\Psi_P \right\|_{\mathcal{H}_{K_0}}^2,
\end{aligned} \tag{8}$$

where

$$\mathcal{A}_{P_0} := \int_{\mathcal{X}} K_0(\cdot, x) \otimes_{\mathcal{H}_{K_0}} K_0(\cdot, x) dP_0(x), \quad \text{and} \quad \Psi_P := \int_{\mathcal{X}} K_0(\cdot, x) dP(x).$$

Based on the form in (8), note that a test statistic would involve estimating Ψ_P from $(X_i)_{i=1}^n$ and then would need either the complete knowledge of P_0 so that \mathcal{A}_{P_0} could be computed or access to independent samples $\mathbb{Y}_m := (Y_i^0)_{i=1}^m$ from P_0 so that \mathcal{A}_{P_0} can be estimated. However, this defeats the purpose of working with KSD and these alternate solutions have already been worked out in the case of MMD (Balasubramanian et al., 2021; Hagrass et al., 2023). Therefore, to exploit the power of KSD, we propose a variation to (8) so that this variation can be completely estimated based on $(X_i)_{i=1}^n$. To this end, define

$$\tilde{D}_\lambda^2(P, P_0) := \left\| g_\lambda^{1/2}(\mathcal{A}_P)\Psi_P \right\|_{\mathcal{H}_{K_0}}^2. \tag{9}$$

We argue that (9) is a valid approximation to (8) since if P is closer to P_0 , then it is clearly a good approximation. On the other hand, if P is far from P_0 , then the testing problem is easy and so P and P_0 could be distinguished without a significant influence from the regularization term.

To this end, we propose a test based on the following estimator of \tilde{D}_λ^2 as the test statistic,

$$\hat{\mathbb{S}}_\lambda^P(\mathbb{X}_n) = \frac{1}{n_1(n_1 - 1)} \sum_{1 \leq i \neq j \leq n_1} \left\langle g_\lambda^{1/2}(\hat{\mathcal{A}}_P)K_0(\cdot, X_i), g_\lambda^{1/2}(\hat{\mathcal{A}}_P)K_0(\cdot, X_j) \right\rangle_{\mathcal{H}_{K_0}}, \tag{10}$$

with

$$\hat{\mathcal{A}}_P := \frac{1}{n_2} \sum_{i=1}^{n_2} K_0(\cdot, Z_i) \otimes_{\mathcal{H}_{K_0}} K_0(\cdot, Z_i),$$

where the given samples $\mathbb{X}_n := (X_i)_{i=1}^n$ are divided into $(X_i)_{i=1}^{n_1}$ and $\mathbb{Z}_{n_2} := (Z_i)_{i=1}^{n_2} := (X_i)_{i=n_1+1}^n$, $n_2 = n - n_1$, with $(X_i)_{i=1}^{n_1}$ being used to estimate Ψ_P and \mathbb{Z}_{n_2} being used to estimate \mathcal{A}_P . The

advantage of the test statistic in (10) is that it is completely computable just using the given samples \mathbb{X}_n and knowing P_0 up to a normalization (as in Examples 2–4). Observe that under the null, we have $\mathbb{E}(\hat{\mathbb{S}}_\lambda^P) = \mathbb{E}(\hat{\mathbb{S}}_\lambda^{P_0}) = 0$. When $P \neq P_0$, but P closer to P_0 , we expect $\hat{\mathbb{S}}_\lambda^P(\mathbb{X}_n)$ to be closer to $\hat{\mathbb{S}}_\lambda^{P_0}(\mathbb{X}_n)$ (defined by replacing $\hat{\mathcal{A}}_P$ by \mathcal{A}_{P_0} or its estimator $\hat{\mathcal{A}}_{P_0}$) and therefore will inherit the minimax optimality of $\hat{\mathbb{S}}_\lambda^{P_0}(\mathbb{X}_n)$, as shown in Corollaries 6 and 7.

The following result (proved in Section 8.3) provides the computational form of the statistic in (10) and shows that it can be computed by solving a finite-dimensional eigensystem.

Theorem 3. *Let $(\hat{\lambda}_i, \hat{\alpha}_i)_i$ be the eigensystem of $\frac{1}{n_2}K_{n_2}$ where $K_{n_2} := [K_0(Z_i, Z_j)]_{i,j \in [n_2]}$. Define*

$$G := \sum_i \left(\frac{g_\lambda(\hat{\lambda}_i) - g_\lambda(0)}{\hat{\lambda}_i} \right) \hat{\alpha}_i \hat{\alpha}_i^\top.$$

Then

$$\hat{\mathbb{S}}_\lambda^P(\mathbb{X}_n) = \frac{1}{n_1(n_1 - 1)} \left(\mathbf{1}_{n_1}^\top M \mathbf{1}_{n_1} - \text{Tr}(M) \right),$$

where

$$M = g_\lambda(0)K_{n_1} + \frac{1}{n_2}K_{n_1, n_2} G K_{n_1, n_2}^\top,$$

with $K_{n_1} := [K_0(X_i, X_j)]_{i,j \in [n_1]}$ and $K_{n_1, n_2} := [K_0(X_i, Z_j)]_{i \in [n_1], j \in [n_2]}$.

The complexity of computing $\hat{\mathbb{S}}_\lambda^P(\mathbb{X}_n)$ is $O(n_2^3 + n_1^2 + n_2^2 n_1)$. It is worth noting that this represents a savings of a factor of order $m^2 + mn_2^2$ compared to the regularized MMD test statistic of Hagrass et al. (2023, Theorem 7), which involves sampling m i.i.d. samples from P_0 , and it also eliminates the time required for sampling those m samples.

To compute the testing threshold, we use the i.i.d. weighted bootstrap approach similar to that introduced for finding the testing threshold of \hat{D}_{KSD} as described in Section 3. Let

$$\hat{\mathbb{S}}_\lambda^\varepsilon(\mathbb{X}_n) := \frac{1}{n_1(n_1 - 1)} \sum_{i \neq j} \varepsilon_i \varepsilon_j \left\langle g_\lambda^{1/2}(\hat{\mathcal{A}}_P) K_0(\cdot, X_i), g_\lambda^{1/2}(\hat{\mathcal{A}}_P) K_0(\cdot, X_j) \right\rangle_{\mathcal{H}_{K_0}},$$

where $\varepsilon \in \{-1, 1\}^{n_1}$ is a random vector of n_1 i.i.d Rademacher random variables. The following result (proved in Section 8.4) shows that the distribution of $\hat{\mathbb{S}}_\lambda^\varepsilon(\mathbb{X}_n)$ can be used to approximate the distribution of $\hat{\mathbb{S}}_\lambda^P(\mathbb{X}_n)$. Thus, the $(1 - \alpha)$ quantile of $\hat{\mathbb{S}}_\lambda^\varepsilon(\mathbb{X}_n)$, defined as

$$q_{1-\alpha}^\lambda := \inf \left\{ q \in \mathbb{R} : P_\varepsilon \{ \hat{\mathbb{S}}_\lambda^\varepsilon(\mathbb{X}_n) \leq q | \mathbb{X}_n \} \geq 1 - \alpha \right\},$$

can be used as a test threshold (see Remark 1 for practical considerations).

Theorem 4. *Suppose $P = P_0$ and \mathbf{A}_0 holds. Let \mathcal{D}_1 be the distribution of $n_1 \hat{\mathbb{S}}_\lambda^\varepsilon(\mathbb{X}_n)$ conditioned on \mathbb{X}_n , and \mathcal{D}_2 be the distribution of $n_1 \hat{\mathbb{S}}_\lambda^P(\mathbb{X}_n)$ conditioned on \mathbb{Z}_{n_2} . Then for any $n_2 > 0$, $\lambda > 0$, \mathcal{D}_1 and \mathcal{D}_2 converge to the same distribution as $n_1 \rightarrow \infty$.*

Using the critical level $q_{1-\alpha}^\lambda$, in the next result (proved in Section 8.5), we investigate the conditions on the separation boundary of the test based on $\hat{\mathbb{S}}_\lambda^P(\mathbb{X}_n)$ so that the power is asymptotically one. To this end, we impose the following assumption on the pair (K_0, P_0) .

Assumption A2. $\mathbb{E}_{P_0}[K_0(X, X)]^v \leq \tilde{\kappa}^v$ for some $v > 2$, $\tilde{\kappa} > 0$.

Note that although \mathbf{A}_2 appears very similar to \mathbf{A}_1 , it is weaker than \mathbf{A}_1 . This means the behavior of the regularized KSD test captured in Theorem 5 is under the weaker condition of \mathbf{A}_2 in contrast to Theorem 3 which studies the behavior of the KSD test under \mathbf{A}_1 .

Define

$$\mathcal{N}_1(\lambda) := \text{Tr}(\mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{A}_{P_0} \mathcal{A}_{P_0, \lambda}^{-1/2}) \text{ and } \mathcal{N}_2(\lambda) := \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{A}_{P_0} \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_{K_0})},$$

which quantify the intrinsic dimensionality or degrees of freedom of \mathcal{H}_{K_0} . Notably, $\mathcal{N}_1(\lambda)$ plays a significant role in the analysis of kernel ridge regression, as discussed in works such as Caponnetto and Vito (2007).

Theorem 5. *Let $n_2 \asymp n$, $\tilde{\theta} := \min\{\theta, \xi\}$, $\Delta_n := \chi^2(P_n, P_0)$, $\sup_{P_n \in \mathcal{P}} \left\| \Upsilon_{P_0}^{-\theta} u \right\|_{L^2(P_0)} < \infty$, with $u := \frac{dP_n}{dP_0} - 1$. Suppose*

$$\lim_{n \rightarrow \infty} \Delta_n \lambda_n^{-2\tilde{\theta}} > d_1, \text{ for some } d_1 > 0, \quad (11)$$

and

$$\lim_{n \rightarrow \infty} \frac{n \Delta_n D_{\lambda_n}^{-1/2}}{1 + \sqrt{C_{\lambda_n} \Delta_n}} \rightarrow \infty, \quad (12)$$

where

$$C_\lambda := \begin{cases} \mathcal{N}_1(\lambda) \sup_i \|\phi_i\|_\infty^2, & \sup_i \|\phi_i\|_\infty^2 < \infty \\ (\mathcal{N}_1(\lambda))^{\frac{v-2}{2(v-1)}} \left(\frac{\kappa}{\lambda}\right)^{\frac{v}{v-1}}, & \text{otherwise} \end{cases}, \text{ and } D_\lambda = \begin{cases} \mathcal{N}_2^2(\lambda), & 4C_\lambda \Delta_n < 1 \\ \lambda^{-1}, & \text{otherwise} \end{cases}.$$

Then for $(P_n)_n \subset \mathcal{P}$, we have

$$\lim_{n \rightarrow \infty} P\{\hat{\mathbb{S}}_\lambda^{P_n}(\mathbb{X}_n) \geq q_{1-\alpha}^\lambda\} = 1$$

under either of the following conditions:

(i) \mathbf{A}_2 holds and

$$\lambda_n \sqrt{n} \rightarrow \infty. \quad (13)$$

Furthermore if $4C_{\lambda_n} \Delta_n < 1$, then (13) can be replaced by

$$\frac{n \lambda_n (\mathcal{N}_1(\lambda_n) + C_{\lambda_n} \sqrt{\Delta_n})^{-1}}{\log n} \rightarrow \infty. \quad (14)$$

(ii) $\sup_i \|\phi_i\|_\infty < \infty$ holds and

$$\frac{n \lambda_n}{\log n} \rightarrow \infty. \quad (15)$$

Furthermore if $4C_\lambda \Delta_n < 1$ and $\inf_{n>0} \mathbb{E}_{P_n} \phi_1^2 > 0$, then (15) can be replaced by

$$\frac{n}{\mathcal{N}_1(\lambda_n) \log \mathcal{N}_1(\lambda_n)} \rightarrow \infty. \quad (16)$$

The following corollaries to Theorem 5 offer explicit expressions for the separation boundary in the cases of polynomially and exponentially decaying eigenvalues of the operator \mathcal{A}_{P_0} . For the sake of simplified presentation, we will provide the results under the following assumption \mathbf{A}'_2 that is stronger than \mathbf{A}_2 , and the more general versions under \mathbf{A}_2 are deferred to Appendix B.

Assumption \mathbf{A}'_2 . $\mathbb{E}_{P_0}[K_0(X, X)]^v \leq \tilde{\kappa}^v$ for all $v > 2$, and some $\tilde{\kappa} > 0$.

Remark 5. (i) The difference between the assumptions \mathbf{A}'_2 and \mathbf{A}_2 lies in assuming that the required condition holds for all $v > 2$, rather than only for some v as assumed in \mathbf{A}_2 . Thus the result presented below is just the limiting result as v goes to infinity of the general result in Corollaries B.1 and B.2. Observe that \mathbf{A}'_2 is stronger than \mathbf{A}_1 , which is stronger than \mathbf{A}_2 .

(ii) Assuming g_λ possesses infinite qualification (examples include Tikhonov and TikMax regularizers), denoted by $\xi = \infty$, then $\tilde{\theta} = \theta$. Under \mathbf{A}'_2 , a comparison between Corollary 6 (or Corollary 7) and Theorem 2 indicates that the spectral regularized test, based on $\hat{\mathbb{S}}_\lambda^P(\mathbb{X}_n)$, achieves minimax optimality with respect to \mathcal{P} within the ranges of θ where the separation boundary of the test coincides with the minimax separation. For a more general scenario, we can align Corollary B.1 (or Corollary B.2) with Theorem 2 under \mathbf{A}_1 and observe that the test is again minimax optimal w.r.t. \mathcal{P} .

Corollary 6. Suppose $\lambda_i \lesssim i^{-\beta}$, $\beta > 1$. Let $\lambda_n \asymp \Delta_n^{\frac{1}{2\tilde{\theta}}}$, with $\Delta_n \rightarrow 0$. Then for $(P_n)_n \subset \mathcal{P}$, we have

$$\lim_{n \rightarrow \infty} P\{\hat{\mathbb{S}}_\lambda^{P_n} \geq q_{1-\alpha}^\lambda\} = 1,$$

if one of the following holds:

(i) \mathbf{A}'_2 holds and $\Delta_n B_n^{-1} \rightarrow \infty$, where for $\beta \geq \frac{3}{2}$

$$B_n = \begin{cases} n^{\frac{-4\tilde{\theta}\beta}{4\tilde{\theta}\beta+1}}, & \tilde{\theta} > \frac{2}{3} \\ \left(\frac{\log n}{n}\right)^{\frac{4\tilde{\theta}\beta}{1+4\beta-2\tilde{\theta}\beta}}, & \frac{1}{4\beta} + \frac{1}{2} \leq \tilde{\theta} \leq \frac{2}{3}, \\ n^{-\tilde{\theta}}, & \text{otherwise} \end{cases},$$

and for $\beta < \frac{3}{2}$

$$B_n = \begin{cases} n^{\frac{-4\tilde{\theta}\beta}{4\tilde{\theta}\beta+1}}, & \tilde{\theta} > \frac{1}{2} + \frac{1}{4\beta} \\ \left(\frac{\log n}{n}\right)^{\frac{2\tilde{\theta}\beta}{\beta+1}}, & 1 - \frac{1}{2\beta} \leq \tilde{\theta} \leq \frac{1}{2} + \frac{1}{4\beta}, \\ n^{-\tilde{\theta}}, & \text{otherwise} \end{cases},$$

with $\tilde{\theta} = \min\{\theta, \xi\}$.

(ii) $\sup_i \|\phi_i\|_\infty < \infty$ and $\Delta_n B_n^{-1} \rightarrow \infty$, where

$$B_n = \begin{cases} n^{\frac{-4\tilde{\theta}\beta}{4\tilde{\theta}\beta+1}}, & \tilde{\theta} \geq \frac{1}{2\beta} \\ n^{\frac{-4\tilde{\theta}\beta}{2\tilde{\theta}\beta+\beta+1}}, & \frac{1}{2} - \frac{1}{2\beta} \leq \tilde{\theta} \leq \frac{1}{2\beta}. \\ \left(\frac{\log n}{n}\right)^{2\tilde{\theta}}, & \text{otherwise} \end{cases}.$$

Corollary 7. Suppose $\lambda_i \lesssim e^{-\tau i}$, $\tau > 0$. Let $\lambda_n \asymp \Delta_n^{\frac{1}{2\tilde{\theta}}}$, with $\Delta_n \rightarrow 0$. Then for any $(P_n)_n \subset \mathcal{P}$, we have

$$\lim_{n \rightarrow \infty} P\{\hat{\mathbb{S}}_\lambda^{P_n} \geq q_{1-\alpha}^\lambda\} = 1,$$

if one of the following holds:

(i) \mathbf{A}'_2 holds and $\Delta_n B_n^{-1} \rightarrow \infty$, where

$$B_n = \begin{cases} \frac{\sqrt{\log n}}{n}, & \tilde{\theta} > \frac{2}{3} \\ \left(\frac{(\log n)^2}{n}\right)^{\frac{2\tilde{\theta}}{2-\tilde{\theta}}}, & \frac{1}{2} \leq \tilde{\theta} \leq \frac{2}{3}, \\ n^{-\tilde{\theta}}, & \text{otherwise} \end{cases}$$

with $\tilde{\theta} = \min\{\theta, \xi\}$.

(ii) $\sup_i \|\phi_i\|_\infty < \infty$ and $\Delta_n B_n^{-1} \rightarrow \infty$, where

$$B_n = \frac{\sqrt{\log n}}{n}.$$

4.1 Adaptation to λ by aggregation

Theorem 5 and Corollaries 6, 7 demonstrate that the optimal λ needed to achieve the minimax separation boundary depends on the unknown parameters, θ (and β in the case of polynomial decay). To make the test practical, the idea is to devise an adaptive test by aggregating multiple tests constructed for various λ taking values in a finite set, Λ . Using similar analysis as in Hagrass et al. (2023, 2022); Balasubramanian et al. (2021) and Schrab et al. (2023), the aggregate test can be shown to yield the optimal separation boundary as shown in Corollaries 6 and 7, but up to a log log factor. In the following, we briefly introduce the aggregate test but skip stating the corresponding theoretical results since they are exactly the same as Corollaries 6 and 7 but with an additional log log factor.

Define $\Lambda := \{\lambda_L, 2\lambda_L, \dots, \lambda_U\}$, where $\lambda_U = 2^b \lambda_L$ for $b \in \mathbb{N}$, such that $|\Lambda| = b + 1 = 1 + \log_2 \frac{\lambda_U}{\lambda_L}$, with $|\Lambda|$ denoting the cardinality of Λ . Let λ^* be the optimal λ for achieving minimax optimality. The crux lies in selecting λ_L and λ_U so that an element in Λ closely approximates λ^* for any θ (and β in the case of polynomially decaying eigenvalues). Define $v^* := \sup\{x \in \Lambda : x \leq \lambda^*\}$. Then, it is straightforward to verify that $v^* \asymp \lambda^*$, meaning v^* also represents an optimal choice for λ within Λ , as for $\lambda_L \leq \lambda^* \leq \lambda_U$, we have $\frac{\lambda^*}{2} \leq v^* \leq \lambda^*$. Motivated by this observation, we construct an adaptive test based on the union of corresponding tests across $\lambda \in \Lambda$, rejecting H_0 if any of the tests does so. Furthermore, we can also adapt across different choices of K , or the kernel bandwidth (or kernel parameter) of a given K , e.g., $K(x, y) = e^{-\|x-y\|_2^2/h}$, $h > 0$. Thus, the resulting test rejects H_0 if $\hat{\mathbb{S}}_\lambda^{P,h}(\mathbb{X}_n) \geq q_{1-\frac{\alpha}{|\Lambda||W|}}^{\lambda,h}$ for any $(\lambda, h) \in \Lambda \times W$, where $\hat{\mathbb{S}}_\lambda^{P,h}(\mathbb{X}_n)$ is the test statistic as defined in (10).

We would like to mention that recently, Schrab et al. (2022) proposed an adaptive version of the KSD test of Chwialkowski et al. (2016) and Liu et al. (2016)—see Example 2—by adapting to kernel parameter of K (here, K is chosen to be the Gaussian kernel with bandwidth h as shown above) by aggregating over many KSD tests constructed for a given bandwidth. They obtained the separation boundary for the aggregate test over restricted Sobolev balls. However, they did not investigate the minimax separation rates for such alternative spaces, leaving open the question of the minimax optimality of their rates.

5 Interpreting the class of alternatives, \mathcal{P}

As aforementioned, the alternative class, \mathcal{P} , involves a smoothness condition defined through $\text{Ran}(\Upsilon_{P_0}^\theta)$, where Υ_{P_0} is based on K_0 and P_0 . Since \mathcal{P} is abstract, in this section, we provide

an interpretation for \mathcal{P} by analyzing the smoothness condition through some concrete examples. First, we present a generic result about $\text{Ran}(\Upsilon_{P_0}^\theta)$ which will be used in all the examples.

Proposition 8. *Suppose*

$$K_0(x, y) = \sum_{k \in I} \lambda_k \gamma_k(x) \gamma_k(y),$$

where $1 \cup (\gamma_k)_{k \in I}$ represent orthonormal basis with respect to $L^2(P_0)$. Then

$$\text{Ran}(\Upsilon_{P_0}^\theta) = \left\{ \sum_{k \in I} a_k \gamma_k(x) : \sum_{k \in I} \frac{a_k^2}{\lambda_k^{2\theta}} < \infty \right\},$$

where $\Upsilon_{P_0} : L^2(P_0) \rightarrow L^2(P_0)$, $f \mapsto \int K_0(\cdot, x) f(x) dP_0(x)$.

Based on the above results, we now discuss a few examples.

Example 5 (Uniform distribution on $[0, 1]$). *Let P_0 be the uniform distribution defined on $[0, 1]$,*

$$K(x, y) = a_0 + \sum_{k \neq 0} |k|^{-\beta} e^{\sqrt{-1}2\pi kx} e^{-\sqrt{-1}2\pi ky}, \quad a_0 \geq 0, \quad \beta > 2,$$

and $K_0(x, y) := \langle \psi_{P_0}(\cdot, x), \psi_{P_0}(\cdot, y) \rangle_{\mathcal{H}}$. Suppose $\psi_{P_0}(\cdot, x) = K(\cdot, x) - \mathbb{E}_{P_0} K(\cdot, X)$. Then

$$R_1 := \text{Ran}(\Upsilon_{P_0}^\theta) = \left\{ \sum_{k \neq 0} a_k e^{\sqrt{-1}2\pi kx} : \sum_{k \neq 0} a_k^2 k^{2\theta\beta} < \infty \right\}.$$

On the other hand, if $\psi_{P_0}(\cdot, x) = K(\cdot, x) \nabla_x \log p_0(x) + \nabla_x K(\cdot, x)$, i.e., K_0 is the Stein kernel, then

$$R_2 := \text{Ran}(\Upsilon_{P_0}^\theta) = \left\{ \sum_{k \neq 0} a_k e^{\sqrt{-1}2\pi kx} : \sum_{k \neq 0} a_k^2 k^{2\theta(\beta-2)} < \infty \right\}.$$

See Section 8.7 for the derivation of R_1 and R_2 . Note that R_1 and R_2 correspond to the range space condition for MMD and KSD tests, respectively. This means that the functions in R_2 are 2θ less smooth than those in R_1 since the Fourier coefficients in R_2 decay at a rate that is 2θ orders slower than those of R_1 . Moreover, since $\sum_{k \neq 0} a_k^2 k^{2\theta(\beta-2)} \leq \sum_{k \neq 0} a_k^2 k^{2\theta\beta}$, for $\beta > 2$, we have $R_1 \subseteq R_2$. Observe that the eigenfunctions satisfy the uniform boundedness condition as $\sup_k |e^{\sqrt{-1}2\pi kx}| = 1$. For $\beta > 3$, it can be demonstrated that both the base kernel K and the Stein kernel K_0 are bounded. Consequently, the entire alternative space \mathcal{P}_Δ associated with the Stein kernel (i.e., the one defined through the condition $u \in R_2$) encompasses the alternative space associated with the base kernel (i.e., the one defined through the condition $u \in R_1$). Moreover, the minimax separation boundary linked with the alternative space defined through the base kernel follows an order of $n^{\frac{-4\theta\beta}{4\theta\beta+1}}$. Meanwhile, the larger alternative space associated with the Stein kernel has a minimax separation boundary order of $n^{\frac{-4\theta\tilde{\beta}}{4\theta\tilde{\beta}+1}}$, which is achieved by our proposed regularized KSD test, where $\tilde{\beta} = \beta - 2$. This adjustment arises from the corresponding eigenvalue decay rate of the Stein kernel, which holds an order of $|k|^{-(\beta-2)}$.

Remark 6. (i) The s -order Sobolev space defined on $[0, 1]$ is given by

$$\mathcal{W}^{s,2} := \left\{ f(x) = \sum_{k \in \mathbb{Z}} a_k e^{i2\pi kx}, \quad x \in [0, 1] : \sum_{k \in \mathbb{Z}} (1 + k^2)^s a_k^2 < \infty \right\}.$$

Since $\sum_{k \neq 0} k^{2\theta\beta} a_k^2 \leq \sum_{k \in \mathbb{Z}} (1+k^2)^{\theta\beta} a_k^2$, it follows that $\mathcal{W}^{s,2} \subset R_1$. This means, if $u := \frac{dP}{dP_0} - 1 \in \mathcal{W}^{s,2}$, then $u \in R_1$ with $\theta = \frac{s}{\beta}$.

(ii) Note that for $d \geq 2$, we can generalize the above example by using a product of kernels over each dimension, which yields

$$R_1 := \left\{ \sum_{k_1, \dots, k_d} a_{k_1, \dots, k_d} \gamma_{k_1, \dots, k_d} : \sum_{k_1, \dots, k_d} a_{k_1, \dots, k_d}^2 (k_1 k_2 \dots k_d)^{2\theta\beta} < \infty \right\},$$

and

$$R_2 := \left\{ \sum_{k_1, \dots, k_d} a_{k_1, \dots, k_d} \gamma_{k_1, \dots, k_d} : \sum_{k_1, \dots, k_d} a_{k_1, \dots, k_d}^2 (k_1^2 + \dots + k_d^2)^{-2\theta} (k_1 k_2 \dots k_d)^{2\theta\beta} < \infty \right\},$$

where $\gamma_{k_1, \dots, k_d}(x) = \prod_{j=1}^d e^{\sqrt{-1}2\pi k_j x_j}$, $x = (x_1, \dots, x_d) \in \mathbb{R}^d$ and for $\beta > \frac{2}{d}$, $R_1 \subset R_2$. Moreover observe that

$$\sum_{k_1, \dots, k_d} a_{k_1, \dots, k_d}^2 (k_1 k_2 \dots k_d)^{2\theta\beta} \leq \sum_{k_1, \dots, k_d} a_{k_1, \dots, k_d}^2 (1 + k_1^2 + \dots + k_d^2)^{d\theta\beta},$$

which yields that $u \in \mathcal{W}^{s,2}$ implies $u \in R_1$ for $\theta = \frac{s}{d\beta}$.

Example 6 (Gaussian distribution with Mehler kernel on \mathbb{R}). Let P_0 be a standard Gaussian distribution on \mathbb{R} and K be the Mehler kernel, i.e.,

$$K(x, y) := \frac{1}{\sqrt{1-\rho^2}} \exp\left(-\frac{\rho^2(x^2 + y^2) - 2\rho xy}{2(1-\rho^2)}\right),$$

for $0 < \rho < 1$. Define $K_0(x, y) := \langle \psi_{P_0}(\cdot, x), \psi_{P_0}(\cdot, y) \rangle_{\mathcal{H}}$, where $\psi_{P_0}(\cdot, x) = K(\cdot, x) - \mathbb{E}_{P_0} K(\cdot, X)$. Then

$$R_1 := \text{Ran}(\Upsilon_{P_0}^\theta) = \left\{ \sum_{k=1}^{\infty} a_k \gamma_k(x) : \sum_{k=1}^{\infty} a_k^2 e^{-2k\theta \log \rho} < \infty \right\},$$

where $\gamma_k(x) = \frac{H_k(x)}{\sqrt{k!}}$, and $H_k(x) = (-1)^k e^{x^2/2} \frac{d^k}{dx^k} e^{-x^2/2}$ is the Hermite polynomial. Furthermore, if $\psi_{P_0}(\cdot, x) = K(\cdot, x) \nabla_x \log p_0(x) + \nabla_x K(\cdot, x)$, then

$$R_2 := \text{Ran}(\Upsilon_{P_0}^\theta) = \left\{ \sum_{k=1}^{\infty} a_k \gamma_k(x) : \sum_{k=1}^{\infty} a_k^2 k^{-2\theta} e^{-2k\theta \log \rho} < \infty \right\}.$$

Note that R_1 and R_2 correspond to the range space condition for MMD and KSD tests, respectively. Since $\sum_{k=1}^{\infty} a_k^2 k^{-2\theta} e^{-2k\theta \log \rho} \leq \sum_{k=1}^{\infty} a_k^2 e^{-2k\theta \log \rho}$, we have $R_1 \subseteq R_2$. See Section 8.8 for details.

Remark 7. (i) Let $\tilde{K}(x, y) := \phi(x-y) := \exp(-\sigma(x-y)^2)$, for $\sigma < \frac{\rho}{4(1-\rho)}$. Then the corresponding RKHS $\mathcal{H}_{\tilde{K}}$ is the Gaussian RKHS, defined by

$$\mathcal{H}_{\tilde{K}} = \left\{ f \in L^2(\mathbb{R}) : \int_{\mathbb{R}} |\hat{f}(\omega)|^2 e^{\frac{\omega^2}{4\sigma}} d\omega < \infty \right\},$$

where \hat{f} is the Fourier transform of $f \in L^2(\mathbb{R})$. It can be shown that $\mathcal{H}_{\tilde{K}} \subset R_1$ for any $\theta \leq \frac{1}{2}$ (see Section 8.9 for details). Therefore, if $u := \frac{dP}{dP_0} - 1 \in \mathcal{H}_{\tilde{K}}$, then $u \in R_1 \subset R_2$, for any $\theta \leq \frac{1}{2}$.

(ii) For $d \geq 2$, the above example can be generalized by using the product of kernels over each dimension. This yields that

$$R_1 := \left\{ \sum_{k_1, \dots, k_d} a_{k_1, \dots, k_d} \gamma_{k_1, \dots, k_d} : \sum_{k_1, \dots, k_d} a_{k_1, \dots, k_d}^2 e^{-2(k_1 + \dots + k_d)\theta \log \rho} < \infty \right\},$$

and

$$R_2 := \left\{ \sum_{k_1, \dots, k_d} a_{k_1, \dots, k_d} \gamma_{k_1, \dots, k_d} : \sum_{k_1, \dots, k_d} a_{k_1, \dots, k_d}^2 (k_1 + \dots + k_d)^{-2\theta} e^{-2(k_1 + \dots + k_d)\theta \log \rho} < \infty \right\},$$

where $\gamma_{k_1, \dots, k_d}(x) = \prod_{j=1}^d \gamma_{k_j}(x_j)$, $x = (x_1, \dots, x_d) \in \mathbb{R}^d$.

Example 7 (Uniform distribution on \mathbb{S}^2). Let P_0 be a uniform distribution on $\mathcal{X} = \mathbb{S}^2$, where \mathbb{S}^2 denotes the unit sphere. Let

$$K_0(x, y) := \sum_{k=1}^{\infty} \sum_{j=-k}^k \lambda_k Y_k^j(\theta_x, \phi_x) Y_k^j(\theta_y, \phi_y), \quad (17)$$

where $x = (\sin \theta_x \cos \phi_x, \sin \theta_x \sin \phi_x, \cos \theta_x)$, $y = (\sin \theta_y \cos \phi_y, \sin \theta_y \sin \phi_y, \cos \theta_y)$ with $0 < \theta_x, \theta_y < \pi$, $0 < \phi_x, \phi_y < 2\pi$, and

$$Y_k^j(\theta, \phi) := \sqrt{\frac{(2k+1)(k-j)!}{4\pi(k+j)!}} p_k^j(\cos \theta) e^{\sqrt{-1}j\phi},$$

with $p_k^j(x) := (-1)^j (1-x^2)^{\frac{j}{2}} \frac{d^j p_k(x)}{dx^j}$ and $p_k(x) := \frac{1}{k!2^k} \frac{d^k (x^2-1)^k}{dx^k}$. If $\sum_{k=1}^{\infty} (2k+1)\lambda_k < \infty$, then

$$\text{Ran}(\Upsilon_{P_0}^\theta) := \left\{ \sum_{k=1}^{\infty} \sum_{j=-k}^k a_{kj} Y_k^j(\theta_x, \phi_x) : \sum_{k=1}^{\infty} \sum_{j=-k}^k a_{kj}^2 \lambda_k^{-2\theta} < \infty \right\}.$$

Remark 8. (i) In Example 7, we directly assume a form for the Stein kernel K_0 , in contrast to Examples 5 and 6, where we first express a base kernel K , and then compute the corresponding Stein kernel K_0 . Therefore, the claim in Example 7 follows directly from Proposition 8. On the other hand, suppose we choose (17) as the base kernel and then apply the appropriate Stein transformation as outlined in Example 3, the transformed eigenfunctions lose orthogonality to P_0 , making it challenging to derive an explicit expression for $\text{Ran}(\Upsilon_{P_0}^\theta)$ using Proposition 8. Conversely, in Examples 5 and 6, the transformation only alters the eigenvalues while preserving the eigenfunctions of the base kernel.

(ii) For more details about kernels that take the form in (17), we refer the reader to Minh et al. (2006). For example, Minh et al. (2006, Theorem 2 and 3) provides explicit expressions for the eigenvalues corresponding to Gaussian and polynomial kernels on the sphere.

The examples above have focused on the smoothness condition when \mathcal{X} is continuous. Suppose now that \mathcal{X} is a finite set with cardinality $|\mathcal{X}|$, on which P_0 and K_0 are supported—refer, for example, to Yang et al. (2018) for a definition of the Stein kernel K_0 for discrete distributions.

Let $\mathcal{X} = \{a_1, \dots, a_{|\mathcal{X}|}\}$ and define $P_0 = \sum_{i=1}^{|\mathcal{X}|} w_i \delta_{a_i}$ where $w_i > 0$ for all i . Then it is easy to verify that $L^2(P_0)$ is isomorphic to $\mathbb{R}^{|\mathcal{X}|}$ since $\|f\|_{L^2(P_0)}^2 = \sum_{i=1}^{|\mathcal{X}|} w_i f^2(a_i)$. Consider $(\Upsilon_{P_0} f)(a_j) = \sum_{i=1}^{|\mathcal{X}|} w_i K_0(a_j, a_i) f(a_i)$, which implies that Υ_{P_0} can be identified with a $|\mathcal{X}| \times |\mathcal{X}|$ matrix. Then $u \in \text{Ran}(\Upsilon_{P_0})$, implies that there exists $h \in \mathbb{R}^{|\mathcal{X}|}$ such that $u = \Upsilon_{P_0} h = \mathbf{K}_0 \text{diag}(w_1, \dots, w_{|\mathcal{X}|}) h$, where $[\mathbf{K}_0]_{ij} = K_0(a_i, a_j)$, $i, j \in [|\mathcal{X}|]$. Therefore, if \mathbf{K}_0 is invertible, then it is clear that there always exists $h \in \mathbb{R}^{|\mathcal{X}|}$ such that $u = \Upsilon_{P_0} h$. Hence, the range space condition is always satisfied since $\text{Ran}(\Upsilon_{P_0}^\theta) = \text{Ran}(\Upsilon_{P_0}) = \mathbb{R}^{|\mathcal{X}|}$ for all $\theta > 0$. This means the alternative space is solely defined by the χ^2 separation condition. A more interesting scenario is when \mathcal{X} is countable, in which case $L^2(P_0)$ is isomorphic to the sequence space, $\ell^2(\mathcal{X})$. We leave a detailed study of interpreting the range space condition in this setting for future work.

6 Experiments

In this section, we conduct an empirical evaluation and comparison of various goodness-of-fit tests:

- $KSD(\text{Tikhonov})$: An adaptive spectral regularized KSD test with a Tikhonov regularizer.
- $KSD(\text{TikMax})$: An adaptive spectral regularized KSD test with a TikMax regularizer.
- $KSD(\text{Tikhonov})^*$: An adaptive spectral regularized KSD test with a Tikhonov regularizer based on an estimator of the regularized discrepancy in (8). The test statistic is similar to that of (10) but with $\hat{\mathcal{A}}_P$ being replaced by $\hat{\mathcal{A}}_{P_0}$, where $\hat{\mathcal{A}}_{P_0}$ is computed based on extra samples from P_0 .
- $KSD(\text{TikMax})^*$: An adaptive spectral regularized KSD test constructed similar to that of $KSD(\text{Tikhonov})^*$ but with a TikMax regularizer.
- KSD : An unregularized KSD test with the median heuristic bandwidth for the kernel parameter.
- $MMD(\text{Tikhonov})$: An adaptive spectral regularized MMD test (as implemented in Hagraas et al. 2023), which uses additional samples from P_0 to compute the test statistic.

Remark 9. $KSD(\text{Tikhonov})^*$ and $KSD(\text{TikMax})^*$ are being considered to compare the performance of the regularized test when using $\hat{\mathcal{A}}_P$ versus $\hat{\mathcal{A}}_{P_0}$ as the estimated covariance operator. Recall that in our proposed test, we opted for $\hat{\mathcal{A}}_P$ over $\hat{\mathcal{A}}_{P_0}$ to create a practical test that circumvents the requirement for additional samples from P_0 .

All these tests are conducted with both Gaussian and inverse multiquadric (IMQ) kernels defined as $K(x, y) = \exp\left(-\frac{1}{2h}\|x - y\|_2^2\right)$ and $K(x, y) = \left(1 + \frac{\|x - y\|_2^2}{h}\right)^{-0.5}$, respectively. For our proposed KSD adaptive tests, we aggregate tests jointly over $\lambda \in \Lambda$ and $h \in W$ (refer to Section 4.1 for details). Unless otherwise specified, all experiments utilize $\Lambda := \{10^{-4}, 10^{-2.5}, 10^{-1}, 10^{0.5}\}$ and $W := \{10^{-2}, 10^{-1.5}, 10^{-1}, 10^{-0.5}, 1, 10^{0.5}\} \times h_m$, where h_m represents the kernel bandwidth determined using the median heuristic defined by $h_m := \text{median}\{\|q - q'\|_2^2 : q, q' \in X \cup Y\}$. Each test is repeated 500 times, and the average power is reported. For all KSD tests, we employed the i.i.d. weighted bootstrap to compute the test threshold with the number of bootstrap samples being $B = 1000$. On the other hand, for the $MMD(\text{Tikhonov})$ test, we used the same values for

Λ and W as mentioned above, along with 150 permutations to compute the test threshold (note that the MMD test is a permutation test).

We examined the performance of these tests across various domains, including \mathbb{R}^d , \mathbb{S}^{d-1} , and infinite-dimensional Hilbert space. Depending on the domain \mathcal{X} , we employed the appropriate Stein operator to compute the kernel Stein discrepancy. Of course, these tests are applied in scenarios where P_0 is known only up to a normalization constant. In Section 6.1, we applied the KSD method of Chwialkowski et al. (2016) and Liu et al. (2016) (see Example 2 for details) to a Gaussian-Bernoulli restricted Boltzmann machine. In Section 6.2, we utilized the directional Stein discrepancy (dKSD) introduced by Xu and Matsuda (2020) (see Example 3 for details) on directional data, and we employed the infinite-dimensional version of KSD as described in Wynne et al. (2023) (see Example 4 for details) on Brownian motion in Section 6.3.

The main takeaway from the experiments could be summarized as follows:

- Overall, the proposed regularized KSD tests, consistently outperform the standard unregularized KSD test.
- Generally, when independent samples from P_0 are readily available, MMD(Tikhonov) demonstrates the best performance, albeit at the highest computational cost. Even in this setting, the regularized KSD tests are competitive and achieve performance matching that of MMD(Tikhonov) in some cases.
- The proposed regularized versions of KSD with respect to $\hat{\mathcal{A}}_P$ and $\hat{\mathcal{A}}_{P_0}$ (KSD*), exhibit similar performance in most cases, with KSD* slightly outperforming the regularized KSD. However, note that KSD* has the benefit of using additional samples from P_0 .

6.1 Gaussian-Bernoulli restricted Boltzmann machine

As considered in Liu et al. (2016) and Schrab et al. (2022), we investigate a Gaussian-Bernoulli restricted Boltzmann machine, a graphical model characterized by a binary hidden variable $h \in \{-1, 1\}^{d_h}$ and a continuous observable variable $x \in \mathbb{R}^d$. These variables share a joint density function defined as

$$f(x, h) = \frac{1}{Z} \exp \left(\frac{1}{2} x^\top B h + b^\top x + c^\top h - \frac{1}{2} \|x\|_2^2 \right),$$

where Z represents an unknown normalization constant. The target density is obtained as

$$\frac{dP_0}{dx} = p_0(x) = \sum_{h \in \{-1, 1\}^{d_h}} f(x, h).$$

We adopt the same settings and implementation as described in Schrab et al. (2022), where $d = 50$, and $d_h = 40$. Note that p_0 is known only up to a normalization constant and the score function admits a closed-form expression:

$$\nabla \log p_0(x) = b - x + B \frac{\exp(2(B^\top x + c)) - 1}{\exp(2(B^\top x + c)) + 1}.$$

Here b and c are drawn from a standard Gaussian, and the elements in B are selected randomly from $\{-1, 1\}$. The alternate distribution P is constructed similarly to P_0 but with each element of




Figure 1: Power of the tests for Gaussian-Bernoulli restricted Boltzmann machine with $d = 50$, $n_2 = 100$ and $n = 1000$.

B being perturbed by Gaussian noise, $N(0, \sigma)$ with $\sigma \in \{0.01, 0.02, 0.03\}$. Differentiating P from P_0 becomes more challenging as σ becomes small.

We draw $n = 1000$ samples from P using a Gibbs sampler, using which all aforementioned KSD tests (except KSD* and MMD) are constructed with $n_2 = 100$. To construct KSD* and MMD tests, we also sample 100 and 1000 samples, respectively, from P_0 using a Gibbs sampler. Figure 1 illustrates the power of the tests under consideration. We note that KSD(Tikhonov)* and KSD(TikMax)* yield matching results and perform the best while being closely followed by our proposed tests KSD(Tikhonov) and KSD(TikMax), which strictly dominate the KSD test. Surprisingly, the MMD(Tikhonov) exhibits the worst performance in this case, despite having access to extra samples from P_0 . This unexpected outcome may be attributed to the nature of the experiment, where the samples from P_0 are obtained using a Gibbs sampler, which might only yield approximate samples or samples that are not independent, while the MMD(Tikhonov) requires extra independent samples from P_0 .

6.2 Directional data

The density for Watson distribution on \mathbb{S}^{d-1} is given by $f(x, \mu, k) = \frac{\Gamma(d/2)}{2\pi^{d/2}M(1/2, d/2, k)} \exp(k(\mu^T x)^2)$, where $x \in \mathbb{S}^{d-1}$, $k \geq 0$ is the concentration parameter, μ is the mean parameter, and M denotes the Kummer's confluent hypergeometric function. In this experiment, we consider a mixture of two multivariate Watson distributions, representing axially symmetric distributions on a sphere. Note that $k = 0$ corresponds to a uniform distribution on \mathbb{S}^{d-1} , which is chosen to be P_0 for this experiment. The alternate distributions are chosen to be a mixture of two Watson distributions with similar concentration parameters k and mean parameters μ_1, μ_2 respectively, where $\mu_1 = (1, \dots, 1) \in \mathbb{R}^d$ and $\mu_2 = (-1, 1, \dots, 1) \in \mathbb{R}^d$ with the first coordinate changed to -1 , where changing k creates a sequence of alternatives. The smaller the k is, the more difficult is to distinguish between P_0 and P .

We generate $n = 500$ samples from P (for different values of k) and construct all tests (except KSD* and MMD) based on them with $n_2 = 100$. We use additionally generated samples of size 100 and 500 from P_0 to construct KSD* and MMD tests, respectively. Figure 2 presents the outcomes




Figure 2: Power for mixture of Watson distributions for different concentration parameter k using $n_2 = 100$ and $n = 500$.

of testing against a spherical uniform distribution across various concentration parameters. With both Gaussian and IMQ kernels, we note that MMD(Tikhonov) demonstrates the most optimal performance, followed by KSD* versions, which can be attributed to its access to extra independent samples from the null distribution. Our proposed tests KSD(Tikhonov) and KSD(TikMax) behave similarly and better than unregularized KSD with IMQ while KSD(Tikhonov) has similar performance to that of KSD with the Gaussian kernel.

Also, KSD(TikMax) outperforms both the unregularized KSD test and the KSD(Tikhonov) test. Interestingly, using the TikMax regularizer generally provides better results than using the Tikhonov regularizer for the Gaussian kernel. This observation underscores the fact that the choice of regularizer and kernel can impact performance in practice.

6.3 Brownian motion

In this section, we adopt the setup outlined in Wynne et al. (2023), which considers $\mathcal{X} = L^2[0, 1]$ and $P_0 = N_C$, a Brownian motion over $[0, 1]$ with covariance operator C that has eigenvalues $\gamma_i = (i - \frac{1}{2})^{-2}\pi^{-2}$ and eigenfunctions $e_i(t) = \sqrt{2}\sin((i - \frac{1}{2})\pi t)$, $t \in [0, 1]$. The following alternate distributions are considered in the experiments, which are variations of those considered in Wynne et al. (2023). Let B_t denote the standard Brownian motion.

- P follows the law of $X_t = (1 + \sin(2\delta\pi t))B_t$, $\delta \in \{0.15, 0.1, 0.05, 0.01\}$.
- P is the law of the Brownian motion clipped to k frequencies, $k \in [4]$, i.e., $\sum_{i=1}^k \sqrt{\gamma_i}\xi_i e_i$, where $\xi_i \stackrel{i.i.d}{\sim} N(0, 1)$.

Note that δ and k generate a sequence of alternatives in the above cases with small δ and large k making P difficult to distinguish from P_0 . For our proposed regularized tests, we use $\Lambda = \{0.01, 0.05, 1\}$ and $W = \{h_m\}$, where h_m denotes the median bandwidth. We use additionally generated samples of size n_2 and n (see the captions of Figures 3 and 4 for details) from P_0 to construct KSD* and MMD tests. We generated the samples using the settings and code provided in Wynne et al. (2023). In this setup, function samples were observed at 100 points uniformly




Figure 3: Power for different values for δ using $n_2 = 35$ and $n = 70$.

distributed across the interval $[0,1]$, and 100 basis functions of the Brownian motion were used for the numerical approximation of X_t .

Figure 3 shows the power of the first experiment for different values of δ with $n_2 = 35$ and $n = 70$. In this experiment, the best performance is achieved by $\text{KSD}(\text{Tikhonov})^*$ and $\text{MMD}(\text{Tikhonov})$, which have access to extra samples, followed by $\text{KSD}(\text{Tikhonov})$, with matching power to $\text{KSD}(\text{Tikhonov})^*$ at $\delta = 0.01$. In this scenario, Tikhonov regularization outperforms TikMax regularization, which can be explained by the fact that X_t is a perturbed version of B_t with high-frequency components being added. Thus, Tikhonov regularization is a more appropriate choice in this case.

Figure 4 illustrates the power of the second experiment with different values of k with $n_2 = 100$ and $n = 200$. In this experiment, once again, the best performance is achieved by $\text{MMD}(\text{Tikhonov})$, followed by the regularized KSD methods with Tikhonov regularization, which outperforms the unregularized KSD and $\text{KSD}(\text{TikMax})$. This can be explained by the fact that X_t differs from B_t at higher frequency components, making the Tikhonov regularizer a more appropriate choice in this case too.

7 Discussion

To summarize, we studied the minimax optimality of goodness-of-fit tests across general domains, using the kernelized Stein discrepancy (KSD) framework—a versatile tool for conducting goodness-of-fit testing that can accommodate diverse data structures beyond Euclidean spaces and operate efficiently with only partial knowledge of the reference distribution. We formulated a comprehensive framework and provided an operator-theoretic representation of KSD, encompassing various existing KSD tests in the literature, each tailored to specific domains. Despite its versatility, we established the non-optimality of the KSD test w.r.t. a suitable class of alternatives with χ^2 -distance as the separation metric. To address this non-optimality issue, we proposed a spectral regularized version of the KSD test and showed it to be minimax optimal. Furthermore, through numerical experiments, we demonstrated the superior performance of our proposed tests across various domains compared to the unregularized KSD test.




Figure 4: Power for different values of k using $n_2 = 100$ and $n = 200$.

There are several promising directions for future research. First, extending our results to handle composite null hypotheses would broaden the applicability of the proposed tests. Next, while our framework encompasses many KSD variations defined across different domains, there remains room for further investigation in terms of interpreting the smoothness condition (defined through the range space of an integral operator) for various specific domains and null distributions. Examples include discrete distributions supported on countable domain and network data, which can be addressed within the proposed general framework but require additional careful examination for any required domain-specific adjustments. Moreover, exploring alternative methods of adaptation could enhance the test performance. For instance, investigating approaches that provide tighter bounds than the conservative union bound, such as studying the asymptotic distribution of the adapted test, holds promise for further enhancing the test power.

8 Proofs

In this section, we present the proofs of all the main results of the paper.

8.1 Proof of Theorem 1

First we will show that if $D_{\text{KSD}}^2(P_n, P_0) = \|\Psi_{P_n}\|_{\mathcal{H}_{K_0}}^2 = o(n^{-1})$, then the desired result holds, where $\Psi_P := \mathbb{E}_P K_0(\cdot, X)$. Observe that

$$\begin{aligned}
\hat{D}_{\text{KSD}}^2 &= \frac{1}{n(n-1)} \sum_{i \neq j} K_0(X_i, X_j) \\
&= \frac{1}{n(n-1)} \sum_{i \neq j} \langle K_0(\cdot, X_i) - \Psi_{P_n}, K_0(\cdot, X_j) - \Psi_{P_n} \rangle_{\mathcal{H}_{K_0}} + \frac{2}{n} \sum_{i=1}^n \langle K_0(\cdot, X_i), \Psi_{P_n} \rangle_{\mathcal{H}_{K_0}} - \|\Psi_{P_n}\|_{\mathcal{H}_{K_0}}^2 \\
&= \frac{1}{n(n-1)} \sum_{i \neq j} \langle K_0(\cdot, X_i) - \Psi_{P_n}, K_0(\cdot, X_j) - \Psi_{P_n} \rangle_{\mathcal{H}_{K_0}} + \frac{2}{n} \sum_{i=1}^n \langle K_0(\cdot, X_i) - \Psi_{P_n}, \Psi_{P_n} \rangle_{\mathcal{H}_{K_0}} \\
&\quad + \|\Psi_{P_n}\|_{\mathcal{H}_{K_0}}^2
\end{aligned}$$

$$:= \textcircled{1} + \textcircled{2} + \|\Psi_{P_n}\|_{\mathcal{H}_{K_0}}^2.$$

Then under H_1 , i.e., $(X_i)_i \stackrel{i.i.d.}{\sim} P_n \in \mathcal{P}$, using standard arguments for the convergence of U-statistics (see Serfling, 1980, Section 5.5.2) yields that

$$n \textcircled{1} \rightsquigarrow \sum_{i=1}^n \lambda_i (a_i^2 - 1) := S,$$

where $a_i \stackrel{i.i.d.}{\sim} \mathcal{N}(0, 1)$. Next, note that

$$n \textcircled{2} = 2\sqrt{n}\mathcal{V}_n \frac{1}{\sqrt{n}} \sum_{i=1}^n \left\langle K_0(\cdot, X_i) - \Psi_{P_n}, \frac{\Psi_{P_n}}{\mathcal{V}_n} \right\rangle_{\mathcal{H}_{K_0}},$$

where $\mathcal{V}_n := \sqrt{\|\mathcal{A}_{P_n}^{1/2} \Psi_{P_n}\|_{\mathcal{H}_{K_0}}^2 - \|\Psi_{P_n}\|_{\mathcal{H}_{K_0}}^4}$ and $\mathcal{A}_{P_n} := \int_{\mathcal{X}} K_0(\cdot, x) \otimes_{\mathcal{H}_{K_0}} K_0(\cdot, x) dP_n(x)$. The central limit theorem yields that $\frac{1}{\sqrt{n}} \sum_{i=1}^n \left\langle K_0(\cdot, X_i) - \Psi_{P_n}, \frac{\Psi_{P_n}}{\mathcal{V}_n} \right\rangle_{\mathcal{H}_{K_0}} \rightsquigarrow Z$, where $Z \sim \mathcal{N}(0, 1)$. Thus, it follows that if $\|\Psi_{P_n}\|_{\mathcal{H}_{K_0}}^2 = o(n^{-1})$ and $\chi^2(P_n, P_0) = O(1)$ as $n \rightarrow \infty$, then

$$\begin{aligned} n\mathcal{V}_n^2 &= n \left\| \mathcal{A}_{P_n}^{1/2} \Psi_{P_n} \right\|_{\mathcal{H}_{K_0}}^2 - n \|\Psi_{P_n}\|_{\mathcal{H}_{K_0}}^4 \leq n \|\mathcal{A}_{P_n}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})} \|\Psi_{P_n}\|_{\mathcal{H}_{K_0}}^2 \leq n \mathbb{E}_{P_n} K_0(X, X) \|\Psi_{P_n}\|_{\mathcal{H}_{K_0}}^2 \\ &\leq n \left(\mathbb{E}_{P_0} K_0(X, X) + \sqrt{\mathbb{E}_{P_0} K_0^2(X, X)} \sqrt{\chi^2(P_n, P_0)} \right) \|\Psi_{P_n}\|_{\mathcal{H}_{K_0}}^2 \rightarrow 0 \end{aligned}$$

as $n \rightarrow \infty$, implying

$$n\hat{D}_{\text{KSD}}^2 \rightsquigarrow S,$$

thereby yielding

$$P_{H_1} \{ \hat{D}_{\text{KSD}}^2 \geq q_{1-\alpha} \} = P_{H_1} \{ n\hat{D}_{\text{KSD}}^2 \geq nq_{1-\alpha} \} \rightarrow P\{S \geq \tilde{d}\} < 1,$$

where $\tilde{d} = \lim_{n \rightarrow \infty} nq_{1-\alpha}$, and the last inequality follows using the facts that $q_{1-\alpha} > 0$ (by the symmetry of the wild bootstrap distribution) S has a non-zero probability of being negative. Thus it remains to show that we can construct a sequence of probability distributions $\{P_n\}_n \in \mathcal{P}$ such that $\|\Psi_{P_n}\|_{\mathcal{H}_{K_0}}^2 = o(n^{-1})$. Suppose $\lambda_i = h(i)$, where h is strictly decreasing continuous function on \mathbb{N} . Let $k_n = \lfloor h^{-1}(\Delta_n^{1/2\theta}) \rfloor$ and define

$$u_n := \lambda_{k_n}^\theta \phi_{k_n}.$$

Observe that $u_n \in \text{Ran}(\Upsilon_{P_0}^\theta)$ and $\|u_n\|_{L^2(P_0)}^2 = \chi^2(P_n, P_0) = \Delta_n$. Thus it remains to ensure there exists P_n such that $u_n = \frac{dP_n}{dP_0} - 1$, for which it is sufficient to bound $|u_n(x)|$ as follows:

Case 1: \mathbf{A}_1 holds and $\theta > 1$

In this case,

$$\begin{aligned} |u_n(x)| &= \lambda_{k_n}^\theta |\langle K_0(\cdot, x), \phi_{k_n} \rangle_{\mathcal{H}_{K_0}}| = \lambda_{k_n}^{\theta-1} |\mathbb{E}_{P_0}(\tilde{\phi}_{k_n}(Y) K_0(x, Y))| \\ &\leq \lambda_{k_n}^{\theta-1} \left(\mathbb{E}_{P_0} \tilde{\phi}_{k_n}^2(Y) \right)^{1/2} \left(\mathbb{E}_{P_0} K_0^2(x, Y) \right)^{1/2} \lesssim \lambda_{k_n}^{\theta-1}, \end{aligned}$$

which implies that $|u_n(x)| < 1$ for large enough n . Then to ensure that $P_n \in \mathcal{P}$, it remains to show $P_n \in \mathcal{S}$. This is indeed the case since

$$\mathbb{E}_{P_n} [K_0(X, X)] = \mathbb{E}_{P_0} [K_0(X, X)(u_n(X) + 1)] \stackrel{(*)}{\leq} 2\mathbb{E}_{P_0} [K_0(X, X)] < \infty,$$

and

$$\mathbb{E}_{P_n} [K_0(X, X)]^r = \mathbb{E}_{P_0} [K_0(X, X)]^r (u_n(X) + 1) \stackrel{(*)}{\leq} 2\mathbb{E}_{P_0} [K_0(X, X)]^r \stackrel{(\dagger)}{\leq} 2cr! \kappa^r,$$

where $(*)$ follows using $|u_n(x)| < 1$ for large n , and (\dagger) follows by **A1**.

Case 2: $\sup_i \|\phi_i\|_\infty < \infty$

In this case,

$$|u_n(x)| \leq \sup_k \|\phi_k\|_\infty \lambda_{k_n}^\theta \leq 1,$$

for n large enough. Then to ensure that $P_n \in \mathcal{P}$, it remains to show $P_n \in \mathcal{S}$ which follows similar to the previous case.

$$\text{Thus in both the cases, } D_{\text{KSD}}^2(P_n, P_0) = \|\Psi_{P_n}\|_{\mathcal{H}_{K_0}}^2 = \left\| \Upsilon_{P_0}^{1/2} u_n \right\|_{L^2(P_0)}^2 = \Delta_n^{\frac{2\theta+1}{2\theta}} = o(n^{-1}).$$

8.2 Proof of Theorem 2

Following standard arguments from Ingster (1987) and Ingster (1993), it is sufficient to find a set of distributions $\{P_k\}_{k=1}^L \subset \mathcal{P}$ such that

$$\mathbb{E}_{P_0^n} \left[\left(\frac{1}{L} \sum_{k=1}^L \frac{dP_k^n}{dP_0^n} \right)^2 \right] < \infty.$$

Then the proof exactly follows the proof of Theorem 2 of Hagrass et al. (2023) by replacing \mathcal{T} with Υ_{P_0} with one main difference, which is bounding

$$\left| \frac{dP_k}{dP_0}(x) - 1 \right| = |u_{n,k}(x)| = \left| a_n \sum_{i=1}^{B_n} \varepsilon_{ki} \phi_i(x) \right|$$

in the case when $\sup_i \|\phi_i\|_\infty$ is not finite, since in this paper we do not assume that K_0 is bounded. To this end, in the following, we obtain a bound on $|u_{n,k}(x)|$ without assuming the boundedness of K_0 . Let $C_n = \lfloor \sqrt{B_n} \rfloor$ and $a_n = \sqrt{\frac{\Delta_n}{C_n}}$. Then

$$\begin{aligned} |u_{n,k}(x)| &= \left| \left\langle K_0(\cdot, x), a_n \sum_{i=1}^{B_n} \varepsilon_{ki} \phi_i \right\rangle_{\mathcal{H}_{K_0}} \right| = \left| \left\langle K_0(\cdot, x), a_n \sum_{i=1}^{B_n} \varepsilon_{ki} \lambda_i^{-1} \mathfrak{J}^* \tilde{\phi}_i \right\rangle_{\mathcal{H}_{K_0}} \right| \\ &= \left| \mathbb{E}_{P_0} \left[K_0(x, Y) a_n \sum_{i=1}^{B_n} \varepsilon_{ki} \lambda_i^{-1} \tilde{\phi}_i(Y) \right] \right| \\ &\leq \left(a_n^2 \sum_{i,j}^{B_n} \varepsilon_{ki} \varepsilon_{kj} \lambda_i^{-1} \lambda_j^{-1} \mathbb{E}_{P_0} [\tilde{\phi}_i(Y) \tilde{\phi}_j(Y)] \right)^{1/2} (\mathbb{E}_{P_0} [K(x, Y)]^2)^{1/2} \end{aligned}$$

$$= a_n \left(\sum_{i=1}^{B_n} \varepsilon_{ki}^2 \lambda_i^{-2} \right)^{1/2} (\mathbb{E}_{P_0} [K(x, Y)]^2)^{1/2}.$$

Thus when $\lambda_i \asymp i^{-\beta}$, the above bound yields

$$|u_{n,k}(x)| \lesssim \sqrt{\Delta_n} B_n^\beta \lesssim \Delta_n^{\frac{1}{2} - \frac{1}{2\theta}},$$

where $B_n \asymp \Delta_n^{\frac{-1}{2\theta\beta}}$, which implies $|u_{n,k}(x)| < 1$ for large enough n and $\theta > 1$. For the case $\lambda_i \asymp e^{-\tau i}$, the bound yields that

$$|u_{n,k}(x)| \lesssim a_n C_n^{1/2} e^{\tau B_n} \lesssim 1,$$

where we used $B_n \asymp \tau^{-1} \log(\Delta_n^{-1/2})$. This means for large n , $|u_{n,k}(x)| < 1$. Moreover, under \mathbf{A}_1 , we have

$$\mathbb{E}_{P_k} [K_0(X, X)] = \mathbb{E}_{P_0} [K_0(X, X)(u_{n,k}(X) + 1)] \leq 2\mathbb{E}_{P_0} [K_0(X, X)] < \infty,$$

and

$$\mathbb{E}_{P_k} [K_0(X, X)]^r = \mathbb{E}_{P_0} [K_0(X, X)^r (u_{n,k}(X) + 1)] \leq 2\mathbb{E}_{P_0} [K_0(X, X)]^r \leq 2cr! \kappa^r.$$

Thus $P_k \in \mathcal{P}$ for any $k \in \{1, \dots, L\}$.

8.3 Proof of Theorem 3

Define

$$S_z : \mathcal{H}_{K_0} \rightarrow \mathbb{R}^{n_2}, \quad f \mapsto \frac{1}{\sqrt{n_2}} (f(Z_1), \dots, f(Z_{n_2}))^\top$$

so that

$$S_z^* : \mathbb{R}^{n_2} \rightarrow \mathcal{H}_{K_0}, \quad \alpha \mapsto \frac{1}{\sqrt{n_2}} \sum_{i=1}^{n_2} \alpha_i K_0(\cdot, Z_i).$$

Then observe that $\hat{\mathcal{A}}_P = S_z^* S_z$ and $n_2 S_z S_z^* = [K_0(Z_i, Z_j)]_{i,j \in [n_2]} := K_{n_2}$. Let $(\hat{\lambda}_i, \hat{\alpha}_i)_i$ be the eigensystem of $\frac{1}{n_2} K_{n_2}$, i.e., $S_z S_z^* \hat{\alpha}_i = \hat{\lambda}_i \hat{\alpha}_i$. By applying S_z^* to both sides on the eigen equation, we can show that $(\hat{\lambda}_i, \hat{\psi}_i)$ are the eigensystem of $\hat{\mathcal{A}}_P$ where $\hat{\psi}_i := \frac{1}{\sqrt{\hat{\lambda}_i}} S_z^* \hat{\alpha}_i$. Thus, using $\hat{\psi}_i$ in the definition of $g_\lambda(\hat{\mathcal{A}}_P)$ yields that

$$g_\lambda(\hat{\mathcal{A}}_P) := g_\lambda(0) \mathbf{I} + S_z^* G S_z.$$

Define $\mathbf{1}_n := (1, \dots, 1)^\top$, and let $\mathbf{1}_n^i$ be a vector of zeros with only the i^{th} entry equal one. Also define S_x similar to that of S_z based on samples $(X_i)_{i=1}^{n_1}$ and observe that $\sqrt{n_1 n_2} S_x S_z^* := K_{n_1, n_2}$. Therefore,

$$\hat{\mathbb{S}}_\lambda^P(\mathbb{X}_n) = \frac{1}{n_1(n_1 - 1)} (\textcircled{1} - \textcircled{2}),$$

where $\textcircled{1} := \sum_{i,j=1}^{n_1} h(X_i, X_j)$, $\textcircled{2} := \sum_{i=1}^{n_1} h(X_i, X_i)$, and

$$h(X_i, X_j) = \left\langle g_\lambda^{1/2}(\hat{\mathcal{A}}_P) K_0(\cdot, X_i), g_\lambda^{1/2}(\hat{\mathcal{A}}_P) K_0(\cdot, X_j) \right\rangle_{\mathcal{H}_{K_0}}.$$

Note that

$$\begin{aligned}
\textcircled{1} &= \left\langle g_\lambda(\hat{\mathcal{A}}_P) \sum_{i=1}^{n_1} K_0(\cdot, X_i), \sum_{j=1}^{n_1} K_0(\cdot, X_j) \right\rangle_{\mathcal{H}_{K_0}} = n_1 \left\langle g_\lambda(\hat{\mathcal{A}}_P) S_x^* \mathbf{1}_{n_1}, S_x^* \mathbf{1}_{n_1} \right\rangle_{\mathcal{H}_{K_0}} \\
&= n_1 \left\langle S_x g_\lambda(\hat{\mathcal{A}}_P) S_x^* \mathbf{1}_{n_1}, \mathbf{1}_{n_1} \right\rangle_2 = n_1 \left\langle g_\lambda(0) \frac{1}{n_1} K_{n_1} + \frac{1}{n_1 n_2} K_{n_1, n_2} G K_{n_1, n_2}^\top \mathbf{1}_{n_1}, \mathbf{1}_{n_1} \right\rangle_2 \\
&= \mathbf{1}_{n_1}^\top \left(g_\lambda(0) K_{n_1} + \frac{1}{n_2} K_{n_1, n_2} G K_{n_1, n_2}^\top \right) \mathbf{1}_{n_1},
\end{aligned}$$

and

$$\begin{aligned}
\textcircled{2} &= \sum_{i=1}^{n_1} \left\langle g_\lambda(\hat{\mathcal{A}}_P) K_0(\cdot, X_i), K_0(\cdot, X_i) \right\rangle_{\mathcal{H}_{K_0}} = n_1 \sum_{i=1}^{n_1} \left\langle g_\lambda(\hat{\mathcal{A}}_P) S_x^* \mathbf{1}_{n_1}^i, S_x^* \mathbf{1}_{n_1}^i \right\rangle_{\mathcal{H}_{K_0}} \\
&= n_1 \sum_{i=1}^{n_1} \left\langle S_x g_\lambda(\hat{\mathcal{A}}_P) S_x^* \mathbf{1}_{n_1}^i, \mathbf{1}_{n_1}^i \right\rangle_2 = n_1 \sum_{i=1}^{n_1} \left\langle g_\lambda(0) \frac{1}{n_1} K_{n_1} + \frac{1}{n_1 n_2} K_{n_1, n_2} G K_{n_1, n_2}^\top \mathbf{1}_{n_1}^i, \mathbf{1}_{n_1}^i \right\rangle_2 \\
&= \text{Tr} \left(g_\lambda(0) K_{n_1} + \frac{1}{n_2} K_{n_1, n_2} G K_{n_1, n_2}^\top \right).
\end{aligned}$$

8.4 Proof of Theorem 4

Let $h(x, y) := \left\langle g_\lambda^{1/2}(\hat{\mathcal{A}}_{P_0}) K_0(\cdot, x), g_\lambda^{1/2}(\hat{\mathcal{A}}_{P_0}) K_0(\cdot, y) \right\rangle_{\mathcal{H}_{K_0}}$. Then observe that h is symmetric and degenerate with respect to P_0 , i.e., $\mathbb{E}_{P_0} h(x, Y) = 0$ and $\mathbb{E}_{P_0} h(X, Y) = 0$. Moreover

$$\mathbb{E}_{P_0}(h^2(X, Y) | \mathbb{Z}_{n_2}) \leq \left\| g_\lambda^{1/2}(\hat{\mathcal{A}}_{P_0}) \right\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^4 [\mathbb{E}_{P_0} K_0(X, X)] [\mathbb{E}_{P_0} K_0(Y, Y)] \lesssim \frac{[\mathbb{E}_{P_0} K_0(X, X)]^2}{\lambda^2} < \infty,$$

and

$$\mathbb{E}_{P_0}(h(X, X) | \mathbb{Z}_{n_2}) \leq \left\| g_\lambda^{1/2}(\hat{\mathcal{A}}_{P_0}) \right\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 [\mathbb{E}_{P_0} K_0(X, X)] \lesssim \frac{1}{\lambda} [\mathbb{E}_{P_0} K_0(X, X)] < \infty.$$

The result therefore follows from Dehling and Mikosch (1994, Theorem 3.1).

8.5 Proof of Theorem 5

We will show that under the conditions stated in the theorem, for any $0 < \delta < 1$, we have

$$\lim_{n \rightarrow \infty} P\{\hat{\mathbb{S}}_\lambda^{P_n}(\mathbb{X}_n) \geq q_{1-\alpha}^\lambda\} > 1 - 4\delta,$$

and thus the desired result holds. Let

$$\gamma := \frac{\log \frac{2}{\alpha}}{n_1 \sqrt{\delta}} \left(\|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})} \sqrt{\zeta} + \|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \sqrt{D_\lambda} + \zeta \right),$$

and

$$\gamma_1 := \frac{1}{\sqrt{\delta}} \left(\frac{\|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})} \sqrt{\zeta}}{\sqrt{n_1}} + \frac{\|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \sqrt{D_\lambda}}{n_1} \right).$$

Then by Lemma A.3, Lemma A.7 and (Hagrass et al., 2022, Lemma A.1), we can deduce that it is sufficient to show

$$P(\gamma \leq \zeta - c_1\gamma_1) \geq 1 - 2\delta,$$

for some constant $c_1 > 0$. Observe that

$$\begin{aligned} P(\gamma \leq \zeta - c_1\gamma_1) &= P(\zeta \geq \gamma + c_1\gamma_1) \\ &\geq P\left(\{\zeta \geq \gamma\} \cap \left\{\zeta \geq \frac{c_1 \|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})} \sqrt{\zeta}}{\sqrt{\delta n_1}}\right\} \cap \left\{\zeta \geq \frac{c_1 \|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \sqrt{D_\lambda}}{\sqrt{\delta n_1}}\right\}\right) \\ &\stackrel{(*)}{\geq} P\left(\{\zeta \geq \gamma\} \cap \left\{\frac{c_2 \|\mathcal{W}^{-1}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^{-2} \Delta_n}{1 + \sqrt{C_\lambda} \sqrt{\Delta_n}} \geq \frac{\|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2}{\delta n_1}\right\} \cap \left\{\|\mathcal{W}^{-1}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \leq \frac{3}{2}\right\}\right. \\ &\quad \left. \cap \left\{\frac{c_2 \|\mathcal{W}^{-1}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^{-2} \Delta_n}{1 + \sqrt{C_\lambda} \sqrt{\Delta_n}} \geq \frac{\|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \sqrt{D_\lambda}}{\sqrt{\delta n_1}}\right\} \cap \left\{\|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \leq 2\right\}\right) \\ &\stackrel{(**)}{\geq} P\left(\{\zeta \geq \gamma\} \cap \left\{\|\mathcal{W}^{-1}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \leq \frac{3}{2}\right\} \cap \left\{\|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \leq 2\right\}\right) \\ &\stackrel{(\dagger)}{\geq} P\left(\left\{\|\mathcal{W}^{-1}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \leq \frac{3}{2}\right\} \cap \left\{\|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \leq 2\right\}\right) \\ &\geq 1 - P\left(\|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \geq 2\right) - P\left(\|\mathcal{W}^{-1}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \geq \frac{3}{2}\right) \stackrel{(\ddagger)}{\geq} 1 - 2\delta, \end{aligned}$$

where $c_2 > 0$, $(*)$ follows from Lemma A.6(i) under the assumption that $\sup_{P \in \mathcal{P}} \|\Upsilon_{P_0}^{-\theta} u\|_{L^2(P_0)} < \infty$, $u \in \text{Ran}(\Upsilon_{P_0}^\theta)$, and for some $c_3 > 0$,

$$\Delta_n \geq c_3 \lambda^{2\tilde{\theta}}. \quad (18)$$

$(**)$ and (\dagger) follows under the dominant condition that

$$\frac{\Delta_n}{1 + \sqrt{C_\lambda} \sqrt{\Delta_n}} \geq \frac{c_4(\delta) \sqrt{D_\lambda}}{n}, \quad (19)$$

for some constant $c_4(\delta) \asymp \delta^{-1}$. (\ddagger) follows from Lemma A.4(i) assuming

$$\lambda \geq \frac{c_5(\delta)}{\sqrt{n}}, \quad (20)$$

where $c_5(\delta) \asymp \log(\frac{2}{\delta})$, and if $4C_\lambda \Delta_n < 1$, then using Lemma A.4(ii), (20) can be replaced by

$$\lambda \geq \frac{c_6(\delta) \log n}{n} \left(\mathcal{N}_1(\lambda) + C_\lambda \sqrt{\Delta_n} \right), \quad (21)$$

for some constant $c_6(\delta) \asymp \log(\frac{2}{\delta})$. When $\sup_i \|\phi_i\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \leq C < \infty$ (which implies that the kernel is bounded) then using Lemma A.4(iii), (20) can be replaced by

$$\lambda \geq \frac{c_7(\delta) \log n}{n}, \quad (22)$$

where $c_7(\delta) \asymp \log(\frac{1}{\delta})$. On the other hand, when $\sup_i \|\phi_i\|_{\mathcal{L}^\infty(\mathcal{H}_K)}^2 \leq C < \infty$ and $4C_\lambda \Delta_n < 1$, define

$$\gamma := \frac{\log \frac{2}{\alpha}}{n_1 \sqrt{\delta}} \left(\|\mathcal{V}\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \sqrt{\zeta} + \|\mathcal{V}\|_{\mathcal{L}^\infty(\mathcal{H}_K)}^2 \sqrt{D_\lambda} + \zeta \right),$$

and

$$\gamma_1 := \frac{1}{\sqrt{\delta}} \left(\frac{\|\mathcal{V}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})} \sqrt{\zeta}}{\sqrt{n_1}} + \frac{\|\mathcal{V}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \sqrt{D_\lambda}}{n_1} \right).$$

Then by Lemma A.3(ii), Lemma A.6(ii) and Lemma A.5 and the exactly using the same approach as above, we can show that

$$P(\gamma \leq \zeta - c_8 \gamma_1) \geq 1 - 2\delta,$$

for some constant $c_8 > 0$, under the same conditions (18), (19) and replacing (20) with

$$n \geq c_9(\delta) \mathcal{N}_1(\lambda) \log \mathcal{N}_1(\lambda), \quad (23)$$

for some constant $c_9(\delta) \asymp \log(\frac{1}{\delta})$, and using $\log \frac{x}{\delta} \leq \log \frac{1}{\delta} \log x$ for large enough $x \in \mathbb{R}$, and assuming $\mathbb{E}_{P_n} \phi_1^2 > 0$. Finally observe that the conditions (11)–(16) given in the theorem statement implies equations (18)–(23) respectively for any $0 < \delta < 1$.

8.6 Proof of Proposition 8

Observe that

$$\Upsilon_{P_0} = \sum_{k \in I} \lambda_k \gamma_k \otimes_{L^2(P_0)} \gamma_k$$

since

$$\Upsilon_{P_0} f = \int K_0(\cdot, x) f(x) dP_0(x) = \int \sum_{k \in I} \lambda_k \gamma_k \gamma_k(x) f(x) dP_0(x) = \sum_{k \in I} \lambda_k \gamma_k \langle \gamma_k, f \rangle_{L^2(P_0)}, \quad f \in L^2(P_0).$$

Therefore, for $f \in L^2(P_0)$,

$$\Upsilon_{P_0}^\theta f = \sum_{k \in I} \lambda_k^\theta b_k \gamma_k,$$

where $b_k := \langle f, \gamma_k \rangle_{L^2(P_0)}$, with $\sum_{k \in I} b_k^2 < \infty$. Thus any function belonging to $\text{Ran}(\Upsilon_{P_0}^\theta)$ can be expressed as $\sum_{k \in I} a_k \gamma_k$, with $(a_k)_{k \in I}$ satisfying $\sum_{k \in I} a_k^2 \lambda_k^{-2\theta} < \infty$.

8.7 Proof of Example 5

If $\psi_{P_0}(\cdot, x) = K(\cdot, x) - \mathbb{E}_{P_0} K(\cdot, X)$, then note that $K_0(x, y) := \sum_{k \neq 0} |k|^{-\beta} e^{\sqrt{-1}2\pi k x} e^{-\sqrt{-1}2\pi k y}$. Since $1 \cup (e^{\sqrt{-1}2\pi k \cdot})_{k \in I}$ form an orthonormal basis in $L^2([0, 1])$, the result follow by applying Proposition 8.

When $\psi_{P_0}(\cdot, x) = K(\cdot, x) \nabla_x \log p_0(x) + \nabla_x K(\cdot, x)$, we have

$$K_0(x, y) = \sum_{k \neq 0} \lambda_k \frac{d\gamma_k(x)}{dx} \frac{\overline{d\gamma_k(y)}}{dy},$$

where $\lambda_k := |k|^{-\beta}$, and $\gamma_k(x) := e^{\sqrt{-1}2\pi k x}$. Then observe that $\frac{d\gamma_k(x)}{dx} = \sqrt{-1}2\pi k \gamma_k(x)$, thus

$$K_0(x, y) = (2\pi)^2 \sum_k \tilde{\lambda}_k \gamma_k(x) \gamma_k(y),$$

with $\tilde{\lambda}_k := k^2 \lambda_k = |k|^{-(\beta-2)}$. Thus the result follows by applying Proposition 8.

8.8 Proof of Example 6

By Mehler's formula (Kibble, 1945), we have

$$K(x, y) = \sum_{k=0}^{\infty} \rho^k \gamma_k(x) \gamma_k(y), \quad (24)$$

where $\gamma_k(x) = \frac{1}{\sqrt{k!}} H_k(x)$, and $H_k(x) = (-1)^k e^{x^2/2} \frac{d^k}{dx^k} e^{-x^2/2}$. Observe that $\gamma_0(x) = 1$ and $1 \cup (\gamma_k)_{k=1}^{\infty}$ form an orthonormal basis with respect to $L^2(P_0)$ where $p_0(x) := \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$.

If $\psi_{P_0}(\cdot, x) = K(\cdot, x) - \mathbb{E}_{P_0} K(\cdot, X)$ then

$$K_0(x, y) = \sum_{k=0}^{\infty} \rho^k \bar{\gamma}_k(x) \bar{\gamma}_k(y) = \sum_{k=1}^{\infty} \rho^k \gamma_k(x) \gamma_k(y),$$

where $\bar{\gamma}_k(x) = \gamma_k(x) - \mathbb{E}_{P_0} \gamma_k(X)$. Thus the result follow by applying Theorem 8. On the other hand, if $\psi_{P_0}(\cdot, x) = K(\cdot, x) \nabla_x \log p_0(x) + \nabla_x K(\cdot, x)$ then

$$K_0(x, y) = \frac{d \log p_0(x)}{dx} \frac{d \log p_0(y)}{dy} K(x, y) + \frac{d \log p_0(x)}{dx} \frac{\partial K(x, y)}{\partial y} + \frac{d \log p_0(y)}{dy} \frac{\partial K(x, y)}{\partial x} + \frac{\partial^2 K(x, y)}{\partial x \partial y}.$$

Observe that

$$\frac{d \log p_0(x)}{dx} = -x, \quad \frac{\partial K(x, y)}{\partial y} = \sum_{n=0}^{\infty} \frac{k \rho^k}{k!} H_{k-1}(y) H_k(x),$$

and

$$\frac{\partial^2 K(x, y)}{\partial x \partial y} = \sum_{n=0}^{\infty} \frac{k^2 \rho^k}{k!} H_{k-1}(y) H_{k-1}(x),$$

where we used that $\frac{dH_k(x)}{dx} = kH_{k-1}(x)$. Then by using $H_{k+1}(x) = xH_k(x) - kH_{k-1}(x)$, it can be shown that

$$K_0(x, y) = \frac{1}{\rho} \sum_{k=1}^{\infty} \lambda_k \gamma_k(x) \gamma_k(y),$$

where $\lambda_k = k \rho^k = k e^{k \log \rho}$. Thus the result follows by applying Proposition 8.

8.9 Proof of Remark 6(i)

Observe that

$$K(x, y) = \frac{1}{\sqrt{1-\rho^2}} \exp\left(-\frac{\rho^2(x^2 + y^2) - 2\rho xy}{2(1-\rho^2)}\right)$$

$$= \frac{1}{\sqrt{1-\rho^2}} \exp\left(\frac{-\rho}{2(1-\rho^2)}(x-y)^2\right) \exp\left(\frac{\rho}{2(1+\rho)}x^2\right) \exp\left(\frac{\rho}{2(1+\rho)}y^2\right).$$

Let $\bar{K}(x, y) = \exp\left(\frac{-\rho}{2(1-\rho^2)}(x-y)^2\right)$. An application of the Mehler formula in (24) yields

$$\begin{aligned} \bar{K}(x, y) &= \sqrt{1-\rho^2} \sum_{k=0}^{\infty} \rho^k \gamma_k(x) \gamma_k(y) \exp\left(\frac{-\rho}{2(1+\rho)}x^2\right) \exp\left(\frac{-\rho}{2(1+\rho)}y^2\right) \\ &= \sum_{k=0}^{\infty} \lambda_k \psi_k(x) \psi_k(y), \end{aligned}$$

where $\lambda_k = (1-\rho)\rho^k$, and $\psi_k(x) = \left(\frac{1+\rho}{1-\rho}\right)^{1/4} \gamma_k(x) \exp\left(\frac{-\rho}{2(1+\rho)}x^2\right)$. Note that $(\psi_k)_{k=0}^{\infty}$ form an orthonormal basis with respect $L^2(Q)$ with Q being a Gaussian distribution with mean zero and variance $\frac{1+\rho}{1-\rho}$. Then the RKHS associated with \bar{K} can be defined as

$$\mathcal{H}_{\bar{K}} := \left\{ \sum_{k=0}^{\infty} a_k \psi_k(x) : \sum_{k=0}^{\infty} a_k^2 \lambda_k^{-1} < \infty \right\}.$$

Hence for any $f \in \mathcal{H}_{\bar{K}}$, the function $\bar{f}(x) := \exp\left(\frac{\rho}{2(1+\rho)}x^2\right) f(x) - a_0 \in R_1$ for any $\theta \leq \frac{1}{2}$. Therefore,

$$S := \left\{ \exp\left(\frac{\rho}{2(1+\rho)}x^2\right) g_f(x) - a_0 : g_f \in \mathcal{H}_{\bar{K}} \right\} \subseteq R_1,$$

for any $\theta \leq \frac{1}{2}$.

Let $\tilde{K}(x, y) = \exp(-\sigma(x-y)^2)$. Then we will show that $\mathcal{H}_{\tilde{K}} \subseteq S$. To this end, we need to show that for any $f \in \mathcal{H}_{\tilde{K}}$ with $\mathbb{E}_{P_0} f = 0$, $\exists g_f \in \mathcal{H}_{\tilde{K}}$ such that $f(x) = \exp\left(\frac{\rho}{2(1+\rho)}x^2\right) g_f(x) - a_0$. Thus it is sufficient to show that

$$g_f(x) := \exp\left(\frac{-\rho}{2(1+\rho)}x^2\right) (f(x) + a_0) \in \mathcal{H}_{\tilde{K}}.$$

Observe that

$$g_f(x) = \exp\left(\frac{-\rho}{2(1+\rho)}x^2\right) f(x) + a_0 \exp\left(\frac{-\rho}{2(1+\rho)}x^2\right) =: g_1(x) + g_2(x).$$

We will show $g_1, g_2 \in \mathcal{H}_{\tilde{K}}$ and therefore the claim follows that $\mathcal{H}_{\tilde{K}} \subseteq R_1$. For that matter, note that

$$\begin{aligned} &|\hat{g}_1(\omega)| \\ &= \left| \int_{-\infty}^{\infty} \hat{f}(t) \exp\left(\frac{-(\omega-t)^2(1+\rho)}{2\rho}\right) dt \right| \\ &\leq \int_{-\infty}^{\infty} |\hat{f}(t)| \exp\left(\frac{-(\omega-t)^2(1+\rho)}{2\rho}\right) \exp\left(\frac{-t^2}{8\sigma}\right) \exp\left(\frac{t^2}{8\sigma}\right) dt \\ &\leq \left(\int_{-\infty}^{\infty} |\hat{f}(t)|^2 \exp\left(\frac{t^2}{4\sigma}\right) dt \right)^{1/2} \left(\int_{-\infty}^{\infty} \exp\left(\frac{-(\omega-t)^2(1+\rho)}{\rho}\right) \exp\left(\frac{-t^2}{4\sigma}\right) dt \right)^{1/2} \end{aligned}$$

$$\begin{aligned}
&= \|f\|_{\mathcal{H}_K} \left(\int_{-\infty}^{\infty} \exp\left(\frac{-(\omega-t)^2(1+\rho)}{\rho}\right) \exp\left(\frac{-t^2}{4\sigma}\right) dt \right)^{1/2} \\
&\lesssim \sqrt{\exp\left(\frac{-(1+\rho)}{4\sigma(1+\rho)+\rho}\omega^2\right)}.
\end{aligned}$$

Then

$$\begin{aligned}
\|g_1\|_{\mathcal{H}_K}^2 &= \int_{-\infty}^{\infty} |\hat{g}_1(\omega)|^2 \exp\left(\frac{\omega^2(1-\rho^2)}{2\rho}\right) d\omega \\
&\lesssim \int_{-\infty}^{\infty} \exp\left(-\omega^2\left(\frac{1+\rho}{4\sigma(1+\rho)+\rho} - \frac{1-\rho^2}{2\rho}\right)\right) d\omega < \infty,
\end{aligned}$$

where the last inequality follows using $\sigma < \frac{\rho}{4(1-\rho)}$ to ensure the exponent argument is negative. Next, observe that $\hat{g}_2(\omega) = a_0 \exp\left(\frac{-\omega^2(1+\rho)}{2\rho}\right)$. Thus

$$\begin{aligned}
\|g_2\|_{\mathcal{H}_K}^2 &= \int_{-\infty}^{\infty} |\hat{g}_2(\omega)|^2 \exp\left(\frac{\omega^2(1-\rho^2)}{2\rho}\right) d\omega \\
&= a_0^2 \int_{-\infty}^{\infty} \exp\left(-\omega^2\left(\frac{1+\rho}{2\rho} - \frac{1-\rho^2}{2\rho}\right)\right) d\omega < \infty.
\end{aligned}$$

Acknowledgments

OH and BKS are partially supported by the National Science Foundation (NSF) CAREER award DMS-1945396. KB is supported by NSF grant DMS-2053918.

References

- M. Arcones and E. Giné. On the bootstrap of U and V statistics. *The Annals of Statistics*, 20(2):655–674, 1992.
- N. Aronszajn. Theory of reproducing kernels. *Trans. Amer. Math. Soc.*, 68:337–404, 1950.
- S. Balakrishnan and L. Wasserman. Hypothesis testing for densities and high-dimensional multinomials: Sharp local minimax rates. *The Annals of Statistics*, 47(4):1893–1927, 2019.
- K. Balasubramanian, T. Li, and M. Yuan. On the optimality of kernel-embedding based goodness-of-fit tests. *Journal of Machine Learning Research*, 22(1):1–45, 2021.
- F. Bauer, S. Pereverzev, and L. Rosasco. On regularization algorithms in learning theory. *Journal of Complexity*, 23(1):52–72, 2007.
- P. J. Bickel and L. Breiman. Sums of functions of nearest neighbor distances, moment bounds, limit theorems and a goodness of fit test. *The Annals of Probability*, 11(1):185–214, 1983.
- G. Boente, D. Rodriguez, and W. G. Manteiga. Goodness-of-fit test for directional data. *Scandinavian Journal of Statistics*, 41(1):259–275, 2014.
- M. V. Burnashev. On the minimax detection of an inaccurately known signal in a white Gaussian noise background. *Theory of Probability & Its Applications*, 24(1):107–119, 1979.

- A. Caponnetto and E. De Vito. Optimal rates for regularized least-squares algorithm. *Foundations of Computational Mathematics*, 7:331–368, 2007.
- K. Chwialkowski, H. Strathmann, and A. Gretton. A kernel test of goodness of fit. In *Proceedings of The 33rd International Conference on Machine Learning*, volume 48 of *Proceedings of Machine Learning Research*, pages 2606–2615, New York, New York, USA, 2016. PMLR.
- V. H. de la Peña and E. Giné. *Decoupling: From Dependence to Independence*. Springer Science and Business Media, 1999.
- H. Dehling and T. Mikosch. Random quadratic forms and the bootstrap for U -statistics. *Journal of Multivariate Analysis*, 51(2):392–413, 1994.
- A. Dvoretzky, J. Kiefer, and J. Wolfowitz. Asymptotic minimax character of the sample distribution function and of the classical multinomial estimator. *The Annals of Mathematical Statistics*, 27(3):642–669, 1956.
- B. Ebner, N. Henze, and J. E. Yukich. Multivariate goodness-of-fit on flat and curved spaces via nearest neighbor distances. *Journal of Multivariate Analysis*, 165:231–242, 2018.
- H. W. Engl, M. Hanke, and A. Neubauer. *Regularization of Inverse Problems*. Kluwer Academic Publishers, Dordrecht, The Netherlands, 1996.
- T. Fernandez, N. Rivera, W. Xu, and A. Gretton. Kernelized Stein discrepancy tests of goodness-of-fit for time-to-event data. In Hal Daumé III and Aarti Singh, editors, *Proceedings of the 37th International Conference on Machine Learning*, volume 119 of *Proceedings of Machine Learning Research*, pages 3112–3122. PMLR, 13–18 Jul 2020.
- A. Fischer and C. Igel. An introduction to restricted Boltzmann machines. In Luis Alvarez, Marta Mejail, Luis Gomez, and Julio Jacobo, editors, *Progress in Pattern Recognition, Image Analysis, Computer Vision, and Applications*, pages 14–36. Springer Berlin Heidelberg, 2012.
- A. Gretton, K. Borgwardt, M. Rasch, B. Schölkopf, and A. Smola. A kernel method for the two-sample problem. In B. Schölkopf, J. Platt, and T. Hoffman, editors, *Advances in Neural Information Processing Systems*, volume 19, pages 513–520. MIT Press, 2006.
- A. Gretton, K. M. Borgwardt, M. J. Rasch, B. Schölkopf, and A. Smola. A kernel two-sample test. *Journal of Machine Learning Research*, 13(25):723–773, 2012.
- O. Hagrass, B. K. Sriperumbudur, and B. Li. Spectral regularized kernel two-sample tests. 2022. <https://arxiv.org/pdf/2212.09201v2.pdf>.
- O. Hagrass, B. K. Sriperumbudur, and B. Li. Spectral regularized kernel goodness-of-fit tests. 2023. <https://arxiv.org/pdf/2308.04561.pdf>.
- M. Hallin, G. Mordant, and J. Segers. Multivariate goodness-of-fit tests based on Wasserstein distance. *Electronic Journal of Statistics*, 15:1328–1371, 2021.
- M. Huskova and P. Janssen. Consistency of the generalized bootstrap for degenerate U -statistics. *The Annals of Statistics*, 21(4):1811–1823, 1993.
- Y. Ingster and I. A. Suslina. *Nonparametric Goodness-of-Fit Testing under Gaussian Models*, volume 169. Springer Science & Business Media, New York, 2003.

- Y. I. Ingster. Minimax testing of nonparametric hypotheses on a distribution density in the L_p metrics. *Theory of Probability & Its Applications*, 31(2):333–337, 1987.
- Y. I. Ingster. Asymptotically minimax hypothesis testing for nonparametric alternatives i, ii, iii. *Mathematical Methods of Statistics*, 2(2):85–114, 1993.
- P. E. Jupp. Data-driven Sobolev tests of uniformity on compact Riemannian manifolds. *The Annals of Statistics*, 36(3):1246–1260, 2008.
- H. Kanagawa, W. Jitkrittum, L. Mackey, K. Fukumizu, and A. Gretton. A kernel Stein test for comparing latent variable models. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 85(3):986–1011, 05 2023.
- E. V. Khmaladze. An innovation approach to goodness-of-fit tests in \mathbb{R}^m . *The Annals of Statistics*, 16(4):1503–1516, 1988.
- E. V. Khmaladze. Goodness of fit problem and scanning innovation martingales. *The Annals of Statistics*, 21(2):798–829, 1993.
- E. V. Khmaladze. Note on distribution free testing for discrete distributions. *The Annals of Statistics*, 41(6):2979–2993, 2013.
- E. V. Khmaladze. Unitary transformations, empirical processes and distribution free testing. *Bernoulli*, 22(1):563–588, 2016.
- W. F. Kibble. An extension of a theorem of Mehler’s on Hermite polynomials. *Mathematical Proceedings of the Cambridge Philosophical Society*, 41(1):12–15, 1945.
- E. L. Lehmann and J. P. Romano. *Testing Statistical Hypotheses*. Springer Science & Business Media, New York, 2006.
- T. Li and M. Yuan. On the optimality of Gaussian kernel based nonparametric tests against smooth alternatives. 2019. <https://arxiv.org/pdf/1909.03302.pdf>.
- Q. Liu, J. Lee, and M. Jordan. A kernelized Stein discrepancy for goodness-of-fit tests. In *Proceedings of The 33rd International Conference on Machine Learning*, volume 48 of *Proceedings of Machine Learning Research*, pages 276–284, New York, New York, USA, 2016. PMLR.
- K. V. Mardia, D. Holmes, and J. T. Kent. A goodness-of-fit test for the von Mises–Fisher distribution. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 46(1):72–78, 1984.
- P. Massart. The tight constant in the Dvoretzky-Kiefer-Wolfowitz Inequality. *The Annals of Probability*, 18(3):1269–1283, 1990.
- H. Q. Minh, P. Niyogi, and Y. Yao. Mercer’s theorem, feature maps, and smoothing. In Gábor Lugosi and Hans Ulrich Simon, editors, *Learning Theory*, pages 154–168, Berlin, 2006. Springer.
- S. Minsker. On some extensions of Bernstein’s inequality for self-adjoint operators. *Statistics & Probability Letters*, 127:111–119, 2017.
- K. Muandet, K. Fukumizu, B. Sriperumbudur, and B. Schölkopf. Kernel mean embedding of distributions: A review and beyond. *Foundations and Trends® in Machine Learning*, 10(1-2): 1–141, 2017.

- X. Qu, X. Fan, and B. Vemuri. Kernel Stein discrepancy on Lie groups: Theory and applications. 2023. <https://arxiv.org/pdf/2305.12551.pdf>.
- M. Reed and B. Simon. *Methods of Modern Mathematical Physics: Functional Analysis I*. Academic Press, New York, 1980.
- M. F. Schilling. Goodness of fit testing in \mathbb{R}^m based on the weighted empirical distribution of certain nearest neighbor statistics. *The Annals of Statistics*, 11(1):1–12, 1983a.
- M. F. Schilling. An infinite-dimensional approximation for nearest neighbor goodness of fit tests. *The Annals of Statistics*, 11(1):13–24, 1983b.
- A. Schrab, B. Guedj, and A. Gretton. KSD aggregated goodness-of-fit test. In . Koyejo, S. Mohamed, A. Agarwal, D. Belgrave, K. Cho, and A. Oh, editors, *Advances in Neural Information Processing Systems*, volume 35, pages 32624–32638. Curran Associates, Inc., 2022.
- A. Schrab, I. Kim, M. Albert, B. Laurent, B. Guedj, and A. Gretton. MMD aggregated two-sample test. *Journal of Machine Learning Research*, 24(194):1–81, 2023.
- R. Serfling. *Approximation Theorems of Mathematical Statistics*. Wiley, New York, 1980.
- C. Simon-Gabriel and B. Schölkopf. Kernel distribution embeddings: Universal kernels, characteristic kernels and kernel metrics on distributions. *Journal of Machine Learning Research*, 19(44):1–29, 2018.
- A. J. Smola, A. Gretton, L. Song, and B. Schölkopf. A Hilbert space embedding for distributions. In Marcus Hutter, Rocco A. Servedio, and Eiji Takimoto, editors, *Algorithmic Learning Theory*, pages 13–31. Springer-Verlag, Berlin, Germany, 2007.
- B. K. Sriperumbudur. On the optimal estimation of probability measures in weak and strong topologies. *Bernoulli*, 22(3):1839 – 1893, 2016.
- B. K. Sriperumbudur and N. Sterge. Approximate kernel PCA using random features: Computational vs. statistical trade-off. *The Annals of Statistics*, 50(5):2713–2736, 2022.
- B. K. Sriperumbudur, A. Gretton, K. Fukumizu, B. Schölkopf, and G. R. G. Lanckriet. Hilbert space embeddings and metrics on probability measures. *Journal of Machine Learning Research*, 11:1517–1561, 2010.
- I. Steinwart and C. Scovel. Mercer’s theorem on general domains: On the interaction between measures, kernels, and RKHSs. *Constructive Approximation*, 35:363–417, 2012.
- G. Wynne, M. Kasprzak, and A. B. Duncan. A Fourier representation of kernel Stein discrepancy with application to goodness-of-fit tests for measures on infinite dimensional Hilbert spaces. 2023. <https://arxiv.org/pdf/2206.04552.pdf>.
- W. Xu and T. Matsuda. A Stein goodness-of-fit test for directional distributions. In Silvia Chiappa and Roberto Calandra, editors, *Proceedings of the Twenty Third International Conference on Artificial Intelligence and Statistics*, volume 108 of *Proceedings of Machine Learning Research*, pages 320–330. PMLR, 2020.

- W. Xu and T. Matsuda. Interpretable Stein goodness-of-fit tests on Riemannian manifold. In Marina Meila and Tong Zhang, editors, *Proceedings of the 38th International Conference on Machine Learning*, volume 139 of *Proceedings of Machine Learning Research*, pages 11502–11513. PMLR, 2021.
- W. Xu and G. Reinert. A Stein goodness-of-test for exponential random graph models. In Arindam Banerjee and Kenji Fukumizu, editors, *Proceedings of the 24th International Conference on Artificial Intelligence and Statistics*, volume 130 of *Proceedings of Machine Learning Research*, pages 415–423. PMLR, 2021.
- W. Xu and G. D. Reinert. AgraSSt: Approximate graph Stein statistics for interpretable assessment of implicit graph generators. *Advances in Neural Information Processing Systems*, 35: 24268–24279, 2022.
- J. Yang, Q. Liu, V. Rao, and J. Neville. Goodness-of-fit testing for discrete distributions via Stein discrepancy. In Jennifer Dy and Andreas Krause, editors, *Proceedings of the 35th International Conference on Machine Learning*, volume 80 of *Proceedings of Machine Learning Research*, pages 5561–5570. PMLR, 2018.
- J. Yang, V. Rao, and J. Neville. A Stein–Papangelou goodness-of-fit test for point processes. In Kamalika Chaudhuri and Masashi Sugiyama, editors, *Proceedings of the 22nd International Conference on Artificial Intelligence and Statistics*, volume 89 of *Proceedings of Machine Learning Research*, pages 226–235. PMLR, 2019.

A Technical results

In the following, we present technical results that are used to prove the main results of the paper. Unless specified otherwise, the notation used in this section matches that of the main paper.

Lemma A.1. *Let \mathcal{H}_K be an RKHS associated with kernel K . Define $\mathcal{A}_Q : \mathcal{H}_K \rightarrow \mathcal{H}_K$,*

$$\mathcal{A}_Q := \int_{\mathcal{X}} K(\cdot, x) \otimes_{\mathcal{H}_K} K(\cdot, x) dQ(x),$$

where Q is a probability measure that can be either P_0 or P ,

$$\mathcal{N}_1(\lambda) := \text{Tr}(\mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{A}_{P_0} \mathcal{A}_{P_0, \lambda}^{-1/2}), \text{ and } \mathcal{N}_2(\lambda) := \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{A}_{P_0} \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_K)},$$

where $\mathcal{A}_{P, \lambda} := \mathcal{A}_P + \lambda \mathbf{I}$. Suppose

$$\int_{\mathcal{X}} K(x, x) dQ(x) < \infty, \text{ and } \int_{\mathcal{X}} [K(x, x)]^v dP_0(x) < \kappa^v,$$

for some $\kappa > 0$ and $v \geq 2$. Then the following hold:

$$(i) \left\| \mathcal{A}_{P, \lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P, \lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_K)}^2 \leq \frac{\text{Tr}(\mathcal{A}_P)}{\lambda};$$

(ii) If $4C_\lambda \|u\|_{L^2(P_0)}^2 \leq 1$, then

$$(a) \left\| \mathcal{A}_{P,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P,\lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_K)}^2 \leq 8\mathcal{N}_2^2(\lambda) + 2,$$

$$(b) \operatorname{Tr}(\mathcal{A}_{P,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P,\lambda}^{-1/2}) \leq 2\mathcal{N}_1(\lambda) + 2C_\lambda \|u\|_{L^2(P_0)},$$

where $u := \frac{dP}{dP_0} - 1$ and

$$C_\lambda = \begin{cases} \mathcal{N}_1(\lambda) \sup_i \|\phi_i\|_\infty^2, & \sup_i \|\phi_i\|_\infty^2 < \infty \\ (\mathcal{N}_1(\lambda))^{\frac{v-2}{2(v-1)}} \left(\frac{\kappa}{\lambda}\right)^{\frac{v}{v-1}}, & \text{otherwise} \end{cases},$$

with $\phi_i := \frac{\alpha_i}{\sqrt{\lambda_i}}$ and $(\lambda_i, \alpha_i)_i$ being the orthonormal eigensystem of \mathcal{A}_{P_0} .

Proof. Observe that $\mathcal{A}_{P_0} = \mathfrak{J}^* \mathfrak{J}$, where $\mathfrak{J} : \mathcal{H}_K \rightarrow L^2(P_0)$, $f \mapsto [f]_\sim$ and $\mathfrak{J}^* : L^2(P_0) \rightarrow \mathcal{H}_K$, $f \mapsto \int K(\cdot, x) f(x) dP_0(x)$. Then it can be verified that $\Upsilon_{P_0} := \mathfrak{J} \mathfrak{J}^* : L^2(P_0) \rightarrow L^2(P_0)$, $f \mapsto \int K(\cdot, x) f(x) dP_0(x)$. Note that

$$\begin{aligned} \operatorname{Tr}(\Upsilon_{P_0}) &= \operatorname{Tr}(\mathcal{A}_{P_0}) = \int_x \operatorname{Tr}(K(\cdot, x) \otimes_{\mathcal{H}_K} K(\cdot, x)) dP_0(x) = \int_x \langle K(\cdot, x), K(\cdot, x) \rangle_{\mathcal{H}_K} dP_0(x) \\ &= \int_{\mathcal{X}} K(x, x) dP_0(x) < \infty, \end{aligned}$$

which implies Υ_{P_0} and \mathcal{A}_{P_0} (similarly \mathcal{A}_P) are trace class operators. Let $(\lambda_i, \tilde{\phi}_i)_i$ be the eigenvalues and eigenfunctions of Υ_{P_0} . Since $\Upsilon_{P_0} = \mathfrak{J} \mathfrak{J}^*$ and $\mathcal{A}_{P_0} = \mathfrak{J}^* \mathfrak{J}$, we have $\mathfrak{J} \mathfrak{J}^* \tilde{\phi}_i = \lambda_i \tilde{\phi}_i$ which implies $\mathfrak{J}^* \mathfrak{J} (\mathfrak{J}^* \tilde{\phi}_i) = \lambda_i (\mathfrak{J}^* \tilde{\phi}_i)$, i.e., $\mathcal{A}_{P_0} \alpha_i = \lambda_i \alpha_i$, where $\alpha_i := \mathfrak{J}^* \tilde{\phi}_i / \sqrt{\lambda_i}$. Note that $(\alpha_i)_i$ form an orthonormal system in \mathcal{H}_K . Define $\phi_i = \frac{\tilde{\phi}_i}{\lambda_i}$, thus $\alpha_i = \sqrt{\lambda_i} \phi_i$. Based on these notations, we present the proofs of the desired results as follows:

(i) Observe that

$$\left\| \mathcal{A}_{P,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P,\lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_K)}^2 \leq \left\| \mathcal{A}_{P,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \operatorname{Tr} \left(\mathcal{A}_{P,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P,\lambda}^{-1/2} \right) \leq \frac{\operatorname{Tr}(\mathcal{A}_P)}{\lambda}.$$

(ii)-(a) Let

$$\mathcal{B} = \int_{\mathcal{X}} K(\cdot, x) \otimes_{\mathcal{H}_K} K(\cdot, x) u(x) dP_0(x).$$

Then it is clear that $\mathcal{A}_P = \mathcal{A}_{P_0} + \mathcal{B}$ and we have

$$\begin{aligned} & \left\| \mathcal{A}_{P,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P,\lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_K)}^2 \leq \left\| \mathcal{A}_{P,\lambda}^{-1/2} \mathcal{A}_{P_0,\lambda}^{1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)}^4 \left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P_0,\lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_K)}^2 \\ & \leq \left\| \mathcal{A}_{P_0,\lambda}^{1/2} \mathcal{A}_{P,\lambda}^{-1} \mathcal{A}_{P_0,\lambda}^{1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)}^2 \left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P_0,\lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_K)}^2 \\ & \leq \left\| \left(\mathbf{I} - \mathcal{A}_{P_0,\lambda}^{-1/2} (\mathcal{A}_{P_0} - \mathcal{A}_P) \mathcal{A}_{P_0,\lambda}^{-1/2} \right)^{-1} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)}^2 \left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P_0,\lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_K)}^2 \\ & \stackrel{(*)}{\leq} \left(1 - \left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{B} \mathcal{A}_{P_0,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \right)^{-2} \left(2\mathcal{N}_2^2(\lambda) + 2 \left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{B} \mathcal{A}_{P_0,\lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_K)}^2 \right) \\ & \stackrel{(\dagger)}{\leq} \left(1 - \sqrt{C_\lambda} \|u\|_{L^2(P_0)} \right)^{-2} \left(2\mathcal{N}_2^2(\lambda) + 2C_\lambda \|u\|_{L^2(P_0)}^2 \right) \end{aligned}$$

$$\leq 8\mathcal{N}_2^2(\lambda) + 2,$$

where (*) follows when $\left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{B} \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} < 1$. By using

$$\left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{B} \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \leq \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{B} \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_K)},$$

(†) follows if

$$\left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{B} \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_K)} \leq \sqrt{C_\lambda} \|u\|_{L^2(P_0)}, \quad (\text{A.1})$$

which we show below for the two cases of C_λ .

Case 1: $\sup_i \|\phi_i\|_\infty^2 < \infty$. Then we have,

$$\begin{aligned} & \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{B} \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_K)}^2 = \text{Tr} \left(\mathcal{A}_{P_0, \lambda}^{-1} \mathcal{B} \mathcal{A}_{P_0, \lambda}^{-1} \mathcal{B} \right) \\ &= \int_{\mathcal{X}} \int_{\mathcal{X}} \text{Tr} \left[\mathcal{A}_{P_0, \lambda}^{-1} K(\cdot, x) \otimes_{\mathcal{H}_K} K(\cdot, x) \mathcal{A}_{P_0, \lambda}^{-1} K(\cdot, y) \otimes_{\mathcal{H}_K} K(\cdot, y) \right] u(x) u(y) dP_0(y) dP_0(x) \\ &\stackrel{(\ddagger)}{=} \sum_{i, j} \left(\frac{\lambda_i}{\lambda_i + \lambda} \right) \left(\frac{\lambda_j}{\lambda_j + \lambda} \right) \left[\int_{\mathcal{X}} \phi_i(x) \phi_j(x) u(x) dP_0(x) \right]^2 \\ &\leq \sum_{i, j} \frac{\lambda_i}{\lambda_i + \lambda} \langle \phi_j, \phi_i u \rangle_{L^2(P_0)}^2 \leq \sum_i \frac{\lambda_i}{\lambda_i + \lambda} \|\phi_i u\|_{L^2(P_0)}^2 \leq \mathcal{N}_1(\lambda) \sup_i \|\phi_i\|_\infty^2 \|u\|_{L^2(P_0)}^2, \end{aligned}$$

where (‡) follows from

$$\begin{aligned} & \text{Tr} \left[\mathcal{A}_{P_0, \lambda}^{-1} K(\cdot, x) \otimes_{\mathcal{H}_K} K(\cdot, x) \mathcal{A}_{P_0, \lambda}^{-1} K(\cdot, y) \otimes_{\mathcal{H}_K} K(\cdot, y) \right] = \left\langle K(\cdot, y), \mathcal{A}_{P_0, \lambda}^{-1} K(\cdot, x) \right\rangle_{\mathcal{H}_K}^2 \\ &= \left\langle K(\cdot, y), \sum_i \frac{1}{\lambda_i + \lambda} \left\langle K(\cdot, x), \sqrt{\lambda_i} \phi_i \right\rangle_{\mathcal{H}_K} \sqrt{\lambda_i} \phi_i \right. \\ &\quad \left. + \frac{1}{\lambda} \left(K(\cdot, x) - \sum_i \left\langle K(\cdot, x), \sqrt{\lambda_i} \phi_i \right\rangle_{\mathcal{H}_K} \sqrt{\lambda_i} \phi_i \right) \right\rangle_{\mathcal{H}_K}^2 \\ &= \left[\sum_i \frac{\lambda_i}{\lambda_i + \lambda} \phi_i(x) \phi_i(y) + \frac{1}{\lambda} \left(\langle K(\cdot, x), K(\cdot, y) \rangle_{\mathcal{H}_K} - \sum_i \lambda_i \phi_i(x) \phi_i(y) \right) \right]^2 \\ &\stackrel{(\star)}{=} \left[\sum_i \frac{\lambda_i}{\lambda_i + \lambda} \phi_i(x) \phi_i(y) \right]^2, \quad (\text{A.2}) \end{aligned}$$

where in (★) we used $\langle K(\cdot, x), K(\cdot, y) \rangle_{\mathcal{H}} = \sum_i \lambda_i \phi_i(x) \phi_i(y)$ which follows by Mercer's theorem (see Steinwart and Scovel 2012, Lemma 2.6).

Case 2: Suppose $\sup_i \|\phi_i\|_\infty^2$ is not finite. Then

$$\begin{aligned} & \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{B} \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_K)}^2 = \text{Tr} \left(\mathcal{A}_{P_0, \lambda}^{-1} \mathcal{B} \mathcal{A}_{P_0, \lambda}^{-1} \mathcal{B} \right) \\ &= \int_{\mathcal{X}} \int_{\mathcal{X}} \text{Tr} \left[\mathcal{A}_{P_0, \lambda}^{-1} K(\cdot, x) \otimes_{\mathcal{H}_K} K(\cdot, x) \mathcal{A}_{P_0, \lambda}^{-1} K(\cdot, y) \otimes_{\mathcal{H}_K} K(\cdot, y) \right] u(x) u(y) dP_0(y) dP_0(x) \\ &= \int_{\mathcal{X}} \int_{\mathcal{X}} \left\langle K(\cdot, y), \mathcal{A}_{P_0, \lambda}^{-1} K(\cdot, x) \right\rangle_{\mathcal{H}_K}^2 u(x) u(y) dP_0(x) dP_0(y) \end{aligned}$$

$$\begin{aligned}
&\leq \left(\int_{\mathcal{X}} \int_{\mathcal{X}} \left\langle K(\cdot, y), \mathcal{A}_{P_0, \lambda}^{-1} K(\cdot, x) \right\rangle_{\mathcal{H}_K}^4 dP_0(x) dP_0(y) \right)^{1/2} \|u\|_{L^2(P_0)}^2 \\
&= \left(\int_{\mathcal{X}} \int_{\mathcal{X}} \left\langle K(\cdot, y), \mathcal{A}_{P_0, \lambda}^{-1} K(\cdot, x) \right\rangle_{\mathcal{H}_K}^{\frac{2v}{v-1}} \left\langle K(\cdot, y), \mathcal{A}_{P_0, \lambda}^{-1} K(\cdot, x) \right\rangle_{\mathcal{H}_K}^{\frac{2(v-2)}{v-1}} dP_0(x) dP_0(y) \right)^{1/2} \|u\|_{L^2(P_0)}^2 \\
&\leq \left(\int_{\mathcal{X}} \int_{\mathcal{X}} \left\| \mathcal{A}_{P_0, \lambda}^{-1} K(\cdot, x) \right\|_{\mathcal{H}_K}^{\frac{2v}{v-1}} \left\| K(\cdot, y) \right\|_{\mathcal{H}_K}^{\frac{2v}{v-1}} \left\langle K(\cdot, y), \mathcal{A}_{P_0, \lambda}^{-1} K(\cdot, x) \right\rangle_{\mathcal{H}_K}^{\frac{2(v-2)}{v-1}} dP_0(x) dP_0(y) \right)^{1/2} \\
&\quad \times \|u\|_{L^2(P_0)}^2 \\
&\leq \frac{\|u\|_{L^2(P_0)}^2}{\lambda^{\frac{v}{v-1}}} \left(\int_{\mathcal{X}} \int_{\mathcal{X}} [K(x, x)K(y, y)]^{\frac{v}{v-1}} \left\langle K(\cdot, y), \mathcal{A}_{P_0, \lambda}^{-1} K(\cdot, x) \right\rangle_{\mathcal{H}_K}^{\frac{2(v-2)}{v-1}} dP_0(x) dP_0(y) \right)^{1/2} \\
&\leq \frac{\|u\|_{L^2(P_0)}^2}{\lambda^{\frac{v}{v-1}}} \left(\int_{\mathcal{X}} \int_{\mathcal{X}} [K(x, x)K(y, y)]^v dP_0(x) dP_0(y) \right)^{\frac{1}{2(v-1)}} \\
&\quad \times \left(\mathbb{E}_{P_0} \left\langle K(\cdot, Y), \mathcal{A}_{P_0, \lambda}^{-1} K(\cdot, X) \right\rangle_{\mathcal{H}_K}^2 \right)^{\frac{v-2}{2(v-1)}} \\
&\leq \|u\|_{L^2(P_0)}^2 \left(\frac{\kappa}{\lambda} \right)^{\frac{v}{v-1}} \left(\mathbb{E}_{P_0} \text{Tr} \left[\mathcal{A}_{P_0, \lambda}^{-1} K(\cdot, X) \otimes_{\mathcal{H}_K} K(\cdot, X) \mathcal{A}_{P_0, \lambda}^{-1} K(\cdot, Y) \otimes_{\mathcal{H}_K} K(\cdot, Y) \right] \right)^{\frac{v-2}{2(v-1)}} \\
&\leq \|u\|_{L^2(P_0)}^2 \left(\frac{\kappa}{\lambda} \right)^{\frac{v}{v-1}} \left(\text{Tr}(\mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{A}_{P_0} \mathcal{A}_{P_0, \lambda}^{-1/2}) \right)^{\frac{v-2}{2(v-1)}} = C_\lambda \|u\|_{L^2(P_0)}^2.
\end{aligned}$$

(ii)-(b) Note that

$$\begin{aligned}
\text{Tr}(\mathcal{A}_{P, \lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P, \lambda}^{-1/2}) &\leq \left\| \mathcal{A}_{P, \lambda}^{-1/2} \mathcal{A}_{P_0, \lambda}^{1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)}^2 \text{Tr}(\mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P_0, \lambda}^{-1/2}) \\
&\leq \left(1 - \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{B} \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \right)^{-1} \left(\mathcal{N}_1(\lambda) + \text{Tr}(\mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{B} \mathcal{A}_{P_0, \lambda}^{-1/2}) \right) \\
&\stackrel{(*)}{\leq} \left(1 - \sqrt{C_\lambda} \|u\|_{L^2(P_0)} \right)^{-1} \left(\mathcal{N}_1(\lambda) + C_\lambda \|u\|_{L^2(P_0)} \right) \\
&\stackrel{(\dagger)}{\leq} 2\mathcal{N}_1(\lambda) + 2C_\lambda \|u\|_{L^2(P_0)},
\end{aligned}$$

where (*) follows using $\left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{B} \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \leq \sqrt{C_\lambda} \|u\|_{L^2(P_0)}$ as in (ii)-(a), and if

$$\text{Tr}(\mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{B} \mathcal{A}_{P_0, \lambda}^{-1/2}) \leq C_\lambda \|u\|_{L^2(P_0)}$$

holds, which we show below.

Case 1: $\sup_i \|\phi_i\|_\infty^2 < \infty$. Then we have,

$$\begin{aligned}
\text{Tr}(\mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{B} \mathcal{A}_{P_0, \lambda}^{-1/2}) &= \int_{\mathcal{X}} \text{Tr}(\mathcal{A}_{P_0, \lambda}^{-1/2} K(\cdot, x) \otimes_{\mathcal{H}_K} K(\cdot, x) \mathcal{A}_{P_0, \lambda}^{-1/2}) u(x) dP_0(x) \\
&\stackrel{(*)}{=} \sum_i \frac{\lambda_i}{\lambda_i + \lambda} \int_{\mathcal{X}} \phi_i^2(x) u(x) dP_0(x) \\
&\leq \mathcal{N}_1(\lambda) \|u\|_{L^2(P_0)} \sup_i \|\phi_i\|_\infty^2 = C_\lambda \|u\|_{L^2(P_0)},
\end{aligned}$$

where (*) follows similar to the proof of (A.2).

Case 2: Suppose $\sup_i \|\phi_i\|_\infty^2$ is not finite. Then

$$\begin{aligned}
& \text{Tr}(\mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{B} \mathcal{A}_{P_0,\lambda}^{-1/2}) \\
&= \int_{\mathcal{X}} \left\langle \mathcal{A}_{P_0,\lambda}^{-1} K(\cdot, x), K(\cdot, x) \right\rangle_{\mathcal{H}_K} u(x) dP_0(x) \\
&\leq \|u\|_{L^2(P_0)} \left(\int_{\mathcal{X}} \left\langle \mathcal{A}_{P_0,\lambda}^{-1} K(\cdot, x), K(\cdot, x) \right\rangle_{\mathcal{H}_K}^2 dP_0(x) \right)^{1/2} \\
&\leq \|u\|_{L^2(P_0)} \left(\int_{\mathcal{X}} \left\langle \mathcal{A}_{P_0,\lambda}^{-1} K(\cdot, x), K(\cdot, x) \right\rangle_{\mathcal{H}_K}^v dP_0(x) \right)^{\frac{1}{2(v-1)}} \left(\mathbb{E}_{P_0} \left\langle \mathcal{A}_{P_0,\lambda}^{-1} K(\cdot, x), K(\cdot, x) \right\rangle_{\mathcal{H}_K} \right)^{\frac{v-2}{2(v-1)}} \\
&\leq \|u\|_{L^2(P_0)} \left(\frac{\kappa}{\lambda} \right)^{\frac{v}{2(v-1)}} \left(\text{Tr}(\mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{A}_{P_0} \mathcal{A}_{P_0,\lambda}^{-1/2}) \right)^{\frac{v-2}{2(v-1)}} \leq C_\lambda \|u\|_{L^2(P_0)}
\end{aligned}$$

and the result follows. \square

Lemma A.2. Let $\mathbb{S}_\lambda^P := \left\| g_\lambda^{1/2}(\mathcal{A}_P) \Psi_P \right\|_{\mathcal{H}_{K_0}}^2$ and $\mathbb{S}_\lambda^{P_0} := \left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \Psi_P \right\|_{\mathcal{H}_{K_0}}^2$, where $\mathcal{A}_{P_0,\lambda} = \mathcal{A}_{P_0} + \lambda \mathbf{I}$ and g_λ satisfies (E₁)–(E₄). Then

$$\mathbb{S}_\lambda^P \geq \frac{C_4 \mathbb{S}_\lambda^{P_0}}{1 + \sqrt{C_\lambda} \|u\|_{L^2(P_0)}},$$

where C_λ is defined in Lemma A.1 with $K = K_0$.

Proof. By applying (Hagrass et al., 2022, Lemma A.10, Lemma A.8(iv))—by replacing Σ_{PQ} with \mathcal{A}_P in those results—, we have

$$\mathbb{S}_\lambda^P \geq \left\| \mathcal{A}_{P,\lambda}^{-1/2} g_\lambda^{-1/2}(\mathcal{A}_P) \right\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^{-2} \left\| \mathcal{A}_{P,\lambda}^{-1/2} \Psi_P \right\|_{\mathcal{H}_{K_0}}^2 \geq C_4 \left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{A}_{P,\lambda}^{1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^{-2} \mathbb{S}_\lambda^{P_0}.$$

Next we upper bound $\left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{A}_{P,\lambda}^{1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2$ as shown below. Let $\mathcal{B} = \mathcal{A}_P - \mathcal{A}_{P_0}$. Then

$$\begin{aligned}
\left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{A}_{P,\lambda}^{1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 &= \left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{A}_{P,\lambda} \mathcal{A}_{P_0,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})} = \left\| \mathbf{I} + \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{B} \mathcal{A}_{P_0,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})} \\
&\leq 1 + \left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{B} \mathcal{A}_{P_0,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})} \leq 1 + \left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{B} \mathcal{A}_{P_0,\lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_{K_0})} \\
&\stackrel{(*)}{\leq} 1 + \sqrt{C_\lambda} \|u\|_{L^2(P_0)},
\end{aligned}$$

where (*) follows from (A.1). \square

Lemma A.3. Let $\zeta = \left\| g_\lambda^{1/2}(\hat{\mathcal{A}}_P) \Psi_P \right\|_{\mathcal{H}_{K_0}}^2$, $\mathcal{W} = \hat{\mathcal{A}}_{P,\lambda}^{-1/2} \mathcal{A}_{P,\lambda}^{1/2}$ and $\mathcal{V} = \hat{\mathcal{A}}_{P,\lambda}^{-1/2} \mathcal{A}_{P_0,\lambda}^{1/2}$, where g_λ satisfies (E₁)–(E₄). Then

(i)

$$\mathbb{E} \left((\hat{\mathbb{S}}_\lambda^P - \zeta)^2 | \mathbb{Z}_{n_2} \right) \leq c \left(\frac{\|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \zeta}{n_1} + \frac{\|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^4 D_\lambda}{n_1^2} \right),$$

where $c > 0$ is a constant, $D_\lambda = \begin{cases} \mathcal{N}_2^2(\lambda), & 4C_\lambda \|u\|_{L^2(P_0)}^2 \leq 1 \\ \lambda^{-1}, & \text{otherwise} \end{cases}$, and C_λ is defined in Lemma A.1 with $K = K_0$;

(ii)

$$\mathbb{E} \left((\hat{\mathbb{S}}_\lambda^P - \zeta)^2 | \mathbb{Z}_{n_2} \right) \leq \tilde{c} \left(\frac{\|\mathcal{V}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \zeta}{n_1} + \frac{\|\mathcal{V}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^4 \mathcal{N}_2^2(\lambda)}{n_1^2} \right),$$

if $4C_\lambda \|u\|_{L^2(P_0)}^2 < 1$, where $\tilde{c} > 0$ is a constant.

Proof. Let $a(x) = \mathcal{B}\mathcal{A}_{P,\lambda}^{-1/2}(K_0(\cdot, x) - \Psi_P)$ and $\mathcal{B} = g_\lambda^{1/2}(\hat{\mathcal{A}}_P)\mathcal{A}_{P,\lambda}^{1/2}$. Then

$$\hat{\mathbb{S}}_\lambda^P = \frac{1}{n_1(n_1 - 1)} \sum_{i \neq j} \langle a(X_i), a(X_j) \rangle_{\mathcal{H}_{K_0}} + \frac{2}{n_1} \sum_{i=1}^{n_1} \left\langle a(X_i), \mathcal{B}\mathcal{A}_{P,\lambda}^{-1/2}\Psi_P \right\rangle_{\mathcal{H}_{K_0}} + \zeta.$$

Thus we have $\hat{\mathbb{S}}_\lambda^P - \zeta = \textcircled{1} + \textcircled{2}$, where $\textcircled{1} := \frac{1}{n_1(n_1-1)} \sum_{i \neq j} \langle a(X_i), a(X_j) \rangle_{\mathcal{H}_{K_0}}$ and $\textcircled{2} := \frac{2}{n_1} \sum_{i=1}^{n_1} \left\langle a(X_i), \mathcal{B}\mathcal{A}_{P,\lambda}^{-1/2}\Psi_P \right\rangle_{\mathcal{H}_{K_0}}$. Next, we bound each term as follows:

(i)

$$\begin{aligned} \mathbb{E}(\textcircled{1}^2 | \mathbb{Z}_{n_2}) &\stackrel{(*)}{=} \frac{2}{n_1^2(n_1 - 1)^2} \sum_{i \neq j} \mathbb{E} \langle a(X_i), a(X_j) \rangle_{\mathcal{H}_{K_0}}^2 \\ &\leq \frac{4}{n_1^2} \left\| \mathcal{B}\mathcal{A}_{P,\lambda}^{-1/2} \mathbb{E}_P[(K_0(\cdot, X) - \Psi_P) \otimes_{\mathcal{H}_{K_0}} (K_0(\cdot, X) - \Psi_P)] \mathcal{A}_{P,\lambda}^{-1/2} \mathcal{B}^* \right\|_{\mathcal{L}^2(\mathcal{H}_{K_0})}^2 \\ &\stackrel{(\dagger)}{\leq} \frac{4}{n_1^2} (C_1 + C_2)^2 \|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^4 \left\| \mathcal{A}_{P,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P,\lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_{K_0})}^2, \end{aligned}$$

where $(*)$ follows from (Hagrass et al., 2022, Lemma A.3(ii)) and in (\dagger) we used (Hagrass et al., 2022, Lemma A.8 (iii)) and that

$$\mathbb{E}_P((K_0(\cdot, X_i) - \Psi_P) \otimes_{\mathcal{H}_{K_0}} (K_0(\cdot, X_i) - \Psi_P)) = \mathcal{A}_P - \Psi_P \otimes_{\mathcal{H}_{K_0}} \Psi_P \preceq \mathcal{A}_P.$$

For term $\textcircled{2}$ we have

$$\begin{aligned} \mathbb{E}(\textcircled{2}^2 | \mathbb{Z}_{n_2}) &\leq \frac{4}{n_1^2} \sum_{i=1}^{n_1} \mathbb{E} \left\langle a(X_i), \mathcal{B}\mathcal{A}_{P,\lambda}^{-1/2}\Psi_P \right\rangle_{\mathcal{H}_{K_0}}^2 \\ &= \frac{4}{n_1^2} \sum_{i=1}^{n_1} \mathbb{E} \left\langle a(X_i) \otimes_{\mathcal{H}_{K_0}} a(X_i), \mathcal{B}\mathcal{A}_{P,\lambda}^{-1/2}\Psi_P \otimes_{\mathcal{H}_{K_0}} \Psi_P \mathcal{A}_{P,\lambda}^{-1/2} \mathcal{B}^* \right\rangle_{\mathcal{L}^2(\mathcal{H}_{K_0})} \\ &\leq \frac{4}{n_1} \text{Tr} \left(\mathcal{A}_{P,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P,\lambda}^{-1/2} \mathcal{B}^* \mathcal{B} \mathcal{A}_{P,\lambda}^{-1/2} \Psi_P \otimes_{\mathcal{H}_{K_0}} \Psi_P \mathcal{A}_{P,\lambda}^{-1/2} \mathcal{B}^* \mathcal{B} \right) \\ &\leq \frac{4}{n_1} \left\| \mathcal{A}_{P,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})} (C_1 + C_2) \|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \text{Tr} \left(\mathcal{B}\mathcal{A}_{P,\lambda}^{-1/2}\Psi_P \otimes_{\mathcal{H}_{K_0}} \Psi_P \mathcal{A}_{P,\lambda}^{-1/2} \mathcal{B}^* \right) \\ &= \frac{4}{n_1} (C_1 + C_2) \|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_K)}^2 \zeta. \end{aligned}$$

Using the above bounds and Lemma A.1 yields the desired result.

(ii) Consider

$$\mathbb{E}(\textcircled{1}^2 | \mathbb{Z}_{n_2}) = \frac{2}{n_1^2(n_1 - 1)^2} \sum_{i \neq j} \mathbb{E} \langle a(X_i), a(X_j) \rangle_{\mathcal{H}_{K_0}}^2$$

$$\begin{aligned}
&\leq \frac{4}{n_1^2} \left\| g_\lambda^{1/2}(\hat{\mathcal{A}}_P) \mathbb{E}_P[(K_0(\cdot, X) - \Psi_P) \otimes_{\mathcal{H}_{K_0}} (K_0(\cdot, X) - \Psi_P)] g_\lambda^{1/2}(\hat{\mathcal{A}}_P) \right\|_{\mathcal{L}^2(\mathcal{H}_{K_0})}^2 \\
&\leq \frac{4}{n_1^2} (C_1 + C_2)^2 \|\mathcal{V}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^4 \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_{K_0})}^2 \\
&\leq \frac{4 \|\mathcal{V}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^4 (C_1 + C_2)^2}{n_1^2} \left(2 \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{A}_{P_0} \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_{K_0})}^2 \right. \\
&\quad \left. + 2 \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} (\mathcal{A}_P - \mathcal{A}_{P_0}) \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_{K_0})}^2 \right) \\
&\stackrel{(*)}{\leq} \frac{4 \|\mathcal{V}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^4 (C_1 + C_2)^2}{n_1^2} \left(2\mathcal{N}_2^2(\lambda) + 2C_\lambda \|u\|_{L^2(P_0)}^2 \right) \\
&\lesssim \frac{4 \|\mathcal{V}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^4 \mathcal{N}_2^2(\lambda)}{n_1^2},
\end{aligned}$$

where (*) follow from (A.1) and

$$\begin{aligned}
\mathbb{E}(\textcircled{2}^2 | \mathbb{Z}_{n_2}) &\leq \frac{4}{n_1} \text{Tr} \left(\mathcal{A}_{P, \lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P, \lambda}^{-1/2} \mathcal{B}^* \mathcal{B} \mathcal{A}_{P, \lambda}^{-1/2} \Psi_P \otimes_{\mathcal{H}_{K_0}} \Psi_P \mathcal{A}_{P, \lambda}^{-1/2} \mathcal{B}^* \mathcal{B} \right) \\
&\leq \frac{4}{n_1} \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})} (C_1 + C_2) \|\mathcal{V}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \text{Tr} \left(\mathcal{B} \mathcal{A}_{P, \lambda}^{-1/2} \Psi_P \otimes_{\mathcal{H}_{K_0}} \Psi_P \mathcal{A}_{P, \lambda}^{-1/2} \mathcal{B}^* \right) \\
&\leq \frac{4 \|\mathcal{V}\|_{\mathcal{L}^\infty(\mathcal{H}_K)}^2 \zeta (C_1 + C_2)}{n_1} \left(\left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{A}_{P_0} \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})} + \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} (\mathcal{A}_P - \mathcal{A}_{P_0}) \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})} \right) \\
&\stackrel{(\dagger)}{\leq} \frac{4 \|\mathcal{V}\|_{\mathcal{L}^\infty(\mathcal{H}_K)}^2 \zeta (C_1 + C_2)}{n_1} \left(1 + \sqrt{C_\lambda} \|u\|_{L^2(P_0)} \right) \lesssim \frac{\|\mathcal{V}\|_{\mathcal{L}^\infty(\mathcal{H}_K)}^2 \zeta}{n_1},
\end{aligned}$$

where we used (A.1) in (\dagger). □

Lemma A.4. *Suppose there exist some $c > 0$ and $\kappa > 0$ such that for any $r > 2$,*

$$\int_{\mathcal{X}} [K(x, x)]^r dP(x) \leq cr! \kappa^r,$$

and one of the following hold:

- (i) $\lambda \geq \frac{16\kappa}{\sqrt{n}} \log \frac{2}{\delta} \max\{\sqrt{c}, 1\}$.
- (ii) $4C_\lambda \|u\|_{L^2(P_0)}^2 < 1$, $n \geq 4$, $\mathbb{E}_{P_0}[K(X, X)]^v < \tilde{\kappa}^v$, and $\lambda \geq \frac{d(\log \frac{2}{\delta})(\log n)}{n} \left(\mathcal{N}_1(\lambda) + C_\lambda \|u\|_{L^2(P_0)} \right)$, for some $d > 0$, $v > 2$ and C_λ is defined in Lemma A.1 (with κ replaced by $\tilde{\kappa}$).
- (iii) $\sup_x K(x, x) \leq \kappa$ and $\frac{d_1 \log \frac{d_2 n}{\delta}}{n} \leq \lambda \leq \|\mathcal{A}_P\|_{\mathcal{L}^\infty(\mathcal{H}_K)}$, for some $d_1 > 0$ and $d_2 > 0$.

Then for $0 < \delta \leq 1$,

- (a) $P \left(\left\| \mathcal{A}_{P, \lambda}^{-1/2} (\mathcal{A}_P - \hat{\mathcal{A}}_P) \mathcal{A}_{P, \lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \leq \frac{1}{2} \right) \geq 1 - \delta$;
- (b) $P \left(\left\| \mathcal{A}_{P, \lambda}^{1/2} \hat{\mathcal{A}}_{P, \lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \leq \sqrt{2} \right) \geq 1 - \delta$;

$$(c) P \left(\left\| \mathcal{A}_{P,\lambda}^{-1/2} \hat{\mathcal{A}}_{P,\lambda}^{1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \leq \sqrt{\frac{3}{2}} \right) \geq 1 - \delta,$$

where \mathcal{A}_P is defined in Lemma A.1.

Proof. Let $\xi_i = \mathcal{A}_{P,\lambda}^{-1/2} ((K(\cdot, Z_i) \otimes_{\mathcal{H}_K} K(\cdot, Z_i)) - \mathcal{A}_P) \mathcal{A}_{P,\lambda}^{-1/2}$. Then $\frac{1}{n} \sum_{i=1}^n \xi_i = \mathcal{A}_{P,\lambda}^{-1/2} (\hat{\mathcal{A}}_P - \mathcal{A}_P) \mathcal{A}_{P,\lambda}^{-1/2}$ and $\mathbb{E}_P(\xi_i) = 0$. The main idea is to apply Bernstein's inequality for Hilbert-valued random elements as stated in Sriperumbudur and Sterge (2022, Theorem D.1).

Let $\theta^2 = 2c^{1-q} \left(\frac{d_q}{\lambda} \right)^{2-q} \left[\text{Tr}(\mathcal{A}_{P,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P,\lambda}^{-1/2}) \right]^q$, $B = \frac{d_q}{\lambda}$ and $d_q = \frac{2\kappa}{1-q}$, for some $0 \leq q < 1$. Then we bound $\mathbb{E} \|\xi_1\|_{\mathcal{L}^2(\mathcal{H}_K)}^r$ as follows.

$$\begin{aligned} \mathbb{E} \|\xi_1\|_{\mathcal{L}^2(\mathcal{H}_K)}^r &\stackrel{(*)}{\leq} \mathbb{E} \left(\left\| \mathcal{A}_{P,\lambda}^{-1/2} (K(\cdot, Z_1) \otimes_{\mathcal{H}_K} K(\cdot, Z_1)) \mathcal{A}_{P,\lambda}^{-1/2} \right\|_{\mathcal{L}^2(\mathcal{H}_K)}^2 \right)^{\frac{r}{2}} \\ &= \mathbb{E} \left(\left\langle \mathcal{A}_{P,\lambda}^{-1/2} K(\cdot, Z_1), \mathcal{A}_{P,\lambda}^{-1/2} K(\cdot, Z_1) \right\rangle_{\mathcal{H}_K}^2 \right)^{\frac{r}{2}} \\ &= \mathbb{E} \left(\text{Tr} \left(\mathcal{A}_{P,\lambda}^{-1} K(\cdot, Z_1) \otimes_{\mathcal{H}_K} K(\cdot, Z_1) \mathcal{A}_{P,\lambda}^{-1} K(\cdot, Z_1) \otimes_{\mathcal{H}_K} K(\cdot, Z_1) \right) \right)^{\frac{r}{2}-q+q} \\ &\stackrel{(\dagger)}{\leq} \left[\mathbb{E} \left(\left\langle \mathcal{A}_{P,\lambda}^{-1} K(\cdot, Z_1), K(\cdot, Z_1) \right\rangle_{\mathcal{H}_K}^2 \right)^{\frac{2(r/2-q)}{1-q}} \right]^{\frac{1-q}{2}} \\ &\quad \times \left[\mathbb{E} \left(\text{Tr} \left(\mathcal{A}_{P,\lambda}^{-1} K(\cdot, Z_1) \otimes_{\mathcal{H}_K} K(\cdot, Z_1) \mathcal{A}_{P,\lambda}^{-1} K(\cdot, Z_1) \otimes_{\mathcal{H}_K} K(\cdot, Z_1) \right) \right)^{\frac{2q}{1+q}} \right]^{\frac{1+q}{2}} \\ &\leq \frac{1}{\lambda^{r-2q}} \left[\mathbb{E}(K(Z_1, Z_1))^{\frac{2(r-2q)}{1-q}} \right]^{\frac{1-q}{2}} \\ &\quad \times \left[\mathbb{E} \left(\left\| \mathcal{A}_{P,\lambda}^{-1/2} K(\cdot, Z_1) \otimes_{\mathcal{H}_K} K(\cdot, Z_1) \mathcal{A}_{P,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \text{Tr} \left(\mathcal{A}_{P,\lambda}^{-1} K(\cdot, Z_1) \otimes_{\mathcal{H}_K} K(\cdot, Z_1) \right) \right)^{\frac{2q}{1+q}} \right]^{\frac{1+q}{2}} \\ &\stackrel{(\ddagger)}{\leq} \frac{1}{\lambda^{r-q}} \left[\mathbb{E}(K(Z_1, Z_1))^{\frac{2(r-2q)}{1-q}} \right]^{\frac{1-q}{2}} \left[\mathbb{E}(K(Z_1, Z_1))^{\frac{2q}{1-q}} \right]^{\frac{1-q}{2}} \left[\text{Tr}(\mathcal{A}_{P,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P,\lambda}^{-1/2}) \right]^q \stackrel{(**)}{\leq} \frac{r!}{2} \theta^2 B^{r-2}, \end{aligned}$$

where $(*)$ follows since \mathcal{A}_P is positive definite operator. (\dagger) and (\ddagger) follow by applying Holder's inequality. $(**)$ follows using the condition $\mathbb{E}_P[K(Z, Z)]^r \leq cr! \kappa^r$ and that for any $k > 1$, $l > 0$, we have $((lk)!)^{\frac{1}{k}} \leq l! k^l$.

Thus using $\left\| \frac{1}{n} \sum_{i=1}^n \xi_i \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \leq \left\| \frac{1}{n} \sum_{i=1}^n \xi_i \right\|_{\mathcal{L}^2(\mathcal{H}_K)}$ and applying Sriperumbudur and Sterge (2022, Theorem D.1) yields that with probability at least $1 - \delta$,

$$\left\| \mathcal{A}_{P,\lambda}^{-1/2} (\hat{\mathcal{A}}_P - \mathcal{A}_P) \mathcal{A}_{P,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \leq \frac{d_q \log \frac{2}{\delta}}{n\lambda} + \sqrt{\frac{4d_q^{2-q} c^{1-q} \left[\text{Tr}(\mathcal{A}_{P,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P,\lambda}^{-1/2}) \right]^q \log \frac{2}{\delta}}{n\lambda^{2-q}}},$$

holds for any $0 < q \leq 1$. Setting $q = 0$, yields that

$$\left\| \mathcal{A}_{P,\lambda}^{-1/2} (\hat{\mathcal{A}}_P - \mathcal{A}_P) \mathcal{A}_{P,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \leq \frac{2\kappa \log \frac{2}{\delta}}{n\lambda} + \sqrt{\frac{16\kappa^2 c \log \frac{2}{\delta}}{n\lambda^2}},$$

which under condition (i) implies the desired bound in (a).

Suppose $q \neq 0$ and let $\varepsilon = 1 - q$, then we have

$$\left\| \mathcal{A}_{P,\lambda}^{-1/2} (\hat{\mathcal{A}}_P - \mathcal{A}_P) \mathcal{A}_{P,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \leq \frac{2\kappa \log \frac{2}{\delta}}{\varepsilon n \lambda} + \sqrt{\frac{4(2\kappa)^{1+\varepsilon} c^\varepsilon \left[2\mathcal{N}_1(\lambda) + 2C_\lambda \|u\|_{L^2(P_0)} \right]^{1-\varepsilon} \log \frac{2}{\delta}}{n\varepsilon^{1+\varepsilon} \lambda^{1+\varepsilon}}},$$

where we used Lemma A.1(ii)-b. The above bound implies the desired result in (a) when

$$\lambda \geq \frac{256\kappa \max\{c, 1\} \left(\mathcal{N}_1(\lambda) + C_\lambda \|u\|_{L^2(P_0)} \right) \log \frac{2}{\delta}}{\varepsilon n^{\frac{1}{1+\varepsilon}}},$$

then by setting $\varepsilon = \frac{1}{\log n}$ and using $n^{\frac{\log n}{1+\log n}} \geq \frac{n}{e}$ yields the condition on λ in (ii).

For (iii), when the kernel is bounded, we use Minsker's inequality (Minsker, 2017) for operators as stated in Sriperumbudur and Sterge (2022, Theorem D.2) instead of Bernstein's inequality. Observe that

$$\|\xi_i(Z)\|_{\mathcal{L}^\infty(\mathcal{H}_K)} = \left\| \mathcal{A}_{P,\lambda}^{-1/2} K(\cdot, Z) \right\|_{\mathcal{H}_K} \leq \frac{\kappa}{\lambda}$$

and

$$\begin{aligned} \mathbb{E}(\xi - \mathbb{E}(\xi))^2 &\preceq \mathbb{E}(\xi^2) = \mathbb{E} \left(\mathcal{A}_{P,\lambda}^{-1/2} (K(\cdot, Z_i) \otimes_{\mathcal{H}_K} K(\cdot, Z_i)) \mathcal{A}_{P,\lambda}^{-1} (K(\cdot, Z_i) \otimes_{\mathcal{H}_K} K(\cdot, Z_i)) \mathcal{A}_{P,\lambda}^{-1/2} \right) \\ &\preceq \sup_Z \|\xi(Z)\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \mathbb{E}(\mathcal{A}_{P,\lambda}^{-1/2} (K(\cdot, Z_i) \otimes_{\mathcal{H}_K} K(\cdot, Z_i)) \mathcal{A}_{P,\lambda}^{-1/2}) \\ &\preceq \frac{\kappa}{\lambda} \mathcal{A}_{P,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P,\lambda}^{-1/2} := S, \end{aligned}$$

where

$$\|S\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \leq \frac{\kappa}{\lambda} := \sigma^2,$$

and

$$d := \frac{\text{Tr}(S)}{\|S\|_{\mathcal{L}^\infty(\mathcal{H}_K)}} = \frac{\text{Tr}(\mathcal{A}_{P,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P,\lambda}^{-1/2})}{\left\| \mathcal{A}_{P,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)}} \leq \frac{\kappa(\|\mathcal{A}_P\| + \lambda)}{\|\mathcal{A}_P\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \lambda} \leq \frac{2\kappa}{\lambda}.$$

Thus applying Sriperumbudur and Sterge (2022, Theorem D.2) yields with probability at least $1 - \delta$ that

$$\left\| \mathcal{A}_{P,\lambda}^{-1/2} (\hat{\mathcal{A}}_P - \mathcal{A}_P) \mathcal{A}_{P,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \leq \frac{\beta\kappa}{\lambda n} + \sqrt{\frac{3\beta\kappa}{\lambda n}},$$

where $\beta := \frac{2}{3} \log \frac{4d}{\delta}$. Then the desired result in (a) follows under the condition on λ in (iii).

Finally, it remains to show that (b) and (c) follows from (a). To this end, define $B_n := \mathcal{A}_{P,\lambda}^{-1/2} (\mathcal{A}_P - \hat{\mathcal{A}}_P) \mathcal{A}_{P,\lambda}^{-1/2}$, and observe that

$$\begin{aligned} \left\| \mathcal{A}_{P,\lambda}^{1/2} \hat{\mathcal{A}}_P^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} &= \left\| \mathcal{A}_{P,\lambda}^{1/2} \hat{\mathcal{A}}_P^{-1} \mathcal{A}_{P,\lambda}^{1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)}^{1/2} \\ &= \left\| (\mathbf{I} - B_n)^{-1} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)}^{1/2} \leq (1 - \|B_n\|_{\mathcal{L}^\infty(\mathcal{H}_K)})^{-1/2}, \end{aligned}$$

and

$$\left\| \mathcal{A}_{P,\lambda}^{-1/2} \hat{\mathcal{A}}_P^{1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} = \left\| \mathbf{I} - B_n \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)}^{1/2} \leq (1 + \|B_n\|_{\mathcal{L}^\infty(\mathcal{H}_K)})^{1/2},$$

the desired results follow. \square

Lemma A.5. Let ϕ_i be as defined in Lemma A.1 with $\sup_i \|\phi_i\|_{\mathcal{L}^\infty(\mathcal{H}_K)}^2 \leq C < \infty$. If

$$4C\mathcal{N}_1(\lambda) \|u\|_{L^2(P_0)}^2 < 1,$$

$n > \frac{200C\mathcal{N}_1(\lambda)}{3} \log\left(\frac{12\mathcal{N}_1(\lambda)}{v_p\delta}\right)$, and $\lambda \leq \|\mathcal{A}_{P_0}\|_{\mathcal{L}^\infty(\mathcal{H}_K)}$, where $v_p := \mathbb{E}_P\phi_1^2 > 0$. Then for $0 < \delta \leq 1$,

$$(i) \ P\left(\left\|\mathcal{A}_{P_0,\lambda}^{-1/2}(\mathcal{A}_P - \hat{\mathcal{A}}_P)\mathcal{A}_{P_0,\lambda}^{-1/2}\right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \leq \frac{1}{4}\right) \geq 1 - \delta;$$

$$(ii) \ P\left(\left\|\mathcal{A}_{P_0,\lambda}^{1/2}\hat{\mathcal{A}}_{P,\lambda}^{-1/2}\right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \leq 2\right) \geq 1 - \delta;$$

$$(iii) \ P\left(\left\|\mathcal{A}_{P_0,\lambda}^{-1/2}\hat{\mathcal{A}}_{P,\lambda}^{1/2}\right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \leq \frac{\sqrt{7}}{2}\right) \geq 1 - \delta,$$

where

$$\hat{\mathcal{A}}_P = \frac{1}{n} \sum_{i=1}^n K(\cdot, Z_i) \otimes_{\mathcal{H}_K} K(\cdot, Z_i), \quad Z_i \stackrel{i.i.d.}{\sim} P_0.$$

Proof. Let $\xi_i := \mathcal{A}_{P_0,\lambda}^{-1/2}(K(\cdot, Z_i) \otimes_{\mathcal{H}_K} K(\cdot, Z_i))\mathcal{A}_{P_0,\lambda}^{-1/2}$. Then $\mathbb{E}(\xi_i) = \mathcal{A}_{P_0,\lambda}^{-1/2}\mathcal{A}_P\mathcal{A}_{P_0,\lambda}^{-1/2}$, and

$$\frac{1}{n} \sum_{i=1}^n \xi_i - \mathbb{E}(\xi) = \mathcal{A}_{P_0,\lambda}^{-1/2}(\hat{\mathcal{A}}_P - \mathcal{A}_P)\mathcal{A}_{P_0,\lambda}^{-1/2}.$$

Observe that

$$\begin{aligned} \|\xi(Z)\|_{\mathcal{L}^\infty(\mathcal{H}_K)} &= \left\|\mathcal{A}_{P_0,\lambda}^{-1/2}(K(\cdot, Z) \otimes_{\mathcal{H}_K} K(\cdot, Z))\mathcal{A}_{P_0,\lambda}^{-1/2}\right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \\ &= \left\|\mathcal{A}_{P_0,\lambda}^{-1/2}K(\cdot, Z)\right\|_{\mathcal{H}_K}^2 = \left\langle K(\cdot, Z), \mathcal{A}_{P_0,\lambda}^{-1}K(\cdot, Z) \right\rangle_{\mathcal{H}_K} \\ &= \left\langle K(\cdot, Z), \sum_{i \geq 1} \frac{1}{\lambda_i + \lambda} \left\langle K(\cdot, Z), \sqrt{\lambda_i}\phi_i \right\rangle_{\mathcal{H}_K} \sqrt{\lambda_i}\phi_i \right. \\ &\quad \left. + \frac{1}{\lambda} \left(K(\cdot, Z) - \sum_i \left\langle K(\cdot, Z), \sqrt{\lambda_i}\phi_i \right\rangle_{\mathcal{H}_K} \sqrt{\lambda_i}\phi_i \right) \right\rangle_{\mathcal{H}_K} \\ &= \sum_i \frac{\lambda_i}{\lambda_i + \lambda} \phi_i^2(Z) + \frac{1}{\lambda} \left(K(Z, Z) - \sum_i \lambda_i \phi_i(Z)\phi_i(Z) \right) \\ &\stackrel{(*)}{=} \sum_i \frac{\lambda_i}{\lambda_i + \lambda} \phi_i^2(Z) \leq \sup_i \|\phi_i\|_{\mathcal{L}^\infty(\mathcal{H}_K)}^2 \mathcal{N}_1(\lambda) \leq C\mathcal{N}_1(\lambda), \end{aligned}$$

where (*) follows by Mercer's theorem (see Steinwart and Scovel 2012, Lemma 2.6). Also, we have

$$\begin{aligned} \mathbb{E}(\xi - \mathbb{E}(\xi))^2 &\preceq \mathbb{E}(\xi^2) = \mathbb{E}\left(\mathcal{A}_{P_0,\lambda}^{-1/2}(K(\cdot, Z_i) \otimes_{\mathcal{H}_K} K(\cdot, Z_i))\mathcal{A}_{P_0,\lambda}^{-1/2}(K(\cdot, Z_i) \otimes_{\mathcal{H}_K} K(\cdot, Z_i))\mathcal{A}_{P_0,\lambda}^{-1/2}\right) \\ &\preceq \sup_Z \|\xi(Z)\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \mathbb{E}(\mathcal{A}_{P_0,\lambda}^{-1/2}(K(\cdot, Z_i) \otimes_{\mathcal{H}_K} K(\cdot, Z_i))\mathcal{A}_{P_0,\lambda}^{-1/2}) \\ &\preceq C\mathcal{N}_1(\lambda)\mathcal{A}_{P_0,\lambda}^{-1/2}\mathcal{A}_P\mathcal{A}_{P_0,\lambda}^{-1/2} := S \end{aligned}$$

so that

$$\|S\|_{\mathcal{L}^\infty(\mathcal{H}_K)} = C\mathcal{N}_1(\lambda) \left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P_0,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \stackrel{(\dagger)}{\leq} C\mathcal{N}_1(\lambda) \left(1 + \sqrt{C\mathcal{N}_1(\lambda)} \|u\|_{L^2(P_0)} \right) := \sigma^2,$$

where in (\dagger) we used the bound in (A.1). Define

$$\begin{aligned} d &:= \frac{\text{Tr}(S)}{\|S\|_{\mathcal{L}^\infty(\mathcal{H}_K)}} = \frac{\text{Tr}(\mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P_0,\lambda}^{-1/2})}{\left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P_0,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)}} \stackrel{(**)}{\leq} \frac{\mathcal{N}_1(\lambda) + C\mathcal{N}_1(\lambda) \|u\|_{L^2(P_0)}}{\left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P_0,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)}} \\ &\stackrel{(\ddagger)}{\leq} \frac{\left(\mathcal{N}_1(\lambda) + C\mathcal{N}_1(\lambda) \|u\|_{L^2(P_0)} \right) \left(\|\mathcal{A}_{P_0}\|_{\mathcal{L}^\infty(\mathcal{H}_K)} + \lambda \right)}{v_p \|\mathcal{A}_{P_0}\|_{\mathcal{L}^\infty(\mathcal{H}_K)}} \\ &\leq \frac{2}{v_p} \mathcal{N}_1(\lambda) (1 + C \|u\|_{L^2(P_0)}), \end{aligned}$$

where the last inequality follows by using $\lambda \leq \|\mathcal{A}_{P_0}\|_{\mathcal{L}^\infty(\mathcal{H}_K)}$. $(**)$ follows using Lemma A.1(ii) and in (\ddagger) we used

$$\left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P_0,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \geq \frac{v_p \|\mathcal{A}_{P_0}\|_{\mathcal{L}^\infty(\mathcal{H}_K)}}{\|\mathcal{A}_{P_0}\|_{\mathcal{L}^\infty(\mathcal{H}_K)} + \lambda},$$

which is proven below.

$$\begin{aligned} \left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P_0,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} &= \sup_{\|f\|_{\mathcal{H}_K} \leq 1} \left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P_0,\lambda}^{-1/2} f \right\|_{\mathcal{H}_K} \\ &\geq \left\| \mathcal{A}_{P_0,\lambda}^{-1/2} \mathcal{A}_P \mathcal{A}_{P_0,\lambda}^{-1/2} \psi_1 \right\|_{\mathcal{H}_K} = \left\| \sum_i \frac{\sqrt{\lambda_1}}{\sqrt{\lambda_1 + \lambda}} \frac{\sqrt{\lambda_i}}{\sqrt{\lambda_i + \lambda}} (\mathbb{E}_P \phi_i \phi_1) \psi_i \right\|_{\mathcal{H}_K} \\ &= \sqrt{\frac{\lambda_1}{\lambda_1 + \lambda} \sum_i \frac{\lambda_i}{\lambda_i + \lambda} (\mathbb{E}_P \phi_i \phi_1)^2} \geq \frac{\lambda_1}{\lambda_1 + \lambda} \mathbb{E}_P \phi_1^2 = \frac{v_p \|\mathcal{A}_{P_0}\|_{\mathcal{L}^\infty(\mathcal{H}_K)}}{\|\mathcal{A}_{P_0}\|_{\mathcal{L}^\infty(\mathcal{H}_K)} + \lambda}, \end{aligned}$$

where $\psi_i = \sqrt{\lambda_i} \phi_i$ are orthonormal eigenfunctions of \mathcal{A}_{P_0} . Finally, we apply Minsker's inequality (Minsker, 2017) for operators as stated in Sriperumbudur and Sterge (2022, Theorem D.2), which yields that with probability at least $1 - \delta$,

$$\left\| \mathcal{A}_{P_0,\lambda}^{-1/2} (\mathcal{A}_P - \hat{\mathcal{A}}_P) \mathcal{A}_{P_0,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \leq \frac{C\beta\mathcal{N}_1(\lambda)}{n} + \sqrt{\frac{3C\beta\mathcal{N}_1(\lambda)(1 + \sqrt{C\mathcal{N}_1(\lambda)}) \|u\|_{L^2(P_0)}}{n}},$$

where $\beta := \frac{2}{3} \log \frac{4d}{\delta}$. Thus, the desired result in (i) follows under the conditions $4C\mathcal{N}_1(\lambda) \|u\|_{L^2(P_0)}^2 < 1$ and $n > \frac{200C\mathcal{N}_1(\lambda)}{3} \log \left(\frac{12\mathcal{N}_1(\lambda)}{v_p \delta} \right)$. Then it remains to show that (ii) and (iii) follows from (i).

For (ii) observe that

$$\begin{aligned} \left\| \mathcal{A}_{P_0,\lambda}^{1/2} \hat{\mathcal{A}}_{P,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)}^2 &= \left\| \mathcal{A}_{P_0,\lambda}^{1/2} \hat{\mathcal{A}}_{P,\lambda}^{-1} \mathcal{A}_{P_0,\lambda}^{1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} = \left\| \mathbf{I} - \mathcal{A}_{P_0,\lambda}^{-1/2} (\mathcal{A}_{P_0} - \hat{\mathcal{A}}_P) \mathcal{A}_{P_0,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \\ &\leq \left(1 - \left\| \mathcal{A}_{P_0,\lambda}^{-1/2} (\mathcal{A}_{P_0} - \hat{\mathcal{A}}_P) \mathcal{A}_{P_0,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \right)^{-1} \leq 4, \end{aligned}$$

where the last inequality follows using

$$\left\| \mathcal{A}_{P_0,\lambda}^{-1/2} (\mathcal{A}_{P_0} - \hat{\mathcal{A}}_P) \mathcal{A}_{P_0,\lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)}$$

$$\begin{aligned}
& \leq \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} (\mathcal{A}_P - \hat{\mathcal{A}}_P) \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} + \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} (\mathcal{A}_{P_0} - \mathcal{A}_P) \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \\
& \stackrel{(*)}{\leq} \frac{1}{4} + \sqrt{C\mathcal{N}_1(\lambda)} \|u\|_{L^2(P_0)} < \frac{3}{4}.
\end{aligned}$$

For (iii) we have

$$\begin{aligned}
\left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \hat{\mathcal{A}}_{P, \lambda}^{1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)}^2 &= \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \hat{\mathcal{A}}_{P, \lambda} \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \\
&= \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} (\hat{\mathcal{A}}_{P, \lambda} - \mathcal{A}_{P, \lambda} + \mathcal{A}_{P, \lambda}) \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \\
&= \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} (\hat{\mathcal{A}}_P - \mathcal{A}_P) \mathcal{A}_{P_0, \lambda}^{-1/2} + \mathcal{A}_{P_0, \lambda}^{-1/2} \mathcal{A}_{P, \lambda} \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \\
&= \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} (\hat{\mathcal{A}}_P - \mathcal{A}_P) \mathcal{A}_{P_0, \lambda}^{-1/2} + \mathcal{A}_{P_0, \lambda}^{-1/2} (\mathcal{A}_P - \mathcal{A}_{P_0}) \mathcal{A}_{P_0, \lambda}^{-1/2} + \mathbf{I} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \\
&\leq 1 + \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} (\hat{\mathcal{A}}_P - \mathcal{A}_P) \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} + \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} (\mathcal{A}_P - \mathcal{A}_{P_0}) \mathcal{A}_{P_0, \lambda}^{-1/2} \right\|_{\mathcal{L}^\infty(\mathcal{H}_K)} \\
&\stackrel{(**)}{\leq} 1 + \frac{1}{4} + \sqrt{C\mathcal{N}_1(\lambda)} \|u\|_{L^2(P_0)} \leq \frac{7}{4},
\end{aligned}$$

where (*) and (**) follow from (A.1). \square

Lemma A.6. Let $\zeta = \left\| g_\lambda^{1/2} (\hat{\mathcal{A}}_P) \Psi_P \right\|_{\mathcal{H}_{K_0}}^2$, $\mathcal{W} = \hat{\mathcal{A}}_{P, \lambda}^{-1/2} \mathcal{A}_{P, \lambda}^{1/2}$, and $\mathcal{V} = \hat{\mathcal{A}}_{P, \lambda}^{-1/2} \mathcal{A}_{P_0, \lambda}^{1/2}$, where g_λ satisfies (E₁)–(E₄). Suppose $u \in \text{Ran}(\Upsilon_{P_0}^\theta)$, $\theta > 0$ and

$$\|u\|_{L^2(P_0)}^2 \geq d_1 \|\Upsilon_{P_0}\|_{\mathcal{L}^\infty(L^2(P_0))}^{2 \max(\theta - \xi, 0)} \lambda^{2\tilde{\theta}} \|\Upsilon_{P_0}^{-\theta} u\|_{L^2(P_0)}^2,$$

where $\tilde{\theta} = \min(\theta, \xi)$, for some constant $d_1 > 0$. Then

$$(i) \quad \zeta \geq d_2 \|\mathcal{W}^{-1}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^{-2} \frac{\|u\|_{L^2(P_0)}^2}{1 + \sqrt{C_\lambda} \|u\|_{L^2(P_0)}};$$

$$(ii) \quad \zeta \geq d_3 \|\mathcal{V}^{-1}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^{-2} \|u\|_{L^2(P_0)}^2,$$

where C_λ is defined in Lemma A.1 with $K = K_0$ and d_2, d_3 are positive constants.

Proof. (i): By applying Hagrass et al. (2022, Lemma A.10, Lemma A.8(v)) (by replacing Σ_{PQ} with \mathcal{A}_P), we have

$$\begin{aligned}
\zeta &\geq \left\| \hat{\mathcal{A}}_{P, \lambda}^{-1/2} g_\lambda^{-1/2} (\hat{\mathcal{A}}_P) \right\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^{-2} \left\| \hat{\mathcal{A}}_{P, \lambda}^{-1/2} \Psi_P \right\|_{\mathcal{H}_{K_0}}^2 \geq C_4 \|\mathcal{W}^{-1}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^{-2} \left\| \mathcal{A}_{P, \lambda}^{-1/2} \Psi_P \right\|_{\mathcal{H}_{K_0}}^2 \\
&\stackrel{(*)}{\geq} C_4^2 \|\mathcal{W}^{-1}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^{-2} \frac{\left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \Psi_P \right\|_{\mathcal{H}_{K_0}}^2}{1 + \sqrt{C_\lambda} \|u\|_{L^2(P_0)}},
\end{aligned}$$

where (*) follows from Lemma A.2. For the bound in (ii) we have,

$$\zeta \geq \left\| \hat{\mathcal{A}}_{P, \lambda}^{-1/2} g_\lambda^{-1/2} (\hat{\mathcal{A}}_P) \right\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^{-2} \left\| \hat{\mathcal{A}}_{P, \lambda}^{-1/2} \Psi_P \right\|_{\mathcal{H}_{K_0}}^2 \geq C_4 \|\mathcal{V}^{-1}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^{-2} \left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \Psi_P \right\|_{\mathcal{H}_{K_0}}^2.$$

Thus, it remains to show

$$\left\| \mathcal{A}_{P_0, \lambda}^{-1/2} \Psi_P \right\|_{\mathcal{H}_{K_0}}^2 \geq \tilde{d} \|u\|_{L^2(P_0)}^2,$$

for some $\tilde{d} > 0$ which follows under the given conditions on u , similar to the proof of Hagrass et al. (2023, Lemma A.2) by replacing \mathcal{T} with Υ_{P_0} , Σ_0 with \mathcal{A}_{P_0} and μ_P with Ψ_P . \square

Lemma A.7. *For any $\delta > 0$, there exists \tilde{C} such that*

$$P_{H_1}(q_{1-\alpha}^\lambda \leq \tilde{C}\gamma) \geq 1 - \delta,$$

where $\gamma := \frac{\log \frac{2}{\alpha}}{n_1 \sqrt{\delta}} \left(\|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})} \sqrt{\zeta} + \|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \sqrt{D_\lambda} + \zeta \right)$. Furthermore the operator \mathcal{W} can be replaced by \mathcal{V} when $4C_\lambda \|u\|_{L^2(P_0)}^2 \leq 1$, where \mathcal{W} , \mathcal{V} , D_λ and ζ are defined in Lemma A.3 and C_λ is defined in Lemma A.1.

Proof. Let $h(x, y) := \left\langle g_\lambda^{1/2}(\hat{\mathcal{A}}_P)K_0(\cdot, x), g_\lambda^{1/2}(\hat{\mathcal{A}}_P)K_0(\cdot, y) \right\rangle_{\mathcal{H}_{K_0}}$. Conditioned on the samples \mathbb{X}_n , it follows from de la Peña and Giné (1999, Corollary 3.2.6) that

$$P_\varepsilon \left(\left| \sum_{i \neq j} \varepsilon_i \varepsilon_j h(X_i, X_j) \right| \geq t \right) \leq 2 \exp \left(\frac{-at}{\sqrt{\sum_{i,j} h^2(X_i, X_j)}} \right),$$

for some constant $a > 0$. This implies that

$$q_{1-\alpha}^\lambda \leq \frac{\tilde{C} \log \frac{2}{\alpha}}{n_1(n_1-1)} \sqrt{\sum_{i,j} h^2(X_i, X_j)}, \quad (\text{A.3})$$

almost surely. Let $I := \frac{1}{n_1(n_1-1)} \sqrt{\sum_{i,j} h^2(X_i, X_j)}$. Then similar to the proof of Lemma A.3, we can show that

$$\mathbb{E}(I^2 | \mathbb{Z}_{n_2}) \lesssim \frac{1}{n_1^2} \left(\|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^2 \zeta + \|\mathcal{W}\|_{\mathcal{L}^\infty(\mathcal{H}_{K_0})}^4 D_\lambda + \zeta^2 \right),$$

where \mathcal{W} can be replaced by \mathcal{V} when $4C_\lambda \|u\|_{L^2(P_0)}^2 \leq 1$. Thus using (A.3) and Markov's inequality, we obtain the desired result. \square

B General version of Corollories 6 and 7

In this appendix, we present a general version of Corollories 6 and 7 under the weaker assumption of **A₂**.

Corollary B.1. *Suppose $\lambda_i \lesssim i^{-\beta}$, $\beta > 1$. Let $\lambda_n \asymp \Delta_n^{\frac{1}{2\beta}}$, with $\Delta_n \rightarrow 0$. Then for $(P_n)_n \subset \mathcal{P}$, we have*

$$\lim_{n \rightarrow \infty} P\{\hat{\mathbb{S}}_\lambda^{P_n} \geq q_{1-\alpha}^\lambda\} = 1,$$

if one of the following holds:

(i) \mathbf{A}_2 holds and $\Delta_n B_n^{-1} \rightarrow \infty$, where

$$B_n = \begin{cases} n^{\frac{-4\tilde{\theta}\beta}{4\tilde{\theta}\beta+1}}, & \tilde{\theta} > b_1 \\ \left(\frac{\log n}{n}\right)^{\frac{2\tilde{\theta}\beta}{\beta+1}}, & b_2 < \tilde{\theta} \leq b_1 \\ \left(\frac{\log n}{n}\right)^{\frac{4\tilde{\theta}\beta}{2\beta + \frac{v}{v-1}(1 - \frac{2}{v} + 2\beta) - 2\tilde{\theta}\beta}}, & \frac{v-2}{4\beta(v-1)} + \frac{v}{2(v-1)} < \tilde{\theta} \leq b_2 \\ n^{\frac{-8\tilde{\theta}\beta}{2(1 + \frac{v}{v-1})\beta + 4\tilde{\theta}\beta + \frac{v-2}{v-1}}}, & \frac{2v-3}{2v-2} - \frac{v-2}{4\beta(v-1)} \leq \tilde{\theta} \leq \frac{v-2}{4\beta(v-1)} + \frac{v}{2(v-1)} \\ n^{-\tilde{\theta}}, & \text{otherwise} \end{cases},$$

with $b_1 = \max\left\{\frac{1}{3} + \frac{2\beta v-1}{6\beta(v-1)}, \frac{v-2}{4\beta(v-1)} + \frac{v}{2(v-1)}\right\}$, $b_2 = \min\left\{\frac{v}{v-1} - \frac{v}{2(v-1)\beta}, b_1\right\}$, and $\tilde{\theta} = \min\{\theta, \xi\}$.

(ii) $\sup_i \|\phi_i\|_\infty < \infty$ and $\Delta_n B_n^{-1} \rightarrow \infty$, where

$$B_n = \begin{cases} n^{\frac{-4\tilde{\theta}\beta}{4\tilde{\theta}\beta+1}}, & \tilde{\theta} > \frac{1}{2\beta} \\ n^{\frac{-4\tilde{\theta}\beta}{2\tilde{\theta}\beta+\beta+1}}, & \frac{1}{2} - \frac{1}{2\beta} < \tilde{\theta} \leq \frac{1}{2\beta} \\ \left(\frac{\log n}{n}\right)^{2\tilde{\theta}}, & \tilde{\theta} \leq \min\left\{\frac{1}{2} - \frac{1}{2\beta}, \frac{1}{2\beta}\right\} \end{cases}.$$

Proof. We will verify that the conditions of Theorem 5 are satisfied. If $\lambda_i \lesssim i^{-\beta}$, for $\beta > 1$ then by Sriperumbudur and Sterge (2022, Lemma B.9), we have $\mathcal{N}_1(\lambda) \lesssim \lambda^{\frac{-1}{\beta}}$ and $\mathcal{N}_2(\lambda) \leq \mathcal{N}_1^{1/2}(\lambda) \lesssim \lambda^{\frac{-1}{2\beta}}$. Since $\lambda_n \asymp \Delta_n^{\frac{1}{2\tilde{\theta}}}$, (11) is directly satisfied.

(i) In this case

$$C_\lambda = \frac{(\mathcal{N}_1(\lambda))^{\frac{v-2}{2(v-1)}}}{\lambda^{\frac{v}{v-1}}} = \lambda^{\frac{-v}{v-1} \left(\frac{1}{2\beta} - \frac{1}{v\beta} + 1\right)} \asymp \Delta_n^{\frac{-v}{2\tilde{\theta}(v-1)} \left(\frac{1}{2\beta} - \frac{1}{v\beta} + 1\right)}.$$

Observe that

$$4C_\lambda \Delta_n \asymp \Delta_n^{1 - \frac{v}{2\tilde{\theta}(v-1)} \left(\frac{1}{2\beta} - \frac{1}{v\beta} + 1\right)} \stackrel{(*)}{\lesssim} 1,$$

for n large enough, where $(*)$ holds since $\Delta_n \rightarrow 0$, and assuming $1 - \frac{v}{2\tilde{\theta}(v-1)} \left(\frac{1}{2\beta} - \frac{1}{v\beta} + 1\right) > 0$, which is equivalent to $\tilde{\theta} > \frac{v-2}{4\beta(v-1)} + \frac{v}{2(v-1)}$. Thus it remains to verify (12) and (14) for this range of $\tilde{\theta}$. Observe that in this range $D_\lambda = \mathcal{N}_2^2(\lambda) \lesssim \lambda^{\frac{-1}{\beta}} \asymp \Delta_n^{\frac{-1}{2\tilde{\theta}\beta}}$, thus (12) is equivalent to $\Delta_n^{\frac{4\tilde{\theta}\beta+1}{4\tilde{\theta}\beta}} n \rightarrow \infty$, which in turn is equivalent to

$$\Delta_n n^{\frac{4\tilde{\theta}\beta}{4\tilde{\theta}\beta+1}} \rightarrow \infty. \quad (\text{B.1})$$

(14) is implied if $\frac{n\lambda_n}{\mathcal{N}_1(\lambda_n) \log n} \rightarrow \infty$ and $\frac{n\lambda_n}{C_{\lambda_n} \sqrt{\Delta_n} \log n} \rightarrow \infty$, where we used that $\frac{1}{a+b} \leq \frac{1}{a} + \frac{1}{b}$, for $a, b > 0$. Then after simplifying the expressions, the first condition $\frac{n\lambda_n}{\mathcal{N}_1(\lambda_n) \log n} \rightarrow \infty$ is implied when

$$\Delta_n \left(\frac{n}{\log n}\right)^{\frac{2\tilde{\theta}\beta}{\beta+1}} \rightarrow \infty, \quad (\text{B.2})$$

and the second condition $\frac{n\lambda_n}{C_{\lambda_n}\sqrt{\Delta_n}\log n} \rightarrow \infty$ is implied if $\Delta_n \frac{2\beta + \frac{v}{v-1}(1 - \frac{2}{v} + 2\beta) - 2\tilde{\theta}\beta}{4\tilde{\theta}\beta} \left(\frac{n}{\log n}\right) \rightarrow \infty$. This is directly satisfied if $2\beta + \frac{v}{v-1}(1 - \frac{2}{v} + 2\beta) - 2\tilde{\theta}\beta < 0$ which is equivalent to $\tilde{\theta} > 1 + \frac{v}{v+1} + \frac{v-2}{(v-1)2\beta}$. On the other hand, if $\tilde{\theta} \leq 1 + \frac{v}{v+1} + \frac{v-2}{(v-1)2\beta}$, then the second condition $\frac{n\lambda_n}{C_{\lambda_n}\sqrt{\Delta_n}\log n} \rightarrow \infty$ is implied if

$$\Delta_n \left(\frac{n}{\log n}\right)^{\frac{4\tilde{\theta}\beta}{2\beta + \frac{v}{v-1}(1 - \frac{2}{v} + 2\beta) - 2\tilde{\theta}\beta}} \rightarrow \infty. \quad (\text{B.3})$$

Since $\max\left\{1 + \frac{v}{v+1} + \frac{v-2}{(v-1)2\beta}, \frac{v-2}{4\beta(v-1)} + \frac{v}{2(v-1)}\right\} = 1 + \frac{v}{v+1} + \frac{v-2}{(v-1)2\beta}$ for all $v > 2$, in the case of $\tilde{\theta} > 1 + \frac{v}{v+1} + \frac{v-2}{(v-1)2\beta}$, the conditions on Δ_n to be satisfied are (B.1) and (B.2), and in the case $\frac{v-2}{4\beta(v-1)} + \frac{v}{2(v-1)} < \tilde{\theta} \leq 1 + \frac{v}{v+1} + \frac{v-2}{(v-1)2\beta}$, then (B.3) has also to be satisfied in addition to (B.1) and (B.2). On the other hand, if $\tilde{\theta} \leq \frac{v-2}{(v-1)4\beta} + \frac{v}{2(v-1)}$, then it is easy to verify that (12) (with $D_\lambda = \lambda^{-1}$) and (13) are implied if

$$\Delta_n n^{\frac{4\tilde{\theta}}{4\tilde{\theta}+1}} \rightarrow \infty, \quad (\text{B.4})$$

$$\Delta_n n^{\frac{8\tilde{\theta}\beta}{2(1 + \frac{v}{v-1})\beta + 4\tilde{\theta}\beta + \frac{v-2}{v-1}}} \rightarrow \infty, \quad (\text{B.5})$$

and

$$\Delta_n n^{\tilde{\theta}} \rightarrow \infty. \quad (\text{B.6})$$

Observe that when $\frac{4\tilde{\theta}\beta}{4\tilde{\theta}\beta+1} \leq \frac{2\tilde{\theta}\beta}{\beta+1} \Leftrightarrow \frac{1}{2} + \frac{1}{4\beta} \leq \tilde{\theta}$, and $\frac{4\tilde{\theta}\beta}{4\tilde{\theta}\beta+1} \leq \frac{4\tilde{\theta}\beta}{2\beta + \frac{v}{v-1}(1 - \frac{2}{v} + 2\beta) - 2\tilde{\theta}\beta} \Leftrightarrow \frac{1}{3} + \frac{2\beta v-1}{6\beta(v-1)} \leq \tilde{\theta}$ then (B.1) dominates (B.2) and (B.3) for large enough n . When $\frac{4\tilde{\theta}\beta}{2\beta + \frac{v}{v-1}(1 - \frac{2}{v} + 2\beta) - 2\tilde{\theta}\beta} < \frac{4\tilde{\theta}\beta}{\beta+1} \Leftrightarrow \tilde{\theta} < \frac{v}{v-1} - \frac{v}{2(v-1)\beta}$, then (B.3) dominates (B.2). If $\frac{8\tilde{\theta}\beta}{2(1 + \frac{v}{v-1})\beta + 4\tilde{\theta}\beta + \frac{v-2}{v-1}} \leq \tilde{\theta} \Leftrightarrow \frac{2v-3}{2v-2} - \frac{v-2}{4\beta(v-1)} \leq \tilde{\theta}$, and $\frac{8\tilde{\theta}\beta}{2(1 + \frac{v}{v-1})\beta + 4\tilde{\theta}\beta + \frac{v-2}{v-1}} \leq \frac{4\tilde{\theta}}{4\tilde{\theta}+1} \Leftrightarrow \tilde{\theta} < \frac{v}{2(v-1)} + \frac{v-2}{4\beta(v-1)}$, then (B.5) dominates both (B.4) and (B.6) for large enough n . In other ranges (B.6) dominates. Thus putting things together yields that

$$\Delta_n B_n^{-1} \rightarrow \infty,$$

with B_n as mention in Corollary B.1 (i).

(ii) $\sup_i \|\phi_i\|_\infty < \infty$: In this case, $C_\lambda \asymp \mathcal{N}_1(\lambda)$ and we need to verify equations (11), (12) and (15), where (15) can be replaced by (16) when $4C_\lambda\Delta_n < 1$. When $\tilde{\theta} > \frac{1}{2\beta}$, it can be verified that $4C_\lambda\Delta_n < 1$ since $\Delta_n \rightarrow 0$. Then the required conditions are implied when

$$\Delta_n \min \left\{ n^{\frac{4\tilde{\theta}\beta}{4\tilde{\theta}\beta+1}}, \left(\frac{n}{\log n}\right)^{2\tilde{\theta}\beta} \right\} \rightarrow \infty.$$

On the other hand, if $\tilde{\theta} \leq \frac{1}{2\beta}$, then the required conditions are implied when

$$\Delta_n \min \left\{ n^{\frac{4\tilde{\theta}}{4\tilde{\theta}+1}}, n^{\frac{4\tilde{\theta}\beta}{2\tilde{\theta}\beta+\beta+1}}, \left(\frac{n}{\log n}\right)^{2\tilde{\theta}} \right\} \rightarrow \infty.$$

Putting the conditions together yields

$$\Delta_n B_n^{-1} \rightarrow \infty,$$

where B_n is mentioned in Corollary B.1(ii). \square

Corollary B.2. *Suppose $\lambda_i \lesssim e^{-\tau i}$, $\tau > 0$. Let $\lambda_n \asymp \Delta_n^{\frac{1}{2\tilde{\theta}}}$ with $\Delta_n \rightarrow 0$. Then for any $(P_n)_n \subset \mathcal{P}$, we have*

$$\lim_{n \rightarrow \infty} P\{\hat{\mathbb{S}}_\lambda^{P_n} \geq q_{1-\alpha}^\lambda\} = 1,$$

if one of the following holds:

(i) **A₂** holds and $\Delta_n B_n^{-1} \rightarrow \infty$, where

$$B_n = \begin{cases} \frac{\sqrt{\log n}}{n}, & \tilde{\theta} > \frac{2v-1}{3(v-1)} \\ \left(\frac{(\log n)^2}{n}\right)^{\frac{2\tilde{\theta}}{v-1-\tilde{\theta}}}, & \frac{v}{2(v-1)} \leq \tilde{\theta} \leq \frac{2v-1}{3(v-1)}, \\ \left(\frac{\log n}{n}\right)^{\frac{4\tilde{\theta}(v-1)}{2\tilde{\theta}(v-1)+2v-1}}, & \frac{2v-3}{2(v-1)} \leq \tilde{\theta} \leq \frac{v}{2(v-1)} \\ n^{-\tilde{\theta}}, & \text{otherwise} \end{cases},$$

where $\tilde{\theta} = \min\{\theta, \xi\}$.

(ii) $\sup_i \|\phi_i\|_\infty < \infty$ and $\Delta_n B_n^{-1} \rightarrow \infty$, where

$$B_n = \frac{\sqrt{\log n}}{n}.$$

Proof. We will verify that the conditions of Theorem 5 are satisfied. If $\lambda_i \asymp e^{-\tau i}$, then (Sriperumbudur and Sterge, 2022, Lemma B.9) yields $\mathcal{N}_1(\lambda) \lesssim \log \frac{1}{\lambda}$ and $\mathcal{N}_2(\lambda) \leq \mathcal{N}_1^{1/2}(\lambda) \lesssim \sqrt{\log \frac{1}{\lambda}}$. Since $\lambda_n \asymp \Delta_n^{1/2\tilde{\theta}}$, (11) is directly satisfied.

$$(i) \text{ In this case } C_\lambda \asymp \frac{(\log \lambda^{-1})^{\frac{v-2}{2(v-1)}}}{\lambda^{\frac{v}{v-1}}}.$$

$\tilde{\theta} > \frac{v}{2(v-1)}$: In this case, it can be verified that $4C_\lambda \Delta_n < 1$ since $\Delta_n \rightarrow 0$. Thus we only need to verify (12) and (14). Equation (12) is implied when

$$\Delta_n \frac{n}{\sqrt{\log n}} \rightarrow \infty.$$

(14) is implied if

$$\Delta_n \left(\frac{n}{(\log n)^2}\right)^{2\tilde{\theta}} \rightarrow \infty,$$

and $\frac{n\lambda_n}{C_\lambda \sqrt{\Delta_n} \log n} \rightarrow \infty$ which is satisfied for large enough n when $\tilde{\theta} \geq \frac{2v-1}{v-1}$.

$\frac{v}{2(v-1)} \leq \tilde{\theta} \leq \frac{2v-1}{v-1}$: In addition to the conditions in the previous case we also have

$$\Delta_n \left(\frac{n}{(\log n)^2}\right)^{\frac{2\tilde{\theta}}{v-1-\tilde{\theta}}} \rightarrow \infty.$$

$\tilde{\theta} \leq \frac{v}{2(v-1)}$: In this case, we need to verify equations (12) (with $D_\lambda = \lambda^{-1}$) and (13) which are implied if

$$\Delta_n \min \left\{ n^{\frac{4\tilde{\theta}}{4\tilde{\theta}+1}}, \left(\frac{n}{\log n} \right)^{\frac{4\tilde{\theta}(v-1)}{2\tilde{\theta}(v-1)+2v-1}}, n^{\tilde{\theta}} \right\} \rightarrow \infty.$$

Putting the conditions together while using

$$\frac{2\tilde{\theta}}{\frac{2v-1}{v-1} - \tilde{\theta}} \geq 1 \Leftrightarrow \tilde{\theta} \geq \frac{2v-1}{3(v-1)},$$

and

$$\frac{4\tilde{\theta}(v-1)}{2\tilde{\theta}(v-1) + 2v-1} > \tilde{\theta} \Leftrightarrow \frac{2v-3}{2(v-1)} > \tilde{\theta},$$

yields the desired result.

(ii) $\sup_i \|\phi_i\|_\infty < \infty$: In this case $C_\lambda \asymp \mathcal{N}_1(\lambda) \asymp \log \frac{1}{\lambda}$, and it can also be verified that for large enough n , $4C_\lambda \Delta_n < 1$ is implied when $\Delta_n \rightarrow 0$. Thus it only remains to verify equations (12) and (16) which are both implied when $\Delta_n \frac{n}{\sqrt{\log n}} \rightarrow \infty$. \square