

---

# MAIN SIGNLESS LAPLACIAN EIGENVALUES OF *quasi*-THRESHOLD GRAPHS

---

A PREPRINT

**Átila Jones**

Instituto Federal Sudeste De Minas  
Campus Juiz De Fora, Juiz De Fora, MG, Brazil.  
atila.jones@ifsudestemg.edu.br

**Vilmar Trevisan**

Instituto de Matemática e Estatística  
Universidade Federal do Rio Grande do Sul Porto Alegre, RS, Brazil.  
trevisan@mat.ufrgs.br

**Cybele T.M.Vinagre**

Instituto de Matemática e Estatística  
Universidade Federal Fluminense, Niterói, RJ, Brazil  
cybele\_vinagre@id.uff.br

April 16, 2024

## ABSTRACT

In this note, we present a structural description of certain connected cographs having  $k \geq 2$  main signless Laplacian eigenvalues. This result allows us to characterize the cographs which are *quasi*-threshold graphs with two main  $\mathbf{Q}$ -eigenvalues. In addition, we describe all the *quasi*-threshold graphs belonging to the subclass of generalized core-satellite graphs with  $k \geq 2$  main  $\mathbf{Q}$ -eigenvalues.

**Keywords** Cograph · *Quasi*-threshold graph · Signless Laplacian matrix · Main signless Laplacian eigenvalue

## 1 Introduction and main results

An eigenvalue  $\lambda$  is a *main eigenvalue* of a square matrix  $\mathbf{M}$  (or a *main  $\mathbf{M}$ -eigenvalue*) if the eigenspace  $\mathcal{E}(\lambda)$  of  $\lambda$  is non-orthogonal to the all ones vector  $\mathbf{j}$ , that is, if  $\mathcal{E}(\lambda)$  contains some eigenvector whose sum of entries is non-zero.

Main eigenvalues of graphs were introduced in [Cvetković, 1970] for the case  $\mathbf{M} = \mathbf{A}(G)$ , the adjacency matrix of the graph  $G$ . In [Chen and Huang., 2013] and Deng and Huang. [2013], the case  $\mathbf{M} = \mathbf{Q}(G)$ , the signless Laplacian matrix of  $G$ , was studied and it was shown that regular graphs were the only ones with exactly one main  $\mathbf{Q}$ -eigenvalue. The problem of characterizing graphs with  $k \geq 2$  main  $\mathbf{Q}$ -eigenvalues was also addressed, in particular the case  $k = 2$  for trees and unicyclic graphs. In [Chen and Huang., 2013], the bicyclic graphs with two main  $\mathbf{Q}$ -eigenvalues were characterized. In addition, trees, unicyclic and bicyclic graphs with three main  $\mathbf{Q}$ -eigenvalues, one of which being zero, were characterized in [Javarsineh and Fath-Tabar, 2017a] and [Javarsineh and Fath-Tabar, 2017b].

In [Vinagre et al., 2020], the authors gave an exact characterization of the threshold graphs having  $k$  main  $\mathbf{Q}$ -eigenvalues, for each  $k \geq 2$ . The present paper is a step forward to generalize this characterization for *quasi*-threshold graphs.

We may recall that threshold graphs form a subclass of cographs characterized as the class of  $\{P_4, C_4, 2K_2\}$ -free graphs, while *quasi*-threshold graphs constitute the class of  $\{C_4, P_4\}$ -free graphs (we refer to the definitions below).

The main result of this paper is a structural characterization of connected *quasi*-threshold graphs having  $k \geq 2$  main  $\mathbf{Q}$ -eigenvalues. In addition, we determine precisely what are the connected *quasi*-threshold graphs having two main  $\mathbf{Q}$ -eigenvalues. We also provide a full characterization of a subclass of *quasi*-threshold graphs. More precisely, for  $k = 2, 3, \dots$ , we determine what are the generalized core-satellite graphs having  $k$  main  $\mathbf{Q}$ -eigenvalues. It is important to note that the graph characterizations shown in this article were obtained with the help of the *Graph Filter* software [Jones et al.].

The paper is organized as follows. In the next section, we provide the definitions and the necessary background for presenting the results. We then start, in Section 3, studying the number  $k$  of main  $\mathbf{Q}$ -eigenvalues of a connected cograph  $G$ , presenting an upper bound on  $k$  based on the cotree representation of  $G$ . This allows us to compute the spectra of some particular *quasi*-threshold graphs, a necessary step to the results of the next section. In fact, in Section 4, we give a structural characterization of the *quasi*-threshold graphs having  $k \geq 2$  main  $\mathbf{Q}$ -eigenvalues. We show they have the form  $G = K_c \oplus H$  for some integer  $c \geq 1$  and a disconnected *quasi*-threshold graph  $H$ . Additionally, we prove that if  $\overline{H}$  is bipartite, then  $k = 2$ . In turn, these results lead to a precise determination of the connected *quasi*-threshold graphs having two main  $\mathbf{Q}$ -eigenvalues given in Theorem 4. Section 5 is devoted to the study of main  $\mathbf{Q}$ -eigenvalues of generalized core-satellite graphs, an important subclass of the class of *quasi*-threshold graphs, which contains, for example, the windmill graphs. For this class, we characterize which graphs have  $k$  main  $\mathbf{Q}$ -eigenvalues, for  $k = 2, 3, \dots$

## 2 Preliminaries

Let  $G = (V, E)$  be a simple graph. We denote by  $d_G(u) = d(u)$  the degree of vertex  $u \in V$ . As usual, we say that  $|E| = m$  is the *size* of  $G$  and  $|V| = n$  is the *order* of  $G$ . The *neighborhood* of a vertex  $v \in V$  is the set  $N_G(v) = \{w \in V; \{v, w\} \in E\}$  of neighbors of  $v$  in  $G$  and the *closed neighborhood* of  $v$  is  $N_G[v] = N_G(v) \cup \{v\}$ .

The complete graph on  $n$  vertices is denoted by  $K_n$ . Its complement  $\overline{K_n}$  is the graph without edges. For an integer  $p \geq 1$  and a graph  $G$ ,  $pG$  denotes the disjoint union of  $p$  copies of  $G$ . For any  $S \subset V$ , the subgraph of  $G$  induced by  $S$  is denoted  $G[S]$ . If  $G[S]$  is a complete subgraph then  $S$  is a *clique* in  $G$ . A vertex  $v$  is said to be *simplicial* (respectively, *universal*) in  $G$  when  $N_G(v)$  is a clique (respectively,  $N_G[v] = V$ ).

### 2.1 Chordal graphs

A *chordal graph* is a graph in which every cycle of length four or greater has a *chord*, that is, an edge between non-consecutive vertices of the cycle. Basic concepts about chordal graphs can be found in [Blair and Peyton, 1993]. The notions which are relevant to this note follow. A subset  $S \subset V$  is a *separator* of a non-complete graph  $G$  if its removal leaves the remaining graph disconnected. The set  $S$  is a *minimal separator* of  $G$  if  $S$  is a separator and no proper subset of  $S$  separates the graph. The minimum cardinality of a separator of  $G$  corresponds to  $\kappa(G)$ , the *vertex connectivity* of the graph. Let  $u, v$  be two non-adjacent vertices in  $V$ . A subset  $S \subset V$  is a *uv-separator vertex* (or a *separator for  $u$  and  $v$* ) if the removal of  $S$  from the graph separates  $u$  and  $v$  into distinct connected components. If no proper subset of  $S$  is a *uv-separator* then  $S$  is a *minimal uv-separator*. When the pair of vertices remains unspecified, we refer to  $S$  as a *minimal vertex separator (mvs)*. A minimal separator is always a minimal vertex separator but the converse is not true.

**Lemma 1** *Suppose that a connected regular graph  $G$  has a simplicial vertex. Then  $G$  is a complete graph.*

**Proof:** Let  $u$  be a simplicial vertex of the graph  $G$  and suppose that  $S = N_G[u]$  is a  $k$ -clique; then  $d(u) = k - 1$ . Suppose there is a vertex  $v$  in  $G \setminus S$ . Since  $G$  is a connected graph, there exists a vertex  $w \in S$ ,  $w \neq u$ , and there exists a path  $vv_1 \dots v_r w$  in  $G$ . Then  $d(w) \geq (k - 1) + 1 = k$ , in contradiction with the regularity of  $G$ . Therefore,  $G \setminus S$  is the empty set and  $G$  is a complete graph.

**Corollary 1** *A connected regular chordal graph is a complete graph.*

**Proof:** It is well known that a graph  $G$  is chordal if and only if every induced subgraph of  $G$  has a simplicial vertex. So, by Lemma 1, the result follows.

### 2.2 Cographs

The class of *complement reducible graphs*, known as *cographs*, is the hereditary class of  $P_4$ -free graphs, denoted by  $\text{Forb}(P_4)$ . This class appeared naturally in various contexts and has been rediscovered several times, but the seminal paper [Corneil, Lerchs, and Burlingham, 1981] is considered the origin of the study of cographs. There are several

equivalent ways in which the elements of this class may be defined. One of the definitions uses disjoint unions and complements recursively, as the authors of [Corneil et al., 1981]. A similar definition may be obtained through unions and joins. Consider the class  $\mathcal{D}$  defined recursively as follows:

- ( $\alpha$ )  $K_1 \in \mathcal{D}$ ;
- ( $\beta$ ) If  $G_1$  and  $G_2$  are vertex disjoint graphs in  $\mathcal{D}$ , then  $G_1 \cup G_2 \in \mathcal{D}$ ;
- ( $\gamma$ ) If  $G_1$  and  $G_2$  lie in  $\mathcal{D}$ , then its join  $G_1 \oplus G_2 \in \mathcal{D}$ .

Using this definition, it is possible to characterize bipartite cographs (important for our purposes in this note) as the following known result shows.

**Lemma 2** [Demange et al., 2005, Lemma 14] *If  $G$  is a connected and bipartite cograph then  $G = \overline{K_a} \oplus \overline{K_b}$  for positive integers  $a, b$ .*

We recall that *threshold graphs* constitute a subclass of cographs characterized as the hereditary class of  $\{P_4, C_4, 2K_2\}$ -free graphs. Since, by definition, chordal graphs are  $C_4$ -free graphs, the class of chordal cographs is the hereditary class of  $\{C_4, P_4\}$ -free graphs whose elements are often called *quasi-threshold graphs* and constitute an important class of graphs (see, for example, [Yan et al., 1996]).

**Remark 1** *For fixed positive integers  $a_1, a_2, b_1, b_2$ , we may note that  $(K_{a_1} \cup K_{b_1}) \oplus (K_{a_2} \cup K_{b_2})$  is a non-chordal cograph. To see this, it suffices to choose one vertex at each different copy of the complete graph to obtain an induced  $C_4$ . In particular, if  $c \geq 2$ , then the graph of the form  $\overline{K_c} \oplus (K_{a_2} \cup K_{b_2})$  is a non-chordal cograph. On the other hand, for  $t \geq 1$  and  $a_i \geq 1, 1 \leq i \leq t$ , the graph  $tK_{a_i}$  is a disconnected chordal cograph.*

### 2.3 Signless Laplacian matrix and their main eigenvalues

For a simple graph  $G$  with vertices  $v_1, \dots, v_n$ , its adjacency matrix  $\mathbf{A}(G) = [a_{ij}]$  is the square matrix of order  $n$  where  $a_{ij} = 1$  if  $v_i$  and  $v_j$  are adjacent and 0, otherwise. The *signless Laplacian matrix*  $\mathbf{Q}(G)$  is defined as  $\mathbf{Q}(G) = \mathbf{D}(G) + \mathbf{A}(G)$ , where  $\mathbf{D}(G)$  is the diagonal matrix whose entries are the degrees of the vertices. The matrix  $\mathbf{Q} = \mathbf{Q}(G)$  is symmetric and positive semidefinite and then it has  $n$  real non-negative eigenvalues, which will be called  *$\mathbf{Q}$ -eigenvalues* of  $G$ . We denote the multiset of these eigenvalues by  $\sigma(G) = \{q_1^{[r_1]}, q_2^{[r_2]}, \dots, q_p^{[r_p]}\}$ , where  $q^{[r]}$  means that the eigenvalue  $q$  has multiplicity  $r \geq 1$ .

The following known property of the matrix  $\mathbf{Q}(G)$  of a bipartite graph  $G$  is stated here for future reference.

**Lemma 3** [Cvetković et al., 2007] *For any connected graph  $G$ , the least  $\mathbf{Q}$ -eigenvalue of  $G$  is 0 if and only if  $G$  is bipartite. In this case, 0 is a simple eigenvalue. In general, the multiplicity of 0 as a  $\mathbf{Q}$ -eigenvalue of  $G$  is the number of bipartite components of  $G$ .*

An eigenvector  $\mathbf{v}$  associated with a  $\mathbf{Q}$ -eigenvalue  $q$  of  $G$  is a *main eigenvector* of  $G$  if  $\mathbf{v}^\top \mathbf{j} \neq 0$ . In this case,  $q$  is said to be a *main  $\mathbf{Q}$ -eigenvalue* of  $G$ . From the Perron-Frobenius' Theorem for non-negative irreducible real symmetric matrices, if  $G$  is a connected graph on  $n \geq 2$  vertices then its largest  $\mathbf{Q}$ -eigenvalue is simple and main, since it has an associated eigenvector whose entries are all positive. The main  $\mathbf{Q}$ -eigenvalues of a graph may be considered as pairwise distinct eigenvalues, since it is possible to find an orthonormal basis for each one of the associated eigenspaces containing only a single eigenvector non-orthogonal to  $\mathbf{j}$ . This fact, whose proof can be found in [Sciriha and Cardoso, 2012, Remark 3], will be taken into account throughout the paper.

The facts about main  $\mathbf{Q}$ -eigenvalues of a graph listed below will be applied in the remaining of the text.

**Lemma 4** [Deng and Huang., 2013], [Chen and Huang., 2013] *A graph  $G$  has exactly one main signless Laplacian eigenvalue if and only if it is regular.*

**Lemma 5** [Vinagre et al., 2020] *A graph  $G$  and its complement  $\overline{G}$  have the same number of main  $\mathbf{Q}$ -eigenvalues.*

The  $\mathbf{Q}$ -spectrum of the union of two graphs is the union of the  $\mathbf{Q}$ -spectra of the original graphs [see Cvetković and Simić, 2009]. It follows that the main  $\mathbf{Q}$ -eigenvalues of the union of two graphs are obtained from the union (here considered as a set, not a multiset) of the main- $\mathbf{Q}$  eigenvalues of each one of the graphs. The following two results are consequences of these facts.

**Lemma 6** [Vinagre et al., 2020] Let  $H$  and  $\overline{K_p}$ , for  $p \geq 1$  an integer, be graphs with disjoint vertex sets. A real number  $q \neq 0$  is a main  $\mathbf{Q}$ -eigenvalue of  $H$  with main eigenvector  $\mathbf{v}$  if and only if  $q$  is a main  $\mathbf{Q}$ -eigenvalue of the union  $G = \overline{K_p} \cup H$  with associated main eigenvector  $\begin{bmatrix} \mathbf{v} \\ \mathbf{0}_p \end{bmatrix}$ , where  $\mathbf{0}_p$  stands for the null vector on  $p$  coordinates.

The following result was proven for a connected  $H$ , but holds for a non-connected graph as well.

**Lemma 7** [Vinagre et al., 2020] Any graph  $G = \overline{K_p} \cup H$ , where  $p \geq 1$  is an integer and  $H$  is a graph, has 0 as a main  $\mathbf{Q}$ -eigenvalue.

### 3 A bound on the number of main $\mathbf{Q}$ -eigenvalues of a cograph

In this section, we use a cograph representation to provide a bound on the number of main  $\mathbf{Q}$ -eigenvalues of cographs. Before doing so, we use Lemma 7 to prove a result that predicts the number of main  $\mathbf{Q}$ -eigenvalues of a particular cograph.

**Theorem 1** Let  $G$  be cograph with  $k \geq 2$  main  $\mathbf{Q}$ -eigenvalues. Then

- (a) For any integer  $c \geq 1$ , the cograph  $K_c \oplus G$  has  $k$  or  $k + 1$  main  $\mathbf{Q}$ -eigenvalues;
- (b) For each integer  $c \geq 1$ , the cograph  $K_c \oplus G$  has  $k + 1$  main  $\mathbf{Q}$ -eigenvalues if and only if  $\overline{G}$  is not bipartite.

**Proof:** Let  $G$  be a cograph with  $k \geq 2$  main  $\mathbf{Q}$ -eigenvalues and let  $c \geq 1$  be an integer. By Lemma 5, we know that  $\overline{G}$  has  $k$  main  $\mathbf{Q}$ -eigenvalues. By invoking Lemma 7, we notice that  $\overline{K_c} \cup \overline{G}$  has 0 as a main  $\mathbf{Q}$ -eigenvalue. Regarding item (a), we need to consider whether 0 is a main  $\mathbf{Q}$ -eigenvalue of  $\overline{G}$  or not. Since we are considering that the main  $\mathbf{Q}$ -eigenvalues are all distinct, in the first case we have that  $\overline{K_c} \cup \overline{G}$  remains with  $k$  main  $\mathbf{Q}$ -eigenvalues and, in the second, that the cograph  $\overline{K_c} \cup \overline{G}$  has  $k + 1$  main  $\mathbf{Q}$ -eigenvalues. Applying again Lemma 5, the assertion of item (a) is proved. Taking into account Lemma 3, the previous argumentation leads us to conclude also that the item (b) is true.

We may represent every cograph on vertex set  $\{1, \dots, n\}$  as a rooted tree, named *cotree*, whose nodes consist of  $n$  leaves labeled 1 through  $n$  and internal vertices that carry either the label “ $\cup$ ” for union or “ $\oplus$ ” for join. Given such a tree  $T$ , we can easily construct the corresponding cograph  $G_T$ . The construction can be made unique (see, for example [Hoppen et al., 2022]) as follows. We say that a cotree is in its *normalized form* if every internal node has at least two children and has a label that differs from the label of its parent. In other words, the children of nodes labeled  $\cup$  are leaves or nodes labeled  $\oplus$ , while the children of nodes labeled  $\oplus$  are leaves or nodes labeled  $\cup$ . In this note, we assume that the cotree representation of a cograph is always in normalized form.

Consider the cotree  $T_G$  of a cograph  $G$ . We group the leaves of the cotree that represent the vertices of  $G$  as follows. Let  $r$  be the number of distinct sets of siblings, which we call the *width* of the cotree. We call each of these sets of sibling vertices a *bag*  $B_i$  and denote the respective cardinality as  $t_i = |B_i|$ ,  $1 \leq i \leq r$ . When the siblings are *co-duplicates* (meaning the parent is  $\oplus$ ), we call it a *J-bag*, and when the siblings are *duplicates* (meaning the parent is  $\cup$ ), we call it a *U-bag*. If the cotree  $T_G$  has  $r$  bags  $B_i$  whose cardinalities are  $t_i$ , we can represent  $T_G$  as the sequence

$$\{B_i^{t_i}\}_{i=1}^r,$$

which we call *bag representation* of  $T_G$ .

We state the following result that uses this representation of cographs and is useful for our purposes.

**Proposition 1** Let  $G$  be a cograph with cotree  $T_G$  having  $r_1$  *J*-bags  $B_i^{t_i}$ ,  $i = 1, \dots, r_1$ , and  $r_2 = r - r_1$  *U*-bags  $B_j^{t_j}$ ,  $j = r_1 + 1, \dots, r$ . Let  $p_i$  be the degree of each of the vertices of  $G$  in the bag  $B_i$ ,  $i = 1, \dots, r$ . Then

- (a)  $q = p_i - 1$  and  $q = p_j$  are eigenvalues of  $\mathbf{Q}(G)$  with multiplicities at least  $t_i - 1$ , for  $i = 1, \dots, r_1$ , and  $t_j - 1$ , for  $j = r_1 + 1, \dots, r$ , respectively.
- (b) All of them are non-main  $\mathbf{Q}$ -eigenvalues of  $G$ .

**Proof:** We notice that item (a) is given in [Jones et al., 2023, Proposition 2(c)]. In any case, the proof of item (b) we give now, also proves item (a). For a cograph  $G$  as above, where each vertex within the bag  $B_i$  has degree equal to  $p_i$ , we consider the matrix  $\mathbf{Q}(G)$ . For each pair of vertices  $u, v$  belonging to the same *J*-bag (respectively, *U*-bag)  $B_i$ , we consider the vector whose entries are all equal to 0 except those corresponding to  $u$  and  $v$ , which are taken to be 1 and

-1, respectively. We obtain a  $\mathbf{Q}(G)$ -eigenvector associated with eigenvalue  $p_i - 1$  (respectively,  $p_i$ ). This reasoning allows one to conclude that, for each  $i = 1, \dots, r_1$ , the integer  $p_i - 1$  is a non-main  $\mathbf{Q}$ -eigenvalue of the graph  $G$  with multiplicity at least  $t_i - 1$  and for each  $j = r_1 + 1, \dots, r$ ,  $p_j$  is a non-main  $\mathbf{Q}(G)$ -eigenvalue with multiplicity at least  $t_j - 1$ .

We observe that, from Proposition 1, we obtain  $n - r$   $\mathbf{Q}$ -eigenvalues of  $G$  that are non-main. Further, as proved in [Jones et al., 2023], the remaining  $r$  eigenvalues are those of the  $r \times r$  *condensed signless Laplacian matrix*, which is the symmetric matrix defined as  $\mathbf{C}(G) = [c_{ij}]$ , where

$$c_{ij} = \begin{cases} p_i + (t_i - 1), & \text{if } i = j \text{ and } 1 \leq i \leq r_1; \\ p_i, & \text{if } i = j \text{ and } r_1 + 1 \leq i \leq r; \\ \sqrt{t_i t_j} z_{ij} & \text{if } i \neq j \text{ and } 1 \leq i, j \leq r, \end{cases}$$

where  $z_{ij} = 1$  or  $0$ , depending on whether or not the vertices of bag  $B_i$  are adjacent to the vertices of  $B_j$ . Thus, the matrix  $\mathbf{C}(G)$  contains the main  $\mathbf{Q}$ -eigenvalues of the graph  $G$  and we obtain the following result as a consequence.

**Theorem 2** *If  $G$  is a cograph with bag representation*

$$\{B_i^{t_i}\}_{i=1}^r,$$

*then  $G$  has at most  $r$  main  $\mathbf{Q}$ -eigenvalues.*

Hence, the number of main  $\mathbf{Q}$ -eigenvalues of a cograph does not exceed the width of its cotree. We notice that, in general, this bound can be far from optimal. For example, as we will see in Lemma 9, the cograph  $K_c \oplus (tK_a)$ , for integers  $a, c \geq 1$  and  $t \geq 2$  has two main  $\mathbf{Q}$ -eigenvalues while its width, which is  $t + 1$ , can be arbitrarily large.

In the sequence, we apply the facts mentioned in the proof of Proposition 1 to identify the main  $\mathbf{Q}$ -eigenvalues of a connected bipartite cograph that, according to Lemma 2, has the form  $\overline{K_a} \oplus \overline{K_b}$  for some integers  $1 \leq a \leq b$ .

Firstly, we may recall from [Grone et al., 1990] that the  $\mathbf{Q}$ -spectrum of the complete graph on  $n$  vertices is

$$\sigma(K_n) = \{(2n - 2)^*, (n - 2)^{[(n-1)]}\},$$

where  $q^*$  indicates that  $q$  is a main  $\mathbf{Q}$ -eigenvalue of the graph.

For bipartite graphs, as the  $\mathbf{Q}$ -characteristic polynomial and the  $\mathbf{L}$ -characteristic polynomial are equal (definition of Laplacian matrix in the next section), it is known (see, for example, [Cvetković et al., 2007] and [Grone et al., 1990]) that  $\sigma(\overline{K_a} \oplus \overline{K_b}) = \{(a + b), b^{[a-1]}, a^{[b-1]}, 0\}$ , for  $a, b$  non-negative integers. If  $a, b \geq 1$  are distinct integers, we claim that the main  $\mathbf{Q}$ -eigenvalues of  $G = \overline{K_a} \oplus \overline{K_b}$  are  $a + b$  and  $0$ . Indeed, from the proof of Proposition 1, the condensed signless Laplacian matrix of this graph is

$$\mathbf{C}(\overline{K_a} \oplus \overline{K_b}) = \begin{bmatrix} b & \sqrt{ab} \\ \sqrt{ab} & a \end{bmatrix},$$

whose eigenvalues are exactly  $a + b$  and  $0$ . From Theorem 2, the number  $k$  of main  $\mathbf{Q}$ -eigenvalues is at most 2. But since  $G$  is not a regular graph, we have that  $k = 2$ , by Lemma 4. In case  $a = b$ , the graph is regular and there is a single main  $\mathbf{Q}$ -eigenvalue, which is  $2a$ . For future reference, we state the following result.

**Lemma 8** *If  $a, b$  are positive integers with  $a \neq b$  then*

$$\sigma(\overline{K_a} \oplus \overline{K_b}) = \{(a + b)^*, b^{[a-1]}, a^{[b-1]}, 0^*\}.$$

*In the particular case when  $a = b$ , the graph is regular and*

$$\sigma(\overline{K_a} \oplus \overline{K_a}) = \{(2a)^*, a^{[2a-2]}, 0\}.$$

From this result, we obtain the number of main  $\mathbf{Q}$ -eigenvalues of two types of cographs which we will see, in Proposition 2 below, that are *quasi*-threshold graphs.

**Lemma 9** *Consider the cographs*

$$\begin{aligned} G_1 &= K_c \oplus (K_a \cup K_b), \text{ for integers } a, b, c \geq 1 \text{ with } a \neq b \text{ and} \\ G_2 &= K_c \oplus (tK_a), \text{ for integers for } t \geq 2, a, c \geq 1. \end{aligned}$$

*The number of main  $\mathbf{Q}$ -eigenvalues of  $G_1$  and  $G_2$  is 2.*

**Proof:** In fact, the signless Laplacian condensed matrix  $\mathbf{C}(G_1)$  is the  $3 \times 3$  matrix

$$\begin{bmatrix} a + b + 2(c - 1) & \sqrt{ca} & \sqrt{cb} \\ \sqrt{ca} & 2(a - 1) + c & 0 \\ \sqrt{cb} & 0 & 2(b - 1) + c \end{bmatrix},$$

for which the eigenvalues are  $a + b + c - 2$  and the two roots of the equation

$$q^2 - (3c + 2b + 2a - 4)q + 2c^2 + (2b + 2a - 6)c + (4a - 4)b - 4a + 4 = 0.$$

From Proposition 1, we see that  $q = a + b + c - 2$  is one of the non-main  $\mathbf{Q}(G_1)$ -eigenvalues and that its multiplicity is at least  $c - 1$ , since it corresponds to the degrees of vertices of the  $J$ -bag with  $c$  vertices. We claim that  $q = a + b + c - 2$  actually has multiplicity equal to  $c$  as a non-main eigenvalue of  $\mathbf{Q}(G_1)$ . Indeed, it is not difficult to see

that  $\mathbf{C}(G)\mathbf{v} = (a + b + c - 2)\mathbf{v}$ , for  $\mathbf{v} = \begin{bmatrix} 1 \\ c/(b - a) \\ -c/(b - a) \end{bmatrix}$ . Let us denote by  $\mathbf{J}_{r \times s}$  (respectively,  $\mathbf{O}_{r \times s}$ ) the all ones matrix (respectively, the null matrix) of order  $r \times s$ , by  $\mathbf{j}_s$  the all ones vector of length  $s$  and by  $\mathbf{I}_r$  the identity matrix of order  $r$ . Then, we can describe  $\mathbf{Q}(G_1)$  as

$$\mathbf{Q}(G_1) = \begin{bmatrix} \mathbf{J}_{c \times c} + (a + b + c - 2)\mathbf{I}_c & \mathbf{J}_{c \times a} & \mathbf{J}_{c \times b} \\ \mathbf{J}_{a \times c} & \mathbf{J}_{a \times a} + (a + c - 2)\mathbf{I}_a & \mathbf{O}_{a \times b} \\ \mathbf{J}_{b \times c} & \mathbf{O}_{b \times a} & \mathbf{J}_{b \times b} + (b + c - 2)\mathbf{I}_b \end{bmatrix}$$

and verify that  $\mathbf{w} = \begin{bmatrix} \mathbf{j}_c \\ c/(b - a)\mathbf{j}_a \\ -c/(b - a)\mathbf{j}_b \end{bmatrix}$  is an eigenvector of  $\mathbf{Q}(G_1)$  associated with the eigenvalue  $q = a + b + c - 2$ , which forms a linearly independent set when grouped with the other eigenvalues associated with  $q$  provided by Proposition 1. Furthermore, the coordinates of  $\mathbf{w}$  add up to zero. This completes our claim. Since  $G_1$  is not regular, the two remaining eigenvalues of  $\mathbf{C}(G_1)$  must be main  $\mathbf{Q}$ -eigenvalues of  $G_1$ . The assertion is proved for  $G_1$ .

For  $G_2$ , by Lemma 5, it suffices to compute the number of main  $\mathbf{Q}$ -eigenvalues of  $\overline{G_2} = \overline{K_c} \cup (\oplus_{i=1}^t \overline{K_a})$ . Now,  $\overline{K_c}$  has 0 as the only main  $\mathbf{Q}$ -eigenvalue. As for  $(\oplus_{i=1}^t \overline{K_a})$ , it is a regular graph having also a unique main  $\mathbf{Q}$ -eigenvalue. The degree of regularity is  $(t - 1)a$ , which is non-zero, since  $t > 1$ , hence its main  $\mathbf{Q}$ -eigenvalue is non-zero, proving the result for  $G_2$ .

#### 4 *Quasi*-threshold graphs with $k$ main $\mathbf{Q}$ -eigenvalues

The *Laplacian matrix* of a graph  $G$  is defined as  $\mathbf{L}(G) = \mathbf{D}(G) - \mathbf{A}(G)$ . As  $\mathbf{L}(G)$  is symmetric and positive semidefinite, its eigenvalues are non-negative real numbers. Fiedler showed in [Fiedler, 1973] that  $G$  is a connected graph if and only if the second smallest eigenvalue of  $\mathbf{L}(G)$  is positive; for this reason, this eigenvalue is called the *algebraic connectivity* of  $G$  and denoted by  $a(G)$ . Moreover, Fiedler proved that for  $G \neq K_n$ ,  $a(G) \leq \kappa(G)$ , where  $\kappa(G)$  is the vertex connectivity of  $G$ .

Graphs for which  $a(G) = \kappa(G)$  were characterized in [Kirkland et al., 2002] taking into account their structures. It is known that  $\kappa(G) = a(G)$  for cographs. A proof of this result can be found in [Lucas et al., 2022] and we state it here for easy reference.

**Lemma 10** [Lucas et al., 2022] *Let  $G$  be a cograph. Then*

$$\kappa(G) = a(G).$$

The particular case of chordal graphs for which  $a(G) = \kappa(G)$  was treated in Abreu et al. [2021] and we state the result here as the following lemma.

**Lemma 11** [Abreu et al., 2021, Theorem 2] *Let  $G$  be a non-complete connected chordal graph. Then  $\kappa(G) = a(G)$  if and only if there is a minimal separator of  $G$  such that all its elements are universal vertices.*

The next result gives a structural characterization of *quasi*-threshold graphs that are connected.

**Proposition 2** *Let  $G$  be a connected quasi-threshold. If  $G$  is a non-complete graph, then  $G = K_c \oplus H$ , for some integer  $c \geq 1$  and a disconnected quasi-threshold graph  $H$ .*

**Proof:** Let  $G$  be a cograph which is connected and chordal. Since it is not complete, it follows, by Lemma 10, that  $a(G) = \kappa(G)$  and thus, following Lemma 11,  $G$  admits a minimal vertex separator  $S$  in which all the vertices are universal, implying that  $S$  is a clique. Let  $|S| = c$ ,  $c \geq 1$ , be the cardinality of  $S$ . We have  $S = K_c$ , and because all vertices of  $S$  are universal, we can describe  $G$  as  $G = K_c \oplus H$ , where  $H$  is a disconnected *quasi*-threshold graph, proving the result.

We now give information about the structure of a connected *quasi*-threshold graph  $G$  having  $k$  main  $\mathbf{Q}$ -eigenvalues. Clearly, Corollary 1 and Lemma 4 imply that  $k = 1$  if and only if  $G$  is a complete graph.

**Theorem 3** *For a connected quasi-threshold graph  $G$ , let  $G = K_c \oplus H$  be its decomposition given in Proposition 2. If  $k \geq 2$  is the number of main  $\mathbf{Q}$ -eigenvalues of  $G$  then, exactly one of the following situations occurs:*

- (a)  $\overline{H}$  is non-bipartite, that is,  $\overline{H}$  has  $k - 1$  non-zero main  $\mathbf{Q}$ -eigenvalues or
- (b)  $\overline{H}$  is bipartite, and in this case,  $k = 2$ .

**Proof:** We notice that  $\overline{H}$  is connected, since  $H$  is disconnected. By Lemma 7, we know that  $\overline{G} = \overline{K_c} \cup \overline{H}$  has 0 as a main  $\mathbf{Q}$ -eigenvalue. Also, Lemma 6 asserts that a real  $q \neq 0$  is a main  $\mathbf{Q}$ -eigenvalue of  $\overline{G}$  if and only if  $q$  is a main  $\mathbf{Q}$ -eigenvalue of  $\overline{H}$ . We may recall that  $\overline{G}$  also has  $k$  main  $\mathbf{Q}$ -eigenvalues, by Lemma 5. Thus, in accordance to all these facts, we need to consider the only two possible cases: (i)  $\overline{H}$  has  $k - 1$  non-zero main  $\mathbf{Q}$ -eigenvalues; and (ii)  $\overline{H}$  has  $k$  main  $\mathbf{Q}$ -eigenvalues, one of which is 0. In case (i), we see that  $\overline{H}$  can not be bipartite by Lemma 3 and, moreover, that all its  $k - 1$  main  $\mathbf{Q}$ -eigenvalues are non-zero. This proves item (a). As for case (ii), it follows, from Lemmas 3 and 11, that  $\overline{H}$  is a connected bipartite chordal cograph, which is of the form  $\overline{K_a} \oplus \overline{K_b}$ , for some positive integers  $a, b$ , by Lemma 2. Moreover, Lemma 8 ensures that  $\overline{H}$  has exactly one non-zero main  $\mathbf{Q}$ -eigenvalue. Hence  $G = K_c \oplus (K_a \cup K_b)$  for some integers  $a, b, c \geq 1$ , which has two main  $\mathbf{Q}$ -eigenvalues, proving item (b).

If the number of main  $\mathbf{Q}$ -eigenvalues of a connected chordal cograph is 2, we can be more precise and determine exactly which graph it is.

**Theorem 4**  *$G$  is a connected quasi-threshold graph with two main  $\mathbf{Q}$ -eigenvalues if and only if  $G$  assumes one of the following expressions:*

- (a)  $G = K_c \oplus (K_a \cup K_b)$  for integers  $a, b, c \geq 1$  and  $a \neq b$ ;
- (b)  $G = K_c \oplus (tK_a)$ , for integers  $t \geq 2$ ,  $a, c \geq 1$ .

**Proof:** Let  $G$  be a cograph which is connected and chordal and has two main  $\mathbf{Q}$ -eigenvalues. By repeating the argument in the proof of Theorem 3, we conclude that  $G = K_c \oplus H$ , for some integer  $c \geq 1$  and  $H$  a disconnected *quasi*-threshold graph. Moreover, the only possible cases are: (i)  $\overline{H}$  has two main  $\mathbf{Q}$ -eigenvalues, one of which is zero; and (ii)  $\overline{H}$  has only one main  $\mathbf{Q}$ -eigenvalue. Concerning case (i), it follows from Lemma 3 that  $\overline{H}$  is a connected bipartite cograph, and it is of the form  $\overline{H} = \overline{K_a} \oplus \overline{K_b}$ , by Lemma 2, for some positive integers  $a \neq b$  (if  $a = b$  then  $\overline{H}$  would be regular and consequently with a single main  $\mathbf{Q}$ -eigenvalue). Hence  $G = K_c \oplus (K_a \cup K_b)$  for some integers  $a, b, c \geq 1$ ,  $a \neq b$ . In case (ii), the assumption on  $\overline{H}$  implies that  $H$  is a regular graph, by Lemmas 4 and 5. So,  $G = K_c \oplus H$ , where  $H$  is a cograph which is disconnected, regular and chordal. In view of Corollary 1 and Remark 1,  $H$  is the union of at least two copies of the same complete graph  $K_a$ , where  $a \geq 1$  is an integer (a unique copy of a complete graph would imply  $G$  to be complete, a contradiction). So  $G = K_c \oplus (tK_a)$ , for integers  $t \geq 2$  and  $a \geq 1$ . Now, combining these two cases, we have that  $G$  has one of the forms of items (a) or (b). Reciprocally, if  $G$  has the forms of item (a) or item (b), Lemma 8 asserts that  $G$  has two main  $\mathbf{Q}$ -eigenvalues.

**Remark 2** *In Theorem 4, for an integer  $c \geq 2$ , the quasi-threshold graphs having the form  $K_c \oplus (K_a \cup K_b)$ , with  $a = 1$  or  $b = 1$ , together with those of the form  $K_c \oplus (tK_a)$ , with  $t \geq 2$  and  $a = 1$ , are exactly the connected threshold graphs with two main  $\mathbf{Q}$ -eigenvalues whose characterization was given in Vinagre et al. [2020]. In the latter case, we have the graphs  $G = K_c \oplus \overline{K_t}$ , the so called complete split graphs.*

## 5 Main $\mathbf{Q}$ -eigenvalues of generalized core-satellite graphs

In [Estrada and Benzi, 2017], the graphs composed by a central clique (the *core*) connected to several other cliques of same size (the *satellites*) were called *core-satellite graphs*; they generalize both the complete split graphs and the windmill graphs. The authors of [Estrada and Benzi, 2017] also considered the generalization of core-satellite graphs to the case where the satellites cliques are not restricted to having all the same order; they called these graphs

*generalized core-satellite*. It is not difficult to see that these graphs are also *quasi*-threshold graphs. This fact was pointed out in [Yan et al., 1996].

Then the general form of a generalized core-satellite graph is

$$G = K_{n_0} \oplus (a_1 K_{n_1} \cup a_2 K_{n_2} \cup \dots \cup a_p K_{n_p}) = K_{n_0} \oplus \left( \bigcup_{i=1}^p a_i K_{n_i} \right),$$

for integers  $p, a_1, a_2, \dots, a_p, n_0, n_1, \dots, n_p \geq 1$  and  $n_i \neq n_j$ , for all  $1 \leq i < j \leq p$ . Thus  $G$  has a total of  $\sum_{i=1}^p a_i$  satellites of  $p$  different orders. In particular, the graphs in Theorem 4 have at least two satellites. The ones of item (a) are generalized core-satellite graphs with  $p = 2$  and  $a_1 = a_2 = 1$ , while the graphs of item (b) are core-satellite graphs having at least two satellites, that is,  $p = 1$  and  $a_1 = t > 1$ . Both types of graphs have two main  $\mathbf{Q}$ -eigenvalues. We notice that a generalized core-satellite with one satellite is a complete graph. Hence, we may assume that either  $p > 1$ , or that  $p = 1$  and  $a_1 \geq 2$ .

In the sequence, for a given generalized core-satellite graph  $G$ , we determine the exact number  $k \geq 2$  of main  $\mathbf{Q}$ -eigenvalues of  $G$ .

**Theorem 5** *Let  $G$  be a generalized core-satellite graph. Then:*

- (a)  $G$  has  $k = 2$  main  $\mathbf{Q}$ -eigenvalues if and only if  $G$  has exactly two satellites, all with different orders, or  $G$  has two or more satellites, all of the same order.
- (b) Let  $k \geq 3$ . Then  $G$  has  $k$  main  $\mathbf{Q}$ -eigenvalues if and only if  $G$  has at least three satellites, divided into  $k - 1$  different orders.

**Proof:** Let  $G$  be a generalized core-satellite graph  $G = K_{n_0} \oplus H$ , where

$$H = \bigcup_{i=1}^p a_i K_{n_i},$$

for integers  $p, a_1, a_2, \dots, a_p, n_0, n_1, n_2, \dots, n_p \geq 1$  and  $n_i \neq n_j$ , for all  $1 \leq i < j \leq p$ .

(a) By Theorem 4,  $G$  has two main  $\mathbf{Q}$ -eigenvalues if and only if  $p = 2$  and  $a_1 = a_2 = 1$  or  $p = 1$  and  $a_1 \geq 2$ .

(b) Assume  $p \geq 2$ . Firstly, we note that the cograph  $H = a_1 K_{n_1} \cup a_2 K_{n_2} \cup \dots \cup a_p K_{n_p}$  has exactly  $p$  main  $\mathbf{Q}$ -eigenvalues, since main eigenvalues are considered to be distinct. Therefore, its complement graph  $\overline{H} = a_1 \overline{K_{n_1}} \oplus a_2 \overline{K_{n_2}} \oplus \dots \oplus a_p \overline{K_{n_p}}$  also has  $p$  main  $\mathbf{Q}$ -eigenvalues, which are all non-zero by Lemmas 3 and 2, since  $\overline{H}$  is not bipartite for  $p \geq 3$  or  $p = 2$  and  $a_1 \geq 2$  or  $a_2 \geq 2$ . Thus,  $\overline{K_{n_0}} \cup \overline{H}$  has  $p + 1$  main  $\mathbf{Q}$ -eigenvalues, and so has its complement, which is the initial graph  $G = K_{n_0} \oplus H$ . We conclude that the generalized core-satellite graph  $G$  has  $k \geq 3$  main  $\mathbf{Q}$ -eigenvalues if and only if  $k = p + 1$ , for  $p = 2$  and  $a_1 \geq 2$  or  $a_2 \geq 2$  or  $p \geq 3$ , that is,  $G$  has at least three satellites of  $k - 1$  different orders.

## 6 Constructing *quasi*-threshold non-generalized-core-satellite graphs with three main $\mathbf{Q}$ -eigenvalues

In [Vinagre et al., 2020], a recursive method to construct all threshold graphs with  $k + 1$  main  $\mathbf{Q}$ -eigenvalues from threshold graphs having  $k$  main  $\mathbf{Q}$ -eigenvalues was presented. It is a natural question to consider whether a similar procedure to construct *quasi*-threshold graphs with  $k + 1$  main  $\mathbf{Q}$ -eigenvalues from *quasi*-threshold graphs having  $k$  main  $\mathbf{Q}$ -eigenvalues may be feasible.

In spite of the fact that we have a complete characterization of the basis case ( $k = 2$ ), as well as the characterization of all generalized core-satellite graphs – a subclass of *quasi*-threshold graphs – having  $k \geq 2$  main  $\mathbf{Q}$ -eigenvalues (see Theorem 5), which are steps towards the investigation of main  $\mathbf{Q}$ -eigenvalues of cographs, a similar construction does not seem possible for *quasi*-threshold graphs in general. To justify this assertion, we present in the sequel several examples of *quasi*-threshold graphs, which are not generalized core-satellite graphs, having three main  $\mathbf{Q}$ -eigenvalues. We understand that these examples show that a complete description of all *quasi*-threshold graphs having  $k > 2$  main  $\mathbf{Q}$ -eigenvalues is a very hard problem.

In order to present our constructions, we start by recalling that, by Proposition 2, a connected non-complete *quasi*-threshold graph can be described as  $G = K_c \oplus H$ , for some integer  $c \geq 1$  and a disconnected *quasi*-threshold graph



$H$ . Also, from Theorem 3, we know that, if such graph  $G$  has three main  $\mathbf{Q}$ -eigenvalues then the graph  $H$  has two main  $\mathbf{Q}$ -eigenvalues, if we take Lemma 5 into account.

To present our constructions, we recall some facts found in the literature that we state as the following remark for further reference.

**Remark 3** *The  $\mathbf{Q}$ -spectrum of the split complete graph  $K_a \oplus \overline{K_b}$ , where  $a \geq 1$  and  $b > 1$  is*

$$\{(a+b-2)^{[a-1]}, a^{[b-1]}\} \cup \{q_1^*, q_2^*\}$$

where  $q_1, q_2$  are the two roots of the equation  $q^2 - (b+3a-2)q + (2a^2-2a) = 0$ . Since  $a+b-2$  and  $a$  are the non-main eigenvalues determined by Proposition 1, the two main  $\mathbf{Q}$ -eigenvalues of the graph are  $q_1$  and  $q_2$  as indicated, considering that the graph is not regular. For our purposes, we give below the signless Laplacian condensed matrix of the graph, which is,

$$\mathbf{C}(K_a \oplus \overline{K_b}) = \begin{bmatrix} 2(a-1)+b & \sqrt{ab} \\ \sqrt{ab} & a \end{bmatrix}.$$

In [de Freitas et al., 2010], the authors give a condition on  $a$  and  $b$  under which the eigenvalues  $q_1$  and  $q_2$  are integers and provide some examples of such graphs, as the infinite family where  $a = 2s - 1$  and  $b = 3s$ , for each integer  $s \geq 1$ : indeed, it holds that  $q_1 = 8s - 4$  and  $q_2 = s - 1$ , for  $s \geq 1$ .

On the other hand, from Lemma 9 we have that the two main  $\mathbf{Q}$ -eigenvalues of the graph  $K_c \oplus (K_{a_1} \cup K_{a_2})$ , where  $c, a_1, a_2$  are positive integers with  $a_1 \neq a_2$ , are the roots  $q_1, q_2$  of a certain quadratic equation. Also in this case, we find in [de Freitas et al., 2010] conditions on parameters  $a_1, a_2, c$  which allow the spectrum of this kind of graph to consist entirely of integers. For example, for each positive integer  $s$ , if  $c = s$ ,  $a_1 = s + 1$  and  $a_2 = s + 2$  then  $q_1 = 5s + 2$  and  $q_2 = 2s$ ; and if  $c = a_1 = a_2 = s$  then  $q_1 = 5s - 2$  and  $q_2 = 2s - 2$ .

In the following, we apply the facts listed in Remark 3 to describe some disconnected *quasi*-threshold graphs with two main  $\mathbf{Q}$ -eigenvalues, satisfying the conditions of item (a) of Theorem 3, that could replace graph  $H$  in decomposition given in Proposition 2. Actually, in each item of the example below, we give an infinite family of such graphs. By using these graphs  $H$ , connected *quasi*-threshold graphs of type  $G = K_c \oplus H$  with three main  $\mathbf{Q}$ -eigenvalues, which are not generalized core-satellite graphs, can be constructed.

### Example 1

(a) For integers  $p, a \geq 1$  and  $b \geq 2$ , the graph

$$H_1(a, b, p) = \overline{K_a} \cup pK_b,$$

has the main  $\mathbf{Q}$ -eigenvalues  $2b - 2$  and  $0$ ;

(b) For integers  $p \geq 2$  and  $a, b \geq 2$ ,

$$H_2(a, b, p) = p(K_a \oplus \overline{K_b})$$

has two main  $\mathbf{Q}$ -eigenvalues in accordance to Remark 3. In the particular case

$$H_2'(b, p) = H_2(1, b, p) = p(K_1 \oplus \overline{K_b}),$$

for integers  $p \geq 2$  and  $b \geq 2$ , we have the union of at least two bipartite chordal cographs with  $1 + b$  and  $0$  as their main  $\mathbf{Q}$ -eigenvalues. And from this latter case, we also obtain

$$H_2''(b, p_1, p_2) = \overline{K_{p_1}} \cup p_2(K_1 \oplus \overline{K_b}),$$

for positive integers  $p_1$  or  $p_2 \geq 2$  and  $b \geq 2$ , with the same main  $\mathbf{Q}$ -eigenvalues.

(c) For integers  $p \geq 2, s \geq 1$  and  $a_1 \geq 2$  or  $a_2 \geq 2$ , the graph

$$H_3(s, a_1, a_2, p) = p(K_s \oplus (K_{a_1} \cup K_{a_2}))$$

has two main  $\mathbf{Q}$ -eigenvalues as we have seen in Lemma 9;

(d) By choosing appropriate values for the parameter  $b$  in the above graph of type  $H_2'(b, p)$ , we obtain, for example, for positive integers  $a \geq 2, p_1, p_2$ , the graph

$$H_4(a, p_1, p_2) = p_1 K_a \cup p_2 (K_1 \oplus \overline{K_{(2a-3)}}),$$

for which the main  $\mathbf{Q}$ -eigenvalues are  $2a - 2$  and  $0$ , by Lemma 8.

(e) From the two previous cases, for positive integers  $a \geq 2, p_1, p_2, p_3$ , we can define

$$H_5(a, p_1, p_2, p_3) = p_1 K_a \cup p_2 (K_1 \oplus \overline{K_{(2a-3)}}) \cup \overline{K_{p_3}},$$

for which the main  $\mathbf{Q}$ -eigenvalues are also  $2a - 2$  and  $0$ ;

(f) By taking into account the conditions presented in Remark 3, we can choose appropriate values for the parameters, to obtain, for example, the graph

$$H_6(s, p_1, p_2, p_3) = p_1 (K_{(2s-1)} \oplus \overline{K_{(3s)}}) \cup p_2 K_{(4s-1)} \cup p_3 K_{(s+1)/2},$$

where  $s, p_1, p_2, p_3$  are positive integers and  $s$  is odd, which has  $8s - 4$  and  $s - 1$  as main  $\mathbf{Q}$ -eigenvalues;

(g) Following the same ideas, we obtain

$$H_7(s, p_1, p_2, p_3) = p_1 (K_s \oplus (K_{s+1} \cup K_{s+2})) \cup p_2 K_{(5s+4)/2} \cup p_3 K_{(s+1)},$$

for integers  $s, p_1, p_2, p_3 \geq 1$  such that  $s$  is even, for which the main  $\mathbf{Q}$ -eigenvalues are  $5s + 2$  and  $2s$ , by Remark 3;

(h) Still from the conditions presented in Remark 3, for integers  $s, p_1, p_2, p_3 \geq 1$  such that  $s$  is even, we obtain the graph

$$H_8(s, p_1, p_2, p_3) = p_1 (K_s \oplus (K_s \cup K_s)) \cup p_2 K_{(5s/2)} \cup p_3 K_s,$$

for which the main  $\mathbf{Q}$ -eigenvalues are  $5s - 2$  and  $2s - 2$ .

With the above example, we hope to have shown that a general characterization of *quasi*-threshold graphs with  $k \geq 3$  main  $\mathbf{Q}$ -eigenvalues, in the way it was done for threshold graphs in [Vinagre et al., 2020] and above, in the case of generalized core-satellites, does not seem to be possible. Even the case of connected cographs with  $k = 2$  main  $\mathbf{Q}$ -eigenvalues remains open. We note that the structural characterization of *quasi*-threshold graphs presented in Proposition 2, which was based on the description of the minimal separators of vertices presented in Lemma 11 of [Abreu et al., 2021], was fundamental for characterizing *quasi*-threshold connected graphs in Theorem 4. Therefore, in a possible approach to the case of connected cographs with  $k = 2$  main  $\mathbf{Q}$ -eigenvalues, it seems important to obtain a similar structural description.

## References

- D. Cvetković. The generating function for variations with restrictions and paths of the graph and self-complementary graphs. *Univ. Beograd Publ. Elektrotehn. Fak. Ser. Mat. Fiz.*, (320-328):27–34, 1970.
- Lin Chen and Qiong Xiang Huang. Trees, unicyclic graphs and bicyclic graphs with exactly two  $\mathbf{Q}$ -main eigenvalues. *Acta Math. Sin. (Engl. Ser.)*, 29(11):2193–2208, 2013.
- Hanyuan Deng and He Huang. On the main signless laplacian eigenvalues of a graph. *Electron. J. Linear Algebra*, 26: 381–393, 2013.
- Mehrnoosh Javarsineh and Gholam Hossein Fath-Tabar. On graphs with exactly three  $\mathbf{Q}$ -main eigenvalues. *Filomat*, 31(6):1803–1812, 2017a.
- Mehrnoosh Javarsineh and Gholam Hossein Fath-Tabar. Unicyclic and bicyclic graphs with exactly three  $\mathbf{Q}$ -main eigenvalues. *Appl. Math. Comput.*, 315:603–614, 2017b.
- Cybele T.M. Vinagre, Vilmar Trevisan, Johann Bolckau, and Rodrigo Chimelli. Characterizing threshold graphs with  $k$  main signless laplacian eigenvalues. *Linear Algebra and its Applications*, 602:33–45, 2020.
- A. Jones, L. de Castro, F. Pimenta, and I. Pinto. Graph filter: software for manipulating and searching graphs. version 3.1.2, july 25, 2023. Available on <http://sistemas.jf.ifsudestemg.edu.br/graphfilter>. Accessed: 09-28-2023.
- Jean RS Blair and Barry Peyton. An introduction to chordal graphs and clique trees. In *Graph theory and sparse matrix computation*, pages 1–29. Springer, 1993.
- D. G. Corneil, H. Lerchs, and L. Stewart Burlingham. Complement reducible graphs. *Discrete Appl. Math.*, 3(3):163–174, 1981. ISSN 0166-218X. doi: 10.1016/0166-218X(81)90013-5. URL [http://dx.doi.org/10.1016/0166-218X\(81\)90013-5](http://dx.doi.org/10.1016/0166-218X(81)90013-5).
- Marc Demange, Tinaz Ekim, and Dominique de Werra. Partitioning cographs into cliques and stable sets. *Discrete Optimization*, 2(2):145–153, 2005.

- Jing-Ho Yan, Jer-Jeong Chen, and Gerard J. Chang. *Quasi*-threshold graphs. *Discrete Appl. Math.*, 69(3):247–255, 1996.
- Dragoř Cvetković, Peter Rowlinson, and Slobodan K. Simić. Signless laplacians of finite graphs. *Linear Algebra and its Applications*, 423(1):155 – 171, 2007. Special Issue devoted to papers presented at the Aveiro Workshop on Graph Spectra.
- I. Sciriha and D. M. Cardoso. Necessary and sufficient conditions for a hamiltonian graph. *Journal of Combinatorial Mathematics and Combinatorial Computing*, 80:127–150, 2012.
- D. Cvetković and S. K. Simić. Towards a spectral theory of graphs based on signless laplacian I. *Publ. Inst. Math. (Beograd)*, 85(99):19–33, 2009.
- Carlos Hoppen, David P Jacobs, and Vilmar Trevisan. *Locating Eigenvalues in Graphs: Algorithms and Applications*. Springer Nature, 2022.
- Átila Jones, Vilmar Trevisan, and Cybele T.M. Vinagre. Exploring symmetries in cographs: Obtaining spectra and energies. *Discrete Applied Mathematics*, 325:120–133, 2023. ISSN 0166-218X.
- R. Grone, R. Merris, and V. S. Sunder. The laplacian spectrum of a graph. *SIAM J. Matrix. Anal. Appl.*, 11(2): 218–238, 1990.
- Miroslav Fiedler. Algebraic connectivity of graphs. *Czechoslovak Math. J.*, 23(98):298–305, 1973.
- Stephen J. Kirkland, Jason J. Molierno, Michael Neumann, and Bryan L. Shader. On graphs with equal algebraic and vertex connectivity. *Linear Algebra Appl.*, 341:45–56, 2002.
- Clarianne Luciola de A. V. M. Lucas, Renata R. Del-Vecchio, Maria Agueiras A. de Freitas, and Joice Santos do Nascimento. Relationships between algebraic connectivity and vertex connectivity. *Comput. Appl. Math.*, 41(3):Paper No. 105, 15, 2022.
- Nair Abreu, Claudia Marcela Justel, and Lilian Markenzon. Integer laplacian eigenvalues of chordal graphs. *Linear Algebra and its Applications*, 614:68–81, 2021. Special Issue ILAS 2019.
- Ernesto Estrada and Michele Benzi. Core–satellite graphs: Clustering, assortativity and spectral properties. *Linear Algebra and its Applications*, 517:30–52, 2017.
- Maria Agueiras A. de Freitas, Nair M.M. de Abreu, Renata R. Del-Vecchio, and Samuel Jurkiewicz. Infinite families of  $\mathbf{Q}$ -integral graphs. *Linear Algebra and its Applications*, 432(9):2352–2360, 2010. Special Issue devoted to Selected Papers presented at the Workshop on Spectral Graph Theory with Applications on Computer Science, Combinatorial Optimization and Chemistry (Rio de Janeiro, 2008).