Finite-sample expansions for the optimal error probability in asymmetric binary hypothesis testing

Valentinian Lungu * Ioannis Kontoyiannis [†]

May 30, 2024

Abstract

This is an expository note, not intended for publication.¹ The problem of binary hypothesis testing between two probability measures is considered. Sharp bounds are derived for the best achievable error probability of such tests based on independent and identically distributed observations. Specifically, the asymmetric version of the problem is examined, where different requirements are placed on the two error probabilities. Accurate nonasymptotic expansions with explicit constants are obtained for the error probability, using tools from large deviations and Gaussian approximation. Examples are shown indicating that, in the asymmetric regime, the approximations suggested by these bounds are significantly more accurate than the approximations provided by either normal approximation or error exponents.

Keywords — Hypothesis testing, error exponent, normal approximation, large deviations, sample complexity

1 Introduction

1.1 Binary hypothesis testing

Let P, Q be two probability measures on a measurable space (A, \mathcal{A}) , and let λ denote a dominating σ -finite measure so that both P and Q are absolutely continuous with respect to λ . Recall that such a λ always exists; e.g., we can take $\lambda = P + Q$.

We revisit the binary hypothesis testing problem between P and Q. The problem formulation is simple and quite elementary, though its importance can hardly be overstated, in view of its application across the sciences and engineering.

A deterministic test between P and Q consists of a decision region $B \subset A$, such that, given a sample $x \in A$, the test declares P to be the 'true' underlying distribution iff $x \in B$. Therefore, a deterministic test can always be expressed as the *deterministic* binary value of a decision function δ ,

$$\delta(x) = \begin{cases} 0 & \text{if } x \in B, \\ 1 & \text{if } x \in B^c. \end{cases}$$

^{*}Statistical Laboratory, Centre for Mathematical Sciences, University of Cambridge, Wilberforce Road, Cambridge CB3 0WB, UK. Email: vml26@cam.ac.uk. Supported by the Heilbronn Institute for Mathematical Research.

[†]Statistical Laboratory, Centre for Mathematical Sciences, University of Cambridge, Wilberforce Road, Cambridge CB3 0WB, UK. Email: yiannis@maths.cam.ac.uk.

¹A preliminary version of this paper appears at ISIT 2024 [20].

More generally, a randomised test is a family of probabilistic kernels $P_{Z|X}(z|x)$, $z \in \{0, 1\}$, $x \in A$, where, for each $x \in A$, $P_{Z|X}(\cdot|x)$ is a probability mass function on $\{0, 1\}$, and where both $P_{Z|X}(0|x)$ and $P_{Z|X}(1|x)$ are measurable functions of x. The test result is now the random binary value of the random variable Z which, conditional on X = x, has distribution $P_{Z|X}(\cdot|x)$, $x \in A$.

The two error probabilities associated with a test $P_{Z|X}$,

$$e_1 = e_1(P_{Z|X}) = \mathbb{P}(Z = 0|X \sim Q) = \int_A P_{Z|X}(0|x) \, dQ(x),$$

$$e_2 = e_2(P_{Z|X}) = \mathbb{P}(Z = 1|X \sim P) = \int_A P_{Z|X}(1|x) \, dP(x),$$

determine the test's performance. The best achievable performance among all deterministic or randomised tests can be described as the smallest possible value, $e_1^*(\epsilon)$, of the first error probability e_1 over all tests whose second error probability, e_2 , satisfies $e_2 \leq \epsilon$,

$$e_1^*(\epsilon) := \inf \{ e_1(P_{Z|X}) : P_{Z|X} \text{ s.t. } e_2(P_{Z|X}) \le \epsilon \}$$

Here, our interest is in the case of hypothesis tests between two product measures P^n and Q^n , corresponding to a sequence $X_1^n = (X_1, X_2, \ldots, X_n)$ of independent and identically (i.i.d.) observations with values in A. We write $e_{1,n}$ and $e_{2,n}$ for the error probabilities of a specific test, and we denote the best achievable value of the first error probability by:

$$e_{1,n}^*(\epsilon) := \inf \{ e_{1,n}(P_{Z|X_1^n}) : P_{Z|X_1^n} \text{ s.t. } e_{2,n}(P_{Z|X_1^n}) \le \epsilon \}.$$

1.2 Background

When the maximum allowed value $\epsilon > 0$ of the second error probability is fixed, the firstorder asymptotic behaviour of $e_{1,n}^*(\epsilon)$ is described by what has been come to be known as *Stein's lemma* [7, 19, 8]. It states that smallest achievable first error probability $e_{1,n}^*(\epsilon)$ decays exponentially with the sample size n,

$$\log e_{1,n}^*(\epsilon) = -nD(P||Q) + o(n), \quad \text{as } n \to \infty,$$

where all logarithms are taken to base e (log = log_e throughout), and D(P||Q) denotes the relative entropy between two probability measures P and Q on (A, A):

$$D(P||Q) := \begin{cases} \int \frac{dP}{dQ} \log\left(\frac{dP}{dQ}\right) dQ, & \text{if } P \ll Q, \\ +\infty, & \text{otherwise.} \end{cases}$$

Stein's lemma was refined by Strassen [26] who claimed that, as $n \to \infty$,

$$\log e_{1,n}^*(\epsilon) = -nD(P||Q) - \sqrt{n\sigma}\Phi^{-1}(\epsilon) - \frac{1}{2}\log n + O(1),$$
(1)

where σ^2 is the variance of the log-likelihood ratio $\log[\frac{dP}{dQ}(X_1)]$ with $X_1 \sim P$, and Φ denotes the standard normal distribution function. Although some issues of rigour were raised in [17] regarding Strassen's proof, an even stronger version of (1) was established by Polyanskiy-Poor-Verdú in [25], where explicit, finite-*n* bounds were obtained for the O(1) term. For the sake of completeness, we state and prove a version of these bounds in Theorems 2.3 and 2.4 in Section 2.

When the maximum allowed value $\epsilon > 0$ of the second error probability is not fixed, but is required to decay to zero exponentially fast, it turns out that, at least for a certain range of exponential rates, it is possible also have first probability of error decay to zero at an exponential rate as $n \to \infty$. Specifically, suppose that the second probability of error is required to be no greater than $e^{-n\delta}$, for some $\delta > 0$, and let $E_{1,n}^*(\delta)$ denote the best achievable $e_{1,n}$ error probability:

$$E_{1,n}^*(\delta) := e_{1,n}^*(e^{-n\delta})$$

In this case, Hoeffding [15] showed that, for any $\delta \in (0, D(Q || P))$,

$$\log E_{1,n}^*(\delta) = -nD(\delta) + o(n), \quad \text{as } n \to \infty,$$

where the optimal error exponent $D(\delta)$ is given by,

$$D(\delta) := \inf_{P': D(P'||P) \le \delta} D(P'||Q).$$

$$\tag{2}$$

A refinement of this, analogous to Strassen's refinement of Stein's lemma, was claimed by Csiszár and Longo [10], who stated that,

$$\log E_{1,n}^*(\delta) = -nD(\delta) - \frac{1}{2(1-\alpha^*)}\log n + O(1),$$
(3)

as $n \to \infty$, where α^* is an appropriate constant in (0, 1) that depends on P, Q and δ . The precise asymptotic limit of the O(1) term was determined in [11] and [28].

All of the works mentioned so far, including the one by Csiszár and Longo [10], are based on classical bounds and expansions related to normal approximation in the regime of the central limit theorem – namely, versions of the Berry-Esséen bound and the first-order Edgeworth expansion [12]. A slightly different approach was recently taken in [28]. There, the authors employ saddlepoint approximations [16] to derive precise expansions for both error probabilities, showing that, at best, we can have,

$$\log e_{1,n} = -nD(\delta) - \frac{1}{2}\log n + O(1),$$
(4)

and
$$\log e_{2,n} = -n\delta - \frac{1}{2}\log n + O(1)$$
, as $n \to \infty$. (5)

Although at first sight the expansions (4) and (5) might seem somewhat different from (3), it is actually easy to show that they are very closely related.

1.3 Finite-n asymptotic approximation

In the same sense in which the finite-*n* bounds of Polyanskiy-Poor-Verdú [25] strengthen the asymptotic expansion (1) of Strassen, it is possible to derive *finite-n* bounds that provide a corresponding stronger version of the asymptotic expansions (3), (4), (5) of [10, 28]. In Theorems 3.1 and 3.2 of Section 3 we provide explicit constants C and C' such that,

$$C' \le \log E_1^*(n,\delta) - \left[-nD(\delta) - \frac{1}{2(1-\alpha^*)} \log n \right] \le C,$$
 (6)

for all n greater than some explicit N_0 . These are exactly analogous to the bounds derived in [17] and [13] for lossless data compression without and with side information, respectively.

Lower bounds in the same spirit as our Theorem 3.2 were previously established for finite alphabets by Altuğ and Wagner [1, 2] and for general spaces by Nakiboğlu [21]. A general upper bound similar to that in our Theorems 3.1 was more recently also proved by Nakiboğlu [22].

Despite their technical nature, the approximations to $e_{1,n}^*$ provided by all these different approaches are of as much practical relevance as they are of mathematical interest. In particular, they each are useful in different regimes of the sample-size and the error probability requirements. For example, Stein's lemma and its refined form in (1) are only relevant for 'moderate' values of ϵ . If ϵ is small, then the second term dominates and the approximation is no longer valid or useful.

For example, consider the case of observations generated by either $P \sim \text{Bern}(0.6)$ or $Q \sim \text{Bern}(0.25)$ on $A = \{0, 1\}$. In the case when ϵ is relatively small, we examine the four approximations to $e_{1,n}^*(\epsilon)$ suggested by the above results. In the 'Stein regime',

Stein:
$$\log e_{1,n}^* \approx -nD(P||Q),$$
 (7)

Strassen:
$$\log e_{1,n}^* \approx -nD(P||Q) - \sqrt{n\sigma}\Phi^{-1}(\epsilon) - \frac{1}{2}\log n,$$
 (8)

and in the 'error exponents regime',

Hoeffding:
$$\log e_{1,n}^* \approx -nD(\delta)$$
 (9)

Thms. 3.1-3.2:
$$\log e_{1,n}^* \approx -nD(\delta) - \frac{1}{2(1-\alpha^*)} \log n.$$
 (10)

Table 1 shows representative results for different small values of ϵ , when the sample size n = 50. These results clearly demonstrate the utility of the approximation (10) as it provides by far the most accurate estimates of the error probability $e_{1,n}^*(\epsilon)$ for the problem parameters considered.

ϵ	$e_{1,n}^*(\epsilon)$	Stein (7)	Strassen (8)	Hoeffding (9)	Thms. 3.1-3.2: (10)
0.00006	0.098	10^{-6}	84.2	0.804	0.083
0.00018	0.055	10^{-6}	18.8	0.639	0.056
0.00052	0.029	10^{-6}	4.23	0.472	0.035
0.00137	0.014	10^{-6}	0.096	0.324	0.019
0.00336	0.006	10^{-6}	0.022	0.208	0.010
0.00762	0.003	10^{-6}	0.050	0.126	0.005
0.01604	0.001	10^{-6}	0.011	0.071	0.002

Table 1: Comparison between the true value of the optimal error probability $e_{1,n}^*(\epsilon)$ and four different approximations. Clearly the approximation suggested by Theorems 3.1 and 3.2 gives the best results in this regime.

We close this introduction with a brief discussion of some related work of a different nature. An important special case of binary hypothesis testing is the fully symmetric regime, where the natural question is: If both error probabilities are required to be no greater than some $\epsilon > 0$, how big does the sample size n need to be in order for a test to exist with this property? It turns out [5, 3, 6], that the smallest such sample size, $n^*(P, Q, \epsilon)$, is,

$$n^*(P,Q,\epsilon) = \Theta\Big(\frac{\log(1/\epsilon)}{H_2^2(P,Q)}\Big)$$

where $H_2(P,Q)$ denotes the Hellinger distance between P and Q. This suggests the rough approximation,

$$\log e_{1,n}^*(\epsilon) \approx -nH_2^2(P,Q). \tag{11}$$

But taking $e_{1,n} = e_{2,n}$, or equivalently $D(\delta) = \delta$ in the error exponents regime, leads to an error exponent C(P,Q) known as the Chernoff information, which, at least for P close to Q, is

approximately equal to $H_2^2(P,Q)$. Therefore, in the symmetric regime, the estimates (9) and (11) are roughly equivalent.

Finally, we mention that extensive discussions of different aspects of binary hypothesis testing can be found in Csiszár and Körner's classic text, [9], Han's book on information spectrum methods [14], and Tan's monograph [27].

2 Preliminaries

Consider a binary hypothesis test between two probability measures P, Q on (A, \mathcal{A}) , absolutely continuous with respect to a σ -finite dominating measure λ . Let p, q denote the corresponding densities, $p = dP/d\lambda$ and $q = dQ/d\lambda$. Although our main object of interest, the optimal error probability $e_1^*(\epsilon)$, cannot be explicitly evaluated in general, the optimal test that achieves $e_1^*(\epsilon)$ is well known and given by the classical Neyman-Pearson region [23]. The following one-shot converse and achievability bounds are simple variations of the Neyman-Pearson lemma. In this form, they were proved in [25]; the are stated here without proof.

Proposition 2.1 (One-shot converse) For any two probability measures P and Q on the same measurable space (A, \mathcal{A}) , for any test $P_{Z|X}$, and for any $\gamma > 0$, we have:

$$e_1 \ge \frac{1}{\gamma} \left[1 - e_2 - \mathbb{P} \left(\log \frac{dP}{dQ}(X) > \log \gamma \right) \right], \quad X \sim P.$$

The following simple achievability result is obtained via deterministic tests.

Proposition 2.2 (One-shot achievability) Let P, Q be two probability measures on the same measurable space (A, A), and let $\epsilon \in (0, 1)$. For any $\gamma > 0$ such that,

$$\mathbb{P}\Big(\log\frac{dP}{dQ}(X) < \log\gamma\Big) \le \epsilon, \quad X \sim P.$$

we have,

$$e_1^*(\epsilon) \le \mathbb{P}\Big(\log \frac{dP}{dQ}(Y) \ge \log \gamma\Big), \quad Y \sim Q.$$

Next, we apply the above one-shot bounds to the hypothesis testing problem with i.i.d. data as described in the Introduction, to derive refined, finite-*n* approximations to $e_{1,n}^*(\epsilon)$ in the Stein regime. Theorems 2.3 and 2.4 are essentially proved in [25]. For the sake of completeness, we include their proofs in the Appendix.

Theorem 2.3 (Stein regime converse) Suppose $P \neq Q$ are two probability measures on the same measurable space (A, A), such that:

$$T = \int \left| \log \frac{dP}{dQ} - D(P ||Q) \right|^3 dP < \infty.$$

Let $\epsilon \in (0, 1)$. Then, for any $\Delta > 0$, we have,

$$\log e_{1,n}^*(\epsilon) \ge -nD(P||Q) - \sqrt{n\sigma}\Phi^{-1}\left(\epsilon + \frac{B+\Delta}{\sqrt{n}}\right) - \frac{1}{2}\log n + \log\Delta,$$

for all $n \ge [(B + \Delta)/(1 - \epsilon)]^2$, where,

$$\sigma^{2} = \int \left[\log \frac{dP}{dQ} - D(P ||Q) \right]^{2} dP \quad and \quad B = \frac{T}{2\sigma^{3}}$$

Theorem 2.4 (Stein regime achievability) Suppose $P \neq Q$ are two probability measures on the same measurable space (A, \mathcal{A}) , and let $\epsilon \in (0, 1)$. Then,

$$\log e_{1,n}^*(\epsilon) \le -nD(P||Q) - \sqrt{n\sigma}\Phi^{-1}\left(\epsilon - \frac{B}{\sqrt{n}}\right) - \frac{1}{2}\log n + \log\left(\frac{1}{\sqrt{2\pi\sigma}} + 2B\right),\tag{12}$$

for all $n > (B/\epsilon)^2$, where σ , B and $T < \infty$ are as in Theorem 2.3.

In the following section we will need some simple results about the error exponent function $D(\delta)$. These are stated in Lemmas 2.5 and 2.6 and Proposition 2.7, and they are proved in the Appendix.

For any two probability measures P, Q on (A, \mathcal{A}) with densities p, q respectively, for all $\alpha \in (0, 1)$ we define:

$$Z(\alpha) = \int p^{\alpha} q^{1-\alpha} \, d\lambda. \tag{13}$$

Lemma 2.5 (Properties of $Z(\alpha)$) (i) If D(P||Q) and D(Q||P) are both finite, then $Z(\alpha)$ has a continuous derivative at each $\alpha \in (0, 1)$, with:

$$\frac{dZ(\alpha)}{d\alpha} = \int p^{\alpha} q^{1-\alpha} \log \frac{p}{q} \, d\lambda. \tag{14}$$

(ii) If, moreover, $\int p |\log(p/q)|^3 d\lambda$ and $\int q |\log(p/q)|^3 d\lambda$ are both finite, then $Z(\alpha)$ is three times differentiable in $\alpha \in (0, 1)$, with:

$$\frac{d^2 Z(\alpha)}{d\alpha^2} = \int p^{\alpha} q^{1-\alpha} \left[\log \frac{p}{q} \right]^2 d\lambda, \tag{15}$$

$$\frac{d^3 Z(\alpha)}{d\alpha^3} = \int p^{\alpha} q^{1-\alpha} \left[\log \frac{p}{q} \right]^3 d\lambda.$$
(16)

Lemma 2.6 (Derivatives of the error exponents) For each $\alpha \in (0,1)$, let Q_{α} denote the probability measure on (A, \mathcal{A}) with density,

$$q_{\alpha} := \frac{p^{\alpha} q^{1-\alpha}}{Z(\alpha)},\tag{17}$$

where $Z(\alpha)$ is defined in (13). Under the assumptions of Lemma 2.5 (ii), both $D(Q_{\alpha}||P)$ and $D(Q_{\alpha}||Q)$ are twice differentiable in $\alpha \in (0,1)$ with:

$$\frac{dD(Q_{\alpha}||P)}{d\alpha} = -(1-\alpha)\operatorname{Var}\Big(\log\frac{dP}{dQ}(X^{(\alpha)})\Big),\tag{18}$$

$$\frac{dD(Q_{\alpha}||Q)}{d\alpha} = \alpha \operatorname{Var}\Big(\log \frac{dP}{dQ}(X^{(\alpha)})\Big),\tag{19}$$

$$\frac{d^2 D(Q_{\alpha} \| P)}{d\alpha^2} = \operatorname{Var}\left(\log \frac{dP}{dQ}(X^{(\alpha)})\right) - (1 - \alpha) \mathbb{E}\left[\left(\log \frac{dP}{dQ}(X^{(\alpha)}) - \mathbb{E}\left(\log \frac{dP}{dQ}(X^{(\alpha)})\right)\right)^3\right],\tag{20}$$

$$\frac{d^2 D(Q_{\alpha} \| Q)}{d\alpha^2} = \operatorname{Var}\left(\log \frac{dP}{dQ}(X^{(\alpha)})\right) + \alpha \mathbb{E}\left[\left(\log \frac{dP}{dQ}(X^{(\alpha)}) - \mathbb{E}\left(\log \frac{dP}{dQ}(X^{(\alpha)})\right)\right)^3\right], \quad (21)$$

where $X^{(\alpha)} \sim Q_{\alpha}$.

Recall the definition of the optimal error exponent $D(\delta)$ in (2). The following representation of $D(\delta)$ is well-known for finite spaces A; see, e.g., [10] or [4].

Proposition 2.7 (Error exponent representation) Let P, Q be two probability measures on (A, \mathcal{A}) satisfying the assumptions of Lemma 2.5 (ii). Then for any $0 < \delta < D(Q||P)$, there exists a unique $\alpha^* \in (0, 1)$ such that, $D(Q^*_{\alpha}||P) = \delta$,

$$D(\delta) := \inf_{P': D(P'||P) \le \delta} D(P'||Q) = D(Q_{\alpha^*}||Q),$$

and,

$$D(\delta) = \left(\frac{\alpha^*}{1-\alpha^*}\right)\delta - \left(\frac{1}{1-\alpha^*}\right)\log Z(\alpha^*),$$

where $Z(\alpha)$ and Q_{α} are defined in (13) and (17), respectively.

Parts of our main arguments later will be based on the classical Berry-Esséen bound [18, 24].

Lemma 2.8 (Berry-Esséen bound) Suppose Z_1, Z_2, \ldots, Z_n are *i.i.d.* random variables with mean $\mu = \mathbb{E}(Z_1)$, variance $\sigma^2 = \operatorname{Var}(Z_1)$, and $\rho = \mathbb{E}(|Z_1 - \mu|^3) < \infty$. Let \overline{Z}_n denote the empirical average, $\overline{Z}_n = \frac{1}{n}(Z_1 + Z_2 + \ldots + Z_n)$. We have:

$$\left| \mathbb{P}\left(\frac{\sqrt{n}(\bar{Z}_n - \mu)}{\sigma} \le x \right) - \Phi(x) \right| \le \frac{\rho}{2\sigma^3 \sqrt{n}}, \quad x \in \mathbb{R}, \ n \ge 1.$$

The following is a simple consequence of the Berry-Esséen bound. It is proved in the Appendix.

Lemma 2.9 With Z_1, Z_2, \ldots, Z_n as in Lemma 2.8, we have:

$$\mathbb{E}\left[e^{-\sum_{i=1}^{n} Z_{i}}\mathbb{I}_{\{\sum_{i=1}^{n} Z_{i} \ge x\}}\right] \le \left(\frac{1}{\sqrt{2\pi}} + \frac{\rho}{\sigma^{2}}\right) \frac{1}{\sqrt{n\sigma}} e^{-x}, \quad x \in \mathbb{R}, \ n \ge 1.$$

3 Finite-sample bounds

We are now ready to state the main results, stated in equation (6) in the Introduction. Recall the definition of the optimal error exponent $D(\delta)$ in (2) and its representation in terms of the family of 'tilted' distributions Q_{α} , $\alpha \in (0, 1)$ in Proposition 2.7.

Theorem 3.1 (Achievability) Let P, Q be two probability measures on the measurable space (A, \mathcal{A}) with densities p, q, respectively, with respect to the σ -finite dominating measure λ . Suppose that $\int p |\log(p/q)|^3 d\lambda$ and $\int q |\log(p/q)|^3 d\lambda$ are both finite. Then, for any $0 < \delta < D(Q||P)$ we have,

$$\log E_1^*(n,\delta) \le -nD(\delta) - \frac{1}{2(1-\alpha^*)}\log n + C,$$

for all $n \ge 1$, where the constant $C = C(\delta, P, Q)$ is given by,

$$C = \log\left(\frac{1}{\sigma^* \alpha^* \sqrt{2\pi}} + \frac{\rho^*}{\sigma^{*3}}\right) + \frac{\alpha^*}{1 - \alpha^*} \log\left(\frac{1}{\sigma^* (1 - \alpha^*)\sqrt{2\pi}} + \frac{\rho^*}{\sigma^{*3}}\right),\tag{22}$$

with α^* being the unique $\alpha \in (0,1)$ that achieves $\delta = D(Q_{\alpha^*} || P)$, and where,

$$\sigma^{*2} = \operatorname{Var}\left(\log \frac{dQ_{\alpha^*}}{dP}(X^{(\alpha^*)})\right),$$

$$\rho^* = \mathbb{E}\left|\log \frac{dQ_{\alpha^*}}{dP}(X^{(\alpha^*)}) - \mathbb{E}\left(\log \frac{dQ_{\alpha^*}}{dP}(X^{(\alpha^*)})\right)\right|^3,$$

with $X^{(\alpha^*)} \sim Q_{\alpha}$.

PROOF. By Proposition 2.7, there exists a unique $\alpha^* \in (0,1)$ such that $D(Q_{\alpha^*} || P) = \delta$. Let $\gamma_n > 0$ be a constant that will be chosen later, and consider the decision region,

$$B_n := \Big\{ \log \frac{p^n}{q^n} < \log \gamma_n \Big\}.$$

In view of the definition of q_{α^*} , and using independence, B_n can also be expressed as,

$$B_{n} = \left\{ \frac{1}{1 - \alpha^{*}} \log \frac{p^{n}}{q_{\alpha^{*}}^{n}} < \frac{n \log Z_{\alpha^{*}}}{1 - \alpha^{*}} + \log \gamma_{n} \right\}$$
$$= \left\{ x_{1}^{n} \in A^{n} : \sum_{i=1}^{n} \log \frac{q_{\alpha^{*}}(x_{i})}{p(x_{i})} > -n \log Z(\alpha^{*}) - (1 - \alpha^{*}) \log \gamma_{n} \right\}.$$
(23)

Equivalently, interchanging the roles of P and Q, we get,

$$B_n = \left\{ x_1^n \in A^n : \sum_{i=1}^n \log \frac{q_{\alpha^*}(x_i)}{q(x_i)} < -n \log Z(\alpha^*) + \alpha^* \log \gamma_n \right\}.$$
 (24)

Now define,

$$\tau_1^2 = \operatorname{Var}\left(\log \frac{dQ_{\alpha^*}}{dP}(X^{(\alpha^*)})\right) \quad \text{and} \quad r_1 = \mathbb{E}\left|\log \frac{dQ_{\alpha^*}}{dP}(X^{(\alpha^*)}) - \mathbb{E}\left(\log \frac{dQ_{\alpha^*}}{dP}(X^{(\alpha^*)})\right)\right|^3.$$

First, we claim that, for the choice of γ_n given by,

$$\log \gamma_n := n(D(\delta) - \delta) + \frac{1}{2(1 - \alpha^*)} \log n - \frac{1}{1 - \alpha^*} \log \left[\left(\frac{1}{\tau_1 \sqrt{2\pi}} + \frac{r_1}{\tau_1^3} \right) \right],$$

we have,

$$\mathbb{P}\Big(\log\frac{dP^n}{dQ^n}(X_1^n) < \log\gamma_n\Big) \le e^{-n\delta},\tag{25}$$

where $X_1^n \sim P^n$. To show that, let $(X_1^{(\alpha^*)}, X_2^{(\alpha^*)}, \dots, X_n^{(\alpha^*)}) \sim Q_{\alpha^*}^n$, and write,

$$\mathbb{P}\left(\log\frac{dP^n}{dQ^n}(X_1^n) < \log\gamma_n\right) = \int_{A^n} \mathbb{I}_{B_n} dP^n$$

$$= \int_{A^n} \exp\left\{\sum_{i=1}^n \log\frac{p(x_i)}{q_{\alpha^*}(x_i)}\right\} \mathbb{I}_{B_n} dQ_{\alpha^*}^n$$

$$= \mathbb{E}\left[\exp\left\{-\sum_{i=1}^n \log\frac{q_{\alpha^*}(X_i^{(\alpha^*)})}{p(X_i^{(\alpha^*)})}\right\} \times \mathbb{I}\left\{\sum_{i=1}^n \log\frac{q_{\alpha^*}(X_i^{(\alpha^*)})}{p(X_i^{(\alpha^*)})} > -n\log Z(\alpha^*) - (1-\alpha^*)\log\gamma_n\right\}\right],$$

where we used the representation of B_n in (23). Note that the terms in the above sums are i.i.d. with variance and absolute third centred moment given by τ_1^2 and r_1 , respectively. Applying Lemma 2.9, then, yields,

$$\mathbb{P}\Big(\log\frac{dP^n}{dQ^n}(X_1^n) < \log\gamma_n\Big) \le \frac{Z(\alpha^*)^n \gamma_n^{1-\alpha^*}}{\sqrt{n}} \Big(\frac{1}{\tau_1\sqrt{2\pi}} + \frac{r_1}{\tau_1^3}\Big).$$

And using the definition of γ_n and Proposition 2.7, we obtain,

$$\mathbb{P}\Big(\log\frac{dP^n}{dQ^n}(X_1^n) < \log\gamma_n\Big) \le \exp\Big(-n(1-\alpha^*)[D(\delta)-\delta] + n(1-\alpha^*)D(\delta) - n\alpha^*\delta)\Big) = e^{-n\delta},$$

which is exactly (25).

Given that γ_n satisfies (25), we can now apply the one-shot achievability result of Proposition 2.2 with P^n, Q^n in place of P, Q, to obtain that,

$$E_1^*(n,\delta) \le \mathbb{P}\Big(\log \frac{dP^n}{dQ^n}(Y_1^n) \ge \log \gamma_n\Big) = \int \mathbb{I}_{B_n^c} dQ^n$$

with $Y_1^n \sim Q^n$, and using the expression for B_n in (24),

$$E_{1}^{*}(n,\delta) \leq \int_{A^{n}} \exp\left\{\sum_{i=1}^{n} \log \frac{q(x_{i})}{q_{\alpha^{*}}(x_{i})}\right\} \mathbb{I}_{B_{n}^{c}}(x_{1}^{n}) dQ_{\alpha^{*}}^{n}(x_{1}^{n}) = \mathbb{E}\left[\exp\left\{-\sum_{i=1}^{n} \log \frac{q_{\alpha^{*}}(X_{i}^{(\alpha^{*})})}{q(X_{i}^{(\alpha^{*})})}\right\} \mathbb{I}\left\{\sum_{i=1}^{n} \log \frac{q_{\alpha^{*}}(X_{i}^{(\alpha^{*})})}{q(X_{i}^{(\alpha^{*})})} \geq -n \log Z_{\alpha^{*}} + \alpha^{*} \log \gamma_{n}\right\}\right].$$

Note that the sums above consist of i.i.d. terms with variance and absolute third centred moment given, respectively, by,

$$\tau_2^2 = \operatorname{Var}\left(\log \frac{q_{\alpha^*}(X^{(\alpha^*)})}{q(X^{(\alpha^*)})}\right) \quad \text{and} \quad r_2 = \mathbb{E}\left|\log \frac{q_{\alpha^*}(X^{(\alpha^*)})}{q(X^{(\alpha^*)})} - \mathbb{E}\left(\log \frac{q_{\alpha^*}(X^{(\alpha^*)})}{q(X^{(\alpha^*)})}\right)\right|^3.$$

Then applying Lemma 2.9, we can further bound,

$$E_1^*(n,\delta) \le \frac{Z(\alpha^*)^n}{\gamma_n^{\alpha^*}\sqrt{n}} \Big(\frac{1}{\tau_2\sqrt{2\pi}} + \frac{r_2}{\tau_2^3}\Big),$$

and using the definition of γ_n and Proposition 2.7,

$$\log E_{1}^{*}(n,\delta) \leq n \log Z(\alpha^{*}) - \alpha^{*} \log \gamma_{n} - \frac{1}{2} \log n + \log \left(\frac{1}{\tau_{2}\sqrt{2\pi}} + \frac{r_{2}}{\tau_{2}^{3}}\right)$$
$$= -nD(\delta) - \frac{1}{2(1-\alpha^{*})} \log n$$
$$+ \log \left(\frac{1}{\tau_{2}\sqrt{2\pi}} + \frac{r_{2}}{\tau_{2}^{3}}\right) + \frac{\alpha^{*}}{1-\alpha^{*}} \log \left(\frac{1}{\tau_{1}\sqrt{2\pi}} + \frac{r_{1}}{\tau_{1}^{3}}\right).$$
(26)

Finally, we show that the sum of the two constants in (26) is exactly equal to C in (22). Using the definition of Q_{α^*} , direct computation gives,

$$\tau_1^2 = \operatorname{Var}\left(\log\frac{q_{\alpha^*}(X^{(\alpha^*)})}{p(X^{(\alpha^*)})}\right) = \operatorname{Var}\left(-(1-\alpha^*)\log\frac{p(X^{(\alpha^*)})}{q(X^{(\alpha^*)})} - \log Z_{\alpha^*}\right)$$

so that,

$$\tau_1^2 = (1 - \alpha^*)^2 \operatorname{Var}\left(\log \frac{p(X^{(\alpha^*)})}{q(X^{(\alpha^*)})}\right) = (1 - \alpha^*)^2 \sigma^{*2},\tag{27}$$

and similarly, we obtain,

$$\tau_2^2 = \alpha^{*2} \sigma^{*2}.$$
 (28)

For the third moments r_1 and r_2 , analogous computations give,

$$r_{1} = \mathbb{E} \left| \log \frac{q_{\alpha^{*}}(X^{(\alpha^{*})})}{p(X^{(\alpha^{*})})} - \mathbb{E} \left(\log \frac{q_{\alpha^{*}}(X^{(\alpha^{*})})}{p(X^{(\alpha^{*})})} \right) \right|^{3}$$

= $(1 - \alpha^{*})^{3} \mathbb{E} \left| \log \frac{p(X^{(\alpha^{*})})}{q(X^{(\alpha^{*})})} - \mathbb{E} \left(\log \frac{p(X^{(\alpha^{*})})}{q(X^{(\alpha^{*})})} \right) \right|^{3}$
= $(1 - \alpha^{*})^{3} \rho^{*},$ (29)

and similarly,

$$r_2 = \alpha^{*3} \rho^*. \tag{30}$$

Substituting (27), (28), (29), and (30) into (26), completes the proof.

Theorem 3.2 (Converse) Under the same assumptions and in the notation of Theorem 3.1, suppose that,

$$\sigma_0^2 := \inf_{\alpha \in (0,1)} \operatorname{Var}\Big(\log \frac{p(X^{(\alpha)})}{q(X^{(\alpha)})}\Big) > 0,$$

and,

$$\rho_0 := \sup_{\alpha \in (0,1)} \mathbb{E} \left| \log \frac{p(X^{(\alpha)})}{q(X^{(\alpha)})} - \mathbb{E} \left(\log \frac{p(X^{(\alpha)})}{q(X^{(\alpha)})} \right) \right|^3 < \infty,$$

where $X^{(\alpha)} \sim Q_{\alpha}$ with Q_{α} as in Proposition 2.7. Then,

$$\log E_1^*(n,\delta) \ge -nD(\delta) - \frac{1}{2(1-\alpha^*)} \log n + C',$$
(31)

where the constant $C' = C'(\delta, P, Q)$ is given by,

$$C' = \frac{\log 2}{(1 - \alpha^*)} - \frac{(\sigma^{*2} + 2\rho_0)(2 - \alpha^*)}{2(1 - \alpha^*)} - \frac{|\sigma_0^2 - \rho_0|}{2} + m,$$

where,

$$m = -2\sqrt{2\pi}(1 - \alpha^*)(\rho_0/\sigma_0^3 + 1)\sqrt{\sigma^{*2} + \rho_0},$$

and the bound (31) holds for all,

$$n > \max\left\{7(\rho_0/\sigma_0^3 + 1)^2, \frac{(\sigma^{*2} + 2\rho_0 - 2m + 2\log 2)^2}{(1 - \alpha^*)^2 \sigma^{*4}}, n_0\right\},\$$

where n_0 is the smallest n such that $\log n \leq (1 - \alpha^*) \sigma^{*2} \sqrt{n}$.

PROOF. With A^n, P^n, Q^n in place of A, P, Q, respectively, and with $X_1^n \sim P^n$, the result will eventually follow by an application of Proposition 2.1 for a particular choice of γ_n :

$$E_1^*(n,\delta) \ge \frac{1}{\gamma_n} \left[\mathbb{P} \left(\log \frac{dP^n}{dQ^n} (X_1^n) \le \log \gamma_n \right) - e^{-n\delta} \right].$$
(32)

Here we choose γ_n via,

$$\log \gamma_n = n(D(\delta_n) - \delta_n),$$

where $\delta_n = D(Q_{\alpha_n} || P)$, and α_n is given defined via the relation,

$$n(\alpha_n - \alpha^*)(1 - \alpha^*)\sigma^2 = \frac{1}{2}\log n + \frac{1}{2}(\sigma^{*2} + 2\rho_0) - q + \log 2,$$
(33)

so that,

$$\alpha_n = \alpha^* + \frac{1}{2(1-\alpha^*)\sigma^{*2}} \times \frac{\log n}{n} + \frac{(\sigma^{*2} + 2\rho_0)/2 - m + \log 2}{(1-\alpha^*)\sigma^{*2}} \times \frac{1}{n},$$

with m, ρ_0 and σ_0^2 as in the statement of theorem. The reason for this choice is that, as we show next, this α_n leads to a γ_n such that,

$$\mathbb{P}\Big(\log\frac{dP^n}{dQ^n}(X_1^n) \le \log\gamma_n\Big) \ge 2e^{-n\delta}.$$
(34)

0

Establishing (34) will occupy most of this proof.

Let,

$$B_n := \Big\{ x_1^n \in A^n : \log \frac{p^n(x_1^n)}{q^n(x_1^n)} \le \log \gamma_n \Big\},\$$

and note that, as in the earlier proof of Theorem 3.1, B_n can also be expressed as,

$$B_n = \left\{ x_1^n \in A^n : \log \frac{p^n(x_1^n)}{q_{\alpha_n}(x_1^n)} \le n \log Z(\alpha_n) + (1 - \alpha_n) \log \gamma_n \right\}$$
$$= \left\{ x_1^n \in A^n : \log \frac{q_{\alpha_n}^n(x_1^n)}{p^n(x_1^n)} \ge n \mathbb{E} \left(\log \frac{q_{\alpha_n}(X^{(\alpha_n)})}{p(X^{(\alpha_n)})} \right) \right\}.$$
(35)

Using the Berry-Esséen bound in Lemma 2.8, we can bound above the probability of B_n^c under $Q_{\alpha_n}^n$ as,

$$Q_{\alpha_n}^n(B_n^c) = \mathbb{P}\left(\sum_{i=1}^n \log \frac{q_{\alpha_n}(X_i^{(\alpha_n)})}{p(X_i^{(\alpha_n)})} < n\mathbb{E}\left(\log \frac{q_{\alpha_n}(X^{(\alpha_n)})}{p(X^{(\alpha_n)})}\right)\right) \le \Phi(0) + \frac{r_n}{2\tau_n^3\sqrt{n}}$$

where $(X_1^{(\alpha_n)}, \ldots, X_n^{(\alpha_n)}) \sim Q_{\alpha_n}^n$,

$$\tau_n^2 = \operatorname{Var}\Big(\log\frac{q_{\alpha_n}(X^{(\alpha_n)})}{p(X^{(\alpha_n)})}\Big),$$

and,

$$r_n = \mathbb{E} \left| \log \frac{q_{\alpha_n}(X^{(\alpha_n)})}{p(X^{(\alpha_n)})} - \mathbb{E} \left(\log \frac{q_{\alpha_n}(X^{(\alpha_n)})}{p(X^{(\alpha_n)})} \right) \right|^3.$$

Using the exact same computations that led to relations (27) and (29) in the previous proof, here we have that $\tau_n = (1 - \alpha_n)\sigma_n$ and $r_n = (1 - \alpha_n)^3\rho_n$ where σ_n^2 and ρ_n are the variance and the absolute third centred moment of $\log[p(X^{(\alpha_n)})/q(X^{(\alpha_n)})]$. Therefore,

$$Q_{\alpha_n}^n(B_n^c) \le \frac{1}{2} + \frac{\rho_0}{2\sigma_0^3 \sqrt{n}}.$$
(36)

Now, with P in place of Q and Q_{α_n} in place of P, and in view of (35), Proposition 2.2 implies that the probability in (34) can be bounded below as,

$$\mathbb{P}\Big(\log\frac{p^n(X_1^n)}{q^n(X_1^n)} \le \log\gamma_n\Big) \ge \tilde{e}_{1,n}\big(Q_{\alpha_n}^n(B_n^c)\big),$$

where $\tilde{e}_{1,n}(\epsilon)$ is the same as the minimal error probability $e_{1,n}^*(\epsilon)$, but with P in place of Q and Q_{α_n} in place of P. Moreover, by (36),

$$\mathbb{P}\Big(\log\frac{p^n(X_1^n)}{q^n(X_1^n)} \le \log\gamma_n\Big) \ge \tilde{e}_{1,n}\Big(\frac{1}{2} + \frac{\rho_0}{2\sigma_0^3\sqrt{n}}\Big).$$

and we can use the Stein regime converse, Theorem 2.3, with $\Delta = 1$, to obtain that,

$$\log \mathbb{P} \left(\log \frac{dP^{n}}{dQ^{n}} (X_{1}^{n}) \leq \log \gamma_{n} \right)$$

$$\geq -nD(Q_{\alpha_{n}} \| P) - \sqrt{n\tau_{n}} \Phi^{-1} \left(\frac{1}{2} + \frac{\rho_{0}}{2\sigma_{0}^{3}\sqrt{n}} + \frac{r_{n}}{2\tau_{n}^{3}\sqrt{n}} + \frac{1}{\sqrt{n}} \right) - \frac{1}{2} \log n$$

$$\geq -nD(Q_{\alpha_{n}} \| P) - \sqrt{n\tau_{n}} \Phi^{-1} \left(\frac{1}{2} + \frac{\rho_{0}}{\sigma_{0}^{3}\sqrt{n}} + \frac{1}{\sqrt{n}} \right) - \frac{1}{2} \log n,$$

for all $n > 4(\rho_0/\sigma_0^3 + 1)^2$. Using the fact that $\alpha_n \ge \alpha^*$ by definition, and substituting a simple two-term Taylor expansions for Φ^{-1} , gives,

$$\log \mathbb{P}\left(\log \frac{dP^{n}}{dQ^{n}}(X_{1}^{n}) \leq \log \gamma_{n}\right) \geq -nD(Q_{\alpha_{n}}\|P) - \sqrt{n\tau_{n}}\Phi^{-1}\left(\frac{1}{2}\right) - \frac{1}{2}\log n - \frac{(\rho_{0}/\sigma_{0}^{3}+1)\tau_{n}}{\phi(\Phi^{-1}(\frac{1}{2}+\frac{\rho_{0}/\sigma_{0}^{3}+1}{\sqrt{n}}))} \\ \geq -nD(Q_{\alpha_{n}}\|P) - \frac{1}{2}\log n - \frac{(1-\alpha^{*})(\rho_{0}/\sigma_{0}^{3}+1)\sigma_{n}}{\phi(\Phi^{-1}(\frac{1}{2}+\frac{\rho_{0}/\sigma_{0}^{3}+1}{\sqrt{n}}))}.$$
(37)

There are two more terms in (37) that will be bounded using a second order Taylor expansion. For σ_n , using (20) from Lemma 2.6 we have,

$$\sigma_n^2 = \operatorname{Var}\left(\log \frac{dP}{dQ}(X^{(\alpha_n)})\right)$$
$$= \operatorname{Var}\left(\log \frac{dP}{dQ}(X^{(\alpha_n)}) + (\alpha_n - \alpha^*) \mathbb{E}\left[\left(\log \frac{dP}{dQ}(X^{(\psi)}) - \mathbb{E}\left(\log \frac{dP}{dQ}(X^{(\psi)})\right)\right)^3\right],$$

for some $\psi \in [\alpha^*, \alpha_n]$. Since the conditions in Lemma 2.6 are fulfilled, the above derivatives are justified. Thus,

$$\sigma_n^2 \le \operatorname{Var}\left(\log \frac{dP}{dQ}(X^{(\alpha^*)})\right) + \mathbb{E}\left|\log \frac{dP}{dQ}(X^{(\psi)}) - \mathbb{E}\left(\log \frac{dP}{dQ}(X^{(\psi)})\right)\right|^3 \le \sigma^{*2} + \rho_0.$$
(38)

And for $D(Q_{\alpha_n}||P)$, using the first and second derivatives of $D(Q_{\alpha}||P)$ in (18) and (20) of Lemma 2.6, we have,

$$D(Q_{\alpha_n} \| P) = D(Q_{\alpha^*} \| P) - (\alpha_n - \alpha^*)(1 - \alpha^*)\sigma^{*2} + \frac{1}{2}(\alpha_n - \alpha^*)^2 S_n(\psi),$$
(39)

where,

$$S_n(\psi) = \operatorname{Var}\left(\log\frac{dP}{dQ}(X^{(\psi)})\right) - (1-\psi)\mathbb{E}\left[\left(\log\frac{dP}{dQ}(X^{(\psi)}) - \mathbb{E}\left(\log\frac{dP}{dQ}(X^{(\psi)})\right)\right)^3\right],$$

for some (possibly different from above) $\psi \in [\alpha^*, \alpha_n]$. Arguing as in (38), $S(\psi)$ and subsequently $D(Q_{\alpha_n} || P)$ can be bounded above as,

$$D(Q_{\alpha_n} \| P) \le D(Q_{\alpha^*} \| P) - (\alpha_n - \alpha^*)(1 - \alpha^*)\sigma^{*2} + \frac{1}{2}(\alpha_n - \alpha^*)^2(\sigma^{*2} + 2\rho_0).$$
(40)

Substituting the bounds (38) and (40) in (37) yields,

$$\log \mathbb{P}\left(\log \frac{dP^{n}}{dQ^{n}}(X_{1}^{n}) \leq \log \gamma_{n}\right) \geq -nD(Q_{\alpha^{*}} ||P) + n(\alpha_{n} - \alpha^{*})(1 - \alpha^{*})\sigma^{*2}$$
$$-\frac{n}{2}(\alpha_{n} - \alpha^{*})^{2}(\sigma^{*2} + 2\rho_{0}) - \frac{1}{2}\log n$$
$$-\frac{(1 - \alpha^{*})(\rho_{0}/\sigma_{0}^{3} + 1)\sqrt{\sigma^{*2} + \rho_{0}}}{\phi(\Phi^{-1}(\frac{1}{2} + \frac{\rho_{0}/\sigma_{0}^{3} + 1}{\sqrt{n}}))}.$$

If we further take $n \ge (\rho_0/\sigma_0^3 + 1)^2/(\Phi(\sqrt{2\log 2}) - 1/2)^2$, and recalling the definition of m, we can further bound the last term above to obtain,

$$\log \mathbb{P} \Big(\log \frac{p^{n}(X_{1}^{n})}{q^{n}(X_{1}^{n})} \leq \log \gamma_{n} \Big) \\ \geq -n\delta + n(\alpha_{n} - \alpha^{*})(1 - \alpha^{*})\sigma^{*2} - \frac{n}{2}(\alpha_{n} - \alpha^{*})^{2}(\sigma^{*2} + 2\rho_{0}) - \frac{1}{2}\log n + m.$$
(41)

It is actually at this point where the exact form of α_n is justified; recalling its definition, if we take n large enough such that the following three conditions are satisfied,

$$\frac{\log n}{\sqrt{n}} \le (1 - \alpha^*)\sigma^{*2},$$

$$n \ge \frac{[(\sigma^{*2} + 2\rho_0) - 2m + 2\log 2]^2}{(1 - \alpha^*)^2\sigma^{*4}},$$

$$n > \frac{1}{(1 - \alpha^*)^2},$$

then $\alpha_n < 1$ and $n(\alpha_n - \alpha^*)^2 \leq 1$. Using this, the third term in the right-hand side of (41) can be bounded by $(\sigma^{*2} + 2\rho_0)/2$, and substituting the expression for α_n in (33) into (41), yields,

$$\log \mathbb{P}\Big(\log \frac{dP^n}{dQ^n}(X_1^n) \le \log \gamma_n\Big) \ge -n\delta + \log 2,$$

which is exactly (34), as desired.

Finally, we can return to the result of Proposition 2.1 in (32), which, after using the bound (34) and the definition of γ_n becomes,

$$\log E_1^*(n,\delta) \ge -nD(Q_{\alpha_n} \| Q) + nD(Q_{\alpha_n} \| P) - n\delta.$$
(42)

For the first term, a three-term Taylor expansion around α^* , where $D(Q_{\alpha^*} || Q) = D(\delta)$, gives,

$$nD(Q_{\alpha_n} || Q) - nD(\delta) \leq n(\alpha_n - \alpha^*)\alpha^* \sigma^{*2} + \frac{n}{2}(\alpha_n - \alpha^*)^2(\sigma^{*2} + 2\rho_0)$$

$$\leq n(\alpha_n - \alpha^*)\alpha^* \sigma^{*2} + \frac{1}{2}(\sigma^{*2} + 2\rho_0)$$

$$\leq \frac{\alpha^*}{(1 - \alpha^*)} \frac{\log n}{2} + \frac{\alpha^*(\sigma^{*2} + 2\rho_0)/2 - \alpha^* m + \alpha^* \log 2}{(1 - \alpha^*)} + \frac{1}{2}(\sigma^{*2} + 2\rho_0)$$

$$= \frac{\alpha^*}{(1 - \alpha^*)} \frac{\log n}{2} + \frac{\alpha^*(\log 2 - m)}{(1 - \alpha^*)} + \frac{1}{2(1 - \alpha^*)}(\sigma^{*2} + 2\rho_0), \quad (43)$$

where we used again the expression for α_n from (33), and the fact that $n(\alpha_n - \alpha^*)^2 \leq 1$. For the second term, using the second order Taylor expansion from (39), we can similarly bound,

$$nD(Q_{\alpha_n} || P) \ge nD(Q_{\alpha^*} || P) - n(\alpha_n - \alpha^*)(1 - \alpha^*)\sigma^{*2} + \frac{n(\alpha_n - \alpha^*)^2}{2}(\sigma_0^2 - \rho_0)$$

$$\ge n\delta - n(\alpha_n - \alpha^*)(1 - \alpha^*)\sigma^{*2} - \frac{1}{2}|\sigma_0^2 - \rho_0|$$

$$\ge n\delta - \frac{\log n}{2} - \frac{\sigma^{*2} + 2\rho_0}{2} + \log 2 - m - \frac{1}{2}|\sigma_0^2 - \rho_0|.$$
(44)

Substituting the bounds (43) and (44) into (42) yields,

$$\begin{split} \log E_1^*(n,\delta) &\geq -nD(\delta) - \frac{\alpha^*}{(1-\alpha^*)} \frac{\log n}{2} - \frac{\alpha^*}{(1-\alpha^*)} (\log 2 - m) - \frac{(\sigma^{*2} + 2\rho_0)}{2(1-\alpha^*)} \\ &\quad - \frac{\log n}{2} - \frac{(\sigma^{*2} + 2\rho_0)}{2} + \log 2 - m - \frac{|\sigma_0^2 - \rho_0|}{2} \\ &\quad = -nD(\delta) - \frac{\log n}{2(1-\alpha^*)} + \frac{(1-2\alpha^*)(\log 2 - m)}{(1-\alpha^*)} - \frac{(\sigma^{*2} + 2\rho_0)(2-\alpha^*)}{2(1-\alpha^*)} - \frac{|\sigma_0^2 - \rho_0|}{2}, \end{split}$$

and using the definition of m in the statement of the theorem yields exactly the bound claimed in (31) and completes the proof.

Appendix

PROOF OF THEOREM 2.3. Our starting point is the one-shot converse in Proposition 2.1. With A^n, P^n, Q^n in place of A, P, Q, respectively, and with $X_1^n \sim P^n$, we have that, for any $\gamma_n > 0$:

$$e_{1,n}^{*}(\epsilon) \geq \frac{1}{\gamma_n} \left[1 - \epsilon - \mathbb{P} \Big(\log \frac{dP^n}{dQ^n}(X_1^n) > \log \gamma_n \Big) \right], \tag{45}$$

with $X_1^n \sim P^n$. To further bound $e_{1,n}^*(\epsilon)$, we examine the probability in the right-hand side of (45). First,

$$\mathbb{P}\left(\log\frac{dP^{n}}{dQ^{n}}(X_{1}^{n}) > \log\gamma_{n}\right) = \mathbb{P}\left(\sum_{i=1}^{n} \left[\log\frac{dP}{dQ}(X_{i}) - D\right] > \log\gamma_{n} - nD\right) \\
= 1 - \mathbb{P}\left(\frac{1}{\sqrt{n\sigma}}\sum_{i=1}^{n} \left[\log\frac{dP}{dQ}(X_{i}) - D\right] \le \frac{1}{\sqrt{n\sigma}}[\log\gamma_{n} - nD]\right) \\
= 1 - F_{n}\left(\frac{\log\gamma_{n} - nD}{\sqrt{n\sigma}}\right),$$
(46)

where we write F_n for the distribution function of $\frac{1}{\sqrt{n\sigma}} \left(\sum_{i=1}^n \left[\log \frac{dP}{dQ}(X_i) - D \right] \right)$, and D for $D(P \| Q)$. By the Berry-Esséen bound in Lemma 2.8, with $Z_i = \log \left[\frac{dP}{dQ}(X_i) \right]$, we have,

$$\Phi\left(\frac{\log\gamma_n - nD}{\sqrt{n\sigma}}\right) - \frac{T}{2\sigma^3\sqrt{n}} \le F_n\left(\frac{\log\gamma_n - nD}{\sqrt{n\sigma}}\right) \le \Phi\left(\frac{\log\gamma_n - nD}{\sqrt{n\sigma}}\right) + \frac{T}{2\sigma^3\sqrt{n}}, \quad (47)$$

and combining (45) with (46) and with the left-hand inequality in (47),

$$\log e_{1,n}^*(\epsilon) \ge -\log \gamma_n + \log \left(-\epsilon - \frac{B}{\sqrt{n}} + \Phi\left(\frac{\log \gamma_n - nD}{\sqrt{n}\sigma}\right) \right)$$

Finally we will choose an appropriate value of γ_n . Taking $\gamma_n = \exp\{nD + \sqrt{n\sigma}\Phi^{-1}(\psi_n)\}$ for some $\psi_n \in (0, 1/2]$, yields,

$$\log e_{1,n}^*(\epsilon) \ge -nD - \sqrt{n\sigma}\Phi^{-1}(\psi_n) + \log\left(-\epsilon - \frac{B}{\sqrt{n}} + \psi_n\right),$$

and choosing $\psi_n = \epsilon + \frac{B+\Delta}{\sqrt{n}}$ for some $\Delta > 0$,

$$\log e_{1,n}^* \ge -nD - \sqrt{n\sigma}\Phi^{-1}\left(\epsilon + \frac{B+\Delta}{\sqrt{n}}\right) - \frac{1}{2}\log n + \log\Delta.$$

This holds as long as ψ_n can be chosen appropriately, that is, as long as we have $\psi_n \in (0, 1)$ so that the inverse function Φ^{-1} is well defined. We obviously always have $\psi_n > 0$, and for $\psi_n < 1$ we need $n > [(B + \Delta)/(1 - \epsilon)]^2$. This completes the proof of (2.3).

PROOF OF THEOREM 2.4. Our starting point now is the one-shot achievability result in Proposition 2.2. With A^n, P^n, Q^n in place of A, P, Q, respectively, let $X_1^n \sim P^n, Y_1^n \sim Q^n$, and let $\gamma_n > 0$ be such that,

$$\mathbb{P}\Big(\log\frac{dP^n}{dQ^n}(X_1^n) < \gamma_n\Big) \le \epsilon.$$
(48)

Then Proposition 2.2 states that,

$$e_{1,n}^*(\epsilon) \le \mathbb{P}\Big(\log \frac{dP^n}{dQ^n}(Y_1^n) \ge \gamma_n\Big).$$

Writing $B_n \subset A^n$ for the decision region,

$$B_n := \Big\{ \log \frac{dP^n}{dQ^n} \ge \log \gamma_n \Big\} = \Big\{ x_1^n \in A^n : \sum_{i=1}^n \log \frac{dP}{dQ}(x_i) \ge \log \gamma_n \Big\},$$

we have,

$$e_{1,n}^*(\epsilon) \leq \int \mathbb{I}_{B_n} dQ^n$$

= $\int \exp\left\{-\sum_{i=1}^n \log \frac{dP}{dQ}(x_i)\right\} \mathbb{I}_{B_n}(x_1^n) dP^n(x_1^n)$
= $\mathbb{E}\left[\exp\left\{-\sum_{i=1}^n \log \frac{dP}{dQ}(X_i)\right\} \mathbb{I}_{B_n}(X_1^n)\right],$

and by Lemma 2.9,

$$e_{1,n}^*(\epsilon) \le \frac{1}{\gamma_n \sqrt{n}} \Big(\frac{1}{\sigma \sqrt{2\pi}} + 2B \Big). \tag{49}$$

Now, we choose a specific γ_n and verify that (48) is satisfied. As in the proof of Theorem 2.3, write D = D(P||Q) and let $\gamma_n = nD + \sqrt{n\sigma}\Phi^{-1}(\psi_n)$, but this time with $\psi_n = \epsilon - B/\sqrt{n}$. From (47) in the proof of Theorem 2.3 we have that,

$$\mathbb{P}\Big(\log\frac{dP^n}{dQ^n}(X_1^n) < \gamma_n\Big) \le F_n\Big(\frac{\log\gamma_n - nD}{\sqrt{n\sigma}}\Big) \le \Phi\Big(\frac{\log\gamma_n - nD}{\sqrt{n\sigma}}\Big) + \frac{B}{\sqrt{n}} = \epsilon$$

so that (48) is satisfied, as long as $\psi_n \in (0, 1)$, i.e., as long as $n > (B/\epsilon)^2$. And substituting the value of γ_n in (49) and taking logarithms, yields exactly (2.4).

PROOF OF LEMMA 2.5. Formally, all three results of the lemma are straightforward computations; the only thing we need to justify is the interchange of the order of differentiation and integration.

Write $h = \log \frac{p}{q}$. Since D(P||Q) and D(Q||P) are both finite, we have that $\int |h| dP$ and $\int |h| dQ$ are both finite. Moreover, we have the simple inequality $0 \le p^{\alpha}q^{1-\alpha} \le p+q$ for all $\alpha \in (0,1)$. Therefore, the expression for the first derivative in the lemma is finite for all $\alpha \in (0,1)$. Also, the two measures P, Q are mutually absolutely continuous, so we can restrict the integral in the definition of $Z(\alpha)$ to $S := \{p > 0\} \cap \{q > 0\}$.

Now let $\alpha \in (0, 1)$. For all ϵ with $0 < |\epsilon| < \min\{\alpha, 1 - \alpha\}$, we have,

$$\frac{Z(\alpha+\epsilon)-Z(\alpha)}{\epsilon} = \int_{S} \frac{1}{\epsilon} \left[p^{\alpha+\epsilon} q^{1-\alpha-\epsilon} - p^{\alpha} q^{1-\alpha} \right] d\lambda = \int_{S} q e^{\alpha h} \left[\frac{e^{\epsilon h} - 1}{\epsilon} \right] d\lambda.$$

By a first-order Taylor expansion, the integrand above is equal to,

$$p^{\alpha+\xi}q^{1-\alpha-\xi}h,$$

for some $\xi = \xi(\epsilon)$ with $|\xi| \le \epsilon$, which is dominated by the λ -integrable function (p+q)|h|, so the desired differentiation under the integral is justified by dominated convergence.

To see that the derivative is continuous in α , take $\alpha \in (0,1)$ and $0 < |\epsilon| < \min\{\alpha, 1 - \alpha\}$. Then,

$$Z'(\alpha + \epsilon) = \int_{S} p^{\alpha + \epsilon} q^{1 - \alpha - \epsilon} h \, d\lambda,$$

where the integrand converges pointwise to $p^{\alpha}q^{1-\alpha}h$ as $\epsilon \to 0$, and it is absolutely dominated by (p+q)|h|, which is λ -integrable. Therefore, by dominated convergence, $Z'(\alpha + \epsilon) \to Z'(\alpha)$ as $\epsilon \to 0$, completing the proof of (i).

For (*ii*) we similarly note that since $\int ph^2 d\lambda$ and $\int qh^2 d\lambda$ are both finite, and since for all α we have $0 \leq p^{\alpha}q^{1-\alpha} \leq p+q$, the expression for the second derivative in the lemma is finite. Then for $\alpha \in (0,1)$ and $0 < |\epsilon| < \min\{\alpha, 1-\alpha\}$,

$$\frac{Z'(\alpha+\epsilon)-Z'(\alpha)}{\epsilon} = \int_{S} \frac{1}{\epsilon} \left[p^{\alpha+\epsilon} q^{1-\alpha-\epsilon} - p^{\alpha} q^{1-\alpha} \right] h \, d\lambda = \int_{S} q e^{\alpha h} \left[\frac{e^{\epsilon h} - 1}{\epsilon} \right] h \, d\lambda$$

where the integrand can be expressed as $p^{\alpha+\xi'}q^{1-\alpha-\xi'}h^2$, for some $\xi' = \xi'(\epsilon)$ with $|\xi'| \leq \epsilon$. Since this is dominated by the λ -integrable function $(p+q)h^2$, the exchange of differentiation and integration is again justified by dominated convergence, justifying differentiating twice under the integral sign.

The proof for the third derivative is perfectly analogous to the two cases above.

PROOF OF LEMMA 2.6. Starting with the definition of the density q_{α} , we have,

$$D(Q_{\alpha}||P) = \int q_{\alpha} \log\left(\frac{p^{\alpha}q^{1-\alpha}}{Z(\alpha)p}\right) d\lambda$$
$$= -\log Z(\alpha) - (1-\alpha) \int q_{\alpha} \log\frac{p}{q} d\lambda$$
$$= -\log Z(\alpha) - \frac{1-\alpha}{Z(\alpha)} \int p^{\alpha}q^{1-\alpha} \log\frac{p}{q} d\lambda$$

By Lemma 2.5, the last expression above is differentiable in $\alpha \in (0, 1)$, and can be expressed as,

$$D(Q_{\alpha}||P) = -\log Z(\alpha) - \frac{(1-\alpha)Z'(\alpha)}{Z(\alpha)}.$$

Therefore,

$$\frac{d}{d\alpha}D(Q_{\alpha}||P) = -(1-\alpha)\frac{Z(\alpha)Z''(\alpha) - Z'(\alpha)^2}{Z(\alpha)^2}$$
$$= -(1-\alpha)\left[\int q_{\alpha}h^2 d\lambda - \left(\int q_{\alpha}h d\lambda\right)^2\right]$$

where $h = \log \frac{p}{q}$. This proves (18). Again by Lemma 2.5 we have that this is differentiable, so that,

$$\frac{d^2}{d\alpha^2} D(Q_{\alpha} \| P) = \left[\frac{Z''(\alpha)}{Z(\alpha)} - \left(\frac{Z'(\alpha)}{Z'(\alpha)} \right) \right] - (1 - \alpha) \left[\frac{Z'''(\alpha)}{Z(\alpha)} - 3 \frac{Z''(\alpha)Z'(\alpha)}{Z(\alpha)^2} + 2 \left(\frac{Z'(\alpha)}{Z(\alpha)} \right)^3 \right]$$
$$= \left[\int q_{\alpha} h^2 d\lambda - \left(\int q_{\alpha} h d\lambda \right)^2 \right] - (1 - \alpha) \left[\int q_{\alpha} h^3 d\lambda - 3 \int q_{\alpha} h^2 d\lambda \int q_{\alpha} h d\lambda + 2 \left(\int q_{\alpha} h d\lambda \right)^3 \right],$$

where the second term in the expression is the centered third moment of $\log \frac{p}{q}$ under Q_{α} . This proves (20). The proofs of (19) and (21) are identical.

PROOF OF PROPOSITION 2.7. Choose and fix a $0 < \delta < D(Q||P)$. Let $g(\alpha) = D(Q_{\alpha}||P)$, $\alpha \in [0,1]$. Since g(0) = D(Q|P) and g(1) = 0, to prove the existence of an $\alpha^* \in (0,1)$ with $g(\alpha^*) = \delta$ it suffices to show that $g(\alpha)$ is continuous on [0,1].

Let $h = \log(p/q)$. First we observe that $Z(\alpha)$ is bounded above and below away from zero: As noted earlier, we have $p^{\alpha}q^{1-\alpha} \leq p+q$ for all $\alpha \in [0,1]$, and hence $Z(\alpha) \leq 2$, $\alpha \in [0,1]$. Also, since D(P||Q) and D(Q||P) are both finite, P, Q are mutually absolutely continuous, and we must have that $m := Q(p \geq q) > 0$. Then, for $\alpha \in (0,1)$:

$$Z(\alpha) = \int p^{\alpha} q^{1-\alpha} \, d\lambda \ge \int_{\{p \ge q\}} p^{\alpha} q^{1-\alpha} \, d\lambda \ge \int_{\{p \ge q\}} q \, d\lambda = m > 0.$$

Recalling Lemma 2.6, we have,

$$g(\alpha) = \int q_{\alpha} \log \frac{q_{\alpha}}{p} d\lambda$$

= $\int q_{\alpha} \Big[\log \frac{1}{Z(\alpha)} + (1 - \alpha) \log \frac{q}{p} \Big] d\lambda$
= $\log \frac{1}{Z(\alpha)} - (1 - \alpha) \frac{Z'(\alpha)}{Z(\alpha)}.$ (50)

so $g(\alpha)$ is continuous for $\alpha \in (0,1)$. For all $\alpha \in [0,1]$ the last integrand above is absolutely bounded by $|p^{\alpha}q^{1-\alpha}h| \leq p|h| + q|h|$, where the first absolute moments $\int p|h| d\lambda$ and $\int q|h| d\lambda$ are both finite. Hence, by dominated convergence, the integral converges to $\int qh d\lambda = -D(Q||P)$ as $\alpha \downarrow 0$. Moreover, dominated convergence implies that $Z(\alpha) \to 1$ and hence $g(\alpha) \to g(0) =$ D(Q||P) as $\alpha \downarrow 0$. Similarly, we have that $\int p^{\alpha}q^{1-\alpha}h d\lambda \to D(P||Q) < \infty$ as $\alpha \uparrow 1$, which, combined with the fact that, again by dominated convergence, we have $Z(\alpha) \to 1$ as $\alpha \uparrow 1$, shows that $g(\alpha) \to g(1) = 0$ as $\alpha \uparrow 1$.

Therefore, g is continuous on all of [0,1], and the existence of an $\alpha^* \in (0,1)$ such that $D(Q_{\alpha^*}||P) = \delta$ follows. If moreover $\int p(\log \frac{p}{q})^2 d\lambda$ and $\int q(\log \frac{p}{q})^2 d\lambda$ are both finite, then, by Lemma 2.6 the first derivative of $g(\alpha)$ exists and is strictly negative for all $\alpha \in (0,1)$, hence proving the uniqueness of $\alpha^* \in (0,1)$.

Next, we prove that $D(\delta) = D(Q_{\alpha^*} || Q)$. Of course, by definition, $D(\delta) \leq D(Q_{\alpha^*} || Q)$, so we only need to prove that,

$$D(W||Q) \ge D(Q_{\alpha^*}||Q)$$
, for any W with $D(W||P) \le \delta$.

Indeed, let W be a probability measure on (A, \mathcal{A}) such that $D(W||P) \leq \delta < \infty$. Therefore, $W \ll P$ and

$$(1 - \alpha^*)[D(W \| Q) - D(Q_{\alpha^*} \| Q)] = D(W \| Q_{\alpha^*}) - \alpha^* D(W \| P) + \alpha^* D(Q_{\alpha^*} \| P)$$

$$\geq D(W \| Q_{\alpha^*}) \geq 0,$$

where we used $D(W||P) \leq \delta$ and $\delta = D(Q_{\alpha^*}||P)$.

PROOF OF LEMMA 2.9. Let F_n and G_n , respectively, denote the distribution functions of S_n and $[S_n - n\mu]/(\sigma\sqrt{n})$ where $S_n = \sum_{i=1}^n Z_i$. Using integration by parts and the definition of G_n ,

$$\mathbb{E}\left[e^{-S_n}\mathbb{I}_{\{S_n \ge x\}}\right] = \int_x^\infty e^{-s} dF_n(s)$$
$$= -e^{-x}F_n(x) + \int_x^\infty e^{-s}F_n(s) ds$$
$$= -e^{-x}G_n\left(\frac{x-n\mu}{\sigma\sqrt{n}}\right) + \int_x^\infty e^{-s}G_n\left(\frac{s-n\mu}{\sigma\sqrt{n}}\right) ds$$

and applying the Berry-Esséen bound in Lemma 2.8,

$$\mathbb{E}\left[e^{-S_n}\mathbb{I}_{\{S_n \ge x\}}\right] \le -e^{-x}\left[\Phi\left(\frac{x-n\mu}{\sigma\sqrt{n}}\right) - \frac{\rho}{2\sigma^3\sqrt{n}}\right] + \int_x^\infty e^{-s}\left[\Phi\left(\frac{s-n\mu}{\sigma\sqrt{n}}\right) + \frac{\rho}{2\sigma^3\sqrt{n}}\right]ds$$
$$= \int_x^\infty \Phi\left(\frac{s-n\mu}{\sigma\sqrt{n}}\right)e^{-s}\,ds - \left[\Phi\left(\frac{x-n\mu}{\sigma\sqrt{n}}\right) - \frac{\rho}{\sigma^3\sqrt{n}}\right]e^{-x}.$$

Finally integrating by parts again,

$$\mathbb{E}\left[e^{-S_n}\mathbb{I}_{\{S_n \ge x\}}\right] \le \frac{1}{\sigma\sqrt{n}} \int_x^\infty e^{-s} \phi\left(\frac{s-n\mu}{\sqrt{n}}\right) ds + \frac{\rho}{\sigma^3\sqrt{n}} e^{-x}$$
$$\le \frac{1}{\sigma\sqrt{n}} e^{-x} \frac{1}{\sqrt{2\pi}} + \frac{\rho}{\sigma^3\sqrt{n}} e^{-x}$$
$$= \left(\frac{1}{\sqrt{2\pi}} + \frac{\rho}{\sigma^2}\right) \frac{1}{\sigma\sqrt{n}} e^{-x},$$

where we used the fact that the standard normal density ϕ is bounded above by $1/\sqrt{2\pi}$.

References

- Y. Altuğ and A.B. Wagner. Refinement of the sphere packing bound for symmetric channels. In Proceedings of the 49th Allerton Conference on Communication, Control, and Computing, pages 30–37, Monticello, IL, September 2011.
- [2] Y. Altuğ and A.B. Wagner. Refinement of the sphere-packing bound: Asymmetric channels. *IEEE Trans. Inform. Theory*, 60(3):1592–1614, March 2014.
- [3] Z. Bar-Yossef. The complexity of massive data set computations. PhD thesis, Department of Computer Sciecen, University of California, Berkeley, Berkeley, CA, 2002.
- [4] R.E. Blahut. Hypothesis testing and information theory. *IEEE Trans. Inform. Theory*, 20(4):405–417, July 1974.
- [5] A.A. Borovkov. *Mathematical statistics*. Gordon and Breach Science Publishers, Amsterdam, The Netherlands, 1998.
- [6] C.L. Canonne. A survey on distribution testing: Your data is big. But is it blue? Number 9 in Graduate Surveys. Theory of Computing Library, 2020.
- [7] H. Chernoff. A measure of asymptotic efficiency for tests of a hypothesis based on the sum of observations. Ann. Math. Statist., 23(4):493–507, 1952.
- [8] T.M. Cover and J.A. Thomas. *Elements of information theory*. John Wiley & Sons, New York, NY, second edition, 2012.
- [9] I. Csiszár and J. Körner. Information theory: Coding theorems for discrete memoryless systems. Cambridge University Press, Cambridge, U.K., second edition, 2011.
- [10] I. Csiszár and G. Longo. On the error exponent for source coding and for testing simple statistical hypotheses. *Studia Sci. Math. Hungar.*, 6:181–191, 1971.
- [11] R.L. Dobrushin. Asymptotic bounds of the probability of error for the transmission of messages over a discrete memoryless channel with a symmetric transition probability matrix. *Teor. Veroyatnost. i Primenen*, 7:283–311, 1962.
- [12] W. Feller. An introduction to probability theory and its applications. Vol. II. John Wiley & Sons, New York, NY, second edition, 1971.
- [13] L. Gavalakis and I. Kontoyiannis. Fundamental limits of lossless data compression with side information. *IEEE Trans. Inform. Theory*, 67(5):2680–2692, May 2021.
- [14] T.S. Han. Information-spectrum methods in information theory. Springer, Berlin, 2003.
- [15] W. Hoeffding. Asymptotically optimal tests for multinomial distributions. Ann. Math. Statist., 36(2):369–408, April 1965.
- [16] J.L. Jensen. Saddlepoint approximations. Oxford University Press, Oxford, U.K., 1995.
- [17] I. Kontoyiannis and S. Verdú. Optimal lossless data compression: Non-asymptotics and asymptotics. *IEEE Trans. Inform. Theory*, 60(2):777–795, February 2014.
- [18] V.Yu. Korolev and I.G. Shevtsova. On the upper bound for the absolute constant in the Berry–Esséen inequality. *Theory Probab. Appl.*, 54(4):638–658, 2010.

- [19] S. Kullback. Information theory and statistics. Dover Publications, Mineola, NY, 1997. Reprint of the second (1968) edition.
- [20] V. Lungu and I. Kontoyiannis. The optimal finite-sample error probability in asymmetric binary hypothesis testing. In 2024 IEEE International Symposium on Information Theory (ISIT), Athens, Greece, July 2024.
- [21] B. Nakiboğlu. A simple derivation of the refined SPB for the constant composition codes. In 2019 IEEE International Symposium on Information Theory (ISIT), pages 2659–2663, Paris, France, July 2019.
- [22] B. Nakiboğlu. A simple derivation of the refined sphere packing bound under certain symmetry hypotheses. Turk. J. Math., 44(3):919–948, 2020.
- [23] J. Neyman and E.S. Pearson. On the problem of the most efficient tests of statistical hypotheses. *Philos. Trans. Roy. Soc. London A*, 231(694-706):289–337, 1933.
- [24] V.V. Petrov. Limit theorems of probability theory. The Clarendon Press, Oxford University Press, New York, NY, 1995.
- [25] Y. Polyanskiy, H.V. Poor, and S. Verdú. Channel coding rate in the finite blocklength regime. *IEEE Trans. Inform. Theory*, 56(5):2307–2359, May 2010.
- [26] V. Strassen. Asymptotische Abschätzungen in Shannons Informationstheorie. In Trans. Third Prague Conf. Information Theory, Statist. Decision Functions, Random Processes (Liblice, 1962), pages 689–723. Publ. House Czech. Acad. Sci., Prague, 1964.
- [27] V.Y.F. Tan. Asymptotic estimates in information theory with non-vanishing error probabilities. Foundations and Trends in Communications and Information Theory, 11(1-2):1–184, September 2014.
- [28] G. Vazquez-Vilar, A. Guillén i Fàbregas, T. Koch, and A. Lancho. Saddlepoint approximation of the error probability of binary hypothesis testing. In 2018 IEEE International Symposium on Information Theory (ISIT), pages 2306–2310, Vail, CO, June 2018.