

CONVOLUTION IDENTITIES OF STIRLING NUMBERS

NADIA N. LI AND WENCHANG CHU

ABSTRACT. By means of the generating function method, a linear recurrence relation is explicitly resolved. The solution is expressed in terms of the Stirling numbers of both the first and the second kind. Two remarkable pairs of combinatorial identities are established as applications, that contain some well-known convolution formulae on Stirling numbers as special cases.

1. INTRODUCTION AND MOTIVATION

Denote by \mathbb{N} the set of natural numbers with $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. For an indeterminate x , define the rising and falling factorials by $(x)_0 = \langle x \rangle_0 \equiv 1$ and

$$\begin{aligned} (x)_n &= x(x+1) \cdots (x+n-1) \quad \text{for } n \in \mathbb{N}, \\ \langle x \rangle_n &= x(x-1) \cdots (x-n+1) \quad \text{for } n \in \mathbb{N}. \end{aligned}$$

Then the unsigned Stirling numbers of the first kind $\left[\begin{smallmatrix} n \\ k \end{smallmatrix} \right]$ are determined as the connection coefficients:

$$\langle x \rangle_n = \sum_{k=0}^n (-1)^{n-k} \left[\begin{smallmatrix} n \\ k \end{smallmatrix} \right] x^k.$$

Analogously, the Stirling numbers of the second kind $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ are given by

$$x^n = \sum_{k=0}^n \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} \langle x \rangle_k$$

which admits the explicit expression

$$\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} = \frac{1}{k!} \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} j^n.$$

These numbers appear frequently in mathematical literatures and have wide applications in combinatorics and number theory (see [2, Chapter 5], [4, §6.2] and [5], just for example).

Recently, Stenlund [6] introduced an interesting bivariate polynomial sequence $\{P_n(x, z)\}_{n \in \mathbb{N}}$ through the following recurrence relation:

$$P_{n+1}(x, z) = x \binom{n+z}{n} - x \sum_{m=1}^n \binom{n-m+z}{n-m+1} P_m(x, z) \quad \text{with } P_1(x, z) = x. \quad (1)$$

2020 *Mathematics Subject Classification.* Primary 11B65, 11B73 Secondary 05A10.

Key words and phrases. Recurrence relation; Convolution formula; Generating function; Stirling number of the first kind; Stirling number of the second kind.

* Corresponding author (N. N. Li): lina20190606@outlook.com.

The same author not only found out an explicit double sum expression

$$P_n(x, z) = \sum_{k=1}^n \sum_{j=1}^k \frac{(-1)^{j-1} (j-1)!}{(n-1)!} \begin{bmatrix} n \\ k \end{bmatrix} \begin{Bmatrix} k \\ j \end{Bmatrix} x^j z^{k-1}, \quad (2)$$

but also explored applications to combinatorial identities and probability theory.

Inspired by the above work of Stenlund [6], we shall examine the extended polynomial sequence $\{Q_n\}_{n \in \mathbb{N}_0}$ defined by the recurrence relation

$$Q_n = \binom{\lambda}{n} (-1)^n - x \sum_{k=1}^n (-1)^k \binom{y}{k} Q_{n-k} \quad \text{with} \quad Q_0 = 1. \quad (3)$$

The first three terms are recorded as follows:

$$\begin{aligned} Q_1 &= xy - \lambda, & Q_2 &= \frac{xy}{2}(1 - y + 2xy) - \frac{\lambda}{2}(1 - \lambda + 2xy), \\ Q_3 &= \frac{xy}{6}(y-1)(y-2) - \frac{\lambda}{6}(\lambda-1)(\lambda-2) - xy(1-y+xy)(\lambda-xy) + \frac{\lambda xy}{2}(\lambda-y). \end{aligned}$$

These Q_n -polynomials in (3) generalize the P_n -polynomials in (1) because it is not hard to verify that Q_n becomes P_{n+1}/x when $\lambda = -1 - z$ and $y \rightarrow -z$.

The rest of the paper will be organized as follows. In the next section, we shall derive the generating function and the explicit formulae for the P_n -polynomials. As applications, two pairs of convolution sums containing both kinds of Stirling numbers will be evaluated in closed forms in Section 3.

2. GENERATING FUNCTION AND EXPLICIT EXPRESSIONS

For the sequence $\{Q_n\}_{n \in \mathbb{N}_0}$ defined by (3), we shall derive, by making use of generating function approach, their generating function and explicit formulae. The informed reader will notice that this treatment is more transparent than the induction approach adopted by Stenlund [6].

Multiplying both sides of (3) by τ^n and then summing over n for $n \geq 0$, we can manipulate the generating function

$$\begin{aligned} Q(\tau) &:= \sum_{n=0}^{\infty} Q_n \tau^n = 1 + \sum_{n=1}^{\infty} Q_n \tau^n \\ &= 1 + \sum_{n=1}^{\infty} \left(\binom{\lambda}{n} (-\tau)^n - x \sum_{n=1}^{\infty} \sum_{k=1}^n (-1)^k \binom{y}{k} Q_{n-k} \right) \tau^n \\ &= (1 - \tau)^\lambda - x \sum_{k=1}^{\infty} (-1)^k \binom{y}{k} \tau^k \sum_{n=k}^{\infty} Q_{n-k} \tau^{n-k}. \end{aligned}$$

This leads us to the functional equation

$$Q(\tau) = (1 - \tau)^\lambda - xQ(\tau)\{(1 - \tau)^y - 1\}$$

and the rational function expression.

Lemma 1 (Generating function).

$$Q(\tau) = \frac{(1 - \tau)^\lambda}{1 - x + x(1 - \tau)^y}.$$

By writing in the geometric series

$$\begin{aligned} Q(\tau) &= \frac{1}{1-x} \times \frac{(1-\tau)^\lambda}{1 - \frac{x}{x-1}(1-\tau)^y} \\ &= \sum_{k=0}^{\infty} \frac{(-x)^k}{(1-x)^{k+1}} (1-\tau)^{\lambda+ky} \end{aligned}$$

and then extracting the coefficient of τ^n , we find the infinite series expression.

Proposition 2 (Single sum formula).

$$Q_n = \sum_{k=0}^{\infty} (-1)^{n+k} \binom{\lambda+ky}{n} \frac{x^k}{(1-x)^{k+1}}.$$

Expanding the rightmost fraction into binomial series, we have further

$$\begin{aligned} Q_n &= \sum_{k=0}^{\infty} (-1)^{n+k} \binom{\lambda+ky}{n} \sum_{j=0}^{\infty} \binom{k+j}{j} x^{k+j} \\ &= \sum_{m=0}^{\infty} (-1)^n x^m \sum_{k=0}^m (-1)^k \binom{m}{k} \binom{\lambda+ky}{n}, \end{aligned}$$

where the last line is justified by the replacement $k+j=m$. Observe that the inner sum with respect to k results substantially in the differences of order m about a polynomial of degree n , which is equal to zero when $m > n$. Therefore, we have derived a finite double sum expression.

Proposition 3 (Double sum formula).

$$Q_n = \sum_{m=0}^n \sum_{k=0}^m (-1)^{n+k} \binom{m}{k} \binom{\lambda+ky}{n} x^m.$$

Expressing the last binomial coefficient in terms of unsigned Stirling numbers and then making use of the binomial theorem, we have

$$\begin{aligned} \binom{\lambda+ky}{n} &= \frac{\langle \lambda+ky \rangle_n}{n!} = \sum_{i=0}^n \frac{(-1)^{n-i}}{n!} \begin{bmatrix} n \\ i \end{bmatrix} (\lambda+ky)^i \\ &= \sum_{i=0}^n \frac{(-1)^{n-i}}{n!} \begin{bmatrix} n \\ i \end{bmatrix} \sum_{j=0}^i \binom{i}{j} \lambda^{i-j} (ky)^j. \end{aligned}$$

By substitutions, Q_n can be written as a four-fold sum

$$\begin{aligned} Q_n &= \sum_{m=0}^n \sum_{k=0}^m \binom{m}{k} x^m \sum_{i=0}^n \frac{(-1)^{k-i}}{n!} \begin{bmatrix} n \\ i \end{bmatrix} \sum_{j=0}^i \binom{i}{j} \lambda^{i-j} (ky)^j \\ &= \sum_{m=0}^n x^m \sum_{i=0}^n \frac{(-1)^{m-i}}{n!} \begin{bmatrix} n \\ i \end{bmatrix} \sum_{j=0}^i \binom{i}{j} \lambda^{i-j} y^j \sum_{k=0}^m (-1)^{m-k} \binom{m}{k} k^j. \end{aligned}$$

Writing the rightmost sum in terms of Stirling number of the second kind, we get a triple sum expression

$$Q_n = \sum_{m=0}^n x^m \sum_{i=0}^n (-1)^{m-i} \frac{m!}{n!} \begin{bmatrix} n \\ i \end{bmatrix} \sum_{j=0}^i \binom{i}{j} \left\{ \begin{matrix} j \\ m \end{matrix} \right\} \lambda^{i-j} y^j. \quad (4)$$

Taking into account that $\left\{ \begin{matrix} j \\ m \end{matrix} \right\} = 0$ for $0 \leq j < m$ and $\binom{i}{j} = 0$ for $0 \leq i < j$, we can restate the last formula as follows.

Proposition 4 (Triple sum formula).

$$Q_n = \sum_{m=0}^n \sum_{i=m}^n \sum_{j=m}^i (-1)^{m-i} \frac{m!}{n!} \begin{bmatrix} n \\ i \end{bmatrix} \binom{i}{j} \left\{ \begin{matrix} j \\ m \end{matrix} \right\} x^m \lambda^{i-j} y^j.$$

By comparing the coefficients of x^m between the two expressions in Proposition 3 and Proposition 4, we derive the following summation formula.

Theorem 5 (Convolution formula).

$$\sum_{i=m}^n \sum_{j=m}^i (-1)^{n+i} \begin{bmatrix} n \\ i \end{bmatrix} \binom{i}{j} \left\{ \begin{matrix} j \\ m \end{matrix} \right\} \lambda^{i-j} y^j = \frac{n!}{m!} \sum_{k=0}^m (-1)^{m+k} \binom{m}{k} \binom{\lambda + ky}{n}.$$

3. APPLICATIONS TO CONVOLUTION FORMULAE

By specifying y to particular values, we shall derive from Theorem 5 two pairs of combinatorial identities, that extend, with an extra parameter “ λ ”, the related known convolution formulae on Stirling numbers.

The first pair of identities are given in the following theorem.

Theorem 6 ($1 \leq m \leq n$).

$$\begin{aligned} \text{(a)} \quad & \sum_{i=m}^n \sum_{j=m}^i (-1)^{n+i} \begin{bmatrix} n \\ i \end{bmatrix} \binom{i}{j} \left\{ \begin{matrix} j \\ m \end{matrix} \right\} \lambda^{i-j} = \frac{n!}{m!} \binom{\lambda}{n-m}, \\ \text{(b)} \quad & \sum_{i=m}^n \sum_{j=m}^i (-1)^{i+j} \begin{bmatrix} n \\ i \end{bmatrix} \binom{i}{j} \left\{ \begin{matrix} j \\ m \end{matrix} \right\} \lambda^{i-j} = \frac{n!}{m!} \binom{\lambda-m}{n-m} (-1)^{m+n}. \end{aligned}$$

They contain the two well-known results below as special cases of $\lambda = 0$.

Corollary 7 ($1 \leq m \leq n$).

$$\begin{aligned} \text{(a)} \quad & \sum_{k=m}^n (-1)^{n-k} \begin{bmatrix} n \\ k \end{bmatrix} \left\{ \begin{matrix} k \\ m \end{matrix} \right\} = \begin{cases} 1, & m = n; \\ 0, & m \neq n; \end{cases} \\ \text{(b)} \quad & \sum_{k=m}^n \begin{bmatrix} n \\ k \end{bmatrix} \left\{ \begin{matrix} k \\ m \end{matrix} \right\} = \frac{n!}{m!} \binom{n-1}{m-1}. \quad \boxed{\text{Lah number}} \end{aligned}$$

Proof of Theorem 6. Letting $y = 1$ in Theorem 5, the corresponding equality becomes

$$\sum_{i=m}^n \sum_{j=m}^i (-1)^{n+i} \begin{bmatrix} n \\ i \end{bmatrix} \binom{i}{j} \left\{ \begin{matrix} j \\ m \end{matrix} \right\} \lambda^{i-j} = \frac{n!}{m!} \sum_{k=0}^m (-1)^{m+k} \binom{m}{k} \binom{\lambda + k}{n}.$$

Then the first formula “(a)” follows by evaluating the above sum on the right

$$\begin{aligned} \sum_{k=0}^m (-1)^{m+k} \binom{m}{k} \binom{\lambda+k}{n} &= \sum_{k=0}^m (-1)^{m+k} \binom{m}{k} \sum_{\ell=0}^n \binom{\lambda}{n-\ell} \binom{k}{\ell} \\ &= \sum_{\ell=0}^n \binom{\lambda}{n-\ell} \binom{m}{\ell} \sum_{k=\ell}^m (-1)^{m+k} \binom{m-\ell}{k-\ell} \\ &= \sum_{\ell=0}^n \binom{\lambda}{n-\ell} \binom{m}{\ell} (1-1)^{m-\ell} = \binom{\lambda}{n-m}, \end{aligned}$$

where the Chu–Vandemonde convolution formula and the binomial theorem have been utilized.

Alternatively for $y = -1$, the corresponding equality in Theorem 5 reads as

$$\sum_{i=m}^n \sum_{j=m}^i (-1)^{i+j} \begin{bmatrix} n \\ i \end{bmatrix} \binom{i}{j} \left\{ \begin{matrix} j \\ m \end{matrix} \right\} \lambda^{i-j} = (-1)^{m-n} \frac{n!}{m!} \sum_{k=0}^m (-1)^k \binom{m}{k} \binom{\lambda-k}{n}.$$

Then we can analogously evaluate the sum on the right as follows:

$$\begin{aligned} \sum_{k=0}^m (-1)^k \binom{m}{k} \binom{\lambda-k}{n} &= \sum_{k=0}^m (-1)^k \binom{m}{k} \sum_{\ell=0}^n \binom{\lambda-m}{n-\ell} \binom{m-k}{\ell} \\ &= \sum_{\ell=0}^n \binom{\lambda-m}{n-\ell} \binom{m}{\ell} \sum_{k=0}^{m-\ell} (-1)^k \binom{m-\ell}{k} \\ &= \sum_{\ell=0}^n \binom{\lambda-m}{n-\ell} \binom{m}{\ell} (1-1)^{m-\ell} = \binom{\lambda-m}{n-m}. \end{aligned}$$

This proves the second formula “(b)” in the theorem. \square

We can further show another pair of identities.

Theorem 8 ($1 \leq m \leq n$).

$$\begin{aligned} \text{(a)} \quad & \sum_{i=m}^n \sum_{j=m}^i (-1)^{n+i} \begin{bmatrix} n \\ i \end{bmatrix} \binom{i}{j} \left\{ \begin{matrix} j \\ m \end{matrix} \right\} \lambda^{i-j} 2^{n-j} \\ &= \frac{n!}{m!} \sum_{\ell=m}^n \frac{m}{\ell} \binom{\lambda}{n-\ell} \binom{-\ell}{\ell-m} 2^{m+n-2\ell}, \\ \text{(b)} \quad & \sum_{i=m}^n \sum_{j=m}^i (-1)^{m+i} \begin{bmatrix} n \\ i \end{bmatrix} \binom{i}{j} \left\{ \begin{matrix} j \\ m \end{matrix} \right\} \lambda^{i-j} 2^{j-m} \\ &= \frac{n!}{m!} \sum_{\ell=m}^n (-1)^{m+n} \binom{\lambda}{n-\ell} \binom{m}{\ell-m} 2^{m-\ell}. \end{aligned}$$

Proof of Theorem 8. For $y = \frac{1}{2}$, the equality in Theorem 5 reduces to

$$\sum_{i=m}^n \sum_{j=m}^i (-1)^{n+i} \begin{bmatrix} n \\ i \end{bmatrix} \binom{i}{j} \left\{ \begin{matrix} j \\ m \end{matrix} \right\} \lambda^{i-j} 2^{n-j} = 2^n \frac{n!}{m!} \sum_{k=0}^m (-1)^{m+k} \binom{m}{k} \binom{\lambda + \frac{k}{2}}{n}.$$

In this case, the first formula “(a)” is confirmed by first reformulating the above sum on the right

$$\begin{aligned} \sum_{k=0}^m (-1)^{m+k} \binom{m}{k} \binom{\lambda + \frac{k}{2}}{n} &= \sum_{k=0}^m (-1)^{m+k} \binom{m}{k} \sum_{\ell=0}^n \binom{\lambda}{n-\ell} \binom{\frac{k}{2}}{\ell} \\ &= \sum_{\ell=0}^n \binom{\lambda}{n-\ell} \sum_{k=0}^m (-1)^{m+k} \binom{m}{k} \binom{\frac{k}{2}}{\ell} \end{aligned}$$

and then evaluating the inner sum by the binomial formula (cf. [3, Equation 3.164])

$$\sum_{k=0}^m (-1)^{m-k} \binom{m}{k} \binom{\frac{k}{2}}{\ell} = \frac{m}{\ell} \binom{-\ell}{\ell-m} 2^{m-2\ell}.$$

When $y = 2$, the corresponding equality in Theorem 5 can be stated as

$$\sum_{i=m}^n \sum_{j=m}^i (-1)^{m+i} \begin{bmatrix} n \\ i \end{bmatrix} \begin{bmatrix} i \\ j \end{bmatrix} \left\{ \begin{matrix} j \\ m \end{matrix} \right\} \lambda^{i-j} 2^{j-m} = 2^{-m} \frac{n!}{m!} \sum_{k=0}^m (-1)^{n+k} \binom{m}{k} \binom{\lambda + 2k}{n}.$$

The above sum on the right can be rewritten as

$$\begin{aligned} \sum_{k=0}^m (-1)^{n+k} \binom{m}{k} \binom{\lambda + 2k}{n} &= \sum_{k=0}^m (-1)^{n+k} \binom{m}{k} \sum_{\ell=0}^n \binom{\lambda}{n-\ell} \binom{2k}{\ell} \\ &= \sum_{\ell=0}^n (-1)^n \binom{\lambda}{n-\ell} \sum_{k=0}^m (-1)^k \binom{m}{k} \binom{2k}{\ell}. \end{aligned}$$

Denote by $[x^n]f(x)$ the coefficient of x^n in the formal power series $f(x)$. Observe that the last inner sum can be expressed and then evaluated as

$$\begin{aligned} \sum_{k=0}^m (-1)^k \binom{m}{k} \binom{2k}{\ell} &= [x^\ell] \sum_{k=0}^m (-1)^k \binom{m}{k} (1+x)^{2k} \\ &= [x^\ell] \left\{ 1 - (1+x)^2 \right\}^m \\ &= (-1)^m [x^{\ell-m}] (2+x)^m \\ &= (-1)^m \binom{m}{\ell-m} 2^{2m-\ell}. \end{aligned}$$

After substitutions, the second formula “(b)” in the theorem is done. □

Finally letting $\lambda = 0$ in Theorem 8, we recover the following two elegant identities due to Yang and Qiao [7], who discovered them by employing the Riordan array and expressed the results in terms of Bessel numbers [1].

Corollary 9 ($1 \leq m \leq n$).

$$\begin{aligned} \text{(a)} \quad \sum_{k=m}^n \begin{bmatrix} n \\ k \end{bmatrix} \left\{ \begin{matrix} k \\ m \end{matrix} \right\} (-2)^{n-k} &= \frac{(n-1)!}{(m-1)!} \binom{-n}{n-m} 2^{m-n}, \\ \text{(b)} \quad \sum_{k=m}^n \begin{bmatrix} n \\ k \end{bmatrix} \left\{ \begin{matrix} k \\ m \end{matrix} \right\} (-2)^{j-m} &= \frac{n!}{m!} \binom{m}{n-m} (-2)^{m-n}. \end{aligned}$$

Acknowledgement: *The first author is partially supported, during this work, by the National Science foundation of China (Youth Grant No.11601543).*

REFERENCES

- [1] G. Cheon, J. Jung, and L. Shapiro, *Generalized Bessel numbers and some combinatorial settings*, Discrete Math. 313:20 (2013), 2127–2138.
- [2] L. Comtet, *Advanced Combinatorics*, Dordrecht–Holland, The Netherlands, 1974.
- [3] H. W. Gould, *Combinatorial identities*, W. V. Morgantown, 1972.
- [4] R. L. Graham, D. E. Knuth, and O. Patashnik, *Concrete Mathematics*, Addison–Wesley Publ. Company, Reading, Massachusetts, 1989.
- [5] D. E. Knuth, *Two notes on notation*, Amer. Math. Monthly 99 (1992), 403–422.
- [6] D. Stenlund, *Some observations on the connection between Stirling numbers and Bessel numbers*, arXiv:2007.11557v1 [math.CO] 22 Jul 2020.
- [7] S. L. Yang and Z. K. Qiao, *The Bessel numbers and Bessel matrices*, J. Math. Res. Expos. 31:4 (2011), 627–636.

N. N. Li: SCHOOL OF MATHEMATICS AND STATISTICS
ZHOUKOU NORMAL UNIVERSITY, ZHOUKOU (HENAN), CHINA

Email address: lina20190606@outlook.com

W. Chu: DEPARTMENT OF MATHEMATICS AND PHYSICS
UNIVERSITY OF SALENTO, LECCE 73100, ITALY

Email address: chu.wenchang@unisalento.it