Parameterized Algorithms for Editing to Uniform Cluster Graph

Ajinkya Gaikwad, Hitendra Kumar and Soumen Maity

Indian Institute of Science Education and Research, Pune, India ajinkya.gaikwad@students.iiserpune.ac.in; soumen@iiserpune.ac.in

Abstract. Given a graph G = (V, E) and an integer $k \in \mathbb{N}$, we investigate the 2-EIGENVALUE VERTEX DELETION (2-EVD) problem. The objective is to remove at most k vertices such that the adjacency matrix of the resulting graph has at most two eigenvalues. It is established that the adjacency matrix of a graph has at most two eigenvalues if and only if the graph is a collection of equal-sized cliques. Thus, the 2-EIGENVALUE VERTEX DELETION amounts to removing a set of at most k vertices to transform the graph into a collection of equal-sized cliques. The 2-EIGENVALUE EDGE EDITING (2-EEE), 2-EIGENVALUE EDGE DELETION (2-EED) and 2-EIGENVALUE EDGE ADDITION (2-EEA) problems are defined analogously. We present a kernel of size $\mathcal{O}(k^3)$ for 2-EVD, along with an FPT algorithm with a running time of $\mathcal{O}^*(2^k)$. For the problem 2-EEE, we provide a kernel of size $\mathcal{O}(k^2)$. Additionally, we present linear kernels of size 5k and 6k for 2-EEA and 2-EED respectively. For the 2-EED, we also construct an algorithm with running time $\mathcal{O}^*(1.47^k)$. These results address open questions posed by Misra et al. (ISAAC 2023) regarding the complexity of these problems when parameterized by the solution size.

Keywords: Parameterized Complexity \cdot FPT \cdot Eigenvalue \cdot Uniform Cluster Graphs

1 Introduction

Numerous algorithmic challenges involving graphs can be framed as tasks of modifying a graph to meet specific criteria. Notably, over the last three decades, these graph modification tasks have been a significant source of inspiration for innovating new methodologies in parameterized algorithms and complexity theory. This paper delves into a particular graph modification problem known as UNIFORM CLUSTER VERTEX DELETION, along with several related variations of the same problem. Previously explored in [7], the complexity of reducing the count of distinct eigenvalues of a graph by either removing vertices or editing edges was examined. Notably, it can be observed that an adjacency matrix of graph possesses at most two distinct eigenvalues if and only if it comprises of disjoint unions of equally sized cliques. Our notation aligns with that of the aforementioned paper. Let G = (V, E) be a simple undirected graph, where V 2

denotes the vertex set of G and E denotes the edge set of G. We use n to denote |V|. For $u \in V$, we define $N(u) = \{v \in V : (u, v) \in E\}$ and $N[u] = N(u) \cup \{u\}$. The degree of $u \in V$ is |N(u)| and denoted by $d_G(u)$. A clique C in an undirected graph G = (V, E) is a subset of the vertices $C \subseteq V$ such that every two distinct vertices are adjacent. A cluster graph is a graph where every component is a clique. Observe that a graph is a cluster graph if and only if it does not have an induced P_3 , that is, an induced path on three vertices. Let $U \subseteq V$ be a subset of vertices of G and $F \subseteq {V \choose 2}$ be a subset of pairs of vertices of G. The subgraph induced by $U \subseteq V$ is denoted by G[U]. We define $G - U = G[V \setminus U]$, $G - F = (V, E \setminus F)$, $G + F = (V, E \cup F)$ and $G \triangle F = (V, E \triangle F)$. Here, $E \triangle F$ is the symmetric difference between E and F. If $U = \{u\}$ or $F = \{e\}$ then we simply write G - u, G - e and G + e for G - U, G - F and G + F, respectively. It is known that the spectrum of the adjacency matrix of a graph can be computed in polynomial time. The following lemma is crucial to our discussions:

Lemma 1. [2,4] The adjacency matrix of a graph G has at most two distinct eigenvalues if and only if G is a disjoint union of equal sized cliques.

We refer to [1,3] for details on parameterized complexity. Our goal is to present the parameterized complexity landscape of the following problems:

2-EIGENVALUE VERTEX DELETION (2-EVD) **Input:** An undirected graph G = (V, E), and a positive integer k. **Question:** Is there a subset $S \subseteq V$ with $|S| \leq k$ such that G-S is a collection of equal sized cliques?

2-EIGENVALUE EDGE EDITING (2-EEE) **Input:** An undirected graph G = (V, E), and a positive integer k. **Question:** Is there a subset $F \subseteq \binom{V(G)}{2}$ with $|F| \leq k$ such that $G \triangle F = (V, E \Delta F)$ is a collection of equal sized cliques?

2-EIGENVALUE EDGE DELETION (2-EED) **Input:** An undirected graph G = (V, E), and a positive integer k. **Question:** Is there a subset $F \subseteq E$ with $|F| \leq k$ such that $G - F = (V, E \setminus F)$ is a collection of equal sized cliques?

2-EIGENVALUE EDGE ADDITION (2-EEA) **Input:** An undirected graph G = (V, E), and a positive integer k. **Question:** Is there a subset $F \subseteq \binom{V(G)}{2}$ with $|F| \leq k$ such that $G + F = (V, E \cup F)$ is a collection of equal sized cliques?

1.1 Our results

In this paper, we study 2-EVD, 2-EEE, 2-EED, and 2-EEA from the parameterized complexity point of view. Clearly, one can study these problems for any

3

fixed number of distinct eigenvalues. Using some simple yet powerful structural observations and non-trivial ideas, we have provided combinatorial algorithms for all considered variants of the problem. We give a kernel of size $\mathcal{O}(k^3)$ for 2-EVD. Additionally, we provide an FPT algorithm that runs in time $\mathcal{O}^*(2^k)$, which improves upon the FPT algorithm given in [7] based on the standard branching method, which has a running time of $\mathcal{O}^*(3^k)$. For the problem 2-EEE, we provide a kernel of size $\mathcal{O}(k^2)$. Furthermore, we provide linear kernels of sizes 6k and 5k for 2-EED and 2-EEA, respectively, improving upon the quadratic kernel given in [7] for 2-EEA. We also present an FPT algorithm running in time $\mathcal{O}^*(1.47^k)$ for 2-EED, further improving the algorithm from [7]. Through this work, we resolve three open questions listed by Misra et al. [7] concerning the complexity of these problems parameterized by solution size. This study provides a clear picture of the parameterized complexity of the studied problems.

Uniform cluster	FPT algorithm	Kernel
Vertex deletion	Theorem (2) $\mathcal{O}^*(2^k)$	Theorem (1) $\mathcal{O}(k^3)$
Edge editing	-	Theorem (3) $\mathcal{O}(k^2)$
Edge deletion	Theorem (5)	Theorem (4) $6k$
	$\mathcal{O}^*(1.47^k)$	
Edge Addition	-	Theorem (6) $5k$

1.2 Review of previous work

As mentioned in [7], the problem of modifying the graph to reduce the count of distinct eigenvalues to r of the corresponding adjacency matrix was first brought up in [6]. In [7], Misra et al. considered classical and parameterized complexity of this problem. The paper studied mainly four possible operations which are vertex deletion, edge deletion, edge addition and edge editing. For the special case of r = 2, the vertex deletion variant was shown to be NP-complete even on triangle-free and 3d-regular graphs for any $d \geq 2$, and also NP-complete on d-regular graphs for any $d \geq 8$. Moreover, the edge deletion, addition, and editing variants were proved to be NP-complete for r = 2 case. Furthermore, for any fixed $r \geq 3$, they showed that r-EVD is NP-complete on bipartite graphs. Also, the 2-EEA was shown to be NP-complete when the input is either a cluster graph, a forest, or a collection of cycles. Apart from studying the classical complexity, the authors also studied and provided numerous results in the realm of parameterized complexity. In particular they gave FPT algorithms for the 2-EVD and 2-EED problems with running time $\mathcal{O}^*(3^k)$ and $\mathcal{O}^*(2^k)$, respectively. We have improved both of these algorithms. The paper also presented a quadratic kernel for 2-EEA.

The clustering problems hold high importance in the complexity theory as they can model multiple scenarios. One more way to view this problem is to modify graph in such a way that the components of the resulting graph have smaller diameter. This is called as s-CLUB CLUSTER VERTEX DELETION problem where the input is a graph with two integers $s \ge 2$ and $k \ge 1$ and the goal is to decide whether it is possible to remove at most k vertices from G such that each connected component of the resulting graph has diameter at most s.

We note that Madathil and Meeks [5] recently explored a generalization of this problem, termed BALANCED CLUSTER EDITING, in which, given a graph G, and two integers $0 \le \eta \le n$ and k, we are permitted to "edit" up to k edges. This editing should result in a cluster graph where the size difference between any two connected components of the resulting cluster graph does not exceed η . One can see that when $\eta = 0$, this problem reduces to the UNIFORM CLUSTER EDIT-ING problem. The authors presented polynomial kernels for BALANCED CLUS-TER COMPLETION, BALANCED CLUSTER DELETION, and BALANCED CLUSTER EDITING. While they provided polynomial kernels for the general cases, we have achieved better bounds in all considered variants when $\eta = 0$. We obtained our results simultaneously with and independent from those by Madathil and Meeks [5].

2 Kernelization and FPT algorithm for 2-EVD parameterized by solution size

In this section, we begin by proving the following theorem.

Theorem 1. 2-EVD parameterized by solution size admits a kernel of size $\mathcal{O}(k^3)$.

A cluster graph is a graph in which every connected component is a clique. Note that a graph is a cluster graph if and only if it does not contain an induced P_3 . This observation serves as a cornerstone for our kernelization technique. The first step of the kernelization process is to compute a maximal set \mathcal{P}_3 of vertex-disjoint induced P_{3s} in G. This can be accomplished using the greedy algorithm shown below:

Algorithm 1

Require: A graph G = (V, E). **Ensure:** A maximal set \mathcal{P}_3 of vertex-disjoint induced P_3 in G. 1: $\mathcal{P}_3 = \emptyset$ 2: while G has an induced P_3 do 3: Identify an induced $P_3 = (u, v, w)$ in G4: $\mathcal{P}_3 = \mathcal{P}_3 \cup \{(u, v, w)\}$ 5: $G = G - \{u, v, w\}$ 6: end while 7: return \mathcal{P}_3 At the end, if $|\mathcal{P}_3| > k$, we have a no-instance. So we assume that $|\mathcal{P}_3| \leq k$, and let S be the vertices of \mathcal{P}_3 . We have $|S| \leq 3k$. Let us denote the set of cliques of G - S by \mathcal{C} . We have the following simple rules:

Reduction EVD 1 If there exists $s \in S$ with neighbours in at least k+2 cliques of C = G - S, then delete s (and its incident edges from G) and decrement the parameter k by 1. The new instance is (G - s, k - 1)

Lemma 2. Reduction Rule EVD 1 is safe.

Proof. Let X be a uniform cluster vertex deletion set of G of size at most k. For the sake of contradiction, let us assume that $s \in S$, which has neighbours in at least k + 2 cliques of $\mathcal{C} = G - S$, is not in X. Therefore, s must be in a clique of G - X. That clique can have vertices from at most one clique, say C_i in \mathcal{C} . Since s has neighbours in at least k + 2 cliques, we would need to include neighbours of s in at least k + 1 cliques other than C_i in the solution X. This contradicts the condition that the solution size |X| is at most k. Hence if (G, k)is a yes-instance (meaning there exists a uniform cluster vertex deletion set of size at most k), then s must be in X.

Next, we partition the set of cliques in \mathcal{C} based on whether they have neighbours in S or not. We define:

$$\mathcal{C}_0 = \{ C \in \mathcal{C} : \text{no vertex in } C \text{ has a neighbour in } S \}$$

and

$$\mathcal{C}_1 = \{ C \in \mathcal{C} : \text{some vertex in } C \text{ has a neighbour in } S \}.$$

Observe that the exhaustive application of reduction rule EVD 1 completely removes the vertices of S that have neighbours in at least k + 2 cliques of C_1 . In other words, after the exhaustive application of reduction rule EVD 1, each vertex of S can have neighbours in at most k + 1 cliques of C_1 . Since $|S| \leq 3k$ and each vertex of S can have neighbours in at most k + 1 cliques of C_1 , there are at most 3k(k + 1) cliques in C_1 . Therefore, we have

$$|\mathcal{C}_1| \le 3k(k+1).$$

An r-clique is a clique of size r. Now, we have the following simple rule.

Reduction EVD 2 If for some $r \in \{1, 2, ..., n\}$ there are more than k + 1r-cliques in C_0 , then remove all but k + 1 r-cliques from G.

Lemma 3. Reduction rule EVD 2 is safe.

Proof. Suppose (G, k) is a yes-instance and X is a uniform cluster vertex deletion set of G of size at most k. If for some $r \in \{1, 2, ..., n\}$ there are more than k + 1r-cliques in \mathcal{C}_0 , then we claim that r is the size of all cliques in G - X. For the sake of contradiction, let us assume that r' < r is the size of all cliques in G - X. In that case, we need to delete at least one vertex from each r-clique.

This means we would have to add at least k + 1 vertices in X, which contradicts the assumption that X is of size at most k. Thus, we conclude that r is the size of all cliques in G - X. Since the solution size is at most k, this allows us to remove all but k + 1 r-cliques from G.

To bound the number of cliques in C_0 , we make the following observation. This observation helps us bound the number of cliques in C_0 by limiting the variety of cliques sizes present.

Observation 1 C_0 contains cliques of at most k + 1 distinct sizes.

This observation is correct because by deleting at most k vertices we can alter the sizes of at most k cliques. Consequently, C_0 can contain cliques of at most k + 1 distinct sizes. According to Reduction Rule EVD 2, there can be at most k + 1 cliques of the same size. Therefore, we have:

$$|\mathcal{C}_0| \le (k+1)(k+1).$$

Combining this with our earlier result for C_1 :

$$|\mathcal{C}| = |\mathcal{C}_0| + |\mathcal{C}_1| \le (k+1)(k+1) + 3k(k+1) = 4k^2 + 5k + 1.$$

Thus, the total number of cliques $|\mathcal{C}|$ is bounded by $4k^2 + 5k + 1$.

Now, we consider two cases based on the maximum size of a clique in C. Let $\omega(C) = \max_{C \in C} \{|C|\}.$

2.1 Case 1: $\omega(\mathcal{C}) < 8k$

In this case, we obtain a kernel of size: $|S| + \sum_{C \in \mathcal{C}} |C| \le 3k + 8k(4k^2 + 5k + 1) = \mathcal{O}(k^3).$

2.2 Case 2: $\omega(\mathcal{C}) \geq 8k$

Let C_0 be a clique in \mathcal{C} such that $\omega(\mathcal{C}) = |C_0|$. Then C_0 becomes a clique of size at least $|C_0| - k \ge 7k$ in G - X, where X is a solution of size at most k. In this case, the resulting graph G - X must be a collection of r-cliques, where $r \ge |C_0| - k \ge 7k$.

Reduction EVD 3 If there exist C_1 and $C_2 \in C$ such that $|C_1| - |C_2| > 4k$, then delete all the vertices in C_2 and decrement the parameter k by $|C_2|$. The new instance is $(G - C_2, k - |C_2|)$.

Lemma 4. Reduction Rule EVD 3 is safe,

Proof. For the sake of contradiction, let us assume that $v \in C_2$ does not belong to some solution X of size at most k. First, since C_2 is a clique, v has at most $|C_2| - 1$ neighbours in C_2 in the resulting graph. Second, v can have at most 3k neighbours in S. Thus, v can be part of a clique of size at most $(|C_2| - 1) + 3k + 1 = |C_2| + 3k$.

On the other hand, there always exist a vertex $u \in C_1$ such that u is not contained in X. Clearly, u will have at least $|C_1| - k - 1$ neighbours in G - X. Therefore, u must be contained in a clique of size at least $|C_1|-k > |C_2|+4k-k = |C_2|+3k$. Therefore, u and v can never be contained in equal-sized cliques. This contradicts the fact that G - X is a collection of equal sized-cliques. Therefore, we must include all vertices of C_2 in every uniform cluster vertex deletion set X of G of size at most k.

Observation 2 After applying Reduction Rule EVD 3, we get that $\min_{C \in \mathcal{C}} |C| \ge 4k$.

This is true because we have assumed that $\omega(\mathcal{C}) \geq 8k$.

We now introduce the concept of a *heavy neighbour* of a vertex $s \in S$ in C, which is a key element for explaining the forthcoming reduction rules.

Definition 1. A clique $C \in C$ is called a heavy neighbour of $s \in S$ if s is adjacent to at least $\max\{|C| - 4k, k + 1\}$ vertices in C.

We observe that if $s \in S$ has no heavy neighbour, then this vertex should belong to every solution of size at most k. Our next reduction rule is as follows:

Reduction EVD 4 If $s \in S$ has no heavy neighbour, remove s from G, and decrement the parameter k by 1. The new instance is (G - s, k - 1).

Lemma 5. Reduction rule EVD 4 is safe.

Proof. For the sake of contradiction, let us assume that $s \in S$ has no heavy neighbour and does not belong to some solution X of size at most k. It means s must belong to some clique in G - X. Since s has no heavy neighbour in C, s is adjacent to at most $\max\{|C|-4k, k+1\}-1$ vertices in some $C \in C$. Additionally, s can have at most 3k - 1 neighbours in S. Now consider two cases:

- 1. If $\max\{|C| 4k, k+1\} 1 = |C| 4k 1$, then s can be part of a clique of size at most (|C| 4k 1) + (3k 1) + 1 = |C| k 1. This contradicts the fact that G X is a collection of r-cliques, where $r \ge |C| k$.
- 2. If $\max\{|C|-4k, k+1\}-1=k$, then s can be part of a clique of size at most k+(3k-1)+1=4k. This contradicts the fact that G-X is a collection of r-cliques, where $r \ge 7k$.

Therefore, s must be part of every solution of size at most k.

Next, we observe that if $s \in S$ has at least k + 1 neighbours in each of two cliques, say $C, C' \in C$, then this vertex should belong to every solution of size at most k. Our next reduction rule is the following:

Reduction EVD 5 If $s \in S$ has at least k + 1 neighbours in more than one clique in C, then remove s from G, and decrement the parameter k by 1. The new instance is (G - s, k - 1).

Lemma 6. Reduction rule EVD 5 is safe.

8

Proof. Suppose $s \in S$ has at least k + 1 neighbours in each of the cliques $C, C' \in C$. For the sake of contradiction, assume that $s \in S$ does not belong to some solution X of size at most k. It implies that s must belong to a clique in G - X. This clique can contain vertices either from C or C', but not both. Therefore, we must include at least k + 1 neighbours of s from either C or C' in X. This contradicts the fact that X is of size at most k. Hence, s must be part of every solution of size at most k. Thus, removing s and decrementing k by 1 is a safe reduction.

Observation 3 After applying reduction rules 4 and 5 exhaustively, each $s \in S$ has exactly one heavy neighbour in C.

Reduction EVD 6 If $s \in S$ has more than k neighbours in cliques other than its heavy neighbour, then remove s from G, and decrement the parameter k by 1. The new instance is (G - s, k - 1).

Lemma 7. Reduction Rule 6 is safe.

Proof. Note that if $s \in S$ has more than k neighbours in cliques other than its heavy neighbour, then we can construct k + 1 induced P_3 's where the only common vertex is s. Therefore, s must be added to the solution.

Let S_C denotes the set of vertices in S whose heavy neighbour is C. For each $C \in \mathcal{C}$, we define

$$\mathcal{N}(C) = \bigcup_{s \in S_C} \left(C \setminus N(s) \right) \cup \bigcup_{s \in S \setminus S_C} \left(N(s) \cap C \right).$$

We also define

$$\mathcal{N}(\mathcal{C}) = \bigcup_{C \in \mathcal{C}} \mathcal{N}(C)$$

Lemma 8. We have $|\mathcal{N}(\mathcal{C})| = \mathcal{O}(k^2)$.

Proof. Let us assume that C_s denotes the unique heavy neighbour of s. We have

$$\mathcal{N}(\mathcal{C}) = \bigcup_{C \in \mathcal{C}} \left[\bigcup_{s \in S_C} \left(C \setminus N(s) \right) \cup \bigcup_{s \in S \setminus S_C} \left(N(s) \cap C \right) \right]$$
$$= \bigcup_{s \in S} \left[\left(C_s \setminus N(s) \right) \cup \bigcup_{C \in \mathcal{C} \setminus C_s} \left(N(s) \cap C \right) \right]$$

Note that by Reduction Rule 6, for any fixed $s \in S$, we have

$$\left| \bigcup_{C \in \mathcal{C} \setminus C_s} \left(N(s) \cap C \right) \right| < (k+1).$$

Also, for each $s \in S$, we have

$$\begin{aligned} |C_s \setminus N(s)| &\leq |C| - (|C| - 4k) = 4k \text{ if } \max\{|C| - 4k, k + 1\} = |C| - 4k \\ &\leq |C| - (k + 1) \leq 4k \text{ if } \max\{|C| - 4k, k + 1\} = k + 1 \end{aligned}$$

As there are at most 3k vertices in S, we get that $|\mathcal{N}(\mathcal{C})| \leq 3k[k+4k] = 15k^2 = \mathcal{O}(k^2)$.

Lemma 9. For each $C \in C$, every pair of adjacent vertices x and y in $C \setminus \mathcal{N}(C)$ are true twins, that is, N[x] = N[y].

Proof. By the definition of $\mathcal{N}(C)$, it is evident that for any vertex $v \in C \setminus \mathcal{N}(C)$, we have $N(v) \cap S = S_C$. Recall that for the vertices in S_C , C is their heavy neighbour. Since $v \in C$ and C is a clique, v is adjacent to all other vertices of C. Therefore, for every $v \in C \setminus \mathcal{N}(C)$, we have $N[v] = C \cup S_C$.

Reduction EVD 7 If $\min_{C \in \mathcal{C}} \{|C \setminus \mathcal{N}(C)|\} > k+1$, then arbitrarily delete exactly one vertex from each $C \setminus \mathcal{N}(C)$ for every $C \in \mathcal{C}$. The new instance is (G-W,k), where W is the set of vertices deleted from G.

Lemma 10. Reduction Rule EVD 7 is safe.

Proof. Let I' = (G', k) be an instance of 2-EVD obtained from I = (G, k) by exhaustively applying Reduction Rule EVD 7. We will show that I is a yesinstance if and only if I' is a yes-instance. Let $X \subseteq V(G)$ be a solution of size at most k for the instance I. We will construct a solution X' of the same size as X for the instance I'. We know G - X is a collection of equal sized cliques Q_1, Q_2, \ldots, Q_r . We observe that there is no clique Q_i such that $Q_i \subseteq S$. This is true because otherwise $|Q_i| \leq 3k$ as $|S| \leq 3k$, and we have already observed that the size of cliques must be at least 7k. We construct X' from X as follows: if $u \in X \cap (C \setminus \mathcal{N}(C))$ for some $C \in \mathcal{C}$ and u is deleted during the execution of Reduction Rule 7, then remove u from X and include u's true twin $v \notin X$.

We have to show that G' - X' is a collection of equal-sized cliques. It is easy to see that G' - X' is a collection of cliques $Q'_i = Q_i \setminus \{u\}$ where $u \in Q_i$ is a vertex deleted by Reduction Rule 7. As $|Q_i| = |Q_j|$, we get $|Q'_i| = |Q'_j|$ for all $i \neq j$. This shows that I' is a yes-instance.

In the other direction, let us assume that X' is a solution of size at most k for the instance I'. Let us denote the equal sized cliques in G' - X' by Q'_1, Q'_2, \ldots, Q'_r . We claim that X := X' is a solution for the instance I. Note that G - X is again a collection of cliques Q_1, Q_2, \ldots, Q_r where $Q_i = Q'_i \cup \{u\}$ where u is a vertex deleted by Reduction Rule 7. Clearly, we have $|Q_i| = |Q_j|$ as we know that $|Q'_i| = |Q'_j|$ for all $i \neq j$.

After exhaustively applying Reduction Rule 7, we know that there exists a clique, say $C_q \in \mathcal{C}$, such that $|C_q| - |\mathcal{N}(C_q)| \le k + 1$. Note that $\max_{C \in \mathcal{C}} |C| - |C_q| \le 4k$ due to Reduction Rule EVD 3. Therefore, we get $\max_{C \in \mathcal{C}} |C| - |\mathcal{N}(C_q)| \le 5k + 1$. Since $|\mathcal{N}(C_q)| = \mathcal{O}(k^2)$, we get $\max_{C \in \mathcal{C}} |C| \le \mathcal{O}(k^2)$. Note that

$$V(G) = S \cup \bigcup_{C \in \mathcal{C}} C$$

= $S \cup \bigcup_{C \in \mathcal{C}} \left(\mathcal{N}(C) \cup (C \setminus \mathcal{N}(C)) \right)$
= $S \cup \mathcal{N}(C) \cup \bigcup_{C \in \mathcal{C}} (C \setminus \mathcal{N}(C))$

To get a cubic kernel, we need to show that the size of the set $\bigcup_{C \in \mathcal{C}} (C \setminus \mathcal{N}(C))$ is bounded by a cubic function of the parameter. To see this, we partition the set of cliques in \mathcal{C} into two parts:

 $\mathcal{H} = \{ C \in \mathcal{C} \mid C \text{ is a heavy neighbour of some } s \in S \}$

and $\mathcal{L} = \mathcal{C} \setminus \mathcal{H}$. As we have seen, every vertex $s \in S$ has exactly one heavy neighbour, which implies that $|\mathcal{H}| \leq 3k$. Therefore, we get that $\bigcup_{C \in \mathcal{H}} |(C \setminus \mathcal{N}(C))| = \mathcal{O}(k^3)$ as $\max_{C \in \mathcal{C}} |C| \leq \mathcal{O}(k^2)$. Now, let us focus on the set \mathcal{L} . Let us denote $\mathcal{L}' = \{C \in \mathcal{L} \mid |C \setminus \mathcal{N}(C)| \geq k+1\}$ and $\mathcal{L}'' = \mathcal{L} \setminus \mathcal{L}'$. It is clear that $\bigcup_{C \in \mathcal{L}''} |(C \setminus \mathcal{N}(C))| \leq \mathcal{O}(k^3)$ as $|\mathcal{L}''| \leq |\mathcal{C}| \leq \mathcal{O}(k^2)$. We can assume that $|\mathcal{L}'| \geq 2k+1$; otherwise, we achieve the required cubic kernel.

Claim. In this case, we can calculate the size c of equal sized cliques obtained after deleting the vertices in a solution in polynomial time, assuming the input is a yes-instance.

Proof. Let us denote $\mathcal{L}' = \{C'_1, C'_2, \ldots, C'_l\}$, where $l \geq 2k+1$. Let X be a solution of size at most k of the input instance. As $|C'_i \setminus \mathcal{N}(C'_i)| \geq k+1$, at least one vertex will survive from each set $C'_i \setminus \mathcal{N}(C'_i)$ in G - X. Because the vertices in $C'_i \setminus \mathcal{N}(C'_i)$ do not have any neighbours in S, the deletion of at most k vertices can change the degree of vertices in at most k sets $C'_i \setminus \mathcal{N}(C'_i)$. This implies that vertices in all but at most k distinct sets $C'_i \setminus \mathcal{N}(C'_i)$ must have same degree. If this is not the case, we can return a no-instance. If it is the case, then let's say the degree is d. One can easily see that c = d + 1. It is important to note that having $|\mathcal{L}'| \geq 2k + 1$ is necessary; otherwise, identifying d is not possible. \Box

Reduction EVD 8 If there is a vertex v such that d(v) < d, then add it to the solution. The new instance is (G - v, k - 1).

Note that the degree of every vertex in G-X must be equal to d. Since d(v) < d and we cannot increase the degree of v through vertex deletion, v must be added to the solution.

Due to Reduction Rule 8, we can assume that all the cliques in \mathcal{L}' have size at least c.

Reduction EVD 9 If there is a clique $C' \in \mathcal{L}'$ such that |C'| = c, then delete all the vertices in N(C') from the graph G. The new instance is (G - N(C'), k - |N(C')|).

Lemma 11. Reduction Rule 9 is safe.

Proof. Note that at least one vertex $u \in C' \setminus \mathcal{N}(C')$ must survive after deleting the vertices in a solution. This vertex must be part of a clique of size c. Therefore, the clique C' must appear unchanged in the resulting graph. Hence, we must delete all the neighbours of C'.

Let us denote by $\mathcal{L}'_{iso,c}$ the set of isolated cliques of size c in \mathcal{L}' obtained after applying Reduction Rule 9. Similarly, denote by $\mathcal{L}'_{>c}$ the set of cliques of size more than c in \mathcal{L}' .

Reduction EVD 10 If $|\mathcal{L}'_{iso,c}| > k+1$, then retain only k+1 cliques from $\mathcal{L}'_{iso,c}$ and discard the rest.

Lemma 12. Reduction Rule 10 is safe.

Proof. Since the size of cliques after deleting the vertices in a solution of size at most k must be c, we can keep k + 1 cliques of size c to ensure this property holds, and delete the rest. Observe that deleting at most k vertices can alter the size of at most k cliques. By keeping k + 1 cliques, we ensure that the size of cliques in the resulting graph is equal to c.

Reduction EVD 11 If $|\mathcal{L}'_{>c}| > k$, then conclude that we are dealing with a no-instance.

If $|\mathcal{L}'_{>c}| > k$, it implies we need to delete more that k vertices to ensure that each clique in $\mathcal{L}'_{>c}$ becomes a clique of size c in the resulting graph. However, since we are constrained to finding a solution of size at most k, achieving this is not feasible. Therefore, $|\mathcal{L}'_{>c}| \leq k$.

Observation 4 After applying Reduction Rule 8, 9, 10 and 11, the size of \mathcal{L}' is bounded by 2k + 1.

From Observation 4, it follows that $\bigcup_{C \in \mathcal{L}'} |C \setminus \mathcal{N}(C)| \leq \mathcal{O}(k^3)$. This completes the proof of Theorem 1. Next, we will present an FPT algorithm.

2.3 An FPT algorithm for 2-EVD

In this section we give an FPT algorithm for the 2-EVD problem. We prove the following theorem:

Theorem 2. The 2-EVD problem parameterized by solution size k can be solved in $2^k \cdot n^{\mathcal{O}(1)}$ time.

We utilize an algorithm from [9] to find a cluster vertex deletion set X of size at most k in $\mathcal{O}^*(1.811^k)$ time. If this algorithm determines that no such set exists, conclude that we are dealing with a no-instance. Otherwise, this algorithm returns a cluster vertex deletion set X of size at most k. To decide whether G contains a uniform cluster vertex deletion set S of size at most k, we proceed as follows:

- We guess the intersection of S with X, that is, we guess the set $X_{in} = X \cap S$, delete X_{in} from G and reduce parameter k by $|X_{in}|$.
- For each guess of X_{in} , we set $X_{out} = X \setminus X_{in}$ and solve DISJOINT 2-EVD on the instance $(G X_{in}, X_{out}, k |X_{in}|)$.

For each guess X_{in} , we try to find a uniform cluster vertex deletion set S' in $G - X_{in}$ of size at most $k - |X_{in}|$ that is disjoint from X_{out} . This problem is called DISJOINT 2-EVD. If for some guess X_{in} , we find a uniform cluster vertex deletion set S' in $G - X_{in}$ of size at most $k - |X_{in}|$ that is disjoint from X_{out} , then we output $S = X_{in} \cup S'$. Otherwise, we conclude that the given instance of the DISJOINT 2-EVD problem is a no-instance. The number of all guesses is bounded by 2^k . Therefore, to obtain an FPT algorithm for 2-EVD, it is sufficient to solve DISJOINT 2-EVD in polynomial time.

2.3.1 An algorithm for DISJOINT 2-EVD: Let $(G - X_{in}, X_{out}, k)$ be an instance of DISJOINT 2-EVD, and let H = G - X, where $X = X_{in} \cup X_{out}$. We denote $G' = G - X_{in}$.

Reduction Disjoint-EDV 1 If $G[X_{out}]$ is not a disjoint union of cliques or G - X is not a disjoint union of cliques, then return that (G', X_{out}, k) is a no-instance.

Reduction Disjoint-EDV 2 If there is a vertex v in G - X which has neighbours in two distinct cliques in $G[X_{out}]$, then delete v from G and decrement the parameter k by 1. The new instance is $(G' - \{v\}, X_{out}, k - 1)$.

The previous reduction rule is correct because we will get an induced P_3 where the only vertex that we are allowed to delete is v. Therefore, we must add v to the solution.

So from now onwards, we assume that $G[X_{out}]$ is a disjoint union of cliques and H = G - X is a disjoint union of cliques. We also guess the exact size of the solution as k'. The possible guesses for k' range from 1 to k. This will be helpful

in the final part of the algorithm. We will solve this problem by guessing the size c of cliques in the resulting graph. At the end, we will compare the solution sizes for each c and return the minimum one. The possible guesses for c ranges from 1 to n.

Reduction Disjoint-EDV 3 If there is a clique of size exactly c in $G[X_{out}]$ then delete N(C) and add all the vertices of the set N(C) to the solution. The new instance is $(G' - N[C], X_{out}, k - |N(C)|)$.

Observation 5 If there is a clique C of size larger than c in $G[X_{out}]$, then we can discard such a guess for c.

Observation 6 Every clique in $G[X_{out}]$ of size less than c must utilize vertices from exactly one clique in G - X to form a clique of size c.

Due to Observation 6, for every clique C of size less than c in $G[X_{out}]$, there is a unique clique in G - X that helps to form a clique of size c containing vertices from C.

We now construct a bipartite graph with bipartition (A, B) in the following way. For a given guess (k', c), observe that the number of equal-sized cliques in the resulting graph is $p = \frac{|V(G')| - k'}{c}$. We need a total of p vertices in A. Suppose the number of cliques in $G[X_{out}]$ is p_1 . We add a vertex $a \in A$ corresponding to each clique in $G[X_{out}]$. If $p_1 < p$, we add $p-p_1$ dummy vertices in A to ensure the number of vertices in A equals p. Similarly, we add a vertex $b \in B$ corresponding to each clique in G-X. Due to Reduction Rule Disjoint-EDV 3 and Observation 5, it follows that every clique in $G[X_{out}]$ has size less than c. We add an edge between $a \in A$ and $b \in B$ if the clique corresponding to a can be transformed into a clique of size c using some vertices from the clique corresponding to b. Additionally, some cliques in G-X can be transformed into cliques of size c by deleting the necessary number of vertices from the respective cliques. We make $b \in B$ adjacent to all dummy vertices in A if the clique corresponding to b can be transformed into a clique of size c by deleting the necessary number of vertices.

We now try to find an A-saturated matching in this bipartite graph. If an Asaturated matching does not exist, we discard such a guess (k', c). Otherwise, we find an A-saturated matching M of size p. We identify p cliques in the resulting graph as follows: If $(a, b) \in M$ and a is not a dummy vertex, then the clique corresponding to $a \in A$ will be transformed into a clique of size c using some vertices from the clique corresponding to b. If $(a, b) \in M$ and a is a dummy vertex, then the clique corresponding to $b \in B$ will be transformed to a clique of size c by deleting the necessary number of vertices. Thus we know exactly the cliques in the final graph and can delete the necessary vertices to form the cliques of size c. If the solution size k' matches the number of vertices that are deleted from G', then we return a yes-instance; otherwise, we return a no-instance. \Box

3 Kernelization algorithm for 2-EEE parameterized by solution size

In this section, we study the following problem: For a given graph G, can we transform G into a uniform cluster graph by editing at most k adjacencies, where editing involves adding or deleting at most k edges? More formally, let G = (V, E) be a graph. Then $F \subseteq V \times V$ is called a *uniform cluster editing set* for G if $G \triangle F$ is a uniform cluster graph. In this section, we prove the following theorem.

Theorem 3. 2-EEE parameterized by solution size admits a kernel of size $\mathcal{O}(k^2)$.

Note that many preprocessing rules applied in the case of 2-EEE are also applicable in the cases of 2-EEA and 2-EED. To maintain conciseness, we will refer to the following preprocessing step as the *preparation step*, which begins here.

A graph is a cluster graph if and only if it does not have an induced P_3 . It is straightforward to compute a maximal set \mathcal{P}_3 of vertex-disjoint induced P_3 s in G using Algorithm 1. If $|\mathcal{P}_3| > k$, then we have a no-instance. Therefore, we assume that $|\mathcal{P}_3| \leq k$, and let S be the vertices of \mathcal{P}_3 . We have $|S| \leq 3k$. Let \mathcal{C} denote the set of cliques of G - S. Let F be a uniform cluster editing set of size at most k for G, and let V_F be the vertices of F. Then we have $|V_F| \leq 2k$.

We observe that if there indeed exists a uniform cluster editing set F of size at most k, then the vertices in $V \setminus V_F$ have equal degree in both G and $G \triangle F$. In other words, for any two distinct vertices $x, y \in V \setminus V_F$, we have $d_G(x) = d_G(y) = d_{G \triangle F}(x) = d_{G \triangle F}(y)$. We assume that G has at least 4k + 1vertices; otherwise, we would have a kernel of size 4k. It is known that if the input is a yes-instance, then at least 2k + 1 vertices have the same degree d, and at most 2k vertices have degrees not equal to d. Therefore, we first check if the input instance (G, k) satisfies this condition. If the input instance does not satisfy this condition, then conclude that we are dealing with a no-instance. If the input instance satisfies this condition, we can calculate the exact value of d in polynomial time. Assuming that the input is a yes-instance, let c be the size of equal sized cliques in $G \triangle F$. It follows that c must be equal to d+1 in $G \triangle F$. This implies that the minimum degree of G is at least d - k, because executing k edge editing operations can increase the degree of a vertex by at most k. Similarly, the maximum degree of G is at most d+k, because executing k edge editing operations can decreases the degree of a vertex by at most k. The preparation step ends here. The following two rules are safe.

Reduction EEE 1 If $\delta(G) < d-k$ or $\Delta(G) > d+k$ then conclude that we are dealing with a no-instance.

Reduction EEE 2 If the number of vertices in V(G) with degrees not equal to d exceeds 2k, then conclude that we are dealing with a no-instance.

We will consider two cases based on the value of d.

Case 1: Let us assume that $d \ge 6k$.

Lemma 13. If (G, k) is a yes-instance, then for each $C \in C$, we have |C| > 2k.

Proof. If there is a clique $C \in \mathcal{C}$ such that $|C| \leq 2k$, then every vertex $v \in C$ has a degree at most 5k-1. This is because v has at most 2k-1 neighbours within C and at most 3k neighbours in S. Given $d \geq 6k$, we have 5k-1 < d-k. This implies $\delta(G) < d-k$. Accordingly to Reduction Rule 1, if $\delta(G) < d-k$, then we conclude that we are dealing with a no-instance. Therefore, we must have |C| > 2k.

Reduction EEE 3 If $s \in S$ is adjacent to at least k + 1 vertices of a clique $C \in C$, then add all missing edges between s and C and decrement the parameter k by |W|, where W is the set of all missing edges between s and C. The resulting instance is (G + W, k - |W|). If $|W| \ge k + 1$, then conclude that we are dealing with a no-instance.

Lemma 14. Reduction rule 3 is safe.

Proof. The vertices of C must belong to the same connected component in the edited graph. Since s has at least k + 1 neighbours in C, s must be in the same connected component as the vertices in C. Otherwise, to place s in a separate component, we would need to delete at least k + 1 edges between s and C, which contradicts our assumption that $|F| \leq k$. Therefore, we should add all the missing edges between s and C.

Reduction EEE 4 If $s \in S$ is not adjacent to at least k+1 vertices of a clique $C \in C$, then delete all edges between s and C, and decrement the parameter k by |W|, where W is the set of edges between s and C. The resulting instance is $(G \triangle W, k - |W|)$. If $|W| \ge k + 1$, then conclude that we are dealing with a no-instance.

Lemma 15. Reduction rule 4 is safe.

Proof. All the vertices of C must belong to the same connected component in the edited graph. Note that s cannot be in the same connected component as the vertices in C, as it would require adding at least k + 1 edges. Therefore, we must delete all the edges between s and C.

Observe that exhaustive application of reductions EEE 3 and EEE 4 ensures that for each $s \in S$ and each $C \in C$, either s is adjacent to every vertex of C or s is adjacent to no vertex of C.

Reduction EEE 5 If $s \in S$ has neighbours in two distinct cliques $C_1, C_2 \in C$, then conclude that we are dealing with a no-instance.

Lemma 16. Reduction rule 5 is safe.

Proof. Due to Reduction rules EEE 3 and EEE 4, s must be adjacent to every vertex of C_1 and C_2 . This implies that the graph contains at least 2k+1 edge disjoint induced P_{3s} . Therefore the input instance is a no-instance.

Reduction EEE 6 If s_1 and s_2 are two non-adjacent vertices of S and have a neighbour in $C \in C$, then add the edge (s_1, s_2) and decrement the parameter k by 1. The resulting instance is $(G + (s_1, s_2), k - 1)$.

Lemma 17. Reduction rule 6 is safe.

Proof. Due to reductions EEE 3 and EEE 4, s_1 and s_2 are adjacent to every vertex of C. Thus, s_1 and s_2 must be in the same connected component as the vertices in C in the edited graph. Therefore, we must add the edge (s_1, s_2) to the solution.

Reduction EEE 7 If s_1 and s_2 are two adjacent vertices of S and have neighbours in $C_1 \in \mathcal{C}$ and $C_2 \in \mathcal{C}$ respectively, then delete the edge (s_1, s_2) and decrement the parameter k by 1. The resulting instance is $(G - (s_1, s_2), k - 1)$.

Lemma 18. Reduction rule 7 is safe.

Proof. The vertices of C_1 and the vertices of C_2 must be in different connected components in the edited graph. Therefore, s_1 and s_2 must also be in different connected components in the edited graph. Therefore, we must delete the edge (s_1, s_2) and add it to the solution.

Lemma 19. If (G, k) is a yes-instance and none of the reduction rules EEE 1 to EEE 7 are applicable to G, then G is a disjoint union of cliques, each of size d + 1.

Proof. When none of the reduction rules EEE 1 to EEE 7 is applicable to G, then G is clearly a disjoint union of cliques. Now, we show that these cliques are of size d+1. If there exists a clique of size more than d+1, then there would be more than 2k vertices whose degree is not equal to d. Similarly, if there exists a clique of size less than d+1, then there would be more than 2k vertices whose degree is not equal to d, because as shown in Lemma 13, all cliques $C \in \mathcal{C}$ have size at least 2k + 1. This scenario contradicts Reduction Rule 2. \square

In Case 1, we obtain a kernel of size $\mathcal{O}(1)$.

Case 2: Let us assume that $d \leq 6k - 1$.

We partition $\mathcal{C} = G - S$ into four parts as follows:

- $C_{\leq} = \{ C \in C : |C| < d+1 \}$
- $-\mathcal{C}_{0,d+1} = \{C \in \mathcal{C} : |C| = d+1 \text{ and no vertex in } C \text{ has a neighbour in } S\}$
- $-\mathcal{C}_{1,d+1} = \{C \in \mathcal{C} : |C| = d+1 \text{ and some vertex in } C \text{ has a neighbour in } S\}$ $-C_{>} = \{C \in C : |C| > d+1\}$

Lemma 20. If (G, k) is a yes-instance and reduction rule EEE 2 is not applicable to G, then $\bigcup_{C \in C_{>}} |V(C)| \leq 2k$.

Proof. Note that every vertex in $\bigcup_{C \in C_{>}} V(C)$ has degree more than d + 1. Due to Reduction rule 2, the number of vertices with degree more than d + 1 is bounded by 2k.

Reduction EEE 8 If $|C_{0,d+1}| > 2k$, then retain only 2k+1 cliques from $C_{0,d+1}$ and discard the rest.

Lemma 21. Reduction Rule 8 is safe.

Proof. An edge editing set of size at most k can change degrees of at most 2k vertices in $\mathcal{C}_{0,d+1}$. In other words, an edge editing set of size at most k can affect at most 2k cliques in $\mathcal{C}_{0,d+1}$. There will still exist at least one clique of size d+1 in the final graph. Therefore, we can retain only 2k + 1 cliques.

Lemma 22. $|C_{1,d+1}| \leq 2k$.

Proof. Each clique $C \in C_{1,d+1}$, by definition, contains at least one vertex that has a neighbour in S. This implies that every clique in $C_{1,d+1}$ contains at least one vertex whose degree is more than d. Due to Reduction Rule 2, there are at most 2k such vertices. Therefore, $|C_{1,d+1}| \leq 2k$.

Therefore, we get $\left| \bigcup_{C \in \mathcal{C}_{d+1}} V(C) \right| \le (4k+1)(6k-1) = 24k^2 + 2k - 1$, where $\mathcal{C}_{d+1} = \mathcal{C}_{0,d+1} \cup \mathcal{C}_{1,d+1}$.

Lemma 23. If (G, k) is a yes-instance and reduction rule EEE 1 is not applicable to G, then we have $\left| \bigcup_{C \in \mathcal{C}_{<}} V(C) \right| < 21k^2 + 5k.$

Proof. Since Reduction Rule 1 is not applicable to G, every vertex in G has a degree at most $d + k \leq 7k - 1$. For the sake of contradiction, assume that $\bigcup_{C \in \mathcal{C}_{<}} |V(C)| \geq 21k^2 + 5k$. Each clique $C \in \mathcal{C}_{<}$ has size less than d + 1, hence

the vertices in C have degree less than d. As (G, k) is a yes-instance, G has a uniform cluster vertex editing set of size at most k that can increase the degree of at most 2k vertices. Therefore, at least $21k^2 + 5k - 2k$ vertices in $\bigcup_{C \in \mathcal{C}_{<}} V(C)$

have neighbours in S. Since $|S| \leq 3k$ and S has at least $21k^2 + 53k$ neighbours in G-S, by the Pigeonhole principle, there is a vertex $s \in S$ with at least 7k + 1 neighbours in G-S, which contradicts the fact that $d_G(v) \leq 7k - 1$ for all $v \in V$. Therefore, we conclude that $\bigcup_{C \in \mathcal{C}_{\leq}} |V(C)| < 21k^2 + 5k$.

Applying the above reduction rules and results, we find that:

$$|V(G)| = |S| + |V \setminus S|$$

= $|S| + \left| \bigcup_{C \in \mathcal{C}_{<}} V(C) \right| + \left| \bigcup_{C \in \mathcal{C}_{d+1}} V(C) \right| + \left| \bigcup_{C \in \mathcal{C}_{>}} V(C) \right|$
 $\leq 3k + (21k^2 + 5k) + (24k^2 + 2k - 1) + 2k$
= $45k^2 + 12k - 1.$

4 Kernelization and FPT algorithm for 2-EED parameterized by solution size

In this section, we shall prove the following theorem.

Theorem 4. 2-EED parameterized by solution size admits a kernel of size at most 6k.

We start with the same preparation step as designed for 2-EEE. However, in this preparation step, we apply edge deletion operations instead of edge editing operations.

Reduction EED 1 If $\delta(G) < d$ or $\Delta(G) > d + k$, then conclude that we are dealing with a no-instance.

Reduction EED 2 If the number of vertices in V(G) with degrees not equal to d exceeds 2k, then conclude that we are dealing with a no-instance.

Reduction EED 3 If there exists a vertex u of degree d in G such that the subgraph induced by N[u] is not a clique, then conclude that we are dealing with a no-instance.

Lemma 24. Reduction Rule 3 is safe.

Proof. Note that if the input is a yes-instance, then every vertex u of degree d must be part of a clique of size d + 1 after deleting the edges of the solution. Since the degree of u is d, this can only happen if the closed neighborhood of u (denoted N[u]) forms a clique. Therefore, if the subgraph induced by N[u] is not a clique, the instance is a no-instance.

Reduction EED 4 Let $u \in V(G)$ be a vertex with degree d. If u has a neighbour v with a degree strictly greater than d+1, then delete all the edges $(v, w) \in E(G)$ such that w is neither u nor a neighbour of u. The set of edges deleted is denoted as W. The resulting instance is G = (V(G), k - |W|). Also if $|W| \ge k + 1$ then conclude that we are dealing with a no-instance.

Lemma 25. Reduction Rule 4 is safe.

Proof. As we have seen from the proof of Lemma 24, N[v] must form a clique component in the final graph. Therefore, we must delete the edges $(v, w) \in E(G)$ such that $v \in N[u]$ but $w \in V(G) \setminus N[u]$. It is also worth noting that the condition $|V(G)| \ge 4k + 1$ ensures that the resulting graph, obtained by deleting the edges in the solution, must be a cluster graph, where every clique is of size exactly d+1.

Reduction EED 5 If there are multiple cliques of size d + 1, then we retain only x cliques of size d + 1 where $x(d + 1) \ge 2k + 1$ and remove the remaining cliques of size d + 1.

Lemma 26. Reduction Rule 5 is safe.

Proof. The condition $x(d+1) \ge 2k+1$ in Reduction Rule EED 5 ensures that at least 2k+1 vertices have degree exactly d, which in turn ensures that the size of cliques, after deleting at most k edges, remains exactly d+1. Therefore deleting the extra cliques does not change the size of a solution.

We make two cases based on the size of d.

Case 1. Assume d > 2k

Reduction EED 6 Let (G, k) be an input instance such that Reduction Rules EED 1-5 are not applicable to (G, k). If $k \ge 0$ and G is a disjoint union of cliques of size d + 1, then conclude that we are dealing with an yes-instance.

Lemma 27. Reduction Rule 6 is safe.

Proof. Suppose there exists a vertex v with a degree of at least d + 1. If N(v) contains a vertex with degree d, then Reduction Rule EED 4 could have been applied. Therefore, we assume that all vertices in N(v) have a degree of at least d + 1. As a result, the number of vertices with degrees not equal to d is at least d+2 > 2k+2 as d > 2k. However, in this scenario, Reduction Rule EED 2 could have been applied because the number of vertices with degrees not equal to d exceeds 2k. Consequently, in a reduced graph, every vertex has a degree exactly d. After exhaustively applying Reduction Rules EED 3 and EED 4, G must be a disjoint union of cliques of size d + 1. □

In Case 1, we obtain a kernel of size $\mathcal{O}(1)$.

Case 2. Let us assume $d \leq 2k$.

After exhaustively applying Reduction Rules EED 3 and EED 4, the components containing a vertex of degree d must form a d + 1-sized clique. After applying EED 5, the cliques of size d+1 collectively contain at most 4k vertices. According to Reduction Rule EED 2, we have at most 2k vertices that do not have degree d. Therefore, we have $|V(G)| \leq 6k$.

This completes the proof of Theorem 4.

Theorem 5. The 2-EED problem, parameterized by solution size, can be solved by an algorithm running in $\mathcal{O}^*(1.47^k)$ time.

To prove Theorem 5, we will establish a simple yet important lemma. A cluster edge deletion set E_s is minimal if it contains no other cluster edge deletion set.

Lemma 28. In a non-cluster graph G, any uniform cluster edge deletion set of size at most k will always contain a minimal cluster edge deletion set of size at most k.

Proof. Let E_s be a uniform cluster edge deletion set. Then, $G \setminus E_s$ is a disjoint union of x-sized cliques denoted by $\mathcal{C} = \{C_1, C_2, \ldots, C_l\}$. If x = 1, then E_s equals E. In this case, we can find a minimal cluster edge deletion set by identifying a P_3 (a path of length 3) and deleting an arbitrary edge from it. It is evident that such a set of edges is contained in E. Now, we assume x > 1. Algorithm 2 provides a method to construct a minimal cluster edge deletion set contained within a given uniform cluster edge deletion set.

Algorithm 2

Require: A graph G = (V, E) and a uniform cluster edge deletion set E_s . **Ensure:** A minimal cluster edge deletion set of G contained in E_s .

- 1: Set $C = G E_s = \{C_1, C_2, ..., C_l\}$ and $E' = E_s$
- 2: while there exist C_i and C_j in \mathcal{C} such that $1 \leq i < j \leq l$ and $G[C_i \cup C_j]$ is a clique do

Remove C_i and C_j from \mathcal{C} , and add $C_i \cup C_j$ to \mathcal{C} . Re-enumerate cliques in \mathcal{C} . 3:

 $E' = E' \setminus E_{ij}$, where E_{ij} is the set of edges between C_i to C_j . 4:

5: end while 6: return E'

Let us prove the correctness of Algorithm 2. Note that $E' \neq \emptyset$, otherwise G would be a cluster graph. Clearly, E' is a cluster edge deletion set. We claim that E' is a minimal cluster edge deletion set.

For the sake of contradiction, assume there is a proper subset F of E' such that F is a cluster edge deletion set. Suppose $(u, v) \in F$ where $u \in C_i$ and $v \in C_i$. Since $G[C_i \cup C_i]$ is not a clique, we find an induced P_3 , which contradicts the fact that E' is a cluster edge deletion set.

Let us proceed with the proof of Theorem 5. To begin with, we will enumerate all minimal cluster edge deletion sets of size at most k. This enumeration can be done in $\mathcal{O}^*(1.47^k)$ time [8]. Next, we will make a guess of a minimal cluster edge deletion set E' that is contained in a potential solution. It is evident that $G \setminus E'$ forms a cluster graph. Now we guess the size of cliques after deleting edges in a solution. Lets denote it as c. It is worth noting that a uniform cluster edge deletion set $E' \cup E''$ can be readily constructed in polynomial time, where the cliques have size c. If such a set does not exist, the guess is discarded. Afterward, we check if the size of set $E' \cup E''$ is less than or equal to k. If it is, we return an

yes-instance, otherwise we discard the guess. If all guesses have been discarded by the end of this process, we return a no-instance.

5 Kernelization algorithm for 2-EEA parameterized by solution size

In this section, we shall prove the following theorem.

Theorem 6. 2-EEA admits a kernel with at most 5k vertices.

We start with the same preparation step as designed for 2-EEE. However, in this preparation step, we apply edge addition operations instead of edge editing operations.

Reduction EEA 1 If $\delta(G) < d - k$ or $\Delta(G) > d$, then conclude that we are dealing with a no-instance.

Reduction EEA 2 If the number of vertices in V(G) with degrees not equal to d exceeds 2k, then conclude that we are dealing with a no-instance.

Reduction EEA 3 If there is a path between two nonadjacent vertices u and v, then make u adjacent to v and decrease k by 1. The new instance becomes (G + (u, v), k - 1).

Lemma 29. Reduction Rule 3 is safe.

Proof. Since there is a path between u and v, they must belong to the same clique in the edited graph. Therefore, we must connect u and v by an edge. \Box

Observe that exhaustive application of Reduction Rule EEA 3 ensures that the new graph is a collection of disjoint cliques.

Reduction EEA 4 If there are multiple cliques of size d + 1, then we retain only x cliques of size d + 1 where $x(d + 1) \ge 2k + 1$ and remove the remaining cliques of size d + 1.

Lemma 30. Reduction Rule 4 is safe.

Proof. Note that Reduction Rule EEA 4 ensures that at least 2k + 1 vertices have a degree exactly d, which in turn ensures that the size of cliques remains exactly d + 1 after adding at most k edges. Therefore, deleting the extra cliques does not change the size of the solution.

Case 1: Let us assume that $d \ge k + 1$.

Reduction EEA 5 If there is a clique of size less than d+1 in G, then conclude that we are dealing with a no-instance.

Proof. Due to Reduction Rule EEA 1, there cannot be a clique of size more than d + 1. Let's assume that we have a clique C of size x < d + 1. To obtain a clique of size d + 1 from C, we would need to add at least x(d + 1 - x) edges. However, this exceeds the allowable number k of edge addition operations, as x(d + 1 - x) > k.

In Case 1, we obtain a kernel of size $\mathcal{O}(1)$.

Case 2: Let us assume that $d \leq k$.

After applying Reduction Rule EEA 4, the cliques of size d + 1 collectively contain at most 3k vertices. Accordingly to Reduction Rule 2, there are at most 2k vertices with degrees not equal to d. Therefore, we have $|V(G)| \leq 5k$.

This completes the proof of Theorem 6.

6 Conclusion and Open Problems

In this paper, we studied the problem of modifying a given graph so that the resulting graph becomes a collection of equal-sized cliques. We provided polynomial kernels for various types of modification problems. It would be interesting to explore whether the kernelization algorithms presented in this paper can be improved to achieve better bounds.

References

- M. Cygan, F. V. Fomin, L. Kowalik, D. Lokshtanov, D. Marx, M. Pilipczuk, M. Pilipczuk, and S. Saurabh. *Parameterized Algorithms*. Springer, 2015.
- 2. M. Doob. On characterizing certain graphs with four eigenvalues by their spectra. Linear Algebra and its Applications, 3:461–482, 1970.
- 3. R. G. Downey and M. R. Fellows. Parameterized Complexity. Springer, 2012.
- F. Goldberg, S. Kirkland, A. Varghese, and A. Vijayakumar. On split graphs with four distinct eigenvalues. *Discrete Applied Mathematics*, 277:163–171, 2020.
- J. Madathil and K. Meeks. Parameterized algorithms for balanced cluster edge modification problems, 2024.
- S. Meesum, P. Misra, and S. Saurabh. Reducing rank of the adjacency matrix by graph modification. *Theoretical Computer Science*, 654:70–79, 2016. Computing and Combinatorics.
- N. Misra, H. Mittal, S. Saurabh, and D. Thakkar. On the Complexity of the Eigenvalue Deletion Problem. In S. Iwata and N. Kakimura, editors, 34th International Symposium on Algorithms and Computation (ISAAC 2023), volume 283 of Leibniz International Proceedings in Informatics (LIPIcs), pages 53:1–53:17, Dagstuhl, Germany, 2023. Schloss Dagstuhl Leibniz-Zentrum für Informatik.
- T. Nolander. On the fixed-parameter enumerability of cluster deletion. Master's thesis, Computing Science, Chalmers (Göteborg), 2006.
- D. Tsur. Faster parameterized algorithm for cluster vertex deletion. Theory of Computing Systems, 65:1–21, 02 2021.