


Computing with Hypergeometric-Type Terms

Bertrand Teguia Tabugua 

Department of Computer Science, University of Oxford, UK
bertrand.teguia@cs.ox.ac.uk

Abstract. Take a multiplicative monoid of sequences in which the multiplication is given by Hadamard product. The set of linear combinations of interleaving monoid elements then yields a ring. We consider such a construction for the monoid of hypergeometric sequences, yielding what we call the ring of hypergeometric-type sequences – a subring of the ring of holonomic sequences. We present two algorithms in this setting: one for computing holonomic recurrence equations from hypergeometric-type normal forms and the other for finding products of hypergeometric-type terms. These are newly implemented commands in our Maple package `HyperTypeSeq` available [here](#), which we also describe.

Keywords: m -fold indicator sequence · P-recursive sequence · interlaced hypergeometric term · symbolic computation

1 Introduction

In [1], Abramov defined m -sparse sequences as those whose frequent non-zero terms have the same index modulo some positive integer m . Any such sequence has a general term of the form

$$c_n \chi_{\{n \equiv j \pmod{m}\}}, \quad (1)$$

where $(c)_n := (c_n)_{n \in \mathbb{N}} = (c_0, c_1, \dots)$ is some non-zero sequence, and $\chi_{\{n \equiv j \pmod{m}\}}$ evaluates the indicator function of the set $\{j + mk, k \in \mathbb{N}\}$ at n , $j \in \mathbb{N}, j < m$. We introduced the latter in [13] under the name m -fold indicator sequence to enable a symbolic treatment of interlacement (also known as interleaving). The formal definition can be stated as follows.

Definition 1 (m -fold indicator sequence). *The m -fold indicator sequence of characteristic m and remainder j , denoted $(\chi_{\{j \pmod{m}\}})_n := (\chi_{\{n \equiv j \pmod{m}\}})_{n \in \mathbb{N}}$, is defined by the general term*

$$\chi_{\{j \pmod{m}\}}(n) = \chi_{\{n \equiv j \pmod{m}\}} = \begin{cases} 1 & \text{if } n \equiv j \pmod{m} \\ 0 & \text{otherwise} \end{cases}, \quad n \in \mathbb{N}, \quad (2)$$

where $n \equiv j \pmod{m}$ means that j is the non-negative remainder of n in its division by m .

Remark 1 (Notation). The general term of a sequence $(s)_n$ is s_n (without the parentheses) or $s(n)$. We usually use the former notation; we use the latter when the subscript is already occupied. For instance, the term $T_1(n)$ is the general term of the sequence $(T_1)_n$. Throughout this paper, \mathbb{K} denotes a field of characteristic zero.

By convention, the zero sequence $(0, 0, \dots)$ is $(\chi_{\{0 \pmod{0}\}})_n$ and the one sequence $(1, 1, \dots)$ is $(\chi_{\{0 \pmod{1}\}})_n$. Although there is a similar understanding of interlacement in the literature (see

[6, Section 2.2]), we are unaware of any algorithms or software to perform automatic computations when interacements occur in the general term of a sequence. We consider linear combinations of interlaced sequences. These linear combinations define a ring when the corresponding sequences are closed under Hadamard product. The proof is ipso facto the same as that of [13, Theorem 1]. For holonomic sequences, this ring is the same as the ring of holonomic sequences. We are interested in the case of hypergeometric sequences.

Definition 2 (Hypergeometric-type sequence). *A sequence $(s)_n$ is said to be of hypergeometric type if there exist finitely many m -fold indicator sequences $(\chi_{\{j_1 \bmod m_1\}})_n, \dots, (\chi_{\{j_l \bmod m_l\}})_n$ such that its general term s_n writes*

$$s_n = H_1(\sigma_1(n)) \cdot \chi_{\{n \equiv j_1 \bmod m_1\}} + H_2(\sigma_2(n)) \cdot \chi_{\{n \equiv j_2 \bmod m_2\}} + \dots + H_l(\sigma_l(n)) \cdot \chi_{\{n \equiv j_l \bmod m_l\}}, \quad (3)$$

where $\sigma_i: \mathbb{N} \rightarrow \mathbb{Q}$ is such that $\sigma_i(m_i \cdot n + j_i) \in \mathbb{N}$, and $H_i(n)$ is a \mathbb{K} -linear combination of hypergeometric terms, $i = 1, \dots, l$. We call the H_i 's the coefficients of s_n .

Example 1 (OEIS [10] A212579 (see [13])). The sequence $(s)_n$ whose general term s_n counts the number of (w, x, y, z) in $\{1, \dots, n\}^4$ such that $\min(|w-x|, |w-y|) = \min(|x-y|, |x-z|)$ is of hypergeometric type. We have

$$s_n = \frac{4}{9} + \frac{31}{12}n - 3n^2 + \frac{67}{36}n^3 - \frac{1}{4}n\chi_{\{n \equiv 2 \bmod 0\}} - \frac{4}{9}n\chi_{\{n \equiv 3 \bmod 0\}} - \frac{8}{9}\chi_{\{n \equiv 3 \bmod 1\}}. \quad (4)$$

■

A hypergeometric-type term written in the form of (3) is in normal form. We denote by (\mathcal{H}_T) the set of hypergeometric-type sequences, and \mathcal{H}_T the one of their general terms in normal forms. We present the Maple software package `HyperTypeSeq` [11] dedicated to computations with hypergeometric-type terms. The package contains the following commands.

- `HolonomicRE(expr, a(n), maxreorder = d)`: adapts `HolonomicDE` from [7] to search for a holonomic recurrence equation from an expression and a given order bound d . For any term s_n for which $s_{n+1} = f(s_n, n)$ for some function f , the procedure looks for a non-trivial $\mathbb{K}[n]$ -linear dependency in $\langle s_{n+j}, 0 \leq j \leq d \rangle$.
- `mfoldInd(n, m, j)`: evaluates an m -fold indicator term or write it symbolically as $\chi_{\{mod p(n,m)=j\}}$.
- `HTSeval(s, n = j)`: evaluates a hypergeometric-type term at index j .
- `HTS(expr, n)`: writes a given expression into a hypergeometric-type normal form whenever possible.
- `REtoHTS(RE, a(n), P)`: aims to decide whether a given holonomic term is of hypergeometric type or not by writing it in hypergeometric-type normal form. The argument P is a procedure to compute the sequence terms from the recurrence or a list of sufficiently many initial terms.

Example 2. Let us give some examples. The next section will give examples for `HolonomicRE`.

```
> with(HyperTypeSeq): #load the package
> mfoldInd(7,3,1); mfoldInd(n,3,1);
s:=n!*mfoldInd(n,4,2)+2^n*mfoldInd(n,2,1); HTSeval(s,n=6);
```

$$1 \quad (5)$$

$$\chi_{\{mod p(n,3)=1\}} \quad (6)$$

$$s := n! \chi_{\{mod p(n,4)=2\}} + 2^n \chi_{\{mod p(n,2)=1\}} \quad (7)$$

$$720 \quad (8)$$

> HTS(cos(n*Pi/4)^2, n)

$$\frac{1}{2} + \frac{(-1)^{\frac{n}{2}} \chi_{\{modp(n,2)=0\}}}{2} \quad (9)$$

Based on Euler's formula, a computer-assisted computation can obtain an equivalent formula for (9). Mathematica finds a representation over \mathbb{C} without interlacement. Unlike our method, this limited strategy may require different *simplifications* for various trigonometric inputs.

```
> RE:= (n + 1)*(n^6 + 14*n^5 + 35*n^4 - 350*n^3 - 2436*n^2 - 5545*n - 4319)*(n -
      2)*a(n) - (n - 1)*(n^6 + 8*n^5 - 20*n^4 - 370*n^3 - 1301*n^2 - 1799*n - 838)*a
      (n + 1) - (n + 1)*(n - 7)*(n^6 + 14*n^5 + 35*n^4 - 350*n^3 - 2436*n^2 - 5545*n
      - 4319)*a(n + 5) + (n^6 + 8*n^5 - 20*n^4 - 370*n^3 - 1301*n^2 - 1799*n - 838)
      *(n - 6)*a(n + 6) = 0:
> L := [1, 7, 2, 6, 24, 120, 721]:
> REtoHTS(RE, a(n), L)
```

$$n! - (n - 7) \chi_{\{modp(n,5)=1\}} \quad (10)$$

■

The commands HTS and REtoHTS rely on the maple command LREtools:-mhypergeomsols from FPS [7], which implements the m -fold hypergeometric term solver from [14], an extension of the algorithms by Petkovšek and van Hoeij [8] [5]. Of course, HTS also uses HolonomicRE to convert expressions into P-recursive equations.

In Section 2, we describe the algorithm used by HolonomicRE to compute holonomic recurrence equations from terms in \mathcal{H}_T . Section 3 presents HTSproduct, the implementation of the product rule in \mathcal{H}_T , i.e, Hadamard product rule in (\mathcal{H}_T) . All computations of this paper can be done with Maple versions 2019-2023.

2 Recurrence Equations

The proof of [13, Proposition 3] is constructive and can be turned into the following algorithm.

Algorithm 1 P-recursive equations for terms in \mathcal{H}_T

Input: $s_n := \sum_{i=1}^l H_i(\sigma_i(n)) \chi_{\{n \equiv j_i \pmod{m_i}\}} \in \mathcal{H}_T$.

Output: A P-recursive equation satisfied by s_n .

1. Set $u_i(m_i n + j_i) := H_i(\sigma_i(m_i n + j_i))$, $i = 1, \dots, l$.
 2. Find a P-recursive m_i -shift recurrence equation satisfied by $u_i(m_i n + j_i)$, $i = 1, \dots, l$.
 3. Deduce a P-recursive equation satisfied by $u_i(n)$ by the substitution $n \rightarrow \frac{n-j_i}{m_i}$, $i = 1, \dots, l$.
 4. Return the P-recursive equation obtained by applying the P-recursive addition algorithm to $\sum_{i=1}^l u_i$.
-

Remark 2.

- If we denote by k_i the number of hypergeometric terms in the linear combinations of the coefficient H_i in s_n , $i = 1, \dots, l$, then Algorithm 1 returns a recurrence equation of order at most $\sum_{i=1}^l m_i k_i$.
- We mention that Algorithm 1 can be efficiently implemented in Maple without using loops. See the recommendations in [4].

- Algorithm 1, step 2 can be done using [12, Section 2] for finding recurrence equations satisfied by linear combinations of hypergeometric terms.
- Algorithm 1, step 2 may recursively apply the addition algorithm from the GFUN package [9]. However, the subpackage `AlgebraHolonomicSeq` of `HyperTypeSeq` provides the command `AddHolonomicRE` that computes a holonomic recurrence equation for linear combinations of holonomic terms from arbitrarily many holonomic equations. This procedure is not recursive; it uses a linear dynamical system defined by the equations and proceeds like our standard `HolonomicRE` to find the desired equation.

Example 3.

```
> with(HyperTypeSeq):
> HolonomicRE(3^n*mfoldInd(n,3,1)+(2^n+n)*mfoldInd(n,2,0),a(n))
```

$$\begin{aligned} & (-353808n^2 - 586764n - 150444) a(n) - 78732 a(n+1) \\ & + (442260n^2 + 25839n - 44901) a(n+2) + (13104n^2 + 21732n + 25255) a(n+3) \\ & + (-88452n^2 + 30213n + 23544) a(n+4) + (-16380n^2 - 957n + 1663) a(n+5) \\ & - 729 a(n+6) + (3276n^2 - 1119n - 764) a(n+7) = 0 \end{aligned} \quad (11)$$

```
> HolonomicRE(n!*mfoldInd(n,4,3) + pochhammer(2,n),a(n))
```

$$\begin{aligned} & (n+4)(n+3)(n+2)(n+1)(n+5)a(n) \\ & - (n+5)(n+4)(n+3)(n+2)a(n+1) + (-n-5)a(n+4) + a(n+5) = 0 \end{aligned} \quad (12)$$

■

Now that `HolonomicRE` can compute recurrence equations for hypergeometric-type terms in normal forms, we can automatically solve the zero-equivalence problem in (\mathcal{H}_T) . We will give an interesting example in the next section.

3 Products

The product algorithm is essentially based on the following proposition, which can be seen as a consequence of the *Chinese Remainder Theorem*.

Proposition 1 (see **Lemma 2** in [13]). *Let $j_1, j_2, m_1, m_2 \in \mathbb{N}$ such that $j_1 < m_1, j_2 < m_2$. Let $\mu = \text{lcm}(m_1, m_2)$ and*

$$\mathcal{N} = \{j < \mu, j = j_1 + m_1 k_1 = j_2 + m_2 k_2, k_1, k_2 \in \mathbb{N}\}.$$

Then $\mathcal{N} = \{j_0\}$ or $\mathcal{N} = \emptyset$ and we have

$$\chi_{\{n \equiv j_1 \pmod{m_1}\}} \cdot \chi_{\{n \equiv j_2 \pmod{m_2}\}} = \begin{cases} \chi_{\{n \equiv j_0 \pmod{\mu}\}}, & \text{if } \mathcal{N} = \{j_0\} \\ 0 = \chi_{\{n \equiv 0 \pmod{0}\}}, & \text{if } \mathcal{N} = \emptyset \end{cases}. \quad (13)$$

An interesting fact resulting from this proposition is that the product of m -fold indicator terms of the same remainder is an m -fold indicator term with that same remainder; in particular, an m -fold indicator term is unchanged by exponentiation.

On account of Proposition 1, one can easily implement an algorithm for computing products of hypergeometric-type terms. Our implementation is the command `HTSproduct` from `HyperTypeSeq`.

Example 4.

```
> with(HyperTypeSeq):
> u1 := (n!)^2*mfoldInd(n, 3, 1) + n^3*mfoldInd(n, 2, 1);
u2 := (n + 1)*mfoldInd(n, 4, 3)/n! + (n + 2)*mfoldInd(n, 2, 0)
```

$$u1 := n!^2 \chi_{\{\text{mod}p(n,3)=1\}} + n^3 \chi_{\{\text{mod}p(n,2)=1\}} \quad (14)$$

$$u2 := \frac{(n+1) \chi_{\{\text{mod}p(n,4)=3\}}}{n!} + (n+2) \chi_{\{\text{mod}p(n,2)=0\}} \quad (15)$$

```
> HTSproduct(u1, u2, n)
```

$$\frac{n^3 (n+1) \chi_{\{\text{mod}p(n,4)=3\}}}{n!} + n!^2 (n+2) \chi_{\{\text{mod}p(n,6)=4\}} + n! (n+1) \chi_{\{\text{mod}p(n,12)=7\}} \quad (16)$$

■

We gave some arguments in [13, Section 3.1] on the reasons why canonical forms are difficult to define in \mathcal{H}_T . However, the algorithm implemented by HTS computes normal forms [3, Chapter 3]. This implies that we can recognize equivalent formulas. Indeed, for any two equivalent hypergeometric-type terms in \mathcal{H}_T , the normal form returned by HTS for their difference is zero.

Example 5. It is known that trigonometric and elliptic functions are good candidates to generate hypergeometric-type terms [2]. We consider $s_n := \sin\left(\frac{n\pi}{3}\right)^2 \cos\left(\frac{n\pi}{4}\right)^2$, and we use HTS and HTSproduct to compute two equivalent representations, here $\mathbf{s1}$ and $\mathbf{s2}$, respectively. Finally we verify that HTS($\mathbf{s1} - \mathbf{s2}, n$) is zero.

```
> with(HyperTypeSeq):
> s1 := HTS(sin(n*Pi/3)^2*cos(n*Pi/4)^2, n)
```

$$s1 := \frac{3}{8} + \frac{3(-1)^{\frac{n}{2}} \chi_{\{\text{mod}p(n,2)=0\}}}{8} - \frac{3\chi_{\{\text{mod}p(n,3)=0\}}}{8} - \frac{3(-1)^{\frac{n}{6}} \chi_{\{\text{mod}p(n,6)=0\}}}{8} \quad (17)$$

```
> u1 := HTS(sin(n*Pi/3)^2, n)
```

$$u1 := \frac{3\chi_{\{\text{mod}p(n,3)=1\}}}{4} + \frac{3\chi_{\{\text{mod}p(n,3)=2\}}}{4} \quad (18)$$

```
> u2 := HTS(cos(n*Pi/4)^2, n)
```

$$u2 := \frac{1}{2} + \frac{(-1)^{\frac{n}{2}} \chi_{\{\text{mod}p(n,2)=0\}}}{2} \quad (19)$$

```
> s2 := HTSproduct(u1, u2, n)
```

$$s2 := \frac{3\chi_{\{\text{mod}p(n,3)=1\}}}{8} + \frac{3\chi_{\{\text{mod}p(n,3)=2\}}}{8} + \frac{3(-1)^{\frac{n}{2}} \chi_{\{\text{mod}p(n,6)=4\}}}{8} + \frac{3(-1)^{\frac{n}{2}} \chi_{\{\text{mod}p(n,6)=2\}}}{8} \quad (20)$$

```
> HTS(s1-s2, n)
```

$$0 \quad (21)$$

■

Acknowledgment. The author was supported by UKRI Frontier Research Grant EP/X033813/1.

References

1. Abramov, S.A.: m-sparse solutions of linear ordinary differential equations with polynomial coefficients. *Discrete Mathematics* **217**(1-3), 3–15 (2000) [1](#)
2. Dreyfus, T.: Hypertranscendence and linear difference equations, the exponential case (2022), preprint [arXiv:2212.00388](https://arxiv.org/abs/2212.00388) [5](#)
3. Geddes, K.O., Czapor, S.R., Labahn, G.: *Algorithms for Computer Algebra*. Springer Science & Business Media (1992) [5](#)
4. Gerhard, J.: Maple conference 2019 - Maple programming: tips and tricks (2019), available at <https://www.youtube.com/watch?v=cUVbMOTQzRA> [3](#)
5. van Hoeij, M.: Finite singularities and hypergeometric solutions of linear recurrence equations. *Journal of Pure and Applied Algebra* **139**(1-3), 109–131 (1999) [3](#)
6. Kauers, M.: *D-Finite Functions*. Springer Cham (2023) [2](#)
7. Koepf, W., Tegua Tabugua, B.: FPS: A Maple package for symbolic computations with power series and linear PDEs. Available at http://www.mathematik.uni-kassel.de/~btegua/FPS_webpage/FPS.htm (2022) [2](#), [3](#)
8. Petkovšek, M.: Hypergeometric solutions of linear recurrences with polynomial coefficients. *Journal of symbolic computation* **14**(2-3), 243–264 (1992) [3](#)
9. Salvy, B., Zimmermann, P.: GFUN: a maple package for the manipulation of generating and holonomic functions in one variable. *ACM Transactions on Mathematical Software (TOMS)* **20**(2), 163–177 (1994) [4](#)
10. Sloane, N.J., et al.: *The online encyclopedia of integer sequences* (2003), website: <https://oeis.org> [2](#)
11. Tegua Tabugua, B.: HyperTypeSeq: A Maple package for symbolic computations with hypergeometric-type sequences. Available at <https://github.com/T3gua/HyperTypeSeq> [2](#)
12. Tegua Tabugua, B.: A variant of van Hoeij’s algorithm to compute hypergeometric term solutions of holonomic recurrence equations. *Journal of Algorithms and Computation* **53**(2), 1–32 (2021) [4](#)
13. Tegua Tabugua, B.: Hypergeometric-type sequences. To appear in the *Journal of Symbolic Computation* (special issue in memory of Marko Petkovšek) (2023) [1](#), [2](#), [3](#), [4](#), [5](#)
14. Tegua Tabugua, B., Koepf, W.: Symbolic conversion of holonomic functions to hypergeometric-type power series. *Programming and Computer Software* **48**(2), 125–146 (2022) [3](#)