

# An upper bound on the number of relevant variables for Boolean functions on the Hamming graph

Alexandr Valyuzhenich\*

## Abstract

The spectrum of a complex-valued function  $f$  on  $\mathbb{Z}_q^n$  is the set  $\{|u| : u \in \mathbb{Z}_q^n \text{ and } \widehat{f}(u) \neq 0\}$ , where  $|u|$  is the Hamming weight of  $u$  and  $\widehat{f}$  is the Fourier transform of  $f$ . Let  $1 \leq d' \leq d \leq n$ . In this work, we study Boolean functions on  $\mathbb{Z}_q^n$ ,  $q \geq 3$ , whose spectrum is a subset of  $\{0\} \cup \{d', \dots, d\}$ . We prove that such functions have at most  $\frac{d}{2} \cdot \frac{q^{d+d'}}{2^{d'(q-1)^{d'}}$  relevant variables for  $d' + d \leq n + 1$ . In particular, we prove that any Boolean function of degree  $d$  on  $\mathbb{Z}_q^n$ ,  $q \geq 3$ , has at most  $\frac{dq^{d+1}}{4(q-1)}$  relevant variables. We also show that any equitable 2-partition of the Hamming graph  $H(n, q)$ ,  $q \geq 3$ , associated with the eigenvalue  $n(q-1) - qd$  has at most  $\frac{d}{2} \cdot \frac{q^{2d}}{2^{d(q-1)^d}$  relevant variables for  $d \leq \frac{n+1}{2}$ .

## 1 Introduction

The *Hamming graph*  $H(n, q)$  is defined as follows. The vertex set of  $H(n, q)$  is  $\mathbb{Z}_q^n$ , and two vertices are adjacent if they differ in exactly one position. The *spectrum* of a function  $f : \mathbb{Z}_q^n \rightarrow \mathbb{C}$  is the set  $\{|u| : u \in \mathbb{Z}_q^n \text{ and } \widehat{f}(u) \neq 0\}$ , where  $|u|$  is the Hamming weight of  $u$  and  $\widehat{f}$  is the Fourier transform of  $f$ . The *degree* of a function  $f : \mathbb{Z}_q^n \rightarrow \mathbb{C}$  is the largest element of the spectrum of  $f$ . Given a function  $f$  on  $\mathbb{Z}_q^n$ , a variable  $x_i$ ,  $1 \leq i \leq n$ , is called *relevant* (or *essential*) if there exist  $a_1, \dots, a_{i-1}, a_{i+1}, \dots, a_n \in \mathbb{Z}_q$  and  $b_1, b_2 \in \mathbb{Z}_q$  such that

$$f(a_1, \dots, a_{i-1}, b_1, a_{i+1}, \dots, a_n) \neq f(a_1, \dots, a_{i-1}, b_2, a_{i+1}, \dots, a_n).$$

A classical theorem of Nisan and Szegedy [13] states that any Boolean function of degree  $d$  on  $\mathbb{Z}_2^n$  has at most  $d \cdot 2^{d-1}$  relevant variables. In 2020, Chiarelli, Hatami and Saks [2] improved the bound to  $6.614 \cdot 2^d$ , and then Wellens [15] improved it further to  $4.394 \cdot 2^d$ . The same problem for Boolean functions of degree  $d$  on the Johnson graph was studied in [3, 5].

---

\*Postdoctoral Research Station of Mathematics, Hebei Normal University, Shijiazhuang 050024, PR China; Email address: graphkiper@mail.ru

In this work, we consider Boolean functions of a bounded degree on  $\mathbb{Z}_q^n$  (or, equivalently, on the Hamming graph  $H(n, q)$ ) for  $q \geq 3$ . Any Boolean function of degree 1 on  $\mathbb{Z}_q^n$  has at most one relevant variable. This fact is a very special case of the result obtained by Meyerowitz in [10]. Another proof of this fact can be found in [4]. Boolean functions of degree 2 on  $\mathbb{Z}_q^n$  whose spectrum is  $\{0, 2\}$  were studied in [11, 12]. In particular, Mogilnykh and Valyuzhenich [11] showed that such functions have at most  $\max(4, \frac{q}{2} + 1)$  relevant variables. The remark of Filmus and Ihringer at the end of [3] and Wellens's result imply that any Boolean function of degree  $d$  on  $\mathbb{Z}_q^n$  has at most  $4.394 \cdot 2^{\lceil \log_2 q \rceil d}$  relevant variables.

Let  $1 \leq d' \leq d \leq n$ . In this work, we study Boolean functions on  $\mathbb{Z}_q^n$ ,  $q \geq 3$ , whose spectrum is a subset of  $\{0\} \cup \{d', \dots, d\}$ . We prove that such functions have at most  $\frac{d}{2} \cdot \frac{q^{d+d'}}{2^{d'(q-1)^{d'}}$  relevant variables for  $d' + d \leq n + 1$ . In particular, we prove that any Boolean function of degree  $d$  on  $\mathbb{Z}_q^n$ ,  $q \geq 3$ , has at most  $\frac{dq^{d+1}}{4(q-1)}$  relevant variables. This bound improves the bound  $4.394 \cdot 2^{\lceil \log_2 q \rceil d}$  in the following cases:

- $q \in \{3, 5, 6, 7\}$ ;
- $2^{k-1} < q \leq 15 \cdot 2^{k-4}$ , where  $k \geq 4$ .

Boolean functions on  $\mathbb{Z}_q^n$  whose spectrum is  $\{0, d\}$  are in one-to-one correspondence with equitable 2-partitions of  $H(n, q)$  associated with the eigenvalue  $n(q-1) - qd$ . Therefore, our result for  $d' = d$  implies that any equitable 2-partition of  $H(n, q)$ ,  $q \geq 3$ , associated with the eigenvalue  $n(q-1) - qd$  has at most  $\frac{d}{2} \cdot \frac{q^{2d}}{2^{d(q-1)^d}$  relevant variables for  $d \leq \frac{n+1}{2}$ .

The paper is organized as follows. In Section 2, we introduce basic definitions. In Section 3, we give preliminary results. In Section 4, we prove the main results.

## 2 Basic definitions

For a positive integer  $n$ , denote  $[n] = \{1, \dots, n\}$ . The weight of a vector  $u \in \mathbb{Z}_q^n$ , denoted by  $|u|$ , is the number of its non-zero coordinates. The support of a function  $f$  is denoted by  $S(f)$ .

The *Hamming graph*  $H(n, q)$  is defined as follows. The vertex set of  $H(n, q)$  is  $\mathbb{Z}_q^n$ , and two vertices are adjacent if they differ in exactly one coordinate. This graph has  $n+1$  distinct eigenvalues  $\lambda_k(n, q) = n(q-1) - q \cdot k$ , where  $0 \leq k \leq n$ . Denote by  $E_{n,q}$  and  $A_{n,q}$  the edge set and the adjacency operator of  $H(n, q)$ .

The set of all complex-valued functions on  $\mathbb{Z}_q^n$ , denoted by  $U(n, q)$ , forms a vector space over  $\mathbb{C}$ . Define an inner product on this vector space as follows:

$$\langle f, g \rangle = \frac{1}{q^n} \sum_{x \in \mathbb{Z}_q^n} f(x) \overline{g(x)}.$$

For every  $0 \leq k \leq n$ , we denote

$$U_k(n, q) = \{f : \mathbb{Z}_q^n \rightarrow \mathbb{C} : A_{n,q}f = \lambda_k(n, q)f\}.$$

In fact,  $U_k(n, q)$  is a complex eigenspace of  $A_{n,q}$  associated with  $\lambda_k(n, q)$ . The direct sum of subspaces

$$U_k(n, q) \oplus U_{k+1}(n, q) \oplus \dots \oplus U_m(n, q),$$

where  $0 \leq k \leq m \leq n$ , is denoted by  $U_{[k,m]}(n, q)$ .

For every  $u \in \mathbb{Z}_q^n$ , define a *character*  $\chi_u$  on  $\mathbb{Z}_q^n$  by  $\chi_u(x) = \omega^{\langle u, x \rangle}$ , where  $\omega = e^{2\pi i/q}$  is a primitive  $q$ th root of unity and  $\langle u, x \rangle = u_1x_1 + \dots + u_nx_n$ . In what follows, we will need the following two properties of the characters.

**Remark 1.** *The following statements hold:*

1. *The set  $\{\chi_u : u \in \mathbb{Z}_q^n\}$  forms an orthonormal basis of the vector space  $U(n, q)$ .*
2. *The equality  $A_{n,q}\chi_u = \lambda_u\chi_u$ , where  $\lambda_u = n(q-1) - q \cdot |u|$ , holds for any vector  $u \in \mathbb{Z}_q^n$ .*

Every function  $f \in U(n, q)$  is uniquely expressible as

$$f = \sum_{u \in \mathbb{Z}_q^n} \widehat{f}(u)\chi_u,$$

where  $\widehat{f}(u) \in \mathbb{C}$  for all  $u \in \mathbb{Z}_q^n$ . The complex numbers  $\widehat{f}(u)$ , where  $u \in \mathbb{Z}_q^n$ , are called the *Fourier coefficients* of  $f$ . The *spectrum* of a function  $f \in U(n, q)$  is the set  $\{|u| : u \in \mathbb{Z}_q^n \text{ and } \widehat{f}(u) \neq 0\}$ . The *degree* of a function  $f \in U(n, q)$  is  $\max\{|u| : u \in \mathbb{Z}_q^n \text{ and } \widehat{f}(u) \neq 0\}$ .

Let  $f \in U(n, q)$ , let  $i \in [n]$  and let  $a, b \in \mathbb{Z}_q$ , where  $a \neq b$ . We define a function  $f_{i,a,b}$  on  $\mathbb{Z}_q^{n-1}$  as follows: for any vector  $y = (y_1, \dots, y_{i-1}, y_{i+1}, \dots, y_n)$  from  $\mathbb{Z}_q^{n-1}$

$$f_{i,a,b}(y) = f(y_1, \dots, y_{i-1}, a, y_{i+1}, \dots, y_n) - f(y_1, \dots, y_{i-1}, b, y_{i+1}, \dots, y_n).$$

Let  $f \in U(n, q)$ , let  $i \in [n]$  and let  $a, b \in \mathbb{Z}_q$ , where  $a \neq b$ . Denote by  $\nu(f)$  the number of edges  $\{x, y\} \in E_{n,q}$  such that  $f(x) \neq f(y)$ ; and denote by  $\nu_{i,a,b}(f)$  the number of edges  $\{x, y\} \in E_{n,q}$  such that  $x_i = a$ ,  $y_i = b$  and  $f(x) \neq f(y)$ .

Note that

$$\nu(f) = \sum_{\substack{i \in [n] \\ a, b \in \mathbb{Z}_q: a < b}} \nu_{i,a,b}(f).$$

On the other hand, we have  $\nu_{i,a,b}(f) = |S(f_{i,a,b})|$ . Therefore,

$$\nu(f) = \sum_{\substack{i \in [n] \\ a, b \in \mathbb{Z}_q: a < b}} |S(f_{i,a,b})|. \quad (1)$$

Given a function  $f$  on  $\mathbb{Z}_q^n$ , a variable  $x_i$ ,  $i \in [n]$ , is called *relevant* if there exist  $a_1, \dots, a_{i-1}, a_{i+1}, \dots, a_n \in \mathbb{Z}_q$  and  $b_1, b_2 \in \mathbb{Z}_q$  such that

$$f(a_1, \dots, a_{i-1}, b_1, a_{i+1}, \dots, a_n) \neq f(a_1, \dots, a_{i-1}, b_2, a_{i+1}, \dots, a_n).$$

Given a function  $f$  on  $\mathbb{Z}_q^n$ , an index  $i \in [n]$  is called *relevant* if the corresponding variable  $x_i$  is relevant.

### 3 Preliminaries

The following two formulas can be proved by standard methods (the proof for  $q = 2$  can be found in [13]). We give the proof for completeness.

**Lemma 1.** *For any function  $f : \mathbb{Z}_q^n \rightarrow \{-1, 1\}$ , the following equalities hold:*

$$\sum_{u \in \mathbb{Z}_q^n} |\widehat{f}(u)|^2 = 1, \quad (2)$$

$$\nu(f) = \frac{q^{n+1}}{4} \sum_{u \in \mathbb{Z}_q^n} |u| \cdot |\widehat{f}(u)|^2. \quad (3)$$

*Proof.* By Remark 1, we have

$$\langle f, f \rangle = \left\langle \sum_{u \in \mathbb{Z}_q^n} \widehat{f}(u) \chi_u, \sum_{u \in \mathbb{Z}_q^n} \widehat{f}(u) \chi_u \right\rangle = \sum_{u \in \mathbb{Z}_q^n} |\widehat{f}(u)|^2.$$

On the other hand,

$$\langle f, f \rangle = \frac{1}{q^n} \sum_{x \in \mathbb{Z}_q^n} |f(x)|^2 = 1.$$

This immediately implies the equality (2). Using Remark 1, we obtain

$$\begin{aligned} \langle A_{n,q} f, f \rangle &= \left\langle \sum_{u \in \mathbb{Z}_q^n} \lambda_u \widehat{f}(u) \chi_u, \sum_{u \in \mathbb{Z}_q^n} \widehat{f}(u) \chi_u \right\rangle = \sum_{u \in \mathbb{Z}_q^n} \lambda_u \cdot |\widehat{f}(u)|^2 = \\ &= n(q-1) - q \sum_{u \in \mathbb{Z}_q^n} |u| \cdot |\widehat{f}(u)|^2. \end{aligned} \quad (4)$$

On the other hand,

$$\langle A_{n,q} f, f \rangle = \frac{2}{q^n} \sum_{\{x,y\} \in E_{n,q}} f(x)f(y) = n(q-1) - \frac{4\nu(f)}{q^n}. \quad (5)$$

Subtracting (5) from (4), we get the equality (3).  $\square$

Using Lemma 1 for Boolean functions of degree at most  $d$ , we immediately obtain the following result.

**Corollary 1.** *Let  $f$  be a Boolean function of degree at most  $d$  on  $\mathbb{Z}_q^n$ . Then  $\nu(f) \leq \frac{d}{4} \cdot q^{n+1}$ .*

The following lemma shows the connection between the eigenspaces of the Hamming graphs  $H(n, q)$  and  $H(n-1, q)$ .

**Lemma 2.** *Let  $f \in U_{[k, m]}(n, q)$ , let  $i \in [n]$ , and let  $a, b \in \mathbb{Z}_q$ ,  $a \neq b$ . Then  $f_{i, a, b} \in U_{[k-1, m-1]}(n-1, q)$ .*

The proof of Lemma 2 can be found, for example, in [14, Lemma 4]. The following lower bound on the support size of functions on  $\mathbb{Z}_q^n$  was proved in [14].

**Theorem 1.** *Let  $f \in U_{[k, m]}(n, q)$ , where  $q \geq 3$ ,  $k + m \leq n$  and  $f \not\equiv 0$ . Then*

$$|S(f)| \geq 2^k \cdot (q-1)^k \cdot q^{n-k-m}.$$

We note that the bound from Theorem 1 is sharp.

## 4 Main results

The main result of this paper is the following.

**Theorem 2.** *Let  $f \in U_0(n, q) \oplus U_{[d', d]}(n, q)$ , where  $q \geq 3$ ,  $1 \leq d' \leq d \leq n$  and  $d' + d \leq n + 1$ . If  $f$  is Boolean, then  $f$  has at most  $\frac{d}{2} \cdot \frac{q^{d+d'}}{2^{d'(q-1)^{d'}}$  relevant variables.*

*Proof.* Suppose  $f$  has  $n'$  relevant variables. Let us consider a relevant index  $i \in [n]$ . Define a partition  $A_1 \cup \dots \cup A_t$  of  $\mathbb{Z}_q$  as follows:  $a, b \in A_r$  for some  $r \in [t]$  if and only if  $f_{i, a, b} \equiv 0$ .

Since  $i$  is relevant, we have  $t \geq 2$ . For every  $r \in [t]$ , denote  $q_r = |A_r|$ . Note that

$$|\{(a, b) : a, b \in \mathbb{Z}_q, a < b, f_{i, a, b} \not\equiv 0\}| = \sum_{1 \leq r < s \leq t} q_r q_s \geq q_1(q_2 + \dots + q_t) \geq q - 1. \quad (6)$$

By Lemma 2, we have  $f_{i, a, b} \in U_{[d'-1, d-1]}(n-1, q)$  for all  $a, b \in \mathbb{Z}_q$ , where  $a \neq b$ . Applying Theorem 1 for  $f_{i, a, b}$ , where  $a \neq b$ , we get

$$|S(f_{i, a, b})| \geq 2^{d'-1} \cdot (q-1)^{d'-1} \cdot q^{n-d'-d+1} \quad (7)$$

or  $f_{i, a, b} \equiv 0$ . Combining (6) and (7), we obtain that the inequality

$$\sum_{a, b \in \mathbb{Z}_q : a < b} |S(f_{i, a, b})| \geq 2^{d'-1} \cdot (q-1)^{d'} \cdot q^{n-d'-d+1} \quad (8)$$

holds for any relevant index  $i \in [n]$ .

Using the equality (1) and the inequality (8), we obtain the following lower bound on  $\nu(f)$ :

$$\nu(f) \geq n' \cdot 2^{d'-1} \cdot (q-1)^{d'} \cdot q^{n-d'-d+1}.$$

On the other hand, we have  $\nu(f) \leq \frac{d}{4} \cdot q^{n+1}$  due to Corollary 1. Therefore,

$$n' \leq \frac{d}{2} \cdot \frac{q^{d+d'}}{2^{d'}(q-1)^{d'}}.$$

□

Applying Theorem 2 for  $d' = 1$  and  $d' = d$  respectively, we obtain the following results.

**Corollary 2.** *Any Boolean function of degree  $d$  on  $\mathbb{Z}_q^n$ ,  $q \geq 3$ , has at most  $\frac{dq^{d+1}}{4(q-1)}$  relevant variables.*

**Corollary 3.** *Let  $f \in U_0(n, q) \oplus U_d(n, q)$ , where  $q \geq 3$  and  $1 \leq d \leq \frac{n+1}{2}$ . If  $f$  is Boolean, then  $f$  has at most  $\frac{d}{2} \cdot \frac{q^{2d}}{2^d(q-1)^d}$  relevant variables.*

Corollary 3 can be reformulated in terms of equitable 2-partitions of  $H(n, q)$ . Firstly, we recall some definitions. Let  $G$  be a graph. An  $r$ -partition  $C_1 \cup \dots \cup C_r$  of the vertex set of  $G$  is called *equitable* if for any  $i, j \in [r]$  there is a constant  $s_{i,j}$  such that any vertex of  $C_i$  has exactly  $s_{i,j}$  neighbors in  $C_j$ . The matrix  $S = (s_{i,j})_{1 \leq i, j \leq r}$  is the *quotient matrix* of the partition. One of the important properties of equitable partitions is the following: the eigenvalues of the quotient matrix of an equitable partition of a graph  $G$  are also eigenvalues of the adjacency matrix of  $G$  (see, for example, [8, Lemma 2.2]). There are many different names in the literature for equitable partitions of graphs: perfect coloring, partition design, regular partition. We also note that every class of an equitable 2-partition of a graph is a completely regular code of radius one. An interesting discussion of equitable partitions and related concepts can be found in [9].

Fon-Der-Flaass initiated the systematic study of equitable 2-partitions of the Hamming graph in [6, 7] (he considered the case  $q = 2$ ). In particular, he proposed several iterative constructions of equitable 2-partitions of  $H(n, 2)$ . He also proved several important necessary conditions for the existence of equitable 2-partitions of  $H(n, 2)$  with a given quotient matrix. For more information on equitable 2-partitions of the Hamming graphs we refer the reader to a recent survey [1]. Suppose  $C_1 \cup C_2$  is an equitable 2-partition of  $H(n, q)$  with the quotient matrix

$$S = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

The matrix  $S$  has two different eigenvalues: the *trivial* eigenvalue  $\theta_0 = n(q-1)$  (the degree of  $H(n, q)$ ) and the *second* eigenvalue  $\theta_1 = a - c$ . Moreover, we

have  $\theta_1 = \lambda_k(n, q)$  for some  $0 \leq k \leq n$ . We say that an equitable 2-partition of  $H(n, q)$  has *degree*  $d$  if the second eigenvalue of its quotient matrix is  $\lambda_d(n, q)$ .

Suppose that  $C_1 \cup C_2$  is an equitable 2-partition of  $H(n, q)$  of degree  $d$ . We have  $\mathbf{1}_{C_1} \in U_0(n, q) \oplus U_d(n, q)$ , where  $\mathbf{1}_{C_1}$  is the characteristic function of  $C_1$  in  $\mathbb{Z}_q^n$ . Applying Corollary 3 to the function  $\mathbf{1}_{C_1}$ , we get the following.

**Corollary 4.** *Let  $q \geq 3$  and  $1 \leq d \leq \frac{n+1}{2}$ . Then any equitable 2-partition of  $H(n, q)$  of degree  $d$  has at most  $\frac{d}{2} \cdot \frac{q^{2d}}{2^d(q-1)^d}$  relevant variables.*

Finally, we compare the bounds from Theorem 2, Corollary 2 and Corollary 3 with the bound  $4.394 \cdot 2^{\lceil \log_2 q \rceil d}$ :

1. If  $d' > \frac{d}{8.788} \cdot \frac{q}{q-2}$ , then the bound from Theorem 2 is better than  $4.394 \cdot 2^{\lceil \log_2 q \rceil d}$  for all  $q \geq 3$  (we assume that  $d' + d \leq n + 1$ ).
2. The bound from Corollary 2 is better than  $4.394 \cdot 2^{\lceil \log_2 q \rceil d}$  in the following cases:
  - $q \in \{3, 5, 6, 7\}$ ;
  - $2^{k-1} < q \leq 15 \cdot 2^{k-4}$ , where  $k \geq 4$ .
3. The bound from Corollary 3 is better than  $4.394 \cdot 2^{\lceil \log_2 q \rceil d}$  for all  $q \geq 3$  (we assume that  $d \leq \frac{n+1}{2}$ ).

## References

- [1] E. Beshpalov, D. Krotov, A. Matushev, A. Taranenko, K. Vorob'ev, Perfect 2-colorings of Hamming graphs, *Journal of Combinatorial Designs* 29(6) (2021) 367–396.
- [2] J. Chiarelli, P. Hatami, M. Saks, An asymptotically tight bound on the number of relevant variables in a bounded degree Boolean function, *Combinatorica* 40 (2020) 237–244.
- [3] Y. Filmus, F. Ihringer, Boolean constant degree functions on the slice are juntas, *Discrete Mathematics* 342(12) (2019) 111614.
- [4] Y. Filmus, F. Ihringer, Boolean degree 1 functions on some classical association schemes, *Journal of Combinatorial Theory, Series A* 162 (2019) 241–270.
- [5] Y. Filmus, Junta threshold for low degree Boolean functions on the slice, *The Electronic Journal of Combinatorics*, 30(1) (2023) #P1.55.
- [6] D. G. Fon-Der-Flaass, Perfect 2-colorings of a hypercube, *Siberian Mathematical Journal* 48(4) (2007) 740–745.

- [7] D. G. Fon-Der-Flaass, A bound on correlation immunity, *Sib. Elektron. Mat. Izv.* 4 (2007) 133–135.
- [8] C. D. Godsil, *Algebraic Combinatorics*, Chapman and Hall Mathematics Series, Chapman Hall, New York, 1993.
- [9] F. Ihringer, Translating terminology: equitable partitions and related concepts, blog entry: <https://ratiobound.wordpress.com/2018/12/02/translating-terminology-equitable-partition>
- [10] A. Meyerowitz, Cycle-balance conditions for distance-regular graphs, *Discrete Mathematics* 264(1–3) (2003) 149–165.
- [11] I. Mogilnykh, A. Valyuzhenich, Equitable 2-partitions of the Hamming graphs with the second eigenvalue, *Discrete Mathematics* 343(11) (2020) 112039.
- [12] I. Mogilnykh, A. Taranenko, K. Vorob'ev, Completely regular codes with covering radius 1 and the second eigenvalue in 3-dimensional Hamming graphs, arXiv:2403.02702, March 2024.
- [13] N. Nisan, M. Szegedy, On the degree of Boolean functions as real polynomials, *Computational Complexity* 4 (1994) 301–313.
- [14] A. Valyuzhenich, K. Vorob'ev, Minimum supports of functions on the Hamming graphs with spectral constraints, *Discrete Mathematics* 342(5) (2019) 1351–1360.
- [15] J. Wellens, Relationships between the number of inputs and other complexity measures of Boolean functions, arXiv:2005.00566v2, 2022.