

INTEGER-VALUED O-MINIMAL FUNCTIONS

NEER BHARDWAJ, RAYMOND MCCULLOCH¹, NANDAGOPAL RAMACHANDRAN,
AND KATHARINE WOO²

ABSTRACT. We study $\mathbb{R}_{\text{an,exp}}$ -definable functions $f : \mathbb{R} \rightarrow \mathbb{R}$ that take integer values at all sufficiently large positive integers. If $|f(x)| = O(2^{(1+10^{-5})x})$, then we find polynomials P_1, P_2 such that $f(x) = P_1(x) + P_2(x)2^x$ for all sufficiently large x . Our result parallels classical theorems of Pólya and Selberg for entire functions and generalizes Wilkie’s classification for the case of $|f(x)| = O(C^x)$, for some $C < 2$.

Let $k \in \mathbb{N}$ and $\gamma_k = \sum_{j=1}^k 1/j$. Extending Wilkie’s theorem in a separate direction, we show that if f is k -concordant and $|f(x)| = O(C^x)$, for some $C < e^{\gamma_k} + 1$, then f must eventually be given by a polynomial. This is an analog of a result by Pila for entire functions.

1. INTRODUCTION

In [13], Wilkie studies functions $f : \mathbb{R} \rightarrow \mathbb{R}$ definable in the o-minimal structure $\mathbb{R}_{\text{an,exp}}$ with the property that $f(a) \in \mathbb{Z}$ for all sufficiently large positive integers a . If there is a constant $C < 2$ such that $|f(x)| \leq C^x$ for all sufficiently large x , then [13, Theorem 5.1] shows that f is eventually given by a polynomial. This parallels a special case of the following seminal theorem by Pólya for entire functions [10].

Theorem 1.1. *Let $g : \mathbb{C} \rightarrow \mathbb{C}$ be an entire function which satisfies $g(a) \in \mathbb{Z}$ for all positive integers a . Suppose there is $M \in \mathbb{N}$ such that*

$$|g(z)| = O(|z|^M 2^{|z|}) \quad \text{as } |z| \rightarrow \infty.$$

Then there are polynomials P_1, P_2 such that $g(z) = P_1(z) + P_2(z)2^z$ for all $z \in \mathbb{C}$.

We show that a similar characterization holds for $\mathbb{R}_{\text{an,exp}}$ -definable functions under a slightly more lenient restriction on growth. Our statement can be seen to correspond to Selberg’s extension of Theorem 1.1 in [11], which covers all entire functions that have growth bounded by $e^{(\ln(2)+1/1500)|z|}$.

Theorem 1.2. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be definable in $\mathbb{R}_{\text{an,exp}}$. Suppose $f(a) \in \mathbb{Z}$ for all sufficiently large positive integers a and that*

$$|f(x)| = O(2^{(1+10^{-5})x}) \quad \text{as } x \rightarrow \infty.$$

Then there are polynomials $P_1, P_2 \in \mathbb{Q}[X]$ such that $f(x) = P_1(x) + P_2(x)2^x$ for all sufficiently large x .

The proof of this theorem broadly follows Wilkie’s strategy for the aforementioned result from [13]. This approach relies on the important fact, established in [13], that an $\mathbb{R}_{\text{an,exp}}$ -definable function on the reals is *closely* approximated by a complex-analytic function in an appropriate right half-plane. We use this ingredient without

¹RM thanks the Heilbronn Institute of Mathematical Research for their support.

²KW is supported by the National Science Foundation under Award No. DGE-2039656.

change, and our main technical contribution lies in the more extensive reworking of Langley's generalization of Theorem 1.1, where the same characterization is achieved for functions analytic in a right-half plane [6].

In fact, our adaptation can be used to obtain a minor improvement to the primary theorem of Langley; see Remark 3.4. This small upgrade would certainly not be a novel result, and indeed more comprehensive classifications are known for integer-valued complex-analytic functions, but via different methods ([2, 14]). At the end of [13], an extensive analog of these classifications is conjectured for $\mathbb{R}_{\text{an,exp}}$ -definable functions of exponential type, and we view Theorem 1.2 as a step towards this larger goal; see Conjecture 3.5.

Our second main theorem is a direct analog of a result by Pila, which studies entire functions satisfying a strong interpolative property. For $k \in \mathbb{N}$, we say that a function $f : \mathbb{R} \rightarrow \mathbb{R}$ is *k-concordant* if for each tuple m_0, \dots, m_k of sufficiently large positive integers, there is a polynomial $P \in \mathbb{Z}[X]$ such that $P(m_i) = f(m_i)$ for all $i = 0, \dots, k$. Set $\gamma_k := \sum_{j=1}^k 1/j$, and $\gamma_0 = 0$ by convention.

Theorem 1.3. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be definable in $\mathbb{R}_{\text{an,exp}}$. Suppose f is k -concordant for some $k \in \mathbb{N}$, and that there is $C < e^{\gamma_k} + 1$ such that*

$$|f(x)| \leq C^x$$

for all large enough x . Then there is a polynomial $P \in \mathbb{Q}[X]$ such that $f(x) = P(x)$ for all sufficiently large x .

Pila's theorem [8, Theorem 1.3] shows that a similar statement holds for entire functions. For $k = 0$, Theorem 1.3 is exactly [13, Theorem 5.1], and for the case of $k = 1$, our result corresponds to a theorem by Perelli and Zannier [7] for entire functions; see Remark 4.5.

The present paper fits into the larger theme of interactions of o-minimality with diophantine issues, and we view [12, Corollary 2.2] as the starting point for the particular trajectory pursued in this article. The results in [12] were succeeded also by the celebrated Pila-Wilkie theorem [9], a specific variant of which is used in fact by Jones, Thomas, and Wilkie in [5] to obtain a precursor to Theorem 1.2. Moreover, Jones and Qiu [4] obtain similar results for $\mathbb{R}_{\text{an,exp}}$ -definable functions which are either *close* to being integer-valued or take integer values on a sufficiently dense subset of the natural numbers.

Notations and conventions. Throughout we work with the conventions that $\mu, \nu, i, j, k \in \mathbb{N} := \{0, 1, \dots\}$, $a, m, n, B, K, N \in \mathbb{N}^{\geq 1}$, p is prime, $\epsilon, c, C \in \mathbb{R}^{>0}$, and $r \in \mathbb{R}$. Any symbol with an added subscript will denote an object in the same domain as the plain symbol. So $B_1 \in \mathbb{N}^{\geq 1}$, p_0 is a prime, $C_* \in \mathbb{R}^{>0}$, and so on.

The variable x varies always over the real numbers and e denotes Euler's number. For any $r \in \mathbb{R}$, we set $\mathcal{H}(r) := \{z \in \mathbb{C} : \text{Re}(z) > r\}$; so $\mathcal{H}(0)$ denotes the standard right-half plane.

We will work with a difference operator Δ given as follows. For any $f : \mathbb{R} \rightarrow \mathbb{R}$, set $\Delta f(x) := f(x+1) - f(x)$. We let Δ^n denote the n^{th} -iterate of the Δ operator, and by a straight-forward induction we have for all x that

$$\Delta^n f(x) = \sum_{k=0}^n (-1)^{n-k} \frac{n!}{k!(n-k)!} f(x+k), \quad (1)$$

which can also be seen in [13]. We shall also work with the $\Delta - 1$ operator, given by $(\Delta - 1)f(x) = f(x + 1) - 2f(x)$, and we consider iterates of this operator as well. If $f(x) = 2^x h(x)$, then for all x we have that $(\Delta - 1)^n f(x) = 2^{x+n} \Delta^n h(x)$.

Acknowledgements. We are deeply grateful to the organisers of the Arizona Winter School 2023 for bringing us together, and making this project possible. We thank Gabriel Dill and Jonathan Pila for introducing us to the problem which evolved into Theorem 1.3; and extend a very special thanks to Gabrielle Scullard, who was part of our working group at the AWS. We express our gratitude also to Gal Binyamini, Gareth Jones, Yuval Salant, and Alex Wilkie for helpful discussions during the preparation of this article.

2. TECHNICAL PRELIMINARIES

In this section, we record some important ingredients to be used in the proofs of Theorems 1.2 and 1.3 in §3 and §4 respectively.

Suppose $f : \mathbb{R} \rightarrow \mathbb{R}$ be given by $f(x) = P_1(x) + P_2(x)2^x$, for some polynomials P_1, P_2 . Then $\Delta^n (\Delta - 1)^k f = 0$ for any $n \geq \deg P_1 + 1$ and any $k \geq \deg P_2 + 1$. The following sharp converse will be crucial for us; the statement may be viewed as an appropriate [13, Lemma 5.2] type analog of [6, Lemma 3].

Lemma 2.1. *Let a function $f : \mathbb{N} \rightarrow \mathbb{R}$ be given. Suppose there are B, K such that for all $a \geq B$ we have that*

$$\Delta^{n-a} (\Delta - 1)^a f(a) = 0$$

for all n satisfying $Ka \leq n \leq K(a + 1)$. Then there are polynomials P_1, P_2 such that $f(a) = P_1(a) + P_2(a)2^a$ for all $a \geq B$. If moreover, $f(a) \in \mathbb{Z}$ for all $a \geq B$, then we get that $P_1, P_2 \in \mathbb{Q}[X]$.

Proof. Let $f : \mathbb{N} \rightarrow \mathbb{R}$ be given, and suppose B, K are such that for all $a \geq B$, we have that $\Delta^{n-a} (\Delta - 1)^a f(a) = 0$ for all n satisfying $Ka \leq n \leq K(a + 1)$. Let P_1 be a polynomial of degree at most $(K - 1)B - 1$ and P_2 be a polynomial of degree at most $B - 1$ such that

$$P_1(m) + P_2(m) \cdot 2^m = f(m)$$

for all $B \leq m \leq KB + B - 1$. Let $h : \mathbb{N} \rightarrow \mathbb{R}$ be given by

$$h(m) = P_1(m) + P_2(m)2^m - f(m).$$

If $n - a \geq (K - 1)B$, then $\Delta^{n-a} P_1(m) = 0$ for all m . Similarly, $a \geq B$ implies $(\Delta - 1)^a (P_2(m)2^m) = 2^{m+a} \Delta^a P_2(m) = 0$ for all m . So we have for all $a \geq B$ that

$$\Delta^{n-a} (\Delta - 1)^a (P_1(a) + P_2(a)2^a) = 0$$

for all $n \geq Ka$. Then by the linearity of Δ , the definition of h , and the starting assumption on f , we have for all $a \geq B$ that

$$\Delta^{n-a} (\Delta - 1)^a h(a) = 0$$

for all n satisfying $Ka \leq n \leq K(a + 1)$.

We shall show that $h(a) = 0$ for all $a \geq B$, which clearly proves the first assertion of the lemma. To achieve this goal, we prove by induction on a that

$$h(m) = 0 \quad \text{for all } B \leq m \leq Ka + a - 1.$$

We have the base case $a = B$ by the definition of h . Fix $a \geq B$ and assume the desired statement holds for a . It remains to prove the statement for $a + 1$, i.e.

$$h(m) = 0 \quad \text{for all} \quad Ka + a \leq m \leq K(a + 1) + a$$

For all n satisfying $Ka \leq n \leq K(a + 1)$, we have that $\Delta^{n-a}(\Delta - 1)^a h(a) = 0$, which gives that

$$h(a + n) = \text{a linear combination of } h(a), h(a + 1), \dots, h(a + n - 1).$$

If $n = Ka$, then $a + j \leq a + Ka - 1$ for all $j = 1, \dots, n - 1$. By the inductive assumption we get $h(a) = \dots = h(a + Ka - 1) = 0$, and it follows that $h(a + Ka) = 0$. Proceeding in the same manner, we immediately conclude that $h(m) = 0$ for all $m \leq a + K(a + 1)$, and the proof of the first part of the lemma is complete.

For the second assertion in the lemma, we need only observe that $f(a) \in \mathbb{Z}$ for all $a \geq B$ gives that the polynomials P_1 and P_2 above have rational coefficients. \square

The following simple fact will help handle the approximation error induced from deploying Proposition 2.5.

Lemma 2.2. *Let $K \geq 2$, and suppose $H : \mathbb{R} \rightarrow \mathbb{R}$ is such that $|H(x)| \leq \exp(-Kx)$ for all sufficiently large x . Then for all sufficiently large a , we have that*

$$|\Delta^{n-a}(\Delta - 1)^a H(a)| < 1/2$$

for all $a \leq n \leq K(a + 1)$.

Proof. Let B be such that $|H(x)| \leq \exp(-Kx)$ for all $x \geq B$. Fix an $a \geq B$ and an $a \leq n \leq K(a + 1)$. Then we have that

$$\begin{aligned} |\Delta^{n-a}(\Delta - 1)^a H(a)| &\leq \sum_{j=0}^a \frac{a!}{j!(a-j)!} \cdot |\Delta^{n-j} H(a)| \\ &\leq \sum_{j=0}^a \frac{a!}{j!(a-j)!} \cdot 2^{n-j} \exp(-Ka) \\ &\leq 2^n \cdot (3/2)^a \cdot \exp(-Ka) \\ &\leq 2^K \cdot (2^K \cdot (3/2) \cdot \exp(-K))^a, \end{aligned}$$

using that $n \leq K(a + 1)$. Since $K \geq 2$, we have that $(3/2) \cdot (2/e)^K < 1$. Hence for all sufficiently large a , we get that $|\Delta^{n-a}(\Delta - 1)^a H(a)| < 1/2$ for all $n \leq K(a + 1)$, as desired. \square

O-minimality. In this paper, as in [13], we work only with a particular o-minimal structure $\mathbb{R}_{\text{an,exp}}$. A rich class of sets and functions are definable in $\mathbb{R}_{\text{an,exp}}$, and this structure appears ubiquitously in applications of o-minimality to diophantine geometry. See [3] for a description of sets definable in $\mathbb{R}_{\text{an,exp}}$.

We shall use an immediate consequence of the defining axioms of o-minimality; see [1, Appendix A].

Fact 2.3. Let $f, g : \mathbb{R} \rightarrow \mathbb{R}$ be definable in $\mathbb{R}_{\text{an,exp}}$. Suppose $f(a) = g(a)$ for all sufficiently large a , then $f(x) = g(x)$ for all sufficiently large x .

This immediately gives the following improvement to Lemma 2.1.

Corollary 2.4. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be definable in $\mathbb{R}_{\text{an,exp}}$. Suppose there is K such that all sufficiently large a , we have that $f(a) \in \mathbb{Z}$ and*

$$\Delta^{n-a}(\Delta - 1)^a f(a) = 0$$

for all n satisfying $Ka \leq n \leq K(a+1)$. Then there are $P_1, P_2 \in \mathbb{Q}[X]$ such that $f(x) = P_1(x) + P_2(x)2^x$ for all sufficiently large x .

Our next result is a version of [13, Corollary 4.8], and is a key ingredient which allows access to the complex-analytic realm; see also [4, Theorem 3.1]. Set

$$\mathcal{R}_{\text{subexp}} := \{f : \mathbb{R} \rightarrow \mathbb{R} \text{ definable in } \mathbb{R}_{\text{an,exp}} : \forall \epsilon \exists r \text{ such that } |f(x)| \leq e^{\epsilon x}, \forall x \geq r\}.$$

Recall that $\mathcal{H}(B) \subset \mathbb{C}$ denotes the right-half plane comprising complex numbers with real part greater than B .

Proposition 2.5. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be definable in $\mathbb{R}_{\text{an,exp}}$ and suppose there are $c > 0$ and $C > 1$ such that*

$$|f(x)| \leq c \cdot C^x$$

for all sufficiently large x . Then for any positive reals ϵ and R , there is B and an analytic function $g : \mathcal{H}(B) \rightarrow \mathbb{C}$ such that

- (i) $|f(x) - g(x)| < e^{-Rx}$ for all $x > B$,
- (ii) $|g(z)| \leq |C^{z+\epsilon|z}|$ for all $z \in \mathcal{H}(B)$.

Proof. Let f be as above, and let $\epsilon, R \in \mathbb{R}^{>0}$ be given. Then [13, Corollary 4.8] gives $N, s_1, \dots, s_N \in \mathbb{R}, f_1, \dots, f_N \in \mathcal{R}_{\text{subexp}}$, and B_0 such that, for all $x > B_0$,

$$\left| f(x) - \sum_{j=1}^N f_j(x) \cdot C^{s_j x} \right| < \exp(-Rx).$$

We can and do assume that the s_j are distinct, and then the growth condition on f implies that $s_j \leq 1$ for all $j = 1, \dots, N$. For each $j \in \{1, \dots, N\}$, set $g_j(x) := f_j(x) \cdot C^{s_j x}$. Since $f_j \in \mathcal{R}_{\text{subexp}}$, [13, Lemma 5.3] gives B_j and a complex analytic continuation $\hat{f}_j : \mathcal{H}(B_j) \rightarrow \mathbb{C}$ of f_j that satisfies $|\hat{f}_j(z)| \leq C^{\epsilon|z|}/N$ for all $z \in \mathcal{H}(B_j)$. By the identity theorem, the function $\hat{g} : \mathcal{H}(B_j) \rightarrow \mathbb{C}$ given by

$$\hat{g}_j(z) := \hat{f}_j(z) C^{s_j z}$$

is an analytic continuation of g_j and $|\hat{g}_j(z)| \leq |C^{z+\epsilon|z}|/N$.

Observe that $B := \max\{B_0, B_1, \dots, B_N\}$ and $g := \hat{g}_1 + \dots + \hat{g}_N$ are as desired. \square

3. INTEGER-VALUED FUNCTIONS DEFINABLE IN $\mathbb{R}_{\text{an,exp}}$

We begin this section by revisiting the function $G(k, x, y)$ from [6, Lemma 6].

Lemma 3.1. *For fixed a , and $k = 0, \dots, a$, set*

$$G(k, x, y) = \prod_{0 \leq q \leq k-1} (1 + x + qy) \prod_{k \leq q \leq a-1} (1 - qy),$$

with the convention that a product over an empty range of q is 1. Let $A_{\mu, \nu}$ denote the coefficients of $\Delta^a G(0, x, y)$, so we have that

$$\Delta^a G(0, x, y) = \sum_{\substack{\mu \leq a, \\ \nu < a}} A_{\mu, \nu} x^\mu y^\nu$$

Then for all $\mu \leq a$ and $\nu < a$, $A_{\mu, \nu} = 0$ if $\mu + 2\nu < a$, and $|A_{\mu, \nu}| \leq 6^a a^\nu$.

Proof. For each $k \in \{0, \dots, a\}$, $G(k, x, y)$ is a polynomial

$$G(k, x, y) = \sum_{\substack{\mu \leq a, \\ \nu < a}} B_{\mu, \nu}(k) x^\mu y^\nu.$$

Fix some $\mu \leq a$ and $\nu < a - 1$. Then [6, Lemma 6 (ii)] implies that $A_{\mu, \nu} = 0$ if $\mu + 2\nu < a$.

Towards the second assertion, note that $|B_{\mu, \nu}(k)| \leq 3^a a^\nu$ for all $k \leq a$. Then the formula (1) gives

$$\Delta^a G(0, x, y) = \sum_{k=0}^a (-1)^{a-k} \frac{a!}{k!(a-k)!} G(k, x, y).$$

It follows that $|A_{\mu, \nu}| \leq \sum_{k=0}^a \frac{a!}{k!(a-k)!} |B_{\mu, \nu}(k)|$, and hence $|A_{\mu, \nu}| \leq 6^a a^\nu$. \square

Next, we have a detailed reworking of [6, Lemma 5].

Lemma 3.2. *For n and any positive real $s < n$, let $\mathcal{C}_{n,s}$ be the contour consisting of the arc $\Omega_{n,s}$ of the circle $|z| = 2n$ from $-s - i\sqrt{4n^2 - s^2}$ to $-s + i\sqrt{4n^2 - s^2}$ described once counter-clockwise, followed by the line segment $T_{n,s}$ from $-s + i\sqrt{4n^2 - s^2}$ to $-s - i\sqrt{4n^2 - s^2}$. Then there are positive integers b, d such that for all $n \geq 4$, $2 \leq s \leq n/2$, and $\mu \leq s$ we have that*

$$I_{n,s} = \int_{\Omega_{n,s}} \frac{n! 2^{|z|}}{|z| \dots |z-n|} \left| \frac{z-2n}{n} \right|^\mu |dz| < d \cdot (bs/n)^{\mu/2},$$

$$J_{n,s} = \int_{T_{n,s}} \frac{n!}{|z| \dots |z-n|} \left| \frac{z-2n}{n} \right|^\mu |dz| < d \cdot (bs/n)^{s-1}.$$

In fact, the choice of $b = d = 100$ suffices.

Proof. Fix $n \geq 4$ and s with $2 \leq s \leq n/2$. We follow the proof of [6, Lemma 5] with the aim of getting more explicit estimates.

First consider $I_{n,s}$. The arc $\Omega_{n,s}$ is parameterized by $z = 2ne^{i\theta}$, $-d_n \leq \theta \leq d_n$, where $d_n < 2$ since $s \leq n/2$. So $1 - \cos(\theta) \geq \frac{1}{3}\theta^2$ for all θ satisfying $-d_n \leq \theta \leq d_n$. This gives for all $z \in \Omega_{n,s}$ and $0 \leq k \leq n$ that

$$|z-k|^2 = (2n-k)^2 + 4nk(1 - \cos \theta) \geq (2n-k)^2 \exp\left(\frac{nk\theta^2}{3(2n-k)^2}\right).$$

It follows for all $z \in \Omega_{n,s}$ that

$$\prod_{k=0}^n |z-k| \geq \exp(n\theta^2/36) \cdot \prod_{k=0}^n (2n-k). \quad (2)$$

Also, we have for $z \in \Omega_{n,s}$ that

$$|z-2n| \leq 2n|\theta|, \quad \text{and} \quad \left| \frac{z-2n}{n} \right|^\mu \leq 2^\mu |\theta|^\mu. \quad (3)$$

Using (2), (3), Stirling's approximation, and a substitution $x = \sqrt{n}\theta/6$, we get that

$$\begin{aligned} I_{n,s} &\leq 2^\mu \cdot \frac{(2n)n!2^{2n}}{2n(2n-1)\dots(2n-n)} \cdot 2 \int_0^\pi \theta^\mu \exp(-n\theta^2/36) d\theta \\ &\leq 2 \cdot 2^\mu \cdot 4\sqrt{n} \cdot (36/n)^{\mu/2+1/2} \cdot \int_0^\infty x^\mu \exp(-x^2) dx \\ &\leq 48 \cdot 2^\mu \cdot (36/n)^{\mu/2} \cdot (\mu/2)^{\mu/2} \\ &\leq 48 \cdot (72\mu/n)^{\mu/2}. \end{aligned}$$

Here we used that $\int_0^\infty x^\mu \exp(-x^2) dx = \Gamma(\mu/2 + 1)/2 \leq (\mu/2)^{\mu/2}$.

This finishes the proof for the first part of the lemma and we turn our attention to $J_{n,s}$. For $z \in T_{n,s}$, we have $z = -s + iy$ for $-\sqrt{4n^2 - s^2} \leq y \leq \sqrt{4n^2 - s^2}$, and

$$|z - k|^2 = (s + k)^2 + y^2$$

for $0 \leq k \leq n$. It follows that

$$J_{n,s} \leq 2 \cdot 4^\mu \cdot \int_0^\infty \frac{n!}{\prod_{k=0}^n \sqrt{(s+k)^2 + y^2}} dy.$$

Next we observe that for D in [6, Equation 24], we have that $D \leq 2s$ and this gives

$$\begin{aligned} \frac{n!}{\prod_{k=0}^n \sqrt{(s+k)^2 + y^2}} &\leq \frac{3n^{n+1/2} e^{-n} (s^2 + y^2)^{(s-1)/2} e^n e^s}{((n+s)^2 + y^2)^{(n+s)/2} \exp(y \arctan((n+s)/y))} \\ &\leq 3^{s+1} \cdot n^{n+1/2} n^{-(n+s)} \cdot (s^2 + y^2)^{(s-1)/2} e^{-y/4}, \end{aligned}$$

using the simple inequality $\arctan(u) \geq 1/4$ for all $u \geq 1/2$. Now $\mu \leq s$ and $n \geq 4$ give us that

$$J_{n,s} \leq 2 \cdot 4^s \cdot 3^{s+1} \cdot (n^{-s+1}/2) \cdot \int_0^\infty (s^2 + y^2)^{(s-1)/2} e^{-y/4} dy.$$

Splitting the integral gives

$$\begin{aligned} \int_0^\infty (s^2 + y^2)^{(s-1)/2} e^{-y/4} dy &\leq 2^{s-1} \left[\int_0^\infty s^{(s-1)} e^{-y/4} dy + \int_0^\infty y^{(s-1)} e^{-y/4} dy \right] \\ &\leq 2^{s-1} [4 \cdot s^{s-1} + (4s)^{s-1}] \end{aligned}$$

Finally using $s \geq 2$ we arrive at

$$J_{n,s} \leq 3^2 \cdot 4 \cdot 2 \cdot (96s/n)^{s-1}.$$

The proof is complete. \square

Fixing K . Throughout the rest of this section we set $K := 10^5$. Hence $K = 10^3 b$, where b is the positive integer given by Lemma 3.2.

Proposition 3.3. *Let $g : \mathcal{H}(-1) \rightarrow \mathbb{C}$ be analytic and suppose we have for all $z \in \mathcal{H}(-1)$ that*

$$|g(z)| \leq |2^{z+2}|z|/K|.$$

Then there is $C_\star < 1$ such that for all sufficiently large a , we have that

$$|\Delta^{n-a}(\Delta - 1)^a g(a)| < C_\star^a$$

for all n satisfying $Ka \leq n \leq K(a + 1)$.

Proof. Suppose we are given an analytic function $g : \mathcal{H}(-1) \rightarrow \mathbb{C}$ such that

$$|g(z)| \leq |2^{z+2|z|/K}|$$

for all $z \in \mathcal{H}(-1)$. For all positive integers a , we consider the translations g_a of g given by complex analytic functions

$$g_a(z) := g(a+z) : \mathcal{H}(-a-1) \rightarrow \mathbb{C}.$$

So we have for all a that $|g_a(z)| \leq 2^{(1+2/K)a} \cdot |2^{z+2|z|/K}|$, and that g_a is analytic on an open set containing the contour $\mathcal{C}_{n,a}$ from Lemma 3.2. Fix an $a \geq 4$, and an n such that $Ka \leq n \leq K(a+1)$. We will obtain small asymptotic upper bounds for

$$D_{a,n} = |\Delta^{n-a}(\Delta-1)^a g(a)| = |\Delta^{n-a}(\Delta-1)^a g_a(0)|.$$

Following [6, Section 4], equation (31) there together with Lemma 3.1 gives

$$D_{a,n} \cdot \prod_{j=0}^{a-1} (1-j/n) \leq \sum_{\substack{\mu \leq a, \\ \nu < a}} |A_{\mu,\nu}| n^{-\nu} \int_{\mathcal{C}_{n,a}} \frac{n! |g_a(z)|}{|z| \dots |z-n|} \left| \frac{z-2n}{n} \right|^\mu |dz|.$$

So we have for all $\mu \leq a$ and $\nu < a$ that

$$A_{\mu,\nu} = 0 \quad \text{if} \quad \mu + 2\nu < a \quad \text{and} \quad |A_{\mu,\nu}| \leq 6^a a^\nu.$$

We have that $|z| \leq 2n$ for all $z \in \mathcal{C}_{n,a}$. Now using $a \geq 4$ and $n \leq K(a+1)$, we note that $|g_a(z)| \leq 2^{7a} 2^{|z|}$ for all $z \in \Omega_{n,a}$, and $|g_a(z)| \leq 2^{6a}$ for all $z \in T_{n,a}$. Combining these observations with an application of Lemma 3.2 for $s = a$ we get

$$\begin{aligned} D_{a,n} \cdot (1-a/n)^a &\leq 2^{7a} 6^a \sum_{\substack{\mu \leq a, \\ \nu < a, \\ \mu+2\nu \geq a}} (a/n)^\nu \cdot (I_{n,a} + J_{n,a}) \\ &\leq 768^a d \sum_{\substack{\mu \leq a, \\ \nu < a, \\ \mu+2\nu \geq a}} (a/n)^\nu (ba/n)^{\mu/2} + (a/n)^\nu (ba/n)^{a-1}, \end{aligned}$$

where b, d are the constants from Lemma 3.2. Now using $a/n \leq 1/K < 1$, we get

$$\begin{aligned} D_{a,n} &\leq (1-1/K)^{-a} \cdot 768^a \cdot d \cdot a^2 \cdot (b/K)^{a-1} \\ &\leq c \cdot (800b/(K-1))^{a-1}, \end{aligned}$$

for a constant c . Since $K = 1000b$, we get for all $a \geq 4$ that

$$D_{a,n} \leq c \cdot (6/5)^{-a+1}$$

for all n satisfying $Ka \leq n \leq K(a+1)$, and the proof is complete. \square

With modest modifications, one can obtain variants of Lemmas 3.2 and 3.3 which can be used together with Lemma 2.1 to slightly liberalize the growth condition in [6, Theorem 2] (and hence also in Theorem 1.1). We do not pursue this relatively minor increment here, and record it merely as an observation as follows.

Remark 3.4. There is an ϵ_0 such that the following holds. Let $g : \mathcal{H}(r) \rightarrow \mathbb{C}$ be analytic for some r , and that $g(a) \in \mathbb{Z}$ for all positive integers a . Suppose we have for all $z \in \mathcal{H}(r)$ that $|g(z)| = O(2^{(1+\epsilon_0)|z|})$. Then there are polynomials P_1, P_2 such that $g(z) = P_1(z) + P_2(z)2^z$ for all $z \in \mathcal{H}(r)$.

We are ready to finish the proof of our first theorem.

Proof of Theorem 1.2. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be as in the statement of the theorem, and let c, B_0 be such that $f(a) \in \mathbb{Z}$ and $f(x) \leq c2^{(1+10^{-5})x} = c2^{(1+1/K)x}$ for all $a \geq B_0$. Apply Proposition 2.5 for $C = 2^{(1+1/K)}$, $\epsilon = 1/K$, and $R = K$ to obtain $B_1 \geq B_0$ and an analytic function $g : \mathcal{H}(B_1) \rightarrow \mathbb{C}$ with

- (i) $|f(x) - g(x)| < e^{-Kx}$ for all $x > B_1$,
- (ii) $|g(z)| \leq |2^{z+2|z|/K}|$ for all $z \in \mathcal{H}(B_1)$.

Then applying Proposition 3.3 to an appropriate translate of g , we get $B_2 > B_1$ such that for all $a \geq B_2$,

$$|\Delta^{n-a}(\Delta - 1)^a g(a)| < 1/2$$

for all n satisfying $Ka \leq n \leq K(a+1)$. Then by the linearity of Δ and Lemma 2.2, we obtain $B_3 \geq B_2$ such that for all $a \geq B_3$ and n satisfying $Ka \leq n \leq K(a+1)$ we have that

$$|\Delta^{n-a}(\Delta - 1)^a f(a)| < 1, \quad \text{and hence} \quad |\Delta^{n-a}(\Delta - 1)^a f(a)| = 0.$$

We use here the simple observation that since $f(a) \in \mathbb{Z}$ for all $a \geq B_0$, also $\Delta^{n-a}(\Delta - 1)^a f(a) \in \mathbb{Z}$ for all $a \geq B_0$. The theorem now follows immediately by Corollary 2.4. \square

Connection to a conjecture of Wilkie. We want to draw attention to the following broad conjecture made at the conclusion of [13].

Conjecture 3.5. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be definable in $\mathbb{R}_{\text{an,exp}}$. Suppose that $f(a) \in \mathbb{Z}$ for all sufficiently large positive integers a and there is $r > 0$ such that $|f(x)| < \exp(rx)$ for all sufficiently large x . Then there is a polynomial $P(x, y_1, \dots, y_m)$ with rational coefficients, and positive real algebraic integers $\alpha_1, \dots, \alpha_m$ such that*

$$f(x) = P(x, \alpha_1^x, \dots, \alpha_m^x)$$

for all sufficiently large x .

Wilkie's result [13, Theorem 5.1] settles the case of $r < \ln(2)$, and Theorem 1.2 upgrades the classification to all $r \leq \ln(2)(1 + 10^{-5})$. Here we remark also that our choice of K earlier in the section is not optimal, and hence micro-improvements in this exponent $\ln(2)(1 + 10^{-5})$ may be easily achieved.

4. CONCORDANT INTEGER-VALUED FUNCTIONS DEFINABLE IN $\mathbb{R}_{\text{an,exp}}$

We begin with a basic Corollary 2.4 type result.

Fact 4.1. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be given.

- (i) Suppose there is B such that $\Delta^n f(B) = 0$ for all sufficiently large n . Then there is a polynomial P such that $f(a) = P(a)$ for all $a \geq B$.
- (ii) If moreover, $f(a) \in \mathbb{Z}$ for all $a \geq B$, then we get that $P \in \mathbb{Q}[X]$.
- (iii) If furthermore, f is definable in $\mathbb{R}_{\text{an,exp}}$, then $f(x) = P(x)$ for all large x .

Proof. Item (i) is a classical fact observed by Pólya, and we sketch a proof as follows. Let N be such that $\Delta^n f(B) = 0$ for all $n \geq N$. Let P be a polynomial of degree at most $N - 1$ such that $f(a) = P(a)$ for all $B \leq m \leq B + N - 1$. Then $\Delta^N P(B) = 0$, and using $\Delta^N f(B) = 0$, we get that $f(B + N) = P(B + N)$. Proceeding in the same manner, we get that $f(a) = P(a)$ for $a \geq B$, as desired.

If moreover $f(a) \in \mathbb{Z}$ for all $a \geq B$, then the polynomial P above has rational coefficients; and (iii) follows by direct application of Fact 2.3. \square

We shall use the following purely combinatorial fact; the statement is perhaps of some independent interest as well.

Lemma 4.2. *Let $k \geq 1$ and p be given. For all $\ell \in \mathbb{N}$ we have that*

$$\sum_{j=0}^k (-1)^{jp} \binom{kp}{jp} (jp)^\ell \equiv 0 \pmod{p^k},$$

and also for any $i \in \{1, \dots, p-1\}$ that

$$\sum_{j=0}^{k-1} (-1)^{jp} \binom{kp}{jp+i} (jp+i)^\ell \equiv 0 \pmod{p^k}.$$

Proof. We fix a primitive root of unity ζ_p . By a mod p reduction of the binomial coefficients we readily observe that

$$(1 - \zeta_p)^{p-1} = py \quad \text{for some } y \in \mathbb{Z}[\zeta_p]. \quad (4)$$

For any $a \in \mathbb{Q}(\zeta_p)$, we let $\text{Tr}(a)$ denote the trace of a with respect to the algebraic field extension $\mathbb{Q}(\zeta_p)/\mathbb{Q}$.

Observe for any $M \in \mathbb{N}$ and $t \in \mathbb{Z}$ that

$$\begin{aligned} \text{Tr}(\zeta_p^t (1 - \zeta_p)^M) &= \sum_{i=1}^{p-1} \zeta_p^{it} (1 - \zeta_p^i)^M = \sum_{i=0}^{p-1} \zeta_p^{it} (1 - \zeta_p^i)^M \\ &= \sum_{i=0}^{p-1} \sum_{\mu=0}^M (-1)^\mu \binom{M}{\mu} \zeta_p^{(\mu+t)i} \\ &= \sum_{\mu=0}^M (-1)^\mu \binom{M}{\mu} \sum_{i=0}^{p-1} (\zeta_p^{\mu+t})^i. \end{aligned}$$

For all $\mu \in \mathbb{N}$, we have that

$$\sum_{i=0}^{p-1} (\zeta_p^{\mu+t})^i = 0 \quad \text{if } p \nmid \mu + t \quad \text{and} \quad \sum_{i=0}^{p-1} (\zeta_p^{\mu+t})^i = p \quad \text{if } p \mid \mu + t,$$

which gives, for $t > -p$, that

$$\text{Tr}(\zeta_p^t (1 - \zeta_p)^M) = p \sum_{j=\max(0, \lceil t/p \rceil)}^{\lfloor (M+t)/p \rfloor} (-1)^{jp-t} \binom{M}{jp-t}. \quad (5)$$

Note this implies, by the linearity of the trace operator, that $p \mid \text{Tr}((1 - \zeta_p)^M y')$ for any $M \in \mathbb{N}^{\geq 1}$ and $y' \in \mathbb{Z}[\zeta_p]$.

We now work out the first assertion. Applying (5) for $M = kp$ and $t = 0$ gives

$$\text{Tr}((1 - \zeta_p)^{kp}) = p \sum_{j=0}^k (-1)^{jp} \binom{kp}{jp}.$$

Then the identity (4) implies that

$$p^k \cdot \text{Tr}((1 - \zeta_p)^k y^k) = p \sum_{j=0}^k (-1)^{jp} \binom{kp}{jp}.$$

Then by the remark just after (5),

$$p \mid \operatorname{Tr}((1 - \zeta_p)^k y^k) \quad \text{and hence} \quad \sum_{j=0}^k (-1)^{jp} \binom{kp}{jp} \equiv 0 \pmod{p^k}.$$

Hence we have the first assertion for the case $\ell = 0$.

Note that this claim is immediate for $\ell \geq k$. For any $1 \leq \ell < k$, consider

$$\sum_{j=0}^k (-1)^{jp} \binom{kp}{jp} (jp)(jp-1) \dots (jp-\ell+1) = \frac{(kp)!}{(kp-\ell)!} \sum_{j=\lceil \ell/p \rceil}^k (-1)^{jp} \binom{kp-\ell}{jp-\ell}.$$

By (5) for $M = kp - \ell$ and $t = \ell$, and (4) we see that

$$p \sum_{j=\lceil \ell/p \rceil}^k (-1)^{jp-\ell} \binom{kp-\ell}{jp-\ell} = \operatorname{Tr}(\zeta_p^\ell (1 - \zeta_p)^{kp-\ell}) = p^k \cdot \operatorname{Tr}(\zeta_p^\ell (1 - \zeta_p)^{k-\ell} y^k).$$

Then as before we use the remark just after (5) to conclude

$$\sum_{j=0}^k (-1)^{jp} \binom{kp}{jp} (jp)(jp-1) \dots (jp-\ell+1) \equiv 0 \pmod{p^k}.$$

From the above equation, the first assertion follows for $\ell = 1$. For $\ell > 1$, note we have integers $b_1, \dots, b_{\ell-1}$ such that

$$jp(jp-1) \dots (jp-\ell+1) = (jp)^\ell + b_1(jp)^{\ell-1} + \dots + b_{\ell-1}(jp),$$

and hence a simple induction on ℓ completes the proof of the first assertion.

We follow a similar process for the proof of the other claim as well, and consider first the case of $\ell = 0$. Fix $i \in \{1, \dots, p-1\}$. Then applying (5) for $M = kp$ and $t = -i$, we get that

$$\operatorname{Tr}(\zeta_p^{-i} (1 - \zeta_p)^{kp}) = p \sum_{j=0}^{k-1} (-1)^{i+jp} \binom{kp}{jp+i}.$$

Then the same arguments as before imply the desired result. We leave the general $\ell \geq 1$ case of the second assertion as an exercise to the reader. \square

This lemma will be used to obtain a strong divisibility implication of concordance. Recall the following definition from the introduction.

Definition 4.3. A function $f : \mathbb{R} \rightarrow \mathbb{R}$ is *k-concordant* if for each tuple m_0, \dots, m_k of sufficiently large positive integers, there is a polynomial $P \in \mathbb{Z}[X]$ such that $P(m_i) = f(m_i)$ for all $i = 0, \dots, k$. Note we do not require the m_0, \dots, m_k in the statement above to be distinct.

Remark 4.4. Clearly, $f : \mathbb{R} \rightarrow \mathbb{R}$ is 0-concordant if and only if $f(a) \in \mathbb{Z}$ for all sufficiently large a . Note that 1-concordance is equivalent to the condition that $f(m_1) \equiv f(m_2) \pmod{m_1 - m_2}$ for all sufficiently large m_1, m_2 . For all k , it follows directly from definition that $k+1$ -concordance implies k -concordance, and hence k -concordant functions take integer values for all sufficiently positive integer inputs.

Remark 4.5. In [7], Perelli and Zannier consider functions satisfying a weaker form of 1-concordance. For an integer-valued entire function $f : \mathbb{C} \rightarrow \mathbb{C}$, they show that if $f(a+p) \equiv f(a) \pmod{p}$ for all positive integers a and large enough primes p , then $|f(z)| = O(C^{|z|})$, for some $C < e+1$, implies that f must be a polynomial.

We remark that the conclusions of the following lemma and Theorem 1.3 for the case of $k = 1$ hold even when the function $f : \mathbb{R} \rightarrow \mathbb{R}$ satisfies only this weaker form of 1-concordance. The details are left to the reader.

Recall $\gamma_k = \sum_{j=1}^k 1/j$. The following corresponds to [8, Theorem 3.5, Lemma 5.1].

Lemma 4.6. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be k -concordant, with $k \geq 1$. Then there are c, B such that for all $a \geq B$ and all n we have that*

$$|\Delta^n f(a)| \geq e^{\gamma_k n - cn/\log(n)} \quad \text{or} \quad \Delta^n f(a) = 0.$$

Proof. Let a k -concordant $f : \mathbb{R} \rightarrow \mathbb{R}$ be given. Fix a B such that for all $m_0, \dots, m_k \in \mathbb{N}^{\geq B}$, there is a polynomial $P \in \mathbb{Z}[X]$ with $P(m_i) = f(m_i)$ for all $i = 0, \dots, k$. We claim that for any $a \geq B$ and prime p that

$$\Delta^{kp} f(a) \equiv 0 \pmod{p^k}. \quad (6)$$

To show this, we fix a p and use (1) to get that

$$\Delta^{kp} f(a) = (-1)^{kp} \sum_{i=0}^{kp} (-1)^i \binom{kp}{i} f(a+i).$$

This can be rearranged as follows

$$(-1)^{kp} \Delta^{kp} f(a) = \sum_{j=0}^k (-1)^{jp} \binom{kp}{jp} f(a+jp) + \sum_{i=1}^{p-1} \sum_{j=0}^{k-1} (-1)^{jp+i} \binom{kp}{jp+i} f(a+jp+i).$$

Consider the first sum on the right hand side. Since $a \geq B$, the k -concordance of f gives a polynomial $P \in \mathbb{Z}[X]$ such that $P(jp) = f(a+jp)$ for all $j = 0, \dots, k$. Now the first assertion of Lemma 4.2 implies that

$$\sum_{j=0}^k (-1)^{jp} \binom{kp}{jp} f(a+jp) = \sum_{j=0}^k (-1)^{jp} \binom{kp}{jp} P(jp) \equiv 0 \pmod{p^k}.$$

A similar argument, now involving the second assertion of Lemma 4.2, gives for each $i \in \{1, \dots, p-1\}$ that

$$\sum_{j=0}^{k-1} (-1)^{jp+i} \binom{kp}{jp+i} f(a+jp+i) \equiv 0 \pmod{p^k},$$

and we have proved the claim set forth in (6).

So we have for any $p \leq \frac{n}{k}$ that $\Delta^n f(a) \equiv 0 \pmod{p^k}$, and hence we get for all $\ell = 1, 2, \dots, k$,

$$\prod_{p \leq \frac{n}{\ell}} p^\ell \mid \Delta^n f(a),$$

with an empty product being 1 by convention. In other words,

$$\prod_{\ell \leq k} \prod_{p \leq \frac{n}{\ell}} p \mid \Delta^n f(a).$$

The prime number theorem gives a constant c such that

$$\sum_{\ell \leq k} \sum_{p \leq \frac{n}{\ell}} \log(p) \geq \gamma_k n - cn/\log(n)$$

for all n . Thus we get for all n that either

$$|\Delta^n f(a)| \geq e^{\gamma_k n - cn/\log(n)} \quad \text{or} \quad \Delta^n f(a) = 0,$$

and the proof is complete. \square

Lemma 4.7. *Let analytic $g : \mathcal{H}(-2) \rightarrow \mathbb{C}$ and $k \geq 1$ be given. Suppose there are $\epsilon < 1$, and C such that*

$$|g(z)| \leq |C^{z+\epsilon}|z|$$

for all $z \in \mathcal{H}(-2)$. If $C^{1+\epsilon} < e^{\gamma_k} + 1$, then there is $C_\star < e^{\gamma_k}$ such that for all sufficiently large n we have that

$$|\Delta^n g(0)| \leq C_\star^n.$$

Proof. Let an analytic function $g : \mathcal{H}(-2) \rightarrow \mathbb{C}$, and positive integer k be given. Let ϵ, C be such that $|g(z)| \leq |C^{z+\epsilon}|z|$ for all $z \in \mathcal{H}(-2)$, and also $C^{1+\epsilon} < e^{\gamma_k} + 1$. If $C \leq 1$, then $|\Delta^n g(0)| \leq 2^n$ and we are done. Hence we assume that $C > 1$.

Fix an n , and let \mathcal{C}_n be the contour formed by traversing once counter-clockwise the boundary of the region inside $\{z \in \mathbb{C} : \operatorname{Re}(z) \geq -1\}$ that is bounded by the circle $|z| = (1 + 1/e^{\gamma_k})n$ and the vertical line $\operatorname{Re}(z) = -1$. We denote by T_n the part of \mathcal{C}_n that is contained in the vertical line, and let Ω_n denote the rest of the contour, so $\mathcal{C}_n = \Omega_n \cup T_n$. Then Cauchy's integral formula and (1) gives that

$$\Delta^n g(0) = \frac{1}{2\pi i} \int_{\mathcal{C}_n} \frac{n!g(z)}{z(z-1)\dots(z-n)} dz.$$

For all $z \in \Omega_n$, $|g(z)| \leq (C^{1+\epsilon})^{(1+1/e^{\gamma_k})n}$, and $|z - k| \geq |z| - k = (1 + 1/e^{\gamma_k})n - k$ for all $k = 0, 1, \dots, n$. A straightforward application of Stirling's approximation gives c_1 such that

$$\frac{n!}{(1 + 1/e^{\gamma_k})n((1 + 1/e^{\gamma_k})n - 1) \dots ((1 + 1/e^{\gamma_k})n - n)} < \frac{c_1 \cdot e^{\gamma_k n}}{(e^{\gamma_k} + 1)^{(1+1/e^{\gamma_k})n}}.$$

Then changing c_1 as required, we get for $C_1 := e^{\gamma_k} \cdot (C^{1+\epsilon}/(e^{\gamma_k} + 1))^{1+1/e^{\gamma_k}}$ that

$$\int_{\Omega_n} \frac{n!|g(z)|}{|z| \cdot |z-1| \dots |z-n|} dz < c_1 \cdot n C_1^n;$$

note that $C_1 < e^{\gamma_k}$. We now turn our attention to the integral over T_n . For all $z \in T_n$, we have that $|g(z)| \leq (C^\epsilon)^{(1+1/e^{\gamma_k})n}$, and $|z - k| \geq k + 1$ for all $k = 0, 1, \dots, n$.

Using $C^{1+\epsilon} < e^{\gamma_k} + 1$ and $\epsilon < 1$ it follows that $C^{2\epsilon} < e^{\gamma_k} + 1$. Then since $x^4 \geq (x+1)^3$ for all $x \geq e$, we get that $C_2 := (C^\epsilon)^{(1+1/e^{\gamma_k})} < C^{3\epsilon/2} < e^{\gamma_k}$. So there is c_2 such that

$$\int_{T_n} \frac{n!|g(z)|}{|z| \cdot |z-1| \dots |z-n|} \leq \int_{T_n} n^{-1} \cdot C_2^n \leq c_2 \cdot C_2^n.$$

Hence it is immediate that the desired conclusion follows for any choice of C_\star satisfying $\max(C_1, C_2) < C_\star < e^{\gamma_k}$. \square

Proof of Theorem 1.3. Let k -concordant $\mathbb{R}_{\text{an,exp}}$ -definable $f : \mathbb{R} \rightarrow \mathbb{R}$ be given, and suppose there is $C < e^{\gamma_k} + 1$ such that $|f(x)| \leq C^x$ for all sufficiently large x . The case is $k = 0$ follows by Theorem 1.2 (or [13, Theorem 5.1]). So we assume $k \geq 1$ and apply Lemma 4.6 to obtain c, B_0 such that

$$|\Delta^n f(a)| \geq e^{\gamma_k n - cn/\log(n)} \quad \text{or} \quad \Delta^n f(a) = 0 \quad (7)$$

for all $a \geq B_0$ and all n . Fix any $\epsilon < 1$ such that $C^{1+\epsilon} < e^{\gamma k} + 1$. Proposition 2.5 to gives $B \geq B_0$ and an analytic function $g : \mathcal{H}(B - 2) \rightarrow \mathbb{C}$ with

- (i) $|f(x) - g(x)| < 1$ for all $x \geq B$,
- (ii) $|g(z)| \leq |C^{z+\epsilon|z|}|$ for all $z \in \mathcal{H}(B - 2)$.

This implies the easy estimate $|\Delta^n(f(B) - g(B))| \leq 2^n$ for all n . Then the linearity of Δ , and Lemma 4.7 applied to an appropriate translate of g , gives $C_\star < e^{\gamma k}$ such that for all large sufficiently large n we have that

$$\Delta^n f(B) \leq C_\star^n.$$

This together with (7) implies that $\Delta^n f(B) = 0$ for all sufficiently large n , and finally the theorem follows by Fact 4.1. \square

REFERENCES

1. N. Bhardwaj and L. van den Dries, *On the Pila-Wilkie theorem*, Expo. Math. **40** (2022), no. 3, 495–542. MR 4475392
2. R. C. Buck, *Integral valued entire functions*, Duke Math. J. **15** (1948), 879–891. MR 29984
3. L. van den Dries, A. Macintyre, and D. Marker, *The elementary theory of restricted analytic fields with exponentiation*, Ann. of Math. (2) **140** (1994), no. 1, 183–205. MR 1289495
4. G. Jones and S. Qiu, *Integer-valued definable functions in $\mathbb{R}_{\text{an,exp}}$* , Int. J. Number Theory **17** (2021), no. 8, 1739–1752. MR 4310204
5. G. O. Jones, M. E. M. Thomas, and A. J. Wilkie, *Integer-valued definable functions*, Bull. Lond. Math. Soc. **44** (2012), no. 6, 1285–1291. MR 3007661
6. J. Langley, *Integer-valued analytic functions in a half-plane*, Comput. Methods Funct. Theory **7** (2007), no. 2, 433–442. MR 2376683
7. A. Perelli and U. Zannier, *On a theorem of Pólya*, Boll. Un. Mat. Ital. A (5) **18** (1981), no. 2, 305–307. MR 618350
8. J. Pila, *Entire functions having a concordant value sequence*, Israel J. Math. **134** (2003), 317–343. MR 1972183
9. J. Pila and A. J. Wilkie, *The rational points of a definable set*, Duke Math. J. **133** (2006), no. 3, 591–616. MR 2228464
10. G. Pólya, *Über ganze ganzwertige funktionen*, Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse (1920), 1–10.
11. A. Selberg, *Über ganzwertige ganze transzendente Funktionen*, Arch. Math. Naturvid. **44** (1941), 45–52. MR 4306
12. A. J. Wilkie, *Diophantine properties of sets definable in o-minimal structures*, J. Symbolic Logic **69** (2004), no. 3, 851–861. MR 2078926
13. ———, *Complex continuations of $\mathbb{R}_{\text{an,exp}}$ -definable unary functions with a diophantine application*, J. Lond. Math. Soc. (2) **93** (2016), no. 3, 547–566. MR 3509953
14. K. Yoshino, *Analytic continuation of arithmetic holomorphic functions on a half-plane*, Tokyo J. Math. **2** (1979), no. 1, 121–128. MR 541901

WEIZMANN INSTITUTE OF SCIENCE, ISRAEL
 Email address: nbhardwaj@msri.org

UNIVERSITY OF MANCHESTER, UK
 Email address: raymond.mcculloch@manchester.ac.uk

UNIVERSITY OF CALIFORNIA SAN DIEGO, USA
 Email address: naramach@ucsd.edu

PRINCETON UNIVERSITY, USA
 Email address: khwoo@princeton.edu