

Two Homomorphisms from the affine Yangian associated with $\widehat{\mathfrak{sl}}(n)$ to the affine Yangian associated with $\widehat{\mathfrak{sl}}(n+1)$

Mamoru Ueda*

Department of Mathematical and Statistical Sciences, University of Alberta,
11324 89 Ave NW, Edmonton, AB T6G 2J5, Canada

Abstract

We construct a homomorphism from the affine Yangian $Y_{\hbar,\varepsilon+\hbar}(\widehat{\mathfrak{sl}}(n))$ to the affine Yangian $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n+1))$ which is different from the one in [20]. By using this homomorphism, we give a homomorphism from $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n)) \otimes Y_{\hbar,\varepsilon+n\hbar}(\widehat{\mathfrak{sl}}(m))$ to $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(m+n))$. As an application, we construct a homomorphism from the affine Yangian $Y_{\hbar,\varepsilon+n\hbar}(\widehat{\mathfrak{sl}}(m))$ to the centralizer algebra of the pair of affine Lie algebras $(\widehat{\mathfrak{gl}}(m+n), \widehat{\mathfrak{sl}}(n))$ and the coset vertex algebra of the pair of rectangular W -algebras $\mathcal{W}^k(\mathfrak{gl}(2m+2n), (2^{m+n}))$ and $\mathcal{W}^{k+m}(\mathfrak{sl}(2n), (2^n))$.

1 Introduction

The Yangian $Y_{\hbar}(\mathfrak{g})$ associated with a finite dimensional simple Lie algebra \mathfrak{g} was introduced by Drinfeld ([4], [5]). The Yangian $Y_{\hbar}(\mathfrak{g})$ is a quantum group which is a deformation of the current algebra $\mathfrak{g} \otimes \mathbb{C}[z]$. The Yangian has several presentations: the RTT presentation, the current presentation, the Drinfeld J presentation and so on.

By using the current presentation of the Yangian, we can extend the definition of the Yangian $Y_{\hbar}(\mathfrak{g})$ to a symmetrizable Kac-Moody Lie algebra \mathfrak{g} . The affine Yangian $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$ was first introduced by Guay ([10] and [11]). The affine Yangian $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$ is a 2-parameter affine Yangian associated with $\widehat{\mathfrak{sl}}(n)$ and a quantum group which is a deformation of the universal enveloping algebra of the central extension of $\mathfrak{sl}(n)[u^{\pm 1}, v]$. Recently, the affine Yangian is applied to the study of a W -algebra. The W -algebra $\mathcal{W}^k(\mathfrak{g}, f)$ is a vertex algebra associated with a finite dimensional reductive Lie algebra \mathfrak{g} , a nilpotent element $f \in \mathfrak{g}$ and a complex number k . In [21], we gave a relationship between the affine Yangian and a rectangular W -algebra. The rectangular W -algebra $\mathcal{W}^k(\mathfrak{gl}(ln), (l^n))$ is a W -algebra associated with $\mathfrak{gl}(ln)$ and a nilpotent element of type (l^n) . The author [21] gave a surjective homomorphism

$$\Phi^n: Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n)) \rightarrow \mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(ln), (l^n))),$$

where $\mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(ln), (l^n)))$ is the universal enveloping algebra of $\mathcal{W}^k(\mathfrak{gl}(ln), (l^n))$.

One of the difference between finite Yangians and affine Yangians is the existence of the RTT presentation. By using the RTT presentation, we find that there exists a natural embedding Ψ_1^f from the finite Yangian associated with $\mathfrak{gl}(n)$ to the finite Yangian associated with $\mathfrak{gl}(n+m)$. In the affine setting, the author [20] constructed a homomorphism corresponding to Ψ_1^f :

$$\Psi_1: Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n)) \rightarrow \widetilde{Y}_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(m+n)),$$

where $\widetilde{Y}_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(m+n))$ is the degreewise completion of $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(m+n))$. In the finite setting, by Ψ_1 , Olshanskii [18] gave a homomorphism from the finite Yangian associated with $\mathfrak{gl}(m)$ to the

*mueda@ualberta.ca

centralizer algebra of the pair of $(\mathfrak{gl}(m+n), \mathfrak{gl}(n))$. Moreover, the finite Yangian associated with $\mathfrak{gl}(m)$ can be embedded into the projective limit of this centralizer algebra.

In this article, we construct a homomorphism

$$\Psi_2: Y_{\hbar, \varepsilon + n\hbar}(\widehat{\mathfrak{sl}}(m)) \rightarrow \widetilde{Y}_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(m+n)),$$

where $\Psi_2(Y_{\hbar, \varepsilon + n\hbar}(\widehat{\mathfrak{sl}}(m)))$ and $\Psi_1(Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n)))$ are commutative with each other. The affine Yangian has a surjective homomorphism called the evaluation map (see Guay [11] and Kodera [14], [13]):

$$\text{ev}_{\hbar, \varepsilon}^n: Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n)) \rightarrow U(\widehat{\mathfrak{gl}}(m+n)).$$

By combining $\text{ev}_{\hbar, \varepsilon}^{m+n}$ and Ψ_2 , we obtain a homomorphism

$$\text{ev}_{\hbar, \varepsilon}^{m+n} \circ \Psi_2: Y_{\hbar, \varepsilon + n\hbar}(\widehat{\mathfrak{sl}}(m)) \rightarrow C(\widehat{\mathfrak{gl}}(m+n), \widehat{\mathfrak{gl}}(n)),$$

where $C(\widehat{\mathfrak{gl}}(m+n), \widehat{\mathfrak{gl}}(n))$ is a centralizer algebra of the pair of $U(\widehat{\mathfrak{gl}}(m+n))$ and $U(\widehat{\mathfrak{gl}}(n))$.

The similar result holds for the rectangular W -algebra. There exists a natural embedding from the rectangular W -algebra $\mathcal{W}^{k+m}(\mathfrak{gl}(2n), (2^n))$ to $\mathcal{W}^k(\mathfrak{gl}(2(m+n)), (2^{m+n}))$. By combining Φ^{m+n} and Ψ_2 , we can obtain a homomorphism

$$\Phi^{m+n} \circ \Psi_2: Y_{\hbar, \varepsilon + n\hbar}(\widehat{\mathfrak{sl}}(m)) \rightarrow \mathcal{U}(C(\mathcal{W}^k(\mathfrak{gl}(2m+2n), (2^{m+n})), \mathcal{W}^{k+m}(\mathfrak{sl}(2n), (2^n)))),$$

where $C(\mathcal{W}^k(\mathfrak{gl}(2m+2n), (2^{m+n})), \mathcal{W}^{k+m}(\mathfrak{sl}(2n), (2^n)))$ is a coset vertex algebra of the pair $\mathcal{W}^k(\mathfrak{gl}(2m+2n), (2^{m+n}))$ and $\mathcal{W}^k(\mathfrak{sl}(2n), (2^n))$. We also conjecture that we can obtain the similar homomorphism for any $l \geq 3$. For proving this conjecture, we only need to show that we can embed the rectangular W -algebra $\mathcal{W}^{k+(l-1)m}(\mathfrak{gl}(ln), (l^n))$ to $\mathcal{W}^k(\mathfrak{gl}(lm+ln), (l^{m+n}))$ naturally.

Kodera-Ueda [15] gave the meaning to Φ^n from the perspective of l . The meaning is that the coproduct for the affine Yangian corresponds to the parabolic induction for a rectangular W -algebra via the homomorphism Φ^n . The result of this article attaches the meaning to Φ^n from the perspective of n . This meaning will be connected to Gaiotto-Rapcak's triality

Gaiotto and Rapcak [8] introduced a kind of vertex algebras called Y -algebras. The Y -algebras are related to twisted $N=4$ supersymmetric gauge theories. Gaiotto-Rapcak [8] conjectured a triality of the isomorphism of Y -algebras. Let us consider a W -algebra associated with $\mathfrak{sl}(m+n)$ and its nilpotent element $f_{n,m}$ of type $(n^1, 1^m)$. The nilpotent element $f_{n,m}$ can be decomposed into two nilpotent elements: one is a principal nilpotent element of $\mathfrak{gl}(n)$ and another is a nilpotent element of type (1^n) , that is, zero. The W -algebra $\mathcal{W}^k(\mathfrak{sl}(m+n), f_{n,m})$ naturally contains the universal affine vertex algebra $V^{k-m-1}(\mathfrak{gl}(m))$, which is corresponding to the latter nilpotent element. It is known that some kinds of Y -algebras can be realized as a coset of the pair of $\mathcal{W}^k(\mathfrak{sl}(m+n), f_{n,m})$ and $V^{k-m-1}(\mathfrak{gl}(m))$ up to Heisenberg algebras. In this case, Creutzig-Linshaw [3] have proved the triality conjecture. This result is the generalization of the Feigin-Frenkel duality and the coset realization of principal W -algebra.

The Y -algebras can be interpreted as a truncation of $\mathcal{W}_{1+\infty}$ -algebra ([7]), whose universal enveloping algebra is isomorphic to the affine Yangian of $\widehat{\mathfrak{gl}}(1)$ up to suitable completions (see [2], [19] and [17]). In this article, we consider the coset of rectangular W -algebras associated with nilpotent elements of types (2^{m+n}) and (2^n) and relate the affine Yangian to this coset. We expect that we can construct a similar relation for general W -algebras and general decompositions of a nilpotent element. We also hope that this result will be connected to obtain the generalization of the Gaiotto-Rapcak's triality.

Acknowledgement

The author expresses his sincere thanks to Thomas Creutzig and Nicolas Guay for the helpful discussion.

2 Affine Yangian

Let us recall the definition of the affine Yangian of type A (Definition 3.2 in [10] and Definition 2.3 in [11]). Hereafter, we sometimes identify $\{0, 1, 2, \dots, n-1\}$ with $\mathbb{Z}/n\mathbb{Z}$. Let us set $\{X, Y\} = XY + YX$ and

$$a_{i,j} = \begin{cases} 2 & \text{if } i = j, \\ -1 & \text{if } j = i \pm 1, \\ 0 & \text{otherwise} \end{cases}$$

for $i \in \mathbb{Z}/n\mathbb{Z}$.

Definition 2.1. Suppose that $n \geq 3$. The affine Yangian $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$ is the associative algebra generated by $X_{i,r}^+, X_{i,r}^-, H_{i,r}$ ($i \in \{0, 1, \dots, n-1\}, r = 0, 1$) subject to the following defining relations:

$$[H_{i,r}, H_{j,s}] = 0, \quad (2.2)$$

$$[X_{i,0}^+, X_{j,0}^-] = \delta_{i,j} H_{i,0}, \quad (2.3)$$

$$[X_{i,1}^+, X_{j,0}^-] = \delta_{i,j} H_{i,1} = [X_{i,0}^+, X_{j,1}^-], \quad (2.4)$$

$$[H_{i,0}, X_{j,r}^\pm] = \pm a_{i,j} X_{j,r}^\pm, \quad (2.5)$$

$$[\tilde{H}_{i,1}, X_{j,0}^\pm] = \pm a_{i,j} (X_{j,1}^\pm), \text{ if } (i, j) \neq (0, n-1), (n-1, 0), \quad (2.6)$$

$$[\tilde{H}_{0,1}, X_{n-1,0}^\pm] = \mp \left(X_{n-1,1}^\pm + \left(\varepsilon + \frac{n}{2}\hbar\right) X_{n-1,0}^\pm \right), \quad (2.7)$$

$$[\tilde{H}_{n-1,1}, X_{0,0}^\pm] = \mp \left(X_{0,1}^\pm - \left(\varepsilon + \frac{n}{2}\hbar\right) X_{0,0}^\pm \right), \quad (2.8)$$

$$[X_{i,1}^\pm, X_{j,0}^\pm] - [X_{i,0}^\pm, X_{j,1}^\pm] = \pm a_{i,j} \frac{\hbar}{2} \{X_{i,0}^\pm, X_{j,0}^\pm\} \text{ if } (i, j) \neq (0, n-1), (n-1, 0), \quad (2.9)$$

$$[X_{0,1}^\pm, X_{n-1,0}^\pm] - [X_{0,0}^\pm, X_{n-1,1}^\pm] = \mp \frac{\hbar}{2} \{X_{0,0}^\pm, X_{n-1,0}^\pm\} + \left(\varepsilon + \frac{n}{2}\hbar\right) [X_{0,0}^\pm, X_{n-1,0}^\pm], \quad (2.10)$$

$$(\text{ad } X_{i,0}^\pm)^{1+|a_{i,j}|} (X_{j,0}^\pm) = 0 \text{ if } i \neq j, \quad (2.11)$$

where $\tilde{H}_{i,1} = H_{i,1} - \frac{\hbar}{2} H_{i,0}^2$.

Remark 2.12. Definition 2.1 is different from Definition 3.2 in [10] and Definition 2.3 in [11]. Guay-Nakajima-Wendlandt [12] gave the minimalistic presentation of the affine Yangian. Definition 2.1 can be derived from the minimalistic presentation (see Section 2 in [20]).

By using the defining relations of the affine Yangian, we find the following relations (see Section 2 in [20]):

$$[X_{i,r}^\pm, X_{j,s}^\pm] = 0 \text{ if } |i-j| > 1, \quad (2.13)$$

$$[X_{i,1}^\pm, [X_{i,0}^\pm, X_{j+1,r}^\pm]] + [X_{i,0}^\pm, [X_{i,1}^\pm, X_{j+1,r}^\pm]] = 0. \quad (2.14)$$

By the definition of the affine Yangian $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$, we find that there exists a natural homomorphism from the universal enveloping algebra of $\widehat{\mathfrak{sl}}(n)$ to $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$. In order to simplify the notation, we denote the image of $x \in U(\widehat{\mathfrak{sl}}(n))$ via this homomorphism by x .

We take one completion of $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$. We set the degree of $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$ by

$$\deg(H_{i,r}) = 0, \quad \deg(X_{i,r}^\pm) = \begin{cases} \pm 1 & \text{if } i = 0, \\ 0 & \text{if } i \neq 0. \end{cases}$$

We denote the standard degreewise completion of $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$ by $\tilde{Y}_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$. Let us set $A_i \in \tilde{Y}_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$ as

$$A_i = \frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ u > v}} E_{u,v} t^{-s} [E_{i,i}, E_{v,u} t^s] + \frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ u < v}} E_{u,v} t^{-s-1} [E_{i,i}, E_{v,u} t^{s+1}]$$

$$\begin{aligned}
&= \frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ u > i}} E_{u,i} t^{-s} E_{i,u} t^s - \frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ i > v}} E_{i,v} t^{-s} E_{v,i} t^s \\
&\quad + \frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ u < i}} E_{u,i} t^{-s-1} E_{i,u} t^{s+1} - \frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ i < v}} E_{i,v} t^{-s-1} E_{v,i} t^{s+1},
\end{aligned}$$

where $E_{i,j}$ is a matrix unit whose (a, b) component is $\delta_{a,i} \delta_{b,j}$. Similarly to Section 3 in [12], we define

$$J(h_i) = \tilde{H}_{i,1} + A_i - A_{i+1} \in \tilde{Y}_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n)).$$

We also set $J(x_i^\pm) = \pm \frac{1}{2} [J(h_i), x_i^\pm]$.

Guay-Nakajima-Wendlandt [12] defined the automorphism of $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n))$ by

$$\tau_i = \exp(\text{ad}(x_{i,0}^+)) \exp(-\text{ad}(x_{i,0}^-)) \exp(\text{ad}(x_{i,0}^+)).$$

Let α be a positive real root. By definition of the Weyl algebra, there is an element w of the Weyl group of $\widehat{\mathfrak{sl}}(n)$ and a simple root α_j such that $\alpha = w\alpha_j$. Then we define a corresponding root vector by

$$x_\alpha^\pm = \tau_{i_1} \tau_{i_2} \cdots \tau_{i_{p-1}} (x_j^\pm),$$

where $w = s_{i_1} s_{i_2} \cdots s_{i_{p-1}}$ is a reduced expression of w . We can define $J(x_\alpha^\pm)$ as

$$J(x_\alpha^\pm) = \tau_{i_1} \tau_{i_2} \cdots \tau_{i_{p-1}} J(x_j^\pm).$$

Lemma 2.15 ((3.14) and Proposition 3.21 in [12]). *1. The following relations hold:*

$$[J(h_i), X_{j,0}^\pm] \pm a_{ij} J(x_j^\pm) \text{ if } (i, j) \neq (0, n-1), (n-1, 0), \quad (2.16)$$

$$[J(x_i^\pm), X_{j,0}^\pm] = [X_{i,0}^\pm, J(x_j^\pm)] \text{ if } (i, j) \neq (0, n-1), (n-1, 0), \quad (2.17)$$

$$[J(x_i^\pm), X_{j,0}^\pm] = 0 \text{ if } |i - j| > 1. \quad (2.18)$$

2. There exists $c_{\alpha, i} \in \mathbb{C}$ satisfying that

$$[J(h_i), x_\alpha^\pm] = \pm(\alpha_i, \alpha) J(x_\alpha^\pm) \pm c_{\alpha, i} x_\alpha^\pm.$$

3 A homomorphism from the affine Yangian $Y_{\hbar, \varepsilon + \hbar}(\widehat{\mathfrak{sl}}(n))$ to the affine Yangian $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n+1))$

In this section, we will construct a homomorphism from the affine Yangian $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n))$ to the degreewise completion of the affine Yangian $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n+1))$, which is different from the one given in [20].

Theorem 3.1. *There exists an algebra homomorphism*

$$\Psi: Y_{\hbar, \varepsilon + \hbar}(\widehat{\mathfrak{sl}}(n)) \rightarrow \tilde{Y}_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n+1))$$

determined by

$$\begin{aligned}
\Psi(H_{i,0}) &= \begin{cases} H_{0,0} + H_{1,0} & \text{if } i = 0, \\ H_{i+1,0} & \text{if } i \neq 0, \end{cases} \\
\Psi(X_{i,0}^+) &= \begin{cases} E_{n+1,2t} & \text{if } i = 0, \\ E_{i+1, i+2} & \text{if } i \neq 0, \end{cases} \quad \Psi(X_{i,0}^-) = \begin{cases} E_{2, n+1} t^{-1} & \text{if } i = 0, \\ E_{i+2, i+1} & \text{if } i \neq 0, \end{cases}
\end{aligned}$$

and

$$\begin{aligned}\Psi(H_{i,1}) &= H_{i+1,1} + \hbar \sum_{s \geq 0} E_{1,i+1} t^{-s-1} E_{i+1,1} t^{s+1} - \hbar \sum_{s \geq 0} E_{1,i+2} t^{-s-1} E_{i+2,1} t^{s+1}, \\ \Psi(X_{i,1}^+) &= X_{i+1,1}^+ + \hbar \sum_{s \geq 0} E_{1,i+2} t^{-s-1} E_{i+1,1} t^{s+1}, \\ \Psi(X_{i,1}^-) &= X_{i+1,1}^- + \hbar \sum_{s \geq 0} E_{1,i+1} t^{-s-1} E_{i+2,1} t^{s+1}\end{aligned}$$

for $i \neq 0$. In particular, we have

$$\Psi(\tilde{H}_{i,1}) = \tilde{H}_{i+1,1} + \hbar \sum_{s \geq 0} E_{1,i+1} t^{-s-1} E_{i+1,1} t^{s+1} - \hbar \sum_{s \geq 0} E_{1,i+2} t^{-s-1} E_{i+2,1} t^{s+1} \text{ for } i \neq 0.$$

Corollary 3.2. *The following relations hold:*

$$\begin{aligned}\Psi(H_{0,1}) &= H_{0,1} + H_{1,1} + \hbar H_{0,0} H_{1,0} + \frac{\hbar}{2} H_{0,0} \\ &\quad - \hbar \sum_{s \geq 0} E_{1,2} t^{-s-1} E_{2,1} t^{s+1} + \hbar \sum_{s \geq 0} E_{1,n+1} t^{-s-1} E_{n+1,1} t^{s+1}, \\ \Psi(X_{0,1}^+) &= [X_{0,0}^+, X_{1,1}^+] + \hbar \sum_{s \geq 0} E_{1,2} t^{-s-1} E_{n+1,1} t^{s+2}, \\ \Psi(X_{i,1}^-) &= [X_{1,0}^-, X_{0,1}^-] + \hbar \sum_{s \geq 0} E_{1,n+1} t^{-s-2} E_{2,1} t^{s+1}.\end{aligned}$$

In particular, we obtain

$$\begin{aligned}\Psi(\tilde{H}_{0,1}) &= \tilde{H}_{0,1} + \tilde{H}_{1,1} + \frac{\hbar}{2} H_{0,0} \\ &\quad - \hbar \sum_{s \geq 0} E_{1,2} t^{-s-1} E_{2,1} t^{s+1} + \hbar \sum_{s \geq 0} E_{1,n+1} t^{-s-1} E_{n+1,1} t^{s+1}.\end{aligned}$$

Proof. First, we show the relation for $\Psi(X_{0,1}^+)$. By (2.6) and the definition of $\Psi(\hat{H}_{1,1})$, we have

$$\begin{aligned}\Psi(X_{0,1}^+) &= -[\tilde{H}_{2,1}, E_{n+1,2}t] \\ &\quad - [\hbar \sum_{s \geq 0} E_{1,2} t^{-s-1} E_{2,1} t^{s+1}, E_{n+1,2}t] + [\hbar \sum_{s \geq 0} E_{1,3} t^{-s-1} E_{3,1} t^{s+1}, E_{n+1,2}t] \\ &= -[\tilde{H}_{2,1}, E_{n+1,2}t] + \hbar \sum_{s \geq 0} E_{1,2} t^{-s-1} E_{n+1,1} t^{s+2}.\end{aligned} \tag{3.3}$$

By (2.6), we obtain

$$-[\tilde{H}_{2,1}, E_{n+1,2}t] = -[\tilde{H}_{2,1}, [X_{0,0}^+, X_{1,0}^+]] = [X_{0,0}^+, X_{1,1}^+].$$

Thus, we have proven the relation for $\Psi(X_{0,1}^+)$. Similarly, we can obtain the relation for $\Psi(X_{0,1}^-)$.

Next, we show the relation for $\Psi(H_{0,1})$. By (2.4) and the relation for $\Psi(X_{0,1}^+)$, we obtain

$$\begin{aligned}\Psi(H_{0,1}) &= [[X_{0,0}^+, X_{1,1}^+], [X_{1,0}^-, X_{0,0}^-]] + [\hbar \sum_{s \geq 0} E_{1,2} t^{-s-1} E_{n+1,1} t^{s+2}, E_{2,n+1} t^{-1}] \\ &= [[X_{0,0}^+, X_{1,1}^+], [X_{1,0}^-, X_{0,0}^-]] \\ &\quad - \hbar \sum_{s \geq 0} E_{1,2} t^{-s-1} E_{2,1} t^{s+1} + \hbar \sum_{s \geq 0} E_{1,n+1} t^{-s-2} E_{n+1,1} t^{s+2}.\end{aligned} \tag{3.4}$$

By (2.2)-(2.6), we can rewrite the first term of (3.4) as follows:

$$\begin{aligned}
[[X_{0,0}^+, X_{1,1}^+], [X_{1,0}^-, X_{0,0}^-]] &= [[X_{0,0}^+, H_{1,1}], X_{0,0}^-] + [X_{1,0}^-, [H_{0,0}, X_{1,1}^+]] \\
&= -[[\tilde{H}_{1,1} + \frac{\hbar}{2}H_{1,0}^2, X_{0,0}^+], X_{0,0}^-] + H_{1,1} \\
&= H_{0,1} + H_{1,1} + \frac{\hbar}{2}[\{H_{1,0}, X_{0,0}^+\}, X_{0,0}^-] \\
&= H_{0,1} + H_{1,1} + \hbar H_{0,0}H_{1,0} + \frac{\hbar}{2}\{X_{0,0}^-, X_{0,0}^+\} \\
&= H_{0,1} + H_{1,1} + \hbar H_{0,0}H_{1,0} + \hbar X_{0,0}^- X_{0,0}^+ + \frac{\hbar}{2}H_{0,0}. \tag{3.5}
\end{aligned}$$

By applying (3.5) to (3.4), we obtain the relation for $\Psi(H_{0,1})$. \square

The proof of Theorem 3.1. The proof of Theorem 3.1 is similar to Theorem 3.1 in [20]. In this article, we will show the compatibility with (2.2) and (2.10). The other cases can be proven in a similar way to [20].

3.1 Compatibility of (2.10)

We only show the + case. The - case can be proven in the same way. By the definition of Ψ , we have

$$\begin{aligned}
[\Psi(X_{n-1,1}^+), \Psi(X_{0,0}^+)] &= [X_{n,1}^+, [X_{0,0}^+, X_{1,0}^+]] + [\hbar \sum_{s \geq 0} E_{1,n+1} t^{-s-1} E_{n,1} t^{s+1}, E_{n+1,2} t] \\
&= [X_{n,1}^+, [X_{0,0}^+, X_{1,0}^+]] + \hbar \sum_{s \geq 0} E_{1,2} t^{-s} E_{n,1} t^{s+1}
\end{aligned}$$

and

$$\begin{aligned}
[\Psi(X_{0,1}^+), \Psi(X_{n-1,0}^+)] &= [[X_{0,0}^+, X_{1,1}^+], X_{n,0}^+] + [\hbar \sum_{s \geq 0} E_{1,2} t^{-s-1} E_{n+1,1} t^{s+2}, E_{n,n+1}] \\
&= [[X_{0,0}^+, X_{1,1}^+], X_{n,0}^+] - \hbar \sum_{s \geq 0} E_{1,2} t^{-s-1} E_{n,1} t^{s+2}.
\end{aligned}$$

Then, by a direct computation, we obtain

$$\begin{aligned}
[\Psi(X_{0,1}^\pm), \Psi(X_{n-1,0}^\pm)] &- [\Psi(X_{n-1,1}^+), \Psi(X_{0,0}^+)] \\
&= [[X_{0,0}^+, X_{1,1}^+], X_{n,0}^+] + [X_{n,1}^+, [X_{0,0}^+, X_{1,0}^+]] + \hbar E_{1,2} E_{n,1} t. \tag{3.6}
\end{aligned}$$

By (2.9), (2.13) and (2.10), we obtain

$$\begin{aligned}
&[X_{n,1}^+, [X_{0,0}^+, X_{1,0}^+]] + [[X_{0,0}^+, X_{1,1}^+], X_{n,0}^+] \\
&= [X_{n,1}^+, [X_{0,0}^+, X_{1,0}^+]] + [[X_{0,1}^+, X_{1,0}^+], X_{n,0}^+] + [\frac{\hbar}{2}\{X_{0,0}^+, X_{1,0}^+\}, X_{n,0}^+] \\
&= [[X_{n,1}^+, X_{0,0}^+] + [X_{0,1}^+, X_{n,0}^+], X_{1,0}^+] + \frac{\hbar}{2}[\{X_{0,0}^+, X_{n,0}^+\}, X_{1,0}^+] \\
&= [-\frac{\hbar}{2}\{X_{0,0}^+, X_{n,0}^+\} + (\varepsilon + \frac{n+1}{2}\hbar)[X_{0,0}^+, X_{n,0}^+], X_{1,0}^+] + \frac{\hbar}{2}[\{X_{0,0}^+, X_{n,0}^+\}, X_{1,0}^+] \\
&= -\frac{\hbar}{2}\{E_{n+1,2} t, E_{n,n+1}\} + (\varepsilon + \frac{n+1}{2}\hbar)[E_{n+1,2} t, E_{n,n+1}] - \frac{\hbar}{2}\{E_{n,1} t, E_{1,2}\}. \tag{3.7}
\end{aligned}$$

By applying (3.7) to (3.6), we obtain

$$\begin{aligned}
[\Psi(X_{0,1}^\pm), \Psi(X_{n-1,0}^\pm)] &- [\Psi(X_{n-1,1}^+), \Psi(X_{0,0}^+)] \\
&= -\frac{\hbar}{2}\{E_{n+1,2} t, E_{n,n+1}\} + (\varepsilon + \frac{n+1}{2}\hbar + \frac{\hbar}{2})[E_{n+1,2} t, E_{n,n+1}].
\end{aligned}$$

Thus, we have proven the compatibility with (2.10).

3.2 The compatibility with (2.2)

By the definition of Ψ , it is enough to show the relation $[\Psi(\tilde{H}_{i,1}), \Psi(\tilde{H}_{j,1})] = 0$. We only show the case that $i, j \neq 0$. The other cases can be proven in a similar way. Let us set

$$R_i = \hbar \sum_{s \geq 0} E_{1,1+i} t^{-s-1} E_{1+i,1} t^{s+1}.$$

By the definition of $J(h_i)$ and Ψ , we have

$$\begin{aligned} & [\Psi(\tilde{H}_{i,1}), \Psi(\tilde{H}_{j,1})] = [\tilde{H}_{i,1} + R_i - R_{i+1}, \tilde{H}_{j,1} + R_j - R_{j+1}] \\ & = 0 + [\tilde{H}_{1+i,1}, R_j - R_{j+1}] + [R_i - R_{i+1}, \tilde{H}_{1+j,1}] + [R_i - R_{i+1}, R_j, R_{j+1}] \\ & = [J(h_{1+i}) - A_{1+i} + A_{2+i}, R_j - R_{j+1}] \\ & \quad + [R_i - R_{i+1}, J(h_{1+j}) - A_{1+j} + A_{2+j}] + [R_i - R_{i+1}, R_j - R_{j+1}] \\ & = [-A_{1+i} + A_{2+i}, R_j - R_{j+1}] + [R_i - R_{i+1}, -A_{1+j} + A_{2+j}] + [R_i - R_{i+1}, R_j - R_{j+1}], \end{aligned} \quad (3.8)$$

where the last equality is due to Lemma 2.15. Thus, it is enough to show the relation

$$-[A_{1+i}, R_j] + [A_{1+j}, R_i] + [R_i, R_j] = 0.$$

We will compute all terms of the right hand side of (3.2). By a direct computation, we obtain

$$\begin{aligned} & [R_i, R_j] \\ & = \hbar^2 \sum_{s, v \geq 0} E_{1,1+i} t^{-s-1} (E_{1+i,1+j} t^{s-v}) E_{1+j,1} t^{v+1} - \hbar^2 \sum_{s, v \geq 0} E_{1,1+j} t^{-v-1} (E_{1+j,1+i} t^{v-s}) E_{1+i,1} t^{s+1}. \end{aligned} \quad (3.9)$$

By the definition of A_i , we obtain

$$\begin{aligned} [A_{1+i}, R_j] & = \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ u > 1+i}} E_{u,1+i} t^{-s} E_{1+i,u} t^s, R_j \right] - \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ 1+i > u}} E_{1+i,u} t^{-s} E_{u,1+i} t^s, R_j \right] \\ & \quad + \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ u < 1+i}} E_{u,1+i} t^{-s-1} E_{1+i,u} t^{s+1}, R_j \right] - \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ 1+i < u}} E_{1+i,u} t^{-s-1} E_{u,1+i} t^{s+1}, R_j \right]. \end{aligned} \quad (3.10)$$

We compute the right hand side of (3.10). By (A.4) and (A.5), we obtain

$$\begin{aligned} (3.10)_1 & = -\frac{\hbar^2}{2} \delta_{i,j} \sum_{\substack{s, v \geq 0 \\ u > 1+i}} E_{u,1+i} t^{-s} E_{1,u} t^{s-v-1} E_{1+j,1} t^{v+1} \\ & \quad - \frac{\hbar^2}{2} \delta(j > i) \sum_{s, v \geq 0} E_{1,1+i} t^{-s-v-1} E_{1+i,1+j} t^s E_{1+j,1} t^{v+1} \\ & \quad + \frac{\hbar^2}{2} \delta(j > i) \sum_{s, v \geq 0} E_{1,1+j} t^{-v-1} E_{1+j,1+i} t^{-s} E_{1+i,1} t^{s+v+1} \\ & \quad + \frac{\hbar^2}{2} \delta_{i,j} \sum_{\substack{s, v \geq 0 \\ u > 1+i}} E_{1,1+j} t^{-v-1} E_{u,1} t^{v-s+1} E_{1+i,u} t^s, \end{aligned} \quad (3.11)$$

$$(3.10)_2 = \frac{\hbar^2}{2} \delta(i > j) \sum_{s, v \geq 0} E_{1+i,1+j} t^{-s} E_{1,1+i} t^{s-v-1} E_{1+j,1} t^{v+1}$$

$$\begin{aligned}
& -\frac{\hbar^2}{2} \sum_{s,v \geq 0} E_{1+i,1+j} t^{-s-v-1} E_{1,1+i} t^s E_{1+j,1} t^{v+1} \\
& + \frac{\hbar^2}{2} \delta_{i,j} \sum_{\substack{s,v \geq 0 \\ 1+i > u}} E_{1,u} t^{-s-v-1} E_{u,1+i} t^s E_{1+j,1} t^{v+1} \\
& - \frac{\hbar^2}{2} \delta_{i,j} \sum_{\substack{s,v \geq 0 \\ 1+i > u}} E_{1,1+j} t^{-v-1} E_{1+i,u} t^{-s} E_{u,1} t^{s+v+1} \\
& + \frac{\hbar^2}{2} \sum_{s,v \geq 0} E_{1,1+j} t^{-v-1} E_{1+i,1} t^{-s} E_{1+j,1+i} t^{s+v+1} \\
& - \frac{\hbar^2}{2} \delta(i > j) \sum_{s,v \geq 0} E_{1,1+j} t^{-v-1} E_{1+i,1} t^{v-s+1} E_{1+j,1+i} t^s, \tag{3.12}
\end{aligned}$$

$$\begin{aligned}
(3.10)_3 & = \frac{\hbar^2}{2} \sum_{s,v \geq 0} E_{1,1+i} t^{-s-1} E_{1+i,1+j} t^{s-v} E_{1+j,1} t^{v+1} \\
& - \frac{\hbar}{2} \delta_{i,j} \sum_{s,v \geq 0} E_{1,1+i} t^{-s-1} E_{1,1} t^{s-v} E_{1+j,1} t^{v+1} \\
& - \frac{\hbar^2}{2} \delta(j < i) \sum_{s,v \geq 0} E_{1,1+i} t^{-s-v-2} E_{1+i,1+j} t^{s+1} E_{1+j,1} t^{v+1} \\
& + \frac{\hbar^2}{2} \delta(j < i) \sum_{s,v \geq 0} E_{1,1+j} t^{-v-1} E_{1+j,1+i} t^{-s-1} E_{1+i,1} t^{s+v+2} \\
& + \frac{\hbar^2}{2} \sum_{\substack{s,v \geq 0 \\ u < 1+i}} E_{1,1+j} t^{-v-1} E_{u,1} t^{v-s} E_{1+i,u} t^{s+1} \\
& - \frac{\hbar^2}{2} \delta_{i,j} \sum_{s,v \geq 0} E_{1,1+j} t^{-v-1} E_{1+j,1+i} t^{v-s} E_{1+i,1} t^{s+1}, \tag{3.13}
\end{aligned}$$

$$\begin{aligned}
(3.10)_4 & = \frac{\hbar^2}{2} \delta(i < j) \sum_{s,v \geq 0} E_{1+i,1+j} t^{-s-1} E_{1,1+i} t^{s-v} E_{1+j,1} t^{v+1} \\
& + \frac{\hbar^2}{2} \delta_{i,j} \sum_{\substack{s,v \geq 0 \\ 1+i < u}} E_{1,u} t^{-s-v-1} E_{u,1+i} t^s E_{1+j,1} t^{v+1} \\
& - \frac{\hbar^2}{2} \delta_{i,j} \sum_{\substack{s,v \geq 0 \\ 1+i < u}} E_{1,1+j} t^{-v-1} E_{1+i,u} t^{-s-1} E_{u,1} t^{s+v+2} \\
& - \frac{\hbar^2}{2} \delta(i < j) \sum_{s,v \geq 0} E_{1,1+j} t^{-v-1} E_{1+i,1} t^{v-s} E_{1+j,1+i} t^{s+1}. \tag{3.14}
\end{aligned}$$

Considering the sum $-[A_{1+i}, R_j] + [A_{1+j}, R_i]$, the terms containing $\delta_{i,j}$ in (3.11)-(3.14) vanish each other. Hereafter, in order to simplify the computation, we will denote the i -th term of the right hand side of the equation (\cdot) by $(\cdot)_i$. We divide the following terms into two pieces:

$$\begin{aligned}
(3.12)_1 & = \frac{\hbar^2}{2} \delta(i > j) \sum_{s,v \geq 0} E_{1+i,1+j} t^{-s} E_{1,1+i} t^{-v-1} E_{1+j,1} t^{s+v+1} \\
& + \frac{\hbar^2}{2} \delta(i > j) \sum_{s,v \geq 0} E_{1+i,1+j} t^{-s-v-1} E_{1,1+i} t^s E_{1+j,1} t^{v+1}, \tag{3.15}
\end{aligned}$$

$$\begin{aligned}
(3.12)_4 &= -\frac{\hbar^2}{2}\delta(i > j) \sum_{s,v \geq 0} E_{1,1+j}t^{-v-1}E_{1+i,1}t^{-s}E_{1+j,1+i}t^{s+v+1} \\
&\quad - \frac{\hbar^2}{2}\delta(i > j) \sum_{s,v \geq 0} E_{1,1+j}t^{-v-s-1}E_{1+i,1}t^{v+1}E_{1+j,1+i}t^s, \tag{3.16}
\end{aligned}$$

$$\begin{aligned}
(3.14)_1 &= \frac{\hbar^2}{2}\delta(i < j) \sum_{s,v \geq 0} E_{1+i,1+j}t^{-s-1}E_{1,1+i}t^{-v-1}E_{1+j,1}t^{s+2+v} \\
&\quad + \frac{\hbar^2}{2}\delta(i < j) \sum_{s,v \geq 0} E_{1+i,1+j}t^{-s-1-v}E_{1,1+i}t^sE_{1+j,1}t^{v+1}, \tag{3.17}
\end{aligned}$$

$$\begin{aligned}
(3.14)_4 &= -\frac{\hbar^2}{2}\delta(i < j) \sum_{s,v \geq 0} E_{1,1+j}t^{-v-s-2}E_{1+i,1}t^{v+1}E_{1+j,i}t^{s+1} \\
&\quad - \frac{\hbar^2}{2}\delta(i < j) \sum_{s,v \geq 0} E_{1,1+j}t^{-v-1}E_{1+i,1}t^{-s}E_{1+j,1+i}t^{v+s+1}. \tag{3.18}
\end{aligned}$$

In this proof, we denote, we denote the result of substituting $i = k$ and $j = l$ into the equation (\cdot) as $(\cdot)_{k,l}$. By a direct computation, we obtain

$$(3.11)_{2,i,j} - (3.16)_{2,j,i} = -\frac{\hbar^2}{2}\delta(j > i) \sum_{s \geq 0} (s+1)E_{1,1+i}t^{-s-1}E_{1+i,1}t^{s+1}, \tag{3.19}$$

$$(3.11)_{3,i,j} - (3.15)_{1,j,i} = \frac{\hbar^2}{2}\delta(j > i) \sum_{s \geq 0} (s+1)E_{1,1+i}t^{-s-1}E_{1+i,1}t^{s+1}, \tag{3.20}$$

$$(3.15)_{2,i,j} + (3.17)_{2,i,j} = \frac{\hbar^2}{2}\delta(i \neq j) \sum_{s,v \geq 0} E_{1+i,1+j}t^{-s-v-1}E_{1,1+i}t^sE_{1+j,1}t^{v+1}, \tag{3.21}$$

$$(3.16)_{1,i,j} + (3.18)_{2,i,j} = -\frac{\hbar^2}{2}\delta(i \neq j) \sum_{s,v \geq 0} E_{1,1+j}t^{-v-1}E_{1+i,1}t^{-s}E_{1+j,1+i}t^{s+v+1}, \tag{3.22}$$

$$(3.13)_{3,i,j} - (3.18)_{1,j,i} = -\frac{\hbar^2}{2}\delta(j < i) \sum_{s \geq 0} (s+1)E_{1,1+i}t^{-s-2}E_{1+i,1}t^{s+2}, \tag{3.23}$$

$$(3.13)_{4,i,j} - (3.17)_{1,j,i} = \frac{\hbar^2}{2}\delta(j < i) \sum_{s \geq 0} E_{1,1+i}t^{-s-2}E_{1+i,1}t^{s+2}. \tag{3.24}$$

Since $(3.19) + (3.20) = 0$, $(3.23) + (3.24) = 0$ and

$$(3.12)_2 + (3.21) = \frac{\hbar^2}{2}\delta_{i,j} \sum_{s,v \geq 0} E_{1+i,1+j}t^{-s-v-1}E_{1,1+i}t^sE_{1+j,1}t^{v+1},$$

$$(3.12)_5 + (3.22) = -\frac{\hbar^2}{2}\delta_{i,j} \sum_{s,v \geq 0} E_{1,1+j}t^{-v-1}E_{1+i,1}t^{-s}E_{1+j,1+i}t^{s+v+1}$$

hold by a direct computation, we have

$$-[A_{1+i}, R_j] + [A_{1+j}, R_i] = -(3.13)_{1,i,j} + (3.13)_{6,j,i} + (3.13)_{1,j,i} - (3.13)_{6,i,j}. \tag{3.25}$$

Since we obtain

$$(3.9)_1 - (3.13)_{1,i,j} + (3.13)_{6,j,i} = 0,$$

$$(3.9)_2 + (3.13)_{1,j,i} - (3.13)_{6,i,j} = 0$$

by a direct computation, we find that the right hand side of (3.2) is equal to zero. \square

4 Two homomorphisms from the affine Yangian associated with $\widehat{\mathfrak{sl}}(n)$ to the affine Yangian associated with $\widehat{\mathfrak{sl}}(n+1)$

In [20], the author constructed a homomorphism from $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$ to $\widetilde{Y}_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n+1))$, which is different from the one in Theorem 3.1.

Theorem 4.1 (Theorem 3.1 in [20]). *There exists a homomorphism*

$$\widetilde{\Psi}: Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n)) \rightarrow \widetilde{Y}_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n+1))$$

defined by

$$\widetilde{\Psi}(X_{i,0}^+) = \begin{cases} E_{n,1}t & \text{if } i = 0, \\ E_{i,i+1} & \text{if } i \neq 0, \end{cases} \quad \widetilde{\Psi}(X_{i,0}^-) = \begin{cases} E_{1,n}t^{-1} & \text{if } i = 0, \\ E_{i+1,i} & \text{if } i \neq 0, \end{cases}$$

and

$$\widetilde{\Psi}(H_{i,1}) = H_{i,1} - \hbar \sum_{s \geq 0} E_{i,n+1}t^{-s-1}E_{n+1,i}t^{s+1} + \hbar \sum_{s \geq 0} E_{i+1,n+1}t^{-s-1}E_{n+1,i+1}t^{s+1}$$

for $i \neq 0$.

Let us take integers $m, n \geq 3$. Combining the homomorphisms $\widetilde{\Psi}$ in Theorem 4.1 for $n, \dots, m+n-1$, we obtain a homomorphism

$$\Psi_1: Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n)) \rightarrow \widetilde{Y}_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(m+n))$$

given by

$$\Psi_1(X_{i,0}^+) = \begin{cases} E_{n,1}t & \text{if } i = 0, \\ E_{i,i+1} & \text{if } i \neq 0, \end{cases} \quad \Psi_1(X_{i,0}^-) = \begin{cases} E_{1,n}t^{-1} & \text{if } i = 0, \\ E_{i+1,i} & \text{if } i \neq 0, \end{cases}$$

and

$$\Psi_1(H_{i,1}) = H_{i,1} - \hbar \sum_{s \geq 0} \sum_{k=n+1}^{m+n} E_{i,k}t^{-s-1}E_{k,i}t^{s+1} + \hbar \sum_{s \geq 0} \sum_{k=n+1}^{m+n} E_{i+1,k}t^{-s-1}E_{k,i+1}t^{s+1}$$

for $i \neq 0$. Similarly, by combining the homomorphisms given in Theorem 3.1 for $m, \dots, m+n-1$, we obtain a homomorphism

$$\Psi_2: Y_{\hbar,\varepsilon+n\hbar}(\widehat{\mathfrak{sl}}(m)) \rightarrow \widetilde{Y}_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(m+n))$$

determined by

$$\Psi_2(X_{i,0}^+) = \begin{cases} E_{m+n,n+1}t & \text{if } i = 0, \\ E_{n+i,n+i+1} & \text{if } i \neq 0, \end{cases} \quad \Psi_2(X_{i,0}^-) = \begin{cases} E_{n+1,m+n}t^{-1} & \text{if } i = 0, \\ E_{n+i+1,n+i} & \text{if } i \neq 0, \end{cases}$$

and

$$\Psi_2(H_{i,1}) = H_{i+n,1} + \hbar \sum_{s \geq 0} \sum_{k=1}^n E_{k,n+i}t^{-s-1}E_{n+i,k}t^{s+1} - \hbar \sum_{s \geq 0} \sum_{k=1}^n E_{k,n+i+1}t^{-s-1}E_{n+i+1,k}t^{s+1}$$

for $i \neq 0$.

Theorem 4.2. *The homomorphisms Ψ_1 and Ψ_2 induce a homomorphism*

$$Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n)) \otimes Y_{\hbar,\varepsilon+n\hbar}(\widehat{\mathfrak{sl}}(m)) \rightarrow \widetilde{Y}_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(m+n)).$$

Proof. Let us set

$$P_i = \hbar \sum_{s \geq 0} \sum_{k=n+1}^{m+n} E_{i,k} t^{-s-1} E_{k,i} t^{s+1}, \quad Q_i = \hbar \sum_{s \geq 0} \sum_{k=1}^n E_{k,n+i} t^{-s} E_{n+i,k} t^s$$

We fix integers $1 \leq i \leq n$ and $1 \leq j \leq m-1$. The affine Yangian $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n))$ (resp. $Y_{\hbar, \varepsilon+n\hbar}(\widehat{\mathfrak{sl}}(m))$) can be generated by $\tilde{H}_{i,1}$ (resp. $\tilde{H}_{j,1}$) and $\{X_{k,0}^{\pm} \mid 0 \leq k \leq n-1\}$ (resp. $\{X_{l,0}^{\pm} \mid 0 \leq l \leq m-1\}$). Thus, it is enough to show the commutativity between $\Psi_1(\tilde{H}_{i,1})$, $\Psi_1(X_{k,0}^{\pm})$ and $\Psi_2(\tilde{H}_{j,1})$, $\Psi_2(X_{l,0}^{\pm})$. The commutativity between $\Psi_1(X_{k,0}^{\pm})$ and $\Psi_2(X_{l,0}^{\pm})$ is obvious. Thus, we will show the other cases in the following three subsections.

4.1 Commutativity between $\Psi_1(\tilde{H}_{i,1})$ and $\Psi_2(X_{l,0}^{\pm})$

We only show the + case. The - case can be proven in a similar way. The case that $l \neq 0$ comes from the definition of $\Psi_1(\tilde{H}_{i,1})$. We will consider the case that $l = 0$. By a direct computation, we obtain

$$[\Psi_1(\tilde{H}_{i,1}), E_{m+n, n+1} t] = [J(h_i), E_{m+n, n+1} t] - [A_i - A_{i+1}, E_{m+n, n+1} t] - [P_i - P_{i+1}, E_{m+n, n+1} t]. \quad (4.3)$$

Since $E_{m+n, n+1} t = [\cdots [X_{0,0}^+, X_{1,0}^+], X_{2,0}^+, \cdots, X_{n,0}^+] = 0$ holds, we find that $[J(h_i), E_{m+n, n+1} t] = 0$ by Lemma 2.15. By the definition of A_i , we obtain

$$\begin{aligned} [A_i, E_{m+n, n+1} t] &= \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ u > i}} E_{u,i} t^{-s} E_{i,u} t^s, E_{m+n, n+1} t \right] - \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ i < v}} E_{i,v} t^{-s-1} E_{v,i} t^{s+1}, E_{m+n, n+1} t \right] \\ &= -\frac{\hbar}{2} E_{m+n, i} t E_{i, n+1} - \frac{\hbar}{2} E_{i, n+1} E_{m+n, i} t. \end{aligned} \quad (4.4)$$

By the definition of P_i , we have

$$[P_i, E_{m+n, n+1} t] = \left[\hbar \sum_{s \geq 0} \sum_{k=n+1}^{m+n} E_{i,k} t^{-s-1} E_{k,i} t^{s+1}, E_{m+n, n+1} t \right] = \hbar E_{i, n+1} E_{m+n, i} t. \quad (4.5)$$

By applying (4.4) and (4.5) to (4.3), we obtain

$$[\Psi(\tilde{H}_{i,1}), E_{m+n, n+1} t] = 0 - \frac{\hbar}{2} [E_{i, n+1}, E_{m+n, i} t] + \frac{\hbar}{2} [E_{i+1, n+1}, E_{m+n, i+1} t] = 0.$$

4.2 Commutativity between $\Psi_2(\tilde{H}_{j,1})$ and $\Psi_1(X_{k,0}^+)$

We only show the + case. The - case can be proven in a similar way. The case that $k \neq 0$ comes from the definition of $\Psi_2(\tilde{H}_{j,1})$. We will consider the case that $k = 0$.

By a direct computation, we obtain

$$[\Psi_2(\tilde{H}_{j,1}), E_{n,1} t] = [J(h_{j+n}), E_{n,1} t] - [A_{j+n} - A_{j+n+1}, E_{n,1} t] + [Q_i - Q_{i+1}, E_{n,1} t]. \quad (4.6)$$

Since $E_{n,1} t = [X_{n,0}^+, [\cdots, [X_{n-1,0}^+, X_{0,0}^+] \cdots]]$ holds, we obtain $[J(h_{j+n}), E_{n,1} t] = 0$ by Lemma 2.15. By the definition of A_i , we have

$$\begin{aligned} [A_{j+n}, E_{n,1} t] &= -\left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ j+n > v}} E_{j+n, v} t^{-s} E_{v, j+n} t^s, E_{n,1} t \right] + \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ u < j+n}} E_{u, j+n} t^{-s-1} E_{j+n, u} t^{s+1}, E_{n,1} t \right] \\ &= -\frac{\hbar}{2} E_{j+n, 1} t E_{n, j+n} - \frac{\hbar}{2} E_{n, j+n} E_{j+n, 1} t. \end{aligned} \quad (4.7)$$

By the definition of Q_i , we obtain

$$[Q_j, E_{n,1}t] = [\hbar \sum_{s \geq 0} \sum_{k=1}^n E_{k,j+n} t^{-s-1} E_{j+n,k} t^{s+1}, E_{n,1}t] = -\hbar E_{n,j+n} E_{j+n,1}t. \quad (4.8)$$

Applying (4.7) and (4.8) to (4.6), we obtain

$$[\Psi_2(\tilde{H}_{i,1}), E_{n,1}t] = 0 + \frac{\hbar}{2}[E_{n,j+n}, E_{j+n,1}t] - \frac{\hbar}{2}[E_{n,j+n+1}, E_{j+n+1,1}t] = 0.$$

4.3 Commutativity between $\Psi_1(\tilde{H}_{i,1})$ and $\Psi_2(\tilde{H}_{j,1})$

By (2.2) and the definition of $J(h_i)$, Ψ_1 and Ψ_2 , we have

$$\begin{aligned} & [\Psi_1(\tilde{H}_{i,1}), \Psi_2(\tilde{H}_{j,1})] \\ &= [\tilde{H}_{i,1}, \tilde{H}_{j+n,1}] - [P_i, \tilde{H}_{j+n,1}] + [\tilde{H}_{i,1}, Q_j] - [P_i - P_{i+1}, Q_j - Q_{j+1}] \\ &= 0 - [P_i - P_{i+1}, J(h_{j+n})] + [P_i - P_{i+1}, A_{j+n} - A_{j+n+1}] \\ &\quad + [J(h_i), Q_j - Q_{j+1}] - [A_i - A_{i+1}, Q_j - Q_{j+1}] - [P_i - P_{i+1}, Q_j - Q_{j+1}]. \end{aligned}$$

By Lemma 2.15, we obtain $-[P_i - P_{i+1}, J(h_{j+n})] + [J(h_i), Q_j - Q_{j+1}] = 0$. Thus, it is enough to show the relation

$$[P_i, Q_j] + [A_i, Q_j] + [A_{j+n}, P_i] = 0. \quad (4.9)$$

We will compute each terms of the left hand side of (4.9). By a direct computation, we obtain

$$\begin{aligned} & [P_i, Q_j] \\ &= \hbar^2 \sum_{s,u \geq 0} \sum_{k=1}^{m+n} E_{i,k} t^{-s-1} E_{k,j+n} t^{-u-1} E_{j+n,i} t^{s+u+2} \\ &\quad - \hbar^2 \sum_{s,u \geq 0} \sum_{k=1}^{m+n} E_{i,j+n} t^{-u-s-2} E_{j+n,k} t^{u+1} E_{k,i} t^{s+1} \\ &\quad + \hbar^2 \sum_{s,u \geq 0} \sum_{k=n+1}^{m+n} E_{i,k} t^{-s-u-1} E_{k,j+n} t^s E_{j+n,i} t^{u+1} - \hbar^2 \sum_{s,u \geq 0} \sum_{l=1}^n E_{i,j+n} t^{-s-1} E_{l,i} t^{-u} E_{j+n,l} t^{s+u+1} \\ &\quad + \hbar^2 \sum_{s,u \geq 0} \sum_{l=1}^n E_{l,j+n} t^{-u-s-1} E_{i,l} t^u E_{j+n,i} t^{s+1} \\ &\quad - \hbar^2 \sum_{s,u \geq 0} \sum_{k=n+1}^{m+n} E_{i,j+n} t^{-u-1} E_{j+n,k} t^{-s} E_{k,i} t^{u+s+1}. \end{aligned} \quad (4.10)$$

By the definition of A_i , we can divide $[A_i, P_j]$ into four pieces:

$$\begin{aligned} [A_{j+n}, P_i] &= \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ u > j+n}} E_{u,j+n} t^{-s} E_{j+n,u} t^s, P_i \right] - \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ j+n > u}} E_{j+n,u} t^{-s} E_{u,j+n} t^s, P_i \right] \\ &\quad + \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ u < j+n}} E_{u,j+n} t^{-s-1} E_{j+n,u} t^{s+1}, P_i \right] - \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ j+n < u}} E_{j+n,u} t^{-s-1} E_{u,j+n} t^{s+1}, P_i \right]. \end{aligned} \quad (4.11)$$

We compute the right hand side of (4.11). By (A.2) and (A.3), we obtain

$$\left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ u > j+n}} E_{u,j+n} t^{-s} E_{j+n,u} t^s, P_i \right]$$

$$= -\frac{\hbar^2}{2} \sum_{\substack{s,v \geq 0 \\ u > j+n}} E_{u,j+n} t^{-s-v-1} E_{i,u} t^s E_{j+n,i} t^{v+1} + \frac{\hbar^2}{2} \sum_{\substack{s,v \geq 0 \\ u > j+n}} E_{i,j+n} t^{-v-1} E_{u,i} t^{-s} E_{j+n,u} t^{s+v+1}, \quad (4.12)$$

$$\begin{aligned} & - \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ j+n > u}} E_{j+n,u} t^{-s} E_{u,j+n} t^s, P_i \right] \\ & = -\frac{\hbar^2}{2} \sum_{s,v \geq 0} \sum_{u=1}^n E_{i,j+n} t^{-v-1} E_{j+n,u} t^{-s} E_{u,i} t^{s+v+1} \\ & \quad - \frac{\hbar^2}{2} \sum_{s,v \geq 0} \sum_{k=j+n}^{m+n} E_{j+n,k} t^{-s-v-1} E_{i,j+n} t^s E_{k,i} t^{v+1} \\ & \quad + \frac{\hbar^2}{2} \sum_{s,v \geq 0} \sum_{u=1}^n E_{i,u} t^{-s-v-1} E_{u,j+n} t^s E_{j+n,i} t^{v+1} \\ & \quad + \frac{\hbar^2}{2} \sum_{s,v \geq 0} \sum_{k=j+n}^{m+n} E_{i,k} t^{-v-1} E_{j+n,i} t^{-s} E_{k,j+n} t^{s+v+1}, \quad (4.13) \end{aligned}$$

$$\begin{aligned} & \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ u < j+n}} E_{u,j+n} t^{-s-1} E_{j+n,u} t^{s+1}, P_i \right] \\ & = \frac{\hbar^2}{2} \sum_{s,v \geq 0} \sum_{k=1}^{m+n} E_{i,j+n} t^{-s-2-v} E_{j+n,k} t^{s+1} E_{k,i} t^{v+1} \\ & \quad - \frac{\hbar^2}{2} \sum_{s,v \geq 0} \sum_{k=1}^{m+n} E_{i,k} t^{-v-1} E_{k,j+n} t^{-s-1} E_{j+n,i} t^{s+v+2} \\ & \quad + \frac{\hbar^2}{2} \sum_{s,v \geq 0} \sum_{k=n+1}^{m+n} E_{i,j+n} t^{-s-1} E_{j+n,k} t^{-v} E_{k,i} t^{s+v+1} - \frac{\hbar^2}{2} \sum_{\substack{s,v \geq 0 \\ u < j+n}} E_{u,j+n} t^{-s-v-1} E_{i,u} t^s E_{j+n,i} t^{v+1} \\ & \quad + \frac{\hbar^2}{2} \sum_{\substack{s,v \geq 0 \\ u < j+n}} E_{i,j+n} t^{-v-1} E_{u,i} t^{-s} E_{j+n,u} t^{s+v+1} - \frac{\hbar^2}{2} \sum_{s,v \geq 0} \sum_{k=n+1}^{m+n} E_{i,k} t^{-s-v-1} E_{k,j+n} t^v E_{j+n,i} t^{s+1} \quad (4.14) \end{aligned}$$

$$\begin{aligned} & - \left[\frac{\hbar}{2} \sum_{\substack{s,v \geq 0 \\ j+n < u}} E_{j+n,u} t^{-s-1} E_{u,j+n} t^{s+1}, P_i \right] \\ & = \frac{\hbar^2}{2} \sum_{\substack{s,v \geq 0 \\ j+n < u}} E_{j+n,u} t^{-s-v-1} E_{i,j+n} t^s E_{u,i} t^{v+1} - \frac{\hbar^2}{2} \sum_{\substack{s,v \geq 0 \\ j+n < u}} E_{i,u} t^{-v-1} E_{j+n,i} t^{-s} E_{u,j+n} t^{s+v+1}. \quad (4.15) \end{aligned}$$

By the definition of Q_i , we have

$$\begin{aligned} [A_i, Q_j] & = \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ u > i}} E_{u,i} t^{-s} E_{i,u} t^s, Q_j \right] - \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ i > u}} E_{i,u} t^{-s} E_{u,i} t^s, Q_j \right] \\ & \quad + \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ u < i}} E_{u,i} t^{-s-1} E_{i,u} t^{s+1}, Q_j \right] - \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ i < u}} E_{i,u} t^{-s-1} E_{u,i} t^{s+1}, Q_j \right]. \quad (4.16) \end{aligned}$$

We compute the right hand side of (4.16). By a direct computation, we obtain

$$\begin{aligned}
& \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ u > i}} E_{u,i} t^{-s} E_{i,u} t^s, Q_j \right] \\
&= -\frac{\hbar^2}{2} \sum_{s,v \geq 0} \sum_{l=1}^i E_{l,i} t^{-s-v-1} E_{i,j+n} t^s E_{j+n,l} t^{v+1} \\
&\quad - \frac{\hbar^2}{2} \sum_{s,v \geq 0} \sum_{u=n+1}^{m+n} E_{i,j+n} t^{-v-1} E_{u,i} t^{-s} E_{j+n,u} t^{s+v+1} \\
&\quad + \frac{\hbar^2}{2} \sum_{s,v \geq 0} \sum_{u=n+1}^{m+n} E_{u,j+n} t^{-s-v-1} E_{i,u} t^s E_{j+n,i} t^{v+1} \\
&\quad + \frac{\hbar^2}{2} \sum_{s,v \geq 0} \sum_{l=1}^i E_{l,j+n} t^{-v-1} E_{j+n,i} t^{-s} E_{i,l} t^{s+v+1}, \tag{4.17}
\end{aligned}$$

$$\begin{aligned}
& - \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ i > u}} E_{i,u} t^{-s} E_{u,i} t^s, Q_j \right] \\
&= -\frac{\hbar^2}{2} \sum_{\substack{s,v \geq 0 \\ i > u}} E_{i,u} t^{-s-v-1} E_{u,j+n} t^s E_{j+n,i} t^{v+1} + \frac{\hbar^2}{2} \sum_{\substack{s,v \geq 0 \\ i > u}} E_{i,j+n} t^{-v-1} E_{j+n,u} t^{-s} E_{u,i} t^{s+v+1}, \tag{4.18}
\end{aligned}$$

$$\begin{aligned}
& \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ u < i}} E_{u,i} t^{-s-1} E_{i,u} t^{s+1}, Q_j \right] \\
&= \frac{\hbar^2}{2} \sum_{\substack{s,v \geq 0 \\ u < i}} E_{u,i} t^{-s-1-v} E_{i,j+n} t^s E_{j+n,u} t^{v+1} - \frac{\hbar^2}{2} \sum_{\substack{s,v \geq 0 \\ u < i}} E_{u,j+n} t^{-v-1} E_{j+n,i} t^{-s} E_{i,u} t^{v+s+1}, \tag{4.19}
\end{aligned}$$

$$\begin{aligned}
& \left[\frac{\hbar}{2} \sum_{\substack{s \geq 0 \\ u < i}} E_{u,i} t^{-s-1} E_{i,u} t^{s+1}, Q_j \right] \\
&= -\frac{\hbar^2}{2} \sum_{s,v \geq 0} \sum_{l=1}^{m+n} E_{l,j+n} t^{-v-1} E_{i,l} t^{-s-1} E_{j+n,i} t^{v+s+2} \\
&\quad + \frac{\hbar^2}{2} \sum_{s,v \geq 0} \sum_{l=1}^{m+n} E_{i,j+n} t^{-s-v-2} E_{l,i} t^{s+1} E_{j+n,l} t^{v+1} \\
&\quad - \frac{\hbar^2}{2} \sum_{\substack{s,v \geq 0 \\ i < u}} E_{i,u} t^{-s-v-1} E_{u,j+n} t^s E_{j+n,i} t^{v+1} + \frac{\hbar^2}{2} \sum_{s,v \geq 0} \sum_{l=1}^n E_{i,j+n} t^{-s-1} E_{l,i} t^{-v} E_{j+n,l} t^{s+v+1} \\
&\quad - \frac{\hbar^2}{2} \sum_{s,v \geq 0} \sum_{l=1}^n E_{l,j+n} t^{-v-s-1} E_{i,l} t^v E_{j+n,i} t^{s+1} + \frac{\hbar^2}{2} \sum_{\substack{s,v \geq 0 \\ i < u}} E_{i,j+n} t^{-v-1} E_{j+n,u} t^{-s} E_{u,i} t^{v+s+1}. \tag{4.20}
\end{aligned}$$

We compute the sum of (4.10), (4.12)-(4.15) and (4.17)-(4.20) into eight pieces as follows:

$$\begin{aligned}
& (4.13)_2 + (4.15)_1 + (4.17)_1 + (4.19)_1 \\
&= -\frac{\hbar^2}{2} \sum_{s,v \geq 0} E_{j+n,j+n} t^{-s-v-1} E_{i,j+n} t^s E_{j+n,i} t^{v+1} - \frac{\hbar^2}{2} \sum_{s,v \geq 0} E_{i,i} t^{-s-v-1} E_{i,j+n} t^s E_{j+n,i} t^{v+1},
\end{aligned}$$

$$\begin{aligned}
(4.10)_2 + (4.14)_1 + (4.20)_2 &= -(m+n) \frac{\hbar^2}{2} \sum_{s \geq 0} (s+1) E_{i,j+n} t^{-s-2} E_{j+n,i} t^{s+2}, \\
(4.10)_4 + (4.12)_2 + (4.14)_5 + (4.17)_2 + (4.20)_4 \\
&= -\frac{\hbar^2}{2} \sum_{s,v \geq 0} E_{i,j+n} t^{-v-1} E_{j+n,i} t^{-s} E_{j+n,j+n} t^{s+v+1}, \\
(4.10)_6 + (4.13)_1 + (4.14)_3 + (4.18)_2 + (4.20)_6 &= -\frac{\hbar^2}{2} \sum_{s,v \geq 0} E_{i,j+n} t^{-v-1} E_{j+n,i} t^{-s} E_{i,i} t^{s+v+1}, \\
(4.10)_1 + (4.14)_2 + (4.20)_1 &= (m+n) \frac{\hbar^2}{2} \sum_{s \geq 0} (s+1) E_{i,j+n} t^{-s-2} E_{j+n,i} t^{s+2}, \\
(4.10)_3 + (4.13)_3 + (4.14)_6 + (4.18)_1 + (4.20)_3 &= \frac{\hbar^2}{2} \sum_{s,u \geq 0} E_{i,i} t^{-s-u-1} E_{i,j+n} t^s E_{j+n,i} t^{u+1}, \\
(4.10)_5 + (4.12)_1 + (4.14)_4 + (4.17)_3 + (4.20)_5 &= \frac{\hbar^2}{2} \sum_{s,u \geq 0} E_{j+n,j+n} t^{-u-s-1} E_{i,j+n} t^u E_{j+n,i} t^{s+1}, \\
(4.13)_4 + (4.15)_2 + (4.17)_4 + (4.19)_1 \\
&= \frac{\hbar^2}{2} \sum_{s \geq 0} E_{i,j+n} t^{-v-1} E_{j+n,i} t^{-s} E_{j+n,j+n} t^{s+v+1} + \frac{\hbar^2}{2} \sum_{s,v \geq 0} E_{i,j+n} t^{-v-1} E_{j+n,i} t^{-s} E_{i,i} t^{s+v+1}.
\end{aligned}$$

Since the sum of the eight equations above is equal to zero, we have shown (4.9). \square

5 Application to the evaluation map for the affine Yangian

The evaluation map for the affine Yangian is a non-trivial homomorphism from the affine Yangian $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$ to the completion of the universal enveloping algebra of the affinization of $\mathfrak{gl}(n)$. We set a Lie algebra

$$\widehat{\mathfrak{gl}}(n) = \mathfrak{gl}(n) \otimes \mathbb{C}[z^{\pm 1}] \oplus \mathbb{C}\tilde{c} \oplus \mathbb{C}z$$

whose commutator relations are given by

$$[x \otimes t^u, y \otimes t^v] = \begin{cases} [x, y] \otimes t^{u+v} + \delta_{u+v,0} \text{utr}(xy) \tilde{c} & \text{if } x, y \in \mathfrak{sl}(n), \\ [e_{a,b}, e_{i,i}] \otimes t^{u+v} + \delta_{u+v,0} \text{utr}(E_{a,b} E_{i,i}) \tilde{c} + \delta_{u+v,0} \delta_{a,b} u z & \text{if } x = e_{a,b}, y = e_{i,i}, \end{cases}$$

z and \tilde{c} are central elements of $\widehat{\mathfrak{gl}}(n)$,

where tr is a trace of $\mathfrak{gl}(n)$, that is, $\text{tr}(E_{i,j} E_{k,l}) = \delta_{i,l} \delta_{j,k}$.

We consider a completion of $U(\widehat{\mathfrak{gl}}(n))/U(\widehat{\mathfrak{gl}}(n))(z-1)$ following [16] and [12]. We take the grading of $U(\widehat{\mathfrak{gl}}(n))/U(\widehat{\mathfrak{gl}}(n))(z-1)$ as $\deg(X(s)) = s$ and $\deg(\tilde{c}) = 0$. We denote the degreewise completion of $U(\widehat{\mathfrak{gl}}(n))/U(\widehat{\mathfrak{gl}}(n))(z-1)$ by $\mathcal{U}(\widehat{\mathfrak{gl}}(n))$.

Theorem 5.1 (Theorem 3.8 in [14] and Theorem 4.18 in [13]). *1. Suppose that $\tilde{c} = \frac{\varepsilon}{\hbar}$. Then, there exists an algebra homomorphism*

$$\text{ev}_{\hbar,\varepsilon}^n : Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n)) \rightarrow \mathcal{U}(\widehat{\mathfrak{gl}}(n))$$

uniquely determined by

$$\text{ev}_{\hbar,\varepsilon}^n(X_{i,0}^+) = \begin{cases} E_{n,1} t & \text{if } i = 0, \\ E_{i,i+1} & \text{if } 1 \leq i \leq n-1, \end{cases} \quad \text{ev}_{\hbar,\varepsilon}^n(X_{i,0}^-) = \begin{cases} E_{1,n} t^{-1} & \text{if } i = 0, \\ E_{i+1,i} & \text{if } 1 \leq i \leq n-1, \end{cases}$$

$$\text{ev}_{\hbar,\varepsilon}^n(H_{i,0}) = \begin{cases} E_{n,n} - E_{1,1} + \tilde{c} & \text{if } i = 0, \\ E_{i,i} - E_{i+1,i+1} & \text{if } 1 \leq i \leq n-1. \end{cases}$$

and

$$\begin{aligned} \text{ev}_{\hbar,\varepsilon}^n(H_{i,1}) &= -\frac{i}{2}\hbar \text{ev}_{\hbar,\varepsilon}^n(H_{i,0}) - \hbar E_{i,i} E_{i+1,i+1} \\ &+ \hbar \sum_{s \geq 0} \sum_{k=1}^i E_{i,k} t^{-s} E_{k,i} t^s + \hbar \sum_{s \geq 0} \sum_{k=i+1}^n E_{i,k} t^{-s-1} E_{k,i} t^{s+1} \\ &- \hbar \sum_{s \geq 0} \sum_{k=1}^i E_{i+1,k} t^{-s} E_{k,i+1} t^s - \hbar \sum_{s \geq 0} \sum_{k=i+1}^n E_{i+1,k} t^{-s-1} E_{k,i+1} t^{s+1} \end{aligned}$$

for $i \neq 0$.

2. In the case that $\varepsilon \neq 0$, the image of the evaluation map is dense in $\mathcal{U}(\widehat{\mathfrak{gl}}(n))$.

We note that $U(\widehat{\mathfrak{gl}}(n))$ can be embedded into $U(\widehat{\mathfrak{gl}}(m+n))$ by $\tilde{c} \mapsto \tilde{c}$ and $E_{i,j} t^s \mapsto E_{i,j} t^s$ for $i \neq j$. By the definition of the evaluation map and Ψ_1 , we obtain the following theorem.

Theorem 5.2. *The following relation holds:*

$$\text{ev}_{\hbar,\varepsilon}^{m+n} \circ \Psi_1 = \text{ev}_{\hbar,\varepsilon}^n.$$

Let us set the centralizer algebra

$$C(\widehat{\mathfrak{gl}}(m+n), \widehat{\mathfrak{gl}}(n)) = \{x \in \mathcal{U}(\widehat{\mathfrak{gl}}(m+n)) \mid [x, U(\widehat{\mathfrak{gl}}(n))] = 0\}.$$

Theorem 4.2 and Theorem 5.1 induces the following corollary.

Corollary 5.3. *In the case that $\varepsilon \neq 0$, we obtain a homomorphism*

$$\text{ev}_{\hbar,\varepsilon}^{m+n} \circ \Psi_2: Y_{\hbar,\varepsilon+n\hbar}(\widehat{\mathfrak{sl}}(m)) \rightarrow C(\widehat{\mathfrak{gl}}(m+n), \widehat{\mathfrak{gl}}(n)).$$

This result can be interpreted from the perspective of a vertex algebra. For a vertex algebra V , we denote the generating field associated with $v \in V$ by $v(z) = \sum_{n \in \mathbb{Z}} v_{(n)} z^{-n-1}$. We also denote the OPE of V by

$$u(z)v(w) \sim \sum_{s \geq 0} \frac{(u_{(s)}v)(w)}{(z-w)^{s+1}}$$

for all $u, v \in V$. We denote the vacuum vector (resp. the translation operator) by $|0\rangle$ (resp. ∂).

Let us recall the definition of a universal enveloping algebra of a vertex algebra in the sense of [6] and [16]. For any vertex algebra V , let $L(V)$ be the Borcharads Lie algebra, that is,

$$L(V) = V \otimes \mathbb{C}[t, t^{-1}] / \text{Im}(\partial \otimes \text{id} + \text{id} \otimes \frac{d}{dt}), \quad (5.4)$$

where the commutation relation is given by

$$[ut^a, vt^b] = \sum_{r \geq 0} \binom{a}{r} (u_{(r)}v) t^{a+b-r}$$

for all $u, v \in V$ and $a, b \in \mathbb{Z}$.

Definition 5.5 (Section 6 in [16]). We set $\mathcal{U}(V)$ as the quotient algebra of the standard degreewise completion of the universal enveloping algebra of $L(V)$ by the completion of the two-sided ideal generated by

$$(u_{(a)}v)t^b - \sum_{i \geq 0} \binom{a}{i} (-1)^i (ut^{a-i}vt^{b+i} - (-1)^a vt^{a+b-i}ut^i), \quad (5.6)$$

$$|0\rangle t^{-1} - 1. \quad (5.7)$$

We call $\mathcal{U}(V)$ the universal enveloping algebra of V .

We denote the universal affine vertex algebra associated with a finite dimensional Lie algebra \mathfrak{g} and its inner product κ by $V^\kappa(\mathfrak{g})$. By the PBW theorem, we can identify $V^\kappa(\mathfrak{g})$ with $U(t^{-1}\mathfrak{g}[t^{-1}])$. In order to simplify the notation, here after, we denote the generating field $(ut^{-1})(z)$ as $u(z)$. By the definition of $V^\kappa(\mathfrak{g})$, the generating fields $u(z)$ and $v(z)$ satisfy the OPE

$$u(z)v(w) \sim \frac{[u, v](w)}{z-w} + \frac{\kappa(u, v)}{(z-w)^2} \quad (5.8)$$

for all $u, v \in \mathfrak{g}$. Let us set an inner product on $\mathfrak{gl}(m) \subset \mathfrak{gl}(m+n)$ by

$$\kappa(E_{i,j}, E_{p,q}) = \delta_{i,q}\delta_{p,j}\tilde{c} + \delta_{i,j}\delta_{p,q}.$$

Then, we find that $\mathcal{U}(V^\kappa(\mathfrak{gl}(m+n)))$ and $\mathcal{U}(V^\kappa(\mathfrak{gl}(n)))$ coincide with $\mathcal{U}(\widehat{\mathfrak{gl}}(m+n))$ and $\mathcal{U}(\widehat{\mathfrak{gl}}(n))$. We denote $E_{i,j}t^{-s} \in U(t^{-1}\mathfrak{gl}(n)[t^{-1}]) = V^\kappa(\mathfrak{gl}(n))$ by $E_{i,j}[-s]$.

For a vertex algebra A and its vertex subalgebra B , we set a coset vertex algebra of the pair (A, B) as follows:

$$C(A, B) = \{v \in A \mid w_{(r)}v = 0 \text{ for } w \in B \text{ and } r \geq 0\}.$$

Theorem 5.9. *The homomorphism induces the homomorphism*

$$\text{ev}_{\hbar, \varepsilon}^{m+n} \circ \Psi_2: Y_{\hbar, \varepsilon+n\hbar}(\widehat{\mathfrak{sl}}(m)) \rightarrow \mathcal{U}(C(V^\kappa(\mathfrak{gl}(m+n)), V^\kappa(\mathfrak{sl}(n)))).$$

Proof. By the definition of the universal affine vertex algebra and coset, we find that $E_{i,j}[-1]$ is contained in $C(V^\kappa(\mathfrak{gl}(m+n)), V^\kappa(\mathfrak{sl}(n)))$ for $i, j \geq n+1$. Thus, the image of $\text{ev}_{\hbar, \varepsilon}^{m+n} \circ \Psi_2$ is contained in $\mathcal{U}(C(V^\kappa(\mathfrak{gl}(m+n)), V^\kappa(\mathfrak{sl}(n))))$ if $\sum_{u=1}^n (E_{u,i}[-1])_{(-1)} E_{j,u}[-1]$ is contained in $C(V^\kappa(\mathfrak{gl}(m+n)), V^\kappa(\mathfrak{sl}(n)))$ for $i, j \geq n+1$. By a direct computation, we obtain

$$x_{(r)} \left(\sum_{u=1}^n (E_{u,i}[-1])_{(-1)} E_{j,u}[-1] \right) = 0 \text{ if } x \in \mathfrak{sl}(n) \text{ and } r \geq 0.$$

Then, we find that $\sum_{u=1}^n (E_{u,i}[-1])_{(-1)} E_{j,u}[-1]$ is contained in $C(V^\kappa(\mathfrak{gl}(m+n)), V^\kappa(\mathfrak{sl}(n)))$ for $i, j \geq n+1$. \square

6 Application to the rectangular W -algebra

The W -algebra $\mathcal{W}^k(\mathfrak{g}, f)$ is a vertex algebra associated with a finite dimensional reductive Lie algebra \mathfrak{g} and a nilpotent element f . We call the W -algebra associated with $\mathfrak{gl}(ln)$ and a nilpotent element of type (l^n) the rectangular W -algebra and denote it by $\mathcal{W}^k(\mathfrak{gl}(ln), (l^n))$. In this article, we only consider the case that $l = 2$. The nilpotent element is

$$f = \sum_{u=1}^n E_{n+u, u} \in \mathfrak{gl}(2n).$$

We set the inner product on $\mathfrak{gl}(n)$ by

$$\kappa(E_{i,j}, E_{p,q}) = \delta_{j,p}\delta_{i,q}\alpha + \delta_{i,j}\delta_{p,q},$$

where $\alpha = k + n$.

By Theorem 3.1 and Corollary 3.2 in [1], we obtain the following theorem.

Theorem 6.1 (Corollary 5.2 in [9], Theorem 3.1 and Corollary 3.2 in [1] and Section 4 in [20]). (1) The rectangular W -algebra $\mathcal{W}^k(\mathfrak{gl}(2n), (2^n))$ can be realized as a vertex subalgebra of $V^\kappa(\mathfrak{gl}(n))^{\otimes 2}$.

(2) The W -algebra $\mathcal{W}^k(\mathfrak{g}, f)$ has the following strong generators:

$$\begin{aligned} W_{i,j}^{(1)} &= E_{i,j}^{(1)}[-1] + E_{i,j}^{(2)}[-1], \\ W_{i,j}^{(2)} &= \sum_{1 \leq u \leq n} E_{u,j}^{(1)}[-1]E_{i,u}^{(2)}[-1] + \alpha E_{i,j}^{(2)}[-1] \end{aligned}$$

for $1 \leq i, j \leq n$, where $E_{i,j}^{(1)}[-1] = E_{i,j}[-1] \otimes 1 \in V^\kappa(\mathfrak{gl}(n))^{\otimes 2}$ and $E_{i,j}^{(2)}[-1] = 1 \otimes E_{i,j}[-1] \in V^\kappa(\mathfrak{gl}(n))^{\otimes 2}$.

(3) There exists the embedding determined by

$$\iota: \mathcal{W}^{k+m}(\mathfrak{gl}(2n), (2^n)) \rightarrow \mathcal{W}^k(\mathfrak{gl}(2m+2n), (2^{m+n})), \quad W_{i,j}^{(u)} \mapsto W_{i,j}^{(u)}.$$

(4) The W -algebra $\mathcal{W}^k(\mathfrak{gl}(2n), (2^n))$ has a subalgebra $\mathcal{W}^k(\mathfrak{gl}(2n), (2^n))$.

In [21] Theorem 5.1, the author constructed a surjective homomorphism from the affine (super) Yangian to the universal enveloping algebra of a rectangular W -(super)algebra.

Theorem 6.2 (Theorem 5.1 in [21] and Theorem 9.2 in [15]). 1. Suppose that $\varepsilon = \hbar\alpha$. There exists an algebra homomorphism

$$\Phi^n: Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n)) \rightarrow \mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(2n), (2^n)))$$

determined by

$$\begin{aligned} \Phi^n(H_{i,0}) &= \begin{cases} W_{n,n}^{(1)} - W_{1,1}^{(1)} + 2\alpha & \text{if } i = 0, \\ W_{i,i}^{(1)} - W_{i+1,i+1}^{(1)} & \text{if } i \neq 0, \end{cases} \\ \Phi^n(X_{i,0}^+) &= \begin{cases} W_{n,1}^{(1)}t & \text{if } i = 0, \\ W_{i,i+1}^{(1)} & \text{if } i \neq 0, \end{cases} \quad \Phi^n(X_{i,0}^-) = \begin{cases} W_{1,n}^{(1)}t^{-1} & \text{if } i = 0, \\ W_{i+1,i}^{(1)} & \text{if } i \neq 0, \end{cases} \end{aligned}$$

and

$$\begin{aligned} \Phi^n(H_{i,1}) &= -\hbar W_{i,i}^{(2)}t + \hbar W_{i+1,i+1}^{(2)}t + \frac{i}{2}\hbar\Phi^n(H_{i,0}) - \hbar W_{i,i}^{(1)}W_{i+1,i+1}^{(1)} \\ &+ \hbar \sum_{s \geq 0} \sum_{u=1}^i W_{i,u}^{(1)}t^{-s}W_{u,i}^{(1)}t^s + \hbar \sum_{s \geq 0} \sum_{u=i+1}^n W_{i,u}^{(1)}t^{-s-1}W_{u,i}^{(1)}t^{s+1} \\ &- \hbar \sum_{s \geq 0} \sum_{u=1}^i W_{i+1,u}^{(1)}t^{-s}W_{u,i+1}^{(1)}t^s - \hbar \sum_{s \geq 0} \sum_{u=i+1}^n W_{i+1,u}^{(1)}t^{-s-1}W_{u,i+1}^{(1)}t^{s+1} \end{aligned}$$

for $i \neq 0$.

2. In the case that $\varepsilon \neq 0$, the image of Φ^n is dense in $\mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(2n), (2^n)))$.

By the definition of Φ^n , we obtain the following relation.

Theorem 6.3 (Theorem 5.6 in [20]). Suppose that $\varepsilon = -(k + (n + m))\hbar$. We obtain the following relation:

$$\Phi^{m+n} \circ \Psi_1 = \iota \circ \Phi^n.$$

We can consider the coset vertex algebra $C(\mathcal{W}^k(\mathfrak{gl}(2m+2n), (2^{m+n}), \mathcal{W}^{k+m}(\widehat{\mathfrak{sl}}(2n), (2^n))))$ by Theorem 6.1. Then, Theorem 4.2 and Theorem 6.3 induce the following corollary.

Corollary 6.4. *In the case that $\varepsilon \neq 0$, we obtain*

$$\Phi_{\hbar,\varepsilon}^{m+n} \circ \Psi_2: Y_{\hbar,\varepsilon+n\hbar}(\widehat{\mathfrak{sl}}(m)) \rightarrow C(\mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(2m+2n), (2^{m+n})), \mathcal{U}(\mathcal{W}^{k+m}(\mathfrak{gl}(2n), (2^n)))),$$

where

$$\begin{aligned} & C(\mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(2m+2n), (2^{m+n})), \mathcal{U}(\mathcal{W}^{k+m}(\mathfrak{gl}(2n), (2^n)))) \\ & = \{x \in \mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(2m+2n), (2^{m+n}))) \mid [x, \mathcal{U}(\mathcal{W}^{k+m}(\mathfrak{gl}(2n), (2^n)))] = 0\}. \end{aligned}$$

We can also

Theorem 6.5. *In the case that $\varepsilon \neq 0$, the homomorphism induces the homomorphism*

$$\Phi_{\hbar,\varepsilon}^{m+n} \circ \Psi_2: Y_{\hbar,\varepsilon+n\hbar}(\widehat{\mathfrak{sl}}(m)) \rightarrow \mathcal{U}(C(\mathcal{W}^k(\mathfrak{gl}(2m+2n), (2^{m+n})), \mathcal{W}^{k+m}(\mathfrak{sl}(2n), (2^n)))).$$

Proof. By the definition, we find that $W_{i,j}^{(1)}[-1]$ is contained in the universal enveloping algebra of the coset for $i, j \geq n+1$. The OPEs of $W_{i,j}^{(1)}$ and $\mathcal{W}^{k+m}(\mathfrak{sl}(2n), (2^n))$ are non-zero for $i, j \geq n+1$ due to the inner product $(E_{i,i}, E_{p,p}) = 1$ for $i \geq n+1, p \leq n$. Since $(E_{i,i} - E_{i+1,i+1}, E_{p,p} - E_{p+1,p+1}) = 0$ for $i \geq n+1, p \leq n-1$, we find that

$$[W_{i,j}^{(1)} t^s, \mathcal{U}(\mathcal{W}^{k+m}(\mathfrak{sl}(2n), (2^n)))] = 0$$

for $i, j \geq n+1$. Thus, the image of $\Phi_{\hbar,\varepsilon}^{m+n} \circ \Psi_2$ is contained in the coset if $W_{i,i}^{(2)} - W_{i+1,i+1}^{(2)}$ is contained in the coset for $i \geq n+1$. By the OPEs of the rectangular W -algebra $(\mathcal{W}^k(\mathfrak{gl}(2m+2n), (2^{m+n})))$ given in [21], we have

$$\begin{aligned} & [((W_{i,i}^{(2)} - W_{i+1,i+1}^{(2)})t, W_{i,i+1}^{(1)} t^{-2}), W_{i+1,i}^{(1)}] \\ & = (W_{i,i}^{(2)} - W_{i+1,i+1}^{(2)})t^{-1} + \text{the elements generated by } W_{p,q}^{(1)} t^s \text{ for } p, q \geq n+1 \text{ and } s \in \mathbb{Z}. \end{aligned} \quad (6.6)$$

By Corollary 6.4, the left hand side of (6.6) is contained in the centralizer of $\mathcal{U}(\mathcal{W}^{k+m}(\mathfrak{gl}(2n), (2^n)))$ with $\mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(2m+2n), (2^{m+n})))$. Thus, we obtain

$$[xt^r, (W_{i,i}^{(2)} - W_{i+1,i+1}^{(2)})t^{-1}] = 0 \text{ if } x \in \mathcal{W}^{k+m}(\mathfrak{sl}(2n), (2^n)).$$

Then, we find that $W_{i,i}^{(2)} - W_{i+1,i+1}^{(2)}$ is contained in the coset. □

A Some formulas for the proof of Theorem 3.1 and Theorem 4.2

For the proof of Theorem 4.2, we note one lemma.

Lemma A.1. *For $a, b \geq 0$, we obtain*

$$\begin{aligned} & [E_{j+n,u} t^{-s-a} E_{u,j+n} t^{s+a}, E_{i,k} t^{-v-1} E_{k,i} t^{v+1}] \\ & = E_{j+n,u} t^{-s-a} (\delta_{i,j+n} E_{u,k} t^{s-v+a-1} - \delta_{k,u} E_{i,j+n} t^{s-v+a-1}) E_{k,i} t^{v+1} \\ & \quad + (\delta_{u,i} E_{j+n,k} t^{-s-v-a-1} - \delta_{j+n,k} E_{i,u} t^{-s-v-a-1}) E_{u,j+n} t^{s+a} E_{k,i} t^{v+1} \\ & \quad + E_{i,k} t^{-v-1} E_{j+n,u} t^{-s-a} (\delta_{j+n,k} E_{u,i} t^{s+v+a+1} - \delta_{u,i} E_{k,j+n} t^{s+v+a+1}) \\ & \quad + E_{i,k} t^{-v-1} (\delta_{u,k} E_{j+n,i} t^{v-s-a+1} - \delta_{i,j+n} E_{k,u} t^{v-s-a+1}) E_{u,j+n} t^{s+a}, \quad (A.2) \\ & [E_{u,j+n} t^{-s-a} E_{j+n,u} t^{s+a}, E_{i,k} t^{-v-1} E_{k,i} t^{v+1}] \\ & = E_{u,j+n} t^{-s-a} (\delta_{u,i} E_{j+n,k} t^{s+a-v-1} - \delta_{k,j+n} E_{i,u} t^{s+a-v-1}) E_{k,i} t^{v+1} \\ & \quad + (\delta_{i,j+n} E_{u,k} t^{-s-v-a-1} - \delta_{k,u} E_{i,j+n} t^{-s-v-a-1}) E_{j+n,u} t^{s+a} E_{k,i} t^{v+1} \end{aligned}$$

$$\begin{aligned}
& + E_{i,k}t^{-v-1}E_{u,j+n}t^{-s-a}(\delta_{u,k}E_{j+n,i}t^{s+v+a+1} - \delta_{i,j+n}E_{k,u}t^{s+v+a+1}) \\
& + E_{i,k}t^{-v-1}(\delta_{j+n,k}E_{u,i}t^{v-s-a+1} - \delta_{i,u}E_{k,j+n}t^{v-s-a+1})E_{j+n,u}t^{s+a}, \tag{A.3}
\end{aligned}$$

$$\begin{aligned}
& [E_{u,i}t^{-s-a}E_{i,u}t^{s+a}, E_{l,j+n}t^{-v-b}E_{j+n,l}t^{v+b}] \\
= & E_{u,i}t^{-s-a}(\delta_{u,l}E_{i,j+n}t^{s-v+a-b} - \delta_{i,j+n}E_{l,u}t^{s-v+a-b})E_{j+n,l}t^{v+b} \\
& + (\delta_{l,i}E_{u,j+n}t^{-s-v-a-b} - \delta_{u,j+n}E_{l,i}t^{-s-v-a-b})E_{i,u}t^{s+a}E_{j+n,l}t^{v+b} \\
& + E_{l,j+n}t^{-v-b}E_{u,i}t^{-s-a}(\delta_{u,j+n}E_{i,l}t^{s+v+a+b} - \delta_{i,l}E_{j+n,u}t^{s+v+a+b}) \\
& + E_{l,j+n}t^{-v-b}(\delta_{i,j+n}E_{u,l}t^{v-s+b-a} - \delta_{u,l}E_{j+n,i}t^{v-s+b-a})E_{i,u}t^{s+a}, \tag{A.4}
\end{aligned}$$

$$\begin{aligned}
& [E_{i,u}t^{-s-a}E_{u,i}t^{s+a}, E_{l,j+n}t^{-v-b}E_{j+n,l}t^{v+b}] \\
= & E_{i,u}t^{-s-a}(\delta_{i,l}E_{u,j+n}t^{s-v+a-b} - \delta_{j+n,u}E_{l,i}t^{s-v+a-b})E_{j+n,l}t^{v+b} \\
& + (\delta_{u,l}E_{i,j+n}t^{-s-v-a-b} - \delta_{i,j+n}E_{l,u}t^{-s-v-a-b})E_{u,i}t^{s+a}E_{j+n,l}t^{v+b} \\
& + E_{l,j+n}t^{-v-b}E_{i,u}t^{-s-a}(\delta_{i,j+n}E_{u,l}t^{s+v+a+b} - \delta_{u,l}E_{j+n,i}t^{s+v+a+b}) \\
& + E_{l,j+n}t^{-v-b}(\delta_{u,j+n}E_{i,l}t^{v-s+b-a} - \delta_{i,l}E_{j+n,u}t^{v-s+b-a})E_{u,i}t^{s+a}. \tag{A.5}
\end{aligned}$$

The proof is due to a direct computation.

References

- [1] T. Arakawa and A. Molev. Explicit generators in rectangular affine W -algebras of type A . *Lett. Math. Phys.*, 107(1):47–59, 2017, <https://doi.org/10.1007/s11005-016-0890-2>.
- [2] N. Arbesfeld and O. Schiffmann. A presentation of the deformed $W_{1+\infty}$ algebra. In *Symmetries, integrable systems and representations*, volume 40 of *Springer Proc. Math. Stat.*, pages 1–13. Springer, Heidelberg, 2013.
- [3] T. Creutzig and A. R. Linshaw. Trialities of \mathcal{W} -algebras. *Camb. J. Math.*, 10(1):69–194, 2022.
- [4] V. G. Drinfeld. Hopf algebras and the quantum Yang-Baxter equation. *Dokl. Akad. Nauk SSSR*, 283(5):1060–1064, 1985, <https://doi.org/10.1142/9789812798336.0013>.
- [5] V. G. Drinfeld. A new realization of Yangians and of quantum affine algebras. *Dokl. Akad. Nauk SSSR*, 296(1):13–17, 1987.
- [6] I. B. Frenkel and Y. Zhu. Vertex operator algebras associated to representations of affine and Virasoro algebras. *Duke Math. J.*, 66(1):123–168, 1992, <https://doi.org/10.1215/S0012-7094-92-06604-X>.
- [7] M. R. Gaberdiel and R. Gopakumar. Triality in minimal model holography. *J. High Energy Phys.*, (7):127, front matter+26, 2012.
- [8] D. Gaiotto and M. Rapčák. Vertex algebras at the corner. *J. High Energy Phys.*, (1):160, front matter+85, 2019.
- [9] N. Genra. Screening operators for \mathcal{W} -algebras. *Selecta Math. (N.S.)*, 23(3):2157–2202, 2017, <https://doi.org/10.1007/s00029-017-0315-9>.
- [10] N. Guay. Cherednik algebras and Yangians. *Int. Math. Res. Not.*, (57):3551–3593, 2005, <https://doi.org/10.1155/IMRN.2005.3551>.
- [11] N. Guay. Affine Yangians and deformed double current algebras in type A . *Adv. Math.*, 211(2):436–484, <https://doi.org/10.1016/j.aim.2006.08.007>, 2007.
- [12] N. Guay, H. Nakajima, and C. Wendlandt. Coproduct for Yangians of affine Kac-Moody algebras. *Adv. Math.*, 338:865–911, 2018, <https://doi.org/10.1016/j.aim.2018.09.013>.

- [13] R. Kodera. Braid group action on affine Yangian. *SIGMA Symmetry Integrability Geom. Methods Appl.*, 15:Paper No. 020, 28, 2019, <https://doi.org/10.3842/SIGMA.2019.020>.
- [14] R. Kodera. On Guay’s evaluation map for affine Yangians. *Algebr. Represent. Theory*, 24(1):253–267, 2021, <https://doi.org/10.1007/s10468-019-09945-w>.
- [15] R. Kodera and M. Ueda. Coproduct for affine Yangians and parabolic induction for rectangular W -algebras. *Lett. Math. Phys.*, 112(1):Paper No. 3, 37, 2022.
- [16] A. Matsuo, K. Nagatomo, and A. Tsuchiya. Quasi-finite algebras graded by Hamiltonian and vertex operator algebras. *London Math. Soc. Lecture Note Ser.*, 372:282–329, 2010, <https://doi.org/10.1017/CBO9780511730054.015>.
- [17] D. Maulik and A. Okounkov. Quantum groups and quantum cohomology. *Astérisque*, (408):ix+209, 2019.
- [18] G. I. Olshanskii. Representations of infinite-dimensional classical groups, limits of enveloping algebras, and Yangians. In *Topics in representation theory*, volume 2 of *Adv. Soviet Math.*, pages 1–66. Amer. Math. Soc., Providence, RI, 1991.
- [19] A. Tsybaliuk. The affine Yangian of \mathfrak{gl}_1 revisited. *Adv. Math.*, 304:583–645, 2017.
- [20] M. Ueda. A homomorphism from the affine Yangian $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$ to the affine Yangian $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n+1))$. arXiv:2312.09933.
- [21] M. Ueda. Affine super Yangians and rectangular W -superalgebras. *J. Math. Phys.*, 63(5):Paper No. 051701, 34, 2022.