GLOBAL SOLUTIONS FOR SEMILINEAR PARABOLIC EVOLUTION PROBLEMS WITH HÖLDER CONTINUOUS NONLINEARITIES

BOGDAN–VASILE MATIOC AND CHRISTOPH WALKER

ABSTRACT. It is shown that semilinear parabolic evolution equations $u' = A + f(t, u)$ featuring Hölder continuous nonlinearities $f = f(t, u)$ with at most linear growth possess global strong solutions for a general class of initial data. The abstract results are applied to a recent model describing front propagation in bushfires and in the context of a reaction-diffusion system.

1. Introduction

For semilinear parabolic problems

$$
u' = Au + f(t, u), \quad t \in (0, T], \qquad u(0) = u^0,
$$
\n(1.1)

involving a generator A of an analytic semigroup on a Banach space E_0 and a locally Lipschitz continuous nonlinearity f , there is a well-established theory for well-posedness based on Banach's fixed point theorem, e.g. see [\[1,](#page-14-0) Section 12] and the references therein. The situation is different if $f = f(t, u)$ is not locally Lipschitz with respect to u and less seems to be known. If compactness properties are available, (versions of) Schauder's fixed point theorem may be used to derive existence – but not uniqueness – results, see e.g. the textbooks $[9,10]$ $[9,10]$ or also $[3]$. We also refer to $[13]$ for the treatment of non-Lipschitz semilinear parabolic scalar equations by comparison principle arguments.

The aim of this paper is to establish the existence of strong solutions to [\(1.1\)](#page-0-0) in the context of nonlinearities $f = f(t, u)$ with at most linear growth in u, which are not necessarily locally Lipschitz continuous with respect to the variable u , under optimal (i.e. low) regularity assumptions on the initial value.

To be more precise, in the following let E_0 and E_1 be two Banach spaces over $\mathbb{K} \in \{ \mathbb{R}, \mathbb{C} \}$ with compact, continuous, and dense embedding

$$
E_1 \stackrel{d}{\iff} E_0\,.
$$

For each $\theta \in (0,1)$, let $(\cdot, \cdot)_{\theta}$ be an arbitrary admissible interpolation functor of exponent θ (see [\[2,](#page-14-5) I. Section 2.11]) and denote by $E_{\theta} := (E_0, E_1)_{\theta}$ the corresponding interpolation space with norm $\|\cdot\|_{\theta}$. Then, the embeddings

$$
E_1 \stackrel{d}{\iff} E_\theta \stackrel{d}{\iff} E_\vartheta \stackrel{d}{\iff} E_0, \qquad 0 < \vartheta < \theta < 1,
$$

are all dense and compact. We fix an infinitesimal generator $A: E_1 \to E_0$ of a strongly continuous analytic semigroup $\{e^{tA} : t \geq 0\} \subset \mathcal{L}(E_0)$ and assume that $f : [0,T] \times E_{\xi} \to E_{\gamma}$ for some $\gamma, \xi \in [0,1)$ and $T \in (0, \infty)$.

We first point out the following result guaranteeing the global existence of (at least) a global strong solution to [\(1.1\)](#page-0-0) for initial data $u_0 \in E_\alpha$ for which $f(0, u_0)$ is defined, that is, for $\alpha \in [\xi, 1)$,

²⁰²⁰ Mathematics Subject Classification. 35A01; 35K58; 35Q92.

Key words and phrases. Global strong solutions; Semilinear parabolic problems; Hölder continuous semilinearity.

under the general assumption that $f(t, u)$ grows at most linearly in u and is continuous (along with an additional Hölder continuity property in the limiting case $\gamma = 0$.

Proposition 1.1. Let $\gamma, \xi \in [0,1)$ and assume that $f \in C([0,T] \times E_{\xi}, E_{\gamma})$ is such that there is a $constant C > 0$ with

$$
|| f(t, u) ||_{\gamma} \le C \left(1 + ||u||_{\xi} \right), \qquad t \in [0, T], \quad u \in E_{\xi}.
$$
 (1.2)

If $\gamma = 0$ assume additionally that there exists $\vartheta_0 \in (0,1)$ such that for each $R > 0$ there is a constant $K(R) > 0$ with

$$
|| f(t, u) - f(s, v) ||_0 \le K(R) \left(|t - s|^{\vartheta_0} + ||u - v||_{\xi}^{\vartheta_0} \right), \qquad t, s \in [0, T], \quad u, v \in \bar{\mathbb{B}}_{E_{\xi}}(0, R). \tag{1.3}
$$

Then, if $u^0 \in E_\alpha$ for some $\alpha \in [\xi, 1)$, the Cauchy problem [\(1.1\)](#page-0-0) admits a solution

 $u \in C^1((0,T], E_0) \cap C((0,T], E_1) \cap C([0,T], E_\alpha)$.

The main purpose of this paper is to extend the existence theory for the evolution problem [\(1.1\)](#page-0-0) by providing a global strong solution to [\(1.1\)](#page-0-0) for a general initial datum u^0 in the ambient space E_0 . To this end we require, additionally to being linearly bounded in u , that f is locally Hölder continuous with respect to u . More precisely, we fix

$$
0 \le \gamma < 1
$$
, $0 < \xi < \min\left\{1, \frac{1}{q}\right\}$, $0 < \vartheta_j \le q_j$, $1 \le j \le m \in \mathbb{N}^*$, (1.4a)

where

 $q := \max\{q_i : 1 \leq j \leq m\} > 0$,

and assume that $f \in C((0,T] \times E_{\xi}, E_{\gamma})$ is such that there exists a positive constant $C > 0$ with

$$
||f(t, u)||_{\gamma} \le C(1 + ||u||_{\xi})
$$
\n(1.4b)

and

$$
||f(t, u) - f(t, v)||_{\gamma} \le C \sum_{j=1}^{m} \left(1 + ||u||_{\xi}^{q_j - \vartheta_j} + ||v||_{\xi}^{q_j - \vartheta_j}\right) ||u - v||_{\xi}^{\vartheta_j}
$$
(1.4c)

for all $t \in (0, T]$ and $u, v \in E_{\xi}$. In the particular case $\gamma = 0$ we additionally require (similarly as in Proposition [1.1,](#page-1-0) but in a slightly weaker form) that there exists $\vartheta_0 \in (0,1)$ such that for each $R > 0$ there is a constant $K(R) > 0$ with

$$
|| f(t, u) - f(s, u) ||_0 \le K(R) |t - s|^{\vartheta_0}, \qquad t, s \in (0, T], \quad u \in E_{\xi} \cap \bar{\mathbb{B}}_{E_0}(0, R). \tag{1.5}
$$

Our main result then reads:

Theorem 1.2. Suppose [\(1.4\)](#page-1-1) and if $\gamma = 0$ also assume [\(1.5\)](#page-1-2). Then, given $u^0 \in E_0$, the Cauchy problem [\(1.1\)](#page-0-0) possesses a global strong solution

$$
u \in C^{1}((0, T], E_{0}) \cap C((0, T], E_{1}) \cap C([0, T], E_{0}).
$$

Furthermore, if $u^0 \in E_\alpha$ for some $\alpha \in [0, \min\{1, 1 + \gamma - \xi\})$, then $C([0, T], E_\alpha)$.

Theorem [1.2](#page-1-3) provides strong solutions assuming minimal regularity on the initial value. In particular, $f(0, u^0)$ needs not be defined under the assumptions of Theorem [1.2](#page-1-3) (in contrast to Proposition [1.1\)](#page-1-0). For this reason, if $u^0 \in E_\alpha \setminus E_\xi$, the continuity in E_α of the solution at $t = 0$ is guaranteed only for possibly restricted α .

Remarks 1.3. (i) If $1 \le \vartheta_j \le q_j$ for all $1 \le j \le m$, then $f = f(t, u)$ is locally Lipschitz continuous in u and the classical theory on well-posedness for semilinear parabolic equations applies, see e.g. $[1]$ for a general treatment (or [\[11\]](#page-14-6) for a similar approach to the one chosen herein requiring minimal regularity assumptions on the initial value).

(ii) In general, however, the nonlinearity $f = f(t, u)$ in [\(1.1\)](#page-0-0) is not locally Lipschitz continuous in u as the exponents ϑ_i in [\(1.4c\)](#page-1-4) are allowed to be less than 1. Thus, our approach to Proposition [1.1](#page-1-0) and Theorem [1.2,](#page-1-3) relying on the Leray-Schauder fixed point theorem, does not provide uniqueness of solutions.

(iii) The linear bound [\(1.4b\)](#page-1-5) is not in contradiction with assumption [\(1.4c\)](#page-1-4) (even if q is possibly larger than 1). This is well exemplified in Section [3](#page-8-0) by application [\(3.4\)](#page-9-0) with $\nu = 1$.

Combining Proposition [1.1](#page-1-0) and Theorem [1.2,](#page-1-3) we may establish a global existence result for $T = \infty$.

Corollary 1.4. Let $\gamma, \xi \in [0,1)$, $f \in C([0,\infty) \times E_{\xi}, E_{\gamma})$, and assume that for each $T > 0$ there exists a constant $C(T) > 0$ with

$$
|| f(t, u) ||_{\gamma} \le c(T) (1 + ||u||_{\xi}), \qquad t \in [0, T], \quad u \in E_{\xi}.
$$
 (1.6)

(a) Let $\gamma > 0$. Then [\(1.1\)](#page-0-0) has a global strong solution

 $u \in C^1((0,\infty),E_0) \cap C((0,\infty),E_1) \cap C([0,\infty),E_\alpha)$

provided either $u^0 \in E_\alpha$ with $\alpha \in [\xi,1)$, or f satisfies [\(1.4\)](#page-1-1) for some $T = T_0 > 0$ and $u^0 \in E_\alpha$ with $\alpha \in [0, \min\{1, 1 + \gamma - \xi\})$.

(b) Let $\gamma = 0$ and suppose there is $\vartheta_0 \in (0,1)$ and for $T, R > 0$ there is $K = K(T, R) > 0$ such that

$$
|| f(t, u) - f(s, v) ||_0 \le K \left(|t - s|^{\vartheta_0} + ||u - v||_{\xi}^{\vartheta_0} \right), \qquad t, s \in [0, T], \quad u, v \in \bar{\mathbb{B}}_{E_{\xi}}(0, R). \tag{1.7}
$$

Then [\(1.1\)](#page-0-0) has a global strong solution

$$
u \in C^1((0,\infty),E_0) \cap C((0,\infty),E_1) \cap C([0,\infty),E_\alpha)
$$

provided either $u^0 \in E_\alpha$ with $\alpha \in [\xi, 1)$, or f satisfies [\(1.4\)](#page-1-1) and [\(1.5\)](#page-1-2) for some $T = T_0 > 0$ and $u^0 \in E_\alpha$ with $\alpha \in [0, 1-\xi)$.

The proofs of Proposition [1.1,](#page-1-0) Theorem [1.2,](#page-1-3) and Corollary [1.4](#page-2-0) are presented in Section [2.](#page-2-1) In Section [3](#page-8-0) we apply the abstract results to models describing front propagation in bushfires and to reaction-diffusion models.

2. Proof of the main results

We provide first the details of the proof Theorem [1.2](#page-1-3) (for which the arguments are a bit more delicate) and subsequently only sketch the proofs of Proposition [1.1](#page-1-0) and Corollary [1.4.](#page-2-0)

Since A is the generator of the strongly continuous analytic semigroup $\{e^{tA} : t \geq 0\}$, the Cauchy problem [\(1.1\)](#page-0-0) can be formulated (under suitable assumptions on u and f, see e.g. [\[2,](#page-14-5) II. Remarks 2.1.2 (a)]) as the fixed point equation

$$
u(t) = e^{tA}u^{0} + \int_{0}^{t} e^{(t-\tau)A} f(\tau, u(\tau)) d\tau, \quad t \in [0, T],
$$
\n(2.1)

which we shall solve using the Leray-Schauder fixed point theorem.

2.1. Proof of Theorem [1.2.](#page-1-3) Assuming the premises of Theorem [1.2,](#page-1-3) we set

$$
\vartheta := \min\{\vartheta_j \, : \, 1 \le j \le m\} > 0
$$

and choose $\mu \in \mathbb{R}$ such that

$$
\xi < \mu < \min\left\{1, \frac{1}{q}\right\}.\tag{2.2}
$$

We denote by

$$
X_T := C_\mu((0,T],E_\xi)
$$

the Banach space consisting of all functions $u \in C((0,T], E_{\xi})$ such that $t^{\mu}u(t) \to 0$ in E_{ξ} as $t \to 0$, which is equipped with the norm

$$
||u||_{X_T} := \sup \{ t^{\mu} ||u(t)||_{\xi} : t \in (0,T] \}.
$$

Our goal is to prove that the function $F: X_T \to X_T$, defined by

$$
F(u)(t) := e^{tA}u^0 + \int_0^t e^{(t-\tau)A} f(\tau, u(\tau)) d\tau, \qquad t \in [0, T], \quad u \in X_T,
$$
\n(2.3)

satisfies the hypotheses of the Leray-Schauder fixed point theorem [\[8,](#page-14-7) Theorem 11.3] and thus admits a fixed point in X_T , which is then shown to be a global strong solution to [\(1.1\)](#page-0-0).

Well-definedness. In order to establish that $F(u) \in X_T$ for each $u \in X_T$, we first recall from [\[2,](#page-14-5) II. Lemma 5.1.3] the estimates

$$
||e^{tA}||_{\mathcal{L}(E_{\theta})} + t^{\alpha - \beta_0}||e^{tA}||_{\mathcal{L}(E_{\beta}, E_{\alpha})} \le M, \qquad 0 \le t \le T,
$$
\n(2.4)

for $\theta \in [0,1]$ and $0 \le \beta_0 \le \beta \le \alpha \le 1$ with $\beta_0 < \beta$ if $0 < \beta < \alpha < 1$, where $M = M(T)$ depends also on these parameters.

Given $u, v \in X_T$ with $||u||_{X_T} \leq L$ and $||v||_{X_T} \leq L$ (for some arbitrary $L > 0$), assumption [\(1.4c\)](#page-1-4) implies

$$
||f(t, u(t)) - f(t, v(t))||_{\gamma} \le c(L) \sum_{j=1}^{m} \left(1 + t^{-\mu(q_j - \vartheta_j)}\right) ||u(t) - v(t)||_{\xi}^{\vartheta_j}
$$

$$
\le c(T, L) t^{-\mu q} ||u - v||_{X_T}^{\vartheta}, \qquad t \in (0, T],
$$
 (2.5)

while [\(1.4b\)](#page-1-5) yields

$$
|| f(t, u(t)) ||_{\gamma} \le c(T, L) t^{-\mu}, \quad t \in (0, T].
$$
 (2.6)

Fix $u \in X_T$ with $||u||_{X_T} \leq L$ (for some arbitrary L). Let $\eta \in (0,1)$ be given and define $\gamma_0 = \gamma_0(\eta)$ as

$$
\gamma_0 := 0 \quad \text{if} \quad \gamma = 0 \qquad \text{and} \qquad \gamma_0 \in (0, \min\{\gamma, \eta\}) \quad \text{if} \quad \gamma > 0. \tag{2.7}
$$

In view of (2.4) and (2.6) we obtain

$$
||F(u)(t)||_{\eta} \leq ||e^{tA}||_{\mathcal{L}(E_0, E_{\eta})} ||u^0||_0 + \int_0^t ||e^{(t-\tau)A}||_{\mathcal{L}(E_{\gamma}, E_{\eta})} ||f(\tau, u(\tau))||_{\gamma} d\tau
$$

\n
$$
\leq ct^{-\eta} ||u^0||_0 + c(T, L) \int_0^t (t-\tau)^{\gamma_0-\eta} \tau^{-\mu} d\tau
$$

\n
$$
\leq c(T, L) \left(t^{-\eta} + t^{1+\gamma_0-\eta-\mu} \mathcal{B}(1+\gamma_0-\eta, 1-\mu) \right)
$$
\n(2.8)

for $t \in (0, T]$, where B denotes the Beta function. Therefore, for $\eta = \xi$, we obtain

$$
t^{\mu} \| F(u)(t) \|_{\xi} \le c(T, L) \left(t^{\mu - \xi} + t^{1 + \gamma_0 - \xi} \right), \quad t \in (0, T], \tag{2.9}
$$

and the right hand side converges to zero as $t \to 0$ since $\xi < \mu < 1$.

Given $\varepsilon \in (0,T)$, set $u_{\varepsilon}(t) := u(t+\varepsilon)$ for $t \in [0,T-\varepsilon]$. Then, $u_{\varepsilon} \in C([0,T-\varepsilon],E_{\xi})$ and the continuity of f implies $f(\cdot + \varepsilon, u_{\varepsilon}) \in C([0, T - \varepsilon], E_{\gamma})$. Moreover, the definition of $F(u)$ entails

$$
F(u)(t+\varepsilon) = e^{tA} F(u)(\varepsilon) + \int_0^t e^{(t-\tau)A} f(\tau + \varepsilon, u_\varepsilon(\tau)) d\tau, \quad t \in [0, T - \varepsilon]. \tag{2.10}
$$

Hence, [\[2,](#page-14-5) II.Theorem 5.3.1] yields

$$
F(u)(\cdot + \varepsilon) \in C\big((0, T - \varepsilon], E_{\theta}\big), \qquad \theta \in (0, 1), \quad \varepsilon \in (0, T),
$$

that is,

$$
F(u) \in C((0, T], E_{\theta}), \quad \theta \in (0, 1).
$$
 (2.11)

Together with [\(2.9\)](#page-4-0) we conclude $F(u) \in X_T$.

Continuity. Given $u, v \in X_T$ with $||u||_{X_T} \leq L$ and $||v||_{X_T} \leq L$, we obtain in view of [\(2.4\)](#page-3-0) and [\(2.5\)](#page-3-2), similarly as above, that

$$
||F(u)(t) - F(v)(t)||_{\xi} \le \int_0^t ||e^{(t-\tau)A}||_{\mathcal{L}(E_{\gamma}, E_{\xi})} ||f(\tau, u(\tau)) - f(\tau, v(\tau))||_{\gamma} d\tau
$$

\n
$$
\le c||u - v||_{X_T}^{\vartheta} B(1 + \gamma_0 - \xi, 1 - \mu q) t^{1+\gamma_0 - \xi - \mu q}, \qquad t \in (0, T], \qquad (2.12)
$$

with $\gamma_0 = \gamma_0(\xi)$ as in [\(2.7\)](#page-3-3). Since $\mu q < 1$ and $\xi < \mu < 1$, we get

$$
||F(u) - F(v)||_{X_T} \le c_1(T, L) ||u - v||_{X_T}^{\vartheta}.
$$

This proves in particular the (Hölder) continuity of F.

Compactness. We show that $F: X_T \to X_T$ is compact. Let therefore $(u_k)_k$ be a bounded sequence in X_T and choose $L > 0$ such that $||u_k||_{X_T} \leq L$ for all $k \in \mathbb{N}$. We prove that $(F(u_k))_k$ is relatively compact in X_T by applying the Arzelà-Ascoli theorem to the sequence $(v_k)_k \subset C([0,T], E_\xi)$ defined as

$$
v_k(t) := t^{\mu} F(u_k)(t), \qquad t \in [0, T], \quad k \in \mathbb{N}.
$$

Note that $v_k(0) = 0$ for $k \in \mathbb{N}$. Since $(v_k(t))_k$ is bounded in E_η for $\eta \in (\xi, \mu]$ by (2.8) , the compactness of the embedding $E_\eta \hookrightarrow E_\xi$ ensures that $(v_k(t))_k$ is relatively compact in E_ξ for each $t \in [0, T]$.

We next prove that the sequence $(v_k)_k$ is equi-continuous. Given $0 < s < t \leq T$ and $k \in \mathbb{N}$, we have

$$
||v_{k}(t) - v_{k}(s)||_{\xi} \leq (t^{\mu} - s^{\mu})||F(u_{k})(t)||_{\xi} + s^{\mu} \int_{s}^{t} ||e^{(t-\tau)A}||_{\mathcal{L}(E_{\gamma}, E_{\xi})} ||f(\tau, u_{k}(\tau))||_{\gamma} d\tau
$$

+ $s^{\mu} \int_{0}^{s} ||e^{(t-\tau)A} - e^{(s-\tau)A}||_{\mathcal{L}(E_{\gamma}, E_{\xi})} ||f(\tau, u_{k}(\tau))||_{\gamma} d\tau$
+ $s^{\mu} ||(e^{tA} - e^{sA})u^{0}||_{\xi} =: I_{1} + I_{2} + I_{3} + I_{4}.$ (2.13a)

In the following, we simply write c for positive constants $c = c(T, L)$ depending on T and L. From [\(2.9\)](#page-4-0), we obtain

$$
||F(u_k)(t)||_{\xi} \le ct^{-\xi}, \qquad t \in (0,T], \quad k \in \mathbb{N},
$$

and therefore

$$
I_1 \le ct^{-\xi}(t^{\mu} - s^{\mu}) \le ct^{-\xi}(t - s)^{\mu} \le c(t - s)^{\mu - \xi}.
$$
 (2.13b)

Using (2.4) and (2.6) , we deduce

$$
I_2 \le cs^{\mu} \int_s^t (t - \tau)^{\gamma_0 - \xi} \tau^{-\mu} d\tau
$$

\n
$$
\le cs^{\mu} (t - s)^{1 + \gamma_0 - \xi} \int_0^1 (1 - r)^{\gamma_0 - \xi} (r(t - s) + s)^{-\mu} d\tau
$$

\n
$$
\le c(t - s)^{1 + \gamma_0 - \xi}.
$$
\n(2.13c)

Let now $\eta \in (\xi, \mu)$ be fixed and $\gamma_0 = \gamma_0(\eta)$ as in [\(2.7\)](#page-3-3). Since

$$
||e^{(t-s)A} - 1||_{\mathcal{L}(E_{\eta}, E_{\xi})} \le c(t-s)^{\eta-\xi}
$$
\n(2.13d)

due to [\[2,](#page-14-5) II. Theorem 5.3.1] (with $f = 0$ therein), we use [\(2.4\)](#page-3-0) and [\(2.6\)](#page-3-1) to derive

$$
I_3 \le s^{\mu} \int_0^s \|e^{(t-s)A} - 1\|_{\mathcal{L}(E_{\eta}, E_{\xi})} \|e^{(s-\tau)A}\|_{\mathcal{L}(E_{\gamma}, E_{\eta})} \|f(\tau, u_k(\tau))\|_{\gamma} d\tau
$$

\n
$$
\le cs^{\mu}(t-s)^{\eta-\xi} \int_0^s (s-\tau)^{\gamma_0-\eta} \tau^{-\mu} d\tau
$$

\n
$$
\le cs^{1+\gamma_0-\eta} B(1+\gamma_0-\eta, 1-\mu) (t-s)^{\eta-\xi}
$$

\n
$$
\le c(t-s)^{\eta-\xi}.
$$
\n(2.13e)

Finally, from $(2.13d)$ and (2.4) we get

$$
I_4 \le s^{\mu} \|e^{(t-s)A} - 1\|_{\mathcal{L}(E_{\eta}, E_{\xi})} \|e^{sA}\|_{\mathcal{L}(E_0, E_{\eta})} \|u^0\|_0
$$

$$
\le cT^{\mu-\eta}(t-s)^{\eta-\xi} \|u^0\|_0.
$$
 (2.13f)

We thus conclude from [\(2.13\)](#page-4-1) that the sequence $(v_k)_k$ is equi-continuous. The Arzelà-Ascoli theorem now ensures (up to a subsequence) that $v_k \to v$ in $C([0,T], E_{\xi})$ for some function v with $v(0) = 0$. Defining $u(t) := t^{-\mu}v(t)$, $t \in (0,T]$, we obtain $F(u_k) \to u$ in X_T . This establishes the compactness of F .

A priori bound. We prove that the set

$$
S := \left\{ u \in X_T \, : \, u = \lambda F(u) \text{ for some } \lambda \in [0,1] \right\}
$$

is bounded in X_T . To this end we infer from [\(1.4b\)](#page-1-5) and [\(2.4\)](#page-3-0) that for $u \in S$ and $t \in (0,T]$ we have

$$
||u(t)||_{\xi} \le ||F(u)(t)||_{\xi} \le ||e^{tA}||_{\mathcal{L}(E_0, E_{\xi})} ||u^0||_0 + \int_0^t ||e^{(t-\tau)A}||_{\mathcal{L}(E_{\gamma}, E_{\xi})} ||f(\tau, u(\tau))||_{\gamma} d\tau
$$

$$
\le ct^{-\xi} + c \int_0^t (t-\tau)^{\gamma_0-\xi} (1 + ||u(\tau)||_{\xi}) d\tau
$$

$$
\le ct^{-\mu} + c \int_0^t (t-\tau)^{\gamma_0-\xi} ||u(\tau)||_{\xi} d\tau,
$$

with $\gamma_0 = \gamma_0(\xi)$ defined as in [\(2.7\)](#page-3-3). The singular Gronwall inequality [\[2,](#page-14-5) II.Corollary 3.3.2] now implies the existence of a constant $K = K(T) > 0$ such that

$$
||u(t)||_{\xi} \leq K t^{-\mu}, \qquad t \in (0, T],
$$

which entails the boundedness of S in X_T .

Existence of a solution. Due to the above considerations, we may apply Leray-Schauder's fixed point theorem [\[8,](#page-14-7) Theorem 11.3] to deduce that there exists $u \in X_T$ such that $F(u) = u$. It remains to prove that u is a strong solution to the Cauchy problem (1.1) .

To this end, we note that if $u_0 \in E_\alpha$ and $\alpha \in [0, \min\{1, 1 + \gamma - \xi\})$, then $u \in C([0, T], E_\alpha)$. Indeed, the continuity of u on $(0,T]$ is established in (2.11) . Since $[t \mapsto e^{tA}u_0] \in C([0,T], E_\alpha)$, it follows from [\(2.8\)](#page-3-4) (with $u_0 = 0$, $\eta = \alpha$, and $\gamma_0 = \gamma_0(\alpha)$ and μ chosen such that $\alpha < 1 + \gamma_0 - \mu$) that $u = F(u) \in C([0, T], E_\alpha)$, in particular $u(0) = u^0$.

Given $\varepsilon \in (0,T)$, we set as before $u_{\varepsilon}(t) := u(t+\varepsilon)$ for $t \in [0,T-\varepsilon]$ and obtain from the continuity of f that $f(\cdot + \varepsilon, u_{\varepsilon}) \in C([0, T - \varepsilon], E_{\gamma}).$

If $\gamma > 0$, we may apply [\[2,](#page-14-5) II.Theorem 1.2.2, II.Remarks 2.1.2 (e)] to conclude from [\(2.10\)](#page-4-3) that

$$
u_{\varepsilon} = F(u)(\cdot + \varepsilon) \in C\big((0, T - \varepsilon], E_1\big) \cap C^1\big((0, T - \varepsilon], E_0\big), \quad \varepsilon \in (0, T),
$$

hence

$$
u \in C\big((0, T], E_1\big) \cap C^1\big((0, T], E_0\big) \tag{2.14}
$$

is a strong solution to [\(1.1\)](#page-0-0).

If $\gamma = 0$, we have $u_{\varepsilon}(0) \in E_{\theta}$ for some $\theta \in (\xi, 1)$ by (2.11) and $f(\cdot + \varepsilon, u_{\varepsilon}) \in C([0, T - \varepsilon], E_0)$ so that [\[2,](#page-14-5) II.Theorem 5.3.1] together with (2.10) imply $u_{\varepsilon} \in C^{\theta-\xi}([0,T-\varepsilon], E_{\xi})$. Along with $(1.4c)$ and [\(1.5\)](#page-1-2) we conclude $f(\cdot + \varepsilon, u_{\varepsilon}) \in C^{\rho}([0, T - \varepsilon], E_0)$ with $\rho := \min\{\vartheta_0, \vartheta(\theta - \xi)\} \in (0, 1)$. Invoking now $[2, II.$ Theorem 1.2.1 we deduce from (2.10)

$$
u_{\varepsilon} \in C\big((0, T - \varepsilon], E_1\big) \cap C^1\big((0, T - \varepsilon], E_0\big)
$$

for each $\varepsilon \in (0,T)$, hence u is a strong solution to [\(1.1\)](#page-0-0) enjoying the regularity properties [\(2.14\)](#page-6-0). This proves Theorem [1.2.](#page-1-3) \Box

2.2. **Proof of Proposition [1.1.](#page-1-0)** We now provide the proof of Proposition [1.1,](#page-1-0) which relies on the same strategy as that of Theorem [1.2,](#page-1-3) the main difference being that the Leray-Schauder fixed point theorem is now applied in the Banach space $C([0,T], E_\alpha)$ of continuous functions in E_α .

Assume the premises of Proposition [1.1.](#page-1-0) Set $X_T := C([0,T], E_\alpha)$ and let $u \in X_T$. Since $\alpha \geq \xi$, the continuity of f ensures $f(\cdot, u) \in C([0,T], E_\gamma)$, so that $F(u)$ in [\(2.3\)](#page-3-5) satisfies

$$
F(u) \in C^{\alpha-\beta}([0,T], E_{\beta}) \cap C((0,T], E_{\theta}), \qquad \beta \in [0,\alpha], \quad \theta \in (0,1),
$$

due to [\[2,](#page-14-5) II.Theorem 5.3.1] and [\(2.11\)](#page-4-2). In particular, $F(u)$ is an element of X_T . Moreover, a straightforward contradiction argument shows that given $\varepsilon > 0$ arbitrary, there is $\delta > 0$ such that

$$
|| f(t, u(t)) - f(t, v(t)) ||_{\gamma} \leq \varepsilon, \qquad t \in [0, T], \quad v \in \bar{\mathbb{B}}_{X_T}(u, \delta).
$$

From this we deduce, analogously to [\(2.12\)](#page-4-4), that

$$
||F(u)(t) - F(v)(t)||_{\alpha} \le c(T)\varepsilon, \qquad t \in [0, T], \quad v \in \bar{\mathbb{B}}_{X_T}(u, \delta),
$$

and thus the continuity of $F: X_T \to X_T$.

To establish the compactness of F, let $(u_k)_k$ be a sequence in X_T with $||u_k||_{X_T} \leq L$ for some $L > 0$. Then, in view of (1.2) , (2.13) reduces to

$$
||F(u_k)(t) - F(u_k)(s)||_{\alpha} \le \int_s^t ||e^{(t-\tau)A}||_{\mathcal{L}(E_{\gamma}, E_{\alpha})} ||f(\tau, u_k(\tau))||_{\gamma} d\tau + \int_0^s ||e^{(t-\tau)A} - e^{(s-\tau)A}||_{\mathcal{L}(E_{\gamma}, E_{\alpha})} ||f(\tau, u_k(\tau))||_{\gamma} d\tau + ||(e^{tA} - e^{sA})u^0||_{\alpha} \le c(T, L)(t-s)^{\eta-\alpha} + ||(e^{tA} - e^{sA})u^0||_{\alpha}, \qquad k \in \mathbb{N}, \quad 0 < s < t \le T,
$$

where $\eta \in (\alpha, 1)$ and $[t \mapsto e^{tA}u_0] \in C([0, T], E_\alpha)$. This establishes the equi-continuity of $(F(u_k))_k$. Moreover, arguing as in [\(2.8\)](#page-3-4), we also have (due to [\(1.2\)](#page-1-6)), for some fixed $\eta \in (\alpha, 1)$,

$$
||F(u_k)(t)||_{\eta} \le c(T, L)t^{\alpha - \eta}, \qquad k \in \mathbb{N}, \quad 0 < t \le T.
$$

The Arzelà-Ascoli theorem now guarantees, in view of the compactness of the embedding $E_\eta \hookrightarrow E_\alpha$, that $(F(u_k))_k$ is relatively compact in X_T .

Finally, the proof of an a priori bound for the set

$$
S := \{ u \in X_T : u = \lambda F(u) \text{ for some } \lambda \in [0,1] \}
$$

is the same as in Theorem [1.2](#page-1-3) (using $u^0 \in E_\alpha$ and [\(1.2\)](#page-1-6)). Thus, F admits a fixed point $u \in X_T$, which is a strong solution to (1.1) by the same arguments as in the last part of the proof of Theorem [1.2,](#page-1-3) taking directly $\varepsilon = 0$ there if $\gamma > 0$, respectively using [\(1.3\)](#page-1-7) in the case $\gamma = 0$. This proves Proposition [1.1.](#page-1-0)

We conclude this section by providing the proof for Corollary [1.4.](#page-2-0)

2.3. **Proof of Corollary [1.4.](#page-2-0)** We first prove the claim for $u^0 \in E_\alpha$ with $\alpha \in [0, \min\{1, 1 + \gamma - \xi\})$. Since f satisfies [\(1.4\)](#page-1-1), and if $\gamma = 0$ also [\(1.5\)](#page-1-2) for some $T_0 > 0$, Theorem [1.2](#page-1-3) guarantees the existence of a strong solution

$$
u_1 \in C^1((0,T_0],E_0) \cap C((0,T_0],E_1) \cap C([0,T_0],E_\alpha) \cap C_\mu((0,T_0],E_\xi)
$$

for some $\mu \in (\xi, 1)$, see (2.2) , to

$$
u_1' = Au_1 + f(t, u_1), \quad t \in (0, T_0], \qquad u_1(0) = u^0. \tag{2.15}
$$

Noticing that $u_1(T_0) \in E_1 \hookrightarrow E_{\xi}$, we infer from Proposition [1.1](#page-1-0) (see [\(1.6\)](#page-2-2) and [\(1.7\)](#page-2-3) if $\gamma = 0$) that there exists a strong solution

$$
u_2 \in C^1((0,T_0],E_0) \cap C((0,T_0],E_1) \cap C([0,T_0],E_{\xi})
$$

to

$$
u_2' = Au_2 + f(t + T_0, u_2), \quad t \in (0, T_0], \qquad u_2(0) = u_1(T_0).
$$
 (2.16)

Setting

$$
u(t) := \begin{cases} u_1(t), & 0 \le t \le T_0, \\ u_2(t - T_0), & T_0 \le t \le 2T_0, \end{cases}
$$

we obtain $u \in C_{\mu}((0, 2T_0], E_{\xi})$. In particular, $f(\cdot, u) \in C((0, 2T_0], E_{\gamma})$ satisfies (2.6) (with $T = 2T_0$ therein). Since u_1 and u_2 are given by the corresponding mild formulation of (2.15) respectively (2.16) (see (2.1)), it readily follows that u satisfies

$$
u(t) = e^{tA}u^{0} + \int_{0}^{t} e^{(t-\tau)A} f(\tau, u(\tau)) d\tau, \qquad t \in [0, 2T_0].
$$

We may now argue as in the proof of Theorem [1.2](#page-1-3) to deduce that

$$
u \in C^1((0, 2T_0], E_0) \cap C((0, 2T_0], E_1) \cap C([0, 2T_0], E_\alpha)
$$

is a strong solution to (1.1) on $[0, 2T_0]$. Consequently, using Proposition [1.1,](#page-1-0) we may inductively extend the solution to $[0, nT_0]$ for $n \geq 2$ and obtain in this way a global strong solution

$$
u \in C^1((0,\infty),E_0) \cap C((0,\infty),E_1) \cap C([0,\infty),E_\alpha)
$$

to (1.1) on $[0,\infty)$.

The assertion for $u^0 \in E_\alpha$ with $\alpha \in [\xi, 1)$ follows by applying Proposition [1.1](#page-1-0) successively on $[n, n+1]$ for $n \in \mathbb{N}$ and gluing the solutions as above. This proves Corollary [1.4.](#page-2-0)

3. Applications

We apply the abstract theory to concrete models. As a starting point we provide in Section [3.1](#page-8-1) a functional analytic framework for the subsequent applications. In Section [3.2](#page-9-1) we then analyze an evolution equation for the spreading of bushfires and in Section [3.3](#page-12-0) we apply the abstract results to a reaction-diffusion system. The examples presented in Section [3.2–](#page-9-1)Section [3.3](#page-12-0) effectively illustrate the significance of both Proposition [1.1](#page-1-0) and Theorem [1.2](#page-1-3) in practical applications.

3.1. Functional Analytic Setting. Let $\Omega \subset \mathbb{R}^n$, $n \in \mathbb{N}^*$, be an open and bounded set with smooth boundary and outward unit normal ν . In order to include Dirichlet and Neumann boundary conditions, we choose $\delta \in \{0,1\}$ and define

$$
\mathcal{B}u := u
$$
 on $\partial\Omega$ if $\delta = 0$, $\mathcal{B}u := \partial_{\nu}u$ on $\partial\Omega$ if $\delta = 1$,

hence B is the trace operator if $\delta = 0$ corresponding to Dirichlet boundary conditions, while $\delta = 1$ refers to Neumann boundary conditions. We introduce

$$
F_0 := L_2(\Omega), \qquad F_1 := H^2_{\mathcal{B}}(\Omega) = \{ v \in H^2(\Omega) : \mathcal{B}v = 0 \text{ on } \partial\Omega \},
$$

with $H^2(\Omega) = W^2_{2,\mathcal{B}}(\Omega)$ denoting the Sobolev space of second-order over $L_2(\Omega)$, and recall that

$$
B_0 := \Delta_{\mathcal{B}} \in \mathcal{H}(H^2_{\mathcal{B}}(\Omega), L_2(\Omega)),
$$

that is, $B_0 = \Delta_B$ with domain $H^2_B(\Omega)$ generates an analytic semigroup on $F_0 = L_2(\Omega)$. Let

$$
\{(F_{\theta}, B_{\theta}) : -1 \leq \theta < \infty\}
$$

be the interpolation-extrapolation scale generated by (F_0, B_0) and the complex interpolation functor $[\cdot, \cdot]_{\theta}$ (see [\[1,](#page-14-0) §6] and [\[2,](#page-14-5) §V.1]), that is,

$$
B_{\theta} \in \mathcal{H}(F_{1+\theta}, F_{\theta}), \quad -1 \le \theta < \infty, \tag{3.1}
$$

and, for $2\theta \neq -\delta - 1/2$, we have (see [\[1,](#page-14-0) Theorem 7.1; Equations (7.4)-(7.5)])

$$
F_{\theta} \doteq H_{\mathcal{B}}^{2\theta}(\Omega) := \begin{cases} \{v \in H^{2\theta}(\Omega) : \mathcal{B}v = 0 \text{ on } \partial\Omega\}, & \delta + \frac{1}{2} < 2\theta \le 2, \\ H^{2\theta}(\Omega), & -\frac{3}{2} + \delta < 2\theta < \delta + \frac{1}{2}. \end{cases} (3.2)
$$

Since Δ_B has bounded imaginary powers, [\[1,](#page-14-0) Remarks 6.1 (d)] provides the following reiteration property

$$
[F_{\alpha}, F_{\beta}]_{\theta} \doteq F_{(1-\theta)\alpha+\theta\beta}.
$$
\n(3.3)

3.2. A model for front propagation in bushfires. In [\[6,](#page-14-8) [7\]](#page-14-9) the authors recently proposed and studied, both from an analytical and numerical point of view, the nonlocal equation

$$
\partial_t u = \Delta u + \int_{\Omega} \left(u(t, y) - \Theta(t, y) \right)_+ K(x, y) \, dy + \left(\left(\omega + \frac{\beta(u)\nabla u}{|\nabla u|^{\nu}} \right) \cdot \nabla u \right)_-, \qquad x \in \Omega, \quad t > 0, \tag{3.4a}
$$

describing front propagation in bushfires, where the unknown $u = u(t, x)$ is the environmental temperature. Here, $r_{\pm} := \max\{0, \pm r\}$ for $r \in \mathbb{R}$ and $\nu \in [1, 2]$. We assume that Ω is on an open, bounded, smooth subset of \mathbb{R}^n with $n \geq 2$ (for an extension of the subsequent results to Lipschitz domains Ω one may use [\[14\]](#page-14-10)). We refer to [\[6,](#page-14-8)[7\]](#page-14-9) for the physical interpretation of the integral kernel K and the functions Θ , ω , and β . Equation [\(3.4a\)](#page-9-2) is supplemented by the Dirichlet boundary condition

$$
u = 0 \quad \text{on } \partial\Omega \,, \qquad t > 0\,, \tag{3.4b}
$$

and is subject to the initial condition

$$
u(0, x) = u^{0}(x), \qquad x \in \Omega.
$$
 (3.4c)

In [\[6\]](#page-14-8) the authors established the existence of local and global weak solutions to [\(3.4\)](#page-9-0) for initial data $u^0 \in H^1(\Omega)$ with $u = 0$ on $\partial\Omega$. The following theorem provides, under slightly more restrictive regularity assumptions on Θ and ω , but less restrictive integrability assumptions on Θ (and no size conditions on β , ω , and Θ when $\nu = 1$) compared to [\[6\]](#page-14-8), global strong solutions to [\(3.4\)](#page-9-0) for the larger class of initial data $u^0 \in L_2(\Omega)$.

In the context of [\(3.4\)](#page-9-0) we define

$$
H_D^{2\theta}(\Omega) := H_{\mathcal{B}}^{2\theta}(\Omega), \qquad 2\theta \in \left(-\frac{3}{2}, 2\right] \setminus \left\{\pm \frac{1}{2}\right\},\,
$$

with Dirichlet boundary conditions $\mathcal{B}u = 0$ on $\partial\Omega$, see [\(3.2\)](#page-8-2) with $\delta = 0$. With this notation our result reads as follows.

Theorem 3.1. Let $\nu \in [1,2]$ and assume that

$$
K \in L_2(\Omega \times \Omega), \quad \beta \in W^1_{\infty}(\mathbb{R}), \quad \Theta \in C([0,\infty), L_2(\Omega)), \quad \omega \in C([0,\infty), L_{\infty}(\Omega, \mathbb{R}^n)),
$$

and $2\varepsilon \in (0,1/7)$. Then, given $u^0 \in L_2(\Omega)$, there exists a global strong solution to [\(3.4\)](#page-9-0) such that

$$
u \in C^{1}((0,\infty), H^{-2\varepsilon}(\Omega)) \cap C((0,\infty), H_{D}^{2-2\varepsilon}(\Omega)) \cap C([0,\infty), L_{2}(\Omega)).
$$
\n(3.5)

Moreover,

- (i) if $\nu = 2$, then the solution is unique;
- (ii) if, in addition, $\Theta \in C^{\vartheta}([0,\infty), L_2(\Omega))$ and $\omega \in C^{\vartheta}([0,\infty), L_{\infty}(\Omega, \mathbb{R}^n))$ for some $\vartheta \in (0,1)$, then the solution satisfies

$$
u \in C^{1}((0, \infty), L_{2}(\Omega)) \cap C((0, \infty), H_{D}^{2}(\Omega)).
$$

Concerning the uniqueness statement for $\nu = 2$ we add the following observation.

Remark 3.2. The solution found in Theorem [3.1](#page-9-3) satisfies additionally $u \in C_{\mu}((0,T], H_D^{1+2\varepsilon}(\Omega))$ for some fixed $\mu \in (1/2 + 2\varepsilon, 1)$ and each $T > 0$. For $\nu = 2$, the uniqueness of the solution, for functions which satisfy [\(3.5\)](#page-9-4) and belong to $C_{\mu}((0,T], H_D^{1+2\varepsilon}(\Omega))$ for each $T > 0$, can be also shown by using directly the fixed point formulation [\(2.1\)](#page-2-4) together with the singular Gronwall inequality [\[2,](#page-14-5)

II.Corollary 3.3.2 (since the nonlinearity $f = f(t, u)$ is locally Lipschitz continuous in u, see [\(3.6\)](#page-10-0) and Lemma [3.3](#page-10-1) below). However, the uniqueness result in Theorem [3.1](#page-9-3) (i) is more general.

Before proving Theorem [3.1](#page-9-3) we note some auxiliary results. The following lemma is related to [\[7,](#page-14-9) Lemma 2.2, Lemma 2.4].

Lemma 3.3. Let $K \in L_2(\Omega \times \Omega)$. Then the mapping $g_0: L_2(\Omega)^2 \to L_2(\Omega)$, given by

$$
g_0(u,\Theta)(x) := \int_{\Omega} \left(u(y) - \Theta(y) \right)_+ K(x,y) \, dy, \qquad x \in \Omega, \quad u, \Theta \in L_2(\Omega),
$$

is well-defined and satisfies

$$
||g_0(u,\Theta)||_2 \le ||K||_2(||\Theta||_2 + ||u||_2), \qquad u,\Theta \in L_2(\Omega),
$$

and

$$
||g_0(u_1,\Theta_1)-g_0(u_2,\Theta_2)||_2 \le ||K||_2(||u_1-u_2||_2+||\Theta_1-\Theta_2||_2), \qquad u_i,\Theta_i \in L_2(\Omega), \quad i=1,2.
$$

Proof. The claims follow by using Hölder's inequality together with the inequality $|a_+ - b_+| \leq |a-b|$ for $a, b \in \mathbb{R}$.

We turn to the second, nonlinear term on the right of [\(3.4a\)](#page-9-2). To this end we define for $\nu \in [1,2]$ the mapping

$$
g_{\nu}: L_2(\Omega) \times L_2(\Omega, \mathbb{R}^n) \times L_{\infty}(\Omega, \mathbb{R}^n) \to L_2(\Omega)
$$

according to

$$
g_{\nu}(u,p,\omega):=\big(\omega\cdot p+\beta(u)|p|^{2-\nu}\big)_-.
$$

The following result was observed in [\[7,](#page-14-9) Lemma 2.1, Lemma 2.3, Lemma 2.5] (the proof therein is valid also for $\nu = 1$).

Lemma 3.4. If $\beta \in W^1_\infty(\mathbb{R})$ and $\nu \in [1,2]$, then for $(u, p, \omega) \in L_2(\Omega) \times L_2(\Omega, \mathbb{R}^n) \times L_\infty(\Omega, \mathbb{R}^n)$ we have

$$
||g_{\nu}(u,p,\omega)||_2\leq \|\omega\|_{\infty}\|p\|_2+|\Omega|^{(\nu-1)/2}\|\beta\|_{\infty}\|p\|_2^{2-\nu}
$$

and for $(u_i, p_i, \omega_i) \in L_2(\Omega) \times L_2(\Omega, \mathbb{R}^n) \times L_\infty(\Omega, \mathbb{R}^n)$, $i = 1, 2$,

$$
||g_{\nu}(u_1, p_1, \omega_1) - g_{\nu}(u_2, p_2, \omega_2)||_2
$$

\n
$$
\leq ||\omega_1||_{\infty} ||p_1 - p_2||_2 + ||p_2||_2 ||\omega_1 - \omega_2||_{\infty} + 2||\beta||_{W^1_{\infty}} ||p_1||_2^{2-\nu} ||u_1 - u_2||_2^{\nu-1}
$$

\n
$$
+ |\Omega|^{(\nu-1)/2} ||\beta||_{\infty} ||p_1 - p_2||_2^{2-\nu}.
$$

In fact, for $\nu = 2$,

$$
||g_2(u_1, p_1, \omega_1) - g_2(u_2, p_2, \omega_2)||_2
$$

\n
$$
\leq ||\omega_1||_{\infty} ||p_1 - p_2||_2 + ||p_2||_2 ||\omega_1 - \omega_2||_{\infty} + ||\beta||_{W^1_{\infty}} ||u_1 - u_2||_2.
$$
\n(3.6)

Proof. The claims follow by using Hölder's inequality together with the inequality $|a - b_-\| \leq |a - b|$ for $a, b \in \mathbb{R}$.

We note for $\nu = 1$ that the estimate in Lemma [3.4](#page-10-2) does not provide local Hölder continuity of g_1 . This is achieved now in Lemma [3.5](#page-11-0) by restricting the range for p.

Lemma 3.5. If $\beta \in W^1_{\infty}(\mathbb{R})$ and $2\varepsilon \in (0, 1/2)$, then there is $c > 0$ with

$$
||g_1(u_1, p_1, \omega_1) - g_1(u_2, p_2, \omega_2)||_2 \leq (||\omega_1||_{\infty} + ||\beta||_{\infty}) ||p_1 - p_2||_2 + ||p_2||_2 ||\omega_1 - \omega_2||_{\infty} + c||\beta||_{W^1_{\infty}} ||p_1||_{H^{2\varepsilon}} ||u_1 - u_2||_2^{4\varepsilon/n}
$$

for all $(u_i, p_i, \omega_i) \in L_2(\Omega) \times H^{2\varepsilon}(\Omega, \mathbb{R}^n) \times L_\infty(\Omega, \mathbb{R}^n), i = 1, 2.$

Proof. The details are identical to that in Lemma [3.4,](#page-10-2) except those used when estimating the term $||p_1(\beta(u_1) - \beta(u_2))||_2$. Letting $r_{\varepsilon}, q_{\varepsilon} \in (2,\infty)$ be given by

$$
\frac{1}{r_{\varepsilon}} := \frac{1}{2} - \frac{2\varepsilon}{n}, \qquad \frac{1}{q_{\varepsilon}} := \frac{1}{2} - \frac{1}{r_{\varepsilon}} = \frac{2\varepsilon}{n},
$$

Hölder's inequality together with the embedding $H^{2\varepsilon}(\Omega) \hookrightarrow L_{r_{\varepsilon}}(\Omega)$ implies

$$
||p_1(\beta(u_1) - \beta(u_2))||_2 \le ||p_1||_{r_{\varepsilon}} ||\beta(u_1) - \beta(u_2)||_{q_{\varepsilon}} \le c||p_1||_{H^{2\varepsilon}} \left(\int_{\Omega} |\beta(u_1) - \beta(u_2)|^{q_{\varepsilon}} dx\right)^{1/q_{\varepsilon}}
$$

$$
\le c||p_1||_{H^{2\varepsilon}} (||\beta||_{\infty})^{\frac{q_{\varepsilon}-2}{q_{\varepsilon}}} \left(\int_{\Omega} |\beta(u_1) - \beta(u_2)|^2 dx\right)^{1/q_{\varepsilon}}
$$

$$
\le c||\beta||_{W_{\infty}^1} ||p_1||_{H^{2\varepsilon}} ||u_1 - u_2||_2^{2/q_{\varepsilon}},
$$

which provides the desired estimate. \Box

We are now in a position to establish Theorem [3.1.](#page-9-3)

Proof of Theorem [3.1.](#page-9-3) Let $2\varepsilon \in (0, 1/7)$ be fixed and set

$$
E_0:=H^{-2\varepsilon}_D(\Omega)\,,\qquad E_1:=H^{2-2\varepsilon}_D(\Omega)\,.
$$

Defining $2\xi := 1 + 4\varepsilon$ and $\gamma := \varepsilon \in (0,1)$, we infer from [\(3.3\)](#page-9-5) that

$$
E_{\xi} = H_D^{1+2\varepsilon}(\Omega) , \qquad E_{\gamma} = L_2(\Omega) .
$$

We may thus recast [\(3.4\)](#page-9-0) as a semilinear parabolic Cauchy problem

$$
u' = \Delta u + f_{\nu}(t, u), \quad t > 0, \qquad u(0) = u^{0}, \tag{3.7}
$$

where $\Delta \in \mathcal{H}(E_1, E_0)$ (with Dirichlet boundary conditions, see [\(3.1\)](#page-8-3)-[\(3.2\)](#page-8-2)) and $f_\nu : [0, \infty) \times E_\xi \to E_\gamma$ for $\nu \in [1, 2]$ is defined by

$$
f_{\nu}(t, u) := g_0(u, \Theta(t)) + g_{\nu}(u, \nabla u, \omega(t)), \qquad (t, u) \in [0, \infty) \times E_{\xi}.
$$

Lemma [3.3–](#page-10-1)Lemma [3.5](#page-11-0) ensure that $f_{\nu} \in C([0,\infty) \times E_{\xi}, E_{\gamma})$ has the property that for each $T > 0$ there exists $C(T) > 0$ such that

$$
||f_{\nu}(t,u)||_{E_{\gamma}} \leq C(T)(1+||u||_{E_{\xi}}), \qquad t \in [0,T], \quad u \in E_{\xi},
$$

and, if $\nu \in (1, 2]$, then for all $t \in [0, T]$ and $u_1, u_2 \in E_{\xi}$,

$$
||f_{\nu}(t, u_1) - f_{\nu}(t, u_2)||_{E_{\gamma}} \leq C(T) \left(||u_1 - u_2||_{E_{\xi}}^{\vartheta_1} + ||u_1||_{E_{\xi}}^{\vartheta_2 - \vartheta_2} ||u_1 - u_2||_{E_{\xi}}^{\vartheta_2} + ||u_1 - u_2||_{E_{\xi}}^{\vartheta_3} \right)
$$

with

$$
\vartheta_1 := 1 =: q_1, \quad 0 \le \vartheta_2 := \nu - 1 \le q_2 := 1, \quad 0 \le \vartheta_3 := 2 - \nu =: q_3 \le 1,
$$

while, for $\nu = 1$, we have

$$
||f_1(t, u_1) - f_1(t, u_2)||_{E_{\gamma}} \leq C(T) \left(||u_1 - u_2||_{E_{\xi}}^{\vartheta_1} + ||u_1||_{E_{\xi}}^{\vartheta_1 - \vartheta_1} ||u_1 - u_2||_{E_{\xi}}^{\vartheta_2} \right)
$$

with

$$
\theta_1 := 1 =: q_1, \quad 0 \le \theta_2 := 4\varepsilon/n \le q_2 := 1 + \theta_2.
$$

The largest value q of the exponents q_i is easily computed as

$$
q := \begin{cases} 1, & \nu \in (1,2], \\ 1 + 4\varepsilon/n, & \nu = 1, \end{cases}
$$

and, since $n \geq 2$ and $2\varepsilon < 1/7$, it holds that $\xi < 1/q = \min\{1, 1/q\}$.

This shows that assumption [\(1.4\)](#page-1-1) is fulfilled in the context of [\(3.7\)](#page-11-1) for each $T > 0$. We may thus apply Corollary [1.4](#page-2-0) (a) with $\alpha := \varepsilon \in [0, 1/2 - \varepsilon) = [0, \min\{1, 1 + \gamma - \xi\}]$ and deduce, for each $u^0 \in E_\alpha = L_2(\Omega)$, the existence of a global strong solution

$$
u \in C^1((0,\infty),E_0) \cap C((0,\infty),E_1) \cap C([0,\infty),E_\alpha)
$$

to [\(3.7\)](#page-11-1).

Regarding the uniqueness claim for $\nu = 2$ stated in (i), let u_1, u_2 be solutions to [\(3.4\)](#page-9-0) as found in Theorem [3.1](#page-9-3) such that $u_1(0) = u_2(0)$ and choose an arbitrary $T > 0$. Using the Lions-Magenes lemma, see [\[4,](#page-14-11) Theorem II.5.12], we have for $0 < \tilde{t} < t \leq T$

$$
||(u_1 - u_2)(t)||_2^2 - ||(u_1 - u_2)(\tilde{t})||_2^2 = -2 \int_{\tilde{t}}^t ||\nabla(u_1 - u_2)(\tau)||_2^2 d\tau
$$

+
$$
2 \int_{\tilde{t}}^t \int_{\Omega} (u_1 - u_2)(\tau) [f_2(\tau, u_1(\tau)) - f_2(\tau, u_2(\tau))] dx d\tau.
$$

Since $||f_2(\tau, u_1(\tau)) - f_2(\tau, u_2(\tau))||_2 \leq C(T) ||(u_1 - u_2)(\tau)||_{H^1}$ for $\tau \in (0, T]$ due to Lemma [3.3](#page-10-1) and Lemma [3.4,](#page-10-2) Hölder's inequality, Young's inequality, and the previous estimate lead, after taking the limit $\tilde{t} \to 0$, to

$$
||(u_1 - u_2)(t)||_2^2 \leq C(T) \int_0^t ||(u_1 - u_2)(\tau)||_2^2 d\tau, \qquad 0 \leq t \leq T.
$$

The desired assertion follows now from Gronwall's lemma.

Finally, let $\Theta \in C^{\vartheta}([0,\infty), L_2(\Omega))$ and $\omega \in C^{\vartheta}([0,\infty), L_\infty(\Omega, \mathbb{R}^n))$ for some $\vartheta \in (0,1)$ and fix $\delta \in (0,1)$. Since $f(\cdot, u) \in C([\delta, \infty), L_2(\Omega))$, $u(\delta) \in H_0^{3/2}$ $\mathcal{O}_0^{3/2}(\Omega)$, and $\Delta \in \mathcal{H}(H_0^2(\Omega), L_2(\Omega))$, we infer from [\[2,](#page-14-5) II. Theorem 5.3.1] that $u \in C^{\rho}([\delta,\infty),E_{\xi})$ for some $\rho \in (0,1)$. This property together with Lemma [3.3-](#page-10-1)Lemma [3.5](#page-11-0) show that $f(\cdot, u) \in C^{\rho'}([\delta, \infty), L_2(\Omega))$ for some $\rho' \in (0, \rho)$. In view of [\[2,](#page-14-5) II. Theorem 1.2.1] we get $u \in C^1((\delta,\infty), L_2(\Omega)) \cap C((\delta,\infty), H_0^2(\Omega))$ and, since $\delta \in (0,1)$ is chosen arbitrarily, this establishes (ii), and the proof of Theorem [3.1](#page-9-3) is complete. \Box

3.3. Dynamics of an isothermal, autocatalytic chemical reaction scheme. We consider a mathematical model for the dynamics of an isothermal, autocatalytic chemical reaction scheme with termination, taking place in an unstirred environment and undergoing molecular diffusion investigated in a one-dimensional setting in [\[12\]](#page-14-12). The governing equations are summarized by the

14 B.–V. MATIOC AND CH. WALKER

system

$$
\partial_t u - \Delta u = -u_+^{\mu} v_+^{\beta} , \qquad \qquad t > 0, \quad x \in \Omega , \qquad (3.8a)
$$

$$
\partial_t v - \Delta v = u_+^{\mu} v_+^{\beta} - a v_+^{\theta} , \qquad t > 0, \quad x \in \Omega , \qquad (3.8b)
$$

where u and v are concentrations of the reactant and the autocatalyst, respectively, a is a (positive) constant, and the constants $\mu, \beta, \theta \in (0, 1]$ are assumed to satisfy $\mu + \beta \leq 1$. As before, we set $r_+ := \max\{0, r\}$ for $r \in \mathbb{R}$. Moreover, $\Omega \subset \mathbb{R}^n$, $n \in \mathbb{N}^*$, is an open and bounded set with smooth boundary and outward unit normal ν .

We supplemented [\(3.8a\)](#page-13-0)-[\(3.8b\)](#page-13-1) by homogeneous Neumann boundary conditions

$$
\partial_{\nu}u = \partial_{\nu}v = 0 \qquad \text{on } \partial\Omega, \quad t > 0,
$$
\n(3.8c)

and impose the initial condition

$$
u(0, x) = u^{0}(x), \quad v(0, x) = v^{0}(x), \qquad x \in \Omega.
$$
 (3.8d)

In the context of the evolution problem [\(3.8\)](#page-13-2) we chose the Hilbert spaces $H_N^{2\theta}(\Omega)$, which are defined by [\(3.2\)](#page-8-2) with $\mathcal{B}u = \partial_{\nu}u$, to establish the following result.

Theorem 3.6. Let $a \in \mathbb{R}$ and μ , β , $\theta \in (0,1]$ satisfy $\mu + \beta \leq 1$. Then, given $u^0 \in L_2(\Omega)$, there exists a global strong solution to [\(3.8\)](#page-13-2) such that

$$
(u, v) \in C^{1}((0, \infty), L_{2}(\Omega, \mathbb{R}^{2})) \cap C((0, \infty), H_{N}^{2}(\Omega, \mathbb{R}^{2})) \cap C([0, \infty), L_{2}(\Omega, \mathbb{R}^{2})) .
$$
 (3.9)

Proof. Set

$$
E_0 := L_2(\Omega, \mathbb{R}^2), \qquad E_1 := H_N^2(\Omega, \mathbb{R}^2),
$$

and $\xi := \gamma := 0$. The evolution problem [\(3.8\)](#page-13-2) may be recast as the semilinear parabolic Cauchy problem

$$
w' = Aw + f(w), \quad t > 0, \qquad w(0) = w^0 := (u^0, v^0), \tag{3.10}
$$

where

$$
A:=\begin{pmatrix} \Delta & 0 \\ 0 & \Delta \end{pmatrix} \in \mathcal{H}(E_1,E_0),
$$

see [\[2,](#page-14-5) I. Theorem 1.6.1] and [\(3.1\)](#page-8-3)-[\(3.2\)](#page-8-2), and $f : E_0 \to E_0$ is given by

$$
f(w) := (-u_+^{\mu}v_+^{\beta}, u_+^{\mu}v_+^{\beta} - av_+^{\theta}), \qquad w = (u, v) \in E_0.
$$

In fact, using Hölder's inequality, for $w = (u, v) \in E_0$ we have

$$
\|v^{\theta}_{+}\|_{2} \leq |\Omega|^{(1-\theta)/2} \|v\|_{2}^{\theta} \leq |\Omega|^{(1-\theta)/2} \|w\|_{E_{0}}^{\theta},
$$

$$
\|u^{\mu}_{+}v^{\beta}_{+}\|_{2} \leq |\Omega|^{(1-\beta-\mu)/2} \|u\|_{2}^{\mu} \|v\|_{2}^{\beta} \leq |\Omega|^{(1-\beta-\mu)/2} \|w\|_{E_{0}}^{\mu+\beta},
$$
\n(3.11)

which shows that f is well-defined and satisfies

$$
||f(w)||_{E_0} \le C\big(1 + ||w||_{E_0}\big), \qquad w \in E_0.
$$
\n(3.12)

Moreover, [\(3.11\)](#page-13-3) together with the inequality

$$
|a^{\vartheta}_+ - b^{\vartheta}_+| \le |a - b|^{\vartheta}, \qquad a, b \in \mathbb{R}, \quad \vartheta \in (0, 1],
$$

shows that there exists a constant $C > 0$ such that for all $w, \tilde{w} \in E_0$ we have

$$
||f(w) - f(\tilde{w})||_{E_0} \le C(||w - \tilde{w}||_{E_0}^{\theta} + ||w||_{E_0}^{\mu}||w - \tilde{w}||_{E_0}^{\beta} + ||w||_{E_0}^{\beta}||w - \tilde{w}||_{E_0}^{\mu}).
$$
\n(3.13)

In view of [\(3.12\)](#page-13-4) and [\(3.13\)](#page-13-5), we may thus apply Corollary [1.4](#page-2-0) (b) (with $\alpha = \xi = \gamma = 0$) in the context of the evolution problem [\(3.10\)](#page-13-6) to deduce, for each $w^0 \in E_0$, the existence of a strong solution to [\(3.8\)](#page-13-2) with the required regularity property [\(3.9\)](#page-13-7). \Box

There are other examples to which the abstract results Theorem [1.2](#page-1-3) and Corollary [1.4](#page-2-0) apply, e.g., to reaction–diffusion equations of the form

 $\partial_t u - \Delta u = u|u|^{p-1}, \quad t > 0, \quad x \in \Omega,$

with $p \in (0,1)$, see [\[5\]](#page-14-13) and the references therein. In this case, the function $f: L_2(\Omega) \to L_2(\Omega)$ with $f(u) := u|u|^{p-1}$ is linearly bounded and uniformly p-Hölder continuous.

REFERENCES

- [1] H. Amann, Nonhomogeneous linear and quasilinear elliptic and parabolic boundary value problems, in Function spaces, differential operators and nonlinear analysis (Friedrichroda, 1992), vol. 133 of Teubner-Texte Math., Teubner, Stuttgart, 1993, p. 9–126.
- [2] $_______________________\._______\._$ Boston, Inc., Boston, MA, 1995. Abstract linear theory.
- [3] W. ARENDT AND R. CHILL, Global existence for quasilinear diffusion equations in isotropic nondivergence form, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5), 9 (2010), pp. 523–539.
- [4] F. Boyer and P. Fabrie, Mathematical tools for the study of the incompressible Navier-Stokes equations and related models, vol. 183 of Applied Mathematical Sciences, Springer, New York, 2013.
- [5] V. Clark and J. C. Meyer, On two-signed solutions to a second order semi-linear parabolic partial differential equation with non-Lipschitz nonlinearity, J. Differential Equations, 269 (2020), pp. 1401–1431.
- [6] S. DIPIERRO, E. VALDINOCI, G. WHEELER, AND V.-M. WHEELER, Existence theory for a bushfire equation. arXiv:2402.15762, 2024.
- \rightarrow , A simple but effective bushfire model: analysis anbd real-time simulations. arXiv:2402.15753, 2024.
- [8] D. GILBARG AND N. S. TRUDINGER, Elliptic Partial Differential Equations of Second Order, Springer Verlag, Berlin, 2001.
- [9] O. A. LADYŽENSKAJA, V. A. SOLONNIKOV, AND N. N. URAL'CEVA, *Linear and quasilinear equations of para*bolic type, vol. Vol. 23 of Translations of Mathematical Monographs, American Mathematical Society, Providence, RI, 1968. Translated from the Russian by S. Smith.
- [10] G. M. LIEBERMAN, Second order parabolic differential equations, World Scientific Publishing Co., Inc., River Edge, NJ, 1996.
- [11] B.-V. Matioc and C. Walker, Well-posedness of quasilinear parabolic equations in time-weighted spaces. arXiv:2312.07974, 2023.
- [12] J. C. Meyer, Theoretical aspects of the Cauchy problem for non-Lipschitz semi-linear parabolic partial differential equations, ProQuest LLC, Ann Arbor, MI, 2013. Thesis (Ph.D.)–University of Birmingham (United Kingdom).
- [13] J. C. MEYER AND D. J. NEEDHAM, The Cauchy problem for non-Lipschitz semi-linear parabolic partial differential equations, vol. 419 of London Mathematical Society Lecture Note Series, Cambridge University Press, Cambridge, 2015.
- [14] I. Wood, Maximal L^p-regularity for the Laplacian on Lipschitz domains, Math. Z., 255 (2007), pp. 855–875.

Fakultät für Mathematik, Universität Regensburg, 93053 Regensburg, Germany. Email address: bogdan.matioc@ur.de

LEIBNIZ UNIVERSITÄT HANNOVER, INSTITUT FÜR ANGEWANDTE MATHEMATIK, WELFENGARTEN 1, 30167 HANnover, Germany.

Email address: walker@ifam.uni-hannover.de