

Two-Dimensional Coulomb Gas on an Elliptic Annulus

Taro Nagao

*Graduate School of Mathematics, Nagoya University, Chikusa-ku,
Nagoya 464-8602, Japan*

Abstract

It is well-known that two-dimensional Coulomb gases at a special inverse temperature $\beta = 2$ can be analyzed by using the orthogonal polynomial method borrowed from the theory of random matrices. In this paper, such Coulomb gas molecules are studied when they are distributed on an elliptic annulus, and the asymptotic forms of the molecule correlation functions in the thermodynamic limit are evaluated. For that purpose, two-dimensional orthogonality relations of the Chebyshev polynomials on an elliptic annulus are utilized.

PACS: 02.50.-r; 05.20.Jj

KEYWORDS: two-dimensional Coulomb gas; orthogonal polynomials;
random matrices

1 Introduction

Let us consider two-dimensional one-component Coulomb gas systems with Hamiltonians

$$\mathcal{H} = \sum_{j=1}^N \mathcal{V}(z_j) - \sum_{j<l}^N \log |z_j - z_l|. \quad (1.1)$$

Here the locations (x_j, y_j) of logarithmically interacting N molecules are denoted by complex numbers $z_j = x_j + iy_j$ ($j = 1, 2, \dots, N$) on the two dimensional complex plane \mathbf{C} . The one-particle potential functions are given by $\mathcal{V}(z)$ equated with $\mathcal{V}(\bar{z})$, where $z = x + iy$ and $\bar{z} = x - iy$ with real numbers x and y .

The equilibrium probability density function at an inverse temperature $\beta = 2$ is known to be

$$P(z_1, z_2, \dots, z_N) \propto e^{-\beta\mathcal{H}} = \prod_{j=1}^N w_{\mathbf{C}}(z_j) \prod_{j<l}^N |z_j - z_l|^2 \quad (1.2)$$

with a one-molecule weight function $w_{\mathbf{C}}(z) = e^{-2\mathcal{V}(z)}$ and an integration measure $\prod_{j=1}^N dz_j = \prod_{j=1}^N dx_j dy_j$. Here $P(z_1, z_2, \dots, z_N)$ is normalized as

$$\int_{\mathbf{C}} dz_1 \int_{\mathbf{C}} dz_2 \cdots \int_{\mathbf{C}} dz_N P(z_1, z_2, \dots, z_N) = 1, \quad (1.3)$$

and we define the k -molecule correlation functions

$$\rho(z_1, z_2, \dots, z_k) = \frac{N!}{(N-k)!} \int_{\mathbf{C}} dz_{k+1} \int_{\mathbf{C}} dz_{k+2} \cdots \int_{\mathbf{C}} dz_N P(z_1, z_2, \dots, z_N), \quad (1.4)$$

which are typical observable physical quantities.

As is well-known, the probability density function (1.2) can be analyzed by using the corresponding orthogonal polynomials on the complex plane. This fact originated in Ginibre's seminal work[1] on random matrices and then extended to general Coulomb gas systems of the above form[2, 3]. Suppose that $M_n(z) = z^n + (\text{lower order terms})$ with non-negative integers n are the monic orthogonal polynomials satisfying the orthogonality relation

$$\int_{\mathbf{C}} \sqrt{w_{\mathbf{C}}(\bar{z})w_{\mathbf{C}}(z)} M_m(\bar{z}) M_n(z) dz = h_n \delta_{mn} \quad (1.5)$$

with $m, n = 0, 1, 2, \dots$. Then the k -molecule correlation functions can be written in determinant forms[4]

$$\rho(z_1, z_2, \dots, z_k) = \det [K_N(z_j, z_\ell)]_{j,\ell=1,2,\dots,k}, \quad (1.6)$$

where

$$K_N(z_j, z_\ell) = \sqrt{w_{\mathbf{C}}(z_j)w_{\mathbf{C}}(\bar{z}_\ell)} \sum_{n=0}^{N-1} \frac{1}{h_n} M_n(z_j) M_n(\bar{z}_\ell). \quad (1.7)$$

Therefore, in order to see the asymptotic behavior of the k -molecule correlation functions in the thermodynamic limit $N \rightarrow \infty$, we only need to evaluate the kernel function $K_N(z_1, z_2)$.

Let us consider an elliptic annulus

$$\mathcal{A} = \left\{ z \mid (x/a_v)^2 + (y/b_v)^2 \leq 1 \text{ and } (x/a_R)^2 + (y/b_R)^2 \geq 1 \right\} \quad (1.8)$$

on the complex plane. Here $z = x + iy$ (x and y are real),

$$a_v = \frac{1}{2} \left(v + \frac{1}{v} \right), \quad b_v = \frac{1}{2} \left(v - \frac{1}{v} \right), \quad (1.9)$$

$$a_R = \frac{1}{2} \left(R + \frac{1}{R} \right), \quad b_R = \frac{1}{2} \left(R - \frac{1}{R} \right) \quad (1.10)$$

and $1 < R < v$. In this paper, we are interested in the Coulomb gas systems confined on the elliptic annulus \mathcal{A} . Namely the one-molecule weight function $w_{\mathbf{C}}(z)$ has a form

$$w_{\mathbf{C}}(z) = \begin{cases} w_{\mathcal{A}}(z), & z \in \mathcal{A}, \\ 0, & z \notin \mathcal{A}. \end{cases} \quad (1.11)$$

Now we moreover suppose that the weight function $w_{\mathcal{A}}(z)$ on the annulus has a flat form

$$w_{\mathcal{A}}(z) = 1 \quad (1.12)$$

or one of the forms

$$w_{\mathcal{A}}(z) = \frac{1}{|1 - z^2|}, \quad \frac{1}{|1 - z|} \text{ and } \frac{1}{|1 + z|}. \quad (1.13)$$

In these cases, we find that the corresponding orthogonal polynomials on the elliptic annulus are proportional to the Chebyshev polynomials. Then we can evaluate the asymptotic formulas of the kernel functions in the limit $N \rightarrow \infty$.

This paper is organized as follows. In §2, the orthogonality of the Chebyshev polynomials on the elliptic annulus is proved and used to obtain kernel function formulas for a finite N . In §3, $N \rightarrow \infty$ limits of the kernel functions are evaluated, when the elliptic annulus is contained in the neighborhood of the outer edge, and when it is in the neighborhood of the real interval $[-1, 1]$. In Appendices, conventional Coulomb gas results on a disc and on a real one-dimensional interval are derived for comparison.

2 Chebyshev polynomials

For a complex coordinate $z = x + iy$ (x and y are real) which is not on the real interval $[-1, 1]$, we introduce a complex parameter ω as

$$z = \frac{1}{2} \left(\omega + \frac{1}{\omega} \right) \quad (2.1)$$

with $|\omega| > 1$. Then the Chebyshev polynomials are defined in the following way. A Chebyshev polynomial with a non-negative integer index n is an n -th degree polynomial of z .

(1) Chebyshev polynomials of the first kind

$$T_n(z) = \frac{1}{2} \left(\omega^n + \omega^{-n} \right). \quad (2.2)$$

The corresponding monic orthogonal polynomials $M_n(z)$ are

$$M_n^{(1)}(z) = \begin{cases} 1, & n = 0, \\ \frac{1}{2^{n-1}} T_n(z), & n > 0. \end{cases} \quad (2.3)$$

(2) Chebyshev polynomials of the second kind

$$U_n(z) = \frac{\omega^{n+1} - \omega^{-n-1}}{\omega - \omega^{-1}} \quad (2.4)$$

with the monic orthogonal polynomials $M_n^{(2)}(z) = (1/2^n)U_n(z)$.

(3) Chebyshev polynomials of the third kind

$$V_n(z) = \frac{\omega^{n+1} + \omega^{-n}}{\omega + 1} \quad (2.5)$$

with the monic orthogonal polynomials $M_n^{(3)}(z) = (1/2^n)V_n(z)$.

(4) Chebyshev polynomials of the fourth kind

$$W_n(z) = \frac{\omega^{n+1} - \omega^{-n}}{\omega - 1} \quad (2.6)$$

with the monic orthogonal polynomials $M_n^{(4)}(z) = (1/2^n)W_n(z)$.

The Chebyshev polynomials are known to satisfy orthogonality relations[5, 6] on the elliptic region

$$\mathcal{E} = \left\{ z = x + iy \mid (x/a_v)^2 + (y/b_v)^2 \leq 1 \right\} \quad (2.7)$$

with the weight functions (1.12) and (1.13). We can readily extend them to the orthogonality relations on the elliptic annulus \mathcal{A} .

2.1 Model I

Let us first consider the Chebyshev polynomials of the first kind with the Model I weight function

$$w_{\mathcal{A}}(z) = \frac{1}{|1 - z^2|}. \quad (2.8)$$

The complex parameter ω can be written in terms of the polar coordinates (r, θ) as

$$\omega = re^{i\theta}, \quad r > 1, \quad \theta \text{ real}. \quad (2.9)$$

It follows from $z = x + iy$ that

$$\begin{aligned} \frac{\partial x}{\partial r} &= \frac{1}{2r} \left(r - \frac{1}{r} \right) \cos \theta, & \frac{\partial x}{\partial \theta} &= -\frac{1}{2} \left(r + \frac{1}{r} \right) \sin \theta, \\ \frac{\partial y}{\partial r} &= \frac{1}{2r} \left(r + \frac{1}{r} \right) \sin \theta, & \frac{\partial y}{\partial \theta} &= \frac{1}{2} \left(r - \frac{1}{r} \right) \cos \theta \end{aligned} \quad (2.10)$$

and

$$\frac{\partial(x, y)}{\partial(r, \theta)} = \frac{\partial x}{\partial r} \frac{\partial y}{\partial \theta} - \frac{\partial x}{\partial \theta} \frac{\partial y}{\partial r} = \frac{1}{4r} \left| \omega - \frac{1}{\omega} \right|^2 = \frac{|1 - z^2|}{r}. \quad (2.11)$$

When $r = v$,

$$\begin{aligned} z &= \frac{1}{2} \left(\omega + \frac{1}{\omega} \right) = \frac{1}{2} \left(ve^{i\theta} + \frac{1}{v} e^{-i\theta} \right) \\ &= a_v \cos \theta + ib_v \sin \theta \end{aligned} \quad (2.12)$$

is on the boundary of the elliptic region \mathcal{E} .

Let us now take an arbitrary function $f(r)$ of r . If relevant integrals exist, the two-dimensional integral

$$I_{mn} = \int_{\mathcal{E}} dx dy \frac{f(r)}{|1 - z^2|} M_m^{(1)}(\bar{z}) M_n^{(1)}(z) \quad (2.13)$$

over the elliptic region \mathcal{E} can be evaluated as

$$\begin{aligned} I_{mn} &= 2^{2-m-n} \int_1^v dr \int_0^{2\pi} d\theta \frac{f(r)}{r} T_m(\bar{z}) T_n(z) \\ &= 2^{-m-n} \int_1^v dr \int_0^{2\pi} d\theta \frac{f(r)}{r} (\bar{\omega}^m + \bar{\omega}^{-m})(\omega^n + \omega^{-n}) \\ &= 2^{-m-n} \int_1^v dr \frac{f(r)}{r} \int_0^{2\pi} d\theta \\ &\quad \times \left(r^{m+n} e^{i(n-m)\theta} + r^{m-n} e^{-i(m+n)\theta} + r^{n-m} e^{i(m+n)\theta} + r^{-m-n} e^{i(m-n)\theta} \right) \\ &= h_n \delta_{mn} \end{aligned} \quad (2.14)$$

for $m, n > 0$. In general, $I_{mn} = h_n \delta_{mn}$ for $m, n \geq 0$ with

$$h_n = \frac{\pi}{2^{2n-1}} \int_1^v dr f(r) (r^{2n-1} + r^{-2n-1}), \quad n > 0 \quad (2.15)$$

and

$$h_0 = 2\pi \int_1^r dr \frac{f(r)}{r}. \quad (2.16)$$

Therefore $M_n^{(1)}(z)$ are monic orthogonal polynomials with a weight function

$$w_{\mathbf{C}}(z) = \begin{cases} \frac{f(r)}{|1-z^2|}, & z \in \mathcal{E}, \\ 0, & z \notin \mathcal{E} \end{cases} \quad (2.17)$$

on the complex plane.

Now we take a special form of $f(r)$ as

$$f(r) = \begin{cases} 1, & r \geq R, \\ 0, & r < R \end{cases} \quad (2.18)$$

with $1 < R < v$. When $r = R$,

$$\begin{aligned} z &= \frac{1}{2} \left(\omega + \frac{1}{\omega} \right) = \frac{1}{2} \left(R e^{i\theta} + \frac{1}{R} e^{-i\theta} \right) \\ &= a_R \cos \theta + i b_R \sin \theta \end{aligned} \quad (2.19)$$

is on the inner boundary of the elliptic annulus \mathcal{A} . Therefore the orthogonality relation

$$\int_{\mathcal{A}} dx dy \frac{1}{|1-z^2|} M_m^{(1)}(\bar{z}) M_n^{(1)}(z) = h_n^{(1)} \delta_{mn}, \quad m, n \geq 0 \quad (2.20)$$

with

$$\begin{aligned} h_n^{(1)} &= \frac{\pi}{2^{2n-1}} \int_R^v dr (r^{2n-1} + r^{-2n-1}) \\ &= \frac{\pi}{2^{2n} n} (v^{2n} - v^{-2n} - R^{2n} + R^{-2n}), \quad n > 0 \end{aligned} \quad (2.21)$$

and

$$h_0^{(1)} = 2\pi \int_R^v dr \frac{1}{r} = 2\pi (\log v - \log R) \quad (2.22)$$

holds. It follows that the kernel function $K_N(z_1, z_2)$ with $N > 1$ is equal to

$$\begin{aligned}
K_N^{(1)}(z_1, z_2) &= \frac{1}{\sqrt{|1 - (z_1)^2||1 - (z_2)^2|}} \sum_{n=0}^{N-1} \frac{1}{h_n^{(1)}} M_n^{(1)}(z_1) M_n^{(1)}(\bar{z}_2) \\
&= \frac{1}{\sqrt{|1 - (z_1)^2||1 - (z_2)^2|}} \\
&\quad \times \left\{ \frac{1}{2\pi \log(v/R)} + \frac{1}{\pi} \sum_{n=1}^{N-1} n \frac{((\omega_1)^n + (\omega_1)^{-n})((\bar{\omega}_2)^n + (\bar{\omega}_2)^{-n})}{v^{2n} - v^{-2n} - R^{2n} + R^{-2n}} \right\},
\end{aligned} \tag{2.23}$$

where

$$z_j = \frac{1}{2} \left(\omega_j + \frac{1}{\omega_j} \right) \tag{2.24}$$

with $|\omega_j| > 1$ and $j = 1, 2$.

2.2 Model II

Next we consider the Chebyshev polynomials of the second kind corresponding to the flat Model II weight function

$$w_{\mathcal{A}}(z) = 1. \tag{2.25}$$

As before, the orthogonality relation for the Chebyshev polynomials of the second kind on the elliptic region \mathcal{E} can be derived as

$$\int_{\mathcal{E}} dx dy f(r) M_m^{(2)}(\bar{z}) M_n^{(2)}(z) = h_n \delta_{mn}, \quad m, n \geq 0, \tag{2.26}$$

where

$$h_n = \frac{\pi}{2^{2n+1}} \int_1^v dr f(r) \left(r^{2n+1} + r^{-2n-3} \right), \quad n \geq 0. \tag{2.27}$$

Focusing on the special case (2.18), we arrive at the orthogonality relation on the annulus \mathcal{A}

$$\int_{\mathcal{A}} dx dy M_m^{(2)}(\bar{z}) M_n^{(2)}(z) = h_n^{(2)} \delta_{mn}, \quad m, n \geq 0, \tag{2.28}$$

where

$$h_n^{(2)} = \frac{\pi}{2^{2n+2}(n+1)} \left(v^{2n+2} - v^{-2n-2} - R^{2n+2} + R^{-2n-2} \right), \quad n \geq 0. \tag{2.29}$$

Therefore the kernel function $K_N(z_1, z_2)$ is equal to

$$\begin{aligned} K_N^{(2)}(z_1, z_2) &= \sum_{n=0}^{N-1} \frac{1}{h_n^{(2)}} M_n^{(2)}(z_1) M_n^{(2)}(\bar{z}_2) \\ &= \frac{4}{\pi} \frac{1}{(\omega_1 - \omega_1^{-1})(\bar{\omega}_2 - \bar{\omega}_2^{-1})} \sum_{n=1}^N n \frac{((\omega_1)^n - (\omega_1)^{-n})((\bar{\omega}_2)^n - (\bar{\omega}_2)^{-n})}{v^{2n} - v^{-2n} - R^{2n} + R^{-2n}}. \end{aligned} \quad (2.30)$$

2.3 Model III

The Chebyshev polynomials of the third kind correspond to the Model III weight function

$$w_A(z) = \frac{1}{|1 - z|}. \quad (2.31)$$

Following the previous procedures, we can similarly obtain the orthogonality relation for the Chebyshev polynomials of the third kind as

$$\int_{\mathcal{E}} dx dy \frac{1}{|1 - z|} f(r) M_m^{(3)}(\bar{z}) M_n^{(3)}(z) = h_n \delta_{mn}, \quad m, n \geq 0, \quad (2.32)$$

where

$$h_n = \frac{\pi}{2^{2n}} \int_1^v dr f(r) (r^{2n} + r^{-2n-2}), \quad n \geq 0. \quad (2.33)$$

We restrict ourselves to the special case (2.18) and find the orthogonality relation

$$\int_{\mathcal{A}} dx dy \frac{1}{|1 - z|} M_m^{(3)}(\bar{z}) M_n^{(3)}(z) = h_n^{(3)} \delta_{mn}, \quad m, n \geq 0, \quad (2.34)$$

where

$$h_n^{(3)} = \frac{\pi}{2^{2n}(2n+1)} (v^{2n+1} - v^{-2n-1} - R^{2n+1} + R^{-2n-1}), \quad n \geq 0. \quad (2.35)$$

Then the kernel function $K_N(z_1, z_2)$ is equated with

$$\begin{aligned} K_N^{(3)}(z_1, z_2) &= \frac{1}{\sqrt{|1 - z_1||1 - z_2|}} \sum_{n=0}^{N-1} \frac{1}{h_n^{(3)}} M_n^{(3)}(z_1) M_n^{(3)}(\bar{z}_2) \\ &= \frac{1}{\pi} \frac{1}{(\omega_1 + 1)(\bar{\omega}_2 + 1)} \frac{1}{\sqrt{|1 - z_1||1 - z_2|}} \\ &\quad \times \sum_{n=0}^{N-1} (2n+1) \frac{((\omega_1)^{n+1} + (\omega_1)^{-n})((\bar{\omega}_2)^{n+1} + (\bar{\omega}_2)^{-n})}{v^{2n+1} - v^{-2n-1} - R^{2n+1} + R^{-2n-1}}. \end{aligned} \quad (2.36)$$

2.4 Model IV

Finally, for the Chebyshev polynomials of the fourth kind associated with the Model IV weight function

$$w_{\mathcal{A}}(z) = \frac{1}{|1+z|}, \quad (2.37)$$

the orthogonality relation turns out to be

$$\int_{\mathcal{E}} dx dy \frac{1}{|1+z|} f(r) M_m^{(4)}(\bar{z}) M_n^{(4)}(z) = h_n \delta_{mn}, \quad m, n \geq 0, \quad (2.38)$$

where

$$h_n = \frac{\pi}{2^{2n}} \int_1^v dr f(r) (r^{2n} + r^{-2n-2}), \quad n \geq 0. \quad (2.39)$$

In the special case (2.18), it is reduced to be

$$\int_{\mathcal{A}} dx dy \frac{1}{|1+z|} M_m^{(4)}(\bar{z}) M_n^{(4)}(z) = h_n^{(4)} \delta_{mn}, \quad m, n \geq 0, \quad (2.40)$$

where

$$h_n^{(4)} = \frac{\pi}{2^{2n}(2n+1)} (v^{2n+1} - v^{-2n-1} - R^{2n+1} + R^{-2n-1}), \quad n \geq 0. \quad (2.41)$$

We then find the equality of the kernel functions $K_N(z_1, z_2)$ and

$$\begin{aligned} K_N^{(4)}(z_1, z_2) &= \frac{1}{\sqrt{|1+z_1||1+z_2|}} \sum_{n=0}^{N-1} \frac{1}{h_n^{(4)}} M_n^{(4)}(z_1) M_n^{(4)}(\bar{z}_2) \\ &= \frac{1}{\pi} \frac{1}{(\omega_1 - 1)(\bar{\omega}_2 - 1)} \frac{1}{\sqrt{|1+z_1||1+z_2|}} \\ &\quad \times \sum_{n=0}^{N-1} (2n+1) \frac{((\omega_1)^{n+1} - (\omega_1)^{-n})(\bar{\omega}_2)^{n+1} - (\bar{\omega}_2)^{-n}}{v^{2n+1} - v^{-2n-1} - R^{2n+1} + R^{-2n-1}}. \end{aligned} \quad (2.42)$$

3 Asymptotic forms

In this section, we discuss the asymptotic forms of the kernel functions in the thermodynamic limit $N \rightarrow \infty$.

3.1 Model I

Our first case is equipped with the Model I weight function (2.8), for which the kernel function $K_N^{(1)}(z_1, z_2)$ is given in (2.23). Now we introduce the polar coordinates $r_1, r_2 > 1$ and real θ_1, θ_2 as

$$\omega_j = r_j e^{i\theta_j}, \quad j = 1, 2. \quad (3.1)$$

When the elliptic annulus is contained in the neighborhood of the outer edge $r_1, r_2 = v > 1$, we introduce the scaling parameters t_1, t_2, ϕ_1, ϕ_2 and T as

$$r_1 = v \left(1 - \frac{t_1}{N}\right), \quad r_2 = v \left(1 - \frac{t_2}{N}\right), \quad (3.2)$$

$$\theta_1 = \psi + \frac{\phi_1}{N}, \quad \theta_2 = \psi + \frac{\phi_2}{N} \quad (3.3)$$

and

$$R = v \left(1 - \frac{T}{N}\right), \quad (3.4)$$

where $0 < t_1, t_2 < T$ and $0 \leq \psi < 2\pi$. Thus the angular coordinates θ_1, θ_2 are in the neighborhood of a constant angle ψ .

Then it is straightforward to obtain asymptotic relations

$$\begin{aligned} & \sqrt{|1 - (z_1)^2| |1 - (z_2)^2|} \\ &= \frac{1}{4} \sqrt{((r_1)^2 + (r_1)^{-2} - 2 \cos 2\theta_1) ((r_2)^2 + (r_2)^{-2} - 2 \cos 2\theta_2)} \\ &\sim \frac{1}{4} (v^2 + v^{-2} - 2 \cos 2\psi), \quad N \rightarrow \infty, \end{aligned} \quad (3.5)$$

$$\begin{aligned} & v^{2n} - v^{-2n} - R^{2n} + R^{-2n} \\ &= v^{2n} - v^{-2n} - v^{2n} \left(1 - \frac{T}{N}\right)^{2n} + v^{-2n} \left(1 - \frac{T}{N}\right)^{-2n} \\ &\sim v^{2n} (1 - e^{-2cT}), \quad N \rightarrow \infty, \end{aligned} \quad (3.6)$$

$$\begin{aligned} & ((\omega_1)^n + (\omega_1)^{-n}) ((\bar{\omega}_2)^n + (\bar{\omega}_2)^{-n}) \\ &= \left(v^n \left(1 - \frac{t_1}{N}\right)^n e^{in(\psi + \frac{\phi_1}{N})} + v^{-n} \left(1 - \frac{t_1}{N}\right)^{-n} e^{-in(\psi + \frac{\phi_1}{N})} \right) \\ & \quad \times \left(v^n \left(1 - \frac{t_2}{N}\right)^n e^{-in(\psi + \frac{\phi_2}{N})} + v^{-n} \left(1 - \frac{t_2}{N}\right)^{-n} e^{in(\psi + \frac{\phi_2}{N})} \right) \\ &\sim v^{2n} e^{-c(t_1+t_2)} e^{ic(\phi_1-\phi_2)}, \quad N \rightarrow \infty \end{aligned} \quad (3.7)$$

with $c = n/N$ fixed. Putting these asymptotic relations into (2.23), we find

$$\begin{aligned} & K_N^{(1)}(z_1, z_2) \\ \sim & \frac{4N^2}{\pi} \frac{1}{v^2 + v^{-2} - 2 \cos 2\psi} \int_0^1 dc \, c \frac{e^{-c(t_1+t_2)} e^{ic(\phi_1-\phi_2)}}{1 - e^{-2cT}}, \quad N \rightarrow \infty. \end{aligned} \quad (3.8)$$

Let us then examine the thin annulus limit $T \rightarrow 0$. Noting that $t_1, t_2 \rightarrow 0$ limit must also be taken because of the inequality $0 < t_1, t_2 < T$, one obtains

$$\begin{aligned} K_N^{(1)}(z_1, z_2) & \sim \frac{2N^2 v}{\pi T} \frac{1}{v^2 + v^{-2} - 2 \cos 2\psi} \int_0^1 dc \, e^{ic(\phi_1-\phi_2)} \\ & = \rho_v^{(1)}(\psi) e^{i(\phi_1-\phi_2)/2} \frac{\sin \{(\phi_1 - \phi_2)/2\}}{(\phi_1 - \phi_2)/2}, \end{aligned} \quad (3.9)$$

where $\rho_v(\psi)$ is the asymptotic molecule density

$$\rho_v^{(1)}(\psi) = \frac{2N^2 v}{\pi T} \frac{1}{v^2 + v^{-2} - 2 \cos 2\psi} \quad (3.10)$$

on the edge of the elliptic region \mathcal{E} . As the phase factor $e^{i(\phi_j-\phi_\ell)/2}$ can be replaced with 1 without changing the determinant forms (1.6) of the k -molecule correlation functions $\rho(z_1, z_2, \dots, z_k)$, we arrive at

$$\frac{\rho(z_1, z_2, \dots, z_k)}{\{\rho_v^{(1)}(\psi)\}^k} \sim \det \left[\frac{\sin \{(\phi_j - \phi_\ell)/2\}}{(\phi_j - \phi_\ell)/2} \right]_{j,\ell=1,2,\dots,k}. \quad (3.11)$$

This formula gives the elliptic version of the sine kernel formula for the 1-dimensional Coulomb gas on a circle or the eigenvalues of the circular unitary ensemble (CUE) of random matrices[7].

On the other hand, when the elliptic annulus is in the neighborhood of the real interval $[-1, 1]$, it is useful to modify the definitions of the scaling parameters t_1, t_2, ϕ_1, ϕ_2 and T as

$$r_1 = 1 + \frac{t_1}{N}, \quad r_2 = 1 + \frac{t_2}{N}, \quad (3.12)$$

$$\theta_1 = \psi + \frac{\phi_1}{N}, \quad \theta_2 = \psi + \frac{\phi_2}{N} \quad (3.13)$$

and

$$R = 1 + \frac{T}{N}. \quad (3.14)$$

We additionally define a parameter u as

$$v = 1 + \frac{u}{N} \quad (3.15)$$

and suppose that $0 < T < t_1, t_2 < u$. The constant angle ψ is in the region $0 \leq \psi < 2\pi$. When the elliptic annulus is in the neighborhood of $[-1, 1]$, only the cases $\psi = 0$, $\psi = \pi/2$ and $\psi = \pi$ are considered.

As before we can derive asymptotic relations

$$\begin{aligned} & \sqrt{|1 - (z_1)^2||1 - (z_2)^2|} \\ &= \frac{1}{4} \sqrt{\left\{ \left(1 + \frac{t_1}{N}\right)^2 + \left(1 + \frac{t_1}{N}\right)^{-2} - 2 \cos 2 \left(\psi + \frac{\phi_1}{N}\right) \right\}} \\ & \quad \times \sqrt{\left\{ \left(1 + \frac{t_2}{N}\right)^2 + \left(1 + \frac{t_2}{N}\right)^{-2} - 2 \cos 2 \left(\psi + \frac{\phi_2}{N}\right) \right\}} \\ & \sim \begin{cases} |s_1||s_2|/N^2, & \psi = 0 \text{ or } \psi = \pi, \\ (1 - \cos 2\psi)/2, & \psi \neq 0 \text{ and } \psi \neq \pi, \end{cases} \quad N \rightarrow \infty, \end{aligned} \quad (3.16)$$

$$v^{2n} - v^{-2n} - R^{2n} + R^{-2n} \sim e^{2cu} - e^{-2cu} - e^{2cT} + e^{-2cT}, \quad N \rightarrow \infty \quad (3.17)$$

and

$$\begin{aligned} & \left((\omega_1)^n + (\omega_1)^{-n} \right) \left((\bar{\omega}_2)^n + (\bar{\omega}_2)^{-n} \right) \\ & \sim (e^{cs_1} e^{in\psi} + e^{-cs_1} e^{-in\psi}) (e^{c\bar{s}_2} e^{-in\psi} + e^{-c\bar{s}_2} e^{in\psi}), \quad N \rightarrow \infty, \end{aligned} \quad (3.18)$$

where

$$s_j = t_j + i\phi_j, \quad j = 1, 2 \quad (3.19)$$

and $c = n/N$ fixed.

In the case $\psi = 0$ or $\psi = \pi$, these asymptotic relations and (2.23) yield

$$K^{(1)}(z_1, z_2) \sim \frac{N^4}{\pi} \frac{1}{|s_1||s_2|} \int_0^1 dc \, c \frac{(e^{cs_1} + e^{-cs_1})(e^{c\bar{s}_2} + e^{-c\bar{s}_2})}{e^{2cu} - e^{-2cu} - e^{2cT} + e^{-2cT}} \quad (3.20)$$

in the limit $N \rightarrow \infty$. Let us consider the thin annulus limit shrinking to the interval $[-1, 1]$. We first put $T = 0$ and then take the limit $u \rightarrow 0$. Keeping

in mind that $u \rightarrow 0$ implies $t_1, t_2 \rightarrow 0$, we obtain

$$\begin{aligned} K_N^{(1)}(z_1, z_2) &\sim \frac{N^4}{4\pi u} \frac{1}{|\phi_1||\phi_2|} \int_0^1 dc (e^{ic\phi_1} + e^{-ic\phi_1})(e^{ic\phi_2} + e^{-ic\phi_2}) \\ &= \frac{N^4}{2\pi u} \frac{1}{|\phi_1\phi_2|} \left(\frac{\sin(\phi_1 - \phi_2)}{\phi_1 - \phi_2} + \frac{\sin(\phi_1 + \phi_2)}{\phi_1 + \phi_2} \right). \end{aligned} \quad (3.21)$$

When $\psi = \frac{\pi}{2}$, noting $e^{2in\psi} = (-1)^n$ and evaluating the sums in (2.23) for odd n and even n separately, we find

$$\begin{aligned} K_N^{(1)}(z_1, z_2) &= \frac{1}{\sqrt{|1 - (z_1)^2||1 - (z_2)^2|}} \\ &\times \left\{ \frac{1}{2\pi \log(v/R)} + \frac{1}{\pi} \sum_{n=1}^{N-1} n \frac{((\omega_1)^n + (\omega_1)^{-n})(\bar{\omega}_2^n + (\bar{\omega}_2)^{-n})}{v^{2n} - v^{-2n} - R^{2n} + R^{-2n}} \right\} \\ &= \frac{1}{\sqrt{|1 - (z_1)^2||1 - (z_2)^2|}} \left\{ \frac{1}{2\pi \log(v/R)} \right. \\ &+ \frac{1}{\pi} \sum_{n=0}^{[N/2-1]} (2n+1) \frac{((\omega_1)^{2n+1} + (\omega_1)^{-2n-1})(\bar{\omega}_2^{2n+1} + (\bar{\omega}_2)^{-2n-1})}{v^{4n+2} - v^{-4n-2} - R^{4n+2} + R^{-4n-2}} \\ &+ \left. \frac{1}{\pi} \sum_{n=1}^{[(N-1)/2]} (2n) \frac{((\omega_1)^{2n} + (\omega_1)^{-2n})(\bar{\omega}_2^{2n} + (\bar{\omega}_2)^{-2n})}{v^{4n} - v^{-4n} - R^{4n} + R^{-4n}} \right\} \\ &\sim \frac{N^2}{2\pi} \int_0^1 dc c \frac{e^{c(s_1+\bar{s}_2)} + e^{-c(s_1+\bar{s}_2)} - e^{c(s_1-\bar{s}_2)} - e^{-c(s_1-\bar{s}_2)}}{e^{2cu} - e^{-2cu} - e^{2cT} + e^{-2cT}} \\ &+ \frac{N^2}{2\pi} \int_0^1 dc c \frac{e^{c(s_1+\bar{s}_2)} + e^{-c(s_1+\bar{s}_2)} + e^{c(s_1-\bar{s}_2)} + e^{-c(s_1-\bar{s}_2)}}{e^{2cu} - e^{-2cu} - e^{2cT} + e^{-2cT}} \\ &= \frac{N^2}{\pi} \int_0^1 dc c \frac{e^{c(s_1+\bar{s}_2)} + e^{-c(s_1+\bar{s}_2)}}{e^{2cu} - e^{-2cu} - e^{2cT} + e^{-2cT}} \end{aligned} \quad (3.22)$$

in the limit $N \rightarrow \infty$. Here $[x]$ is the largest integer less than or equal to x . Taking the thin annulus limit $u \rightarrow 0$ after putting $T = 0$ results in

$$K_N^{(1)}(z_1, z_2) \sim \frac{N^2}{2\pi u} \int_0^1 dc \cos(c(\phi_1 - \phi_2)) = \rho_u^{(1)} \frac{\sin(\phi_1 - \phi_2)}{\phi_1 - \phi_2}, \quad (3.23)$$

where $\rho_u^{(1)} = \frac{N^2}{2\pi u}$. The k -molecule correlation functions $\rho(z_1, z_2, \dots, z_k)$ ac-

cordingly satisfy

$$\frac{\rho(z_1, z_2, \dots, z_k)}{\{\rho_u^{(1)}\}^k} \sim \det \left[\frac{\sin(\phi_j - \phi_\ell)}{\phi_j - \phi_\ell} \right]_{j, \ell=1, 2, \dots, k}. \quad (3.24)$$

This formula reproduces the sine kernel formula for the Coulomb gas on a 1-dimensional line originally known from the theory of the Gaussian unitary ensemble (GUE) of random matrices[7].

3.2 Model II

Next we consider Model II with the weight function (2.25) and the kernel function $K_N^{(2)}(z_1, z_2)$ in (2.30). The polar coordinates r_1, r_2, θ_1 and θ_2 are introduced in (3.1) as before.

If one is interested in the elliptic annulus in the neighborhood of the outer edge $r_1, r_2 = v > 1$, the scaling parameters t_1, t_2, ϕ_1, ϕ_2 and T defined in (3.2), (3.3) and (3.4) are again useful. Let us first derive asymptotic relations

$$\begin{aligned} & (\omega_1 - \omega_1^{-1})(\bar{\omega}_2 - \bar{\omega}_2^{-1}) \\ &= \left(v \left(1 - \frac{t_1}{N} \right) e^{i(\psi + \frac{\phi_1}{N})} - v^{-1} \left(1 - \frac{t_1}{N} \right)^{-1} e^{-i(\psi + \frac{\phi_1}{N})} \right) \\ & \quad \times \left(v \left(1 - \frac{t_2}{N} \right) e^{-i(\psi + \frac{\phi_2}{N})} - v^{-1} \left(1 - \frac{t_2}{N} \right)^{-1} e^{i(\psi + \frac{\phi_2}{N})} \right) \\ & \sim v^2 + v^{-2} - 2 \cos \psi, \quad N \rightarrow \infty \end{aligned} \quad (3.25)$$

and

$$\begin{aligned} & ((\omega_1)^n - (\omega_1)^{-n}) ((\bar{\omega}_2)^n - (\bar{\omega}_2)^{-n}) \\ &= \left(v^n \left(1 - \frac{t_1}{N} \right)^n e^{in(\psi + \frac{\phi_1}{N})} - v^{-n} \left(1 - \frac{t_1}{N} \right)^{-n} e^{-in(\psi + \frac{\phi_1}{N})} \right) \\ & \quad \times \left(v^n \left(1 - \frac{t_2}{N} \right)^n e^{-in(\psi + \frac{\phi_2}{N})} - v^{-n} \left(1 - \frac{t_2}{N} \right)^{-n} e^{in(\psi + \frac{\phi_2}{N})} \right) \\ & \sim v^{2n} e^{-c(t_1+t_2)} e^{ic(\phi_1-\phi_2)}, \quad N \rightarrow \infty \end{aligned} \quad (3.26)$$

with $c = n/N$ fixed. We put these asymptotic relations and (3.6) into (2.30) and arrive at

$$K_N^{(2)}(z_1, z_2)$$

$$\sim \frac{4N^2}{\pi} \frac{1}{v^2 + v^{-2} - 2 \cos 2\psi} \int_0^1 dc c \frac{e^{-c(t_1+t_2)} e^{ic(\phi_1-\phi_2)}}{1 - e^{-2cT}}, \quad N \rightarrow \infty. \quad (3.27)$$

This is exactly equal to the asymptotic result in (3.8) for Model I. We thus expect it to be a universal asymptotic behavior. As before it yields the elliptic version of the sine kernel formula in the thin annulus limit $T \rightarrow 0$.

We then consider an elliptic annulus in the neighborhood of the real interval $[-1, 1]$. The scaling parameters t_1, t_2, ϕ_1, ϕ_2 and T are modified and an additional parameter u is introduced according to (3.12), (3.13), (3.14) and (3.15). Resulting asymptotic relations are

$$\begin{aligned} & (\omega_1 - \omega_1^{-1})(\bar{\omega}_2 - \bar{\omega}_2^{-1}) \\ &= \left(\left(1 + \frac{t_1}{N}\right) e^{i(\psi + \frac{\phi_1}{N})} - \left(1 + \frac{t_1}{N}\right)^{-1} e^{-i(\psi + \frac{\phi_1}{N})} \right) \\ & \quad \times \left(\left(1 + \frac{t_2}{N}\right) e^{-i(\psi + \frac{\phi_2}{N})} - \left(1 + \frac{t_2}{N}\right)^{-1} e^{i(\psi + \frac{\phi_2}{N})} \right) \\ &\sim \begin{cases} 4s_1\bar{s}_2/N^2, & \psi = 0 \text{ or } \psi = \pi, \\ 2 - 2 \cos 2\psi, & \psi \neq 0 \text{ and } \psi \neq \pi, \end{cases} \quad N \rightarrow \infty \end{aligned} \quad (3.28)$$

and

$$\begin{aligned} & \left((\omega_1)^n - (\omega_1)^{-n} \right) \left((\bar{\omega}_2)^n - (\bar{\omega}_2)^{-n} \right) \\ &\sim (e^{cs_1} e^{in\psi} - e^{-cs_1} e^{-in\psi}) (e^{c\bar{s}_2} e^{-in\psi} - e^{-c\bar{s}_2} e^{in\psi}), \quad N \rightarrow \infty, \end{aligned} \quad (3.29)$$

where $s_j = t_j + i\phi_j$, $j = 1, 2$ and $c = n/N$ fixed.

In the case $\psi = 0$ or $\psi = \pi$, it follows from these asymptotic relations, (2.30) and (3.17) that

$$K_N^{(2)}(z_1, z_2) \sim \frac{N^4}{\pi} \frac{1}{s_1\bar{s}_2} \int_0^1 dc c \frac{(e^{cs_1} - e^{-cs_1})(e^{c\bar{s}_2} - e^{-c\bar{s}_2})}{e^{2cu} - e^{-2cu} - e^{2cT} + e^{-2cT}} \quad (3.30)$$

in the limit $N \rightarrow \infty$. The difference between the asymptotic limit (3.20) of Model I correlation functions and the corresponding limit (3.30) of Model II reveals a breakdown of the universality around the edges of the interval $[-1, 1]$. In order to examine the thin annulus limit, let us put $T = 0$ and then take the limit $u \rightarrow 0$. We obtain

$$\begin{aligned} K_N^{(2)}(z_1, z_2) &\sim \frac{N^4}{4\pi u} \frac{1}{\phi_1\phi_2} \int_0^1 dc (e^{ic\phi_1} - e^{-ic\phi_1})(e^{ic\phi_2} - e^{-ic\phi_2}) \\ &= \frac{N^4}{2\pi u} \frac{1}{\phi_1\phi_2} \left(\frac{\sin(\phi_1 - \phi_2)}{\phi_1 - \phi_2} - \frac{\sin(\phi_1 + \phi_2)}{\phi_1 + \phi_2} \right), \end{aligned} \quad (3.31)$$

which shows the striking sign difference of Model I and Model II results.

When $\psi = \frac{\pi}{2}$, as before we can derive

$$K_N^{(2)}(z_1, z_2) \sim \frac{N^2}{\pi} \int_0^1 dc c \frac{e^{c(s_1+\bar{s}_2)} + e^{-c(s_1+\bar{s}_2)}}{e^{2cu} - e^{-2cu} - e^{2cT} + e^{-2cT}} \quad (3.32)$$

in the limit $N \rightarrow \infty$. This is exactly equal to the corresponding limit (3.22) of Model I result. Thus the universal behavior is recovered.

3.3 Model III

Model III has the weight function (2.31) and the kernel function $K_N^{(3)}(z_1, z_2)$ in (2.36). We use the polar coordinates r_1, r_2, θ_1 and θ_2 introduced in (3.1).

In order to study the elliptic annulus in the neighborhood of the outer edge $r_1, r_2 = v > 1$, we again employ the scaling parameters t_1, t_2, ϕ_1, ϕ_2 and T defined in (3.2), (3.3) and (3.4). Then we find asymptotic relations

$$\begin{aligned} |1 - z_j| &= \left| 1 - \frac{1}{2} \left(\omega_j + \frac{1}{\omega_j} \right) \right| = \left| 1 - \frac{1}{2} \left(r_j e^{i\theta_j} + \frac{1}{r_j} e^{-i\theta_j} \right) \right| \\ &= \frac{1}{2} \left| \left(\sqrt{r_j} e^{i\theta_j/2} - \frac{1}{\sqrt{r_j}} e^{-i\theta_j/2} \right)^2 \right| = \frac{1}{2} \left| \sqrt{r_j} e^{i\theta_j/2} - \frac{1}{\sqrt{r_j}} e^{-i\theta_j/2} \right|^2 \\ &= \frac{1}{2} (r_j + r_j^{-1} - 2 \cos \theta_j) \\ &\sim \frac{1}{2} (v + v^{-1} - 2 \cos \psi), \quad N \rightarrow \infty \end{aligned} \quad (3.33)$$

with $j = 1, 2$,

$$\begin{aligned} (\omega_1 + 1)(\bar{\omega}_2 + 1) &= (r_1 e^{i\theta_1} + 1)(r_2 e^{-i\theta_2} + 1) \\ &\sim (v e^{i\psi} + 1)(v e^{-i\psi} + 1) = v(v + v^{-1} + 2 \cos \psi), \quad N \rightarrow \infty \end{aligned} \quad (3.34)$$

and

$$\begin{aligned} & \left((\omega_1)^{n+1} + (\omega_1)^{-n} \right) \left((\bar{\omega}_2)^{n+1} + (\bar{\omega}_2)^{-n} \right) \\ & \sim v^{2(n+1)} e^{-c(t_1+t_2)} e^{ic(\phi_1-\phi_2)}, \quad N \rightarrow \infty \end{aligned} \quad (3.35)$$

with $c = n/N$ fixed. Putting these asymptotic relations and

$$v^{2n+1} - v^{-2n-1} - R^{2n+1} + R^{-2n-1} \sim v^{2n+1} (1 - e^{-2cT}), \quad N \rightarrow \infty \quad (3.36)$$

into (2.36) yields

$$\begin{aligned} & K_N^{(3)}(z_1, z_2) \\ \sim & \frac{4N^2}{\pi} \frac{1}{v^2 + v^{-2} - 2 \cos 2\psi} \int_0^1 dc c \frac{e^{-c(t_1+t_2)} e^{ic(\phi_1-\phi_2)}}{1 - e^{-2cT}}, \quad N \rightarrow \infty. \end{aligned} \quad (3.37)$$

This should be a universal asymptotic behavior, because it is again identical to the results in (3.8) and (3.27) for Model I and Model II.

Let us modify the scaling parameters t_1, t_2, ϕ_1, ϕ_2 and T , and introduce an additional parameter u as written in (3.12), (3.13), (3.14) and (3.15). Then, in the limit $N \rightarrow \infty$, the elliptic annulus shrinks into the neighborhood of the real interval $[-1, 1]$. We require asymptotic formulas

$$\begin{aligned} |1 - z_j| &= \frac{1}{2}(r_j + r_j^{-1} - 2 \cos \theta_j) \\ &= \frac{1}{2} \left(\left(1 + \frac{t_j}{N}\right) + \left(1 + \frac{t_j}{N}\right)^{-1} - 2 \cos \left(\psi + \frac{\phi_j}{N}\right) \right) \\ &\sim \begin{cases} |s_j|^2/(2N^2), & \psi = 0, \\ (2 - 2 \cos \psi)/2, & \psi \neq 0, \end{cases} \quad N \rightarrow \infty, \end{aligned} \quad (3.38)$$

$$\begin{aligned} \omega_j + 1 &= r_j e^{i\theta_j} + 1 \\ &= \left(1 + \frac{t_j}{N}\right) e^{i(\psi + \phi_j/N)} + 1 \\ &\sim \begin{cases} -s_j/N & \psi = \pi, \\ e^{i\psi} + 1, & \psi \neq \pi, \end{cases} \quad N \rightarrow \infty \end{aligned} \quad (3.39)$$

with $j = 1, 2$,

$$\begin{aligned} & \left((\omega_1)^{n+1} + (\omega_1)^{-n} \right) \left((\bar{\omega}_2)^{n+1} + (\bar{\omega}_2)^{-n} \right) \\ \sim & (e^{cs_1} e^{i(n+1)\psi} + e^{-cs_1} e^{-in\psi}) (e^{c\bar{s}_2} e^{-i(n+1)\psi} + e^{-c\bar{s}_2} e^{in\psi}), \\ & N \rightarrow \infty \end{aligned} \quad (3.40)$$

and

$$v^{2n+1} - v^{-2n-1} - R^{2n+1} + R^{-2n-1} \sim e^{2cu} - e^{-2cu} - e^{2cT} + e^{-2cT}, \quad N \rightarrow \infty. \quad (3.41)$$

Here $s_j = t_j + i\phi_j$, $j = 1, 2$ and $c = n/N$ fixed.

In the case $\psi = 0$, these asymptotic relations and (2.36) lead to

$$K_N^{(3)}(z_1, z_2) \sim \frac{N^4}{\pi} \frac{1}{|s_1||s_2|} \int_0^1 dc c \frac{(e^{cs_1} + e^{-cs_1})(e^{c\bar{s}_2} + e^{-c\bar{s}_2})}{e^{2cu} - e^{-2cu} - e^{2cT} + e^{-2cT}} \quad (3.42)$$

in the limit $N \rightarrow \infty$. This is equal to the asymptotic limit (3.20) of $K_N^{(1)}(z_1, z_2)$ with $\psi = 0$. On the other hand, in the case $\psi = \pi$, we obtain

$$K_N^{(3)}(z_1, z_2) \sim \frac{N^4}{\pi} \frac{1}{s_1\bar{s}_2} \int_0^1 dc c \frac{(e^{cs_1} - e^{-cs_1})(e^{c\bar{s}_2} - e^{-c\bar{s}_2})}{e^{2cu} - e^{-2cu} - e^{2cT} + e^{-2cT}} \quad (3.43)$$

in the limit $N \rightarrow \infty$. This is equal to the asymptotic limit (3.30) of $K_N^{(2)}(z_1, z_2)$ with $\psi = \pi$. Around the right edge of the interval $[-1, 1]$, Model III behaves like Model I, and around the left edge, it behaves like Model II. This is expected from the forms of the weight functions.

Putting $\psi = \frac{\pi}{2}$, we find

$$K_N^{(3)}(z_1, z_2) \sim \frac{N^2}{\pi} \int_0^1 dc c \frac{e^{c(s_1+\bar{s}_2)} + e^{-c(s_1+\bar{s}_2)}}{e^{2cu} - e^{-2cu} - e^{2cT} + e^{-2cT}} \quad (3.44)$$

in the limit $N \rightarrow \infty$. This is again equal to the corresponding limits (3.22) and (3.32). It is the universal behavior expected when $\psi = \pi/2$.

3.4 Model IV

We finally consider Model IV. The weight function is (2.37) and the kernel function $K_N^{(4)}(z_1, z_2)$ is (2.42). Let us as before use the polar coordinates r_1, r_2, θ_1 and θ_2 introduced in (3.1).

The scaling parameters t_1, t_2, ϕ_1, ϕ_2 and T in (3.2), (3.3) and (3.4) are again used in order to magnify the region in the neighborhood of the outer edge $r_1, r_2 = v > 1$. The asymptotic relations

$$|1 + z_j| = \frac{1}{2}(r_j + r_j^{-1} + 2 \cos \theta_j) \sim \frac{1}{2}(v + v^{-1} + 2 \cos \psi), \quad N \rightarrow \infty \quad (3.45)$$

with $j = 1, 2$,

$$(\omega_1 - 1)(\bar{\omega}_2 - 1) = (r_1 e^{i\theta_1} - 1)(r_2 e^{-i\theta_2} - 1) \sim v(v + v^{-1} - 2 \cos \psi), \quad N \rightarrow \infty \quad (3.46)$$

and

$$\begin{aligned} & \left((\omega_1)^{n+1} - (\omega_1)^{-n} \right) \left((\bar{\omega}_2)^{n+1} - (\bar{\omega}_2)^{-n} \right) \\ & \sim v^{2(n+1)} e^{-c(t_1+t_2)} e^{ic(\phi_1-\phi_2)}, \quad N \rightarrow \infty \end{aligned} \quad (3.47)$$

follow with $c = n/N$ fixed. One puts these relations and (3.36) into (2.42). Then one obtains

$$\begin{aligned} & K_N^{(4)}(z_1, z_2) \\ & \sim \frac{4N^2}{\pi} \frac{1}{v^2 + v^{-2} - 2 \cos 2\psi} \int_0^1 dc \, c \frac{e^{-c(t_1+t_2)} e^{ic(\phi_1-\phi_2)}}{1 - e^{-2cT}}, \quad N \rightarrow \infty. \end{aligned} \quad (3.48)$$

We see that it is again identified with the universal results in (3.8), (3.27) and (3.37) for Model I, Model II and Model III, respectively.

On the other hand, in order to magnify the region in the neighborhood of the real interval $[-1, 1]$, we redefine the scaling parameters $t_1, t_2, \phi_1, \phi_2, T$ and u as written in (3.12), (3.13), (3.14) and (3.15). Then, in the limit $N \rightarrow \infty$, asymptotic formulas

$$\begin{aligned} |1 + z_j| &= \frac{1}{2}(r_j + r_j^{-1} + 2 \cos \theta_j) \\ &= \frac{1}{2} \left(\left(1 + \frac{t_j}{N} \right) + \left(1 + \frac{t_j}{N} \right)^{-1} + 2 \cos \left(\psi + \frac{\phi_j}{N} \right) \right) \\ &\sim \begin{cases} |s_j|^2 / (2N^2), & \psi = \pi, \\ (2 + 2 \cos \psi) / 2, & \psi \neq \pi, \end{cases} \quad N \rightarrow \infty, \end{aligned} \quad (3.49)$$

$$\begin{aligned} \omega_j - 1 &= r_j e^{i\theta_j} - 1 \\ &= \left(1 + \frac{t_j}{N} \right) e^{i(\psi + \phi_j/N)} - 1 \\ &\sim \begin{cases} s_j/N & \psi = 0, \\ e^{i\psi} - 1, & \psi \neq 0, \end{cases} \quad N \rightarrow \infty \end{aligned} \quad (3.50)$$

with $j = 1, 2$ and

$$\begin{aligned} & \left((\omega_1)^{n+1} - (\omega_1)^{-n} \right) \left((\bar{\omega}_2)^{n+1} - (\bar{\omega}_2)^{-n} \right) \\ & \sim (e^{cs_1} e^{i(n+1)\psi} - e^{-cs_1} e^{-in\psi}) (e^{c\bar{s}_2} e^{-i(n+1)\psi} - e^{-c\bar{s}_2} e^{in\psi}), \\ & N \rightarrow \infty \end{aligned} \quad (3.51)$$

are obtained. Here $s_j = t_j + i\phi_j$, $j = 1, 2$ and $c = n/N$ fixed.

In the case $\psi = 0$, from these asymptotic relations, (2.42) and (3.41), we find

$$K_N^{(4)}(z_1, z_2) \sim \frac{N^4}{\pi} \frac{1}{s_1 \bar{s}_2} \int_0^1 dc c \frac{(e^{cs_1} - e^{-cs_1})(e^{c\bar{s}_2} - e^{-c\bar{s}_2})}{e^{2cu} - e^{-2cu} - e^{2cT} + e^{-2cT}} \quad (3.52)$$

in the limit $N \rightarrow \infty$. This is equal to the asymptotic limit (3.30) of $K_N^{(2)}(z_1, z_2)$ with $\psi = 0$. In the case $\psi = \pi$, an asymptotic formula

$$K_N^{(4)}(z_1, z_2) \sim \frac{N^4}{\pi} \frac{1}{|s_1||s_2|} \int_0^1 dc c \frac{(e^{cs_1} + e^{-cs_1})(e^{c\bar{s}_2} + e^{-c\bar{s}_2})}{e^{2cu} - e^{-2cu} - e^{2cT} + e^{-2cT}} \quad (3.53)$$

is obtained in the limit $N \rightarrow \infty$. This is equal to the asymptotic limit (3.20) of $K_N^{(1)}(z_1, z_2)$ with $\psi = \pi$.

In the remaining case $\psi = \frac{\pi}{2}$, one arrives at

$$K_N^{(4)}(z_1, z_2) \sim \frac{N^2}{\pi} \int_0^1 dc c \frac{e^{c(s_1+\bar{s}_2)} + e^{-c(s_1+\bar{s}_2)}}{e^{2cu} - e^{-2cu} - e^{2cT} + e^{-2cT}} \quad (3.54)$$

in the limit $N \rightarrow \infty$. This is equal to the universal limits (3.22), (3.32) and (3.44), as expected.

Summary and discussion

Two-dimensional Coulomb gases distributed on an elliptic annulus at a special inverse temperature $\beta = 2$ were studied, and asymptotic forms of the correlation functions in the thermodynamic limit were evaluated. For an elliptic annulus contained in the neighborhood of the outer edge, we found universal asymptotic forms of the correlation functions. When the annulus shrinks into the real interval $[-1, 1]$, non-universal behavior was observed around the edges -1 and 1 , where the weight functions $w_{\mathcal{A}}(z)$ on the annulus have singular points. The models considered in this paper are related to the Chebyshev polynomials, and their orthogonality on the complex plane were employed in the analysis. More general polynomials related to the Gegenbauer polynomials are also known to be orthogonal on the complex plane[8, 9]. There are also Coulomb gas models associated with such more general polynomials, and it is interesting to check the universal behavior by using such more general models.

Acknowledgement

The author acknowledges support by the Japan Society for the Promotion of Science (KAKENHI 20K03764).

References

- [1] J. Ginibre, *J. Math. Phys.* 6 (1965) 440.
- [2] P. J. Forrester, *Nucl. Phys.* B904 (2016) 253.
- [3] P. J. Forrester, *Log-Gases and Random Matrices*. (Princeton University Press, 2010).
- [4] B.A. Khoruzhenko and H.-J. Sommers, *The Oxford Handbook of Random Matrix Theory* (ed. by G. Akemann, J. Baik and P. Di Francesco, Oxford University Press, 2011) chapter 18.
- [5] P. Henrici, *Applied and Computational Complex Analysis. Vol. 3* (Wiley-Interscience, 1986).
- [6] J.C. Mason and D.C. Handscomb, *Chebyshev Polynomials* (Chapman and Hall/CRC, 2002).
- [7] M.L. Mehta, *Random Matrices* (Elsevier, 3rd edition, 2004).
- [8] T. Nagao, G. Akemann, M. Kieburg and I. Parra, *J. Phys. A: Math. Theor.* 53 (2020) 075201.
- [9] G. Akemann, T. Nagao, I. Parra and G. Vernizzi, *Constr. Approx.* 53 (2021) 441.
- [10] D. Fox and P.B. Kahn, *Phys. Rev.* 134 (1964) B1151.
- [11] T. Nagao and M. Wadati, *J. Phys. Soc. Jpn.* 60 (1991) 3298.
- [12] T. Nagao and K. Slevin, *J. Math. Phys.* 34 (1993) 2075.
- [13] G. Szegő, *Orthogonal Polynomials* (American Mathematical Society, 4th edition, 1975).

Appendix A Radially symmetric case

In this Appendix, we study a conventional radially symmetric case with a probability density function

$$P(z_1, z_2, \dots, z_N) \propto \prod_{j=1}^N w_{\mathbf{C}}(z_j) \prod_{j<l}^N |z_j - z_l|^2 \quad (\text{A.1})$$

with a one-molecule weight function

$$w_{\mathbf{C}}(z) = \begin{cases} w_{\mathcal{D}}(z), & z \in \mathcal{D}, \\ 0, & z \notin \mathcal{D}, \end{cases} \quad (\text{A.2})$$

where \mathcal{D} is an annulus

$$\mathcal{D} = \{z \mid |z|^2 \leq (v/2)^2 \text{ and } |z|^2 \geq (R/2)^2\} \quad (\text{A.3})$$

on the complex plane. Here $1 < R < v$. Moreover the weight function is supposed to be

$$w_{\mathcal{D}}(z) = |z|^\gamma, \quad (\text{A.4})$$

where γ is an arbitrary real number. Then $w_{\mathbf{C}}(z)$ is fully determined by the radius $|z|$. Such a radially symmetric case is conventional, because the corresponding monic orthogonal polynomials $M_n(z)$ are known to be monomials z^n . Here asymptotic limit of the kernel function is evaluated for comparison with the Coulomb gas on an elliptic annulus. The orthogonality constant h_n for $M_n(z) = z^n$ is

$$h_n = \frac{2\pi}{2n + \gamma + 2} \frac{v^{2n+\gamma+2} - R^{2n+\gamma+2}}{2^{2n+\gamma+2}}, \quad (\text{A.5})$$

because of the integral

$$\begin{aligned} & \int_{\mathbf{C}} \sqrt{w_{\mathbf{C}}(\bar{z})w_{\mathbf{C}}(z)} M_m(\bar{z}) M_n(z) dx dy \\ &= \int_{R/2}^{v/2} dr \int_0^{2\pi} d\theta r^{m+n+\gamma+1} e^{i(n-m)\theta} \\ &= \delta_{mn} \frac{2\pi}{2n + \gamma + 2} \frac{v^{2n+\gamma+2} - R^{2n+\gamma+2}}{2^{2n+\gamma+2}}. \end{aligned} \quad (\text{A.6})$$

Let us calculate the asymptotic limit of the kernel function

$$\begin{aligned} K_{N,\gamma}^{(RS)}(z_1, z_2) &= \sqrt{w_{\mathcal{D}}(z_1)w_{\mathcal{D}}(\bar{z}_2)} \sum_{n=0}^{N-1} \frac{1}{h_n} M_n(z_1)M_n(\bar{z}_2) \\ &= \frac{(r_1 r_2)^{\gamma/2}}{2\pi} \sum_{n=0}^{N-1} (2n + \gamma + 2) \frac{(r_1 r_2)^n e^{in(\theta_1 - \theta_2)}}{(v/2)^{2n+\gamma+2} - (R/2)^{2n+\gamma+2}}, \end{aligned} \quad (\text{A.7})$$

where $z_j = r_j e^{i\theta_j}$ with a positive r_j and a real θ_j ($j = 1, 2$). Introducing the scaling parameters t_1, t_2, ϕ_1, ϕ_2 and T as

$$r_1 = \frac{v}{2} \left(1 - \frac{t_1}{N}\right), \quad r_2 = \frac{v}{2} \left(1 - \frac{t_2}{N}\right), \quad (\text{A.8})$$

$$\theta_1 = \psi + \frac{\phi_1}{N}, \quad \theta_2 = \psi + \frac{\phi_2}{N} \quad (\text{A.9})$$

and

$$R = \frac{v}{2} \left(1 - \frac{T}{N}\right), \quad (\text{A.10})$$

where $0 < t_1, t_2 < T$ and $0 \leq \psi < 2\pi$, we find

$$K_{N,\gamma}^{(RS)}(z_1, z_2) \sim \frac{4N^2}{\pi v^2} \int_0^1 dc \, c \frac{e^{-c(t_1+t_2)} e^{ic(\phi_1-\phi_2)}}{1 - e^{-2cT}} \quad (\text{A.11})$$

in the limit $N \rightarrow \infty$ with $c = n/N$ and γ fixed. This is equal to the $v \rightarrow \infty$ limit of the universal formula (3.8), (3.27), (3.37) and (3.48) as expected, and itself shows a universal behavior because it is independent of γ .

Appendix B Bessel kernel

In this Appendix, we consider another conventional model with a probability density function

$$P(x_1, x_2, \dots, x_N) \propto \prod_{j=1}^N w_{\mathbf{R}}(x_j) \prod_{j<l}^N |x_j - x_l|^2. \quad (\text{B.1})$$

It is a one-dimensional model and real numbers x_1, x_2, \dots, x_N are molecule positions. A one-molecule weight function is

$$w_{\mathbf{R}}(x) = \begin{cases} w_{[-1,1]}(x), & x \in [-1, 1], \\ 0, & x \notin [-1, 1], \end{cases} \quad (\text{B.2})$$

where x is real and $w_{[-1,1]}(x) = (1-x)^a(1+x)^b$ with $a, b > -1$. This model is called the Jacobi unitary ensemble in random matrix theory[10, 11, 12]. In the limit that the elliptic annulus shrinks into the interval $[-1, 1]$, the two-dimensional Coulomb gas model is reduced to this one-dimensional model.

The monic orthogonal polynomials corresponding to the Jacobi unitary ensemble is

$$M_n(x) = 2^n n! \frac{\Gamma(a+b+n+1)}{\Gamma(a+b+2n+1)} P_n^{(a,b)}(x) \quad (\text{B.3})$$

and the orthogonality constant h_n is

$$h_n = 2^{2n+a+b+1} n! \frac{\Gamma(a+n+1)\Gamma(b+n+1)\Gamma(a+b+n+1)}{\Gamma(a+b+2n+1)\Gamma(a+b+2n+2)}. \quad (\text{B.4})$$

Here $\Gamma(x)$ is the Gamma function and $P_n^{(a,b)}(x)$ are the Jacobi polynomials

$$P_n^{(a,b)}(x) = \frac{(-1)^n}{2^n n!} (1-x)^{-a} (1+x)^{-b} \frac{d^n}{dx^n} \left\{ (1-x)^a (1+x)^b (1-x^2)^n \right\}. \quad (\text{B.5})$$

Using an asymptotic form of the Jacobi polynomials[13]

$$P_n^{(a,b)} \left(1 - \frac{\phi^2}{2N^2} \right) \sim \left(\frac{\phi}{2N} \right)^{-a} J_a(c\phi) \quad (\text{B.6})$$

in the limit $N \rightarrow \infty$ with $c = n/N$ and $\phi > 0$ fixed, we can derive the asymptotic form of the kernel function[12]

$$\begin{aligned} K_{N,a,b}^{Jacobi}(x_1, x_2) &= \sqrt{w_{[-1,1]}(x_1)w_{[-1,1]}(x_2)} \sum_{n=0}^{N-1} \frac{1}{h_n} M_n(x_1)M_n(x_2) \\ &\sim K_a^{Bessel}(\phi_1^+, \phi_2^+) = N^2 \int_0^1 dc c J_a(c\phi_1^+) J_a(c\phi_2^+) \end{aligned} \quad (\text{B.7})$$

in the limit $N \rightarrow \infty$, where

$$x_1 = 1 - \frac{(\phi_1^+)^2}{2N^2}, \quad x_2 = 1 - \frac{(\phi_2^+)^2}{2N^2} \quad (\text{B.8})$$

with $\phi_1^+, \phi_2^+ > 0$ fixed. This asymptotic form $K_a^{Bessel}(\phi_1^+, \phi_2^+)$ is called the Bessel kernel, as $J_a(x)$ is the Bessel function. The asymptotic correlation functions of this one-dimensional model have determinant forms in terms of the Bessel kernel.

The Chebyshev polynomials are special cases of the Jacobi polynomials. In particular the cases $(a, b) = (-1/2, -1/2)$ and $(1/2, 1/2)$ are the Chebyshev polynomials of the first and second kind, respectively. Now we put $a = -1/2$ in the Bessel kernel, we obtain

$$K_{-1/2}^{Bessel}(\phi_1^+, \phi_2^+) = \frac{N^2}{\pi} \frac{1}{\sqrt{\phi_1^+ \phi_2^+}} \left(\frac{\sin(\phi_1^+ - \phi_2^+)}{\phi_1^+ - \phi_2^+} + \frac{\sin(\phi_1^+ + \phi_2^+)}{\phi_1^+ + \phi_2^+} \right). \quad (\text{B.9})$$

This special case of Bessel kernel can be derived from a similar asymptotic form (3.21) for $K_N^{(1)}(z_1, z_2)$ of Model I, when it is integrated over the transversal coordinates.

Let us explain the transversal integration in the case $\psi = 0$. Using (2.11) and the notations for (3.20), we find

$$\frac{\partial(x_j, y_j)}{\partial(r_j, \theta_j)} = \frac{|1 - z_j|^2}{r_j} \sim \frac{|s_j|^2}{N^2 r_j}, \quad N \rightarrow \infty \quad (\text{B.10})$$

for $j = 1, 2$. Therefore a factor $K_N^{(1)}(z_1, z_2)K_N^{(1)}(z_2, z_1)$ in the determinant forms of the correlation functions is integrated over the transversal coordinates t_1, t_2 as

$$\begin{aligned} & \int K_N^{(1)}(z_1, z_2)K_N^{(1)}(z_2, z_1)dx_1dy_1dx_2dy_2 \\ &= \int K_N^{(1)}(z_1, z_2)K_N^{(1)}(z_2, z_1)\frac{|1 - z_1^2|^2}{r_1}\frac{|1 - z_2^2|^2}{r_2}dr_1d\theta_1dr_2d\theta_2 \\ &\sim \frac{1}{N^4} \sum_{\pm} \int_T^u dt_1 \int_T^u dt_2 K_N^{(1)}(z_1, z_2)K_N^{(1)}(z_2, z_1) \frac{|s_1 s_2|^2}{|\phi_1 \phi_2|} dx_1 dx_2, \quad N \rightarrow \infty \end{aligned} \quad (\text{B.11})$$

with $\phi_1, \phi_2 \neq 0$. Here a sum

$$\sum_{\pm} = \sum_{\epsilon_1 = \pm 1} \sum_{\epsilon_2 = \pm 1} \quad (\text{B.12})$$

with $\phi_j = \epsilon_j |\phi_j|$ should be included in the transversal integration, because both of $\epsilon_j = \pm 1$ with the same $|\phi_j|$ yield the same x_j ($j = 1, 2$). In the derivation of the last line of (B.11), we use the relation

$$x_j = \frac{1}{2} \left(r_j + \frac{1}{r_j} \right) \cos \frac{\phi_j}{N} \sim 1 + \frac{t_j^2 - \phi_j^2}{2N^2}, \quad N \rightarrow \infty \quad (\text{B.13})$$

($j = 1, 2$). Now we put $T = 0$, insert the corresponding asymptotic formula (3.21) and take the limit $u \rightarrow 0$. Then the sum \sum_{\pm} gives a factor 4 and we obtain

$$\begin{aligned} & \lim_{u \rightarrow 0} \frac{N^4}{(\pi u)^2} \int_0^u dt_1 \int_0^u dt_2 \frac{|s_1 s_2|^2}{|\phi_1 \phi_2|^3} \left(\frac{\sin(\phi_1 - \phi_2)}{\phi_1 - \phi_2} + \frac{\sin(\phi_1 + \phi_2)}{\phi_1 + \phi_2} \right)^2 dx_1 dx_2 \\ & = K_r^{(1)}(\phi_1, \phi_2) K_r^{(1)}(\phi_2, \phi_1) dx_1 dx_2, \end{aligned} \quad (\text{B.14})$$

where

$$K_r^{(1)}(\phi_1, \phi_2) = \frac{N^2}{\pi} \frac{1}{\sqrt{|\phi_1 \phi_2|}} \left(\frac{\sin(\phi_1 - \phi_2)}{\phi_1 - \phi_2} + \frac{\sin(\phi_1 + \phi_2)}{\phi_1 + \phi_2} \right). \quad (\text{B.15})$$

In general we can similarly show

$$\begin{aligned} & \int K_N^{(1)}(z_1, z_2) K_N^{(1)}(z_2, z_3) \cdots K_N^{(1)}(z_{j-1}, z_j) K_N^{(1)}(z_j, z_1) \\ & \quad \times dx_1 dy_1 dx_2 dy_2 \cdots dx_j dy_j \\ \sim & K_r^{(1)}(\phi_1, \phi_2) K_r^{(1)}(\phi_2, \phi_3) \cdots K_r^{(1)}(\phi_{j-1}, \phi_j) K_r^{(1)}(\phi_j, \phi_1) dx_1 dx_2 \cdots dx_j, \\ & \quad j = 2, 3, \dots, \kappa \end{aligned} \quad (\text{B.16})$$

and

$$\int K_N^{(1)}(z_1, z_1) dx_1 dy_1 \sim K_r^{(1)}(\phi_1, \phi_1) dx_1 \quad (\text{B.17})$$

for integrals over transversal coordinates with an integer $\kappa \geq 2$. As a determinant can be defined as a sum over permutations and a permutation has a cycle decomposition, it follows that

$$\begin{aligned} & \int \det[K_N^{(1)}(z_j, z_\ell)]_{j,\ell=1,2,\dots,k} dx_1 dy_1 dx_2 dy_2 \cdots dx_k dy_k \\ \sim & \det[K_r^{(1)}(\phi_j, \phi_\ell)]_{j,\ell=1,2,\dots,k} dx_1 dx_2 \cdots dx_k \end{aligned} \quad (\text{B.18})$$

for the transversal integration of the determinant formula (1.6). Here $z_j = x_j + iy_j$,

$$x_j = \frac{1}{2} \left(r_j + \frac{1}{r_j} \right) \cos \theta_j, \quad y_j = \frac{1}{2} \left(r_j - \frac{1}{r_j} \right) \sin \theta_j \quad (\text{B.19})$$

with $\theta_j = \phi_j/N$ ($r_j > 1$, ϕ_j real) for $j = 1, 2, \dots, k$. Therefore $K_r^{(1)}(\phi_j, \phi_\ell)$ with $j, \ell = 1, 2, \dots, k$ gives the one-dimensional kernel function after the transversal integration. Since

$$K_r^{(1)}(\phi_j, \phi_\ell) = K_r^{(1)}(-\phi_j, \phi_\ell) = K_r^{(1)}(\phi_j, -\phi_\ell) = K_r^{(1)}(-\phi_j, -\phi_\ell), \quad (\text{B.20})$$

we find $K_r^{(1)}(\phi_j, \phi_\ell) = K_r^{(1)}(|\phi_j|, |\phi_\ell|)$. Because of (B.8) and (B.13), ϕ_j^+ can be equated with $|\phi_j|$ in the limit $u \rightarrow 0$, so that the one-dimensional kernel functions (B.9) and (B.15) satisfy

$$K_{-1/2}^{Bessel}(\phi_j^+, \phi_\ell^+) \sim K_r^{(1)}(|\phi_j|, |\phi_\ell|) = K_r^{(1)}(\phi_j, \phi_\ell), \quad (\text{B.21})$$

as expected.

Let us next put $a = 1/2$ in the Bessel kernel (B.7). Then we find

$$K_{1/2}^{Bessel}(\phi_1^+, \phi_2^+) = \frac{N^2}{\pi} \frac{1}{\sqrt{\phi_1^+ \phi_2^+}} \left(\frac{\sin(\phi_1^+ - \phi_2^+)}{\phi_1^+ - \phi_2^+} - \frac{\sin(\phi_1^+ + \phi_2^+)}{\phi_1^+ + \phi_2^+} \right). \quad (\text{B.22})$$

As before, we can derive this special case of Bessel kernel from a similar asymptotic form (3.31) for Model II. That is, a transversal integration of the determinant formula (1.6) leads to

$$\begin{aligned} & \int \det[K_N^{(2)}(z_j, z_\ell)]_{j,\ell=1,2,\dots,k} dx_1 dy_1 dx_2 dy_2 \cdots dx_k dy_k \\ & \sim \det[K_r^{(2)}(\phi_j, \phi_\ell)]_{j,\ell=1,2,\dots,k} dx_1 dx_2 \cdots dx_k, \end{aligned} \quad (\text{B.23})$$

where

$$K_r^{(2)}(\phi_j, \phi_\ell) = \frac{N^2}{\pi} \frac{1}{\sqrt{|\phi_j \phi_\ell|}} \left(\frac{\sin(\phi_j - \phi_\ell)}{\phi_j - \phi_\ell} - \frac{\sin(\phi_j + \phi_\ell)}{\phi_j + \phi_\ell} \right) \quad (\text{B.24})$$

with $j, \ell = 1, 2, \dots, k$. Owing to a relation

$$K_r^{(2)}(\phi_j, \phi_\ell) = -K_r^{(2)}(-\phi_j, \phi_\ell) = -K_r^{(2)}(\phi_j, -\phi_\ell) = K_r^{(2)}(-\phi_j, -\phi_\ell), \quad (\text{B.25})$$

$K_r^{(2)}(\phi_j, \phi_\ell)$ can be replaced with $K_r^{(2)}(|\phi_j|, |\phi_\ell|)$ in (B.23) without changing the determinant. As before the one-dimensional kernel functions (B.22) and (B.24) satisfy

$$K_{1/2}^{Bessel}(\phi_j^+, \phi_\ell^+) \sim K_r^{(2)}(|\phi_j|, |\phi_\ell|) \quad (\text{B.26})$$

by equating ϕ_j^+ with $|\phi_j|$.

Note that a similar argument can be applied to the sine kernel formula (3.23). Because both of

$$\theta_j = \pm \left(\frac{\pi}{2} + \frac{\phi_j}{N} \right) \quad (\text{B.27})$$

give the same

$$x_j = \frac{1}{2} \left(r_j + \frac{1}{r_j} \right) \cos \theta_j = \frac{1}{2} \left(r_j + \frac{1}{r_j} \right) \cos \left(\frac{\pi}{2} + \frac{\phi_j}{N} \right) \quad (\text{B.28})$$

and an asymptotic relation

$$x_j \sim -\frac{\phi_j}{N}, \quad N \rightarrow \infty \quad (\text{B.29})$$

holds ($j = 1, 2, \dots, k$), we can derive a transversal integration formula

$$\begin{aligned} & \int \det[K_N^{(1)}(z_j, z_\ell)]_{j,\ell=1,2,\dots,k} dx_1 dy_1 dx_2 dy_2 \cdots dx_k dy_k \\ & \sim \det[K_c(\phi_j, \phi_\ell)]_{j,\ell=1,2,\dots,k} dx_1 dx_2 \cdots dx_k \end{aligned} \quad (\text{B.30})$$

for the determinant (1.6). Here $K_c(\phi_j, \phi_\ell)$ is a one-dimensional kernel function

$$K_c(\phi_j, \phi_\ell) = \frac{N \sin(\phi_j - \phi_\ell)}{\pi \phi_j - \phi_\ell} \quad (\text{B.31})$$

with $j, \ell = 1, 2, \dots, k$. This is a known result [10, 11, 12] for the Jacobi unitary ensemble.