

POST-POISSON ALGEBRAS AND POISSON YANG-BAXTER EQUATION VIA BIMODULE ALGEBRA DEFORMATIONS

SIYUAN CHEN, CHENGMING BAI, AND LI GUO

ABSTRACT. A fundamental construction of Poisson algebras is as the quasiclassical limits (QCLs) of associative algebra deformations of commutative associative algebras. This paper extends this construction to the relative context with the notion of (bi)module algebras over another algebra for a given algebraic structure. In this language, a module Poisson algebras can be realized as the QCLs of bimodule associative deformations of module commutative associative algebras. Moreover, the notion of the scalar deformation of an \mathcal{O} -operator is introduced so that the process of bimodule algebras deformations to QCLs is endowed with \mathcal{O} -operators in a consistent manner. As an explicit illustration of this process, post-Poisson algebras are realized as the QCLs of bimodule associative deformations of module commutative associative algebras with the identity maps as \mathcal{O} -operators, recovering the known fact that post-Poisson algebras are the QCLs of tridendriform algebra deformations of commutative tridendriform algebras. Furthermore, the notion of scalar deformations of solutions of the associative Yang-Baxter equation (AYBE) is applied to realize solutions of the Poisson Yang-Baxter equation (PYBE) in Poisson algebras as solutions of the AYBE in commutative associative algebras, giving a YBE version of Poisson algebras as the QCLs of associative deformations of the commutative associative algebras. Finally, concrete solutions of the PYBE are obtained from the aforementioned tridendriform deformation-to-QCLs process.

CONTENTS

1. Introduction	2
1.1. Deformations of algebras and quasiclassical limits	2
1.2. \mathcal{O} -operators and classical Yang-Baxter type equations	2
1.3. Deformations in the relative contexts and for classical Yang-Baxter type equations	3
1.4. Organization of the paper	4
2. Bimodule algebra deformations and \mathcal{O} -operators	4
2.1. Deformations of bimodule algebras and the corresponding quasiclassical limits	5
2.2. \mathcal{O} -operators in bimodule algebra deformations	8
3. Bimodule algebra perspective of tridendriform deformations	10
3.1. Deformations of tridendriform algebras and the corresponding QCLs	10
3.2. Constructions of bimodule algebra deformations and tridendriform deformations by derivations	15
4. Scalar deformations of solutions of AYBE	17
4.1. Scalar deformations of solutions of AYBE in associative deformations	17
4.2. Topological dual bimodules	19
4.3. AYBE and PYBE via bimodule algebra deformations: tridendriform deformations	22
4.4. AYBE and PYBE via bimodule deformations: dendriform deformations	27
References	29

2020 *Mathematics Subject Classification.* 13D10, 16W60, 17B38, 17B63, 53D55, 13N15 .

Key words and phrases. Bimodule algebra deformation; quasiclassical limit; \mathcal{O} -operator; classical Yang-Baxter equation; post-Poisson algebra; dendriform algebra; tridendriform algebra; derivation.

1. INTRODUCTION

An importance of the Poisson algebra is as the quasiclassical limit (QCL) of an associative algebra deformation of a commutative associative algebra. This paper extends this process to the relative context, through the bimodule algebra deformations. Then by applying \mathcal{O} -operators, the above classical process is extended to post-Poisson algebras as the QCLs of tridendriform deformation of commutative tridendriform algebras, as well as classical Yang-Baxter type equations in commutative associative algebras, deformed associative algebras and Poisson algebras.

1.1. Deformations of algebras and quasiclassical limits. The deformation theory of associative algebras was initiated by Gerstenhaber [18]. For an associative algebra (A, \cdot) , its formal deformation is an associative product \cdot_h on $A[[\hbar]]$ such that

$$x \cdot_h y = x \cdot y + \sum_{s=1}^{\infty} x \cdot_s y \hbar^s, \quad x, y \in A,$$

where $x \cdot_s y \in A$. When (A, \cdot) is commutative, then with the bracket $[\cdot, \cdot]$ defined by

$$[x, y] = x \cdot_1 y - y \cdot_1 x,$$

the triple $(A, [\cdot, \cdot], \cdot)$ is called the **quasiclassical limit (QCL)** of the deformation.¹ The triple $(A, [\cdot, \cdot], \cdot)$ is a Poisson algebra, and the (topological) algebra $(A[[\hbar]], \cdot_h)$ is called a quantization of the Poisson algebra $(A, [\cdot, \cdot], \cdot)$. The quantization procedure establishes a transition from Poisson bracket to commutator, which corresponds to the transition from classical mechanics to quantum mechanics. The problem of quantizing the Poisson algebra of smooth functions on a Poisson manifold was resolved in the well-known work of Kontsevich [23].

The study for deformation and QCL has been extended to dendriform algebras and tridendriform algebras, both of which are associative algebras whose products split into several operations. Dendriform algebras were introduced by Loday [26] in his study of algebraic K -theory and tridendriform algebras were introduced by Loday and Ronco [27] in their study of the Koszul duality and polytopes. The QCL of the dendriform algebra deformation of a Zinbiel algebra, which is the commutative version of a dendriform algebra, was shown by Aguiar [2] to be a pre-Poisson algebra. It is further shown by Ospel, Panaite and Vanhaecke [35] that the QCL corresponding to the tridendriform algebra deformation of a commutative tridendriform algebra is a post-Poisson algebra, a notion introduced in [7] from an operadic point of view and in [34] related to Poisson bialgebras.

1.2. \mathcal{O} -operators and classical Yang-Baxter type equations. The concept of an \mathcal{O} -operator was introduced by Kupershmidt [24] for Lie algebras to study classical Yang-Baxter equation (CYBE). The \mathcal{O} -operators on associative algebras, also under the names of relative Rota-Baxter operators [11] and generalized Rota-Baxter operators [38], have been studied extensively recently [9, 13], in the contexts of Poisson structures, deformations and cohomology. As in the Lie algebra case, an \mathcal{O} -operator on an associative algebra can be regarded as the operator form of the solution of the associative Yang-Baxter equation (AYBE), introduced by Aguiar [1, 3] and by Polishchuk [36] in the form with spectral parameters. Numerous applications of the AYBE can be found in bialgebra theory [3], mirror symmetry [36], algebraic geometry [37], integrable systems [33] and magnetization dynamics [4].

Defined in a Poisson algebra, the Poisson Yang-Baxter equation (PYBE) is a combination of AYBE and CYBE. PYBE arose from the study of Poisson bialgebras [34]. The relationship between the \mathcal{O} -operator on a Poisson algebra and PYBE has been studied in [25, 34] and is

¹Quasiclassical limit also bears the name classical limit or semiclassical limit. We follow the convention of [15].

parallel to the associative case studied in [9, 10]. Since the Poisson algebra is the quasiclassical limit of an associative deformation, the similarities between the operator characterizations of the AYBE and the PYBE motivate us to consider the bimodule deformation and further the bimodule algebra deformation with an \mathcal{O} -operator.

1.3. Deformations in the relative contexts and for classical Yang-Baxter type equations.

This study extends the above QCL construction of Poisson algebras by deformations to the contexts of relative pairs of algebras, of \mathcal{O} -operators and pre-structures, and of solutions of classical Yang-Baxter type equations.

Deformations for relative commutative associative algebras. The commutative algebra is by definition commutative rings in the relative context: an A -algebra is a commutative ring B over the base commutative ring A given by the structure map $A \rightarrow B$. This is the affine case of studying algebraic varieties with a fixed base variety which is the hallmark of Grothendieck's relative viewpoint on algebraic geometry, e.g. in the Grothendieck-Riemann-Roch Theorem [12, 17].

This paper develops deformations and their QCLs in the relative context. Thus starting with a relative pair of commutative rings, their deformations are expected to be a relative pair of noncommutative rings whose QCL is a relative pair of Poisson algebras. However, the natural analog of relative commutative associative algebras in terms of morphisms between a pair of objects is not working. Indeed a noncommutative associative algebra, with its base another noncommutative associative algebra, does not function well in the classical interpretation. For a noncommutative associative A -algebra, either the base algebra A is still assumed to be commutative or its action on the main algebra require strong conditions (see for example [16]). This obstacle becomes more prominent in studying other algebraic structures in the relative contexts.

This study addresses this difficulty via a representation theoretic approach, by regarding a relative pair of an algebraic structure as a triple of the algebraic structure, one for each of the main object, the base object and their semidirect product (see Definition 2.1). In this approach, the desired interconnection between the main object and the base object is naturally encoded in the requirement that the semidirect product has the same structure. This is similar in spirit to the Manin triple approach to Lie bialgebras [14] and several other bialgebra theories developed recently (e.g. [5, 34]).

For example, with this approach, a relative pair (R, A) of (noncommutative) associative algebras is for R to be the A -bimodule algebra, a term introduced in [9] though whose study can be tracked back to [29], instead of the usually considered associative ring R on which the action of A is commutative. Since a bimodule algebra (R, A) is a triple of associative algebra structures, on A , R and $A \oplus R$, the deformation of a bimodule algebra involves the deformations of these three associative algebras. As a consequence, the QCLs have a compatible triple of Poisson algebras, giving rise to the structure of relative pair of Poisson algebras, as expected (Theorem 2.15).

Scalar deformations of \mathcal{O} -operators. To deform a bimodule algebra with an \mathcal{O} -operator, it is possible to deform either the associative algebra, or its bimodule algebra, or the \mathcal{O} -operator. The approach that the \mathcal{O} -operator was deformed while the associative algebra and its bimodule algebra were undeformed was taken in [13], where cohomologies of \mathcal{O} -operators were studied. In this paper, we study the deformations of an associative algebra and its bimodule algebra, with a scalar deformation of the \mathcal{O} -operator still being an \mathcal{O} -operator. In this case, the \mathcal{O} -operator is also an \mathcal{O} -operator associated to the QCL module Poisson algebra (Proposition 2.21).

Deformations of commutative tridendriform algebras. It is usually difficult to determine whether the scalar deformation of an \mathcal{O} -operator is an \mathcal{O} -operator associated to a given bimodule algebra

deformation. Exploiting the equivalent characterization of a tridendriform algebra as a bimodule algebra obtained in [11], from any tridendriform deformation of a commutative tridendriform algebra whose QCL is a post-Poisson algebra [35], we show the scalar deformation of the identity operator is always an \mathcal{O} -operator associated to the induced bimodule algebra deformation (Proposition 3.16), illustrating the aforementioned bimodule algebra deformation-QCL process endowed with \mathcal{O} -operators explicitly.

On the other hand, [11] shows that any bimodule algebra together with an \mathcal{O} -operator induces a tridendriform algebra. Extending to the context of scalar deformations of \mathcal{O} -operators yields post-Poisson algebras (Proposition 3.15).

Scalar deformations for solutions of the AYBE. We finally study how the noncommutative deformations and their QCLs as Poisson algebras behave as the solution spaces of suitable classical Yang-Baxter type equations (AYBE and PYBE). Here we first take the direct approach of treating the solutions of the equations in tensor forms. We then take the operator approach by borrowing the previous results on \mathcal{O} -operators and the relations between the tensor forms and operator forms of the solutions for the AYBE and PYBE. We show that the solutions of the PYBE from the two approaches also agree.

In parallel, determining whether the scalar deformation of a solution of the AYBE is a solution of the AYBE in a given associative deformation is also challenging. However, on any tridendriform algebra, the identity operator is an \mathcal{O} -operator. So we consider tridendriform deformations and the induced scalar deformation of the \mathcal{O} -operator as well as the related scalar deformations of the solutions of the AYBE. Then utilizing the operator approach of the solutions of the AYBE, we finally obtain explicit solutions of the PYBE from any tridendriform deformation of a commutative tridendriform algebra.

1.4. Organization of the paper. With the main ideas above, the paper is outlined as follows.

In Section 2, we give the notion of deformation of a bimodule algebra and determine the corresponding QCL. Then we show that a bimodule algebra deformation with a scalar deformation of an \mathcal{O} -operator provides an \mathcal{O} -operator associated to the QCL.

In Section 3, we apply bimodule algebra deformations with scalar deformations of \mathcal{O} -operators to tridendriform deformations of commutative tridendriform algebras. We give an explicit deformation formula via derivations.

In Section 4, we apply bimodule algebra deformations with scalar deformations of \mathcal{O} -operators to establish the links between certain solutions of the AYBE and the PYBE, which are illustrated by the commutative diagrams.

Notations. Throughout this paper, we fix a ground field \mathbf{k} of characteristic 0 for vector spaces, algebras and tensor products. We also fix the power series ring $\mathbf{K} = \mathbf{k}[[\hbar]]$ as the ground algebra for topological modules, topological algebras and topological tensor products. For a vector space A with an operation \circ , the linear maps $L_\circ, R_\circ : A \rightarrow \text{End}_{\mathbf{k}}(A)$ are respectively defined by

$$L_\circ(x)y := x \circ y, \quad R_\circ(x)y := y \circ x, \quad x, y \in A.$$

Let \mathbb{N} and \mathbb{Z}^+ denote the set of natural numbers and the set of positive integers respectively.

2. BIMODULE ALGEBRA DEFORMATIONS AND \mathcal{O} -OPERATORS

This section introduces the notions of a bimodule algebra deformation and the corresponding quasiclassical limit (QCL). The scalar deformation of an \mathcal{O} -operator on a commutative associative algebra is also introduced to give an \mathcal{O} -operator on the Poisson algebra as the QCL.

2.1. Deformations of bimodule algebras and the corresponding quasiclassical limits. We introduce a general notion of bimodule algebras in order to provide a uniform framework for several structures considered in this paper.

Let \mathcal{P} be a binary quadratic operad [28]. For convenience, we also let \mathcal{P} denote the category of \mathcal{P} -algebras.

Definition 2.1. Let \mathcal{P} be a binary quadratic operad with binary operations $\{\mu_i\}_{i \in I}$. Let $(A, \{\circ_i\}_{i \in I})$ and $(V, \{\cdot_i\}_{i \in I})$ be \mathcal{P} -algebras. Let $l_i, r_i : A \rightarrow \text{End}_{\mathbf{k}}(V), i \in I$ be linear maps. If the vector space direct sum $A \oplus V$ is again a \mathcal{P} -algebra with the multiplications \odot_i defined by

$$(x, u) \odot_i (y, v) = (x \circ_i y, l_i(x)v + r_i(y)u + u \cdot_i v), \quad x, y \in A, u, v \in V, i \in I, \quad (1)$$

then the quadruple $(V, \{\cdot_i\}, \{l_i\}_i, \{r_i\}_i)$ is called an **A -bimodule \mathcal{P} -algebra** and the \mathcal{P} -algebra $(A \oplus V, \{\odot_i\})$ is denoted by $A \rtimes_{l_i, r_i} V$. If the operations μ_i are either symmetric (meaning commutative) or antisymmetric, implying $r_i = \pm l_i, i \in I$, then the triple $(V, \{\cdot_i\}, \{l_i\}_i)$ is called an **A -module \mathcal{P} -algebra**. The prefixes A - or \mathcal{P} - will be suppressed if the context is clear.

If V is just a vector space, equip it with a \mathcal{P} -algebra structure by taking \cdot_i to be the trivial operation. Then the resulting A -bimodule algebra $(V, \{\cdot_i\}, \{l_i\}, \{r_i\})$ is called an **A -bimodule** and is simply denoted by $(V, \{l_i\}, \{r_i\})$. In the case when each \circ_i is either symmetric or antisymmetric, then V is also called an **A -module**. The resulting \mathcal{P} -algebra $(A \oplus V, \{\odot_i\})$ is still denoted by $A \rtimes_{l_i, r_i} V$.

When \mathcal{P} is the operad of associative algebras, we recover the notion of a bimodule algebra in [9], originally formulated by relations among the binary operations and maps as follows. See also [32] for a similar construction.

Proposition 2.2. [9] *Let (A, \circ) and (V, \cdot) be associative algebras. Let $l, r : A \rightarrow \text{End}_{\mathbf{k}}(V)$ be linear maps. Then (V, \cdot, l, r) is an A -bimodule algebra if and only if the following conditions are satisfied.*

$$l(x \circ y) = l(x)l(y), \quad l(x)r(y) = r(y)l(x), \quad r(x \circ y) = r(y)r(x), \quad (2)$$

$$l(x)(u \cdot v) = (l(x)u) \cdot v, \quad (r(x)u) \cdot v = u \cdot (l(x)v), \quad r(x)(u \cdot v) = u \cdot (r(x)v), \quad x, y \in A, u, v \in V. \quad (3)$$

Let A be an associative algebra and let V be a vector space. Then the resulting A -bimodule algebra (V, \cdot, l, r) coincides with the usual **A -bimodule** defined by Eq. (2).

Now apply the general formulation to the operad \mathcal{P} of commutative associative algebras. As noted above, $l(x) = r(x) := \rho_{\circ}(x)$ for $x \in A$. Then (2) and (3) reduce to

$$\rho_{\circ}(x \circ y) = \rho_{\circ}(x)\rho_{\circ}(y) \quad \text{and} \quad \rho_{\circ}(x)(u \cdot v) = (\rho_{\circ}(x)u) \cdot v, \quad x, y \in A, u, v \in V, \quad (4)$$

respectively. This means that an **A -module algebra** is simply a commutative A -algebra. For notational consistency in the paper, we will use the former term as in [34].

For the operad of Lie algebras, the notion of module \mathcal{P} -algebras will be detailed in Section 4.3 (see the paragraph below Theorem 4.16). At this moment, for the sake of later references, we fix some notations when \mathcal{P} is the operad of Poisson algebras. Recall that a **Poisson algebra** $(P, \{\cdot, \cdot\}, \circ)$ consists of a Lie algebra $(P, \{\cdot, \cdot\})$ and a commutative associative algebra (P, \circ) such that

$$\{x, y \circ z\} = \{x, y\} \circ z + y \circ \{x, z\}, \quad x, y, z \in P. \quad (5)$$

Definition 2.3. [34] Let $(P, \{\cdot, \cdot\}, \circ)$ and $(Q, [\cdot, \cdot], \cdot)$ be Poisson algebras, and let $\rho_{\{\cdot, \cdot\}}, \rho_{\circ} : P \rightarrow \text{End}_{\mathbf{k}}(Q)$ be linear maps. If the direct sum of vector spaces $P \oplus Q$ is turned into a Poisson algebra with the bracket $[[\cdot, \cdot]]$ and the multiplication \odot respectively defined by

$$[[(x, u), (y, v)]] := (\{x, y\}, \rho_{\{\cdot, \cdot\}}(x)v - \rho_{\{\cdot, \cdot\}}(y)u + [u, v]), \quad (6)$$

$$(x, u) \odot (y, v) := (x \circ y, \rho_{\circ}(x)v + \rho_{\circ}(y)u + u \cdot v), \quad x, y \in P, u, v \in Q, \quad (7)$$

then we call $(Q, [,], \cdot, \rho_{(\cdot)}, \rho_{\circ})$ a **P -module Poisson algebra** and denote the resulting Poisson algebra $(P \oplus Q, \llbracket, \rrbracket, \odot)$ by $P \ltimes_{\rho_{(\cdot)}, \rho_{\circ}} Q$.

The same notion of **P -module** is used if Q is only a vector space but equipped with the trivial operations as in the general construction in Definition 2.1.

Remark 2.4. There is also an equivalent characterization of a P -module Poisson algebra as well as a P -module in [34] in terms of a system of equations similar to (2) and (3). We will not list them since they are not needed in the rest of this paper.

We next recall the needed notations for (formal) deformations as follows [15]. Let $\mathbf{K} = \mathbf{k}[[h]]$ be the formal power series algebra and let V be a vector space. Consider the set of formal series

$$V_h := V[[h]] := \left\{ \sum_{i=0}^{\infty} v_i h^i \mid v_i \in V \right\}.$$

For $v' = \sum_{i=0}^{\infty} v_i h^i, w' = \sum_{i=0}^{\infty} w_i h^i \in V[[h]]$ and $\lambda' = \sum_{i=0}^{\infty} \lambda_i h^i \in \mathbf{K}$, set

$$v' + w' := \sum_{i=0}^{\infty} (v_i + w_i) h^i, \quad \lambda' v' := \sum_{i=0}^{\infty} \left(\sum_{m+n=i} \lambda_m v_n \right) h^i.$$

Then $V[[h]]$ becomes a \mathbf{K} -module.

For a fixed real number $C > 1$, define the **h -adic norm** $\| \cdot \|$ on $V[[h]]$ by

$$\left\| \sum_{i=0}^{\infty} v_i h^i \right\| := C^{-m},$$

where m is the smallest integer such that $v_m \neq 0$. The topology induced by this norm is called the **h -adic topology**. A \mathbf{K} -module that is \mathbf{K} -linearly isomorphic to $V[[h]]$ is called a **topologically free \mathbf{K} -module**. In particular, V_h is a topologically free \mathbf{K} -module.

Let A_h be a topologically free \mathbf{K} -module. If there is a \mathbf{K} -bilinear operation \cdot_h on A_h such that

$$(x \cdot_h y) \cdot_h z = x \cdot_h (y \cdot_h z), \quad x, y, z \in A_h,$$

then (A_h, \cdot_h) is called a **topologically free associative algebra**.

Definition 2.5. A **deformation of an associative algebra** (A, \cdot) is a topologically free associative algebra (A_h, \cdot_h) such that

- (a) $A_h = A[[h]]$ as \mathbf{K} -modules;
- (b) $x \cdot_h y \equiv x \cdot y \pmod{h}$ for $x, y \in A$.

Remark 2.6. (a) In [15], Conditions (a) and (b) in the above definition are presented equivalently as the condition $A = A_h/hA_h$.

- (b) To describe the associativity for a topologically free associative algebra, we need the notion of a **topological tensor product**. Let $V[[h]]$ and $W[[h]]$ be any topologically free \mathbf{K} -modules. The topological tensor product $V[[h]] \hat{\otimes} W[[h]]$ is the completion of $V[[h]] \otimes_{\mathbf{K}} W[[h]]$ under the h -adic norm. So the topological tensor product is the completion of the \mathbf{K} -bilinear extension of the usual (algebraic) tensor product.

A deformation (A_h, \cdot_h) of an associative algebra (A, \cdot) is also called an **associative algebra deformation** or simply an **associative deformation** of (A, \cdot) .

Definition 2.7. Let (A_h, \cdot_h) be an associative deformation of a commutative associative algebra (A, \cdot) . Let $[,]$ be an operation on A such that

$$[x, y] \equiv \frac{x \cdot_h y - y \cdot_h x}{h} \pmod{h}, \quad x, y \in A.$$

Then $(A, [,], \cdot)$ is called the **quasiclassical limit (QCL) of the associative deformation** (A_h, \cdot_h) .

The following result is well known.

Theorem 2.8. [15] *Let (A_h, \cdot_h) be an associative deformation of a commutative associative algebra (A, \cdot) . Then its QCL $(A, [\cdot], \cdot)$ is a Poisson algebra.*

We next extend the associative deformations to bimodule algebras.

Definition 2.9. Let (A_h, \circ_h) and (V_h, \cdot_h) be topologically free associative algebras. Let $l_h, r_h : A_h \rightarrow \text{End}_{\mathbf{K}}(V_h)$ be \mathbf{K} -linear maps. If (V_h, \cdot_h, l_h, r_h) is an A_h -bimodule algebra, then we call (V_h, \cdot_h, l_h, r_h) a **topologically free A_h -bimodule algebra** and denote the topologically free associative algebra structure on $A_h \oplus V_h$ by $A_h \ltimes_{l_h, r_h} V_h$.

Let (A, \circ) be an associative algebra and (V, \cdot, l, r) be an A -bimodule algebra. Let (A_h, \circ_h) be a topologically free associative algebra and (V_h, \cdot_h, l_h, r_h) be a topologically free A_h -bimodule algebra. If $A_h \ltimes_{l_h, r_h} V_h$ is an associative deformation of $A \ltimes_{l, r} V$, then we call (V_h, \cdot_h, l_h, r_h) an **A_h -bimodule algebra deformation of the A -bimodule algebra (V, \cdot, l, r)** .

Remark 2.10. Let $\iota : A_h \oplus V_h \rightarrow (A \oplus V)_h$ be the map defined by

$$\iota \left(\sum_{i=0}^{\infty} x_i h^i, \sum_{i=0}^{\infty} v_i h^i \right) := \sum_{i=0}^{\infty} (x_i, v_i) h^i, \quad i \in \mathbb{N}, x_i \in A, v_i \in V.$$

Then ι is a \mathbf{K} -linear isomorphism. With $A_h \oplus V_h$ identified with $(A \oplus V)_h$ this way, it makes sense to say that $A_h \ltimes_{l_h, r_h} V_h$ is an associative deformation of $A \ltimes_{l, r} V$.

Proposition 2.11. *Let (A, \circ) be an associative algebra and (V, \cdot, l, r) be an A -bimodule algebra. Let (A_h, \circ_h) be a topologically free associative algebra and (V_h, \cdot_h, l_h, r_h) be a topologically free A_h -bimodule algebra. Then (V_h, \cdot_h, l_h, r_h) is an A_h -bimodule algebra deformation of the A -bimodule algebra (V, \cdot, l, r) if and only if the following congruences hold:*

$$x \circ_h y \equiv x \circ y \pmod{h}, \quad u \cdot_h v \equiv u \cdot v \pmod{h}, \quad (8)$$

$$l_h(x)v \equiv l(x)v \pmod{h}, \quad r_h(y)u \equiv r(y)u \pmod{h}, \quad x, y \in A, u, v \in V. \quad (9)$$

Proof. Let \odot_h denote the associative product of the topologically free associative algebra $A_h \ltimes_{l_h, r_h} V_h$. Then $A_h \ltimes_{l_h, r_h} V_h$ is an associative deformation of the associative algebra $A \ltimes_{l, r} V$ if and only if

$$(x, u) \odot_h (y, v) \equiv (x, u) \odot (y, v) \pmod{h}, \quad x, y \in A, u, v \in V.$$

This holds if and only if

$$(x \circ_h y, l_h(x)v + r_h(y)u + u \cdot_h v) \equiv (x \circ y, l(x)v + r(y)u + u \cdot v) \pmod{h}, \quad x, y \in A, u, v \in V. \quad (10)$$

By suitable choice of x, y, u and v , (10) holds if and only if (8) and (9) hold, yielding the conclusion. \square

Remark 2.12. The condition (8) holds if and only if (A_h, \circ_h) is an associative deformation of (A, \circ) and (V_h, \cdot_h) is an associative deformation of (V, \cdot) .

Definition 2.13. Let (A, \circ) be a commutative associative algebra and (V, \cdot, ρ_\circ) be an A -module algebra. Let (A_h, \circ_h) and (V_h, \cdot_h) be the associative deformations of (A, \circ) and (V, \cdot) respectively, and $(A, \{\cdot\}, \circ)$ and $(V, [\cdot], \cdot)$ be the QCL. Let (V_h, \cdot_h, l_h, r_h) be an A_h -bimodule algebra deformation of the A -module algebra (V, \cdot, ρ_\circ) . Let $\rho_{\{\cdot\}} : A \rightarrow \text{End}_{\mathbf{K}}(V)$ be a linear map satisfying

$$\rho_{\{\cdot\}}(x)(u) \equiv \frac{l_h(x)u - r_h(x)u}{h} \pmod{h}, \quad x \in A, u \in V. \quad (11)$$

We call $(V, [\cdot], \cdot, \rho_{\{\cdot\}}, \rho_\circ)$ the **quasiclassical limit (QCL) of the A_h -bimodule algebra deformation (V_h, \cdot_h, l_h, r_h)** .

- Remark 2.14.** (a) The fraction $\frac{l_h(x)u - r_h(x)u}{h}$ is a well-defined element in V_h for each $x \in A$ and $u \in V$, since $l_h \equiv r_h \pmod{h}$. Moreover, $\rho_{\{\cdot\}}(x)(u)$ is the zeroth order term of $\frac{l_h(x)u - r_h(x)u}{h}$ and the linear map $\rho_{\{\cdot\}}$ in (11) is well defined.
- (b) In the same way or by taking \cdot and \cdot_h being null operations in the above definitions, we can give the notions of a **topologically free bimodule**, an A_h -**bimodule deformation** and the **quasiclassical limit (QCL) of the A_h -bimodule deformation**.

Theorem 2.15. *Let (A, \circ) be a commutative associative algebra, (A_h, \circ_h) be an associative deformation of (A, \circ) and $(A, \{\cdot\}, \circ)$ be the Poisson algebra as the QCL. Let (V, \cdot, ρ_\circ) be an A -module algebra and (V_h, \cdot_h, l_h, r_h) be an A_h -bimodule algebra deformation. Then the corresponding QCL $(V, [\cdot, \cdot], \cdot, \rho_{\{\cdot\}}, \rho_\circ)$ is an $(A, \{\cdot\}, \circ)$ -module Poisson algebra. Moreover, the QCL of the associative deformation $A_h \ltimes_{l_h, r_h} V_h$ of the commutative associative algebra $A \ltimes_{\rho_\circ} V$ is exactly the Poisson algebra $A \ltimes_{\rho_{\{\cdot\}}, \rho_\circ} V$.*

Proof. As the QCL of (V_h, \cdot_h) , $(V, [\cdot, \cdot], \cdot)$ is a Poisson algebra. Define operations $[[\cdot, \cdot]]$ and \odot on the direct sum of vector spaces $A \oplus V$ by (6) and (7) respectively, that is,

$$\begin{aligned} [[(x, u), (y, v)]] &=: (\{x, y\}, \rho_{\{\cdot\}}(x)v - \rho_{\{\cdot\}}(y)u + [u, v]), \\ (x, u) \odot (y, v) &=: (x \circ y, \rho_\circ(x)v + \rho_\circ(y)u + u \cdot v), \quad x, y \in A, u, v \in V. \end{aligned}$$

Then $(A \oplus V, \odot)$ is a commutative associative algebra. Let $(A_h \oplus V_h, \odot_h) := A_h \ltimes_{l_h, r_h} V_h$ be the associative deformation. For $x, y \in A$ and $u, v \in V$, we have

$$\begin{aligned} (x, u) \odot_h (y, v) - (y, v) \odot_h (x, u) &= (x \circ_h y - y \circ_h x, l_h(x)v - r_h(x)v - l_h(y)u + r_h(y)u + u \cdot_h v - v \cdot_h u) \\ &\equiv (\{x, y\}, \rho_{\{\cdot\}}(x)v - \rho_{\{\cdot\}}(y)u + [u, v])h \pmod{h^2}. \end{aligned}$$

Then $(A \oplus V, [[\cdot, \cdot]], \odot)$ is the QCL of the associative deformation $(A_h \oplus V_h, \odot_h)$. Therefore, $(A \oplus V, [[\cdot, \cdot]], \odot)$ is a Poisson algebra and hence $(V, [\cdot, \cdot], \cdot, \rho_{\{\cdot\}}, \rho_\circ)$ is an $(A, \{\cdot\}, \circ)$ -module Poisson algebra by Definition 2.3. The last statement holds since the two Poisson algebras $A \ltimes_{\rho_{\{\cdot\}}, \rho_\circ} V$ and $(A \oplus V, [[\cdot, \cdot]], \odot)$ coincide. \square

Regarding a bimodule as a bimodule algebra where the multiplication is trivial, we have the following consequence of Theorem 2.15.

Corollary 2.16. *Let (A, \circ) be a commutative associative algebra and (V, ρ_\circ) be an A -module. Let (A_h, \circ_h) be an associative deformation of (A, \circ) and $(A, \{\cdot\}, \circ)$ be the QCL which is a Poisson algebra. Let (V_h, l_h, r_h) be an A_h -bimodule deformation of (V, ρ_\circ) . Then the corresponding QCL $(V, \rho_{\{\cdot\}}, \rho_\circ)$ is an $(A, \{\cdot\}, \circ)$ -module. Accordingly, the associative deformation $A_h \ltimes_{l_h, r_h} V_h$ of the commutative associative algebra $A \ltimes_{\rho_\circ} V$ is exactly the Poisson algebra $A \ltimes_{\rho_{\{\cdot\}}, \rho_\circ} V$.*

2.2. Θ -operators in bimodule algebra deformations. We give a unified notion of Θ -operators on various algebras.

Definition 2.17. Let \mathcal{P} be a binary quadratic operad with binary operations $\{\mu_i\}_{i \in I}$. Let $(A, \{\circ_i\}_{i \in I})$ and $(V, \{\cdot_i\}_{i \in I})$ be \mathcal{P} -algebras. Let $l_i, r_i : A \rightarrow \text{End}_k(V)$, $i \in I$, be linear maps such that the quadruple $(V, \{\cdot_i\}, \{l_i\}, \{r_i\})$ is an A -bimodule \mathcal{P} -algebra as given in Definition 2.1. A linear map $T : V \rightarrow A$ is called an Θ -operator of weight λ on the \mathcal{P} -algebra (A, \circ) associated to the A -bimodule \mathcal{P} -algebra (V, \cdot, l, r) if T satisfies

$$T(u) \circ_i T(v) = T(l_i(T(u))v) + T(r_i(T(v))u) + \lambda T(u \cdot_i v), \quad u, v \in V. \quad (12)$$

When an A -bimodule (V, l_i, r_i) is regarded as an A -bimodule \mathcal{P} -algebra equipped with the trivial multiplications, the weight is irrelevant, and T is called an Θ -operator on the \mathcal{P} -algebra $(A, \{\circ_i\}_{i \in I})$ associated to the A -bimodule $(V, \{l_i\}, \{r_i\})$.

When \mathcal{P} is the operad of associative algebras, that is, when (A, \circ) is an associative algebra and (V, \cdot, l, r) is an A -bimodule algebra, we recover the notion of an \mathcal{O} -operator of weight λ on (A, \circ) associated to (V, \cdot, l, r) [11].

Remark 2.18. For an associative algebra (A, \circ) , the regular representation gives an A -bimodule algebra $(A, \circ, L_\circ, R_\circ)$. Then an \mathcal{O} -operator of weight λ on the associative algebra (A, \circ) associated to this A -bimodule algebra gives the notion of a **Rota-Baxter operator** of the same weight, defined by

$$T(x) \circ T(y) = T(T(x) \circ y) + T(x \circ T(y)) + \lambda T(x \circ y), \quad x, y \in A. \quad (13)$$

Thus an \mathcal{O} -operator of weight λ is also called a **relative Rota-Baxter operator of weight λ** .

For later reference of notations, we recall the details when \mathcal{P} is the operad of Poisson algebras.

Definition 2.19. [34] Let $(P, \{, \}, \circ)$ be a Poisson algebra, $(Q, [,], \cdot, \rho_{\{, \}}, \rho_\circ)$ be a P -module Poisson algebra and $\lambda \in \mathbf{k}$. A linear map $T : Q \rightarrow P$ is called an **\mathcal{O} -operator of weight λ on $(P, \{, \}, \circ)$ associated to $(Q, [,], \cdot, \rho_{\{, \}}, \rho_\circ)$** if T satisfies

$$\{T(u), T(v)\} = T(\rho_{\{, \}}(T(u))v) - T(\rho_{\{, \}}(T(v))u) + \lambda T([u, v]) \quad (14)$$

$$T(u) \circ T(v) = T(\rho_\circ(T(u))v) + T(\rho_\circ(T(v))u) + \lambda T(u \cdot v), \quad u, v \in Q. \quad (15)$$

In particular, when the operations $[,]$ and \cdot on Q are trivial, that is, $(Q, \rho_{\{, \}}, \rho_\circ)$ is a P -module, T is simply called an **\mathcal{O} -operator on $(P, \{, \}, \circ)$ associated to $(Q, \rho_{\{, \}}, \rho_\circ)$** .

Remark 2.20. Similar to the case of associative algebras, a Rota-Baxter operator T of weight λ on a Poisson algebra $(P, \{, \}, \circ)$ is defined by the following equations:

$$\{T(x), T(y)\} = T(\{T(x), y\}) + T(\{x, T(y)\}) + \lambda T(\{x, y\}), \quad (16)$$

$$T(x) \circ T(y) = T(T(x) \circ y) + T(x \circ T(y)) + \lambda T(x \circ y), \quad x, y \in P. \quad (17)$$

Let (A, \circ) be an associative algebra and (V, \cdot, l, r) be an A -bimodule algebra. Let (A_h, \circ_h) be an associative deformation of (A, \circ) and (V_h, \cdot_h, l_h, r_h) be an A_h -bimodule algebra deformation of (V, \cdot, l, r) . For $T \in \text{Hom}_{\mathbf{k}}(V, A)$, define the **scalar deformation** of T to be the \mathbf{K} -linear extension

$$T_{\mathbf{K}} : V_h \rightarrow A_h, \quad T_{\mathbf{K}}\left(\sum_{i=0}^{\infty} v_i h^i\right) := \sum_{i=0}^{\infty} T(v_i) h^i, \quad \sum_{i=0}^{\infty} v_i h^i \in V_h. \quad (18)$$

Then $T_{\mathbf{K}}$ is an \mathcal{O} -operator of weight λ on (A_h, \circ_h) associated to (V_h, \cdot_h, l_h, r_h) means that

$$T_{\mathbf{K}}(u) \circ_h T_{\mathbf{K}}(v) = T_{\mathbf{K}}(l_h(T_{\mathbf{K}}(u))v) + T_{\mathbf{K}}(r_h(T_{\mathbf{K}}(v))u) + \lambda T_{\mathbf{K}}(u \cdot_h v), \quad u, v \in V. \quad (19)$$

For $x, y \in A$ and $u \in V$, consider the expansions

$$x \circ_h y = \sum_{s=0}^{\infty} (x \circ_s y) h^s, \quad l_h(x)u = \sum_{s=0}^{\infty} l_s(x)u h^s \quad \text{and} \quad r_h(x)u = \sum_{s=0}^{\infty} r_s(x)u h^s,$$

with $x \circ_s y \in A$ and $l_s(x), r_s(x) \in \text{End}_{\mathbf{k}}(V)$. Then (19) holds if and only if

$$T(u) \circ_s T(v) = T(l_s(T(u))v) + T(r_s(T(v))u) + \lambda T(u \cdot_s v), \quad u, v \in V, s \geq 0. \quad (20)$$

When (V, \cdot, l, r) is the A -bimodule algebra $(A, \circ, L_\circ, R_\circ)$, we replace \cdot_s by \circ_s and (20) becomes

$$T(x) \circ_s T(y) = T(T(x) \circ_s y) + T(x \circ_s T(y)) + \lambda T(x \circ_s y), \quad x, y \in A, s \geq 0. \quad (21)$$

Theorem 2.21. *Let (A, \circ) be a commutative associative algebra and (V, \cdot, ρ_\circ) be an A -module algebra. Let (A_h, \circ_h) be an associative deformation of (A, \circ) and (V_h, \cdot_h, l_h, r_h) be an A_h -bimodule algebra deformation of (V, \cdot, ρ_\circ) . Suppose that $(A, \{, \}, \circ)$, $(V, [,], \cdot)$ and $(V, [,], \cdot, \rho_{\{, \}}, \rho_\circ)$ are the QCLs of (A_h, \circ_h) , (V_h, \cdot_h) and (V_h, \cdot_h, l_h, r_h) respectively. Let $T \in \text{Hom}_{\mathbf{k}}(V, A)$ be an \mathcal{O} -operator of weight λ on (A, \circ) associated to (V, \cdot, ρ_\circ) . If $T_{\mathbf{K}}$ defined by (18) is an \mathcal{O} -operator of weight λ on (A_h, \circ_h) associated to (V_h, \cdot_h, l_h, r_h) , then T is an \mathcal{O} -operator of weight λ on the Poisson algebra $(A, \{, \}, \circ)$ associated to the $(A, \{, \}, \circ)$ -module Poisson algebra $(V, [,], \cdot, \rho_{\{, \}}, \rho_\circ)$.*

Proof. Since T is an \mathcal{O} -operator of weight λ on (A, \circ) associated to (V, \cdot, ρ_\circ) , it is direct to see that (15) is satisfied. Applying (20) and exchanging u and v , we obtain

$$T(v) \circ_s T(u) = T(l_s(T(v))u) + T(r_s(T(u))v) + \lambda T(v \cdot_s u), \quad u, v \in V, s \geq 0. \quad (22)$$

Taking $s = 1$ and subtracting (22) from (20), we deduce that the equation

$$T(u) \circ_1 T(v) - T(v) \circ_1 T(u) = T((l_1 - r_1)(T(u))v) - T((l_1 - r_1)(T(v))u) + \lambda T(u \cdot_1 v - u \cdot_1 v), \quad u, v \in V,$$

holds if and only if (14) holds. Thus T is an \mathcal{O} -operator of weight λ on $(A, \{\cdot, \cdot\}, \circ)$ associated to $(V, [\cdot, \cdot], \cdot, \rho_{\{\cdot, \cdot\}}, \rho_\circ)$. \square

3. BIMODULE ALGEBRA PERSPECTIVE OF TRIDENDRIFORM DEFORMATIONS

In this section, we study tridendriform deformations of commutative tridendriform algebras in terms of bimodule algebra deformations. This provides an application of Theorem 2.21 where the deformations and QCLs with \mathcal{O} -operators can be constructed explicitly. We also give constructions of bimodule algebra deformations and tridendriform deformations via derivations

3.1. Deformations of tridendriform algebras and the corresponding QCLs.

Definition 3.1. [27] A **tridendriform algebra** (A, \succ, \prec, \cdot) consists of a vector space A with three binary operations \succ, \prec and \cdot such that

$$(x \prec y) \prec z = x \prec (y \circ z), \quad (x \succ y) \prec z = x \succ (y \prec z), \quad (x \circ y) \succ z = x \succ (y \succ z), \quad (23)$$

$$(x \succ y) \cdot z = x \succ (y \cdot z), \quad (x \prec y) \cdot z = x \cdot (y \succ z), \quad (x \cdot y) \prec z = x \cdot (y \prec z), \quad (24)$$

$$(x \cdot y) \cdot z = x \cdot (y \cdot z), \quad x, y, z \in A, \quad (25)$$

where

$$x \circ y := x \prec y + x \succ y + x \cdot y, \quad x, y \in A. \quad (26)$$

If in addition, $x \succ y = y \prec x$ and $x \cdot y = y \cdot x$ for $x, y \in A$, then we call the tridendriform algebra **commutative** and denote it by (A, \succ, \cdot) .

Remark 3.2. We recall that a **dendriform algebra** (A, \succ, \prec) consists of a vector space A with binary operations \succ and \prec such that the identities in (23) are satisfied for $x, y, z \in A$, where $x \circ y := x \prec y + x \succ y$. Moreover, if $x \succ y = y \prec x$ for $x, y \in A$, we call it a **Zinbiel algebra** and denote it by (A, \succ) . So we may regard a dendriform algebra (A, \succ, \prec) as a tridendriform algebra (A, \succ, \prec, \cdot) in which the operation \cdot is zero.

There is a bimodule algebra characterization of a tridendriform algebra.

Proposition 3.3. [11] *Let \succ, \prec and \cdot be binary operations on a vector space A . Define a binary operation \circ on A by (26). Then (A, \succ, \prec, \cdot) is a tridendriform algebra if and only if (A, \circ) is an associative algebra and $(A, \cdot, L_\succ, R_\prec)$ is an (A, \circ) -bimodule algebra. In this case, id_A is an \mathcal{O} -operator of weight 1 on (A, \circ) associated to $(A, \cdot, L_\succ, R_\prec)$. Moreover, (A, \succ, \cdot) is a commutative tridendriform algebra if and only if (A, \circ) is a commutative associative algebra and (A, \cdot, L_\succ) is an (A, \circ) -module algebra.*

The notion of post-Lie algebras arose from the study of operads [39] with applications to numerical analysis [31] and specializes to **pre-Lie algebras** [6] when the Lie algebras are abelian.

Definition 3.4. [39] A **post-Lie algebra** $(\mathfrak{g}, [\cdot, \cdot], \triangleright)$ consists of a Lie algebra $(\mathfrak{g}, [\cdot, \cdot])$ and a binary operation \triangleright such that

$$x \triangleright [y, z] = [x \triangleright y, z] + [y, x \triangleright z], \quad (27)$$

$$[x, y] \triangleright z = x \triangleright (y \triangleright z) - (x \triangleright y) \triangleright z - y \triangleright (x \triangleright z) + (y \triangleright x) \triangleright z, \quad x, y, z \in \mathfrak{g}. \quad (28)$$

Definition 3.5. [7] A **post-Poisson algebra** $(P, [,], \triangleright, \succ, \cdot)$ consists of a post-Lie algebra $(P, [,], \triangleright)$ and a commutative tridendriform algebra (P, \succ, \cdot) such that (5) and the following equations hold.

$$[x, y \succ z] = y \succ [x, z] - z \cdot (y \triangleright x), \quad x \triangleright (y \cdot z) = (x \triangleright y) \cdot z + y \cdot (x \triangleright z), \quad (29)$$

$$(x \circ y) \triangleright z = x \succ (y \triangleright z) + y \triangleright (z \prec x), \quad x \triangleright (y \succ z) = y \succ (x \triangleright z) + \{x, y\} \succ z, \quad (30)$$

for $x, y, z \in P$, where

$$x \circ y := x \succ y + y \succ x + x \cdot y, \quad \{x, y\} := x \triangleright y - y \triangleright x + [x, y], \quad x, y \in P. \quad (31)$$

Remark 3.6. Introduced by Aguiar [2], a **pre-Poisson algebra** $(A, \triangleright, \succ)$ consists of a Zinbiel algebra (A, \succ) and a pre-Lie algebra (A, \triangleright) such that identities in (30) are satisfied for $x, y, z \in A$, where $\{x, y\} := x \triangleright y - y \triangleright x$ and $x \circ y := x \succ y + y \succ x$. So a pre-Poisson algebra is a post-Poisson algebra where the operations \cdot and $[,]$ are trivial.

Similar to Proposition 3.3, the following statement holds.

Proposition 3.7. [34] Let $\triangleright, [,], \succ$ and \cdot be four binary operations on a vector space P . Define binary operations \circ and $\{, \}$ on P by (31). Then $(P, \triangleright, [,], \succ, \cdot)$ is a post-Poisson algebra if and only if $(P, \{, \}, \circ)$ is a Poisson algebra and $(P, [,], \cdot, L_{\triangleright}, L_{\succ})$ is a $(P, \{, \}, \circ)$ -module Poisson algebra. Moreover, in this case, id_P is an \mathcal{O} -operator of weight 1 on $(P, \{, \}, \circ)$ associated to $(P, [,], \cdot, L_{\triangleright}, L_{\succ})$.

Definition 3.8. A **topologically free tridendriform algebra** $(A_h, \succ_h, \prec_h, \cdot_h)$ consists of a topologically free \mathbf{K} -module A_h and \mathbf{K} -bilinear operations \succ_h, \prec_h and \cdot_h such that $(A_h, \succ_h, \prec_h, \cdot_h)$ is a tridendriform algebra.

A **deformation of a tridendriform algebra** (A, \succ, \prec, \cdot) is a topologically free tridendriform algebra $(A_h, \succ_h, \prec_h, \cdot_h)$ such that

- (a) $A_h = A[[h]]$ as topologically free \mathbf{K} -modules;
- (b) For $x, y \in A$, we have

$$x \succ_h y \equiv x \succ y \pmod{h}, \quad x \prec_h y \equiv x \prec y \pmod{h}, \quad x \cdot_h y \equiv x \cdot y \pmod{h}. \quad (32)$$

We also call $(A_h, \succ_h, \prec_h, \cdot_h)$ a **tridendriform algebra deformation** or simply a **tridendriform deformation** of (A, \succ, \prec, \cdot) .

Let $(A_h, \succ_h, \prec_h, \cdot_h)$ be a tridendriform deformation of (A, \succ, \prec, \cdot) . For each $*_h \in \{\succ_h, \prec_h, \cdot_h, \circ_h\}$, we can write

$$x *_h y = \sum_{s=0}^{\infty} (x *_s y) h^s, \quad x, y \in A, \quad (33)$$

where $x *_s y \in A$. By (32), we have $x *_0 y = x *_s y$ for $x, y \in A$.

Definition 3.9. Let $(A_h, \succ_h, \prec_h, \cdot_h)$ be a tridendriform deformation of a commutative tridendriform algebra (A, \succ, \cdot) . Let $[,]$ and \triangleright be the binary operations on A such that

$$[x, y] \equiv \frac{x \cdot_h y - y \cdot_h x}{h} \pmod{h}, \quad x \triangleright y \equiv \frac{x \succ_h y - y \prec_h x}{h} \pmod{h}, \quad x, y \in A. \quad (34)$$

Then we call $(A, [,], \triangleright, \succ, \cdot)$ the **quasiclassical limit (QCL) of the tridendriform deformation** $(A_h, \succ_h, \prec_h, \cdot_h)$.

Remark 3.10. By setting $\cdot = \cdot_h = 0$ in the above definitions, one can define the notions of a **topologically free dendriform algebra**, a **dendriform deformation** and the **quasiclassical limit of a dendriform deformation**.

For the QCL $(A, [,], \triangleright, \succ, \cdot)$, first we note that, as the QCL of the associative deformation (A_h, \cdot_h) , the triple $(A, [,], \cdot)$ is a Poisson algebra. Define a binary operation on A_h by

$$x_h \circ_h y_h := x_h \succ_h y_h + x_h \prec_h y_h + x_h \cdot_h y_h, \quad x_h, y_h \in A_h. \quad (35)$$

Next, with the binary operation $\{, \}$ on A defined by

$$\{x, y\} \equiv \frac{x \circ_h y - y \circ_h x}{h} \pmod{h}, \quad x, y \in A, \quad (36)$$

and \circ defined by (26), the triple $(A, \{, \}, \circ)$ is the QCL of the associative deformation (A_h, \circ_h) and hence is also a Poisson algebra. Let $x, y \in A$. Using the expansion in (33), we have

$$x \triangleright y = x \succ_1 y - y \prec_1 x, \quad [x, y] = x \cdot_1 y - y \cdot_1 x, \quad \{x, y\} = x \circ_1 y - y \circ_1 x.$$

It is direct to see that

$$\{x, y\} = x \triangleright y - y \triangleright x + [x, y]. \quad (37)$$

Hence the operation $\{, \}$ also splits into three operations.

Theorem 3.11. [35] *Let $(A_h, \succ_h, \prec_h, \cdot_h)$ be a tridendriform deformation of a commutative tridendriform algebra (A, \succ, \cdot) . Then the corresponding QCL $(A, [,], \triangleright, \succ, \cdot)$ is a post-Poisson algebra.*

By taking \cdot and \cdot_h in Theorem 3.11 as trivial operations, we obtain

Corollary 3.12. [2] *Let (A_h, \succ_h, \prec_h) be a dendriform deformation of a Zinbiel algebra (A, \succ) . Then the corresponding QCL $(A, \triangleright, \succ)$ is a pre-Poisson algebra.*

We give the following example of tridendriform deformations constructed from an arbitrary associative deformation.

Example 3.13. Let (A^n, \cdot) be the direct product of n copies of an associative algebra (A, \diamond) , equipped with the product

$$a \cdot b = (a_1 \diamond b_1, \dots, a_i \diamond b_i, \dots, a_n \diamond b_n), \quad a = (a_1, \dots, a_n), b = (b_1, \dots, b_n) \in A^n. \quad (38)$$

Set

$$a \succ b := (0, a_1 \diamond b_2, \dots, \sum_{j=1}^{n-1} a_j \diamond b_n), \quad a \prec b := (0, a_2 \diamond b_1, \dots, \sum_{j=1}^{n-1} a_n \diamond b_j). \quad (39)$$

Then $(A^n, \succ, \prec, \cdot)$ is a tridendriform algebra. If (A_h, \diamond_h) is an associative deformation of (A, \diamond) , then the following equations induce a tridendriform deformation $(A_h^n, \succ_h, \prec_h, \cdot_h)$ of $(A^n, \succ, \prec, \cdot)$:

$$a \succ_h b := \sum_{s=0}^{\infty} (0, a_1 \diamond_s b_2, \dots, \sum_{j=1}^{i-1} a_j \diamond_s b_i, \dots, \sum_{j=1}^{n-1} a_j \diamond_s b_n) h^s, \quad (40)$$

$$a \prec_h b := \sum_{s=0}^{\infty} (0, a_2 \diamond_s b_1, \dots, \sum_{j=1}^{i-1} a_i \diamond_s b_j, \dots, \sum_{j=1}^{n-1} a_n \diamond_s b_j) h^s, \quad (41)$$

$$a \cdot_h b := \sum_{s=0}^{\infty} (a_1 \diamond_s b_1, \dots, a_i \diamond_s b_i, \dots, a_n \diamond_s b_n) h^s. \quad (42)$$

Here, we use the infinite series expansion

$$a_i \diamond_h b_j = \sum_{s=0}^{\infty} (a_i \diamond_s b_j) h^s, \quad 1 \leq i, j \leq n.$$

If (A, \diamond) is commutative and $(A, [,], \diamond)$ is the QCL of (A_h, \diamond_h) , then the QCL $(A^n, [,], \triangleright, \succ, \cdot)$ of $(A_h^n, \succ_h, \prec_h, \cdot_h)$ is determined by (38), (39) and the equations

$$[a, b] = ([a_1, b_1], \dots, [a_i, b_i], \dots, [a_n, b_n]), \quad a \triangleright b = (0, [a_1, b_2], \dots, \sum_{j=1}^{i-1} [a_j, b_i], \dots, \sum_{j=1}^{n-1} [a_j, b_n]).$$

We recall the following results on derived structures from \mathcal{O} -operators.

Theorem 3.14. [11, 34] *Let (A, \circ) (resp. $(P, \{\cdot, \cdot\}, \circ)$) be an associative algebra (resp. a Poisson algebra) and (V, \cdot, l, r) (resp. $(Q, [\cdot, \cdot], \cdot, \rho_{\{\cdot, \cdot\}}, \rho_\circ)$) be an A -bimodule algebra (resp. a P -module Poisson algebra). Let $T : V \rightarrow A$ (resp. $T : Q \rightarrow P$) be an \mathcal{O} -operator of weight λ on (A, \circ) associated to (V, \cdot, l, r) (resp. on $(P, \{\cdot, \cdot\}, \circ)$ associated to $(Q, [\cdot, \cdot], \cdot, \rho_{\{\cdot, \cdot\}}, \rho_\circ)$). Set*

$$u \triangleright v := l(T(u))v, \quad u \triangleleft v := r(T(v))u, \quad u, v \in V.$$

$$\text{(resp. } u \triangleright v := \rho_{\{\cdot, \cdot\}}(T(u))v, \quad u \triangleright v := \rho_\circ(T(u))v, \quad u, v \in Q).$$

Then $(V, \triangleright, \triangleleft, \lambda \cdot)$ is a tridendriform algebra (resp. $(Q, \lambda[\cdot, \cdot], \triangleright, \triangleleft, \lambda \cdot)$ is a post-Poisson algebra).

Let (A, \circ) be a commutative associative algebra and (V, \cdot, ρ_\circ) be an A -module algebra. Let (A_h, \circ_h) be an associative deformation of (A, \circ) and (V_h, \cdot_h, l_h, r_h) be an A_h -bimodule algebra deformation of (V, \cdot, ρ_\circ) . Let T be an \mathcal{O} -operator of weight λ on (A, \circ) associated to (V, \cdot, ρ_\circ) . By Theorem 3.14, $(V, \triangleright, \triangleleft, \lambda \cdot)$ is a commutative tridendriform algebra, where $u \triangleright v := \rho_\circ(T(u))v, u, v \in V$. Suppose that $T_{\mathbf{K}}$ defined by (18) is an \mathcal{O} -operator of weight λ on (A_h, \circ_h) associated to (V_h, \cdot_h, l_h, r_h) . Set

$$u_h \triangleright_h v_h := l_h(T_{\mathbf{K}}(u_h))v_h, \quad u_h \triangleleft_h v_h := r_h(T_{\mathbf{K}}(v_h))u_h, \quad u_h, v_h \in V_h.$$

By Theorem 3.14 again, $(V_h, \triangleright_h, \triangleleft_h, \lambda \cdot_h)$ is a tridendriform algebra. Since (V_h, \cdot_h) is an associative deformation of (V, \cdot) , we deduce that $\lambda x \cdot_h y \equiv \lambda x \cdot y \pmod{h}$. Because (V_h, \cdot_h, l_h, r_h) is an A_h -bimodule algebra deformation of (V, \cdot, ρ_\circ) , by Proposition 2.11 we deduce that

$$u \triangleright_h v = l_h(T_{\mathbf{K}}(u))v \equiv \rho_\circ(T(u))v \pmod{h} \equiv u \triangleright v \pmod{h},$$

$$u \triangleleft_h v = r_h(T_{\mathbf{K}}(v))u \equiv \rho_\circ(T(v))u \pmod{h} \equiv v \triangleright u \pmod{h} \equiv u \triangleleft v \pmod{h}, \quad u, v \in V.$$

Thus we conclude that $(V_h, \triangleright_h, \triangleleft_h, \lambda \cdot_h)$ is a tridendriform deformation of $(V, \triangleright, \triangleleft, \lambda \cdot)$. By Theorem 3.11, let $(V, [\cdot, \cdot]', \triangleright', \triangleleft', \lambda \cdot)$ be its QCL which is a post-Poisson algebra, where

$$[u, v]' := \lambda(u \cdot_1 v - v \cdot_1 u), \quad u \triangleright' v := u \triangleright_1 v - v \triangleleft_1 u, \quad u, v \in V.$$

Here we use the expansion in (33).

On the other hand, suppose that $(A, \{\cdot, \cdot\}, \circ)$, $(V, [\cdot, \cdot], \cdot)$ and $(V, [\cdot, \cdot], \cdot, \rho_{\{\cdot, \cdot\}}, \rho_\circ)$ are the QCLs of (A_h, \circ_h) , (V_h, \cdot_h) and (V_h, \cdot_h, l_h, r_h) respectively. By Theorems 2.15 and 2.21, T is an \mathcal{O} -operator of weight λ on the Poisson algebra $(A, \{\cdot, \cdot\}, \circ)$ associated to $(V, [\cdot, \cdot], \cdot, \rho_{\{\cdot, \cdot\}}, \rho_\circ)$. By Theorem 3.14, there is a post-Poisson algebra $(V, \lambda[\cdot, \cdot], \triangleright, \triangleleft, \lambda \cdot)$. Note that

$$[u, v] = u \cdot_1 v - v \cdot_1 u, \quad u \triangleright_1 v - v \triangleleft_1 u = l_1(T(u))v - r_1(T(v))u = \rho_{\{\cdot, \cdot\}}(T(u))v, \quad u, v \in V.$$

Hence $[u, v]' = \lambda[u, v]$ and $u \triangleright' v = u \triangleright v$ for $u, v \in V$. That is, the two post-Poisson algebras $(V, [\cdot, \cdot]', \triangleright', \triangleleft', \lambda \cdot)$ and $(V, \lambda[\cdot, \cdot], \triangleright, \triangleleft, \lambda \cdot)$ coincide. In summary, we obtain the following conclusion.

Proposition 3.15. *With the notations above, starting from an \mathcal{O} -operator T , the topologically free tridendriform algebra $(V_h, \triangleright_h, \triangleleft_h, \lambda \cdot_h)$ induced by the scalar deformation \mathcal{O} -operator $T_{\mathbf{K}}$ of the \mathcal{O} -operator T coincides with the tridendriform deformation of the commutative tridendriform algebra $(V, \triangleright, \triangleleft, \lambda \cdot)$ induced by the \mathcal{O} -operator T . Moreover, the QCL post-Poisson algebra of the deformation tridendriform algebra $(V_h, \triangleright_h, \triangleleft_h, \lambda \cdot_h)$ coincides with the post-Poisson algebra induced by the \mathcal{O} -operator T on the QCL Poisson algebra $(A, \{\cdot, \cdot\}, \circ)$. In other words, the following diagram commutes.*

\mathcal{O} -operator T on (A, \circ) associated to (V, \cdot, ρ_\circ)	deformation	\mathcal{O} -operator $T_{\mathbf{K}}$ on $\triangleright(A_h, \circ_h)$ associated to (V_h, \cdot_h, l_h, r_h)	$\xrightarrow{\text{Theorem 2.15}}$ $\xrightarrow{\text{Theorem 2.21}}$	\mathcal{O} -operator T on $\triangleright(A, \{\cdot, \cdot\}, \circ)$ associated to $(V, [\cdot, \cdot], \cdot, \rho_{\{\cdot, \cdot\}}, \rho_\circ)$
$\xrightarrow{\text{Theorem 3.14}}$		$\xrightarrow{\text{Theorem 3.14}}$		$\xrightarrow{\text{Theorem 3.14}}$
\vee commutative tridendriform algebra $(V, \triangleright, \triangleleft, \lambda \cdot)$	deformation	\vee topologically free \triangleright -tridendriform algebra $(V_h, \triangleright_h, \triangleleft_h, \lambda \cdot_h)$	$\xrightarrow{\text{QCL}}$ $\xrightarrow{\text{Theorem 3.11}}$	\vee post-Poisson algebra $(V, \lambda[\cdot, \cdot], \triangleright, \triangleleft, \lambda \cdot)$

Conversely, from a tridendriform deformation of a commutative tridendriform algebra, we obtain the same \mathcal{O} -operator on a Poisson algebra associated to its module Poisson algebra by two ways as follows.

Let (A, \succ, \cdot) be a commutative tridendriform algebra, $(A_h, \succ_h, \prec_h, \cdot_h)$ be a tridendriform deformation and $(A, [,], \triangleright, \succ, \cdot)$ be the QCL post-Poisson algebra. Define binary operations \circ and $\{ , \}$ on A by (31). Define a binary operation \circ_h on A_h by (35). By Propositions 3.3 and 3.7 respectively, we obtain the following data.

- a commutative associative algebra (A, \circ) , an (A, \circ) -module algebra (A, \cdot, L_\succ) and an \mathcal{O} -operator id_A on (A, \circ) of weight 1 associated to (A, \cdot, L_\succ) ;
- a topologically free associative algebra (A_h, \circ_h) , a topologically free (A_h, \circ_h) -bimodule algebra $(A_h, \cdot_h, L_{\succ_h}, R_{\prec_h})$ and an \mathcal{O} -operator id_{A_h} of weight 1 on (A_h, \circ_h) associated to $(A_h, \cdot_h, L_{\succ_h}, R_{\prec_h})$;
- a Poisson algebra $(A, \{ , \}, \circ)$, an $(A, \{ , \}, \circ)$ -module Poisson algebra $(A, [,], \cdot, L_\triangleright, L_\succ)$ and an \mathcal{O} -operator id_A of weight 1 on $(A, \{ , \}, \circ)$ associated to $(A, [,], \cdot, L_\triangleright, L_\succ)$.

Proposition 3.16. *Keeping the notations and assumptions as above, we have the following results.*

- The topologically free (A_h, \circ_h) -bimodule algebra $(A_h, \cdot_h, L_{\succ_h}, R_{\prec_h})$ is an A_h -bimodule deformation of the (A, \circ) -module algebra (A, \cdot, L_\succ) and the QCL is the $(A, \{ , \}, \circ)$ -module Poisson algebra $(A, [,], \cdot, L_\triangleright, L_\succ)$.*
- Starting from the commutative tridendriform algebra (A, \succ, \cdot) , the \mathcal{O} -operator id_{A_h} on (A_h, \circ_h) associated to $(A_h, \cdot_h, L_{\succ_h}, R_{\prec_h})$ induced by the tridendriform deformation $(A_h, \succ_h, \prec_h, \cdot_h)$ of (A, \succ, \cdot) coincides with the scalar deformation $(\text{id}_A)_\mathbb{K}$ of the \mathcal{O} -operator id_A on (A, \circ) associated to (A, \cdot, L_\succ) induced by (A, \succ, \cdot) , that is, $\text{id}_{A_h} = (\text{id}_A)_\mathbb{K}$.*
- Starting from the topologically tridendriform algebra $(A_h, \succ_h, \prec_h, \cdot_h)$, the \mathcal{O} -operator id_A on $(A, \{ , \}, \circ)$ associated to $(A, [,], \cdot, L_\triangleright, L_\succ)$ induced by the QCL post-Poisson algebra $(A, [,], \triangleright, \succ, \cdot)$ of $(A_h, \succ_h, \prec_h, \cdot_h)$ coincides with the \mathcal{O} -operator induced by $\text{id}_{A_h} = (\text{id}_A)_\mathbb{K}$ via Theorem 2.21.*

Then we have the following commutative diagram.

$$\begin{array}{ccccc}
 \begin{array}{c} \text{commutative} \\ \text{tridendriform algebra} \\ (A, \succ, \cdot) \end{array} & \begin{array}{c} \text{deformation} \\ \triangleright \end{array} & \begin{array}{c} \text{topologically free} \\ \triangleright \text{tridendriform algebra} \\ (A_h, \succ_h, \prec_h, \cdot_h) \end{array} & \begin{array}{c} \text{QCL} \\ \text{Theorem 3.11} \end{array} & \begin{array}{c} \triangleright \\ \text{post-Poisson} \\ \text{algebra} \\ (A, [,], \triangleright, \succ, \cdot) \end{array} \\
 \text{Proposition 3.3} & & \text{Proposition 3.3} & & \text{Proposition 3.7} \\
 \begin{array}{c} \mathcal{O}\text{-operator } \text{id}_A \text{ on} \\ (A, \circ) \text{ associated to} \\ (A, \cdot, L_\succ) \end{array} & \begin{array}{c} \text{deformation} \\ \triangleright \end{array} & \begin{array}{c} \mathcal{O}\text{-operator } \text{id}_{A_h} = (\text{id}_A)_\mathbb{K} \\ \triangleright \text{ on } (A_h, \circ_h) \text{ associated to} \\ (A_h, \cdot_h, L_{\succ_h}, R_{\prec_h}) \end{array} & \begin{array}{c} \text{Theorem 2.15} \\ \text{Theorem 2.21} \end{array} & \begin{array}{c} \mathcal{O}\text{-operator } \text{id}_A \text{ on} \\ \triangleright (A, \{ , \}, \circ) \text{ associated to} \\ (A, [,], \cdot, L_\triangleright, L_\succ) \end{array}
 \end{array}$$

Proof. (a). By (32), we deduce

$$\begin{aligned}
 x \cdot_h y &\equiv x \cdot y \pmod{h}, \\
 x \circ_h y &= x_h \succ_h y_h + x_h \prec_h y_h + x_h \cdot_h y_h \equiv x \circ y \pmod{h}, \\
 L_{\succ_h}(x)y &= x \succ_h y \equiv x \succ y \pmod{h} \equiv L_\succ(x)y \pmod{h}, \\
 R_{\prec_h}(y)x &= x \prec_h y \equiv x \prec y \pmod{h} \equiv R_\prec(y)x \pmod{h}, \quad x, y \in A.
 \end{aligned}$$

Thus the (A_h, \circ_h) -bimodule algebra $(A_h, \cdot_h, L_{\succ_h}, R_{\prec_h})$ is an A_h -bimodule deformation of the (A, \circ) -module algebra (A, \cdot, L_\succ) by Proposition 2.11.

By (37) and (34), the QCLs of (A_h, \circ_h) and (A_h, \cdot_h) are the Poisson algebras $(A, \{ , \}, \circ)$ and $(A, [,], \circ)$ respectively. Note that

$$\frac{L_{>_h}(x)y - R_{<_h}(x)y}{h} = \frac{x >_h y - y <_h x}{h} \equiv x \triangleright y \pmod{h} \equiv L_{\triangleright}(x)y \pmod{h}, \quad x, y \in A.$$

Thus the conclusion follows.

Because

$$\text{id}_{A_h} \left(\sum_{i=0}^{\infty} x_i h^i \right) = \sum_{i=0}^{\infty} x_i h^i = \sum_{i=0}^{\infty} \text{id}_A(x_i) h^i, \quad \sum_{i=0}^{\infty} x_i h^i \in A_h,$$

we obtain that id_{A_h} coincides with the scalar deformation $(\text{id}_A)_{\mathbf{K}}$ of id_A . Hence (b) and (c) hold. \square

Remark 3.17. As an application of Proposition 3.16, we give another proof of Theorem 3.11 as follows. By Proposition 3.16 (a), $(A, [\cdot, \cdot], \cdot, L_{\triangleright}, L_{\triangleright})$ is exactly the QCL of $(A_h, \cdot_h, L_{>_h}, R_{<_h})$. Then $(A, [\cdot, \cdot], \cdot, L_{\triangleright}, L_{\triangleright})$ is an $(A, \{\cdot, \cdot\}, \circ)$ -module Poisson algebra by Theorem 2.15. Hence by Proposition 3.7, $(A, [\cdot, \cdot], \triangleright, \triangleright, \cdot)$ is a post-Poisson algebra.

Also note that Proposition 3.16 provides the scalarly deformed \mathcal{O} -operator id_{A_h} as an explicit example to illustrate Theorem 2.21.

3.2. Constructions of bimodule algebra deformations and tridendriform deformations by derivations. We recall a deformation construction for associative algebras induced by derivations.

Lemma 3.18. [19] *Let d_1 and d_2 be commuting derivations on an associative algebra (A, \circ) . Set*

$$x \circ_h y := \sum_{s=0}^{\infty} (d_1^s(x) \circ d_2^s(y)) \frac{h^s}{s!}, \quad x, y \in A, \quad (43)$$

and extend it to A_h by \mathbf{K} -bilinearity. Then (A_h, \circ_h) is an associative deformation of (A, \circ) .

We extend this construction to A_h -bimodule algebra deformations.

Proposition 3.19. *Let (A, \circ) be an associative algebra and (V, \cdot, l, r) an A -bimodule algebra. Let d_1 and d_2 be commuting derivations on the associative algebra $A \ltimes_{l,r} V$ such that $d_i(A) \subset A$ and $d_i(V) \subset V$, $i = 1, 2$. Define*

$$\begin{aligned} x \circ_h y &:= \sum_{s=0}^{\infty} (d_1^s(x) \circ d_2^s(y)) \frac{h^s}{s!}, & u \cdot_h v &:= \sum_{s=0}^{\infty} (d_1^s(u) \cdot d_2^s(v)) \frac{h^s}{s!}, \\ l_h(x)v &:= \sum_{s=0}^{\infty} (l(d_1^s(x))d_2^s(v)) \frac{h^s}{s!}, & r_h(x)v &:= \sum_{s=0}^{\infty} (r(d_1^s(x))d_2^s(v)) \frac{h^s}{s!}, \end{aligned} \quad x, y \in A, u, v \in V.$$

Extend \mathbf{K} -bilinearly the above operations to A_h and V_h . Then (V_h, \cdot_h, l_h, r_h) is an A_h -bimodule algebra deformation of (V, \cdot, l, r) .

Proof. Obviously, d_1 and d_2 are commuting derivations on the subalgebras (A, \circ) and (V, \cdot) respectively. By Lemma 3.18, (A_h, \circ_h) and (V_h, \cdot_h) are associative deformations of (A, \circ) and (V, \cdot) respectively. We only need to show that $(A_h \oplus V_h, \odot_h)$ is an associative deformation of $A \ltimes_{l,r} V$. Note that

$$\begin{aligned} (x, u) \odot_h (y, v) &= (x \circ_h y, l_h(x)v + r_h(y)u + u \cdot_h v) \\ &= \sum_{s=0}^{\infty} (d_1^s(x) \circ d_2^s(y), l(d_1^s(x))d_2^s(v) + r(d_1^s(y))d_2^s(u) + d_1^s(u) \cdot d_2^s(v)) \frac{h^s}{s!} \\ &= \sum_{s=0}^{\infty} d_1^s((x, u)) \odot d_2^s((y, v)) \frac{h^s}{s!}, \quad x, y \in A, u, v \in V. \end{aligned}$$

By Lemma 3.18 again, $(A_h \oplus V_h, \odot_h)$ is an associative deformation of $A \ltimes_{l,r} V$. Thus (V_h, \cdot_h, l_h, r_h) is an A_h -bimodule algebra deformation of (V, \cdot, l, r) . \square

Let (A, \succ, \prec, \cdot) be a tridendriform algebra. If $d \in \text{End}_{\mathbf{k}}(A)$ satisfies

$$d(x * y) = d(x) * y + x * d(y), \quad x, y \in A, * \in \{\succ, \prec, \cdot\},$$

then d is called a **derivation on the tridendriform algebra** (A, \succ, \prec, \cdot) . Applying Proposition 3.19 to the (A, \circ) -bimodule algebra $(A, \cdot, L_{\succ}, R_{\prec})$ gives the following consequence.

Corollary 3.20. *If d_1 and d_2 are commuting derivations on a tridendriform algebra (A, \succ, \prec, \cdot) , then the operations*

$$x *_h y := \sum_{s=0}^{\infty} (d_1^s(x) * d_2^s(y)) \frac{h^s}{s!}, \quad x, y \in A, \quad (44)$$

for each $* \in \{\succ, \prec, \cdot\}$, give a tridendriform deformation $(A_h, \succ_h, \prec_h, \cdot_h)$ of (A, \succ, \prec, \cdot) .

Proposition 3.21. *With the assumptions in Proposition 3.19. Let T be an \mathcal{O} -operator of weight λ on (A, \circ) associated to (V, \cdot, l, r) . If $Td_1|_V = d_1T$ and $Td_2|_V = d_2T$, then $T_{\mathbf{K}}$ defined by (18) is an \mathcal{O} -operator of weight λ on (A_h, \circ_h) associated to (V_h, \cdot_h, l_h, r_h) .*

Proof. Let $u, v \in V$. Then for $s \in \mathbb{N}$, we have

$$\begin{aligned} T(u) \cdot_s T(v) &= \frac{d_1^s(T(u)) \cdot d_2^s(T(v))}{s!} \\ &= \frac{T(d_1^s(u)) \cdot T(d_2^s(v))}{s!} \\ &= \frac{T(l(d_1^s(u))d_2^s(v))}{s!} + \frac{T(r(Td_2^s(v))d_1^s(u))}{s!} + \lambda \frac{T(d_1^s(u) \cdot d_2^s(v))}{s!} \\ &= \frac{T(l(d_1^s(T(u))d_2^s(v))}{s!} + \frac{T(r(d_2^s(T(v)))d_1^s(u))}{s!} + \lambda \frac{T(d_1^s(u) \cdot d_2^s(v))}{s!} \\ &= T(l_s(T(u))v) + T(r_s(T(v))) + \lambda T(u \cdot_s v). \end{aligned}$$

Therefore $T_{\mathbf{K}}$ is an \mathcal{O} -operator of weight λ on (A_h, \circ_h) associated to (V_h, \cdot_h, l_h, r_h) . \square

Proposition 3.22. *Let d_1 and d_2 be commuting derivations on an associative algebra (A, \cdot) and T be a Rota-Baxter operator of weight 1 on (A, \cdot) that commutes with d_1 and d_2 . Define*

$$\begin{aligned} x > y &:= T(x) \cdot y, \quad x < y := x \cdot T(y), \\ x >_h y &:= \sum_{s=0}^{\infty} (d_1^s(T(x)) \cdot d_2^s(y)) \frac{h^s}{s!}, \quad x <_h y := \sum_{s=0}^{\infty} (d_1^s(x) \cdot d_2^s(T(y))) \frac{h^s}{s!}, \\ x \cdot_h y &= \sum_{s=0}^{\infty} (d_1^s(x) \cdot d_2^s(y)) \frac{h^s}{s!}, \quad x, y \in A. \end{aligned}$$

Extend \mathbf{K} -bilinearly the above operations $>_h, <_h$ and \cdot_h to A_h . Then $(A_h, \succ_h, \prec_h, \cdot_h)$ is a tridendriform deformation of the tridendriform algebra (A, \succ, \prec, \cdot) . If in addition (A, \cdot) is commutative, then the QCL of the tridendriform deformation is the post-Poisson algebra $(A, \triangleright, \{, \}, \succ, \cdot)$, where \triangleright and $\{, \}$ are respectively defined by

$$x \triangleright y := d_1(T(x)) \cdot d_2(y) - d_1(y) \cdot d_2(T(x)), \quad \{x, y\} := d_1(x) \cdot d_2(y) - d_1(y) \cdot d_2(x), \quad x, y \in A.$$

Proof. Applying Proposition 3.21 to the \mathcal{O} -operator T of weight 1 on (A, \cdot) associated to the A -bimodule algebra (A, \cdot, L, R) , we find that $T_{\mathbf{K}}$ is an Rota-Baxter operator of weight 1 on (A_h, \cdot_h) . Note that (32) also holds. Thus $(A_h, \succ_h, \prec_h, \cdot_h)$ is a tridendriform deformation of (A, \succ, \prec, \cdot) . The last conclusion follows from a direct verification. \square

We apply the above construction to give an example of the tridendriform deformation.

Let (A, \cdot) be the (nonunital) subalgebra of $\mathbf{k}[x_1, x_2]$ generated by x_1 and x_2 , that is, $A = x_1\mathbf{k}[x_1, x_2] + x_2\mathbf{k}[x_1, x_2]$. It is known [20] that, if $q \in \mathbf{k}$ is not a root of unit, then a Rota-Baxter operator of weight 1 on the algebra $x_1\mathbf{k}[x_1]$ is given by

$$T(x_1^{i_1}) = \frac{q^{i_1} x_1^{i_1}}{1 - q^{i_1}}, \quad i_1 \in \mathbb{Z}^+.$$

Let $q_1, q_2 \in \mathbf{k}$ be such that $q_1^{i_1} q_2^{i_2} \neq 1$ for $i_1, i_2 \geq 0$ with $i_1 i_2 \neq 0$. Define a linear map

$$T_{q_1, q_2} : A \rightarrow A, \quad T_{q_1, q_2}(x_1^{i_1} x_2^{i_2}) := \frac{q_1^{i_1} q_2^{i_2}}{1 - q_1^{i_1} q_2^{i_2}} x_1^{i_1} x_2^{i_2}, \quad i_1, i_2 \in \mathbb{N}, i_1 i_2 \neq 0.$$

Noting that

$$\frac{q_1^{i_1+j_1} q_2^{i_2+j_2}}{(1 - q_1^{i_1} q_2^{i_2})(1 - q_1^{j_1} q_2^{j_2})} = \frac{q_1^{i_1} q_2^{i_2}}{1 - q_1^{i_1} q_2^{i_2}} \frac{q_1^{i_1+j_1} q_2^{i_2+j_2}}{1 - q_1^{i_1+j_1} q_2^{i_2+j_2}} + \frac{q_1^{j_1} q_2^{j_2}}{1 - q_1^{j_1} q_2^{j_2}} \frac{q_1^{i_1+j_1} q_2^{i_2+j_2}}{1 - q_1^{i_1+j_1} q_2^{i_2+j_2}} + \frac{q_1^{i_1+j_1} q_2^{i_2+j_2}}{1 - q_1^{i_1+j_1} q_2^{i_2+j_2}},$$

where $i_1, i_2, j_1, j_2 \in \mathbb{N}$, $i_1 i_2 \neq 0$, $j_1 j_2 \neq 0$, we deduce that

$$\begin{aligned} & T_{q_1, q_2}(x_1^{i_1} x_2^{i_2}) \cdot T_{q_1, q_2}(x_1^{j_1} x_2^{j_2}) \\ &= T_{q_1, q_2}(T_{q_1, q_2}(x_1^{i_1} x_2^{i_2}) \cdot x_1^{j_1} x_2^{j_2}) + T_{q_1, q_2}(x_1^{i_1} x_2^{i_2} \cdot T_{q_1, q_2}(x_1^{j_1} x_2^{j_2})) + T_{q_1, q_2}(x_1^{i_1} x_2^{i_2} \cdot x_1^{j_1} x_2^{j_2}). \end{aligned}$$

Hence T_{q_1, q_2} is a Rota-Baxter operator of weight 1 on A . We also set

$$d_1 := x_1 \frac{\partial}{\partial x_1}, \quad d_2 := x_2 \frac{\partial}{\partial x_2}.$$

Then T_{q_1, q_2}, d_1 and d_2 are pairwise commuting. Therefore we obtain the following example.

Example 3.23. Let $A = x_1 \mathbf{k}[x_1, x_2] + x_2 \mathbf{k}[x_1, x_2]$. Fix $q_1, q_2 \in \mathbf{k}$ such that $q_1^{i_1} q_2^{i_2} \neq 1$ for $i_1, i_2 \geq 0$ with $i_1 i_2 \neq 0$. A commutative tridendriform algebra structure $(A, >, \cdot)$ is defined by

$$x_1^{i_1} x_2^{i_2} > x_1^{j_1} x_2^{j_2} := \frac{q_1^{i_1} q_2^{i_2}}{1 - q_1^{i_1} q_2^{i_2}} x_1^{i_1+j_1} x_2^{i_2+j_2}, \quad (45)$$

$$x_1^{i_1} x_2^{i_2} \cdot x_1^{j_1} x_2^{j_2} := x_1^{i_1+j_1} x_2^{i_2+j_2}, \quad i_1, i_2, j_1, j_2 \in \mathbb{N}, i_1 i_2 \neq 0, j_1 j_2 \neq 0. \quad (46)$$

A tridendriform deformation $(A, >_h, <_h, \cdot_h)$ of $(A, >, \cdot)$ is given by

$$\begin{aligned} x_1^{i_1} x_2^{i_2} >_h x_1^{j_1} x_2^{j_2} &:= \sum_{s=0}^{\infty} \frac{i_1^s j_2^s q_1^{i_1} q_2^{i_2}}{1 - q_1^{i_1} q_2^{i_2}} x_1^{i_1+j_1} x_2^{i_2+j_2} \frac{h^s}{s!}, \quad x_1^{i_1} x_2^{i_2} <_h x_1^{j_1} x_2^{j_2} := \sum_{s=0}^{\infty} \frac{i_1^s j_2^s q_1^{i_1} q_2^{i_2}}{1 - q_1^{i_1} q_2^{i_2}} x_1^{i_1+j_1} x_2^{i_2+j_2} \frac{h^s}{s!}, \\ x_1^{i_1} x_2^{i_2} \cdot_h x_1^{j_1} x_2^{j_2} &:= \sum_{s=0}^{\infty} i_1^s j_2^s x_1^{i_1+j_1} x_2^{i_2+j_2} \frac{h^s}{s!}. \end{aligned}$$

The post-Poisson algebra $(A, [,], \triangleright, >, \cdot)$ for the corresponding QCL is given by (45), (46) and

$$[x_1^{i_1} x_2^{i_2}, x_1^{j_1} x_2^{j_2}] := (i_1 j_2 - i_2 j_1) x_1^{i_1+j_1} x_2^{i_2+j_2}, \quad x_1^{i_1} x_2^{i_2} \triangleright x_1^{j_1} x_2^{j_2} := (i_1 j_2 - i_2 j_1) \frac{q_1^{i_1} q_2^{i_2}}{1 - q_1^{i_1} q_2^{i_2}} x_1^{i_1+j_1} x_2^{i_2+j_2}.$$

4. SCALAR DEFORMATIONS OF SOLUTIONS OF AYBE

In this section, we construct solutions of the Poisson Yang-Baxter equation (PYBE) by our general method of bimodule algebra deformations. For a commutative associative algebra (A, \circ) and an associative deformation (A_h, \circ_h) whose QCL is a Poisson algebra $(A, \{, \}, \circ)$, we show that a scalarly deformed solution of associative Yang-Baxter equation (AYBE) in (A_h, \circ_h) with invariant symmetric part induces a solution of PYBE in $(A, \{, \}, \circ)$. We finally obtain solutions of the PYBE from tridendriform deformations of commutative tridendriform algebras.

4.1. Scalar deformations of solutions of AYBE in associative deformations. We recall the notions of associative Yang-Baxter equations.

Definition 4.1. [9] Let (A, \circ) be an associative algebra and $r = \sum_i a_i \otimes b_i \in A \otimes A$. Set

$$r_{12} \circ r_{13} := \sum_{i,j} a_i \circ a_j \otimes b_i \otimes b_j, \quad r_{13} \circ r_{23} := \sum_{i,j} a_i \otimes a_j \otimes b_i \circ b_j, \quad r_{23} \circ r_{12} := \sum_{i,j} a_j \otimes a_i \circ b_j \otimes b_i. \quad (47)$$

We say that r is a solution of the **associative Yang-Baxter equation (AYBE)** in (A, \circ) if

$$\mathbf{A}(r) := r_{12} \circ r_{13} + r_{13} \circ r_{23} - r_{23} \circ r_{12} = 0. \quad (48)$$

Remark 4.2. In [1], another form of the AYBE is defined by

$$\mathbf{A}^{\text{op}}(r) := r_{13} \circ r_{12} + r_{23} \circ r_{13} - r_{12} \circ r_{23} = 0, \quad (49)$$

where

$$r_{13} \circ r_{12} := \sum_{i,j} a_i \circ a_j \otimes b_j \otimes a_j, \quad r_{23} \circ r_{13} := \sum_{i,j} a_j \otimes a_i \otimes b_i \circ b_j, \quad r_{12} \circ r_{23} := \sum_{i,j} a_i \otimes b_i \circ a_j \otimes b_j. \quad (50)$$

By [3, Lemma 3.4], when the symmetric part

$$\beta := \frac{1}{2}(r + \sigma(r)) \quad (51)$$

of r is (A, \circ) -invariant, that is,

$$(\text{id}_A \otimes L_\circ(x) - R_\circ(x) \otimes \text{id}_A)\beta = 0, \quad x \in A, \quad (52)$$

then $\mathbf{A}^{\text{op}}(r) = 0$ if and only if $\mathbf{A}(r) = 0$. Here σ is the **twisting operator**

$$\sigma : A \otimes A \rightarrow A \otimes A, \quad \sigma(x \otimes y) = y \otimes x, \quad x, y \in A.$$

The AYBE can be defined on a topologically free associative algebra by replacing the usual tensor product in the AYBE with the topological tensor product. See [21, 22] for details. In fact, let V and W be vector spaces and $V_h = V[[h]]$ and $W_h = W[[h]]$ be topologically free \mathbf{K} -modules. Then $V_h \hat{\otimes} W_h \cong (V \otimes W)[[h]]$.

Let (A, \circ) be an associative algebra and (A_h, \circ_h) be an associative deformation. Any element $r = \sum_i a_i \otimes b_i \in A \otimes A$ is naturally an element in $(A \otimes A)[[h]]$. Then under the isomorphism $(A \otimes A)[[h]] \cong A_h \hat{\otimes} A_h$, r is also naturally an element in $A_h \hat{\otimes} A_h$ and so takes the form $\sum_i a_i \hat{\otimes} b_i$. We let \hat{r} denote the image of $r \in A \otimes A$ under the linear embedding of $A \otimes A$ into $A_h \hat{\otimes} A_h$ and call \hat{r} the **scalar deformation** of r .

We can also extend the twisting operator $\sigma : A \otimes A \rightarrow A \otimes A$ to the topological tensor product by \mathbf{K} -bilinear extension to get the **topological twisting operator**

$$\sigma_{\mathbf{K}} : A_h \hat{\otimes} A_h \rightarrow A_h \hat{\otimes} A_h, \quad \sigma_{\mathbf{K}}(x_h \hat{\otimes} y_h) = y_h \hat{\otimes} x_h, \quad x_h, y_h \in A_h.$$

Let $(\mathfrak{g}, [,])$ be a Lie algebra and $r = \sum_i y_i \otimes y'_i \in \mathfrak{g} \otimes \mathfrak{g}$. Set

$$[r_{12}, r_{13}] := \sum_{i,j} [y_i, y_j] \otimes y'_i \otimes y'_j, \quad [r_{12} \circ r_{23}] := \sum_{i,j} y_i \otimes [y'_i, y_j] \otimes y'_j, \quad [r_{13}, r_{23}] := \sum_{i,j} y_i \otimes y_j \otimes [y'_i, y'_j]. \quad (53)$$

Recall that the equation

$$\mathbf{C}(r) := [r_{12}, r_{13}] + [r_{12}, r_{23}] + [r_{13}, r_{23}] = 0 \quad (54)$$

is called the **classical Yang-Baxter equation (CYBE)**. For the adjoint representation of \mathfrak{g} :

$$\text{ad} : \mathfrak{g} \rightarrow \text{End}_{\mathbf{K}}(\mathfrak{g}), \quad \text{ad}(x)(y) = [x, y], \quad x, y \in \mathfrak{g},$$

an element $r \in \mathfrak{g} \otimes \mathfrak{g}$ is called $(\mathfrak{g}, [,])$ -invariant if

$$(\text{ad}(x) \otimes \text{id} + \text{id} \otimes \text{ad}(x))r = 0, \quad x \in \mathfrak{g}.$$

Definition 4.3. [34] Let $(A, \{, \}, \circ)$ be a Poisson algebra and $r \in A \otimes A$. The equation

$$\mathbf{A}(r) = \mathbf{C}(r) = 0 \quad (55)$$

is called the **Poisson Yang-Baxter equation (PYBE)** in $(A, \{, \}, \circ)$.

Theorem 4.4. *Let (A, \circ) be a commutative associative algebra, (A_h, \circ_h) be an associative deformation and $(A, \{, \}, \circ)$ be the corresponding QCL. Let $r = \sum_i a_i \otimes b_i \in A \otimes A$ be a solution of the AYBE in (A, \circ) and $r + \sigma(r)$ be (A, \circ) -invariant. Let \hat{r} denote the image of r under the linear embedding of $A \otimes A$ into $A_h \hat{\otimes} A_h$. Suppose that \hat{r} is a solution of the AYBE in (A_h, \circ_h) and $\hat{r} + \sigma_{\mathbf{K}}(\hat{r})$ is (A_h, \circ_h) -invariant. Then r is a solution of the PYBE in $(A, \{, \}, \circ)$ and $r + \sigma(r)$ is $(A, \{, \}, \circ)$ -invariant.*

Proof. Since \hat{r} is a solution of the AYBE in (A_h, \circ_h) and $\hat{r} + \sigma_{\mathbf{K}}(\hat{r})$ is (A_h, \circ_h) -invariant, it follows that $\mathbf{A}^{\text{op}}(\hat{r}) = 0$ in (A_h, \circ_h) . Note that

$$\begin{aligned} \hat{r}_{12} \circ_h \hat{r}_{13} - \hat{r}_{13} \circ_h \hat{r}_{12} &= \sum_{i,j} a_i \circ_h a_j \hat{\otimes} b_i \hat{\otimes} b_j - \sum_{i,j} a_i \circ_h a_j \hat{\otimes} b_j \hat{\otimes} b_i \\ &= \sum_{i,j} a_i \circ_h a_j \hat{\otimes} b_i \hat{\otimes} b_j - \sum_{i,j} a_j \circ_h a_i \hat{\otimes} b_i \hat{\otimes} b_j \\ &= \sum_{i,j} (a_i \circ_h a_j - a_j \circ_h a_i) \hat{\otimes} b_i \hat{\otimes} b_j \\ &\equiv \sum_{i,j} \{a_i, a_j\} \hat{\otimes} b_i \hat{\otimes} b_j \ h \pmod{h^2}. \end{aligned}$$

Similarly, we obtain

$$\begin{aligned} \hat{r}_{13} \circ_h \hat{r}_{23} - \hat{r}_{23} \circ_h \hat{r}_{13} &\equiv \sum_{i,j} a_i \hat{\otimes} a_j \hat{\otimes} \{b_i b_j\} \ h \pmod{h^2}, \\ \hat{r}_{12} \circ_h \hat{r}_{23} - \hat{r}_{23} \circ_h \hat{r}_{12} &\equiv \sum_{i,j} a_i \hat{\otimes} \{b_i, a_j\} \hat{\otimes} b_j \ h \pmod{h^2}. \end{aligned}$$

So

$$\frac{\mathbf{A}(\hat{r}) - \mathbf{A}^{\text{op}}(\hat{r})}{h} \equiv \mathbf{C}(r)h \pmod{h}.$$

Then we obtain $\mathbf{C}(r) = 0$ in $(A, \{, \})$. Note that r is a solution of the AYBE in (A, \circ) . Hence r is also a solution of the PYBE in $(A, \{, \}, \circ)$.

Since $\hat{r} + \sigma_{\mathbf{K}}(\hat{r})$ is (A_h, \circ_h) -invariant, we have

$$(\text{id}_{A_h} \hat{\otimes} L_{\circ_h}(x) - R_{\circ_h}(x) \hat{\otimes} \text{id}_{A_h})(\hat{r} + \sigma_{\mathbf{K}}(\hat{r})) = 0, \quad x \in A. \quad (56)$$

Applying $\sigma_{\mathbf{K}}$ to both sides of (56), we deduce that

$$(L_{\circ_h}(x) \hat{\otimes} \text{id}_{A_h} - \text{id}_{A_h} \hat{\otimes} R_{\circ_h}(x))(\hat{r} + \sigma_{\mathbf{K}}(\hat{r})) = 0, \quad x \in A. \quad (57)$$

So adding (56) and (57) gives

$$((L_{\circ_h}(x) - R_{\circ_h}(x)) \hat{\otimes} \text{id}_{A_h} + \text{id}_{A_h} \hat{\otimes} (L_{\circ_h}(x) - R_{\circ_h}(x)))(\hat{r} + \sigma_{\mathbf{K}}(\hat{r})) = 0, \quad x \in A.$$

Since $\frac{L_{\circ_h}(x) - R_{\circ_h}(x)}{h} \equiv \text{ad}_{\{, \}}(x) \pmod{h}$ for $x \in A$, we see that $r + \sigma(r)$ is also $(A, \{, \})$ -invariant, implying its $(A, \{, \}, \circ)$ -invariance due to the already assumed (A, \circ) -invariance. \square

Remark 4.5. In fact, without the assumption that \hat{r} is a solution of the AYBE in (A_h, \circ_h) , the conclusion involving the invariance is still available. Explicitly, for any symmetric and (A, \circ) -invariant $\beta \in A \otimes A$, if $\hat{\beta}$ is (A_h, \circ_h) -invariant, then β is $(A, \{, \}, \circ)$ -invariant.

4.2. Topological dual bimodules. Let V and W be finite-dimensional vector spaces. Let $V^* = \text{Hom}_{\mathbf{K}}(V, \mathbf{K})$ and $W^* = \text{Hom}_{\mathbf{K}}(W, \mathbf{K})$ be the dual space of V and W respectively. For $r \in W \otimes V$, let r^{\sharp} be the image under the linear isomorphism

$$W \otimes V \cong \text{Hom}_{\mathbf{K}}(V^*, W). \quad (58)$$

It is characterized by

$$\langle w^*, r^{\sharp}(v^*) \rangle = \langle w^* \otimes v^*, r \rangle, \quad v^* \in V^*, w^* \in W^*, \quad (59)$$

where $\langle \cdot, \cdot \rangle$ is the natural pairing between the dual space of a vector space and the vector space. For $f \in \text{Hom}_{\mathbf{K}}(V^*, W)$, we also let \tilde{f} denote its image in $W \otimes V$. For a vector space A and a linear map $\varrho : A \rightarrow \text{End}_{\mathbf{K}}(V)$, we let $\varrho^* : A \rightarrow \text{End}_{\mathbf{K}}(V^*)$ be the unique linear map satisfying

$$\langle \varrho^*(x)u^*, v \rangle = -\langle u^*, \varrho(x)v \rangle, \quad x \in A, u^* \in V^*, v \in V. \quad (60)$$

Let (A, \circ) be an associative algebra and $(V, l_{\circ}, r_{\circ})$ be an A -bimodule. Then $(V, -r_{\circ}^*, -l_{\circ}^*)$ is an A -bimodule, called the **dual bimodule** of $(V, l_{\circ}, r_{\circ})$. Similarly, if $(P, [,], \cdot)$ is a Poisson algebra and $(W, \rho_{[,]}, \rho_{\cdot})$ is a $(P, [,], \cdot)$ -module, then $(W^*, \rho_{[,]}^*, -\rho_{\cdot}^*)$ is a $(P, [,], \cdot)$ -module, called the **dual module** of the $(P, [,], \cdot)$ -module $(W, \rho_{[,]}, \rho_{\cdot})$.

For topologically free \mathbf{K} -modules V_h and W_h , we have \mathbf{K} -module isomorphisms

$$\text{Hom}_{\mathbf{K}}(V^*, W)[[h]] \cong (W \otimes V)[[h]] \cong W_h \hat{\otimes} V_h.$$

On the other hand, by [22, XVI.7 Exercise 2], there is the \mathbf{K} -module isomorphism

$$\Phi_{V,W} : \text{Hom}_{\mathbf{k}}(V, W)[[h]] \rightarrow \text{Hom}_{\mathbf{K}}(V_h, W_h), \quad \Phi_{V,W} \left(\sum_{s=0}^{\infty} \psi_s h^s \right) := \sum_{s=0}^{\infty} (\psi_s)_{\mathbf{K}} h^s, \quad (61)$$

where $\psi_s \in \text{Hom}_{\mathbf{k}}(V, W)$ for $s \in \mathbb{N}$ and $(\psi_s)_{\mathbf{K}}$ is defined as in (18). In particular, taking $W = \mathbf{k}$, we obtain

$$\text{Hom}_{\mathbf{K}}(V_h, \mathbf{K}) \cong \text{Hom}_{\mathbf{k}}(V, \mathbf{k})[[h]] = (V^*)_h.$$

Then the \mathbf{K} -linear map $\Phi_{V,\mathbf{k}} : \text{Hom}_{\mathbf{k}}(V, \mathbf{k})[[h]] \rightarrow \text{Hom}_{\mathbf{K}}(V_h, \mathbf{K})$ is given by

$$\Phi_{V,\mathbf{k}} \left(\sum_{s=0}^{\infty} v_s^* h^s \right) := \sum_{s=0}^{\infty} (v_s^*)_{\mathbf{K}} h^s, \quad (62)$$

where $v_s^* \in \text{Hom}_{\mathbf{k}}(V, \mathbf{k})$ for any $s \in \mathbb{N}$. For clarity, we use the notation

$$V_h^* := (V^*)_h.$$

Then we obtain

$$\text{Hom}_{\mathbf{K}}(V^*, W)[[h]] \cong \text{Hom}_{\mathbf{K}}(V_h^*, W_h) \cong W_h \hat{\otimes} V_h \cong (W \otimes V)[[h]]. \quad (63)$$

Under these isomorphisms, the images of $f \in \text{Hom}_{\mathbf{k}}(V^*, W)$ in $\text{Hom}_{\mathbf{K}}(V_h^*, W_h)$, $W_h \hat{\otimes} V_h$ and $(W \otimes V)[[h]]$ are denoted by $f_{\mathbf{K}}$, \hat{f} and \tilde{f} respectively. Also, the images of $r \in W \otimes V$ in $\text{Hom}_{\mathbf{k}}(V^*, W)[[h]]$, $\text{Hom}_{\mathbf{K}}(V_h^*, W_h)$ and $W_h \hat{\otimes} V_h$ are denoted by r^{\sharp} , $(r^{\sharp})_{\mathbf{K}}$ and \hat{r} respectively.

Next we define the pairing

$$\langle \cdot, \cdot \rangle_{\mathbf{K}} : V_h^* \hat{\otimes} V_h \rightarrow \mathbf{K}, \quad \langle u^*, v' \rangle_{\mathbf{K}} := \Phi_{V,\mathbf{k}}(u^*)(v'), \quad u^* \in V_h^*, v' \in V_h. \quad (64)$$

Since $\Phi_{V,\mathbf{k}}(v^*) = (v^*)_{\mathbf{K}}$, $v^* \in V^*$, we get

$$\langle v^*, u \rangle_{\mathbf{K}} = (v^*)_{\mathbf{K}}(u) = \langle v^*, u \rangle, \quad v^* \in V^*, u \in V. \quad (65)$$

Hence, $\langle \cdot, \cdot \rangle_{\mathbf{K}}$ is precisely the \mathbf{K} -bilinear extension of the usual pairing $\langle \cdot, \cdot \rangle$. Since there are natural isomorphisms of \mathbf{K} -modules

$$W_h^* \hat{\otimes} V_h^* \cong (W^* \otimes V^*)[[h]] \text{ and } W_h \hat{\otimes} V_h \cong (W \otimes V)[[h]],$$

the pairing $\langle \cdot, \cdot \rangle_{\mathbf{K}}$ can be turned into a pairing between $W_h^* \hat{\otimes} V_h^*$ and $W_h \hat{\otimes} V_h$ by

$$\langle w^* \hat{\otimes} v^*, w_h \hat{\otimes} v_h \rangle_{\mathbf{K}} = \langle w^*, w_h \rangle_{\mathbf{K}} \langle v^*, v_h \rangle_{\mathbf{K}}, \quad w^* \hat{\otimes} v^* \in W_h^* \hat{\otimes} V_h^*, w_h \hat{\otimes} v_h \in W_h \hat{\otimes} V_h.$$

For any $r_h \in W_h \hat{\otimes} V_h$, let $(r_h)^{\sharp}$ be the image of r_h in the \mathbf{K} -linear isomorphism

$$W_h \hat{\otimes} V_h \cong \text{Hom}_{\mathbf{K}}(V_h^*, W_h). \quad (66)$$

It is characterized by

$$\langle w^*, r_h^{\sharp}(v^*) \rangle_{\mathbf{K}} = \langle w^* \hat{\otimes} v^*, r_h \rangle_{\mathbf{K}}, \quad v^* \in V_h^*, w^* \in W_h^*. \quad (67)$$

For $f_h \in \text{Hom}_{\mathbf{K}}(V_h^*, W_h)$, we also let \tilde{f}_h denote its image in $W_h \hat{\otimes} V_h$. This agrees with the \mathbf{K} -linear isomorphism $W_h \hat{\otimes} V_h \cong \text{Hom}_{\mathbf{K}}(V_h^*, W_h)$ given in (63) as follows.

Proposition 4.6. *Let r be an element in $W \otimes V$. Then we have*

$$(\hat{r})^{\sharp} = (r^{\sharp})_{\mathbf{K}}. \quad (68)$$

Accordingly, for $f \in \text{Hom}_{\mathbf{k}}(V, W)$, we have

$$\tilde{f}_{\mathbf{K}} = \hat{f}. \quad (69)$$

Proof. Let $r = \sum_i w_i \otimes v_i \in W \otimes V$ with $v_i \in V$ and $w_i \in W$. Let $v^* \in V^*$, $w^* \in W_h^*$. Then we have

$$\begin{aligned} \langle w^*, (\hat{r})^{\sharp}(v^*) \rangle_{\mathbf{K}} &= \langle w^* \hat{\otimes} v^*, \hat{r} \rangle_{\mathbf{K}} = \sum_i \langle w^*, w_i \rangle_{\mathbf{K}} \langle v^*, v_i \rangle_{\mathbf{K}} = \sum_i \langle w^*, \langle v^*, v_i \rangle_{\mathbf{K}} w_i \rangle_{\mathbf{K}} \\ &= \langle w^*, \sum_i \langle v^*, v_i \rangle_{\mathbf{K}} w_i \rangle_{\mathbf{K}} = \langle w^*, r^{\sharp}(v^*) \rangle_{\mathbf{K}}. \end{aligned}$$

So $(\hat{r})^{\sharp}(v^*) = r^{\sharp}(v^*)$ and hence

$$(\hat{r})^{\sharp} \left(\sum_{s=0}^{\infty} v_s^* h^s \right) = \sum_{s=0}^{\infty} r^{\sharp}(v_s^*) h^s, \quad \sum_{s=0}^{\infty} v_s^* h^s \in V_h^*.$$

Thus $(\hat{r})^\sharp$ coincides with the \mathbf{K} -linear extension of r^\sharp , which proves (68). By (68), we have $(\hat{f})^\sharp = ((\tilde{f})^\sharp)_\mathbf{K}$. Then the last conclusion follows from $(\tilde{f})^\sharp = f$. \square

For topologically free \mathbf{K} -modules A_h and V_h , and a \mathbf{K} -linear map $\varrho_h : A_h \rightarrow \text{End}_\mathbf{K}(V_h)$, let $\varrho_h^\star : A_h \rightarrow \text{End}_\mathbf{K}(V_h^\star)$ be the unique \mathbf{K} -linear map such that

$$\langle \varrho_h^\star(x')u^\star, v' \rangle_\mathbf{K} = -\langle u^\star, \varrho_h(x')v' \rangle_\mathbf{K}, \quad x' \in A_h, v' \in V_h, u^\star \in V_h^\star. \quad (70)$$

Let (A_h, \circ_h) be a topologically free associative algebra and $(V_h, l_{\circ_h}, r_{\circ_h})$ be a topologically free (A_h, \circ_h) -bimodule. Then $(V_h^\star, -r_{\circ_h}^\star, -l_{\circ_h}^\star)$ is a topologically free (A_h, \circ_h) -bimodule, called the **topological dual bimodule** of $(V_h, l_{\circ_h}, r_{\circ_h})$.

Proposition 4.7. *Let (A, \circ) be an associative algebra and (V, l_\circ, r_\circ) be an (A, \circ) -bimodule. Let (A_h, \circ_h) be an associative deformation of (A, \circ) and $(V_h, l_{\circ_h}, r_{\circ_h})$ be an (A_h, \circ_h) -bimodule deformation of (V, l_\circ, r_\circ) . Then $(V_h^\star, -r_{\circ_h}^\star, -l_{\circ_h}^\star)$ is an (A_h, \circ_h) -bimodule deformation of the (A, \circ) -bimodule $(V, -r_\circ^\star, -l_\circ^\star)$.*

Proof. Let $x \in A, u_h \in V_h$ and $v^\star \in V^\star$. Then we have

$$\begin{aligned} \langle -r_{\circ_h}^\star(x)v^\star, u_h \rangle_\mathbf{K} &= \langle v^\star, r_{\circ_h}(x)u_h \rangle_\mathbf{K} \\ &\equiv \langle v^\star, r_\circ(x)u \rangle \pmod{h} \\ &\equiv \langle -r_\circ^\star(x)v^\star, u \rangle \pmod{h} \\ &\equiv \langle -r_\circ^\star(x)v^\star, u_h \rangle_\mathbf{K} \pmod{h}, \end{aligned}$$

where u is the element in V such that $u_h \equiv u \pmod{h}$. Then we deduce that $-r_{\circ_h}^\star(x) \equiv -r_\circ^\star(x) \pmod{h}$. Similarly, we can deduce that $-l_{\circ_h}^\star(x) \equiv -l_\circ^\star(x) \pmod{h}$. Thus $(V_h^\star, -r_{\circ_h}^\star, -l_{\circ_h}^\star)$ is an (A_h, \circ_h) -bimodule deformation of the (A, \circ) -bimodule $(V^\star, -r_\circ^\star, -l_\circ^\star)$. \square

The following conclusion shows that the processes of taking the dual modules and taking the QCLs commute.

Proposition 4.8. *Let (A, \circ) be a commutative associative algebra and (V, ρ_\circ) be an (A, \circ) -module. Let (A_h, \circ_h) be an associative deformation of (A, \circ) and $(A, \{, \}, \circ)$ be the Poisson algebra as the QCL. Let $(V_h, l_{\circ_h}, r_{\circ_h})$ be the A_h -bimodule deformation of (V, ρ_\circ) and $(V, \rho_{\{, \}}, \rho_\circ)$ be the $(A, \{, \}, \circ)$ -module as the corresponding QCL. Then the QCL of the A_h -bimodule deformation $(V_h^\star, -r_{\circ_h}^\star, -l_{\circ_h}^\star)$ of the (A, \circ) -module $(V^\star, -\rho_\circ^\star)$ is exactly the $(A, \{, \}, \circ)$ -module $(V^\star, \rho_{\{, \}}^\star, -\rho_\circ^\star)$ from taking the dual module of the $(A, \{, \}, \circ)$ -module $(V, \rho_{\{, \}}, \rho_\circ)$*

Proof. Let $x \in A, u^\star \in V^\star, v_h \in V_h$. Then we have

$$\begin{aligned} \langle -r_{\circ_h}^\star(x)u^\star, v_h \rangle_\mathbf{K} - \langle -l_{\circ_h}^\star(x)u^\star, v_h \rangle_\mathbf{K} &= \langle u^\star, r_{\circ_h}(x)v_h - l_{\circ_h}(x)v_h \rangle_\mathbf{K} \\ &\equiv \langle u^\star, -\rho_{\{, \}}(x)v \rangle h \pmod{h^2} \\ &\equiv \langle \rho_{\{, \}}^\star(x)u^\star, v \rangle h \pmod{h^2} \\ &\equiv \langle \rho_{\{, \}}^\star(x)u^\star, v_h \rangle_\mathbf{K} h \pmod{h^2}, \end{aligned}$$

where v is the element in V such that $v_h \equiv v \pmod{h}$. Hence we obtain

$$\frac{-r_{\circ_h}^\star(x) - (-l_{\circ_h}^\star(x))}{h} \equiv \rho_{\{, \}}^\star(x) \pmod{h}.$$

Thus we conclude that the QCL of the A_h -bimodule deformation $(V_h^\star, -r_{\circ_h}^\star, -l_{\circ_h}^\star)$ is exactly the $(A, \{, \}, \circ)$ -module $(V^\star, \rho_{\{, \}}^\star, -\rho_\circ^\star)$. \square

4.3. AYBE and PYBE via bimodule algebra deformations: tridendriform deformations.

We first recall a basic relation between solutions of the AYBE and \mathcal{O} -operators.

Theorem 4.9. [9, Corollary 3.6] *Let (A, \circ) be a finite-dimensional associative algebra and $r \in A \otimes A$. Let $\beta := (r + \sigma(r))/2$ be (A, \circ) -invariant in the sense of (52). Let r^\sharp and β^\sharp be the images of r and β under $A \otimes A \cong \text{Hom}_{\mathbf{K}}(A^*, A)$ (see (58)). Then*

$$R_{\circ}^*(\beta^\sharp(a^*))b^* = L_{\circ}^*(\beta^\sharp(b^*))a^*, \quad a^*, b^* \in A^*.$$

Moreover, define a binary operation \cdot on A^* by

$$a^* \cdot b^* := 2R_{\circ}^*(\beta^\sharp(a^*))b^*, \quad a^*, b^* \in A^*. \quad (71)$$

Then (A^*, \cdot) is an associative algebra and $(A^*, \cdot, -R_{\circ}^*, -L_{\circ}^*)$ is an (A, \circ) -bimodule algebra. Furthermore, r is a solution of the AYBE in (A, \circ) if and only if r^\sharp is an \mathcal{O} -operator of weight 1 on (A, \circ) associated to $(A^*, \cdot, -R_{\circ}^*, -L_{\circ}^*)$.

Remark 4.10. When the associative algebra (A, \circ) in the above statement is commutative, then (A^*, \cdot) is commutative and $(A^*, \cdot, -R_{\circ}^*)$ is an (A, \circ) -module algebra. Note that in this case, $L_{\circ}^* = R_{\circ}^*$.

The following is the analogue of Theorem 4.9 for topologically free associative algebras and can be proved in the same way (see the proof of [9, Corollary 3.6]).

Theorem 4.11. *Let A be a finite-dimensional vector space and (A_h, \circ_h) be a topologically free associative algebra. Let $r_h \in A_h \hat{\otimes} A_h$ and $\beta_h := (r_h + \sigma(r_h))/2$ be the symmetric part of r_h . Denote the isomorphic images of r_h and β_h in $\text{Hom}_{\mathbf{K}}(A_h^*, A_h)$ by $(r_h)^\sharp$ and $(\beta_h)^\sharp$ respectively. Suppose that β_h is (A_h, \circ_h) -invariant. Then*

$$R_{\circ_h}^*((\beta_h)^\sharp(a^*))b^* = L_{\circ_h}^*((\beta_h)^\sharp(b^*))a^*, \quad a^*, b^* \in A_h^*. \quad (72)$$

Moreover, define a binary operation \cdot_h on A_h^* by

$$a^* \cdot_h b^* := 2R_{\circ_h}^*((\beta_h)^\sharp(a^*))b^*, \quad a^*, b^* \in A_h^*. \quad (73)$$

Then (A_h^*, \cdot_h) is a topologically free associative algebra and $(A_h^*, \cdot_h, -R_{\circ_h}^*, -L_{\circ_h}^*)$ is a topologically free (A_h, \circ_h) -bimodule algebra. Furthermore, r_h is a solution of the AYBE in (A_h, \circ_h) if and only if $(r_h)^\sharp$ is an \mathcal{O} -operator of weight 1 on (A_h, \circ_h) associated to $(A_h^*, \cdot_h, -R_{\circ_h}^*, -L_{\circ_h}^*)$.

The following is the PYBE analogue of Theorem 4.9.

Theorem 4.12. [34] *Let $(A, \{, \}, \circ)$ be a finite-dimensional Poisson algebra and $r \in A \otimes A$. Suppose that $\beta := (r + \sigma(r))/2$ is $(A, \{, \}, \circ)$ -invariant. Then*

$$[a^*, b^*] := -2\text{ad}_{\{, \}}^*(\beta^\sharp(a^*))b^*, \quad a^* \cdot b^* := 2R_{\circ}^*(\beta^\sharp(a^*))b^*, \quad a^*, b^* \in A^*, \quad (74)$$

define a Poisson algebra $(A^*, [,], \cdot)$, and $(A^*, [,], \cdot, \text{ad}_{\{, \}}^*, -R_{\circ}^*)$ is an $(A, \{, \}, \circ)$ -module Poisson algebra. Furthermore, r is a solution of the PYBE in $(A, \{, \}, \circ)$ if and only if r^\sharp is an \mathcal{O} -operator of weight 1 on $(A, \{, \}, \circ)$ associated to the $(A, \{, \}, \circ)$ -module $(A^*, [,], \cdot, \text{ad}_{\{, \}}^*, -R_{\circ}^*)$.

Let (A, \circ) be a finite-dimensional commutative associative algebra, (A_h, \circ_h) be an associative deformation and $(A, \{, \}, \circ)$ be the corresponding QCL. Let r be in $A \otimes A$ and $\beta = (r + \sigma(r))/2$ be its symmetric part. Denote the isomorphic images of r and β in $\text{Hom}_{\mathbf{K}}(A^*, A)$ by r^\sharp and β^\sharp respectively. Denote the isomorphic images of r and β under the linear embedding of $A \otimes A$ into $A_h \hat{\otimes} A_h$ by \hat{r} and $\hat{\beta}$ respectively. Suppose that $\hat{\beta}$ is (A_h, \circ_h) -invariant. By Theorem 4.11, (A_h^*, \cdot_h) is a topologically free associative algebra and $(A_h^*, \cdot_h, -R_{\circ_h}^*, -L_{\circ_h}^*)$ is a topologically free (A_h, \circ_h) -bimodule algebra, where

$$a^* \cdot_h b^* := 2R_{\circ_h}^*((\hat{\beta})^\sharp(a^*))b^*, \quad a^*, b^* \in A_h^*.$$

By Remark 4.5, β is $(A, \{, \}, \circ)$ -invariant. Thus by Remark 4.10, $(A^*, \cdot, -R_{\circ}^*)$ is an (A, \circ) -module algebra, where

$$a^* \cdot b^* := 2R_{\circ}^*(\beta^\sharp(a^*))b^*, \quad a^*, b^* \in A^*.$$

By Theorem 4.12, $(A^*, [,], \cdot, \text{ad}_{\{, \}}^*, -R_{\circ}^*)$ is an $(A, \{, \}, \circ)$ -module Poisson algebra, where

$$[a^*, b^*] := -2\text{ad}_{\{, \}}^*(\beta^\sharp(a^*))b^*, \quad a^* \cdot b^* := 2R_{\circ}^*(\beta^\sharp(a^*))b^*, \quad a^*, b^* \in A^*.$$

Proposition 4.13. *Keep the assumptions and notations as above.*

- (a) *The topologically free (A_h, \circ_h) -bimodule algebra $(A_h^*, \cdot_h, -R_{\circ_h}^*, -L_{\circ_h}^*)$ is an (A_h, \circ_h) -bimodule algebra deformation of the (A, \circ) -module algebra $(A^*, \cdot, -R_{\circ}^*)$, and its QCL is exactly the $(A, \{, \}, \circ)$ -module Poisson algebra $(A^*, [,], \cdot, \text{ad}_{\{, \}}^*, -R_{\circ}^*)$.*
- (b) *Take the downward arrows in the following diagram. Then starting from a solution r of the AYBE in (A, \circ) , the left square in the diagram commutes. Moreover, starting from a solution \hat{r} of the AYBE in (A_h, \circ_h) , the right square in the diagram commutes.*
- (c) *Take the upward arrows in the following diagram. Then starting from the \mathcal{O} -operator r^\sharp of weight 1 on (A, \circ) associated to $(A^*, \cdot, -R_{\circ}^*)$, the left square of the diagram commutes. Moreover, starting from the \mathcal{O} -operator $(\hat{r})^\sharp$ of weight 1 on (A_h, \circ_h) associated to $(A_h^*, \cdot_h, -R_{\circ_h}^*, -L_{\circ_h}^*)$, the right square of the diagram commutes.*

$$\begin{array}{ccccc}
 \begin{array}{c} \text{solution } r \text{ of} \\ \text{AYBE in } (A, \circ), \\ (A, \circ)\text{-invariant } \beta \\ \wedge \\ \text{Theorem 4.9} \end{array} & \begin{array}{c} \text{deformation} \\ > \end{array} & \begin{array}{c} \text{solution } \hat{r} \text{ of} \\ \text{AYBE in } (A_h, \circ_h), \\ (A_h, \circ_h)\text{-invariant } \hat{\beta} \\ \wedge \\ \text{Theorem 4.11} \end{array} & \begin{array}{c} \text{Theorem 4.4} \\ > \end{array} & \begin{array}{c} \text{solution } r \text{ of} \\ \text{PYBE in } (A, \{, \}, \circ), \\ (A, \{, \}, \circ)\text{-invariant } \beta \\ \wedge \\ \text{Theorem 4.12} \end{array} \\
 \\
 \begin{array}{c} \vee \\ \mathcal{O}\text{-operator } r^\sharp \text{ on} \\ (A, \circ) \text{ associated} \\ \text{to } (A^*, \cdot, -R_{\circ}^*) \end{array} & \begin{array}{c} \text{deformation} \\ > \end{array} & \begin{array}{c} \vee \\ \mathcal{O}\text{-operator } (\hat{r})^\sharp = (r^\sharp)_\mathbf{K} \text{ on} \\ (A_h, \circ_h) \text{ associated} \\ \text{to } (A_h^*, \cdot_h, -R_{\circ_h}^*, -L_{\circ_h}^*) \end{array} & \begin{array}{c} \text{Theorem 2.15} \\ \text{Theorem 2.21} \end{array} & \begin{array}{c} \vee \\ \mathcal{O}\text{-operator } r^\sharp \text{ on} \\ (A, \{, \}, \circ) \text{ associated} \\ \text{to } (A^*, [,], \cdot, \text{ad}_{\{, \}}^*, -R_{\circ}^*). \end{array}
 \end{array}$$

Proof. (a). Note that the topologically free (A_h, \circ_h) -bimodule $(A_h^*, -R_{\circ_h}^*, -L_{\circ_h}^*)$ is the topological dual bimodule of the (A_h, \circ_h) -bimodule $(A_h, L_{\circ_h}, R_{\circ_h})$, and $(A_h, L_{\circ_h}, R_{\circ_h})$ is an (A_h, \circ_h) -bimodule deformation of the (A, \circ) -module (A, L_{\circ}) . Then by Proposition 4.7 and $R_{\circ}^* = L_{\circ}^*$, we obtain

$$-R_{\circ_h}^*(x) \equiv -L_{\circ_h}^*(x) \equiv -R_{\circ}^*(x) \pmod{h}, x \in A.$$

By (68), we have $(\hat{\beta})^\sharp = (\beta^\sharp)_\mathbf{K}$. Hence, for $a^*, b^* \in A^*$, we have

$$a^* \cdot_h b^* = 2R_{\circ_h}^*((\hat{\beta})^\sharp(a^*))b^* = 2R_{\circ_h}^*((\beta^\sharp)_\mathbf{K}(a^*))b^* \equiv 2R_{\circ}^*(\beta^\sharp(a^*))b^* \pmod{h} \equiv a^* \cdot b^* \pmod{h}.$$

Thus the topologically free (A, \circ_h) -bimodule algebra $(A_h^*, \cdot_h, -R_{\circ_h}^*, -L_{\circ_h}^*)$ is an (A, \circ_h) -bimodule algebra deformation of the (A, \circ) -module algebra $(A^*, \cdot, -R_{\circ}^*)$.

Let $(A^*, [,], \cdot, \rho'_{\{, \}}, -R_{\circ}^*)$ denote the QCL of $(A_h^*, -R_{\circ_h}^*, -L_{\circ_h}^*)$. By Proposition 4.8, the QCL of the (A, \circ_h) -bimodule deformation $(A_h^*, -R_{\circ_h}^*, -L_{\circ_h}^*)$ is exactly the $(A, \{, \}, \circ)$ -module $(A^*, \cdot, \text{ad}_{\{, \}}^*, -R_{\circ}^*)$.

Thus $\text{ad}_{\{, \}}^* = \rho'_{\{, \}}$. By (72), $R_{\circ_h}^*((\hat{\beta})^\sharp(a^*))b^* = L_{\circ_h}^*((\hat{\beta})^\sharp(b^*))a^*$ for $a^*, b^* \in A_h^*$. Then we deduce

$$\begin{aligned}
 \frac{a^* \cdot_h b^* - b^* \cdot_h a^*}{h} &= \frac{2R_{\circ_h}^*((\hat{\beta})^\sharp(a^*))b^* - 2R_{\circ_h}^*((\hat{\beta})^\sharp(b^*))a^*}{h} \\
 &= \frac{2R_{\circ_h}^*(\beta^\sharp(a^*))b^* - 2L_{\circ_h}^*(\beta^\sharp(a^*))b^*}{h} \\
 &\equiv -2\text{ad}_{\{, \}}^*(\beta(a^*))b^* \pmod{h}, \\
 &\equiv [a^*, b^*] \pmod{h} \quad a^*, b^* \in A^*.
 \end{aligned}$$

Hence $[a^*, b^*] = [a^*, b^*]'$. Therefore, the two $(A, \{, \}, \circ)$ -module Poisson algebras $(A^*, [,], \cdot, \text{ad}_{\{, \}}^*, -R_{\circ}^*)$ and $(A^*, [,], \cdot, \rho'_{\{, \}}, -R_{\circ}^*)$ coincide.

Since $(\hat{r})^\sharp$ coincides with the scalar deformation $(r^\sharp)_\mathbf{K}$ of r^\sharp by Proposition 4.6, the statements in (b) and (c) follow. \square

Tridendriform algebras correspond to A -bimodule algebras equipped with the \mathcal{O} -operator id_A of weight 1 and hence give rise to solutions of the AYBE as in the following theorem. First for vector spaces V and W , we have the canonical embedding

$$\eta : V \otimes W \rightarrow (V \oplus W) \otimes (V \oplus W), \quad \sum_i v_i \otimes w_i \mapsto \sum_i (v_i, 0) \otimes (0, w_i) \in (V \oplus W) \otimes (V \oplus W). \quad (75)$$

Similarly, we can embed $V_h \hat{\otimes} W_h$ into $(V_h \oplus W_h) \hat{\otimes} (V_h \oplus W_h)$. Then identifying $V_h \oplus W_h$ with $(V \oplus W)_h$ by Remark 2.10, we obtain the following \mathbf{K} -linear embedding:

$$\eta_h : V_h \hat{\otimes} W_h \rightarrow (V \oplus W)_h \hat{\otimes} (V \oplus W)_h, \quad \sum_{s \geq 0} v_s h^s \hat{\otimes} \sum_{l \geq 0} w_l h^l \mapsto \sum_{s, l \geq 0} (v_s, 0) h^s \hat{\otimes} (0, w_l) h^l. \quad (76)$$

Theorem 4.14. [10, Corollary 6.19] *Let (A, \succ, \prec, \cdot) be a finite-dimensional tridendriform algebra and $A^{\oplus 2}$ be the underlying vector space of the associative algebra $(\hat{A} := A \rtimes_{L_\succ, R_\prec} A, \odot)$. Define the operators $\alpha_1, \alpha_2, \alpha_3, \alpha_4 \in \text{End}_{\mathbf{K}}(A^{\oplus 2})$ by*

$$\alpha_1(x, y) = (-x + y, 0), \quad \alpha_2(x, y) = -(y, y), \quad \alpha_3(x, y) = -(x, x), \quad \alpha_4(x, y) = (0, x - y), \quad x, y \in A. \quad (77)$$

Denote the identity operator on $A^{\oplus 2}$ by id . For $i \in \{1, 2, 3, 4\}$, embed $\tilde{\alpha}_i$ and $\tilde{\text{id}}$ into $(A^{\oplus 2} \oplus (A^{\oplus 2})^)^{\otimes 2}$ by the embedding η in (75). Then $\eta(\tilde{\alpha}_i) - \sigma(\eta(\tilde{\alpha}_i)) + \eta(\tilde{\text{id}})$ and $\eta(\tilde{\alpha}_i) - \sigma(\eta(\tilde{\alpha}_i)) - \sigma(\eta(\tilde{\text{id}}))$ are solutions of the AYBE in the associative algebra $\hat{A} \rtimes_{-R_\odot^*, -L_\odot^*} (A^{\oplus 2})^*$.*

Remark 4.15. The above constructions of solutions of the AYBE are special cases of more general results where the constructions of solutions of the AYBE utilize Rota-Baxter operators (see [10, Proposition 3.9(e)]). In fact, since id_A is an invertible \mathcal{O} -operator on (A, \circ) of weight 1 associated to $(A, \cdot, L_\succ, R_\prec)$, each α_i is a Rota-Baxter operator of weight 1 on $(\hat{A} = A \rtimes_{L_\succ, R_\prec} A, \odot)$ (see [10, Propositions 6.1 and 6.8]).

The following result gives the topologically free tridendriform algebra analogue of Theorem 4.14 and can be proved in a similar manner (see the proof of [10, Corollary 6.19]).

Theorem 4.16. *Let A be a finite-dimensional vector space and $(A_h, \succ_h, \prec_h, \cdot_h)$ be a topologically free tridendriform algebra. Denote the underlying topologically free \mathbf{K} -module of the topologically free associative algebra $\hat{A}_h := (A_h \rtimes_{L_{\succ_h}, R_{\prec_h}} A_h, \odot_h)$ by $(A^{\oplus 2})_h$. Let $\alpha_{1,h}, \alpha_{2,h}, \alpha_{3,h}, \alpha_{4,h} \in \text{End}_{\mathbf{K}}((A^{\oplus 2})_h)$ be the operators defined by*

$$\begin{aligned} \alpha_{1,h}(x_h, y_h) &= (-x_h + y_h, 0), \quad \alpha_{2,h}(x_h, y_h) = -(y_h, y_h), \\ \alpha_{3,h}(x_h, y_h) &= -(x_h, x_h), \quad \alpha_{4,h}(x_h, y_h) = (0, x_h - y_h), \quad x_h, y_h \in A_h. \end{aligned}$$

Denote the identity operator on $(A^{\oplus 2})_h$ by id_h . For $i \in \{1, 2, 3, 4\}$, embed $\widetilde{\alpha}_{i,h}$ and $\widetilde{\text{id}}_h$ into $(A^{\oplus 2} \oplus (A^{\oplus 2})^)_h \hat{\otimes} (A^{\oplus 2} \oplus (A^{\oplus 2})^*)_h$ by the embedding η_h in (76). Then*

$$\eta_h(\widetilde{\alpha}_{i,h}) - \sigma_{\mathbf{K}}(\eta_h(\widetilde{\alpha}_{i,h})) + \eta_h(\widetilde{\text{id}}_h) \quad \text{and} \quad \eta_h(\widetilde{\alpha}_{i,h}) - \sigma_{\mathbf{K}}(\eta_h(\widetilde{\alpha}_{i,h})) - \sigma_{\mathbf{K}}(\eta_h(\widetilde{\text{id}}_h))$$

are solutions of the AYBE in $\hat{A}_h \rtimes_{-R_{\odot_h}^, -L_{\odot_h}^*} (A^{\oplus 2})_h^*$.*

Let $(\mathfrak{g}, \{, \})$ be a Lie algebra. Taking the operad of Lie algebras in Definition 2.1, we recover the notion of a \mathfrak{g} -module Lie algebra [8, 30]. Let $(\mathfrak{h}, [,], \rho_{\{, \}})$ be a \mathfrak{g} -module Lie algebra. Then taking the operad of Lie algebras in Definition 2.17, we recover the notion of an \mathcal{O} -operator of weight λ on \mathfrak{g} associated to $(\mathfrak{h}, [,], \rho_{\{, \}})$ [8]. A linear map $T \in \text{End}_{\mathbf{K}}(\mathfrak{g})$ is called a Rota-Baxter operator of weight λ on \mathfrak{g} if it satisfies (16).

Theorem 4.17. *Let $(A, [,], \triangleright, \succ, \cdot)$ be a finite-dimensional post-Poisson algebra. Denote the underlying vector space of the Poisson algebra $(\check{A} := A \rtimes_{L_\triangleright, L_\succ} A, \llbracket, \rrbracket, \odot)$ by $A^{\oplus 2}$. Let $\alpha_i, \text{id}, \tilde{\alpha}_i, \tilde{\text{id}}$ and η be as defined in Theorem 4.14. Then $\eta(\tilde{\alpha}_i) - \sigma(\eta(\tilde{\alpha}_i)) + \eta(\tilde{\text{id}})$ and $\eta(\tilde{\alpha}_i) - \sigma(\eta(\tilde{\alpha}_i)) - \sigma(\eta(\tilde{\text{id}}))$ are solutions of the PYBE in the Poisson algebra $\check{A} \rtimes_{\text{ad}_{\llbracket, \rrbracket}^*, -R_\odot^*} (A^{\oplus 2})^*$.*

Proof. Since (A, \succ, \cdot) is a finite-dimensional tridendriform algebra, Theorem 4.14 shows that $\eta(\tilde{\alpha}_i) - \sigma(\eta(\tilde{\alpha}_i)) + \eta(\tilde{\text{id}})$ and $\eta(\tilde{\alpha}_i) - \sigma(\eta(\tilde{\alpha}_i)) - \sigma(\eta(\tilde{\text{id}}))$ are solutions of the AYBE in $(A \rtimes_{L_\succ} A) \rtimes_{-R_\odot^*} (A^{\oplus 2})^*$.

By Proposition 3.7, $(A, [,], L_\triangleright)$ is an $(A, \{, \})$ -module Lie algebra for $\{, \}$ defined in (31), and id_A is an \mathcal{O} -operator of weight 1 on $(A, \{, \})$ associated to $(A, [,], L_\triangleright)$. Thus by [8, Proposition 4.1], α_1 and α_2 in (77) are Rota-Baxter operators of weight 1 on the Lie algebra $A \ltimes_{L_\triangleright} A$ respectively. It is straightforward to show that α_3 and α_4 are also Rota-Baxter operators of weight 1 on the Lie algebra $A \ltimes_{L_\triangleright} A$ respectively. That is, α_i is a Rota-Baxter operator of weight 1 on the Lie algebra $A \ltimes_{L_\triangleright} A$ for any $\alpha_i \in \{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$. Thus by [8, Corollary 2.16 (d)], $\eta(\tilde{\alpha}_i) - \sigma(\eta(\tilde{\alpha}_i)) + \eta(\widetilde{\text{id}})$ and $\eta(\tilde{\alpha}_i) - \sigma(\eta(\tilde{\alpha}_i)) - \sigma(\eta(\widetilde{\text{id}}))$ are also solutions of the CYBE in the Lie algebra $(A \ltimes_{L_\triangleright} A) \ltimes_{\text{ad}_{[\cdot, \cdot]}}^* (A^{\oplus 2})^*$. \square

Lemma 4.18. *Let A be a finite-dimensional vector space and (A_h, \circ_h) be a topologically free associative algebra. Denote the isomorphic image of id_A in $A^* \otimes A$ by $\widetilde{\text{id}}_A$ and embed it to $(A \oplus A^*)^{\otimes 2}$ by η in (75). Denote the isomorphic image of $\eta(\widetilde{\text{id}}_A)$ under the linear embedding of $(A \oplus A^*)^{\otimes 2}$ into $(A \oplus A^*)_h \hat{\otimes} (A \oplus A^*)_h$ by $\widehat{\eta}(\widetilde{\text{id}}_A)$. Then the symmetric part of $\widehat{\eta}(\widetilde{\text{id}}_A)$ is $A_h \ltimes_{-R_{\circ_h}^*, -L_{\circ_h}^*} A_h^*$ -invariant.*

Proof. Let $\{e_1, \dots, e_n\}$ be a basis of A and $\{e_1^*, \dots, e_n^*\}$ be the dual basis of A^* , where $n = \dim A$. Then

$$\widehat{\eta}(\widetilde{\text{id}}_A) + \sigma_{\mathbf{K}}(\widehat{\eta}(\widetilde{\text{id}}_A)) = \sum_{i=1}^n (e_i, 0) \hat{\otimes} (0, e_i^*) + \sum_{i=1}^n (0, e_i^*) \hat{\otimes} (e_i, 0).$$

Denote the identity operator on $A_h \oplus A_h^*$ by Id and let \circ_h be the associative product of the associative algebra $A_h \ltimes_{-R_{\circ_h}^*, -L_{\circ_h}^*} A_h^*$. Let $x_h \in A_h, a^* \in A_h^*$. Then we have

$$\begin{aligned} & \text{Id} \hat{\otimes} L_{\circ_h}((x_h, a^*)) \sum_{i=1}^n (e_i, 0) \hat{\otimes} (0, e_i^*) - R_{\circ_h}((x_h, a^*)) \hat{\otimes} \text{Id} \sum_{i=1}^n (e_i, 0) \hat{\otimes} (0, e_i^*) \\ &= \sum_{i=1}^n (e_i, 0) \hat{\otimes} (0, -R_{\circ_h}^*(x_h)e_i^*) - \sum_{i=1}^n (e_i \circ_h x_h, -R_{\circ_h}^*(e_i)a^*) \hat{\otimes} (0, e_i^*) \\ &= \sum_{i=1}^n (e_i, 0) \hat{\otimes} (0, -R_{\circ_h}^*(x_h)e_i^*) - \sum_{i=1}^n \sum_{j=1}^n \langle e_j^*, e_i \circ_h x_h \rangle_{\mathbf{K}} (e_j, 0) \hat{\otimes} (0, e_i^*) + \sum_{i=1}^n (0, \sum_{j=1}^n \langle a^*, -e_j \circ_h e_i \rangle_{\mathbf{K}} e_j^*) \hat{\otimes} (0, e_i^*) \\ &= \sum_{i=1}^n (e_i, 0) \hat{\otimes} (0, -R_{\circ_h}^*(x_h)e_i^*) + \sum_{i=1}^n \sum_{j=1}^n \langle R_{\circ_h}^*(x_h)e_j^*, e_i \rangle_{\mathbf{K}} (e_j, 0) \hat{\otimes} (0, e_i^*) + \sum_{i=1}^n (0, \sum_{j=1}^n \langle L_{\circ_h}^*(e_j)a^*, e_i \rangle_{\mathbf{K}} e_j^*) \otimes (0, e_i^*) \\ &= \sum_{i=1}^n (e_i, 0) \hat{\otimes} (0, -R_{\circ_h}^*(x_h)e_i^*) + \sum_{j=1}^n (e_j, 0) \hat{\otimes} (0, R_{\circ_h}^*(x_h)e_j^*) + \sum_{j=1}^n (0, e_j^*) \hat{\otimes} (0, L_{\circ_h}^*(e_j)a^*) \\ &= \sum_{j=1}^n (0, e_j^*) \hat{\otimes} (0, L_{\circ_h}^*(e_j)a^*). \end{aligned}$$

Similarly, we obtain

$$\text{Id} \hat{\otimes} L_{\circ_h}((x_h, a^*)) \sum_{i=1}^n (0, e_i^*) \hat{\otimes} (e_i, 0) - R_{\circ_h}((x_h, a^*)) \hat{\otimes} \text{Id} \sum_{i=1}^n (0, e_i^*) \hat{\otimes} (e_i, 0) = \sum_{i=1}^n (0, e_i^*) \hat{\otimes} (0, -L_{\circ_h}^*(e_i)a^*).$$

Thus $\widehat{\eta}(\widetilde{\text{id}}_A) + \sigma_{\mathbf{K}}(\widehat{\eta}(\widetilde{\text{id}}_A))$ is $A_h \ltimes_{-R_{\circ_h}^*, -L_{\circ_h}^*} A_h^*$ -invariant and hence the conclusion holds. \square

Let $(A, >, \cdot)$ be a finite-dimensional commutative tridendriform algebra, $(A, >_h, <_h, \cdot_h)$ be a tridendriform deformation and $(A, \triangleright, [,], >, \cdot)$ be its QCL which is a post-Poisson algebra. Define binary operations \circ and $\{, \}$ on A by (31). Define a binary operation \circ_h on A_h by (35). Thus by Proposition 3.16, the topologically free (A_h, \circ_h) -bimodule algebra $(A_h, \cdot_h, L_{>_h}, R_{<_h})$ is a deformation of the (A, \circ) -module algebra $(A, \cdot, L_{>})$ and its QCL is the $(A, \{, \}, \circ)$ -module Poisson algebra $(A, [,], \cdot, L_{\triangleright}, L_{>})$. So we have the following data:

- two commutative associative algebras $(\hat{A} := A \ltimes_{L_{>}} A, \odot)$ and $\hat{A} \ltimes_{-R_{\odot}^*} (A^{\oplus 2})^*$;
- two topologically free associative algebras $(\hat{A}_h := A_h \ltimes_{L_{>_h}, R_{<_h}} A_h, \odot_h)$ and $\hat{A}_h \ltimes_{-R_{\odot_h}^*, -L_{\odot_h}^*} (A^{\oplus 2})_h^*$;
- two Poisson algebras $(\check{A} := A \ltimes_{L_{>}, L_{>}} A, \llbracket, \rrbracket, \odot)$ and $\check{A} \ltimes_{\text{ad}_{\llbracket, \rrbracket}^*, -R_{\odot}^*} (A^{\oplus 2})^*$,

where $A^{\oplus 2}$ is the underlying vector space of $A \ltimes_{L_{>}} A$. Fix an $i \in \{1, 2, 3, 4\}$. Let α_i, id and η be the linear maps and $\tilde{\alpha}_i$ and id be tensors given in Theorem 4.14. Let $\alpha_{i,h}, \text{id}_h$ and η_h be the \mathbf{K} -linear maps and $\tilde{\alpha}_{i,h}$ and id_h be tensors given in Theorem 4.16. Let

$$r \in \{\eta(\tilde{\alpha}_i) - \sigma(\eta(\tilde{\alpha}_i)) + \eta(\text{id}), \eta(\tilde{\alpha}_i) - \sigma(\eta(\tilde{\alpha}_i)) - \sigma(\eta(\text{id}))\}$$

and denote the isomorphic image of r in $(A^{\oplus 2} \oplus (A^{\oplus 2})^*)_h \hat{\otimes} (A^{\oplus 2} \oplus (A^{\oplus 2})^*)_h^*$ by \hat{r} . It is clear that \hat{r} is in $\{\widehat{\eta(\tilde{\alpha}_i) - \sigma(\eta(\tilde{\alpha}_i)) + \eta(\text{id})}, \widehat{\eta(\tilde{\alpha}_i) - \sigma(\eta(\tilde{\alpha}_i)) - \sigma(\eta(\text{id}))}\}$.

Proposition 4.19. *Retain the above assumptions and notations.*

- The topologically free associative algebra $\hat{A}_h \ltimes_{-R_{\odot_h}^*, -L_{\odot_h}^*} (A^{\oplus 2})_h^*$ is an associative deformation of the commutative associative algebra $\hat{A} \ltimes_{-R_{\odot}^*} (A^{\oplus 2})^*$ and the corresponding QCL is exactly the Poisson algebra $\check{A} \ltimes_{\text{ad}_{\llbracket, \rrbracket}^*, -R_{\odot}^*} (A^{\oplus 2})^*$.
- Starting with a commutative tridendriform algebra $(A, >, \cdot)$, the left square of the following diagram commutes.
- Starting from a topologically free tridendriform algebra $(A_h, >_h, <_h, \cdot_h)$, the right square of the following diagram commutes.

commutative tridendriform algebra $(A, >, \cdot)$	deformation	topologically free >tridendriform algebra $(A_h, >_h, <_h, \cdot_h)$	QCL Theorem 3.11	>	post-Poisson algebra $(A, \{, \}, \triangleright, >, \cdot)$
Theorem 4.14 $r = \eta(\tilde{\alpha}_i) - \sigma(\eta(\tilde{\alpha}_i)) + \eta(\text{id})$ $(r = \eta(\tilde{\alpha}_i) - \sigma(\eta(\tilde{\alpha}_i)) - \sigma(\eta(\text{id})))$		Theorem 4.16 $r_h = \eta_h(\tilde{\alpha}_{i,h}) - \sigma_{\mathbf{K}}(\eta_h(\tilde{\alpha}_{i,h})) + \eta_h(\text{id}_h)$ $(r_h = \eta_h(\tilde{\alpha}_{i,h}) - \sigma_{\mathbf{K}}(\eta_h(\tilde{\alpha}_{i,h})) - \eta_h(\text{id}_h))$			Theorem 4.17 $r = \eta(\tilde{\alpha}_i) - \sigma(\eta(\tilde{\alpha}_i)) + \eta(\text{id})$ $(r = \eta(\tilde{\alpha}_i) - \sigma(\eta(\tilde{\alpha}_i)) - \sigma(\eta(\text{id})))$
solution r of AYBE in $\hat{A} \ltimes_{-R_{\odot}^*} (A^{\oplus 2})^*$	deformation	solution r_h of AYBE in $\hat{A}_h \ltimes_{-R_{\odot_h}^*, -L_{\odot_h}^*} (A^{\oplus 2})_h^*$	Theorem 4.4 Lemma 4.18	>	solution r of PYBE in $\check{A} \ltimes_{\text{ad}_{\llbracket, \rrbracket}^*, -R_{\odot}^*} (A^{\oplus 2})^*$

Proof. (a) It is clear that $A_h \ltimes_{L_{>_h}, R_{<_h}} A_h$ is an associative deformation of $A \ltimes_{L_{>}} A$ since the topologically free (A_h, \circ_h) -bimodule algebra $(A_h, \cdot_h, L_{>_h}, R_{<_h})$ is a deformation of the (A, \circ) -module algebra $(A, \cdot, L_{>})$. Thus the topologically free \hat{A}_h -bimodule $((A^{\oplus 2})_h^*, -R_{\odot_h}^*, -L_{\odot_h}^*)$ which is the topological dual bimodule of the \hat{A}_h -bimodule $((A^{\oplus 2})_h, L_{\odot_h}, R_{\odot_h})$ is a deformation of the \hat{A} -module $((A^{\oplus 2})^*, -R_{\odot}^*)$ which is the dual module of the \hat{A} -module $(A^{\oplus 2}, R_{\odot})$ by Proposition 4.7. Therefore, $\hat{A}_h \ltimes_{-R_{\odot_h}^*, -L_{\odot_h}^*} (A^{\oplus 2})_h^*$ is an associative deformation of $\hat{A} \ltimes_{-R_{\odot}^*} (A^{\oplus 2})^*$.

By Theorem 2.15, the QCL of the associative deformation $A_h \ltimes_{L_{>_h}, R_{<_h}} A_h$ of the commutative associative algebra $A \ltimes_{L_{>}} A$ is exactly the Poisson algebra $A \ltimes_{L_{>}, L_{>}} A$. Then the QCL of the \hat{A}_h -bimodule $((A^{\oplus 2})_h, L_{\odot_h}, R_{\odot_h})$ is the \check{A} -module $(A^{\oplus 2}, \text{ad}_{\llbracket, \rrbracket}, R_{\odot})$. Thus by Proposition 4.8, the QCL corresponding to the \hat{A}_h -bimodule $((A^{\oplus 2})_h^*, -R_{\odot_h}^*, -L_{\odot_h}^*)$ is exactly the \check{A} -module $((A^{\oplus 2})^*, \text{ad}_{\llbracket, \rrbracket}^*, -R_{\odot}^*)$. We conclude that the corresponding QCL of $\hat{A}_h \ltimes_{-R_{\odot_h}^*, -L_{\odot_h}^*} (A^{\oplus 2})_h^*$ is exactly the Poisson algebra $\check{A} \ltimes_{\text{ad}_{\llbracket, \rrbracket}^*, -R_{\odot}^*} (A^{\oplus 2})^*$ by Corollary 2.16.

Note that

$$\alpha_{i,h}((x, y)) = \alpha_i((x, y)) = (\alpha_i)_{\mathbf{K}}((x, y)), \quad \forall x, y \in A.$$

Since $\alpha_{i,h}$ and $(\alpha_i)_{\mathbf{K}}$ are in $\text{End}_{\mathbf{K}}((A^{\oplus 2})_h)$ and their restrictions to $A^{\oplus 2}$ coincide, we conclude that $\alpha_{i,h}$ coincides with $(\alpha_i)_{\mathbf{K}}$. Thus we obtain $\widetilde{\alpha}_{i,h} = \widehat{\alpha}_i$ by (69). Let $\{e_1, \dots, e_n\}$ be a basis of $A^{\oplus 2}$ and $\{e_1^*, \dots, e_n^*\}$ be the dual basis of $(A^{\oplus 2})^*$, where $n = \dim A^{\oplus 2}$. So $\widetilde{\alpha}_i = \sum_{j=1}^n \alpha_i(e_j) \otimes e_j^*$. Then

$$\eta_h(\widetilde{\alpha}_{i,h}) = \eta_h(\widehat{\alpha}_i) = \sum_{j=1}^n (\alpha_i(e_j), 0) \widehat{\otimes} (0, e_j^*) = \widehat{\eta}(\widehat{\alpha}_i). \quad (78)$$

Similarly, we have

$$\eta_h(\widetilde{\text{id}}_h) = \widehat{\eta}(\widehat{\text{id}}). \quad (79)$$

Also note that

$$\sigma_{\mathbf{K}}(\eta_h(\widetilde{\alpha}_{i,h})) = \sigma_{\mathbf{K}}(\eta_h(\widehat{\alpha}_i)) = \sum_{j=1}^n (0, e_j^*) \widehat{\otimes} (\alpha_i(e_j), 0) = \sigma(\widehat{\eta}(\widehat{\alpha}_i)). \quad (80)$$

Similarly, we obtain $\sigma_{\mathbf{K}}(\eta_h(\widetilde{\text{id}}_h)) = \sigma(\widehat{\eta}(\widehat{\text{id}}))$. Thus we have

$$\begin{aligned} \eta_h(\widetilde{\alpha}_{i,h}) - \sigma_{\mathbf{K}}(\eta_h(\widetilde{\alpha}_{i,h})) + \eta_h(\widetilde{\text{id}}_h) &= \widehat{\eta}(\widehat{\alpha}_i) - \sigma(\widehat{\eta}(\widehat{\alpha}_i)) + \widehat{\eta}(\widehat{\text{id}}), \\ \eta_h(\widetilde{\alpha}_{i,h}) - \sigma_{\mathbf{K}}(\eta_h(\widetilde{\alpha}_{i,h})) - \sigma_{\mathbf{K}}(\eta_h(\widetilde{\text{id}}_h)) &= \widehat{\eta}(\widehat{\alpha}_i) - \sigma(\widehat{\eta}(\widehat{\alpha}_i)) - \sigma(\widehat{\eta}(\widehat{\text{id}})). \end{aligned}$$

We conclude that r_h coincides with the scalar deformation \widehat{r} of r . The symmetric parts of $\eta_h(\widetilde{\alpha}_{i,h}) - \sigma_{\mathbf{K}}(\eta_h(\widetilde{\alpha}_{i,h})) + \eta_h(\widetilde{\text{id}}_h)$ and $\eta_h(\widetilde{\alpha}_{i,h}) - \sigma_{\mathbf{K}}(\eta_h(\widetilde{\alpha}_{i,h})) - \sigma_{\mathbf{K}}(\eta_h(\widetilde{\text{id}}_h))$ are

$$\frac{\eta_h(\widetilde{\text{id}}_h) + \sigma_{\mathbf{K}}(\eta_h(\widetilde{\text{id}}_h))}{2} \quad \text{and} \quad -\frac{\eta_h(\widetilde{\text{id}}_h) + \sigma_{\mathbf{K}}(\eta_h(\widetilde{\text{id}}_h))}{2}$$

respectively. By (79), we have

$$\frac{\eta_h(\widetilde{\text{id}}_h) + \sigma_{\mathbf{K}}(\eta_h(\widetilde{\text{id}}_h))}{2} = \frac{\widehat{\eta}(\widehat{\text{id}}) + \sigma_{\mathbf{K}}(\widehat{\eta}(\widehat{\text{id}}))}{2}.$$

So we deduce that the symmetric part of r_h is $\widehat{A}_h \ltimes_{-R_{\circ_h}^*, -L_{\circ_h}^*} (A^{\oplus 2})_h^*$ -invariant by applying Lemma 4.18 to the topologically free associative algebra \widehat{A}_h . Then it is clear that (b) and (c) hold. \square

4.4. AYBE and PYBE via bimodule deformations: dendriform deformations. As a special case, an A -bimodule is an A -bimodule algebra equipped with the trivial operation. So we consider the special case of the previous subsection when A -bimodule algebras are replaced by A -bimodules. Note that in this case, accordingly, tridendriform algebras are replaced by dendriform algebras and the solutions of AYBE and PYBE are skew-symmetric.

Taking the tensor r in Proposition 4.13 to be skew-symmetric leads to the following commutative diagram.

$$\begin{array}{ccccc} \text{skew-symmetric} & & \text{skew-symmetric} & & \text{skew-symmetric} \\ \text{solution } r \text{ of} & & \text{solution } \widehat{r} \text{ of} & & \text{solution } r \text{ of} \\ \text{AYBE in } (A, \circ) & \xrightarrow{\text{deformation}} & \text{AYBE in } (A_h, \circ_h) & \xrightarrow{\text{deformation}} & \text{PYBE in } (A, \{\cdot, \cdot\}, \circ) \\ \downarrow \vee & & \downarrow \vee & & \downarrow \vee \\ \text{\textcircled{O}}\text{-operator } r^\# \text{ on} & & \text{\textcircled{O}}\text{-operator } (\widehat{r})^\# = (r^\#)_{\mathbf{K}} \text{ on} & & \text{\textcircled{O}}\text{-operator } r^\# \text{ on} \\ (A, \circ) \text{ associated} & \xrightarrow{\text{deformation}} & (A_h, \circ_h) \text{ associated} & \xrightarrow{\text{deformation}} & (A, \{\cdot, \cdot\}, \circ) \text{ associated} \\ \text{to } (A^*, -R_\circ^*) & & \text{to } (A_h^*, -R_{\circ_h}^*, -L_{\circ_h}^*) & & \text{to } (A^*, \text{ad}_{\{\cdot, \cdot\}}^*, -R_\circ^*). \end{array}$$

There is a general construction of solutions of skew-symmetric solutions of AYBE or PYBE from arbitrary $\text{\textcircled{O}}$ -operators.

Theorem 4.20. [9] *Let (A, \circ) be a finite-dimensional associative algebra and (V, l_\circ, r_\circ) be a finite-dimensional A -bimodule. Let $T : V \rightarrow A$ be a linear map. Denote its isomorphic image in $A \otimes V^*$ by \widetilde{T} and embed \widetilde{T} into $(A \oplus V^*)^{\otimes 2}$ by η defined by (75). Then $r = \eta(\widetilde{T}) - \sigma(\eta(\widetilde{T}))$ is a skew-symmetric solution of the AYBE in the associative algebra $A \ltimes_{-r_\circ^*, -l_\circ^*} V^*$ if and only if T is an $\text{\textcircled{O}}$ -operator on the associative algebra (A, \circ) associated to the (A, \circ) -bimodule (V, l_\circ, r_\circ) .*

The topologically free associative algebra analogue of the above conclusion is given as follows by omitting the proof.

Theorem 4.21. *Let A and V be finite-dimensional vector spaces. Let (A_h, \circ_h) be a topologically free associative algebra and $(V_h, l_{\circ_h}, r_{\circ_h})$ be a topologically free A_h -bimodule. Let $T_h \in \text{Hom}_{\mathbf{K}}(V_h, A_h)$ be a \mathbf{K} -linear map. Embed $\tilde{T}_h \in A_h \otimes V_h^*$ into $(A \oplus V^*)_h \hat{\otimes} (A \oplus V^*)_h$ by η_h defined by (76). Set that $r_h = \eta_h(\tilde{T}_h) - \sigma_{\mathbf{K}}(\eta_h(\tilde{T}_h))$. Then r_h is a skew-symmetric solution of AYBE in $A_h \ltimes_{-r_{\circ_h}^*, -l_{\circ_h}^*} V_h^*$ if and only if T_h is an \mathcal{O} -operator on (A_h, \circ_h) associated to $(V_h, l_{\circ_h}, r_{\circ_h})$.*

There is also a Poisson algebra analogue of Theorem 4.20 as follows.

Theorem 4.22. [25] *Let $(A, \{, \}, \circ)$ be a finite-dimensional Poisson algebra and $(V, \rho_{\{, \}}, \rho_{\circ})$ be a finite-dimensional $(A, \{, \}, \circ)$ -module. Let $T : V \rightarrow A$ be a linear map. Denote its isomorphic image in $A \otimes V^*$ by \tilde{T} and embed \tilde{T} into $(A \oplus V^*)^{\otimes 2}$ by η defined by (75). Then $r = \eta(\tilde{T}) - \sigma(\eta(\tilde{T}))$ is a skew-symmetric solution of the PYBE in the Poisson algebra $A \ltimes_{\rho_{\{, \}}, -\rho_{\circ}^*} V^*$ if and only if T is an \mathcal{O} -operator on the Poisson algebra $(A, \{, \}, \circ)$ associated to the $(A, \{, \}, \circ)$ -module $(V, \rho_{\{, \}}, \rho_{\circ})$.*

Let (A, \circ) be a finite-dimensional commutative associative algebra and (V, ρ_{\circ}) be a finite dimensional (A, \circ) -module. Let (A_h, \circ_h) be an associative deformation of (A, \circ) and $(A, \{, \}, \circ)$ be the corresponding QCL. Let $(V_h, l_{\circ_h}, r_{\circ_h})$ be an A_h -bimodule deformation of (V, ρ_{\circ}) and $(V, \rho_{\{, \}}, \rho_{\circ})$ be the corresponding QCL, which is an $(A, \{, \}, \circ)$ -module by Corollary 2.16. Then we have the following data:

- the (A, \circ) -module (V^*, ρ_{\circ}^*) which is the dual module of the (A, \circ) -module (V, ρ_{\circ}) and a commutative associative algebra $A \ltimes_{-\rho_{\circ}} V^*$;
- the topologically free (A_h, \circ_h) -bimodule $(V_h^*, -r_{\circ_h}^*, -l_{\circ_h}^*)$ which is the topological dual module of the (A_h, \circ_h) -bimodule $(V_h, l_{\circ_h}, r_{\circ_h})$ and a topologically free associative algebra $A_h \ltimes_{-r_{\circ_h}^*, -l_{\circ_h}^*} V_h^*$;
- the $(A, \{, \}, \circ)$ -module $(V^*, \rho_{\{, \}}^*, -\rho_{\circ}^*)$ which is the dual module of the $(A, \{, \}, \circ)$ -module $(V, \rho_{\{, \}}, \rho_{\circ})$. and a Poisson algebra $A \ltimes_{\rho_{\{, \}}, -\rho_{\circ}^*} V^*$.

Let $T \in \text{Hom}_{\mathbf{K}}(V, A)$ and $T_{\mathbf{K}}$ be defined by (18). Embed $\tilde{T} \in A \otimes V^*$ into $(A \oplus V^*)^{\otimes 2}$ via η defined in (75). Then set $r := \eta(\tilde{T}) - \sigma(\eta(\tilde{T}))$. So r is skew-symmetric. Denote the isomorphic images of \tilde{T} and r in $(A \oplus V^*)_h \hat{\otimes} (A \oplus V^*)_h$ by $\hat{\tilde{T}}$ and \hat{r} respectively. We obtain $\hat{r} = \widehat{\eta(\tilde{T})} - \widehat{\sigma(\eta(\tilde{T}))}$. Embed $\widetilde{T_{\mathbf{K}}} \in A_h \otimes V_h^*$ into $(A \oplus V^*)_h \hat{\otimes} (A \oplus V^*)_h$ via η_h defined in (76). Set

$$r_h := \eta_h(\widetilde{T_{\mathbf{K}}}) - \sigma_{\mathbf{K}}(\eta_h(\widetilde{T_{\mathbf{K}}}))$$

Proposition 4.23. *Keep the above assumptions and notations. We have the following results.*

- The topologically free associative algebra $A_h \ltimes_{-r_{\circ_h}^*, -l_{\circ_h}^*} V_h^*$ is an associative deformation of the commutative associative algebra $A \ltimes_{-\rho_{\circ}} V^*$ and the corresponding QCL is exactly the Poisson algebra $A \ltimes_{\rho_{\{, \}}, -\rho_{\circ}^*} V^*$.*
- Assume T and $T_{\mathbf{K}}$ are \mathcal{O} -operators on (A, \circ) and (A_h, \circ_h) associated to $(V, l_{\circ}, r_{\circ})$ and $(V_h, l_{\circ_h}, r_{\circ_h})$ respectively. Starting with an \mathcal{O} -operator T on (A, \circ) associated to $(V, l_{\circ}, r_{\circ})$, the left square of the following diagram commutes. Moreover, starting from the \mathcal{O} -operator $T_{\mathbf{K}}$ on (A_h, \circ_h) associated to $(V_h, l_{\circ_h}, r_{\circ_h})$, the right square of the diagram commutes.*

\mathcal{O} -operator T on (A, \circ) associated to (V, ρ_\circ) <i>Theorem 4.20</i> $r = \eta(\widetilde{T}) - \sigma(\eta(\widetilde{T}))$ skew-symmetric solution r of AYBE in $A \ltimes_{-\rho_\circ} V^*$	<i>deformation</i> \succ	\mathcal{O} -operator $T_{\mathbf{K}}$ on $\succ(A_h, \circ_h)$ associated to (V_h, l_h, r_h) <i>Theorem 4.21</i> $r_h = \eta_h(\widetilde{T}_{\mathbf{K}}) - \sigma_{\mathbf{K}}(\eta_h(\widetilde{T}_{\mathbf{K}}))$ skew-symmetric solution r_h of AYBE in $A_h \ltimes_{-r_h^*, -l_h^*} V_h^*$	<i>Corrolary 2.16</i> <i>Theorem 2.21</i> <i>Theorem 4.4</i> \succ	\mathcal{O} -operator T on $\succ(A, \{, \}, \circ)$ associated to $(V, \rho_{\{, \}}, \rho_\circ)$ <i>Theorem 4.22</i> $r = \eta(\widetilde{T}) - \sigma(\eta(\widetilde{T}))$ skew-symmetric solution r of PYBE in $A \ltimes_{\rho_{\{, \}}, -\rho_\circ} V^*$
---	-------------------------------	---	---	--

Proof. (a). The proof is similar to the one for Proposition 4.19 (a).

(b). Since $T_{\mathbf{K}}$ is the scalar deformation of T , similar to the proofs of (78) and (80), we obtain

$$\eta_h(\widetilde{T}_{\mathbf{K}}) = \widehat{\eta}(\widetilde{T}) \text{ and } \sigma_{\mathbf{K}}(\eta_h(\widetilde{T}_{\mathbf{K}})) = \sigma(\widehat{\eta}(\widetilde{T}))$$

respectively. Thus r_h coincides with the scalar deformation \widehat{r} of r , proving the conclusion. \square

At the end of this paper, we consider dendriform deformations, by regarding dendriform algebras as the special case of \mathcal{O} -operators being identity maps, thanks to Proposition 3.3.

Let (A, \succ) be a finite-dimensional Zinbiel algebra, (A_h, \succ_h, \prec_h) be a dendriform deformation and $(A, \triangleright, \succ)$ be the QCL which is a pre-Poisson algebra. By Propositions 3.16 and 4.23, we obtain the following commutative diagram. Note that in applying Proposition 3.16, we take \cdot, \cdot_h and $[\cdot]$ in Proposition 3.16 as trivial operations.

Zinbiel algebra (A, \succ) <i>Proposition 3.3</i> \mathcal{O} -operator id_A on (A, \circ) associated to (A, L_\succ) <i>Theorem 4.20</i> skew-symmetric solution $r := \eta(\widetilde{\text{id}}_A) - \sigma(\eta(\widetilde{\text{id}}_A))$ of AYBE in $A \ltimes_{-L_\succ} A^*$	<i>deformation</i> \succ	topologically free \succ dendriform algebra (A_h, \succ_h, \prec_h) <i>Proposition 3.3</i> \mathcal{O} -operator id_{A_h} on $\succ(A_h, \circ_h)$ associated to $(A_h, L_{\succ_h}, R_{\prec_h})$ <i>Theorem 4.21</i> skew-symmetric solution $r_h := \eta_h(\widetilde{\text{id}}_{A_h}) - \sigma_{\mathbf{K}}(\eta_h(\widetilde{\text{id}}_{A_h}))$ of AYBE in $A_h \ltimes_{-R_{\prec_h}^*, -L_{\succ_h}^*} A_h^*$	QCL <i>Corollary 3.12</i> <i>Theorem 4.4</i> \succ	\succ pre-Poisson algebra $(A, \triangleright, \succ)$ <i>Proposition 3.7</i> \mathcal{O} -operator id_A on $\succ(A, \{, \}, \circ)$ associated to $(A, L_\triangleright, L_\succ)$ <i>Theorem 4.22</i> skew-symmetric solution $r := \eta(\widetilde{\text{id}}_A) - \sigma(\eta(\widetilde{\text{id}}_A))$ of PYBE in $A \ltimes_{L_\triangleright, -L_\succ} A^*$
---	-------------------------------	--	---	--

Acknowledgements. This work is supported by NSFC (11931009, 12271265, 12261131498), Fundamental Research Funds for the Central Universities and Nankai Zhide Foundation.

Declaration of interests. The authors have no conflicts of interest to disclose.

Data availability. Data sharing is not applicable as no new data were created or analyzed.

REFERENCES

[1] M. Aguiar, Infinitesimal Hopf algebras. *Contemp. Math.* **267**, 1-29, 2000. [2](#), [18](#)
[2] M. Aguiar, Pre-Poisson algebras. *Lett. Math. Phys.* **54** (2000), 263-277. [2](#), [11](#), [12](#)
[3] M. Aguiar, On the associative analog of Lie bialgebras. *J. Algebra.* **244** (2001), 492-532. [2](#), [18](#)
[4] K. Atalikov and A. Zotov, Higher rank 1 + 1 integrable Landau-Lifshitz field theories from the associative Yang-Baxter equation. *JETP Lett.* **115** (2022), 757-762. [2](#)
[5] C. Bai, Double constructions of Frobenius algebras, Connes cocycles and their duality. *J. Noncommut. Geom.* **4** (2010), 475-530. [3](#)
[6] C. Bai, An introduction to pre-Lie algebras. In: *Algebra and Applications 1: Nonassociative Algebras and Categories*, Wiley Online Library (2021), 245-273. [10](#)
[7] C. Bai, O. Bellier, L. Guo and X. Ni, Splitting of operations, Manin products and Rota-Baxter operators. *Int. Math. Res. Not.* **3** (2013), 485-524. [2](#), [11](#)

- [8] C. Bai, L. Guo and X. Ni, Generalizations of the classical Yang-Baxter equation and \mathcal{O} -operators. *J. Math. Phys.* **52** (2011), 063515. [24](#), [25](#)
- [9] C. Bai, L. Guo and X. Ni, \mathcal{O} -operators on associative algebras and associative Yang-Baxter equation. *Pacific J. Math.* **256** (2012), 257-289. [2](#), [3](#), [5](#), [17](#), [22](#), [27](#)
- [10] C. Bai, L. Guo and X. Ni, \mathcal{O} -operators on associative algebras, associative Yang-Baxter equations and dendriform algebras. In: *Quantized Algebra and Physics*. World Scientific, 2012, 10-51. [3](#), [24](#)
- [11] C. Bai, L. Guo and X. Ni, Relative Rota-Baxter operators and tridendriform algebras. *J. Algebra Appl.* **12** (2013), 1350027. [2](#), [4](#), [9](#), [10](#), [13](#)
- [12] A. Borel and J.-P. Serre, Le théorème de Riemann-Roch. *Bull. Soc. Math. France* **86** (1958), 97-136. [3](#)
- [13] A. Das, Cohomology and deformations of weighted Rota-Baxter operators. *J. Math. Phys.* **63** (2022), 091703. [2](#), [3](#)
- [14] V. G. Drinfeld, (1987), Quantum groups. Proceedings of the International Congress of Mathematicians (Berkeley, Calif., 1986), vol. 1, Amer. Math. Soc., 1987, 798-820. [3](#)
- [15] P. Etingof and O. Schiffmann, Lectures on Quantum Groups, 2nd ed. International Press, 2010. [2](#), [6](#), [7](#)
- [16] B. Farb and R. Dennis, Noncommutative Algebra. Springer-Verlag, 1993. [3](#)
- [17] W. Fulton, Intersection Theory. Springer, 1998. [3](#)
- [18] M. Gerstenhaber, On the deformation of rings and algebras. *Ann. Math.* **79** (1964), 59-103. [2](#)
- [19] M. Gerstenhaber, On the deformation of rings and algebras: III. *Ann. Math.* **88** (1968), 1-34. [15](#)
- [20] L. Guo, An Introduction to Rota-Baxter Algebra. International Press, Somerville, MA, 2012. [16](#)
- [21] Y. Hong, C. Bai and L. Guo, Infinite-dimensional Lie bialgebras via affinization of Novikov bialgebras and Koszul duality. *Comm. Math. Phys.* **401** (2023), 2011-2049. [18](#)
- [22] C. Kassel, Quantum groups. Springer-Verlag, 1995. [18](#), [20](#)
- [23] M. Kontsevich, Deformation quantization of Poisson manifolds I. *Lett. Math. Phys.* **66** (2003), 157-216. [2](#)
- [24] B. A. Kupersmidt, What a classical r-matrix really is. *J. Nonlinear Math. Phys.* **6** (1999), 448-488. [2](#)
- [25] J. Liu, C. Bai and Y. Sheng, Noncommutative Poisson bialgebras. *J. Algebra.* **556** (2020), 35-66. [2](#), [28](#)
- [26] J.-L. Loday, Dialgebras. In: *Dialgebras and Related Operads. Lect. Notes in Math* **1763**, 7-66, 2002. [2](#)
- [27] J.-L. Loday and M. Ronco, Trialgebras and families of polytopes. *Contemp. Math.* **346**, 369-398, 2004. [2](#), [10](#)
- [28] J.-L. Loday and B. Vallette, Algebraic Operads. Springer, 2012. [5](#)
- [29] S.-T. Lue, Crossed homomorphisms of Lie algebras. *Proc. Cambridge Philos. Soc.* **62** (1966), 577-581. [3](#)
- [30] S. -T. Lue, Non-abelian cohomology of associative algebras, *J. Math. Oxford* **19** (1968), 159-180. [24](#)
- [31] A. Lundervold and H. Munthe-Kaas, On post-Lie algebras, Lie-Butcher series and moving frames. *Found. Comput. Math.* **13** (2013), 583-613. [10](#)
- [32] Z. Mesyan, The ideals of an ideal extension. *J. Algebra Its Appl.* **9** (2010), 407-431. [5](#)
- [33] A.-V. Mikhailov and V.-V. Sokolov, Integrable ODEs on Associative Algebras. *Comm. Math. Phys.* **211** (2000), 231-251. [2](#)
- [34] X. Ni and C. Bai, Poisson bialgebras. *J. Math. Phys.* **54** (2013), 023515. [2](#), [3](#), [5](#), [6](#), [9](#), [11](#), [13](#), [18](#), [22](#)
- [35] C. Ospel, F. Panaite and P. Vanhaecke, Polarization and deformations of generalized dendriform algebras. *J. Noncommut. Geom.* **16** (2022), 561-594. [2](#), [4](#), [12](#)
- [36] A. Polishchuk, Classical Yang-Baxter equation and the A_∞ -constraint. *Adv. Math.* **168** (2002), 56-95. [2](#)
- [37] A. Polishchuk, Massey products on cycles of projective lines and trigonometric solutions of the Yang-Baxter equations. In: *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II. Prog. Math.* **270**, 573-617, 2009. [2](#)
- [38] K. Uchino, Quantum analogy of Poisson geometry, related dendriform algebras and Rota-Baxter operators. *Lett. Math. Phys.* **85** (2008), 91-109. [2](#)
- [39] B. Vallette, Homology of generalized partition posets. *J. Pure Appl. Algebra.* **208** (2007), 699-725. [10](#)

CHERN INSTITUTE OF MATHEMATICS, NANKAI UNIVERSITY, TIANJIN 300071 CHINA
 Email address: 1120210010@mail.nankai.edu.cn

CHERN INSTITUTE OF MATHEMATICS, NANKAI UNIVERSITY, TIANJIN 300071 CHINA
 Email address: baicm@nankai.edu.cn

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, RUTGERS UNIVERSITY, NEWARK, NJ 07102, USA
 Email address: liguo@rutgers.edu