

# COMPUTING RENORMALIZED CURVATURE INTEGRALS ON POINCARÉ–EINSTEIN MANIFOLDS

JEFFREY S. CASE, AYUSH KHAITAN, YUEH-JU LIN, AARON J. TYRRELL,  
AND WEI YUAN

**ABSTRACT.** We describe a general procedure for computing renormalized curvature integrals on Poincaré–Einstein manifolds. In particular, we explain the connection between the Gauss–Bonnet-type formulas of Albin and Chang–Qing–Yang for the renormalized volume, and explicitly identify a scalar conformal invariant in the latter formula. Our approach constructs scalar conformal invariants that are divergences at any Einstein manifold; these imply that the scalar invariant in the Chang–Qing–Yang formula is not unique in dimension at least eight. Our procedure also produces explicit conformally invariant Gauss–Bonnet-type formulas for compact Einstein manifolds.

## 1. INTRODUCTION

Poincaré–Einstein manifolds are complete Einstein manifolds with a well-defined conformal boundary; the model example is the Poincaré ball model of hyperbolic space. The rich interplay between the conformal geometry of the boundary of a Poincaré–Einstein manifold and the Riemannian geometry of its interior, which is at the heart of the AdS/CFT correspondence [19, 20], leads to many constructions of local and nonlocal conformal invariants (e.g. [5, 15, 17, 18]). Nevertheless, it remains a fundamental open problem to understand the moduli space of Poincaré–Einstein manifolds with a given conformal boundary (cf. [3, 8–10]).

A first step in studying this moduli space is to develop a good understanding of global invariants of Poincaré–Einstein manifolds. In even dimensions, renormalized curvature integrals [1] give many such invariants: Let  $r$  be a geodesic defining function for an even-dimensional Poincaré–Einstein manifold  $(M^n, g_+)$ . A **renormalized curvature integral**

$$\int^R I \, dV := \text{fp} \int_{r>\varepsilon} I^{g_+} \, dV_{g_+}$$

is the constant term in the Laurent expansion of the total integral of a scalar Riemannian invariant  $I$  of degree at least  $-n$ ; i.e.  $I$  is a linear combination of complete contractions  $I_j$  of tensor products of covariant derivatives of the Riemann curvature tensor, each of which has the property that there is a weight  $w_j \geq -n$  such that  $I_j^{c^2 g} = c^{w_j} I_j^g$  for all metrics  $g$  and all constants  $c > 0$ . The restriction that  $I$  has degree at least  $-n$  ensures that the corresponding scalar  $\tilde{I}$  in the ambient space is well-defined (cf. [14]). We say that the renormalized curvature integral  $\int^R I \, dV$  is **even** if  $I$  is even; i.e. if  $I$  is invariant under orientation-reversing diffeomorphisms.

---

2020 *Mathematics Subject Classification.* Primary 53C25.

*Key words and phrases.* renormalized volume; Gauss–Bonnet; Poincaré–Einstein manifold.

There are many renormalized curvature integrals of interest. The most commonly studied such integral is the **renormalized volume** [16]

$$\mathcal{V} := \int^R dV;$$

i.e. the renormalized integral of the constant scalar 1. Another example is the renormalized integral of the Pfaffian  $\text{Pf}$  (of the Riemann curvature tensor). As a generalization of the Gauss–Bonnet–Chern Theorem [12], Albin showed [1] that the Euler characteristic is proportional to this renormalized curvature integral:

$$(1.1) \quad \int^R \text{Pf} dV = (2\pi)^{n/2} \chi(M).$$

These are both even renormalized curvature integrals. A parity argument shows that the renormalized curvature integral of an odd scalar Riemannian invariant is the renormalized integral of a top-degree Pontrjagin form [5, p. 502]. Such forms are conformally invariant [4], and hence  $\int^R I dV = \int_M I dV$  in this case. For this reason, we restrict our attention to even renormalized curvature integrals.

In dimension four, the Poincaré–Einstein condition implies that any even renormalized curvature integral is a linear combination of the renormalized volume and the renormalized  $L^2$ -norm of the Weyl tensor  $W$ . Moreover,  $|W|^2 dV$  is conformally invariant in this dimension, and hence  $\int |W|^2 dV$  is finite. For example, expanding the Pfaffian in Equation (1.1) recovers Anderson’s formula [3]

$$(1.2) \quad 4\pi^2 \chi(M) = 3\mathcal{V} + \frac{1}{8} \int_M |W|^2 dV$$

for the renormalized volume.

In higher dimensions there are many additional renormalized curvature integrals. While Equation (1.1) still relates the renormalized volume to the Euler characteristic, it now does so through the renormalized integral of a scalar polynomial in the Weyl tensor. In a different direction, Chang, Qing, and Yang showed [11], in analogy with Equation (1.2), that there is a scalar conformal invariant  $\mathcal{W}_n$  of weight  $-n$  such that if  $(M^n, g_+)$  is Poincaré–Einstein, then

$$(1.3) \quad (2\pi)^{n/2} \chi(M) = (-1)^{n/2} (n-1)!! \mathcal{V} + \int_M \mathcal{W}_n dV.$$

An explicit formula for  $\mathcal{W}_n$  is only known in dimension at most six [16]; in higher dimensions, the existence of  $\mathcal{W}_n$  follows from Alexakis’ classification of conformally invariant integrals of scalar Riemannian invariants [2]. For this reason it is unclear whether one can derive one of Equation (1.1) or (1.3) from the other.

The main purpose of this paper is to describe a procedure which is independent of Alexakis’ classification and explicitly computes many renormalized curvature integrals. Applying our procedure to the Pfaffian produces an ambient formula for  $\mathcal{W}_n$  and explains the relationship between Equations (1.1) and (1.3):

**Theorem 1.1.** *Let  $(M^n, g_+)$  be an even-dimensional Poincaré–Einstein manifold. Then*

$$(2\pi)^{n/2} \chi(M) = (-1)^{n/2} (n-1)!! \mathcal{V} + \sum_{\ell=2}^{n/2} (-2)^{\ell-n/2} \frac{(\ell-1)!}{(n/2-1)!} \int_M \mathcal{P}_{\ell,n} dV,$$

where  $\mathcal{P}_{\ell,n} := i^* (\widetilde{\Delta}^{n/2-\ell} \text{Pf}_\ell(\widetilde{\text{Rm}}))$ .

Given a section  $T$  of  $\otimes^4 T^*M$  on a pseudo-Riemannian manifold  $(M^n, g)$ , we define

$$(1.4) \quad \text{Pf}_\ell(T) := 2^{-\ell}(2\ell-1)!! \delta_{b_1 \dots b_{2\ell}}^{a_1 \dots a_{2\ell}} T_{a_1 a_2}^{b_1 b_2} \dots T_{a_{2\ell-1} a_{2\ell}}^{b_{2\ell-1} b_{2\ell}},$$

$$(1.5) \quad \delta_{b_1 \dots b_k}^{a_1 \dots a_k} := \delta_{[b_1}^{[a_1} \dots \delta_{b_k]}^{a_k]},$$

with the convention  $\text{Pf}_0(T) := 1$ . Thus  $\text{Pf}_{n/2}(\text{Rm})$  is the Pfaffian and  $k! \delta_{b_1 \dots b_k}^{a_1 \dots a_k}$  is the generalized Kronecker delta. With this notation,  $\text{Pf}_\ell(\widetilde{\text{Rm}})$  is the corresponding scalar invariant determined by the Riemann curvature tensor  $\widetilde{\text{Rm}}$  of the ambient space of  $(M^n, [g])$ , and  $\mathcal{P}_{\ell,n}$  is the evaluation at  $(M^n, g)$  of the conformal invariant of weight  $-n$  determined by the scalar invariant  $\widetilde{\Delta}^{n/2-\ell} \text{Pf}_\ell(\widetilde{\text{Rm}})$  built using also the Laplacian  $\widetilde{\Delta}$  in the ambient space.

Direct computation gives  $\mathcal{P}_{2,4} = \frac{1}{8}|W|^2$ , and hence Theorem 1.1 recovers Anderson's formula (1.2); see Section 5 for this and related computations. More generally, producing a local formula for  $\mathcal{P}_{\ell,n}$  requires knowing the ambient metric to order  $\rho^{n/2-\ell+1}$ . In particular, computations of Fefferman and Graham [14] can be used to produce explicit local expressions for  $\mathcal{P}_{\ell,n}$  when  $n \leq 8$ .

Our proof of Theorem 1.1 also produces a formula for the Euler characteristic of a compact even-dimensional Einstein manifold as a linear combination of its volume and the integral of a scalar conformal invariant:

**Theorem 1.2.** *Let  $(M^n, g)$  be a compact even-dimensional Einstein manifold with  $\text{Ric}_g = 2\lambda(n-1)g$ . Then*

$$(2\pi)^{n/2} \chi(M) = (2\lambda)^{n/2} (n-1)!! \text{Vol}_g(M) + \sum_{\ell=2}^{n/2} (-2)^{\ell-n/2} \frac{(\ell-1)!}{(n/2-1)!} \int_M \mathcal{P}_{\ell,n} dV,$$

where  $\mathcal{P}_{\ell,n}$  is as in Theorem 1.1.

Theorems 1.1 and 1.2 are special cases of a general procedure for computing renormalized curvature integrals. This procedure begins with a distinguished class of scalar Riemannian invariants that are well-suited to computations in the Fefferman–Graham ambient space:

**Definition 1.3.** A scalar Riemannian invariant  $I$  of weight  $w$  is **straightenable** if  $\widetilde{I} = \tau^w \pi^* I^g$  whenever  $(\widetilde{\mathcal{G}}, \widetilde{g})$ ,

$$(1.6) \quad \begin{aligned} \widetilde{\mathcal{G}} &:= (0, \infty)_t \times M \times (-\varepsilon, \varepsilon)_\rho, \\ \widetilde{g} &:= 2\rho dt^2 + 2t dt d\rho + \tau^2 g, \\ \tau &:= t(1 + \lambda\rho), \end{aligned}$$

is the straight and normal ambient space associated to an Einstein manifold  $(M^n, g)$  with  $\text{Ric} = 2\lambda(n-1)g$ , where  $\pi: \widetilde{\mathcal{G}} \rightarrow M$  is the canonical projection determined by  $g$  and  $\widetilde{I}$  is the evaluation of  $I$  at  $(\widetilde{\mathcal{G}}, \widetilde{g})$ .

Computations of Fefferman and Graham [14] imply that scalar polynomials in the Weyl tensor, such as  $\text{Pf}_\ell(W)$ , are straightenable. However, not all scalar Riemannian invariants are straightenable; e.g.  $|\text{Rm}|^2$ . In Section 3 we define and construct additional straightenable tensor Riemannian invariants. Remarkably, these constructions show that the invariant  $\mathcal{W}_n$  in Equation (1.3) is *not* uniquely determined in dimension  $n \geq 8$ ; see Remark 3.11 for further discussion. In Section 5

we describe how to use straightenable invariants to compute many renormalized curvature integrals, including all such integrals in dimension at most eight.

While straightenable scalar Riemannian invariants are not necessarily conformally invariant, they are readily identified with conformal invariants. Indeed, if  $I$  is a straightenable scalar Riemannian invariant of weight  $w$ , then  $i^*\tilde{I}$  defines a scalar conformal invariant of weight  $w$  which agrees with  $I$  when evaluated at an Einstein manifold. See Remark 3.7 below for an example of a straightenable scalar Riemannian invariant that is not conformally invariant.

Our main result is a formula for the renormalized integral of a straightenable scalar Riemannian invariant in terms of the convergent integral of a scalar conformal invariant.

**Theorem 1.4.** *Let  $n \in \mathbb{N}$  be an even integer and let  $I$  be a straightenable scalar Riemannian invariant of weight  $-2k \geq -n$ . If  $(M^n, g_+)$  is a Poincaré–Einstein manifold, then*

$$\int_M^R I \, dV = \frac{2^{k-n/2}(k-1)!}{(n/2-1)!(n-2k-1)!!} \int_M \mathcal{I}_{n/2-k} \, dV,$$

where  $\mathcal{I}_{n/2-k} := i^*(\tilde{\Delta}^{n/2-k}\tilde{I})$ .

Note that  $\mathcal{I}_{n/2-k}$  is a scalar conformal invariant of weight  $-n$ .

Theorem 1.4 gives a new proof (cf. [1]) that  $\int_M^R I \, dV$  is independent of the choice of geodesic defining function when  $I$  is straightenable. It also applies to a wide range of invariants; see Section 5 for examples.

There are two key observations needed to prove Theorem 1.4. First, given a straightenable scalar Riemannian invariant  $I$ , there is a constant  $c$  depending only on the dimension and the weight of  $I$  such that  $\Delta I + cRI$  is straightenable, where  $R$  is the scalar curvature. Iteratively applying this to a straightenable scalar Riemannian invariant  $I$  of weight  $-2k \geq -n$  produces a straightenable scalar Riemannian invariant of weight  $-n$  which, when evaluated at an Einstein manifold, is a linear combination of  $I$  and divergences. Second, a parity argument of Albin [1] can be adapted to prove that the renormalized integral of a divergence is always zero.

Applying Theorem 1.4 and its proof to  $\text{Pf}_\ell(W)$  recovers Theorems 1.1 and 1.2.

This paper is organized as follows:

In Section 2 we fix our conventions and recall some necessary facts about the Fefferman–Graham ambient space.

In Section 3 we produce many examples of straightenable scalar Riemannian invariants. As an application, we construct scalar conformal invariants that are divergences when evaluated at Einstein manifolds.

In Section 4 we prove Theorem 1.4.

In Section 5 we prove Theorems 1.1 and 1.2, derive more explicit formulas for the invariants  $\mathcal{P}_{\ell,n}$  in dimensions at most eight, and discuss how to apply Theorem 1.4 to compute more general renormalized curvature integrals.

## 2. CONVENTIONS AND BACKGROUND

Let  $M^n$  be a smooth manifold of dimension  $n \geq 3$ . We typically perform computations using abstract index notation, using lowercase Latin letters to denote factors of  $T^*M$  (when subscripts) or its dual  $TM$  (when superscripts), with repeated indices denoting a contraction via the canonical pairing of  $TM$  and  $T^*M$ .

For example, we often write  $T_{abc}$  to denote a section  $T \in \Gamma(\otimes^3 T^*M)$  and  $X^a$  to denote a vector field  $X \in \Gamma(TM)$ . If also  $Y, Z \in \Gamma(TM)$ , then

$$T(X, Y, Z) = X^a Y^b Z^c T_{abc}.$$

We use square brackets and round parentheses to denote skew-symmetrization and symmetrization, respectively; e.g.

$$\begin{aligned} T_{[abc]} &:= \frac{1}{6}(T_{abc} - T_{acb} + T_{bca} - T_{bac} + T_{cab} - T_{cba}), \\ T_{(abc)} &:= \frac{1}{6}(T_{abc} + T_{acb} + T_{bca} + T_{bac} + T_{cab} + T_{cba}). \end{aligned}$$

Let  $g$  be a pseudo-Riemannian metric on  $M$ . Denote by  $g^{-1}$  or  $g^{ab}$  the inverse of  $g$ . We denote by  $\text{Rm}$  or  $R_{abcd}$  the Riemann curvature tensor, defined by the convention

$$\nabla_a \nabla_b \tau_c - \nabla_b \nabla_a \tau_c = R_{abc}{}^d \tau_d.$$

We use  $g_{ab}$  and  $g^{ab}$  to lower and raise indices; e.g.  $R_{abcd} = R_{abc}{}^e g_{ed}$ . The **Schouten tensor** is

$$P_{ab} := \frac{1}{n-2} (R_{ab} - J g_{ab}),$$

where  $R_{ab} := R_{acb}{}^c$  is the Ricci tensor,  $R := R_a{}^a$  is the scalar curvature, and  $J := P_a{}^a$  is the trace of the Schouten tensor. Recall that the **Weyl tensor**

$$W_{abcd} := R_{abcd} - 2P_{a[c} g_{d]b} + 2P_{b[c} g_{d]a}$$

is conformally covariant:  $W^{e^{2u}g} = e^{2u} W^g$ .

A **tensor Riemannian invariant**  $I$  of rank  $k$  on  $n$ -manifolds is an assignment to each pseudo-Riemannian  $n$ -manifold  $(M^n, g)$  of a section  $I^g \in \Gamma(\otimes^k T^*M)$  that can be expressed as an  $\mathbb{R}$ -linear combination of partial contractions of

$$(2.1) \quad \nabla^{N_1} \text{Rm} \otimes \cdots \otimes \nabla^{N_j} \text{Rm} \otimes g^{\otimes \ell},$$

where we regard  $\nabla^N \text{Rm} \in \Gamma(\otimes^{N+4} T^*M)$  and  $g \in \Gamma(\otimes^2 T^*M)$ , so that all contractions are made using the inverse metric  $g^{-1}$ , and by convention the empty tensor product is the constant scalar 1. We say that  $I$  has **weight**  $w \in \mathbb{R}$  if  $I^{c^2g} = c^w I^g$  for all pseudo-Riemannian  $n$ -manifolds  $(M^n, g)$  and all constants  $c > 0$ ; in this case we say that  $I$  is **homogeneous** and we define its **tensor weight** to be  $w - k$ . A **scalar Riemannian invariant** is a tensor Riemannian invariant of rank 0. We say that  $I$  is a **scalar conformal invariant** of weight  $w \in \mathbb{R}$  if additionally  $I^{u^2g} = u^w I^g$  for all positive  $u \in C^\infty(M)$ .

The **straight and normal ambient space**  $(\tilde{\mathcal{G}}, \tilde{g})$  associated to  $(M^n, g)$  is the smooth manifold  $\tilde{\mathcal{G}} := (0, \infty)_t \times M \times (-\varepsilon, \varepsilon)_\rho$  together with the pseudo-Riemannian metric

$$(2.2) \quad \tilde{g} = 2\rho dt^2 + 2t dt d\rho + t^2 g_\rho,$$

where  $g_\rho$  is a one-parameter family of metrics on  $M$  such that  $g_0 = g$  and

- (i) if  $n$  is odd, then  $\widetilde{\text{Ric}} = O(\rho^\infty)$  along  $\{\rho = 0\}$ ; while
- (ii) if  $n$  is even, then  $\widetilde{\text{Ric}} = O(\rho^{n/2-1})$  and  $\tilde{R} = O(\rho^{n/2})$  along  $\{\rho = 0\}$ .

Here  $\tilde{T} = O(\rho^k)$  along  $\{\rho = 0\}$  if  $\partial_\rho^\ell|_{\rho=0} \tilde{T} = 0$  for all  $\ell < k$ . Note that  $\delta_s^* \tilde{g} = s^2 \tilde{g}$  for  $\delta_s: \tilde{\mathcal{G}} \rightarrow \tilde{\mathcal{G}}$ ,  $s > 0$ , the one-parameter family of dilations  $\delta_s(t, x, \rho) := (st, x, \rho)$ . Fefferman and Graham proved [14, Theorem 2.9] the existence and uniqueness of the straight and normal ambient space. Uniqueness means that if  $n$  is odd, then

$g_\rho$  is uniquely determined modulo  $O(\rho^\infty)$ ; while if  $n$  is even, then  $g_\rho$  and  $\text{tr}_g g_\rho$  are uniquely determined modulo  $O(\rho^{n/2})$  and  $O(\rho^{n/2+1})$ , respectively. Denote by  $i_g: M \rightarrow \tilde{\mathcal{G}}$  the inclusion  $i_g(x) := (1, x, 0)$ , and by  $\pi_g: \tilde{\mathcal{G}} \rightarrow M$  the projection  $\pi_g(t, x, \rho) := x$ . Fefferman and Graham also showed [14, Theorem 2.3] that if  $(\tilde{\mathcal{G}}_j, \tilde{g}_j)$ ,  $j \in \{1, 2\}$  are straight and normal ambient metrics for conformally related metrics  $g_2 = e^{2u}g_1 \in [g]$ , then, after shrinking  $\tilde{\mathcal{G}}_j$  if necessary, there is a  $\delta_s$ -equivariant diffeomorphism  $\Phi: \tilde{\mathcal{G}}_1 \rightarrow \tilde{\mathcal{G}}_2$  such that  $\Phi(t, x, 0) = (e^{-u(x)}t, x, 0)$  and  $\Phi^*\tilde{g}_2 \equiv \tilde{g}_1$  modulo the undetermined terms in the expansions of  $g_\rho$ .

One application of the straight and normal ambient space is to the construction of scalar conformal invariants: If  $I$  is a homogeneous scalar Riemannian invariant of weight  $w \geq -n$ , then  $\mathcal{I}^g := i_g^* \tilde{I}$  defines a scalar conformal invariant of weight  $w$ , where  $\tilde{I} \in C^\infty(\tilde{\mathcal{G}})$  is the function determined by replacing  $\nabla$ ,  $\text{Rm}$ , and  $g$  in Display (2.1) by  $\tilde{\nabla}$ ,  $\tilde{\text{Rm}}$ , and  $\tilde{g}$ , respectively, contracting as specified in the definition of  $I$ , and treating the coefficients of the linear combination of complete contractions as fixed real numbers, rather than dimension-dependent numbers; cf. Example 5.5. We say that  $\mathcal{I}$  is **determined ambiently** by  $I$ . All even scalar conformal invariants of weight  $w \geq -n$  arise in this way [5]. Note that  $\mathcal{I} \neq I$  in general; e.g. the scalar curvature determines the scalar conformal invariant 0.

When computing with the ambient metric, we use capital Latin letters to denote factors of  $T\tilde{\mathcal{G}}$  and its dual; and we use the symbols 0,  $a$ , and  $\infty$  to denote projections onto  $T(0, \infty)_t$ ,  $TM$ , and  $T(-\varepsilon, \varepsilon)_\rho$ , respectively, and their duals. For example, Fefferman and Graham observed [14, Equation (6.1)] that the nonvanishing projections of the Riemann curvature tensor  $\tilde{\text{Rm}}$  of  $\tilde{g}$  can be computed from the formulas

$$\begin{aligned} \tilde{R}_{abcd} &= t^2 \left( R_{abcd} - g_{a[c} g'_{d]b} + g_{b[c} g'_{d]a} + \rho g'_{a[c} g'_{d]b} \right), \\ \tilde{R}_{\infty abc} &= t^2 \nabla_{[c} g'_{b]a}, \\ \tilde{R}_{\infty ab\infty} &= \frac{t^2}{2} \left( g''_{ab} - \frac{1}{2} g^{cd} g'_{ac} g'_{bd} \right), \end{aligned} \tag{2.3}$$

where  $' := \partial_\rho$  and the geometric invariants on the right-hand side are computed with respect to  $g_\rho$ .

### 3. STRAIGHTENABLE SCALAR RIEMANNIAN INVARIANTS

In this section we construct many straightenable scalar Riemannian invariants. The main point is that there is an explicit straight and normal ambient space associated to an Einstein manifold which is in fact an *exact* solution of  $\tilde{\text{Ric}} = 0$ .

**Lemma 3.1** ([14, Equation (7.12)]). *The straight and normal ambient space  $(\tilde{\mathcal{G}}, \tilde{g})$  associated to an Einstein manifold  $(M^n, g)$ ,  $n \geq 3$ , with  $\text{Ric}_g = 2\lambda(n-1)g$  is given by Equations (1.6). Moreover,  $\tilde{\text{Ric}} = 0$ .*

Thus straightenable scalar Riemannian invariants are characterized by their evaluations at straight and normal ambient spaces associated to an Einstein manifold. While a straightenable scalar Riemannian invariant need not be conformally invariant (see Remark 3.7 below), it necessarily is equal to a conformal invariant when evaluated at an Einstein manifold.

**Lemma 3.2.** *Fix  $n \in \mathbb{N}$ . Let  $I$  be a straightenable scalar Riemannian invariant of weight  $w \geq -n$ . Let  $\mathcal{I}$  be the scalar conformal invariant determined ambiently by  $I$ . If  $(M^n, g)$  is Einstein, then  $I^g = \mathcal{I}^g$ .*

*Proof.* Let  $(\tilde{\mathcal{G}}, \tilde{g})$  be the straight and normal ambient space associated to the Einstein manifold  $(M^n, g)$ . Since  $i_g^* \tau = 1$  and  $\pi_g \circ i_g = \text{Id}$ , we compute that

$$\mathcal{I}^g = i_g^* \tilde{\mathcal{I}} = i_g^* (\tau^w \pi_g^* I^g) = I^g. \quad \square$$

For our general construction of straightenable scalar Riemannian invariants, it is convenient to extend Definition 1.3 to tensor Riemannian invariants.

**Definition 3.3.** A tensor Riemannian invariant  $T$  of weight  $w$  is **straightenable** if  $\tilde{T} = \tau^w \pi^* T^g$  whenever  $(\tilde{\mathcal{G}}, \tilde{g})$  is the straight and normal ambient space associated to an Einstein manifold  $(M^n, g)$ .

Straightenable tensor Riemannian invariants exist:

**Lemma 3.4.** *The Weyl tensor is straightenable and of tensor weight  $-2$ .*

*Proof.* Let  $(\tilde{\mathcal{G}}, \tilde{g})$  be the straight and normal ambient space of an Einstein manifold  $(M^n, g)$ . Equation (2.3) and Lemma 3.1 imply that  $\widetilde{\text{Rm}} = \tau^2 \pi_g^* W^g$  as sections of  $\otimes^4 T^* \tilde{\mathcal{G}}$ . Since  $\widetilde{\text{Rm}} = \widetilde{W}$ , we conclude that the Weyl tensor is straightenable and has tensor weight  $-2$ .  $\square$

Partial contraction of tensor products yields additional straightenable tensor Riemannian invariants.

**Lemma 3.5.** *Suppose that  $S$  and  $T$  are straightenable tensor Riemannian invariants of tensor weight  $s$  and  $t$ , respectively. Then any partial contraction of  $S \otimes T$  is straightenable and of tensor weight  $s + t$ . In particular, any complete contraction of  $W^{\otimes k}$  is a straightenable scalar Riemannian invariant of weight  $-2k$ .*

*Proof.* It follows immediately from Definition 3.3 that  $S \otimes T$  is a straightenable tensor Riemannian invariant of tensor weight  $s + t$ . Since  $\tilde{g}^{ab} = \tau^{-2} g^{ab}$  when  $g$  is Einstein, the same is true of any partial contraction of  $S \otimes T$ . The final conclusion follows from Lemma 3.4.  $\square$

A nice property of the ambient Laplacian on functions allows us to produce higher-order straightenable scalar Riemannian invariants:

**Proposition 3.6.** *Let  $I$  be a straightenable scalar Riemannian invariant of weight  $-2k$ . Given  $\ell \in \mathbb{N}$ , set*

$$(3.1) \quad I_\ell := \left( \prod_{j=0}^{\ell-1} \left( \Delta - \frac{4(k+j)(n-2k-2j-1)}{n} \mathbf{J} \right) \right) I.$$

*Then  $I_\ell$  is a straightenable scalar Riemannian invariant of weight  $-2k - 2\ell$ . Moreover, if  $(M^n, g)$  is Einstein, then*

$$(3.2) \quad I_\ell \equiv \left( -\frac{4}{n} \mathbf{J} \right)^\ell \frac{(k+\ell-1)!(n-2k-1)!!}{(k-1)!(n-2k-2\ell-1)!!} I \pmod{\Delta I, \dots, \Delta^\ell I}.$$

*Remark 3.7.* Lemma 3.4 implies that  $|W|^2$  is a straightenable scalar Riemannian invariant of weight  $-4$ . Set  $I := \Delta|W|^2 - \frac{8(n-5)}{n}J|W|^2$ . On the one hand, Proposition 3.6 implies that  $I$  is straightenable. On the other hand, direct computation shows that  $I$  is not conformally invariant in dimension  $n \neq 10$  (cf. [6, Section 8.4]). Therefore straightenable scalar Riemannian invariants need not be conformally invariant.

*Proof.* Let  $(\tilde{\mathcal{G}}, \tilde{g})$  be the straight and normal ambient space associated to an Einstein manifold  $(M^n, g)$  with  $\text{Ric} = 2\lambda(n-1)g$ . Direct computation shows [7, Lemma 5.1] that if  $u \in C^\infty(M)$  and  $w \in \mathbb{R}$ , then

$$(3.3) \quad \tilde{\Delta}(\tau^w \pi^* u) = \tau^{w-2} \pi^* ((\Delta + 2\lambda w(n+w-1))u).$$

Let  $I$  be a straightenable scalar Riemannian invariant of weight  $-2k$  and define  $I_\ell$  by Equation (3.1). Since  $\tilde{g}$  is scalar-flat,

$$\tilde{I}_\ell = \tilde{\Delta}^\ell \tilde{I}.$$

Repeated application of Equation (3.3) and the fact  $J = n\lambda$  yield

$$\tilde{\Delta}^\ell \tilde{I} = \tilde{\Delta}^\ell (\tau^{-2k} \pi^* I) = \tau^{-2k-2\ell} \pi^* I_\ell.$$

Finally, since  $J$  is constant at an Einstein manifold, we compute that

$$\begin{aligned} I_\ell &\equiv \left(-\frac{4}{n}J\right)^\ell \left(\prod_{j=0}^{\ell-1} (k+j)\right) \left(\prod_{j=0}^{\ell-1} (n-2k-2j-1)\right) I \pmod{\Delta I, \dots, \Delta^\ell I} \\ &\equiv \left(-\frac{4}{n}J\right)^\ell \frac{(k+\ell-1)!(n-2k-1)!!}{(k-1)!(n-2k-2\ell-1)!!} I \pmod{\Delta I, \dots, \Delta^\ell I}. \end{aligned} \quad \square$$

A nice property of the ambient divergence on symmetric tensors allows us to produce higher-order straightenable tensor Riemannian invariants:

**Proposition 3.8.** *Let  $T_{a_1 \dots a_k}$  be a straightenable symmetric tensor Riemannian invariant of weight  $w$ . Then*

$$U_{a_1 \dots a_{k-1}} := \nabla^b T_{a_1 \dots a_{k-1} b} + \frac{k-1}{w-2k+2} \nabla_{(a_1} T_{a_2 \dots a_{k-1}) b}{}^b$$

*is a straightenable symmetric tensor Riemannian invariant of weight  $w-2$ .*

*Proof.* Clearly  $U_{a_1 \dots a_{k-1}}$  is symmetric.

Let  $(M^n, g)$  be such that  $\text{Ric} = 2\lambda(n-1)g$  and let  $(\tilde{\mathcal{G}}, \tilde{g})$  be its straight and normal ambient space. Denote  $\sigma := 1 + \lambda\rho$ . Specializing the formulas [14, Equation (3.16)] for the Christoffel symbols of a metric of the form (2.2) to  $\tilde{g}$  yields

$$\begin{aligned} \Gamma_{AB}^0 &= \begin{pmatrix} 0 & 0 & 0 \\ 0 & -\lambda\tau g_{ab} & 0 \\ 0 & 0 & 0 \end{pmatrix}, \\ \Gamma_{AB}^c &= \begin{pmatrix} 0 & t^{-1}\delta_b^c & 0 \\ t^{-1}\delta_a^c & \Gamma_{ab}^c & \lambda\sigma^{-1}\delta_a^c \\ 0 & \lambda\sigma^{-1}\delta_b^c & 0 \end{pmatrix}, \\ \Gamma_{AB}^\infty &= \begin{pmatrix} 0 & 0 & t^{-1} \\ 0 & \sigma(\lambda\rho-1)g_{ab} & 0 \\ t^{-1} & 0 & 0 \end{pmatrix}. \end{aligned}$$



Combining this with the assumption  $\tilde{T} = \tau^w \pi^* T^g$  yields

$$\begin{aligned}\tilde{\nabla}^B \tilde{T}_{a_1 \dots a_{k-1} B} &= \tau^{w-2} \nabla^b T_{a_1 \dots a_{k-1} b}, \\ \tilde{\nabla}^B \tilde{T}_{0 a_2 \dots a_{k-1} B} &= -\frac{\tau^{w-2}}{t} T_{a_2 \dots a_{k-1} b}{}^b, \\ \tilde{\nabla}^B \tilde{T}_{\infty a_2 \dots a_{k-1} B} &= -\frac{\lambda \tau^{w-2}}{\sigma} T_{a_2 \dots a_{k-1} b}{}^b,\end{aligned}$$

and all other components vanish. We also compute that

$$\begin{aligned}\tilde{\nabla}_{(a_1} \tilde{T}_{a_2 \dots a_{k-1}) B}{}^B &= \tau^{w-2} \nabla_{(a_1} T_{a_2 \dots a_{k-1}) b}{}^b, \\ \tilde{\nabla}_{(0} \tilde{T}_{a_2 \dots a_{k-1}) B}{}^B &= \frac{w-2k+2}{k-1} \frac{\tau^{w-2}}{t} T_{a_2 \dots a_{k-1} b}{}^b, \\ \tilde{\nabla}_{(\infty} \tilde{T}_{a_2 \dots a_{k-1}) B}{}^B &= \frac{w-2k+2}{k-1} \frac{\lambda \tau^{w-2}}{\sigma} T_{a_2 \dots a_{k-1} b}{}^b,\end{aligned}$$

and all other components vanish. The conclusion readily follows.  $\square$

Applying Proposition 3.8 twice to straightenable symmetric tensor Riemannian invariants of rank 2 produces nontrivial scalar conformal invariants which are divergences at Einstein manifolds in dimension at least eight.

**Corollary 3.9.** *Let  $T_{ab}$  be a straightenable symmetric tensor Riemannian invariant of weight  $w$ . Then*

$$\nabla^a \nabla^b T_{ab} + \frac{1}{w-2} \nabla^a \nabla_a T_b{}^b$$

*is a straightenable scalar Riemannian invariant of weight  $w-4$ .*

*Remark 3.10.* The largest weight for which Corollary 3.9 produces nontrivial invariants is  $w = -8$ . Indeed, there are no nontrivial straightenable symmetric tensor Riemannian invariants formed from a partial contraction of  $W_{abcd}$ , while applying Corollary 3.9 to  $W_{acde} W_b{}^{cde}$  yields the weight  $-6$  invariant

$$D = (n-4) \nabla^a (W_{abcd} C^{cd}{}_b),$$

which vanishes at all Einstein manifolds. There are two straightenable symmetric tensor Riemannian invariants of rank 2 formed from partial contractions of  $W^{\otimes 3}$ . Applying Corollary 3.9 to these yields the nontrivial straightenable invariants

$$\begin{aligned}\nabla^a \nabla^b (W_{abcd} W^{cefg} W^d{}_{efg}), \\ \nabla^a \nabla^b (W_{acde} W_b{}^c{}_{fg} W^{defg}) - \frac{1}{6} \Delta (W_{ab}{}^{cd} W_{cd}{}^{ef} W_{ef}{}^{ab}),\end{aligned}$$

of weight  $-8$ .

*Remark 3.11.* Every invariant produced by Corollary 3.9 is a divergence. It then follows from Lemma 3.2 and Remark 3.10 that in all even dimensions  $n \geq 8$  there are nontrivial scalar conformal invariants whose total integrals vanish at *all* compact Einstein manifolds.

*Proof.* Proposition 3.8 implies that

$$U_a := \nabla^b T_{ab} + \frac{1}{w-2} \nabla_a T_b{}^b \quad \text{and} \quad \nabla^a U_a$$

are straightenable tensor Riemannian invariants of weight  $w-2$  and weight  $w-4$ , respectively.  $\square$

## 4. PROOF OF THEOREM 1.4

We begin by proving that the renormalized integral of a natural divergence vanishes. We do so by adapting an argument used by Albin [1, Sections 3 and 4] to study the asymptotic expansions of scalar Riemannian invariants on Poincaré–Einstein manifolds.

**Lemma 4.1.** *Fix an even integer  $n$ . Let  $F$  be a tensor Riemannian invariant of rank 1 and weight at least  $2 - n$ . If  $(M^n, g_+)$  is a Poincaré–Einstein manifold, then*

$$\int^R \operatorname{div} F \, dV = 0.$$

*Proof.* Pick a representative  $h$  of the conformal boundary of  $(M^n, g_+)$ . Chrusciel, Delay, Lee, and Skinner showed [13, Theorem A] that there is a collar diffeomorphism  $\Phi$  from  $[0, \varepsilon_0)_r \times \partial M$  onto a neighborhood of  $\partial M$  in  $M$  such that  $\Phi(0, \cdot)$  is the inclusion  $\partial M \hookrightarrow M$  and  $\Phi^* g_+ = r^{-2}(dr^2 + g_r)$  for a one-parameter family  $g_r$  of metrics on  $\partial M$  such that

$$(4.1) \quad g_r = h + h_{(2)}r^2 + \cdots + h_{(n-2)}r^{n-2} + kr^{n-1} + O(r^n)$$

has an even expansion to order  $O(r^{n-1})$  and  $k$  is trace-free with respect to  $h$ .

Let  $\varepsilon \in (0, \varepsilon_0)$ . The Divergence Theorem implies that

$$(4.2) \quad \int_{r>\varepsilon} (\operatorname{div} F)^{g_+} dV_{g_+} = \oint_{r=\varepsilon} \langle F, \mathbf{n} \rangle^{g_+} dA_{g_+},$$

where  $\mathbf{n}$  is the outward-pointing unit normal and  $dA_{g_+}$  is the induced volume form on the level set  $\{r = \varepsilon\}$ . Note that

$$dA_{g_+} = \varepsilon^{1-n} dV_{g_\varepsilon}.$$

Since  $\operatorname{tr}_h k = 0$ , we deduce from Equation (4.1) that the asymptotic expansion

$$(4.3) \quad dA_{g_+} = (a_0\varepsilon^{1-n} + a_2\varepsilon^{3-n} + \cdots + a_{n-2}\varepsilon^{-1} + a_n\varepsilon + O(\varepsilon^2)) dA_h$$

is odd to order  $O(\varepsilon^2)$ . We will show that the asymptotic expansion

$$(4.4) \quad \langle F, \mathbf{n} \rangle^{g_+} = f_0 + f_2\varepsilon^2 + \cdots + f_n\varepsilon^n + O(\varepsilon^{n+1})$$

is even to order  $O(\varepsilon^{n+1})$ . Combining Equations (4.2), (4.3), and (4.4) then gives the final conclusion.

Pick a point  $p \in \partial M$ . Let  $\{x^\alpha\}_{\alpha=1}^{n-1}$  be geodesic normal coordinates for  $h$  near  $p$  in  $\partial M$ . Extend these to coordinates on a neighborhood of  $p$  in  $M$  via the collar diffeomorphism  $\Phi$  and let  $\{\partial_r, \partial_{x^\alpha}\}$  denote the resulting coordinate vector fields. Set  $Y_0 := r\partial_r$  and  $Y_\alpha := r\partial_{x^\alpha}$  for  $\alpha \in \{1, \dots, n-1\}$ . Unless otherwise specified, in the remainder of this proof we use  $a \in \{0, \dots, n-1\}$  and  $\alpha \in \{1, \dots, n-1\}$  to denote components of Riemannian tensors for  $g_+$  defined with respect to the frame  $\{Y_0, Y_\alpha\}$ . For example,  $(g_+)_{00} = 1$ ,  $(g_+)_{0\alpha} = 0$ , and  $(g_+)_{\alpha\beta} = g_{\alpha\beta}$ , where  $g_{\alpha\beta}$  are the components of  $g := r^2 g_+$  with respect to  $\{\partial_r, \partial_{x^\alpha}\}$ . Thus  $(g_+^{-1})^{00} = 1$ ,  $(g_+^{-1})^{0\alpha} = 0$ , and  $(g_+^{-1})^{\alpha\beta} = g^{\alpha\beta}$ , where  $g^{\alpha\beta}$  are the components of  $g^{-1}$  with respect to  $\{\partial_r, \partial_{x^\alpha}\}$ .

Define  $\gamma_{ab}^c$  by  $\nabla_{Y_a}^{g_+} Y_b = \gamma_{ab}^c Y_c$ . Albin showed [1, Equation (3.5)] that

$$\gamma_{ab}^c = r\Gamma_{ab}^c - \delta_b^0 \delta_a^c + \delta_0^c g_{ab},$$

where  $\Gamma_{ab}^c$  denotes the Christoffel symbols of  $g$ . In particular,

$$(4.5) \quad \begin{aligned} \gamma_{\alpha\beta}^\epsilon &= r\Gamma_{\alpha\beta}^\epsilon, & \gamma_{\alpha\beta}^0 &= -\frac{1}{2}r\partial_r g_{\alpha\beta} + g_{\alpha\beta}, \\ \gamma_{\alpha 0}^\epsilon &= \frac{1}{2}r g^{\epsilon\zeta} \partial_r g_{\alpha\zeta} - \delta_\alpha^\epsilon, & \gamma_{0\alpha}^\epsilon &= \frac{1}{2}r g^{\epsilon\zeta} \partial_r g_{\alpha\zeta}, \end{aligned}$$

and all other components of  $\gamma_{ab}^c$  vanish. Denote by  $\mathcal{F}((-1)^\ell)$  the set of all functions  $u \in C^\infty(X)$  such that the asymptotic expansion

$$r^{(-1+(-1)^\ell)/2}u = u_{(0)} + u_{(2)}r^2 + \cdots + u_{(n-2)}r^{n-2} + u_{(n-1)}r^{n-1} + O(r^n)$$

is even to order  $O(r^{n-1})$ . That is,  $\mathcal{F}(1)$  consists of the functions that have an even expansion in  $r$  to order  $O(r^{n-1})$ , and  $\mathcal{F}(-1)$  consists of the functions that have an odd expansion in  $r$  to order  $O(r^n)$ . Note that if  $u \in \mathcal{F}((-1)^\ell)$  and  $v \in \mathcal{F}((-1)^m)$ , then  $uv \in \mathcal{F}((-1)^{\ell+m})$ . Since  $g_{ab}, g^{ab} \in \mathcal{F}(1)$ , we may freely raise and lower indices within  $\mathcal{F}((-1)^\ell)$ . Equations (4.5) imply that  $\gamma_{ab}^c \in \mathcal{F}((-1)^{\#T(a,b,c)})$ , where

$$\#Te := |\{1 \leq j \leq k : e_j \neq 0\}|$$

denotes the number of nonzero terms in the multi-index  $e \in \{0, \dots, n-1\}^k$ .

Consider now the components  $R_{ab}{}^c{}_d$  of the Riemann curvature tensor of  $g_+$ ,

$$R_{ab}{}^c{}_d Y_c = \nabla_{Y_a}^{g_+} \nabla_{Y_b}^{g_+} Y_d - \nabla_{Y_b}^{g_+} \nabla_{Y_a}^{g_+} Y_d - \nabla_{[Y_a, Y_b]}^{g_+} Y_d.$$

Albin showed [1, Equation (3.6)] that

$$(4.6) \quad R_{ab}{}^c{}_d = \gamma_{ae}^c \gamma_{bd}^e - \gamma_{be}^c \gamma_{ad}^e + Y_a(\gamma_{bd}^c) - Y_b(\gamma_{ad}^c) - \delta_a^0 \gamma_{bd}^c + \delta_b^0 \gamma_{ad}^c.$$

Since  $Y_0(\mathcal{F}((-1)^m)) \subseteq \mathcal{F}((-1)^m)$  and  $Y_\alpha(\mathcal{F}((-1)^m)) \subseteq \mathcal{F}((-1)^{m+1})$ , we deduce that  $R_{abcd} \in \mathcal{F}((-1)^{\#T(a,b,c,d)})$ .

Denote the components of  $\nabla^\ell \text{Rm}$  by

$$(4.7) \quad R_{abcd;e_1 \dots e_\ell} := Y_{e_\ell}(R_{abcd;e_1 \dots e_{\ell-1}}) - \gamma_{e_\ell a}^f R_{fbcd;e_1 \dots e_{\ell-1}} \\ - \cdots - \gamma_{e_\ell e_{\ell-1}}^f R_{abcd;e_1 \dots e_{\ell-2} f}.$$

It follows from induction that

$$R_{abcd;e_1 \dots e_\ell} \in \mathcal{F}((-1)^{\#T(a,b,c,d,e)}).$$

Now, since  $F$  is a tensor Riemannian invariant of rank 1, the function  $\langle F, \mathbf{n} \rangle$  is a linear combination of complete contractions of

$$(4.8) \quad \nabla^{N_1} \text{Rm} \otimes \cdots \otimes \nabla^{N_p} \text{Rm} \otimes \mathbf{n},$$

where  $N_1, \dots, N_p$  denote powers. Since  $\mathbf{n} = -Y_0$  along  $\partial M$ , any such complete contraction is a linear combination of products of  $R_{abcd;e_1 \dots e_\ell}$ , and each product has an even number of tangential indices. From the previous discussion, we conclude that  $\langle F, \mathbf{n} \rangle^{g_+} \in \mathcal{F}(1)$ . Moreover, the coefficient of  $r^{n-1}$  of  $\langle F, \mathbf{n} \rangle^{g_+}$  is computed by multiplying the coefficient of  $r^{n-1}$  in one factor  $R_{abcd;e_1 \dots e_\ell}$  by the coefficients of  $r^0$  in the remaining factors.

Given  $v \in C^\infty(M)$ , denote by  $v^{(0)}$  and  $v^{(n-1)}$  the coefficients of  $r^0$  and  $r^{n-1}$ , respectively, in the Taylor expansion of  $v$  near  $\partial M$ . Equations (4.5) and (4.6) yield

$$\begin{aligned} (\gamma_{\alpha\beta}^0)^{(0)} &= h_{\alpha\beta}, & (\gamma_{\alpha\beta}^0)^{(n-1)} &= -\frac{n-3}{2}k_{\alpha\beta}, \\ (\gamma_{\alpha 0}^\beta)^{(0)} &= -\delta_\alpha^\beta, & (\gamma_{\alpha 0}^\beta)^{(n-1)} &= \frac{n-1}{2}h^{\beta\epsilon}k_{\alpha\epsilon}, \\ (\gamma_{0\alpha}^\beta)^{(0)} &= 0, & (\gamma_{0\alpha}^\beta)^{(n-1)} &= \frac{n-1}{2}h^{\beta\epsilon}k_{\alpha\epsilon}, \\ (R_{\alpha\beta\epsilon\zeta})^{(0)} &= -2h_{\alpha[\epsilon}h_{\zeta]\beta}, & (R_{\alpha\beta\epsilon\zeta})^{(n-1)} &= (n-3)(h_{\alpha[\epsilon}k_{\zeta]\beta} - h_{\beta[\epsilon}k_{\zeta]\alpha}), \\ (R_{\alpha 0\beta 0})^{(0)} &= -h_{\alpha\beta}, & (R_{\alpha 0\beta 0})^{(n-1)} &= -\frac{n^2-4n+5}{2}k_{\alpha\beta}. \end{aligned}$$

Combining this with Equation (4.7), the previous paragraph, and naturality implies that the coefficient of  $r^{n-1}$  of  $\langle F, \mathbf{n} \rangle^{g+}$  must be proportional to  $\text{tr}_h k = 0$ .  $\square$

Combining Proposition 3.6 and Lemma 4.1 yields our formula for the renormalized integral of a straightenable scalar Riemannian invariant.

*Proof of Theorem 1.4.* Let  $I$  be a straightenable scalar Riemannian invariant of weight  $-2k \geq -n$ . Set  $\ell := n/2 - k$  and let  $I_\ell$  be the straightenable scalar Riemannian invariant determined by Proposition 3.6. Let  $\mathcal{I}_\ell$  be the conformal invariant determined ambiently by  $I_\ell$ . On the one hand, since  $\mathcal{I}_\ell$  is conformally invariant of weight  $-n$ , we see that

$$\int_M^R \mathcal{I}_\ell dV = \int_M \mathcal{I}_\ell dV.$$

On the other hand, Lemma 3.2, Equation (3.2), and Lemma 4.1 imply that

$$\int_M^R \mathcal{I}_\ell dV = \int_M^R I_\ell dV = 2^{n/2-k} \frac{(n/2-1)!(n-2k-1)!!}{(k-1)!} \int_M^R I dV. \quad \square$$

## 5. APPLICATIONS

Theorem 1.4 and Lemma 3.4 together compute the renormalized integral of any scalar polynomial in the Weyl tensor. This allows us to derive an explicit formula expressing the renormalized volume of an even-dimensional Poincaré–Einstein manifold in terms of its Euler characteristic and the convergent integral of a scalar conformal invariant. The key observation is that the Pfaffian  $\text{Pf} = \text{Pf}_{n/2}(\text{Rm})$  of an Einstein manifold is a scalar polynomial in the Weyl tensor (cf. [1, Lemma 4.4]).

**Lemma 5.1.** *Let  $(M^n, g)$  be an even-dimensional Einstein manifold. Then*

$$\text{Pf} = \sum_{\ell=0}^{n/2} (n-2\ell-1)!! \left(\frac{2}{n}\mathbf{J}\right)^{n/2-\ell} \text{Pf}_\ell(W).$$

*Proof.* The assumption that  $(M^n, g)$  is Einstein implies that

$$R_{a_1 a_2}{}^{b_1 b_2} = W_{a_1 a_2}{}^{b_1 b_2} + \frac{4}{n} \mathbf{J} \delta_{a_1 a_2}^{b_1 b_2}.$$

We deduce from Equation (1.4) that

$$\begin{aligned} \text{Pf} &= 2^{-n/2} (n-1)!! \sum_{\ell=0}^{n/2} \binom{n/2}{\ell} \left(\frac{4}{n}\mathbf{J}\right)^{n/2-\ell} \\ &\quad \times \delta_{b_1 \dots b_n}^{a_1 \dots a_n} W_{a_1 a_2}^{b_1 b_2} \dots W_{a_{2\ell-1} a_{2\ell}}^{b_{2\ell-1} b_{2\ell}} \delta_{a_{2\ell+1}}^{b_{2\ell+1}} \dots \delta_{a_n}^{b_n}. \end{aligned}$$

The conclusion follows from the identity

$$\delta_{b_1 \dots b_k}^{a_1 \dots a_k} \delta_{a_k}^{b_k} = \frac{n-k+1}{k} \delta_{b_1 \dots b_{k-1}}^{a_1 \dots a_{k-1}}. \quad \square$$

Since  $\text{Pf}_\ell(W)$  is a straightenable scalar Riemannian invariant of weight  $-2\ell$ , we can apply Theorem 1.4 to produce an explicit ambient formula for the scalar conformal invariant in Equation (1.3). More generally, we produce a local formula for the Pfaffian of an Einstein manifold modulo divergences.

**Corollary 5.2.** *Let  $(M^n, g)$  be an even-dimensional Einstein manifold. Then*

$$\text{Pf} \equiv (n-1)!! \left(\frac{2}{n}\mathbf{J}\right)^{n/2} + \sum_{\ell=2}^{n/2} (-2)^{\ell-n/2} \frac{(\ell-1)!}{(n/2-1)!} \mathcal{P}_{\ell,n} \pmod{\text{im } \Delta},$$

where  $\mathcal{P}_{\ell,n} := i^* \left( \tilde{\Delta}^{n/2-\ell} \text{Pf}_\ell(\tilde{W}) \right)$ .

*Proof.* First note that  $\text{Pf}_1(W) = 0$ . We then deduce from Lemma 5.1 that

$$\text{Pf} = (n-1)!! \left(\frac{2}{n}\mathbf{J}\right)^{n/2} + \sum_{\ell=2}^{n/2} (n-2\ell-1)!! \left(\frac{2}{n}\mathbf{J}\right)^{n/2-\ell} \text{Pf}_\ell(W).$$

Combining this with Proposition 3.6 yields

$$(\text{Pf}_\ell(W))_{n/2-\ell} \equiv \left(-\frac{4}{n}\mathbf{J}\right)^{n/2-\ell} \frac{(n/2-1)!(n-2\ell-1)!!}{(\ell-1)!} \text{Pf}_\ell(W) \pmod{\text{im } \Delta}.$$

The final conclusion follows from Lemma 3.2.  $\square$

Corollary 5.2 yields our explicit formula for the (renormalized) volume of an even-dimensional (Poincaré-)Einstein manifold.

*Proof of Theorem 1.1.* Apply Equation (1.1), Lemma 4.1, and Corollary 5.2.  $\square$

*Proof of Theorem 1.2.* Apply Corollary 5.2.  $\square$

By direct computation, any complete contraction of  $W^{\otimes 2}$  is proportional to

$$(5.1a) \quad \mathcal{W}_{2,1} := W_{abcd} W^{abcd},$$

any complete contraction of  $W^{\otimes 3}$  is a linear combination of

$$(5.1b) \quad \mathcal{W}_{3,1} := W_{ab}{}^{cd} W_{cd}{}^{ef} W_{ef}{}^{ab}, \quad \mathcal{W}_{3,2} := W_a{}^c{}_b{}^d W_c{}^e{}_d{}^f W_e{}^a{}_f{}^b;$$

and any complete contraction of  $W^{\otimes 4}$  is a linear combination of

$$(5.1c) \quad \begin{aligned} \mathcal{W}_{4,1} &:= W_{abcd} W^{abcd} W_{efgh} W^{efgh}, & \mathcal{W}_{4,2} &:= W_{ab}{}^{cd} W_{cd}{}^{ef} W_{ef}{}^{gh} W_{gh}{}^{ab}, \\ \mathcal{W}_{4,3} &:= W_{acde} W_b{}^{cde} W_a{}^{fg} W^{bfgh}, & \mathcal{W}_{4,4} &:= W_{abcd} W_{ef}{}^{cd} W_a{}^e{}_g{}^h W^{bgfh}, \\ \mathcal{W}_{4,5} &:= W_{abcd} W_{ef}{}^{cd} W_a{}^e{}_g{}^h W^{bfgh}, & \mathcal{W}_{4,6} &:= W_a{}^c{}_b{}^d W_c{}^e{}_d{}^f W_e{}^g{}_f{}^h W_g{}^a{}_h{}^b, \\ \mathcal{W}_{4,7} &:= W_a{}^c{}_b{}^d W_{ecfd} W_a{}^e{}_g{}^h W^{bgfh}. \end{aligned}$$

We express  $\text{Pf}_\ell(W)$ ,  $\ell \leq 4$ , in terms of these contractions:

**Lemma 5.3.** *Let  $(M^n, g)$  be a Riemannian manifold. Then*

$$\begin{aligned}\text{Pf}_2(W) &= \frac{1}{8}\mathcal{W}_{2,1}, \\ \text{Pf}_3(W) &= \frac{1}{12}\mathcal{W}_{3,1} - \frac{1}{6}\mathcal{W}_{3,2}, \\ \text{Pf}_4(W) &= \frac{1}{128}\mathcal{W}_{4,1} + \frac{1}{64}\mathcal{W}_{4,2} - \frac{1}{8}\mathcal{W}_{4,3} - \frac{1}{4}\mathcal{W}_{4,4} + \frac{1}{8}\mathcal{W}_{4,5} + \frac{1}{8}\mathcal{W}_{4,6} - \frac{1}{4}\mathcal{W}_{4,7}.\end{aligned}$$

*Proof.* Recall that the Weyl tensor is totally trace-free. Direct computation from the definition of  $\text{Pf}_\ell$  yields

$$\begin{aligned}\text{Pf}_2(W) &= \frac{1}{222!}W_{abcd}W^{abcd}, \\ \text{Pf}_3(W) &= \frac{1}{233!}(2W_{ab}{}^{ef}W_{cd}{}^{ab}W_{ef}{}^{cd} - 8W_{ab}{}^{ce}W_{cd}{}^{af}W_{ef}{}^{bd}), \\ \text{Pf}_4(W) &= \frac{1}{244!}\left(3W_{ab}{}^{cd}W_{cd}{}^{ab}W_{ef}{}^{gh}W_{gh}{}^{ef} + 6W_{ab}{}^{cd}W_{cd}{}^{ef}W_{ef}{}^{gh}W_{gh}{}^{ab} \right. \\ &\quad \left. - 48W_{ab}{}^{cg}W_{cd}{}^{ab}W_{ef}{}^{dh}W_{gh}{}^{ef} - 96W_{ab}{}^{eg}W_{cd}{}^{ab}W_{ef}{}^{ch}W_{gh}{}^{df} \right. \\ &\quad \left. + 48W_{ab}{}^{ce}W_{cd}{}^{ag}W_{ef}{}^{bh}W_{gh}{}^{df} - 96W_{ab}{}^{eg}W_{cd}{}^{ah}W_{ef}{}^{bc}W_{gh}{}^{df}\right).\end{aligned}$$

The conclusion now follows from the Bianchi identity  $W_{[abc]d} = 0$ :

$$\begin{aligned}W_{ab}{}^{ce}W_{cd}{}^{af}W_{ef}{}^{bd} &= W_{abc}{}^eW^{afcd}W^b{}_{fed} - W_{abc}{}^eW^{afcd}W^b{}_{efd} \\ &= W_{abc}{}^eW^{afcd}W^b{}_{fed} - \frac{1}{4}W_{acb}{}^eW^{acfd}W^b{}_{efd}\end{aligned}$$

and

$$\begin{aligned}W_{ab}{}^{eg}W_{cd}{}^{ab}W_{ef}{}^{ch}W_{gh}{}^{df} &= W_{cd}{}^{ab}W_{abeg}W^{chef}W^d{}_{fgh} \\ &= W_{cd}{}^{ab}W_{abeg}W^{chef}W^d{}_{hgf} + W_{cd}{}^{ab}W_{abeg}W^{chef}W^{dg}{}_{fh} \\ &= W_{cd}{}^{ab}W_{abeg}W^{chef}W^d{}_{hgf} - \frac{1}{2}W_{cd}{}^{ab}W_{abeg}W^{cefh}W^{dg}{}_{fh}.\end{aligned}\quad \square$$

Lemma 5.3 allows us to compute  $\mathcal{P}_{\ell,n}$  for all integers  $2 \leq \ell \leq 4$ .

**Corollary 5.4.** *Let  $(M^n, g)$  be an even-dimensional Riemannian manifold. Then*

$$\begin{aligned}\mathcal{P}_{2,n} &= \frac{1}{8}i^*(\tilde{\Delta}^{n/2-2}\widetilde{\mathcal{W}}_{2,1}), & \text{if } n \geq 4, \\ \mathcal{P}_{3,n} &= \frac{1}{12}i^*(\tilde{\Delta}^{n/2-3}(\widetilde{\mathcal{W}}_{3,1} - 2\widetilde{\mathcal{W}}_{3,2})), & \text{if } n \geq 6, \\ \mathcal{P}_{4,n} &= \frac{1}{128}i^*(\tilde{\Delta}^{n/2-4}(\widetilde{\mathcal{W}}_{4,1} + 2\widetilde{\mathcal{W}}_{4,2} - 16\widetilde{\mathcal{W}}_{4,3} - 32\widetilde{\mathcal{W}}_{4,4} \\ &\quad + 16\widetilde{\mathcal{W}}_{4,5} + 16\widetilde{\mathcal{W}}_{4,6} - 32\widetilde{\mathcal{W}}_{4,7})), & \text{if } n \geq 8.\end{aligned}$$

*Proof.* Combine Lemma 5.3 with the definition of  $\mathcal{P}_{\ell,n}$  in Theorem 1.1.  $\square$

We conclude this paper by computing three additional examples of renormalized curvature integrals which indicate the applicability of our results.

Our first example, when combined with the second Bianchi identity, computes the renormalized integral of any complete contraction of  $(\nabla W)^{\otimes 2}$ .

**Example 5.5.** Let  $(M^n, g_+)$ ,  $n \geq 6$ , be an even-dimensional Poincaré–Einstein manifold. Then

$$\int^R |\nabla W|^2 dV = \frac{2^{3-n/2} 2!}{(n/2 - 1)!(n - 7)!!} \int_M i^*(\tilde{\Delta}^{n/2-3} \tilde{L}) dV,$$

where

$$\tilde{L} := \frac{n-1}{2(n-5)} \tilde{\Delta} \tilde{\mathcal{W}}_{2,1} + \tilde{\mathcal{W}}_{3,1} + 4\tilde{\mathcal{W}}_{3,2}.$$

Indeed, Lemma 4.1 yields

$$\int^R |\nabla W|^2 dV = - \int^R \langle W, \Delta W \rangle dV,$$

while the second Bianchi identity and the assumption  $\text{Ric}_{g_+} = -(n-1)g_+$  yield

$$(5.2) \quad \Delta W_{abcd} = -2(n-1)W_{abcd} - W_{ab}{}^{ef}W_{efcd} - 2W_{aecf}W_b{}^e{}_d{}^f + 2W_{aedf}W_b{}^e{}_c{}^f.$$

Combining these observations yields

$$\int^R |\nabla W|^2 dV = \int^R (2(n-1)\mathcal{W}_{2,1} + \mathcal{W}_{3,1} + 4\mathcal{W}_{3,2}) dV.$$

The claim now follows from Theorem 1.4.

The explanation of our second example indicates how to compute the renormalized integral of a complete contraction of  $(\nabla^2 W)^{\otimes 2}$  or  $W \otimes (\nabla W)^{\otimes 2}$ .

**Example 5.6.** Let  $(M^n, g_+)$ ,  $n \geq 8$ , be an even-dimensional Poincaré–Einstein manifold. Then

$$\int^R |\nabla^2 W|^2 dV = \frac{2^{4-n/2} 3!}{(n/2 - 1)!(n - 9)!!} \int_M i^*(\tilde{\Delta}^{n/2-4} \tilde{L}) dV,$$

where

$$\begin{aligned} \tilde{L} := & \frac{(n-1)(3n+1)}{12(n-5)(n-7)} \tilde{\Delta}^2 \tilde{\mathcal{W}}_{2,1} + \frac{5n+3}{6(n-7)} \tilde{\Delta} \tilde{\mathcal{W}}_{3,1} + \frac{2(5n+3)}{3(n-7)} \tilde{\Delta} \tilde{\mathcal{W}}_{3,2} \\ & + \tilde{\mathcal{W}}_{4,2} + 4\tilde{\mathcal{W}}_{4,3} + 8\tilde{\mathcal{W}}_{4,4} - 6\tilde{\mathcal{W}}_{4,5} + 8\tilde{\mathcal{W}}_{4,6} - 8\tilde{\mathcal{W}}_{4,7}. \end{aligned}$$

Indeed, the Ricci identity and the assumption  $\text{Ric}_{g_+} = -(n-1)g_+$  yield

$$\begin{aligned} \Delta \nabla_e W_{abcd} = & \nabla_e \Delta W_{abcd} - (n+3) \nabla_e W_{abcd} \\ & - 4W_{feh[a} \nabla^f W^h{}_{b]cd} - 4W_{feh[c} \nabla^f W^h{}_{d]ab}. \end{aligned}$$

Combining this with Lemma 4.1 yields

$$\begin{aligned} \int^R |\nabla^2 W|^2 dV = & \int^R (|\Delta W|^2 - (n+3) \langle W, \Delta W \rangle \\ & + 8W^{abcd} W_{efha} \nabla^e \nabla^f W^h{}_{bcd}) dV. \end{aligned}$$

On the one hand, the first Bianchi identity yields

$$\begin{aligned} W_{aecf} W_b{}^e{}_d{}^f W^{agdh} W_g{}^b{}_c{}^h &= W_{aecf} W_b{}^e{}_d{}^f W^{agdh} (W_g{}^c{}_h{}^b + W^{bc}{}_{gh}) \\ &= W_{aecf} (W_d{}^e{}_b{}^f + W_{bd}{}^{ef}) W^{agdh} W_g{}^c{}_h{}^b + \frac{1}{2} W_{aecf} W_b{}^e{}_d{}^f W^{agdh} W^{bc}{}_{gh} \\ &= \mathcal{W}_{4,7} - \mathcal{W}_{4,4} + \frac{1}{4} \mathcal{W}_{4,5}. \end{aligned}$$

Combining this with Equation (5.2) yields

$$\begin{aligned} \int^R (|\Delta W|^2 - (n+3)\langle W, \Delta W \rangle) dV &= \int^R \left( 2(n-1)(3n+1)\mathcal{W}_{2,1} \right. \\ &\quad \left. + (5n-1)\mathcal{W}_{3,1} + 4(5n-1)\mathcal{W}_{3,2} \right. \\ &\quad \left. + \mathcal{W}_{4,2} + 16\mathcal{W}_{4,4} - 2\mathcal{W}_{4,5} + 8\mathcal{W}_{4,6} - 8\mathcal{W}_{4,7} \right) dV. \end{aligned}$$

On the other hand, the Ricci identity yields

$$\begin{aligned} \int^R 8W^{abcd}W_{efha}\nabla^e\nabla^f W^h{}_{bcd} dV \\ = \int^R 4\left(\mathcal{W}_{3,1} + 4\mathcal{W}_{3,2} + \mathcal{W}_{4,3} - 2\mathcal{W}_{4,4} - \mathcal{W}_{4,5}\right) dV. \end{aligned}$$

Combining these formulas with Theorem 1.4 yields the final claim.

Our final example illustrates a more general approach to applying Propositions 3.6 and 3.8 to compute renormalized curvature integrals.

**Example 5.7.** Let  $(M^n, g_+)$ ,  $n \geq 10$ , be an even-dimensional Poincaré–Einstein manifold. Then

$$\int^R |\nabla|W|^2|^2 dV = \frac{2^{5-n/2}4!}{(n/2-1)!(n-11)!!} \int_M i^*(\tilde{\Delta}^{n/2-5}\tilde{L}) dV,$$

where

$$\tilde{L} := -|\tilde{W}|^2\tilde{\Delta}|\tilde{W}|^2 + \frac{(n-5)}{2(n-9)}\tilde{\Delta}|\tilde{W}|^4.$$

Indeed, Lemma 4.1 yields

$$\begin{aligned} \int^R |\nabla|W|^2|^2 dV &= \int^R (-|W|^2\Delta|W|^2) dV \\ &= \int^R (-|W|^2(|W|^2)_1 + 4(n-5)|W|^4) dV, \end{aligned}$$

where  $(|W|^2)_1 := \Delta|W|^2 + 4(n-5)|W|^2$  is the specialization to  $(M^n, g_+)$  of the straightenable scalar Riemannian invariant produced by Proposition 3.6. The conclusion now follows from Theorem 1.4.

The general principle is that Propositions 3.6 and 3.8 allow one to write certain divergences as a linear combination of a straightenable and a lower-order tensor Riemannian invariant. We speculate that, when combined with the Bianchi identity, this procedure allows one to compute *any* renormalized curvature integral.

#### ACKNOWLEDGEMENTS

This project is a part of the AIM SQuaRE “Global invariants of Poincaré–Einstein manifolds and applications”. We thank the American Institute for Mathematics for their support. We also thank Matthew Gursky for useful comments on an early draft of this paper.



## FUNDING

JSC acknowledges support from a Simons Foundation Collaboration Grant for Mathematicians, ID 524601. WY acknowledges support from the NSFC (Grant No. 12071489, No. 12025109).

## DECLARATIONS

The authors have no relevant financial or non-financial interests to disclose.

## DATA AVAILABILITY

Data sharing not applicable to this article as no datasets were generated or analyzed during the study.

## REFERENCES

- [1] P. Albin, *Renormalizing curvature integrals on Poincaré-Einstein manifolds*, Adv. Math. **221** (2009), no. 1, 140–169. MR2509323
- [2] S. Alexakis, *The decomposition of global conformal invariants*, Annals of Mathematics Studies, vol. 182, Princeton University Press, Princeton, NJ, 2012. MR2918125
- [3] M. T. Anderson,  *$L^2$  curvature and volume renormalization of AHE metrics on 4-manifolds*, Math. Res. Lett. **8** (2001), no. 1-2, 171–188. MR1825268
- [4] A. Avez, *Characteristic classes and Weyl tensor: Applications to general relativity*, Proc. Nat. Acad. Sci. U.S.A. **66** (1970), 265–268. MR263098
- [5] T. N. Bailey, M. G. Eastwood, and C. R. Graham, *Invariant theory for conformal and CR geometry*, Ann. of Math. (2) **139** (1994), no. 3, 491–552. MR1283869
- [6] J. S. Case, Y.-J. Lin, and W. Yuan, *Conformally variational Riemannian invariants*, Trans. Amer. Math. Soc. **371** (2019), no. 11, 8217–8254. MR3955546
- [7] J. S. Case, Y.-J. Lin, and W. Yuan, *Curved Versions of the Ovsienko–Redou Operators*, Int. Math. Res. Not. IMRN **19** (2023), 16904–16929. MR4651902
- [8] S.-Y. A. Chang and Y. Ge, *Compactness of conformally compact Einstein manifolds in dimension 4*, Adv. Math. **340** (2018), 588–652. MR3886176
- [9] S.-Y. A. Chang, Y. Ge, and J. Qing, *Compactness of conformally compact Einstein 4-manifolds II*, Adv. Math. **373** (2020), 107325, 33. MR4130465
- [10] ———, *Compactness of conformally compact Einstein 4-manifolds II*, Adv. Math. **373** (2020), 107325, 33. MR4130465
- [11] S.-Y. A. Chang, J. Qing, and Paul. Yang, *On the renormalized volumes for conformally compact Einstein manifolds*, J. Math. Sciences (N. Y.) **149** (2008), no. 6, 1755–1769. MR2336463
- [12] S.-S. Chern, *On curvature and characteristic classes of a Riemann manifold*, Abh. Math. Sem. Univ. Hamburg **20** (1955), 117–126. MR0075647
- [13] P. T. Chruściel, E. Delay, J. M. Lee, and D. N. Skinner, *Boundary regularity of conformally compact Einstein metrics*, J. Differential Geom. **69** (2005), no. 1, 111–136. MR2169584
- [14] C. Fefferman and C. R. Graham, *The ambient metric*, Annals of Mathematics Studies, vol. 178, Princeton University Press, Princeton, NJ, 2012. MR2858236
- [15] M. d. M. González and J. Qing, *Fractional conformal Laplacians and fractional Yamabe problems*, Anal. PDE **6** (2013), no. 7, 1535–1576. MR3148060
- [16] C. R. Graham, *Volume and area renormalizations for conformally compact Einstein metrics*, The Proceedings of the 19th Winter School “Geometry and Physics” (Srń, 1999), 2000, pp. 31–42. MR1758076
- [17] C. R. Graham and M. Zworski, *Scattering matrix in conformal geometry*, Invent. Math. **152** (2003), no. 1, 89–118. MR1965361
- [18] A. Juhl, *Explicit formulas for GJMS-operators and Q-curvatures*, Geom. Funct. Anal. **23** (2013), no. 4, 1278–1370. MR3077914
- [19] J. Maldacena, *The large  $N$  limit of superconformal field theories and supergravity*, Adv. Theor. Math. Phys. **2** (1998), no. 2, 231–252. MR1633016
- [20] E. Witten, *Anti de Sitter space and holography*, Adv. Theor. Math. Phys. **2** (1998), no. 2, 253–291. MR1633012

DEPARTMENT OF MATHEMATICS, PENN STATE UNIVERSITY, UNIVERSITY PARK, PA 16802, USA  
*Email address:* `jscase@psu.edu`

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, HILL CENTER FOR THE MATHEMATICAL SCIENCES, 110 FRELINGHUYSEN RD., PISCATAWAY, NJ 08854  
*Email address:* `ayush.khaitan@rutgers.edu`

DEPARTMENT OF MATHEMATICS, STATISTICS, AND PHYSICS, WICHITA STATE UNIVERSITY, WICHITA, KS 67260, USA  
*Email address:* `yueh-ju.lin@wichita.edu`

DEPARTMENT OF MATHEMATICS AND STATISTICS, TEXAS TECH UNIVERSITY, LUBBOCK, TX 79409-1042, USA  
*Email address:* `aatyrrel@ttu.edu`

DEPARTMENT OF MATHEMATICS, SUN YAT-SEN UNIVERSITY, GUANGZHOU, GUANGDONG 510275, CHINA  
*Email address:* `yuanw9@mail.sysu.edu.cn`