FRACTIONAL-DIFFRACTION-OPTICS CAUCHY PROBLEM: RESOLVENT-FUNCTION SOLUTION OF THE MATRIX INTEGRAL EQUATION

Murat O. Mamchuev ¹, Felix N. Chukhovskii ¹

Abstract. The fractional diffraction optics theory has been elaborated using the Green function technique. The optics-fractional equation describing the diffraction X-ray scattering by imperfect crystals has been derived as the fractional matrix integral Fredholm–Volterra equation of the second kind. In the paper, to solve the Cauchy problems, the Liouville–Neumann-type series formalism has been used to build up the matrix Resolvent-function solution. In the case when the imperfect crystal-lattice elastic displacement field is the linear function $f(\mathbf{R}) = ax + b$, a, b = const, the explicit solution of the diffraction-optics Cauchy problem has been obtained and analyzed for arbitrary fractional-order-parameter $\alpha, \alpha \in (0, 1]$.

MSC 2010: Primary 35F35;

Secondary 35F40, 35A08, 35C15, 35L40, 45F05, 35Q70, 35Q92;

Key Words and Phrases: Diffraction optics system of fractional differential equations, the Gerasimov–Caputo differential operator, the Cauchy problem, matrix Resolvent-function solution, matrix Fredholm–Volterra integral equation of the second kind, the Liouville–Neumann-type series;

1 Introduction

Usually, the diffraction optics (DO) has been formulated and based on the known differential Takagi–Taupin (TT) equations when the fractional-order-parameter $\alpha=1$ (see, e.g., [1], [2], [3], [4], [5], [6], [7], [8] for details). At the same time, in the last decades, substantial progress has been achieved in mathematical physics using equations with fractional-order derivatives [9], [10], [11], [12], [13], [14], [15], [16]. Indeed, the different theory-physics models for solving the Cauchy problems using systems of fractional differential equations have been treated in [17], [18], [19], [20], [21], [22], [23], [24], [25].

Following this logic, one can push one step further in the DO now founded on the TT-type equations with the fractional derivatives of the arbitrary order $\alpha \in (0, 1]$ along the crystal depth.

In the paper, using Green's function technique, one derives the matrix Fredholm-Volterra integral equation of the second kind and builds up the matrix Resolvent-function solution of the DO Cauchy problem. A goal of working is to develop the integral formalism of the DO theory earlier proposed

by authors [26] and to build up the matrix Resovent-function solution of the fractional Cauchy problem.

As an example, for an arbitrary fractional-order-parameter (FOP) α , $\alpha \in (0,1]$, when the crystal-lattice displacement field function $f(\mathbf{R})$ is a linear function, namely: $f(\mathbf{R}) = ax + b$, a, b = const, one finds out an explicit analytical solution of the DO Cauchy problem.

Accordingly, the original system of fractional DO equations has the form (cf. [26])

$$\begin{pmatrix}
\partial_{0t}^{\alpha} - \frac{\partial}{\partial x} & 0 \\
0 & \partial_{0t}^{\alpha} + \frac{\partial}{\partial x}
\end{pmatrix}
\begin{pmatrix}
E_{0}(x, t) \\
E_{h}(x, t)
\end{pmatrix} = i \begin{pmatrix}
\gamma & \sigma \exp[if(x, t)] \\
\sigma \exp[-if(x, t)] & \gamma
\end{pmatrix}
\begin{pmatrix}
E_{0}(x, t) \\
E_{h}(x, t)
\end{pmatrix}, (1.1)$$

with the DO Cauchy problem's condition

$$\begin{pmatrix} E_0(x,0) \\ E_h(x,0) \end{pmatrix} = \begin{pmatrix} \varphi_0(x) \\ \varphi_h(x) \end{pmatrix}, \quad -\infty < x < \infty, \tag{1.2}$$

where $\varphi_0(x)$ and $\varphi_h(x)$ are the given real-valued functions.

2 Preliminaries

Following up to [13], the Gerasimov-Caputo fractional derivative beginning at point a is determined as (cf. [13])

$$\partial_{at}^{\nu}g(t) = \operatorname{sgn}^{n}(t-a)D_{at}^{\nu-n}\frac{d^{n}}{dt^{n}}g(t), \quad n-1 < \nu \le n, \quad n \in \mathbb{N},$$
 (2.1)

where D_{ay}^{ν} is the Riemann–Liouville fractional integro-differential operator of order ν is equal to

$$D_{ay}^{\nu}g(y) = \frac{\operatorname{sgn}(y-a)}{\Gamma(-\nu)} \int_{a}^{y} \frac{g(s)ds}{|y-s|^{\nu+1}}, \quad \nu < 0,$$

for $\nu \geq 0$ the operator D_{ay}^{ν} can be determined by recursive relation

$$D_{ay}^{\nu}g(y) = \operatorname{sgn}(y - a)\frac{d}{dy}D_{ay}^{\nu - 1}g(y), \quad \nu \ge 0, \tag{2.2}$$

 $\Gamma(z)$ is the Euler gamma-function.

Note that in the limit case of the FOP $\alpha = 1$ the operator $\partial_{0t}^{\alpha} g(t)$ reduces to the standard derivative $\frac{d}{dt}g(t)$.

Formula for the composition of operators of fractional integration is valid as [13]

$$D_{0t}^{\nu}D_{0t}^{\delta}g(t) = D_{0t}^{\nu+\delta}g(t), \quad \nu < 0, \quad \delta < 0.$$
 (2.3)

There is a formula for fractional integration by parts, namely [13]:

$$\int_{0}^{t} g(t,\xi) D_{0\xi}^{\nu} h(\xi) d\xi = \int_{0}^{t} h(\xi) D_{t\xi}^{\nu} g(t,\xi) d\xi, \quad \nu < 0.$$
 (2.4)

Further, we denote the Fourier transform of the function f(x) by $(f(x))_k$, the Laplace transform of the function g(t) by $(g(t))_p$, and respectively, the double Fourier-Laplace transform of the function h(x,t) by $(h(x,t))_{k,p}$.

By using the following formula [14, p. 98]

$$[\partial_{0t}^{\alpha} H(x,t)]_p = p^{\alpha} [H(x,t)]_p - p^{\alpha-1} H(x,0),$$

one can get

$$[O_{\pm}^{\alpha}H(x,t)]_{k,p} = (p^{\alpha} \pm ik)[H(x,t)]_{k,p} - p^{\alpha-1}[H(x,0)]_k, \tag{2.5}$$

The following series

$$\phi(\beta, \rho; z) = \sum_{k=0}^{\infty} \frac{z^k}{k!\Gamma(\beta k + \rho)}, \quad \beta > -1, \quad \rho \in \mathbb{C}$$

defines the Wright function [27] which depends on two parameters ρ and μ . Accordingly, the following differentiation formula is valid [27]:

$$\frac{d}{dz}\phi(\beta,\rho;z) = \phi(\beta,\beta+\rho;z), \quad \beta > -1.$$
 (2.6)

Let $\beta \in (0,1)$, and $\mu, \nu \in \mathbb{R}$, and the inequality takes place

$$\beta \in (0,1), \quad 0 \le |\arg \lambda| < \frac{1-\beta}{2}\pi,$$

then the formula

$$D_{0y}^{\nu} y^{\mu-1} \phi(-\beta, \mu; -\lambda y^{-\beta}) = y^{\mu-\nu-1} \phi(-\beta, \mu - \nu; -\lambda y^{-\beta})$$
 (2.7)

is valid |28|.

The following formula take place [29]

$$(y^{\delta-1}\phi(-\beta,\mu;-ty^{-\beta}))_{p} = p^{-\mu}e^{-p^{\beta}t}.$$
 (2.8)

Correspondingly, the integrals

$$\int_{0}^{\infty} \frac{\cos kx}{k^2 + \rho^2} dk = \frac{\pi}{2\rho} e^{-\rho x},\tag{2.9}$$

$$\frac{1}{2\pi i} \int_{-i\infty}^{i\infty} \frac{e^{-x\sqrt{p^2+\sigma^2}}e^{pt}dp}{\sqrt{p^2+\sigma^2}} = J_0\left(\sigma\sqrt{t^2-x^2}\right)\Theta(t-|x|),\tag{2.10}$$

are held on, where $\Theta(x)$ is the Heaviside function, and $J_0(x)$ is the zero-order Bessel function of real argument.

The properties-in-details of the function

$$G_{\alpha,\gamma}(x,t) = \frac{1}{2} \int_{|x|}^{\infty} e^{i\gamma\tau} J_0\left(\sigma\sqrt{\tau^2 - x^2}\right) \frac{1}{t} \phi\left(-\alpha, 0; -\frac{\tau}{t^{\alpha}}\right) d\tau$$

have been analyzed in the [30].

Note, in limit of the FOP $\alpha = 1$, the function $G_{\alpha,\gamma}(x,t)$ can be cast into the form

$$\lim_{\alpha \to 1} G_{\alpha,\gamma}(x,t) = e^{i\gamma|x|} J_0\left(\sigma\sqrt{t^2 - x^2}\right) \Theta(t - |x|).$$

The estimate

$$\left| \frac{\partial^m}{\partial x^m} D_{0t}^{\nu} G_{\alpha,\gamma}(x,t) \right| \le C|x|^{-\theta} y^{\alpha(1-m+\theta)-\nu-1}, \quad \theta \ge 0$$
 (2.11)

holds on for arbitrary $m \in \mathbb{N} \cup \{0\}$ and $\nu \in \mathbb{R}$, where C is the positive constant [30].

Accordingly, the following estimate

$$\left| \frac{\partial^m}{\partial x^m} D_{0t}^{\nu} G_{\alpha,\gamma}(x,t) \right| \le C \exp\left(-\sigma_0 |x|^{\varepsilon} t^{-\alpha \varepsilon} \right), \tag{2.12}$$

holds on as $|x| \to \infty$ for all $t < \infty$ [30], and for arbitrary $m \in \mathbb{N} \cup \{0\}$, $\nu \in \mathbb{R}$; here and further $\sigma_0 < (1 - \alpha)\alpha^{\alpha\varepsilon}$, $\varepsilon = \frac{1}{1-\alpha}$; C is the positive constant.

Lemma 2.1. [21, p. 172]. For $0 < \mu, \nu < 1$ the estimate

$$\int_{r-\delta}^{x+\delta} |x-y|^{\mu-1} |y-\xi|^{\nu-1} dy \le C_{\gamma} |x-\xi|^{\gamma-1}, \tag{2.13}$$

holds on, where

$$\gamma = \min\{\mu, \nu\}, \quad C_{\gamma} = \left[\frac{2}{\gamma} + B(\mu, \nu)\right] (\delta_1 + \delta_2)^{\mu + \nu - \gamma}.$$

Lemma 2.2. [21, p. 176]. Let $0 < \eta < s < t < T$, $\sigma_2 < \sigma_3 < \sigma_1$, then the inequality

$$\int_{x+\delta}^{\pm\infty} \exp\left[-\frac{\sigma_1|x-y|^{\varepsilon}}{(t-s)^{\alpha\varepsilon}}\right] \exp\left[-\frac{\sigma_2|y-\xi|^{\varepsilon}}{(s-\eta)^{\alpha\varepsilon}}\right] dy \le C \exp\left[-\frac{\sigma_2|x-\xi|^{\varepsilon}}{(t-\eta)^{\alpha\varepsilon}}\right], (2.14)$$

takes place, where

$$C = \frac{\sigma_1 - \sigma_3}{\varepsilon T^{\varepsilon \alpha}} \delta^{1-\varepsilon} \exp \left[-\frac{\sigma_1 - \sigma_3}{T^{\varepsilon \alpha}} \delta^{\varepsilon} \right].$$

Furthermore, one needs the following formulae (see [31, p. 201])

$$\int_{0}^{\tau} \frac{\eta}{\sqrt{\tau^2 - \eta^2}} \cos\left(\rho\sqrt{\tau^2 - \eta^2}\right) J_0\left(\sigma\eta\right) d\eta = \frac{1}{k} \sin\left(k\tau\right), \tag{2.15}$$

$$\int_{0}^{\tau} \frac{J_1(\sigma\eta)}{\sqrt{\tau^2 - \eta^2}} \cos\left(\rho\sqrt{\tau^2 - \eta^2}\right) d\eta = \frac{1}{\sigma\tau} \cos(\rho\tau) - \frac{1}{\sigma\tau} \cos(k\tau), \quad (2.16)$$

where $k = \sqrt{\sigma^2 + \rho^2}$.

Under formula (2.16), one can obtain

$$\int_{0}^{\tau} \sin\left(\rho\sqrt{\tau^{2}-\eta^{2}}\right) J_{1}\left(\sigma\eta\right) d\eta = \rho \int_{0}^{\tau} \left(\xi \int_{0}^{\xi} \frac{\cos\left(\rho\sqrt{\xi^{2}-\eta^{2}}\right)}{\sqrt{\xi^{2}-\eta^{2}}} J_{1}\left(\sigma\eta\right) d\eta\right) d\xi = 0$$

$$= \frac{\rho}{\sigma} \int_{0}^{\tau} \left[\cos(\rho \xi) - \cos(k \xi) \right] d\xi = \frac{1}{\sigma} \sin(\rho \tau) - \frac{\rho}{\sigma k} \sin(k \tau). \tag{2.17}$$

The Stankovic's transformation integral (see [32, p. 84])

$$\int_{0}^{\infty} \exp(\lambda \tau) t^{\nu - 1} \phi\left(-\mu, \nu; -\tau t^{-\mu}\right) d\tau = t^{\mu + \nu - 1} E_{\mu}\left(-\lambda t^{\mu}; \mu + \nu\right), \qquad (2.18)$$

takes place for any $\lambda \in \mathbb{C}$, $\mu \in (0,1)$, $\nu \in \mathbb{R}$, where

$$E_{\rho}(z;\mu) = \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\mu + \rho k)}$$

is the Mittag-Leffler-type function [33, c. 117].

3 Matrix integral formalism of the DO Cauchy problem

Let us convert the DO Cauchy problem in the 'differential form' (1.1)–(1.2) to the matrix Fredholm–Volterra-type integral equation of the second kind. The latter takes a special sense to build up the Resolvent-function solution of the DO Cauchy problem in terms of the Liouville–Neumann-type series, which in turn is very important to computer-aided modeling and recovering the crystal-lattice displacement field function $f(\mathbf{R})$ from the X-ray DO microtomography data.

The system of differential TT-type equations (1.1) may be rewritten into the form

$$\begin{pmatrix} O_{-}^{\alpha} - i\gamma & 0\\ 0 & O_{+}^{\alpha} - i\gamma \end{pmatrix} \mathbf{E} = i\sigma \mathbf{K} \mathbf{E}, \tag{3.1}$$

where

$$O_{+}^{\alpha} = \partial_{0t}^{\alpha} + \frac{\partial}{\partial x}, \quad O_{-}^{\alpha} = \partial_{0t}^{\alpha} - \frac{\partial}{\partial x},$$

$$\mathbf{E} \equiv \mathbf{E}(x,t) = \begin{pmatrix} E_{0}(x,t) \\ E_{h}(x,t) \end{pmatrix}, \quad \mathbf{K} \equiv \mathbf{K}(x,t) = \begin{pmatrix} 0 & e^{if(x,t)} \\ e^{-if(x,t)} & 0 \end{pmatrix}.$$

Acting onto both sides of (3.1) by the operator $diag(O_+^{\alpha} - i\gamma, O_-^{\alpha} - i\gamma)$, one obtains

$$\begin{pmatrix} O_{+}^{\alpha} - i\gamma & 0 \\ 0 & O_{-}^{\alpha} - i\gamma \end{pmatrix} \begin{pmatrix} O_{-}^{\alpha} - i\gamma & 0 \\ 0 & O_{+}^{\alpha} - i\gamma \end{pmatrix} \mathbf{E} =$$

$$= i\sigma \begin{pmatrix} O_{+}^{\alpha} - i\gamma & 0 \\ 0 & O_{-}^{\alpha} - i\gamma \end{pmatrix} \mathbf{KE}$$
(3.2)

and after some straightforward routine calculations, one finds out

$$\begin{pmatrix} O_{+}^{\alpha} - i\gamma & 0 \\ 0 & O_{-}^{\alpha} - i\gamma \end{pmatrix} \mathbf{K} \mathbf{E} = \begin{bmatrix} \partial_{0t}^{\alpha} + \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \frac{\partial}{\partial x} - i\gamma \end{bmatrix} (\mathbf{K} \mathbf{E}) = \partial_{0t}^{\alpha} (\mathbf{K} \mathbf{E}) + \\ + \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{bmatrix} \left(\frac{\partial}{\partial x} \mathbf{K} \right) \mathbf{E} + \mathbf{K} \left(\frac{\partial}{\partial x} \mathbf{E} \right) \end{bmatrix} + \mathbf{K} (\partial_{0t}^{\alpha} \mathbf{E}) - \mathbf{K} (\partial_{0t}^{\alpha} \mathbf{E}) - i\gamma \mathbf{K} \mathbf{E} = \\ = \mathbf{K} \begin{pmatrix} O_{-}^{\alpha} - i\gamma & 0 \\ 0 & O_{+}^{\alpha} - i\gamma \end{pmatrix} \mathbf{E} + \partial_{0t}^{\alpha} (\mathbf{K} \mathbf{E}) + \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \left(\frac{\partial}{\partial x} \mathbf{K} \right) \mathbf{E} - \mathbf{K} (\partial_{0t}^{\alpha} \mathbf{E}) = \\ = \mathbf{K} \begin{pmatrix} O_{-}^{\alpha} - i\gamma & 0 \\ 0 & O_{+}^{\alpha} - i\gamma \end{pmatrix} \mathbf{E} + (\partial_{0t}^{\alpha} + if_{x}') (\mathbf{K} \mathbf{E}) - \mathbf{K} (\partial_{0t}^{\alpha} \mathbf{E}). \tag{3.3}$$

The column vector $\mathbf{E} = \mathbf{E}(x,t)$ is nothing else the solution of Eq. (3.1) and \mathbf{K}^2 is equal to Unit matrix, from Eq. (3.3) it directly follows

$$i\sigma \begin{pmatrix} O_{+}^{\alpha} - i\gamma & 0 \\ 0 & O_{-}^{\alpha} - i\gamma \end{pmatrix} \mathbf{K}\mathbf{E} = -\sigma^{2}\mathbf{E} + i\sigma \left(\partial_{0t}^{\alpha} + if_{x}'\right) \left(\mathbf{K}\mathbf{E}\right) - i\sigma\mathbf{K} \left(\partial_{0t}^{\alpha}\mathbf{E}\right).$$

The last allows us to rewrite down Eq. (3.2) into the form

$$\begin{pmatrix}
(O_{+}^{\alpha} - i\gamma)(O_{-}^{\alpha} - i\gamma) + \sigma^{2} & 0 \\
0 & (O_{-}^{\alpha} - i\gamma)(O_{+}^{\alpha} - i\gamma) + \sigma^{2}
\end{pmatrix} \mathbf{E} = i\sigma \left(\partial_{0t}^{\alpha} + if_{x}'\right) (\mathbf{K}\mathbf{E}) - i\sigma \mathbf{K} \left(\partial_{0t}^{\alpha} \mathbf{E}\right). \tag{3.4}$$

Keeping in mind Eq.(2.5), and applying the double Fourier-Laplace transform to Eq. (3.4), one can obtain

$$\left[(O_{+}^{\alpha} - i\gamma)(O_{-}^{\alpha} - i\gamma)E_{0}(x, t) \right]_{k,p} = (p^{\alpha} - i\gamma + ik)\left[(O_{-}^{\alpha} - i\gamma)E_{0}(x, t) \right]_{k,p} - p^{\alpha-1} \left\{ \left[(O_{-}^{\alpha} - i\gamma)E_{0}(x, t) \right]_{t=0} \right\}_{k} = (p^{\alpha} - i\gamma + ik)(p^{\alpha} - i\gamma - ik)\left[E_{0}(x, t) \right]_{k,p} - p^{\alpha-1} \left\{ (p^{\alpha} - i\gamma + ik)\left[E_{0}(x, 0) \right]_{k} + i\sigma \left[e^{if(x, 0)}E_{h}(x, 0) \right]_{k} \right\},$$
(3.5)

and

$$\left[(O_{-}^{\alpha} - i\gamma)(O_{+}^{\alpha} - i\gamma)E_{h}(x, t) \right]_{k,p} = (p^{\alpha} - i\gamma - ik)(p^{\alpha} - i\gamma + ik)[E_{h}(x, t)]_{k,p} - p^{\alpha-1} \left\{ (p^{\alpha} - i\gamma - ik)[E_{h}(x, 0)]_{k} + i\sigma[e^{-if(x, 0)}E_{0}(x, 0)]_{k} \right\}.$$
(3.6)

Here, one has used the relationships (cf. (3.1))

$$[(O_{-}^{\alpha} - i\gamma)E_{0}(x,t)]_{t=0} = i\sigma e^{if(x,0)}E_{h}(x,0),$$

$$[(O_+^{\alpha} - i\gamma)E_h(x,t)]_{t=0} = i\sigma e^{-if(x,0)}E_0(x,0).$$

Thus from Eqs.(3.4) – (3.6) it directly follows

$$\begin{pmatrix} E_{0}(x,t) \\ E_{h}(x,t) \end{pmatrix}_{k,p} = \frac{p^{\alpha-1}}{(p^{\alpha} - i\gamma)^{2} + k^{2} + \sigma^{2}} \times$$

$$\times \left\{ \begin{pmatrix} p^{\alpha} - i\gamma + ik & 0 \\ 0 & p^{\alpha} - i\gamma - ik \end{pmatrix} \begin{pmatrix} E_{0}(x,0) \\ E_{h}(x,0) \end{pmatrix}_{k} + i\sigma \begin{pmatrix} e^{if(x,0)} E_{h}(x,0) \\ e^{-if(x,0)} E_{0}(x,0) \end{pmatrix}_{k} \right\} +$$

$$+ \frac{1}{(p^{\alpha} - i\gamma)^{2} + k^{2} + \sigma^{2}} \left\{ \begin{pmatrix} \partial_{0t}^{\alpha} + if'_{x} & 0 \\ 0 & \partial_{0t}^{\alpha} + if'_{x} \end{pmatrix} \begin{pmatrix} i\sigma e^{if} E_{h}(x,t) \\ i\sigma e^{-if} E_{0}(x,t) \end{pmatrix} -$$

$$- \begin{pmatrix} 0 & i\sigma e^{if} \\ i\sigma e^{-if} & 0 \end{pmatrix} \begin{pmatrix} \partial_{0t}^{\alpha} E_0(x,t) \\ \partial_{0t}^{\alpha} E_h(x,t) \end{pmatrix} \right\}_{k,n}.$$
 (3.7)

Applying the Efros's theorem for operational calculus [34, p. 512], the equalities (2.6)–(2.10), and using the inverse double Fourier–Laplace transform, the following elations take place

$$\left(\frac{1}{(p^{\alpha}-i\gamma)^{2}+k^{2}+\sigma^{2}}\right)_{x,t} = \frac{1}{i(2\pi)^{2}} \int_{-i\infty}^{i\infty} dp \int_{-\infty}^{\infty} \frac{e^{pt+ikx}}{(p^{\alpha}-i\gamma)^{2}+k^{2}+\sigma^{2}} dk =$$

$$= \frac{1}{2} \int_{|x|}^{\infty} e^{i\gamma\tau} J_{0} \left(\sigma\sqrt{\tau^{2}-x^{2}}\right) \frac{1}{t} \phi \left(-\alpha,0;-\frac{\tau}{t^{\alpha}}\right) d\tau = G_{\alpha,\gamma}(x,t),$$

$$\left(\frac{p^{\alpha-1}}{(p^{\alpha}-i\gamma)^{2}+k^{2}+\sigma^{2}}\right)_{x,t} = \frac{1}{i(2\pi)^{2}} \int_{-i\infty}^{i\infty} dp \int_{-\infty}^{\infty} \frac{p^{\alpha-1}e^{pt+ikx}}{(p^{\alpha}-i\gamma)^{2}+k^{2}+\sigma^{2}} dk =$$

$$= D_{0t}^{\alpha-1} G_{\alpha,\gamma}(x,t) = \frac{1}{2} \int_{-\infty}^{\infty} e^{i\gamma\tau} J_{0} \left(\sigma\sqrt{\tau^{2}-x^{2}}\right) t^{-\alpha} \phi \left(-\alpha,1-\alpha;-\frac{\tau}{t^{\alpha}}\right) d\tau.$$

As a result, from (3.7) one obtains the DO integral matrix equation

$$\mathbf{E}(x,t) = (\mathbf{A}^{\alpha,\gamma}\mathbf{E}(x,t))(x,t) + (\mathbf{B}^{\alpha,\gamma}\mathbf{E}(x,0))(x,t), \tag{3.8}$$

where the following notations are introduced

$$(\mathbf{A}^{\alpha,\gamma}\mathbf{E}(x,t))(x,t) = -i\sigma \int_{0}^{t} dv \int_{-\infty}^{\infty} G_{\alpha,\gamma}(x-u,t-v) \cdot \left\{ D_{0v}^{\alpha-1} \frac{\partial}{\partial v} [\mathbf{K}(u,v)\mathbf{E}(u,v)] + \left(\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array} \right) \left[\frac{\partial}{\partial u} \mathbf{K}(u,v) \right] \mathbf{E}(u,v) - \mathbf{K}(u,v) D_{0v}^{\alpha-1} \frac{\partial}{\partial v} \mathbf{E}(u,v) \right\} du,$$

$$(\mathbf{B}^{\alpha,\gamma}\mathbf{E}(x,0))(x,t) = \int_{0}^{t} dv \int_{-\infty}^{\infty} D_{tv}^{\alpha-1} G_{\alpha,\gamma}(x-u,t-v) \times$$

$$\times \left(\begin{array}{cc} O_{+}^{\alpha} - i\gamma & i\sigma e^{if(u,0)} \\ i\sigma e^{-if(u,0)} & O_{-}^{\alpha} - i\gamma \end{array} \right) \mathbf{E}(u,0) \delta(v) du,$$

where $\delta(v)$ is the Dirac delta-function.

Using the formula for integration by parts and its fractional analogue (2.4), the estimates (2.11) and (2.12), definition of the fractional Riemann–Liouville derivative, one finds out

$$(\mathbf{A}^{\alpha,\gamma}\mathbf{E}(x,t))(x,t) = -i\sigma \int_{0}^{t} dv \int_{-\infty}^{\infty} \mathbf{K}_{1}(x,t;u,v)\mathbf{E}(u,v)du - i\sigma \int_{-\infty}^{\infty} D_{0t}^{\alpha-1}G_{\alpha,\gamma}(x-u,t)\mathbf{K}(u,0)\mathbf{E}(u,0)du,$$
(3.9)

where

$$\mathbf{K}_{1}(x,t;u,v) = D_{vt}^{\alpha}G_{\alpha,\gamma}(x-u,t-v)\cdot\mathbf{K}(u,v) +$$

$$+if'_{u}(u,v)\cdot G_{\alpha,\gamma}(x-u,t-v)\mathbf{K}(u,v) + D_{vt}^{\alpha}\left[G_{\alpha,\gamma}(x-u,t-v)\mathbf{K}(u,v)\right].$$
In view of (2.3) the $(\mathbf{B}^{\alpha,\gamma}\mathbf{E}(x,0))(x,t)$ may be rewritten as

 $\overset{\infty}{f}$

$$(\mathbf{B}^{\alpha,\gamma}\mathbf{E}(x,0))(x,t) = -\int_{-\infty}^{\infty} D_{0t}^{2\alpha-1}G_{\alpha,\gamma}(x-u,t)\mathbf{E}(u,0)du -$$

$$-i\gamma\int_{-\infty}^{\infty} D_{0t}^{\alpha-1}G_{\alpha,\gamma}(x-u,t)\mathbf{E}(u,0)du +$$

$$+\left(\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right)\int_{-\infty}^{\infty} D_{0t}^{\alpha-1}G_{\alpha,\gamma}(x-u,t)\mathbf{E}'(u,0)du +$$

$$+i\sigma\int_{-\infty}^{\infty} D_{0t}^{\alpha-1}G_{\alpha,\gamma}(x-u,t)\mathbf{K}(u,0)\mathbf{E}(u,0)du. \tag{3.10}$$

Taking into account Eqs. (3.9), (3.10), the integral matrix equation (3.8) may be to reduce

$$\mathbf{E}(x,t) + i\sigma \int_{0}^{t} dv \int_{-\infty}^{\infty} \mathbf{K}_{1}(x,t;u,v) \mathbf{E}(u,v) du = \mathbf{F}(x,t), \qquad (3.11)$$

where the following notations are introduced

$$\mathbf{F}(x,t) = (\mathbf{B}^{\alpha,\gamma}\mathbf{E}(x,0))(x,t) - i\sigma \int_{-\infty}^{\infty} D_{0t}^{\alpha-1}G_{\alpha,\gamma}(x-u,t)\mathbf{K}(u,0)\mathbf{E}(u,0)du.$$

4 Matrix Resolvent-function solution of the DO Cauchy problem

Let A(x) be the matrix with entries $a_{ij}(x)$. By notation $|A(x)|_*$ one denotes a scalar function taking the maximum absolute value of entries $a_{ij}(x)$ of the matrix A(x) for each x; i.e., $|A(x)|_* = \max_{i,j} |a_{ij}(x)|$.

Let f(x,t) and $f_t(x,t)$ be the continuous, bounded functions.

From the Eqs. (2.11) - (2.14) one obtains following estimates for the matrix kernel function \mathbf{K}_1

$$|\mathbf{K}_1(x,t;\xi,\eta)|_* \le C|x-\xi|^{-\theta}(t-\eta)^{\beta-1},$$
 (4.1)

$$|\mathbf{K}_1(x,t;\xi,\eta)|_* \le C(t-\eta)^{\beta-1} \exp\left[-\frac{\sigma_0|x-\xi|^{\varepsilon}}{(t-\eta)^{\alpha\varepsilon}}\right],$$
 (4.2)

where $\beta = \alpha \theta$, $\theta \in (0, 1)$, $\sigma_0 < (1 - \alpha)\alpha^{\alpha \varepsilon}$, $\varepsilon = \frac{1}{1 - \alpha}$.

Further, let us find out the corresponding estimates for iterative kernels functions, namely:

$$\mathbf{K}_{n}(x,t;\xi,\eta) = \int_{\eta}^{t} dv \int_{-\infty}^{\infty} \mathbf{K}_{n-1}(x,t;u,v) \mathbf{K}_{1}(u,v;\xi,\eta) du.$$

From estimates (4.1) and (4.2) it follows

$$|\mathbf{K}_2(x,t;\xi,\eta)|_* \le$$

$$\leq \int_{\eta}^{t} dv \left(\int_{x-\delta_{1}}^{x+\delta_{2}} + \int_{-\infty}^{x-\delta_{1}} + \int_{x+\delta_{2}}^{\infty} \right) |\mathbf{K}_{1}(x,t;u,v)\mathbf{K}_{1}(u,v;\xi,\eta)|_{*} du \leq
\leq CC_{2,\beta,\theta}(t-\eta)^{2\beta-1} \left(|x-\xi|^{-\theta} + \exp\left[-\frac{\sigma_{0}|x-\xi|^{\varepsilon}}{(t-\eta)^{\alpha\varepsilon}} \right] \right), \tag{4.3}$$

where

$$C_{2,\beta,\theta} = \frac{\Gamma^2(\beta)C_{1-\theta}}{\Gamma(2\beta)}.$$

Accordingly, from (4.3), keeping in mind $0 < e^{-s} < 1$, for s > 0, one obtains

$$|\mathbf{K}_{2}(x,t;\xi,\eta)|_{*} \le CC_{2,\beta,\theta}|x-\xi|^{-\theta}(t-\eta)^{2\beta-1},$$
 (4.4)

$$|\mathbf{K}_{2}(x,t;\xi,\eta)|_{*} \leq CC_{2,\beta,\theta}(t-\eta)^{2\beta-1} \exp\left[-\frac{\sigma_{0}|x-\xi|^{\varepsilon}}{(t-\eta)^{\alpha\varepsilon}}\right]. \tag{4.5}$$

Under the above, the following estimates take place

$$|\mathbf{K}_n(x,t;\xi,\eta)|_* \le CC_{n,\beta,\theta}\Gamma(n\beta)|x-\xi|^{-\theta}(t-\eta)^{n\beta-1},\tag{4.6}$$

$$|\mathbf{K}_n(x,t;\xi,\eta)|_* \le CC_{n,\beta,\theta}(t-\eta)^{n\beta-1} \exp\left[-\frac{\sigma_0|x-\xi|^{\varepsilon}}{(t-\eta)^{\alpha\varepsilon}}\right].$$
 (4.7)

where

$$C_{n,\beta,\theta} = \frac{\Gamma^n(\beta)C_{1-\theta}^{n-1}}{\Gamma(n\beta)}.$$

The matrix Resolvent-function solution of the Eq. (3.11) takes the form

$$\mathbf{R}(x,t;\xi,\eta) = \sum_{n=1}^{\infty} \mathbf{K}_n(x,t;\xi,\eta). \tag{4.8}$$

From estimates (4.1)–(4.7), it follows the Liouville–Neumann-type series (4.8) converges and can be estimated as a value of

$$|\mathbf{R}(x,t;\xi,\eta)|_{*} \leq C\Gamma(\beta)(t-\eta)^{\beta-1}E_{\beta}\left[C_{1-\theta}\Gamma(\beta)(t-\eta)^{\beta};\beta\right] \times \left(|x-\xi|^{-\theta} + \exp\left[-\frac{\sigma_{0}|x-\xi|^{\varepsilon}}{(t-\eta)^{\alpha\varepsilon}}\right]\right).$$

On the other hand, since the Mittag-Leffler-type function $E_{\beta}(s;\beta)$ is bounded on the interval [0,T], the estimates of (4.1)–(4.2) are proved and valid also for the matrix Resolvent-function solution as a whole.

So, one can state the unique solution of the matrix integral Eq. (3.11) takes the form

$$\mathbf{E}(x,t) = \mathbf{F}(x,t) + \int_{0}^{t} \int_{-\infty}^{+\infty} \mathbf{R}(x,t;\xi,\eta) \mathbf{F}(\xi,\eta) d\xi d\eta.$$

5 The DO Cauchy problem. The FOP $\alpha = 1$

In the case when FOP $\alpha = 1$, the basic system (3.1) reduces to

$$\begin{pmatrix} \frac{\partial}{\partial t} - \frac{\partial}{\partial x} & 0\\ 0 & \frac{\partial}{\partial t} + \frac{\partial}{\partial x} \end{pmatrix} \begin{pmatrix} E_0(x,t)\\ E_h(x,t) \end{pmatrix} =$$

$$= i \begin{pmatrix} \gamma & \sigma \exp[if(x,t)]\\ \sigma \exp[-if(x,t)] & \gamma \end{pmatrix} \begin{pmatrix} E_0(x,t)\\ E_h(x,t) \end{pmatrix}.$$

As was shown earlier [26], one can find out that for $\alpha \to 1$ integral equation (3.8) can be written down into the form

$$\mathbf{E}(x,t) = (\mathbf{A}^{1,\gamma}\mathbf{E}(x,t))(x,t) + (\mathbf{B}^{1,\gamma}\mathbf{E}(x,0))(x,t),$$

where

$$(\mathbf{A}^{1,\gamma}\mathbf{E}(x,t))(x,t) = i\sigma \int_{0}^{t} dv \int_{x-t+v}^{x+t-v} G_{1,\gamma}(x-u,t-v) \times$$

$$\times \begin{pmatrix} 0 & O_{+} e^{if(u,v)} \\ O_{-} e^{-if(u,v)} & 0 \end{pmatrix} \mathbf{E}(u,v) du,$$

$$(\mathbf{B}^{1,\gamma}\mathbf{E}(x,0))(x,t) = \int_{0}^{t} dv \int_{x-t+v}^{x+t-v} G_{1,\gamma}(x-u,t-v) \times$$

$$\times \begin{pmatrix} O_{+} - i\gamma & i\sigma e^{if(u,0)} \\ i\sigma e^{-if(u,0)} & O_{-} - i\gamma \end{pmatrix} \delta(v) \mathbf{E}(u,0) du,$$

and respectively,

$$G_{1,\gamma}(x,t) = \frac{1}{2}e^{i\gamma t}J_0\left(\sigma\sqrt{t^2 - x^2}\right)\Theta[t - |x|].$$

6 The DO Cauchy problem. The crystal-lattice displacement field function $f(\mathbf{R}) = ax + b$

Here we will build up a solution of the basic fractional DO Cauchy problem when the crystal-lattice displacement field function $f(\mathbf{R}) = ax + b$.

After trivial exponentional substitutions for the wave amplitudes $E_0(x,t)$, $E_h(x,t)$, the system (1.1) can written down as (for simplicity, further, the same notations for the wave amplitudes $E_0(x,t)$, $E_h(x,t)$, are to be saved)

$$\left(\partial_{0t}^{\alpha} - \frac{\partial}{\partial x}\right) E_0(x,t) = i\gamma E_0(x,t) + i\sigma e^{iax} E_h(x,t),
\left(\partial_{0t}^{\alpha} + \frac{\partial}{\partial x}\right) E_h(x,t) = i\sigma e^{-iax} E_0(x,t) + i\gamma E_h(x,t).$$
(6.1)

Substituting the functions $E_0(x,t)$, $E_h(x,t)$ as

$$E_0(x,t) = \exp\left(i\frac{ax}{2}\right)\mathcal{E}_0(x,t), \quad E_h(x,t) = \exp\left(-i\frac{ax}{2}\right)\mathcal{E}_h(x,t),$$

one obtains

$$\left(\partial_{0t}^{\alpha} - \frac{\partial}{\partial x}\right) \mathcal{E}_{0}(x,t) = i\left(\gamma + \frac{a}{2}\right) \mathcal{E}_{0}(x,t) + i\sigma \mathcal{E}_{h}(x,t),
\left(\partial_{0t}^{\alpha} + \frac{\partial}{\partial x}\right) \mathcal{E}_{h}(x,t) = i\sigma \mathcal{E}_{0}(x,t) + i\left(\gamma + \frac{a}{2}\right) \mathcal{E}_{h}(x,t),$$
(6.2)

and the initial condition

$$\begin{pmatrix} \mathcal{E}_0(x,0) \\ \mathcal{E}_h(x,0) \end{pmatrix} = \begin{pmatrix} \psi_0(x) \\ \psi_h(x) \end{pmatrix} = \begin{pmatrix} e^{-iax/2}\varphi_0(x) \\ e^{iax/2}\varphi_h(x) \end{pmatrix}, \quad -\infty < x < \infty. \quad (6.3)$$

Then, let us introduce the notations

$$\Gamma(x,t) = \frac{1}{2} \int_{|x|}^{\infty} g(t,\tau) \mathbf{Q}(x,\tau) d\tau + \Gamma_0(x,t),$$
 (6.4)

$$\mathbf{Q}(x,\tau) = \begin{bmatrix} -\sigma \frac{\tau - x}{\sqrt{\tau^2 - x^2}} J_1(\sigma \sqrt{\tau^2 - x^2}) & i\sigma J_0(\sigma \sqrt{\tau^2 - x^2}) \\ i\sigma J_0(\sigma \sqrt{\tau^2 - x^2}) & -\sigma \frac{\tau + x}{\sqrt{\tau^2 - x^2}} J_1(\sigma \sqrt{\tau^2 - x^2}) \end{bmatrix}, (6.5)$$

$$\Gamma_0(x,t) = g(t,|x|) \begin{bmatrix} \Theta(-x) & 0 \\ 0 & \Theta(x) \end{bmatrix}, \tag{6.6}$$

$$g(t,\tau) = \frac{e^{i(\gamma + a/2)\tau}}{t} \phi\left(-\alpha, 0; -\tau t^{-\alpha}\right),\,$$

 $\Theta(x)$ is the Heaviside function, $J_m(z)$ is the m-order Bessel function of the argument z.

According to [25], the basic matrix solution of the DO Cauchy problem (6.2) - (6.3) has the form

$$\mathcal{E}(x,t) = \int_{-\infty}^{\infty} D_{0t}^{\alpha-1} \Gamma(x-\xi,t) \psi(\xi) d\xi, \tag{6.7}$$

in the class of function

$$\mathcal{E}(x,t) \in C(\overline{\Omega}), \quad \partial_{0t}^{\alpha} \mathcal{E}(x,t), \frac{\partial}{\partial x} \mathcal{E}(x,t) \in C(\Omega),$$

where $\psi(x) = [\psi_0(x), \psi_h(x)]^{tr} \in C(-\infty, \infty)$ is the function, which satisfies the Hölder condition, and the following relation as $|x| \to \infty$

$$\psi(x) = O(\exp(\rho|x|^{\varepsilon})) \quad \varepsilon = \frac{1}{1-\alpha}, \quad \rho < (1-\alpha)(\alpha T^{-1})^{\frac{\alpha}{1-\alpha}}.$$

Underline the solution to the Cauchy problem (6.2) - (6.3) is unique in the class of functions, which satisfy the condition

$$\mathcal{E}(x,t) = O(\exp(k|x|^{\varepsilon})), \quad \text{as} \quad |x| \to \infty,$$

for some k > 0.

From Eqs. (6.3) - (6.7) it follows the solution of the Cauchy problem (6.1), (1.2) can be cast into the form

$$\mathbf{E}(x,t) = \int_{-\infty}^{\infty} D_{0t}^{\alpha-1} \mathbf{G}(x,\xi,t) \varphi(\xi) d\xi, \tag{6.8}$$

where the following notations are introduced

$$\mathbf{G}(x,\xi,t) = \frac{1}{2} \int_{|x-\xi|}^{\infty} g(t,\tau)\mathbf{S}(x,\xi,\tau)d\tau + \mathbf{G}_0(x,\xi,t), \tag{6.9}$$

$$\mathbf{S}(x,\xi,\tau) = \begin{bmatrix} -\sigma e^{iaX_1/2} \frac{\tau - X_1}{\sqrt{\tau^2 - X_1^2}} h_1(\tau, X_1) & i\sigma e^{iaX_2/2} h_0(\tau, X_1) \\ i\sigma e^{-iaX_2/2} h_0(\tau, X_1) & -\sigma e^{-iaX_1/2} \frac{\tau + X_1}{\sqrt{\tau^2 - X_1^2}} h_1(\tau, X_1) \end{bmatrix},$$

$$h_0(\tau, X_1) = J_0(\sigma \sqrt{\tau^2 - X_1^2}), \quad h_1(\tau, X_1) = J_1(\sigma \sqrt{\tau^2 - X_1^2})$$

$$\mathbf{G}_{0}(x,\xi,t) = \begin{bmatrix} e^{-i\gamma X_{1}} \Theta(-X_{1}) & 0\\ 0 & e^{i\gamma X_{1}} \Theta(X_{1}) \end{bmatrix} \frac{1}{t} \phi\left(-\alpha,0; -|X_{1}|t^{-\alpha}\right), \quad (6.10)$$

$$X_{1} = x - \xi, \quad X_{2} = x + \xi.$$

7 Case of the function
$$f(\mathbf{R}) = ax + b$$
 and $E_0(x, 0) = 1$, $E_h(x, 0) = 0$

Let us consider the case when the initial conditions for the wavefield amplitudes $E_0(x,0)$ and $E_h(x,0)$ are constant and equal to

$$E_0(x,0) = \varphi_0(x) \equiv 1, \quad E_h(x,0) = \varphi_h(x) \equiv 0.$$
 (7.1)

Then, the formulae (6.8)–(6.10) can be simplified and expressed in terms of the Mittag–Leffler-type functions.

Keeping in mind Eq. (7.1), Eq. (6.8) for $\mathbf{E}(x,t)$ can be written down as

$$\mathbf{E}(x,t) = \int_{-\infty}^{\infty} D_{0t}^{\alpha-1} \mathbf{G}(x,\xi,t) \begin{pmatrix} 1 \\ 0 \end{pmatrix} d\xi =$$

$$= \frac{1}{2} \int_{-\infty}^{\infty} d\xi \int_{|x-\xi|}^{\infty} D_{0t}^{\alpha-1} g(t,\tau) \mathbf{S}(x,\xi,\tau) \begin{pmatrix} 1\\0 \end{pmatrix} d\tau + \int_{-\infty}^{\infty} D_{0t}^{\alpha-1} \mathbf{G}_{0}(x,\xi,t) \begin{pmatrix} 1\\0 \end{pmatrix} d\xi \equiv \mathbf{I}_{1}(x,t) + \mathbf{I}_{2}(x,t).$$
 (7.2)

By changing the integration order, let us evaluate the integral $I_1(x,t)$

$$\mathbf{I}_{1}(x,t) = \frac{1}{2} \int_{0}^{\infty} D_{0t}^{\alpha-1} g(t,\tau) d\tau \int_{x-\tau}^{x+\tau} \mathbf{S}(x,\xi,\tau) \begin{pmatrix} 1\\0 \end{pmatrix} d\xi =$$

$$= \frac{1}{2} \int_{0}^{\infty} D_{0t}^{\alpha-1} g(t,\tau) d\tau \int_{0}^{\tau} \mathbf{S}(x,x-\eta,\tau) \begin{pmatrix} 1\\0 \end{pmatrix} d\eta. \tag{7.3}$$

From the Eqs. (2.15)–(2.17), it directly follows up

$$\int_{-\tau}^{\tau} e^{\pm ia\eta/2} J_0(\sigma\sqrt{\tau^2 - \eta^2}) d\eta = \frac{2}{k} \sin(k\tau), \tag{7.4}$$

$$\int_{-\tau}^{\tau} e^{\pm ia\eta/2} \frac{\tau \mp \eta}{\sqrt{\tau^2 - \eta^2}} J_1(\sigma \sqrt{\tau^2 - \eta^2}) d\eta =$$

$$= -\frac{1}{\sigma} \left[2\cos(k\tau) - i\frac{a}{k}\sin(k\tau) - 2e^{-ia\tau/2} \right], \tag{7.5}$$

where $k = \sqrt{a^2/4 + \sigma^2}$.

Accordingly, from Eqs. (7.4), (7.5) one obtains

$$\int_{-\tau}^{\tau} S_{11,22}(x, x - \eta, \tau) d\eta = -\sigma \int_{-\tau}^{\tau} e^{\pm ia\eta/2} \frac{\tau \mp \eta}{\sqrt{\tau^2 - \eta^2}} J_1(\sigma \sqrt{\tau^2 - \eta^2}) d\eta =
= 2\cos(k\tau) - i\frac{a}{k}\sin(k\tau) - 2e^{-ia\tau/2},$$
(7.6)
$$\int_{-\tau}^{\tau} S_{12,21}(x, x - \eta, \tau) d\eta = i\sigma e^{\pm iax} \int_{-\tau}^{\tau} e^{\mp ia\eta/2} J_0(\sigma \sqrt{\tau^2 - \eta^2}) d\eta =
= i\sigma \frac{2}{k} e^{\pm iax} \sin(k\tau),$$
(7.7)

where $S_{ij}(x, x - \eta, \tau)$ (i, j = 1, 2) are the elements of matrix $\mathbf{S}(x, x - \eta, \tau)$. Following equalities

$$e^{ia_1\tau} \left[2\cos(k\tau) - i\frac{a}{k}\sin(k\tau) \right] =$$

$$= \left(1 - \frac{a}{2k} \right) e^{i(a_1+k)\tau} + \left(1 + \frac{a}{2k} \right) e^{i(a_1-k)\tau}, \tag{7.8}$$

$$-2e^{ia_1\tau}e^{ia\tau/2} = -2e^{i\gamma\tau},\tag{7.9}$$

$$e^{ia_1\tau}\sin(k\tau) = \frac{i}{2}e^{i(a_1-k)\tau} - \frac{i}{2}e^{i(a_1+k)\tau},$$
 (7.10)

hold, where

$$a_1 = \gamma + \frac{a}{2}, \quad k = \sqrt{a^2/4 + \sigma^2}.$$

Next, exploiting calculations (7.6)–(7.10), from (7.3) one obtains

$$\mathbf{I}_{1}(x,t) = \int_{0}^{\infty} t^{-\alpha} \phi\left(-\alpha, 1 - \alpha; -\tau t^{-\alpha}\right) \mathbf{N}(x,\tau) \begin{pmatrix} 1\\ 0 \end{pmatrix} d\tau, \tag{7.11}$$

where

$$\mathbf{N}(x,\tau) = \mathbf{N}_1(x)e^{i(a_1+k)\tau} + \mathbf{N}_2(x)e^{i(a_1-k)\tau} - \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}e^{i\gamma\tau},$$
$$\mathbf{N}_{1,2}(x) = \frac{1}{2} \begin{pmatrix} 1 \mp \frac{a}{2k} & \pm \frac{\sigma}{k}e^{iax} \\ \pm \frac{\sigma}{k}e^{-iax} & 1 \mp \frac{a}{2k} \end{pmatrix}.$$

From (6.10) one finds out

$$\mathbf{I}_{2}(x,t) = \int_{0}^{\infty} t^{-\alpha} \phi\left(-\alpha, 1 - \alpha; -\eta t^{-\alpha}\right) e^{i\gamma\eta} \begin{pmatrix} 1\\ 0 \end{pmatrix} d\eta. \tag{7.12}$$

Applying formula (2.18) to equalities (7.11) and (7.12), the total solution (7.2) can be cast into the form

$$\mathbf{E}(x,t) = \frac{1}{2} \begin{pmatrix} 1 - \frac{a}{2k} & 1 + \frac{a}{2k} \\ \frac{\sigma}{k} e^{-iax} & -\frac{\sigma}{k} e^{-iax} \end{pmatrix} \begin{pmatrix} E_{\alpha,1} \left(i(a_1 + k)t^{\alpha} \right) \\ E_{\alpha,1} \left(i(a_1 - k)t^{\alpha} \right) \end{pmatrix}. \tag{7.13}$$

In the case when the FOP $\alpha \to 1$, the total solution (7.13) is reduced to

$$\mathbf{E}(x,t) = \frac{1}{2} \begin{pmatrix} 1 - \frac{a}{2k} & 1 + \frac{a}{2k} \\ \frac{\sigma}{k} e^{-iax} & -\frac{\sigma}{k} e^{-iax} \end{pmatrix} \begin{pmatrix} e^{i(a_1+k)t} \\ e^{i(a_1-k)t} \end{pmatrix}. \tag{7.14}$$

Finally, putting on the parameters $\gamma = 0$ and a = 0, the solution (7.14) goes into the pendulum solutions of the canonic TT equations regarding the Bloch wavefield amplitudes in the perfect crystal (cf. [26])

$$E_0 = \cos \sigma t$$
, $E_h = i \sin \sigma t$.

8 Conclusion

In this paper, a goal of our study is to elaborate the novel mathematics framework for processing the DO data based on a concept of the matrix integral equation to solve the inverse Radon problem that arose in the computer microtomography (see, e.g., [5], [8]).

In the case when FOP $\alpha = 1$, the above results obtained can be applied to the standard X-ray DO using the computer diffraction microtjmography technique (cf. [26]).

In contrast to [7], [8], some advantage of our study is to open a window to develop the theory models of the X-ray and electron diffraction scattering by imperfect crystals. The fractional DO theory operates based on the matrix Fredholm–Volterra integral equation of the second kind. The matrix Resolvent-function solution of the Fredholm–Volterra integral equation of the second kind allows us to model the X-ray diffraction scattering by imperfect crystals. Besides, it can modify the optimizing procedure of the χ^2 -target function in trial to solve the inverse Radon problem in the DO microtomography investigating the crystal-lattice defects structure.

The common approach of the DO imaging, the FOP $\alpha = 1$, has to be rethinked using the fractional derivatives formalism $\partial_{0t}^{\alpha}E$ in the sense of the Gerasimov–Caputo approach (2.1). Accordingly, in the fractional DO, it is also worth noting alternative interest due to the possibility of using the fractional Riemann–Liouville derivative $D_{0t}^{\alpha}E$ (2.2). The latter is a good topic for future work in the fractional DO.

Список литературы

- [1] S. Takagi, Dynamical theory of diffraction applicable to crystals with any kind of small distortion. *Acta Crystallogr.* **15**, (1962), 1311–1312.
- [2] S. Takagi, A Dynamical Theory of Diffraction for a Distorted Crystal. J. Phys. Soc. Jpn., 26, (1969), 1239–1253.
- [3] D. Taupin, Théorie dynamique de la diffraction des rayon X par les cristaux déformés, Bull. Soc. Jpn. Fr. Miner. Crist., 87, (1964), 469–511
- [4] D. K. Bowen, B. K. Tanner, *High Resolution X-Ray Diffractometry and Topography*. Taylor & Francis, New York (1998).
- [5] A. Authier, *Dynamical Theory of X-ray Diffraction*. Oxford University Press, New York (2001).

- [6] F.N. Chukhovskii, P.V. Petrashen'. A general dynamical theory of the X-ray Laue diffraction from a homogeneously bent crystal, Acta Cryst., A33, (1977), 311–319.
- [7] A.P. Honkanen, C. Ferrero, J.P. Guigay, V. Mocella, A finite-element approach to dynamical diffraction problems in reflection geometry. *J. Appl. Crystallogr.* **51(2)**, (2018). 511–525.
- [8] F.N. Chukhovskii, P.V. Konarev, V.V. Volkov, Towards a solution of the inverse X-ray diffraction tomography challenge: theory and iterative algorithm for recovering the 3D displacement field function of Coulombtype point defects in a crystal. Acta Cryst., A76, (2020), 16–25. https://doi.org/10.1107/S2053273320000145
- [9] K.B. Oldham, J. Spanier, The Fractional Calculus: Theory and Applications of Differentiation and Integration to Arbitrary Order, N.Y., London: Academic; Press, (1974).
- [10] K.S. Miller, B. Ross, An Introduction to the Fractional Calculus and Fractional Differential Equations, Wiley and Sons, New York, (1993).
- [11] I. Podlubny, Fractional Differential Equations, Academic Press, San Diego, (1999).
- [12] R. Hilfer, Applications of fractional calculus in physics, World Scientific, New Jersey, (2000).
- [13] A.M. Nakhushev, Fractional calculus and its applications, Moscow: Fizmatlit, (2003). (In Russian).
- [14] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, *Theory and Applications of Fractional Differential Equations*, in: North Holland Mathematics Studies, 204, Elsevier Science, Publishers BV, Amsterdam, (2006).
- [15] V.V. Uchaikin, The method of fractional derivatives, Artishok, Ulianovsk, (2008). (In Russian).
- [16] T.M. Atanackovic, S. Pilipovic, B.Stankovic and D. Zorica, Fractional calculus with applications in mechanics: wave propagation, impact and variational principles, John Wiley and Sons, (2014).
- [17] M.O. Mamchuev, Fundamental Solution of a System of Fractional Partial Differential Equations. *Differential Equations*, **46**, No. 8. (2010), 1123–1134.

- [18] M.O. Mamchuev, Cauchy problem in non-local statement for a system of fractional partial differential equations. *Differential Equations*, **48**, No. 3. (2012), 354–361.
- [19] A. Heibig, Existence of solutions for a fractional derivative system of equations, *Integral Equation and Operator Theory*, **72**, (2012), 483–508.
- [20] A.N. Kochubei, Fractional-hiperbolic systems, Fractional Calculus and Applied Analysis 16, No 4. (2013), 860–873. DOI: 10.2478/s13540-013-0053-4
- [21] M.O. Mamchuev, Boundary value problems for equations and systems with the partial derivatives of fractional order, Publishing house KBSC of RAS, Nalchik, (2013). (In Russian).
- [22] M.O. Mamchuev, Mixed problem for loaded system of equations with Riemann-Liouville derivatives, *Mathematical notes*, **97**, No. 3. (2015), 412–222.
- [23] M.O. Mamchuev, Mixed problem for a system of fractional partial differential equations, *Differential Equations*, **52**, No. 1. (2016), 133–138.
- [24] M.O. Mamchuev, Non-local boundary value problem for a system of equations with the partial derivatives of fractional order, *Mathematical notes of NEFU*, **26**, No. 1. (2019), 23-31. (In Russian).
- [25] M.O. Mamchuev, Cauchy problem for a system of equations with the partial Gerasimov Caputo derivatives, *Reports of Circassian International Academy of Sciences*, **21**, No. 4. (2021), 15-22. (In Russian).
- [26] M.O. Mamchuev, F.N. Chukhovskii, Towards to solution of the fractional Takagi – Taupin equations. The Green function method, Fract. Calc. Appl. Anal., 26, No. 2. (2023), 851—863, DOI: 10.1007/s13540-023-00137-4.
- [27] E. M. Wright, The asymptotic expansion of the generalized Bessel function. *Proc. London Math. Soc. Ser. II*, **38**, (1934), 257–270.
- [28] A. Pskhu, Fundamental solutions and Cauchy problems for an oddorder partial differential equation with fractional derevative, *Electronic Journal of Differential Equations*, **21**, (2019), 1–13.

- [29] B. Stanković, On the function of E. M. Wright. *Publications de l'Institut Mathe matique*, **10**, No 24, (1970), 113–124.
- [30] M.O. Mamchuev, Solutions of the main boundary value problems for the time-fractional telegraph equation by the Green function method, *Fract. Calc. Appl. Anal.*, **20**, No. 1. (2017), 190–211, DOI: 10.1515/fca-2017-0010.
- [31] A.P. Prudnikov, Yu.A. Brychkov, O.I. Marichev, *Integrals and series*. Special functions, Nauka, Moscow, Main editorial office of physical and mathematical literature, (1983). (In Russian).
- [32] A.V. Pskhu, Fractional Partial Differential Equations, Nauka, Moscow (2005). (in Russian).
- [33] M.M. Dzhrbashyan, Integral transformations and representations of functions in the complex domain, Nauka, Moscow, (1966), (In Russian).
- [34] M.A. Lavrentiev, B.V. Shabat, Methods of the theory of functions of a complex variable, Nauka, Moscow, (1973). (In Russian).

e man. mamenaev@ramoier.ra (Corr. aa

e-mail: f chukhov@yahoo.ca

Received: March 20, 2024

¹ Institute of Applied Mathematics and Automation, Kabardino-Balkarian Scientific Center RAS, "Shortanov"Str., 89A 360000 Nal'chik, Russian Federation e-mail: mamchuev@rambler.ru (Corr. author)