

On the spectrum of a differential operator on a Hilbert-Pólya space

Xian-Jin Li

Department of Mathematics
Brigham Young University
Provo, Utah 84602 USA
xianjin@byu.edu

Abstract

In this paper we study the spectrum of a fundamental differential operator on a Hilbert-Pólya space. A number is an eigenvalue of this differential operator if and only if it is a nontrivial zero of the Riemann zeta function. An explicit formula is given for the eigenfunction associated with each nontrivial zero of the zeta function. Every eigenfunction is characterized via the Poisson summation formula by a sequence of mysterious functions whose explicit formulas are given.

Subject Class: Primary 30A99

Key Words: Differential operator, zeta zero

1 Introduction

Let $S(\mathbb{R}_+^\times)$ be the set of all infinitely differentiable functions f on \mathbb{R}_+^\times with $\sup_{x \in (0, \infty)} |(\log x)^a f^{(b)}(x)| < \infty$ for $a, b = 0, 1, 2, \dots$. The fundamental differential operator on $S(\mathbb{R}_+^\times)$ is

$$Df(x) = -xf'(x).$$

We denote $Zf(x) = \sum_{n=1}^{\infty} f(nx)$,

$$\mathcal{H}_- = \{f \mid \mathbb{R}_+^\times \rightarrow \mathbb{C} \text{ such that } x^\alpha f(x) \in S(\mathbb{R}_+^\times) \text{ for every real number } \alpha\},$$

and \mathcal{H}_\cap is the set of all even functions $f \in S(\mathbb{R})$ such that $f(0) = 0$ and $\mathfrak{F}f(0) = 0$ where the Fourier transform of f is

$$\mathfrak{F}f(y) = \int_{-\infty}^{\infty} f(x)e^{2\pi ixy}dx.$$

The the quotient space

$$\mathcal{H} = \mathcal{H}_- / Z\mathcal{H}_\cap$$

is called a Hilbert-Pólya space. Let D_- be the operator induced by D on \mathcal{H} .

If ρ is a nontrivial zero of $\zeta(s)$, the author proved in [3] that

$$F_\rho(x) = \int_1^\infty Z\eta(tx)t^{\rho-1}dt$$

is an eigenfunction of D_- in \mathcal{H} associated with the eigenvalue ρ , where $\eta(x) = 8\pi x^2 (\pi x^2 - \frac{3}{2}) e^{-\pi x^2}$. That is,

$$DF_\rho(x) = \rho F_\rho(x) + Z\eta(x), \quad \eta \in \mathcal{H}_\cap. \quad (1)$$

It is also proved in [4] that a complex number ρ is an eigenvalue of D_- on \mathcal{H} if and only if ρ is a nontrivial zero of the zeta function.

In his research on global trace formula [2], the author noticed a mysterious function associated with each function in \mathcal{H} that prevents the Poisson summation formula to hold for that function. In [3] he was able to identify this mysterious function and obtained the following result.

Theorem 1.1 ([3, Theorem 1.3]) *Let $\delta_{l,F}(x) = \int_0^\infty Z^{-1}F(xt) \frac{\sin(2l+1)\pi t}{\pi t} dt$. Then*

$$\lim_{l \rightarrow \infty} \left(\frac{1}{x} \sum_{m=1}^l \mathfrak{Z}Z^{-1}F\left(\frac{m}{x}\right) - \delta_{l,F}(x) \right) = F(x)$$

for every $F \in \mathcal{H}_-$.

Because of the appearance of the unbounded operator Z^{-1} inside $\delta_{l,F}(x)$, it is difficult to know what the function $\delta_{l,F}(x)$ is exactly. In this paper, author was able to get an exact simple formula for $\delta_{l,F_\rho}(x)$ for every nontrivial zero ρ of the Riemann zeta function. More precisely, the author obtained the following formula.

Theorem 1.2 *For every nontrivial zero ρ of the Riemann zeta function, we have*

$$\begin{aligned} \delta_{l,F_\rho}(x) &= \frac{\widehat{\eta}(1-\rho)(l+\frac{1}{2})^\rho}{\rho x^\rho} + 4\pi\left(\frac{l+\frac{1}{2}}{x}\right)^\rho \int_{\frac{l+\frac{1}{2}}{x}}^\infty e^{-\pi u^2} u^{2-\rho} du \\ &= -4\pi\left(\frac{l+\frac{1}{2}}{x}\right)^3 \int_0^1 u^{2-\rho} e^{-\pi(\frac{l+\frac{1}{2}}{x}u)^2} du. \end{aligned}$$

When $\Re\rho < 1/2$, the following improvement of Theorem 1.1 is also obtained in Section 3.

Theorem 1.3 *If $\Re\rho < 1/2$, then*

$$\left\| \frac{1}{x} \sum_{m=1}^l \mathfrak{Z}Z^{-1}F_\rho\left(\frac{m}{x}\right) - F_\rho(x) - \delta_{l,F_\rho}(x) \right\|_{L^2(0,\infty)} \ll \frac{1}{2l+1}$$

for each positive integer l .

2 Proof of Theorem 1.2

Lemma 2.1 ([3, Lemma 3.2]) For every element $F \in \mathcal{H}_-$ we have

$$\delta_{l,F}(x) = \int_0^{\frac{l+\frac{1}{2}}{x}} \mathfrak{F} Z^{-1} F(t) dt.$$

Lemma 2.2 We can write

$$\delta_{l,F_\rho}(x) = \frac{\widehat{\eta}(1-\rho)(l+\frac{1}{2})^\rho}{\rho x^\rho} - \int_0^{\frac{l+\frac{1}{2}}{x}} \left(\int_1^\infty \eta(tu) u^{-\rho} du \right) dt.$$

Proof. Since $\eta \in \mathcal{H}_\cap$, for any fixed $x > 0$ and for all $t \geq 1$ we have $|\eta(tx)| \leq c_x(tx)^{-2}$ for a constant c_x depending on x . Thus,

$$\int_1^\infty \left| \sum_{n \geq X+1} \eta(ntx) \right| t^{\Re\rho-1} dt \leq \frac{c_x}{x^2 X (2 - \Re\rho)} \rightarrow 0$$

as $X \rightarrow \infty$. This implies that we can change the order of summation and integration to write

$$F_\rho(x) = Z \int_1^\infty \eta(tx) t^{\rho-1} dt.$$

By Lemma 2.1,

$$\begin{aligned} \delta_{l,F_\rho}(x) &= \int_0^\infty Z^{-1} F_\rho(t) \frac{\sin \frac{l+\frac{1}{2}}{x} 2\pi t}{\pi t} dt \\ &= 2 \int_0^\infty dt \int_0^{\frac{l+\frac{1}{2}}{x}} \cos 2\pi v t dv \int_1^\infty \eta(ut) u^{\rho-1} du \\ &= 2 \int_0^{\frac{l+\frac{1}{2}}{x}} dv \int_0^\infty \left(\int_1^\infty \eta(ut) u^{\rho-1} du \right) \cos 2\pi v t dt \\ &= 2 \int_0^{\frac{l+\frac{1}{2}}{x}} dv \int_1^\infty u^{\rho-1} du \int_0^\infty \eta(ut) \cos 2\pi v t dt \end{aligned}$$

where changes of order of integration are permissible because the triple integral is absolute integrable. Since $\mathfrak{F}\eta = \eta$, we have

$$2 \int_0^\infty \eta(ut) \cos 2\pi v t dt = \mathfrak{F}_t \{ \eta(ut) \}(v) = \frac{1}{u} \eta\left(\frac{v}{u}\right)$$

Thus

$$\begin{aligned}
\delta_{l,F_\rho}(x) &= \int_0^{\frac{l+\frac{1}{2}}{x}} dv \int_1^\infty \eta\left(\frac{v}{u}\right) u^{\rho-2} du \\
&= \int_0^{\frac{l+\frac{1}{2}}{x}} v^{\rho-1} dv \int_0^\infty \eta\left(\frac{1}{u}\right) u^{\rho-2} du - \int_0^{\frac{l+\frac{1}{2}}{x}} dv \int_0^1 \eta\left(\frac{v}{u}\right) u^{\rho-2} du \\
&= \frac{\hat{\eta}(1-\rho)(l+\frac{1}{2})^\rho}{\rho x^\rho} - \int_0^{\frac{l+\frac{1}{2}}{x}} dt \int_1^\infty \eta(tu) u^{-\rho} du.
\end{aligned}$$

This completes the proof of the lemma. \square

Proof of Theorem 1.2. Since

$$\frac{d}{dt} \left(-4\pi t^3 e^{-\pi t^2} \right) = -4\pi(3t^2 - 2\pi t^4) e^{-\pi t^2} = 8\pi t^2 (\pi t^2 - \frac{3}{2}) e^{-\pi t^2} = \eta(t),$$

we have

$$\int_0^A \eta(t) dt = -4\pi \frac{A^3}{e^{\pi A^2}}.$$

Thus

$$\begin{aligned}
\int_0^{\frac{l+\frac{1}{2}}{x}} \left(\int_1^\infty \eta(tu) u^{-\rho} du \right) dt &= \int_1^\infty u^{-\rho} du \int_0^{\frac{l+\frac{1}{2}}{x}} \eta(tu) dt \\
&= \int_1^\infty u^{-\rho-1} du \int_0^{\frac{l+\frac{1}{2}}{x}u} \eta(t) dt = -4\pi \left(\frac{l+\frac{1}{2}}{x}\right)^3 \int_1^\infty u^{2-\rho} e^{-\pi(\frac{l+\frac{1}{2}}{x}u)^2} du \\
&= -4\pi \left(\frac{l+\frac{1}{2}}{x}\right)^\rho \int_{\frac{l+\frac{1}{2}}{x}}^\infty e^{-\pi u^2} u^{2-\rho} du \\
&= \frac{\hat{\eta}(1-\rho)(l+\frac{1}{2})^\rho}{\rho x^\rho} + 4\pi \left(\frac{l+\frac{1}{2}}{x}\right)^\rho \int_0^{\frac{l+\frac{1}{2}}{x}} e^{-\pi u^2} u^{2-\rho} du.
\end{aligned}$$

By Lemma 2.2,

$$\delta_{l,F_\rho}(x) = \frac{\hat{\eta}(1-\rho)(l+\frac{1}{2})^\rho}{\rho x^\rho} - \int_0^{\frac{l+\frac{1}{2}}{x}} \left(\int_1^\infty \eta(tu) u^{-\rho} du \right) dt.$$

It follows that

$$\delta_{l,F_\rho}(x) = \frac{\hat{\eta}(1-\rho)(l+\frac{1}{2})^\rho}{\rho x^\rho} + 4\pi \left(\frac{l+\frac{1}{2}}{x}\right)^\rho \int_{\frac{l+\frac{1}{2}}{x}}^\infty e^{-\pi u^2} u^{2-\rho} du = -4\pi \left(\frac{l+\frac{1}{2}}{x}\right)^3 \int_0^1 u^{2-\rho} e^{-\pi(\frac{l+\frac{1}{2}}{x}u)^2} du.$$

This completes the proof of Theorem 1.2. \square

3 Proof of Theorem 1.3

Lemma 3.1 ([3, Lemma 4.2]) Let $f \in L^1(\mathbb{R})$ be an even function. If $\mathfrak{F}f(0) = 0$, then

$$\frac{1}{x} \sum_{m=1}^l \mathfrak{F}f\left(\frac{m}{x}\right) = \sum_{m=1}^{\infty} \frac{f(mx+0) + f(mx-0)}{2} + \int_0^{\frac{1}{2}} f(xt) \frac{\sin(2l+1)\pi t}{\sin \pi t} dt + R(f, x)$$

where

$$\begin{aligned} R(f, x) &= \sum_{m=1}^{\infty} \int_m^{m+\frac{1}{2}} \{f(tx) - f(mx+0)\} \frac{\sin(2l+1)\pi t}{\sin \pi t} dt \\ &\quad + \sum_{m=1}^{\infty} \int_{m-\frac{1}{2}}^m \{f(tx) - f(mx-0)\} \frac{\sin(2l+1)\pi t}{\sin \pi t} dt \end{aligned}$$

for every positive integer l .

Lemma 3.2 Let $f(t) = Z^{-1}F_{\rho}(t)$ with $\Re \rho < 1/2$. Then

$$\int_0^{\infty} \left| \sum_{m=1}^{\infty} \int_{-\frac{1}{2}}^{\frac{1}{2}} (f(tx+mx) - f(mx)) \frac{\sin(2l+1)\pi t}{\sin \pi t} dt \right|^2 dx \ll \frac{1}{(2l+1)^4}$$

Proof. By Mellin's inversion formula,

$$f(t) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{\widehat{F}_{\rho}(s)}{\zeta(s)} t^{-s} ds$$

for $c \geq 1$. We write

$$(tx+mx)^{-s} - (mx)^{-s} = -s \int_{mx}^{tx+mx} \frac{du}{u^{s+1}} = \frac{-s}{(mx)^s} \int_1^{1+\frac{t}{m}} \frac{du}{u^{s+1}}$$

Since

$$\begin{aligned} &\left| (f(tx+mx) - f(mx)) \frac{\sin(2l+1)\pi t}{\sin \pi t} \right| \\ &\leq \left| \left(\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{-s\widehat{F}_{\rho}(s)}{\zeta(s)} \left[\int_{mx}^{tx+mx} u^{-s-1} du \right] ds \right) \frac{\sin(2l+1)\pi t}{\sin \pi t} \right| \\ &\leq \frac{(2l+1)\pi t}{2\pi \cdot 2t} \frac{t}{m^{1+c} x^c} \int_{c-i\infty}^{c+i\infty} \left| \frac{-s\widehat{F}_{\rho}(s)}{\zeta(s)} \right| |ds| \ll \frac{1}{m^{1+c}} \end{aligned}$$

for $|t| \leq 1/2$ with the implied constant depending only on x and l not on t , the series

$$\sum_{m=1}^{\infty} (f(tx+mx) - f(mx)) \frac{\sin(2l+1)\pi t}{\sin \pi t}$$

is uniformly convergent for $|t| \leq 1/2$. Thus

$$\begin{aligned} & \sum_{m=1}^{\infty} \int_{-\frac{1}{2}}^{\frac{1}{2}} (f(tx + mx) - f(mx)) \frac{\sin(2l+1)\pi t}{\sin \pi t} dt \\ &= \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{-s\widehat{F}_{\rho}(s)}{x^s \zeta(s)} \left\{ \sum_{m=1}^{\infty} \frac{1}{m^s} \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{\sin(2l+1)\pi t}{\sin \pi t} dt \int_1^{1+\frac{t}{m}} \frac{du}{u^{s+1}} \right\} ds \end{aligned}$$

where the change of order of integration is permissible as the triple integral is absolute integrable.

By (1), $\widehat{F}_{\rho}(s) = \xi(s)/(s - \rho)$ for all complex s . Hence,

$$\frac{\widehat{F}_{\rho}(s)}{\zeta(s)} = \frac{s(s-1)\pi^{-s/2}\Gamma(s/2)}{s-\rho}$$

for all complex $s \neq \rho$. An extension of Stirling's formula [6, line 13, p. 151] is that for any fixed value of σ

$$|\Gamma(\sigma + it)| \sim \sqrt{2\pi} e^{-\frac{\pi|t|}{2}} |t|^{\sigma - \frac{1}{2}}$$

as $|t| \rightarrow \infty$. Thus,

$$\frac{\widehat{F}_{\rho}(s)}{\zeta(s)} \ll |s|^{-k}$$

for any positive integer k when $0 \leq \Re s \leq 1$ and $|s| \rightarrow \infty$. It follows that we can move the line of integration $\Re s = c \geq 1$ to $\Re s = \Re \rho + \epsilon < 1/2$ when $0 < x < 1$ and derive

$$\begin{aligned} & \sum_{m=1}^{\infty} \int_{-\frac{1}{2}}^{\frac{1}{2}} (f(tx + mx) - f(mx)) \frac{\sin(2l+1)\pi t}{\sin \pi t} dt \\ &= \begin{cases} \frac{1}{2\pi i} \int_{\Re s=\Re \rho+\epsilon} \frac{-s\widehat{F}_{\rho}(s)}{x^s \zeta(s)} \left\{ \sum_{m=1}^{\infty} \frac{1}{m^s} \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{\sin(2l+1)\pi t}{\sin \pi t} dt \int_1^{1+\frac{t}{m}} \frac{du}{u^{s+1}} \right\} ds & \text{if } 0 < x < 1 \\ \frac{1}{2\pi i} \int_{\Re s=1} \frac{-s\widehat{F}_{\rho}(s)}{x^s \zeta(s)} \left\{ \sum_{m=1}^{\infty} \frac{1}{m^s} \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{\sin(2l+1)\pi t}{\sin \pi t} dt \int_1^{1+\frac{t}{m}} \frac{du}{u^{s+1}} \right\} ds & \text{if } x \geq 1. \end{cases} \end{aligned}$$

We write

$$\begin{aligned}
& \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{\sin(2l+1)\pi t}{\sin \pi t} dt \int_1^{1+\frac{t}{m}} \frac{du}{u^{s+1}} = \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{\cos(2l+1)\pi t}{s(2l+1)\pi} \frac{d}{dt} \left\{ \frac{\frac{1-(1+\frac{t}{m})^{-s}}{t}}{\frac{\sin \pi t}{t}} \right\} dt \\
&= \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{\cos(2l+1)\pi t}{(2l+1)\pi} \left\{ \frac{\frac{t}{m}(1+\frac{t}{m})^{-s-1} - \frac{1-(1+\frac{t}{m})^{-s}}{s}}{t \sin \pi t} + \frac{d}{dt} \left(\frac{t}{\sin \pi t} \right) \frac{1}{t} \int_1^{1+\frac{t}{m}} \frac{du}{u^{s+1}} \right\} dt \\
&= \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{\cos(2l+1)\pi t}{(2l+1)\pi} \left\{ \frac{\frac{(1+\frac{t}{m})^{-s-1}}{t} \int_1^{1+\frac{t}{m}} \frac{1-(\frac{1+\frac{t}{m}}{t})^{s+1}}{t} du}{\frac{\sin \pi t}{t}} + \frac{d}{dt} \left(\frac{t}{\sin \pi t} \right) \frac{1}{t} \int_1^{1+\frac{t}{m}} \frac{du}{u^{s+1}} \right\} dt \\
&= \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{\cos(2l+1)\pi t}{(2l+1)\pi} \left\{ \frac{\frac{s+1}{t^{2(1+\frac{t}{m})^{s+1}}} \int_1^{1+\frac{t}{m}} du \int_1^{\frac{1+t}{m}} \frac{dv}{v^{s+2}}}{\frac{\sin \pi t}{t}} + \frac{d}{dt} \left(\frac{t}{\sin \pi t} \right) \frac{1}{t} \int_1^{1+\frac{t}{m}} \frac{du}{u^{s+1}} \right\} dt \\
&= \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{\cos(2l+1)\pi t}{(2l+1)\pi} \left\{ \frac{\frac{s+1}{m^2(1+\frac{t}{m})^{s+1}} \int_0^1 du \int_0^{\frac{1-u}{m}} \frac{dv}{(1+\frac{t}{m}v)^{s+2}}}{\frac{\sin \pi t}{t}} + \frac{d}{dt} \left(\frac{t}{\sin \pi t} \right) \frac{1}{m} \int_0^1 \frac{du}{(1+\frac{t}{m}u)^{s+1}} \right\} dt
\end{aligned}$$

Since $\sin t \geq 2t/\pi$ for $t \in [0, \pi/]$, for $|t| \leq 1/2$ we have $|\sin \pi t/t| \geq 2$. Applying partial integration to the last term of the above identity we obtain that

$$\int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{\sin(2l+1)\pi t}{\sin \pi t} dt \int_1^{1+\frac{t}{m}} \frac{du}{u^{s+1}} \ll \frac{(|s|+1)^2}{m(2l+1)^2}.$$

It follows that

$$\left| \sum_{m=1}^{\infty} \int_{-\frac{1}{2}}^{\frac{1}{2}} (f(tx+mx) - f(mx)) \frac{\sin(2l+1)\pi t}{\sin \pi t} dt \right| \ll \begin{cases} \frac{x^{-\Re \rho - \epsilon}}{(2l+1)^2} & \text{if } 0 < x < 1 \\ \frac{x^{-1}}{(2l+1)^2} & \text{if } x \geq 1. \end{cases}$$

Therefore

$$\int_0^{\infty} \left| \sum_{m=1}^{\infty} \int_{-\frac{1}{2}}^{\frac{1}{2}} (f(tx+mx) - f(mx)) \frac{\sin(2l+1)\pi t}{\sin \pi t} dt \right|^2 dx \ll \frac{1}{(2l+1)^4}.$$

This completes the proof of the lemma. \square

Lemma 3.3 *Let $f(t) = Z^{-1}F_{\rho}(t)$. Then*

$$\int_0^{\infty} \left| \int_0^{\frac{1}{2}} f(xt) \left(\frac{1}{\pi t} - \frac{1}{\sin \pi t} \right) \sin(2l+1)\pi t dt \right|^2 dx \ll \frac{1}{(2l+1)^2}.$$

Proof. Note that

$$f(t) = \frac{1}{2\pi i} \int_{\Re s=c} \frac{\widehat{F}_{\rho}(s)}{\zeta(s)} t^{-s} ds$$

for $c \geq 1$. Because the following double integral is absolute integrable, we can change order of integration to derive

$$\begin{aligned} & \int_0^{\frac{1}{2}} f(xt) \left(\frac{1}{\pi t} - \frac{1}{\sin \pi t} \right) \sin(2l+1)\pi t dt \\ &= \frac{1}{2\pi i} \int_{\Re s=c} \frac{\widehat{F}_\rho(s)}{\zeta(s)} \left\{ \int_0^{\frac{1}{2}} t^{-s} \left(\frac{1}{\pi t} - \frac{1}{\sin \pi t} \right) \sin(2l+1)\pi t dt \right\} \frac{ds}{x^s}. \end{aligned}$$

Similarly as in proof of Lemma 3.2 we write

$$\begin{aligned} & \int_0^{\frac{1}{2}} f(xt) \left(\frac{1}{\pi t} - \frac{1}{\sin \pi t} \right) \sin(2l+1)\pi t dt \\ &= \begin{cases} \frac{1}{2\pi i} \int_{\Re s=\Re\rho+\epsilon} \frac{\widehat{F}_\rho(s)}{\zeta(s)} \left\{ \int_0^{\frac{1}{2}} t^{-s} \left(\frac{1}{\pi t} - \frac{1}{\sin \pi t} \right) \sin(2l+1)\pi t dt \right\} \frac{ds}{x^s} & \text{if } 0 < x < 1 \\ \frac{1}{2\pi i} \int_{\Re s=\frac{1}{2}+\epsilon} \frac{\widehat{F}_\rho(s)}{\zeta(s)} \left\{ \int_0^{\frac{1}{2}} t^{-s} \left(\frac{1}{\pi t} - \frac{1}{\sin \pi t} \right) \sin(2l+1)\pi t dt \right\} \frac{ds}{x^s} & \text{if } x \geq 1 \end{cases} \end{aligned}$$

for a sufficiently small positive ϵ .

By [1, (13), p. 190],

$$\frac{1}{t} - \frac{\pi}{\sin(\pi t)} = 2t \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2 - t^2} = 2t \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} \sum_{k=0}^{\infty} \left(\frac{t}{n} \right)^{2k}$$

for $0 < t < 1/2$. Note that

$$2 \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} = -\frac{\pi^2}{6}$$

Applying partial integration twice we derive that

$$\begin{aligned} & \int_0^{\frac{1}{2}} t^{-s} \left\{ 2t \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} \sum_{k=1}^{\infty} \left(\frac{t}{n} \right)^{2k} \right\} \sin(2l+1)\pi t dt \\ &= - \int_0^{\frac{1}{2}} t^{3-s} \left\{ 2 \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} \sum_{k=1}^{\infty} \left(\frac{t}{n} \right)^{2k-2} \right\} d \frac{\cos(2l+1)\pi t}{(2l+1)\pi} \ll \frac{(|s|+3)^2}{(2l+1)^2} \end{aligned}$$

for $0 < \Re s < 1$.

Also, by partial integration we have

$$\begin{aligned} & \int_0^{\frac{1}{2}} t^{-s} \left\{ 2t \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} \right\} \sin(2l+1)\pi t dt = -\frac{\pi^2}{6} \int_0^{\frac{1}{2}} t^{1-s} \sin(2l+1)\pi t dt \\ &= \frac{\pi(s-1)}{6(2l+1)} \int_0^{\frac{1}{2}} t^{-s} \cos(2l+1)\pi t dt = \frac{\pi^s(s-1)}{6(2l+1)^{2-s}} \int_0^{l+\frac{1}{2}} t^{-s} \cos t dt. \end{aligned}$$

for $0 < \Re s < 1$. By [6, Example 10, p. 162],

$$\int_0^{\infty} t^{-s} \cos t dt = \Gamma(1-s) \sin \frac{\pi s}{2}$$

for $0 < \Re s < 1$. As

$$\int_{l+\frac{1}{2}}^{\infty} t^{-s} \cos t dt = -\frac{\sin(l + \frac{1}{2})}{(l + \frac{1}{2})^s} + s \int_{l+\frac{1}{2}}^{\infty} t^{-s-1} \sin t dt \ll \frac{|s|+1}{(l + \frac{1}{2})^{\Re s}}$$

we have

$$\int_0^{\frac{1}{2}} t^{-s} \left\{ 2t \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} \right\} \sin(2l+1)\pi t dt = -\frac{\pi^s \Gamma(2-s) \sin \frac{\pi s}{2}}{6(2l+1)^{2-s}} + O\left(\frac{(|s|+1)^2}{(2l+1)^2}\right)$$

Thus, we have obtained that

$$\int_0^{\frac{1}{2}} t^{-s} \left(\frac{1}{\pi t} - \frac{1}{\sin \pi t} \right) \sin(2l+1)\pi t dt = -\frac{\pi^s \Gamma(2-s) \sin \frac{\pi s}{2}}{6(2l+1)^{2-s}} + O\left(\frac{(|s|+3)^2}{(2l+1)^2}\right)$$

Hence

$$\begin{aligned} & \int_0^{\frac{1}{2}} f(xt) \left(\frac{1}{\pi t} - \frac{1}{\sin \pi t} \right) \sin(2l+1)\pi t dt \\ &= \frac{1}{2\pi i} \int_{\Re s=c} \frac{\widehat{F}_\rho(s)}{\zeta(s)} \left\{ -\frac{\pi^s \Gamma(2-s) \sin \frac{\pi s}{2}}{6(2l+1)^{2-s}} + O\left(\frac{(|s|+3)^2}{(2l+1)^2}\right) \right\} \frac{ds}{x^s}. \end{aligned}$$

By the proof of Lemma 3.2,

$$\frac{\widehat{F}_\rho(s)}{\zeta(s)} \ll |s|^{-k}$$

for any positive integer k when $0 \leq \Re s \leq 1$ and $|s| \rightarrow \infty$. Thus, by estimations

$$\frac{1}{2\pi i} \int_{\Re s=c} \frac{\widehat{F}_\rho(s)}{\zeta(s)} \left\{ -\frac{\pi^s \Gamma(2-s) \sin \frac{\pi s}{2}}{6(2l+1)^{2-s}} + O\left(\frac{(|s|+3)^2}{(2l+1)^2}\right) \right\} \frac{ds}{x^s} \ll \begin{cases} \frac{x^{-\Re \rho - \epsilon}}{(2l+1)^{2-\Re \rho - \epsilon}} & \text{if } 0 < x < 1 \\ \frac{x^{-1}}{2l+1} & \text{if } x \geq 1. \end{cases}$$

Therefore,

$$\int_0^{\frac{1}{2}} f(xt) \left(\frac{1}{\pi t} - \frac{1}{\sin \pi t} \right) \sin(2l+1)\pi t dt = O\left(\begin{cases} \frac{x^{-\Re \rho - \epsilon}}{(2l+1)^{2-\Re \rho - \epsilon}} & \text{if } 0 < x < 1 \\ \frac{x^{-1}}{2l+1} & \text{if } x \geq 1 \end{cases}\right).$$

It follows that

$$\int_0^{\infty} \left| \int_0^{\frac{1}{2}} f(xt) \left(\frac{1}{\pi t} - \frac{1}{\sin \pi t} \right) \sin(2l+1)\pi t dt \right|^2 dx \ll \frac{1}{(2l+1)^2}.$$

This completes the proof of the lemma. \square

Lemma 3.4 Let $f(t) = Z^{-1}F_\rho(t)$ with $\Re \rho < 1/2$. Then

$$\int_0^{\infty} \left| \int_{\frac{1}{2}}^{\infty} f(xt) \frac{\sin(2l+1)\pi t}{\pi t} dt \right|^2 dx \ll \frac{1}{(2l+1)^4}$$

Proof. As

$$f(t) = \frac{1}{2\pi i} \int_{\Re s=c} \frac{\widehat{F}_\rho(s)}{\zeta(s)} t^{-s} ds$$

for $c \geq 1$, we can write

$$\int_{\frac{1}{2}}^{\infty} f(xt) \frac{\sin(2l+1)\pi t}{\pi t} dt = \frac{1}{2\pi i} \int_{\Re s=c} \frac{\widehat{F}_\rho(s)}{\zeta(s)} \left\{ \int_{\frac{1}{2}}^{\infty} \frac{\sin(2l+1)\pi t}{\pi t^{1+s}} dt \right\} \frac{ds}{x^s}$$

where change of order of integration is permissible as the double integral is absolute integrable.

By partial integration,

$$\int_{\frac{1}{2}}^{\infty} f(xt) \frac{\sin(2l+1)\pi t}{\pi t} dt = \begin{cases} \frac{-1}{2\pi i} \int_{\Re s=1} \frac{(1+s)\widehat{F}_\rho(s)}{\zeta(s)} \left\{ \int_{\frac{1}{2}}^{\infty} \frac{\cos(2l+1)\pi t}{(2l+1)\pi^2 t^{2+s}} dt \right\} \frac{ds}{x^s} & \text{if } x \geq 1 \\ \frac{-1}{2\pi i} \int_{\Re s=\Re\rho+\epsilon} \frac{(1+s)\widehat{F}_\rho(s)}{\zeta(s)} \left\{ \int_{\frac{1}{2}}^{\infty} \frac{\cos(2l+1)\pi t}{(2l+1)\pi^2 t^{2+s}} dt \right\} \frac{ds}{x^s} & \text{if } x < 1 \end{cases}$$

where ϵ is a sufficiently small positive satisfying $\Re\rho + \epsilon < \frac{1}{2}$. Also,

$$\int_{\frac{1}{2}}^{\infty} \frac{\cos(2l+1)\pi t}{(2l+1)t^{2+s}} dt = -\frac{2^{2+s} \sin(l+\frac{1}{2})\pi}{(2l+1)^2 \pi} + \frac{2+s}{\pi(2l+1)^2} \int_{\frac{1}{2}}^{\infty} \frac{\sin(2l+1)\pi t}{t^{3+s}} dt \ll \frac{|s|+2}{(2l+1)^2}$$

for $0 < \Re s < 1$. It follows that

$$\left| \int_{\frac{1}{2}}^{\infty} f(xt) \frac{\sin(2l+1)\pi t}{\pi t} dt \right| \ll \begin{cases} \frac{x^{-\Re\rho-\epsilon}}{(2l+1)^2} & \text{if } 0 < x \leq 1 \\ \frac{x^{-1}}{(2l+1)^2} & \text{if } x > 1. \end{cases}$$

This inequality implies that

$$\int_0^{\infty} \left| \int_{\frac{1}{2}}^{\infty} f(xt) \frac{\sin(2l+1)\pi t}{\pi t} dt \right|^2 dx \ll \frac{1}{(2l+1)^4}.$$

This completes the proof of the lemma. \square

Proof of Theorem 1.3. For $f(t) = Z^{-1}F_\rho(t)$, we have $f(0) = 0$ and $\mathfrak{F}f(0) = 0$. Also, f is an even function on \mathbb{R} . Thus, by Lemma 3.1

$$\frac{1}{x} \sum_{m=1}^l \mathfrak{F}f\left(\frac{m}{x}\right) = \sum_{m=1}^{\infty} \frac{f(mx+0) + f(mx-0)}{2} + \int_0^{\frac{1}{2}} f(xt) \frac{\sin(2l+1)\pi t}{\sin \pi t} dt + R(f, x) \quad (2)$$

where

$$R(f, x) = \sum_{m=1}^{\infty} \int_{-\frac{1}{2}}^{\frac{1}{2}} (f(tx+mx) - f(mx)) \frac{\sin(2l+1)\pi t}{\sin \pi t} dt.$$

We rewrite (2) as

$$\begin{aligned}
& \frac{1}{x} \sum_{m=1}^l \mathfrak{F}Z^{-1}F_\rho\left(\frac{m}{x}\right) - \delta_{l,F_\rho}(x) - F_\rho(x) \\
&= \int_0^{\frac{1}{2}} f(xt)\left(\frac{1}{\sin \pi t} - \frac{1}{\pi t}\right) \sin(2l+1)\pi t dt - \int_{\frac{1}{2}}^\infty f(xt) \frac{\sin(2l+1)\pi t}{\pi t} dt \quad (3) \\
&+ \sum_{m=1}^{\infty} \int_{-\frac{1}{2}}^{\frac{1}{2}} (f(tx+mx) - f(mx)) \frac{\sin(2l+1)\pi t}{\sin \pi t} dt.
\end{aligned}$$

By (3), the Minkowski's inequality [5, Theorem 3.5, p. 63], and Lemmas 3.2, 3.3 and 3.4 we obtain that

$$\left\| \left(\frac{1}{x} \sum_{m=1}^l \mathfrak{F}Z^{-1}F_\rho\left(\frac{m}{x}\right) - \delta_{l,F_\rho}(x) \right) - F_\rho(x) \right\|_{L^2(0,\infty)} \ll \frac{1}{2l+1}.$$

This completes the proof of Theorem 1.3. \square

References

- [1] L. V. Ahlfors, Complex Analysis, Third Edition, McGraw-Hill, 1979
- [2] Xian-Jin Li, On Weil's explicit formula, *Sci. China Math.* **58** (2015), 915–982
- [3] Xian-Jin Li, On spectral theory of the Riemann zeta function, *Sci. China Math.* **62** (2019), no. 11, 2317–2330
- [4] L. Ge, X.-J. Li, D. Wu, and B. Xue, Eigenvalues of a differential operator and zeros of the Riemann ζ -function, *Anal. Theory Appl.* **36** (2020), no. 3, 283–294
- [5] W. Rudin, Real and Complex Analysis, Third Edition, McGraw-Hill, 1987
- [6] E. C. Titchmarsh, The Theory of Functions, Second Edition, Oxford, 1939