

Cosmological Inflation and Dark Sector from 11D Supergravity

Jiaming Shi^{a,b} and Zehua Xiao^{c,1,2}

^a*School of Fundamental Physics and Mathematical Sciences, Hangzhou Institute for Advanced Study, University of Chinese Academy of Sciences, Hangzhou 310024, China*

^b*School of Physical Sciences, University of Chinese Academy of Sciences, No.19A Yuquan Road, Beijing 100049, China*

^c*Yau Mathematical Sciences Center, Tsinghua University, Beijing 100084, China*

E-mail: xiaozh20@mails.tsinghua.edu.cn, shijiaming@ucas.ac.cn

ABSTRACT: We explore compactifications of the form $\mathcal{M}^{3,1} \times T_{q_1}^2 \times T_{q_2}^2 \times T_{q_3}^2 \times S^1$ in the framework of 11D supergravity. By imposing suitable gauge conditions and boundary conditions, we find that the four-dimensional FRW universe emerges as a solution representing cosmological D3-branes in the eleven-dimensional bulk. These specific compactification methods can produce cosmological inflation that aligns with the observational constraints set by the 2021 BICEP/Keck [1] and Planck 2018 results [2]. In the cosmological inflation models we construct, the inflaton can be interpreted as the conformal vibrations of extra dimensions with a size around 10^5 times the reduced Planck length. Additionally, we offer the theoretical predictions for the mass of the inflaton, and the tree-level Newton's gravity law between two massive point particles surrounded by a spherically symmetric distribution of the inflaton, which can reproduce the Tully-Fisher relation and explain the flat rotation curves of galaxies.

¹Co-first author

²Corresponding author

Contents

1	Introduction	1
2	Compactification Towards Four Dimensions	4
2.1	A Special Case	4
2.2	General Case	10
3	Cosmological Inflation	12
3.1	Numerical Results	12
3.2	Exit Inflation	16
4	Dark Sector from Extra Dimensions	16
4.1	Dark Matter	19
4.2	Dark Energy	21
5	Conclusions and Discussion	22

1 Introduction

The eleven-dimensional supergravity is widely believed to be the tree-level low-energy effective theory of M-theory. As an ultraviolet complete theory, M-theory possesses inherent advantages in addressing extremely high-energy scenarios, such as the inflationary period of the early universe. However, our understanding of M-theory remains limited, see [3–5] for remarkable progress in M-theory. A comprehensive theoretical description is still necessary for understanding physics in the extremely early universe by M-theory. From the perspective of string theory, it is conjectured that by compactifying M-theory on a special manifold of G_2 holonomy, it could yield the supersymmetric version of Standard Model of Particle Physics and General Relativity [6, 7]. This conjecture has generated some discussions on M-theory cosmology in recent years, particularly regarding G_2 compactification, see [8–10] for examples. While this paper does not aim to fully realize the ambitious goals of M-theory cosmology, it seeks to explore other potential ways for dimensional reduction that could lead to cosmological inflation in the early universe.

In contrast to M-theory, its low-energy effective theory, eleven-dimensional supergravity, has been more extensively studied. Comprehensive reviews, such as [11, 12], provide a valuable resource for understanding this field. It is believed that eleven-dimensional supergravity serves as the mother theory for lower-dimensional supergravity theories. Numerous papers have delved into various lower-dimensional supergravity theories and their cosmological models [13–19], including discussions on no-scale inflation as discussed in references [20–23]. However, cosmological inflation models based on the original eleven-dimensional

supergravity remain rare. Early discussions have touched upon the inflationary scenario involving the compactification of seven dimensions as S^7 [24, 25]. Other approaches from M-theory to string cosmology such as Calabi-Yau compactification $CY^3 \times S^1$ were studied in [26, 27].

After inflation, dark matter plays a vital role in cosmic expansion and galaxy formation which can be interpreted as a scalar field, specifically the axion or dilaton field [28–34]. The dark scalar field non-minimally coupled to gravity was studied in Refs. [35–37]. When considering galactic scales, a scalar field is interpreted as a condensate of ultra-light bosonic dark matter (UBDM) that interacts with general relativity, adhering to the Klein-Gordon equations. In the non-relativistic realm, the equations of motion reduce to the Schrödinger-Poisson system. It is postulated that, at galactic scales, this dark matter forms power-law matter density profiles described by $\rho = \rho_0 r^{-\alpha}$ [38–40] or the Navarro-Frenk-White matter density profiles [41, 42]

$$\rho(r) = \frac{\rho_0}{r/r_s(1 + r/r_s)^2}, \quad (1.1)$$

where r_s is the characteristic scale radius. On the other hand, MODified Newtonian Dynamics (MOND) [43–46] is a possible and promising alternative to the dark matter problem which can explain the flat rotation curves of galaxies and the Tully-Fisher relation $v^4 \propto M$ [47–50], where v is the rotation speed of a galaxy and M is the total baryonic mass (stars plus gas) in the galaxy. However, it is challenged by the observation of Bullet cluster [51] since the modification to dynamics can not reveal the dark matter distribution. To consider the advantages of the MOND theory, we expect that some hidden matter could induce results that are similar to the implications of MOND.

Another mystery in the dark sector of the universe is dark energy [52, 53]. The dark energy with negative pressure dominates the evolution of our current universe and explains the accelerated expansion of the universe. In the standard cosmological model (Λ CDM model), the cosmological constant Λ as the dark energy contributes about 70% of the energy density of the current universe. Unfortunately, the physical interpretation of the dark energy suffers from the cosmological constant problem or the fine-tuning problem since the value of the energy density of cosmological constant $\rho_\Lambda \simeq 10^{-120} m_p^4$, where $m_p \sim 10^{18} \text{GeV}$ is a Planck mass scale [54, 55]. This problem may be alleviated by the mechanism of small-scale spacetime fluctuations [56] or spacetime foam [57]. In fact, dark energy is not necessary to be a cosmological constant. Recent observations from the Dark Energy Spectroscopic Instrument (DESI), along with other datasets including CMB, PantheonPlus, Union3, and DESY5, support the time-varying dark energy equation of state [58].

In this study, we use the tree-level eleven-dimensional supergravity, which we will abbreviate as “11D supergravity”, as the starting point. The action of the 11D supergravity

is given by [11]

$$\begin{aligned}
S = & \frac{m_p^{9(11)}}{2} \int d^{11}x \sqrt{-G_{11}} \left[e^{aM} e^{bN} R_{MNab}(\omega) - \bar{\psi}_A \Gamma^{ABC} D_B \left(\frac{\omega + \hat{\omega}}{2} \right) \psi_C - \frac{1}{24} F^{ABCD} F_{ABCD} \right. \\
& - \frac{\sqrt{2}}{192} \bar{\psi}_M (\Gamma^{ABCDMN} + 12\Gamma^{AB} g^{CM} g^{DN}) \psi_N \left(2F_{ABCD} + \frac{3}{2} \sqrt{2} \bar{\psi}_{[A} \Gamma_{BC} \psi_{D]} \right) \\
& \left. - \frac{2\sqrt{2}}{(144)^2} \varepsilon^{A'B'C'D'ABCDMNP} F_{A'B'C'D'} F_{ABCD} A_{MNP} \right], \tag{1.2}
\end{aligned}$$

where the indices $A, B, C, \dots = 0, 1, 2, \dots, 10$ and $m_p^{(11)}$ is the 11 dimensional reduced Planck mass. e_M^a is the frame fields with $a = 0, 1, 2, \dots, 10$. There are 11D graviton G_{AB} , 11D gravitino ψ_A and 3-form A_{MNP} as elementary fields in 11D supergravity. The field strength F_{ABCD} for the 3-form is defined as $F = dA$. The 3-form possesses a gauge invariance given by $A' = A + d\theta$, in which θ_{MN} serves as a 2-form gauge parameter. The 32 dimensional gamma matrices are $\Gamma^{A_1 A_2 \dots A_k} = \Gamma^{[A_1} \Gamma^{A_2} \dots \Gamma^{A_k]}$. $\varepsilon^{A'B'C'D'ABCDMNP}$ is the totally antisymmetric rank 11 tensor with $\varepsilon^{012\dots 10} = -1$. The covariant derivative is defined as

$$D_M(\tilde{\omega}) = \nabla_M + \frac{1}{4} \tilde{\omega}_{Mab} \Gamma^{ab}, \tag{1.3}$$

where the connection $\tilde{\omega}$ can be

$$\omega_{Mab} = \omega_{Mab}(e) + K_{Mab}, \tag{1.4}$$

$$\hat{\omega}_{Mab} = \omega_{Mab}(e) - \frac{1}{4} (\bar{\psi}_M \Gamma_b \psi_a - \bar{\psi}_a \Gamma_M \psi_b + \bar{\psi}_b \Gamma_a \psi_M), \tag{1.5}$$

$$\omega_M^{ab}(e) = 2e^{N[a} \nabla_{[M} e_N^{b]} - e^{N[a} e^{b]P} e_{Mc} \nabla_N e_P^c, \tag{1.6}$$

$$K_{Mab} = -\frac{1}{4} (\bar{\psi}_M \Gamma_b \psi_a - \bar{\psi}_a \Gamma_M \psi_b + \bar{\psi}_b \Gamma_a \psi_M) + \frac{1}{8} \bar{\psi}_N \Gamma^{NP}{}_{Mab} \psi_P. \tag{1.7}$$

The action (1.2) is invariant under supersymmetry transformation

$$\delta e_M^a = \frac{1}{2} \bar{\epsilon} \Gamma^a \psi_M, \tag{1.8}$$

$$\delta \psi_M = D_M(\hat{\omega}) \epsilon + \frac{\sqrt{2}}{288} (\Gamma^{ABCD}{}_M - 8\Gamma^{BCD} \delta_M^A) \left(F_{ABCD} + \frac{3\sqrt{2}}{2} \bar{\psi}_{[A} \Gamma_{BC} \psi_{D]} \right) \epsilon, \tag{1.9}$$

$$\delta A_{MNP} = -\frac{3\sqrt{2}}{4} \bar{\epsilon} \Gamma_{[MN} \psi_{P]}, \tag{1.10}$$

where Γ^a , $a = 0, 1, 2, \dots, 10$ is the gamma matrix in flat eleven-dimensional tangent space. ϵ , with its Dirac conjugation $\bar{\epsilon}$, is the spinor gauge parameter of eleven-dimensional supersymmetry. In this paper, we define the four-dimensional reduced Planck mass $m_p = \frac{1}{\sqrt{8\pi G}}$, where G is the four-dimensional Newtonian gravitational constant.

In Section 2, we propose that by compactifying seven dimensions in the form of $\mathcal{M}^{3,1} \times T_{q_1}^2 \times T_{q_2}^2 \times T_{q_3}^2 \times S^1$, the action (1.2) can give rise to the complete evolution of the four-dimensional visible universe, encompassing matter described by the Standard Model of

Particle Physics as well as dark matter and dark energy. The four-dimensional extended spacetime, denoted by $\mathcal{M}^{3,1}$, is described using the Friedmann-Robertson-Walker (FRW) metric. The gauge conditions for the gauge parameters θ_{MN} and ϵ of the 11D supergravity can be chosen to prevent the extra dimensions from decompactifying as the universe evolves. Through numerical analysis, in Section 3, we will see that a contraction of the 3 tori and an expansion of the eleventh dimension gives rise to cosmological inflation in the four-dimensional FRW universe. The inflaton is interpreted as the conformal vibrations of the extra dimensions. Furthermore, in Section 4, we discussed the quantum effects coming from the compactified extra dimensions, where the inflaton is supposed to be dark matter or dark energy. We will see that the leading order contribution from the four-point matter scattering amplitude results in a Yukawa-like modification to Newton's law of gravity. Furthermore, this leading order relativistic contribution can explain the Tully-Fisher relation $v^4 \sim M$ observed in galaxies.

2 Compactification Towards Four Dimensions

The universe we see comprises four extended dimensions: three spatial dimensions and one temporal dimension. In the theory of eleven-dimensional supergravity with all loop-order corrections, the remaining seven dimensions of the theory should be compactified to an extremely small scale. These compactified dimensions are presumably microscopic and their vibrations are influenced by quantum effects. At the beginning of the universe, scattering processes involving gravitons, gravitinos, and 3-forms are prevalent, resulting in a boiling spacetime that can not be perceived as a background of the tree-level eleven-dimensional supergravity (1.2). In this scenario, the supergravity necessitates the inclusion of an infinite number of loop-order quantum corrections, which lie beyond our current analytical capabilities. Hence, we postulate that the inflationary period commences when the loop-order graviton scattering processes become a negligible contribution to the universe's total S-matrix, rendering all the loop-order quantum corrections of the graviton field insignificant. In our inflation model, matter fields of four-dimensional spacetime effective action are assumed temporarily inactive during the inflation period, resulting in the graviton and inflaton dominating the universe's dynamics. This assumption of the inflationary period can be interpreted as selecting suitable gauge conditions for the gauge parameters θ_{MN} and ϵ and boundary conditions for the cosmological D3-branes in eleven-dimensional bulk.

2.1 A Special Case

In this subsection, we will focus on a specific dimensional reduction method from 11D to 4D. It involves seven compactified extra dimensions: 2 dimensions form a 2D torus $T_{q_1}^2$ with genus q_1 , 2 dimensions form a 2D torus $T_{q_2}^2$ with genus q_2 , the other 2 dimensions form another 2D torus $T_{q_3}^2$ with genus q_3 , the last 1 dimension forms a circle. The 11D bulk metric is given by

$$ds_{11}^2 = G_{AB} dx^A dx^B. \quad (2.1)$$

Four extended dimensions within the 11D bulk exhibit the FRW universe:

$$ds_4^2 = g_{\mu\nu} dx^\mu dx^\nu = -dt^2 + a^2(t) \delta_{ij} dx^i dx^j, \quad (2.2)$$

where $\mu, \nu = 0, 1, 2, 3$. The metrics of the three 2D tori are given as follows

$$\begin{aligned} ds_{2(1)}^2 &= \Phi(x^\mu) g_{a_1 b_1}(x^{a_1}) dx^{a_1} dx^{b_1}, \\ ds_{2(2)}^2 &= h_{a_2 b_2}(x^{a_2}) dx^{a_2} dx^{b_2}, \\ ds_{2(3)}^2 &= \Phi^2(x^\mu) \gamma_{a_3 b_3}(x^{a_3}) dx^{a_3} dx^{b_3}, \end{aligned} \quad (2.3)$$

where $a_1, b_1 = 4, 5$, $a_2, b_2 = 6, 7$, and $a_3, b_3 = 8, 9$. The scalar $\Phi(x^\mu) > 0$ is the scale function of the tori. It is dynamical, allowing the tori to conformally transform into larger or smaller sizes by changing the value of the function $\Phi(x^\mu)$. The metrics above with scale function $\Phi(x^\mu) = 1$ can be used to describe the standard sizes of these three tori.

The 11th dimension is compactified into a circle by

$$ds_1^2 = \Phi^{-6}(x^\mu) dx^{10} dx^{10}. \quad (2.4)$$

Hence, the metric of the 11D supergravity can be written as

$$ds_{11}^2 = ds_4^2 + ds_{2(1)}^2 + ds_{2(2)}^2 + ds_{2(3)}^2 + ds_1^2. \quad (2.5)$$

The 11D Ricci scalar of G_{AB} after dimensional reduction is given by

$$R_{(11)} = R_{(4)} - \frac{23}{2\Phi^2} \nabla_\mu \Phi \nabla^\mu \Phi + \frac{R_{T_{q_1}^2}}{\Phi} + R_{T_{q_2}^2} + \frac{R_{T_{q_3}^2}}{\Phi^2}, \quad (2.6)$$

where $R_{(4)}$ is the Ricci scalar of the metric $g_{\mu\nu}$ and $R_{T_{q_i}^2}, i = 1, 2, 3$ are the Ricci scalars of the metric $g_{a_1 b_1}, h_{a_2 b_2}, \gamma_{a_3 b_3}$ respectively. The non-zero equations of motion of G_{AB} are provided by Einstein's equations

$$\mathcal{G}_{\mu\nu} - \frac{23}{2\Phi^2} \nabla_\mu \Phi \nabla_\nu \Phi - \frac{1}{2} g_{\mu\nu} \left(\frac{R_{T_{q_1}^2}}{\Phi} + R_{T_{q_2}^2} + \frac{R_{T_{q_3}^2}}{\Phi^2} - \frac{23}{2\Phi^2} \nabla_\rho \Phi \nabla^\rho \Phi \right) = \frac{1}{m_{p(11)}^9} T_{\mu\nu}^{(A,\psi)}, \quad (2.7)$$

$$\mathcal{G}_{a_1 b_1} - \frac{1}{2} g_{a_1 b_1} \left(\Phi R_{(4)} + \Phi R_{T_{q_2}^2} + \frac{R_{T_{q_3}^2}}{\Phi} + \nabla_\mu \nabla^\mu \Phi - \frac{25}{2\Phi} \nabla_\mu \Phi \nabla^\mu \Phi \right) = \frac{1}{m_{p(11)}^9} T_{a_1 b_1}^{(A,\psi)}, \quad (2.8)$$

$$\mathcal{G}_{a_2 b_2} - \frac{1}{2} h_{a_2 b_2} \left(R_{(4)} + \frac{R_{T_{q_1}^2}}{\Phi} + \frac{R_{T_{q_3}^2}}{\Phi^2} - \frac{23}{2\Phi^2} \nabla_\mu \Phi \nabla^\mu \Phi \right) = \frac{1}{m_{p(11)}^9} T_{a_2 b_2}^{(A,\psi)}, \quad (2.9)$$

$$\mathcal{G}_{a_3 b_3} - \frac{1}{2} \gamma_{a_3 b_3} \left(\Phi^2 R_{(4)} + \Phi R_{T_{q_1}^2} + \Phi^2 R_{T_{q_2}^2} + \Phi \nabla_\mu \nabla^\mu \Phi - \frac{27}{2} \nabla_\mu \Phi \nabla^\mu \Phi \right) = \frac{1}{m_{p(11)}^9} T_{a_3 b_3}^{(A,\psi)}, \quad (2.10)$$

$$-\frac{1}{2} \left(\frac{R_{(4)}}{\Phi^6} + \frac{R_{T_{q_1}^2}}{\Phi^7} + \frac{R_{T_{q_2}^2}}{\Phi^6} + \frac{R_{T_{q_3}^2}}{\Phi^8} - \frac{6}{\Phi^7} \nabla_\mu \nabla^\mu \Phi - \frac{11}{2\Phi^8} \nabla_\mu \Phi \nabla^\mu \Phi \right) = \frac{1}{m_{p(11)}^9} T_{10,10}^{(A,\psi)}, \quad (2.11)$$

where $\mathcal{G}_{\mu\nu}, \mathcal{G}_{a_i b_i}, i = 1, 2, 3$, are the Einstein tensor for metric $g_{\mu\nu}, g_{a_1 b_1}, h_{a_2 b_2}, \gamma_{a_3 b_3}$ respectively. The energy-momentum tensor, denoted as $T_{AB}^{(A,\psi)}$, is defined for the gravitino ψ_A and 3-form A_{MNP} of 11D supergravity. Note that there are gauge parameters θ_{MN} and ϵ in 11D supergravity. It is imperative to incorporate this gauge condition when solving the equations of motion for the gravitino and the 3-form. Because the gauge parameter θ_{MN} has 55 degrees of freedom, we can impose gauge conditions as follows:

1. For a given function $V(\Phi)$ and a symmetric tensor $\mathcal{T}_{\mu\nu}^{(\text{matter})}$, the equations

$$T_{\mu\nu}^{(A,\psi)} = \frac{m_{p(11)}^9}{m_p^2} \mathcal{T}_{\mu\nu}^{(\text{matter})} - m_{p(11)}^9 \left[\frac{V(\Phi)}{m_p^2} + \frac{1}{2} \left(\frac{R_{T_{q_1}^2}}{\Phi} + R_{T_{q_2}^2} + \frac{R_{T_{q_3}^2}}{\Phi^2} \right) \right] g_{\mu\nu}, \quad (2.12)$$

expect for $\mu = \nu = 0$ case, restrict 9 degrees of freedom.

2. Particular block diagonal configuration of 11D bulk metric (2.5) restricts 46 degrees of freedom

$$G_{a_i A} = 0, (A \neq a_i \text{ and } i = 1, 2, 3) \quad (2.13)$$

$$G_{\mu A} = 0, (A \neq \mu) \quad (2.14)$$

$$G_{10, A} = 0, (A \neq 10) \quad (2.15)$$

By considering the gauge conditions (2.12)(2.13)(2.14)(2.15), boundary conditions, and the equations of motion for the graviton G_{AB} , gravitino ψ_A and 3-form A_{MNP} collectively, we can determine the solutions for the graviton, gravitino, and 3-form. These solutions are expressed as functions of the scale factor $a(t)$ and the scale function $\Phi(x^\mu)$ by $\psi_A(a(t), \Phi(x^\mu), x^A)$ and $A_{MNP}(a(t), \Phi(x^\mu), x^A)$. To ensure the representation of spinors in four dimensions, the matrix Γ^M can be expressed as the tensor product among a 4×4 matrix $\hat{\gamma}^\mu$, three 2×2 matrices $\hat{\gamma}^{a_i}, i = 1, 2, 3$, and one 2×2 matrix $\hat{\gamma}^{10}$

$$\Gamma^\mu = \hat{\gamma}^\mu \otimes \mathbf{1}_2 \otimes \mathbf{1}_2 \otimes \mathbf{1}_2, \quad (2.16)$$

$$\Gamma^{a_1} = \hat{\gamma}^* \otimes \hat{\gamma}^{a_1} \otimes \mathbf{1}_2 \otimes \mathbf{1}_2, \quad (2.17)$$

$$\Gamma^{a_2} = \hat{\gamma}^* \otimes \hat{\gamma}_1^* \otimes \hat{\gamma}^{a_2} \otimes \mathbf{1}_2, \quad (2.18)$$

$$\Gamma^{a_3} = \hat{\gamma}^* \otimes \hat{\gamma}_1^* \otimes \hat{\gamma}_2^* \otimes \hat{\gamma}^{a_3}, \quad (2.19)$$

$$\Gamma^{10} = \hat{\gamma}^* \otimes \hat{\gamma}_1^* \otimes \hat{\gamma}_2^* \otimes \hat{\gamma}^{10}, \quad (2.20)$$

where $\hat{\gamma}^* = \hat{\gamma}^0 \hat{\gamma}^1 \hat{\gamma}^2 \hat{\gamma}^3, \hat{\gamma}_1^* = \hat{\gamma}^5 \hat{\gamma}^6, \hat{\gamma}_2^* = \hat{\gamma}^7 \hat{\gamma}^8$. Therefore, supersymmetry transformations introduce a 32-dimensional spinor gauge parameter denoted by $\epsilon_{\dot{\alpha}\dot{\alpha}_1\dot{\alpha}_2\dot{\alpha}_3}$, where $\dot{\alpha} = 1, 2, 3, 4$ and $\dot{\alpha}_i = 1, 2$ for $i = 1, 2, 3$ are the four-dimensional spinor indices and two-dimensional spinor indices respectively. Given a specific classical solution of metric, as defined in (2.5), we anticipate that its geometric configuration remains unaltered under the supersymmetry transformation outlined in (1.8)(1.9)(1.10). Consequently, this imposes certain gauge conditions, designated as

$$\delta g_{\mu\nu} = \delta g_{a_1 b_1} = \delta h_{a_2 b_2} = \delta \gamma_{a_3 b_3} = 0, \quad (2.21)$$

where the operator $\delta(\dots)$ is the variation respect to supersymmetry. The supersymmetric transformation law for the metric G_{AB} can be expressed as

$$\delta G_{AB} = \frac{1}{2} \eta_{ab} \bar{\epsilon} \left(\Gamma^a \psi_A e_B^b + e_A^a \Gamma^b \psi_B \right). \quad (2.22)$$

Additionally, its components are subject to gauge conditions (2.21), denoted by

$$\delta e_\mu^{a_0} = \frac{1}{2} \bar{\epsilon} \Gamma^{a_0} \psi_\mu = 0, a_0 = 1, 2, 3, 4 \quad (2.23)$$

$$\delta G_{a_1 b_1} = g_{a_1 b_1} \delta \Phi, \quad (2.24)$$

$$\delta G_{a_2 b_2} = 0, \quad (2.25)$$

$$\delta G_{a_3 b_3} = 2g_{a_3 b_3} \Phi \delta \Phi, \quad (2.26)$$

$$\delta G_{10,10} = -6\Phi^{-7} \delta \Phi. \quad (2.27)$$

The gauge condition (2.23) indicates the absence of gravitino, $\psi_\mu = 0$, in the four-dimensional spacetime. Consequently, supersymmetry of gravity in the four-dimensional spacetime is spontaneous breaking. As there is no strong observable signal from supersymmetric field theory in 4D, we postulate that Φ has no superpartner, thus assuming $\delta \Phi = 0$. Hence, the compactification methods spontaneously break all the $\mathcal{N} = 32$ supersymmetries present in the eleven-dimensional bulk, where the classical solutions of (2.23)(2.24)(2.25)(2.26)(2.27) are $\psi_M = 0$. The non-trivial solution to the gauge parameter ϵ can be derived from the equation

$$\nabla_M \epsilon + \frac{\sqrt{2}}{288} (\Gamma^{ABCD}{}_M - 8\Gamma^{BCD} \delta_M^A) F_{ABCD} \epsilon = 0. \quad (2.28)$$

Note that there are trivial solutions $\epsilon = f$, with constant Grassman numbers f , when the 3-form takes trivial solution $A_{MNP} = 0$. After imposing the gauge conditions (2.12)(2.13)(2.14) and (2.15)(2.21)(2.28), the only two remaining undetermined functions are the scale factor $a(t)$ and the scale function $\Phi(x^\mu)$ in (2.5).

In the four-dimensional spacetime, we anticipate that Einstein's gravity predominantly governs the laws of classical gravity as seen in (2.6). Moreover, gauge conditions (2.12) imply that it is possible to obtain a four-dimensional effective action, in which the solutions to the equations of motion are represented by $\Phi(x^\mu)$ and $a(t)$. While the equations of (2.12), except for $\mu = \nu = 0$, have already been utilized as gauge conditions to determine the solution of 3-form, these equations do not establish the relation between $\Phi(x^\mu)$ and $a(t)$. To clarify the relationship between $\Phi(x^\mu)$ and $a(t)$, we introduce an additional equation, which is the same as (2.12) but only for the case $\mu = \nu = 0$. Subsequently, the solution for $\mathcal{T}_{00}^{(\text{matter})}[\psi_A, A_{MNP}, a(t), \Phi(x^\mu), x^A]$ obtained from this equation uniquely defines the Lorentz-invariant symmetric tensor $\mathcal{T}_{\mu\nu}^{(\text{matter})}[\psi_A, A_{MNP}, a(t), \Phi(x^\mu), x^A]$, thereby imposing 9 gauge conditions specified in (2.12). These 9 gauge conditions, along with (2.13)(2.14)(2.15)(2.21)(2.28), constrain the solutions $\psi_A(a(t), \Phi(x^\mu), x^A)$ and $A_{MNP}(a(t), \Phi(x^\mu), x^A)$ in such a way that they determine $\mathcal{T}_{\mu\nu}^{(\text{matter})}[a(t), \Phi(x^\mu), x^A]$ uniquely. This tensor can be derived by the equations of motion for $\Phi(x^\mu)$ and $a(t)$ in four-dimensional spacetime. The

equations (2.12) for $\mu, \nu = 0, 1, 2, 3$ serves to define an effective energy-stress tensor $\mathcal{T}_{\mu\nu}$ as

$$\mathcal{T}_{\mu\nu} = \frac{23m_p^2}{2\Phi^2} \nabla_\mu \Phi \nabla_\nu \Phi + g_{\mu\nu} \left(-\frac{1}{2} \frac{23m_p^2}{2\Phi^2} \nabla_\rho \Phi \nabla^\rho \Phi - V(\Phi) \right) + \mathcal{T}_{\mu\nu}^{(\text{matter})}, \quad (2.29)$$

where m_p is the four-dimensional reduced Planck mass and $\mathcal{T}_{\mu\nu}^{(\text{matter})}$ is the stress tensor of some four-dimensional matter. Hence, in the four-dimensional spacetime, the equation of motion (2.7) gives Einstein's equation

$$\mathcal{G}_{\mu\nu} = \frac{1}{m_p^2} \mathcal{T}_{\mu\nu}, \quad (2.30)$$

and the associated effective action is given by

$$S_{\text{eff}} = \int d^4x \sqrt{-g_4} \left[\frac{m_p^2}{2} R_{(4)} - \frac{1}{2} \frac{23m_p^2}{2\Phi^2} \nabla_\mu \Phi \nabla^\mu \Phi - V(\Phi) + \mathcal{L}_{\text{matter}} \right], \quad (2.31)$$

where

$$\mathcal{T}_{\mu\nu}^{(\text{matter})} = -2 \frac{\delta \mathcal{L}_{\text{matter}}}{\delta g^{\mu\nu}} + g_{\mu\nu} \mathcal{L}_{\text{matter}} \quad (2.32)$$

The dynamics in the 4D spacetime may be influenced by the coordinates associated with the extra dimensions, as reflected in the energy stress tensor $\mathcal{T}_{\mu\nu}^{(\text{matter})}$. To address this problem, it is natural to introduce N coincident four-dimensional cosmological branes with Dirichlet boundary conditions, such as $x^{10} = c_0$ and $x^{a_i} = c_{a_i}$ for $i = 1, 2, 3$, where c_0 and c_{a_i} are constants. The gravitino field $\psi_{A\dot{\alpha}\dot{\alpha}_1\dot{\alpha}_2\dot{\alpha}_3}$ gives rise to 56 fermion fields $\psi_{i\dot{\alpha}\dot{\alpha}_1\dot{\alpha}_2\dot{\alpha}_3}$ for $i = 4, 5, \dots, 10$ on cosmological D3-branes, which contains the fermion fields in the Standard Model of Particle Physics. Additionally, there are 21 spin-1 boson fields $A_{\mu ij}$, 35 spin-0 scalar fields A_{ijk} , and 7 spin-0 tensor fields $A_{\mu\nu k}$ for $i, j, k = 4, 5, \dots, 10$. These fields contain ordinary matter, dark matter, and dark energy. Then, the lagrangian for matter on the cosmological D3-branes can encompass ordinary matter from the Standard Model of Particle Physics \mathcal{L}_{SM} , as well as dark matter \mathcal{L}_{DM} and dark energy \mathcal{L}_{DE} ,

$$\mathcal{L}_{\text{matter}}^{(\text{Dirichlet})} = \mathcal{L}_{\text{SM}} + \mathcal{L}_{\text{DM}} + \mathcal{L}_{\text{DE}}, \quad (2.33)$$

and the associated effective action of the cosmological brane is

$$S_{\text{brane}} = \int d^4x \sqrt{-g_4} \left[\frac{m_p^2}{2} R_{(4)} - \frac{1}{2} \frac{23m_p^2}{2\Phi^2} \nabla_\mu \Phi \nabla^\mu \Phi - V(\Phi) + \mathcal{L}_{\text{matter}}^{(\text{Dirichlet})} \right]. \quad (2.34)$$

This allows us to derive a cosmological inflation model based on a specified driven potential $V(\Phi)$, where the potential can be set as

$$V(\Phi) = -\frac{m_p^2}{2} \left(\frac{R_{T_{q_1}^2}}{\Phi} + R_{T_{q_2}^2} + \frac{R_{T_{q_3}^2}}{\Phi^2} \right). \quad (2.35)$$

As an example, concentrate on the trivial solutions to the equations of motion for gravitino ψ_A and 3-form A_{MNP} , where the solutions are $\psi_A = 0$ and $A_{MNP} = 0$. The physical

significance of these two solutions is that both the gravitino and the 3-form remain inactive during the entire inflationary period. Consequently, the energy-stress tensor simplifies to $T_{AB}^{(A,\psi)} = 0$, leaving the 11D Einstein-Hilbert action as the outcome for action (1.2). By adopting specific gauge conditions, where $T_{\mu\nu}^{(A,\psi)} = 0$ in equation (2.12), and imposing suitable Dirichlet boundary conditions on the cosmological D3-brane with definition of driven potential (2.35). we can simplify our calculations by setting $\mathcal{L}_{\text{matter}}^{(\text{Dirichlet})} = 0$. This means the four-dimensional matter fields on the branes are temporarily inactive during the inflation period.

Interestingly, one can deduce a suitable potential $V(\Phi)$ by integrating out the physics arising from various configurations of the extra dimensions within the 11D Einstein-Hilbert action

$$S_{EH} = \int d^{11}x \sqrt{-G_{11}} \frac{m_{p(11)}^9}{2} R_{(11)}. \quad (2.36)$$

Due to the block diagonal property of the 11D metric (2.5), the integral over the 11D bulk can be decomposed into multiple integrals over the 4D spacetime and the compactified extra dimensions, corresponding to each block of the 11D metric

$$\int d^{11}x \sqrt{-G_{11}}[\dots] = \int_{V_{\text{FRW}}} d^4x \sqrt{-g_4} \int_{T_{q_1}^2} d^2x \sqrt{g_2} \int_{T_{q_2}^2} d^2x \sqrt{h} \int_{T_{q_3}^2} d^2x \sqrt{\gamma} \int_{S^1} dx[\dots]. \quad (2.37)$$

After integrating the 11D Ricci scalar (2.6), the 11 dimensional Einstein-Hilbert action is now represented by

$$S_{EH} = \int d^4x \sqrt{-g_4} \left[\frac{m_p^2}{2} R_{(4)} - \frac{1}{2} \frac{23m_p^2}{2\Phi^2} \nabla_\mu \Phi \nabla^\mu \Phi + \frac{4\pi(1-q_1)m_p^2}{A_{T_{q_1}^2} \Phi} + \frac{4\pi(1-q_2)m_p^2}{A_{T_{q_2}^2}} + \frac{4\pi(1-q_3)m_p^2}{A_{T_{q_3}^2} \Phi^2} \right], \quad (2.38)$$

where we have defined the 4D reduced Planck mass m_p by utilizing the 11D reduced Planck mass $m_{p(11)}$, the areas of the three tori of standard sizes $A_{T_{q_i}^2}$, $i = 1, 2, 3$, and the standard radius of the eleventh dimension $\tilde{\mathcal{R}}_{10}$ as follows

$$\frac{2\pi \tilde{\mathcal{R}}_{10} A_{T_{q_1}^2} A_{T_{q_2}^2} A_{T_{q_3}^2} m_{p(11)}^9}{2} = \frac{m_p^2}{2}. \quad (2.39)$$

The last three terms in (2.38), which function as the cosmological inflation potential, originate from the Gauss-Bonnet theorem

$$\int_{T_{q_i}^2} R_{T_{q_i}^2} = 8\pi(1-q_i), \quad i = 1, 2, 3. \quad (2.40)$$

The second term in (2.38) is a kinetic term, which becomes more apparent through the redefinition of the field Φ as

$$\phi = \sqrt{\frac{23}{2}} m_p \ln \Phi, \quad (2.41)$$

$$\Phi = \exp\left(\sqrt{\frac{2}{23}} \frac{\phi}{m_p}\right). \quad (2.42)$$

Restrict $q_1 = 0, q_2 = 2, q_3 = 2$ and $2A_{T_{q_1}^2} = A_{T_{q_2}^2} = A_{T_{q_3}^2} = A_0$, we derive a single-field cosmological inflation action

$$S_{\text{brane}} = S_{EH} = \int d^4x \sqrt{-g_4} \left[\frac{m_p^2}{2} R_{(4)} - \frac{1}{2} \nabla_\mu \phi \nabla^\mu \phi - \frac{4\pi m_p^2}{A_0} \left(1 - e^{-\sqrt{\frac{2}{23}} \frac{\phi}{m_p}}\right)^2 \right], \quad (2.43)$$

where ϕ is the inflaton. It is important to note that assigning specific values to the genus q_1, q_2, q_3 and the areas $A_{T_{q_1}^2}, A_{T_{q_2}^2}, A_{T_{q_3}^2}$ essentially imposes boundary conditions on the metric G_{AB} . The trick presented in (2.37)(2.38) illustrates one potential method of defining the driven potential $V(\Phi(\phi))$ as

$$V(\Phi(\phi)) = \frac{4\pi m_p^2}{A_0} \left(1 - e^{-\sqrt{\frac{2}{23}} \frac{\phi}{m_p}}\right)^2, \quad (2.44)$$

When combining equations (2.35)(2.44), the Dirichlet boundary conditions for the cosmological D3-brane can be derived from equations

$$R_{T_{q_1}^2}[c_{a_1}] = \frac{16\pi}{A_0}, \quad R_{T_{q_2}^2}[c_{a_2}] = -\frac{8\pi}{A_0}, \quad R_{T_{q_3}^2}[c_{a_3}] = -\frac{8\pi}{A_0}. \quad (2.45)$$

Therefore, the technique presented in equations (2.37) and (2.38) is equivalent to imposing specific Dirichlet boundary conditions. In this approach, the relative stretched sizes of the three tori are represented as $(\Phi, 1, \Phi^2)$. While the specific construction of extra dimensions is given in (2.3), it should be noted that other configurations of extra dimensions are also conceivable. Before conducting numerical analysis, we intend to explore scenarios involving various sizes combinations of the three tori.

2.2 General Case

In this subsection, we will explore the scenario involving general relative stretched sizes of the three tori, specifically represented as $(\Phi^{2s_1}, \Phi^{2s_2}, \Phi^{2s_3})$ with parameters $s_1, s_2, s_3 \geq 0$. The corresponding metrics for these three tori are provided by

$$ds_{2(1)}^2 = \Phi(x^\mu)^{2s_1} g_{a_1 b_1}(x^{a_1}) dx^{a_1} dx^{b_1}, \quad (2.46)$$

$$ds_{2(2)}^2 = \Phi(x^\mu)^{2s_2} h_{a_2 b_2}(x^{a_2}) dx^{a_2} dx^{b_2}, \quad (2.47)$$

$$ds_{2(3)}^2 = \Phi(x^\mu)^{2s_3} \gamma_{a_3 b_3}(x^{a_3}) dx^{a_3} dx^{b_3}. \quad (2.48)$$

Since we anticipate the presence of Einstein's Gravity in a four-dimensional spacetime, the 11th dimension is compacted into a circular shape as

$$ds_1^2 = \Phi^{-2(s_1+s_2+s_3)} dx^{10} dx^{10}. \quad (2.49)$$

Its radius is specifically defined to prevent any coupling between Φ and the 4D Ricci scalar $R_{(4)}$. The Ricci scalar in 11 dimensions, after dimensional reduction, is expressed as follows:

$$R_{(11)} = R_{(4)} + \frac{R_{T_{q_1}^2}}{\Phi^{2s_1}} + \frac{R_{T_{q_2}^2}}{\Phi^{2s_2}} + \frac{R_{T_{q_3}^2}}{\Phi^{2s_3}} - \frac{\lambda}{\Phi^2} g^{\mu\nu} \nabla_\mu \Phi \nabla_\nu \Phi, \quad (2.50)$$

where

$$\lambda = 6 \sum_i s_i^2 + 8 \sum_{i>j} s_i s_j, \quad (2.51)$$

with the indices $i, j = 1, 2, 3$. Analogous to the special case presented in Subsection 2.1, the four-dimensional effective action for cosmological D3-branes can be derived through the imposition of appropriate gauge conditions and boundary conditions. Unsurprisingly, the effective action can be written as

$$S_{\text{brane}} = \int d^4x \sqrt{-g_4} \left[\frac{m_p^2}{2} R_{(4)} - \frac{1}{2} \frac{\lambda m_p^2}{\Phi^2} \nabla_\mu \Phi \nabla^\mu \Phi - U(\Phi) + \mathcal{L}_{\text{matter}}^{(\text{Dirichlet})} \right]. \quad (2.52)$$

By utilizing the technique demonstrated in equations (2.37) and (2.38), which imposes specific Dirichlet boundary conditions for the cosmological D3-brane, the potential can be defined as follows:

$$U(\Phi) = - \left(\frac{4\pi(1-q_1)m_p^2}{A_{T_{q_1}^2} \Phi^{2s_1}} + \frac{4\pi(1-q_2)m_p^2}{A_{T_{q_2}^2} \Phi^{2s_2}} + \frac{4\pi(1-q_3)m_p^2}{A_{T_{q_3}^2} \Phi^{2s_3}} \right). \quad (2.53)$$

The corresponding Dirichlet boundary conditions can be derived from the equations

$$R_{T_{q_i}^2} [c_{a_i}] = \frac{8\pi(1-q_i)}{A_{T_{q_i}^2}}, \quad i = 1, 2, 3. \quad (2.54)$$

The redefinition of the field is as stated below:

$$\phi = \sqrt{\lambda} m_p \ln |\Phi|, \quad (2.55)$$

$$\Phi = \pm \exp \left(\sqrt{\lambda^{-1}} \frac{\phi}{m_p} \right). \quad (2.56)$$

To compare our findings with the discussions presented in subsection 2.1, the equations written as $2A_{T_{q_1}^2} = A_{T_{q_2}^2} = A_{T_{q_3}^2} = A_0$ remain applicable. As the inflaton ϕ rolls down to a true vacuum, we can establish conditions $U(\Phi_{\text{min}}) = 0, U'(\Phi_{\text{min}}) = 0$ on the potential. The solution is given by

$$q_1 = \frac{\Phi_{\text{min}}^{2s_1-2s_3} (s_2 - s_3) (1 - q_3)}{2(s_2 - s_1)} + 1, \quad (2.57)$$

$$q_2 = \frac{\Phi_{\text{min}}^{2s_2-2s_3} (s_1 - s_3) (1 - q_3)}{(s_1 - s_2)} + 1. \quad (2.58)$$

To ensure the genus of the three tori $q_1, q_2, q_3 \in \mathbb{N}$, we can set $\Phi_{min} = -\frac{1}{n}, s_1 = \frac{1}{2}, s_2 = 0$, then

$$q_1 = 1 + n^{2s_3-1}(q_3 - 1)s_3, \quad (2.59)$$

$$q_2 = n^{2s_3} [1 + n^{-2s_3} + 2s_3(q_3 - 1) - q_3], \quad (2.60)$$

where $q_3 \geq 2, s_3 > 0, s_3 \in \mathbb{Z}$. In this case

$$U(\Phi) = \frac{4\pi m_p^2 (q_3 - 1) \Phi^{-2s_3-1}}{A_0 n} \left[n\Phi + (n\Phi)^{2s_3} (2s_3 + (2s_3 - 1)n\Phi) \right], \quad (2.61)$$

with $\Phi = -e^{\sqrt{\lambda^{-1}}\phi/m_p}$ and $n, q_3 \in \mathbb{N}$. The pictures of this potential are shown in Figure 1. It can be seen from the listed potential functions that the lowest point of potential energy is located at $U(\Phi_{min}) = 0$. When $\phi \gg 1$, the potential energy tends to be a flat potential, which has the characteristics of the potential function of the Starobinsky model. An important property of the potential $U(\Phi)$ is that the spectral index n_s and tensor-to-scalar ratio r are independent of the factor $(q_3 - 1)$ and parameter n . Therefore, the only constraint imposed on the third torus is $q_3 \geq 2$. For more details, please refer to the [Appendix](#). Moreover, we rediscover the special case in Subsection 2.1 by choosing $\Phi_{min} = 1, s_1 = \frac{1}{2}, s_2 = 0, s_3 = 1, q_1 = 0, q_2 = q_3 = 2$.

3 Cosmological Inflation

3.1 Numerical Results

In this section, we aim to numerically compute the inflation parameters corresponding to the inflation models discussed in Section 2. The inflaton ϕ is clearly associated with the size function $\Phi(x^\mu)$ of the extra dimensions. Therefore, the vibrations of the inflaton ϕ can be considered as the conformal vibrations of the extra dimensions.

The slow-roll parameters ϵ and η , along with the spectral index n_s and tensor-to-scalar ratio r , are determined as follows:

$$\epsilon(\phi) = \frac{m_p^2}{2} \left(\frac{U_\phi}{U} \right)^2, \quad (3.1)$$

$$\eta(\phi) = m_p^2 \left(\frac{U_{\phi\phi}}{U} \right), \quad (3.2)$$

$$n_s = 1 - 6\epsilon(\phi_i) + 2\eta(\phi_i), \quad (3.3)$$

$$r = 16\epsilon(\phi_i). \quad (3.4)$$

The e-folding number is

$$N = -\frac{1}{m_p^2} \int_{\phi_i}^{\phi_e} \frac{U}{U_\phi} d\phi, \quad (3.5)$$

where ϕ_i represents the initial value of the inflaton and ϕ_e represents its end value. The numerical results of the general case with potential (2.61) are shown in Figure 2. The lowest




Figure 1: Pictures for inflation potential. The first figure depicts the potential of the special case given by equation (2.43), where $(q_1, q_2, q_3) = (0, 2, 2)$. The subsequent figures illustrate potentials $U(\Phi)$ in equation (2.61) for varying parameter sets of (q_1, q_2, q_3, n) . Specifically, the second picture in the first row corresponds to the parameter set $(q_1, q_2, q_3, n) = (3, 3, 3, 1)$. In the second row, the first picture represents the potential for $(q_1, q_2, q_3, n) = (5, 9, 3, 2)$, while the last picture corresponds to $(q_1, q_2, q_3, n) = (7, 19, 3, 3)$.

three lines in Figure 2 correspond to the cases of $s_3 = 1, 2, 3$, respectively. They are consistent with the stronger $n_s - r$ constraints by Planck TT,TE,EE+lowE+lensing+BK18+BAO (blue region). With the increasing s_3 , the convergence of $n_s - r$ happens inside the green region of weak $n_s - r$ constraints. As a result, the inflation models from the compactification methods $\mathcal{M}^{3,1} \times T_{q_1}^2 \times T_{q_2}^2 \times T_{q_3}^2 \times S^1$ can be in good accordance with the latest observations. The optimal value for s_3 is $s_3 = 1$, as its corresponding $n_s - r$ line approaches the center of the observational constraints, which is the same as the result of the special case discussed in Subsection 2.1.

For the special case (2.43), the inflation exits as $\epsilon(\phi_e) = 1$, $\phi_e = 1.182m_p$. For e-folding $N = 60$, the solutions are $(\phi_i, \epsilon, \eta, n_s, r) = (8.986m_p, 0.0010, -0.012, 0.9695, 0.016)$. For e-folding $N = 50$, $(\phi_i, \epsilon, \eta, n_s, r) = (8.503m_p, 0.0013, -0.014, 0.9637, 0.022)$. These results match the Planck 2018 data [2]. Taking into account the observational constraint imposed on the power spectrum [59]

$$P_s = \frac{U(\phi_i)}{24\pi^2 m_p^4 \epsilon(\phi_i)} \approx 2.1 \times 10^{-9}. \quad (3.6)$$

When e-folding $N = 60$, the standard areas of the three extra-dimensional tori can be




Figure 2: The $n_s - r$ constraints for potential (2.61). The green contours are from Planck 2018 data [2]. The blue contours are from 2021 BICEP/Keck data [1]. Here, we choose $s_3 = 1, 2, 3, 4, 7, 10, \dots, 37, 40$. The straight lines of varying colors represent distinct values of s_3 . The lowest red straight line represents $s_3 = 1$. As s_3 increases, the corresponding $n_s - r$ line progressively shifts beyond the blue observational region (See the yellow region.). However, as s_3 continues to grow, these lines tend to cluster around the purple line at the top.

obtained as follows:

$$A_0 = \frac{4\pi}{\zeta} l_p^2 = 1.43 \times 10^{-58} [\text{m}]^2, \quad (3.7)$$

where $\zeta = 5.79 \times 10^{-10}$ is coupling constant of potential

$$U(\phi) = \zeta m_p^4 \left(1 - e^{-\sqrt{\frac{3}{23}} \frac{\phi}{m_p}} \right)^2, \quad (3.8)$$

and l_p is the reduced Planck length. The symbol [m] is used to represent the unit of meters. To approximate the size of the extra dimensions, we can assume that the shape of the extra dimensions $T_{q_1}^2$ is roughly spherical. Consequently, the radius of this sphere would be

$$\mathcal{R} = \sqrt{\frac{A_0}{8\pi}} = 29387 l_p = 2.38 \times 10^{-30} [\text{m}]. \quad (3.9)$$




Figure 3: The dynamics of the inflaton and scale function for e-folding $N = 60$. The first figure depicts the scalar field rolling down to 0 at $N = 60$, while the second figure shows the scale function converging to 1. During inflation, the dynamics of the scale function Φ exhibit a decreasing trend.

The range of the scalar field $\phi \in (\phi_i, \phi_e)$, $\phi_i \in (8.5025m_p, 8.986m_p)$ determines the range of the scale function of the extra dimensions during inflation, as specified below:

$$1.417 \leq \Phi \leq 12.271 \quad (\text{for } N = 50) \quad (3.10)$$

$$1.417 \leq \Phi \leq 14.153 \quad (\text{for } N = 60) \quad (3.11)$$

It shows that the scale of the extra dimensions is considerably small at order $O(10^5 l_p)$. In our compactification methods, the eleventh dimension is compactified into a circle, indicating that the theory resulting from compactification is Type IIA string theory in 10 dimensions. Subsequently, the string length l_s relates to the radius of the eleventh dimension \mathcal{R}_{10} as follows [60]:

$$l_s = \frac{l_{p(11)}}{\sqrt{\frac{\mathcal{R}_{10}}{l_{p(11)}}}}. \quad (3.12)$$

In today's universe, where $\Phi = 1$, we postulate that the string length is represented by $l_s = \xi l_{p(11)}$ and its corresponding eleventh dimension radius is given by $\tilde{\mathcal{R}}_{10} = \xi^{-2} l_{p(11)}$. The 11D Newtonian gravitational constant $G_N^{(11)}$ and the 4D Newtonian gravitational constant G are related through the volume of the extra dimensions (2.39) as follows:

$$G_N^{(11)} = \pi \xi^{-2} l_{p(11)} A_0^3 G. \quad (3.13)$$

Consequently, the relationship between the 4D reduced Planck length l_p and the 11D reduced Planck length $l_{p(11)}$ is expressed by equations

$$l_p = \frac{l_{p(11)}^{\frac{9}{2}}}{(\pi \xi^{-2} l_{p(11)} A_0^3)^{\frac{1}{2}}}, \quad (3.14)$$

$$l_{p(11)} = (\pi A_0^3 \xi^{-2})^{\frac{1}{8}} l_p^{\frac{1}{4}} = l_p \left[\frac{(4\pi)^4}{4\zeta^3 \xi^2} \right]^{\frac{1}{8}} = 8676.7 \xi^{-\frac{1}{4}} l_p. \quad (3.15)$$

However, during the inflationary period, the eleventh dimension experiences a scaling factor of Φ^{-3} . Hence, for e-folding $N = 60$, the radius adjusts to $\mathcal{R}_{10} = \xi^{-2}\Phi^{-3}l_{p(11)} \in \xi^{-2}[0.000352l_{p(11)}, 0.351l_{p(11)}]$. The string length during this period can be determined using equation (3.12) as follows:

$$l_s \in \xi [1.7l_{p(11)}, 53.2l_{p(11)}]. \quad (3.16)$$

The standard tori extra dimensions have a size scale of $\mathcal{R} = 29387l_p = 3.39\xi^{\frac{1}{4}}l_{p(11)}$. Assuming the loop-order quantum corrections of eleven-dimensional supergravity [61] can be neglected due to their small coupling constant $\alpha' \sim O(l_s^2)$, the string length and the size scale of the tori must satisfy the condition $\frac{l_s}{\mathcal{R}} \ll 1$. This limitation imposes a constraint on the parameter $\xi \ll 0.025$. Therefore, we anticipate that the current string length meets the criterion $l_s = 8676.7\xi^{\frac{3}{4}}l_p \ll 551.9l_p$.

The dynamics of the inflaton and the scale function are shown in figure 3. It demonstrates that the three tori contract conformally, while the eleventh dimension expands. The scale function oscillates around the point where $\Phi(x^\mu)$ is equal to 1 after the process of inflation.

3.2 Exit Inflation

There exist additional matter terms within the four-dimensional effective action of 11D Supergravity (2.31). Because the 11D metric G_{AB} couples to gravitino ψ_A and 3-form A_{MNP} , the inflaton ϕ , associated to the size function Φ of extra dimensions, will couple to some components of ψ_A and A_{MNP} after compactification. When the inflaton ϕ begins to decay into the gravitino and the 3-form, the inflation exits. In this period, we keep the gauge condition $\delta g_{\mu\nu} = 0$ and release $\delta g_{a_1 b_1}, \delta h_{a_2 b_2}, \delta \gamma_{a_3 b_3} \neq 0$. This decay process gives rise to a non-zero $\mathcal{L}_{\text{matter}}^{(\text{Dirichlet})}$ term in the four-dimensional effective action, as seen in equation (2.34). For example, matter in the universe can be generated by introducing the interactions between the inflaton and the Higgs field H , expressed as $\xi\phi^2 H^2$, into the Lagrangian $\mathcal{L}_{\text{matter}}^{(\text{Dirichlet})}$.

4 Dark Sector from Extra Dimensions

In this section, we concentrate on the quantum effects from the inflaton in the four-dimensional spacetime specifically for the case presented in Subsection 2.1. As shown in Equation (3.9), the extra dimensions exist at the sub-Planck scale $O(10^5 l_p)$, and their conformal vibrations can be interpreted as the vibrations of the inflaton ϕ . Referring to Figure 3, it becomes evident that the inflaton field arrives at a significantly small value, namely $\phi \ll m_p$, in today's universe. Hence, the potential given in (2.44) can be expanded up to the second order as

$$V(\phi) \approx \frac{1}{2}m_\phi^2\phi^2, \quad (4.1)$$

where

$$m_\phi = \sqrt{\frac{4\zeta}{23}}m_p = 10^{-5}m_p = 2.4 \times 10^{10} \frac{\text{TeV}}{c^2}. \quad (4.2)$$

The rest mass of the inflaton is so heavy that detecting it exceeds the capabilities of current accelerators. The energy scale at which the inflaton appears is approximately $\Lambda_\phi = m_\phi c^2 \approx 2.4 \times 10^{10} \text{TeV}$. This energy scale is significantly higher than those achievable by the Large Hadron Collider (LHC), making it challenging to directly observe the inflaton in accelerators. However, gravity is a long-range force, and Newton's law with the modification by the inflaton can be verified through astronomical observations.

Following the second quantization method, the inflaton field emerges as a massive scalar quantum field. The tree-level propagator for graviton with momentum $k^\mu = (k_0, \mathbf{k})$ is given by [62]

$$\Pi_{\mu\nu\rho\sigma}(x - x') = \int \frac{dk_0}{2\pi} \frac{d^3k}{(2\pi)^3} \frac{-iP_{\mu\nu\rho\sigma}}{k_0^2 - |\mathbf{k}|^2 + i\varepsilon} e^{-ik_0(t-t')} e^{i\mathbf{k}\cdot(\mathbf{x}-\mathbf{x}')}, \quad (4.3)$$

where

$$P_{\mu\nu\rho\sigma} = \frac{1}{2} (\eta_{\mu\rho}\eta_{\nu\sigma} + \eta_{\mu\sigma}\eta_{\nu\rho} - \eta_{\mu\nu}\eta_{\rho\sigma}). \quad (4.4)$$

The metric is expanded to first order using a perturbation tensor $h_{\mu\nu}$ as follows:

$$g^{\mu\nu} = \eta^{\mu\nu} + \frac{2}{m_p} h^{\mu\nu}, \quad (4.5)$$

where the Minkowski metric $\eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1)$ in the local reference frames. Consider the action (2.43) along with the matter term $\mathcal{L}_{\text{matter}}^{(\text{Dirichlet})}$, the energy stress tensor can be represented as

$$\mathcal{T}^{\mu\nu} = \mathcal{T}_\phi^{\mu\nu} + \mathcal{T}_{(\text{matter})}^{\mu\nu}, \quad (4.6)$$

where the energy stress tensor of the inflaton $\mathcal{T}_\phi^{\mu\nu}$, as well as the density ρ and the pressure p , are defined by

$$\mathcal{T}_\phi^{\mu\nu} = \partial^\mu \phi \partial^\nu \phi - \frac{1}{2} \eta^{\mu\nu} \partial^\alpha \phi \partial_\alpha \phi - \eta^{\mu\nu} V(\phi), \quad (4.7)$$

$$\rho = \frac{1}{2} \dot{\phi}^2 + \frac{1}{2} \partial_i \phi \partial^i \phi + V(\phi), \quad (4.8)$$

$$p = \frac{1}{2} \dot{\phi}^2 - \frac{1}{2} \partial_i \phi \partial^i \phi - V(\phi). \quad (4.9)$$

The inflaton fields can now be regarded as a form of dark energy for $\rho + 3p < 0$, or cold dark matter for $p/\rho \approx 0$, pervading the entire spacetime. In classical mechanics, when considering two massive point particles with masses M_1 and M_2 , Newton's gravity law has modification due to the interactions from the inflaton. The four-point tree-level scattering amplitude for the matter-graviton-matter process can be expressed as the effective action of the second order [62]

$$iS^{(2)} = \frac{1}{m_p^2} \int d^4x d^4x' \mathcal{T}_1^{\mu\nu}(x) \Pi_{\mu\nu\rho\sigma}(x - x') \mathcal{T}_2^{\rho\sigma}(x'), \quad (4.10)$$

where

$$\mathcal{T}_{(\text{matter}1)}^{\mu\nu} = M_1 u_1^\mu u_1^\nu \delta^3(\mathbf{x} - \mathbf{q}_1(t)), \quad (4.11)$$

$$\mathcal{T}_{(\text{matter}2)}^{\mu\nu} = M_2 u_2^\mu u_2^\nu \delta^3(\mathbf{x}' - \mathbf{q}_2(t)). \quad (4.12)$$

The four-vector $u^\mu = (1 - \mathbf{v}^2)^{-\frac{1}{2}}(1, \mathbf{v}) = I^\mu + O(\mathbf{v})$, where $I^\mu = (1, 0, 0, 0)$. This four-point tree-level scattering amplitude contains both the Newton potential part between M_1 and M_2 , denoted as [62]

$$iS_{(\text{matter})}^{(2)} = \frac{1}{m_p^2} \int d^4x d^4x' \mathcal{T}_{(\text{matter}1)}^{\mu\nu}(x) \Pi_{\mu\nu\rho\sigma}(x - x') \mathcal{T}_{(\text{matter}2)}^{\rho\sigma}(x'), \quad (4.13)$$

and the self-gravity interaction term of inflation, denoted as

$$iS_\phi^{(2)} = \frac{1}{m_p^2} \int d^4x d^4x' \mathcal{T}_\phi^{\mu\nu}(x) \Pi_{\mu\nu\rho\sigma}(x - x') \mathcal{T}_\phi^{\rho\sigma}(x'), \quad (4.14)$$

The remaining two terms in (4.10) are the modification for Newton's gravity law between M_1 and M_2

$$\begin{aligned} i \left(S^{(2)} - S_{(\text{matter})}^{(2)} - S_\phi^{(2)} \right) = & i \frac{M_1}{m_p^2} \int dt \int \frac{d^3\mathbf{k}}{(2\pi)^3} \frac{1}{|\mathbf{k}|^2 - i\varepsilon} \int d^3\mathbf{x} F(t, \mathbf{x}) e^{i\mathbf{k}\cdot(\mathbf{q}_1(t) - \mathbf{x})} \\ & + i \frac{M_2}{m_p^2} \int dt \int \frac{d^3\mathbf{k}}{(2\pi)^3} \frac{1}{|\mathbf{k}|^2 - i\varepsilon} \int d^3\mathbf{x}' F(t, \mathbf{x}') e^{i\mathbf{k}\cdot(\mathbf{x}' - \mathbf{q}_2(t))} \\ & + O(\mathbf{v}_1, \mathbf{v}_2), \end{aligned} \quad (4.15)$$

where

$$F(t, \mathbf{x}) = \left(\frac{\partial\phi(t, \mathbf{x})}{\partial t} \right)^2 - V(\phi(t, \mathbf{x})), \quad (4.16)$$

and we have taken $k_0 \sim \frac{v}{r} \rightarrow 0$ for the massless graviton in $\mathbf{v} \rightarrow 0$ limit. The modified Newton potential is now given by

$$\begin{aligned} W(\mathbf{q}_1(t), \mathbf{q}_2(t)) = & - \frac{M_1 M_2}{8\pi m_p^2} \frac{1}{|\mathbf{q}_1(t) - \mathbf{q}_2(t)|} - \frac{M_1}{m_p^2} \int \frac{d^3\mathbf{k}}{(2\pi)^3} \frac{1}{|\mathbf{k}|^2 - i\varepsilon} \int d^3\mathbf{x} F(t, \mathbf{x}) e^{i\mathbf{k}\cdot(\mathbf{q}_1(t) - \mathbf{x})} \\ & - \frac{M_2}{m_p^2} \int \frac{d^3\mathbf{k}}{(2\pi)^3} \frac{1}{|\mathbf{k}|^2 - i\varepsilon} \int d^3\mathbf{x}' F(t, \mathbf{x}') e^{i\mathbf{k}\cdot(\mathbf{x}' - \mathbf{q}_2(t))} + O(\mathbf{v}_1, \mathbf{v}_2). \end{aligned} \quad (4.17)$$

After doing the calculation on the integrals with respect to momentum \mathbf{k} , the modified Newton potential can be expressed as

$$\begin{aligned} W(\mathbf{q}_1(t), \mathbf{q}_2(t)) = & - \frac{M_1 M_2}{8\pi m_p^2} \frac{1}{|\mathbf{q}_1(t) - \mathbf{q}_2(t)|} - \frac{M_2}{4\pi m_p^2} \int d^3\mathbf{x} \frac{F(t, \mathbf{x})}{|\mathbf{q}_2(t) - \mathbf{x}|} \\ & - \frac{M_1}{4\pi m_p^2} \int d^3\mathbf{x}' \frac{F(t, \mathbf{x}')}{|\mathbf{q}_1(t) - \mathbf{x}'|}, \end{aligned} \quad (4.18)$$

The equation of motion for inflaton $\phi(t, \mathbf{x})$ is given by $\nabla_\mu T_\phi^{\mu\nu}(x) = 0$ in the FRW metric, which is

$$\ddot{\phi}(t, \mathbf{x}) + 3H\dot{\phi}(t, \mathbf{x}) - \frac{1}{a^2(t)}\partial_i\partial^i\phi(t, \mathbf{x}) + m_\phi^2\phi(t, \mathbf{x}) = 0, \quad (4.19)$$

where $H = \frac{\dot{a}(t)}{a(t)}$ is the Hubble constant. As we know, the current universe enters a de Sitter phase since the dark energy dominates the cosmic evolution, therefore H is almost a constant. When the inflaton field $\phi(t, \mathbf{x})$ has spherical symmetry, there is $\partial_i\partial^i\phi(t, r) = \frac{1}{r^2}\partial_r(r^2\partial_r\phi(t, r))$. Solving the equation of motion (4.19) yields the expression for $F(t, \mathbf{x})$, where the FRW metric goes to the Minkowski spacetime as $a(t) \simeq 1$.

4.1 Dark Matter

We consider the solution in time-splitting form $\phi(t, \mathbf{x}) = \phi_1(t)\phi_2(\mathbf{x})$. Assuming that the spatial part $\phi_2(\mathbf{x})$ has spherical symmetry, we can deduce the following equations:

$$\ddot{\phi}_1(t) + 3H\dot{\phi}_1(t) + m_\phi^2\phi_1(t) = 0, \quad (4.20)$$

$$\frac{1}{r^2}\partial_r(r^2\partial_r\phi_2(r)) = 0. \quad (4.21)$$

Note that the Hubble constant is approximately $H \approx 6.1 \times 10^{-61}m_p$. Therefore, there always exists a relation $m_\phi = 10^{-5}m_p \gg H$. In the limit $m_\phi \gg H$ and $a(t) \approx 1$ for today's universe, the solutions to the equations of motion, given by (4.21) and (4.20), can be described by

$$\phi_1(t) \approx 2\phi_0 a^{-\frac{3}{2}} \cos(m_\phi t), \quad (4.22)$$

$$\phi_2(r) = C_1 + \frac{C_2}{r}, \quad (4.23)$$

where ϕ_0, C_1, C_2 are constants and $m_\phi = \sqrt{\frac{4\zeta}{23}}$ in Planck unit. The time average of the function $F(t, \mathbf{x})$ can now be reformulated as

$$\langle F(t, \mathbf{x}) \rangle_t = \frac{m_\phi}{2\pi} \int_0^{\frac{2\pi}{m_\phi}} \left(\dot{\phi}^2(t, r) - \frac{1}{2}m_\phi^2\phi^2(t, r) \right) dt = \phi_2^2(r)a^{-3}m_\phi^2\phi_0^2. \quad (4.24)$$

The function $F(t, \mathbf{x})$ with a background evolution at the time-averaged level is given by

$$\langle F(t, \mathbf{x}) \rangle_t = \frac{1}{2}\phi_2^2(r)\overline{\rho_{DM}(t)}, \quad (4.25)$$

where $\overline{\rho_{DM}(t)} = 2a^{-3}m_\phi^2\phi_0^2 = 3m_p^2H^2 \times \Omega_{DM}$, here Ω_{DM} denotes the fraction parameter of dark matter. The time evolution of $\overline{\rho_{DM}(t)}$ can tell the coldness of the dark matter, i.e. its equation of state $w_{DM} \simeq 0$, which has been already accepted in the standard cosmological model.

Now, setting $\mathbf{q}_1(t) = 0$ and locating the center of the dark matter distribution at the origin $\mathbf{q}_1(t)$, we follow the method in Ref. [63] to calculate Eq. (4.18). Hence Newton's law of gravity for M_2 is given as follows

$$F_2(R) = -G\frac{M_1M_2}{R^2} - 4\pi GM_2\overline{\rho_{DM}(t)} \left(\frac{C_1^2}{3}R + \frac{C_2^2}{R} + C_1C_2 \right), \quad (4.26)$$

where we define $R = |\mathbf{q}_2(t)|$. Here $F_2(R)$ is the gravitational force exerted on M_2 and G is the Newtonian gravitational constant. The modified portion of Newton's gravity law describes a spherically symmetric distribution of the inflaton, which can be interpreted as a form of dark matter located around M_1 . The constants C_1, C_2 can be determined through astronomical observations at the galactic scale. If $C_1 = 0$, the flat rotation curves can be realized because the speed of object M_2 becomes $v_2^2 \equiv -F_2 R/M_2 \simeq GK$ at a very large R with $K = 4\pi C_2^2 \overline{\rho_{DM}(t)}$.

Note that we solely focus on the leading order contribution in the limit $m_\phi \gg H$. Assuming $m_\phi^2 = m_1^2 + m_2^2$, and considering the limits $m_1 \gg H$ and $m_1 \gg m_2$ for both the leading and sub-leading contributions, the equations for $\phi_1(t)$ and $\phi_2(r)$ can be described by equations

$$\ddot{\phi}_1(t) + 3H\dot{\phi}_1(t) + m_1^2\phi_1(t) = 0, \quad (4.27)$$

$$-\frac{1}{r^2}\partial_r\left(r^2\frac{\partial\phi_2(r)}{\partial r}\right) + a^2m_2^2\phi_2(r) = 0. \quad (4.28)$$

The solutions to these two equations are provided in

$$\phi_1(t) \approx 2\phi_0 a^{-\frac{3}{2}} \cos(m_1 t), \quad (4.29)$$

$$\phi_2(r) = C\frac{e^{-am_2 r}}{r} + D\frac{e^{am_2 r}}{r}, \quad (4.30)$$

where ϕ_0, C is constant and $D = 0$ since we should cancel the blow-up term containing $e^{am_2 r}$ when r goes to infinity. Under these circumstances, the function $F(t, \mathbf{x})$ of the inflaton retains the form given in (4.25), but with m_ϕ replaced by m_1 and $\phi_2(r)$ as specified in (4.30). Therefore, the modified Newton potential is still given by (4.18) and the corresponding Newton's gravity law for M_2 is given by

$$F_2(R) = -\frac{GM_1 M_2}{R^2} - \frac{2\pi GM_2 \overline{\rho_{DM}(t)} C^2 (1 - e^{-2am_2 R})}{am_2 R^2}. \quad (4.31)$$

The characteristic length of the Yukawa term, given by $\lambda = \frac{\hbar}{2am_2 c}$, constrains the value of m_2 based on experimental results on Yukawa gravitational potential [64, 65] and observation results on galaxies. Considering m_2 correction, when $m_2 R \gg 1$, the velocity decays in the manner of $v_2^2 \propto R^{-1}$ which reflects the original Newton's gravity law still holds at an extremely large distance but additional mass from dark matter to M_1 should be included as the central gravitational source. When $m_2 R \ll 1$, Eq. (4.26) in the case of $C_1 = 0$ is recovered, and the velocity of M_2 is directly expressed by equation

$$v_2^2 \approx \frac{GM_1}{R} + GK, \quad (4.32)$$

$$K = 4\pi C^2 \overline{\rho_{DM}(t)}, \quad (4.33)$$

where R can be determined using Newtonian acceleration a_N

$$R = \sqrt{\frac{GM_1}{a_N}}. \quad (4.34)$$

Additionally, the acceleration of M_2 is provided by

$$a_2 = \frac{v_2^2}{R} = a_N \left(1 + \sqrt{\frac{a_0}{a_N}} \right), \quad (4.35)$$

where the characteristic acceleration is defined as

$$a_0 = \frac{K^2 G}{M_1}. \quad (4.36)$$

Equation (4.35) recovers the formulation of the MOND theory [43–46]. Then, we find the Tully-Fisher relation from (4.32)(4.36) as $R \rightarrow \infty$

$$v_2^4 = GM_1 a_0. \quad (4.37)$$

These are two of the ways in which extra dimensions affect Newton’s gravity law. In this scenario, the extra dimensions are viewed as a spherically symmetric distribution of dark matter. Astronomical observations, as referenced in [66], reveal that the characteristic acceleration is of the order of $a_0 \sim 10^{-8}[\text{cm/s}^2]$ and the radii of galaxies are approximately $R_{\text{max}} \sim 10^2[\text{kpc}] \approx 3.8 \times 10^{55} l_p$. Consequently, the characteristic mass becomes $m_{2c} = 1/R_{\text{max}} \approx 2.6 \times 10^{-56} m_p$. Therefore, we observe the relations $m_1 \gg m_{2c} \gg m_2$. The modified portion of Newton’s gravity law is associated with both the mass of the inflaton m_ϕ and the strength of the inflaton field

$$C^2 = \frac{\sqrt{\frac{a_0 M_1}{G}}}{4\pi \rho_{DM}(t)}. \quad (4.38)$$

It is worth noting that certain galaxies, as observed in astronomy [67–71], lack dark matter. An explanation for this phenomenon could be that their characteristic acceleration a_0 and mass M_1 are small enough to result in $C^2 \sim 0$. In such galaxies, the inflaton distribution appears to be relatively uniform and sparse so that the scale function of extra dimensions $\Phi(x^\mu) \approx 1$. The relativistic contribution at the leading order to Newton’s gravity law discussed in this subsection differs from the one presented in [66]. The results outlined in (4.18) indicate an effective dark matter density profile

$$\rho_{\text{eff}}(r) = \phi_2^2(r) \overline{\rho_{DM}(t)}, \quad (4.39)$$

distinct from the Navarro-Frenk-White profile given in (1.1). It’s important to mention that our focus is solely on the tree-level contribution to Newton’s gravity law. For a more precise determination of the effective dark matter density profile, higher-order quantum corrections for the four-point matter scattering process can be incorporated. These loop corrections will provide a more refined contribution to the Tully-Fisher relations.

4.2 Dark Energy

In the late universe, assuming $a(t) \approx 1$, we consider the constraints imposed on $\phi_1(t)$ and $\phi_2(r)$ by equations

$$\ddot{\phi}_1(t) + 3H\dot{\phi}_1(t) = 0, \quad (4.40)$$

$$-\frac{1}{r^2} \partial_r \left(r^2 \frac{\partial \phi_2(r)}{\partial r} \right) + m_\phi^2 \phi_2(r) = 0. \quad (4.41)$$

The solutions to these equations representing dark energy are given by

$$\phi_1(t) = C_1 + C_2 a^{-3}(t), \quad (4.42)$$

$$\phi_2(r) = C_3 \frac{e^{-m_\phi r}}{r}, \quad (4.43)$$

where C_1, C_2, C_3 are constants. The the function $F(t, \mathbf{x})$ are described by

$$\langle F(t, \mathbf{x}) \rangle_t = -\phi_2^2(r) \overline{\rho_{DE}(t)}, \quad (4.44)$$

where $\phi_0 = C_1 + C_2$ as $a \simeq 1$ and $\overline{\rho_{DE}(t)} = \frac{1}{2} m_\phi^2 \phi_0^2 = 3 m_p^2 H^2 \times \Omega_{DE}$. Here Ω_{DE} is the fraction parameter of dark energy. After applying the Fourier transformation in (4.17), we discover that the spherically symmetric distribution of the inflaton generates a repulsive force between two massive objects. The Newton's gravity law for M_2 is given by

$$F_2(R) = -\frac{GM_1 M_2}{R^2} + \frac{4\pi G M_2 \overline{\rho_{DE}(t)} C_3^2 (1 - e^{-2m_\phi R})}{m_\phi R^2}. \quad (4.45)$$

This solution suggests that the extra dimensions can be interpreted as a spherically symmetric distribution of dark energy. This result aligns with the experimental constraints on the Yukawa gravitational potential [64, 65], where the characteristic length $\lambda = \frac{\hbar}{2m_\phi c} \approx 4.04 \times 10^{-30}[\text{m}]$, but it is far beyond the current experimental accuracy to detect the gravitational effect (4.45) at centimeter scale. It's worth noting that in this scenario, $w = \frac{p_{DE}}{\rho_{DE}} \approx -1$ indicating that the inflaton might be just one component of dark energy. However, the effective action (2.34) represents the tree-level effective action for eleven-dimensional supergravity. Higher-order derivatives of the inflaton, such as $\nabla^\mu \nabla^\nu \phi \nabla_\mu \nabla_\nu \phi$, can be found in the 1-loop order action of eleven-dimensional supergravity. We leave further discussions on the role of the inflaton as dark energy for future study.

5 Conclusions and Discussion

In this paper, we have discussed multiple potential methods for compactifying 11D supergravity, which can be represented as $\mathcal{M}^{3,1} \times T_{q_1}^2 \times T_{q_2}^2 \times T_{q_3}^2 \times S^1$. As a low-energy effective theory derived from the ultraviolet-complete M-theory, the inherent completeness of quantum gravity in the ultraviolet regime does not forbid the existence of dark matter and dark energy in a four-dimensional FRW universe. The dynamics of the inflaton, Standard Model of Particle Physics, dark matter, and dark energy can be all included in the effective action in four-dimensional spacetime. Meanwhile, the inflaton can be interpreted as the conformal vibrations of extra dimensions.

In our examples of driven potential (2.61), the dynamics of sub-Planck scale extra dimensions govern the cosmological inflation that occurs in four-dimensional spacetime. When the scale function has minimum value $\Phi_{min} = -\frac{1}{n}$ and s_1, s_2, s_3 take values $s_1 = \frac{1}{2}, s_2 = 0, s_3 \geq 1$, the theoretical predictions align with the observational range derived from the 2021 BICEP/Keck results, see Figure 2. The special case addressed in Subsection 2.1 satisfies all the observational constraints from both 2021 BICEP/Keck and 2018 Planck

results. In this scenario, the geometric configuration of the extra dimensions is $S^2 \times T_2^2 \times T_2^2 \times S^1$. As inflation progresses, the three tori undergo a contraction in size. Once the inflaton ϕ initiates its decay into gravitino ψ_A and 3-form A_{MNP} , inflation exits. During this decay process, particles encompassed by the Standard Model of Particle Physics, as well as dark matter and dark energy, are generated. Then the extra dimensions become inactive and stabilize into a fixed shape, where the standard sizes of the extra dimensions are around $10^5 l_p$.

As mentioned in Section 4, the quantum effects from extra dimensions are detectable in principle through high-energy experiments and astronomy observations. However, detecting the inflaton in accelerators is extremely challenging in practice. Nevertheless, there are other possible ways to indirectly infer the existence of extra dimensions, such as observations of the power spectrum of Cosmic Microwave Background (CMB) and the rotation curve of galaxies. On the one hand, in our compactification methods, the supersymmetry of gravity in four-dimensional spacetime is spontaneous breaking. This theoretical prediction aligns with experimental observations, as there is currently no conclusive evidence for superpartners of the graviton and the particles in the Standard Model of Particle Physics. On the other hand, the geometric configuration of the extra dimensions is reflected by the inflaton field, which can be interpreted as a component of dark matter or dark energy in the late universe. The modified Newton's gravity law, taking into account a spherically symmetric distribution of the inflaton, is presented in (4.26)(4.31)(4.45). The equation (4.31), which depicts a point particle surrounded by dark matter, can elucidate the Tully-Fisher relation given in (4.37). The range of the parameter m_2 in (4.31) can be constrained through observations for galaxies. It is worth mentioning that our method to derive the properties of the dark sector is fixed on the Minkowski background and the backreaction sourced by matter to spacetime is neglected since the gravitational field is very weak. To study the case in the strong gravitational field, more precise calculations regarding the backreaction should be performed, which is left for future work.

APPENDIX

We are going to prove the spectral index n_s and tensor-to-scalar ratio r are independent of the factor $(q_3 - 1)$ and parameter n for the potential(2.61). Perform a rescaling by

$$\tilde{n} = e^\alpha n, \quad (5.1)$$

$$\tilde{\Phi} = e^{-\alpha} \Phi, \quad (5.2)$$

$$\tilde{\phi} = -\alpha/\sqrt{\lambda^{-1}} + \phi, \quad (5.3)$$

where the parameter $\alpha = -\frac{1}{2s_3} \ln k$, $k \in \mathbb{N}_+$. The potential differs only by a factor

$$\tilde{U}(\tilde{\Phi}) = U(\tilde{\Phi}) = e^{2s_3\alpha} U(\Phi). \quad (5.4)$$

The slow-roll parameter $\epsilon(\phi)$ transforms as

$$\tilde{\epsilon}(\tilde{\phi}) = \frac{m_p^2}{2} \left(\frac{\tilde{U}_{\tilde{\phi}}}{\tilde{U}} \right)^2 = \frac{m_p^2}{2} \left(\frac{e^{2s_3\alpha} U_\phi d\phi/d\tilde{\phi}}{e^{2s_3\alpha} U} \right)^2 = \epsilon(\phi). \quad (5.5)$$

Hence, we find the parameter α does not affect the slow-roll parameter $\epsilon(\phi)$. This is the same for $\eta(\phi)$. Therefore the spectral index n_s and tensor-to-scalar ratio r are independent of α . As a result, we can select $\tilde{n} = e^\alpha$ such that $n = 1$. The genus of the third torus can be chosen to be $q_3 = 1 + k$, then the potential $\tilde{U}(\tilde{\Phi})$ dropped factor $(q_3 - 1)$.

ACKNOWLEDGEMENTS

We would like to thank Chi-Ming Chang, Zhengyuan Du, Wen-Xin Lai, Wei Song, Yi-xiao Tao, Dongjian Wu, and Tianqing Zhu for useful discussions. The work is partially supported by the national key research and development program of China NO. 2020YFA0713000.

References

- [1] BICEP, KECK collaboration, Improved Constraints on Primordial Gravitational Waves using Planck, WMAP, and BICEP/Keck Observations, [Phys. Rev. Lett.](#) **127** (2021) 151301 [[2110.00483](#)].
- [2] PLANCK collaboration, Planck 2018 results. X. Constraints on inflation, [Astron. Astrophys.](#) **641** (2020) A10 [[1807.06211](#)].
- [3] T. Banks, W. Fischler, S.H. Shenker and L. Susskind, M theory as a matrix model: A Conjecture, [Phys. Rev. D](#) **55** (1997) 5112 [[hep-th/9610043](#)].
- [4] J. Polchinski, M theory and the light cone, [Prog. Theor. Phys. Suppl.](#) **134** (1999) 158 [[hep-th/9903165](#)].
- [5] W. Taylor, M(atr)ix Theory: Matrix Quantum Mechanics as a Fundamental Theory, [Rev. Mod. Phys.](#) **73** (2001) 419 [[hep-th/0101126](#)].
- [6] B.S. Acharya, K. Bobkov, G.L. Kane, P. Kumar and J. Shao, Explaining the Electroweak Scale and Stabilizing Moduli in M Theory, [Phys. Rev. D](#) **76** (2007) 126010 [[hep-th/0701034](#)].
- [7] G. Kane, P. Kumar, R. Lu and B. Zheng, Higgs Mass Prediction for Realistic String/M Theory Vacua, [Phys. Rev. D](#) **85** (2012) 075026 [[1112.1059](#)].
- [8] G. Kane and M.W. Winkler, Deriving the Inflaton in Compactified M-theory with a De Sitter Vacuum, [Phys. Rev. D](#) **100** (2019) 066005 [[1902.02365](#)].
- [9] R. Brustein, S.P. de Alwis and E.G. Novak, M theory moduli space and cosmology, [Phys. Rev. D](#) **68** (2003) 043507 [[hep-th/0212344](#)].
- [10] A.A. Coley and R.J. van den Hoogen, Inhomogeneous M theory cosmologies, [J. Math. Phys.](#) **43** (2002) 4849 [[hep-th/0209022](#)].
- [11] D.Z. Freedman and A. Van Proeyen, [Supergravity](#), Cambridge Univ. Press, Cambridge, UK (5, 2012), [10.1017/CBO9781139026833](#).
- [12] M.B. Green, J.H. Schwarz and E. Witten, [Superstring Theory Vol. 2: 25th Anniversary Edition](#), Cambridge Monographs on Mathematical Physics, Cambridge University Press (11, 2012), [10.1017/CBO9781139248570](#).

- [13] P.D. D'Eath, Supersymmetric quantum cosmology, Cambridge Monographs on Mathematical Physics, Cambridge University Press (1, 2011), [10.1017/CBO9780511524424](https://doi.org/10.1017/CBO9780511524424).
- [14] M. Yamaguchi, Supergravity based inflation models: a review, [*Class. Quant. Grav.* **28** \(2011\) 103001](#) [[1101.2488](#)].
- [15] S. Ferrara, A. Kehagias and A. Sagnotti, Cosmology and Supergravity, [*Int. J. Mod. Phys. A* **31** \(2016\) 1630044](#) [[1605.04791](#)].
- [16] E. Lauria and A. Van Proeyen, $\mathcal{N} = 2$ Supergravity in $D = 4, 5, 6$ Dimensions, vol. 966 (3, 2020), [10.1007/978-3-030-33757-5](https://doi.org/10.1007/978-3-030-33757-5), [[2004.11433](#)].
- [17] S. Ferrara, R. Kallosh, A. Linde and M. Porrati, Minimal Supergravity Models of Inflation, [*Phys. Rev. D* **88** \(2013\) 085038](#) [[1307.7696](#)].
- [18] E. Halyo, Hybrid inflation from supergravity D terms, [*Phys. Lett. B* **387** \(1996\) 43](#) [[hep-ph/9606423](#)].
- [19] R. Kallosh, A. Linde and T. Rube, General inflaton potentials in supergravity, [*Phys. Rev. D* **83** \(2011\) 043507](#) [[1011.5945](#)].
- [20] J. Ellis, D.V. Nanopoulos and K.A. Olive, No-Scale Supergravity Realization of the Starobinsky Model of Inflation, [*Phys. Rev. Lett.* **111** \(2013\) 111301](#) [[1305.1247](#)].
- [21] J. Ellis, D.V. Nanopoulos and K.A. Olive, Starobinsky-like Inflationary Models as Avatars of No-Scale Supergravity, [*JCAP* **10** \(2013\) 009](#) [[1307.3537](#)].
- [22] L. Wu, T. Li and J. Pei, α -Generalized No-Scale Inflation, [2403.07333](#).
- [23] L. Wu and T. Li, Generic no-scale inflation inspired from string theory compactifications, [*Phys. Rev. D* **106** \(2022\) 043514](#) [[2205.14639](#)].
- [24] S.W. Hawking and H.S. Reall, Inflation, singular instantons and eleven-dimensional cosmology, [*Phys. Rev. D* **59** \(1999\) 023502](#) [[hep-th/9807100](#)].
- [25] M.S. Bremer, M.J. Duff, H. Lu, C.N. Pope and K.S. Stelle, Instanton cosmology and domain walls from M theory and string theory, [*Nucl. Phys. B* **543** \(1999\) 321](#) [[hep-th/9807051](#)].
- [26] A.P. Billyard, A.A. Coley, J.E. Lidsey and U.S. Nilsson, Dynamics of M theory cosmology, [*Phys. Rev. D* **61** \(2000\) 043504](#) [[hep-th/9908102](#)].
- [27] M. Cicoli, C.P. Burgess and F. Quevedo, Fibre Inflation: Observable Gravity Waves from IIB String Compactifications, [*JCAP* **03** \(2009\) 013](#) [[0808.0691](#)].
- [28] P. Svrcek and E. Witten, Axions In String Theory, [*JHEP* **06** \(2006\) 051](#) [[hep-th/0605206](#)].
- [29] H.-Y. Schive, T. Chiueh and T. Broadhurst, Cosmic Structure as the Quantum Interference of a Coherent Dark Wave, [*Nature Phys.* **10** \(2014\) 496](#) [[1406.6586](#)].
- [30] D.J.E. Marsh, Nonlinear hydrodynamics of axion dark matter: Relative velocity effects and quantum forces, [*Phys. Rev. D* **91** \(2015\) 123520](#) [[1504.00308](#)].

- [31] Y.V. Stadnik and V.V. Flambaum, Can dark matter induce cosmological evolution of the fundamental constants of Nature?, [Phys. Rev. Lett.](#) **115** (2015) 201301 [[1503.08540](#)].
- [32] T. Tenkanen, Dark matter from scalar field fluctuations, [Phys. Rev. Lett.](#) **123** (2019) 061302 [[1905.01214](#)].
- [33] T. Appelquist, J. Ingoldby and M. Piai, Dilaton Forbidden Dark Matter, [2404.07601](#).
- [34] C.A.J. O’Hare, Cosmology of axion dark matter, [2403.17697](#).
- [35] C. Cosme, J.a.G. Rosa and O. Bertolami, Scale-invariant scalar field dark matter through the Higgs portal, [JHEP](#) **05** (2018) 129 [[1802.09434](#)].
- [36] G. Alonso-Álvarez and J. Jaeckel, Lightish but clumpy: scalar dark matter from inflationary fluctuations, [JCAP](#) **10** (2018) 022 [[1807.09785](#)].
- [37] J. Shi, T. Katsuragawa and T. Qiu, Dark matter candidate induced by Horndeski theory: dark matter halo and cosmological evolution, [Phys. Rev. D](#) **101** (2020) 024046 [[1911.04741](#)].
- [38] G. Pierobon, J. Redondo, K. Saikawa, A. Vaquero and G.D. Moore, Miniclusters from axion string simulations, [2307.09941](#).
- [39] B. Eggemeier, A.K. Anilkumar and K. Dolag, Evidence for axion miniclusters with an increased central density, [2402.18221](#).
- [40] E. Bertschinger, Self - similar secondary infall and accretion in an Einstein-de Sitter universe, [Astrophys. J. Suppl.](#) **58** (1985) 39.
- [41] J.F. Navarro, C.S. Frenk and S.D.M. White, The Structure of cold dark matter halos, [Astrophys. J.](#) **462** (1996) 563 [[astro-ph/9508025](#)].
- [42] J.R. Primack, J.S. Bullock, A.A. Klypin and A.V. Kravtsov, Formation of dark matter halos, [ASP Conf. Ser.](#) **182** (1999) 517 [[astro-ph/9812241](#)].
- [43] M. Milgrom, A modification of the Newtonian dynamics: implications for galaxy systems, [Astrophys. J.](#) **270** (1983) 384.
- [44] M. Milgrom, A Modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis, [Astrophys. J.](#) **270** (1983) 365.
- [45] J. Bekenstein and M. Milgrom, Does the missing mass problem signal the breakdown of Newtonian gravity?, [Astrophys. J.](#) **286** (1984) 7.
- [46] M. Milgrom, Solutions for the modified Newtonian dynamics field equation, [Astrophys. J.](#) **302** (1986) 617.
- [47] R.B. Tully and J.R. Fisher, A New method of determining distances to galaxies, [Astron. Astrophys.](#) **54** (1977) 661.
- [48] S.S. McGaugh, J.M. Schombert, G.D. Bothun and W.J.G. de Blok, The Baryonic Tully-Fisher relation, [Astrophys. J. Lett.](#) **533** (2000) L99 [[astro-ph/0003001](#)].

- [49] S. Torres-Flores, B. Epinat, P. Amram, H. Plana and C.M. de Oliveira, GHASP: an H α kinematic survey of spiral and irregular galaxies – IX. The NIR, stellar and baryonic Tully [Mon. Not. Roy. Astron. Soc. **416** \(2011\) 1936 \[1106.0505\]](#).
- [50] G. Alestas, I. Antoniou and L. Perivolaropoulos, Hints for a Gravitational Transition in Tully–Fisher Data, [Universe **7** \(2021\) 366 \[2104.14481\]](#).
- [51] D. Clowe, M. Bradac, A.H. Gonzalez, M. Markevitch, S.W. Randall, C. Jones et al., A direct empirical proof of the existence of dark matter, [Astrophys. J. Lett. **648** \(2006\) L109 \[astro-ph/0608407\]](#).
- [52] SUPERNOVA SEARCH TEAM collaboration, Observational evidence from supernovae for an accelerating universe and a cosmological constant, [Astron. J. **116** \(1998\) 1009 \[astro-ph/9805201\]](#).
- [53] SUPERNOVA COSMOLOGY PROJECT collaboration, Measurements of Ω and Λ from 42 High Redshift Supernovae, [Astrophys. J. **517** \(1999\) 565 \[astro-ph/9812133\]](#).
- [54] S.M. Carroll, W.H. Press and E.L. Turner, The Cosmological constant, [Ann. Rev. Astron. Astrophys. **30** \(1992\) 499](#).
- [55] J. Martin, Everything You Always Wanted To Know About The Cosmological Constant Problem (But Were Afraid To Ask), [Comptes Rendus Physique **13** \(2012\) 566 \[1205.3365\]](#).
- [56] Q. Wang, Reformulation of the Cosmological Constant Problem, [Phys. Rev. Lett. **125** \(2020\) 051301 \[1904.09566\]](#).
- [57] S. Carlip, Hiding the cosmological constant, [Phys. Rev. Lett. **123** \(2019\) 131302 \[1809.08277\]](#).
- [58] DESI collaboration, DESI 2024 VI: Cosmological Constraints from the Measurements of Baryon Acoustic Oscillations, [2404.03002](#).
- [59] PLANCK collaboration, Planck 2018 results. VI. Cosmological parameters, [Astron. Astrophys. **641** \(2020\) A6 \[1807.06209\]](#).
- [60] B. Ydri, Matrix Models of String Theory, IOP Publishing, Bristol (2018), [10.1088/978-0-7503-1726-9](#).
- [61] Y. Hyakutake and S. Ogushi, Higher derivative corrections to eleven dimensional supergravity via local supersymmetry, [JHEP **02** \(2006\) 068 \[hep-th/0601092\]](#).
- [62] S. Mohanty, Gravitational Waves from a Quantum Field Theory Perspective, .
- [63] R. Fitzpatrick, An Introduction to Celestial Mechanics, Cambridge University Press (2012).
- [64] W.-H. Tan et al., Improvement for Testing the Gravitational Inverse-Square Law at the Submillimeter Range, [Phys. Rev. Lett. **124** \(2020\) 051301](#).
- [65] J.G. Lee, E.G. Adelberger, T.S. Cook, S.M. Fleischer and B.R. Heckel, New Test of the Gravitational $1/r^2$ Law at Separations down to 52 μm , [Phys. Rev. Lett. **124** \(2020\) 101101 \[2002.11761\]](#).

- [66] S. Das, Aspects of Machian Gravity (II): Testing Theory against Rotation Curves of 175 SPARC Galaxies, [2309.00057](#).
- [67] S. Comerón, I. Trujillo, M. Cappellari, F. Buitrago, L.E. Garduño, J. Zaragoza-Cardiel et al., The massive relic galaxy NGC 1277 is dark matter deficient. From dynamical models of integral-field stellar kinematics, [Astron. Astrophys. **675** \(2023\) A143 \[2303.11360\]](#).
- [68] P.E. Piña Mancera, F. Fraternali, T. Oosterloo, E.A.K. Adams, K.A. Oman and L. Leisman, No need for dark matter: resolved kinematics of the ultra-diffuse galaxy AGC 114905, [Mon. Not. Roy. Astron. Soc. **512** \(2022\) 3230 \[2112.00017\]](#).
- [69] P. van Dokkum et al., A galaxy lacking dark matter, [Nature **555** \(2018\) 629 \[1803.10237\]](#).
- [70] P. van Dokkum et al., A trail of dark-matter-free galaxies from a bullet-dwarf collision, [Nature **605** \(2022\) 435 \[2205.08552\]](#).
- [71] S. Danieli, P. van Dokkum, C. Conroy, R. Abraham and A.J. Romanowsky, Still Missing Dark Matter: KCWI High-Resolution Stellar Kinematics of NGC1052-DF2, [Astrophys. J. Lett. **874** \(2019\) L12 \[1901.03711\]](#).