# Regularity results for Hölder minimizers to functionals with non-standard growth

Antonio Giuseppe Grimaldi<sup>∗</sup> and Erica Ipocoana †

April 19, 2024

#### Abstract

We study the regularity properties of Hölder continuous minimizers to non-autonomous functionals satisfying  $(p, q)$ -growth conditions, under Besov assumptions on the coefficients. In particular, we are able to prove higher integrability and higher differentiability results for solutions to our minimum problem.

Keywords: Besov spaces, higher differentiability, non-standard growth, higher integrability, Hölder minimizers.

MSC 2020: 35J50, 35J70, 35J86, 35J87.

## 1 Introduction

In this work, we aim to present some regularity results for the solutions to the problem

<span id="page-0-0"></span>
$$
\min\left\{\mathcal{F}(u,\Omega) := \int_{\Omega} F(x,Du) \, \mathrm{d}x : u \in W^{1,p}(\Omega,\mathbb{R}^N)\right\},\tag{1.1}
$$

where  $N \geq 1$ ,  $\Omega$  is a bounded open subset of  $\mathbb{R}^n$ , for  $n \geq 2$ , and  $F : \Omega \times \mathbb{R}^{N \times n} \to [0, +\infty)$ is a Carathéodory function. Besides, we assume that there exist positive constants  $\nu, L$  and exponents  $1 < p < q < +\infty$  such that

$$
\nu |z|^p \le F(x, z) \le L(1 + |z|^q)
$$
 (F1)

$$
\nu(\mu^2 + |z_1|^2 + |z_2|^2)^{\frac{p-2}{2}} \le \langle \partial_z F(x, z_1) - \partial_z F(x, z_2), z_1 - z_2 \rangle \tag{F2}
$$

for a.e.  $x \in \Omega$  and all  $z, z_1, z_2 \in \mathbb{R}^{N \times n}$ , with  $\mu \in [0, 1]$ . Note that the ellipticity assumption (F2) and (F1) imply

$$
|\partial_z F(x,z)| \le c(1+|z|^{q-1})\tag{F3}
$$

<sup>∗</sup>Dipartimento di Scienze Matematiche "G. L. Lagrange", Politecnico di Torino, Corso Duca degli Abruzzi, 24, 10129 Torino, Italy. E-mail: antonio.grimaldi@polito.it

<sup>†</sup>Freie Universität Berlin, Department of Mathematics and Computer Science, Arnimallee 9, 14195 Berlin, Germany. E-mail: erica.ipocoana@fu-berlin.de

for a.e.  $x \in \Omega$  and every  $z \in \mathbb{R}^{N \times n}$ , with a constant  $c := c(L, q)$ . Moreover, we assume that there exists a non-negative function  $g \in L^{\sigma}_{loc}(\Omega)$ , where  $\sigma$  is finite and its precise expression will be introduced in [\(1.9\)](#page-3-0), such that

$$
|\partial_z F(x, z) - \partial_z F(y, z)| \le |x - y|^{\alpha} (g(x) + g(y))(1 + |z|^{q-1})
$$
 (F4)

for  $\alpha \in (0,1)$ , a.e.  $x, y \in \Omega$  and every  $z \in \mathbb{R}^{N \times n}$ .

Note that, according to standard results recalled in Section [2.2](#page-6-0) and in particular Theorem [2.8,](#page-8-0) assumption (F4) states that the function  $x \mapsto \partial_z F(x, z)$  belongs to  $B^{\alpha}_{\sigma, \infty}$ .

Functionals satisfying assumption  $(F1)$  above are called  $(p, q)$ -growth functionals, as first referred by Marcellini [\[41,](#page-23-0) [42,](#page-23-1) [43\]](#page-23-2). As a necessary and sufficient condition for regularity of minima, these non-standard growth functionals satisfy gap bounds of the type

$$
\frac{q}{p} < 1 + o(n), \qquad o(n) \approx \frac{1}{n},
$$

according to [\[25,](#page-22-0) [43\]](#page-23-2).

A very interesting model case of functionals with  $(p, q)$ -growth is the so called double phase, whose energy density has the form

<span id="page-1-0"></span>
$$
F(x, z) := |z|^p + a(x)|z|^q
$$
\n(1.2)

with  $a(x) \in C^{0,\alpha}(\Omega)$ . This model case, first introduced by Zhikov [\[49,](#page-24-0) [50,](#page-24-1) [51\]](#page-24-2) in order to describe the behaviour of strongly anisotropic materials in the context of homogenisation, nonlinear elasticity and Lavrentiev phenomenon, has been widely investigated in [\[6,](#page-21-0) [25,](#page-22-0) [26\]](#page-22-1) with the aim of identifying sufficient and necessary conditions on the relation between p, q and  $\alpha$  to establish the Hölder continuity of the gradient of the local minimizers. The sharp condition on the gap is given by

$$
\frac{q}{p} \le 1 + \frac{\alpha}{n}.
$$

On the other hand, in [\[16\]](#page-22-2) the authors showed that assuming that the minimizers are a priori bounded yields a gap bound of the form

$$
q \le p + \alpha,
$$

that is sharp as shown in [\[25,](#page-22-0) [26\]](#page-22-1). This is independent of the dimension and this phenomenon holds true for a priori bounded minimizers for functionals with  $(p, q)$ -growth without the structure as [\(1.2\)](#page-1-0) (see for instance [\[6,](#page-21-0) [9,](#page-21-1) [14,](#page-21-2) [36\]](#page-23-3)).

The fact that the a priori boundedness of minima leads to non-dimensional bounds on the distance  $q-p$  has been the motivation to the study of a priori more regular minimizers. Actually, in [\[16\]](#page-22-2) it has been proved that in the case of the double phase functional the gap improves with the a priori  $\mathcal{C}^{0,\gamma}$ -regularity of the minima, for  $0 < \gamma < 1$ . Indeed, the gap bound takes the form

$$
q < p + \frac{\alpha}{1 - \gamma},
$$

that is still independent of the dimension n and is sharp, as shown in [\[5\]](#page-21-3). Moreover, for  $\gamma \to 1$ , it holds

<span id="page-1-1"></span>
$$
q < p + o(\gamma), \qquad o(\gamma) \approx \frac{1}{1 - \gamma}, \tag{1.3}
$$

which is in particular coherent with the fact that Lipschitz continuity of minima is a main issue for  $(p, q)$ -growth problems. An additional step regarding a priori Hölder continuous minimizers is made in [\[20\]](#page-22-3), where the authors are able to improve the regularity of minima of non-autonomous functionals from  $W^{1,p}$  to  $W^{1,q}$ , taking into account the bound [\(1.3\)](#page-1-1). Here, we further extend the results obtained in [\[20\]](#page-22-3) by considering weaker assumptions on the function g, in particular, when g is not bounded (see  $(F3)$ ).

In the recent paper [\[36\]](#page-23-3), it has been proved that a priori bounded solutions to obstacle problems with non-standard growth satisfy extra fractional differentiability properties, under a free dimensional condition between the ellipticity and the growth exponents of the functional. Here, we continue the study of regularity theory of a priori more regular minimizers to integral functionals of Calculus of Variations with  $(p, q)$ -growth. The novelty of this work consists in showing that, assuming a Besov regularity on the coefficients of the problem, the higher integrability and differentiability of a priori Hölder continuous minima hold under a relation between p and q of type  $(1.3)$ .

For precise definition and properties of Besov spaces we refer to Section [2.2.](#page-6-0) We note that, indeed, recently, there have been many research activities regarding the analysis of higher differentiability (of integer and fractional order) of solutions to variational problems; see  $\ket{2, 3, 4}$  $\ket{2, 3, 4}$  $\ket{2, 3, 4}$ . [15,](#page-21-7) [32,](#page-23-4) [33,](#page-23-5) [44,](#page-23-6) [45,](#page-23-7) [46\]](#page-23-8), for the unconstrained case, and [\[8,](#page-21-8) [12,](#page-21-9) [13,](#page-21-10) [18,](#page-22-4) [22,](#page-22-5) [23,](#page-22-6) [27,](#page-22-7) [28,](#page-22-8) [29,](#page-22-9) [30,](#page-22-10) [31,](#page-22-11) [35,](#page-23-9) [36,](#page-23-3) [37,](#page-23-10) [38,](#page-23-11) [48\]](#page-24-3), for the obstacle problem.

As stated above, according to [\(1.1\)](#page-0-0), we are here dealing with a non-autonomous functional. Therefore, the Lavrentiev phenomenon has to be taken into account, i.e.

$$
\inf_{w \in W^{1,p}(\Omega)} \int_{\Omega} F(x, Dw) \, dx < \inf_{w \in W^{1,q}(\Omega)} \int_{\Omega} F(x, Dw) \, dx.
$$

This is a clear obstruction to regularity, since it prevents minimizers to belong to  $W^{1,q}$  (see [\[41,](#page-23-0) [49,](#page-24-0) [51\]](#page-24-2) for more details).

Here, we are interested in the case of a priori Hölder minimizers to the problem [\(1.1\)](#page-0-0). Hence, we will adopt suitable definitions of relaxed and gap functionals, aiming at exploiting this fact. Let us fix constants  $H > 0$  and  $\gamma \in (0, 1)$ . Let  $B \subset \Omega$  be a ball, we can consider the relaxed functional

$$
\bar{\mathcal{F}}_{H,\gamma}(u,B) := \inf_{\{u_j\} \in \mathcal{C}_{H,\gamma}(u,B)} \left\{ \liminf_{j \to \infty} \int_B F(x, Du_j) \, dx \right\} \tag{1.4}
$$

for any  $u \in W^{1,1}(B,\mathbb{R}^N)$ , where

$$
\mathcal{C}_{H,\gamma}(u,B) := \left\{ \{w_j\} \subset W^{1,\infty}(B,\mathbb{R}^N) : w_j \rightharpoonup u \text{ in } W^{1,p}(B,\mathbb{R}^N), \sup_j [w_j]_{0,\gamma;B} \le H \right\} \tag{1.5}
$$

and where we used the standard notation

$$
[v]_{0,\gamma;A} := \sup_{x,y \in A, x \neq y} \frac{|v(x) - v(y)|}{|x - y|^{\gamma}},
$$

whenever  $A \subset \mathbb{R}^n$  and  $v : A \to \mathbb{R}^N$ . By lower semicontinuity of the functional  $\mathcal F$  with respect to the weak convergence in  $W^{1,p}$ , we have  $\tilde{\mathcal{F}}_{H,\gamma}(u, B) \geq \mathcal{F}(u, B)$ . We define for all  $u \in W^{1,1}(B, \mathbb{R}^N)$  the Lavrentiev gap functional relative to  $\mathcal F$  as follows

$$
\mathcal{L}_{F,H,\gamma}(u,B) := \begin{cases} \bar{\mathcal{F}}_{H,\gamma}(u,B) - \mathcal{F}(u,B) & \text{if } \mathcal{F}(u,B) < +\infty \\ 0 & \text{if } \mathcal{F}(u,B) = +\infty. \end{cases}
$$
(1.6)

From the previous definitions, we can easily derive

**Proposition 1.1.** Let  $w \in W^{1,1}(B,\mathbb{R}^N)$  be such that  $\mathcal{F}(w, B)$  is finite, for some ball  $B \in \Omega$ . Then,  $\bar{\mathcal{F}}_{H,\gamma}(w, B) = \mathcal{F}(w, B)$ , for some  $H > 0$  if, and only if, there exists a sequence  $\{w_j\} \in$  $\mathcal{C}_{H,\gamma}(w, B)$  such that

$$
\mathcal{F}(w_j, B) \to \mathcal{F}(w, B).
$$

In particular, in the sequel we need to impose that the Lavrentiev phenomenon does not occur, as a necessary condition for the the regularity of minima [\[1,](#page-21-11) [19,](#page-22-12) [25\]](#page-22-0).

We may now state the two main results contained in this work, concerning higher integrability and higher differentiability of solutions to [\(1.1\)](#page-0-0), respectively.

<span id="page-3-2"></span>**Theorem 1.2** (Higher integrability). Let  $\alpha, \gamma \in (0,1)$  and let  $F(x, z)$  satisfy (F1) and (F2) for exponents  $1 < p < q$  such that

<span id="page-3-3"></span>
$$
q < p + \frac{\min\{\alpha, 2\gamma\}}{\Theta(1-\gamma)},\tag{1.7}
$$

where

<span id="page-3-4"></span>
$$
\Theta := \begin{cases} 1 & \text{if } p \ge 2 \\ \frac{2}{p} & \text{if } 1 < p < 2. \end{cases} \tag{1.8}
$$

Moreover, assume that there exists a non-negative function  $g \in L^{\sigma}_{loc}(\Omega)$ , with

<span id="page-3-0"></span>
$$
\sigma = \frac{p(1-\gamma) + \xi}{(p-q)(1-\gamma) + \xi} \tag{1.9}
$$

and  $(q-p)(1-\gamma) < \xi < \frac{\min\{\alpha, 2\gamma\}}{\gamma}$ Θ , such that  $(F4)$  holds.

Let  $u \in W^{1,p}(\Omega,\mathbb{R}^N)$  be a solution to the minimum problem [\(1.1\)](#page-0-0). Assuming that

<span id="page-3-1"></span>
$$
\mathcal{L}_{F,H,\gamma}(u, B_r) = 0,\tag{1.10}
$$

holds for a ball  $B_r \in \Omega$  with  $r \leq 1$  and for some  $H > 0$ , then

$$
Du \in L^{\tilde{q}}_{loc}(\Omega, \mathbb{R}^{N \times n}),\tag{1.11}
$$

for all  $q < \tilde{q} < p +$  $\min\{\alpha, 2\gamma\}$  $\frac{\dim \{ \alpha, 2\gamma \}}{\Theta(1-\gamma)}$ . In particular, for every  $B_{\rho} \in B_r$ , we have

$$
||Du||_{L^{\tilde{q}}(B_{\rho})} \leq \frac{C}{(r-\rho)^{\kappa_1}} ( [\mathcal{F}(u,B_r)]^{\frac{1}{p}} + ||g||_{L^{\sigma}(B_r)}^{\sigma} + H + 1)^{\kappa_2}, \qquad (1.12)
$$

for constants  $C := C(n, p, q, \tilde{q}, \nu, L, \alpha, \gamma, \sigma)$  and  $\kappa_1, \kappa_2 := \kappa_1, \kappa_2(n, p, q, \tilde{q}, \alpha, \gamma, \sigma)$ .

Remark 1.3. Assumption [\(1.10\)](#page-3-1) in Theorem [1.2](#page-3-2) ensures that the minimizer u is Hölder continuous. By minimality we have  $\mathcal{F}(u, B) < +\infty$ , for every ball  $B \in \Omega$ . Note that  $\mathcal{C}_{H,\gamma}(w, B)$ cannot be empty, since it would imply  $\bar{\mathcal{F}}(u, B) = +\infty$ . Therefore,  $\mathcal{C}_{H,\gamma}(w, B) \neq \emptyset$ , which yields  $u \in C^{0,\gamma}_{loc}(\Omega,\mathbb{R}^N)$ , with  $[u]_{0,\gamma;B} \leq H$ .

Besides, we point out that the gap condition  $(1.7)$  is of type  $(1.3)$ .

It is convenient to introduce the auxiliary function

$$
V_p(z) = (\mu^2 + |z|^2)^{\frac{p-2}{4}} z
$$

defined for all  $z \in \mathbb{R}^k$ ,  $k \in \mathbb{N}$ , in order to state the following result and as it will be largely used in the sequel.

<span id="page-4-0"></span>**Theorem 1.4** (Higher differentiability). Let  $\alpha, \gamma \in (0,1)$  and let  $\sigma$  be the exponent defined in [\(1.9\)](#page-3-0). Let  $F(x, z)$  satisfy  $(F1)$ – $(F4)$  for exponents  $1 < p < q$  such that [\(1.7\)](#page-3-3) holds. Let  $u \in W^{1,p}(\Omega,\mathbb{R}^N)$  be a solution to the minimum problem [\(1.1\)](#page-0-0). Assuming that [\(1.10\)](#page-3-1) holds for a ball  $B_r \in \Omega$  with  $r \leq 1$  and for some  $H > 0$ , then

$$
V_p(Du) \in B_{2,\infty,loc}^{\frac{\alpha}{2}}(\Omega, \mathbb{R}^{N \times n})
$$
\n(1.13)

and the following estimate

$$
\int_{B_{\rho}} |\tau_h V_p(Du)|^2 dx \le C|h|^{\alpha} \left\{ \int_{B_r} (1+|Du|^p) dx + \|g\|_{L^{\sigma}(B_r)}^{\sigma} + H \right\}^{\vartheta} \tag{1.14}
$$

holds true for every  $B_{\rho} \in B_r$ , for constants  $C := C(n, p, q, \nu, L, \alpha, \gamma, \sigma, r, \rho)$  and  $\vartheta := \vartheta(n, p, q, \alpha, \gamma, \sigma)$ .

The strategy of the proof of Theorem [1.2](#page-3-2) is to approximate the original minimizer  $u$  of  $(1.1)$  with a sequence  $\{u_i\}$  of  $W^{1,q}$ -regular solutions to suitable approximating problems. The possibility of this approximation relies on the assumptions [\(1.10\)](#page-3-1). The main difficulty of this work is to build the approximating minimizers  $\{u_i\}$  such that they are uniformly bounded in  $\mathcal{C}^{0,\gamma}$ . Actually, we build the sequence  $\{u_j\}$  in a way that it is bounded in a fractional Sobolev space  $W^{s,2d}$ , which is a good approximation of  $\mathcal{C}^{0,\gamma}$ , for a suitable choice of parameters s and d. In order to deal with this issue, we follow the strategy first proposed by De Filippis and Mingione in [\[20\]](#page-22-3), i.e. we employ a method based on approximating the original local functional with mixed local and non-local functionals. We remark that the weaker assumptions on  $q$ imply the use of different techniques and strategy when controlling the term  $(III)$  in [\(3.20\)](#page-12-0). In particular, we need a higher integrability condition, which is reflected by suitable hypotheses on exponents contained at the beginning of Section [3.1.](#page-9-0)

Next, taking advantage of the higher integrability of solutions to [\(1.1\)](#page-0-0), we will be able to prove the higher differentiability result. The proof of Theorem [1.4](#page-4-0) is achieved by means of difference quotient method, that is quite natural when trying to establish higher differentiabilty results and local gradient estimates (see for instance [\[42\]](#page-23-1)).

Remark 1.5. We remark that assuming a stronger Besov regularity on the coefficients of  $\partial_zF(x, z)$ , namely hypothesis (F4') below, we are able to prove that solutions to [\(1.1\)](#page-0-0) inherit a Besov regularity of the type  $B_{2,t}^{\frac{\alpha}{2}}$ , with  $t \geq 1$ .

More precisely, let  $\alpha, \gamma \in (0, 1)$  and let  $\sigma$  be the exponent defined in [\(1.9\)](#page-3-0). Let  $F(x, z)$  satisfy

(F1) and (F2) for exponents  $1 < p < q$  verifying [\(1.7\)](#page-3-3). Moreover, assume that there exists a sequence of measurable non-negative functions  $g_k \in L^{\sigma}_{loc}(\Omega)$  such that

$$
\sum_{k=1}^{\infty} \|g_k\|_{L^{\sigma}(\Omega)}^t < \infty,
$$

for some  $t > 1$ , and at the same time

$$
|\partial_z F(x, z) - \partial_z F(y, z)| \le |x - y|^{\alpha} (g_k(x) + g_k(y))(1 + |z|^{q-1})
$$
 (F4')

for a.e.  $x, y \in \Omega$  such that  $2^{-k}$  diam( $\Omega$ )  $\leq |x - y| < 2^{-k+1}$  diam( $\Omega$ ) and for every  $z \in \mathbb{R}^{N \times n}$ . Let  $u \in W^{1,p}(\Omega,\mathbb{R}^N)$  be a solution to the minimum problem [\(1.1\)](#page-0-0). Assuming that [\(1.10\)](#page-3-1) holds for a ball  $B_r \in \Omega$  with  $r \leq 1$  and for some  $H > 0$ , then

(i) 
$$
Du \in L^{\tilde{q}}_{loc}(\Omega, \mathbb{R}^{N \times n})
$$
, for all  $q \leq \tilde{q} < p + \frac{\min\{\alpha, 2\gamma\}}{\Theta(1-\gamma)}$ ;

(ii)  $V_p(Du) \in B_{2,t,loc}^{\frac{\alpha}{2}}(\Omega,\mathbb{R}^{N \times n})$ , for all  $t \geq 1$ .

The higher integrability result goes exactly in the same way of that of Theorem [1.2.](#page-3-2) On the other hand, for the proof of the last part we refer to [\[22,](#page-22-5) [29,](#page-22-9) [35,](#page-23-9) [37\]](#page-23-10).

The paper is organized as follows. After recalling some notation and preliminary results in Section [2,](#page-5-0) we prove our main results, Theorems [1.2](#page-3-2) and [1.4](#page-4-0) in Section [3.](#page-9-1)

### <span id="page-5-0"></span>2 Notation and preliminary results

For the rest of the paper, we denote by  $C$ , c general positive constants. Different occurrences from line to line will be still denoted using the same letters. Relevant dependencies on parameters will be emphasized using parentheses or subscripts. We denote by  $B(x,r) = B_r(x) = \{y \in \mathbb{R}^n : |y - x| < r\}$  the ball centered at x of radius r. We shall omit the dependence on the center and on the radius when no confusion arises.

For the auxiliary function  $V_p$ , we recall the following estimate (see the proof of [\[34,](#page-23-12) Lemma 8.3]):

<span id="page-5-2"></span>**Lemma 2.1.** Let  $1 < p < +\infty$ . There exists a constant  $c = c(n, p) > 0$  such that

$$
c^{-1}(\mu^2+|\xi|^2+|\eta|^2)^{\frac{p-2}{2}}\leq \frac{|V_p(\xi)-V_p(\eta)|^2}{|\xi-\eta|^2}\leq c(\mu^2+|\xi|^2+|\eta|^2)^{\frac{p-2}{2}}
$$

for any  $\xi, \eta \in \mathbb{R}^k$ ,  $\xi \neq \eta$ ,  $k \in \mathbb{N}$ .

Now we state a well-known iteration lemma (see [\[34,](#page-23-12) Lemma 6.1] for the proof).

<span id="page-5-1"></span>**Lemma 2.2.** Let  $\Phi : [\frac{R}{2}, R] \to \mathbb{R}$  be a bounded nonnegative function, where  $R > 0$ . Assume that for all  $\frac{R}{2} \leq r < s \leq R$  it holds

$$
\Phi(r) \le \theta \Phi(s) + A + \frac{B}{(s-r)^2} + \frac{C}{(s-r)^{\gamma}}
$$

where  $\theta \in (0,1), A, B, C \geq 0$  and  $\gamma > 0$  are constants. Then there exists a constant  $c = c(\theta, \gamma)$ such that

$$
\Phi\left(\frac{R}{2}\right) \le c\bigg(A + \frac{B}{R^2} + \frac{C}{R^{\gamma}}\bigg).
$$

#### 2.1 Difference quotient

We recall some properties of the finite difference quotient operator that will be needed in the sequel. For every function  $F: \mathbb{R}^n \to \mathbb{R}^k$ ,  $k \in \mathbb{N}$ , let

$$
\tau_h F(x) := F(x+h) - F(x),
$$

where  $h \in \mathbb{R}^n$ .

We start with the description of some elementary properties that can be found, for example, in [\[34\]](#page-23-12).

**Proposition 2.3.** Let  $F \in W^{1,p}(\Omega,\mathbb{R}^k)$ , with  $p \geq 1$ , and let us consider the set

 $\Omega_{|h|} = \{x \in \Omega : \text{dist}(x, \partial \Omega) > |h|\}.$ 

Then (i)  $\tau_h F \in W^{1,p}(\Omega_{|h|}, \mathbb{R}^k)$  and

$$
D_i(\tau_h F) = \tau_h(D_i F).
$$

(ii) If at least one of the functions F or G has support contained in  $\Omega_{|h|}$ , then

$$
\int_{\Omega} F \tau_h G \, \mathrm{d}x = - \int_{\Omega} G \tau_{-h} F \, \mathrm{d}x.
$$

(iii) We have

$$
\tau_h(FG)(x) = F(x+h)\tau_h G(x) + G(x)\tau_h F(x).
$$

The next result about finite difference operator is a kind of integral version of Lagrange Theorem.

<span id="page-6-1"></span>**Lemma 2.4.** If 
$$
0 < \rho < R
$$
,  $|h| < \frac{R-\rho}{2}$ ,  $1 < p < +\infty$  and  $F \in W^{1,p}(B_R, \mathbb{R}^k)$ , then  

$$
\int_{B_{\rho}} |\tau_h F(x)|^p dx \le c(n, p)|h|^p \int_{B_R} |DF(x)|^p dx.
$$

Moreover,

$$
\int_{B_{\rho}} |F(x+h)|^p \, \mathrm{d}x \le \int_{B_R} |F(x)|^p \, \mathrm{d}x.
$$

#### <span id="page-6-0"></span>2.2 Function spaces

Let  $v : \mathbb{R}^n \to \mathbb{R}^k$  be a function,  $k \in \mathbb{N}$ . As in [\[47,](#page-23-13) Section 2.2.2], given  $0 < \alpha < 1$  and  $1 \leq p, s < \infty$ , we say that v belongs to the Besov space  $B^{\alpha}_{p,s}(\mathbb{R}^n, \mathbb{R}^k)$  if  $v \in L^p(\mathbb{R}^n, \mathbb{R}^k)$  and

$$
[v]_{B_{p,s}^{\alpha}(\mathbb{R}^n)} := \left(\int_{\mathbb{R}^n} \left(\int_{\mathbb{R}^n} \frac{|\tau_h v(x)|^p}{|h|^{\alpha p}} \, \mathrm{d}x\right)^{\frac{s}{p}} \frac{\mathrm{d}h}{|h|^n}\right)^{\frac{1}{s}} < \infty.
$$

Equivalently, we could simply say that  $v \in L^p(\mathbb{R}^n, \mathbb{R}^k)$  and  $\frac{\tau_h v}{|h|^{\alpha}} \in L^s(\frac{dh}{|h|^n}; L^p(\mathbb{R}^n))$ . Similarly, we say that  $v \in B^{\alpha}_{p,\infty}(\mathbb{R}^n, \mathbb{R}^k)$  if  $v \in L^p(\mathbb{R}^n, \mathbb{R}^k)$  and

$$
[v]_{B_{p,\infty}^{\alpha}(\mathbb{R}^n)} := \sup_{h \in \mathbb{R}^n} \left( \int_{\mathbb{R}^n} \frac{|\tau_h v(x)|^p}{|h|^{\alpha p}} \, \mathrm{d}x \right)^{\frac{1}{p}} < \infty.
$$

We also define the following norm for  $B^{\alpha}_{p,s}(\mathbb{R}^n, \mathbb{R}^k)$ 

$$
||v||_{B_{p,s}^{\alpha}(\mathbb{R}^n)} := ||v||_{L^p(\mathbb{R}^n)} + [v]_{B_{p,s}^{\alpha}(\mathbb{R}^n)}.
$$
\n(2.1)

Moreover, if one integrates for  $h \in B(0, \delta)$  for a fixed  $\delta > 0$  then an equivalent norm is obtained, because

$$
\left(\int_{\{|h|\geq\delta\}}\left(\int_{\mathbb{R}^n}\frac{|\tau_h v(x)|^p}{|h|^{\alpha p}}\,\mathrm{d}x\right)^{\frac{s}{p}}\frac{\mathrm{d}h}{|h|^n}\right)^{\frac{1}{s}} \leq c(n,\alpha,p,s,\delta)\|v\|_{L^p(\mathbb{R}^n)}.\tag{2.2}
$$

When  $s = \infty$ , one can simply take supremum over  $|h| \leq \delta$  and obtain an equivalent norm.

By construction, one has  $B_{p,s}^{\alpha}(\mathbb{R}^n,\mathbb{R}^k) \subset L^p(\mathbb{R}^n,\mathbb{R}^k)$ . One also has the following version of Sobolev embeddings (a proof can be found at [\[39,](#page-23-14) Proposition 7.12]).

<span id="page-7-0"></span>**Lemma 2.5.** Suppose that  $0 < \alpha < 1$ . (a) If  $1 < p < \frac{n}{\alpha}$  and  $1 \le s \le p^*_{\alpha} = \frac{np}{n-c}$  $\frac{np}{n-\alpha p}$ , then there is a continuous embedding  $B^{\alpha}_{p,s}(\mathbb{R}^n) \subset$  $L^{p^*_{\alpha}}(\mathbb{R}^n)$ . (b) If  $p = \frac{n}{\alpha}$  $\frac{n}{\alpha}$  and  $1 \leq s \leq \infty$ , then there is a continuous embedding  $B^{\alpha}_{p,s}(\mathbb{R}^n) \subset BMO(\mathbb{R}^n)$ ,

where BMO denotes the space of functions with bounded mean oscillations [\[34,](#page-23-12) Chapter 2].

<span id="page-7-1"></span>We recall the following inclusions ([\[39,](#page-23-14) Proposition 7.10 and Formula (7.35)]).

**Lemma 2.6.** Suppose that  $0 < \beta < \alpha < 1$ . (a) If  $1 < p < \infty$  and  $1 \le s \le t \le \infty$ , then  $B^{\alpha}_{p,s}(\mathbb{R}^n) \subset B^{\alpha}_{p,t}(\mathbb{R}^n)$ . (b) If  $1 < p < \infty$  and  $1 \le s, t \le \infty$ , then  $B^{\alpha}_{p,s}(\mathbb{R}^n) \subset B^{\beta}_{p,t}(\mathbb{R}^n)$ .

Given a domain  $\Omega \subset \mathbb{R}^n$ , we say that v belongs to the local Besov space  $B_{p,s,loc}^{\alpha}$  if  $\varphi$  v  $\in$  $B_{p,s}^{\alpha}(\mathbb{R}^n)$  whenever  $\varphi \in \mathcal{C}_c^{\infty}(\Omega)$ . It is worth noticing that one can prove suitable version of Lemmas [2.5](#page-7-0) and [2.6,](#page-7-1) by using local Besov spaces.

We have the following characterization for local Besov spaces (see [\[4\]](#page-21-6) for the proof).

**Lemma 2.7.** A function  $v \in L_{loc}^p(\Omega, \mathbb{R}^k)$  belongs to the local Besov space  $B_{p,s,loc}^{\alpha}$ , with  $0 < \alpha < 1$ , if, and only if,

$$
\left\|\frac{\tau_h v}{|h|^{\alpha}}\right\|_{L^s\left(\frac{dh}{|h|^n};L^p(B)\right)} < \infty,
$$

for any ball  $B \subset 2B \subset \Omega$  with radius  $r_B$ , where

$$
\left\|\frac{\tau_h v}{|h|^{\alpha}}\right\|_{L^s\left(\frac{dh}{|h|^n};L^p(B)\right)} := \left(\int_{\{|h|\geq \delta\}} \left(\int_B \frac{|\tau_h v(x)|^p}{|h|^{\alpha p}} \, \mathrm{d}x\right)^{\frac{s}{p}} \frac{\mathrm{d}h}{|h|^n}\right)^{\frac{1}{s}}.
$$

Here the measure  $\frac{dh}{|h|^n}$  is restricted to the ball  $B(0, r_B)$  on the h-space.

It is known that Besov-Lipschitz spaces of fractional order  $\alpha \in (0,1)$  can be characterized in pointwise terms. Adopting the terminology of [\[40\]](#page-23-15), given a measurable function  $v : \mathbb{R}^n \to \mathbb{R}$ , a fractional  $\alpha$ -Hajlasz gradient for v is a sequence  $\{g_k\}_k$  of measurable, non-negative functions  $g_k: \mathbb{R}^n \to \mathbb{R}$ , together with a null set  $N \subset \mathbb{R}^n$ , such that the inequality

$$
|v(x) - v(y)| \le (g_k(x) + g_k(y))|x - y|^{\alpha}
$$

holds whenever  $k \in \mathbb{Z}$  and  $x, y \in \mathbb{R}^n \setminus N$  are such that  $2^{-k} \leq |x - y| < 2^{-k+1}$ . We say that  ${g_k}_k \in l^s(\mathbb{Z}; L^p(\mathbb{R}^n))$  if

$$
\|\{g_k\}_k\|_{l^s(L^p)} = \left(\sum_{k\in\mathbb{Z}} \|g_k\|_{L^p(\mathbb{R}^n)}^s\right)^{\frac{1}{s}} < \infty.
$$

In fact,  ${g_k}_k$  above is not really a gradient. One should view it as a maximal function of the usual gradient.

The following result was proved in [\[40\]](#page-23-15).

<span id="page-8-0"></span>**Theorem 2.8.** Let  $0 < \alpha < 1$ ,  $1 \leq p < \infty$  and  $1 \leq s \leq \infty$ . Let  $v \in L^p(\mathbb{R}^n)$ . One has  $v \in B^{\alpha}_{p,s}(\mathbb{R}^n)$  if, and only if, there exists a fractional  $\alpha$ -Hajlasz gradient  $\{g_k\}_k \in l^s(\mathbb{Z}; L^p(\mathbb{R}^n))$ for *v*. Moreover,

$$
||v||_{B^{\alpha}_{p,s}(\mathbb{R}^n)} \simeq \inf ||\{g_k\}_k||_{l^s(L^p)},
$$

where the infimum runs over all possible fractional  $\alpha$ -Hajlasz gradients for v.

Let  $\sigma \in (0, +\infty) \setminus \mathbb{N}, p \in [1, +\infty), k \in \mathbb{N}, n \ge 2$ . If  $\sigma \in (0, 1)$ , we say that a function  $v: \Omega \to \mathbb{R}^k$  belongs to the fractional Sobolev space  $W^{\sigma,p}(\Omega,\mathbb{R}^k)$  if, and only if, the following Gagliardo type norm is finite:

$$
||v||_{W^{\sigma,p}(\Omega)} := ||v||_{L^p(\Omega)} + \left(\int_{\Omega} \int_{\Omega} \frac{|v(x) - v(y)|^p}{|x - y|^{n + \sigma p}} dxdy\right)^{\frac{1}{p}}
$$
  
=:  $||v||_{L^p(\Omega)} + [v]_{\sigma,p;\Omega}.$ 

On the other hand, if  $\sigma = [\sigma] + \{\sigma\} \in \mathbb{N} + (0, 1) > 1$ , we have  $v \in W^{\sigma, p}(\Omega, \mathbb{R}^k)$  if, and only if,

$$
||v||_{W^{\sigma,p}(\Omega)} := ||v||_{W^{[\sigma],p}(\Omega)} + [D^{[\sigma]}v]_{\{\sigma\},p;\Omega}
$$

is finite.

From Lemma [2.6,](#page-7-1) we deduce the following fundamental embedding property between Besov and fractional Sobolev spaces.

<span id="page-8-1"></span>**Lemma 2.9.** We consider radii  $0 < \rho < R$  and a function  $v \in L^p(B_R, \mathbb{R}^k)$ . Suppose that there exist  $\sigma \in (0,1)$  and  $M > 0$  such that

$$
\|\tau_h v\|_{L^p(B_\rho)} \le M|h|^\sigma
$$

for every  $h \in \mathbb{R}^n$  with  $|h| < \frac{R-\rho}{2}$  $\frac{-\rho}{2}$ . Then  $v \in W^{\beta,q}(B_\rho, \mathbb{R}^k)$  for every  $\beta \in (0, \sigma)$ . Moreover,

$$
||v||_{W^{\beta,p}(B_{\rho},\mathbb{R}^k)} \leq \frac{c}{(R-\rho)^{\delta}}\left(M + ||v||_{L^p(B_R,\mathbb{R}^k)}\right),
$$

holds true for constants c,  $\delta$  depending on n, p,  $\beta$  and  $\sigma$ .

We conclude this section with a Gagliardo-Nirenberg type inequality in fractional Sobolev spaces (see [\[20\]](#page-22-3) for the proof).

<span id="page-9-3"></span>**Lemma 2.10.** Let  $B_\rho \in B_r \in \mathbb{R}^n$  be concentric balls with  $r \leq 1$ . Let  $0 \leq s_1 < 1 < s_2 < 2, 1 <$  $a, t < \infty, \tilde{p} > 1$  and  $\theta \in (0, 1)$  be such that

$$
\theta s_1 + (1 - \theta)s_2 = 1,
$$
  $\frac{1}{\tilde{p}} = \frac{\theta}{a} + \frac{1 - \theta}{t}.$ 

Then every function  $v \in W^{s_1,a}(B_r, \mathbb{R}^k) \cap W^{s_2,t}(B_r, \mathbb{R}^k)$  belongs to  $W^{1,\tilde{p}}(B_\rho, \mathbb{R}^N)$  and the inequality

$$
||Dv||_{L^{\tilde{p}}(B_{\rho})} \leq \frac{c}{(r-\rho)^{\kappa}} [v]_{s_1,a;B_r}^{\theta} ||Dv||_{W^{s_2-1,t}(B_r)}^{1-\theta}
$$

<span id="page-9-1"></span>holds for constants  $c, \kappa = c, \kappa(n, s_1, s_2, a, t)$ .

### 3 Proof of Theorems [1.2](#page-3-2) and [1.4](#page-4-0)

In the following we prove the main results of this work, namely the higher integrability result, Theorem [1.2,](#page-3-2) and the higher differentiability one, Theorem [1.4.](#page-4-0) The strategy proposed to prove Theorem [1.2](#page-3-2) follows the one introduced in [\[20\]](#page-22-3). However, we remark that we are here dealing with a weaker assumption on the summability of the map  $x \mapsto \partial_z F(x, z)$  (see [\(1.9\)](#page-3-0)). We denote by  $u \in W^{1,p}(\Omega,\mathbb{R}^N)$  a solution to [\(1.1\)](#page-0-0). Throughout this section, we will assume that [\(1.10\)](#page-3-1) holds for a ball  $B_r \in \Omega$  with  $r \leq 1$  and for some  $H > 0$ .

#### <span id="page-9-0"></span>3.1 Higher integrability

We start by considering parameters s, d and  $\beta$  such that

$$
0 \le s < \gamma, \qquad 2d > \max\{q, n\}, \qquad 0 < \beta < \min\{\alpha, 2\gamma\}. \tag{3.1}
$$

We define the function  $\tilde{p} = \tilde{p}(s, d, \beta)$  as

<span id="page-9-4"></span>
$$
\tilde{p} := \begin{cases}\n\frac{2d(p(1-s) + \beta)}{\beta + 2d(1-s)} & \text{if } p \ge 2 \\
\frac{2dp[2(1-s) + \beta]}{p\beta + 4d(1-s)} & \text{if } 1 < p < 2\n\end{cases}
$$
\n(3.2)

By definition, we have

 $d > \frac{p}{2}$ 2  $(3.3)$ 

which yields

$$
\tilde{p} < 2d. \tag{3.4}
$$

The function  $\tilde{p}$  is increasing in all its variables and

$$
\lim_{s \to \gamma, d \to \infty, \beta \to \min\{\alpha, 2\gamma\}} \tilde{p}(s, d, \beta) = p + \frac{\min\{\alpha, 2\gamma\}}{\Theta(1 - \gamma)}.
$$
\n(3.5)

We take d large such that

<span id="page-9-2"></span>
$$
d > \frac{\max\{q, n\}}{2} \Rightarrow \beta_0 := s - \frac{n}{2d} > 0.
$$
 (3.6)

Now we choose s, d, satisfying [\(3.6\)](#page-9-2), and  $\beta$  such that

<span id="page-10-2"></span>
$$
\beta < \alpha_0 := \min\{\alpha, 2\beta_0\} < \min\{\alpha, 2\gamma\} \tag{3.7}
$$

and such that

<span id="page-10-3"></span>
$$
p + \frac{\xi}{1 - \gamma} < \tilde{p}(s, d, \beta) < p + \frac{\min\{\alpha, 2\gamma\}}{1 - \gamma}.\tag{3.8}
$$

Now, we want to approximate, on a fixed ball  $B_r \in \Omega$ , the original minimizer u of F with a sequence  $\{u_i\}$  of  $W^{1,q}$ -regular solutions to approximating variational problems. To find such  $\{u_i\}$ , we define a sequence of suitable functionals  $\mathcal{F}_i$  of mixed local/non-local type as follows. Consider the parameters s,  $\gamma$  and d defined above. For every  $j \in \mathbb{N}$ , let  $\bar{v}_j \in$  $W^{1,2d}(B_r,\mathbb{R}^N)\cap W^{s,2d}(\mathbb{R}^n,\mathbb{R}^N)$  and let introduce the non-local Dirichlet class

$$
\mathbb{X}(\bar{v}_j, B_r) := \left\{ v \in \left( \bar{v}_j + W_0^{1,2d}(B_r, \mathbb{R}^N) \right) \cap W^{s,2d}(\mathbb{R}^n, \mathbb{R}^N) : v \equiv \bar{v}_j \text{ on } \mathbb{R}^n \setminus B_r \right\}.
$$

This subset of  $W^{1,2d}(B_r, \mathbb{R}^N) \cap W^{s,2d}(\mathbb{R}^n, \mathbb{R}^N)$  is convex and closed. Besides, it is non-empty, since  $\bar{v}_j \in \mathbb{X}(\bar{v}_j, B_r)$  by definition. Then, we define

$$
\mathcal{F}_j(w, B_r) := \mathcal{F}(w, B_r) + \varepsilon_j \int_{B_r} (\mu^2 + |Dw|^2)^d dx \n+ \int_{B_r} (|w|^2 - M_0^2)_+^d dx + \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{(|w(x) - w(y)|^2 - M^2 |x - y|^{2\gamma})_+^d}{|x - y|^{n+2sd}} dx dy, (3.9)
$$

for every  $w \in \mathbb{X}(\bar{v}_j, B_r)$  and for some positive parameters  $\varepsilon_j$ ,  $M_0$  and  $M$ . Thanks to assumption [\(1.10\)](#page-3-1), that ensures the absence of the Lavrentiev phenomenon, we are able to prove the existence of good boundary values for which the minimizers  $u_j$  of [\(3.9\)](#page-10-0) converge to u and are bounded in the fractional Sobolev space  $W^{s,2d}$ , exactly as u. Indeed, the following result holds (for the proof we refer to [\[20,](#page-22-3) Sections 3.2 and 3.4]).

<span id="page-10-1"></span>**Proposition 3.1.** There exists a sequence  ${\lbrace \bar{v}_j \rbrace} \in W^{1,2d}(B_r, \mathbb{R}^N) \cap W^{s,2d}(\mathbb{R}^n, \mathbb{R}^N)$  such that  $[\bar{v}_j]_{0,\gamma;\mathbb{R}^n} \leq c_*H$ , with a constant  $c_* := c_*(n,d,\gamma,s)$ . Moreover, choosing in [\(3.9\)](#page-10-0) parameters

<span id="page-10-0"></span>
$$
\begin{cases}\n\varepsilon_j := \frac{1}{\left( \|D\bar{v}_j\|_{L^{2d}(B_r)}^{4d} + j + 1 \right)} \\
M_0 := 16c_*r^{\gamma}H \\
M := 16c_*H\n\end{cases}
$$
\n(3.10)

we have that, for every  $j \in \mathbb{N}$ , there exists  $u_j \in \mathbb{X}(\bar{v}_j, B_r)$  solution to

$$
\mathcal{F}_j(u_j, B_r) \le \mathcal{F}_j(\omega, B_r) \qquad \text{for all } \omega \in \mathbb{X}(\bar{v}_j, B_r)
$$

such that

<span id="page-10-5"></span>
$$
u_j \rightharpoonup u \quad weakly \ in \quad W^{1,p}\big(B_r, \mathbb{R}^N\big) \tag{3.11}
$$

and for every  $j \in \mathbb{N}$  it holds

<span id="page-10-4"></span>
$$
[u_j]_{s,2d;\mathbb{R}^n}^{2d} \le cr^{n+2d(\gamma-s)} H^{2d},\tag{3.12}
$$

for a constant  $c := c(n, p, q, \alpha, \gamma, \sigma)$ . Moreover,

<span id="page-11-2"></span>
$$
\varepsilon_j \int_{B_r} \left(\mu^2 + |Du_j|^2\right)^d \mathrm{d}x \le \mathcal{F}(u, B_r) + 1 \tag{3.13}
$$

and

<span id="page-11-3"></span>
$$
||u_j||_{W^{1,p}(B_r)} \le c[\mathbb{F}(u, B_r)]^{\frac{1}{p}},
$$
\n(3.14)

where

<span id="page-11-1"></span>
$$
\mathbb{F}(u, B_r) := \mathcal{F}(u, B_r) + r^n H^{2d} + 1,
$$
\n(3.15)

for  $c := c(n, N, p, q, \nu, L, \alpha, \gamma, \sigma)$ .

We are now in position to give the proof of the higher integrability result.

*Proof of Theorem [1.2.](#page-3-2)* For every  $j \in \mathbb{N}$ , let  $u_j \in \mathbb{X}(\bar{v}_j, B_r)$  be the minimizer of [\(3.9\)](#page-10-0), where  $\{\bar{v}_j\} \in W^{1,2d}(B_r,\mathbb{R}^N) \cap W^{s,2d}(\mathbb{R}^n,\mathbb{R}^N)$  and parameters  $\varepsilon_j$ ,  $M_0$  and M are as in Proposition [3.1.](#page-10-1) We derive the Euler-Lagrange system of the functional  $\mathcal{F}_j$  defined in [\(3.9\)](#page-10-0). For every  $j \in \mathbb{N}$  and  $t \in (-1,1)$ , we consider the variation  $u_j + t\varphi$ , where  $\varphi \in W_0^{1,2d}$  $b_0^{1,2d}(B_r,\mathbb{R}^N)$  has compact support in  $B_r$ . Note that, from the embedding  $W^{1,2d}(B_r,\mathbb{R}^N) \subset W^{s,2d}(B_r,\mathbb{R}^N)$ , we have  $\varphi \in W^{s,2d}(B_r,\mathbb{R}^N)$ . Moreover, since  $\varphi$  has compact support in  $B_r$ , setting  $\varphi = 0$  outside  $B_r$  yields that  $\varphi \in W^{s,2d}(\mathbb{R}^n,\mathbb{R}^N)$ . Therefore, we have that  $\varphi \in W_0^{1,2d}$  $\chi_0^{1,2d} (B_r,\mathbb{R}^N) \cap W_0^{s,2d}$  $L_0^{s,2d}(\mathbb{R}^n,\mathbb{R}^N)$ and  $u_j + t\varphi \in \mathbb{X}(\bar{v}_j, B_r)$ . By minimality of  $u_j$ , we get

$$
\left. \frac{\mathrm{d} \mathcal{F}_j(u_j + t\varphi, B_r)}{\mathrm{d} t} \right|_{t=0} = 0,
$$

that is

$$
0 = \int_{B_r} \partial_z F_j(x, Du_j) \cdot D\varphi \, dx + 2d \int_{B_r} (|u_j|^2 - M_0^2)_+^{d-1} u_j \cdot \varphi \, dx + \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{(|u_j(x) - u_j(y)|^2 - M^2 |x - y|^{2\gamma})_+^{d-1}}{|x - y|^{n+2sd}} (u_j(x) - u_j(y)) \cdot (\varphi(x) - \varphi(y)) \, dxdy,
$$
(3.16)

where we used the notation

$$
F_j(x, z) := F(x, z) + \varepsilon_j (\mu^2 + z^2)^d.
$$

Now, we fix radii

<span id="page-11-0"></span> $0 < \rho < \tau_1 < \tau_2 < r$ 

and set

$$
\varphi:=\tau_{-h}(\eta^2\tau_h u_j)
$$

with  $\eta \in C_c^1(B_{(3\tau_2+\tau_1)/4})$  verifying  $1_{B_{(\tau_2+\tau_1)/2}} \leq \eta \leq 1_{B_{(3\tau_2+\tau_1)/4}}$  and  $|D\eta| \leq \frac{c}{\tau_2-\tau_1}$ , and  $h \in \mathbb{R}^n \setminus \{0\}$ is such that  $\tau_2 - \tau_1$ 

$$
0<|h|<\frac{\tau_2-\tau_1}{2^{10}}
$$

Testing  $(3.16)$  with  $\varphi$  and using integration by parts for finite difference operators, we obtain

$$
\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{\tau_h[(|u_j(x) - u_j(y)|^2 - M^2|x - y|^{2\gamma})_+^{d-1}(u_j(x) - u_j(y))]}{|x - y|^{n+2sd}} \cdot (\eta^2(x)\tau_h u_j(x) - \eta^2(y)\tau_h u_j(y)) \mathrm{d}x \mathrm{d}y
$$

$$
+2d\int_{B_r} \eta^2 \tau_h((|u_j|^2 - M_0^2)_+^{d-1} u_j) \cdot \tau_h u_j dx + \int_{B_r} \tau_h \partial_z F_j(x, Du_j) \cdot [\eta^2 \tau_h Du_j + 2\eta \tau_h u_j \otimes D\eta] dx := (I) + (II) + (III) = 0.
$$
 (3.17)

Concerning the term  $(I)$ , we take advantage of estimate in [\[20,](#page-22-3) Section 3.6], in particular it holds

<span id="page-12-3"></span>
$$
d\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \int_0^1 \frac{\eta^2(x) B_1(x, y, \lambda)}{|x - y|^{n + 2sd}} d\lambda dx dy \le (I) + \frac{cr^{\frac{n}{d}} \mathbb{F}(u, B_r)}{(\tau_2 - \tau_1)^2} |h|^{2\beta_0},\tag{3.18}
$$

where we defined

$$
B_1(x,y,\lambda):=\left[\frac{d}{d\lambda}(|U_{\lambda,h}(x,y)|^2-M^2|x-y|^{2\gamma})_+^{d-1}U_{\lambda,h}(x,y)\right]\cdot(\tau_hu_j(x)-\tau_hu_j(y)),
$$

with

$$
U_{\lambda,h}(x,y) := u_j(x) - u_j(y) + \lambda \tau_h u_j(x) - \lambda \tau_h u_j(y)
$$

and where **F** was introduced in [\(3.15\)](#page-11-1) and  $\beta_0 = s - n/2d$ . Next, we estimate  $(II)$  as follows

$$
(II) = 2d \int_{B_r} \eta^2 \tau_h u_j \int_0^1 \frac{d}{d\lambda} \left( \left( |u_j + \lambda \tau_h u_j|^2 - M_0^2 \right)_+^{d-1} (u_j + \lambda \tau_h u_j) \right) d\lambda dx
$$
  
=  $4d(d-1) \int_{B_r} \eta^2 \int_0^1 \left( |u_j + \lambda \tau_h u_j|^2 - M^2 \right)_+^{d-2} \left( (u_j + \lambda \tau_h u_j) \cdot \tau_h u_j \right)^2 d\lambda dx$   
+  $2d \int_{B_r} \eta^2 |\tau_h u_j|^2 \int_0^1 \left( |u_j + \lambda \tau_h u_j|^2 - M_0^2 \right)_+^{d-1} d\lambda dx \ge 0.$  (3.19)

Now we take care of the integral  $(III)$ . Recalling the definition  $(3.1)$ , we have

$$
(III) = \int_{B_r} \tau_h \partial_z F(x, Du_j) \cdot [\eta^2 \tau_h Du_j + 2\eta \tau_h u_j \otimes D\eta] dx
$$
  
+ 
$$
2d\varepsilon_j \int_{B_r} ((\mu^2 + |Du_j|^2)^{d-1} Du_j) \cdot [\eta^2 \tau_h Du_j + 2\eta \tau_h u_j \otimes D\eta] dx =: (III)_1 + (III)_2.
$$
  
(3.20)

For  $(III)_1$ , we decompose

$$
(III)_1 = \int_{B_r} [\partial_z F(x+h, Du_j(x+h)) - \partial_z F(x+h, Du_j(x))] \cdot (\eta^2 \tau_h Du_j + 2\eta \tau_h u_j \otimes D\eta) dx + \int_{B_r} [\partial_z F(x+h, Du_j(x)) - \partial_z F(x, Du_j(x))] \cdot (2\eta \tau_h u_j \otimes D\eta + \eta^2 \tau_h Du_j) dx =: (III)_{1,1} + (III)_{1,2} + (III)_{1,3} + (III)_{1,4}
$$
 (3.21)

By the ellipticity assumption (F2), we have

<span id="page-12-2"></span><span id="page-12-1"></span><span id="page-12-0"></span>
$$
(III)_{1,1} \ge \frac{1}{c} \int_{B_r} \eta^2 (\mu^2 + |Du_j(x+h)|^2 + |Du_j(x)|^2)^{\frac{p-2}{2}} |\tau_h Du_j|^2 dx.
$$
 (3.22)

Using growth assumption (F1) and then Hölder's inequality, we obtain

$$
(III)_{1,2} + (III)_{1,3} \leq c \int_{B_r} \eta (1 + |Du_j(x+h)|^2 + |Du_j(x)|^2)^{\frac{q-1}{2}} |D\eta| |\tau_h u_j| \, dx
$$
  
\n
$$
\leq c ||D\eta||_{L^{\infty}} \left( \int_{B_r} \eta (1 + |Du_j(x+h)|^2 + |Du_j(x)|^2)^{\frac{q}{2}} \, dx \right)^{\frac{q-1}{q}}
$$
  
\n
$$
\cdot \left( \int_{B_r} \eta |\tau_h u_j|^q \, dx \right)^{\frac{1}{q}}
$$
  
\n
$$
\leq \frac{c}{\tau_2 - \tau_1} |h| \int_{B_{\tau_2}} (1 + |Du_j|^2)^{\frac{q}{2}} \, dx
$$
  
\n
$$
\leq \frac{c}{\tau_2 - \tau_1} |h| \int_{B_{\tau_2}} (1 + |Du_j|)^{p + \frac{\xi}{1 - \gamma}} \, dx,
$$
\n(3.23)

where  $\xi$  was introduced in [\(1.9\)](#page-3-0) and we have also exploited the bounds  $q < p + \frac{\xi}{1-\xi}$  $\frac{\xi}{1-\gamma}$  and  $(1.7)$ and hypotheses on  $\eta$  and  $D\eta$ .

According to (F3) we can estimate  $(III)_{1,4}$  as

<span id="page-13-1"></span><span id="page-13-0"></span>
$$
(III)_{1,4} \le c|h|^{\alpha} \int_{B_r} \eta^2 (g(x+h) + g(x))(1+|Du_j|^2)^{\frac{q}{2}} dx.
$$

We now use Hölder's inequality with exponents  $\sigma$  and  $\frac{\sigma}{\sigma}$  $\sigma-1$ , where  $\sigma$  was introduced in [\(1.9\)](#page-3-0). Then, by [\(1.7\)](#page-3-3) and Young's inequality we infer

$$
(III)_{1,4} \le |h|^{\alpha} \left( \int_{B_r} g^{\sigma} dx \right)^{\frac{1}{\sigma}} \left( \int_{B_{\tau_2}} (1 + |Du_j|^{p + \frac{\xi}{1 - \gamma}}) dx \right)^{\frac{q(1 - \gamma)}{p(1 - \gamma) + \xi}}.
$$
 (3.24)

We notice that in the last in integral in the right hand side, we exploited the fact that the choices of the parameters ensure  $\frac{\sigma}{\sigma-1}q = p + \frac{\xi}{1-\xi}$  $\frac{\xi}{1-\gamma}$ . Putting [\(3.22\)](#page-12-1), [\(3.23\)](#page-13-0) and [\(3.24\)](#page-13-1) in [\(3.21\)](#page-12-2), we get

$$
(III)_1 \ge \frac{1}{c} \int_{B_r} \eta^2 (\mu^2 + |Du_j(x+h)|^2 + |Du_j(x)|^2)^{\frac{p-2}{2}} |\tau_h Du_j|^2 dx
$$

$$
- c \int_{B_r} g^{\sigma} dx - \frac{c}{\tau_2 - \tau_1} |h|^{\alpha} \int_{B_{\tau_2}} (1 + |Du_j|)^{p + \frac{\xi}{1 - \gamma}} dx.
$$
(3.25)

On the other hand, similarly we might estimate

$$
(III)_2 \ge \frac{\varepsilon_j}{c} \int_{B_r} \eta^2 \left(\mu^2 + |Du_j(x+h)|^2 + |Du_j(x)|^2\right)^{d-1} |\tau_h Du_j|^2 dx
$$

$$
- \frac{c\varepsilon_j|h|}{\tau_2 - \tau_1} \int_{B_{\tau_2}} \left(1 + |Du_j|^2\right)^d dx,
$$
(3.26)

where we recall that  $\varepsilon_j$  was defined in Proposition [3.1.](#page-10-1) Using the estimates for  $(I), (II)$  and  $(III)$  in [\(3.17\)](#page-12-3) and exploiting [\(3.7\)](#page-10-2) and the hypotheses on  $D\eta$  together with [\(3.13\)](#page-11-2) and [\(3.15\)](#page-11-1), we conclude

$$
\int_{B_{\frac{\tau_2+\tau_1}{2}}} \int_{B_{\frac{\tau_2+\tau_1}{2}}} \int_0^1 \frac{B_1(x,y,\lambda)}{|x-y|^{n+2sd}} d\lambda dx dy
$$

$$
+\int_{B_{\frac{\tau_2+\tau_1}{2}}} (\mu^2 + |Du_j(x+h)|^2 + |Du_j(x)|^2)^{\frac{p-2}{2}} |\tau_h Du_j|^2 dx
$$
  

$$
\leq \frac{c}{(\tau_2-\tau_1)^2} |h|^{\alpha_0} \left( \mathbb{F}(u, B_r) + ||g||_{L^{\sigma}(B_r)}^{\sigma} + ||Du_j||_{L^{p+\frac{\xi}{1-\gamma}}(B_{\tau_2})}^{\sigma} + 1 \right),
$$
(3.27)

for a constant  $c := c(n, N, \nu, L, p, q)$  and where  $\alpha_0$  is defined in [\(3.7\)](#page-10-2). From the definition of  $B_1$ , we can easily derive that  $B_1(x, y, \lambda) \geq 0$ . Hence, we can discard the first integral on the left hand side of [\(3.27\)](#page-14-0) and obtain

<span id="page-14-1"></span><span id="page-14-0"></span>
$$
\int_{B_{\frac{\tau_2+\tau_1}{2}}} (\mu^2 + |Du_j(x+h)|^2 + |Du_j(x)|^2)^{\frac{p-2}{2}} |\tau_h Du_j|^2 dx
$$
\n
$$
\leq \frac{c}{(\tau_2 - \tau_1)^2} |h|^{\alpha_0} \left( \tilde{\mathbb{F}}(u, B_r) + ||Du_j||_{L^{p+\frac{\xi}{1-\gamma}}(B_{\tau_2})}^{p+\frac{\xi}{1-\gamma}} + 1 \right),
$$
\n(3.28)

where we set  $\tilde{\mathbb{F}}(u, B_r) := \mathbb{F}(u, B_r) + ||g||_{L^{\sigma}(B_r)}^{\sigma}.$ 

Now, we distinguish the case  $p \geq 2$  and the case  $1 < p < 2$ . Let  $p \geq 2$ . Then, from [\(3.28\)](#page-14-1) we obtain

$$
\int_{B_{\frac{\tau_2+\tau_1}{2}}} |\tau_h D u_j|^p \le \frac{c}{(\tau_2-\tau_1)^2} |h|^{\alpha_0} \left( \tilde{\mathbb{F}}(u, B_r) + \|D u_j\|_{L^{p+\frac{\xi}{1-\gamma}}(B_{\tau_2})}^{p+\frac{\xi}{1-\gamma}} + 1 \right). \tag{3.29}
$$

From Lemma [2.9,](#page-8-1) since  $0 < |h| < \frac{\tau_2 - \tau_1}{\sigma^{10}}$  $\frac{1}{2^{10}}$ , according to [\(3.29\)](#page-14-2) we read the following regularity of  $u_i$ 

<span id="page-14-2"></span>
$$
u_j \in W^{1+\frac{\beta}{p},p}(B_{\frac{\tau_2+\tau_1}{2}},\mathbb{R}^n) \cap W^{s,2d}(B_r,\mathbb{R}^n),
$$

for  $\beta \in (0, \alpha_0)$  satisfying [\(3.8\)](#page-10-3). Hence, by Lemma [2.9](#page-8-1) and inequality [\(3.14\)](#page-11-3), we infer

$$
||u_j||_{W^{1+\frac{\beta}{p},p}\left(B_{\frac{\tau_2+\tau_1}{2}}\right)} \leq \frac{c}{(\tau_2-\tau_1)^{\delta}}||u_j||_{W^{1,p}(B_{\tau_2})} + \frac{c}{(\tau_2-\tau_1)^{\frac{2}{p}+\delta}}\left([\tilde{\mathbb{F}}(u,B_r)]^{\frac{1}{p}}+||Du_j||_{L^{p+\frac{\xi}{1-\gamma}}(B_{\tau_2})}^{1+\frac{\xi}{p-p\gamma}}+1\right) \leq \frac{c}{(\tau_2-\tau_1)^{\frac{2}{p}+\delta}}\left([\tilde{\mathbb{F}}(u,B_r)]^{\frac{1}{p}}+||Du_j||_{L^{p+\frac{\xi}{1-\gamma}}(B_{\tau_2})}^{1+\frac{\xi}{p-p\gamma}}+1\right),
$$
(3.30)

with  $c, \delta = c, \delta(n, p, \beta, \alpha_0)$ . Now, applying Lemma [2.10,](#page-9-3) we get

<span id="page-14-4"></span>
$$
||Du_j||_{L^{\tilde{p}}(B_{\tau_1})} \leq \frac{c}{(\tau_2 - \tau_1)^{\kappa}} [u_j]_{s,2d,B_{\tau_2}}^{\theta} ||Du_j||_{W^{\frac{\beta}{p},p}\left(B_{\frac{\tau_2 + \tau_1}{2}}\right)}^{1-\theta},\tag{3.31}
$$

for  $\theta =$ β  $p(1-s)+\beta$  $\in (0, 1)$ , where c and  $\kappa$  are the constants from Lemma [2.10.](#page-9-3) Moreover, we observe that from [\(3.12\)](#page-10-4) it follows

<span id="page-14-5"></span><span id="page-14-3"></span>
$$
[u_j]_{s,2d,\mathbb{R}^n} \le c [\mathbb{F}(u, B_r)]^{\frac{1}{2d}}.
$$
\n(3.32)

Therefore, combining  $(3.30)$ ,  $(3.31)$  and  $(3.32)$ , we obtain

$$
||Du_j||_{L^{\tilde{p}}(B_{\tau_1})} \leq \frac{c}{(\tau_2 - \tau_1)^{\frac{2}{p} + \delta + \kappa}} [\tilde{F}(u, B_r)]^{\frac{1-\theta}{p}} [\mathbb{F}(u, B_r)]^{\frac{\theta}{2d}} + \frac{c}{(\tau_2 - \tau_1)^{\frac{2}{p} + \delta + \kappa}} [\mathbb{F}(u, B_r)]^{\frac{\theta}{2d}} ||Du_j||^{\frac{(p + \frac{\xi}{1 - \gamma})^{\frac{1-\theta}{p}}}{L^{p + \frac{\xi}{1 - \gamma}}(B_{\tau_2})} + \frac{c}{(\tau_2 - \tau_1)^{\frac{2}{p} + \delta + \kappa}} [\mathbb{F}(u, B_r)]^{\frac{\theta}{2d}}.
$$
(3.33)

We now recall that by assumption (see  $(3.2)$  and  $(3.8)$ )

<span id="page-15-4"></span><span id="page-15-1"></span>
$$
p < p + \frac{\xi}{1 - \gamma} < \tilde{p}.\tag{3.34}
$$

Therefore, we can use the following interpolation inequality

<span id="page-15-0"></span>
$$
||Du_j||_{L^{p+\frac{\xi}{1-\gamma}}} \le ||Du_j||_{L^{\tilde{p}}}^{\theta_1} ||Du_j||_{L^p}^{1-\theta_1},
$$
\n(3.35)

where  $\theta_1 \in (0, 1)$  is such that

<span id="page-15-2"></span>
$$
\frac{1-\gamma}{p(1-\gamma)+\xi} = \frac{\theta_1}{\tilde{p}} + \frac{1-\theta_1}{p},
$$
  

$$
\theta_1 = \frac{\xi \tilde{p}}{(\tilde{p}-p)(p(1-\gamma)+\xi)}.
$$
 (3.36)

hence

Inserting  $(3.35)$  in  $(3.33)$ , we infer

$$
||Du_j||_{L^{\tilde{p}}(B_{\tau_1})} \leq c [\tilde{F}(u, B_r)]^{\frac{1-\theta}{p}} [\mathbb{F}(u, B_r)]^{\frac{\theta}{2d}} + c [\mathbb{F}(u, B_r)]^{\frac{\theta}{2d}} ||Du_j||_{L^{\tilde{p}}(B_{\tau_2})}^{\theta_1(p+\frac{\xi}{1-\gamma})\frac{1-\theta}{p}} ||Du_j||_{L^p(B_{\tau_2})}^{(1-\theta_1)(p+\frac{\xi}{1-\gamma})\frac{1-\theta}{p}} + c [\mathbb{F}(u, B_r)]^{\frac{\theta}{2d}}
$$

Recalling the definition of  $\tilde{p}$  in [\(3.2\)](#page-9-4), we note that when  $p \geq 2$ , it holds

$$
\lim_{s \to \gamma, d \to \infty, \beta \to \min\{\alpha, 2\gamma\}} \frac{\tilde{p}\xi}{\tilde{p} - p} \frac{1}{1 - \gamma} \frac{1 - s}{p(1 - s) + \beta}
$$
\n
$$
\lim_{s \to \gamma, d \to \infty, \beta \to \min\{\alpha, 2\gamma\}} \frac{2d[p(1 - s) + \beta]}{\beta + 2d(1 - s)} \frac{\beta + 2d(1 - s)}{2d[p(1 - s) + \beta] - p\beta - 2pd(1 - s)}
$$
\n
$$
\cdot \frac{\xi(1 - s)}{(1 - \gamma)[p(1 - s) + \beta]}
$$
\n
$$
= \lim_{s \to \gamma, d \to \infty, \beta \to \min\{\alpha, 2\gamma\}} \frac{2d\xi}{1 - \gamma} \frac{1 - s}{\beta(2d - p)} = \frac{\xi}{\min\{\alpha, 2\gamma\}} < 1,
$$

where we remark that  $\frac{\xi}{\min\{\alpha,2\gamma\}}$  can be estimated by 1 according to (F3). Therefore, we increase  $s, d, \beta$  in order to have

<span id="page-15-3"></span>
$$
\frac{\tilde{p}\xi}{\tilde{p}-p}\frac{1}{1-\gamma}\frac{1-s}{p(1-s)+\beta} < 1. \tag{3.37}
$$

Now, owing to  $(1.8)$  and  $(3.36)$ ,

$$
\theta_1\left(p+\frac{\xi}{1-\gamma}\right)\frac{1-\theta}{p}=\frac{\tilde{p}\xi}{\tilde{p}-p}\frac{1}{1-\gamma}\frac{1-s}{p(1-s)+\beta}<1
$$

where we exploited [\(3.37\)](#page-15-3). Applying Young's inequality and reabsorbing the term  $||Du_i||_{L^{\tilde{p}}}$ according to Lemma [2.2,](#page-5-1) we get

$$
||Du_j||_{L^{\tilde{p}}(B_{\tau_1})}\leq \frac{c}{(\tau_2-\tau_1)^{\kappa_1}}[\tilde{\mathbb{F}}(u,B_r)]^{\kappa_2},
$$

for some exponents  $\kappa_1$  and  $\kappa_2$ . From [\(3.11\)](#page-10-5) and weak lower semicontinuity, letting  $j \to +\infty$  we get

$$
Du\in L^{\tilde{p}}(B_{\tau_1}),
$$

hence

$$
Du \in L^{\tilde{q}}_{\text{loc}}(\Omega),
$$

for every  $q < \tilde{q} < p +$  $\min\{\alpha, 2\gamma\}$  $1-\gamma$ . This proves the theorem in the case  $p \geq 2$ . For the higher integrability in the case  $1 < p < 2$  we proceed analogously. Being  $\tilde{p}$  defined as in [\(3.2\)](#page-9-4), for  $1 < p < 2$  it holds

$$
\lim_{s \to \gamma, d \to \infty, \beta \to \min\{\alpha, 2\gamma\}} \frac{\tilde{p}\xi}{\tilde{p} - p} \frac{1}{1 - \gamma} \frac{2(1 - s)}{p[2(1 - s) + \beta]}
$$
\n
$$
= \lim_{s \to \gamma, d \to \infty, \beta \to \min\{\alpha, 2\gamma\}} \frac{\xi}{1 - \gamma} \frac{4d(1 - s)}{p\beta(2d - p)} = \frac{2\xi}{\min\{\alpha, 2\gamma\}} < 1,
$$

where we used the assumption  $\xi < \frac{\min\{\alpha, 2\gamma\}}{\Omega}$ Θ and the definition of  $\Theta$  in [\(1.8\)](#page-3-4) for  $1 < p < 2$ . Hence, we can find  $s, d, \beta$  sufficiently large, such that

<span id="page-16-0"></span>
$$
\frac{\tilde{p}\xi}{(\tilde{p}-p)(1-\gamma)}\frac{2(1-s)}{p[2(1-s)+\beta]} < 1. \tag{3.38}
$$

Now, we note that

$$
\int_{B_{\frac{\tau_2+\tau_1}{2}}} |\tau_h Du_j|^p \, dx = \int_{B_{\frac{\tau_2+\tau_1}{2}}} |\tau_h Du_j|^p (\mu^2 + |Du_j(x+h)|^2 + |Du_j(x)|^2)^{\frac{(p-2)p}{4}} \cdot (\mu^2 + |Du_j(x+h)|^2 + |Du_j(x)|^2)^{\frac{(2-p)p}{4}} \, dx.
$$

Using Hölder's inequality we derive

$$
\int_{B_{\frac{\tau_2+\tau_1}{2}}} |\tau_h Du_j|^p \, dx \le \left( \int_{B_{\frac{\tau_2+\tau_1}{2}}} (\mu^2 + |Du_j(x+h)|^2 + |Du_j(x)|^2)^{\frac{p-2}{2}} |\tau_h Du_j|^2 \, dx \right)^{\frac{p}{2}}
$$

$$
\cdot \left( \int_{B_{\frac{\tau_2+\tau_1}{2}}} (\mu^2 + |Du_j(x+h)|^2 + |Du_j(x)|^2)^{\frac{p}{2}} \, dx \right)^{1-\frac{p}{2}}.
$$

Inserting estimates [\(3.14\)](#page-11-3) and [\(3.28\)](#page-14-1) above, we obtain

$$
\int_{B_{\frac{r_2+r_1}{2}}} |\tau_h Du_j|^p dx \le \left( \int_{B_{\frac{r_2+r_1}{2}}} (\mu^2 + |Du_j(x+h)|^2 + |Du_j(x)|^2)^{\frac{p}{2}} dx \right)^{\frac{p}{2}} [\mathbb{F}(u, B_r)]^{1-\frac{p}{2}}
$$
  

$$
\le \frac{c}{(\tau_2 - \tau_1)^p} |h|^{\frac{\alpha_0 p}{2}} \left( \mathbb{F}(u, B_r) + ||Du_j||_{L^{p+\frac{\xi}{1-\gamma}}(B_{\tau_2})}^{p+\frac{\xi}{1-\gamma}} + 1 \right).
$$

Hence, from Lemma [2.9](#page-8-1) and  $0 < |h| < \frac{\tau_2 - \tau_1}{2.010}$  $\frac{1}{2^{10}}$ , we infer

$$
u_j \in W^{1+\frac{\beta}{2},p}(B_{\frac{\tau_2+\tau_1}{2}},\mathbb{R}^n) \cap W^{s,2d}(B_r,\mathbb{R}^n),
$$

for  $\beta \in (0, \alpha_0)$  satisfying [\(3.8\)](#page-10-3) and [\(3.38\)](#page-16-0). Therefore, from Lemma [2.10,](#page-9-3) it follows

$$
||u_j||_{W^{1+\frac{\beta}{p},p}\left(B_{\frac{\tau_2+\tau_1}{2}}\right)} \leq \frac{c}{(\tau_2-\tau_1)^{\delta}}||u_j||_{W^{1,p}(B_{\tau_2})} +\frac{1}{(\tau_2-\tau_1)^{1+\delta}}\left([\tilde{\mathbb{F}}(u,B_r)]^{\frac{1}{p}}+||Du_j||_{L^{p+\frac{\xi}{1-\gamma}}(B_{\tau_2})}^{p+\frac{\xi}{1-\gamma}}+1\right) \leq \frac{1}{(\tau_2-\tau_1)^{1+\delta}}\left([\tilde{\mathbb{F}}(u,B_r)]^{\frac{1}{p}}+||Du_j||_{L^{p+\frac{\xi}{1-\gamma}}(B_{\tau_2})}^{p+\frac{\xi}{1-\gamma}}+1\right).
$$

Now, applying Lemma [2.10,](#page-9-3) we get

$$
||Du_j||_{L^{\tilde{p}}(B_{\tau_1})}\leq \frac{c}{(\tau_2-\tau_1)^{\kappa}}[u_j]_{s,2d,B_{\tau_2}}^{\theta}||u_j||_{W^{\frac{\beta}{p},p}\left(B_{\frac{\tau_2+\tau_1}{2}}\right)}^{1-\theta},
$$

where in this case  $\theta =$ β  $2(1-s)+\beta$  $\in (0, 1)$ . By  $(3.32)$ , we thus have

<span id="page-17-0"></span>
$$
||Du_j||_{L^{\tilde{p}}(B_{\tau_1})} \leq \frac{c}{(\tau_2 - \tau_1)^{1+\delta+\kappa}} [\tilde{F}(u, B_r)]^{\frac{1-\theta}{p}} [\mathbb{F}(u, B_r)]^{\frac{\theta}{2d}} + \frac{c}{(\tau_2 - \tau_1)^{1+\delta+\kappa}} [\mathbb{F}(u, B_r)]^{\frac{\theta}{2d}} ||Du_j||^{\frac{(p+\frac{\xi}{1-\gamma})^{\frac{1-\theta}{p}}}{L^{p+\frac{\xi}{1-\gamma}}(B_{\tau_2})}}_{L^{p+\frac{\xi}{1-\gamma}}(B_{\tau_2})}
$$
(3.39)

By [\(3.34\)](#page-15-4), we are able to exploit the interpolation inequality [\(3.35\)](#page-15-0), where  $\theta_1$  is defined in the same way, with the difference that the expression of  $\tilde{p}$  is different owing to [\(3.2\)](#page-9-4).

Then, inserting [\(3.35\)](#page-15-0) in [\(3.39\)](#page-17-0) implies

$$
||Du_j||_{L^{\tilde{p}}(B_{\tau_1})} \leq c [\tilde{F}(u, B_r)]^{\frac{1-\theta}{p}} [\mathbb{F}(u, B_r)]^{\frac{\theta}{2d}} + c [\mathbb{F}(u, B_r)]^{\frac{\theta}{2d}} ||Du_j||_{L^{\tilde{p}}(B_{\tau_2})}^{\theta_1(p+\frac{\xi}{1-\gamma})\frac{1-\theta}{p}} ||Du_j||_{L^p(B_{\tau_2})}^{(1-\theta_1)(p+\frac{\xi}{1-\gamma})\frac{1-\theta}{p}} + c [\mathbb{F}(u, B_r)]^{\frac{\theta}{2d}}.
$$

Now we observe that, by  $(3.36)$  and  $(3.38)$  we have

$$
\theta_1\left(p+\frac{\xi}{1-\gamma}\right)\frac{1-\theta}{p}=\frac{\tilde{p}\xi}{\tilde{p}-p}\frac{1}{1-\gamma}\frac{1-s}{p(1-s)+\beta}<1.
$$

Applying Young's inequality and reabsorbing the term  $||Du_j||_{L^{\tilde{p}}}$ , we get

$$
||Du_j||_{L^{\tilde{p}}(B_{\tau_1})}\leq c[\mathbb{F}(u,B_r)]^{\kappa_1}[\tilde{\mathbb{F}}(u,B_r)]^{\kappa_2}.
$$

Then, from [\(3.11\)](#page-10-5) and weak lower semicontinuity, letting  $j \to +\infty$  we can conclude that

$$
Du \in L^{\tilde{q}}_{\text{loc}}(\Omega), \quad \text{for every } q < \tilde{q} < p + \frac{\min\{\alpha, 2\gamma\}}{\Theta(1-\gamma)}.
$$

This completes the proof of the theorem.

#### 3.2 Higher differentiability

In this section, we give the proof of Theorem [1.4](#page-4-0) taking advantage of the higher integrability result proved in Section [3.1.](#page-9-0) According to Theorem [1.2,](#page-3-2) we deduce that

<span id="page-18-0"></span>
$$
Du \in L_{loc}^{p+\frac{\xi}{1-\gamma}}(\Omega, \mathbb{R}^{N \times n}),\tag{3.40}
$$

where  $\xi$  was introduced in equation [\(1.9\)](#page-3-0), hence  $q < p + \frac{\xi}{1-\xi}$  $\frac{\xi}{1-\gamma}$ . This implies that we do not need the approximating problems to study the higher fractional differentiability of  $Du$ .

Proof of Theorem [1.4.](#page-4-0) According to [\(3.40\)](#page-18-0), a solution to problem [\(1.1\)](#page-0-0) satisfies  $u \in W^{1,q}_{loc}(\Omega, R^N)$ , therefore u solves the equation

<span id="page-18-3"></span>
$$
\int_{B_R} \partial_z F(x, Du) \cdot D\varphi \, dx = 0, \quad \forall \varphi \in W_0^{1,q}(B_R, \mathbb{R}^N). \tag{3.41}
$$

Let us fix a ball  $B_R$  such that  $B_{2R} \in \Omega$  and a cut-off function  $\eta \in C_0^1(B_R)$  such that  $0 \leq \eta \leq 1$ ,  $\eta = 1$  on  $B_{R/2}$  and  $|D\eta| \leq \frac{C}{R}$ . Now, for  $|h| \leq R/4$ , we consider test functions

<span id="page-18-2"></span><span id="page-18-1"></span>
$$
\varphi_1(x) = \eta^2(x)\tau_h u(x) \tag{3.42}
$$

and

<span id="page-18-4"></span>
$$
\varphi_2(x) = \eta^2(x - h)\tau_{-h}u(x). \tag{3.43}
$$

Inserting  $(3.42)$  and  $(3.43)$  in  $(3.41)$ , we obtain

$$
\int_{\Omega} \partial_z F(x, Du(x)) \cdot D(\eta^2(x)\tau_h u(x)) \, \mathrm{d}x + \int_{\Omega} \partial_z F(x, Du(x)) \cdot D(\eta^2(x - h)\tau_{-h} u(x)) \, \mathrm{d}x = 0. \tag{3.44}
$$

By means of a simple change of variable, we can write the second integral on the left hand side of the previous inequality as follows

$$
\int_{\Omega} \partial_z F(x+h, Du(x+h)) \cdot D(-\eta^2(x)\tau_h u(x)) \, \mathrm{d}x \tag{3.45}
$$

and so inequality [\(3.44\)](#page-18-4) becomes

$$
\int_{\Omega} \left[\partial_z F(x+h, Du(x+h)) - \partial_z F(x, Du(x))\right] \cdot D(\eta^2 \tau_h u(x)) \, \mathrm{d}x = 0 \tag{3.46}
$$

We can write previous inequality as follows

$$
0 = \int_{\Omega} \left[ \partial_z F(x+h, Du(x+h)) - \partial_z F(x+h, Du(x)) \right] \cdot \eta^2(x) D\tau_h u(x) dx
$$
  
+ 
$$
\int_{\Omega} \left[ \partial_z F(x+h, Du(x+h)) - \partial_z F(x+h, Du(x)) \right] \cdot 2\eta(x) D\eta(x) \tau_h u(x) dx
$$
  
+ 
$$
\int_{\Omega} \left[ \partial_z F(x+h, Du(x)) - \partial_z F(x, Du(x)) \right] \cdot \eta^2(x) D\tau_h u(x) dx
$$
  
+ 
$$
\int_{\Omega} \left[ \partial_z F(x+h, Du(x)) - \partial_z F(x, Du(x)) \right] \cdot 2\eta(x) D\eta(x) \tau_h u(x) dx
$$
  
=:  $I_1 + I_2 + I_3 + I_4$ , (3.47)

that yields

<span id="page-19-2"></span><span id="page-19-1"></span><span id="page-19-0"></span>
$$
I_1 \leq |I_2| + |I_3| + |I_4|. \tag{3.48}
$$

The ellipticity assumption (F2) and Lemma [2.1](#page-5-2) imply

$$
I_1 \ge \nu \int_{\Omega} \eta^2(x) |\tau_h Du|^2 (\mu^2 + |Du(x+h)|^2 + |Du(x)|^2)^{\frac{p-2}{2}} dx
$$
  
 
$$
\ge C(\nu) \int_{B_{R/2}} |\tau_h V_p (Du)|^2 dx,
$$
 (3.49)

where in the last inequality we used that  $\eta = 1$  on the ball  $B_{R/2}$ .

From the growth condition (F4), Hölder's inequality and Lemma [2.4,](#page-6-1) we get

$$
|I_2| \le 2L \int_{\Omega} |D\eta(x)|\eta(x)(1+|Du(x+h)|^2 + |Du(x)|^2)^{\frac{q-1}{2}}|\tau_h u(x)| \, dx
$$
  
\n
$$
\le \frac{C(L)}{R} \left(\int_{B_R} |\tau_h u(x)|^q \, dx\right)^{\frac{1}{q}} \left(\int_{B_{2R}} (1+|Du|)^q \, dx\right)^{\frac{q-1}{q}}
$$
  
\n
$$
\le \frac{C(L)}{R} |h| \int_{B_{2R}} (1+|Du|)^q \, dx. \tag{3.50}
$$

In order to estimate the integral  $I_3$ , we use assumption  $(F3)$ , Hölder's inequality as follows

$$
|I_3| \leq \int_{\Omega} \eta^2 |\tau_h Du||h|^\alpha (g(x+h) + g(x))(1 + |Du(x)|^2)^{\frac{q-1}{2}} dx
$$

$$
\leq |h|^{\alpha} \left( \int_{B_{2R}} g^{\sigma} dx \right)^{\frac{1}{\sigma}} \left( \int_{B_{R}} |\tau_{h} Du|^{\frac{\sigma}{\sigma-1}} (1+|Du|)^{\frac{(q-1)\sigma}{\sigma-1}} dx \right)^{\frac{\sigma-1}{\sigma}}
$$

$$
\leq |h|^{\alpha} \left( \int_{B_{2R}} g^{\sigma} dx \right)^{\frac{1}{\sigma}} \left( \int_{B_{2R}} (1+|Du|)^{\frac{q\sigma}{\sigma-1}} dx \right)^{\frac{\sigma-1}{\sigma}} . \tag{3.51}
$$

We note that

<span id="page-20-1"></span><span id="page-20-0"></span>
$$
\frac{q\sigma}{\sigma - 1} = p + \frac{\xi}{1 - \gamma},
$$

which ensures that the second integral on the right hand side of the previous inequality is finite. Arguing analogously, we infer the following estimate for the integral I4.

$$
|I_{4}| \leq \frac{C}{R} |h|^{\alpha} \int_{B_{R}} |\tau_{h}u|(g(x+h) + g(x))(1 + |Du|^{2})^{\frac{q-1}{2}} dx
$$
  
\n
$$
\leq \frac{C}{R} |h|^{\alpha} \left( \int_{B_{2R}} g^{\sigma} dx \right)^{\frac{1}{\sigma}} \left( \int_{B_{R}} |\tau_{h}u|^{\frac{q\sigma}{\sigma-1}} dx \right)^{\frac{\sigma-1}{q\sigma}} \left( \int_{B_{R}} (1 + |Du|)^{\frac{q\sigma}{\sigma-1}} dx \right)^{\frac{(q-1)(\sigma-1)}{q\sigma}}
$$
  
\n
$$
\leq \frac{C}{R} |h|^{\alpha+1} \left( \int_{B_{2R}} g^{\sigma} dx \right)^{\frac{1}{\sigma}} \left( \int_{B_{2R}} (1 + |Du|)^{\frac{q\sigma}{\sigma-1}} dx \right)^{\frac{\sigma-1}{\sigma}}
$$
(3.52)

Inserting estimates  $(3.49)$ ,  $(3.50)$ ,  $(3.51)$  and  $(3.52)$  in  $(3.48)$ , we infer

$$
C(\nu) \int_{B_{R/2}} |\tau_h V_p(Du)|^2 dx
$$
  
\n
$$
\leq \frac{C(L)}{R} |h| \int_{B_{2R}} (1+|Du|)^q dx
$$
  
\n
$$
+ |h|^{\alpha} \left( \int_{B_{2R}} g^{\sigma} dx \right)^{\frac{1}{\sigma}} \left( \int_{B_{2R}} (1+|Du|)^{\frac{q\sigma}{\sigma-1}} dx \right)^{\frac{\sigma-1}{\sigma}}
$$
  
\n
$$
+ \frac{C}{R} |h|^{\alpha+1} \left( \int_{B_{2R}} g^{\sigma} dx \right)^{\frac{1}{\sigma}} \left( \int_{B_{2R}} (1+|Du|)^{\frac{q\sigma}{\sigma-1}} dx \right)^{\frac{\sigma-1}{\sigma}}, \qquad (3.53)
$$

that yields

$$
\int_{B_{R/2}} |\tau_h V_p(Du)|^2 dx \le C|h|^\alpha \left\{ \int_{B_{2R}} g^\sigma dx + \int_{B_{2R}} (1 + |Du|)^{p + \frac{\xi}{1 - \gamma}} dx \right\},\tag{3.54}
$$

for a constant  $C := C(p, q, \gamma, \sigma, L, \nu, R)$ .

### Acknowledgements

The authors would like to thank Prof. Giuseppe Mingione for fruitful discussion. The authors thank the anonymous referee for the careful reading of the manuscript and for valuable comments. The authors are supported by GNAMPA (Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni) of INdAM (Istituto Nazionale di Alta Matematica). A.G. Grimaldi has been partially supported by the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM) through INdAM-GNAMPA project (CUP\_E53C22001930001).

## <span id="page-21-11"></span>References

- <span id="page-21-4"></span>[1] E. Acerbi, G. Bouchitté and I. Fonseca, Relaxation of convex functionals: the gap problem, Ann. Inst. H. Poincaré Anal. Non Linéaire 20 (2003), 359–390
- <span id="page-21-5"></span>[2] P. Ambrosio, Besov regularity for a class of singular or degenerate elliptic equations, J. Math. anal. Appl. 505(2) (2022)
- [3] P. Ambrosio, Fractional Sobolev regularity for solutions to a strongly degenerate parabolic equation, Forum Math.. https://doi.org/10.1515/forum-2022-0293
- <span id="page-21-6"></span>[4] A. L. Baison, A. Clop, R. Giova, J. Orbitg and A. Passarelli di Napoli, Fractional differentiability for solutions of nonlinear elliptic equations, *Potential Anal.* 46 (3) (2017), 403–430
- <span id="page-21-3"></span><span id="page-21-0"></span>[5] A. Balci, L. Diening and M. Surnachev, New examples on Lavrentiev gap using fractals, Calc. Var. PDE, 59 (2020) 180
- [6] P. Baroni, M. Colombo and G. Mingione, Regularity for general functionals with double phase, Calc. Var. PDE, 57 (2018) 62
- <span id="page-21-8"></span>[7] S.S. Byun, Y. Cho and J. Ok, Global gradient estimates for nonlinear obstacle problems with non-standard growth, Forum Math.  $28(4)$  (2016), 729–747
- [8] S.s. Byun and C. Namkyeong, Higher differentiability for solutions of a general class of nonlinear elliptic obstacle problems with Orlicz growth, NoDEA 29(73) (2022). https://doi.org/10.1007/s00030-022-00807-x
- <span id="page-21-1"></span>[9] M. Carozza, J. Kristensen, and A. Passarelli di Napoli, Higher differentiability of minimizers of convex variational integrals, Ann. Inst. H. Poincaré Anal. Non Linéaire, 28 (2011), 395– 411
- [10] M. Carozza, J. Kristensen, and A. Passarelli di Napoli, Regularity of minimizers of autonomous convex variational integrals, Ann. Sc. Norm. Super. Pisa Cl. Sci.  $13(4)$  (2014), 1065–1089
- [11] M. Carozza, J. Kristensen, and A. Passarelli di Napoli, On the validity of the Euler-Lagrange system, *Comm. Pure Appl. Anal.*,  $14(1)$  (2018),  $51-62$
- <span id="page-21-9"></span>[12] M. Caselli, A. Gentile and R. Giova, Regularity results for solutions to obstacle problems with Sobolev coefficients, *J. Differential Equations* 269, 8308–8330 (2020)
- <span id="page-21-10"></span>[13] I. Chlebicka and C. De Filippis, Removable sets in non-uniformly elliptic problems, Annali di Matematica Pura ed Applicata 199 (2) (2020), 619-649
- <span id="page-21-2"></span>[14] H.J. Choe, Interior behaviour of minimizers for certain functionals with nonstandard growth, Nonlinear Anal. 19 (1992), 933–945
- <span id="page-21-7"></span>[15] A. Clop, R. Giova and A. Passarelli di Napoli, Besov regularity for solutions of p-harmonic equations, Adv. Nonlinear Anal. (2017). https://doi.org/10.1515/anona-2017-0030
- <span id="page-22-2"></span>[16] M. Colombo and G. Mingione, Bounded minimisers of double phase variational integrals, Arch. Ration. Mech. Anal. 218, 219–273 (2015)
- [17] G. Cupini, P. Marcellini, E. Mascolo and A. Passarelli di Napoli, Lipschitz regularity for degenerate elliptic integrals with p, q-growth, Advances in Calculus of Variations (2021). https://doi.org/10.1515/acv-2020-0120
- <span id="page-22-12"></span><span id="page-22-4"></span>[18] C. De Filippis, Regularity results for a class of non-autonomous obstacle problems with  $(p, q)$ -growth, J. Math. Anal. Appl. 501 (2021) 123450
- <span id="page-22-3"></span>[19] C. De Filippis and G. Mingione, On the regularity of minima of non-autonomous functionals, J. Geom. Anal., 30 (2020), 1584–1626
- [20] C. De Filippis and G. Mingione, Interpolative gap bounds for nonautonomous integrals, Analysis and Mathematical Physics,  $11$  (2021), 117 https://doi.org/10.1007/s13324-021-00534-z.
- <span id="page-22-5"></span>[21] M. Eleuteri, Regularity results for a class of obstacle problems, Applications of Mathematics, 52 (2) (2007), 137–170
- <span id="page-22-6"></span>[22] M. Eleuteri, A. Passarelli di Napoli, Higher differentiability for solutions to a class of obstacle problems, Calc. Var., 57 (2018), 115 https://doi.org/10.1007/s00526-018-1387-x
- [23] M. Eleuteri and A. Passarelli di Napoli, Regularity results for a class of non-differentiable obstacle problems, Nonlinear Analysis 194 (2020) 111434
- [24] M. Eleuteri and A. Passarelli di Napoli, On the validity of variational inequalities for obstacle problems with non-standard growth, Annales Fennici Mathematici, 47 (1) (2021), 395–416 https://doi.org/10.54330/afm.114655
- <span id="page-22-0"></span>[25] L. Esposito, F. Leonetti, G. Mingione, Sharp regularity for functionals with  $(p, q)$  growth, J. Differ. Equ., 204 (2004), 5–55
- <span id="page-22-1"></span>[26] I. Fonseca, J. Maly and G. Mingione, Scalar minimizers with fractal singular sets, Arch. Ration. Mech. Anal. 172, 295–307 (2004)
- <span id="page-22-7"></span>[27] C. Gavioli, A priori estimates for solutions to a class of obstacle problems under  $p, q$ -growth conditions, Journal of Elliptic and Parabolic Equations 5 (2) (2019), 325-347
- <span id="page-22-8"></span>[28] C. Gavioli, Higher differentiability of solutions to a class of obstacle problems under non-standard growth conditions, Forum Mathematicum (2019), 31(6), 1501–1516
- <span id="page-22-9"></span>[29] A. Gentile, Higher differentiability results for solutions to a class of non-autonomous obstacle problems with sub-quadratic growth conditions, Forum Mathematicum, 33 (3) (2021), 669–695
- <span id="page-22-10"></span>[30] A. Gentile and R. Giova, Regularity results for solutions to a class of non-autonomous obstacle problems with sub-quadratic growth conditions, Preprint [arXiv:2201.07679v](http://arxiv.org/abs/2201.07679)1
- <span id="page-22-11"></span>[31] A. Gentile, R. Giova and A. Torricelli, Regularity results for bounded solutions to obstacle problems with non-standard growth conditions, Preprint [arXiv:2110.09586v](http://arxiv.org/abs/2110.09586)1
- <span id="page-23-5"></span><span id="page-23-4"></span>[32] R. Giova, Higher differentiability for n-harmonic systems with Sobolev coefficients, J. Differ. Equ. 259(11) (2015), 5667–5687
- [33] R. Giova and A. Passarelli di Napoli, Regularity results for a priori bounded minimizers of non-autonomous functionals with discontinuous coefficients, Adv. Calc. Var. (2018). https://doi.org/10.1515/acv-2016
- <span id="page-23-12"></span><span id="page-23-9"></span>[34] E. Giusti, Direct methods in the calculus of variations, World scientific publishing Co., Singapore (2003)
- <span id="page-23-3"></span>[35] A.G. Grimaldi, Regularity results for solutions to a class of obstacle problems, Nonlinear Analysis: Real World Applications, 62 (2021), 103377
- <span id="page-23-10"></span>[36] A.G. Grimaldi, Higher differentiability for bounded solutions to a class of obstacle problems with  $(p, q)$ -growth, Forum Mathematicum, **35** (2) (2023), 457-485
- [37] A.G. Grimaldi and E. Ipocoana, Higher fractional differentiability for solutions to a class of obstacle problems with non-standard growth conditions, Advances in Calculus of Variations, DOI: 10.1515/acv-2021-0074
- <span id="page-23-11"></span>[38] A.G. Grimaldi and E. Ipocoana, Higher differentiability results in the scale of Besov spaces to a class of double-phase obstacle problems, ESAIM: COCV 28 (2022) 51, doi: 10.1051/cocv/2022050
- <span id="page-23-14"></span>[39] D. Haroske, Envelopes and sharp embeddings of function spaces, Chapman and Hall CRC, Boca Raton (2006)
- <span id="page-23-15"></span>[40] P. Koskela, D. Yang and Y. Zhou, Pointwise characterizations of Besov and Triebel-Lizorkin spaces and quasiconformal mappings,  $Adv. Math. 226 (4) (2011), 3579-3621$
- <span id="page-23-0"></span>[41] P. Marcellini, On the definition and the lower semicontinuity of certain quasiconvex integrals, Ann. Inst. H. Poincaré Anal. Non Linéaire, 3(5) (1986), 391–409
- <span id="page-23-1"></span>[42] P. Marcellini, Regularity of minimizers of integrals of the calculus of variations with nonstandard growth conditions, Arch. Ration. Mech. Anal. 105(3) (1989), 267–284
- <span id="page-23-2"></span>[43] P. Marcellini, Regularity and existence of solutions of elliptic equations with  $p, q$ -growth conditions, *J. Differential Equations*,  $90(1)$  (1991), 1-30
- <span id="page-23-6"></span>[44] A. Passarelli di Napoli, Higher differentiability of minimizers of variational integrals with Sobolev coefficients, Adv. Calc. Var.  $7(1)$   $(2014)$ , 59–89
- <span id="page-23-7"></span>[45] A. Passarelli di Napoli, Higher differentiability of solutions of elliptic systems with Sobolev coefficients: the case  $p = n = 2$ , Potential Anal. 41(3) (2014), 715–735
- <span id="page-23-8"></span>[46] A. Passarelli di Napoli, Regularity results for non-autonomous variational integrals with discontinuous coefficients, Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl  $26(4)$  (2015), 475–496
- <span id="page-23-13"></span>[47] H. Triebel, Theory of function spaces, Birkhäuser-Verlag, Basel (1983)
- <span id="page-24-3"></span><span id="page-24-0"></span>[48] X. Zhang and S. Zheng, Besov regularity for the gradients of solutions to non-uniformly elliptic obstacle problems, Journal of Mathematical Analysis and Applications 505 (2) (2021)
- <span id="page-24-1"></span>[49] V.V. Zhikov, On Lavrentiev's phenomenon, Russ. J. Math. Phys. 3, 249–269 (1995)
- <span id="page-24-2"></span>[50] V.V. Zhikov, On some variational problems, Russ. J. Math. Phys. 5, 105–116 (1997)
- [51] V.V. Zhikov, Lavrentiev phenomenon and homogenization for some variational problems, C. R. Acad. Sci. Paris Sér. I Math. 316, 435–439 (1993)