Conjugate Operators of 1D-harmonic Oscillator

Fumio Hiroshima*and Noriaki Teranishi[†]

May 1, 2024

Abstract

Conjugate operator T of 1D-harmonic oscillator $N = \frac{1}{2}(p^2 + q^2 - 1)$ is defined by an operator satisfying canonical commutation relation [N, T] = -i1 on some domain but not necessarily dense. The angle operator $T_A = \frac{1}{2}(\arctan q^{-1}p + \arctan pq^{-1})$ and Galapon operator $T_G = i \sum_{n=0}^{\infty} \left(\sum_{m \neq n} \frac{(v_m, \cdot)}{n - m} v_n \right)$ are examples of conjugate operators, where $\{v_n\}$ denotes the set of normalized eigenvectors of N. Let \mathscr{T} be a subset of conjugate operators of N. A classification of \mathscr{T} are given as $\mathscr{T} = \mathscr{T}_{\{0\}} \cup \mathscr{T}_{\mathbb{D} \setminus \{0\}} \cup \mathscr{T}_{\partial \mathbb{D}}$, and $T_A \in \mathscr{T}_{\{0\}}$ and $T_G \in \mathscr{T}_{\partial \mathbb{D}}$ are shown. Here the classification is specified by a pair of parameters $(\omega, m) \in \mathbb{C} \times \mathbb{N}$. Finally the time evolution $T_{\omega,m}(t) = e^{itN}T_{\omega,m}e^{-itN}$ for $T_{\omega,m} \in \mathscr{T}$ is investigated, and show that $T_{\omega,m}(t)$ is periodic.

1 Introduction

Let us consider conjugate operator T of 1D-harmonic oscillator

$$N = \frac{1}{2}(p^2 + q^2 - 1).$$

Here $p = -i\frac{d}{dx}$ and $q = M_x$ be the multiplication by x. Both are self-adjoint operators on $L^2(\mathbb{R})$. It satisfies that $[N, T] = -i\mathbb{1}$. The commutator [A, B] of linear operators A and B is a linear operator defined by

$$[A,B] = AB - BA$$

on $D(AB) \cap D(BA)$. Here D(A) denotes the domain of A. In physics, the relationship between position and momentum and that between energy and time are believed to be formally complementary. Since the quantized position and momentum, q and p, satisfy the canonical commutation relation $[q, p] = -i\mathbb{1}$, if N is regarded as the energy of a quantum

^{*}Faculty of Mathematics, Kyushu University

[†]Faculty of Science, Department of Mathematics, Hokkaido University

system, T seems to be a quantization of time. Then T is often times called the time operator associated with N. This paper does not however give any philosophical discussions on *time*. It should be emphasized that time operator is just a name of convenience. We give mathematical definitions of time operators and conjugate operators in Definition 1.1.

In the previous paper [10] we consider two special conjugate operators, angle operator T_A and Galapon operator T_G , of N from a mathematical point of view. T_A and T_G have been mainly investigated so far from a physical point of view, e.g., [13, 1, 18, 7, 6, 11, 12, 17] and as far as we know, there is no firm and robust results on relationships between T_A and T_G in a purely mathematical setting. In this paper we comprehensively investigate conjugate operators including T_A and T_G , and give a relationship between T_A and T_G . Furthermore we classify conjugate operators by the pair of parameters $(\omega, m) \in \mathbb{C} \times \mathbb{N}$ and define disjoint three classes of conjugate operators.

1.1 Angle operators and Galapon operators

Define the annihilation operator a and the creation operator a^* in $L^2(\mathbb{R})$ by

$$a = \frac{1}{\sqrt{2}}(q + ip), \quad a^* = \frac{1}{\sqrt{2}}(q - ip),$$

respectively. The canonical commutation relation (CCR)

$$[a, a^*] = 1, \quad [a, a] = 0 = [a^*, a^*]$$
(1.1)

holds true on a dense subspace of $L^2(\mathbb{R})$. The number operator is defined by a^*a which is actually the harmonic oscillator

 $a^*a = N,$

and N is self-adjoint on $D(N) = D(p^2) \cap D(q^2)$. The normalized ground state of N is given by

$$v(x) = \pi^{-1/4} e^{-x^2/2}.$$

Note that av = 0 and Nv = 0. The normalized eigenvectors v_n are given by

$$v_n = \frac{1}{\sqrt{n!}} a^{*n} v \quad n \ge 1.$$

$$(1.2)$$

Here we write a^{*n} for $(a^*)^n$. It satisfies that $Nv_n = nv_n$ and $\operatorname{Spec}(N) = \mathbb{N}$. Here $\operatorname{Spec}(N)$ denotes the spectrum of N. Note that notation \mathbb{N} describes $\{0, 1, 2, \ldots, \}$ including zero in this paper.

We recall the definitions of conjugate operators and time operators.

Definition 1.1 (Conjugate operators and time operators) If a self-adjoint operator A in Hilbert space \mathcal{H} admits an operator B satisfying the canonical commutation relation:

$$[A,B] = -i\mathbb{1} \tag{1.3}$$

on $D_{A,B} \subset D(AB) \cap D(BA)$ but $D_{A,B} \neq \{0\}$, then B is called a conjugate operator of A and $D_{A,B}$ a CCR-domain of (1.3). Furthermore if B is symmetric, then B is called a time operator of A.

Remark 1.2 By the definitions, conjugate operators are not necessarily densely defined, but the time operators are densely defined.

We shall construct a conjugate operator T of N, i.e.,

$$[N,T] = -i\mathbb{1} \tag{1.4}$$

on some domain. In the previous paper [10] we investigate the angle operator e.g., [15] given by

$$T_A = -\frac{1}{2} \left(\arctan\left(q^{-1}p\right) \oplus \arctan\left(pq^{-1}\right) \right)$$

and the so-called Galapon operator [5, 3, 9, 8] defined by

$$T_G = i \sum_{n=0}^{\infty} \left(\sum_{m \neq n} \frac{(v_m, \cdot)}{n - m} v_n \right).$$

It is known that T_G is a bounded self-adjoint operator, and hence T_G is a time operator of N. However T_A is unbounded and not densely defined. In particular it is not symmetric. Moreover $[N, T_G] = -i\mathbb{1}$ holds on a dense domain, but $[N, T_A] = -i\mathbb{1}$ on a non-dense domain.

Lots of conjugate operators of N appeared in this paper are not densely defined and hence not symmetric. So, we extend relation (1.4) to the so-called ultra-weak canonical commutation relation. See Definition 4.6. For the sesqui-linear form \mathcal{T} , the ultra-weak canonical commutation relation is defined by

$$\mathcal{T}[N\psi,\varphi] - \overline{\mathcal{T}[N\varphi,\psi]} = -i(\psi,\varphi).$$
(1.5)

Here $\mathcal{T}[\psi, \varphi]$ is linear in φ and anti-linear in ψ , and \mathcal{T} is not necessarily densely defined on $L^2(\mathbb{R}) \oplus L^2(\mathbb{R})$. We can define an ultra-weak time operator \mathcal{T}_A associated with T_A , and it is shown in [10] that \mathcal{T}_A satisfies (1.5) on a dense domain. See (4.6).

1.2 Outline of the paper

In this paper, for notational simplicity, we use symbol ℓ^2 to denote the space $\ell^2(\mathbb{N})$ of square-summable complex-valued sequences on \mathbb{N} . We also identify $L^2(\mathbb{R})$ with ℓ^2 and consider conjugate operators of N not on $L^2(\mathbb{R})$ but on ℓ^2 . The key ingredients are shift operators L and L^* on ℓ^2 . Here L denotes the left shift, and L^* is the adjoint of L and hence the right shift on ℓ^2 . The main part of this paper is Sections 4-7, where we investigate (1)-(5) below.

(1) We show that T_A and T_G can be represented in terms of L, L^* and N as

$$T_G = i \left\{ \log(\mathbb{1} - L) + \log(\mathbb{1} - L^*) \right\},$$

$$T_A = \frac{i}{2} \left\{ \log\left(\sqrt{\frac{N+2\mathbb{1}}{N+1}}L^2\right) \oplus \log\left(\sqrt{\frac{N+1}{N+2\mathbb{1}}}L^2\right) \right\}$$

in Theorems 4.9 and 5.4.

(2) We generalize T_G and T_A as

$$i\left\{\log(\mathbb{1}-g_NL) + \log\left(\mathbb{1}-L^*g_N^{-1}\right)\right\},\$$
$$\frac{i}{2}\left\{\log\left(g_{N+2}L^2\right) \oplus \log\left(\tilde{g}_{N+2}L^2\right)\right\},\$$

respectively, in Sections 4 and 5. Here g_{\cdot} is a map from \mathbb{N} to \mathbb{C} . (3) We construct a general class of conjugate operators in Section 6. Let

$$L_{\omega,m} = \omega \mathbb{1} - L^m, \quad (\omega,m) \in \mathbb{C} \times \mathbb{N}.$$

We define $T_{\omega,m}$ by

$$T_{\omega,m} = \frac{i}{m} \log L_{\omega,m}.$$
(1.6)

We can see that $T_{\omega,m}$ is conjugate operators of N. Let \mathbb{D} be the open unit disc on the complex plan \mathbb{C} . We can see that $T_{\omega,m}$ with $\omega \in \partial \mathbb{D}$ is bounded and there exists a dense CCR-domain, and we say that

$$T_{\omega,m}, \quad m \ge 1, \ \omega \in \partial \mathbb{D}$$

are general Galapon operators. It is actually seen that $T_G = T_{1,1} + T_{1,1}^*$. On the other hand general angle operators are introduced by

$$T_{\omega,m}, \quad m \ge 1, \; \omega = 0. \tag{1.7}$$

Let

 $\mathscr{T} = \{ T_{\omega,m} \mid \omega \in \overline{\mathbb{D}}, m \ge 1 \}.$

We divide \mathscr{T} into three disjoint families by $\omega \in \overline{\mathbb{D}}$. We have

$$\mathscr{T} = \mathscr{T}_{\{0\}} \cup \mathscr{T}_{\mathbb{D} \setminus \{0\}} \cup \mathscr{T}_{\partial \mathbb{D}}.$$

Here

$$\begin{aligned} \mathscr{T}_{\{0\}} &= \{ T_{\omega,m} \mid \omega = 0, m \ge 1 \}, \\ \mathscr{T}_{\mathbb{D} \setminus \{0\}} &= \{ T_{\omega,m} \mid 0 < |\omega| < 1, m \ge 1 \}, \\ \mathscr{T}_{\partial \mathbb{D}} &= \{ T_{\omega,m} \mid |\omega| = 1, m \ge 1 \}. \end{aligned}$$

We study CCR-domains for each conjugate operators in \mathscr{T} .

(5) Let H be a self-adjoint operator. Then the strong time operator T_H is defined by the weak Weyl relation

$$T_H e^{-itH} \supset e^{-itH} (T_H + t), \quad t \in \mathbb{R}.$$

It is established that if T_H is a strong time operator for H, then the spectrum of H is purely continuous spectrum. We can show that conjugate operators in \mathscr{T} satisfies a discrete version of weak Weyl relations in Section 7. Furthermore we show that $T_{\omega,m}(t) = e^{itN}T_{\omega,m}e^{-itN}$ is periodic with period $2\pi/m$.

2 Technical tools

2.1 Super coherent vectors

We define the exponent operator e^A for a non self-adjoint operator A in a Hilbert space \mathcal{H} by the geometric series:

$$D(e^{A}) = \left\{ f \in \bigcap_{k=0}^{\infty} D(A^{k}) \mid \lim_{K \to \infty} \sum_{k=0}^{K} \frac{1}{k!} A^{k} f \text{ exists} \right\},\$$
$$e^{A} f = \sum_{k=0}^{\infty} \frac{1}{k!} A^{k} f, \quad f \in D(e^{A}).$$

It should be written as $\sum_{k \in \mathbb{N}} \frac{1}{k!} A^k$ for e^A , but we write e^A for the notational simplicity. We define the set \mathcal{C} consisting of coherent vectors by

$$\mathcal{C} = \operatorname{LH}\left\{ e^{\beta a^*} v \mid \beta \in \mathbb{C} \right\}.$$

Note that $e^{\beta a^*}$ is an unbounded operator, but $v \in D(e^{\beta a^*})$ for any $\beta \in \mathbb{C}$, and it is given by

$$e^{\beta a^*}v(x) = \pi^{-1/4}e^{\beta^2/2}\exp\left(-\frac{(x-\sqrt{2}\beta)^2}{2}\right).$$

It is well known that C is dense in $L^2(\mathbb{R})$. We set $w_{\beta} = e^{-|\beta|^2/2} e^{\beta a^*} v$, where $e^{-|\beta|^2/2}$ is the normalize constant such that $||w_{\beta}|| = 1$. It is shown that $w_{\beta} \in D(a^{*n})$ and $a^{*n}w_{\beta} = e^{-|\beta|^2/2} e^{\beta a^*} a^{*n} v$. Moreover one obtains that

$$aw_{\beta} = \beta w_{\beta}.$$

Thus w_{β} is an eigenvector of a with eigenvalue $\beta \in \mathbb{C}$. We also see that $e^{\gamma a}w_{\beta} = e^{\gamma\beta}w_{\beta}$ for any $\beta, \gamma \in \mathbb{C}$.

Next let us consider vectors of the form $e^{\beta a^{*2}/2}v$. Exponent of a^{*2} is also unbounded. Although $e^{\beta a^{*}}v \in L^{2}(\mathbb{R})$ for any $\beta \in \mathbb{C}$, it is not clear that $e^{\beta a^{*2}/2}v \in L^{2}(\mathbb{R})$. **Lemma 2.1** Let $\beta \in \mathbb{C}$. Then $v \in D\left(e^{\beta a^{*2}/2}\right)$ if and only if $|\beta| < 1$. Moreover, $e^{\beta a^{*2}/2}v \in D(a^{*n})$ for any $|\beta| < 1$ and $n \in \mathbb{N}$.

Proof: Let $|\beta| < 1$. By the Rodrigues formula and orthogonality of the Hermite polynomials H_k ,

$$\left\|\sum_{k=0}^{K} \frac{1}{k!} \left(\frac{\beta}{2}\right)^{k} a^{*2k} v\right\|^{2} = \left\|\sum_{k=0}^{K} \frac{\beta^{k}}{2^{2k} k!} H_{2k} v\right\|^{2} = \sum_{k=0}^{K} \frac{|\beta|^{2k}}{(2^{2k} k!)^{2}} 2^{2k} (2k)! = \sum_{k=0}^{K} \frac{(2k-1)!!}{(2k)!!} |\beta|^{2k}.$$

Let $K \to \infty$. Then $\|e^{\beta a^{*2}/2}v\|^2 = (1 - |\beta|^2)^{-1/2}$ holds. This shows that $v \in D\left(e^{\beta a^{*2}/2}\right)$ if and only if $|\beta| < 1$. The statement that $e^{\beta a^{*2}/2}v \in D(a^{*n})$ is similarly proven.

We call $e^{\beta a^{*2}/2}v$ a super coherent vector and denote the linear hull of super coherent vectors by

$$\mathcal{S} = \operatorname{LH}\left\{ e^{\beta a^{*2}/2} v \mid |\beta| < 1 \right\}.$$

The vectors contained in \mathcal{S} play an important role for studying the angle operator.

2.2 Wiener-Itô decomposition

As is mentioned in Section 1 we shall investigate conjugate operators on ℓ^2 for $L^2(\mathbb{R})$. The first task is to make the identification between ℓ^2 and $L^2(\mathbb{R})$ clear, and we transform a and a^* on $L^2(\mathbb{R})$ to operators on ℓ^2 . By the Wiener-Itô decomposition we have

$$L^2(\mathbb{R}) = \bigoplus_{n=0}^{\infty} L_n,$$

where $L_n = \text{LH}\{v_n\}$ is the one-dimensional linear subspace spanned by eigenvectors v_n of N with eigenvalue $n \in \mathbb{N}$. For each $f \in L^2(\mathbb{R})$, there exists $(c_n)_{n \in \mathbb{N}} \in \ell^2$ so that $f = \sum_{n=0}^{\infty} c_n v_n$. Actually it is given by $c_n = (v_n, f)$. We often use (c_0, c_1, c_2, \ldots) for $(c_n)_{n \in \mathbb{N}}$. By the unitary map

$$Uf = (c_n)_{n \in \mathbb{N}},\tag{2.1}$$

we can identify $L^2(\mathbb{R})$ with ℓ^2 . We set $\xi_n = Uv_n \in \ell^2$. It is seen that

$$UaU^* \colon \xi_n \mapsto \sqrt{n}\xi_{n-1}, \quad n \ge 1,$$
$$Ua^*U^* \colon \xi_n \mapsto \sqrt{n+1}\xi_{n+1}, \quad n \ge 0.$$

For the notational convenience, we write a for UaU^* . Furthermore Uv is denoted by Ω . Then $\Omega = (1, 0, 0, ...)$ and $\xi_n = (0, ..., 0, \overset{n+1_{th}}{1}, 0, ...)$ for $n \ge 0$. Note that $\xi_0 = \Omega$. In the following, our investigation is transformed on ℓ^2 instead of on $L^2(\mathbb{R})$ and we fix $\{\xi_n\}_{n\in\mathbb{N}}$ as a complete orthonormal system of ℓ^2 . Let ℓ_{fin}^2 be the finite particle subspace defined by

$$\ell_{\text{fin}}^2 = \left\{ \sum_{n=0}^m c_n \xi_n \ \middle| \ m \in \mathbb{N}, \ c_n \in \mathbb{C} \right\}.$$

It is dense, and commutator $[a, a^*]$ and polynomials in a and a^* are well defined on this space. Thus algebraic computations on a and a^* can be done on ℓ_{fin}^2 .

2.3 Shift operators L and L^* on ℓ^2

Let L be the left-shift and the adjoint L^* the right-shift on ℓ^2 , which are defined by

$$L\xi_n = \begin{cases} \xi_{n-1} & n \ge 1, \\ 0 & n = 0, \end{cases}$$
$$L^*\xi_n = \xi_{n+1}.$$

We can see that

$$LL^* = 1,$$

 $L^*L = 1 - P_{\{\Omega\}},$

where $P_{\{\Omega\}}$ is the projection to the one-dimensional subspace spanned by Ω . In general an arbitrary closed operator A can be decomposed as A = V|A|, where $|A| = (A^*A)^{1/2}$ is a positive self-adjoint operator and V a partial isometry such that the initial space is $(\text{Ker } A)^{\perp}$ and the final space $\overline{\text{Ran}A}$. A = V|A| is called the polar decomposition of A. V is uniquely determined by these properties together with the addition condition Ker |A| = Ker V. In terms of L, L^* and N, both of a and a^* can be represented as

$$a = L\sqrt{N} = \sqrt{N + \mathbb{1}}L,$$

$$a^* = L^*\sqrt{N + \mathbb{1}} = \sqrt{N}L^*.$$

Note that $N = a^*a$ and $N + 1 = aa^*$. These are just the polar decompositions of closed operators a and a^* . Note that

$$\operatorname{Ker} a = \{ c \Omega \mid c \in \mathbb{C} \} = \operatorname{Ker} \sqrt{N},$$
$$\operatorname{Ker} a^* = \{ 0 \} = \operatorname{Ker} \sqrt{N+1}.$$

L is a partial isometry such that the initial space is $(\text{Ker } a)^{\perp} = \{c \Omega \mid c \in \mathbb{C}\}^{\perp} \cong \bigoplus_{n=1}^{\infty} L_n$ and the final space ℓ^2 . We also see that

$$[N, L^*] \subset L^*, \quad [N, L] \subset -L.$$

$$(2.2)$$

Then $[N, L^{*k}] \subset kL^{*k}$ and $[N, L^k] \subset -kL^k$ hold true. Let $\mathbb{D} = \{z \in \mathbb{C} \mid |z| < 1\}$ be the open unit disc in \mathbb{C} . Notice that

$$e^{\alpha NL^*} \Omega = (1, \alpha, \alpha^2, \alpha^3, \ldots), \quad \alpha \in \mathbb{D}.$$

It can be seen that

$$Le^{\alpha NL^*}\Omega = \alpha e^{\alpha NL^*}\Omega, \quad \alpha \in \mathbb{D}.$$
(2.3)

Thus $e^{\alpha NL^*}\Omega$ is an eigenvector of L with the eigenvalue α .

Lemma 2.2 We have $\operatorname{Spec}(L) = \operatorname{Spec}(L^*) = \overline{\mathbb{D}}$, $\operatorname{Spec}_p(L) = \mathbb{D}$ and $\operatorname{Spec}_p(L^*) = \emptyset$.

Proof: Since $||L|| \leq 1$, $\operatorname{Spec}(L) \cup \operatorname{Spec}(L^*) \subset \overline{\mathbb{D}}$. By (2.3), we see that $\mathbb{D} \subset \operatorname{Spec}_p(L)$ and $\operatorname{Spec}(L) = \operatorname{Spec}(L^*) = \overline{\mathbb{D}}$. Let $|\alpha| = 1$. The relation $L\varphi = \alpha\varphi$ implies $\varphi = (\alpha^n)_{n\in\mathbb{N}}$ but $\varphi \notin \ell^2$. Hence $\operatorname{Spec}_p(L) = \mathbb{D}$. Let $|\alpha| \leq 1$ and $L^*\varphi = \alpha\varphi$. Then $(L^*\varphi)_0 = 0 = \alpha\varphi_0$ and hence $(L^*\varphi)_1 = 0$. Repeating this procedure, we see that $\varphi = 0$. Thus we have $\operatorname{Spec}_p(L^*) = \emptyset$.

Let $k \in \mathbb{N}$. Similarly to the proof of Lemma 2.2 we can see that, for any $k\alpha \in \mathbb{D}$, $e^{\alpha N L^{*k}}\Omega$ is an eigenvector of L^k . Since

$$e^{\alpha N L^{*k}} \Omega = (1, \underbrace{0, \dots, 0}_{k-1}, k\alpha, \underbrace{0, \dots, 0}_{k-1}, (k\alpha)^2, \underbrace{0, \dots, 0}_{k-1}, (k\alpha)^3, \dots),$$

it is actually given by

$$L^{k}e^{\alpha N L^{*k}}\Omega = k\alpha e^{\alpha N L^{*k}}\Omega.$$
(2.4)

We can also extend this to a general version. Let f be a complex-valued function on \mathbb{N} . Set $f(N) = f_N$. Then

$$e^{\alpha f_N L^{*k}} \Omega = \left(1, \underbrace{0, \dots, 0}_{k-1}, f(k)\alpha, \underbrace{0, \dots, 0}_{k-1}, \frac{f(2k)f(k)}{2!}\alpha^2, \underbrace{0, \dots, 0}_{k-1}, \frac{f(3k)f(2k)f(k)}{3!}\alpha^3, \dots\right).$$
(2.5)

For a given function f, we shall find g such that $e^{\alpha f_N L^{*k}} \Omega$ is an eigenvector of $g_{N+k} L^k$ in Sections 4 and 5.

3 Phase operators

We give the definition of $\log A$ for a linear operator A. It is emphasised that A is not necessarily self-adjoint.

Definition 3.1 (log A) Let A be a linear operator on a Hilbert space \mathcal{H} . We define log A by

$$D(\log A) = \left\{ f \in \bigcap_{k=0}^{\infty} D(A^k) \mid \lim_{K \to \infty} \sum_{k=1}^{K} \frac{1}{k} (\mathbb{1} - A)^k f \text{ exists} \right\},$$
$$\log Af = -\sum_{k=1}^{\infty} \frac{1}{k} (\mathbb{1} - A)^k f, \quad f \in D(\log A).$$

Note that in general

$$\log(AB) \neq \log A + \log B,\tag{3.1}$$

$$\log \alpha A \neq \log \alpha + \log A, \quad \alpha \in \mathbb{C}.$$
(3.2)

We define the so-called phase operator ρ on ℓ^2 by

$$\rho = -\frac{i}{2}(\log a - \log a^*).$$
(3.3)

The phase operator ρ has been studied in the large number of literatures e.g., [18, 11, 16] as the conjugate operator of N, but it does not appear to be obvious in the definition of $\log a$ and $\log a^*$.

Lemma 3.2 We have $D(\log a) \cap \ell_{fin}^2 = \{0\}$ and $D(\log a^*) = \{0\}$. In particular, $D(\rho) = \{0\}$.

Proof: Let $\psi = \sum_{l=0}^{m} c_l \xi_l \in \ell_{\text{fin}}^2$ with $c_m \neq 0$. Since

$$\lim_{K \to \infty} \left| \left(\xi_m, \sum_{k=1}^K \frac{1}{k} (\mathbb{1} - a)^k \psi \right) \right| = \lim_{K \to \infty} \left| \sum_{k=1}^K \frac{1}{k} \left((\mathbb{1} - a^*)^k \xi_m, \psi \right) \right| = \lim_{K \to \infty} \left| (\xi_m, \psi) \right| \sum_{k=1}^K \frac{1}{k} = \infty,$$

we see that $D(\log a) \cap \ell_{fin}^2 = \{0\}$. Let $\psi' = \sum_{l=m'}^{\infty} c_l \xi_l \in \ell^2$ with $c_{m'} \neq 0$. Similarly, it follows that

$$\lim_{K \to \infty} \left| \left(\xi_{m'}, \sum_{k=1}^{K} \frac{1}{k} (\mathbb{1} - a^*)^k \psi' \right) \right| = \lim_{K \to \infty} \left| (\xi_{m'}, \psi') \right| \sum_{k=1}^{K} \frac{1}{k} = \infty.$$

Hence $D(\log a^*) = \{0\}.$

From Lemma 3.2 we can see that ρ is not well defined on ℓ_{fin}^2 . This fact is fatal to consider the phase operator ρ on ℓ^2 . Therefore, when we investigate ρ , we have to consider the domain of ρ carefully.

4 Angle operators

4.1 Angle operators by a and a^*

It is said that a relationship between the angle operator T_A and the phase operator ρ is given as $T_A = (\pi/2)\mathbb{1} - \rho + G(N)$ in e.g., [11, (32)]. However it seems to be formal and as was seen in Lemma 3.2 the phase operator ρ is not well defined on ℓ_{fin}^2 .

In order to see a rigorous relationship between phase operator ρ and the angle operator T_A , we express T_A in terms of the creation operators and the annihilation operators. Define

$$\phi = \frac{1}{\sqrt{2}}(a+a^*).$$

We also define two disjoint subspaces of $L^2(\mathbb{R})$ by

$$\mathcal{C}_0 = \operatorname{LH}\left\{ e^{-\alpha x^2/2} \mid \alpha \in (0,1) \right\},$$

$$\mathcal{C}_1 = \operatorname{LH}\left\{ x e^{-\alpha x^2/2} \mid \alpha \in (0,1) \right\}.$$

Note that $C_0 \perp C_1$ and $C_0 + C_1$ is dense. In this paper, # denotes 0 or 1. We transform $C_{\#}$ to the subspaces of ℓ^2 by U defined by (2.1).

Lemma 4.1 Suppose that $0 < \alpha < 1$. Then

$$Ue^{-\alpha x^{2}/2} = \pi^{1/4} \sqrt{\frac{2}{1+\alpha}} \exp\left(\frac{1-\alpha}{1+\alpha} \frac{a^{*2}}{2}\right) \Omega,$$
$$Uxe^{-\alpha x^{2}/2} = \pi^{1/4} \sqrt{\frac{2}{1+\alpha}} \frac{\sqrt{2}}{1+\alpha} a^{*} \exp\left(\frac{1-\alpha}{1+\alpha} \frac{a^{*2}}{2}\right) \Omega.$$

Proof: We set $\beta = (1 - \alpha)/2$. Since $UxU^* = \phi$ and $Ue^{-x^2/2} = \pi^{1/4}\Omega$, we have

$$\begin{split} Ue^{-\alpha x^{2}/2} &= Ue^{-(\alpha-1)x^{2}/2}U^{*}Ue^{-x^{2}/2} = \pi^{1/4}\sum_{n=0}^{\infty}\frac{1}{n!}\beta^{n}\phi^{2n}\Omega\\ &= \pi^{1/4}\sum_{k=0}^{\infty}\left(\sum_{n\geq k}\frac{1}{(n-k)!}\frac{(2n-1)!!}{2^{n-k}(2k-1)!!}\beta^{n}\right)\frac{1}{k!}\frac{a^{*2k}}{2^{k}}\Omega\\ &= \pi^{1/4}\sum_{k=0}^{\infty}\frac{1}{\sqrt{1-\beta}}\left(\frac{\beta}{1-\beta}\right)^{k}\frac{1}{k!}\frac{a^{*2k}}{2^{k}}\Omega\\ &= \pi^{1/4}\sqrt{\frac{2}{1+\alpha}}\exp\left(\frac{1-\alpha}{1+\alpha}\frac{a^{*2}}{2}\right)\Omega. \end{split}$$

Hence we have

$$Uxe^{-\alpha x^{2}/2} = \pi^{1/4} \sqrt{\frac{2}{1+\alpha}} \phi \exp\left(\frac{1-\alpha}{1+\alpha} \frac{a^{*2}}{2}\right) \Omega = \pi^{1/4} \sqrt{\frac{2}{1+\alpha}} \frac{\sqrt{2}}{1+\alpha} a^{*} \exp\left(\frac{1-\alpha}{1+\alpha} \frac{a^{*2}}{2}\right) \Omega.$$

Then the lemma is proven.

Disjoint sets of super coherent vectors is defined by

$$S_0 = \operatorname{LH} \left\{ e^{\beta a^{*2}/2} \Omega \mid \beta \in (0,1) \right\},$$

$$S_1 = \operatorname{LH} \left\{ a^* e^{\beta a^{*2}/2} \Omega \mid \beta \in (0,1) \right\}.$$

Note that $S_0 \perp S_1$ and $S_0 \cup S_1$ is dense. Lemma 4.1 also shows that $UC_{\#} = S_{\#}$. In order to express T_A in terms of a and a^* , we need the inverse of a^* . The operator a^{*-1} is defined by

$$D(a^{*-1}) = \left\{ \sum_{n=0}^{\infty} c_n \xi_n \in \ell^2 \mid c_0 = 0 \right\},\a^{*-1} \sum_{n=1}^{\infty} c_n \xi_n = \sum_{n=1}^{\infty} \frac{c_n}{\sqrt{n}} \xi_{n-1}.$$

We consider $\log(a^{*-1}a)$ and $\log(aa^{*-1})$. Note that a^{*-1} is well defined on $\operatorname{Ran} L^*$.

Lemma 4.2 We have (1) and (2).

(1)
$$\mathcal{S}_0 \subset D\left(\log(a^{*-1}a)\right)$$
 and
 $\log\left(a^{*-1}a\right)e^{\beta a^{*2}/2}\Omega = (\log\beta)e^{\beta a^{*2}/2}\Omega, \quad 0 < \beta < 1.$

(2)
$$S_1 \subset D\left(\log(aa^{*-1})\right)$$
 and
 $\log\left(aa^{*-1}\right)a^*e^{\beta a^{*2}/2}\Omega = (\log\beta)a^*e^{\beta a^{*2}/2}\Omega, \quad 0 < \beta < 1$

Proof: Since $e^{\beta a^{*2}/2}\Omega$ is an eigenvector of $a^{*-1}a$ with the eigenvalue β , for all $n \in \mathbb{N}$, we see that $(\mathbb{1} - a^{*-1}a)^n e^{\beta a^{*2}/2}\Omega = (1 - \beta)^n e^{\beta a^{*2}/2}\Omega$. Hence by the definition of $\log(a^{*-1}a)$ we obtain that $\log(a^{*-1}a) e^{\beta a^{*2}/2}\Omega = (\log \beta) e^{\beta a^{*2}/2}\Omega$. The proof of (2) is the same as that of (1).

Lemma 4.3 Both $\log(a^{*-1}a)$ and $\log(aa^{*-1})$ are unbounded.

Proof: By Lemma 4.2, we can see that $e^{\beta a^{*2}/2}\Omega$ is an eigenvector of log $(a^{*-1}a)$ with eigenvalues log β . Similarly $a^*e^{\beta a^{*2}/2}\Omega$ is an eigenvector of log (aa^{*-1}) with eigenvalues log β . Since $0 < \beta < 1$, we have

Spec
$$\left(\log\left(a^{*-1}a\right)\right) \cap$$
 Spec $\left(\log\left(aa^{*-1}\right)\right) \supset (-\infty, 0).$

Hence the lemma follows.

We recall a definition of $\arctan(A)$ for a linear operator A.

Definition 4.4 Let A be a linear operator on a Hilbert space \mathcal{H} . We define a linear operator $\operatorname{arctan}(A)$ as follows:

$$\begin{split} \mathbf{D}(\arctan(A)) &= \left\{ \varphi \in \bigcap_{k=0}^{\infty} \mathbf{D}\left(A^{2k+1}\right) \ \left| \ \lim_{K \to \infty} \sum_{k=0}^{K} \frac{(-1)^k}{2k+1} A^{2k+1} \varphi \ exists \right\}, \\ \arctan(A)\varphi &= -\sum_{k=0}^{\infty} \frac{(-1)^k}{2k+1} A^{2k+1} \varphi, \quad \varphi \in \mathbf{D}(\arctan(A)). \end{split} \end{split}$$

Now we transform T_A to an operator on ℓ^2 . Let

$$\mathcal{M}_0 = \operatorname{LH}\left\{x^{2n}e^{-\alpha x^2/2} \in L^2(\mathbb{R}) \mid n \in \mathbb{N}, \ \alpha \in (0,1)\right\},$$
$$\mathcal{M}_1 = \operatorname{LH}\left\{x^{2n+1}e^{-\alpha x^2/2} \in L^2(\mathbb{R}) \mid n \in \mathbb{N}, \ \alpha \in (0,1)\right\}.$$

Subspaces $\mathcal{M}_{\#}$ are useful subspaces to consider the commutation relation between the harmonic oscillator and the angle operator:

$$\begin{bmatrix} N, -\arctan(q^{-1}p) \end{bmatrix} = -i\mathbb{1} \quad \text{on } \mathcal{M}_0, \\ \begin{bmatrix} N, -\arctan(pq^{-1}) \end{bmatrix} = -i\mathbb{1} \quad \text{on } \mathcal{M}_1. \end{cases}$$

We define $\mathcal{N}_{\#} \subset \ell^2$ by

$$\mathcal{N}_{0} = \operatorname{LH}\left\{a^{*2n}e^{-\beta a^{*2}/2}\Omega \in \ell^{2} \mid n \in \mathbb{N}, \ \beta \in (0,1)\right\},$$
$$\mathcal{N}_{1} = \operatorname{LH}\left\{a^{*2n+1}e^{-\beta a^{*2}/2}\Omega \in \ell^{2} \mid n \in \mathbb{N}, \ \beta \in (0,1)\right\}.$$

It is immediate to see that $U\mathcal{M}_{\#} = \mathcal{N}_{\#}$. We show algebraic relations we used often in this paper. Let A and B be linear operators. Then the algebraic relation

$$ABf = BAf - ad_B(A)f = (B - ad_B)Af$$

holds for $f \in D(AB) \cap D(BA)$. Hence $AB^2 = (B - ad_B)^2 A$ on $D(AB^2) \cap D(BAB) \cap D(B^2A)$ and then for any $n \in \mathbb{N}$ and any $f \in \bigcap_{k=0}^n D(B^k A B^{n-k})$,

$$AB^n f = (B - \mathrm{ad}_B)^n A f. \tag{4.1}$$

Lemma 4.5 The following relations hold:

$$U \arctan (q^{-1}p) U^* = -\frac{i}{2} \log (a^{*-1}a) \quad on \mathcal{N}_0,$$
$$U \arctan (pq^{-1}) U^* = -\frac{i}{2} \log (aa^{*-1}) \quad on \mathcal{N}_1.$$

Proof: We see that by [10, Lemma 3.6]

$$\arctan\left(q^{-1}p\right)x^{2n}e^{-\alpha x^2/2} = \frac{i}{2}\left\{\left(x^2 - 2\frac{d}{d\alpha}\right)^n \log\left(\frac{1+\alpha}{1-\alpha}\right)\right\}e^{-\alpha x^2/2}.$$

Then, by Lemma 4.1,

$$U \arctan\left(q^{-1}p\right) x^{2n} e^{-\alpha x^2/2} = -\frac{i}{2} \left\{ \left(\phi^2 - 2\frac{d}{d\alpha}\right)^n \log\left(\frac{1-\alpha}{1+\alpha}\right) \right\} \pi^{1/4} \sqrt{\frac{2}{1+\alpha}} \exp\left(\frac{1-\alpha}{1+\alpha}\frac{a^{*2}}{2}\right) \Omega.$$
(4.2)

Let $Z = a^{*-1}a$. Since $\operatorname{ad}_{\phi^2}(Z) = -(1+Z)^2$ on \mathcal{N}_0 , we have for any $m \in \mathbb{N}$ and any analytic function F,

$$\operatorname{ad}_{\phi^2}^m(F(Z)) = \left\{ \left(-(1+x)^2 \frac{d}{dx} \right)^m F(x) \right\} \Big|_{x=Z}$$

on \mathcal{N}_0 . Since $Z \exp(\beta a^{*2}/2) \Omega = \beta \exp(\beta a^{*2}/2) \Omega$, we see that $\exp(\beta a^{*2}/2) \Omega$ is also an eigenvector of $\operatorname{ad}_{\phi^2}^m(F(Z))$ and

$$\left(\operatorname{ad}_{\phi^2}^m F(Z)\right) \exp\left(\frac{\beta}{2}a^{*2}\right) \Omega = \left\{ \left(-(1+\beta)^2 \frac{d}{d\beta}\right)^m F(\beta) \right\} \exp\left(\frac{\beta}{2}a^{*2}\right) \Omega.$$

In particular

$$\left(\operatorname{ad}_{\phi^{2}}^{m}F(Z)\right)\exp\left(\frac{1-\alpha}{1+\alpha}\frac{a^{*2}}{2}\right)\Omega = \left\{\left(2\frac{d}{d\alpha}\right)^{m}F\left(\frac{1-\alpha}{1+\alpha}\right)\right\}\exp\left(\frac{1-\alpha}{1+\alpha}\frac{a^{*2}}{2}\right)\Omega.$$

Moreover replacing F(Z) with $\log Z$, it is obtained that

$$\left(\operatorname{ad}_{\phi^{2}}^{m}\log Z\right)\exp\left(\frac{1-\alpha}{1+\alpha}\frac{a^{*2}}{2}\right)\Omega = \left\{\left(2\frac{d}{d\alpha}\right)^{m}\log\left(\frac{1-\alpha}{1+\alpha}\right)\right\}\exp\left(\frac{1-\alpha}{1+\alpha}\frac{a^{*2}}{2}\right)\Omega.$$
 (4.3)

By (4.2) and (4.3), we have

$$U \arctan(q^{-1}p) x^{2n} e^{-\alpha x^2/2} = -\frac{i}{2} \left\{ \left(\phi^2 - \mathrm{ad}_{\phi^2} \right)^n \log Z \right\} \pi^{1/4} \sqrt{\frac{2}{1+\alpha}} \exp\left(\frac{1-\alpha}{1+\alpha} \frac{a^{*2}}{2}\right) \Omega.$$

From (4.1), we can furthermore see that

$$U \arctan(q^{-1}p) x^{2n} e^{-\alpha x^2/2} = -\frac{i}{2} (\log Z) \phi^{2n} \pi^{1/4} \sqrt{\frac{2}{1+\alpha}} \exp\left(\frac{1-\alpha}{1+\alpha} \frac{a^{*2}}{2}\right) \Omega$$
$$= -\frac{i}{2} (\log Z) U\left(x^{2n} e^{-\alpha x^2/2}\right).$$

Then the first equality is proven. The second equality is similarly proven.

Let

$$S_{0} = \frac{i}{2} \log (a^{*-1}a),$$

$$S_{1} = \frac{i}{2} \log (aa^{*-1}).$$

It is clear that $D(S_0) \cap D(S_1) = \{0\}$, and hence $D(S_0 + S_1) = \{0\}$. Then we define the ultra-weak time operator of $S_0 + S_1$. For the self-consistency we show the definition of ultra-weak time operators below.

Definition 4.6 (Ultra-weak time operator [4]) Let H be a self-adjoint operator on \mathcal{H} and D_1 and D_2 be non-zero subspaces of \mathcal{H} . A sesqui-linear form

$$\mathcal{T}: D_1 \times D_2 \to \mathbb{C}, \quad D_1 \times D_2 \ni (\varphi, \psi) \mapsto \mathcal{T}[\varphi, \psi] \in \mathbb{C}$$

with domain $D(\mathcal{T}) = D_1 \times D_2$ ($\mathcal{T}[\varphi, \psi]$ is antilinear in φ and linear in ψ) is called an ultra-weak time operator of H if there exist non-zero subspaces D and E of $D_1 \cap D_2$ such that (1)–(3) below hold:

(1)
$$E \subset D(H) \cap D$$
.

- (2) $\overline{\mathcal{T}[\varphi,\psi]} = \mathcal{T}[\psi,\varphi] \text{ for all } \varphi,\psi \in D.$
- (3) $HE \subset D_1$ and, for all $\psi, \varphi \in E$,

$$\mathcal{T}[H\varphi,\psi] - \overline{\mathcal{T}[H\psi,\varphi]} = -i(\varphi,\psi).$$
(4.4)

We call E an ultra-weak CCR-domain and D a symmetric domain of \mathcal{T} .

Let us define an ultra-weak time operator of N through S_0 and S_1 . We define

$$\mathcal{T}_0[\varphi, \psi] = \frac{1}{2} \left\{ (S_0 \varphi, \psi) + (\varphi, S_0 \psi) \right\}, \quad \varphi, \psi \in \mathcal{N}_0,$$

$$\mathcal{T}_1[\varphi, \psi] = \frac{1}{2} \left\{ (S_1 \varphi, \psi) + (\psi, S_1 \varphi) \right\}, \quad \varphi, \psi \in \mathcal{N}_1$$

and

$$\mathcal{T} = \mathcal{T}_0 \oplus \mathcal{T}_1 \tag{4.5}$$

with symmetric domain $(D(S_0) \times D(S_0)) \oplus (D(S_1) \times D(S_1))$. On the other hand let us define the ultra-weak time operator \mathcal{T}_A associated with angle operator \mathcal{T}_A below. Let $h_0 = \arctan(q^{-1}p)$ and $h_1 = \arctan(pq^{-1})$. We define

$$\mathcal{T}_{A0}[\varphi,\psi] = \frac{1}{2} \left\{ (h_0\varphi,\psi) + (\varphi,h_0\psi) \right\}, \quad \varphi,\psi \in \mathcal{N}_0,$$

$$\mathcal{T}_{A1}[\varphi,\psi] = \frac{1}{2} \left\{ (h_1\varphi,\psi) + (\psi,h_1\varphi) \right\}, \quad \varphi,\psi \in \mathcal{N}_1$$

and

$$\mathcal{T}_A = \mathcal{T}_{A0} \oplus \mathcal{T}_{A1} \tag{4.6}$$

with symmetric domain $(D(h_0) \times D(h_0)) \oplus (D(h_1) \times D(h_1))$.

Theorem 4.7 The ultra-weak time operator \mathcal{T}_A of the harmonic oscillator on $L^2(\mathbb{R})$ is unitary equivalent to \mathcal{T} on ℓ^2 :

$$\mathcal{T}_A[\varphi,\psi] = \mathcal{T}[U\varphi,U\psi]. \tag{4.7}$$

Proof: This follows from Lemma 4.5.

Remark 4.8 (4.7) in Theorem 4.9 can be interpreted as a rigorous justification of

$$\frac{1}{2} \left(\arctan\left(q^{-1}p\right) + \arctan\left(pq^{-1}\right) \right) \cong \frac{i}{2} \left\{ \log\left(a^{*-1}a\right) + \log\left(aa^{*-1}\right) \right\}.$$

In some literature however the following description is stated:

$$\frac{1}{2}\left(\arctan\left(q^{-1}p\right) + \arctan\left(pq^{-1}\right)\right) \cong \frac{i}{2}\left\{\log a + \log a^*\right\}.$$

The right-hand side can not be however defined as an operator by Lemma 3.2. Even if it is a formal argument, it is dangerous to base the argument on this equivalent relation.

4.2 Angle operator by shift operators

Let L be the left shift operator defined in Section 2.3. The angle operator T_A can be represented by L and N. Let us define

$$L_{A0} = \frac{i}{2} \log \left(\sqrt{\frac{N+2\mathbb{1}}{N+1\mathbb{1}}} L^2 \right),$$
$$L_{A1} = \frac{i}{2} \log \left(\sqrt{\frac{N+1}{N+2\mathbb{1}}} L^2 \right).$$

Theorem 4.9 We have

$$S_0 = L_{A0} \quad on \ \mathcal{N}_0, \tag{4.8}$$

$$S_1 = L_{A1} \quad on \ \mathcal{N}_1. \tag{4.9}$$

In particular

$$U \arctan\left(q^{-1}p\right) U^* = L_{A0} \quad on \ \mathcal{N}_0, \tag{4.10}$$

$$U \arctan(pq^{-1}) U^* = L_{A1} \quad on \ \mathcal{N}_1.$$
 (4.11)

Proof: We see that

$$a^{*-1}a = \left(\sqrt{N}L^*\right)^{-1}\sqrt{N+1}L = L\sqrt{\frac{N+1}{N}}L = \sqrt{\frac{N+21}{N+1}}L^2$$

on $\operatorname{Ran}(\mathbb{1} - P_{\{a^*\Omega\}})$. Similarly

$$aa^{*-1} = L\sqrt{N}\left(L^*\sqrt{N+1}\right)^{-1} = L\sqrt{\frac{N}{N+1}}L = \sqrt{\frac{N+1}{N+21}}L^2$$

on Ran $(\mathbb{1} - P_{\{\Omega\}})$. From this we have (4.8) and (4.9). Relations (4.10) and (4.11) follow from Theorem 4.5.

4.3 Generalization of angle operator T_A

We generalize L_{A0} and L_{A1} to the form of $i \log (g(N)L^2)$. A fundamental idea is to find eigenvectors φ of linear operator $ad_N (\log (g(N)L^2))$ so that $ad_N (\log (g(N)L^2))\varphi = a\varphi$ with $a \neq 0$. Then it follows that

$$\left[N, -\frac{i}{a}\log\left(g(N)L^2\right)\right] = -i\mathbb{1}$$

on CCR-domain LH{eigenvectors φ }. Hereafter we write g_N for g(N). Let $P_{\geq 2}$ be the projection onto $\overline{\text{LH}\{\xi_n \mid n \geq 2\}}$, and P_0 the projection onto $\overline{\text{LH}\{\xi_{2n} \mid n \in \mathbb{N}\}}$.

Lemma 4.10 Let f and g be complex-valued functions on \mathbb{N} , and $\alpha \in \mathbb{C}$. Suppose that there exists a constant $\beta \in \mathbb{C}$ such that

$$g_{N+2}f_{N+2} - g_N f_N P_{\geq 2} = \beta \mathbb{1} \quad on \ P_0 \ell_{\text{fin}}^2.$$
(4.12)

Then, for all $n \in \mathbb{N}$, we see that $\operatorname{ad}_{g_{N+2}L^2}(f_N L^{*2})^n = n\beta (f_N L^{*2})^{n-1}$ on $P_0 \ell_{\operatorname{fin}}^2$.

Proof: The relation (4.12) is equivalent to $[g_{N+2}L^2, f_NL^{*2}] = \beta \mathbb{1}$ on $P_0\ell_{\text{fin}}^2$. This implies that $\operatorname{ad}_{g_{N+2}L^2}(f_NL^{*2})^n = n\beta (f_NL^{*2})^{n-1}$ on $P_0\ell_{\text{fin}}^2$.

Lemma 4.11 Let f be a complex-valued function on \mathbb{N} . Suppose that |f(2n)| > 0 for all natural number $n \ge 1$ and there exists a limit (including infinity)

$$M_f = \lim_{n \to \infty} \frac{n}{|f(2n)|} \le \infty.$$

Then, for all $l, m \in \mathbb{N}$ and $\alpha \in \mathbb{C}$ such that $|\alpha| < M_f$,

$$\Omega \in \mathcal{D}\left(N^l \left(f_N L^{*2}\right)^m e^{\alpha f_N L^{*2}}\right).$$

Proof: Note that

$$\left\| N^{l} \left(f_{N} L^{*2} \right)^{m} e^{\alpha f_{N} L^{*2}} \Omega \right\|^{2} = \sum_{n=0}^{\infty} \frac{|\alpha|^{2n}}{(n!)^{2}} \left\| N^{l} \left(f_{N} L^{*2} \right)^{(n+m)} \Omega \right\|^{2}.$$

Since

$$N^{l} (f_{N} L^{*2})^{(n+m)} \Omega = (2(n+m))^{l} \prod_{j=1}^{n+m} f(2j) L^{*2(n+m)} \Omega,$$

the radius of the convergence of above infinite series is given by $\lim_{n\to\infty} n/|f(2n+2m)| = M_f$. Then the lemma follows.

Lemma 4.12 Let f and g be complex-valued functions on \mathbb{N} such that (4.12) is satisfied. Then, for all $l, m, n \in \mathbb{N}$ and $\alpha \in \mathbb{C}$ such that $|\alpha| < M_f$,

$$\Omega \in \mathcal{D}\left(\left(g_{N+2}L^2\right)^l N^m \left(f_N L^{*2}\right)^n e^{\alpha f_N L^{*2}}\right).$$

Moreover, $e^{\alpha f_N L^{*2}} \Omega$ is an eigenvector of $g_{N+2} L^2$ such that

$$g_{N+2}L^2 e^{\alpha f_N L^{*2}} \Omega = \alpha \beta e^{\alpha f_N L^{*2}} \Omega.$$
(4.13)

Proof: We have

$$\left\| \left(g_{N+2}L^2 \right)^l N^m \left(f_N L^{*2} \right)^m e^{\alpha f_N L^{*2}} \Omega \right\|^2 = \sum_{n=0}^{\infty} \frac{|\alpha|^{2n}}{(n!)^2} \left\| \left(g_{N+2}L^2 \right)^l N^m \left(f_N L^{*2} \right)^{(n+m)} \Omega \right\|^2.$$

Since

$$(g_{N+2}L^2)^l N^m (f_N L^{*2})^{(n+m)} \Omega = \begin{cases} \left(\prod_{j=n+m-l+1}^{n+m} g_{2j}\right) (2(n+m))^m \left(\prod_{j=1}^{n+m} f_{2j}\right) L^{*2(n+m-l)}\Omega, & n+m \ge l, \\ 0, & n+m < l, \end{cases}$$

the right-hand side above converges for $\alpha \in \mathbb{C}$ such that $|\alpha| < M_f$. From Lemma 4.10, we can also see that $e^{\alpha f_N L^{*2}} \Omega$ is an eigenvector of $g_{N+2}L^2$ and (4.13) follows. \Box

We define the super coherent vector $\xi_{\alpha,f}$ by

$$\xi_{\alpha,f} = e^{\alpha f_N L^{*2}} \Omega.$$

We set

$$\mathbb{D}_{f,\beta} = \left\{ \alpha \in \mathbb{C} \mid |1 - \alpha\beta| < 1, \ |\alpha| < M_f \right\}.$$
(4.14)

Lemma 4.13 Let f and g be complex-valued functions on \mathbb{N} such that (4.12) is satisfied. Then for all $n \in \mathbb{N}$ and $\alpha \in \mathbb{D}_{f,\beta}$, we have $(f_N L^{*2})^n \xi_{\alpha,f} \in D(\log(g_{N+2}L^2))$ and

$$\log\left(g_{N+2}L^2\right)\left(f_NL^{*2}\right)^n\xi_{\alpha,f} = \left\{\left(f_NL^{*2} + \frac{d}{d\alpha}\right)^n\left(\log\alpha\beta\right)\right\}\xi_{\alpha,f}.$$

Proof: By Lemma 4.12 we have $(f_N L^{*2})^n \xi_{\alpha,f} \in \bigcap_{m=0}^{\infty} D((1 - g_{N+2}L^2)^m)$. From (4.1) and (4.3) it follows that

$$\log (g_{N+2}L^2) (f_N L^{*2})^n \xi_{\alpha,f} = -\sum_{k=1}^K \frac{1}{k} (1 - g_{N+2}L^2)^k (f_N L^{*2})^n \xi_{\alpha,f}$$

= $-\sum_{k=1}^K \frac{1}{k} \left\{ (f_N L^{*2} - \operatorname{ad}_{f_N L^{*2}})^n (1 - g_{N+2}L^2)^k \right\} \xi_{\alpha,f}$
= $-\sum_{k=1}^K \frac{1}{k} \left\{ (f_N L^{*2} + \frac{d}{d\alpha})^n (1 - \alpha\beta)^k \right\} \xi_{\alpha,f} \rightarrow \left\{ (f_N L^{*2} + \frac{d}{d\alpha})^n (\log \alpha\beta) \right\} \xi_{\alpha,f}$

as $K \to \infty$. Then the proof is complete.

The next theorem is a generalization of (1) of Lemma 4.2.

Theorem 4.14 Let f and g be complex-valued functions on \mathbb{N} such that (4.12) is satisfied. Then,

$$\left[N, \frac{i}{2}\log\left(g_{N+2}L^2\right)\right] = -i\mathbb{1}$$
(4.15)

holds on CCR-domain LH $\{(f_N L^{*2})^n \xi_{\alpha,f} \mid n \in \mathbb{N}, \alpha \in \mathbb{D}_{f,\beta}\}.$

Proof: In this proof we set $X = f_N L^{*2}$ and $Y = g_{N+2}L^2$. From Lemmas 4.11, 4.12 and 4.13, we see that, for all $n \in \mathbb{N}$,

$$X^n \xi_{\alpha, f} \in \mathcal{D}\left(N(\log Y)\right) \cap \mathcal{D}\left((\log Y)N\right).$$

Since $\beta N = 2\overline{XY}$ on $P_0 D(N)$, we have

$$(\log Y)NX^{n}\xi_{\alpha,f} = \frac{2}{\beta}(\log Y)\left(YX - \beta \mathbb{1}\right)X^{n}\xi_{\alpha,f}$$
$$= \frac{2}{\beta}\left(Y(\log Y)X^{n+1} - \beta(\log Y)X^{n}\right)\xi_{\alpha,f}$$

By Lemma 4.13 we have

$$(\log Y)NX^{n}\xi_{\alpha,f} = \frac{2}{\beta}\left\{ (Y(X - \operatorname{ad}_{X}) - \beta)(\log Y)X^{n} \right\} \xi_{\alpha,f}$$
$$= \left\{ \left(N - \frac{2}{\beta}Y \operatorname{ad}_{X} \right) (\log Y)X^{n} \right\} \xi_{\alpha,f}$$
$$= \left\{ N(\log Y)X^{n} - \frac{2}{\beta}Y(X - \operatorname{ad}_{X})^{n} \operatorname{ad}_{X}(\log Y) \right\} \xi_{\alpha,f}$$
$$= \left\{ N(\log Y)X^{n} - \frac{2}{\beta}(X - \operatorname{ad}_{X})^{n}Y \operatorname{ad}_{X}(\log Y) \right\} \xi_{\alpha,f}$$

Here we used $Y(X - \mathrm{ad}_X)Z\varphi = (X - \mathrm{ad}_X)YZ\varphi$ for any linear operator Z and any $\varphi \in D(XYZ) \cap D(YZZ) \cap D(YZX)$. In a similar way to (4.3), we can obtain that

$$(\log Y)NX^{n}\xi_{\alpha,f} = \left\{ N(\log Y)X^{n} + \frac{2}{\beta} \left(X + \frac{d}{d\alpha} \right)^{n} \alpha\beta \frac{d}{d\alpha} \log\left(\alpha\beta\right) \right\} \xi_{\alpha,f}$$
$$= \left(N(\log Y) + 2\mathbb{1} \right) X^{n}\xi_{\alpha,f}.$$

Hence $[N, \frac{i}{2} \log Y] X^n \xi_{\alpha, f} = -i X^n \xi_{\alpha, f}$ follows and (4.15) is proven.

We can also see a generalization of (2) of Lemma 4.2.

Theorem 4.15 Let \tilde{h} be a complex-valued function on \mathbb{N} such that \tilde{h}_{N+1}^{-1} is bounded and \tilde{f} and \tilde{g} be complex-valued functions on \mathbb{N} such that

$$\tilde{h}_{N+1}^{-1}\tilde{g}_{N+3}\tilde{h}_{N+3}\tilde{f}_{N+2} - \tilde{h}_{N-1}^{-1}\tilde{g}_{N+1}\tilde{h}_{N+1}\tilde{f}_{N}P_{\geq 2} = \tilde{\beta}\mathbb{1}$$

on $P_0\ell_{\text{fin}}^2$ with some $\tilde{\beta} \in \mathbb{C}$. Let

$$\mathbb{D}_{\tilde{f},\tilde{h},\tilde{\beta}} = \left\{ \alpha \in \mathbb{C} \mid \alpha \in \mathbb{D}_{\tilde{f},\tilde{\beta}}, \ \xi_{\alpha,\tilde{f}} \in \bigcap_{n=0}^{\infty} \mathcal{D}\left(\tilde{g}_{N+3}\tilde{h}_{N+3}\left(\tilde{f}_{N}L^{*2}\right)^{n}\right) \right\}.$$
(4.16)

Then $\tilde{h}_N L^* \xi_{\alpha,\tilde{f}}$ is an eigenvector of $\tilde{g}_{N+2} L^2$ such that

$$\tilde{g}_{N+2}L^2\tilde{h}_N L^* \xi_{\alpha,\tilde{f}} = \alpha \tilde{\beta} \tilde{h}_N L^* \xi_{\alpha,\tilde{f}}.$$
(4.17)

Furthermore it follows that

$$\left[N, \frac{i}{2}\log\left(\tilde{g}_{N+2}L^2\right)\right] = -i\mathbb{1}$$
(4.18)

on CCR-domain LH $\left\{ \tilde{h}_N L^* \left(\tilde{f}_N L^{*2} \right)^n \xi_{\alpha, \tilde{f}} \mid n \in \mathbb{N}, \ \alpha \in \mathbb{D}_{\tilde{f}, \tilde{h}, \tilde{\beta}} \right\}.$

Proof: By the following equalities

$$\tilde{g}_{N+2}L^2\tilde{h}_NL^* = \tilde{g}_{N+2}\tilde{h}_{N+2}L = \tilde{h}_N\tilde{h}_N^{-1}\tilde{g}_{N+2}\tilde{h}_{N+2}L^*L^2 = \tilde{h}_NL^*\left(\tilde{h}_{N+1}^{-1}\tilde{g}_{N+3}\tilde{h}_{N+3}L^2\right)$$

on $P_0\ell^2 \cap D(\tilde{g}_{N+1}\tilde{h}_{N+1})$, one can show (4.17) in a similar manner to the proof of (4.12) by replacing g_{N+2} of (4.12) with $\tilde{h}_{N+1}^{-1}\tilde{g}_{N+3}\tilde{h}_{N+3}$. The proof of (4.18) is also similar to those of Lemma 4.12 and Theorem 4.14.

By Theorems 4.14 and 4.15 we can also construct an ultra-weak time operator of N. This is a generarization of \mathcal{T} in (4.5). Let $f, \tilde{f}, g, \tilde{g}$ and h be functions on \mathbb{N} given in Theorems 4.14 and 4.15. Let

$$\mathcal{K}_{0} = \operatorname{LH}\left\{\left(f_{N}L^{*2}\right)^{n}\xi_{\alpha,f} \mid n \in \mathbb{N}, \ \alpha \in \mathbb{D}_{f,\beta}\right\},\$$
$$\mathcal{K}_{1} = \operatorname{LH}\left\{\tilde{h}_{N}L^{*}\left(\tilde{f}_{N}L^{*2}\right)^{n}\xi_{\alpha,\tilde{f}} \mid n \in \mathbb{N}, \ \alpha \in \mathbb{D}_{\tilde{f},\tilde{h},\tilde{\beta}}\right\}.$$

Note that $\mathcal{K}_0 \oplus \mathcal{K}_1$ is dense if $\mathbb{D}_{f,\beta} \neq \emptyset$ and $\mathbb{D}_{\tilde{f},\tilde{h},\tilde{\beta}} \neq \emptyset$. Let

$$S_{0g} = \frac{i}{2} \log \left(g_{N+2} L^2 \right),$$

$$S_{1\tilde{g}} = \frac{i}{2} \log \left(\tilde{g}_{N+2} L^2 \right).$$

These are generalization of S_0 and S_1 . Define

$$\mathcal{T}_{0g}[\varphi,\psi] = \frac{1}{2} \{ (S_{0g}\varphi,\psi) + (\varphi,S_{0g}\psi) \}, \quad \varphi,\psi\in\mathcal{K}_0,$$

$$\mathcal{T}_{1\tilde{g}}[\varphi,\psi] = \frac{1}{2} \{ (S_{1\tilde{g}}\varphi,\psi) + (\varphi,S_{1\tilde{g}}\psi) \}, \quad \varphi,\psi\in\mathcal{K}_1$$

and set

$$\mathcal{T}_{g,\tilde{g}} = \mathcal{T}_{0g} \oplus \mathcal{T}_{1\tilde{g}}.$$
(4.19)

Theorem 4.16 Suppose that $f, \tilde{f}, g, \tilde{g}$ and \tilde{h} satisfy the assumptions given in Theorem 4.14 and Lemma 4.15. We also assume that $\mathbb{D}_{f,\beta} \neq \emptyset$ and $\mathbb{D}_{\tilde{f},\tilde{h},\tilde{\beta}} \neq \emptyset$. Then $\mathcal{T}_{g,\tilde{g}}$ is an ultra-weak time operator of N with a dense CCR-domain $\mathcal{K}_0 \oplus \mathcal{K}_1$.

Proof: This follows from (4.15) and (4.18).

4.4 Conjugate operators of the form $\log p(L)$

In the previous section we consider the conjugate operator of the form $\log(g_N L^2)$. Another direction of generalizations is to consider conjugate operators of the form $\log(g_N L^n)$. In

special cases it can be reduced to consider conjugate operators of $\log(g_N L^n)$ to those of $\log(L^n)$. See Lemma 4.17 below. Let f be a function on \mathbb{N} and

$$f!_k(N)\xi_n = \left(\prod_{m=0}^{[n/k]} f_{n-km}\right)\xi_n, \quad n \in \mathbb{N}$$

We extend $f!_k(N)$ to ℓ_{fin}^2 by the linearity and denote the closure of $f!_k(N) \lceil_{\ell_{\text{fin}}^2}$ by the same symbol $f!_k(N)$.

Lemma 4.17 Let f and g be complex-valued functions on \mathbb{N} such that $f_n g_n = n$ for all $n \in \mathbb{N}$. Suppose that $\alpha \in (0, 1)$ and $\Omega \in D(e^{\alpha f_N L^{*k}})$. Then

$$\log\left(g_{N+k}L^{k}\right)e^{\alpha f_{N}L^{*k}}\Omega = f!_{k}(N)\log\left((N+k\mathbb{1})L^{k}\right)e^{\alpha L^{*k}}\Omega.$$
(4.20)

In particular, setting $h_n = n$ for all $n \in \mathbb{N}$, we have

$$\left(\log L^{k}\right)e^{\alpha N L^{*k}}\Omega = h!_{k}(N)\log\left((N+k\mathbb{1})L^{k}\right)e^{\alpha L^{*k}}\Omega,$$
(4.21)

$$\log(g_{N+k}L^{k}) e^{\alpha f_{N}L^{*k}} \Omega = h!_{k}(N)^{-1} f!_{k}(N) (\log L^{k}) e^{\alpha N L^{*k}} \Omega.$$
(4.22)

Proof: It is straightforward to see that

$$\log (g_{N+k}L^k) e^{\alpha f_N L^{*^k}} \Omega = \log (g_{N+k}L^k) f!_k(N) e^{\alpha L^{*^k}} \Omega$$
$$= f!_k(N) \log (g_{N+k}f_{N+k}L^k) e^{\alpha L^{*^k}} \Omega$$
$$= f!_k(N) \log ((N+k1)L^k) e^{\alpha L^{*^k}} \Omega.$$

Putting $g_n = 1$ for all $n \in \mathbb{N}$ in (4.20), we can derive (4.21). Combining (4.20) and (4.21) we can also see (4.22). Then the lemma follows.

By (4.22), $\log(g_{N+k}L^k)$ can be represented as $C(N)\log L^k$ on $LH\{e^{\alpha NL^{*k}}\Omega\}$ with

$$C(N) = h!_k(N)^{-1} f!_k(N).$$

Since the operator C(N) commutes with N, the investigation of conjugate operators of the form $\log (g_{N+k}L^k)$ can be reduced to that of $\log L^k$ on $LH\{L^{*nk}e^{\alpha NL^{*k}}\Omega \mid n \in \mathbb{N}, |\alpha| < 1\}$.

Example 4.18 We see that $T_A \cong L_{A0} \oplus L_{A1}$ with $L_{A0} = \frac{i}{2} \log \left(\sqrt{\frac{N+21}{N+1}} L^2 \right)$ and $L_{A1} = \frac{i}{2} \log \left(\sqrt{\frac{N+1}{N+21}} L^2 \right)$. By Lemma 4.17 to investigate the CCR-domain of L_{A0} and L_{A1} can be reduced to investigate the CCR-domain of $\frac{i}{2} \log L^2$.

Thus from now on we study the conjugate operator of the form $\log p(L)$. The idea to find a CCR-domain of $\log p(L)$ is similar to that of the previous subsection. Formally

$$[N, \log p(L)] \subset -p(L)^{-1}p'(L)L,$$

and $e^{\alpha NL^*}\Omega$ is an eigenvector of L and that of p(L). Thus $[N, \log p(L)] = -p(\alpha)^{-1}p'(\alpha)\alpha$ on LH $\{e^{\alpha NL^*}\Omega\}$. To solve $[N, \log p(L)] = c$ we study the algebraic equation of the form

$$\alpha p'(\alpha) + cp(\alpha) = 0$$

with some $c \in \mathbb{C}$. Then we set

$$D(p,c) = LH\{\varphi \in D(N) \mid \exists \alpha \in \mathbb{C} \ s.t. \ |p(\alpha) - 1| < 1, \ \alpha p'(\alpha) + cp(\alpha) = 0, \ \varphi \in Ker(L - \alpha \mathbb{1})\}$$

It may be possibly $D(p,c) = \{0\}.$

Theorem 4.19 Let p be a polynomial and $c \in \mathbb{C} \setminus \{0\}$. If there exists $\alpha \in \mathbb{C}$ such that $|\alpha| < 1$, $\alpha p'(\alpha) + cp(\alpha) = 0$ and $|p(\alpha) - 1| < 1$, then $D(p, c) \neq \{0\}$ and

$$\left[N, -\frac{i}{c}\log p(L)\right] = -i\mathbb{1}$$

with CCR-domain D(p, c). Moreover, in $D(\sum k^{-1}N(\mathbb{1}-p(L))^k)$, there is no infinite dimensional CCR-domain for N and $c\log p(L)$ unless $p(x) = cx^m$ for some $c \in \mathbb{C}$ and $m \in \mathbb{N}$.

Proof: We can see that

$$[N, \log p(L)] = -\sum_{k=1}^{\infty} \frac{1}{k} \left[N, (\mathbb{1} - p(L))^k \right] = -\sum_{k=1}^{\infty} \left(\mathbb{1} - p(L) \right)^{k-1} Lp'(L)$$
(4.23)

on $D(\sum k^{-1}N(\mathbb{1}-p(L))^k) \cap D(N\log p(L)) \cap D(\log p(L)N)$. Let $\varphi \in Ker(L-\alpha\mathbb{1})$. Then $[N, \log p(L)]\varphi = -\alpha p'(\alpha)p(\alpha)^{-1}\varphi.$

By $\alpha p'(\alpha) + cp(\alpha) = 0$ we see that

$$[N, \log p(L)]\varphi = c\varphi.$$

Next we shall show that the dimension of the CCR-domain in $\operatorname{Ran}(p(L))$ is finite. For any

$$\varphi \in \mathcal{D}\left(\sum_{k \ge 1} \frac{1}{k} N(\mathbb{1} - p(L))^k\right)$$

in the CCR-domain, there exists $c \in \mathbb{C}$ such that $[N, \log p(L)]\varphi = c\varphi$. From (4.23), we see that $\lim_{k\to\infty} (\mathbb{1} - p(L))^k Lp'(L)\varphi = 0$. Therefore,

$$cp(L)\varphi = p(L)[N, \log p(L)]\varphi = -\sum_{k=1}^{\infty} p(L) \left(\mathbb{1} - p(L)\right)^{k-1} Lp'(L)\varphi = -Lp'(L)\varphi.$$

Thus $\varphi \in \operatorname{Ker}(Lp'(L) + cp(L))$. This implies that the CCR-domain is at most finite dimensional space, since dim $\operatorname{Ker}(L - \lambda \mathbb{1})^k \leq k$ for all $\lambda \in \mathbb{C}$ and $k \in \mathbb{N}$, and p is a polynomial with $Lp'(L) + cp(L) \neq 0$. This fact can be derived from a fundamental lemma below. Then the theorem follows. \Box

Lemma 4.20 Let A be a bounded operator on a Banach space \mathcal{K} . Let $\{n_1, \ldots, n_k\} \subset \mathbb{N}$, $\{\lambda_1, \ldots, \lambda_k\} \subset \mathbb{C}$ such that $\lambda_i \neq \lambda_j$ for $i \neq j$. Then it follows that

$$\operatorname{Ker}\left(\prod_{j=1}^{k} (\lambda_j \mathbb{1} - A)^{n_j}\right) = \bigoplus_{j=1}^{k} \operatorname{Ker}(\lambda_j \mathbb{1} - A)^{n_j}.$$

Proof: We refer to [2, Lemma 1.76].

5 Galapon operators

5.1 Galapon operators by shift operators

In this section we investigate Galapon operator which is a bounded self-adjoint time operator. Recall that $\operatorname{Spec}(N) = \{n\}_{n \in \mathbb{N}}$ and $\{v_n\}_{n \in \mathbb{N}}$ the set of normalized eigenvectors of Nin $L^2(\mathbb{R})$. We define T_G by

$$D(T_G) = LH\{v_n \mid n \in \mathbb{N}\},\$$

$$T_G \varphi = i \sum_{n=0}^{\infty} \left(\sum_{m \neq n} \frac{(v_m, \varphi)}{n - m} v_n \right), \quad \varphi \in D(T_G).$$
(5.1)

It is known that T_G is bounded and

$$[N, T_G] = -i\mathbb{1} \tag{5.2}$$

on a dense CCR-domain LH $\{v_n - v_m \mid n, m \in \mathbb{N}\}$ (see [5]). In particular, T_G is not equal to the angle operator T_A . The angle operator T_A is expressed on ℓ^2 by Theorem 4.9. On the other hand the Galapon operator is given by (5.1). As it is, no comparison of T_A and T_G can be made, so we shall transform T_G to an operator on ℓ^2 . Let us consider $\log (\mathbb{1} - L)$ and $\log (\mathbb{1} - L^*)$.

Lemma 5.1 We have $\ell_{\text{fin}}^2 \subset D(\log(\mathbb{1} - L)) \cap D(\log(\mathbb{1} - L^*)).$

Proof: By the definition of $\log(1 - L)$, we see that

$$\log(1 - L)\xi_n = -\sum_{k=1}^{\infty} \frac{1}{k} L^k \xi_n = -\sum_{k=1}^n \frac{1}{k} \xi_{n-k}.$$

Thus $\xi_n \in D(\log(\mathbb{1}-L))$ and hence $\ell_{\text{fin}}^2 \subset D(\log(\mathbb{1}-L))$. We also see that

$$\log(\mathbb{1} - L^*)\xi_n = -\sum_{k=1}^{\infty} \frac{1}{k} L^{*k} \xi_n = -\sum_{k=1}^{\infty} \frac{1}{k} \xi_{n+k}$$

23

and

$$\|\log(1 - L^*)\xi_n\|^2 = \sum_{k=1}^{\infty} \frac{1}{k^2} < \infty.$$

Thus $\xi_n \in \mathcal{D}(\log(\mathbb{1} - L^*))$ and $\ell_{\text{fin}}^2 \subset \mathcal{D}(\log(\mathbb{1} - L^*))$.

Remark 5.2 By the proof of Lemma 5.1 we see that

- (1) $\ell_{\text{fin}}^2 \subset \mathcal{D}(\log(\mathbb{1} L^*)), \text{ but } \mathcal{D}(\log L^*) \cap \ell_{\text{fin}}^2 = \{0\},\$
- (2) $\log(\mathbb{1} L)\ell_{\text{fin}}^2 \subset \ell_{\text{fin}}^2$.

We consider the following operator L_G .

Definition 5.3 We define a linear operator L_G on ℓ^2 by

$$D(L_G) = \left\{ \varphi \in \ell^2 \ \left| \ \lim_{K \to \infty} \sum_{k=1}^K \frac{1}{k} \left(L^{*k} - L^k \right) \varphi \ exists \right\}, \\ L_G \varphi = i \sum_{k=1}^\infty \frac{1}{k} \left(L^{*k} - L^k \right) \varphi, \quad \varphi \in D(L_G). \right\}$$

Note that $\ell_{\text{fin}}^2 \subset D(L_G)$ and $i\{\log(\mathbb{1}-L) - \log(\mathbb{1}-L^*)\} \subset L_G$.

Theorem 5.4 (1) It holds that $L_G \cong T_G$ on ℓ_{fin}^2 .

(2) L_G is a bounded operator.

Proof: (1) Let U be the unitary operator defined by $Uv_n = \xi_n$ for each $n \in \mathbb{N}$. For arbitrary $\varphi \in D(T_G)$, we see that

$$UT_G\varphi = i\sum_{n=0}^{\infty} \left(\sum_{m < n} \frac{(\xi_m, U\varphi)}{n - m} + \sum_{m > n} \frac{(\xi_m, U\varphi)}{n - m}\right) \xi_n$$
$$= i\sum_{n=0}^{\infty} \left(\sum_{m < n} \frac{(L^{n-m}\xi_n, \varphi)}{n - m} - \sum_{m > n} \frac{(L^{*m-n}\xi_n, U\varphi)}{m - n}\right) \xi_n$$
$$= i\sum_{n=0}^{\infty} \left(\sum_{k=1}^{\infty} \frac{(L^k\xi_n, U\varphi)}{k} - \sum_{k=1}^{\infty} \frac{(L^{*k}\xi_n, U\varphi)}{k}\right) \xi_n.$$

From Lemma 5.1, we have $UD(T_G) = \ell_{\text{fin}}^2 \subset D(L_G)$. Then

$$UT_G\varphi = i\sum_{n=0}^{\infty} \left(\sum_{k=1}^{\infty} \frac{1}{k} \left(L^k - L^{*k}\right) \xi_n, U\varphi\right) \xi_n = \sum_{n=0}^{\infty} \left(\xi_n, i\sum_{k=1}^{\infty} \frac{1}{k} \left(L^{*k} - L^k\right) U\varphi\right) \xi_n.$$

This implies that $UT_G \varphi = L_G U \varphi$ for $\varphi \in D(T_G)$.

(2) From the Hilbert inequality, we see that, for any $\varphi = \sum_{n} c_n \xi_n \in D(L_G)$,

$$|(\varphi, L_G \varphi)| = \left| \sum_{n=0}^{\infty} \sum_{m \neq n} \frac{c_n c_m}{n - m} \right| \le \pi ||\varphi||^2.$$

Thus we have $||L_G\varphi|| \leq \pi ||\varphi||$. Then the theorem is proven.

Theorem 5.5 We have $[N, L_G] = -i\mathbb{1}$ on $D(NL_G) \cap D(L_GN)$.

Proof: We obtain that, for any $\varphi \in D(NL_G) \cap D(L_GN)$,

$$[N, L_G]\varphi = \sum_{n=0}^{\infty} \left(\xi_n, (NL_G - L_G N)\varphi\right)\xi_n$$

= $i\sum_{n=0}^{\infty}\sum_{k=1}^{\infty} \frac{1}{k} \left(\xi_n, \left(\left[N, L^{*k}\right] - \left[N, L^k\right]\right)\varphi\right)\xi_n$
= $i\sum_{n=0}^{\infty}\sum_{k=1}^{\infty} \left(\xi_n, \left(L^{*k} + L^k\right)\varphi\right)\xi_n.$ (5.3)

Let $\varphi = \sum_{n=0}^{\infty} c_n \xi_n$. Since $\sum_{n=0}^{\infty} \sum_{k=1}^{\infty} \left(\xi_n, \left(L^{*k} + L^k \right) \varphi \right) \xi_n = \sum_{n=0}^{\infty} \left(\sum_{m \neq n} c_m \right) \xi_n$, we have

$$\infty > \|[N, L_G]\varphi\|^2 = \sum_{n=0}^{\infty} |(\sum_{m \neq n} c_m)|^2.$$

In particular

$$\left\|\sum_{n=0}^{\infty} \left(\sum_{m\neq n} c_m\right) \xi_n + \varphi\right\|^2 = \sum_{n=0}^{\infty} |\sum_{m=0}^{\infty} c_m|^2 < \infty,$$

which implies that $\varphi \in \left\{ \sum_{n=0}^{\infty} c_n \xi_n \in \ell^2 \mid \sum_{n=0}^{\infty} c_n = 0 \right\}$, and

$$\sum_{n=0}^{\infty} \sum_{k=1}^{\infty} \left(\xi_n, \left(L^{*k} + L^k \right) \varphi \right) \xi_n = -\varphi.$$

Hence $[N, L_G]\varphi = -i\varphi$ and the theorem is proven.

It is immediate to see that $[N, T_G]\phi = -i\phi$ does not hold true for $\phi = v_n, n \in \mathbb{N}$. It is however shown in [9] that a CCR-domain of T_G is LH $\{v_n - v_m \mid n, m \in \mathbb{N}\}$. Note that

$$U(LH\{v_n - v_m \mid n, m \in \mathbb{N}\}) = (1 - L^*)\ell_{fin}^2.$$

Hence $[N, L_G] = -i\mathbb{1}$ holds on $(\mathbb{1} - L^*)\ell_{\text{fin}}^2$ and $(\mathbb{1} - L^*)\ell_{\text{fin}}^2 \subset D(NL_G) \cap D(L_GN)$.

Remark 5.6 To our knowledge, it was previously unknown that $[N, L_G] = -i\mathbb{1}$ holds on a strictly larger subspace than $(\mathbb{1} - L^*)\ell_{\text{fin}}^2$. It can be shown however that the CCR-domain $D(NL_G) \cap D(L_GN)$ of L_G is strictly larger than $(\mathbb{1} - L^*)\ell_{\text{fin}}^2$. It can be actually seen that $(\mathbb{1} - L^*)e^{\alpha L^*}\Omega \in (D(L_GN) \cap D(NL_G)) \setminus (\mathbb{1} - L^*)\ell_{\text{fin}}^2$ for $\alpha \neq 0$.

Remark 5.7 In (5.3), if $\varphi \in \operatorname{Ran}(\mathbb{1} - L^*)$, then we have

$$[N, L_G]\varphi = i\sum_{n=0}^{\infty}\sum_{k=1}^{\infty} \left(\xi_n, \left(L^{*k} + L^k\right)\varphi\right)\xi_n = i\left((\mathbb{1} - L^*)^{-1} + (\mathbb{1} - L)^{-1} - 2\mathbb{1}\right)\varphi.$$
(5.4)

(5.4) can be extended in (5.8).

5.2 Generalization of Galapon operator T_G

In Theorem 5.4 we show that T_G can be expressed as $i\{\log(\mathbb{1}-L) - \log(\mathbb{1}-L^*)\}$ on ℓ_{fin}^2 . We shall generalize this. We set

$$L_g = i \left\{ \log \left(\mathbb{1} - g_N L \right) - \log \left(\mathbb{1} - L^* g_N^{-1} \right) \right\}$$

Lemma 5.8 Let g be a complex-valued function on \mathbb{N} such that $|g_n| = 1$ for all $n \in \mathbb{N}$. Then L_g is unitary equivalent to L_G on ℓ_{fin}^2 .

Proof: We can construct the unitary operator V on ℓ^2 such that $g_N L = V^* L V$. It is actually given by

$$V(c_0, c_1, c_2, c_3, \dots,) = (c_0, g_0 c_1, g_0 g_1 c_2, g_0 g_1 g_2 c_3, \dots,).$$

Since $|g_n| = 1$, V is unitary. Thus L_g is unitary equivalent to L_G .

Let $\ell_1^2(g) = \left\{ \sum_{n=0}^{\infty} c_n \xi_n \in \ell^2 \mid \sum_{n=0}^{\infty} g!(n) c_n = 0 \right\}$, where $g!(n) = \prod_{k \le n} g_k$.

Theorem 5.9 Let g be a complex-valued function on \mathbb{N} .

- (1) Suppose that $|g_n| > 0$ for all $n \in \mathbb{N}$. Then $[N, L_g] = -i\mathbb{1}$ on $D(NL_g) \cap D(L_gN)$.
- (2) Suppose that $|g_n| = 1$ for all $n \in \mathbb{N}$. Then L_g is a bounded time operator with CCRdomain $D(NL_q) \cap D(L_qN)$.

Proof: As in the proof of Theorem 5.5, we see that $D(NL_g) \cap D(L_gN) \subset \ell_1^2(g)$. For any $\varphi \in D(NL_g) \cap D(L_gN)$, we have

$$[N, L_g]\varphi = i\sum_{n=0}^{\infty}\sum_{k=1}^{\infty} \left(\xi_n, \left(\left(L^* g_N^{-1}\right)^k + \left(g_N L\right)^k\right)\varphi\right)\xi_n = -i\varphi.$$

Then (1) is proven. Since L_G is self-adjoint and bounded, $\overline{L_g}$ is also self-adjoint and bounded from Lemma 5.8. By (1) L_g satisfies the canonical commutation relation.

 L_g is no longer a symmetric operator when $|g| \neq 1$. Therefore, in order to construct a time operator from L_g with $|g| \neq 1$, we need to symmetrize it.

Corollary 5.10 Let g be a complex-valued function on \mathbb{N} . Suppose that there exist $n_0 \in \mathbb{N}$, $c_1, c_2 \in (0, \infty)$ and $d_1, d_2 \in (0, 1/2)$ such that, for all $n > n_0$,

$$c_1 n^{-d_1} \le \prod_{k=0}^n |g_k| \le c_2 n^{d_2}.$$
 (5.5)

Then $(L_g + L_{\bar{g}^{-1}})/2$ is a time operator of N.

Proof: The assumption $c_1 n^{-d_1} \leq \prod_{k \leq n} |g_k|$ implies that for any $m \in \mathbb{N}$

$$\begin{aligned} \left\| \sum_{k \ge 1} \frac{1}{k} \left(L^* g_N^{-1} \right)^k \xi_m \right\| &= \left\| \sum_{k \ge 1} \frac{1}{k} \left(\prod_{l=0}^{k-1} g_{m+l} \right)^{-1} \xi_{m+k} \right\| \\ &\leq \left(\sum_{1 \le k \le n_0} \frac{1}{k^2} \left| \prod_{l=0}^{k-1} g_{m+l} \right|^{-2} + \left| c_1^{-1} g! (m-1) \right|^2 \sum_{k > n_0} k^{-2(1-d_1)} \right)^{1/2} < \infty \end{aligned}$$

Thus $\xi_m \in \mathcal{D}(\log(\mathbb{1}-L^*g_N^{-1}))$ and $\xi_m \in \mathcal{D}(L_g)$. In the same way, $\prod_{k=0}^n |g_k| \leq c_2 n^{d_2}$ implies that $\xi_n \in \mathcal{D}(\log(\mathbb{1}-L^*\bar{g}_N))$ and then $\xi_n \in \mathcal{D}(L_{\bar{g}^{-1}})$. Thus (5.5) implies that $\xi_n \in \mathcal{D}(L_g) \cap \mathcal{D}(L_{\bar{g}^{-1}})$. Since $L_g^* \supset -i \{\log(\mathbb{1}-L^*\bar{g}_N) - \log(\mathbb{1}-\bar{g}_N^{-1}L)\} = L_{\bar{g}^{-1}}$, the operator $L_g + L_{\bar{g}^{-1}}$ is symmetric. The canonical commutation relation $[N, (L_g + L_{\bar{g}^{-1}})/2] = -i\mathbb{1}$ is proven in the same way as Theorem 5.9.

5.3 Conjugate operators of the form $\log(1-p(L)) - \log(1-p(L^*))$

In Section 4.4 conjugate operators of the form $\log p(L)$ is considered. Let $Y_p = \mathbb{1} - p(L)$. In this section we consider conjugate operators of the form

$$X_p = \log(Y_p) - \log(Y_p^*),$$

under some conditions on polynomials p. X_p is a generalization of L_G . Note that $\text{Ker}(Y_p^*) = \{0\}$ if $p \neq 1$. Hence the inverse of Y_p^* exists.

Lemma 5.11 Let p be a polynomial. Suppose that Y_p is injective and $\lim_{k\to\infty} p(L)^k = 0$. Then

$$[N, \log Y_p] = Lp'(L)Y_p^{-1}$$
(5.6)

on $\operatorname{Ran}(Y_p) \cap \operatorname{D}(N \log Y_p) \cap \operatorname{D}(\log(Y_p)N)$ and

$$\left[N, \log(Y_p^*)\right] = -L^* p'(L^*) (Y_p^*)^{-1}$$
(5.7)

on $\operatorname{Ran}(Y_p^*) \cap \operatorname{D}\left(N \log(Y_p^*)\right) \cap \operatorname{D}\left(\log(Y_p^*)N\right).$

Proof: Let $\varphi \in D(N \log Y_p) \cap D(\log(Y_p)N)$ and $\varphi = Y_p \psi$. Inserting the expansion $\varphi = \sum_{n=0}^{\infty} (\xi_n, \varphi) \xi_n$ to $[N, \log Y_p] \varphi$, we see that

$$[N, \log Y_p] \varphi = -\sum_{n=0}^{\infty} \sum_{k=1}^{\infty} \frac{1}{k} \left(\xi_n, \left[N, p(L)^k \right] \varphi \right) \xi_n.$$

Since $[N, p(L)^k] \subset -kLp'(L)p(L)^{k-1}$, we have

$$[N, \log Y_p] \varphi = \sum_{n=0}^{\infty} \sum_{k=1}^{\infty} \left(\xi_n, Lp'(L)p(L)^{k-1}\varphi\right) \xi_n$$
$$= \sum_{n=0}^{\infty} \sum_{k=1}^{\infty} \left(\xi_n, Lp'(L)p(L)^{k-1}(\mathbb{1} - p(L))\psi\right) \xi_n.$$

By assumption $\lim_{k\to\infty} p(L)^k = 0$, we have

$$[N, \log Y_p]\varphi = -\sum_{n=0}^{\infty} \lim_{k \to \infty} \left(\xi_n, Lp'(L)p(L)^k\psi\right)\xi_n + \sum_{n=0}^{\infty} \left(\xi_n, Lp'(L)\psi\right)\xi_n = Lp'(L)Y_p^{-1}\varphi.$$

Then (5.6) follows. (5.7) can be shown in a similar way to (5.6).

From Lemma 5.11, if $Lp'(L)Y_p^{-1} + L^*p'(L^*)(Y_p^*)^{-1}$ has a non-zero eigenvalue, then N has a conjugate operator on the eigenvector space. Let $\mathbb{S}^1 = \{\alpha \in \mathbb{C} \mid |\alpha| = 1\}$. Thus, we have the following theorem.

Theorem 5.12 Let p be a polynomial with degree m such that

(1)
$$1 - p(L)$$
 is injective,
(2) $\lim_{k \to \infty} p(L)^k = 0$,
(3) $\{z \in \mathbb{C} \mid 1 - p(z) = 0\} \subset \mathbb{S}^1$

Let $\alpha_1, \ldots, \alpha_m \in \mathbb{C}$ be all roots of $1 - p(\alpha) = 0$ within the multiplicity. Then

$$\left[N,\frac{i}{m}X_p\right] = -i\mathbb{1}$$

with CCR-domain $\operatorname{Ran}\left(\prod_{i=1}^{m} (\alpha_i L^* - \mathbb{1})\right) \cap \operatorname{D}(X_p N) \cap \operatorname{D}(N X_p).$

Proof: By Lemma 5.11 we have

$$[N, X_p] = Lp'(L)Y_p^{-1} + L^*p'(L^*)(Y_p^*)^{-1}$$

= $\sum_{i=1}^m \left(\alpha_i (\alpha_i \mathbb{1} - L)^{-1} + \alpha_i (\alpha_i \mathbb{1} - L^*)^{-1} - 2\mathbb{1} \right).$ (5.8)

Note that (5.8) corresponds to (5.4). It suffices to show that the right-hand side above has an eigenvector in $D(NX_p) \cap D(X_pN)$. Since $\alpha_i \in \mathbb{S}^1$, we see that

$$\sum_{i=1}^{m} \left(\alpha_i (\alpha_i \mathbb{1} - L)^{-1} + \alpha_i (\alpha_i \mathbb{1} - L^*)^{-1} \right) = \sum_{i=1}^{m} \left(\alpha_i (\alpha_i \mathbb{1} - L)^{-1} + \alpha_i^{-1} (\alpha_i^{-1} \mathbb{1} - L^*)^{-1} \right).$$

Let $\psi \in \ell^2 \setminus \{0\}$ and $\varphi = (\alpha L^* - \mathbb{1})\psi$. Since

$$\left(\alpha(\alpha\mathbb{1} - L)^{-1} + \alpha^{-1}(\alpha^{-1}\mathbb{1} - L^*)^{-1}\right)\varphi = \alpha L^*\psi - \psi = \varphi,$$

 φ is an eigenvector of $\alpha(\alpha \mathbb{1} - L)^{-1} + \alpha^{-1}(\alpha^{-1}\mathbb{1} - L^*)^{-1}$ with an eigenvalue 1. Hence $\prod_{i=1}^{m} (\alpha_i L^* - \mathbb{1}) \psi$ is an eigenvector of $Lp'(L)Y_p^{-1} + L^*p'(L^*)(Y_p^*)^{-1}$ with the eigenvalue -1 and

$$[N, X_p] \prod_{i=1}^m (\alpha_i L^* - 1) \psi = -m \prod_{i=1}^m (\alpha_i L^* - 1) \psi.$$

Then a CCR-domain is given by $\operatorname{Ran}(\prod_{i=1}^{m}(\alpha_i L^* - 1)) \cap D(X_p N) \cap D(NX_p)$, and the theorem is proven.

Corollary 5.13 Let $|\omega| = 1$. Then $im^{-1} \{ \log(1 - \omega L^m) - \log(1 - \omega^* L^{*m}) \}$ is a time operator of N.

Proof: Let $p(z) = \omega z^m$. Then the corollary follows from Theorem 5.12.

6 Classification of conjugate operators

In this section we focus on investigating conjugate operators of the form

$$T_{\omega,m} = \frac{i}{m} \log(\omega \mathbb{1} - L^m) \quad (\omega,m) \in \overline{\mathbb{D}} \times (\mathbb{N} \setminus \{0\})$$

This type of operators have appeared in Lemma 5.1 and Corollary 5.13. Furthermore note that $T_A \cong L_{A0} \oplus L_{A1}$. In Example 4.18 we mentioned that to investigate the CCR-domain of L_{A0} and L_{A1} can be reduced to investigate the CCR-domain of

$$\frac{i}{2}\log L^2 = T_{0,2}$$

Let us define the family of operators with parameters ω and m by

$$\mathscr{T} = \{T_{\omega,m} \mid \omega \in \overline{\mathbb{D}}, m \ge 1\}.$$

We divide \mathscr{T} into three disjoint families:

$$\mathscr{T} = \mathscr{T}_{\{0\}} \cup \mathscr{T}_{\mathbb{D} \setminus \{0\}} \cup \mathscr{T}_{\partial \mathbb{D}}.$$

Here

$$\begin{aligned} \mathscr{T}_{\{0\}} &= \{ T_{\omega,m} \mid \omega = 0, m \ge 1 \}, \\ \mathscr{T}_{\mathbb{D} \setminus \{0\}} &= \{ T_{\omega,m} \mid 0 < |\omega| < 1, m \ge 1 \}, \\ \mathscr{T}_{\partial \mathbb{D}} &= \{ T_{\omega,m} \mid |\omega| = 1, m \ge 1 \}. \end{aligned}$$

In what follows we discuss CCR-domains of $T_{\omega,m}$.

We define

$$\ell_m^2 = \left\{ \varphi \in \ell^2 \ \Big| \ \lim_{k \to \infty} (\mathbb{1} - L^m)^k \varphi = 0 \right\},$$

Theorem 6.1 $(\mathscr{T}_{\{0\}})$ We have

$$\left[N, \frac{i}{m} \log L^m\right] = -i\mathbb{1}$$

 $on \ \ell_m^2 \cap \mathcal{D}(N \log L^m) \cap \mathcal{D}(\log(L^m)N), \ and \ \dim\left(\ell_m^2 \cap \mathcal{D}(N \log L^m) \cap \mathcal{D}(\log(L^m)N)\right) = \infty.$

Proof: This is a special case of Theorem 4.19. Let $p(z) = z^m$. Then zp'(z) - mp(z) = 0 for all $z \in \mathbb{C}$. In a similar way of the proof of Theorem 4.19, for any $\varphi \in \ell_m^2 \cap D(N \log L^m) \cap D(\log(L^m)N)$, we see that $[N, \log L^m]\varphi = -m\sum_{k=1}^{\infty} (\mathbb{1} - L^m)^{k-1} L^m \varphi = -m\varphi$. Then the theorem follows.

Next we consider $\mathscr{T}_{\mathbb{D}\setminus\{0\}}$. Let $c \in \mathbb{C}$ and we recall that

$$D(\omega - z^m, c) = LH\{\varphi \mid \exists \alpha \in \mathbb{C} \ s.t. \ |1 - \omega + \alpha^m| < 1, \ \alpha^m = c(m + c)^{-1}\omega, \ \varphi \in Ker(L - \alpha \mathbb{1})\}$$

Theorem 6.2 $(\mathscr{T}_{\mathbb{D}\setminus\{0\}})$ Let $\omega \in \mathbb{D}\setminus\{0\}$. Then for $c \in \mathbb{C}\setminus\{0\}$ such that |c| is sufficiently small, we see that $D(\omega - z^m, c) \neq \{0\}$, dim $D(\omega - z^m, c) < \infty$ and

$$\left[N, -\frac{i}{c}\log(\omega\mathbb{1} - L^m)\right] = -i\mathbb{1}$$

on $D(\omega - z^m, c)$. Moreover, for any $c \in \mathbb{C} \setminus \{0\}$, there is no infinite dimensional CCRdomain for N and $c \log(\omega - L^m)$

Proof: Let $p(z) = \omega - z^m$ with $|\omega| < 1$ and $|\omega - 1| < 1$. Suppose that c is a sufficiently small positive number. Then $zp'(z) + cp(z) = -(c+m)z^m + c\omega = 0$ has the roots

$$\alpha_k = \left| \frac{c}{c+m} \right| \omega^{1/m} e^{2\pi i k/m}, \quad k = 0, \dots, m-1.$$

By Theorem 4.19, since $|p(\alpha_k) - 1| = |\omega - 1 - \alpha_k^m| < 1$ and $|\alpha_k| < 1$, $k = 0, \ldots, m - 1$, we have $[N, \frac{i}{c} \log(\omega \mathbb{1} - L^m)] = i\mathbb{1}$ on $\operatorname{Ker}(L - \alpha_k \mathbb{1})$ for $k = 0, \ldots, m - 1$. Moreover the dimension of the CCR-domain $\operatorname{LH}\{\operatorname{Ker}(L - \alpha_k \mathbb{1}), k = 0, \ldots, m - 1\}$ is finite. \Box

We consider conjugate operators of the form $i\{\log(\omega \mathbb{1} - L) - \log(\bar{\omega} \mathbb{1} - L^*)\}$ with $\omega \in \partial \mathbb{D}$. We consider $\log(\omega \mathbb{1} - A)$ for $\log(\omega \mathbb{1} - A)$ to avoid a singularity. **Definition 6.3** $(Log(\omega 1 - A))$ Let A be a linear operator on a Hilbert space \mathcal{H} . We define

$$D(Log(\omega 1 - A)) = \left\{ \varphi \in \bigcap_{k=0}^{\infty} D(A^k) \ \left| \ \lim_{K \to \infty} \sum_{k=1}^{K} \frac{1}{k} \left(\frac{1}{\omega} A \right)^k \varphi \ exists \right\}, \\ Log(\omega 1 - A)\varphi = log(\omega)\varphi - \sum_{k=1}^{\infty} \frac{1}{k} \left(\frac{1}{\omega} A \right)^k \varphi, \quad \varphi \in D(Log(\omega 1 - A)).$$
(6.1)

Theorem 6.4 $(\mathscr{T}_{\partial \mathbb{D}})$ Let $\omega \in \partial \mathbb{D}$. Then $L_{m,\omega} = im^{-1}(\operatorname{Log}(\omega \mathbb{1} - L^m) - \operatorname{Log}(\bar{\omega} \mathbb{1} - L^{*m}))$ is a time operator with the dense CCR-domain $D(NL_{m,\omega}) \cap D(L_{m,\omega}N)$.

Proof: As in the proof of Theorem 5.5, we see that

$$\mathcal{D}(NL_{m,\omega}) \cap \mathcal{D}(L_{m,\omega}N) \subset \left\{ \sum_{n=0}^{\infty} c_n \xi_n \right| \sum_{n=0}^{\infty} \bar{\omega}^n c_{l+mn} = 0 \text{ for all } l < m \right\}$$

and for any $\varphi \in D(NL_{m,\omega}) \cap D(L_{m,\omega}N)$,

$$[N, L_{m,\omega}]\varphi = i\sum_{n=0}^{\infty}\sum_{k=1}^{\infty} \left(\xi_n, \left((\bar{\omega}L^m)^k + (\omega L^{*m})^k\right)\varphi\right)\xi_n = -i\varphi.$$

Then the theorem follows.

	$\mathscr{T}_{\{0\}}$	$\mathscr{T}_{\mathbb{D}\setminus\{0\}}$	$\mathscr{T}_{\partial\mathbb{D}}$
$T_{\omega,m}$	$\omega = 0$	$\omega \in \mathbb{D} \setminus \{0\}$	$\omega\in\partial\mathbb{D}$
boundedness	unbounded	unbounded	bounded
CCR-domain	infinite dim.	finite dim.	dense
example	T_A		$T_G = T_{1,1} + T_{1,1}^*$
log	Def 3.1	Def 3.1	Def 6.3

Table 1: Classification of $T_{\omega,m}$

We can also introduce operator $T_{\omega,m}$ with $|\omega| > 1$ by a Dunford integral. See (A.1) in Appendix.

7 Weak Wyle relations for N

In this section we consider the time evolution of conjugate operators. The Weyl relation [19]

$$e^{-isp}e^{-itq} = e^{ist}e^{-itq}e^{-isp}, \quad s, t \in \mathbb{R}.$$

-		-
L		
L		
L		┛

holds true. From this we can derive the so-called weak Weyl relation [14]:

$$qe^{-itp} = e^{-itp}(q+t), \quad t \in \mathbb{R},$$
(7.1)

on D(q). The strong time operator T is defined as an operator satisfying (7.1) with q and p replaced by T and Hamiltonian H, respectively. More precisely, we explain the weak Weyl relation (7.1) as follows.

Definition 7.1 We say that the pair (H, A) consisting of a self-adjoint operator H and a symmetric operator A on a Hilbert space \mathcal{H} obeys the weak Weyl relation if for all $t \in \mathbb{R}$,

(1) $e^{-itH} D(A) \subset D(A);$ (2) $Ae^{-itH} \varphi = e^{-itH} (A+t) \varphi$ for all $\varphi \in D(A).$

Here A is referred to as a strong time operator associated with H. Note that a strong time operator is not unique. If strong time operator A is self-adjoint, then it is known that

$$e^{-isA}e^{-itH} = e^{-ist}e^{-itH}e^{-isA}, \quad s, t \in \mathbb{R}$$

holds. In particular when Hilbert space is separable, by the von Neumann uniqueness theorem the Weyl relation above implies that H and A are unitarily equivalent to $\bigoplus^n p$ and $\bigoplus^n q$ with some $n \leq \infty$, respectively. As was mentioned above, although a strong time operator is automatically a time operator, the converse is not true. It is remarkable that when the pair (H, A) obeys the weak Weyl relation, H has purely absolutely continuous spectrum. For example since the spectrum of N is purely discrete, there is no strong time operator associated with N. Weak Weyl relation can be understood as the time evolution of A:

$$e^{itH}Ae^{-itH} \supset A+t \tag{7.2}$$

We can see the time evolution of the Galapon operator directly. It can be seen that

$$T_G(t)\varphi = i\sum_{n=0}^{\infty}\sum_{m\neq n}\frac{e^{+it(n-m)}(v_m,\varphi)}{n-m}v_n.$$

Hence $T_G(t)$ is periodic so that

$$T_G(t+2\pi n) = T_G(t), \quad n \in \mathbb{N}, t \in \mathbb{R}.$$

Let us consider the time evolutions of general conjugate operators. We define

$$T_{\omega,m}(t) = e^{itN} T_{\omega,m} e^{-itN}.$$

Theorem 7.2 Let $T_{\omega,m} \in \mathscr{T}$. Then

$$T_{\omega,m}(t) = \frac{i}{m}\log(\omega - e^{-itm}L^m)$$

and $T_{\omega,m}(t)$ is periodic with period $2\pi/m$, i.e.,

$$T_{\omega,m}(t) = T_{\omega,m}(t + (2\pi/m)n), \quad n \in \mathbb{N}, t \in \mathbb{R}.$$

Proof: Since $e^{itN}Le^{-itN} = e^{-it}L$, we see that $e^{itN}\log(\omega\mathbb{1} - L^m)e^{-itN} = \log(\omega - e^{-itm}L^m)$.

Example 7.3 It is shown that the angle operator $T_A = \frac{1}{2}(\arctan q^{-1}p \oplus \arctan pq^{-1})$ can be represented as

$$\frac{1}{2}\arctan\left(q^{-1}p\right) \cong \frac{i}{2}\log\left(\sqrt{\frac{N+2\mathbb{1}}{N+1\mathbb{1}}}L^2\right),$$
$$\frac{1}{2}\arctan\left(pq^{-1}\right) \cong \frac{i}{2}\log\left(\sqrt{\frac{N+1\mathbb{1}}{N+2\mathbb{1}}}L^2\right)$$

in Theorem 4.9. Thus the period of the time evolutions of both of $\arctan(q^{-1}p)$ and $\arctan(pq^{-1})$ are π .

Finally we show an example of (3.2). Let $\omega = 0$ and consider the time evolution of $T_{0,m}(t)$.

Corollary 7.4 Let $m \in \mathbb{N}$ and $\arg e^{-itm} \neq 0$. Then

$$\log(-e^{-itm}L^m) \neq \log\left(e^{-itm}\right) + \log(-L^m).$$

Proof: Suppose that $\log(-e^{-itm}L^m) = \log(e^{-itm}) + \log(-L^m)$. Then we see that

$$e^{itN}T_{0,m}e^{-itN} = \frac{i}{m}\log(-e^{-itm}L^m) = -\frac{1}{m}\arg e^{-itm} + \frac{i}{m}\log(-L^m) = -\frac{1}{m}\arg e^{-itm} + T_{0,m}.$$

Hence we see that for any eigenvector v_n of N

$$(v_n, T_{0,m}v_n) = (v_n, e^{itN}T_{0,m}e^{-itN}v_n) = -\frac{1}{m}(v_n, \arg e^{-itm}v_n) + (v_n, T_{0,m}v_n).$$
(7.3)

Since $(v_n, \arg e^{-itm}v_n) \neq 0$, (7.3) is contradiction. Then the corollary follows. \Box

A CCR domains for $\mathscr{T}_{\overline{\mathbb{D}}^c}$

We consider conjugate operators of the form $i \log(\omega \mathbb{1} - L^m)$ with $|\omega| > 1$. To avoid a singularity, we need to modify the definition of $i \log(\omega \mathbb{1} - L^m)$. Let $r\mathbb{D} = \{rz \mid z \in \mathbb{D}\}$ for r > 0 and f be an analytic function such that $f(\overline{r\mathbb{D}}) \not\supseteq 0$ for some r > 1. Then $\log f$ is analytic on $r\mathbb{D}$. Let ||A|| < 1. Since $(z - A)^{-1}$ is bounded for any $z \not\in \overline{\mathbb{D}}$, it can be seen that the Dunford integral $\oint_{r\partial \mathbb{D}} \log f(z)(z - A)dz$ defines the sesquilinear form

$$(\varphi, \psi) \mapsto Q(\varphi, \psi) = \oint_{r \partial \mathbb{D}} \log f(z)(\varphi, (z - A)^{-1}\psi) dz.$$
 (A.1)

(A.1) also defines the bounded operator such that $Q(\varphi, \psi) = (\varphi, B\psi)$. *B* is also denoted by the same symbol $\log f(A)$ as that of Definition 3.1. Set $f(z) = \omega - z^m$ and A = L. Then we can define $\log(\omega \mathbb{1} - L^m)$ by (A.1).

Lemma A.1 There is no polynomial p satisfying the following conditions:

- (1) $0 \notin \operatorname{Spec}(p(L)),$
- (2) $[N, i\{\log p(L) \log p(L^*)\}] = -i\mathbb{1}$ on some infinite dimensional subspace \mathcal{D} .

Proof: Let $p(x) = \sum_{k=0}^{n} a_k x^k$ be a real-valued polynomial with $a_n \neq 0$. We see that

$$p(L)p(L^*)L^{*n} = \left(\sum_{k=0}^n a_k L^{*n-k}\right) \left(\sum_{k=0}^n a_k L^{*k}\right) = \sum_{k,l\ge 0}^n a_k a_l L^{*n-l+k}.$$
 (A.2)

It is easy to see that

$$[N, \log p(L) - \log p(L^*)] \subset -Lp'(L)p(L)^{-1} - L^*p'(L^*)p(L^*)^{-1}.$$
 (A.3)

Note that $\operatorname{Ran}(L^{*n}) = \{(c_k)_{k \in \mathbb{N}} \in \ell^2 \mid c_0 = \cdots = c_{n-1} = 0\}$. Hence $\mathcal{D} \cap \operatorname{Ran}(L^{*n}) = \{0\}$ implies that $\dim \mathcal{D} < \infty$. Then $\mathcal{D} \cap \operatorname{Ran}(L^{*n}) \neq \{0\}$, since $\dim \mathcal{D} = \infty$. Notice that, for any $k \leq n$, $L^k L^{*k} = L^{*k} L^k = 1$ on $\operatorname{Ran}(L^{*n})$. Thus we also have the following relation on the non-trivial domain by (A.3):

$$p(L)p(L^{*})L^{*n}$$

$$= -p(L)p(L^{*})[N, \log p(L) - \log p(L^{*})]L^{*n}$$

$$= Lp'(L)p(L^{*})L^{*n} + L^{*}p'(L^{*})p(L)L^{*n}$$

$$= \left(\sum_{k=1}^{n} ka_{k}L^{*n-k}\right) \left(\sum_{k=0}^{n} a_{k}L^{*k}\right) + \left(\sum_{k=1}^{n} ka_{k}L^{*k}\right) \left(\sum_{k=0}^{n} a_{k}L^{*n-k}\right)$$

$$= \sum_{k,l\geq 1}^{n} (ka_{k}a_{l} + la_{l}a_{k})L^{*n-l+k} + \sum_{k=1}^{n} ka_{k}a_{0}L^{*n-k} + \sum_{k=1}^{n} ka_{k}a_{0}L^{*n+k}.$$
(A.4)

Let $L^{*n}\varphi \in \mathcal{D} \setminus \{0\}$ and $n_0 = \inf \operatorname{supp}(\varphi)$. Comparing the coefficients of L^{*0} in (A.2) and (A.4), we have $na_0a_n = a_0a_n$. From (1) we have $a_0 \neq 0$. Since $a_0a_n \neq 0$, we see that n = 1. Hence p(x) = ax + b. By (A.4) we also have $a = \pm b$, and then $p(x) = a(1 \pm x)$. We conclude that $p(L) = a(1 \pm L)$ and $\operatorname{Spec}(p(L)) = p(\operatorname{Spec}(L)) \ni 0$. This contradicts $0 \notin \operatorname{Spec}(p(L))$. Then the lemma follows.

Theorem A.2 Let $|\omega| > 1$. The operator $i\{\log(\omega \mathbb{1} - L^m) - \log(\bar{\omega} \mathbb{1} - L^{*m})\}$ has no infinite dimensional CCR-domain for N.

Proof: Let $p(z) = \omega - z^m$ with $|\omega| > 1$. Then $\operatorname{Spec}(p(L)) \not\supseteq 0$. Thus the theorem follows from Lemma A.1.

Remark A.3 The existence of a non zero CCR domain of $i\{\log(\omega \mathbb{1} - L) - \log(\bar{\omega} \mathbb{1} - L^*)\}$ with $|\omega| > 1$ is unknown.

Acknowledgements: FH is financially supported by JSPS KAKENHI 20K20886 and JSPS KAKENHI 20H01808.

References

- Y. Aharonov and D. Bohm. Time in the quantum theory and the uncertainty relation for time and energy, *Phys. Rev.*, 122:1649–1658, 1961.
- [2] P. Aiena. Fredholm and local spectral theory, with applications to multipliers. Kluwer Academic Publishers, Dordrecht, 2004.
- [3] A. Arai. Necessary and sufficient conditions for a Hamiltonian with discrete eigenvalues to have time operators. *Lett. Math. Phys.*, 87:67–80, 2009.
- [4] A. Arai and F. Hiroshima. Ultra-weak time operators of Schrödinger operators. Ann. Henri Poincaré, 18:2995–3033, 2017.
- [5] A. Arai and Y. Matsuzawa. Time operators of a Hamiltonian with purely discrete spectrum. *Rev. Math. Phys.*, 20:951–978, 2008.
- [6] M. Bauer. A time operator in quantum mechanics. Ann. Phys., 150:1–21, 1983.
- [7] P. Carruthers and M.M. Niet. Phase and angle variable in quantum mechanics. *Rev. Mod. Phys.*, 40:411–440, 1968.
- [8] E.A. Galapon. Pauli's theorem and quantum mechanical pairs: the consistency of bounded, self-adjoint time operator canonically conjugate to a hamiltonian with non-empty point spectrum. Proc. R. Soc. Lond. A, 458:451–472, 2002.
- [9] E.A. Galapon. Self-adjoint time operator is the rule for discrete semi-bounded Hamiltonians. Proc. R. Soc. Lond. A, 458:2761–2689, 2002.

- [10] F. Hiroshima and N. Teranishi. Time operators of harmonic oscillators and their representations. to be published in J. Math. Phys., 2024.
- [11] H.R. Lewis, W.E. Lawrence, and J.D. Harris. Quantum Action-Angle Variables for the Harmonic Oscillator. Phys. Rev. Lett., 77:5157–5159, 1996.
- [12] H.R. Lewis, W.E. Lawrence, and J.D. Harris. A Reply to the Comment by T. B. Smith and John A. Vaccaro. *Phys. Rev. Lett.*, 80:2746–2746, 1998.
- [13] L. Mandelstam and Ig. Tamm. The uncertainty relation between energy and time in nonrelativistic quantum mechanics. J. Phys., 1945.
- [14] M. Miyamoto. A generalized Weyl relation approach to the time operator and its connection to the survival probability. J. Math. Phys., 42:1038–1052, 2001.
- [15] G. Muga, R.S. Mayato, and I. Egusquiza. *Time in quantum mechanics-vol.1*. Springer, 2007.
- [16] D.T. Pegg and S.M. Barnett. Phase of properties the quantized single-mode electromagnetic field. Phys. Rev. A, 39:1665–1675, 1989.
- [17] T.B. Smith and John.A. Vaccaro. Comment on "Quantum Action-Angle Variables for the Harmonic Oscillator". Phys. Rev. Lett., 80:2745–2745, 1998.
- [18] L. Susskind and J. Glogower. Quantum mechanical phase and time operator. *Physics*, 1:49–61, 1964.
- [19] H. Weyl. Gruppentheorie und Quantenmechanik. Hirzel, Leipzig, 1928.