

# GENERIC DOUBLING OF RECTANGULAR PEGS

ZHEN GAO

With an appendix joint with Urs Frauenfelder

ABSTRACT. In this paper, we prove a multiplicity result for rectangular pegs that there is a generic class of smooth Jordan curves in which every curve admits two geometrically distinct similar inscribed rectangles with aspect angle in  $(0, \frac{\pi}{2})$ , based on the a priori existence result of rectangular peg by Greene and Lobb [GL21], in which the rectangular peg problem is translated into finding intersection points of two Lagrangian tori. The key insights are certain generic geometric transversality of the two tori, and the correspondence between the intersection points and the inscribed rectangles. We present two distinct proofs: one involves the Lagrangian Floer homology, and the other is differential topological in nature which employs a novel computation formula for the algebraic intersection number. Moreover, such generic doubling result can be further extended to cyclic quadrilateral pegs based on [GL23]. In Appendix B, written jointly with Urs Frauenfelder, we provide the detailed derivation of this computation formula.

## CONTENTS

1. Introduction	2
2. Inscribed rectangles and intersection points	4
3. Geometric transversality	8
4. Proof of the main theorem 1.6	13
5. Comments on a priori existence of rectangular peg	17
6. Generic doubling of cyclic quadrilateral pegs	19
Appendix A. Lagrangian Floer homology	22
A.1. Floer homology of monotone Lagrangian submanifolds	22
A.2. Morse-Bott Floer homology	27
A.3. Liouville manifold	28
Appendix B. Algebraic intersection number for clean intersections	30
B.1. Introduction	30
B.2. The clean normal bundle	32
B.3. Euler characteristic and algebraic intersection number	35
B.4. Categorification	38
References	39

## 1. INTRODUCTION

The inscribed square problem a.k.a. square peg problem firstly posed by Otto Toeplitz in 1911, c.f. [Toe11], asks the following question:

*Does every continuous simple closed curve a.k.a. Jordan curve inscribe a square?*

More precisely, a square is inscribed on a Jordan curve means that there are four points in the Jordan curve as four vertices of the square. The square peg problem has been affirmatively answered in various cases, yet remains open in its full generality to this day. We refer to [Mat14] and [GL21, §2] for historical survey and recent developments. There are many variants of the square peg problem, see e.g. [Mat14, p.349-351] and [Mat22]. One of the closely related problems to the square peg problem is the so-called rectangular peg problem. We list some important breakthroughs: In 1977, Vaughan showed that every continuous Jordan curve inscribes a rectangle (note that there is no knowledge about its aspect angle which is the angle between the two diagonals); from 2018 to 2019, Hugelmeyer showed the existence of rectangular peg on any smooth Jordan curve with any aspect angle lying in a subset of  $(0, \frac{\pi}{2}]$  whose Lebesgue measure is at least  $\frac{\pi}{6}$ ; in 2020, the smooth rectangular peg problem in full generality, i.e. for any smooth Jordan curve and with any aspect angle in  $(0, \frac{\pi}{2}]$ , was proved by Greene and Lobb, c.f. [GL21]. In particular, the result of Greene and Lobb in [GL21] solves the square peg problem in smooth case. Further study by Greene and Lobb generalizes such result to cyclic quadrilateral pegs, c.f. [GL23]. The most important and interesting feature of their proofs is involving highly non-trivial symplectic topology. We review the geometric setup in their paper [GL21] below, and state the main result that we obtain based on their work.

Let  $\gamma$  be a smooth simple closed curve in  $\mathbb{C}$ , i.e. a smooth Jordan curve. This is obviously a Lagrangian closed curve in  $(\mathbb{C}, \omega_{\text{std}})$  with  $\omega_{\text{std}} = \frac{i}{2} dz \wedge d\bar{z}$  in terms of coordinate  $z \in \mathbb{C}$ . The product  $\gamma \times \gamma$  is then a smooth Lagrangian torus in  $(\mathbb{C}^2, \omega_{\text{std}})$  where

$$\omega_{\text{std}} := \frac{i}{2}(dz_1 \wedge d\bar{z}_1 + dz_2 \wedge d\bar{z}_2)$$

in terms of coordinate  $(z_1, z_2) \in \mathbb{C}^2$ . Now we consider the map

$$(1.1) \quad \begin{aligned} l : \mathbb{C}^2 &\rightarrow \mathbb{C}^2 \\ (z_1, z_2) &\mapsto \left( \frac{z_1 + z_2}{2}, \frac{z_1 - z_2}{2} \right) \end{aligned}$$

which is a diffeomorphism and  $l^*\omega_{\text{std}} = \frac{1}{2}\omega_{\text{std}}$ . This yields that  $L := l(\gamma \times \gamma)$  is also a smooth Lagrangian torus in  $(\mathbb{C}^2, \omega_{\text{std}})$ .

Applying the rotation on the second factor

$$(1.2) \quad \begin{aligned} R_\phi : \mathbb{C}^2 &\rightarrow \mathbb{C}^2 \\ (z_1, z_2) &\mapsto (z_1, z_2 e^{i\phi}) \end{aligned}$$

to the Lagrangian torus  $L$  we obtain another Lagrangian torus  $L_\phi := R_\phi(L)$ . They intersect cleanly along  $\gamma \times \{0\}$ :

$$T_p L \cap T_p L_\phi = T_p(\gamma \times \{0\}) \quad \forall p \in \gamma \times \{0\},$$

which is  $z_1 = z_2$  with  $z_1, z_2 \in \gamma$  in  $l(\gamma \times \gamma)$ . There is a symplectic involution

$$(1.3) \quad \begin{aligned} \varrho : \mathbb{C}^2 &\rightarrow \mathbb{C}^2 \\ (z_1, z_2) &\mapsto (z_1, -z_2) \end{aligned}$$

preserving both  $L$  and  $L_\phi$  and fixing  $\gamma \times \{0\}$ .

The goal is to find an intersection point outside  $\gamma \times \{0\}$ , i.e. one needs to show that

$$(1.4) \quad \#(L \cap L_\phi \setminus (\gamma \times \{0\})) \geq 1.$$

Suppose such intersection point exists and reads in the coordinate as  $(z_1, z_2 e^{i\phi})$ , then it follows that four points  $z_1 \pm z_2, z_1 \pm z_2 e^{i\phi}$  all lie on the curve  $\gamma$  and are four vertices of a rectangle whose diagonals meet at the angle  $\phi \in (0, \frac{\pi}{2}]$  in  $\mathbb{C}$ .

The union  $L \cup L_\phi$  has at least  $\gamma \times \{0\}$  in the self-intersection set. After a Lagrangian smoothing surgery of  $L \cup L_\phi$  along  $\gamma \times \{0\}$  (c.f. [GL21, Proposition 1.1]), one gets a smoothly immersed Lagrangian torus  $L \#_{\gamma \times \{0\}} L_\phi$  in  $\mathbb{C}^2$  that coincides with  $L \cup L_\phi$  away from a small open neighborhood  $\mathcal{O}p(\gamma \times \{0\})$ , and it is disjoint from  $\mathbb{C} \times \{0\}$ . The involution  $\varrho$  now acts as a fixed-point free involution on the immersed Lagrangian torus  $L \#_{\gamma \times \{0\}} L_\phi$ .

In [GL21], the following map is considered:

$$g : \mathbb{C}^2 \rightarrow \mathbb{C}^2 \\ (z_1, z_2) \mapsto \left( z_1, \frac{1}{\sqrt{2}} \frac{z_2^2}{|z_2|} \right)$$

which is a symplectomorphism away from  $\mathbb{C} \times \{0\}$ . Applying to the Lagrangian tori  $L$  and  $L_\phi$ , their image  $g(L)$  and  $g(L_\phi)$  are both homeomorphic to a Möbius strip  $\text{Sym}^2(\gamma)$ , and are smooth Lagrangian away from  $\mathbb{C} \times \{0\}$ . The image  $g(L \#_{\gamma \times \{0\}} L_\phi)$  is therefore a smoothly immersed Lagrangian Klein bottle  $K \looparrowright \mathbb{C}^2$  such that

$$g(L \#_{\gamma \times \{0\}} L_\phi \setminus \mathcal{O}p(\gamma \times \{0\})) \cong g(L \cup L_\phi \setminus \mathcal{O}p(\gamma \times \{0\}))$$

where  $\mathcal{O}p(\gamma \times \{0\})$  is a small open neighborhood of  $\gamma \times \{0\}$  on which the immersion restricted is a smooth embedding in  $\mathbb{C}^2$ . Using the result of non-existence of embedded Lagrangian Klein bottle in  $\mathbb{C}^2$ , c.f. [She09, Nem09], they conclude that the set of self-intersection points of the immersed Lagrangian Klein bottle must be non-empty, and hence the existence of rectangular peg is a corollary.

**Theorem 1.5** ([GL21]). *For every smooth Jordan curve and rectangle with any aspect angle  $\phi \in (0, \frac{\pi}{2}]$  in the  $\mathbb{R}^2$ , there exists a similar rectangle whose vertices land on the curve.*

In the work [GL23], which generalizes to arbitrary cyclic quadrilateral albeit in different geometric setup. Instead of using the Lagrangian Klein bottle argument, they compute the minimal Maslov number of the surgered immersed Lagrangian torus. See §6.

Beyond addressing the existence problem, the investigation into the multiplicity of inscribed rectangles remains largely unexplored. Leveraging the aforementioned a priori existence result of the rectangular pegs by Greene and Lobb, our primary contribution lies in the following main theorem.

**Theorem 1.6.** *For any fixed  $\phi \in (0, \frac{\pi}{2})$ , there is a generic class of smooth Jordan curves such that any rectangle with the fixed aspect angle  $\phi$  in  $\mathbb{R}^2$  admits two similar rectangles whose vertices land distinctly on any curve in this generic class.*

**Remark 1.7.** A counter example of  $\phi = \frac{\pi}{2}$  for the statement: Consider an ellipse (noticing that ellipses are in the generic class of curves, see Example 3.4) and it is obvious there is only one inscribed square.

**Remark 1.8.** In fact, such generic doubling result can be further extended to inscribed cyclic quadrilaterals by adopting the geometric setup in [GL23]. See §6, Theorem 6.2.

We present two distinct proofs for Theorem 1.6 in §4. These proofs rely on two essential components: The correspondence between intersection points in  $L \cap L_\phi$  away from  $\mathbb{C} \times \{0\}$ , as described in §2, and a generic geometric transversality property for  $L$  and  $L_\phi$ , elaborated in §3. We give the sketch of the proofs as follows. The first proof involves computing the Euler characteristic of the Lagrangian Floer homology for the Lagrangian tori  $L$  and  $L_\phi$ , generated from a generic curve  $\gamma$ , with the geometric transversality property. This approach applies the invariance of the Lagrangian Floer homology to draw a contradiction when assuming the existence of only one intersection point away from  $\mathbb{C} \times \{0\}$ . The second proof follows a similar path of contradiction but employs a novel formula for the algebraic intersection number of cleanly intersecting Lagrangian submanifolds. In §6, we further extend the main result to inscribed cyclic quadrilaterals. In Appendix A, we provide an overview of the Lagrangian Floer homology suited to our geometric setup. Additionally, Appendix B furnishes a detailed derivation of the computation formula for the algebraic intersection number of cleanly intersecting submanifolds.

*Acknowledgment.* I am deeply indebted to my Ph.D. supervisor, Urs Frauenfelder, for the enlightening conversations and insightful ideas provided throughout this research. I am grateful to Kai Cieliebak and Felix Schmäschke for the valuable comments and fruitful discussions. I also thank Marián Poppr for pointing out the reference [Kar13] to me. The authors acknowledge the partial support from DFG grant FR 2637/4-1.

## 2. INSCRIBED RECTANGLES AND INTERSECTION POINTS

Intersection points in  $L \cap L_\phi$  away from  $\mathbb{C} \times \{0\}$  do not one-to-one correspond to distinguished inscribed rectangle on the smooth Jordan curve  $\gamma$ .

The tori  $L$  and  $L_\phi$  in  $\mathbb{C}^2$  are parameterized by  $S^1 \times S^1$ :

$$\begin{aligned} S^1 \times S^1 &\rightarrow L \subset \mathbb{C}^2 \\ (t_1, t_2) &\mapsto \left( \frac{\gamma(t_1) + \gamma(t_2)}{2}, \frac{\gamma(t_1) - \gamma(t_2)}{2} \right), \\ S^1 \times S^1 &\rightarrow L_\phi \subset \mathbb{C}^2 \\ (t_3, t_4) &\mapsto \left( \frac{\gamma(t_3) + \gamma(t_4)}{2}, \frac{\gamma(t_3) - \gamma(t_4)}{2} e^{i\phi} \right). \end{aligned}$$

An intersection point outside of  $\gamma \times \{0\}$  is determined by the following condition for  $t_1 \neq t_2 \neq t_3 \neq t_4$  in  $S^1$ :

$$(2.1) \quad \gamma(t_1) + \gamma(t_2) = \gamma(t_3) + \gamma(t_4),$$


$$(2.2) \quad \gamma(t_1) - \gamma(t_2) = (\gamma(t_3) - \gamma(t_4))e^{i\phi}.$$

The ordered quadruple  $(t, t', t'', t''')$  solving above equations gives rise to the intersection point

$$p = (z_1, z_2 e^{i\phi}) = \left( \frac{\gamma(t'') + \gamma(t''')}{2}, \frac{\gamma(t'') - \gamma(t''')}{2} e^{i\phi} \right) \in L \cap L_\phi$$


and parameterizes the four points on the curve  $\gamma$  which completely determine a geometrically distinct inscribed rectangle as their vertices  $(A, B, C, D)$  in the counterclockwise order:

$$(z_1 + z_2, z_1 + z_2 e^{i\phi}, z_1 - z_2, z_1 - z_2 e^{i\phi}).$$



Suppose we have two intersection points with respect to a fixed aspect angle  $\phi$ :  $(z_1, z_2 e^{i\phi})$  and  $(w_1, w_2 e^{i\phi})$ , and they constitute the same geometric distinct inscribed rectangle on  $\gamma$ . There are following three cases:


- (1)  $w_1 + w_2 = z_1 + z_2$ : By the counterclockwise ordering of vertices, it yields  $w_1 = z_1, w_2 = z_2$ , i.e. these two intersection points are the same one. This means the intersection point  $(w_1, w_2 e^{i\phi})$  is the image of identity map of  $(z_1, z_2 e^{i\phi})$ .



- (2)  $w_1 + w_2 = z_1 - z_2$ : By the counterclockwise ordering of vertices, it yields  $w_1 = z_1, w_2 = -z_2$ , This means the intersection point  $(w_1, w_2 e^{i\phi})$  is the image of

$$\varrho : (z_1, z_2) \mapsto (z_1, -z_2)$$

of the intersection point  $(z_1, z_2 e^{i\phi}) \in L \cap L_\phi$  in the reflected Lagrangian tori  $\varrho(L) \cap \varrho(L_\phi)$ , and since  $L$  and  $L_\phi$  is invariant under  $\varrho$ , so it is another different intersection point in  $L \cap L_\phi$ , and it gives literally the same inscribed rectangle.



- (3)  $w_1 + w_2 = z_1 \pm z_2 e^{i\phi}$ : By the counterclockwise ordering of vertices, this is only possible when  $\phi = \frac{\pi}{2}$ , otherwise

$$(A, B, C, D) = (z_1 + z_2, z_1 + z_2 e^{i\phi}, z_1 - z_1, z_1 - z_2 e^{i\phi})$$

and

$$(A', B', C', D') = (w_1 + w_2, w_1 + w_2 e^{i\phi}, w_1 - w_1, w_1 - w_2 e^{i\phi})$$


would constitute two geometrically distinct rectangles in  $\mathbb{C}$ , in which  $(A, B, C, D)$  inscribes on  $\gamma$  whereas  $(A', B', C', D')$  does not. In the case of  $\phi = \frac{\pi}{2}$ , there are  $w_1 = z_1, w_2 = \pm z_2 e^{i\frac{\pi}{2}}$ .

- $(w_1, w_2) = (z_1, iz_2)$ : It means the intersection point  $(w_1, w_2 e^{i\phi})$  is the image of the map

$$\iota : (z_1, z_2) \mapsto (z_1, iz_2)$$

and  $\iota^2 = \rho$ .

$$C = z_1 - z_2 \quad z_1 + iz_2 = B \quad C' = z_1 - iz_2 \quad z_1 - z_2 = B'$$




$$D = z_1 - iz_2 \quad z_1 + z_2 = A \quad D' = z_1 + z_2 \quad z_1 + iz_2 = A'$$

- $(w_1, w_2) = (z_1, -iz_2)$ : It means the intersection point  $(w_1, w_2 e^{i\phi})$  is the image of the map

$$\rho \circ \iota : (z_1, z_2) \mapsto (z_1, -iz_2)$$

and  $\iota \circ \rho = \rho \circ \iota$ .

$$C = z_1 - z_2 \quad z_1 + iz_2 = B \quad C' = z_1 + iz_2 \quad z_1 + z_2 = B'$$



$$D = z_1 - iz_2 \quad z_1 + z_2 = A \quad D' = z_1 - z_2 \quad z_1 - iz_2 = A'$$

The above discussion in turns proves the following statement.

**Lemma 2.3.** *Let  $\text{Rect}_\phi(\gamma)$  be the set of inscribed rectangle with fixed aspect angle  $\phi$  on the curve  $\gamma$ .*

- (i) *When  $\phi \in (0, \frac{\pi}{2})$ , there is a bijection between sets*

$$(L \cap L_\phi \setminus (\gamma \times \{0\})) / C_2 \rightarrow \text{Rect}_\phi(\gamma)$$

*where  $C_2$  is presented by*

$$\langle \text{id}, \rho \mid \rho^2 = \text{id} \rangle \cong \mathbb{Z}/2\mathbb{Z}.$$

- (ii) *When  $\phi = \frac{\pi}{2}$ , there is a bijection between sets*

$$(L \cap L_\phi \setminus (\gamma \times \{0\})) / C_4 \rightarrow \text{Rect}_\phi(\gamma)$$

*where  $C_4$  is presented by*

$$\langle \text{id}, \rho, \iota \mid \rho^2 = \text{id}, \iota^2 = \rho \rangle \cong \mathbb{Z}/4\mathbb{Z}.$$

**Remark 2.4.** Recall the ordered tuple  $(t, t', t'', t''')$  determining a geometric distinct rectangle inscribed on the curve  $\gamma$  with vertices ordered as

$$(A, B, C, D) = (z_1 + z_2, z_1 + z_2e^{i\phi}, z_1 - z_1, z_1 - z_2e^{i\phi})$$

mentioned at the beginning of this section. The automorphism group of  $\{t, t', t'', t'''\}$  as a set is the symmetric group  $S_4$ . Some of the elements in  $S_4$  actually give rise to the reparameterizations of the vertices of the fixed geometrically distinct inscribed rectangle on  $\gamma$ , and further correspond to the symmetries of the intersection points in  $L \cap L_\phi \setminus (\gamma \times \{0\})$ , i.e. the elements in  $C_2$  and  $C_4$  above. We describe these symmetries in the following:

- (1)  $\begin{pmatrix} 1 & 2 & 3 & 4 \\ 1 & 2 & 3 & 4 \end{pmatrix} : (t, t', t'', t''') \mapsto (t, t', t'', t''')$  obviously corresponds to the identity map.
- (2)  $\begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 1 & 4 & 3 \end{pmatrix} : (t, t', t'', t''') \mapsto (t', t, t''', t'')$ . This corresponds to the symplectic involution  $\varrho$  acting on the intersection point  $p \in L \cap L_\phi$ , and the image  $q = \varrho(p)$  is another intersection point since  $(t', t, t''', t'')$  also solves the equations (2.1), (2.2) as  $(t, t', t'', t''')$  does. The point  $q \in L \cap L_\phi$  determines four point in  $\mathbb{C}$  in counterclockwise order

$$(A', B', C', D') = (z_1 - z_2, z_1 - z_2e^{i\phi}, z_1 + z_2, z_1 + z_2e^{i\phi}).$$

This form the exact same rectangle as the original one  $(A, B, C, D)$  inscribed on the curve  $\gamma$  with reparameterized vertices.

- (3)  $\bullet \begin{pmatrix} 1 & 2 & 3 & 4 \\ 3 & 4 & 1 & 2 \end{pmatrix} : (t, t', t'', t''') \mapsto (t'', t''', t, t')$  corresponds to  $\iota$  acting on  $p \in L \cap L_\phi$  when  $\phi = \frac{\pi}{2}$ . This can be seen as follows. Since  $t \neq t' \neq t'' \neq t'''$ , the pairs  $(t'', t''')$  and  $(t, t')$  give rise to two points  $p' \in L$  and  $p'' \in L_\phi$  respectively. In particular,  $p' = R_{-\phi}(p)$ ,  $p'' = R_\phi(p)$ , and  $p' \neq p''$ . When  $\phi \in (0, \frac{\pi}{2})$ , this symmetric group element maps the original inscribed rectangle to a new rectangle in the plane which is no longer inscribed on the curve  $\gamma$ . When  $\phi = \frac{\pi}{2}$ , i.e. the rectangle is actually a square, we obtain two additional intersection points  $p'' = R_{\frac{\pi}{2}}(p) \in R_{\frac{\pi}{2}}(L \cap L_{\frac{\pi}{2}}) = L_{\frac{\pi}{2}} \cap R_\pi(L) = L_{\frac{\pi}{2}} \cap L$  and  $p' = R_{-\frac{\pi}{2}}(p) \in R_{-\frac{\pi}{2}}(L \cap L_{\frac{\pi}{2}}) = R_\pi(L_{\frac{\pi}{2}}) \cap L = L_{\frac{\pi}{2}} \cap L$  so that  $p'' = R_\pi(p') = \varrho(p')$ , they commonly determine four points in  $\mathbb{C}$  in counterclockwise order:

$$(A', B', C', D') = (z_1 + iz_2, z_1 - z_2, z_1 - iz_2, z_1 + z_2).$$

This form the exact same square as the original one  $(A, B, C, D)$  inscribed on the curve  $\gamma$  with reparameterized vertices.

- $\bullet \begin{pmatrix} 1 & 2 & 3 & 4 \\ 4 & 3 & 2 & 1 \end{pmatrix} : (t, t', t'', t''') \mapsto (t''', t'', t, t')$  corresponds to  $\varrho \circ \iota$  acting on  $p \in L \cap L_\phi$  when  $\phi = \frac{\pi}{2}$ , and note that

$$\begin{aligned} \begin{pmatrix} 1 & 2 & 3 & 4 \\ 4 & 3 & 2 & 1 \end{pmatrix} &= \begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 1 & 4 & 3 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 \\ 3 & 4 & 1 & 2 \end{pmatrix}, \\ \begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 1 & 4 & 3 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 \\ 3 & 4 & 1 & 2 \end{pmatrix} &= \begin{pmatrix} 1 & 2 & 3 & 4 \\ 3 & 4 & 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 1 & 4 & 3 \end{pmatrix}, \end{aligned}$$

such relation amounts to  $\varrho \circ \iota = \iota \circ \varrho$ . This can be seen as follows. Since  $t \neq t' \neq t'' \neq t'''$ , the pairs  $(t''', t'')$  and  $(t', t)$  give rise to two points  $q' \in L$

and  $q'' \in L_\phi$  respectively. In particular,  $q' = R_{-\pi-\phi}(p)$ ,  $q'' = R_{\pi+\phi}(p)$ , and  $q' \neq q''$ . When  $\phi \in (0, \frac{\pi}{2})$ , this symmetric group element maps the original inscribed rectangle to a new rectangle in the plane which is no longer inscribed on the curve  $\gamma$ . When  $\phi = \frac{\pi}{2}$ , i.e. the rectangle is actually a square, the point  $q'' = R_{\pi+\frac{\pi}{2}}(p) = R_\pi(p'') = p' \in L_{\frac{\pi}{2}} \cap L$  determines four points in  $\mathbb{C}$  in counterclockwise order:

$$(A', B', C', D') = (z_1 - iz_2, z_1 + z_2, z_1 + iz_2, z_1 - z_2).$$

This form the exact same square as the original one  $(A, B, C, D)$  inscribed on the curve  $\gamma$  with reparameterized vertices.

A caveat is that the above set-theoretic bijection of the symmetric group elements in  $S_4$ , which form a Klein 4 group  $K_4$ , to the group elements in  $C_4$  does not constitute a group homomorphism, since  $K_4$  is not isomorphic to  $C_4$ .

It is worth noting that the symmetry of rectangular pegs is also addressed in [Mat22, §4.1].

### 3. GEOMETRIC TRANSVERSALITY

In general, we have no knowledge of the intersection set of  $L$  and  $L_\phi$  outside of  $\gamma \times \{0\}$ . It can be badly degenerate as the example below shows.

**Example 3.1.** In the most special case of  $\gamma$  being a standard circle, the tori  $L$  and  $L_\phi$  have badly degenerate (not clean) intersection in  $\mathbb{C}^2$  outside  $\mathcal{O}p(\gamma \times \{0\})$ . This can be seen explicitly via the polar coordinate of the torus:

$$\begin{aligned} \left( \frac{re^{i\theta_1} + re^{i\theta_2}}{2}, \frac{re^{i\theta_1} - re^{i\theta_2}}{2} \right) &\in L, \\ \left( \frac{re^{i\theta_3} + re^{i\theta_4}}{2}, \frac{re^{i\theta_3} - re^{i\theta_4}}{2} e^{i\phi} \right) &\in L_\phi. \end{aligned}$$

where  $\theta_1, \theta_2, \theta_3, \theta_4 \in [0, 2\pi]$ , and  $r$  is the radius of the standard circle  $\gamma$ .

We show in the following that  $L$  and  $L_\phi$  intersect transversely outside of  $\gamma \times \{0\}$  under certain generic condition. It is the key result of this paper.

**Theorem 3.2.** *For any fixed  $\phi \in (0, \frac{\pi}{2}]$ , there is a residual set of smooth Jordan curves such that for every  $\gamma$  in the set, the corresponding smooth Lagrangian tori  $L$  and  $L_\phi$  constructed in above way has only finitely many transversal intersection points outside a small open neighborhood of  $\gamma \times \{0\}$ .*

*Proof.* Let  $\text{Emb}^k(S^1, \mathbb{C})$  be the space of  $C^k$ -smooth embedded closed curves in  $\mathbb{C}$  for  $k \in \mathbb{N}$ . There are open inclusions  $\text{Emb}^k(S^1, \mathbb{C}) \subset \text{Imm}^k(S^1, \mathbb{C}) \subset C^k(S^1, \mathbb{C})$ . We consider the  $C^k$ -smooth map between  $C^\infty$ -smooth Banach manifolds

$$\begin{aligned} \mathcal{F} : \text{Emb}^k(S^1, \mathbb{C}) \times ((S^1)^4 \setminus \Delta_{(1,2,3,4)}^{(S^1)^4}) &\rightarrow \mathbb{C}^2 \times \mathbb{C}^2 \\ (\gamma, t_1, t_2, t_3, t_4) &\mapsto (z_L(t_1, t_2), w_L(t_1, t_2), z_{L_\phi}(t_3, t_4), w_{L_\phi}(t_3, t_4)) \end{aligned}$$



where

$$\begin{aligned} z_L(t_1, t_2) &= \frac{\gamma(t_1) + \gamma(t_2)}{2}, \\ w_L(t_1, t_2) &= \frac{\gamma(t_1) - \gamma(t_2)}{2}, \\ z_{L_\phi}(t_3, t_4) &= \frac{\gamma(t_3) + \gamma(t_4)}{2}, \\ w_{L_\phi}(t_3, t_4) &= \frac{\gamma(t_3) - \gamma(t_4)}{2} e^{i\phi}, \end{aligned}$$

and

$$\Delta_{(1,2,3,4)}^{(S^1)^4} = \{(t_1, t_2, t_3, t_4) \in (S^1)^4 : t_1 = t_2 = t_3 = t_4\}$$

is the small diagonal of the 4-torus  $(S^1)^4$ . Consider the diagonal in  $\mathbb{C}^2 \times \mathbb{C}^2$ :

$$\Delta_{\mathbb{C}^2} = \{(z_1, z_2, z_3, z_4) \in \mathbb{C}^2 \times \mathbb{C}^2 : (z_1, z_2) = (z_3, z_4)\}$$

We want to show that  $\mathcal{F}^{-1}(\Delta_{\mathbb{C}^2})$  is a manifold, and then further project to  $C^k(S^1, \mathbb{C})$  in order to obtain the genericity result.

The linearization of map  $\mathcal{F}$  at any point  $(\gamma, \mathbf{t}) = (\gamma, t_1, t_2, t_3, t_4)$  away from the small diagonal  $\Delta_{(1,2,3,4)}^{(S^1)^4}$ :

$$\begin{aligned} D_{(\gamma, \mathbf{t})} \mathcal{F} : T_\gamma C^k(S^1, \mathbb{C}) \times T_{\mathbf{t}}(S^1)^4 &\rightarrow T_{\mathcal{F}(\gamma, \mathbf{t})} \mathbb{C}^4 \\ (\xi, v_1, v_2, v_3, v_4) &\mapsto (V_1, V_2, V_3, V_4) \end{aligned}$$

is given by

$$\begin{aligned} V_1 &= \frac{1}{2}(D_{t_1} \gamma(v_1) + D_{t_2} \gamma(v_2) + \xi_{\gamma(t_1)} + \xi_{\gamma(t_2)}), \\ V_2 &= \frac{1}{2}(D_{t_1} \gamma(v_1) - D_{t_2} \gamma(v_2) + \xi_{\gamma(t_1)} - \xi_{\gamma(t_2)}), \\ V_3 &= \frac{1}{2}(D_{t_3} \gamma(v_3) + D_{t_4} \gamma(v_4) + \xi_{\gamma(t_3)} + \xi_{\gamma(t_4)}), \\ V_4 &= \frac{1}{2}(D_{t_3} \gamma(v_3) - D_{t_4} \gamma(v_4) + \xi_{\gamma(t_3)} - \xi_{\gamma(t_4)}), \end{aligned}$$

in which we regard  $\xi \in T_\gamma C^k(S^1, \mathbb{C})$  as a section in  $C^k(S^1, \gamma^* T\mathbb{C})$  via identification  $C^k(S^1, \mathbb{C}) \cong C^k(S^1, \gamma^* T\mathbb{C})$ , thus  $\xi_{\gamma(t)} \in T_{\gamma(t)} \mathbb{C}$  for all  $t \in S^1$ . We observe that at any point  $(\gamma, \mathbf{t}) \in \mathcal{F}^{-1}(\Delta_{\mathbb{C}^2})$ , the composition of continuous linear maps

$$T_\gamma C^k(S^1, \mathbb{C}) \times T_{\mathbf{t}}(S^1)^4 \xrightarrow{D_{(\gamma, \mathbf{t})} \mathcal{F}} T_{\mathcal{F}(\gamma, \mathbf{t})} \mathbb{C}^4 \twoheadrightarrow T_{\mathcal{F}(\gamma, \mathbf{t})} \mathbb{C}^4 / T_{\mathcal{F}(\gamma, \mathbf{t})} \Delta_{\mathbb{C}^2}$$

between Banach spaces is clearly surjective, and the kernel is closed since it is the preimage of the closed subspace  $T_{\mathcal{F}(\gamma, \mathbf{t})} \Delta_{\mathbb{C}^2} \subset T_{\mathcal{F}(\gamma, \mathbf{t})} \mathbb{C}^4$ , thus it has complementary subspace in  $T_\gamma C^k(S^1, \mathbb{C}) \times T_{\mathbf{t}}(S^1)^4$ . This implies that  $\mathcal{F} \pitchfork \Delta_{\mathbb{C}^2}$  in  $\mathbb{C}^4$ . By implicit function theorem on Banach manifolds (c.f. [MS12, Theorem A.3.3]), we conclude that the real codimensional 4 submanifold  $\Delta_{\mathbb{C}^2} \subset \mathbb{C}^2 \times \mathbb{C}^2$  has preimage  $\mathcal{F}^{-1}(\Delta_{\mathbb{C}^2})$  as a  $C^k$ -smooth Banach submanifold in  $\text{Emb}^k(S^1 \times \mathbb{C}) \times ((S^1)^4 \setminus \Delta_{(1,2,3,4)}^{(S^1)^4})$  of codimensional 4.

We next consider the projection map

$$\pi : \text{Emb}^k(S^1, \mathbb{C}) \times ((S^1)^4 \setminus \Delta_{(1,2,3,4)}^{(S^1)^4}) \rightarrow \text{Emb}^k(S^1, \mathbb{C})$$

restricting to Banach submanifold  $\mathcal{F}^{-1}(\Delta_{\mathbb{C}^2})$  which is  $C^k$ -smooth:

$$\begin{aligned} \pi|_{\mathcal{F}^{-1}(\Delta_{\mathbb{C}^2})} : \mathcal{F}^{-1}(\Delta_{\mathbb{C}^2}) &\rightarrow \text{Emb}^k(S^1, \mathbb{C}) \\ (\gamma, \mathbf{t}) &\mapsto \gamma \end{aligned}$$

The linearized map reads

$$\begin{aligned} D_{(\gamma, \mathbf{t})}\pi : T_{(\gamma, \mathbf{t})}\mathcal{F}^{-1}(\Delta_{\mathbb{C}^2}) &\rightarrow T_{\gamma}C^k(S^1, \mathbb{C}) \\ (\xi, \mathbf{v}) &\mapsto \xi \end{aligned}$$

where

$$\begin{aligned} T_{(\gamma, \mathbf{t})}\mathcal{F}^{-1}(\Delta_{\mathbb{C}^2}) &= \{(\xi, \mathbf{v}) \in T_f^k(S^1, \mathbb{C}) \times T_{\mathbf{t}}(S^1)^4 : \mathbf{V} = D_{(\gamma, \mathbf{t})}\mathcal{F}(\xi, \mathbf{v}) \in T_{\mathcal{F}(\gamma, \mathbf{t})}\Delta_{\mathbb{C}^2}\} \\ &= \{(\xi, \mathbf{v}) \in T_f^k(S^1, \mathbb{C}) \times T_{\mathbf{t}}(S^1)^4 : (V_1, V_2) = (V_3, V_4), (\gamma, \mathbf{t}) \in \mathcal{F}^{-1}(\Delta_{\mathbb{C}^2})\}. \end{aligned}$$

A curve  $\gamma$  being regular value of  $\pi|_{\mathcal{F}^{-1}(\Delta_{\mathbb{C}^2})}$  is equivalent to the transversality condition  $\mathcal{F}_{\gamma} \pitchfork \Delta_{\mathbb{C}^2}$  where  $\mathcal{F}_{\gamma} := \mathcal{F}(\gamma, \cdot) : (S^1)^4 \setminus \Delta_{(1,2,3,4)}^{(S^1)^4} \rightarrow \mathbb{C}^4$ , i.e.

$$D_{(\gamma, \mathbf{t})}\mathcal{F}(\{0\} \times T_{\mathbf{t}}(S^1)^4) + T_{\mathcal{F}(\gamma, \mathbf{t})}\Delta_{\mathbb{C}^2} = T_{\mathcal{F}(\gamma, \mathbf{t})}\mathbb{C}^4.$$

This corresponds to the geometric condition in the statement:  $\text{im}(\mathcal{F}_{\gamma}) \pitchfork \Delta_{\mathbb{C}^2}$  means  $L \times L_{\phi} \pitchfork \Delta_{\mathbb{C}^2}$  in  $\mathbb{C}^2 \times \mathbb{C}^2$  outside  $(\gamma \times \{0\}) \times (\gamma \times \{0\})$ , which is equivalent to  $L \pitchfork L_{\phi}$  in  $\mathbb{C}^2$  outside  $\gamma \times \{0\}$ , and these transversal intersection points are isolated and hence finite due to compactness of the domain.

The linear map  $D_{(\gamma, \mathbf{t})}\pi$  at every  $(\gamma, \mathbf{t}) \in \mathcal{F}^{-1}(\Delta_{\mathbb{C}^2})$  is a Fredholm operator: The kernel

$$\begin{aligned} \ker(D_{(\gamma, \mathbf{t})}\pi) &= \left\{ (\xi, \mathbf{v}) \in T_f C^k(S^1, \mathbb{C}) \times T_{\mathbf{t}}(S^1)^4 \left| \begin{array}{l} D_{t_1} \gamma(v_1) = D_{t_3} \gamma(v_3) \\ D_{t_2} \gamma(v_2) = D_{t_4} \gamma(v_4) \end{array} \right. \right\} \\ &= T_{(\gamma, \mathbf{t})}\mathcal{F}^{-1}(\Delta_{\mathbb{C}^2}) \cap (\{0\} \times T_{\mathbf{t}}(S^1)^4) \\ &\cong D_{(\gamma, \mathbf{t})}\mathcal{F}(\{0\} \times T_{\mathbf{t}}(S^1)^4) \cap T_{\mathcal{F}(\gamma, \mathbf{t})}\Delta_{\mathbb{C}^2} \end{aligned}$$

is finite dimensional. The cokernel is also finite dimensional by the following isomorphism

$$\text{coker}(D_{(\gamma, \mathbf{t})}\pi) = \frac{T_f C^k(S^1, \mathbb{C})}{\text{im}(D_{(\gamma, \mathbf{t})}\pi)} \cong \frac{T_{\mathcal{F}(\gamma, \mathbf{t})}\mathbb{C}^4}{D_{(\gamma, \mathbf{t})}\mathcal{F}(\{0\} \times T_{\mathbf{t}}(S^1)^4) + T_{\mathcal{F}(\gamma, \mathbf{t})}\Delta_{\mathbb{C}^2}}$$

due to the first isomorphism theorem:

$$\begin{array}{ccc} T_{\gamma}C^k(S^1, \mathbb{C}) \times T_{\mathbf{t}}(S^1)^4 & \xrightarrow{D_{(\gamma, \mathbf{t})}\mathcal{F}} & T_{\mathcal{F}(\gamma, \mathbf{t})}\mathbb{C}^4 & \xrightarrow{q_1} & \frac{T_{\mathcal{F}(\gamma, \mathbf{t})}\mathbb{C}^4}{D_{(\gamma, \mathbf{t})}\mathcal{F}(\{0\} \times T_{\mathbf{t}}(S^1)^4) + T_{\mathcal{F}(\gamma, \mathbf{t})}\Delta_{\mathbb{C}^2}} \\ \downarrow D_{(\gamma, \mathbf{t})}\pi & & & \searrow & \uparrow \sim \\ T_{\gamma}C^k(S^1, \mathbb{C}) & & & & \\ \downarrow q_2 & & & & \\ \frac{T_{\gamma}C^k(S^1, \mathbb{C})}{\text{im}(D_{(\gamma, \mathbf{t})}\pi)} & \xleftarrow{\sim} & \frac{T_{\gamma}C^k(S^1, \mathbb{C}) \times T_{\mathbf{t}}(S^1)^4}{T_{(\gamma, \mathbf{t})}\mathcal{F}^{-1}(\Delta_{\mathbb{C}^2}) + \{0\} \times T_{\mathbf{t}}(S^1)^4} & & \end{array}$$

since

$$\ker(q_1 \circ D_{(\gamma, \mathbf{t})}\mathcal{F}) = T_{(\gamma, \mathbf{t})}\mathcal{F}^{-1}(\Delta_{\mathbb{C}^2}) + \{0\} \times T_{\mathbf{t}}(S^1)^4 = \ker(q_2 \circ D_{(\gamma, \mathbf{t})}\pi).$$

Consequently, the set  $\text{im}(D_{(\gamma, \mathbf{t})}\pi)$  is closed in  $T_\gamma C^k(S^1, \mathbb{C})$ . Thus, the linear map  $D_{(\gamma, \mathbf{t})}\pi$  is a Fredholm operator with index zero:

$$\begin{aligned}
 \text{ind}(D_{(\gamma, \mathbf{t})}\pi) &= \dim \ker(D_{(\gamma, \mathbf{t})}\pi) - \dim \text{coker}(D_{(\gamma, \mathbf{t})}\pi) \\
 &= \dim(D_{(\gamma, \mathbf{t})}\mathcal{F}(\{0\} \times T_{\mathbf{t}}(S^1)^4) \cap T_{\mathcal{F}(\gamma, \mathbf{t})}\Delta_{\mathbb{C}^2}) - \dim T_{\mathcal{F}(\gamma, \mathbf{t})}\mathbb{C}^4 \\
 &\quad + \dim(D_{(\gamma, \mathbf{t})}\mathcal{F}(\{0\} \times T_{\mathbf{t}}(S^1)^4) + T_{\mathcal{F}(\gamma, \mathbf{t})}\Delta_{\mathbb{C}^2}) \\
 &= \dim(D_{(\gamma, \mathbf{t})}\mathcal{F}(\{0\} \times T_{\mathbf{t}}(S^1)^4) \cap T_{\mathcal{F}(\gamma, \mathbf{t})}\Delta_{\mathbb{C}^2}) - \dim T_{\mathcal{F}(\gamma, \mathbf{t})}\mathbb{C}^4 \\
 &\quad + \dim D_{(\gamma, \mathbf{t})}\mathcal{F}(\{0\} \times T_{\mathbf{t}}(S^1)^4) + \dim T_{\mathcal{F}(\gamma, \mathbf{t})}\Delta_{\mathbb{C}^2} \\
 &\quad - \dim(D_{(\gamma, \mathbf{t})}\mathcal{F}(\{0\} \times T_{\mathbf{t}}(S^1)^4) \cap T_{\mathcal{F}(\gamma, \mathbf{t})}\Delta_{\mathbb{C}^2}) \\
 &= \dim D_{(\gamma, \mathbf{t})}\mathcal{F}(\{0\} \times T_{\mathbf{t}}(S^1)^4) + \dim T_{\mathcal{F}(\gamma, \mathbf{t})}\Delta_{\mathbb{C}^2} - \dim T_{\mathcal{F}(\gamma, \mathbf{t})}\mathbb{C}^4 \\
 &= 4 + 4 - 8 \\
 &= 0
 \end{aligned}$$

In conclusion,  $\pi|_{\mathcal{F}^{-1}(\Delta_{\mathbb{C}^2})}$  is a Fredholm map of index 0. By Sard-Smale theorem (c.f. [MS12, Theorem A.5.1]), the set of regular values of  $\pi|_{\mathcal{F}^{-1}(\Delta_{\mathbb{C}^2})}$  is residual in  $\text{Emb}^k(S^1, \mathbb{C})$ . Thus, we have proved that the set

$$\text{Emb}_{\text{reg}}^k(S^1, \mathbb{C}) := \left\{ \text{regular values of } \pi|_{\mathcal{F}^{-1}(\Delta_{\mathbb{C}^2})} \right\}$$

is generic in  $\text{Emb}^k(S^1, \mathbb{C})$  with respect to  $C^k$ -topology.

Now for passing from  $C^k$ -genericity to  $C^\infty$ -genericity, we employ the argument in [St24, Lemma 6.2] adapting to our setting here. Note that such type of argument is also known to Clifford Taubes earlier, see e.g. [MS12, p.54-55] and [Fra03, Appendix C.2, p.106]. Since  $\text{Emb}_{\text{reg}}^k(S^1, \mathbb{C}) \subseteq \text{Emb}^k(S^1, \mathbb{C})$  is a residual subset with respect to  $C^k$ -topology for each fixed  $k \geq 1$ , and a residual set can be expressed as a countable intersection of open dense sets, c.f. [BD18, Lemma 1.6.2 (vi)], we can therefore write

$$\text{Emb}_{\text{reg}}^k(S^1, \mathbb{C}) = \bigcap_{\ell \in \mathbb{N}} \mathcal{V}_\ell^k$$

where each  $\mathcal{V}_\ell^k$  is open and dense in  $\text{Emb}^k(S^1, \mathbb{C})$  with the  $C^k$ -topology. For each integer  $j \geq k$ , we put

$$\mathcal{V}_\ell^j := \mathcal{V}_\ell^k \cap \text{Emb}^j(S^1, \mathbb{C}),$$

and we have

$$\begin{aligned}
 \bigcap_{\ell \in \mathbb{N}} \mathcal{V}_\ell^j &= \left( \bigcap_{\ell \in \mathbb{N}} \mathcal{V}_\ell^k \right) \cap \text{Emb}^j(S^1, \mathbb{C}) \\
 &= \text{Emb}_{\text{reg}}^k(S^1, \mathbb{C}) \cap \text{Emb}^j(S^1, \mathbb{C}) \\
 &= \text{Emb}_{\text{reg}}^j(S^1, \mathbb{C}).
 \end{aligned}$$

Due to the fact that  $\text{Emb}_{\text{reg}}^j(S^1, \mathbb{C})$  is residual in  $\text{Emb}^j(S^1, \mathbb{C})$ , by the Baire category theorem (c.f. [BD18, Theorem 1.6.4]), it is dense and thus each  $\mathcal{V}_\ell^j$  is dense and open in  $\text{Emb}^j(S^1, \mathbb{C})$ , since the  $C^j$ -topology is finer than the  $C^k$ -topology. We

further put

$$\begin{aligned}\mathcal{V}_\ell^\infty &:= \mathcal{V}_\ell^k \cap \text{Emb}^\infty(S^1, \mathbb{C}) \\ &= \mathcal{V}_\ell^k \cap \left( \bigcap_{j \geq k} \text{Emb}^j(S^1, \mathbb{C}) \right) \\ &= \bigcap_{j \geq k} \mathcal{V}_\ell^j.\end{aligned}$$

We claim that  $\mathcal{V}_\ell^\infty$  is dense in  $\text{Emb}^\infty(S^1, \mathbb{C})$ : Pick an arbitrary  $\gamma_0 \in \text{Emb}^\infty(S^1, \mathbb{C})$ , and an open neighbourhood  $\mathcal{U}_0$  of it in  $\text{Emb}^\infty(S^1, \mathbb{C})$ . Let  $j \in \mathbb{N}_{\geq k}$  and  $\delta > 0$  such that

$$\{\gamma \in \text{Emb}^\infty(S^1, \mathbb{C}) : d_{C^j}(\gamma, \gamma_0) < \delta\} \subseteq \mathcal{U}_0$$

where  $d_{C^j}$  is the metric inducing  $C^j$ -topology on  $\text{Emb}^j(S^1, \mathbb{C})$ . Due to the density of  $\mathcal{V}_\ell^j$  in  $\text{Emb}^j(S^1, \mathbb{C})$ ,

$$\exists \gamma_1 \in \mathcal{V}_\ell^j : d_{C^j}(\gamma_1, \gamma_0) < \frac{\delta}{2}.$$

We know that  $\text{Emb}^\infty(S^1, \mathbb{C})$  is dense in  $\text{Emb}^j(S^1, \mathbb{C})$  with respect to  $C^j$ -topology, and  $\mathcal{V}_\ell^j$  is open in  $\text{Emb}^j(S^1, \mathbb{C})$ , then

$$\exists \gamma_2 \in \text{Emb}^\infty(S^1, \mathbb{C}) \cap \mathcal{V}_\ell^j : d_{C^j}(\gamma_2, \gamma_1) < \frac{\delta}{2}.$$

By the triangle inequality of the metric  $d_{C^j}$ , we have

$$\gamma_2 \in \{\gamma \in \text{Emb}^\infty(S^1, \mathbb{C}) : d_{C^j}(\gamma, \gamma_0) < \delta\} \cap \mathcal{V}_\ell^\infty \subseteq \mathcal{U}_0 \cap \mathcal{V}_\ell^\infty.$$

This proves the claim that  $\mathcal{V}_\ell^\infty$  is dense in  $\text{Emb}^\infty(S^1, \mathbb{C})$ . It is also open in  $\text{Emb}^\infty(S^1, \mathbb{C})$  since the  $C^\infty$ -topology is finer than the  $C^k$ -topology. Moreover,

$$\begin{aligned}\bigcap_{\ell \in \mathbb{N}} \mathcal{V}_\ell^\infty &= \left( \bigcap_{\ell \in \mathbb{N}} \mathcal{V}_\ell^k \right) \cap \text{Emb}^\infty(S^1, \mathbb{C}) \\ &= \text{Emb}_{\text{reg}}^k(S^1, \mathbb{C}) \cap \left( \bigcap_{j \geq k} \text{Emb}^j(S^1, \mathbb{C}) \right) \\ &= \bigcap_{j \geq k} \text{Emb}_{\text{reg}}^j(S^1, \mathbb{C}) \\ &= \text{Emb}_{\text{reg}}^\infty(S^1, \mathbb{C}).\end{aligned}$$

All together implies that  $\text{Emb}_{\text{reg}}^\infty(S^1, \mathbb{C})$  is residual in  $\text{Emb}^\infty(S^1, \mathbb{C})$  with respect to the  $C^\infty$ -topology. As  $\text{Emb}_{\text{reg}}^\infty(S^1, \mathbb{C})$  is precisely the set of smooth Jordan curves with desired property in the statement, this completes the proof.  $\square$

**Remark 3.3.** One should note that this statement does not imply the existence of an intersection point outside  $\gamma \times \{0\}$ . As the ‘empty intersection outside of  $\gamma \times \{0\}$ ’ also stays in the context of ‘transversal intersection outside of  $\gamma \times \{0\}$ ’. In other words, the curves resulting empty intersection outside of  $\gamma \times \{0\}$  also belong to the residual set in the statement.

**Example 3.4.** By Lemma 2.3 and Theorem 3.2, one can easily tell that all ellipses except the standard circle lie in such residual set in the statement of Theorem 3.2. And one could produce more such generic curves from ellipses via small perturbation.

#### 4. PROOF OF THE MAIN THEOREM 1.6

For convenience reason, here we abuse notations, and consider

$$l : (z_1, z_2) \mapsto (z_1 + z_2, z_1 - z_2)$$

which is a symplectomorphism on  $(\mathbb{C}^2, \omega_{\text{std}})$ , i.e.  $l^*\omega_{\text{std}} = \omega_{\text{std}}$ , where

$$\omega_{\text{std}} = d\lambda_{\text{std}}, \quad \lambda_{\text{std}} = -\frac{i}{4} \sum_{i=1,2} (z_i dz_i + \bar{z}_i d\bar{z}_i - z_i d\bar{z}_i - \bar{z}_i dz_i).$$

We still put  $L := l(\gamma \times \gamma)$  and  $L_\phi := R_\phi(L)$ . The important point is that all the previous discussions on  $L$  and  $L_\phi$  as well as their intersection points carry over verbatim. We observe that  $L$  and  $L_\phi$  are monotone Lagrangian tori in  $(\mathbb{C}^2, \omega_{\text{std}})$  with minimal Maslov number  $N_L = N_{L_\phi} = 2$ . The description of monotone Lagrangian submanifolds is referred to A.1. This can be seen as follows. We assume with out lost of generality that  $\gamma$  bounds the domain  $\bar{\gamma}$  with the same area of unit disk  $D^2$ , i.e.  $\text{Area}(\bar{\gamma}) = \text{Area}(D^2) = \pi$ . The torus  $\gamma \times \gamma$  is a monotone Lagrangian torus with minimal Maslov index 2 and monotonicity constant  $\tau = \pi$ . The verification of (A.1) follows analogously from the computation in [Oh15, §16.3, Exmaple 16.3.6 (1)] for the Clifford torus

$$T_{\text{Cliff}} := S^1(1) \times S^1(1) = \{(z_1, z_2) \in \mathbb{C}^2 : |z_1| = |z_2| = 1\},$$

since  $\bar{\gamma}$  is biholomorphic to  $D^2(1)$  by Riemann mapping theorem and the computation is topological. These properties are preserved by symplectomorphisms. In particular, according to [Kar13], there is an area preserving isotopy (i.e. symplectic isotopy) on  $\mathbb{C}$  that take  $\gamma$  to  $D^2$ , further due to  $H_c^1(\mathbb{R}^2; \mathbb{R}) \cong \{0\}$  and by [PR14], this can be turned into a Hamiltonian isotopy. Such result is also shown in [LM14, Lemma 2.3]. Therefore, the two Lagrangian tori  $L$  and  $L_\phi$  are actually Hamiltonian isotopic to the Clifford torus  $T_{\text{Cliff}}$ . The fact that  $L$  and  $L_\phi$  have minimal Maslov number  $N_L = N_{L_\phi} = 2$  can also be deduced from the results [Vit90, Pol91, Oh96] that an embedded Lagrangian torus in  $\mathbb{C}^2$  has minimal Maslov number 2, which can be regarded as a special case of Audin's conjecture, see e.g. [Dam12].

To situate ourselves in the framework of Lagrangian Floer homology, c.f. Appendix A, we now assume that the smooth Jordan curve  $\gamma$  lies in the generic class in Theorem 3.2. We regard  $L_\phi := R_\phi(L)$  as a Hamiltonian diffeomorphism acting on  $L$  generated by the autonomous Hamiltonian function

$$H_\phi = \frac{1}{2}\phi r_2^2 = \frac{1}{2}\phi |z_2|^2$$

where  $\phi \in (0, \frac{\pi}{2})$ , whose Hamiltonian vector field is  $\phi \partial_{\theta_2}$  in terms of polar coordinate  $z_2 = r_2 e^{i\theta_2}$  generating the time-1 map of Hamiltonian flow  $\varphi_{H_\phi} = R_\phi$  as the rotation around  $\mathbb{C} \times \{0\}$  by angle  $\phi$ , such that  $L$  and  $L_\phi = \varphi_{H_\phi}(L) = R_\phi(L)$  intersect cleanly along  $\gamma \times \{0\}$  and transversely outside of  $\gamma \times \{0\}$ . So we appeal for the Floer homology involving the Morse-Bott degeneracy, c.f. Appendix A.2, by choosing an arbitrary auxiliary perfect Morse functions  $f$  on the Morse-Bott circle  $\gamma \times \{0\}$  and zero functions on other transversal intersection points. According to Appendix A.1

and Appendix A.2, the generators of the Floer complex  $CF_*(L, L_\phi, f)$  are the two critical points  $x, y \in \text{Crit}(f)$  in  $\gamma \times \{0\}$  and other transversal intersection points in  $L \cap L_\phi \setminus (\gamma \times \{0\})$ . Due to Theorem 1.5 by Greene and Lobb, there is an intersection point  $p \in L \cap L_\phi \setminus (\gamma \times \{0\})$ , and by Lemma 2.3 there is another distinct intersection point  $\varrho(p) \in L \cap L_\phi \setminus (\gamma \times \{0\})$ .

We next determine a suitable absolute grading for the generator of the Floer complex via the Maslov-Morse index introduced in Appendix A.1 and Appendix A.2. Due to (A.9), the grading differs by multiple of integers up to capping. Let for the moment  $x$  be any point in  $L \cap L_\phi$  equipped with two cappings  $\bar{x}$  and  $\tilde{x}$ , then we have

$$\mu([x, \bar{x}]) - \mu([x, \tilde{x}]) = I_\mu([- \tilde{x} \# \bar{x}])$$

where  $[- \tilde{x} \# \bar{x}] \in \Gamma_{\ell_0}$  in which  $\ell_0 \in \mathcal{P}(L, L_\phi)$  is a based path, and  $\Gamma_{\ell_0}$  is defined by (A.3). Since  $\mathcal{P}(L, L_\phi)$  is clearly path connected, we can choose with out lost of generality that  $\ell_0 \subset \mathbb{C} \times \{0\}$ , i.e. it lies in  $\text{Fix}(\varrho)$ . We show that  $I_\mu([- \tilde{x} \# \bar{x}])$  takes value in  $2\mathbb{Z}$ : Let  $u : S^1 \times [0, 1] \rightarrow \mathbb{C}^2$  be a representative map of the class  $[- \tilde{x} \# \bar{x}] \in \pi_2(\mathbb{C}^2, L \cup L_\phi)$  satisfying  $u(\cdot, 0) \in L$ ,  $u(\cdot, 1) \in L_\phi$ ,  $u(0, \cdot) = x$ ,  $u(\frac{1}{2}, \cdot) = \ell_0$ . And  $v : D^2 \rightarrow \mathbb{C}^2$  is a disk map with  $v|_{\partial D^2} = u|_{S^1 \times \{1\}}$ . Then  $[v] \in \pi_2(\mathbb{C}^2, L_\phi)$  and  $[u \# v] \in \pi_2(\mathbb{C}^2, L)$ . By the gluing property of Maslov index,

$$\mu([u]) + \mu([v]) = \mu([u \# v]),$$

and due to the minimal Maslov number  $N_L = N_{L_\phi} = 2$ , we have

$$\mu([u]) = I_\mu([- \tilde{x} \# \bar{x}]) \in 2\mathbb{Z}.$$

By (A.14) and (A.15), we fix a common capping for any point in the clean intersection  $\gamma \times \{0\}$  and let  $x$  be of Morse index 1 while  $y$  has Morse index 0 so that

$$|x| = 1 \quad \text{and} \quad |y| = 0.$$

We assign similarly the absolute grading  $|p| := \mu(p, \bar{p})$  for  $p$  with respect to a fixed cap  $\bar{p}$  such that

$$|p| = 0 \quad \text{or} \quad 1.$$

It remains to determine the grading  $\varrho(p)$ . We put  $\overline{\varrho(p)} := \varrho(\bar{p})$ , and claim that

$$|\varrho(p)| := \mu(\varrho(p), \varrho(\bar{p})) = \mu(p, \bar{p}) =: |p|.$$

This can be shown as follows. By (A.7), we have

$$\mu(p, \varrho(p); B) = \mu([\varrho(p), \varrho(\bar{p})]) - \mu([p, \bar{p}])$$

where  $\varrho(\bar{p}) = \bar{p} \# B$  with  $B = [- \bar{p} \# \varrho(\bar{p})] \in \pi_2(p, \varrho(p))$ . Similarly,

$$\mu(\varrho(p), \varrho(\varrho(p)); \varrho_*(B)) = \mu(\varrho(p), p; -B) = \mu([p, \bar{p}]) - \mu([\varrho(p), \varrho(\bar{p})])$$

where  $\varrho_*(B) = [- \varrho(\bar{p}) \# \bar{p}] = -B \in \pi_2(\varrho(p), p)$ . Since  $\varrho$  is a symplectomorphism and  $L, L_\phi$  are both  $\varrho$ -invariant, by the invariance of the Maslov index of Lagrangian pair under symplectic action, c.f. [Oh15, Appendix C.1 (1)], we have

$$\mu(\varrho(p), \varrho(\varrho(p)); \varrho_*(B)) = \mu(p, \varrho(p); B).$$

This verifies the claim.

The complex space  $(\mathbb{C}^2, d\lambda_{\text{std}})$  is a Liouville manifold, see Example A.17 (1). By Appendix A.3, we can choose a generic admissible  $d\lambda_{\text{std}}$ -compatible almost complex structure  $J \in \mathcal{J}(\mathbb{C}^2, d\lambda_{\text{std}})$  so that the  $\mathbb{F}_2$ -coefficient Morse-Bott Floer homology  $HF_*(L, L_\phi, f)$  of monotone Lagrangian tori  $L$  and  $L_\phi$  with minimal Maslov index

2 is well-defined as a  $\mathbb{Z}/2\mathbb{Z}$ -graded module, whereas with respect to the Novikov ring  $\Lambda$ -coefficient,  $HF_*(L, L_\phi, f; \Lambda)$  is a bounded  $\mathbb{Z}$ -graded  $\Lambda$ -module concentrated in degree 0 and 1.

*Proof of Theorem 1.6.* Due to Lemma 2.3, we aim to find another  $\varrho$ -invariant pair of intersection points in  $L \cap L_\phi \setminus (\gamma \times \{0\})$  different from the  $\{p, \varrho(p)\}$ . Primarily, we observe that the Lagrangian torus  $L := l(\gamma \times \gamma)$  in  $\mathbb{C}^2$  can be evidently displaced by translation, e.g. translating in  $z_1$ -direction in  $\mathbb{C} \times \mathbb{C}$ :

$$(z_1, z_2) \mapsto (z_1, z_2) + \mathbf{z} = (z_1 + z, z_2)$$

which is a Hamiltonian diffeomorphism for the corresponding translation Hamiltonian function in  $(\mathbb{C}^2, \omega_{\text{std}})$ . The image of  $L$  under translation  $\mathbf{z} = (z, 0) \in \mathbb{C} \times \{0\}$  is denoted by  $L_{\mathbf{z}}$ , when  $|z|$  is very large, then  $L \cap L_{\mathbf{z}} = \emptyset$ . Therefore, the Lagrangian Floer homology vanishes:

$$HF_*(L, L_{\mathbf{z}}; \Lambda) \cong \{0\}.$$

By the Theorem A.16, we have isomorphisms:

$$(4.1) \quad HF_*(L, L_\phi, f; \Lambda) \cong HF_*(L, L_{\mathbf{z}}; \Lambda).$$


Suppose there is only one  $\varrho$ -invariant pair of intersection points  $\{p, \varrho(p)\}$  outside  $\gamma \times \{0\}$  in  $L \cap L_\phi$ . Then the Euler characteristic can be computed as

$$\begin{aligned} \chi(HF_*(L, L_\phi, f; \Lambda)) &:= \sum_{i=0,1} (-1)^i \text{rank}_\Lambda HF_i(L, L_\phi, f; \Lambda) \\ &= \sum_{i=0,1} (-1)^i \text{rank}_\Lambda CF_i(L, L_\phi, f; \Lambda) \\ &= \begin{cases} 2 & \text{if } |p| = |\varrho(p)| = 0 \\ -2 & \text{if } |p| = |\varrho(p)| = 1 \end{cases} \end{aligned}$$

and meanwhile

$$\begin{aligned} \chi(HF_*(L, L_{\mathbf{z}}; \Lambda)) &:= \sum_{i=0,1} (-1)^i \text{rank}_\Lambda HF_i(L, L_{\mathbf{z}}; \Lambda) \\ &= \sum_{i=0,1} (-1)^i \text{rank}_\Lambda CF_i(L, L_{\mathbf{z}}; \Lambda) \\ &= 0 \end{aligned}$$

which contradicts to (4.1). Thus, there must be also another pair  $\{q, \varrho(q)\}$  of intersection points outside  $\gamma \times \{0\}$  in  $L \cap L_\phi$ .



This finishes the proof.  $\square$

**Remark 4.2.** The Theorem 1.6 can also be proved in a purely differential topological way via algebraic intersection number.<sup>1</sup> We present the proof as follows: Recall the algebraic intersection number  $X \bullet Y$  of two closed oriented embedded submanifolds  $X, Y$  in an oriented smooth manifold  $M$  such that  $X \pitchfork Y$  is defined to be the signed count of the intersection points, see Appendix B.1. It is well-known that the sign of each intersection point does not depend on the choice of the positive bases, and the number  $X \bullet Y$  is well-defined as a homotopy invariant. Now take  $X = L$  and  $Y = L_\phi = R_\phi(L)$  in  $M = \mathbb{C}^2$ . For a curve  $\gamma$  in the generic class,  $L$  intersects with  $L_\phi$  cleanly along  $\gamma \times \{0\}$  and transversally outside of that part. To put it in the totally transverse scenario, we perturb the autonomous Hamiltonian function generating the rotation  $R_\phi$  by a small Morse function:

$$H_\phi^\varepsilon = H_\phi + \varepsilon\beta f$$

where  $f$  is a perfect Morse function on  $\gamma \times \{0\}$ , and  $\beta : \mathbb{C}^2 \rightarrow \mathbb{R}$  is a bump function supported in a small tubular neighborhood  $\mathcal{U}$  of  $\gamma \times \{0\}$  containing no intersection points in  $L \cap L_\phi \setminus (\gamma \times \{0\})$ , and  $\varepsilon > 0$  is sufficiently small. The symplectic gradient vector field of  $H_\phi^\varepsilon$  is given by

$$-J_0 \text{grad}_{g_0} H_\phi - J_0 \text{grad}_{g_0} \varepsilon\beta f$$

from which we can tell that the zeros of the vector field in  $\mathcal{U}$  is exactly the two critical points of  $f$  on  $\gamma \times \{0\}$ . The time-1-flow yields a diffeomorphism  $\varphi_{H_\phi^\varepsilon}$  on  $L$  whose image  $L_\phi^\varepsilon := \varphi_{H_\phi^\varepsilon}(L)$  coincides with  $L_\phi$  outside of a small tubular neighborhood  $\mathcal{U}$  of  $\gamma \times \{0\}$ , and  $L \cap L_\phi^\varepsilon$  has now only two transversal intersection points  $x_0, x_1$  in  $\mathcal{U}$  and  $L \cap L_\phi^\varepsilon \setminus \mathcal{U} = L \cap L_\phi \setminus (\gamma \times \{0\})$ , namely  $L \pitchfork L_\phi^\varepsilon$ . There is an obvious homotopy between  $L_\phi$  and  $L_\phi^\varepsilon$ , and due to the homotopy invariance of algebraic intersection number, we put

$$L \bullet L_\phi := L \bullet L_\phi^\varepsilon$$

which is clearly independent from the choices of  $f, \varepsilon, \beta$ , etc. Assume there is only one pair of intersection points  $(p, \varrho(p))$  in  $L \cap L_\phi \setminus (\gamma \times \{0\})$ , then according to the perturbation above, up to choice of orientation, we have

$$\begin{aligned} L \bullet L_\phi &:= L \bullet L_\phi^\varepsilon = (\text{sign}(x_0) + \text{sign}(x_1)) + (\text{sign}(p) + \text{sign}(\varrho(p))) \\ &= \pm\chi(\gamma \times \{0\}) \pm (1 + 1) \\ &= \pm\chi(S^1) \pm 2 \\ &= 0 \pm 2 \\ &= \pm 2. \end{aligned}$$

The first bracket sum is due to the following: First noticing the orientation signs of  $x_0, x_1 \in L \cap L_\phi^\varepsilon \subset \mathbb{C}^2$  coincide with the orientation signs of them as intersection point of sections  $(\gamma \times \{0\}) \cap J_0 \text{grad}_{g_0} \varepsilon\beta f$  in vector bundle  $J_0 T(\gamma \times \{0\})$ . This is, up to a uniform sign, equal to the orientation signs of  $x_0$  and  $x_1$  as intersection points of sections  $(\gamma \times \{0\}) \cap \text{grad}_{g_0} f$  in  $T(\gamma \times \{0\})$ , which in fact are further equal to the indices of  $x_0$  and  $x_1$  as zeros of gradient vector field  $\text{grad}_{g_0} f$  in  $T(\gamma \times \{0\})$ . Hence the sum of the orientation signs of  $x_0, x_1 \in L \cap L_\phi^\varepsilon$  coincides up to a uniform sign with the sum of indices of zeros  $x_0$  and  $x_1$  of  $\text{grad}_{g_0} f$  in  $T(\gamma \times \{0\})$ , which is the Euler characteristic by the Poincaré-Hopf theorem. More detail and general

<sup>1</sup>This was suggested by Kai Cieliebak to the author after his talk in the *Oberseminar Differentialgeometrie*, Wintersemester 2023-2024, at the Universität Augsburg



treatment of computing algebraic intersection number provided appearance of clean intersection is referred to Appendix B. The second bracket sum is due to the fact that  $\varrho$  is orientation preserving diffeomorphism, therefore  $p$  and  $\varrho(p)$  must have the same orientation signs. On the other hand, we know that  $L$  can be displaced away from itself by some translational smooth isotopy, hence it is apparent that

$$L \bullet L_\phi = \text{sign}(\varnothing) = 0.$$

This leads to a contradiction! Hence, the assumption must be wrong, and there must be another pair of intersection point  $(q, \varrho(q))$  in  $L \cap L_\phi \setminus (\gamma \times \{0\})$ . In fact, the Lagrangian Floer homology can be thought of as a ‘categorification’ of intersection number in the sense that

$$\chi(HF_*(L, L_\phi, f; \Lambda)) = L \bullet L_\phi$$

See Appendix B.4. In this spirit, these two proofs are well consistent.

### 5. COMMENTS ON A PRIORI EXISTENCE OF RECTANGULAR PEG

Our original goal is to prove the a priori existence (1.4) of inscribed rectangle with aspect angle  $\phi$  in any smooth Jordan curve  $\gamma$  using Floer theory circumventing the non-existence of embedded Lagrangian Klein bottle in  $\mathbb{C}^2$  used by Greene and Lobb in [GL21]. Suppose  $\gamma$  lies in the generic class such that  $L \pitchfork L_\phi$ . Initially, there might be an inclination to compute and anticipate a non-vanishing result

$$HF_*(L, L_\phi, f) \neq \{0\}$$

in the Morse-Bott setting, leading one to anticipate a potential contradiction to

$$HF_*(L, L_\mathbf{z}) = \{0\}$$

due to the invariance of Floer homology. A convenient choice of the  $\omega_{\text{std}}$ -compatible almost complex structure for computation is the  $\varrho$ -invariant ones, i.e.

$$\varrho^* J = \varrho_* \circ J \circ \varrho_* = J.$$

The set of such almost complex structures is denoted by  $\mathcal{J}^{\mathbb{Z}/2\mathbb{Z}}(\mathbb{C}^2, \omega_{\text{std}})$ , and it is non-empty and contractible. Then a holomorphic strip  $u$  which does not lie entirely in  $\mathbb{C} \times \{0\}$  will give rise to another one by applying the involution  $\varrho$ , i.e.  $u^\varrho := \varrho \circ u$  solves the non-linear Cauchy-Riemann equation:

$$\begin{aligned} \bar{\partial}_J u^\varrho &:= \frac{1}{2}(d(\varrho \circ u) + J \circ d(\varrho \circ u) \circ i) \\ &= \frac{1}{2}(\varrho_* \circ du + J \circ \varrho_* \circ du \circ i) \\ &= \frac{1}{2}(\varrho_* \circ du + \varrho_* \circ J \circ du \circ i) \\ &= \varrho_* \circ \bar{\partial}_J u \\ &= 0 \end{aligned}$$

Then one would get zero contribution from such pair of holomorphic strips in the boundary operator

$$\partial : CF_*(L, L_\phi, f) \rightarrow CF_{*-1}(L, L_\phi, f).$$

However, it seems highly convincing that the holomorphic strip  $u_0$  filling the curve  $\gamma \times \{0\}$  in  $\mathbb{C} \times \{0\}$  with asymptotics  $x, y \in \text{Crit}(f)$  is regular for a generic choice

of admissible  $\{J_t\}_{t \in [0,1]} \subset \mathcal{J}^{\mathbb{Z}/2\mathbb{Z}}(\mathbb{C}^2, \omega_{\text{std}})$ . E.g. one considers the standard almost complex structure

$$J_0 = \begin{pmatrix} i & 0 \\ 0 & i \end{pmatrix}$$

which is clearly  $\varrho$ -invariant  $\omega_{\text{std}}$ -compatible and admissible. One can show that  $u_0$  as a  $J_0$ -holomorphic strip as above has Fredholm index  $\text{ind}(D_{u_0} \bar{\partial}_{J_0}) = 1$  and  $D_{u_0} \bar{\partial}_{J_0}$  is surjective, i.e.  $u_0$  is regular  $J_0$ -holomorphic strip. If this is true, then there would be a non-eliminatable contribution to the boundary operator by a regular  $J_t$ -holomorphic strip  $u_0$  for a generic admissible  $\{J_t\}_{t \in [0,1]} \subset \mathcal{J}^{\mathbb{Z}/2\mathbb{Z}}(\mathbb{C}^2, \omega_{\text{std}})$  with  $\text{im}(u_0) \subset \mathbb{C} \times \{0\}$  and  $\partial \text{im}(u_0) = \gamma \times \{0\}$ . Assuming there is no intersection point outside of  $\gamma \times \{0\}$ , the Morse-Bott Floer complex regarded as a  $\mathbb{Z}/2\mathbb{Z}$ -graded chain complex over  $\mathbb{F}_2$  reads

$$\cdots \xrightarrow{y \mapsto x} \mathbb{F}_2 \langle x \rangle \xrightarrow{x \mapsto 2y} \mathbb{F}_2 \langle y \rangle \xrightarrow{y \mapsto x} \mathbb{F}_2 \langle x \rangle \xrightarrow{x \mapsto 2y} \mathbb{F}_2 \langle y \rangle \xrightarrow{y \mapsto x} \cdots$$

which is schematically depicted in the following picture:



and thus the Morse-Bott Floer homology turns out to be also vanishing:

$$HF_*(L, L_\phi, f, J_t) = \{0\}.$$

Hence, we may not obtain the non-vanishing result in such a way, and the strategy via contradiction argument sketched at the beginning would fail. Moreover, our goal extends beyond merely establishing a prior existence result (1.4) for those smooth Jordan curves in the generic class; we aspire to achieve it for general smooth Jordan curve. One potential avenue is to show a degenerate Arnold conjecture-like statement for the Lagrangian tori by studying the quantitative information of the Lagrangian torus, e.g. the displacement energy.

Very recently, we promptly noticed a new preprint [GL24] by Greene and Lobb, in which they have researched in a similar direction albeit in a different geometric

setup. The Lagrangian tori considered therein are  $\gamma \times \gamma$  and  $\Phi_{h_t}^1(R_\theta(\gamma \times \gamma))$  where  $R_\theta$  is the time- $\theta$  flow of Hamiltonian function  $H = \frac{1}{4}|z_1 - z_2|^2$  which is rotation around the diagonal  $\Delta_{\mathbb{C}} \subset \mathbb{C}^2$  by angle  $\theta$ , and  $\Phi_{h_t}^1$  is the time-1 flow of a generic non-degenerate time-dependent Hamiltonian function  $h_t$  perturbing the Lagrangian torus  $R_\theta(\gamma \times \gamma)$  so that  $\gamma \times \gamma$  and  $\Phi_{h_t}^1(R_\theta(\gamma \times \gamma))$  intersect transversely away from  $\Delta_{\mathbb{C}}$ , on which these two tori intersect cleanly and the clean intersection is  $\Delta_\gamma := \{(p, p) \in \gamma \times \gamma\} \cong \gamma$ . We make a remark here that  $\gamma \times \gamma$  and  $R_\theta(\gamma \times \gamma)$  are invariant under the symplectic involution

$$(z_1, z_2) \mapsto (z_2, z_1)$$

whose fixed point set is  $\Delta_{\mathbb{C}}$ , and  $(\gamma \times \gamma) \cap R_\theta(\gamma \times \gamma) \cap \Delta_{\mathbb{C}} = \Delta_\gamma$ . Inspired from the construction of Heegaard Floer homology, Greene and Lobb manage to establish a version of Lagrangian Floer homology for these two Lagrangian tori as  $\mathbb{Z}$ -graded  $\mathbb{F}_2$ -module whose boundary operator only counts the holomorphic strips avoiding the diagonal  $\Delta_{\mathbb{C}}$ . This is similar to certain Floer homology of the complement of suitable symplectic divisor (or codimension 2 symplectic submanifold) in a symplectic manifold. We expect that the Floer homology in [GL24] can be established analogously in the geometric setup we have considered in this paper for the Lagrangian tori  $L$  and  $L_\phi$ , i.e. the ones in [GL21], where  $\mathbb{C} \times \{0\}$  plays the role of  $\Delta_\gamma$ , and yields the same non-vanishing result as [GL24, Theorem 2.23]. When  $\gamma$  belongs to the generic class in Theorem 3.2, the triple  $(\gamma, \theta, 0)$  would be admissible in the sense of [GL24, Definition 2.3] adapted to our geometric setup. Then the invariance of Floer homology plus the non-vanishing result imply the a priori existence of intersection point in  $L \cap L_\phi$  away from  $\mathbb{C} \times \{0\}$  for such  $\gamma$ , which would weakly achieve the goal we describe at the beginning of this commentary section.

## 6. GENERIC DOUBLING OF CYCLIC QUADRILATERAL PEGS

In [GL23], Greene and Lobb consider a different geometric setup to obtain two embedded Lagrangian tori in  $\mathbb{C}^2$  with respect to  $\omega_{\text{std}}$  from a smooth Jordan curve  $\gamma \subset \mathbb{C}$  as following:

$$T_1 := R_\phi(F_s(\gamma \times \gamma)), \quad T_2 := F_t(\gamma \times \gamma),$$

where  $s, t \in (0, \frac{1}{2}]$ . The map

$$F_r : \mathbb{C}^2 \rightarrow \mathbb{C}^2$$

$$(z_1, z_2) \mapsto \left( (1-r)z_1 + rz_2, \sqrt{r(1-r)}(z_1 - z_2) \right)$$

with  $r \in (0, \frac{1}{2}]$  is a  $\mathbb{C}$ -linear automorphism of  $\mathbb{C}^2$ , and

$$R_\phi : \mathbb{C}^2 \rightarrow \mathbb{C}^2$$


$$(z_1, z_2) \mapsto (z_1, z_2 e^{i\phi})$$

with  $\phi \in (0, \pi)$ . Note that the maps  $F_s$  and  $F_t$  do not preserve  $\omega_{\text{std}}$ , the two embedded Lagrangian tori  $T_1$  and  $T_2$  are not related by Hamiltonian diffeomorphism. In particular,  $T_1$  and  $T_2$  are in general no longer invariant under symplectic involution  $\varrho$  unless  $s = t = \frac{1}{2}$ . Moreover, there is no apparent symmetry for  $T_1$  and  $T_2$  when  $s \neq \frac{1}{2}$  or  $t \neq \frac{1}{2}$ . Nevertheless, they have minimal Maslov number 2 by [Vit90, Pol91], and intersect again cleanly along  $\gamma \times \{0\}$ , c.f. [GL23, Lemma 2].

*Intersection points and pegs.* By [GL23, Lemma 1], an intersection point in  $T_1 \cap T_2 \setminus (\gamma \times \{0\})$ , which is a solution of the equation

$$R_\phi(F_s(A, C)) = F_t(B, D), \quad A \neq C \ (\Leftrightarrow B \neq D),$$

for  $A, B, C, D \in \mathbb{C}$  gives rise to an inscribed cyclic quadrilateral on the curve  $\gamma$



with

$$s = \frac{|AX|}{|AC|}, \quad t = \frac{|BX|}{|BD|}, \quad \phi = \angle AXB.$$

The triple  $(s, t, \phi)$  determines the counterclockwise oriented cyclic quadrilateral uniquely up to similarity, unless  $s = \frac{1}{2}$  or  $t = \frac{1}{2}$  in which case  $(s, t, \phi)$  and  $(t, s, \pi - \phi)$  determine the same oriented cyclic quadrilateral up to similarity. When  $s = t = \frac{1}{2}$ , this recovers the geometric setup in [GL21] as described in §1.

The correspondence between intersection points in  $T_1 \cap T_2 \setminus (\gamma \times \{0\})$  and cyclic quadrilaterals with data  $(s, t, \phi)$  inscribed on  $\gamma$  is described as follows.

- (1) If  $s \neq \frac{1}{2}$  or  $t \neq \frac{1}{2}$ , the intersection points are bijective to the inscribed cyclic quadrilaterals.
- (2) If  $s = t = \frac{1}{2}$ , this is the case as discussed in §2, Lemma 2.3.

*Existence of pegs.* After performing a Lagrangian surgery [GL23, p.933-934], the resulting object  $T_1 \#_{\gamma \times \{0\}} T_2$  is an immersed Lagrangian torus with minimal Maslov number 4, c.f. [GL23, Proposition, p.434]. The self-intersection points in  $T_1 \#_{\gamma \times \{0\}} T_2$  are one-to-one corresponding to intersection points in  $T_1 \cap T_2 \setminus (\gamma \times \{0\})$ . Thus, if there were no self-intersection points in  $T_1 \#_{\gamma \times \{0\}} T_2$ , it would be an embedded Lagrangian torus in  $\mathbb{C}^2$ , whose minimal Maslov number must be 2 by [Vit90, Pol91]. This contradiction shows the a priori existence of inscribed cyclic quadrilateral with any  $(s, t, \phi)$  inscribed on any smooth Jordan curve  $\gamma$ , c.f. [GL23, Theorem, p.931].

*Transversality.* Mimicking the discussion in §3, a similar geometric transversality result holds as follows.

**Theorem 6.1.** *For any fixed data  $s, t \in (0, \frac{1}{2}]$  and  $\phi \in (0, \pi)$ , there is a residual set of smooth Jordan curves such that for every  $\gamma$  in the set, the corresponding smooth Lagrangian tori  $T_1$  and  $T_2$  constructed in above way has only finitely many transversal intersection points outside a small open neighborhood of  $\gamma \times \{0\}$ .*

We provide a sketch of proof which is completely similar to that of Theorem 3.2. The only degenerate cyclic quadrilaterals, which needs to be circumvented, is  $A = B = C = D$ . These amount to the clean intersection points  $\gamma \times \{0\}$ . A map  $\mathcal{F}$  in analogy to that in the proof of Theorem 3.2 can be constructed as:

$$\begin{aligned} \mathcal{F} : \text{Emb}^k(S^1, \mathbb{C}) \times ((S^1)^4 \setminus \Delta_{(1,2,3,4)}^{(S^1)^4}) &\rightarrow \mathbb{C}^2 \times \mathbb{C}^2 \\ (\gamma, t_1, t_2, t_3, t_4) &\mapsto (R_\phi(F_s(\gamma(t_1) \times \gamma(t_2))), F_t(\gamma(t_3) \times \gamma(t_4))) \end{aligned}$$

The set  $\mathcal{F}^{-1}(\Delta_{\mathbb{C}^2})$  can be shown to be a Banach submanifold similarly by implicit function theorem, and the projection

$$\begin{aligned} \pi|_{\mathcal{F}^{-1}(\Delta_{\mathbb{C}^2})} : \mathcal{F}^{-1}(\Delta_{\mathbb{C}^2}) &\rightarrow \text{Emb}^k(S^1, \mathbb{C}) \\ (\gamma, t_1, t_2, t_3, t_4) &\mapsto \gamma \end{aligned}$$

is a Fredholm map with Fredholm index 0, whose regular values amount to the geometric transversal condition  $T_1 \times T_2 \pitchfork \Delta_{\mathbb{C}^2}$  in  $\mathbb{C}^2 \times \mathbb{C}^2$  outside  $(\gamma \times \{0\}) \times (\gamma \times \{0\})$ , in other words  $T_1 \pitchfork T_2$  in  $\mathbb{C}^2$  outside  $\gamma \times \{0\}$ . By Sard-Samle theorem, the  $C^k$ -genericity statement follows. Further via the same argument at the end of the proof of Theorem 3.2, the  $C^k$ -genericity can be promoted to  $C^\infty$ -genericity.

*Doubling.* In the current situation, the two embedded smooth Lagrangian tori  $T_1$  and  $T_2$  are not Hamiltonian isotopic to each other and even not monotone. The Floer homology of monotone Lagrangian submanifolds introduced in Appendix A is not applicable. Nevertheless, the computational formula for algebraic intersection number of cleanly intersecting Lagrangian submanifolds, i.e. (B.4), is still valid. Suppose now  $\gamma$  lies in the generic class in Theorem 6.1. Assume that there is only one intersection point in  $T_1 \cap T_2 \setminus (\gamma \times \{0\})$  due to the a priori existence result in [GL23], then by (B.4), we have

$$\begin{aligned} T_1 \bullet T_2 &= \pm \chi(\gamma \times \{0\}) \pm 1 \\ &= \pm 1. \end{aligned}$$

However,  $T_1$  and  $T_2$  can be displaced from each other by translation in  $\mathbb{C}^2$ , thus we also have

$$T_1 \bullet T_2 = \text{sign}(\emptyset) = 0.$$

This contradiction leads to the existence of a second intersection point distinct from the first one in  $T_1 \cap T_2 \setminus (\gamma \times \{0\})$ . This shows that for a generic class of smooth Jordan curve there are two distinct cyclic quadrilateral pegs with  $(s, t, \phi)$  such that  $s \neq \frac{1}{2}$  or  $t \neq \frac{1}{2}$  by Theorem 6.1 together with the previous discussion on intersection points and inscribed cyclic quadrilaterals. Combining with Theorem 1.6, we have proved the following extended statement.

**Theorem 6.2.** *For any fixed data  $s, t \in (0, \frac{1}{2}]$  and  $\phi \in (0, \pi)$  except  $s = t = \frac{1}{2}$  and  $\phi = \frac{\pi}{2}$ , there is a generic class of smooth Jordan curves such that any cyclic quadrilateral in  $\mathbb{R}^2$  with the fixed data  $(s, t, \phi)$  admits at least two similar ones whose vertices land distinctly on any curve in this generic class.*

At the end, we make a comment similar to §5. We may consider whether there could be a version of Lagrangian Floer homology for  $T_1$  and  $T_2$  in order to prove the existence of cyclic quadrilateral pegs on smooth Jordan curves. The Floer homology in [GL24] cannot be established for  $T_1$  and  $T_2$  straightforwardly due to the fact that they are not monotone in general. The fact that  $T_1$  and  $T_2$  are not Hamiltonian isotopic to each other in general causing much more difficulty for computing intersection Lagrangian Floer homology to obtain a non-vanishing result and conclude

$$T_1 \cap T_2 \setminus (\gamma \times \{0\}) \neq \emptyset,$$

presumably we could manage to establish the intersection Lagrangian Floer homology from  $T_1$  and  $T_2$  in certain symplectic divisor complement removing  $\mathbb{C} \times \{0\}$  which contains the clean intersection  $\gamma \times \{0\}$ . Achieving such goal might heavily rely on the study of the algebraic structure of Floer homology.

## APPENDIX A. LAGRANGIAN FLOER HOMOLOGY

The Floer homology of Lagrangian submanifolds was first introduced by Andreas Floer, c.f. [Flo88I, Flo88II, Flo88III] in aspherical setting, and further extended to more general setting, e.g. [Oh93, Oh95, Oh96], see also [FOOO09, Oh15]. In this appendix, we give an overview of the Lagrangian Floer homology within the scope of this paper.

**A.1. Floer homology of monotone Lagrangian submanifolds.** We consider a *monotone symplectic manifold*  $(M, \omega)$ , i.e.

$$I_{c_1} = \kappa I_\omega$$

for some constant  $\kappa > 0$ , where  $I_\omega : \pi_2(M) \rightarrow \mathbb{R}$  is the evaluation map of symplectic area, and  $I_{c_1} : \pi_2(M) \rightarrow \mathbb{R}$  is the evaluation map of the first Chern class  $c_1(TM) \in H^2(M; \mathbb{Z})$ . Suppose there are two compact Lagrangian submanifolds  $L_0$  and  $L_1$  which are *monotone*, namely

$$(A.1) \quad I_\omega = \tau I_\mu$$

for some constant  $\tau > 0$ , where  $I_\omega : \pi_2(M, L_i) \rightarrow \mathbb{R}$  is the evaluation map of the symplectic area, and  $I_\mu : \pi_2(M, L_i) \rightarrow \mathbb{Z}$  is the evaluation map of the Maslov class  $\mu \in H^1(\Lambda(\mathbb{C}^n); \mathbb{Z})$ , for each  $i = 0, 1$ .

The symplectic action functional

$$(A.2) \quad \mathcal{A}([x, \bar{x}]) = - \int_{[0,1] \times [0,1]} \bar{x}^* \omega$$

is well-defined on the Novikov covering space

$$\tilde{\mathcal{P}}(L_0, L_1)$$

of the path space with Lagrangian boundary condition:

$$\mathcal{P}(L_0, L_1) := \{x : [0, 1] \rightarrow M : x(0) \in L_0, x(1) \in L_1\}.$$

C.f. [FOOO09, §2.2], [Oh15, §13.4]. The Novikov covering space  $\tilde{\mathcal{P}}(L_0, L_1)$  consists of equivalence classes of pairs  $[x, \bar{x}]$  where  $\bar{x} : [0, 1] \times [0, 1] \rightarrow M$  called the *cap* of  $x$  satisfies  $\bar{x}(1, \cdot) = x$ ,  $\bar{x}(0, \cdot) = \ell_0 \in \mathcal{P}(L_0, L_1)$  (i.e. based point in path space),  $\bar{x}(\cdot, 0) \in L_0$ ,  $\bar{x}(\cdot, 1) \in L_1$ , and the equivalence relation is given by

$$(x, \bar{x}) \sim (x, \tilde{x}) :\Leftrightarrow I_\omega(-\bar{x} \# \tilde{x}) = 0 = I_\mu(-\bar{x} \# \tilde{x}).$$

The Deck transformation group is

$$(A.3) \quad \Gamma_{\ell_0} := \Gamma(L_0, L_1; \ell_0) := \frac{\pi_1(\mathcal{P}(L_0, L_1), \ell_0)}{\ker(I_\omega) \cap \ker(I_\mu)},$$

where  $I_\omega : \pi_1(\mathcal{P}(L_0, L_1), \ell_0) \rightarrow \mathbb{R}$  and  $I_\mu : \pi_1(\mathcal{P}(L_0, L_1), \ell_0) \rightarrow \mathbb{Z}$ , c.f. [FOOO09, §2.2.1], [Oh15, §13.4]. The group  $\Gamma_{\ell_0}$  acts on  $\tilde{\mathcal{P}}(L_0, L_1)$  transitively and effectively. The critical points set of the action functional  $\text{Crit}(\mathcal{A})$  is the set of intersection points between these two Lagrangian submanifolds together with capping, i.e.

$$\text{Crit}(\mathcal{A}) \cong (L_0 \cap L_1) \times \pi_1(\mathcal{P}(L_0, L_1), \ell_0).$$

The Floer complex is defined as follows. The Floer chain module is a free  $\mathbb{F}_2$ -module generated by equivalence classes

$$[x, \bar{x}] \in \frac{\text{Crit}(\mathcal{A})}{\ker(I_\omega) \cap \ker(I_\mu)} \cong (L_0 \cap L_1) \times \Gamma_{\ell_0},$$

i.e.

$$CF_*(L_0, L_1, \ell_0) := \bigoplus_{[x, \bar{x}] \in (L_0 \cap L_1) \times \Gamma_{\ell_0}} \mathbb{F}_2 \langle [x, \bar{x}] \rangle$$

and

$$CF_*(L_0, L_1) := \bigoplus_{\ell_{0,i}} CF_*(L_0, L_1, \ell_{0,i}),$$

in which  $\ell_{0,i}$  runs over the set of base points of connected components of  $\mathcal{P}(L_0, L_1)$ . This is an infinitely generated  $\mathbb{F}_2$ -vector space. There is a natural  $\Gamma_{\ell_0}$ -action endowing  $CF_*(L_0, L_1, \ell_0)$  a  $\mathbb{F}_2[\Gamma_{\ell_0}]$ -module structure: For any  $g \in \Gamma_{\ell_0}$ ,

$$(A.4) \quad g \cdot [x, \bar{x}] = [x, g\#\bar{x}]$$

where the element  $g$  can be regarded as a map  $S^1 \times [0, 1] \rightarrow M$  satisfying  $g(\cdot, 0) \in L_0$ ,  $g(\cdot, 1) \in L_1$ . The boundary operator is given by counting rigid holomorphic strips which is described as follows. We pick a smooth family  $\{J_t\}_{t \in [0, 1]}$  of  $\omega$ -compatible almost complex structures whose set is denoted by  $\mathcal{J}(M, \omega)$ . This induces a smooth family of Riemannian metrics  $g_{J_t}(\cdot, \cdot) = \omega(\cdot, J_t \cdot)$ , and further a  $L^2$ -inner product

$$\langle \cdot, \cdot \rangle_J := \int_0^1 \omega(\cdot, J_t \cdot) dt$$

on  $\mathcal{P}(L_0, L_1)$  and hence on  $\tilde{\mathcal{P}}(L_0, L_1)$ . The negative  $L^2$ -gradient flow equation of the symplectic action functional  $\mathcal{A}$  reads: Smooth map  $u : \mathbb{R} \times [0, 1] \rightarrow M$  solving

$$(A.5) \quad \begin{cases} \partial_s u + J_t \partial_t u = 0, \\ u(s, 0) \in L_0, u(s, 1) \in L_1. \end{cases}$$

This is literally the non-linear Cauchy-Riemann equation

$$\bar{\partial}_{J_t} u = 0$$

for  $J$ -holomorphic strip with boundary condition on  $L_0$  and  $L_1$  in terms of coordinate  $(s, t) \in \mathbb{R} \times [0, 1]$ . The energy or the  $L^2$ -norm of the  $J$ -holomorphic strip is equal to its symplectic area:

$$\begin{aligned} E(u) &:= \int_{\mathbb{R} \times [0, 1]} \|\partial_s u\|_{J_t}^2 ds \wedge dt \\ &= \frac{1}{2} \int_{\mathbb{R} \times [0, 1]} (\|\partial_s u\|_{J_t}^2 + \|\partial_t u\|_{J_t}^2) ds \wedge dt \\ &= \int_{\mathbb{R} \times [0, 1]} u^* \omega \end{aligned}$$

When the energy of  $J$ -holomorphic strip is finite, i.e.

$$E(u) < \infty,$$

then the solution has asymptotic behavior:

$$\lim_{s \rightarrow \pm\infty} u(s, \cdot) = x_{\pm} \in L_0 \cap L_1$$

and

$$E(u) = \mathcal{A}([x_-, \bar{x}_-]) - \mathcal{A}([x_+, \bar{x}_+])$$

if  $[x_+, \bar{x}_+] = [x_+, \bar{x}_- \# u]$ , vice versa. C.f. [FOOO09, §2.3.1], [Oh15, §13.9.1]. We denote by

$$\mathcal{M}(x_-, x_+; J_t) := \tilde{\mathcal{M}}(x_-, x_+; J_t) / \mathbb{R}$$

the (unparameterized) moduli space of  $J$ -holomorphic strips with  $x_{\pm} \in L_0 \cap L_1$  as asymptotics, where

$$\widetilde{\mathcal{M}}(x_-, x_+; J_t) := \left\{ u \in C^\infty(\mathbb{R} \times [0, 1], M) : u \text{ solves (A.5) with } \lim_{s \rightarrow \pm\infty} u(s, \cdot) = x_{\pm} \right\}$$

is the parameterized moduli space which carries a proper and transitive  $\mathbb{R}$ -action by shift of  $s$ -parameter. There is a decomposition

$$\widetilde{\mathcal{M}}(x_-, x_+; J_t) = \bigcup_{B \in \pi_2(x_-, x_+)} \widetilde{\mathcal{M}}(x_-, x_+; B, J_t)$$

where  $\pi_2(x_-, x_+)$  is the set of homotopy classes of maps  $u : [0, 1] \times [0, 1] \rightarrow M$  relative to boundary  $u(0, \cdot) = x_-$ ,  $u(1, \cdot) = x_+$ ,  $u(\cdot, 0) \in L_0$ ,  $u(\cdot, 1) \in L_1$ , it is a subset in the relative homotopy group  $\pi_2(M, L_0 \cup L_1)$ . The space  $\widetilde{\mathcal{M}}(x_-, x_+; B, J_t)$  consists of those  $J$ -holomorphic strips  $u$  with homotopy class  $[u] = B$ . The virtual dimension of the moduli space is computed by Fredholm index of the Cauchy-Riemann operator  $\bar{\partial}_{J_t}$  at  $u$  (c.f. [Oh15, Theorem 15.3.2]):

$$(A.6) \quad \dim \widetilde{\mathcal{M}}(x_-, x_+; B, J_t) = \text{ind}(u) = \mu(x_-, x_+; B)$$

where

$$\text{ind}(u) := \text{ind}(\bar{\partial}_{J_t}|_u) := \text{ind}(D_u \bar{\partial}_{J_t})$$

is the Fredholm index of the Fredholm operator  $\bar{\partial}_{J_t}|_u$  associated to  $u$ , and

$$\mu(x_-, x_+; B) \in \mathbb{Z}$$

is the Maslov-Viterbo index of  $u$  introduced by [Vit87], see also [Flo88II], [FOOO09, Definition 2.3.9], [Oh15, Definition 13.6.2]. The Maslov-Viterbo index of  $u$  can be expressed as the difference of the Maslov-Morse index (definition refers to [FOOO09, Definition 2.2.12], [Oh15, Definition 13.6.2]) of the pairs  $[x_-, \bar{x}_-]$  and  $[x_+, \bar{x}_+]$ :

$$(A.7) \quad \mu(x_-, x_+; B) = \mu([x_+, \bar{x}_+]) - \mu([x_-, \bar{x}_-])$$

with  $\bar{x}_+ = \bar{x}_- \# B$ . C.f. [FOOO09, Proposition 2.3.9], [Oh15, Proposition 13.6.3]. This formula is independent of the choice of the cap  $\bar{x}_-$ . In some nice situations, one has transversality which would imply that, for a generic choice of  $\{J_t\}_{t \in [0, 1]} \subset \mathcal{J}(M, \omega)$ , the parameterized moduli space  $\widetilde{\mathcal{M}}(x_-, x_+; B, J_t)$  is a smooth manifold of expected dimension (A.6), and hence the (unparameterized) moduli space  $\mathcal{M}(x_-, x_+; B, J_t)$  is a smooth manifold of dimension

$$(A.8) \quad \dim \mathcal{M}(x_-, x_+; B, J_t) = \mu(x_-, x_+; B) - 1.$$

E.g. refer to the statement [Oh93, Proposition 3.3], [FOOO09, Theorem 2.3.11], [Oh15, Theorem 15.1.3]. In general, the moduli space  $\mathcal{M}(x_-, x_+; J_t)$  can be further compactified due to Floer-Gromov convergence, by adding broken  $J$ -holomorphic strips,  $J$ -holomorphic disks and  $J$ -holomorphic spheres (these lateral two are also known as disk and sphere bubbling), into a compact space  $\overline{\mathcal{M}}(x_-, x_+; J_t)$  with respect to Floer-Gromov topology. Provided above transversality, for a generic choice of  $\{J_t\}_{t \in [0, 1]} \subset \mathcal{J}(M, \omega)$ , the compact space  $\overline{\mathcal{M}}(x_-, x_+; J_t)$  is a stratified space, whose strata are supposed to carry manifold structures. E.g. for the purpose of defining Floer complex, we would like to have enough transversality so that the 0-dimensional stratum  $\mathcal{M}^0$  is compact 0-dimensional manifold (i.e. finite point set), and the 1-dimensional stratum  $\overline{\mathcal{M}}^1$  is a compact 1-dimensional manifold with boundary  $\partial \overline{\mathcal{M}}^1 = \mathcal{M}^0$ . The details for these topological structures of moduli



spaces are rather technical, for nice expositions, we refer to [FOOO09, Chapter 2], [Oh15, §14, §15, §16]. Now with all the above necessary ingredients, the boundary operator is written as

$$\begin{aligned} \partial : CF_*(L_0, L_1) &\rightarrow CF_{*-1}(L_0, L_1) \\ [x, \bar{x}] &\mapsto \sum_{\substack{y \in L_0 \cap L_1 \\ B \in \pi_2(x, y) \\ \mu(x, y; B) = 1}} \#_2 \mathcal{M}^0(x, y; B, J_t)[y, \bar{y}] \end{aligned}$$

where  $\#_2$  stands for mod 2 count of the 0-dimensional moduli space  $\mathcal{M}^0(x, y; B, J_t)$ . In general, due to the formula (A.7), one can put a relative grading or an absolute grading modulo some integer (i.e. a periodic grading) on the Floer complex provided by the Maslov-Morse index (c.f. [Oh15, §13.9, p.65]): For the pair  $[x, \bar{x}]$ , due to (A.4), the grading differs by

$$(A.9) \quad \mu([x, g\#\bar{x}]) = \mu([x, \bar{x}]) - I_\mu(g)$$

for  $g \in \Gamma_{\ell_0}$ , note that we can view the difference of various caps, i.e.  $(-\bar{x})\#\bar{x}$  as such an element  $g$ . See also [Flo88II], [Oh93, Lemma 4.7]. When the boundary operator satisfies the condition

$$\partial \circ \partial = 0,$$

we say that the Lagrangian submanifolds are *unobstructed*. In such case, the Lagrangian Floer homology is well-defined:

$$HF_*(L_0, L_1) := \frac{\ker(\partial : CF_* \rightarrow CF_{*-1})}{\text{im}(\partial : CF_{*+1} \rightarrow CF_*)}.$$

Otherwise, we say the Lagrangian submanifolds are *obstructed*, and there is *anomaly*. The appearance of anomaly often concerns the disk bubbling, c.f. [Oh95], [FOOO09, §2.4], [Oh95, 16.4.2].

In the monotone case, as we have indicated at the beginning of this section, the Lagrangian Floer homology is established by Yong-Geun Oh.


**Theorem A.10** ([Oh93],[Oh95],[Oh15, §16]). *Let  $L_0, L_1$  be compact monotone Lagrangian submanifolds in a compact monotone symplectic manifold  $(M, \omega)$  with minimal Maslov number  $N_{L_0}, N_{L_1}$ , and put  $N := \min\{N_{L_0}, N_{L_1}\}$ .*

- (i) *Suppose  $L_0 \pitchfork L_1$  and  $N \geq 3$ . Then the Floer homology  $HF_*(L_0, L_1)$  is well-defined as a  $\mathbb{F}_2$ -module with  $\mathbb{Z}/N\mathbb{Z}$ -grading.*
- (ii) *Suppose  $N = 2$ , and  $L_1 = \varphi_H(L_0)$  with  $\varphi_H$  a Hamiltonian diffeomorphism generated by a non-degenerate Hamiltonian, i.e.  $L_0 \pitchfork \varphi_H(L_0)$ . Then the Floer homology  $HF_*(L_0, \varphi_H(L_0))$  is well-defined as a  $\mathbb{F}_2$ -module with  $\mathbb{Z}/N\mathbb{Z}$ -grading.*


*In particular, the Floer homology  $HF_*(L_0, L_1)$  is independent of the choice of  $\omega$ -compatible almost complex structures and invariant under Hamiltonian diffeomorphisms.*

**Remark A.11.** Since the compactified moduli space  $\overline{\mathcal{M}}^1$  of dimension 1 is involved in the definition of boundary operator  $\partial$  and the boundary stratum  $\mathcal{M}^0 = \partial\overline{\mathcal{M}}^1$  is relevant for the condition  $\partial^2 = 0$ . These are all smooth manifolds in monotone case, c.f. [Oh15, 16.4.2]. When  $N \geq 3$ , there are only once-broken  $J$ -holomorphic

strips, whose each component is of Maslov index 1, in the boundary stratum due to dimensional reason.



When  $N = 2$ , there are additionally  $J$ -holomorphic disks of Maslov index 2, i.e. disk bubbling, in the boundary stratum. C.f. [Oh15, Theorem 16.4.4].



In particular, there is obstruction due the appearance of disk bubbling, namely the boundary operator satisfies

$$(A.12) \quad \partial \circ \partial = \Phi_{L_0}(\text{pt.}) - \Phi_{L_1}(\text{pt.}).$$

where  $\Phi_{L_i}(\text{pt.})$  is the one-point open Gromov-Witten invariant for Lagrangian submanifolds  $L_i$ ,  $i = 0, 1$  and  $\text{pt.}$  represents an arbitrary point in  $L_i$  respectively, c.f. [Oh15, Example 15.4.3, p.163]. If  $L_1 = \varphi_H(L_0)$  for some Hamiltonian diffeomorphism  $\varphi_H$ , then it is shown that the right hand side of (A.12) cancels, hence the obstruction vanishes. C.f. [Oh95], [Oh15, §16.4.2].

It is more delightful to work with a  $\mathbb{Z}$ -graded bounded chain complex and homology instead of the  $N$ -periodic unbounded chain complex and homology. For this purpose, we introduce the universal Novikov ring over  $\mathbb{F}_2$  and define the Floer homology with such coefficient.

$$(A.13) \quad \Lambda := \left\{ \sum_{i=0}^{\infty} c_i T^{a_i} \mid c_i \in \mathbb{F}_2, a_i \in \mathbb{R}_{\geq 0}, \lim_{i \rightarrow \infty} a_i = \infty \right\}$$

where  $T$  is a formal parameter, and in the monotone case with  $N := \min\{N_{L_0}, N_{L_1}\}$ , one can put the grading  $|T| = -N$  on it. A systematic discussion of Novikov rings can be found in [FOOO09, §2.2.1, §2.4.3, §5.1], [Oh15, §13.4, §13.8]. The Floer complex is now defined as the free  $\Lambda$ -module finitely generated by  $L_0 \cap L_1$ , i.e.

$$CF_*(L_0, L_1; \Lambda) = \bigoplus_{x \in L_0 \cap L_1} \mathbb{F}_2 \langle x \rangle \otimes_{\mathbb{F}_2} \Lambda.$$

The boundary operator is

$$\begin{aligned} \partial : CF_*(L_0, L_1; \Lambda) &\rightarrow CF_{*-1}(L_0, L_1; \Lambda) \\ x &\mapsto \sum_{\substack{y \in L_0 \cap L_1 \\ B \in \pi_2(x, y) \\ \mu(x, y; B) = 1}} \#_2 \mathcal{M}^0(x, y; B, J_t) y \otimes T^{\omega(B)} \end{aligned}$$

given by the mod 2 count of 0-dimensional moduli space  $\mathcal{M}^0(x, y; B)$  of  $J$ -holomorphic strips of homotopy class  $B \in \pi_2(x, y)$  with intersection points  $x$  and  $y$  as asymptotics. By assuming  $\langle c_1(TM), \pi_2(M) \rangle = 0$ , due to [Flo88II, Theorem 4], and fixing a cap for each  $x \in L_0 \cap L_1$ , we provide an absolute  $\mathbb{Z}$ -grading of the Floer complex. In this sense, we obtain a  $\mathbb{Z}$ -graded Floer homology

$$HF_*(L_0, L_1; \Lambda) := H_*(CF_*(L_0, L_1; \Lambda), \partial)$$

as  $\Lambda$ -module which is also invariant under Hamiltonian diffeomorphisms. In particular, when the set of intersection points  $L_0 \cap L_1$  is finite, the Floer complex is a bounded  $\mathbb{Z}$ -graded chain complex over  $\Lambda$ , and so is the Floer homology a bounded  $\mathbb{Z}$ -graded  $\Lambda$ -module.

**A.2. Morse-Bott Floer homology.** Lagrangian Floer theory in the Morse-Bott setting was introduced and developed in [Po94, Fra03, FOOO09, Sch16]. We follow the cascade approach in [Fra03, Sch16]. Assume  $L_0$  and  $L_1$  intersect cleanly, namely  $L_0 \cap L_1$  consists of a disjoint union of connected submanifolds. Suppose  $C \subseteq L_0 \cap L_1$ , then  $T_p L_0 \cap T_p L_1 = T_p C \forall p \in C$ . The action functional (A.2) is a Morse-Bott function on the path space in the sense that

$$\ker(\text{Hess } \mathcal{A}|_{T_p \mathcal{P}(L_0, L_1)}) = T_p C, \quad \forall p \in C.$$

Suppose

$$L_0 \cap L_1 = \bigsqcup_{i \in I} C_i$$

where each  $C_i$  is a connected closed submanifold. We choose auxiliary Morse functions  $f_i$  on  $C_i$ . The Floer chain  $\Lambda$ -module is now freely generated by critical points of Morse function  $x \in \text{Crit}(f_i)$ ,  $i \in I$ :

$$CF_*(L_0, L_1, \{f_i\}; \Lambda) := \bigoplus_{i \in I} \bigoplus_{x \in \text{Crit}(f_i)} \Lambda \langle x \rangle.$$

The grading of  $x \in \text{Crit}(f_i)$  is now given by


$$(A.14) \quad |x| = \text{ind}_{f_i}(x) + \mu([x, \bar{x}]) - \frac{1}{2} \dim C_i, \quad i \in I,$$

where  $\text{ind}_{f_i}(x)$  is the Morse index of  $x$ , and  $\bar{x}$  is a fixed capping of  $x$  in a canonical way (c.f. [FOOO09, §3.7 (3.7.49)]) such that  $\mu([x, \bar{x}])$  stays the same for all points  $x \in C_i$ . Therefore, one can assign an integral grading  $\mu(C_i)$  for each Morse-Bott component  $C_i$  in  $L_0 \cap L_1$  by [Fra03, Appendix C, p.177], [Sch16, §9.1.2]:

$$(A.15) \quad \mu(C_i) := \mu([x, \bar{x}]) - \frac{1}{2} \dim C_i, \quad i \in I.$$

To define the boundary operator, a generalization of Floer trajectories called *cascades* is counted. The formal definition of trajectories with cascades can be found in [Fra03, Definition C.22], [Sch16, §9.1.1]. These are tuples of  $J$ -holomorphic strips (or Floer trajectories) and Morse trajectories connecting the critical points

in sequel.



For generic choice of  $\{J_t\}_{t \in [0,1]} \subset \mathcal{J}(M, \omega)$  and Riemannian metric  $g$  on  $L_0 \cap L_1$ , the (unparameterized) moduli space  $\mathcal{M}(x, y; J_t, g)$  of cascades connecting  $x \in \text{Crit}(f_i)$  and  $y \in \text{Crit}(f_j)$  is a smooth manifold of dimension  $|x| - |y| - 1$ . C.f. [Fra03, Theorem C.23], [Sch16, §9.2.1]. There is also a Floer-Gromov compactification for the moduli space of cascades, c.f. [Sch16, §9.2.2]. And in the monotone setting, despite the broken trajectories, disk bubbling can also appear, refer to Remark A.11. The boundary operator

$$\partial : CF_*(L_0, L_1, \{f_i\}; \Lambda) \rightarrow CF_{*-1}(L_0, L_1, \{f_i\}; \Lambda)$$

is defined similarly as before by mod 2 count of  $\mathcal{M}^0(x, y; J_t, g)$ .

In the monotone setting as before, we obtain the following.

**Theorem A.16** ([Fra03, Appendix C.4], [Sch16, §9.3]). *The Floer homology*

$$HF_*(L_0, L_1, \{f_i\}, J_t, g; \Lambda) := H_*(CF_*(L_0, L_1, \{f_i\}; \Lambda), \partial)$$

is well-defined, independent of the choice of  $(\{f_i\}, J_t, g)$ . Moreover it is invariant under Hamiltonian diffeomorphism. Namely, there is a canonical isomorphism

$$HF_*(L_0, L_1, \{f_i\}, J_t, g; \Lambda) \cong HF_*(L_0, \varphi_H(L_1), \{\tilde{f}_j\}, \tilde{J}_t, \tilde{g}; \Lambda)$$

for any  $\varphi_H \in \text{Ham}(M, \omega)$  such that  $L$  intersects with  $\varphi_H(L)$  cleanly,  $(\{f_i\}, J_t, g)$  and  $(\{\tilde{f}_j\}, \tilde{J}_t, \tilde{g})$  are two arbitrary generic choices of collection of admissible data.

**A.3. Liouville manifold.** The previous discussions for Floer homology are all settled in compact (closed) symplectic manifolds. There is suitable generalization to non-compact symplectic manifolds, e.g. exact symplectic manifolds. In particular, we focus on the most studied case called the Liouville manifold. A *Liouville manifold* is an exact symplectic manifold  $(M, d\lambda)$  with a primitive  $\lambda$  called Liouville 1-form, and a complete Liouville vector field  $Z$ , i.e.  $i_Z d\lambda = \lambda$ , and an exhaustion by Liouville domains. More precisely, a *Liouville domain* is a compact exact symplectic manifold  $(W, d\lambda)$  with contact type boundary  $\partial W$  (i.e.  $\alpha := \lambda|_{\partial W}$  is a contact 1-form) such that the Liouville vector field  $Z$ , which is uniquely determined by the Liouville 1-form, is pointing outwards along  $\partial W$ . The *completion of a Liouville domain*  $(W, d\lambda)$  denoted by  $(\widehat{W}, d\widehat{\lambda})$  is obtained by gluing the *symplectization* of

$\partial W$  to the Liouville domain  $W$  in the following way: The negative part of symplectization of  $\partial W$  is embedded onto a tubular neighborhood of  $\partial W$  in  $W$  via the flow of Liouville vector field

$$\begin{aligned} ((0, 1] \times \partial W, d(r\alpha)) &\hookrightarrow (W, d\lambda) \\ (r, x) &\mapsto \varphi_{\ln r}^Z(x), \end{aligned}$$

while the positive part  $([1, \infty) \times \partial W, d(r\alpha))$  is attached along the boundary  $\partial W$  by identifying  $\partial W$  with  $\{1\} \times \partial W$ . The  $\widehat{\lambda}$  denotes the 1-form globally constructed as above in  $\widehat{W}$  such that  $\widehat{\lambda}|_W = \lambda$  and  $\widehat{\lambda}|_{[1, \infty) \times \partial W} = d(r\alpha)$ , and the Liouville vector field  $Z$  on  $[1, \infty) \times \partial W$  is given by  $r\partial_r$ . Then Liouville manifold is nothing else than the completion of a Liouville domain.

**Example A.17.** The most common examples are the following.

- (1)  $(\mathbb{R}^{2n}, d\lambda_{\text{std}})$  has Liouville domains as  $2n$ -disk  $(D^{2n}, d\lambda_{\text{std}})$  with Liouville 1-form

$$\lambda_{\text{std}} = \frac{1}{2} \sum_{i=1}^n (x_i dy_i - y_i dx_i),$$

and Liouville vector field

$$Z = \frac{1}{2} \sum_{i=1}^n (x_i \partial_{x_i} + y_i \partial_{y_i}),$$

where  $(x_1, y_1, \dots, x_n, y_n) \in \mathbb{R}^{2n}$ .

- (2)  $(T^*Q, d\lambda_{\text{can}})$  the cotangent bundle of a connected closed Riemannian  $n$ -manifold  $Q$  has Liouville domain as codisk bundle  $(D^*Q, d\lambda_{\text{can}})$  with Liouville 1-form

$$\lambda_{\text{can}} = \sum_{i=1}^n p_i dq_i,$$

and Liouville vector field

$$Z = \sum_{i=1}^n p_i \partial_{p_i},$$

where  $(q_1, \dots, q_n, p_1, \dots, p_n) \in T^*Q$ .

Lagrangian Floer theory for Liouville manifolds are introduced and established in [AS10] with certain exact Lagrangian submanifolds as objects. One of the most important feature is that they provide a natural choice of  $d\widehat{\lambda}$ -compatible almost complex structure  $J$ , which is said to be of *contact type* on  $[r_0, \infty) \times \partial W$  for some  $r_0 \geq 1$ :

$$\widehat{\lambda} \circ J = r dr$$

where  $r \in [1, \infty)$ . This is equivalent to  $J(r\partial_r) = R$  where  $R$  is the Reeb vector field of the contact form  $\alpha$  on  $\partial W$ , i.e.  $i_R\alpha = 1$  and  $i_R d\alpha = 0$ . Then an  $d\widehat{\lambda}$ -compatible almost complex structure  $J$  is *admissible* if it is of contact type for  $r_0 \gg 1$ . By maximum principle and similar type of argument, c.f. [AS10, §7], there is  $C^0$ -bound of the Floer trajectories or  $J$ -holomorphic strips, which is required to obtain the Floer-Gromov compactness for the moduli spaces. See also [Oh15, §12.3] for the treatment in cotangent bundles specially. Another important feature in Liouville manifold is that there is no non-constant  $J$ -holomorphic sphere due to exactness of the symplectic manifold, hence sphere bubbling is forbidden by geometric reason.

The Floer theory for compact monotone Lagrangian submanifolds in a Liouville manifold instead of exact Lagrangian submanifolds can be established similarly in the geometric setting with convex type compatible almost complex structure. This is studied e.g. in  $T^*S^n$  [AD23], in which Morse-Bott technique is also involved. For our purpose, we do not need a full Lagrangian Floer theory with higher algebraic structures, instead we only need to established the Lagrangian Floer homology for closed monotone Lagrangian submanifolds in Liouville manifold. Due to above discussions, there is a generic choice of admissible compatible almost complex structure such that all the statements for moduli spaces in Appendix A.1 and Appendix A.2 still hold in Liouville manifold, and the desired Lagrangian Floer homology is well-defined.

## APPENDIX B. ALGEBRAIC INTERSECTION NUMBER FOR CLEAN INTERSECTIONS

Urs Frauenfelder, Zhen Gao

In this appendix, we introduce the notion of clean normal bundle over a clean intersection. The algebraic intersection number can be computed directly in terms of the Euler number of the clean normal bundle. In the special case of Lagrangian intersections the clean normal bundle coincides with the tangent bundle of the intersection.

**B.1. Introduction.** Suppose throughout the appendix that  $M$  is an oriented manifold and  $L_1$  and  $L_2$  are two closed connected oriented submanifolds of  $M$  of complementary dimension, i.e.

$$\dim M = \dim L_1 + \dim L_2.$$

If  $L_1$  and  $L_2$  intersect transversally in the sense that for each intersection point  $p \in L_1 \cap L_2$  it holds that

$$T_p M = T_p L_1 + T_p L_2$$

then one defines the algebraic intersection number

$$L_1 \bullet L_2 \in \mathbb{Z}$$

as the signed count of intersection points of  $L_1$  and  $L_2$ . The sign is determined as follows. If  $p \in L_1 \cap L_2$  choose positively ordered basis  $[v_1, \dots, v_{l_1}]$  of  $T_p L_1$  and  $[w_1, \dots, w_{l_2}]$  of  $T_p L_2$ , where we abbreviate by  $l_1$  the dimension of  $L_1$  and by  $l_2$  the dimension of  $L_2$ . If  $[v_1, \dots, v_{l_1}, w_1, \dots, w_{l_2}]$  is a positively oriented basis of  $T_p M$ , we put  $\text{sign}(p) = 1$  and otherwise in the case where it is a negatively oriented basis we put  $\text{sign}(p) = -1$ . Then we define

$$(B.1) \quad L_1 \bullet L_2 = \sum_{p \in L_1 \cap L_2} \text{sign}(p).$$

The algebraic intersection number is a homotopy invariant. C.f. [Hir12, Chapter 5, Theorem 2.1]. Therefore one can define it as well for non-transverse intersections. Namely after small perturbation one can achieve that the two submanifolds intersect transversally so that one can define it for a transverse perturbation which by homotopy invariance does not depend on the choice of the perturbation.

Although generically two submanifolds intersect transversally in practise one quite often faces the situation where this is not the case. The reason for this is that we often consider the situation where a Lie group  $G$  acts on the manifold  $M$ . If the two submanifolds are invariant under the action of the Lie group, their

intersection  $L_1 \cap L_2$  is invariant as well and therefore intersection points are usually not isolated but come as whole  $G$ -orbits. For that reason one frequently has to deal with the case of *clean intersections*, in the sense that the intersection  $L_0 \cap L_1$  is itself a disjoint union of connected submanifolds (possibly of various dimensions) in  $M$  with the property that for each  $p \in L_0 \cap L_1$  one has

$$T_p(L_0 \cap L_1) = T_p L_1 \cap T_p L_2.$$

In this note we want to discuss a formula, how to compute the algebraic intersection number directly in the case of a clean intersection without perturbing. For that purpose we introduce the *clean normal bundle* (refer to section section B.2)

$$\nu_C^{\text{cl}} \rightarrow C$$

for each connected component  $C$  in  $L_0 \cap L_1$ . The rank of the vector bundle  $\nu_C^{\text{cl}}$  satisfies

$$(B.2) \quad \text{rank}(\nu_C^{\text{cl}}) = \dim C.$$

Moreover, the total space of the clean normal bundle has a canonical orientation induced from the orientations of  $L_1$ ,  $L_2$ , and  $M$ . The the total space of the clean normal bundle is canonically oriented as manifold, however, the clean normal bundle does not need to be orientable as vector bundle, since we do not assume that the intersection  $C$  is orientable.<sup>2</sup> We can in view of (B.2) define the Euler characteristic

$$\chi(\nu_C^{\text{cl}}) \in \mathbb{Z}$$

for every connected component  $C \subset L_0 \cap L_1$ . We do not need for that an orientation of  $C$ . See §B.3 for more detail. We derive a formula for algebraic intersection number in terms of this Euler characteristic

$$(B.3) \quad L_1 \bullet L_2 = \sum_{C \in \mathcal{C}(L_1 \cap L_2)} \chi(\nu_C^{\text{cl}}).$$

In particular, when  $L_1$  and  $L_2$  are two Lagrangian submanifolds intersecting cleanly in a symplectic manifold  $(M, \omega)$ , we obtain:

$$(B.4) \quad L_1 \bullet L_2 = \sum_{C \in \mathcal{C}(L_1 \cap L_2)} \text{sign}(C) \chi(C).$$

---

<sup>2</sup>A (real) vector bundle over a manifold  $E \xrightarrow{\pi} B$  is orientable if and only if  $w_1(E) \in H^1(B; \mathbb{Z}/2\mathbb{Z})$  vanishes. The total space  $E$  is oriented as a manifold means that its tangent bundle  $TE$  is an oriented vector bundle over the space  $E$ . The tangent bundle of  $E$  has non-canonical splitting

$$TE \cong \pi^*TB \oplus \pi^*E.$$

By the Whitney product formula:

$$\begin{aligned} w_1(TE) &= w_0(\pi^*TB) \smile w_1(\pi^*E) + w_1(\pi^*TB) \smile w_0(\pi^*E) \\ &= \pi^*w_1(E) + \pi^*w_1(TB). \end{aligned}$$

If  $B$  is orientable, i.e.  $w_1(TB) = 0$ , then  $w_1(TE) = \pi^*w_1(E)$ , as  $TE$  is an oriented vector bundle, we have  $w_1(TE) = 0 = w_1(E)$ , hence  $E$  is an oriented vector bundle over  $B$ . If  $B$  is non-orientable, then we have  $0 = w_1(E) + w_1(TB)$ , this implies that  $w_1(E) \equiv w_1(TB) \pmod{2}$ , thus  $E$  is a non-orientable vector bundle over  $B$ .

**B.2. The clean normal bundle.** We assume that  $L_1$  and  $L_2$  intersect cleanly, i.e.  $L_1 \cap L_2$  is a disjoint union of connected submanifolds. We denote by  $T_C L_1$  and  $T_C L_2$  the restriction vector bundle of  $TL_1$  and  $TL_2$  to  $C$ , i.e.

$$T_C L_1 := TL_1|_C = \bigsqcup_{p \in C} T_p L_1,$$

and

$$T_C L_2 = TL_2|_C = \bigsqcup_{p \in C} T_p L_2;$$

one can also view them as pullback bundles of  $TL_1$  and  $TL_2$  along the inclusions  $C \hookrightarrow L_1$  and  $C \hookrightarrow L_2$ . Over  $C \in \mathcal{C}(L_1 \cap L_2)$  we have the following three vector bundles

$$\begin{array}{ccccc} T_C L_1 & \longleftarrow & TC & \longrightarrow & T_C L_2 \\ & \searrow & \downarrow & \swarrow & \\ & & C & & \end{array}$$

We perform the fiber sum (distinguishing from Whitney sum) of the vector bundles  $T_C L_1$  and  $T_C L_2$  over  $C$  to obtain a new vector bundle

$$\begin{array}{c} T_C L_1 + T_C L_2 \\ \downarrow \\ C \end{array}$$

whose fiber over a point  $p \in C$  is given by the sum of vector spaces

$$T_p L_1 + T_p L_2 \subset T_p M.$$

In particular, the vector bundle  $T_C L_1 + T_C L_2$  is a subbundle of the vector bundle  $T_C M := TM|_C$ . We define the *clean normal bundle* as the quotient bundle:

$$(B.5) \quad \nu_C^{\text{cl}} := \frac{T_C M}{T_C L_1 + T_C L_2}.$$

It is clear that  $\nu_C^{\text{cl}} \subseteq \nu_C$  where  $\nu_C = T_C M/TC$  is the ordinary normal bundle of  $C$  in  $M$ . Abbreviating the dimensions as

$$n = \dim C, \quad m = \dim M, \quad l_1 = \dim L_1, \quad l_2 = \dim L_2$$

we have

$$\text{rank}(T_C L_1 + T_C L_2) = l_1 + l_2 - n$$

and therefore using that  $L_1$  and  $L_2$  are of complementary dimension in  $M$ , so that  $l_1 + l_2 = m$ , we obtain for the rank of the clean normal bundle over  $C$ :

$$\begin{aligned} \text{rank}(\nu_C^{\text{cl}}) &= \text{rank}(T_C M) - \text{rank}(T_C L_1 + T_C L_2) \\ &= m - (l_1 + l_2 - n) \\ &= n. \end{aligned}$$

In order to avoid the necessity to discuss trivial cases, we have assumed that

$$l_1 > 0, \quad l_2 > 0.$$



*Orientation.* We next explain how the orientations of  $L_1$ ,  $L_2$  and  $M$  induce an orientation on the total space of  $\nu_C^{\text{cl}}$ . For that purpose we assume that  $p \in C$ . We consider several cases:

**Case 1:** Assume that  $0 < n < \min\{l_1, l_2\}$ .

In other words,  $n$  is positive and smaller than  $l_1$  as well as  $l_2$ . Choose an ordered basis of  $T_p C$ :

$$\mathfrak{B}(T_p C) = [v_1, \dots, v_n].$$

Extend this basis to a positively oriented basis of  $T_p L_1$ :

$$\mathfrak{B}_p(T_p L_1) = [v_1, \dots, v_n, w_1, \dots, w_{l_1-n}],$$

and to a positively oriented basis of  $T_p L_2$ :

$$\mathfrak{B}(T_p L_2) = [u_1, \dots, u_{l_2-n}, v_1, \dots, v_n].$$

In contrast to  $T_p L_1$ , where the additional basis vectors are added after the basis vectors of  $T_p C$ , the basis vectors for  $T_p L_2$  are added before the ones of  $T_p C$ . Now extend the union of these two bases to a positively oriented basis of  $T_p M$ :

$$\mathfrak{B}(T_p M) = [u_1, \dots, u_{l_2-n}, v_1, \dots, v_n, w_1, \dots, w_{l_1-n}, \xi_1, \dots, \xi_n].$$

Abbreviate by

$$\pi : T_p M \rightarrow \nu_C^{\text{cl}}|_p$$

the projection map of  $T_p M$  onto the space  $\nu_C^{\text{cl}}|_p$ . Then

$$\pi : (u_1, \dots, u_{l_2-n}, v_1, \dots, v_n, w_1, \dots, w_{l_1-n}, \xi_1, \dots, \xi_n) \mapsto (\xi_1, \dots, \xi_n)$$

and

$$\mathfrak{B}(T_{(p,0)}\nu_C^{\text{cl}}) = [v_1, \dots, v_n, \xi_1, \dots, \xi_n]$$

is an ordered basis of  $T_{(p,0)}\nu_C^{\text{cl}} \cong T_p C \oplus \nu_C^{\text{cl}}|_p$ . We define an orientation on the total space of  $\nu_C^{\text{cl}}$  by declaring this basis to be positively oriented. This does not depend on the choices involved. Indeed, let us change the orientation of  $T_p C$  by replacing  $v_i$  by  $-v_i$  for any  $i \in \{1, \dots, n\}$ . In order to meet the requirement that  $\mathfrak{B}(T_p L_1)$  and  $\mathfrak{B}(T_p L_2)$  are ordered bases with respect to the given orientations on  $T_p L_1$  respectively  $T_p L_2$  we replace  $w_i$  by  $-w_i$  respectively  $u_i$  by  $-u_i$ , so that two of the basis vectors change sign. In order that  $\mathfrak{B}(T_p M)$  is still an ordered basis of  $T_p M$  we replace in additional  $\xi_i$  by  $-\xi_i$  so that four of the basis vectors change sign. Now in the basis  $\mathfrak{B}(T_{(p,0)}\nu_C^{\text{cl}})$ , the vectors  $v_i$  and  $\xi_i$  change sign, so that the orientation of  $T_{(p,0)}\nu_C^{\text{cl}}$  does not change.

**Case 2:** Assume that  $n = l_1 < l_2$ .

In this case the orientation of  $L_1$  induces an orientation of  $C$ , so that we start with a positively oriented basis of  $T_p L_1 = T_p C$ . We extend this basis to a positively oriented basis of  $T_p L_2$  as in Case 1 by adding the additional basis vector before the ones of  $T_p L_1 = T_p C$ . We extend this basis further to a positively oriented basis of  $T_p M$ . As in Case 1 this induces an orientation on  $T_{(p,0)}\nu_C^{\text{cl}}$ .

**Case 3:** Assume that  $n = l_2 < l_1$ .

This case is analogous to Case 2 just with the roles of  $L_1$  and  $L_2$  interchanged.

**Case 4:** Assume that  $n = l_1 = l_2$ .

In this case  $\nu_C^{\text{cl}}$  coincides with the usual normal bundle  $\nu_C$  of  $C$  in  $M$ . An orientation of  $C$  together with the orientation of  $M$  induces an orientation on the normal bundle which together with the orientation on  $C$  gives an orientation on the total space of the normal bundle. This orientation of the total space is independent of the choice of the orientation of  $C$  since flipping the orientation of  $C$  also flips the orientation of the normal bundle so that the orientation on the total space does not change. If the orientation of  $L_1$  and  $L_2$  along  $C$  agree, endow the total space of the normal bundle with its canonical orientation. Otherwise in the case where the orientations of  $L_1$  and  $L_2$  along  $C$  are opposite, endow the total space with the orientation which is opposite to the canonical one.

**Case 5:** Assume that  $n = 0$ .

In this case  $C = \{p\}$  is a single point at which the manifolds  $L_0$  and  $L_1$  intersect transversely. The clean normal bundle  $\nu_C^{\text{cl}}$  as well just consists of the point  $\{(p, 0)\}$ . An orientation of a point is just the assignment of a sign to the point. We endow it with the sign  $\text{sign}(p)$  as discussed before (B.1). This finishes the discussion of the orientation of the total space of  $\nu_{\{p\}}^{\text{cl}}$ .

Suppose now that  $(M, \omega)$  is a symplectic manifold and  $L_1$  and  $L_2$  are oriented Lagrangian submanifolds of  $M$  intersecting cleanly. Because the symplectic form is nondegenerate, the normal bundle of any submanifold can be identified with the symplectic orthogonal complement. Moreover, since  $L_1$  and  $L_2$  are Lagrangian, we have in particular

$$TL_1 = TL_1^{\perp\omega}, \quad TL_2 = TL_2^{\perp\omega}.$$

**Lemma B.6.** *Let  $L_1$  and  $L_2$  be Lagrangian submanifolds intersecting cleanly. There is a vector bundle isomorphism:*

$$\nu_C^{\text{cl}} \cong TC$$

for each  $C \in \mathcal{C}(L_1 \cap L_2)$ .

*Proof.* We first show that there is vector bundle isomorphism

$$\nu_C^{\text{cl}} \cong (TC L_1 + TC L_2)^{\perp\omega}.$$

Note that in general at each  $p \in C$ :

$$(T_p L_1 + T_p L_2) \cap (T_p L_1 + T_p L_2)^{\perp\omega} \neq \{0\}.$$

Recall for a symplectic vector space  $(V, \omega)$  and subspace  $W \subset V$ , we have

$$\dim V = \dim W + \dim W^{\perp\omega}.$$

In particular,

$$V = W \oplus W^{\perp\omega},$$

i.e.

$$W \cap W^{\perp\omega} = \{0\}$$

holds if and only if  $W$  is symplectic subspace. Note that this is not a necessary condition for above dimension formula. However,  $T_p L_1 + T_p L_2 \subset T_p M$  is clearly not symplectic subspace in general. So the way we show this is by making use of tangent-cotangent isomorphisms via  $\omega$ :

$$T_p M \cong T_p^* M, \quad T_p L_1 \cong T_p^* L_1, \quad T_p L_2 \cong T_p^* L_2.$$

Thus, we have isomorphism

$$\nu_C^{\text{cl}} \cong \frac{T_C^* M}{T_C^* L_1 + T_C^* L_2}.$$

At each  $p \in C$ :

$$\begin{aligned} T_p^* M &= \{\omega(X, \cdot) : X \in T_p M\}, \\ T_p^* L_1 + T_p^* L_2 &= \{\omega(X, \cdot) : X \in T_p L_1 + T_p L_2\}. \end{aligned}$$

Therefore

$$\frac{T_p^* M}{T_p^* L_1 + T_p^* L_2} = \{\omega(X, \cdot) : X \in T_p M, \omega(Y, \cdot) = 0 \forall Y \in T_p L_1 + T_p L_2\},$$

and applying cotangent-tangent isomorphism via  $\omega^{-1}$ ,

$$\begin{aligned} \frac{T_p^* M}{T_p^* L_1 + T_p^* L_2} &\cong \{X \in T_p M : \omega(X, Y) = 0 \forall Y \in T_p L_1 + T_p L_2\} \\ &= (T_p L_1 + T_p L_2)^{\perp \omega}. \end{aligned}$$

Hence, we obtain the desired isomorphism at the beginning, and

$$\begin{aligned} \nu_C^{\text{cl}} &\cong (T_C L_1 + T_C L_2)^{\perp \omega} \\ &= (T_C L_1)^{\perp \omega} \cap (T_C L_2)^{\perp \omega} \\ &= T_C L_1 \cap T_C L_2 \\ &= T_C. \end{aligned}$$

This completes the proof.  $\square$

**Remark B.7.** Alternatively, Lemma B.6 can be also proved by using compatible almost complex structure. By definition, at each  $p \in C$ , we have splitting:

$$T_p M = (T_p L_1 + T_p L_2) \oplus \nu_C^{\text{cl}}|_p.$$

On the other hand, let  $J$  be a  $\omega$ -compatible almost complex structure, we have

$$T_p L_1 \cap J T_p L_1 = \{0\}, \quad T_p L_2 \cap J T_p L_2 = \{0\}.$$

This yields that

$$(T_p L_1 + T_p L_2) \cap J T_p C = \{0\}.$$

Thus, by dimension reason, we have

$$T_p M = (T_p L_1 + T_p L_2) \oplus J T_p C.$$

Therefore, we have the identification of vector bundles:

$$\nu_C^{\text{cl}} = J T C.$$

We conclude that there is an isomorphism of vector bundles

$$\nu_C^{\text{cl}} \cong T C$$

given by the almost complex structure  $J$ .

**B.3. Euler characteristic and algebraic intersection number.**

*Euler characteristic of vector bundles.* Let  $E \rightarrow B$  be a real vector bundle (not necessarily orientable) over a closed connected manifold  $B$  satisfying

$$\text{rank}(E) = \dim B =: n.$$

Assume in addition that the total space  $E$  is oriented as manifold. We then define the Euler characteristic  $\chi(E) \in \mathbb{Z}$  as follows: If  $n = 0$ , the total space  $E$  is just a single point and an orientation is the assignment of a sign to the point. In this case the Euler characteristic is just this sign. Suppose now that  $n > 0$ . Choose a smooth section

$$\sigma : B \rightarrow E$$

transverse to the zero section, i.e.  $\sigma \pitchfork B$ . This is in fact a generic property for  $\sigma \in \Gamma(E)$  and the zero locus  $\sigma^{-1}(0)$  is a discrete finite set given  $B$  compact. If  $p \in B$  is a zero of  $\sigma$ , define the sign of  $p$ ,  $\text{sign}(p)$ , as follows. Choose an ordered basis of  $T_p B$

$$\mathfrak{B}(T_p B) = [v_1, \dots, v_n].$$

Since  $s$  is transverse to the zero section, the ordered set

$$\mathfrak{B}(T_p E) = [v_1, \dots, v_n, D\sigma_p(v_1), \dots, D\sigma_p(v_n)]$$

is a basis of  $T_p E$ . If this ordered basis is positively oriented with respect to the given orientation on  $E$  we define  $\text{sign}(p) = +1$  and otherwise  $\text{sign}(p) = -1$ . This definition does not depend on the choice of the basis  $\mathfrak{B}(T_p B)$ . Indeed, pick any  $i \in \{1, \dots, n\}$ , if  $v_i$  changes sign, then as well  $D\sigma_p(v_i)$  changes sign, so that we have two sign changes in  $\mathfrak{B}(E_p)$ . In particular, we need only the orientation of the total space  $E$  to define its Euler characteristic but no orientation of the base  $B$ . The Euler characteristic of  $E$  is given by the formula

$$(B.8) \quad \chi(E) := \sum_{p \in \sigma^{-1}(0)} \text{sign}(p) = B \bullet S$$

where  $S := \text{im}(\sigma)$ . Due to homotopy invariance of algebraic intersection number, this quantity is a well-defined homotopy invariant and independent of choice of transversal section. This is a mild generalization of [Hir12, Chapter 5, §2]. The rank  $n$  vector bundle over a smooth  $n$ -manifold is only required to have oriented total space instead of being oriented vector bundle. In particular, when  $E \rightarrow B$  is an oriented (real) vector bundle, we have the expression in terms of characteristic class:

$$\chi(E) = \langle e(E), [B] \rangle$$

where  $e(E) \in H^n(B; \mathbb{Z})$  is the Euler class of  $E$  and  $[B] \in H_n(B; \mathbb{Z})$  is the fundamental class of  $B$  given it is compact oriented. C.f. [Bre93, Chapter VI, §12].

*Algebraic intersection number.* We are now in position to write down a formula for the algebraic intersection number for a clean intersection. Abbreviate by  $\mathcal{C}(L_0 \cap L_1)$  the set of connected components of  $L_0 \cap L_1$ . Then the formula is

$$(B.3) \quad L_1 \bullet L_2 = \sum_{C \in \mathcal{C}(L_1 \cap L_2)} \chi(\nu_C^{\text{cl}}).$$

Using a transverse section of the clean normal bundle, one can perturb  $L_2$  locally around  $C$  to make the intersection transverse so that the signed count of intersection points of  $L_1$  with the perturbed  $L_2$  corresponds to the signed count of zeros of this transverse section which defines the Euler characteristic of clean normal bundle,

referring to the discussion of orientation of the clean normal bundle determined by that of  $L_1, L_2, M$  in §B.2. Therefore, one sees that the right hand side of the formula above indeed computes the algebraic intersection number.

**Example B.9.** We first confirm that this formula coincides with (B.1) in the special case of a transversal intersection. For a transversal intersection we have

$$\mathcal{C}(L_1 \cap L_2) = \{\{p\} : p \in L_1 \cap L_2\}$$

and for every  $p \in L_1 \cap L_2$  it holds that

$$\nu_{\{p\}}^{\text{cl}} = \{p\} \times \{0\}$$

So that the clean normal bundle is just a signed point for which one gets

$$\chi(\nu_{\{p\}}^{\text{cl}}) = \text{sign}(p).$$

**Example B.10.** Another well-known case of formula (B.3) is the case  $L_1 = L_2 = L$  giving rise to the algebraic self-intersection number. In this case the clean normal bundle is just the usual normal bundle endowed with its canonical orientation if the orientations of  $L_1$  and  $L_2$  agree and otherwise endowed with the opposite orientation. In this case (B.3) simplifies to

$$L_1 \bullet L_2 = L \bullet L = \chi(\nu_L^{\text{cl}}) = \chi(\nu_L).$$

In particular, when  $L_1 = L_2 = \Delta_M \subset M \times M$ , we have

$$\chi(TM) = \Delta_M \bullet \Delta_M = \chi(\nu_{\Delta_M}^{\text{cl}}) = \chi(\nu_{\Delta_M}).$$

In this case, one can see that actually

$$\nu_{\Delta_M}^{\text{cl}} = \nu_{\Delta_M} = \bigsqcup_{(p,p) \in M \times M} \frac{T_p M \times T_p M}{T_{(p,p)} \Delta_M} \cong \bigsqcup_{p \in M} T_p M = TM$$

by referring to section B.2 for the definition of clean normal bundle in this case.

We finally discuss the case of Lagrangian intersections. We assume that  $(M, \omega)$  is a symplectic manifold and  $L_1$  and  $L_2$  are oriented Lagrangian submanifolds of  $M$  intersecting cleanly. As a symplectic manifold  $M$  is endowed with a canonical orientation determined by the volume form  $\frac{1}{n!} \omega^n$ . In this case, the clean normal bundle coincides with the tangent bundle of  $C$ . This correspondence is not necessarily an isomorphism of oriented manifolds. Note that the tangent bundle of any manifold  $C$  carries a canonical orientation as manifold. If  $C$  is closed, the Euler characteristic of the canonically oriented tangent bundle coincides with the Euler characteristic of the manifold, i.e.  $\chi(TC) = \chi(C)$ .

Assume now all connected components of the intersection of  $L_1$  and  $L_2$  are closed. Let  $C \in \mathcal{C}(L_1 \cap L_2)$ , we associate to  $C$  a sign

$$\text{sign}(C) \in \{\pm 1\}$$

as follows. If the orientations of  $\nu_C^{\text{cl}}$  agrees with the canonical orientation of  $TC$  under the isomorphism in Lemma B.6, we put  $\text{sign}(C) = +1$ , and otherwise  $\text{sign}(C) = -1$ . Thus, we have

$$(B.11) \quad \chi(\nu_C^{\text{cl}}) = \text{sign}(C) \chi(TC).$$

With these conventions, (B.3) simplifies in the case of a clean Lagrangian intersection to

$$(B.4) \quad L_1 \bullet L_2 = \sum_{C \in \mathcal{C}(L_1 \cap L_2)} \text{sign}(C) \chi(C).$$

**B.4. Categorification.** We give a direct proof of the formula

$$(B.12) \quad \chi(HF_*(L_1, L_2, \{f_i\}; \Lambda)) = L_1 \bullet L_2$$

utilizing (B.4) whenever the Lagrangian Floer homology  $HF_*(L_1, L_2, \{f_i\}; \Lambda)$  is well-defined for the pair of oriented Lagrangian submanifolds  $L_1$  and  $L_2$  intersecting cleanly whose chain complex  $CF_*(L_1, L_2, \{f_i\}; \Lambda)$  is  $\mathbb{Z}$ -graded via Maslov index and finitely generated over  $\Lambda$  in each degree.

Suppose

$$L_1 \cap L_2 = \bigsqcup_{i \in I} C_i$$

where  $I$  is a finite set, and each  $C_i$  is connected oriented closed submanifold. The Floer complex in the Morse-Bott setting is generated by the critical points of auxiliary Morse functions  $f_i$  defined on each  $C_i$ :

$$CF_*(L_1, L_2, \{f_i\}; \Lambda) := \bigoplus_{k \in \mathbb{Z}} \bigoplus_{\substack{x \in \bigsqcup_{i \in I} \text{Crit}(f_i) \\ |x|=k}} \Lambda \langle x \rangle$$

where we recall the grading is given by (A.14) and (A.15). The Euler characteristic of the homology, due to finiteness assumption, is computed by

$$\begin{aligned} \chi(HF_*(L_1, L_2, \{f_i\}; \Lambda)) &:= \sum_{k \in \mathbb{Z}} (-1)^k \text{rank}_\Lambda HF_k(L_1, L_2, \{f_i\}; \Lambda) \\ &= \sum_{k \in \mathbb{Z}} (-1)^k \text{rank}_\Lambda CF_k(L_1, L_2, \{f_i\}; \Lambda) \\ &= \sum_{k \in \mathbb{Z}} \sum_{\substack{x \in \bigsqcup_{i \in I} \text{Crit}(f_i) \\ |x|=k}} (-1)^{|x|} \\ &= \sum_{i \in I} (-1)^{\mu(C_i)} \sum_{x \in \text{Crit}(f_i)} (-1)^{\text{ind}_{f_i}(x)} \end{aligned}$$

It is well-known that

$$(B.13) \quad \sum_{x \in \text{Crit}(f_i)} (-1)^{\text{ind}_{f_i}(x)} = \chi(C_i).$$

The parity of the Maslov-Morse index  $(-1)^{\mu(C_i)}$  is equal to the orientation sign of  $C_i$  up to a global sign, i.e.

$$(B.14) \quad (-1)^{\mu(C_i)} = \pm \text{sign}(C_i).$$

This can be seen as follows. According to [FOOO09, §2.2.2], when the Lagrangian submanifolds  $L_1$  and  $L_2$  are oriented, the definition of the Maslov-Morse index involves a choice of section of the oriented Lagrangian Grassmannian bundle fitting in the orientation of  $L_1$  and  $L_2$ , therefore the Maslov-Morse index at a intersection point in  $L_1 \cap L_2$  is the orientation sign of the intersection point up to a global sign depending on the convention, this is also discussed by Paul Seidel, see [Sei00, §2.d,

(vi)]. Further due to [FOOO09, §3.7 (3.7.49)], the parity of the Maslov-Morse index for every point in Morse-Bott component  $C_i$  is the same, thus we obtain the desired formula (B.14). The formulas (B.13) and (B.14) together imply that

$$\chi(HF_*(L_1, L_2, \{f_i\}; \Lambda)) = \pm \sum_{i \in I} \text{sign}(C_i) \chi(C_i)$$

which is (B.12) by (B.4).

## REFERENCES

- [AD23] Mohammed Abouzaid and Luís Diogo. *Monotone Lagrangians in cotangent bundles of spheres*. Adv. in Math. **427** (2023), 109114.
- [AS10] Mohamed Abouzaid and Paul Seidel. *An open string analogue of Viterbo functoriality*. Geom. Topol. **14**(2) (2010) 627–718.
- [Bre93] Glen E. Bredon. *Topology and Geometry*. Graduate Texts in Mathematics 139, Springer, 1993.
- [BD18] Theo Bühler and Dietmar A. Salamon. *Functional Analysis*. Graduate Studies in Mathematics, American Mathematical Society, 2018.
- [Dam12] Mihai Damian. *Floer homology on the universal cover, Audin’s conjecture and other constraints on Lagrangian submanifolds*. Comment. Math. Helv. **87**(2012), no. 2, 433–462.
- [Flo88I] Andreas Floer. *Morse theory for Lagrangian intersections*. J. Differential Geom., **28**(3):513–547, 1988.
- [Flo88II] Andreas Floer. *A relative Morse index for the symplectic action*. Comm. Pure Appl. Math., **41**(4):393–407, 1988.
- [Flo88III] Andreas Floer. *The unregularized gradient flow of the symplectic action*. Comm. Pure Appl. Math., **41**(6):775–813, 1988.
- [Fra03] Urs Frauenfelder. *Floer homology of symplectic quotients and the Arnold-Givental conjecture*. PhD thesis, ETH Zürich, 2003.
- [FOOO09] Kenji Fukaya, Yong-Geun Oh, Hiroshi Ohta, and Kaoru Ono. *Lagrangian Intersection Floer Theory – Anomaly and Obstruction*. Volume 46 of AMS/IP Studies in Advanced Mathematics. American Mathematical Society, Providence, RI, 2009.
- [GL21] Joshua Evan Greene and Andrew Lobb. *The rectangular peg problem*. Annals of Mathematics **194** (2021), 509–517.
- [GL23] Joshua Evan Greene and Andrew Lobb, *Cyclic quadrilaterals and smooth Jordan curves*, Invent. Math. **234** (2023), no. 3, 931–935.
- [GL24] Joshua Evan Greene and Andrew Lobb. *Floer homology and square pegs*. [arXiv:2404.05179](https://arxiv.org/abs/2404.05179), 2024.
- [Hir12] Morris W. Hirsch. *Differential Topology*. Graduate Texts in Mathematics, 2012, Springer New York.
- [Kar13] Cecilia Karlsson. *Area-preserving isotopies of self-transverse immersions of  $S^1$  in  $\mathbb{R}^2$* . Ark. Mat., **51** (2013), 85–97.
- [LM14] Yankı Lekili and Maksim Maydanskiy. *The symplectic topology of some rational homology balls*. Comment. Math. Helv. **89** (2014), 571–596.
- [Mat14] Benjamin Matschke. *A survey on the square peg problem*. Notices Amer. Math. Soc. **61** (2014), no. 4, 346–352.
- [Mat22] Benjamin Matschke. *On the square peg problem and its relatives*. Trans. Amer. Math. Soc. **375** (2022), 6255–6280.
- [MS12] Dusa McDuff and Dietmar Salamon. *J-holomorphic Curves and Symplectic Topology*. 2nd edition, American Mathematical Soc., 2012.
- [Nem09] S. Yu. Nemirovski. *Homology class of a Lagrangian Klein bottle*. Izv. Ross. Akad. Nauk Ser. Mat. **73** no. 4 (2009), 37–48.
- [Oh93] Yong-Geun Oh. *Floer cohomology of Lagrangian intersections and pseudoholomorphic disks I*. Comm. Pure Appl. Math. **46** (1993), 949–993.
- [Oh95] Yong-Geun Oh. *Addendum to: “Floer cohomology of Lagrangian intersections and pseudoholomorphic disks I”*. Comm. Pure Appl. Math. **48** (1995), no. 11, 1299–1302.
- [Oh96] Yong-Geun Oh. *Floer cohomology, spectral sequences, and the Maslov class of Lagrangian embeddings*. Int. Math. Res. Not. IMRN 1996, no. **7**, 305–346.

- [Oh15] Yong-Geun Oh. *Symplectic Topology and Floer Homology: Volume 2, Floer Homology and its Applications*. New Mathematical Monographs, 2015, Cambridge University Press.
- [Pol91] Leonid V. Polterovich. *The Maslov class of the Lagrange surfaces and Gromov's pseudo-holomorphic curves*. Trans. Am. Math. Soc. **325**(1), 241–248, 1991.
- [Po94] Marcin Poźniak. *Floer homology, Novikov rings and clean intersections*. Ph.D. Thesis, University of Warwick, 1994.
- [PR14] Leonid Polterovich, Daniel Rosen. *Function Theory on Symplectic Manifolds*. Volume 34 of CRM monograph series, Centre de Recherches Mathématiques, American Mathematical Soc., 2014.
- [Toe11] Otto Toeplitz. *Über einige Aufgaben der Analysis situs*. Verhandlungen der Schweizerischen Naturforschenden Gesellschaft in Solothurn, **4** (1911), no. 197.
- [Sch16] Felix Schmäschke. *Floer homology of Lagrangians in clean intersection*. [arXiv:1606.05327](https://arxiv.org/abs/1606.05327), 2016.
- [Sei00] Paul Seidel. *Graded Lagrangian submanifolds*. Bull. Soc. Math. France **128**, 103–146, 2000.
- [She09] V. V. Shevchishin. *Lagrangian embeddings of the Klein bottle and the combinatorial properties of mapping class groups*. Izv. Ross. Akad. Nauk Ser. Mat. **73** no. 4 (2009), 153–224.
- [St24] Bruno Staffa. *Bumpy metrics theorem for geodesic nets*. Journal of Topology and Analysis, 2024.
- [Vit87] Claude Viterbo. *Intersection de sous-variétés lagrangiennes, fonctionnelles d'action et indice des systèmes hamiltoniens*. Bull. Soc. Math. France **115** (1987), 361–390.
- [Vit90] Claude Viterbo. *A new obstruction to embedding Lagrangian tori*. Invent. Math. **100**(2), 301–320, 1990.

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT AUGSBURG, AUGSBURG, GERMANY  
Email address: [zhen.gao@math.uni-augsburg.de](mailto:zhen.gao@math.uni-augsburg.de)