

# The coproduct for the affine Yangian and the parabolic induction for non-rectangular $W$ -algebras

Mamoru Ueda\*

Department of Mathematical and Statistical Sciences, University of Alberta,  
11324 89 Ave NW, Edmonton, AB T6G 2J5, Canada

## Abstract

By using the coproduct and evaluation map for the affine Yangian and the Miura map for non-rectangular  $W$ -algebras, we construct a homomorphism from the affine Yangian associated with  $\widehat{\mathfrak{sl}}(n)$  to the universal enveloping algebra of a non-rectangular  $W$ -algebra of type  $A$ , which is an affine analogue of the one given in De Sole-Kac-Valeri [6]. As a consequence, we find that the coproduct for the affine Yangian is compatible with the parabolic induction for non-rectangular  $W$ -algebras via this homomorphism.

## 1 Introduction

The Yangian  $Y_{\hbar}(\mathfrak{g})$  associated with a finite dimensional simple Lie algebra  $\mathfrak{g}$  was introduced by Drinfeld ([7], [8]). The Yangian  $Y_{\hbar}(\mathfrak{g})$  is a quantum group which is a deformation of the current algebra  $\mathfrak{g} \otimes \mathbb{C}[z]$ . The affine Yangian associated with  $\widehat{\mathfrak{sl}}(n)$  was first introduced by Guay ([13], [14]). The affine Yangian  $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n))$  is a 2-parameter Yangian and is the deformation of the universal enveloping algebra of the central extension of  $\mathfrak{sl}(n)[u^{\pm 1}, v]$ . Guay-Nakajima-Wendlandt [15] gave the coproduct for the Yangian associated with the Kac-Moody Lie algebra of affine type. The coproduct for the affine Yangian  $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n))$  is a homomorphism satisfying the coassociativity:

$$\Delta: Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n)) \rightarrow Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n)) \widehat{\otimes} Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n)),$$

where  $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n)) \widehat{\otimes} Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n))$  is the standard degreewise completion of  $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n)) \otimes Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n))$ . In [24] and [26], we also constructed two kinds of a homomorphism:

$$\begin{aligned} \Psi_{n, m+n}: Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n)) &\rightarrow Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(m+n)), \\ \Psi^{m, m+n}: Y_{\hbar, \varepsilon+n\hbar}(\widehat{\mathfrak{sl}}(m)) &\rightarrow Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(m+n)). \end{aligned}$$

The images of  $\Psi_{n, m+n}$  and  $\Psi^{m, m+n}$  are commutative with each other and the image of  $\Psi_{n, m+n} \otimes \Psi^{m, m+n}$  is the affine version of the Levi subalgebra of the finite Yangian of type  $A$  given in [2].

Recently, the relationships between affine Yangians and  $W$ -algebras have been studied. The  $W$ -algebra  $\mathcal{W}^k(\mathfrak{g}, f)$  is a vertex algebra associated with a finite dimensional reductive Lie algebra  $\mathfrak{g}$ , a nilpotent element  $f \in \mathfrak{g}$  and a complex number  $k$ . In [27], we constructed a surjective homomorphism from the affine Yangian to the universal enveloping algebra of a rectangular  $W$ -algebra of type  $A$ , which is a  $W$ -algebra associated with  $\mathfrak{gl}(ln)$  and a nilpotent element of type  $(l^n)$ . Kodera and the author [19] gave another proof of the construction of this homomorphism by using the coproduct and evaluation map for the affine Yangian. The evaluation map for the

---

\*mueda@ualberta.ca

affine Yangian is a non-trivial homomorphism from the affine Yangian  $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n))$  to the standard degreewise completion of the universal enveloping algebra of  $\widehat{\mathfrak{gl}}(n)$ :

$$\text{ev}^n: Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n)) \rightarrow \mathcal{U}(\widehat{\mathfrak{gl}}(n)),$$

where  $\mathcal{U}(\widehat{\mathfrak{gl}}(n))$  is the standard degreewise completion of the universal enveloping algebra of  $\widehat{\mathfrak{gl}}(n)$ .

In [28], we gave a homomorphism from the affine Yangian  $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n))$  to the universal enveloping algebra of a non-rectangular  $W$ -algebra of type  $A$ . However, the proof of [28] is so complicated that we cannot understand the meaning of the construction way of [28]. In this article, by the similar way to [19], we give the another proof to [28]. Let us fix a positive integer and its partition:

$$N = q_1 + q_2 + \cdots + q_l, \quad q_1 \leq q_2 \leq \cdots \leq q_v, \quad q_{v+1} \geq q_{v+2} \geq \cdots \geq q_l.$$

Similarly to [3], we correspond a nilpotent element of  $\mathfrak{gl}(N)$  to a diagram. We take a nilpotent element  $f \in \mathfrak{gl}(N)$  whose column heights of the diagram corresponding to  $f$  is  $(q_1, \dots, q_l)$ . By using the Miura map (Kac-Wakimoto[16]), we obtain an embedding, we obtain

$$\mathcal{W}^k(\mathfrak{gl}(N), f) \hookrightarrow \bigotimes_{1 \leq i \leq l} V^{\kappa_i}(\mathfrak{gl}(q_i)),$$

where  $V^{\kappa_i}(\mathfrak{gl}(q_i))$  is the universal affine vertex algebra associated with  $\mathfrak{gl}(q_i)$  and its inner product  $\kappa_i$ . The Miura map induces the injective map

$$\tilde{\mu}: \mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(N), f)) \hookrightarrow \widehat{\bigotimes}_{1 \leq i \leq l} U(\widehat{\mathfrak{gl}}(q_i)),$$

where  $\widehat{\bigotimes}_{1 \leq i \leq l} U(\widehat{\mathfrak{gl}}(q_i))$  is the standard degreewise completion of  $\bigotimes_{1 \leq i \leq l} U(\widehat{\mathfrak{gl}}(q_i))$ .

Let us set  $q_{Min}$  as the minimum number of  $\{q_i\}_{1 \leq i \leq l}$ . By using  $\Psi^{m, m+n}$ , we can construct a homomorphism

$$\Delta^l: Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(q_{Min})) \rightarrow \bigotimes_{1 \leq i \leq l} Y_{\hbar, \varepsilon-(q_i-q_{Min})\hbar}(\widehat{\mathfrak{sl}}(q_i)).$$

In the case that  $q_1 \geq q_2 \geq \cdots \geq q_l$ , we obtain

$$\Delta^l = \prod_{1 \leq i \leq l-1} ((\Psi^{q_{i+1}, q_i} \otimes \text{id}) \circ \Delta) \otimes \text{id}^{\otimes l-i-1}.$$

**Theorem 1.1.** *There exists a homomorphism*

$$\Phi: Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(q_{Min})) \rightarrow \mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(N), f))$$

uniquely determined by

$$\tilde{\mu} \circ \Phi = \bigotimes_{1 \leq i \leq l} \text{ev}^{q_i} \circ \Delta^l.$$

In [28], we only consider the case that  $q_1 \geq q_2 \geq \cdots \geq q_l$ . Theorem 1.1 can be applied to more general cases. One of the application of Theorem 1.1 is the compatibility with the coproduct for the affine Yangian and the parabolic induction for a  $W$ -algebra.

Let us consider two  $W$ -algebras associated with  $(\mathfrak{gl}(N_1), f_1)$  and  $(\mathfrak{gl}(N_2), f_2)$ , where

$$N_1 = q_1 + \cdots + q_w, \quad N_2 = q_{w+1} + \cdots + q_l$$

and nilpotent elements  $f_1$  and  $f_2$  are defined in the same way as  $f$ . In [12], Genra constructed a homomorphism called the parabolic induction:

$$\Delta_W: \mathcal{W}^k(\mathfrak{gl}(N), f) \rightarrow \mathcal{W}^{k+N_2}(\mathfrak{gl}(N_1), f_1) \otimes \mathcal{W}^{k+N_1}(\mathfrak{gl}(N_2), f_2).$$

Genra [12] conjectured that the parabolic induction for a  $W$ -algebra is compatible with the coproduct for the affine Yangian.

Theorem 1.1 gives a relationship between the coproduct for the affine Yangian and the parabolic induction for a non-rectangular  $W$ -algebra. By Theorem 1.1, we obtain two homomorphisms

$$\begin{aligned}\Phi_1 &: Y_{\hbar, \varepsilon + (\min(q_1, q_w) - q_{Min})\hbar}(\widehat{\mathfrak{sl}}(\min(q_1, q_w))) \rightarrow \mathcal{U}(\mathcal{W}^{k+N_2}(\mathfrak{gl}(N_1), f_1)), \\ \Phi_2 &: Y_{\hbar, \varepsilon + (\min(q_{w+1}, q_l) - q_{Min})\hbar}(\widehat{\mathfrak{sl}}(\min(q_{w+1}, q_l))) \rightarrow \mathcal{U}(\mathcal{W}^{k+N_1}(\mathfrak{gl}(N_2), f_2)).\end{aligned}$$

By the construction way in Theorem 1.1, we obtain the following relations for some  $A, b \in \mathbb{C}$ :

$$\begin{aligned}(\Phi_1 \circ \tau_a \circ \Psi^{q_l, \min(q_1, q_w)}) \otimes \Phi_2 \circ \Delta &= \Delta_W \circ \Phi \text{ if } q_1 \geq q_l, \\ (\Phi_1 \circ \tau_b) \otimes (\Phi_2 \circ \Psi^{q_1, \min(q_l, q_{w+1})}) \circ \Delta &= \Delta_W \circ \Phi \text{ if } q_1 < q_l.\end{aligned}$$

where  $\tau_x$  is a shift operator of the affine Yangian. This means that the parabolic induction  $\Delta_W$  is compatible with the coproduct  $\Delta$  via the homomorphism  $\Phi$ .

The motivation of this theorem is the generalization of the AGT conjecture. The AGT conjecture suggests that there exists a representation of the principal  $W$ -algebra of type  $A$  on the equivariant homology space of the moduli space of  $U(r)$ -instantons. Schiffmann and Vasserot [22] gave this representation by using an action of the Yangian associated with  $\widehat{\mathfrak{gl}}(1)$  on this equivariant homology space. It is conjectured by Crutzig-Diaconescu-Ma [4] that an action of an iterated  $W$ -algebra of type  $A$  on the equivariant homology space of the affine Laumon space will be given through an action of an affine shifted Yangian constructed in [9]. For the resolution of the Crutzig-Diaconescu-Ma's conjecture, we need to construct a homomorphism from the shifted affine Yangian to the universal enveloping algebra of a iterated  $W$ -algebra. However, the shifted affine Yangian is so complicated that we cannot directly construct this homomorphism.

In finite setting, Brundan-Kleshchev [3] gave a surjective homomorphism from a shifted Yangian, which is a subalgebra of the finite Yangian associated with  $\mathfrak{gl}(n)$ , to a finite  $W$ -algebra ([21]) of type  $A$  for its general nilpotent element. A finite  $W$ -algebra  $\mathcal{W}^{\text{fin}}(\mathfrak{g}, f)$  is an associative algebra associated with a reductive Lie algebra  $\mathfrak{g}$  and its nilpotent element  $f$  and is a finite analogue of a  $W$ -algebra  $\mathcal{W}^k(\mathfrak{g}, f)$  ([5] and [1]). In [6], De Sole, Kac and Valeri constructed a homomorphism from the finite Yangian of type  $A$  to the finite  $W$ -algebras of type  $A$  by using the Lax operator, which is a restriction of the homomorphism given by Brundan-Kleshchev in [3]. Actually, the homomorphism  $\Phi$  is the affine analogue of the De Sole-Kac-Valeri's homomorphism (see [25]). Thus, we expect that we can extend  $\Phi$  to the shifted affine Yangian and solve the Crutzig-Diaconescu-Ma's conjecture.

## 2 Affine Yangian

Let us recall the definition of the affine Yangian of type  $A$  (Definition 3.2 in [13] and Definition 2.3 in [14]). Hereafter, we sometimes identify  $\{0, 1, 2, \dots, n-1\}$  with  $\mathbb{Z}/n\mathbb{Z}$ . we also set  $\{X, Y\} = XY + YX$  and

$$a_{i,j} = \begin{cases} 2 & \text{if } i = j, \\ -1 & \text{if } j = i \pm 1, \\ 0 & \text{otherwise} \end{cases}$$

for  $i \in \mathbb{Z}/n\mathbb{Z}$ .

**Definition 2.1.** Suppose that  $n \geq 3$ . The affine Yangian  $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n))$  is the associative algebra generated by  $X_{i,r}^+, X_{i,r}^-, H_{i,r}$  ( $i \in \{0, 1, \dots, n-1\}$ ,  $r = 0, 1$ ) subject to the following defining relations:

$$[H_{i,r}, H_{j,s}] = 0, \tag{2.2}$$

$$[X_{i,0}^+, X_{j,0}^-] = \delta_{i,j} H_{i,0}, \tag{2.3}$$

$$[X_{i,1}^+, X_{j,0}^-] = \delta_{i,j} H_{i,1} = [X_{i,0}^+, X_{j,1}^-], \quad (2.4)$$

$$[H_{i,0}, X_{j,r}^\pm] = \pm a_{i,j} X_{j,r}^\pm, \quad (2.5)$$

$$[\tilde{H}_{i,1}, X_{j,0}^\pm] = \pm a_{i,j} (X_{j,1}^\pm), \text{ if } (i,j) \neq (0,n-1), (n-1,0), \quad (2.6)$$

$$[\tilde{H}_{0,1}, X_{n-1,0}^\pm] = \mp \left( X_{n-1,1}^\pm + (\varepsilon + \frac{n}{2}\hbar) X_{n-1,0}^\pm \right), \quad (2.7)$$

$$[\tilde{H}_{n-1,1}, X_{0,0}^\pm] = \mp \left( X_{0,1}^\pm - (\varepsilon + \frac{n}{2}\hbar) X_{0,0}^\pm \right), \quad (2.8)$$

$$[X_{i,1}^\pm, X_{j,0}^\pm] - [X_{i,0}^\pm, X_{j,1}^\pm] = \pm a_{ij} \frac{\hbar}{2} \{X_{i,0}^\pm, X_{j,0}^\pm\} \text{ if } (i,j) \neq (0,n-1), (n-1,0), \quad (2.9)$$

$$[X_{0,1}^\pm, X_{n-1,0}^\pm] - [X_{0,0}^\pm, X_{n-1,1}^\pm] = \mp \frac{\hbar}{2} \{X_{0,0}^\pm, X_{n-1,0}^\pm\} + (\varepsilon + \frac{n}{2}\hbar) [X_{0,0}^\pm, X_{n-1,0}^\pm], \quad (2.10)$$

$$(\text{ad } X_{i,0}^\pm)^{1+|a_{i,j}|}(X_{j,0}^\pm) = 0 \text{ if } i \neq j, \quad (2.11)$$

where we set  $\tilde{H}_{i,1} = H_{i,1} - \frac{\hbar}{2} H_{i,0}^2$ .

By (2.3), (2.4), (2.6) and (2.8), we find that the affine Yangian  $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$  is generated by  $X_{i,0}^\pm$  for  $0 \leq i \leq n-1$  and  $H_{j,1}$  for  $1 \leq j \leq n-1$ .

We set a Lie algebra

$$\widehat{\mathfrak{gl}}(n) = \mathfrak{gl}(n) \otimes \mathbb{C}[z^{\pm 1}] \oplus \mathbb{C}\tilde{c} \oplus \mathbb{C}z$$

whose commutator relations are given by

$$z \text{ and } \tilde{c} \text{ are central elements of } \widehat{\mathfrak{gl}}(n),$$

$$[E_{i,j} \otimes t^u, E_{p,q} \otimes t^v] = (\delta_{j,p} E_{i,q} - \delta_{i,q} E_{p,j}) \otimes t^{u+v} + \delta_{u+v,0} u (\delta_{i,q} \delta_{j,p} \tilde{c} + \delta_{i,j} \delta_{p,q} z).$$

Let us take a Lie subalgebra  $\widehat{\mathfrak{sl}}(n) = \mathfrak{sl}(n) \otimes \mathbb{C}[z^{\pm 1}] \oplus \mathbb{C}\tilde{c}$  and Chevalley generators of  $\widehat{\mathfrak{sl}}(n)$  as

$$h_i = \begin{cases} E_{i,i} - E_{i+1,i+1} & \text{if } i \neq 0, \\ E_{n,n} - E_{1,1} + \tilde{c} & \text{if } i = 0, \end{cases}$$

$$x_i^+ = \begin{cases} E_{i,i+1} & \text{if } i \neq 0, \\ E_{n,1}t & \text{if } i = 0, \end{cases} \quad x_i^- = \begin{cases} E_{i+1,i} & \text{if } i \neq 0, \\ E_{1,n}t^{-1} & \text{if } i = 0. \end{cases}$$

By the defining relations (2.2)-(2.11), we can give a homomorphism from the universal enveloping algebra of  $\widehat{\mathfrak{sl}}(n)$  to the affine Yangian  $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$  given by  $h_i \mapsto H_{i,0}$  and  $x_i^\pm \mapsto X_{i,0}^\pm$ . We denote the image of  $x \in Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$  via this homomorphism by  $x$ .

We take one completion of  $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$ . We set the degree of  $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$  by

$$\deg(H_{i,r}) = 0, \quad \deg(X_{i,r}^\pm) = \begin{cases} \pm 1 & \text{if } i = 0, \\ 0 & \text{if } i \neq 0. \end{cases}$$

This degree is compatible with the natural degree on the universal enveloping algebra of  $\widehat{\mathfrak{sl}}(n)$ . We denote the standard degreewise completion of  $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$  by  $\widetilde{Y}_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$ .

### 3 The coproduct for the affine Yangian

By using the minimalistic presentation of the Yangian, Guay-Nakajima-Wendlandt [15] gave a coproduct for the Yangian associated with a Kac-Moody Lie algebra of the affine type.

**Theorem 3.1** (Theorem 5.2 in [15]). *There exists an algebra homomorphism*

$$\Delta^\pm: Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n)) \rightarrow Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n)) \widehat{\otimes} Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$$

determined by

$$\begin{aligned}\Delta^\pm(X_{j,0}^\pm) &= X_{j,0}^\pm \otimes 1 + 1 \otimes X_{j,0}^\pm \text{ for } 0 \leq j \leq n-1, \\ \Delta^\pm(\tilde{H}_{i,1}) &= \tilde{H}_{i,1} \otimes 1 + 1 \otimes \tilde{H}_{i,1} + A_i^\pm \text{ for } 1 \leq i \leq n-1,\end{aligned}$$

where  $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))\widehat{\otimes} Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$  is the standard degreewise completion of  $\otimes^2 Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$  and

$$\begin{aligned}A_i^+ &= -\hbar(E_{i,i} \otimes E_{i+1,i+1} + E_{i+1,i+1} \otimes E_{i,i}) \\ &\quad + \hbar \sum_{s \geq 0} \sum_{u=1}^i (-E_{u,i} t^{-s-1} \otimes E_{i,u} t^{s+1} + E_{i,u} t^{-s} \otimes E_{u,i} t^s) \\ &\quad + \hbar \sum_{s \geq 0} \sum_{u=i+1}^n (-E_{u,i} t^{-s} \otimes E_{i,u} t^s + E_{i,u} t^{-s-1} \otimes E_{u,i} t^{s+1}) \\ &\quad - \hbar \sum_{s \geq 0} \sum_{u=1}^i (-E_{u,i+1} t^{-s-1} \otimes E_{i+1,u} t^{s+1} + E_{i+1,u} t^{-s} \otimes E_{u,i+1} t^s) \\ &\quad - \hbar \sum_{s \geq 0} \sum_{u=i+1}^n (-E_{u,i+1} t^{-s} \otimes E_{i+1,u} t^s + E_{i+1,u} t^{-s-1} \otimes E_{u,i+1} t^{s+1}), \\ A_i^- &= -\hbar(E_{i,i} \otimes E_{i+1,i+1} + E_{i+1,i+1} \otimes E_{i,i}) \\ &\quad + \hbar \sum_{s \geq 0} \sum_{u=1}^i (-E_{i,u} t^{s+1} \otimes E_{u,i} t^{-s-1} + E_{u,i} t^s \otimes E_{i,u} t^{-s}) \\ &\quad + \hbar \sum_{s \geq 0} \sum_{u=i+1}^n (-E_{i,u} t^s \otimes E_{u,i} t^{-s} + E_{u,i} t^{s+1} \otimes E_{i,u} t^{-s-1}) \\ &\quad - \hbar \sum_{s \geq 0} \sum_{u=1}^i (-E_{i+1,u} t^{s+1} \otimes E_{u,i+1} t^{-s-1} + E_{u,i+1} t^s \otimes E_{i+1,u} t^{-s}) \\ &\quad - \hbar \sum_{s \geq 0} \sum_{u=i+1}^n (-E_{i+1,u} t^s \otimes E_{u,i+1} t^{-s} + E_{u,i+1} t^{s+1} \otimes E_{i+1,u} t^{-s-1}).\end{aligned}$$

The homomorphism  $\Delta^\pm$  is said to be the coproduct for the affine Yangian since  $\Delta^\pm$  satisfies the coassociativity.

## 4 The evaluation map for the affine Yangian

The evaluation map for the affine Yangian is a non-trivial homomorphism from the affine Yangian  $Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n))$  to the completion of the universal enveloping algebra of the affinization of  $\mathfrak{gl}(n)$ . We take the grading of  $U(\widehat{\mathfrak{gl}}(n))/U(\widehat{\mathfrak{gl}}(n))(z-1)$  as  $\deg(Xt^s) = s$  and  $\deg(\tilde{c}) = 0$ . We denote the degreewise completion of  $U(\widehat{\mathfrak{gl}}(n))/U(\widehat{\mathfrak{gl}}(n))(z-1)$  by  $\mathcal{U}(\widehat{\mathfrak{gl}}(n))$ .

**Theorem 4.1** (Theorem 3.8 in [18] and Theorem 4.18 in [17]). *Suppose that  $\hbar \neq 0$  and  $\tilde{c} = \frac{\varepsilon}{\hbar}$ . For  $a \in \mathbb{C}$ , there exists an algebra homomorphism*

$$\text{ev}_{\hbar,\varepsilon}^{n,a}: Y_{\hbar,\varepsilon}(\widehat{\mathfrak{sl}}(n)) \rightarrow \mathcal{U}(\widehat{\mathfrak{gl}}(n))$$

uniquely determined by

$$\text{ev}_{\hbar,\varepsilon}^{n,a}(X_{i,0}^+) = \begin{cases} E_{n,1} t & \text{if } i = 0, \\ E_{i,i+1} & \text{if } 1 \leq i \leq n-1, \end{cases} \quad \text{ev}_{\hbar,\varepsilon}^{n,a}(X_{i,0}^-) = \begin{cases} E_{1,n} t^{-1} & \text{if } i = 0, \\ E_{i+1,i} & \text{if } 1 \leq i \leq n-1, \end{cases}$$

and

$$\begin{aligned} \text{ev}_{\hbar, \varepsilon}^{n,a}(H_{i,1}) &= (a - \frac{i}{2}\hbar) \text{ev}_{\hbar, \varepsilon}^n(H_{i,0}) - \hbar E_{i,i} E_{i+1,i+1} \\ &\quad + \hbar \sum_{s \geq 0} \sum_{k=1}^i E_{i,k} t^{-s} E_{k,i} t^s + \hbar \sum_{s \geq 0} \sum_{k=i+1}^n E_{i,k} t^{-s-1} E_{k,i} t^{s+1} \\ &\quad - \hbar \sum_{s \geq 0} \sum_{k=1}^i E_{i+1,k} t^{-s} E_{k,i+1} t^s - \hbar \sum_{s \geq 0} \sum_{k=i+1}^n E_{i+1,k} t^{-s-1} E_{k,i+1} t^{s+1} \end{aligned}$$

for  $i \neq 0$ .

*Remark 4.2.* The universal enveloping algebra of the universal affine vertex algebra associated with  $\mathfrak{gl}(n)$  coincides with the standard degreewise completion of the universal enveloping algebra of  $\widehat{\mathfrak{gl}}(n)$ . Also, the universal affine vertex algebra is a  $W$ -algebra associated with a nilpotent element 0. Thus, the evaluation map is the easiest example of a homomorphism from the affine Yangian to the universal enveloping algebra of a  $W$ -algebra.

## 5 A homomorphism from the affine Yangian $Y_{\hbar, \varepsilon+m\hbar}(\widehat{\mathfrak{sl}}(n))$ to the affine Yangian $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n+m))$

In [24], we gave a homomorphism from the affine Yangian  $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n))$  to the affine Yangian  $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n+1))$ . In [26], we gave another homomorphism from the affine Yangian  $Y_{\hbar, \varepsilon+\hbar}(\widehat{\mathfrak{sl}}(n))$  to the affine Yangian  $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n+1))$ . In this article, we only use the latter homomorphism.

**Theorem 5.1** (Theorem 3.1 in [26]). *For  $m \geq 3$  and  $n \geq 1$ , there exists a homomorphism*

$$\Psi^{m,m+n}: Y_{\hbar, \varepsilon+n\hbar}(\widehat{\mathfrak{sl}}(m)) \rightarrow \widetilde{Y}_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(m+n))$$

determined by

$$\Psi^{m,m+n}(X_{i,0}^+) = \begin{cases} E_{m+n,n+1}t & \text{if } i = 0, \\ E_{n+i,n+i+1} & \text{if } i \neq 0, \end{cases} \quad \Psi^{m,m+n}(X_{i,0}^-) = \begin{cases} E_{n+1,m+n}t^{-1} & \text{if } i = 0, \\ E_{n+i+1,n+i} & \text{if } i \neq 0, \end{cases}$$

and

$$\begin{aligned} \Psi^{m,m+n}(H_{i,1}) &= H_{i+n,1} + \hbar \sum_{s \geq 0} \sum_{k=1}^n E_{k,n+i} t^{-s-1} E_{n+i,k} t^{s+1} \\ &\quad - \hbar \sum_{s \geq 0} \sum_{k=1}^n E_{k,n+i+1} t^{-s-1} E_{n+i+1,k} t^{s+1} \end{aligned}$$

for  $i \neq 0$ .

## 6 $W$ -algebras of type $A$

We fix some notations for vertex algebras. For a vertex algebra  $V$ , we denote the generating field associated with  $v \in V$  by  $v(z) = \sum_{n \in \mathbb{Z}} v_{(n)} z^{-n-1}$ . We also denote the OPE of  $V$  by

$$u(z)v(w) \sim \sum_{s \geq 0} \frac{(u_{(s)}v)(w)}{(z-w)^{s+1}}$$

for all  $u, v \in V$ . We denote the vacuum vector (resp. the translation operator) by  $|0\rangle$  (resp.  $\partial$ ). We denote the universal affine vertex algebra associated with a finite dimensional Lie algebra  $\mathfrak{g}$  and its inner product  $\kappa$  by  $V^\kappa(\mathfrak{g})$ . By the PBW theorem, we can identify  $V^\kappa(\mathfrak{g})$  with  $U(t^{-1}\mathfrak{g}[t^{-1}])$ . In order to simplify the notation, here after, we denote the generating field  $(ut^{-1})(z)$  as  $u(z)$ . By the definition of  $V^\kappa(\mathfrak{g})$ , the generating fields  $u(z)$  and  $v(z)$  satisfy the OPE

$$u(z)v(w) \sim \frac{[u, v](w)}{z - w} + \frac{\kappa(u, v)}{(z - w)^2} \quad (6.1)$$

for all  $u, v \in \mathfrak{g}$ .

We take a positive integer and its partition:

$$N = \sum_{i=1}^l q_i, \quad q_1 \leq q_2 \leq \cdots \leq q_v, \quad q_{v+1} \geq q_{v+2} \geq \cdots \geq q_l.$$

We also set

$$q_{Max} = \max(q_v, q_{v+1}), \quad q_{Min} = \min(q_1, q_l).$$

We also fix an inner product of  $\mathfrak{gl}(N)$  determined by

$$(E_{i,j}|E_{p,q}) = k\delta_{i,q}\delta_{p,j} + \delta_{i,j}\delta_{p,q}.$$

For  $1 \leq i \leq N$ , we set  $1 \leq \text{col}(i) \leq l$  and  $q_{Max} - q_{\text{col}(i)} < \text{row}(i) \leq q_{Max}$  satisfying

$$\text{col}(i) = s \text{ if } \sum_{j=1}^{s-1} q_j < i \leq \sum_{j=1}^s q_j, \quad \text{row}(i) = i - \sum_{j=1}^{\text{col}(i)-1} q_j + q_{Max} - q_{\text{col}(i)}.$$

We take a nilpotent element  $f$  as

$$f = \sum_{1 \leq j \leq N} e_{\hat{j},j},$$

where the integer  $1 \leq \hat{j} \leq N$  is determined by

$$\text{col}(\hat{j}) = \text{col}(j) + 1, \quad \text{row}(\hat{j}) = \text{row}(j).$$

The  $W$ -algebra is a vertex algebra associated with the reductive Lie algebra  $\mathfrak{g}$  and a nilpotent element  $f$ . We denote the  $W$ -algebra associated with  $\mathfrak{gl}(N)$  and  $f$  by  $\mathcal{W}^k(\mathfrak{gl}(N), f)$ .

We set the inner product on  $\mathfrak{gl}(q_s)$  by

$$\kappa_s(E_{i,j}, E_{p,q}) = \delta_{j,p}\delta_{i,q}\alpha_s + \delta_{i,j}\delta_{p,q},$$

where  $\alpha_s = k + N - q_s$ . Then, by Corollary 5.2 in [11], the  $W$ -algebra  $\mathcal{W}^k(\mathfrak{gl}(N), f)$  can be embedded into  $\bigotimes_{1 \leq s \leq l} V^{\kappa_s}(\mathfrak{gl}(q_s))$ . This embedding is called the Miura map. Hereafter, we denote  $E_{i,j}t^{-u} \in U(t^{-1}\mathfrak{gl}(q_s)[t^{-1}]) = V^\kappa(\mathfrak{gl}(q_s))$  by  $e_{q_{Max}-q_s+i, q_{Max}-q_s+j}[-u]$ .

We can prove the following theorem in the same way as Theorem 4.6 in [28].

**Theorem 6.2.** *We set  $\gamma_a = \sum_{u=a+1}^l \alpha_u$ . For positive integers  $p, q > q_{Max} - q_{Min}$ , the following elements of  $\bigotimes_{1 \leq s \leq l} V^{\kappa_s}(\mathfrak{gl}(q_s))$  are contained in  $\mathcal{W}^k(\mathfrak{gl}(N), f)$ .*

$$\begin{aligned} W_{p,q}^{(1)} &= \sum_{1 \leq r \leq l} e_{p,q}^{(r)}[-1], \\ W_{p,q}^{(2)} &= - \sum_{1 \leq r \leq l} \gamma_r e_{p,q}^{(r)}[-2] + \sum_{\substack{r_1 < r_2 \\ u > q_{Max} - q_{Min}}} e_{u,q}^{(r_1)}[-1]e_{p,u}^{(r_2)}[-1] \\ &\quad - \sum_{\substack{r_1 \geq r_2 \\ q_{Max} - \min(q_{r_1}, q_{r_2}) < u \leq q_{Max} - q_{Min}}} e_{u,q}^{(r_1)}[-1]e_{p,u}^{(r_2)}[-1], \end{aligned}$$

where  $x^{(s)} = 1^{\otimes s-1} \otimes x \otimes 1^{l-s}$ .

Let us recall the definition of a universal enveloping algebra of a vertex algebra in the sense of [10] and [20]. For any vertex algebra  $V$ , let  $L(V)$  be the Borcherds Lie algebra, that is,

$$L(V) = V \otimes \mathbb{C}[t, t^{-1}] / \text{Im}(\partial \otimes \text{id} + \text{id} \otimes \frac{d}{dt}), \quad (6.3)$$

where the commutation relation is given by

$$[ut^a, vt^b] = \sum_{r \geq 0} \binom{a}{r} (u_{(r)} v) t^{a+b-r}$$

for all  $u, v \in V$  and  $a, b \in \mathbb{Z}$ .

**Definition 6.4** (Section 6 in [20]). We set  $\mathcal{U}(V)$  as the quotient algebra of the standard degreewise completion of the universal enveloping algebra of  $L(V)$  by the completion of the two-sided ideal generated by

$$(u_{(a)} v) t^b - \sum_{i \geq 0} \binom{a}{i} (-1)^i (ut^{a-i} vt^{b+i} - (-1)^a vt^{a+b-i} ut^i), \quad (6.5)$$

$$|0\rangle t^{-1} - 1. \quad (6.6)$$

We call  $\mathcal{U}(V)$  the universal enveloping algebra of  $V$ .

Induced by the Miura map  $\mu$ , we obtain the embedding

$$\tilde{\mu}: \mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(N), f)) \rightarrow \widehat{\bigoplus}_{1 \leq s \leq l} U(\widehat{\mathfrak{gl}}(q_s)),$$

where  $\widehat{\bigoplus}_{1 \leq s \leq l} U(\widehat{\mathfrak{gl}}(q_s))$  is the standard degreewise completion of  $\bigotimes_{1 \leq s \leq l} U(\widehat{\mathfrak{gl}}(q_s))$ .

By the definition of  $W_{i,j}^{(1)}$  and  $W_{i,j}^{(2)}$ , we have

$$\tilde{\mu}(W_{i,j}^{(1)} t^s) = \sum_{1 \leq r \leq l} e_{i,j}^{(r)} t^s, \quad (6.7)$$

$$\begin{aligned} \tilde{\mu}(W_{i,j}^{(2)} t) &= \sum_{r=1}^n \gamma_r e_{i,j}^{(r)} + \sum_{s \in \mathbb{Z}} \sum_{r_1 < r_2} \sum_{u > q_{Max} - q_{Min}} e_{u,j}^{(r_1)} t^{-s} e_{i,u}^{(r_2)} t^s \\ &\quad - \sum_{s \geq 0} \sum_{r \geq 0} \sum_{q_{Max} - q_r \leq u \leq q_{Max} - q_{Min}} (e_{u,j}^{(r)} t^{-s-1} e_{i,u}^{(r)} t^{s+1} + e_{i,u}^{(r)} t^{-s} e_{u,j}^{(r)} t^s) \\ &\quad - \sum_{s \in \mathbb{Z}} \sum_{r_1 < r_2} \sum_{q_{Max} - \min(q_{r_1}, q_{r_2}) < u \leq q_{Max} - q_{Min}} e_{i,u}^{(r_1)} t^{-s} e_{u,j}^{(r_2)} t^s. \end{aligned} \quad (6.8)$$

## 7 A parabolic induction for a $W$ -algebra of type $A$

Let us recall the parabolic induction for  $W$ -algebras of type  $A$  (see Theorem 6.1 in [12]). Let us take two positive integers

$$N_1 = q_1 + \cdots + q_w, \quad N_2 = q_{w+1} + \cdots + q_l.$$

and  $f_1 \in \mathfrak{gl}(N_1)$  and  $f_2 \in \mathfrak{gl}(N_2)$  are nilpotent elements defined by the same way as  $f$ . Let us denote the Miura map associated with  $\mathcal{W}^{k+N_2}(\mathfrak{gl}(N_1), f_1)$  and  $\mathcal{W}^{k+N_1}(\mathfrak{gl}(N_2), f_2)$  by  $\mu_1$  and  $\mu_2$ .

**Theorem 7.1** (Theorem 6.1 in [12]). *There exists a homomorphism*

$$\Delta_W: \mathcal{W}^k(\mathfrak{gl}(N), f) \rightarrow \mathcal{W}^{k+N_2}(\mathfrak{gl}(N_1), f_1) \otimes \mathcal{W}^{k+N_1}(\mathfrak{gl}(N_2), f_2) \quad (7.2)$$

satisfying that

$$\mu = (\mu_1 \otimes \mu_2) \circ \Delta_W. \quad (7.3)$$

We call this homomorphism the parabolic induction for a  $W$ -algebra. By (7.3), we find that

$$\begin{aligned}\Delta_W(W_{i,j}^{(1)}) &= W_{i,j}^{(1)} \otimes 1 + 1 \otimes W_{i,j}^{(1)}, \\ \Delta_W(W_{i,j}^{(2)}) &= \begin{cases} W_{i,j}^{(2)} \otimes 1 + 1 \otimes W_{i,j}^{(2)} - \gamma_{w+1} \partial W_{i,j}^{(1)} \otimes 1 \\ \quad - \sum_{q_{Max} - \min(q_1, q_w) \leq u \leq q_{Max} - q_{Min}} (W_{u,j}^{(1)})_{(-1)} W_{i,u}^{(1)} \otimes 1 & \text{if } q_1 \geq q_l, \\ W_{i,j}^{(2)} \otimes 1 + 1 \otimes W_{i,j}^{(2)} - \gamma_{w+1} \partial W_{i,j}^{(1)} \otimes 1 \\ \quad - 1 \otimes \sum_{q_{Max} - \min(q_{w+1}, q_l) \leq u \leq q_{Max} - q_{Min}} (W_{u,j}^{(1)})_{(-1)} W_{i,u}^{(1)} & \text{if } q_1 < q_l. \end{cases}\end{aligned}$$

We also denote by  $\Delta_W$  a homomorphism

$$\Delta_W: \mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(N), f)) \rightarrow \mathcal{U}(\mathcal{W}^{k+N_2}(\mathfrak{gl}(N_1), f_1)) \otimes \mathcal{U}(\mathcal{W}^{k+N_1}(\mathfrak{gl}(N_2), f_2)),$$

which is induced by  $\Delta^W$  in (7.2).

## 8 A homomorphism from the affine Yangian to the universal enveloping algebra of $\mathcal{W}^k(\mathfrak{gl}(N), f)$

In this section, we will give the another proof to Theorem 6.1 in [28]. The following theorem is the same as Theorem 6.1 in [28].

**Theorem 8.1.** *Assume that  $q_{Min} \geq 3$  and  $\varepsilon = \hbar(k + N - q_{Min})$ . Then, there exists an algebra homomorphism*

$$\Phi: Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(q_{Min})) \rightarrow \mathcal{U}(\mathcal{W}^k(\mathfrak{gl}(N), f))$$

determined by

$$\begin{aligned}\Phi(X_{i,0}^+) &= \begin{cases} W_{q_{Max}, q_{Max} - q_{Min} + 1}^{(1)} t & \text{if } i = 0, \\ W_{q_{Max} - q_{Min} + i, q_{Max} - q_{Min} + i + 1}^{(1)} & \text{if } i \neq 0, \end{cases} \\ \Phi(X_{i,0}^-) &= \begin{cases} W_{q_{Max} - q_{Min} + 1, q_{Max}}^{(1)} t^{-1} & \text{if } i = 0, \\ W_{q_{Max} - q_{Min} + i + 1, q_{Max} - q_{Min} + i}^{(1)} & \text{if } i \neq 0, \end{cases}\end{aligned}$$

and

$$\begin{aligned}\Phi(H_{i,1}) &= -\hbar(W_{q_{Max} - q_{Min} + i, q_{Max} - q_{Min} + i}^{(2)} t - W_{q_{Max} - q_{Min} + i + 1, q_{Max} - q_{Min} + i + 1}^{(2)} t) \\ &\quad - \frac{i}{2} \hbar(W_{q_{Max} - q_{Min} + i, q_{Max} - q_{Min} + i}^{(1)} - W_{q_{Max} - q_{Min} + i + 1, q_{Max} - q_{Min} + i + 1}^{(1)}) \\ &\quad + \hbar W_{q_{Max} - q_{Min} + i, q_{Max} - q_{Min} + i}^{(1)} W_{q_{Max} - q_{Min} + i + 1, q_{Max} - q_{Min} + i + 1}^{(1)} \\ &\quad + \hbar \sum_{s \geq 0} \sum_{u=1}^i W_{q_{Max} - q_{Min} + i, q_{Max} - q_{Min} + u}^{(1)} t^{-s} W_{q_{Max} - q_{Min} + u, q_{Max} - q_{Min} + i}^{(1)} t^s \\ &\quad + \hbar \sum_{s \geq 0} \sum_{u=i+1}^n W_{q_{Max} - q_{Min} + i, q_{Max} - q_{Min} + u}^{(1)} t^{-s-1} W_{q_{Max} - q_{Min} + u, q_{Max} - q_{Min} + i}^{(1)} t^{s+1} \\ &\quad - \hbar \sum_{s \geq 0} \sum_{u=1}^i W_{q_{Max} - q_{Min} + i + 1, q_{Max} - q_{Min} + u}^{(1)} t^{-s} W_{q_{Max} - q_{Min} + u, q_{Max} - q_{Min} + i + 1}^{(1)} t^s \\ &\quad - \hbar \sum_{s \geq 0} \sum_{u=i+1}^n W_{q_{Max} - q_{Min} + i + 1, q_{Max} - q_{Min} + u}^{(1)} t^{-s-1} W_{q_{Max} - q_{Min} + u, q_{Max} - q_{Min} + i + 1}^{(1)} t^{s+1}\end{aligned}$$

for  $i \neq 0$ .

*Remark 8.2.* We note that the Definition 2.1 is different from the one in [28]. The parameter  $\varepsilon$  in this article is equal to  $-n\hbar - \varepsilon$  in [28]. This is the reason that the assumption of  $\varepsilon$  is different from the one in [28].

In [28], we gave the proof to Theorem 8.1 by a direct computation. In this article, we prove it by showing the following theorem. Let us take the permutation of  $\sigma \in S_l$  satisfying that

$$q_{\sigma(1)} \geq q_{\sigma(2)} \geq \cdots \geq q_{\sigma(l)}.$$

Let us set a homomorphism

$$\Delta^l = \left( \prod_{s=1}^{l-1} \Delta^{q_{\sigma(s)}, q_{\sigma(s+1)}} \otimes \text{id}^{l-s-1} \right) : Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(q_{Min})) \rightarrow \bigotimes_{1 \leq i \leq l} Y_{\hbar, \varepsilon - \hbar(q_{\sigma(i)} - q_{Min})}(\widehat{\mathfrak{sl}}(q_{\sigma(i)}))$$

where we set

$$\Delta^{q_{\sigma(s)}, q_{\sigma(s+1)}} = \begin{cases} \Delta^+ & \text{if } \sigma(s) < \sigma(s+1), \\ \Delta^- & \text{if } \sigma(s) > \sigma(s+1). \end{cases}$$

We also define a natural homomorphism

$$\Sigma : \bigotimes_{1 \leq i \leq l} Y_{\hbar, \varepsilon - \hbar(q_{\sigma(i)} - q_{Min})}(\widehat{\mathfrak{sl}}(q_{\sigma(i)})) \rightarrow \bigotimes_{1 \leq i \leq l} Y_{\hbar, \varepsilon - \hbar(q_i - q_{Min})}(\widehat{\mathfrak{sl}}(q_i)).$$

**Theorem 8.3.** *We obtain the following relation:*

$$\bigotimes_{1 \leq s \leq l} \text{ev}_{\hbar, \varepsilon - (q_s - q_{Min})\hbar}^{q_s, (\gamma_s - \frac{q_s - q_{Min}}{2})\hbar} \circ \Sigma \circ \Delta^l = \tilde{\mu} \circ \Phi.$$

Since  $\tilde{\mu}$  is an injective homomorphism ([11]) and  $\text{ev}_{\hbar, \varepsilon + (q_s - q_{Min})\hbar}^{q_s, \gamma_s \hbar}$ ,  $\Delta^l$  and  $\tilde{\mu}$  are homomorphisms, we obtain Theorem 8.3.

*The proof of Theorem 8.3.* In order to simplify the notation, we only show the case that  $q_1 \geq q_2 \geq \cdots \geq q_l$ . The other cases can be proven in a similar way.

It is enough to show the relations:

$$\bigotimes_{1 \leq s \leq l} \text{ev}_{\hbar, \varepsilon - (q_s - q_{Min})\hbar}^{q_s, (\gamma_s - \frac{q_s - q_{Min}}{2})\hbar} \circ \Delta^l(X_{j,0}^\pm) = \tilde{\mu} \circ \Phi(X_{j,0}^\pm) \quad (8.4)$$

for  $0 \leq j \leq n-1$  and

$$\bigotimes_{1 \leq s \leq l} \text{ev}_{\hbar, \varepsilon - (q_s - q_{Min})\hbar}^{q_s, (\gamma_s - \frac{q_s - q_{Min}}{2})\hbar} \circ \Delta^l(H_{i,1}) = \tilde{\mu} \circ \Phi(H_{i,1}) \quad (8.5)$$

for  $1 \leq i \leq n-1$ . It is trivial that (8.4) holds. We only show the relation (8.5). For  $A \in \widetilde{Y}_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n))$ ,  $\mathcal{U}(\widehat{\mathfrak{gl}}(q_s))$ , we denote  $A^{(s)} = 1^{\otimes s-1} \otimes A \otimes 1^{\otimes l-s}$ . By the definition of  $\Psi^{n,m+n}$  and  $\Delta^n$ , we obtain

$$\Delta^l(H_{i,1}) = \sum_{1 \leq s \leq l} H_{i+q_s-q_l}^{(s)} + B_i + C_i,$$

where we set

$$\begin{aligned} B_i &= -\hbar \sum_{r_1 < r_2} E_{q_{r_1}-q_l+i, q_{r_1}-q_l+i}^{(r_1)} E_{q_{r_2}-q_l+i+1, q_{r_2}-q_l+i+1}^{(r_2)} \\ &\quad - \hbar \sum_{r_1 < r_2} E_{q_{r_1}-q_l+i+1, q_{r_1}-q_l+i+1}^{(r_1)} E_{q_{r_2}-q_l+i, q_{r_2}-q_l+i}^{(r_2)} \end{aligned}$$

$$\begin{aligned}
& - \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-q_l+i} E_{q_{r_1}-q_{r_2}+u, q_{r_1}-q_l+i}^{(r_1)} t^{-s-1} E_{q_{r_2}-q_l+i, u}^{(r_2)} t^{s+1} \\
& + \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-q_l+i} E_{q_{r_1}-q_l+i, q_{r_1}-q_{r_2}+u}^{(r_1)} t^{-s} E_{u, q_{r_2}-q_l+i}^{(r_2)} t^s \\
& - \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=q_{r_2}-q_l+i+1}^{q_{r_2}} E_{q_{r_1}-q_{r_2}+u, q_{r_1}-q_l+i}^{(r_1)} t^{-s} E_{q_{r_2}-q_l+i, u}^{(r_2)} t^s \\
& + \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=q_{r_2}-q_l+i+1}^{q_{r_2}} E_{q_{r_1}-q_l+i, q_{r_1}-q_{r_2}+u}^{(r_1)} t^{-s-1} E_{u, q_{r_2}-q_l+i}^{(r_2)} t^{s+1} \\
& + \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-q_l+i} E_{q_{r_1}-q_{r_2}+u, q_{r_1}-q_l+i+1}^{(r_1)} t^{-s-1} E_{q_{r_2}-q_l+i+1, u}^{(r_2)} t^{s+1} \\
& - \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-q_l+i} E_{q_{r_1}-q_l+i+1, q_{r_1}-q_{r_2}+u}^{(r_1)} t^{-s} E_{u, q_{r_2}-q_l+i+1}^{(r_2)} t^s \\
& + \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=q_{r_2}-q_l+i+1}^{q_{r_2}} E_{q_{r_1}-q_l+i+1, q_{r_1}-q_{r_2}+u}^{(r_1)} t^{-s} E_{q_{r_2}-q_l+i+1, u}^{(r_2)} t^s \\
& - \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=q_{r_2}-q_l+i+1}^{q_{r_2}} E_{q_{r_1}-q_l+i+1, q_{r_1}-q_{r_2}+u}^{(r_1)} t^{-s-1} E_{u, q_{r_2}-q_l+i+1}^{(r_2)} t^{s+1}
\end{aligned}$$

and

$$\begin{aligned}
C_i = & \hbar \sum_{r=1}^l \sum_{s \geq 0} \sum_{u=1}^{q_r-q_l} E_{u, q_r-q_l+i}^{(r)} t^{-s-1} E_{q_r-q_l+i, u}^{(r)} t^{s+1} \\
& - \hbar \sum_{s \geq 0} \sum_{u=1}^{q_r-q_l} E_{u, q_r-q_l+i+1}^{(r)} t^{-s-1} E_{q_r-q_l+i+1, u}^{(r)} t^{s+1} \\
& + \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-q_l} E_{q_{r_1}-q_{r_2}+u, q_{r_1}-q_l+i}^{(r_1)} t^{-s-1} E_{q_{r_2}-q_l+i, u}^{(r_2)} t^{s+1} \\
& - \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-q_l} E_{q_{r_1}-q_{r_2}+u, q_{r_1}-q_l+i+1}^{(r_1)} t^{-s-1} E_{q_{r_2}-q_l+i+1, u}^{(r_2)} t^{s+1} \\
& + \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-q_l} E_{u, q_{r_2}-q_l+i}^{(r_2)} t^{-s-1} E_{q_{r_1}-q_l+i, q_{r_1}-q_{r_2}+u}^{(r_1)} t^{s+1} \\
& - \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-q_l} E_{u, q_{r_2}-q_l+i+1}^{(r_2)} t^{-s-1} E_{q_{r_1}-q_l+i+1, q_{r_1}-q_{r_2}+u}^{(r_1)} t^{s+1}.
\end{aligned}$$

We note that  $B_i$  comes from the coproduct for the affine Yangian and  $C_i$  comes from the homomorphism  $\Psi^{m.m+n}$ .

By the definition of the evaluation map, we obtain

$$\begin{aligned}
& \bigotimes_{1 \leq s \leq l} \text{ev}_{\hbar, \varepsilon - (q_s - q_{Min})\hbar}^{q_s, (\gamma_s - \frac{q_s - q_{Min}}{2})\hbar} \left( \bigotimes_{1 \leq s \leq l} H_{i+q_s-q_l}^{(s)} \right) \\
& = \sum_{r=1}^l \left( \gamma_r - \frac{i+q_s-q_l}{2} \hbar \right) (e_{q_1-q_l+i, q_1-q_l+i}^{(r)} - e_{q_1-q_l+i+1, q_1-q_l+i+1}^{(r)})
\end{aligned}$$

$$\begin{aligned}
& - \sum_{r=1}^l \hbar e_{q_1-q_l+i, q_1-q_l+i}^{(r)} e_{q_1-q_l+i+1, q_1-q_l+i+1}^{(r)} \\
& + \hbar \sum_{r=1}^l \sum_{s \geq 0} \sum_{u=q_1-q_r+1}^{q_1-q_l+i} e_{q_1-q_l+i, u}^{(r)} t^{-s} e_{u, q_1-q_l+i}^{(r)} t^s \\
& + \hbar \sum_{r=1}^l \sum_{s \geq 0} \sum_{u=q_1-q_l+i+1}^{q_1} e_{q_1-q_l+i, u}^{(r)} t^{-s-1} e_{u, q_1-q_l+i}^{(r)} t^{s+1} \\
& - \hbar \sum_{r=1}^l \sum_{s \geq 0} \sum_{u=q_1-q_r+1}^{q_1-q_l+i} e_{q_1-q_l+i+1, u}^{(r)} t^{-s} e_{u, q_1-q_l+i+1}^{(r)} t^s \\
& - \hbar \sum_{r=1}^l \sum_{s \geq 0} \sum_{u=q_1-q_l+i+1}^{q_1} e_{q_1-q_l+i+1, u}^{(r)} t^{-s-1} e_{u, q_1-q_l+i+1}^{(r)} t^{s+1}, \tag{8.6}
\end{aligned}$$

$$\begin{aligned}
& \bigotimes_{1 \leq s \leq l} \text{ev}_{\hbar, \varepsilon + (q_s - q_l) \hbar}^{q_s, (\gamma_s - \frac{q_s - q_{Min}}{2}) \hbar} (B_i) \\
& = -\hbar \sum_{r_1 < r_2} e_{q_1-q_l+i, q_1-q_l+i}^{(r_1)} e_{q_1-q_l+i+1, q_1-q_l+i+1}^{(r_2)} - \hbar \sum_{r_1 < r_2} e_{q_1-q_l+i+1, q_1-q_l+i+1}^{(r_1)} e_{q_1-q_l+i, q_1-q_l+i}^{(r_2)} \\
& - \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-q_l+i} e_{q_1-q_{r_2}+u, q_1-q_l+i}^{(r_1)} t^{-s-1} e_{q_1-q_{r_2}+i, u}^{(r_2)} t^{s+1} \\
& + \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-q_l+i} e_{q_1-q_l+i, q_1-q_{r_2}+u}^{(r_1)} t^{-s} e_{q_1-q_{r_2}+u, q_1-q_l+i}^{(r_2)} t^s \\
& - \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=q_{r_2}-q_l+i+1}^{q_{r_2}} e_{q_1-q_{r_2}+u, q_1-q_l+i}^{(r_1)} t^{-s} e_{q_1-q_l+i, q_1-q_{r_2}+u}^{(r_2)} t^s \\
& + \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=q_{r_2}-q_l+i+1}^{q_{r_2}} e_{q_1-q_l+i, q_1-q_{r_2}+u}^{(r_1)} t^{-s-1} e_{q_1-q_{r_2}+u, q_1-q_l+i}^{(r_2)} t^{s+1} \\
& + \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-q_l+i} e_{q_1-q_{r_2}+u, q_1-q_l+i+1}^{(r_1)} t^{-s-1} e_{q_1-q_l+i+1, q_1-q_{r_2}+u}^{(r_2)} t^{s+1} \\
& - \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-q_l+i} e_{q_1-q_l+i+1, q_1-q_{r_2}+u}^{(r_1)} t^{-s} e_{q_1-q_{r_2}+u, q_1-q_l+i+1}^{(r_2)} t^s \\
& + \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=q_{r_2}-q_l+i+1}^{q_{r_2}} e_{q_1-q_{r_2}+u, q_1-q_l+i+1}^{(r_1)} t^{-s} e_{q_1-q_l+i+1, q_1-q_{r_2}+u}^{(r_2)} t^s \\
& - \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=q_{r_2}-q_l+i+1}^{q_{r_2}} e_{q_1-q_l+i+1, q_1-q_{r_2}+u}^{(r_1)} t^{-s-1} e_{q_1-q_{r_2}+u, q_1-q_l+i+1}^{(r_2)} t^{s+1}, \tag{8.7}
\end{aligned}$$

$$\begin{aligned}
& \bigotimes_{1 \leq s \leq l} \text{ev}_{\hbar, \varepsilon - (q_s - q_{Min}) \hbar}^{q_s, (\gamma_s - \frac{q_s - q_{Min}}{2}) \hbar} (C_i) \\
& = \hbar \sum_{r=1}^l \sum_{s \geq 0} \sum_{u=1}^{q_r-q_l} e_{q_1-q_r+u, q_1-q_l+i}^{(r)} t^{-s-1} e_{q_1-q_l+i, q_1-q_r+u}^{(r)} t^{s+1} \\
& - \hbar \sum_{s \geq 0} \sum_{u=1}^{q_r-q_l} e_{q_1-q_r+u, q_1-q_l+i+1}^{(r)} t^{-s-1} e_{q_1-q_l+i+1, q_1-q_r+u}^{(r)} t^{s+1}
\end{aligned}$$

$$\begin{aligned}
& + \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-q_l} e_{q_1-q_{r_2}+u, q_1-q_l+i}^{(r_1)} t^{-s-1} e_{q_1-q_l+i, q_1-q_{r_2}+u}^{(r_2)} t^{s+1} \\
& - \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-q_l} e_{q_1-q_{r_2}+u, q_1-q_l+i+1}^{(r_1)} t^{-s-1} e_{q_1-q_l+i+1, q_1-q_{r_2}+u}^{(r_2)} t^{s+1} \\
& + \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-q_l} e_{q_1-q_{r_2}+u, q_1-q_l+i}^{(r_2)} t^{-s-1} e_{q_1-q_l+i, q_1-q_{r_2}+u}^{(r_1)} t^{s+1} \\
& - \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=1}^{q_{r_2}-q_l} e_{q_1-q_{r_2}+u, q_1-q_l+i+1}^{(r_2)} t^{-s-1} e_{q_1-q_l+i+1, q_1-q_{r_2}+u}^{(r_1)} t^{s+1}. \tag{8.8}
\end{aligned}$$

By the definition of  $\tilde{\mu}$ , we have

$$\begin{aligned}
& \hbar \tilde{\mu}(W_{q_1-q_l+i, q_1-q_l+i}^{(2)}) - \hbar \tilde{\mu}(W_{q_1-q_l+i+1, q_1-q_l+i+1}^{(2)}) \\
& = \hbar \sum_{r=1}^n \gamma_r e_{q_1-q_l+i, q_1-q_l+i}^{(r)} + \hbar \sum_{s \in \mathbb{Z}} \sum_{r_1 < r_2} \sum_{u > q_1-q_l} e_{u, q_1-q_l+i}^{(r_1)} t^{-s} e_{q_1-q_l+i, u}^{(r_2)} t^s \\
& - \hbar \sum_{s \geq 0} \sum_{r \geq 0} \sum_{1 \leq u \leq q_r - q_l} e_{q_1-q_r+u, q_1-q_l+i}^{(r)} t^{-s-1} e_{q_1-q_l+i, q_1-q_r+u}^{(r)} t^{s+1} \\
& - \hbar \sum_{s \geq 0} \sum_{r \geq 0} \sum_{1 \leq u \leq q_r - q_l} e_{q_1-q_l+i, q_1-q_r+u}^{(r)} t^{-s} e_{q_1-q_r+u, q_1-q_l+i}^{(r)} t^s \\
& - \hbar \sum_{s \in \mathbb{Z}} \sum_{r_1 < r_2} \sum_{1 \leq u \leq q_{r_2} - q_l} e_{q_1-q_l+i, q_1-q_{r_2}+u}^{(r_1)} t^{s+1} e_{q_1-q_{r_2}+u, q_1-q_l+i}^{(r_2)} t^{-s-1} \\
& - \hbar \sum_{r=1}^n \gamma_r e_{q_1-q_l+i+1, q_1-q_l+i+1}^{(r)} - \hbar \sum_{s \in \mathbb{Z}} \sum_{r_1 < r_2} \sum_{u > q_1-q_l} e_{u, q_1-q_l+i+1}^{(r_1)} t^{-s} e_{q_1-q_l+i+1, u}^{(r_2)} t^s \\
& + \hbar \sum_{s \geq 0} \sum_{r \geq 0} \sum_{1 \leq u \leq q_r} e_{q_1-q_r+u, q_1-q_l+i+1}^{(r)} t^{-s-1} e_{q_1-q_l+i+1, q_1-q_r+u}^{(r)} t^{s+1} \\
& + \hbar \sum_{s \geq 0} \sum_{r \geq 0} \sum_{1 \leq u \leq q_r} e_{q_1-q_l+i+1, q_1-q_r+u}^{(r)} t^{-s} e_{q_1-q_r+u, q_1-q_l+i+1}^{(r)} t^s \\
& + \hbar \sum_{s \in \mathbb{Z}} \sum_{r_1 < r_2} \sum_{1 \leq u \leq q_{r_2} - q_l} e_{q_1-q_l+i+1, q_1-q_{r_2}+u}^{(r_1)} t^{s+1} e_{q_1-q_{r_2}+u, q_1-q_l+i+1}^{(r_2)} t^{-s-1}. \tag{8.9}
\end{aligned}$$

Then, by a direct computation, the sum of (8.6)-(8.9) is equal to

$$\bigotimes_{1 \leq s \leq l} \text{ev}_{\hbar, \varepsilon - (q_s - q_{Min})\hbar}^{q_s, (\gamma_s - \frac{q_s - q_l}{2})\hbar} \circ \Delta^l(H_{i,1}) + \hbar \tilde{\mu}(W_{q_1-q_l+i, q_1-q_l+i}^{(2)}) - \hbar \tilde{\mu}(W_{q_1-q_l+i+1, q_1-q_l+i+1}^{(2)}).$$

For simplifying the notation, we denote the  $i$ -th term of the equation  $(\cdot)$  by  $(\cdot)_i$ . We divide the sum into 8 pieces:

$$(8.6)_1 + (8.9)_1 + (8.9)_6 = -\frac{i}{2} \hbar \tilde{\mu}(W_{q_1-q_l+i, q_1-q_l+i}^{(1)} - W_{q_1-q_l+i+1, q_1-q_l+i+1}^{(1)}), \tag{8.10}$$

$$(8.6)_2 + (8.7)_1 + (8.7)_2 = -\hbar \tilde{\mu}(W_{q_1-q_l+i, q_1-q_l+i}^{(1)} W_{q_1-q_l+i+1, q_1-q_l+i+1}^{(1)}), \tag{8.11}$$

$$(8.6)_3 + (8.6)_4 + (8.8)_1 + (8.9)_3 + (8.9)_4$$

$$\begin{aligned}
& = \hbar \sum_{r=1}^l \sum_{s \geq 0} \sum_{u=q_1-q_l+1}^{q_1-q_l+i} e_{q_1-q_l+i, u}^{(r)} t^{-s} e_{u, q_1-q_l+i}^{(r)} t^s \\
& + \hbar \sum_{r=1}^l \sum_{s \geq 0} \sum_{u=q_1-q_l+i+1}^{q_1} e_{q_1-q_l+i, u}^{(r)} t^{-s-1} e_{u, q_1-q_l+i}^{(r)} t^{s+1}, \tag{8.12}
\end{aligned}$$

$$\begin{aligned}
& (8.6)_5 + (8.6)_6 + (8.8)_2 + (8.9)_8 + (8.9)_9 \\
&= -\hbar \sum_{r=1}^l \sum_{s \geq 0} \sum_{u=q_1-q_l+1}^{q_1-q_l+i} e_{q_1-q_l+i+1,u}^{(r)} t^{-s} e_{u,q_1-q_l+i+1}^{(r)} t^s \\
&\quad - \hbar \sum_{r=1}^l \sum_{s \geq 0} \sum_{u=q_1-q_l+i+1}^n e_{q_1-q_l+i+1,u}^{(r)} t^{-s-1} e_{u,q_1-q_l+i+1}^{(r)} t^{s+1}, \tag{8.13}
\end{aligned}$$

$$\begin{aligned}
& (8.7)_3 + (8.7)_5 + (8.8)_3 + (8.9)_2 \\
&= \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=q_{r_2}-q_l+1}^{q_{r_2}-q_l+i} e_{q_1-q_{r_2}+u,q_1-q_l+i}^{(r_1)} t^s e_{q_1-q_{r_2}+i,u}^{(r_2)} t^{-s} \\
&\quad + \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=q_{r_2}-q_l+i+1}^{q_{r_2}} e_{q_1-q_{r_2}+u,q_1-q_l+i}^{(r_1)} t^{s+1} e_{q_1-q_l+i,q_1-q_{r_2}+u}^{(r_2)} t^{-s-1}, \tag{8.14}
\end{aligned}$$

$$\begin{aligned}
& (8.7)_4 + (8.7)_6 + (8.8)_5 + (8.9)_5 \\
&= \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=q_{r_2}-q_l+1}^{q_{r_2}-q_l+i} e_{q_1-q_l+i,q_1-q_{r_2}+u}^{(r_1)} t^{-s} e_{q_1-q_{r_2}+u,q_1-q_l+i}^{(r_2)} t^s \\
&\quad + \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=q_{r_2}-q_l+i+1}^{q_{r_2}} e_{q_1-q_l+i,q_1-q_{r_2}+u}^{(r_1)} t^{-s-1} e_{q_1-q_{r_2}+u,q_1-q_l+i}^{(r_2)} t^{s+1}, \tag{8.15}
\end{aligned}$$

$$\begin{aligned}
& (8.7)_7 + (8.7)_9 + (8.8)_4 + (8.9)_7 \\
&= -\hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=q_{r_2}-q_l+1}^{q_{r_2}-q_l+i} e_{q_1-q_{r_2}+u,q_1-q_l+i+1}^{(r_1)} t^s e_{q_1-q_l+i+1,q_1-q_{r_2}+u}^{(r_2)} t^{-s} \\
&\quad - \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=q_{r_2}-q_l+i+1}^{q_{r_2}} e_{q_1-q_{r_2}+u,q_1-q_l+i+1}^{(r_1)} t^{s+1} e_{q_1-q_l+i+1,q_1-q_{r_2}+u}^{(r_2)} t^{-s-1}, \tag{8.16} \\
& (8.7)_8 + (8.7)_{10} + (8.8)_6 + (8.9)_{10} \\
&= -\hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=q_{r_2}-q_l+1}^{q_{r_2}-q_l+i} e_{q_1-q_l+i+1,q_1-q_{r_2}+u}^{(r_1)} t^{-s} e_{q_1-q_{r_2}+u,q_1-q_l+i+1}^{(r_2)} t^s \\
&\quad - \hbar \sum_{r_1 < r_2} \sum_{s \geq 0} \sum_{u=q_{r_2}-q_l+i+1}^{q_{r_2}} e_{q_1-q_l+i+1,q_1-q_{r_2}+u}^{(r_1)} t^{-s-1} e_{q_1-q_{r_2}+u,q_1-q_l+i+1}^{(r_2)} t^{s+1}. \tag{8.17}
\end{aligned}$$

By a direct computation, we have

$$\begin{aligned}
(8.12) + (8.14) + (8.15) &= \tilde{\mu}(\hbar \sum_{s \geq 0} \sum_{u=1}^i W_{q_1-q_l+i,q_1-q_l+u}^{(1)} t^{-s} W_{q_1-q_l+u,q_1-q_l+i}^{(1)} t^s \\
&\quad + \hbar \sum_{s \geq 0} \sum_{u=i+1}^n W_{q_1-q_l+i,q_1-q_l+u}^{(1)} t^{-s-1} W_{q_1-q_l+u,q_1-q_l+i}^{(1)} t^{s+1})
\end{aligned}$$

and

$$\begin{aligned}
(8.13) + (8.16) + (8.17) &= \tilde{\mu}(-\hbar \sum_{s \geq 0} \sum_{u=1}^i W_{q_1-q_l+i+1,q_1-q_l+u}^{(1)} t^{-s} W_{q_1-q_l+u,q_1-q_l+i+1}^{(1)} t^s \\
&\quad - \hbar \sum_{s \geq 0} \sum_{u=i+1}^n W_{q_1-q_l+i+1,q_1-q_l+u}^{(1)} t^{-s-1} W_{q_1-q_l+u,q_1-q_l+i+1}^{(1)} t^{s+1}).
\end{aligned}$$

Thus, we have proven the relation (8.5).  $\square$

By using Theorem 8.1, we can show the compatibility with the coproduct for the affine Yangian and the parabolic induction for a  $W$ -algebra of type  $A$ . Let us take  $N_1, N_2$  and nilpotent elements of  $f_1, f_2$  in the same way as Section 7. By Theorem 8.1, we can construct two homomorphisms

$$\begin{aligned}\Phi_1 &: Y_{\hbar, \varepsilon + (\min(q_1, q_w) - q_{Min})\hbar}(\widehat{\mathfrak{sl}}(\min(q_1, q_w))) \rightarrow \mathcal{U}(\mathcal{W}^{k+N_2}(\mathfrak{gl}(N_1), f_1)), \\ \Phi_2 &: Y_{\hbar, \varepsilon + (\min(q_{w+1}, q_l) - q_{Min})\hbar}(\widehat{\mathfrak{sl}}(\min(q_{w+1}, q_l))) \rightarrow \mathcal{U}(\mathcal{W}^{k+N_1}(\mathfrak{gl}(N_2), f_2)).\end{aligned}$$

For a complex number  $a$ , we set a homomorphism called the shift operator of the affine Yangian:

$$\tau_a : Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n)) \rightarrow Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n))$$

determined by  $X_{i,0}^\pm \mapsto X_{i,0}^\pm$  and  $H_{i,1} \mapsto H_{i,1} + aH_{i,0}$ .

**Corollary 8.18.** *We obtain the following relations:*

$$\begin{aligned}(\Phi_1 \circ \tau_{-(\gamma_{w+1} - \frac{\min\{q_1, q_w\} - q_l}{2})\hbar} \circ \Psi^{q_1, \min(q_1, q_w)}) \otimes \Phi_2 \circ \Delta^+ &= \Delta_W \circ \Phi \text{ if } q_1 \geq q_l, \\ .(\Phi_1 \circ \tau_{-\gamma_{w+1}\hbar}) \otimes (\Phi_2 \circ \tau_{\frac{\min\{q_{w+1}, q_l\} - q_1}{2}\hbar} \circ \Psi^{q_1, \min(q_l, q_{w+1})}) \circ \Delta^- &= \Delta_W \circ \Phi \text{ if } q_1 < q_l.\end{aligned}$$

This corollary follows directly from Theorem 8.3 and the relation (7.3).

## References

- [1] T. Arakawa. Representation theory of  $W$ -algebras. *Invent. Math.*, 169(2):219–320, 2007, <https://doi.org/10.1007/s00222-007-0046-1>.
- [2] J. Brundan and A. Kleshchev. Parabolic presentations of the Yangian  $Y(\mathfrak{gl}_n)$ . *Comm. Math. Phys.*, 254(1):191–220, 2005.
- [3] J. Brundan and A. Kleshchev. Shifted Yangians and finite  $W$ -algebras. *Adv. Math.*, 200(1):136–195, 2006, <https://doi.org/10.1016/j.aim.2004.11.004>.
- [4] T. Creutzig, D. E. Diaconescu, and M. Ma. Affine Laumon spaces and iterated  $W$ -algebras. *Comm. Math. Phys.*, 402(3):2133–2168, 2023.
- [5] A. De Sole and V. G. Kac. Finite vs affine  $W$ -algebras. *Jpn. J. Math.*, 1(1):137–261, 2006, <https://doi.org/10.1007/s11537-006-0505-2>.
- [6] A. De Sole, V. G. Kac, and D. Valeri. A Lax type operator for quantum finite  $W$ -algebras. *Selecta Math. (N.S.)*, 24(5):4617–4657, 2018.
- [7] V. G. Drinfeld. Hopf algebras and the quantum Yang-Baxter equation. *Dokl. Akad. Nauk SSSR*, 283(5):1060–1064, 1985, <https://doi.org/10.1142/9789812798336.0013>.
- [8] V. G. Drinfeld. A new realization of Yangians and of quantum affine algebras. *Dokl. Akad. Nauk SSSR*, 296(1):13–17, 1987.
- [9] M. Finkelberg and A. Tsymbaliuk. Multiplicative slices, relativistic Toda and shifted quantum affine algebras. In *Representations and nilpotent orbits of Lie algebraic systems*, volume 330 of *Progr. Math.*, pages 133–304. Birkhäuser/Springer, Cham, 2019.
- [10] I. B. Frenkel and Y. Zhu. Vertex operator algebras associated to representations of affine and Virasoro algebras. *Duke Math. J.*, 66(1):123–168, 1992, <https://doi.org/10.1215/S0012-7094-92-06604-X>.
- [11] N. Genra. Screening operators for  $W$ -algebras. *Selecta Math. (N.S.)*, 23(3):2157–2202, 2017, <https://doi.org/10.1007/s00029-017-0315-9>.

- [12] N. Genra. Screening operators and parabolic inductions for affine  $\mathcal{W}$ -algebras. *Adv. Math.*, 369:107179, 62, 2020, <https://doi.org/10.1016/j.aim.2020.107179>.
- [13] N. Guay. Cherednik algebras and Yangians. *Int. Math. Res. Not.*, (57):3551–3593, 2005, <https://doi.org/10.1155/IMRN.2005.3551>.
- [14] N. Guay. Affine Yangians and deformed double current algebras in type A. *Adv. Math.*, 211(2):436–484, <https://doi.org/10.1016/j.aim.2006.08.007>, 2007.
- [15] N. Guay, H. Nakajima, and C. Wendlandt. Coproduct for Yangians of affine Kac-Moody algebras. *Adv. Math.*, 338:865–911, 2018, <https://doi.org/10.1016/j.aim.2018.09.013>.
- [16] V. G. Kac and M. Wakimoto. Quantum reduction and representation theory of superconformal algebras. *Adv. Math.*, 185(2):400–458, 2004, <https://doi.org/10.1016/j.aim.2003.12.005>.
- [17] R. Kodera. Braid group action on affine Yangian. *SIGMA Symmetry Integrability Geom. Methods Appl.*, 15:Paper No. 020, 28, 2019, <https://doi.org/10.3842/SIGMA.2019.020>.
- [18] R. Kodera. On Guay’s evaluation map for affine Yangians. *Algebr. Represent. Theory*, 24(1):253–267, 2021, <https://doi.org/10.1007/s10468-019-09945-w>.
- [19] R. Kodera and M. Ueda. Coproduct for affine Yangians and parabolic induction for rectangular  $W$ -algebras. *Lett. Math. Phys.*, 112(1):Paper No. 3, 37, 2022.
- [20] A. Matsuo, K. Nagatomo, and A. Tsuchiya. Quasi-finite algebras graded by Hamiltonian and vertex operator algebras. *London Math. Soc. Lecture Note Ser.*, 372:282–329, 2010, <https://doi.org/10.1017/CBO9780511730054.015>.
- [21] A. Premet. Special transverse slices and their enveloping algebras. *Adv. Math.*, 170(1):1–55, 2002, <https://doi.org/10.1006/aima.2001.2063>. With an appendix by Serge Skryabin.
- [22] O. Schiffmann and E. Vasserot. Cherednik algebras,  $W$ -algebras and the equivariant cohomology of the moduli space of instantons on  $\mathbf{A}^2$ . *Publ. Math. Inst. Hautes Études Sci.*, 118:213–342, 2013, <https://doi.org/10.1007/s10240-013-0052-3>.
- [23] M. Ueda. An Example of Homomorphisms from the Guay’s affine Yangians to non-rectangular  $W$ -algebras. *Transformation Groups*(2023).
- [24] M. Ueda. A homomorphism from the affine Yangian  $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n))$  to the affine Yangian  $Y_{\hbar, \varepsilon}(\widehat{\mathfrak{sl}}(n+1))$ . arXiv:2312.09933.
- [25] M. Ueda. Notes on a homomorphism from the affine Yangian associated with  $\widehat{\mathfrak{sl}}(n)$  to the affine Yangian associated with  $\widehat{\mathfrak{sl}}(n)$ . arXiv:2402.01870.
- [26] M. Ueda. Two homomorphisms from the affine Yangian associated with  $\widehat{\mathfrak{sl}}(n)$  to the affine Yangian associated with  $\widehat{\mathfrak{sl}}(n)$ . arXiv:2404.10923.
- [27] M. Ueda. Affine super Yangians and rectangular  $W$ -superalgebras. *J. Math. Phys.*, 63(5):Paper No. 051701, 34, 2022.
- [28] M. Ueda. Guay’s affine Yangians and non-rectangular  $W$ -algebras. *Adv. Math.*, 438:Paper No. 109468, 44, 2024.