

Numerical study of the properties of a holographic superconductor from an anti-de Sitter-Einstein-Born-Infeld black hole with backreaction

Uriel Elinos Calderón* and Marco Maceda†

*Departamento de Física, Universidad Autónoma Metropolitana-Iztapalapa,
Av. Ferrocarril San Rafael Atlixco 186, Ciudad de México 09310, México*

Edgar Guzmán-González‡

*School of Physics and Optoelectronic Engineering, Hainan University, 570228 Haikou, P. R. China and
Departamento de Física, Universidad Autónoma Metropolitana-Iztapalapa,
Av. Ferrocarril San Rafael Atlixco 186, Ciudad de México 09310, México*

(Dated: April 26, 2024)

We study numerically an s -wave holographic superconductor from an anti-de Sitter-Einstein-Born-Infeld black hole with backreaction in the context of the AdS/CFT correspondence. By introducing a parameter G to tune the effects of the backreaction, we can study non-perturbatively how the condensation and conductivity of the superconductor change as the backreaction increases. We find that for small values of G , increasing the nonlinearity of the Born-Infeld model makes the formation of the condensate harder — consistent with previous results reported in the literature — while for values of G close to one, the opposite effect occurs; increasing the nonlinearity slightly facilitates its formation. We also determine how the ratio ω_g/T_c varies for different intensities of nonlinearity and backreaction, showing in particular that large deviations from the so-called universal value arise as backreaction becomes stronger.

I. INTRODUCTION

By this point in time, it is difficult to overstate the huge and fruitful impact that the AdS/CFT correspondence has on various areas of theoretical physics, like quantum information, nuclear physics [1–3] and condensed matter physics [4–8]. Said impact comes mainly from the fact that it establishes a duality between a string theory and a conformal quantum field theory (CFT). In certain cases, it is possible to work with models of string theory below 5 dimensions in an anti-de Sitter (AdS) background, within a sector that can be described as classical gravity. Even though these models have some limitations, they capture important phenomenological aspects of the dual system. In this setting, localized excitations in the CFT are described by data from the boundary on the gravity side; nevertheless a specific string theory is dual to a CFT at any depth of the bulk of the string theory.

Many systems in condensed matter physics can be studied using this duality [9]. An important example in this field that has been extensively studied using the correspondence is the one of superconductors, as it is believed that they can provide the first steps to understand theoretically the so-called high temperature superconductors [10–13]; these are strongly interacting systems that can not be described using the standard BCS theory, as it presupposes weak interactions [14–16]. In this particular case, the duality is between a superconductor and a black hole in the gravitational theory; for obvious reasons, the superconductors obtained from this correspondence are termed holographic superconductors.

One point of particular interest in this kind of analysis, is to determine how the properties of the spacetime of the gravitational theory leave their imprint in the ones of the superconductor. For this reason, different holographic superconductors have been studied previously in the literature, analyzing several of their key properties like the expectation value of the superconducting condensate, its critical temperature and its electrical conductivity. Most of these studies follow a bottom-up approach, where one puts the basic ingredients of the theory of gravity by hand, without worrying about the details of how such theory can be embedded in supergravity.

The motivation for holographic superconductor models is at least twofold: first, theoretical, to study superconductors with strong interactions in a controlled way, and secondly, practical, to see how to change the parameters of the model to engineer a superconductor with desirable properties, like having high critical temperature, high magnetic critical field or high values for the condensate. The existence of excited states or more realistic scenarios where impurities are present may arise from models where modifications in the properties of spacetime are considered; among those that have been considered in the literature we may mention the inclusion of the Ricci curvature of the metric [17], different nonrelativistic theories for the underlying gravity [18–22], nonlinear electrodynamic theories [23, 24], and noncommutative effects [25–27].

It is important to remark that when studying the gravitational theory, the resulting differential equations for the fields and the metric are nonlinear, making an exact analytical treatment next to impossible, so one has to resort to approximations, numerical

* elinosur@gmail.com

† mmac@xanum.uam.mx

‡ edgar.guzman@hainanu.edu.cn

methods or a combination of both. Some examples of these types of techniques include the Sturm-Liouville method [28, 29] or a matching method, where asymptotic solutions to the differential equations near the horizon of the black hole and near the boundary are pasted consistently [17, 30, 31]. Due to the complexity of the resulting field equations, most of these works neglect the backreaction of the different matter fields on the metric, so that the fields are on a fixed background. Even though this approximation greatly simplifies the analysis and gives results that correctly capture the essential physics, to study temperatures considerably smaller than the critical one, or to obtain more precise quantitative results, numeric methods are necessary.

In this work, we use the theoretical framework developed in [11] to numerically analyse an holographic superconductor with backreaction, but instead of considering Maxwell electrodynamics, we consider Born-Infeld electrodynamics [32, 33]. In Born-Infeld electrodynamics the self-energy of a point charge is regularized by introducing a maximum value b for the magnitude of the electric field. In the limits where either b or the distance from the area of study to the sources goes to infinity, Maxwell's theory is recovered. This makes holographic studies particularly simple, as the *dictionary* used to translate the properties of the QFT to the ones of the AdS are essentially the same as in the Maxwell case, as the boundary is far away from the sources. Previous studies of holographic superconductors with Born-Infeld electrodynamics in the probe limit include [34, 35]. Studies beyond the probe limit were done for $(1+1)$ spacetimes [36], for $(4+1)$ spacetimes [37] and close to the probe limit [38, 39]. Here we consider $(3+1)$ spacetimes further away from the probe limit and explore more in detail the formation of a condensate with a full backreacted metric. Studies in the backreacted case in other gravitational backgrounds have also been carried out [40–45] but employing techniques as the matching method and with a limited range of validity. As mentioned previously, with our numerical analysis, we gain insight into a fully back-reacted Born-Infeld holographic superconductor; by its own nature, our approach covers non-perturbative aspects of holographic superconductors.

The structure of the paper is as follows, in Section II we review the basic equations and formalism to read the properties of the superconductor from the ones of the gravitational theory, as well as how to set the initial conditions for the numerical methods. We also discuss the effect of increasing the backreaction and the nonlinearity in the electrodynamical theory on the condensation properties and critical temperature of the holographic superconductor. In Section III we make a similar study to the previous one but for AC conductivity of the superconductor near the critical temperature; we discuss there the behaviour of the ratio ω_g/T_c as function of the Born-Infeld parameter and backreaction. Finally, in Section IV we present our general conclusions, a summary of our results and some possible directions for further research.

II. MODEL FOR THE HOLOGRAPHIC SUPERCONDUCTOR

First, we briefly describe the bulk theory that we are considering. The dynamics of the model, in natural units, is given by the following action (assuming a sum over repeated indices running from 0 to 3)

$$\begin{aligned} S &= \int d^{3+1}x \sqrt{-g} \mathcal{L}, \\ \mathcal{L} &= \frac{1}{16\pi} \left(G^{-1} (R - 2\Lambda) + \mathcal{L}_B \right) - \frac{m^2}{L^2} \psi \psi^* \\ &\quad - (\partial_\alpha \psi - iq A_\alpha \psi) g^{\alpha\beta} (\partial_\beta \psi^* + iq A_\beta \psi^*), \end{aligned} \quad (1)$$

where g is the metric, R is its Ricci scalar, L is the AdS radius and Λ is a cosmological constant; ψ is a complex scalar field with charge q and mass m . A_μ is the electromagnetic potential associated with the Born-Infeld Lagrangian \mathcal{L}_B defined in terms of the electromagnetic tensor $F_{\alpha\beta} = \partial_\alpha A_\beta - \partial_\beta A_\alpha$ as follows,

$$\mathcal{L}_B = 4b^2 \left(1 - \sqrt{1 + \frac{F_{\alpha\beta} F^{\alpha\beta}}{2b^2}} \right), \quad (2)$$

with b the Born-Infeld parameter, that physically is the maximum value for the magnitude of the electric field, and G is the universal gravitational constant. Clearly, at regions where $|F_{\alpha\beta}| \ll b$ we recover Maxwell's linear theory, $L_B(F) \approx -F^{\alpha\beta} F_{\alpha\beta}$.

Note that, as done in [9], G can be regarded as a parameter that controls the effects of the backreaction — when $G = 0$ the dynamics of the metric is given only by Einstein's term $R - 2\Lambda$; as G keeps increasing, the effects on the metric of the other fields also increase and $G = 1$ gives the *fully backreacted case*.

The variation of S with respect to the fields g , ψ , ψ^* and A_μ produces all the equations of motion for the system. This theory allows us to have a system that admits hairy solutions at low temperatures and not at high ones and is dual to a CFT that has a conserved current and a charged operator to break the $U(1)$ symmetry. Also in order to obtain a theory that is dual to an s -wave superconductor on the CFT, we may consider a plane-symmetric metric (other options for the metric's symmetry include spherical or hyperbolic, but all of them are related to different radial foliations associated with the metric where the CFT resides, resulting in different approaches to duality with similar results to the ones presented here [1]) with a charged black hole (say,

at the origin) that is asymptotically anti-de Sitter [10]. The simplest metric that satisfies these requirements is of the following form,

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)n(r)} + r^2(dx^2 + dy^2), \quad (3)$$

where t, r, x, y , are coordinates of the $4D$ spacetime, being r a radial coordinate, $f(r)$ is the metric function and $n(r)$ measures the effects of the backreaction — indeed, $n(r) \equiv 1$ in the probe limit. The boundary of the AdS, where the CFT “lives” is located at $r \rightarrow \infty$. The value r_+ where f vanishes is the horizon radius of the black hole. The temperature of the black hole is given by Hawking’s celebrated expression (we assume that $n(r_+) \neq 0$ and that $f(r), n(r)$ are positive for $r > r_+$),

$$T = \frac{\sqrt{n(r_+)f'(r_+)}}{4\pi}. \quad (4)$$

Since we have a plane-symmetric metric, we assume that ψ only depends on r , and, since we will assume that there is no magnetic field, we initially work on a gauge where the potential takes the following simple form, $A_t = \phi(r)$, $A_r = A_x = A_y = 0$. In terms of this *Ansatz*, Einstein’s equations for the components tt and rr , become (where a prime $'$ denotes the derivative with respect to r),

$$\begin{aligned} 0 &= 16\pi G q^2 r \psi \psi^* \phi^2 + f^2(n' + 16\pi G r n \psi' \psi'^*), \\ 0 &= \frac{4b^2 r^2 G}{\sqrt{1 - \frac{n\phi'^2}{b^2}}} - 2n(rf' + f) - rfn' \\ &\quad - 16\pi G m^2 r^2 \psi \psi^* - 2\Lambda r^2 + 4b^2 r^2 G, \end{aligned} \quad (5)$$

while Born-Infeld’s equation for A_t and the one for the scalar field ψ become,

$$\begin{aligned} 0 &= 8\pi q^2 r^2 \psi \psi^* \phi - f\sqrt{n} \left(\frac{r^2 \sqrt{n} \phi'}{\sqrt{1 - \frac{n\phi'^2}{b^2}}} \right)', \\ 0 &= (q^2 \phi^2 - m^2 f) r^2 \psi + f\sqrt{n} \left(r^2 f \sqrt{n} \psi' \right)'. \end{aligned} \quad (6)$$

The remaining Einstein’s and Born-Infeld’s equations can be proved to be a combination of these four, so they are satisfied automatically if (5) and (6) are. It is also clear from (6) that, since we can assume real initial conditions for ψ (because of the invariance of the action (1) under the transformation $\psi \rightarrow e^{i\gamma}\psi$, with γ real), we can assume that ψ is real throughout all the solution.

Once we have obtained the equations of motion, our goal is to solve them in full generality for arbitrary values of the Born-Infeld parameter; moreover, we want to analyse the behaviour of the solutions when backreaction becomes important. For this purpose, we have to make use of numerical solutions that allow us to explore the full moduli space of the model; in this way, we will obtain a complete characterization of the associated holographic superconductor, in contrast to previous treatments in the literature where effects linear on the Born-Infeld parameter have been discussed and where backreaction has also been considered perturbatively.

The approach we use to solve the previous system of equations numerically is the *shooting method*. As a first step towards its implementation, we fix the parameters of the model to the values $L = 1$, $q = 1$, $m^2 = -2$; next, we impose some regularity and physical conditions at the horizon r_+ and at the boundary; specifically, we begin discussing the asymptotics near the boundary since several interesting observables of the QFT can be read from the asymptotic behaviour of the fields at the boundary, as we remind the reader below.

By studying the equations of motion when $r \rightarrow \infty$, we see that the fields have the following asymptotic form

$$\begin{aligned} \psi(r) &= \frac{\mathcal{J}_O}{r} + \frac{\langle O_2 \rangle}{\sqrt{2}r^2} + \dots, \\ \phi(r) &= \mu - \frac{\rho}{r} + \dots, \\ f(r) &= \lambda^{-1}r^2 + f^0 + \frac{f^1}{r} + \dots, \\ n(r) &= \lambda n_\infty^0 + \frac{n_\infty^1}{r} \dots \end{aligned} \quad (7)$$

where, according to the AdS/CFT dictionary [8, 10, 46], with the gauge we are considering, $\langle O_2 \rangle$ is the expectation value of the superconducting condensate, \mathcal{J}_O , the one of its source, μ is its chemical potential and ρ is its charge density. f^0 is related with \mathcal{J}_O and f^1 is related with the free energy of the system; the particular relationship will not be of interest for our work. To study unsourced spontaneous condensation, we set $\mathcal{J}_O = 0$. This is a condition that has to be implemented in the shooting method.

In order for the temperature of the black hole (4) to coincide with the one of the CFT, it is required that $\lambda = 1$ [11]. In principle, this would be another condition for the shooting method, however, we can simplify this step by fixing $n(r_+)$ to an arbitrary value (say, $n(r_+) = 1$), solve for the remaining conditions using the shooting method and then apply the following scaling symmetry of the field equations that map solutions of the equations of motion to solutions (note that this scaling is the one induced by the coordinate transformation $t' = \lambda t$),

$$f \rightarrow \lambda f, n \rightarrow n/\lambda, \phi \rightarrow \sqrt{\lambda}\phi, \psi \rightarrow \psi. \quad (8)$$

As can be easily verified, this scaling satisfies the remaining conditions $\mathcal{J}_O = 0$ and (10) down below that we enforce on the shooting method. By imposing regularity of the fields in (7), the value $\Lambda = -3$ also gets fixed.

To ensure a divergence-free QFT dual to a charged black hole, it is necessary that the fields f , n , Ψ , and ϕ are analytic at the horizon [47]. Thus, we propose Taylor expansions for these fields around $r = r_+$ and find the relationships among the coefficients of the expansion that ensure their analyticity. These relationships are also important for the numerical implementation of the shooting method. Since $f(r_+) = 0$, the equations (6) are singular at r_+ , so we cannot shoot solutions from r_+ . To circumvent this limitation, we utilize the obtained expansions to evaluate the fields and their derivatives at $r_+ + \epsilon$, with $\epsilon \ll r_+$, providing the necessary initial conditions to shoot solutions from $r + \epsilon$, where the system is not singular. Therefore, if we assume an expansion of the form,

$$\begin{aligned} f(r) &= f_1(r - r_+) + f_2(r - r_+)^2 + \dots, \\ n(r) &= n_0 + n_1(r - r_+) + n_2(r - r_+)^2 + \dots, \\ \psi(r) &= \psi_0 + \psi_1(r - r_+) + \psi_2(r - r_+)^2 + \dots, \\ \phi(r) &= \phi_0 + \phi_1(r - r_+) + \phi_2(r - r_+)^2 + \dots, \end{aligned} \quad (9)$$

and impose analyticity, we conclude the following relations,

$$\begin{aligned} f_1 &= \frac{r_+}{n_0} \left(2b^2 G - \frac{2b^2 G}{\sqrt{1 - \frac{n_0 \phi_1^2}{b^2}}} - 8\pi G m^2 \psi_0^2 - \Lambda \right), \\ \phi_0 &= 0, \quad \psi_1 = \frac{m^2 \psi_0}{f_1 n_0}. \end{aligned} \quad (10)$$

As already mentioned, we can set $n_0 = 1$ and then apply a scaling if necessary. The previous equations fix the remaining necessary conditions in terms of just ψ_0 and ϕ_1 (fixing b , q , G and m), making these two the only parameters we need to vary when applying the shooting method.

So in summary, to apply the shooting method, we do the following: First, we chose a value for r_+ , in our case, we set $r_+ = 1$. Then, we solve the system of equations (5), (6), (10), $f(r_+) = 0$ and $n_0 = 1$, keeping ψ_0 and ϕ_1 as free parameters. Once we have solved the system of equations numerically for some pair (ψ_0, ϕ_1) , we can do a fit for $r \gg r_+$ to compute \mathcal{J}_O and the remaining observables of (7). By varying the values of ψ_0 and ϕ_1 , we can find a curve in the ψ_0 - ϕ_1 plane such that $\mathcal{J}_O = 0$. After applying the scaling (8) to the fields obtained by this procedure, we can use their data to study different properties of the unsourced superconducting condensate.

Since in a QFT only dimensionless quantities have physical meaning, we focus on the ratio $T/\sqrt{\rho}$ (we could consider other dimensionless ratios like T/μ , but they all produce similar results). The parameters ψ_0 , ϕ_1 , where $\langle O_2 \rangle$ becomes zero, are the ones that correspond to the solution related to the critical temperature T_c ; by finding them numerically, we can compute the ratio $T_c/\sqrt{\rho}$ as a function of G and b . In Fig. 1, we fix the values $b = 7$, $b = 3$ and plot $T_c/\sqrt{\rho}$ as a function of G . Note that, when G increases, the ratio $T_c/\sqrt{\rho}$ becomes smaller; this means that, at fixed ρ , increasing G implies a lower critical temperature (i.e. the formation of the superconducting condensate becomes harder). For low values of G , we see a similar behaviour for b , since the $b = 7$ curve is above the $b = 3$ one; we see that decreasing the values of b (increasing the nonlinearity on the electrodynamic theory) makes the condensation harder, however, there is a value G^* (somewhere between $G = 0.6$ and $G = 0.7$) such that, for $G > G^*$, increasing the nonlinearity of the electrodynamic theory makes the condensation slightly easier.

With the proportionality constant between T_c and ρ determined, we can obtain similar conclusions to the previous ones by considering plots of T/T_c vs $\sqrt{\langle O_2 \rangle}/T_c$ for different values of b and G . The results are presented in Fig. 2. In these types of curves, the higher the values of $\sqrt{\langle O_2 \rangle}/T_c$, the harder the formation of superconducting condensate; this behaviour can be properly explained by noticing that the value of the critical temperature T_c gets lower and this indicates that the superconducting




FIG. 1. Plot of the ratio $T_c/\sqrt{\rho}$ as a function of G with $q = 1$ and different values of b . From the left side, the upper curve is for $b = 7$ while the lower one is for $b = 3$. Notice how, for G close to one, the situation reverses, the $b = 3$ curve is now above the $b = 7$ one, indicating that the formation of the condensate is easier for $b = 3$.




FIG. 2. Plot of T/T_c vs $\sqrt{\langle O_2 \rangle}/T_c$ for different values of G . The dashed curves correspond to $b = 3$, while continuous lines are for $b = 7$. The values of G considered are, from top to bottom, $G = 1, 0.7, 0.3, 0$. Notice how, for $G = 0, 0.3$ (small backreaction), the $b = 7$ curves are below the $b = 3$ curves, indicating that the formation of the condensate is harder for $b = 3$, however, for $G = 0.7, 1$ the situation reverses, condensation becomes easier for $b = 3$.

state has been destabilized. This also can be analyzed noticing that the value of the condensate is proportional to the energy gap of the superconducting state, and from these results we can see that for a fixed gap energy, increasing the backreaction requires lower critical temperature to achieve it. From here we confirm the previous conclusions, for $G \in \{0, 0.3\}$, condensation is easier for $b = 7$, while, for $G \in \{0.7, 1.0\}$ the condensation is easier for $b = 3$. We also mention that, as found in previous works, [30, 45], from our numerical analysis we observe the critical exponent of $1/2$ characteristic of second order phase transitions, $\langle O_2 \rangle \propto (1 - T/T_c)^{1/2}$, independently of b and G .

III. CONDUCTIVITY

With the AdS/CFT correspondence, we can study the AC conductivity of the superconducting condensate in the usual way — we turn on a small harmonic electric field with frequency ω and study the linear response of the system [48]. In our setting, the simplest electric field is one that has only either the x or y component — for concreteness we chose the x component — and zero momentum (implying that there will be no heat flow in the superconductor [11]), so we set $A_x(r, t) = \epsilon e^{-i\omega t} A_x(r)$, and study the system up to first order in ϵ .

With $A_x \neq 0$, the Ansatz for the metric (3) is no longer consistent, as Einstein's equation for the tx component is independent

from the remaining equations, producing a system with more independent equations than functions and hence, with no solutions generically. To the first order in ϵ , this can be easily solved by adding the term $\epsilon g_{xt}(r)e^{-i\omega t} dt dx$ to the Ansatz (3). To this order, Eqs. (6) and (7) for the original fields remain the same, while we find the following new independent equations,

$$\begin{aligned}
0 &= \frac{4GA_x\phi'}{\sqrt{1-\frac{n\phi'^2}{b^2}}} + g'_{xt} - \frac{2g_{xt}}{r}, \\
0 &= \left(\frac{\omega^2}{fn} - \frac{4G\phi'^2}{\sqrt{1-\frac{n\phi'^2}{b^2}}} - \frac{8\pi q^2\psi^2\sqrt{1-\frac{n\phi'^2}{b^2}}}{n} \right) \frac{A_x}{f} \\
&\quad + \left(\frac{n'}{2n} + \frac{f'}{f} + \frac{8\pi q^2\psi^2\phi\phi'\sqrt{1-\frac{n\phi'^2}{b^2}}}{b^2f} - \frac{2n\phi'^2}{b^2r} \right) A'_x \\
&\quad + A''_x,
\end{aligned} \tag{11}$$

where we have written g_{tx} in terms of A_x and its derivatives in the last equation, so it is not necessary to find g_{tx} to determine the remaining fields. Therefore, it is enough to just discuss the conditions for A_x at r_+ and at the boundary.

At r_+ , we require that A_x describes an ingoing wave (reflecting the fact that no electromagnetic wave gets out of a black hole [47]), so we assume an expansion near r_+ of the form,

$$A_x(r) = (r - r_+)^{-i\kappa} (A_{x0} + A_{x1} + \dots), \tag{12}$$

with $\kappa > 0$. By considering this expansion in (11) we conclude,

$$\kappa = \frac{\omega}{f_1\sqrt{n_0}}, \tag{13}$$

together with a linear relation between A_{x0} and A_{x1} that we do not present here because of its considerably big size. From this analysis we conclude that we only have to specify a single condition for A_x ; since (11) is linear in A_x by construction, and we are interested in the ratio of two coefficients related to A_x (see Eq. (14)), we can simply set $A_{x0} = 1$.

By considering (11), we see that at the boundary, A_x has the following asymptotic expression,

$$A_x(r) = -\frac{i\mathcal{E}}{\omega} + \frac{\mathcal{J}_\mathcal{E}}{r} + \dots, \tag{14}$$

where \mathcal{E} is the x component in the dual theory of the electric field produced by the time dependent electromagnetic potential and $\mathcal{J}_\mathcal{E}$ is the corresponding current. The previous expression can be obtained either by naive considerations or by using the GKPW prescription [49]. Although the latter computation for the Born-Infeld's case is similar to the one of Maxwell's [11], it is instructive to derive it, so we include it in appendix A.

Either way, by using Ohm's law, we can directly compute the conductivity σ from $\mathcal{J}_\mathcal{E} = \sigma\mathcal{E}$ [28]. Similar techniques can be used to compute other transport properties, like the heat current [47, 50] or the correlation length [51].

In Figure 3, we present a plot of the frequency ω (measured in units of $\langle\mathcal{O}_2\rangle$) vs. the real part of the conductivity $\text{Re}[\sigma]$ for a fixed temperature $T = 0.9T_c$ and different values of b and G . The behavior of $\text{Re}[\sigma]$ can be explained by describing the superconductor in terms of the two-fluid model and studying the response of the “superconducting” and the “normal” electrons to alternating electromagnetic fields. According to the BCS theory, at $T = 0$, there are no normal electrons for small ω , and for $T > 0$, their presence is related to thermal excitations [15]. Precisely at $T = 0$, the only mechanism available for the system to absorb an incident electromagnetic wave is by creating pairs of electrons. Therefore, for frequencies below the gap frequency ω_g —the frequency associated with the necessary energy to break a Cooper pair into a pair of electrons [47]— $\text{Re}[\sigma]$ is zero [52, 53]. At ω_g , there is an absorption edge, and the conductivity quickly increases from zero to unity, resembling the behavior of a holographic conductor in the normal state [54]. For $T > 0$, although most of the current is supercurrent, the nonzero $\text{Re}[\sigma]$ is related to the presence of normal electrons mentioned earlier. In this case, ω_g can be detected as the frequency where $\text{Re}[\sigma]$ starts growing abruptly, and it coincides with the frequency where $\text{Im}[\sigma]$ attains a minimum value [55]. With this in mind, we can observe in Figure 3 how the energy gap increases as we increase G , while it remains almost unchanged when we vary from $b = 3$ to $b = 7$. This can be explained by recalling that higher values of G imply lower critical temperatures, so the energy necessary to break a Cooper pair increases.

In [55], it was seen that for the Einstein-Maxwell case, the universal value $\omega_g/T_c \approx 8$ holds for different values of the remaining parameters. It also holds in other settings, even when noncommutative effects of the underlying spacetime are present [27]. However, it was seen that when changing the electrodynamics or gravitational theories, the value of this ratio can change significantly [17, 56]. In Fig. 4 we plot it for different values of G and b . We see that for strong backreaction, the ratio can vary considerably from 8. Nonetheless, the ratio is still different from the value $\omega_g/T_c \approx 3.5$ predicted by the BCS theory [13, 15], confirming that this is still a superconductor with strong interactions.




FIG. 3. Plot of $\omega/\sqrt{\langle O_2 \rangle}$ vs $\text{Re}[\sigma]$ for different values of b and G at a fixed ratio $T/T_c = 0.9$. Dashed curves are for $b = 3$, while the continuous one are for $b = 7$. The values of G considered are, from left to right, $G = 0, 0.3, 0.7, 1.0$. Notice how ω_g , the frequency where $\text{Re}[\sigma]$ grows abruptly, is very sensible to the value of G .




FIG. 4. Plot of the ratio ω_g/T_c for different values of G and b , at fixed ratio $T/T_c = 0.9$. Dashed curves are for $b = 3$, and continuous curves, for $b = 7$. Large deviations from the universal value $\omega_g/T_c \approx 8$ seen in the Einstein-Maxwell case are observed. Notice that, like in previous figures, for G close to zero, the $b = 3$ curve is above the $b = 7$ one, but at a certain G between 0.6 and 0.7, the situation reverses.

IV. SUMMARY AND CONCLUDING REMARKS

In this work we have studied numerically a holographic superconductor in a Maxwell-Born-Infeld setting numerically, taking into account backreaction. Our results coincide qualitatively with previous results close to the probe limit [24, 35], showing that increasing the nonlinearity of the Born-Infeld theory difficulties the production of the superconducting condensate. However, we found that near the fully backreacted case, the situation reverses, an increase in the nonlinearity slightly enhances the production on the condensate. We found a similar behaviour when studying the AC conductivity of the system. An important point to be stressed is that backreaction modifies the so-called universal ratio $\omega_g/T_c \sim 8$, making it unstable.

The results reported here are similar to those in [27], where it was seen that in the probe limit, by introducing a noncommutative spacetime and tuning the respective noncommutativity parameter with the Born-Infeld parameter in a particular way, the formation of the condensate becomes easier, despite the fact that increasing noncommutativity tends to difficult the formation of the condensate [25, 26]. It is then interesting to make a numerical study similar to ours for the situation discussed in [27] to see how the backreaction changes these conclusions; this analysis, currently under way, will be reported elsewhere.

Along these remarks, another interesting line of investigation would be to consider the Hořava-Lifshitz or other theories of gravity instead of the one of Einstein. In some settings, it was found that Hořava-Lifshitz gravity difficulties the formation of the condensate [57, 58] while in others, it does not [59]. It would be interesting to gain a better understanding of higher derivatives

effects, for example, in holographic superconductors and see what happens in the backreacted case.

Finally, we could also study other observables of this superconductor, like its heat transport or its critical magnetic field [7], to pinpoint why nonlinearity changes from making the condensation harder to easier, when the backreaction parameter is increased. Such analysis will certainly broaden our knowledge of strongly correlated superconductors.

ACKNOWLEDGMENTS

E.G.-G. was funded by a CONAHCYT Scholarship with CVU 494426. U.E. is supported by Mexico's National Council of Humanities, Science and Technology (CONAHCYT) PhD grant.

AUTHOR CONTRIBUTIONS

All the authors contributed equally to this article.

Appendix A: Conductivity of a Born-Infeld holographic superconductor using the GKPW prescription

We recall that for a holographic superconductor based on the Maxwell Lagrangian $\mathcal{L}_M = -\frac{1}{16\pi}F_{\mu\nu}F^{\mu\nu}$, the relevant quantity used to calculate the conductivity is the following volume integral that is the contribution of \mathcal{L}_M that involves A'_x [11],

$$-\frac{1}{16\pi} \int_{r_+}^{\infty} dr \int d^3x \frac{r^2}{\sqrt{n}} \times \frac{n}{2fr^2} (2fA'_x + g_{tx}\phi')^2 \quad (\text{A1})$$

which is quadratic on the perturbations A_x and g_{tx} of the background gauge field and metric; the above integral appears in the regularisation procedure of the perturbed hairy black hole action on shell. After performing an integration by parts on the variable r , together with the asymptotic forms (7) and $A_x = A_x^0 + A_x^1/r + \dots$ to leading order, we get

$$\frac{1}{8\pi} \int d^3x A_x^0 A_x^1 \Big|_{r=\infty} + \dots \quad (\text{A2})$$

Here the \dots denote contributions independent of the gauge field. Variation of the above integral with respect to A_x^0 , together with $A_x^1 \sim A_x^0$, gives $J_x \equiv 4\pi(\delta S/\delta A_x^0) = A_x^1$; in consequence the conductivity is $\sigma \equiv \mathcal{J}_\mathcal{E}/\mathcal{E} = -iA_x^1/\omega A_x^0$ [11].

In the case of Born-Infeld electrodynamics defined by the Lagrangian density

$$\mathcal{L}_{BI} = 4b^2 \left(1 - \sqrt{1 + \frac{1}{2b^2} F_{\mu\nu} F^{\mu\nu}} \right), \quad (\text{A3})$$

the corresponding surface integral of interest is

$$\frac{1}{16\pi} \int_{r_+}^{\infty} dr \int d^3x \frac{r^2}{\sqrt{n}} \times \frac{n}{2r^2 f} \frac{(2fA'_x + g_{tx}\phi')^2}{\sqrt{1 - nb^{-2}\phi'^2}}, \quad (\text{A4})$$

up to second order on the perturbations A_x and g_{tx} ; the square root in the denominator of the above expression is characteristic of Born-Infeld nonlinear electrodynamics. Following the same procedure as for Maxwell electrodynamics and exploiting the fact that $\phi'|_{r=\infty} = 0$ holds as before, a straightforward calculation gives for the previous expression

$$\frac{1}{8\pi} \int d^3x A_x^0 A_x^1 \Big|_{r=\infty} + \dots \quad (\text{A5})$$

It follows that we also have $\sigma = -iA_x^1/\omega A_x^0$; this result is expected since far away from a source, Born-Infeld nonlinear electrodynamics behaves similar to Maxwell electrodynamics.

- [2] M. Natsuume, AdS/CFT duality user guide, *Lecture Notes in Physics* [10.1007/978-4-431-55441-7](#) (2015).
- [3] J. D. Edelstein, J. P. Shock, and D. Zoakos, The AdS/CFT correspondence and non-perturbative QCD, *Particles and fields: Proceedings of the XIII Mexican School of Particles and Fields*, 265 (2009).
- [4] S. A. Hartnoll, P. K. Kovtun, M. Müller, and S. Sachdev, Theory of the Nernst effect near quantum phase transitions in condensed matter and in dyonic black holes, *Physical Review B* **76**, 144502 (2007).
- [5] S. Sachdev, Condensed matter and AdS/CFT, *Lecture Notes in Physics*, 273–311 (2011).
- [6] J. McGreevy, Holographic duality with a view toward many-body physics, *Advances in High Energy Physics* **2010**, 1–54 (2010).
- [7] S. A. Hartnoll, Lectures on holographic methods for condensed matter physics, *Classical and Quantum Gravity* **26**, 224002 (2009).
- [8] J. Zaanen, Y.-W. Sun, Y. Liu, and K. Schalm, *Holographic Duality in Condensed Matter Physics* (Cambridge Univ. Press, 2015).
- [9] M. Ammon, J. Erdmenger, V. Grass, P. Kerner, and A. O’Bannon, On holographic p -wave superfluids with back-reaction, *Physics Letters B* **686**, 192–198 (2010).
- [10] S. A. Hartnoll, C. P. Herzog, and G. T. Horowitz, Building a holographic superconductor, *Physical Review Letters* **101**, 031601 (2008).
- [11] S. A. Hartnoll, C. P. Herzog, and G. T. Horowitz, Holographic superconductors, *Journal of High Energy Physics* **2008**, 015–015 (2008).
- [12] D. Tong, Holographic conductivity, *Acta Physica Polonica B* **44**, 2579 (2013).
- [13] C. P. Herzog, Lectures on holographic superfluidity and superconductivity, *Journal of Physics A: Mathematical and Theoretical* **42**, 343001 (2009).
- [14] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Microscopic theory of superconductivity, *Physical Review* **106**, 162–164 (1957).
- [15] M. Tinkham, *Introduction to superconductivity*, 2nd ed. (Dover Publication, 2004).
- [16] R. D. Parks, *Superconductivity*, 1st ed. (CRC Press, 1969).
- [17] R. Gregory, S. Kanno, and J. Soda, Holographic superconductors with higher curvature corrections, *Journal of High Energy Physics* **2009**, 010–010 (2009).
- [18] M. Taylor, Non-relativistic holography, *arXiv e-prints*, [arXiv:0812.0530](#) (2008).
- [19] K. Balasubramanian and J. McGreevy, Gravity duals for nonrelativistic conformal field theories, *Physical Review Letters* **101**, 061601 (2008).
- [20] E. J. Bagnasco, U. H. Danielsson, L. Thorlacius, and T. Zingg, Holographic superconductors with Lifshitz scaling, *Journal of Physics A: Mathematical and Theoretical* **43**, 065401 (2010).
- [21] D. Momeni, M. R. Setare, and N. Majd, Holographic superconductors in a model of non-relativistic gravity, *Journal of High Energy Physics* **2011**, 118 (2011).
- [22] M. Taylor, Lifshitz holography, *Classical and Quantum Gravity* **33**, 033001 (2016).
- [23] Q. Pan, J. Jing, and B. Wang, Holographic superconductor models with the Maxwell field strength corrections, *Physical Review D* **84**, 126020 (2011).
- [24] Z. Zhao, Q. Pan, S. Chen, and J. Jing, Notes on holographic superconductor models with the nonlinear electrodynamics, *Nuclear Physics B* **871**, 98–110 (2013).
- [25] D. Ghorai and S. Gangopadhyay, Noncommutative effects of spacetime on holographic superconductors, *Physics Letters B* **758**, 106–112 (2016).
- [26] S. Pramanik and S. Ghosh, AdS-CFT correspondence in noncommutative background, related thermodynamics and holographic superconductor in magnetic field, *General Relativity and Gravitation* **51**, 7 (2018).
- [27] M. Maceda and S. Patiño-López, Holographic superconductor from a noncommutative-inspired anti-de Sitter–Einstein–Born–Infeld black hole, *International Journal of Modern Physics D* **29**, 2050003 (2019).
- [28] G. Siopsis and J. Therrien, Analytic calculation of properties of holographic superconductors, *Journal of High Energy Physics* **2010**, 13 (2010).
- [29] L. Wang and F. He, Holographic superconductor in AdS3 spacetime, *arXiv e-prints*, [arXiv:2007.05926](#) (2020).
- [30] S. Gangopadhyay and D. Roychowdhury, Analytic study of properties of holographic superconductors in Born-Infeld electrodynamics, *Journal of High Energy Physics* **2012**, 2 (2012).
- [31] D. Roychowdhury, Effect of external magnetic field on holographic superconductors in presence of nonlinear corrections, *Physical Review D* **86**, 106009 (2012).
- [32] M. Born and L. Infeld, Foundations of the new field theory, *Proceedings of the Royal Society of London. Series A* **144**, 425–451 (1934).
- [33] R. Kerner, Topics in Born-Infeld electrodynamics, *AIP Conference Proceedings*, 377 (2001).
- [34] J. Jing and S. Chen, Holographic superconductors in the Born–Infeld electrodynamics, *Physics Letters B* **686**, 68–71 (2010).
- [35] S. I. Kruglov, Holographic superconductor with nonlinear Born–Infeld-type electrodynamics, *International Journal of Modern Physics A* **34**, 1950019 (2019).
- [36] M. Mohammadi, A. Sheykhi, and M. K. Zangeneh, Analytical and numerical study of backreacting one-dimensional holographic superconductors in the presence of Born–Infeld electrodynamics, *The European Physical Journal C* **78**, 654 (2018).
- [37] Y. Liu, Y. Peng, and B. Wang, Gauss-Bonnet holographic superconductors in Born-Infeld electrodynamics with backreactions, *arXiv e-prints*, [arxiv:1202.3586](#) (2012).
- [38] A. Sheykhi and F. Shaker, Analytical study of holographic superconductor in Born–Infeld electrodynamics with backreaction, *Physics Letters B* **754**, 281–287 (2016).
- [39] D. Ghorai and S. Gangopadhyay, Conductivity of holographic superconductors in Born–Infeld electrodynamics, *Nuclear Physics B* **933**, 1–13 (2018).
- [40] Y. Brihaye and B. Hartmann, Holographic superconductors in 3+1 dimensions away from the probe limit, *Physical Review D* **81**, 126008 (2010).
- [41] Y. Liu, Q. Pan, and B. Wang, Holographic superconductor developed in BTZ black hole background with backreactions, *Physics Letters B* **702**, 94–99 (2011).
- [42] D. Momeni, M. Raza, M. R. Setare, and R. Myrzakulov, Analytical holographic superconductor with backreaction using AdS 3/CFT 2,

[International Journal of Theoretical Physics](#) **52**, 2773–2783 (2013).

- [43] Y. Peng and Q.-Y. Pan, Stückelberg holographic superconductor models with backreactions, [Communications in Theoretical Physics](#) **59**, 110 (2013).
- [44] W. Yao and J. Jing, Analytical study on holographic superconductors for Born-Infeld electrodynamics in Gauss-Bonnet gravity with backreactions, [Journal of High Energy Physics](#) **2013**, 101 (2013).
- [45] P. Chaturvedi and G. Sengupta, p -wave holographic superconductors from Born-Infeld black holes, [Journal of High Energy Physics](#) **2015**, 1 (2015).
- [46] I. R. Klebanov and E. Witten, AdS/CFT correspondence and symmetry breaking, [Nuclear Physics B](#) **556**, 89–114 (1999).
- [47] D. Musso, Introductory notes on holographic superconductors, [arXiv e-prints](#), [arXiv:1401.1504](#) (2014).
- [48] D. T. Son and A. O. Starinets, Minkowski-space correlators in AdS/CFT correspondence: recipe and applications, [Journal of High Energy Physics](#) **2002**, 042–042 (2002).
- [49] S. Gubser, I. Klebanov, and A. Polyakov, Gauge theory correlators from non-critical string theory, [Physics Letters B](#) **428**, 105–114 (1998).
- [50] A. Donos, J. P. Gauntlett, T. Griffin, and L. Melgar, DC conductivity and higher derivative gravity, [Classical and Quantum Gravity](#) **34**, 135015 (2017).
- [51] K. Maeda, M. Natsuume, and T. Okamura, Universality class of holographic superconductors, [Physical Review D](#) **79**, 126004 (2009).
- [52] D. C. Mattis and J. Bardeen, Theory of the anomalous skin effect in normal and superconducting metals, [Physical Review](#) **111**, 412–417 (1958).
- [53] H. Chen, Theory of optical conductivity in BCS superconductors, [Physical Review Letters](#) **71**, 2304–2306 (1993).
- [54] C. P. Herzog, P. Kovtun, S. Sachdev, and D. T. Son, Quantum critical transport, duality, and M theory, [Physical Review D](#) **75**, 085020 (2007).
- [55] G. T. Horowitz and M. M. Roberts, Holographic superconductors with various condensates, [Physical Review D](#) **78**, 126008 (2008).
- [56] A. Sheykhi, D. Hashemi Asl, and A. Dehyadegari, Conductivity of higher dimensional holographic superconductors with nonlinear electrodynamics, [Physics Letters B](#) **781**, 139–154 (2018).
- [57] K. Lin, E. Abdalla, and A. Wang, Holographic superconductors in Hořava–Lifshitz gravity, [International Journal of Modern Physics D](#) **24**, 1550038 (2015).
- [58] J.-W. Lu, Y.-B. Wu, J. Xiao, C.-J. Lu, and M.-L. Liu, Holographic superconductors in IR modified Hořava–Lifshitz gravity, [International Journal of Modern Physics A](#) **31**, 1650110 (2016).
- [59] K. Lin, X.-M. Kuang, W.-L. Qian, Q. Pan, and A. B. Pavan, Analysis of s -wave, p -wave and d -wave holographic superconductors in Hořava–Lifshitz gravity, [Modern Physics Letters A](#) **33**, 1850147 (2018).