

Detecting unresolved lensed SNe Ia in LSST using blended light curves

Satadru Bag^{1,2*}, Simon Huber^{1,2}, Sherry H. Suyu^{1,2,3}, Nikki Arendse⁴, Irham Taufik Andika^{1,2}, Raoul Cañameras^{2,1,5}, Alex Kim⁶, Eric Linder^{6,7}, Kushal Lodha^{8,9}, Alejandra Melo^{1,2}, Anupreeta More^{10,11}, Stefan Schuldt^{12,13} and Arman Shafieloo^{8,9}

- ¹ Technical University of Munich, TUM School of Natural Sciences, Physics Department, James-Frank-Straße 1, 85748 Garching, Germany
- ² Max-Planck-Institut für Astrophysik, Karl-Schwarzschild Straße 1, 85748 Garching, Germany
- ³ Academia Sinica Institute of Astronomy and Astrophysics (ASIAA), 11F of ASMA, No. 1, Section 4, Roosevelt Road, Taipei 10617, Taiwan
- ⁴ Oskar Klein Centre, Department of Physics, Stockholm University, SE-106 91 Stockholm, Sweden
- ⁵ Aix Marseille Univ, CNRS, CNES, LAM, Marseille, France
- ⁶ Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA 94720, USA
- ⁷ Berkeley Center for Cosmological Physics, University of California, Berkeley, CA 94720, USA
- ⁸ Korea Astronomy and Space Science Institute (KASI), 776 Daedeok-daero, Yuseong-gu, Daejeon 34055, Korea
- ⁹ KASI Campus, University of Science and Technology, 217 Gajeong-ro, Yuseong-gu, Daejeon 34113, Korea
- ¹⁰ Inter-University Centre for Astronomy and Astrophysics, Post Bag 4, Ganeshkhind, Pune 411007, India
- ¹¹ Kavli Institute for the Physics and Mathematics of the Universe (IPMU), 5-1-5 Kashiwanoha, Kashiwa-shi, Chiba 277-8583, Japan
- ¹² Dipartimento di Fisica, Università degli Studi di Milano, via Celoria 16, I-20133 Milano, Italy
- ¹³ INAF - IASF Milano, via A. Corti 12, I-20133 Milano, Italy

Preprint online version: April 25, 2024

ABSTRACT

Strong-gravitationally lensed supernovae (LSNe) are promising probes for providing absolute distance measurements using gravitational lens time delays. Spatially unresolved LSNe offer an opportunity to enhance the sample size for precision cosmology. We predict that there will be approximately 3 times more unresolved than resolved LSNe Ia in the Legacy Survey of Space and Time (LSST) by the Rubin Observatory. In this article, we explore the feasibility of detecting unresolved LSNe Ia from the shape of the observed blended light curves using deep learning techniques, and we find that $\sim 30\%$ can be detected with a simple 1D CNN using well-sampled *rizy*-band light curves (with a false-positive rate of $\sim 3\%$). Even when the light curve is well-observed in only a single band among *r*, *i*, and *z*, detection is still possible with false-positive rates ranging from $\sim 4 - 7\%$, depending on the band. Furthermore, we demonstrate that these unresolved cases can be detected at an early stage using light curves up to ~ 20 days from the first observation, with well-controlled false-positive rates, providing ample opportunities for triggering follow-up observations. Additionally, we demonstrate the feasibility of time-delay estimations using solely LSST-like data of unresolved light curves, particularly for doubles, when excluding systems with low time delay and magnification ratio. However, the abundance of such systems among those unresolved in LSST poses a significant challenge. This approach holds potential utility for upcoming wide-field surveys, and overall results could significantly improve with enhanced cadence and depth in the future surveys.

Key words. Gravitational lensing: strong, micro – methods: data analysis – supernovae: Type Ia supernova

1. Introduction

Gravitationally lensed transients, such as quasars (QSOs) and supernovae (SNe), have emerged as novel powerful cosmic probes capable of delivering cosmological information independently of other probes (see, e.g., Treu et al. (2022) for a review). For instance, one can estimate the Hubble constant (H_0) using lensed transients without relying on additional datasets (Refsdal 1964; Refsdal & Bondi 1964; Saha et al. 2006; Oguri 2007; Bonvin et al. 2017; Wong et al. 2020; Birrer et al. 2020; Birrer & Treu 2021; Kelly et al. 2023; Grillo et al. 2024; Pascale et al. 2024). Importantly, the value of H_0 remains fairly insensitive to the chosen

cosmological model. This capability can address the significant tension observed in H_0 calculations between early universe probes (inverse distance ladder, Aghanim et al. 2020) and late universe probes (distance ladder, Riess et al. 2022). With physically motivated lens mass models, the H0LiCOW team has further demonstrated that H_0 can be tightly constrained by combining constraints from multiple strongly lensed QSOs (Wong et al. 2020), thus underscoring the importance of a large sample size.

Among variable sources, QSOs and SNe are currently the most prominent types, each possessing distinct advantages. For instance, lensed QSOs are more abundant and do not fade away, unlike SNe, making them more reliable for time-delay cosmography at present. On the other hand,

* satadru.bag@tum.de

lensed SNe (LSNe) are poised to take the forefront in the coming decade due to better understanding of their light curves, feasibility of improved stellar kinematic and SN host galaxy measurements after the SN fades away (Ding et al. 2021; Suyu et al. 2024). Additionally, type Ia SNe serve as standard candles, offering direct luminosity distances to the sources, provided that magnification effects of microlensing (lensing by stars in the foreground lens galaxy) are mitigated (Foxley-Marrable et al. 2018; Weisenbach et al. 2021). However, a significant challenge stems from the rarity of LSNe, with only a handful confirmed detection to date (Kelly et al. 2015; Goobar et al. 2017; More et al. 2017; Rodney et al. 2021; Chen et al. 2022; Goobar et al. 2023; Frye et al. 2023; Pierel et al. 2024). Furthermore, many LSNe will remain unresolved in the wide-field surveys due to their angular resolution being limited by seeing constraints (Quimby et al. 2014; Goldstein et al. 2018). Detecting these unresolved LSNe could notably enhance the sample size, thereby aiding in achieving precision cosmology. Therefore, the importance of detecting unresolved LSNe in time-domain wide-field surveys, such as the Legacy Survey of Space and Time (LSST) by the Vera C. Rubin Observatory, cannot be overstated.

Various techniques, such as the magnification method (Goldstein & Nugent 2017; Goobar et al. 2017; Wojtak et al. 2019), colour-magnitude diagram (Quimby et al. 2014; Arendse et al. 2023), and shifting centroid (Kodi Ramanah et al. 2022), can detect unresolved lensed SNe from photometry. Since the observed light curve of an unresolved LSN will be the sum of the individual image light curves, the former contains distinct features in its shape and can also be used to discern the lensed systems from the unlensed ones. This constitutes the central focus of our work.¹ We developed several methods based on forward modeling for this purpose (Bag et al. 2021; Denissenya et al. 2021). In this article, inspired by Denissenya & Linder (2022), we employ a simple 1D convolutional neural network (CNN) that leverages the shape of the light curves to detect lensed cases. We apply the CNNs on light curves observations consistent with LSST by Vera C. Rubin Observatory. LSST, which is deeper than the ongoing Zwicky Transient Facility (ZTF) survey (Bellm et al. 2019), is expected to observe approximately 20,000 square degrees over a 10-year span (LSST Science Collaboration et al. 2009, 2017; The LSST Dark Energy Science Collaboration et al. 2018), thereby making the potential for discovering new unresolved LSNe notably significant.

Our primary emphasis is on Type Ia supernovae (SNe Ia), given their pivotal role in cosmology as standardizable candles. We simulate mock LSST light curves using the `baseline v3.2` observing strategy. The sample comprises joint light curves of unresolved lensed SNe Ia as positive examples and unlensed (i.e. intrinsic) SNe Ia light curves as negative examples. We employ two types of CNNs: one for detection and the other intended to estimate time delays directly from the joint light curves in cases when a follow up observation is not possible or conclusive. It's worth noting that this approach relies on the precise photometric clas-

sification of SNe Ia from other transients, e.g., using the algorithms proposed by Lochner et al. (2016); Alves et al. (2022), prior to applying.

The article is organized as follows. In Section 2, we define the unresolved cases in LSST and compare their forecasted numbers and properties against the resolved ones. Next, in Section 3, we explain the process of simulating mock LSST light curves. Section 4 presents a brief overview of the method and data processing. Our main results are explained in Section 5. Finally, in Section 6, we conclude and discuss further scopes in this direction.

2. Unresolved SNe Ia in LSST

2.1. Unresolved light curves

The blended joint light curve of an unresolved lensed SN with N images is the sum of the individual image fluxes (neglecting microlensing for the moment),

$$F(t) = \sum_{i \in \text{images}}^N |\tilde{\mu}_i| \mathcal{F}(t - \tilde{t}_i), \quad (1)$$

where $\tilde{\mu}_i$ and \tilde{t}_i are the magnification (which can be positive or negative depending upon the parity of an image) and time delay of the i th image. Note that the light curve of all the images can be described by the same function $\mathcal{F}(t)$ (the unlensed light curve; here we ignored microlensing and differential dust extinction for simplicity) but with different magnifications and time delays. It is useful to recast the above expression in terms of the image that arrives the earliest which we refer to as the ‘first’ image,

$$F(t) = \sum_i^N \mu_i f(t - \Delta t_i), \quad (2)$$

where $f(t) = |\tilde{\mu}_1| \mathcal{F}(t - \tilde{t}_1)$ is the light curve of the first image and

$$\mu_i \equiv |\tilde{\mu}_i / \tilde{\mu}_1| \quad \text{and} \quad \Delta t_i \equiv \tilde{t}_i - \tilde{t}_1 \quad (3)$$

represent the magnification ratio and time delay of the i -th image with respect to the first image. By definition Δt_i , $\mu_i \geq 0$ for all the images and naturally, $\mu_1 = 1$ and $\Delta t_1 = 0$.

In this work, we only consider the light curve information and aim for detecting if the system is lensed or not, i.e. if the observed light curve stems from the blending of multiple images or corresponds to an unlensed SN. It is important to note that the task becomes challenging when dealing with extreme values of μ or small values of Δt . To understand this issue let us first consider the double systems for simplicity. When the magnification ratio deviates significantly from unity (with Δt not being too high), one image dominates over the other – $\mu_2 \ll 1$ ($\mu_2 \gg 1$) implies that the second (first) image has negligible brightness compared to the first (second) image. On the other hand, if $\Delta t_2 \rightarrow 0$, the two images coincide with each other. Therefore, in both of these conditions, it is mathematically impossible to detect the lensing nature in the unresolved light curve. This can be easily generalised for quads; if two images have extreme magnification ratios or very small time delay, then they cannot be discerned as two images from the unresolved light curve. In simpler terms, to detect lensed systems, it

¹ Recently, similar efforts have been undertaken to detect unresolved lensed QSOs from light curve observations (Shu et al. 2021; Bag et al. 2022; Bag et al. 2023). However, it's important to note that the techniques employed are entirely different due to the distinct nature of QSO light curves.




Fig. 1: Number of unresolved lensed SNe Ia (blue) is compared with the number of resolved cases (orange) for one cumulative year of LSST observation at different critical angular resolution (θ_{crit}) above which the images can be resolved. The solid curves are calculated assuming the macro magnifications only whereas the dashed curves consider the microlensing effects. LSST can resolve images up to $\theta_{\text{th}} \approx 0.5 \text{ arcsec} \sim 2/3\theta_{\text{psf}}$. For this threshold, we obtain approximately 71.1 unresolved (blue dot) and 27.8 resolved (orange dot) lensed SNe Ia based solely on macro magnification calculation, indicating roughly 2.6 times more unresolved systems. The ratio increases to approximately 3.2 when we account for microlensing, as evident from the dashed curves. We emphasize that these numbers correspond to one effective year of LSST observation over 20,000 deg^2 in i -band. LSST will observe a sky location cumulatively for ~ 4 years on the average during its 10 year long run. Therefore, it is essential to scale these figures by a factor of ~ 4 to get the realistic predictions.

is essential to have at least one pair of images (which is the only pair in the case of a double) with $\mu \sim \mathcal{O}(1)$ and $\Delta t > \text{a few days}$. Furthermore, to differentiate quads from doubles, multiple such pairs of images are required.

2.2. OM10 simulation for LSST

In this work, we concentrate on samples of type Ia SNe due to their significant cosmological utility, well-understood light curves, and ease of simulation. Using the Oguri and Marshall 2010 simulations (OM10 hereafter, [Oguri & Marshall \(2010\)](#)) with minor updates ([Oguri 2018](#)), we forecast the number of lensed type Ia supernovae (LSNe Ia) that will remain unresolved in LSST. A key observation from Equation (1) is that (lensed but) unresolved sources will appear brighter than its individual images (if they were resolved) because light from different images can be accumulated for the unresolved cases. This, in turn, suggests that unresolved lensed sources could be visible at greater luminosity distances (or redshifts). Our primary focus is on the LSST i -band since LSNe Ia are at such redshifts that they are usually observed with highest signal-to-noise ratio (S/N) in this filter ([Oguri & Marshall 2010](#); [Huber et al. 2022](#)). Additionally, the LSST i -band presumably offers a more regular cadence, less affected by the moon phase, and typically provides better seeing conditions com-

pared to shorter wavelengths. In the simulation, we allow all image separations and source brightness up to 30 mag in i -band to consider all possible unresolved lensed SNe Ia that could be observed in LSST.

We define the unresolved lensed SNe, detectable in LSST, with the following criteria: (i) image separation has to be smaller than a critical value above which one can resolve the images: $\theta < \theta_{\text{crit}}$ and (ii) peak brightness (of the joint light curve) in i -band has to be brighter than the critical magnitude, m_{crit} . The 5σ depth (m_5) of LSST i -band is 23.3. However, we need the SN to be brighter than this limiting magnitude in order to have sufficiently good quality light curve observations that is essential for the time-delay measurements with adequate accuracy. Therefore, we set the critical magnitude to be 0.7 magnitude brighter (i.e. lower) than the limiting magnitude ([Oguri & Marshall 2010](#); [Dimitriadis et al. 2019](#)), i.e. $m_{\text{crit}} = 22.6$.

On the other hand, we define the resolved systems which have at least one pair of images with angular separation larger than the critical value, $\theta > \theta_{\text{crit}}$ and both images are brighter than $m_{\text{crit}} = 22.6$ ². Figure 1 illustrates how the number of resolved (orange) and unresolved (blue) lensed SNe Ia varies with the critical angular separation (θ_{crit}) for one cumulative year of LSST observation. Therefore, this figure presents the theoretical prediction of number of resolved and unresolved LSNe Ia expected to occur within a year (irrespective of the observing strategy) across the LSST footprint of 20,000 deg^2 . Not all of these events will be observed due to seasonal gaps. Over its 10-year operation, LSST will, on average, cumulatively monitor each sky location for approximately 4 years. Therefore, for a realistic prediction, it is necessary to scale all the curves by a factor of approximately ~ 4 .

The solid curves in Figure 1 assume only the macro magnifications whereas the dashed curves are obtained by considering (rough) microlensing effects in the light curves³. For this range of θ_{crit} , we always get a few times more unresolved systems than resolved ones. Note that the sum of the number of unresolved and resolved cases is not constant, as the latter must meet stricter criteria. Specifically, for resolved cases, at least one pair of lensed images must be brighter than m_{crit} , while for unresolved cases, it's the combined flux that has to be brighter. In other words, when we lower θ_{crit} by a certain amount, some of the previously unresolved systems will now have $\theta > \theta_{\text{crit}}$ and the number of unresolved cases will decrease. But all of them won't be considered as resolved resulting in smaller growth in the resolved case number.

The angular resolution of wide field surveys, such as LSST, is limited by seeing. For LSST, [Oguri & Marshall](#)

² To have an apples to apples comparison, we don't care if it's a double or a quad at this moment; this can be established by a follow up.

³ OM10 simulation for 10 years produces too many lensed SNe Ia for it to be computationally feasible to calculate microlensing for all. Therefore, we take a representative subset of systems to generate the microlensing magnification maps, and for each system in the OM10, we pick the most similar system in this representative subset to approximate the microlensing effect. For each system, we overpopulate it by a factor of 100 and take 100 random positions on its corresponding microlensing magnification maps to scale the fluxes of the multiple images in the system. After repeating this procedure for all systems in OM10, we renormalize by the factor of 100.

(2010) adopts angular resolution $\theta_{\text{th}} = 0.5$ arcsec, which is roughly $2/3$ of the median $\theta_{\text{psf}} \approx 0.7$ arcsec, as the minimum separation that could be resolved in LSST. Following this, we also set $\theta_{\text{crit}} = \theta_{\text{th}} = 0.5$ arcsec for our estimations of resolved and unresolved cases. Final definitions of unresolved and resolved lensed SNe Ia are given below,

- **resolved cases:** if any pair of images have (i) angular separation $\theta > 0.5$ arcsec, and (ii) i -band magnitude $m < 22.6$ for each of the two images,
- **unresolved cases:** if the angular separation between all the pairs is below 0.5 arcsec and (ii) the combined blended light curve has peak brightness $m < 22.6$ in i -band.

Let us first focus on the macro lensing based calculation; we will get back to the estimations considering microlensing shortly. Following the above definitions, we get 27.8 resolved and 71.1 unresolved lensed SNe Ia per cumulative year of LSST observation – these numbers are marked by filled circles on the respective solid curves in Figure 1. Therefore, we expect that ~ 2.6 times more unresolved lensed SNe Ia will be seen in LSST than resolved lensed cases. Therefore, detection of these unresolved cases is an important task.

In addition to introducing fluctuations in light curves, microlensing also either suppresses or boosts the brightness of images, while maintaining the same ensemble average as the macro lensing magnification. Intriguingly, microlensing tends to suppress brightness more frequently than boosting (Goldstein et al. 2018; Arendse et al. 2023), resulting in fewer systems being brighter than the critical magnitude (m_{crit}) as compared to the estimations based on solely macro magnification. This observation is evident as the dashed curves consistently lie below their macro-lensed-only counterparts. However, since unresolved sources are permitted to be fainter than resolved sources (while both still need to be brighter than the limiting magnitude), the impact of microlensing effects on the number of unresolved lensed systems is less pronounced. In other words, summing up light curves of multiple images, each affected differently by microlensing, reduces the impact of microlensing on the peak brightness of unresolved cases. Thus the ratio of unresolved to resolved cases is further enhanced when one considers microlensing. E.g., with $\theta_{\text{crit}} = 0.5$, we get approximately 21.5 resolved and 68.2 unresolved systems per cumulative year, boosting the ratio to ~ 3.2 . However, while exploring the comparison of resolved vs unresolved systems in terms of multiple statistics we only stick to systems obtained using macro magnification only. Note that in the upcoming sections, where we aim to detect unresolved systems, we incorporate comprehensive microlensing effects into the light curves.

Now, let us go back to the forecast with macro-magnification-only based calculation that predicts 27.8 resolved and 71.1 unresolved lensed SNe Ia systems per effective year of LSST observation. Figure 2 compares the 95% CLs of the (κ, γ) distributions for the unresolved (blue contour) and resolved samples (orange contour). Note that the requirement of lower θ for the unresolved cases does not affect the (κ, γ) distribution as compared to that of the resolved cases. However, since unresolved sources are allowed to be fainter, they can accommodate lower magnifications and thus their (κ, γ) distribution extends towards higher values. We need to use this distribution when simulating

the microlensed light curves in the next section. We present the source and lens redshift distributions of the unresolved systems in Appendix A, comparing them against those of the resolved cases.

Next, we compare the $\mu - \Delta t$ distribution for the resolved and unresolved cases in Figure 3. We remind the reader that we always sort the images chronologically and define the magnification ratios and time delays of the images with respect to the reference image which arrives first in time (i.e. the first image). Consequently, a double system possesses only one set of μ and Δt , while a quad system entails three sets of μ and Δt . However, in the case of quads, the image with the largest time delay holds most significant importance in the detection process. Therefore, we exclusively plot the μ and Δt of the largest time-delay image for each quad system in Figure 3.

The left panel of Figure 3 illustrates the 68% and 95% CLs of the $\mu - \Delta t$ distribution. In the middle panel, 1D histogram of μ (specifically for the image with the largest time delay relative to the first image for quads) is presented. This histogram reveals a median μ of 0.40, with a range of (0.17, 0.61) at the 68% percentile and (0.11, 2.46) at the 95% percentile. Given that the first image, which acts as our reference, is usually the brightest for doubles and brighter than the last appearing image for quads, the majority of the μ values fall below unity. The right panel displays the cumulative time-delay distribution, indicating how many systems have the largest time delay above a given t and comparing it for resolved (orange) and unresolved (blue) samples. (We emphasize that the number of systems presented here corresponds to one cumulative year of observation, ensuring a fair and straightforward comparison between resolved and unresolved systems.)

As anticipated, unresolved systems generally exhibit smaller time delays compared to their resolved counterparts, rendering them less suitable for time-delay cosmography. For the unresolved systems, the median gives $\Delta t_{\text{max}} = 2.03$ days. Nevertheless, if a mechanism can be devised to detect the majority of unresolved systems, regardless of their small time delays, their sheer abundance could contribute significantly to achieving precision cosmology. Although detecting these low time-delay systems solely through light curve data is challenging, this article aims to provide a method to accomplish that.

3. Simulating mock blended joint light curves

In this section, we detail the process of constructing the training, validation, and test sets comprising of light curves of unresolved LSNe Ia, consistent with the OM10 simulations, as well as unlensed SNe Ia light curves, across multiple bands.

3.1. Microlensed light curves

In the previous section, we neglected microlensing when exploring the statistics of the forecasted unresolved lensed SNe Ia samples. Microlensing can affect the lensed light curves of individual images differently, and these effects may also differ across filters⁴. Therefore, we must take microlensing into the account when training the models.

⁴ However, multiple studies (e.g. Goldstein et al. 2018; Huber et al. 2021) have shown that there is an achromatic phase in the




Fig. 2: The 95% confidence level of the (κ, γ) distribution is compared for the images of resolved systems (orange contour) and unresolved systems (blue contour). The greyscale represents the macro magnification in log-scale, clipped between $[0.1, 10]$. Notably, the (κ, γ) distribution for the unresolved systems extends towards higher values (compared to that of the resolved systems), accommodating lower magnifications. The crosses mark the (κ, γ) values of the images used to simulate the microlensed light curves, with green and red crosses distinguishing images in doubles and quads, respectively. The diagonal dashed line denotes the $\kappa = \gamma$ line.

To calculate microlensed light curves of lensed SNe Ia, we need (i) specific intensity profiles, which we get from theoretical SN Ia models, and (ii) microlensing magnification maps. Details on the calculation of microlensed light curves are presented in Huber et al. (2019). We use as in Huber et al. (2019); Suyu et al. (2020); Huber et al. (2021, 2022) theoretical SN Ia models from ARTIS (Kromer & Sim 2009) and magnification maps from Gerlumph (Vernardos & Fluke 2014; Vernardos et al. 2014, 2015; Chan et al. 2021). For the microlensing maps we assume as in Huber et al. (2022) and Huber & Suyu (2024) a Salpeter initial mass function (IMF) with a mean mass of the microlenses of $0.35M_{\odot}$ and a resolution of 20000×20000 pixels with a total size of $20 R_{\text{ein}} \times 20 R_{\text{ein}}$ on the source plane.

Another important factor for generating microlensing maps is the smooth matter fraction ($s \in [0, 1]$) that determines what fraction of the total matter is smooth while the rest is in the form of stars at an image location; lower s would lead to more severe microlensing modifications to the macro lensed light curve of an image. The saddle images, that often lie closer to the lens centres especially for doubles, tend to have lower smooth matter fractions. Conversely, the minima images often lie further away from the lens centres and thus typically have higher values of s . Therefore, to maintain a balance between realism and simplicity, we assign $s = 0.4$ for all the saddle images and randomly choose either 0.6 or 0.8 for the minima images.

Finally, we simulate the microlensed light curves for a total of 250 images, stemming from randomly selected 49 doubles and 38 quads. The κ, γ values of these 250 images are shown by cross marks in Figure 2; with green

and red crosses indicating images of doubles and quads, respectively. It is evident that they fairly cover the (κ, γ) distribution of all unresolved systems. For a given image, we generate a microlensing map using its (κ, γ, s) values. Subsequently, we simulate 100 microlensing realizations corresponding to 100 random locations on the microlensing map. The selection of SN Ia model can also play an important role. To maintain a model-agnostic approach, we should incorporate a mix of different SN Ia models during the training process. Among the four theoretical models investigated in Huber et al. (2022), the merger model (Pakmor et al. 2011, 2012) exhibits the most diverse behavior compared to the other three, namely W7 (carbon deflagration, Nomoto et al. (1984)), N100 (delayed detonation, Röpke et al. (2012); Seitenzahl et al. (2013)), and sub-Ch (sub-Chandrasekhar mass detonation, Sim et al. (2010)). For simplicity, while still retaining diversity in the SN Ia models, we opt to randomly choose between the W7 and merger models when simulating the light curves for each system.

When we construct the blended light curve of a lensed unresolved SN, we randomly select a system from the corresponding sample (double or quad) with microlensing simulations available. We then randomly choose one microlensing realisation for each of the images and sum up the microlensed light curves, separately in each filter.

3.2. Unlensed light curves

While simulating the light curves for an unlensed SN Ia, we randomly select one of the lensed systems for which the simulated light curves are available. Subsequently, we obtain the intrinsic light curve by dividing the macrolensed light curve of any of the images by the corresponding macro magnification.

3.3. LSST cadence distribution and noise implementation

We implement a realistic LSST cadence using publicly available simulations from the Rubin Operations Simulator (OpSim), which emulates the field selection and image acquisition process of LSST over its 10-year duration (Delgado et al. 2014; Delgado & Reuter 2016; Naghib et al. 2019). The main survey mode of LSST is the Wide-Fast-Deep (WFD) programme, which will cover around 18,000 square degrees, corresponding to $\sim 90\%$ of observing time. Following the latest recommendations from the Survey Cadence Optimization Committee⁵, the WFD survey will employ a rolling cadence, in which alternating areas of the sky will receive more frequent visits than others. As a result, objects discovered in the high-cadence region will have more observations and, hence, a better light curve coverage. In our simulations, we consider lensed SNe Ia in the WFD region as well as in the remaining sky regions: the Galactic plane, polar regions and the Deep Drilling Fields, encompassing a total footprint of 20,000 square degrees. We implement the **baseline v3.2** observing strategy to simulate our mock light curves. Among the six LSST filters, we consider only ‘ r ’, ‘ i ’, ‘ z ’ and ‘ y ’ bands, as the remaining two, namely ‘ u ’ and ‘ g ’ bands, have much “lower” cadence.

To sample the (joint) light curve according to the observing strategy, we randomly choose 7, 110, 000 sky loca-

microlensing effects up to roughly three rest-frame weeks from the explosion.

⁵ <https://pstn-055.lsst.io/>




Fig. 3: Distribution of the largest time delay (Δt_{\max}) and the corresponding magnification ratio in all the unresolved systems is shown in the left panel; the lighter and darker shades of blue show the 68% and 95% credible regions respectively. The middle panel presents the 1D marginalised distribution of the magnification ratio. The right panel shows the cumulative distribution of Δt_{\max} , i.e. how many systems (in one cumulative year) have Δt_{\max} higher than a given value – resolved (orange) vs unresolved (blue). It is evident that unresolved systems typically exhibit low time delays, with a median value of Δt_{\max} being ~ 2.03 days and approximately 10% having values above 10 days.

bands	systems individually well-observed in 4 bands (4 sets)				systems well-observed in all 4 bands (intersection of the 4 sets)			
	r	i	z	y	r	i	z	y
good systems	13.0%	14.0%	12.2%	9.8%	6.2 %			
avg no. of epochs	21.4	21.6	20.8	19.9	24.9	26.1	24.2	22.3
cadence								
mean & median (days)	7.3 & 4.0	7.6 & 5.0	8.4 & 5.7	11.6 & 5.8	6.7 & 4.0	6.7 & 4.8	7.6 & 5.0	10.8 & 5.0
largest gap								
mean & median (days)	28.3 & 24.9	27.9 & 23.9	29.1 & 25.8	33.3 & 28.9	24.5 & 23.0	23.9 & 21.9	25.9 & 23.9	31.6 & 28.8
mean m_5	23.9	23.3	22.7	21.9	same as in the individual 4 sets corresponding to 4 bands			

Table 1: Statistics from observing strategy **baseline v3.2** in the four LSST filters – r , i , z , and y . We have a set of ‘good systems’ that meet the criteria for ‘good observation’, as described in Section 3.3, for each of these four filters. The table is divided into two parts, each with four columns. The left part presents various statistics, derived from the observing strategy, considering these four sets of systems well-observed in the four bands. The right half presents the same statistics but for the systems well-observed in all four bands, i.e., for the intersection of the four sets. For example, the left part of the second row shows the fraction of unresolved systems that would be well-observed in different bands, whereas the right part indicates that 6.2% of the systems will be well-observed in all four bands. Note that, in comparison, $\sim 40\%$ of the total resolved systems will be detected due to the season gaps in baseline observing strategy (Huber et al. 2019). The next three rows present the average number of observation epochs, mean & median of cadence and the largest gap in it, respectively, in the four bands considering the four sets of systems individually in the left part and their intersection in the right part. The last row provides the mean of 5σ depth (m_5) in the four bands; these numbers should remain the same for the intersection part as well.

tions (which corresponds to an oversampling of 100,000 times the number of unresolved SNe per cumulative year of LSST observation) within the LSST footprint. We use **OpSimSummary** (Biswas et al. 2020) to retrieve all observation times, filters, and 5σ depths corresponding to a specific sky location within the field of view of the telescope. Now for each sky location we randomly choose a time within the 10 year run of LSST that represents the epoch of SN explosion. Many of these SNe will remain unobserved due to season gaps and some will be observed only for a few epochs leading to partially monitored light curves. To filter out the suitable systems that can have light curves observed for sufficient epochs we apply the following criteria,

1. no. of observation > 10 within 120 days from the explosion,
2. the first observation has to be within 20 days from explosion,
3. there has to be at least one observation after 40 days from explosion.

When a SN satisfies the above criteria in a filter we consider that the system has ‘good’ observation in that particular filter. Therefore, we have four sets of well-observed systems in the four LSST filters that we consider – r , i , z and y . Various statistics for these sets, derived from the observation strategy, are presented in Table 1 in two parts (each with four columns corresponding to four bands). The left part considers the 4 sets of good systems individually

whereas the right part presents the same statistics but for the subset of the systems that are well-observed in all four bands, i.e. for the intersection of the 4 sets. For example, the second row presents what percentage of the unresolved systems will be well-observed in different bands in the left part whereas the right part informs that 6.2% will be well-observed in all four bands. The union of good systems covers 18.8% of the total sample. The third row shows the mean no. of observed epochs (till 120 days from the explosion) in different bands. The next two rows present the distributions of the cadence and the largest gap in observation epochs, respectively, in terms of mean and median of the respective quantities. The last row indicates the mean 5σ depth in different bands, these numbers should remain the same for the intersection set as well.

All in all, we find that r and i bands provide better cadence distributions and smaller maximum gaps in observational epochs, as compared to the other two remaining bands. Therefore, the number of ‘good systems’ is higher in these two (r and i) bands. Fortunately, SNe Ia exhibit higher S/N in these two bands due to their higher values of 5σ depth (m_5), making them ideal for this study.

3.4. Uncertainties

Baseline v3.2 observing strategy provides the 5σ depth (m_5) in each filter at every epoch. The bottom row of Table 1 presents the mean m_5 in each filter. Following [LSST Science Collaboration et al. \(2009\)](#), we simulate the uncertainties on the magnitudes as

$$\sigma_{\text{tot}}^2 = \sigma_{\text{sys}}^2 + \sigma_{\text{rand}}^2 \quad (4)$$

where we set the systematic noise $\sigma_{\text{sys}} = 0.005$ (the upper limit) for all filters and the random noise is given by

$$\sigma_{\text{rand}}^2 = (0.04 - \gamma)x + \gamma x^2, \text{ where } x = 10^{m-m_5} \quad (5)$$

and the filter dependent quantity $\gamma = 0.039$ (r and i bands), $= 0.040$ (z and y bands); see [LSST Science Collaboration et al. \(2009\)](#) for details.

Note that when the true (or theoretical) magnitude is much larger than the m_5 , i.e. the SN light curve is too faint compared to the 5σ depth, the σ_{rand} can be quite high. In these cases, a simple normal distribution can often incorrectly produces very bright observation (significantly brighter than the m_5 itself). To avoid this, we generate the observed magnitude by randomly choosing a value $m \sim \text{Uniform}(m_5, m_{\text{true}})$ (see [Huber et al. \(2022\)](#)); however this does not matter much as these are typically the faintest part of the SN light curves. The left panel of Fig. 4 presents the unresolved light curves of a typical quad system in four bands as an example.

4. Methodology: 1d CNN

Inspired by [Denissenya & Linder \(2022\)](#) we also employ 1D CNNs to detect the lensed cases from the light curve data.

4.1. Data preprocessing

In reality, we don’t know the explosion epoch of any lensed SN. To avoid providing any hint for the explosion time to the network, we set the time to zero at the epoch of the

first observation of each light curve. The SNe Ia light curve sample from LSST will contain a different number of observations per system, as well as different observation epochs. Nevertheless, to be able to employ a simple 1D CNN on the mock observed light curves we need the input to the networks to have the same dimensions. To achieve this, in addition to considering observation noise, we adopt the following data processing.

- We first smooth the observed magnitude vs time light curves using the iterative smoothing process of [Shafieloo et al. \(2006\)](#) which takes the uncertainties into account. Then we convert the smoothed magnitudes into the fluxes (in arbitrary unit) to get flux vs time light curves. Next we interpolate (cubic spline) the smoothed light curves and extrapolate them at the tails if necessary to have the same data length, e.g. N_j for the j -th filter. Next we normalize the light curves by scaling all the fluxes from 0 to 1 (for all systems, separately in all filters). At first, we want to test the method on long light curves by empirically choosing $N_j = 100$ for all the 4 filters; that means we use light curve data till 100 days from the first observation for all the systems across all bands. Later, we reduce N_j while trying to detect the lensed system early enough so that the SNe Ia could be still visible in the follow up.
- Then we concatenate (i.e. stack side by side) the normalized smoothed light curves for different bands (if available) to have an 1D array of $N_{\text{tot}} = \sum_{j \in \text{filters}} N_j$ data points comprising light curves for all the 4 bands. Note that we test the classification results using both stacked multi-band light curves and single-band observations.
- Furthermore, we use three different smoothing scales – 8, 13 and 16 days – and pass the 1D processed data (each having N_{tot} length) to the CNNs in three different channels. This choice of the smoothing scale is consistent with the cadence distribution (considering the mean value and the gaps) and is kept the same across all filters.⁶ The smoothed magnitude vs time light curves, corresponding to the three smoothing scales, are shown in each of the four panels of Figure 4(a). We expect that each channel will provide features at three different time scales, if any, in the smoothed light curves.
- Therefore, at the end, our input data end up having the shape $(N_{\text{tot}}, 3)$. Figure 4(b) illustrates how we arranged the input data for the example quad system. The three rows show the three 1D input arrays, each having the length of $N_{\text{tot}} = 400$, corresponding to three different smoothing scale, $\delta = 8, 13, 16$ days.

In addition to accounting for different number of observational epochs for different systems, this choice of data pre-processing handles observation noise which would influence the final smoothed light curves at different smoothing scales.

⁶ The selection of the smoothing scales is important because larger scales can wipe out small-scale features, if present, while smaller scales may overfit and struggle with long gaps in the observation epochs. We have tested multiple combinations of smoothing scales and found this choice to perform the best for the LSST cadence.




Fig. 4: **(a)** Blended light curves of a typical unresolved system are shown in the left panel across four LSST bands. We adopt LSST-like cadence and noise to generate mock light curves (as detailed in Sec. 3.3). Data processing begins by smoothing these light curves using different smoothing scales. The grey curves, delineated with dashed, dash-dotted, and dotted line styles, represent the smoothed light curves corresponding to smoothing scales of 8, 13, and 16 days, respectively. Subsequently, we convert the smoothed magnitudes into fluxes and sample the light curves on a predefined set of epochs, kept fixed for each band. For simplicity, we opt for 100 epochs with 1 day spacing, starting from the initial observation epoch (set as $t = 0$), for all the bands. We then normalize every light curve separately by scaling the fluxes in the range $[0, 1]$. **(b)** Finally, for each smoothing scale, we stack the normalized flux versus time light curves corresponding to the four bands side by side, thereby recasting the data into an array of length 400. Subsequently, we arrange three such arrays, corresponding to the three choices of smoothing scales, into three channels. Consequently, the final data size for each system becomes $(400, 3)$, as depicted in the right panel. This data, of fixed size, is then fed into the CNNs.

set name	Δt_{true}	use in classification	use in Δt estimation
set A	$U[5.0, 24.0]$	train + test	test only
set B	$U[4.0, 24.0]$		
set C	$U[3.0, 24.0]$		
set D	$U[0.5, 30.0]$	–	train only
set LSST	LSST like	test only	test only

Table 2: Description of different sets used for training and testing. Note that, except for set LSST, in rest of the controlled sets we use $\mu \sim U[1/3, 3.0]$ to have a balanced training. Set D is only used to train model for predicting time delays. Set LSST represents the whole sample of unresolved systems in LSST without any cuts being imposed. We use this set to test our models trained on different sets depending on our purpose of classification or time-delay estimation. Note that we randomly choose between W7 and merger models while generating the light curves for each system (lensed or unlensed) in all the sets.

4.2. Training and test sets

Figure 3 shows the distribution of maximum time delay and the corresponding magnification ratio for the unresolved systems in LSST. However, instead of using these distributions we use simple uniform distributions for μ and Δt to construct balanced training sets. As explained in Section 2, low values of time delays ($\Delta t \rightarrow 0$) or extreme values of magnification ratios ($\mu \gg 1$ or $\mu \ll 1$) makes our goal of detecting the lensed systems using blended joint light curves challenging.

At first, we set a reasonable limit on the magnification ratio, $\mu \sim U[1/3, 3.0]$. However, to test the limit on minimum time delay this approach can work on, we construct three different controlled sets, each with a different minimum Δt : (i) $\Delta t \in U[5.0, 24.0]$ for set A, (ii) $\Delta t \in U[4.0, 24.0]$ for set B and (iii) $\Delta t \in U[3.0, 24.0]$ for set C. These sets are summarized in Table 2. We also create a separate test set (along with the validation set) for each training set with the same distributions of μ and Δt for initial testings. We keep a ratio of $0.5 : 0.25 : 0.25$ between the population of unlensed, doubles and quads respectively in all training and test sets used for classification.

Note that even when the macro magnification ratio, μ , is non-extreme (e.g. around unity), microlensing can stochastically make it extreme in some cases. Inclusion of these joint light curves (comprising extreme μ 's) in the training set can confuse the training process, as the light curves can be very similar to an unlensed case. This can eventually result in lower accuracy in classification (as well as time-delay estimations). Therefore, we always scale the microlensed light curves of the images of a system such that their μ 's match the input values (drawn from a uniform distribution), and the i -band peak brightness matches the corresponding value in the OM10 simulation. By scaling with a constant factor, we preserve the microlensing effects (i.e., those that put wiggles) on each light curve. However, although microlensing should roughly preserve the ensemble average of μ , we can expect some cases to have extreme μ 's due to microlensing. We anticipate that these systems

will simply be classified as unlensed cases by the model which is trained on a set that excludes extreme μ 's.

While predicting the time delays for a test set, if we use the model trained on the corresponding training set with the same Δt interval, say $\Delta t \in (\Delta t_{\min}, \Delta t_{\max})$, we will inevitably face bias in the results at the Δt boundaries. In this case the model tends to over (under) predict for smaller (larger) time delays. This is simply because the model won't learn predicting lower than Δt_{\min} and higher than Δt_{\max} values. To overcome this apparent bias issues, we train the model on systems with $\Delta t \sim U[0.5, 30.0]$ days (set D, which covers larger range of Δt than that of set A, B, C) to predict the time delays for the test sets – A, B and C. In this training set for time-delay estimation, we keep the populations of doubles and quads the same. Note that, the test sets also share the same ratio.

Finally, to assess how our models, which are trained on the above sets, performs in realistic scenarios, we test them on a set with LSST like unresolved systems (set LSST) without any restrictions on μ and Δt_{true} .

4.3. Network architectures

As mentioned before, this work focuses on two distinct goals – (i) classification: discern the lensed systems from the unlensed ones (if possible classify between unlensed, doubles and quads) and (ii) estimation of the time delays. Therefore, we employ two types of 1D CNNs designed for the above two purposes: classification and time-delay regression. However, for simplicity, we keep the architecture of the two types of network quite similar until the last layer – starting with two 1D convolution layers (each is followed by a max-pooling layer) followed by 2 dense layers.⁷ The last layer is adapted according to the purpose of classification or regression. We emphasize that we only use simple network architectures in this work; however, a completely different (e.g. long-short term memory or LSTM) network can result in better results. Exploring all possible network architectures is beyond the scope of this work demonstrating the overall feasibility of classifying and analysing unresolved SN light curves with CNNs, but it could be a nice follow up project.

5. Results

5.1. Classification using long light curves

Initially, we use the long light curves, till 100 days from the first observation in each band.

5.1.1. 3×3 classification: unlensed vs doubles vs quads

First we try to classify the three types of systems – unlensed, doubles and quads. The results are summarised in Table 3 for the three sets (A, B and C in Table 2) with minimum Δt being 5.0, 4.0 and 3.0 days from the top. The first row of a confusion matrix provides what percentage of

	Predictions		
	unlensed	double	quad
Truth	unlensed all : 98.4% r: 93.8% i: 96.8%	all : 1.3% r: 4.7% i: 2.6%	all : 0.3% r: 1.5% i: 0.6%
	double all : 14.0% r: 25.8% i: 20.2%	all : 70.2% r: 54.5% i: 55.1%	all : 15.8% r: 19.7% i: 24.7%
	quad all : 2.4% r: 9.2% i: 8.2%	all : 9.9% r: 18.5% i: 19.5%	all : 87.7% r: 72.3% i: 72.3%

set A: $\Delta t_{\text{true}} \sim U[5.0, 24.0]$ and $\mu_{\text{true}} \sim U[1/3, 3.0]$

	Predictions		
	unlensed	double	quad
Truth	unlensed all : 97.0% r: 93.9% i: 96.6%	all : 2.6% r: 4.2% i: 2.7%	all : 0.4% r: 1.9% i: 0.7%
	double all : 14.0% r: 28.3% i: 21.9%	all : 69.6% r: 52.8% i: 52.0%	all : 16.4% r: 18.9% i: 26.1%
	quad all : 3.4% r: 10.3% i: 9.6%	all : 11.1% r: 19.5% i: 19.5%	all : 85.5% r: 70.2% i: 70.9%

set B: $\Delta t_{\text{true}} \sim U[4.0, 24.0]$ and $\mu_{\text{true}} \sim U[1/3, 3.0]$

	Predictions		
	unlensed	double	quad
Truth	unlensed all : 95.2% r: 93.4% i: 96.0%	all : 3.9% r: 4.7% i: 2.8%	all : 0.9% r: 1.9% i: 1.2%
	double all : 16.8% r: 30.0% i: 24.2%	all : 65.0% r: 49.1% i: 47.9%	all : 18.2% r: 20.9% i: 27.9%
	quad all : 3.6% r: 12.0% i: 9.8%	all : 11.8% r: 17.4% i: 17.6%	all : 84.6% r: 70.6% i: 72.6%

set C: $\Delta t_{\text{true}} \sim U[3.0, 24.0]$ and $\mu_{\text{true}} \sim U[1/3, 3.0]$

Table 3: Confusion matrices for the classification of unlensed, double and quad systems. The results obtained from multi-band (*rizy*) light curves are labeled as 'all,' while those from individual *r* and *i* bands are labeled accordingly.

	Predictions	
	unlensed	lensed
Truth	unlensed all : 97.1% r: 92.9% i: 95.8% z: 93.5% y: 86.1%	all : 2.9% r: 7.1% i: 4.2% z: 6.5% y: 15.9%
	lensed all : 7.7% r: 16.6% i: 12.5% z: 17.7% y: 25.1%	all : 92.3% r: 83.4% i: 87.5% z: 82.3% y: 74.9%

Table 4: set A: $\Delta t_{\text{true}} \sim U[5.0, 24.0]$ and $\mu_{\text{true}} \sim U[1/3, 3.0]$

⁷ When we try to optimize the architecture parameters (such as the number of convolution layers, filter sizes and numbers and the number of dense layers) using `keras.tuner` keeping the overall structure the same, we could only find minute improvements. Therefore, for simplicity, in this article we keep the architecture parameters the same in two types of networks until the last layer.

the true unlensed systems are classified as unlensed, double and quads in the three respective columns; similarly the next two rows provide the classification percentages of true doubles and quads. In each cell, we provide the result for all band data, considering only *r*-band and only *i*-band data. As expected the classification results are at the best when

the multi-band data are used, however, r or i band only data also results in reasonably good classification matrices.

Since the lensing probability is low (as we expect only one lensed systems out of $10^4 - 10^5$ randomly chosen SNe Ia), we need to control the false-positive rate (FPR) very well. E.g. if the lensing probability is as good as 0.01%, a FPR of 10% gives roughly 1000 false alarms per one true detection. We find that for the set A (with $\Delta t_{\min} = 5.0$ days), the networks can classify with a FPR as low as $\sim 1.6\%$. However, FPR increase to $\sim 4.8\%$ for the set C where Δt_{\min} is 3 days. The results mentioned above show a slight inferiority compared to those presented in Denissenya & Linder (2022), primarily attributed to differences in cadence, uncertainties, and the distribution of $(\mu, \Delta t)$ between LSST observation conditions considered here and the setup employed in Denissenya & Linder (2022).

We readily notice (in the four bottom-right cells) of Table 3 that the network confuses between doubles and quads very often. This confusion arises because we do not keep any minimum separation between the time delays of the later three images (with respect to the first image) for the quads. As a consequence, the joint light curve of a quad with three images having similar time delays, say $\Delta t'$ from the first image, will be similar to a double with the time delay $\Delta t'$ but with an aggregated magnification. Therefore, it is expected that the network confuses between certain quads and doubles. Also, owing to this confusion, the FPRs in Table 3 exhibit a variation of $\sim 1\text{--}2\%$ (additional variation on the FPR, not percentage of FPR) due to stochasticity in the network training, i.e., with different network initializations. Therefore, it makes more sense to classify the systems as unlensed vs lensed.

5.1.2. 2×2 classification: lensed vs unlensed

Here we consider all doubles and quads having the same truth label – ‘lensed’. The classification results are presented in Tables 4, 5 and 6 for the three sets – A, B and C. The top number (in boldface) in each cell corresponds to the scenario with complete four-band light curve data. The subsequent four numbers are obtained using a single-band light curve data as labeled. The true-positive rate (TPR) stays $\gtrsim 90\%$, with the FPR around 3%, for all three sets when we consider the four-band data. However, excluding the lower time-delay systems (going from set C to A) leads to enhanced overall accuracy, as expected.

Note that except for the y -band, the rest of the three bands— r , i , and z —individually produce well-controlled FPR of approximately 4–7%, while maintaining acceptable TPR of approximately 80–92%. However, it is evident that i -band light curves provide the most reliable classification because of better cadence and signal to noise ratio. Therefore, even with single-band light curve data from any of these three bands, we can achieve reasonably accurate classification.

5.2. Early detection

In the previous section, we could classify the systems well but we had to use light curves till 100 days from the first observation in each band. However, we want to detect them early enough, preferably in the light curve rising phase, so that the follow up can resolve it and subsequently measure

		Predictions	
		unlensed	lensed
Truth	unlensed	all : 96.8%	all : 3.2%
		r : 91.6%	r : 8.4%
		i : 95.2%	i : 4.8%
		z : 92.6%	z : 7.4%
		y : 83.8%	y : 16.2%
Truth	lensed	all : 8.9%	all : 91.1%
		r : 17.2%	r : 82.8%
		i : 13.9%	i : 86.1%
		z : 18.9%	z : 81.1%
		y : 24.8%	y : 75.2%

Table 5: set B: $\Delta t_{\text{true}} \sim U[4.0, 24.0]$ and $\mu_{\text{true}} \sim U[1/3, 3.0]$

		Predictions	
		unlensed	lensed
Truth	unlensed	all : 96.7%	all : 3.3%
		r : 91.5%	r : 8.5%
		i : 95.5%	i : 4.5%
		z : 92.7%	z : 7.3%
		y : 84.3%	y : 15.7%
Truth	lensed	all : 10.7%	all : 89.3%
		r : 19.1%	r : 80.9%
		i : 15.8%	i : 84.2%
		z : 21.0%	z : 79.0%
		y : 26.6%	y : 73.4%

Table 6: set C: $\Delta t_{\text{true}} \sim U[3.0, 24.0]$ and $\mu_{\text{true}} \sim U[1/3, 3.0]$




Fig. 5: Classification results for the light curves truncated at different t_{cut} values: true-positive rates (TPR; solid curves) and true-negative rates (TNR=1–FPR; dotted curves) are plotted for 4 different choices of t_{cut} (note that the far right points corresponds to the results already described in Section 5.1 and Tables 4–6). The results for set A, B and C are shown by the blue, orange and green colours respectively. As we truncate the light curves at lower epochs, both TPR and TNR decrease; therefore FPR increases.

the time delays. However, the nature of our method limits how early we can detect. In this section, we explore the minimum length of light curve observations required after the first detection to confidently determine that the SN is lensed. We refer to this duration as t_{cut} . (Previous section provides results for $t_{\text{cut}} = 100$ days.) Therefore, we cut the light curves at an earlier epoch, t_{cut} days from the epoch

		model trained on		
		set A	set B	set C
recovery	all	28.0%	28.1%	30.2%
	double	29.5%	29.2%	31.3%
	quad	21.9%	23.3%	25.9%
Δt_{med} of recovered systems	all	4.5	4.4	4.1
	double	4.8	4.7	4.6
	quad	2.6	2.1	1.6

Table 7: The top part presents the percentage of mock LSST unresolved systems recovered by models trained on different sets – A, B and C. The three sub-rows show the recovery rate for all recovered sample and then divided it into doubles and quads. The bottom part shows the median time delays of the recovered systems, again divided into three sets – all, doubles and quads from top to bottom. For comparison, the input LSST sample has $\Delta t_{\text{med}} = 2.0, 2.4, 1.1$ days respectively for the full sample, doubles and quads. It is evident that the models can recover the doubles better than the quads since the former systems have typically larger time delays.

of first observation. We keep t_{cut} the same for all bands⁸. We try to perform the classification separately for different choices of t_{cut} , e.g. at = 50, 40, 30, 20 and 10 days, to check how early we can detect the lensed systems with sufficient accuracy. For our goal of triggering a follow up, we only need the unlensed vs lensed classification, instead of the 3×3 classification (unlensed vs double vs quad) which is more difficult.

Instead of the confusion matrix for individual sets, here we present the results in Figure 5 which shows true positive (solid curves) and true negative (dotted curves) detection rates as a function of t_{cut} . Indeed, the classification accuracy remains reasonably well till $t_{\text{cut}} \sim 20$ days with TPR above 82% while maintaining a FPR below 9% for the most difficult set C. Therefore, this method shows the potential for detecting the lensed cases early enough for triggering a follow up.

5.3. Classifying LSST set

Finally, we test the models, which are trained on the controlled sets A, B and C, on an LSST-like sample of unresolved systems (i.e. without any restrictions on μ and Δt). We focus on unlensed vs lensed classification to avoid any confusion in detection of doubles and quads. The FPR of these models are already given in Tables 4–6 (top right element); e.g. FPR for these models is $\sim 3\%$ when all four band data are used. This exercise unambiguously determines what fraction of unresolved LSNe Ia in LSST these models can detect.

For simplicity, here we limit our analysis to light curves observed over a period of 100 days, specifically focusing on systems that have been well-monitored across all four LSST bands – r , i , z and y . The top part of Table 7 summarises

⁸ Cutting the long processed light curves (as in Figure 4(b)) at t_{cut} and feeding it to the CNN would give a hint of the peak location which might actually be beyond t_{cut} . Therefore, for a given t_{cut} , we always smooth the mag vs time light curves considering data up to the first observation after t_{cut} and then cut the smoothed light curve at t_{cut} . Finally we normalise the flux vs time light curves and feed it to the CNN.




Fig. 6: Recovery rate for the unresolved systems in set LSST, which have good multi-band light curves available, divided into 8 time-delay bins. Three colours – blue, orange and green – represent the results obtained using the model trained on the control set A, B and C, respectively.

what fractions of the unresolved systems are recovered by the model trained on set A, B and C (i.e. TPR as all the test samples are lensed). The first three rows presents the TPR considering all systems, only doubles and only quads respectively. For all three sets, roughly $\sim 30\%$ of all the systems are recovered. However, the recovery rate slightly improves left to right (from set A to C) when the training set includes lower time-delay systems as expected since most of the LSST samples have $\Delta t \lesssim 2$ days.⁹ Training on set C ($\Delta t_{\text{min}} = 3$ days) produces the best recovery rate 31.0%. Interestingly, for all three sets, doubles are recovered at a better rate than the quads since the former typically have larger time delays.

The bottom part presents the median time delay (Δt_{med}) of the systems that are recovered by the models trained on different sets. Again three rows here consider all systems, doubles and quads separately. For comparison, the input LSST sample has $\Delta t_{\text{med}} = 2.03, 2.37, 1.10$ days for all systems, doubles and quads. It is clear that the recovered systems have higher time delays as compared to the total input sample.

Figure 6 explicitly shows the recovery rate of the systems across different Δt bins. The results for models trained on three controlled sets, shown by three colours, exhibit a generally consistent pattern with each other. However, the model trained on set C demonstrates slightly superior performance on low time-delay systems. As anticipated from Table 7, the recovery rate overall improves for systems with higher time delays. E.g., for systems with $\Delta t < 3$ days, the recovery rate ranges between 16 – 20%, whereas it exceeds $\sim 60\%$ for systems with Δt between 9 and 12 days. Moreover, approximately 80 – 90% of the systems with $\Delta t > 15$ days could be recovered. Note that the recovery rate for the LSST set, even for higher time-delay systems, is

⁹ When tested with models trained on a few other sets where lower Δt and/or μ systems are included, we found minimal improvements in the recovery rate but at an expense of greater FPR. Therefore, we do not report the results for these additional sets.

typically slightly lower than in the control sets. This difference arises because LSST samples include systems with μ smaller than the lower limit on the control sets. These systems are generally more challenging to detect solely based on the shape of the light curves.

5.4. Time-delay estimation

Ideally, our goal is to detect the lensed systems using the light curves early enough (not too far from the peak of the first image) so that the follow ups can resolve the multiple images (and thus determine if it's a double or a quad) and measure the time delays. However, since we need long light curve measurements to confidently detect the lensed cases, the time-delay measurements might not be possible for many systems. In such cases, the ability to measure time delays directly from the light curves becomes valuable, and that is the focus of this section.

We train separate networks for predicting time delays in double and quad samples. This assumes proper classification of the system (as a double or quad), either through follow-up observations or a near perfect classification process, before directing it to the corresponding model. As previously mentioned, both networks share the same architecture until the last layer, which has three nodes for quads and one node for doubles, employing linear activation (or no activation) function.

Figure 7 compares the estimated time delays against the true values for the three sets – A, B and C – placed in three panels. In each panel, individual Δt estimations are represented by small dots, with orange and blue dots corresponding to time delays in doubles and quads, respectively. The green dashed line signifies $\Delta t_{\text{pred}} = \Delta t_{\text{true}}$, and it is evident that the predicted time delays roughly align with this line. This indicates the feasibility of estimating time delays solely from joint light curves, even with LSST-like cadence and noise.

We further segment the time delays into 10 equispaced bins, separately for doubles and quads. For each bin, we calculate the median of the predicted time delays and the 68% confidence level around it, depicted by error bars in green for doubles and red for quads. As anticipated, predictions are more accurate for higher time delays.

It's worth noting that, even after training on a set with a broader Δt range (set D), a slight over-prediction is observed at the lower end of Δt , particularly visible in the right panel of Figure 7. This is somewhat inevitable as, when approaching very small time delays, predictions become biased due to the inability to accommodate negative Δt .

Importantly, predictions are notably more accurate for double systems compared to quads. Double systems, having only one time delay, exhibit better predictions, while quads, with three time delays, pose a greater challenge for the network. Unlike Denissenya & Linder (2022), we do not impose a minimum time-delay separation between successive images, guided by the distribution of time delays in unresolved systems. This choice makes time-delay estimations for quads considerably more challenging than for doubles.

In each panel, we also provide the standard deviation (σ) of the residuals and the mean of their absolute values ($\langle \epsilon \rangle$), separately for the double and quad samples. Across all three sets, we observe that σ falls within the range of 3.1

to 3.2 days, and $\langle \epsilon \rangle$ is approximately 2.4 days for quads. For doubles, σ is around 2.5 days, with $\langle \epsilon \rangle$ at approximately 1.8 days. In summary, based on the controlled sets, we show in this section that indeed time-delay estimation is possible from only the joint light curves for the controlled sets.

In our final assessment, we test the model on the LSST set, where the majority of systems exhibit low values of Δt and μ (the median values of Δt_{max} and corresponding μ are 2.0 days and 0.4 respectively). The predicted time delays are plotted against the ground truths in Figure 8. The estimated time delays for the quads (blue dots) deviates substantially from the true values, reflecting the challenge in accurate prediction, since quads often entail very small time delays (with a median of $\Delta t_{\text{max}} \approx 1.1$ day). Consequently, we abstain from presenting statistics across different Δt bins for the quads. On the other hand, the predictions for doubles (orange dots) fare somewhat better. Nonetheless, the model exhibits a tendency to overpredict for low time-delay systems, as expected from the above analysis. Even for moderate time-delay systems, a notable bias is observed, with more values underpredicted than otherwise. This bias is visually evident from the error bars, which depict the median and 68% confidence level around it across different Δt bins. Notably, the error bars for the LSST dataset are somewhat larger than those for the controlled sets shown in Figure 7. This discrepancy can be attributed to the distribution of μ in the LSST dataset, which significantly falls below the lower limit of training set D. Overall, this evaluation underscores the inherent difficulty in estimating time delays solely from the shape of joint light curves of unresolved systems in LSST, as these systems often exhibit low time delays and magnification ratios.

6. Summary and discussion

Gravitationally lensed supernovae (LSNe) hold significant promise as vital cosmic probes in the near future. The primary challenge of their rarity can be overcome by identifying the abundant unresolved LSNe anticipated in wide-field surveys. This article leverages the lensing features in the observed light curves to discern between unresolved LSNe Ia and their unlensed counterparts in LSST, utilizing a straightforward 1D CNN approach. Joint light curves of unresolved SNe Ia comprise the positive examples, while the negative examples are represented by the intrinsic light curves of (unlensed) SNe Ia. To simulate LSST-like mock light curves, we employ the cadence distribution and 5σ depth from the `baseline v3.2` observing strategy. Details of the simulation process are provided in Section 3. Out of six LSST filters, we consider four, namely r , i , z and y , due to their better cadences.

Based on OM10 simulations we show that LSST will observe approximately three times more unresolved LSNe Ia than the resolved ones. However, for nearly half of these systems, the maximum time delay (Δt) is expected to be less than 2 days. It is crucial to avoid the limit as Δt approaches zero or extreme magnification ratios (μ) in the training process, as in both scenarios, the distinctive lensing signature within the joint light curves diminishes, rendering it nearly impossible for any method to distinguish lensed cases based solely on light curve shapes. To ensure robustness in our training process, we exclude instances with low values Δt and μ . Initially, we focus on controlled datasets denoted as A, B, and C. These datasets adhere




Fig. 7: Time-delay estimates from blended joint light curves were compared with truths for sets A, B, and C. All predictions comes from the model trained on set D, which covers a larger Δt range ($[0.5, 30]$ days), to minimize apparent bias at Δt boundaries. Orange and blue dots in the plots represent individual predictions for double and quad systems, respectively. The standard deviation of residuals (σ) and mean of their absolute values ($\langle \epsilon \rangle$) are quoted in each plot, separately for doubles and quads. Samples are split into 10 bins, and the 68% CL around the median is shown by error bars, again separately for doubles (black) and quads (red); the two kind of errorbars are plotted with a slight offset along x-axis for a better visualization. It is evident that time-delay predictions are more accurate for doubles compared to quads or at higher time delays, as expected.




Fig. 8: Time-delay estimations for the set LSST are compared against the ground truths. Statistics of Δt_{pred} for the quads (depicted by blue dots) at different bins are omitted, as the Δt values for quads are typically very low, and we lack sufficient numbers at higher Δt . Therefore, we present the median and 68% confidence level of Δt_{pred} at different Δt_{true} bins only for the doubles.

to specific criteria: both Δt and μ are uniformly sampled, with μ following a uniform distribution in the range $[1/3, 3]$ in all sets, and the minimum allowed Δt is 5, 4, and 3 days

respectively for set A, B, and C. The outcomes for these controlled sets are provided below.

- We have observed that the classification between unlensed, double, and quad can maintain a well-controlled false-positive rate (FPR). However, there is notable confusion while distinguishing some quads from doubles. This confusion arises due to the fact that joint light curves of certain quads may closely resemble those of doubles, particularly when the separations between the time delays are not significant enough.
- Hence, the classification between unlensed and lensed cases appears more meaningful. For systems with ‘well-observed’ light curves across all four bands, we achieve a recovery rate of $\sim 90\%$ with a FPR $\sim 3\%$ for all three sets. Even when considering only a single-band light curve observed among the filters r , i and z , the true-positive rate (TPR) remains above 80%, with the FPR staying $\sim 4 - 7\%$. Consequently, confident detection of unresolved lensed system is feasible even with just a single-band light curve.
- Early detection plays a pivotal role in initiating follow-up observations. Our demonstration illustrates that even with light curves truncated at 20 days from the initial observation epoch, reliable detection of lensed cases is feasible with FPR $\lesssim 8\%$ for all three sets. This underscores ample opportunity for triggering follow-up observations aimed at first confirming and then precisely measuring the time delays.
- In scenarios where follow-up observations are not possible, we can still estimate the time delays directly from the joint light curves, provided the systems are accurately classified as doubles or quads. Time-delay esti-

mations are notably more precise for doubles, as they possess a unique time delay.¹⁰

Finally, we test the models, which are trained on these controlled sets (A, B and C), on the full LSST sample (which has no restriction on the Δt and μ). We manage to recover nearly 30% of these systems, provided their light curves are well-observed in the four bands. Based on the testing conducted on the controlled sets, we are already aware that the FPR would remain around 3% for these cases. As expected, the recovery rate is higher for larger time-delay systems. For instance, approximately 60% and 80 – 90% of the systems with time delays ranging between 9 and 12 days, and greater than 15 days, respectively, could be recovered. Subsequently, we apply the model trained on set D to the set LSST for time-delay estimation. The result reveals challenges in achieving accurate time-delay measurements, characterized by biases and significant uncertainties, even for doubles. This difficulty stems from the predominance of low Δt and μ values in the LSST unresolved samples. Consequently, relying solely on joint light curve data for time-delay estimations may not yield reliable results.

Note that our method is specifically applicable to SN Ia light curves for distinguishing between lensed and unlensed SNe Ia. Therefore, this approach relies on the accurate identification of SN Ia events beforehand. Current algorithms, such as [Lochner et al. \(2016\)](#); [Alves et al. \(2022\)](#), demonstrate a high level of reliability in detecting SNe Ia among other transients based on photometry.

All in all, our approach of detecting the unresolved LSNe Ia solely based on the shape of the blended light curves shows great potential for the upcoming LSST survey. Consequently, the recovery rate (of all unresolved systems) is fundamentally limited by two factors. Firstly, we need well sampled light curves. Therefore, we follow the set of ‘good observation’ criteria, as described in Section 3.3, to select the well monitored systems and leave out the rest those could not be sufficiently monitored, e.g. because of season gaps. Since these criteria were mostly ad hoc, we have the flexibility to relax them, allowing for the inclusion of a larger fraction of unresolved systems. This adjustment enables us to explore the limit on these criteria, without significantly sacrificing the accuracy of the results.

Secondly, as expected, this method struggles to detect systems with low time delays, which is significant considering that most unresolved systems in LSST will likely have low time delays. Fortunately, lower time-delay systems often exhibit higher total magnification, where the traditional magnification method (and possibly the colour magnitude diagram too) is expected to perform better. On the contrary, this method performs well for systems with time delays exceeding 5-6 days. Therefore, this approach complements the magnification method.

Another promising way to significantly improve the classification efficiency is by incorporating imaging and/or spectroscopic data alongside the light curves. For instance, in lensed cases, we anticipate the presence of the lensing galaxy near the SN. Moreover, in many instances, the host galaxy might be evident in the reference image.

Additionally, the shape of an unresolved SN would not necessarily appear point-like; rather, it could be elongated in the difference imaging, possibly with its centroid shifting over time as different image dominates the observed light. Furthermore, for type Ia SNe, estimating the magnification is feasible as they can serve as standardizable candles. This allows us to verify whether the observed brightness aligns well with the photo- z estimation of the lens. Incorporating this supplementary information should lead to a better classification efficiency with lower FPR. Another potential path for improvement lies in the network architecture. Instead of employing a CNN, which necessitates fixed input data, we can utilize a long short-term memory (LSTM) network. LSTM networks have the advantage of being able to handle variable-length input sequences and can also accommodate flux uncertainties directly as inputs. These aspects will be further explored in subsequent research endeavors.

It’s worth noting that unresolved supernovae can be observed up to higher luminosity distances, presenting an opportunity to study the properties of supernovae at high redshifts. For instance, they can be utilized to investigate whether there is any evolution in the light curves of type Ia supernovae.

In summary, this study showcases the viability of identifying unresolved LSNe Ia through light curve observations in LSST using deep learning techniques. Additionally, we illustrate the potential for extracting time delays solely from these unresolved light curves, enabling the prompt utilization of such systems delivered by LSST, which may be overlooked by most other detection methods. Although the achieved false-positive rate is below 3% in the optimal scenario, further exploration with more complex network architectures and inclusion of additional datasets may aid in further reducing it.

Appendix A: Source and lens redshift of the unresolved systems in LSST

In Figure A.1 we compare between the resolved and unresolved LSNe Ia in LSST in terms of their source and lens redshifts in the left and right panel respectively. Unresolved systems, being observable at greater distances, can accommodate larger source redshifts compared to resolved ones. This presents a unique opportunity to investigate the properties of SNe at high redshifts, such as potential evolution in the light curve of SNe Ia. Understanding such evolution is crucial for cosmology, as Type Ia SNe serve as fundamental tools for studying the universe’s expansion. On the other hand, their smaller angular separations require larger lens redshifts compared to resolved systems.

Acknowledgement

We thank Masamune Oguri for providing the updated OM10 simulation code and for offering valuable comments. We also thank Taeho Ryu and Kaitlyn Szekerczes for their assistances in setting up the code. SB acknowledges the insightful discussions with Georgios Vernardos, Mikhail Denissenya and Wuhyun Sohn on various aspects of this project. SB acknowledges the funding provided by the Alexander von Humboldt Foundation. SHS thanks the Max Planck Society for support through the Max Planck Fellowship. This project has received funding from the European Research Council (ERC) under the

¹⁰ Note that all the results presented above, listed as separate points, are confined to the controlled sets where we exclude low Δt and μ systems from training and testing; see Table 2.




Fig. A.1: The unresolved and resolved LSNe Ia in LSST have been compared in terms of their distributions of source redshift (left panel) and lens redshift (right panel). Since unresolved systems can be observed up to larger luminosity distances, they can accommodate larger source redshifts as compared to the resolved ones. Conversely, their smaller angular separations necessitate larger lens redshifts compared to the resolved systems.

European Union's Horizon 2020 research and innovation programme (grant agreement No 771776). This research is supported in part by the Excellence Cluster ORIGINS which is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC-2094 – 390783311. SS has received funding from the European Union's Horizon 2022 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 101105167 - FASTIDIoUS. AS would like to acknowledge the support by National Research Foundation of Korea NRF-2021M3F7A1082056, and the support of the Korea Institute for Advanced Study (KIAS) grant funded by the government of Korea.

Data availability

The catalog of unresolved lensed SNe Ia simulated using OM10 simulation pipeline will be made public upon acceptance.

References

Aghanim, N. et al. 2020, *Astron. Astrophys.*, 641, A6
 Alves, C. S., Peiris, H. V., Lochner, M., et al. 2022, *ApJS*, 258, 23
 Arendse, N., Dhawan, S., Sagués Carracedo, A., et al. 2023, *arXiv e-prints*, arXiv:2312.04621
 Bag, S., Kim, A. G., Linder, E. V., & Shafieloo, A. 2021, *Astrophys. J.*, 910, 65
 Bag, S., Shafieloo, A., Liao, K., & Treu, T. 2022, *Astrophys. J.*, 927, 191
 Bag, S., Sohn, W., Shafieloo, A., & Liao, K. 2023, *ApJ*, 950, 158
 Bellm, E. C., Kulkarni, S. R., Graham, M. J., et al. 2019, *PASP*, 131, 018002
 Birrer, S., Shajib, A. J., Galan, A., et al. 2020, *A&A*, 643, A165
 Birrer, S. & Treu, T. 2021, *A&A*, 649, A61
 Biswas, R., Daniel, S. F., Hložek, R., et al. 2020, *ApJS*, 247, 60
 Bonvin, V., Courbin, F., Suyu, S. H., et al. 2017, *MNRAS*, 465, 4914
 Chan, J. H. H., Rojas, K., Millon, M., et al. 2021, *A&A*, 647, A115
 Chen, W., Kelly, P. L., Oguri, M., et al. 2022, *Nature*, 611, 256
 Delgado, F. & Reuter, M. A. 2016, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 9910, *Observatory Operations: Strategies, Processes, and Systems VI*, ed. A. B. Peck, R. L. Seaman, & C. R. Benn, 991013

Delgado, F., Saha, A., Chandrasekharan, S., et al. 2014, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 9150, *Modeling, Systems Engineering, and Project Management for Astronomy VI*, ed. G. Z. Angeli & P. Dierickx, 915015
 Denissenya, M., Bag, S., Kim, A. G., Linder, E. V., & Shafieloo, A. 2021, *arXiv e-prints*
 Denissenya, M. & Linder, E. V. 2022, *MNRAS*, 515, 977
 Dimitriadis, G., Rojas-Bravo, C., Kilpatrick, C. D., et al. 2019, *ApJ*, 870, L14
 Ding, X., Liao, K., Birrer, S., et al. 2021, *Monthly Notices of the Royal Astronomical Society*, 504, 5621
 Foxley-Marrable, M., Collett, T. E., Vernardos, G., Goldstein, D. A., & Bacon, D. 2018, *MNRAS*, 478, 5081
 Frye, B., Pascale, M., Cohen, S., et al. 2023, *Transient Name Server AstroNote*, 96, 1
 Goldstein, D. A. & Nugent, P. E. 2017, *ApJ*, 834, L5
 Goldstein, D. A., Nugent, P. E., Kasen, D. N., & Collett, T. E. 2018, *ApJ*, 855, 22
 Goobar, A., Johansson, J., Schulze, S., et al. 2023, *Nature Astronomy*, 7, 1098
 Goobar, A. et al. 2017, *Science*, 356, 291
 Grillo, C., Pagano, L., Rosati, P., & Suyu, S. H. 2024, *arXiv e-prints*
 Huber, S. & Suyu, S. H. 2024, *arXiv e-prints*, arXiv:2403.08029
 Huber, S., Suyu, S. H., Ghoshdastidar, D., et al. 2022, *A&A*, 658, A157
 Huber, S., Suyu, S. H., Noebauer, U. M., et al. 2019, *A&A*, 631, A161
 Huber, S., Suyu, S. H., Noebauer, U. M., et al. 2021, *A&A*, 646, A110
 Kelly, P. L. et al. 2015, *Science*, 347, 1123
 Kelly, P. L. et al. 2023, *Science*, 380, abh1322
 Kodi Ramanah, D., Arendse, N., & Wojtak, R. 2022, *MNRAS*, 512, 5404
 Kromer, M. & Sim. 2009, *MNRAS*, 398, 1809
 Lochner, M., McEwen, J. D., Peiris, H. V., Lahav, O., & Winter, M. K. 2016, *ApJS*, 225, 31
 LSST Science Collaboration, Abell, P. A., Allison, J., et al. 2009, *arXiv e-prints*, arXiv:0912.0201
 LSST Science Collaboration, Marshall, P., Anguita, T., et al. 2017, *arXiv e-prints*, arXiv:1708.04058
 More, A., Suyu, S. H., Oguri, M., More, S., & Lee, C.-H. 2017, *ApJ*, 835, L25
 Naghib, E., Yoachim, P., Vanderbei, R. J., Connolly, A. J., & Jones, R. L. 2019, *AJ*, 157, 151
 Nomoto, K., Thielemann, F. K., & Yokoi, K. 1984, *ApJ*, 286, 644
 Oguri, M. 2007, *The Astrophysical Journal*, 660, 1
 Oguri, M. 2018, *MNRAS*, 480, 3842
 Oguri, M. & Marshall, P. J. 2010, *MNRAS*, 405, 2579
 Pakmor, R., Hachinger, S., Röpke, F. K., & Hillebrandt, W. 2011, *A&A*, 528, A117
 Pakmor, R., Kromer, M., Taubenberger, S., et al. 2012, *ApJ*, 747, L10

- Pascale, M., Frye, B. L., Pierel, J. D. R., et al. 2024, arXiv e-prints, arXiv:2403.18902
- Pierel, J. D. R., Newman, A. B., Dhawan, S., et al. 2024, arXiv e-prints, arXiv:2404.02139
- Quimby, R. M., Oguri, M., More, A., et al. 2014, *Science*, 344, 396
- Refsdal, S. 1964, *Monthly Notices of the Royal Astronomical Society*, 128, 307
- Refsdal, S. & Bondi, H. 1964, *Monthly Notices of the Royal Astronomical Society*, 128, 295
- Riess, A. G. et al. 2022, *Astrophys. J. Lett.*, 934, L7
- Rodney, S. A., Brammer, G. B., Pierel, J. D. R., et al. 2021, *Nature Astronomy*
- Röpke, F. K., Kromer, M., Seitzzahl, I. R., et al. 2012, *ApJ*, 750, L19
- Saha, P., Coles, J., Macciò, A. V., & Williams, L. L. R. 2006, *ApJ*, 650, L17
- Seitzzahl, I. R., Ciaraldi-Schoolmann, F., Röpke, F. K., et al. 2013, *MNRAS*, 429, 1156
- Shafieloo, A., Alam, U., Sahni, V., & Starobinsky, A. A. 2006, *Mon. Not. Roy. Astron. Soc.*, 366, 1081
- Shu, Y., Belokurov, V., & Evans, N. W. 2021, *MNRAS*, 502, 2912
- Sim, S. A., Röpke, F. K., Hillebrandt, W., et al. 2010, *ApJ*, 714, L52
- Suyu, S. H., Goobar, A., Collett, T., More, A., & Vernardos, G. 2024, *Space Sci. Rev.*, 220, 13
- Suyu, S. H., Huber, S., Cañameras, R., et al. 2020, *A&A*, 644, A162
- The LSST Dark Energy Science Collaboration, Mandelbaum, R., Eifler, T., et al. 2018, arXiv e-prints, arXiv:1809.01669
- Treu, T., Suyu, S. H., & Marshall, P. J. 2022, *Astron. Astrophys. Rev.*, 30, 8
- Vernardos, G. & Fluke, C. J. 2014, *Astron. Comput.*, 6, 1
- Vernardos, G., Fluke, C. J., Bate, N. F., & Croton, D. 2014, *ApJS*, 211, 16
- Vernardos, G., Fluke, C. J., Bate, N. F., Croton, D., & Vohl, D. 2015, *ApJS*, 217, 23
- Weisenbach, L., Schechter, P. L., & Pontula, S. 2021, *ApJ*, 922, 70
- Wojtak, R., Hjorth, J., & Gall, C. 2019, *MNRAS*, 487, 3342
- Wong, K. C. et al. 2020, *Mon. Not. Roy. Astron. Soc.*, 498, 1420