

FOCK SPACE OF LOCAL FIELDS OF THE DISCRETE GFF AND ITS SCALING LIMIT BOSONIC CFT

David Adame - Carrillo^{*}, Delara Behzad[†], and Kalle Kytölä[‡]

^{* ‡} *Department of Mathematics and Systems Analysis, Aalto University,
P.O. Box 11100, FI-00076 Aalto, Finland*

[†] *Mathematical Institute, Utrecht University,
Budapestlaan 6, 3584 CD Utrecht, The Netherlands*

Abstract

To connect conformal field theories (CFT) to probabilistic lattice models, recent works [HKV22, Ada23] have introduced a novel definition of local fields of the lattice models. Local fields in this picture are probabilistically concrete: they are built from random variables in the model. The key insight is that discrete complex analysis ideas allow to equip the space of local fields with the main structure of a CFT: a representation of the Virasoro algebra.

In this article, for the first time, we fully analyze the structure of the space of local fields of a lattice model as a representation, and use this to establish a one-to-one correspondence between the local fields of a lattice model and those of a conformal field theory. The CFT we consider is probabilistically realized in terms of the gradient of the Gaussian Free Field (GFF). Its space of local fields is just a bosonic Fock space for two chiral symmetry algebras. The corresponding lattice model is the discrete Gaussian Free Field. Our first main result is that the space of local fields of polynomials in the gradient of the discrete GFF is isomorphic to the Fock space. Notably, local fields in this setup make sense with both Dirichlet and Neumann boundary conditions. Our second main result is that with the appropriate renormalization, correlation functions of local fields of the discrete GFF converge in the scaling limit to the correlation functions of the CFT. The renormalization needed is, conceptually correctly, according to the eigenvalue of the Virasoro generator $L_0 + \bar{L}_0$ on the local field.

^{*}david.adamecarrillo@aalto.fi

[†]d.behzad@uu.nl

[‡]kalle.kytola@aalto.fi

Contents

1	Introduction	4
1.1	Lattice models as discretizations of conformal field theories	4
1.2	The massless free boson and the Gaussian Free Field	5
1.3	Conformal field theory: local fields and their correlation functions	6
1.4	Local fields in the lattice model	8
1.5	Main results	9
1.6	Organization of the article and outlines of the proofs	11
1.7	Scaling limit results in the literature	13
2	Preliminaries: the Fock space	15
2.1	Heisenberg algebra and Fock space	16
2.2	Virasoro algebra and the Sugawara construction	18
2.3	Primary fields in the Fock space	19
3	Preliminaries: discrete complex analysis	20
3.1	Square grids and discrete differential operators	21
3.2	Discrete domains and Green's functions	23
3.3	Scaling limits of discrete Green's functions	24
3.4	Discrete contour integration and discrete monomial functions	27
4	Discrete Gaussian Free Field and its local fields	34
4.1	DGFF with Dirichlet and Neumann boundary conditions	34
4.2	Discrete holomorphic and antiholomorphic currents	36
4.3	Field polynomials	37
4.4	Local fields	38
4.5	Current modes	40
4.6	Sugawara construction and homogeneous local fields	46
5	Linear local fields of the DGFF	47
5.1	Linear local fields	48
5.2	Filtration by radius	49
5.3	Dimension upper bounds	51
5.4	Homogeneous linear local fields	55

6	Local fields of higher degree	56
6.1	Normal ordering	57
6.2	Local fields form a Fock space	61
6.3	Concrete examples of higher degree local fields	64
7	Scaling limits of local field correlations	65
7.1	Gaussian Free Fields	66
7.2	Current correlation kernels of the GFF	69
7.3	Correlation functions of CFT fields from current OPEs	70
7.4	Renormalized limits of correlations of lattice local fields	75

1 Introduction

1.1 Lattice models as discretizations of conformal field theories. Making mathematical sense of quantum field theories in continuum space(-time) is often very challenging. Many of the physicists' commonly used ways (such as path integrals) of specifying field theories are not as such well-defined in continuum space. One typically needs to introduce a short-distance cutoff (ultraviolet cutoff) to write down proper mathematical definitions, and then one should analyze what happens as the cutoff is removed. One standard way of implementing the cutoff is to discretize the theory to a lattice. The mesh size of the lattice then serves as the short-distance cutoff length scale. The advantage is that defining the discretized field theories as probabilistic models in (finite) lattice domains is usually straightforward. The difficulty then lies in removing the ultraviolet cutoff by forming a scaling limit of these well-defined models, i.e., proving that relevant limits exist when one lets the lattice mesh size tend to zero and proving field-theoretically desired properties of these limits. Despite still being challenging, the approach of lattice discretizations and their scaling limits is one of the main robust and standard ways to construct field theories.

Since the directly physically important quantities in a field theory are its correlation functions, for a scaling limit to be viewed as a construction of a field theory, it should address the convergence of correlation functions in the discretized models as the lattice mesh size tends to zero. In a field theory, correlation functions are assigned to any so-called local fields at any number of spatial positions. Note that the terminology here can be slightly confusing: "local fields" does not simply refer to the "basic fields", in terms of which for example the path integral of the field theory would be written. Local fields are meant to represent *all* of the locally observable quantities in the theory. The basic fields are typically among those, but the notion of local fields is vastly more general.

Both the physics and mathematics research places a lot of emphasis on two-dimensional conformal field theories (CFT), i.e., field theories in two-dimensional space(-time) with certain conformal invariance constraints. One reason is that by the renormalization group philosophy, under physically reasonable assumptions, universal macroscopic (large-scale) behavior of general theories should be governed by renormalization group fixed points which are CFTs. For example statistical physics models at their critical points should renormalize exactly to CFTs, and their universal behavior near the critical point should be governed by the linearized renormalization group flow near the fixed point, which in turn can be analyzed in terms of the CFT fixed point itself. Another reason is that two-dimensional CFTs have remarkable structures that make them both interesting and significantly more tractable than most other field theories, and therefore they are great example cases to study in detail. In particular, by virtue of conformal invariance, the state space of a two-dimensional CFT carries a representation of (two commuting copies of) the Virasoro algebra. And by what is known as the state-field correspondence, the state space is exactly identified with the space of local fields. Therefore the rich algebraic structure of the CFT is present in the observable quantities which are to be inserted in correlation functions. The Virasoro algebra action on local fields describes the (infinitesimal) changes of correlation functions as

(infinitesimal) conformal transformations are applied. Simple but important special cases of conformal transformations are scaling transformations, and consequently the eigenvalues of the corresponding Virasoro generators ($L_0 + \bar{L}_0$) describe exactly how local fields should be renormalized as functions of the lattice mesh when forming the scaling limit (or more generally renormalization with any ultra-violet cutoff scale). This is crucial to the constructive field theory approach via scaling limits, and it furthermore contains the information about the renormalization group transformation in the vicinity of the CFT fixed point, and via that, the critical exponents that govern the universal large-scale behavior of the model near the critical point. The physicists' exact (albeit not always rigorous) determination of critical exponents in a number of interesting models via the use of the representation theory of the Virasoro algebra has indeed been one of the spectacular successes of two-dimensional conformal field theories.

The objective in this article is to make concrete sense of the above picture in the case of one particular conformal field theory and its probabilistic lattice model discretization. The field theory is essentially the simplest imaginable one, the massless free boson in two dimensions, and it correspondingly provides the simplest theory in which the general question of (re)constructing a CFT as a scaling limit is meaningful. The lattice model is also arguably its simplest discretization, the discrete Gaussian Free Field. What is noteworthy is how much detailed structure we can, in this case, match between the CFT and the discrete probabilistic model. In fact, we view our results as the first ones to establish a full conformal field theory as the scaling limit of a probabilistic lattice model. We believe that it will eventually be possible to obtain similar complete CFT scaling limit results also for at least a few other theories — among which the most promising candidates are the Ising model and discrete symplectic fermions, building on [HKV22] and [Ada23], respectively.

1.2 The massless free boson and the Gaussian Free Field. The massless free boson is, first of all, a fundamental example among conformal field theories, and it is common to start discussing CFTs with the free boson as the prototype, see, e.g., [DMS97, Gaw99, KaMa13]. The physics description of the free boson field theory is usually given in terms of path integral, so that for example the partition function (in the Euclidean signature) in a planar domain Ω would be written as

$$“ \int e^{-S(\varphi)} \mathcal{D}\varphi ”, \quad \text{where} \quad S(\varphi) = \frac{1}{8\pi} \int_{\Omega} \|\nabla \varphi(z)\|^2 \, d^2z, \quad (1.1)$$

and with the path integral taken formally over field configurations $\varphi: \Omega \rightarrow \mathbb{R}$. The quantity $S(\varphi)$ in the exponential — the action of the theory — is a quadratic form in φ , the Dirichlet energy of φ . By virtue of this gaussianity, it is actually not very difficult to give a precise meaning to the massless free boson path integral as a probability measure [Gaw99, She07, WePo21, BePo23]. In this probabilistic incarnation, the theory is referred to as the Gaussian Free Field (GFF). If the domain Ω has boundaries, then in the path-integral and in the corresponding Gaussian probability measure, boundary conditions for φ should be specified. Dirichlet and Neumann boundary conditions are the most

common choices. Also unless there are at least some Dirichlet boundary conditions, the zero-mode (“average”) of φ requires special treatment; the GFF φ is then only well-defined up to additive constants. We will actually work with the gradient $\nabla\varphi$ of the GFF (see Section 7 for the precise definition), which gets rid of the additive constant ambiguity, and which is well-defined with both Dirichlet and Neumann boundary conditions.

More complicated field theories can be obtained building on the simple case of the free boson. For example, allowing multiple components for the bosonic fields and compactifying the target space gives rise to interesting CFTs [Gaw99]. The GFF is furthermore at the heart of constructive field theory beyond free fields, because one can view its well-defined Gaussian measure as a reference and then include interactions by adding a potential, see [GlJa81].¹ One prominent recent example is giving a mathematical meaning to the intricate Liouville conformal field theory via a path integral [DKRV16], and proving that the path-integral construction indeed gives a conformal field theory [GKRV21], see [GKR24] for a review.

The GFF fundamentally underlies also many fascinating aspects of random geometry. Via probabilistic couplings, the GFF is intimately related to SLE-type random curves [Dub09, ScSh09, IzKy13, ScSh13, MiSh16, PeWu19] and conformal loop ensembles [QiWe18, ASW19, WePo21]. These random curves can be studied in terms of the GFF and vice versa.

The ubiquity of the GFF, especially its relations to various discrete models, may appear surprising. This is, however, at least partly explained by its simplicity and universality. The articles [BPR20, BPR21, ArPo21] indeed show how modest assumptions suffice to characterize the GFF.

1.3 Conformal field theory: local fields and their correlation functions. To properly specify a field theory, it is not sufficient to just give an action — one also has to address what are the (locally) observable quantities in it, i.e., what is the space of local fields of the theory. By the state-field correspondence, this is the same data as the state space of the theory, or its “spectrum”.

Typically in a field theory there are some basic degrees of freedom, or basic fields, in terms of which for example the action of the theory is written in path integral formulations. The free boson path integral (1.1) involves a single scalar field φ in this role of a basic field. To illustrate the notion of local fields, it is easiest to start with examples. It seems natural that the value $\varphi(z)$ of the basic field at a point z would be an observable quantity located at z . Also the values of the derivatives $\partial_\mu\varphi(z)$, $\partial_\mu\partial_\nu\varphi(z)$, \dots , of any order, are determined by the configuration of the field φ in an infinitesimal neighborhood of z , so they can be viewed as locally observable quantities at z . Furthermore, one could consider polynomials (with suitable regularization) in any of the above, such as $\varphi(z)^4$ or $(\partial_\mu\varphi(z))^2$, etc., or even more complicated functions (again suitably regularized) such as $e^{i\alpha\varphi(z)}$. All of these examples are indeed quantities that one might want to assign correlation functions to, and the informal examples illustrate the idea that a local field is an observable quantity

¹Often such interacting field theories would be treated perturbatively around the GFF. We emphasize, however, that both in principle and in practice, also nonperturbative constructive treatments of interacting theories build on the GFF in this way.

whose value is determined locally (only accessing an infinitesimal neighborhood of a point) from the appropriate basic degrees of freedom.

Note, however, that the basic degrees of freedom are not necessarily themselves observable quantities — for example in gauge theories, observable quantities must be independent of the (unphysical) gauge choices, whereas the basic fields in the path integral depend on the gauge. Similarly, for the GFF with Neumann boundary conditions (as well as for the GFF on Riemann surfaces without boundaries), there is an ambiguity of an additive constant in φ , so it makes sense to exclude the field value $\varphi(z)$ while allowing the values of its derivatives as local fields. We will indeed do so, we will consider a bosonic CFT in which the local fields are the *polynomials in the derivatives of φ* . There are two reasons for this choice. First, this allows us to talk about the same CFT with different boundary conditions (in this article we explicitly state results for Dirichlet and Neumann boundary conditions, but for example mixtures of these could be handled similarly). Second, this choice leads to a simple and concrete CFT. The space of local fields of this CFT will simply be the chargeless Fock space for two chiral Heisenberg algebras², which we will denote by

$$\mathcal{F} \otimes \overline{\mathcal{F}} \quad (\text{the precise definition will be given in Section 2}).$$

In any domain Ω , with Dirichlet or Neumann boundary conditions (abbreviated D/N), any local fields $F_1, \dots, F_n \in \mathcal{F} \otimes \overline{\mathcal{F}}$ inserted at any distinct points $z_1, \dots, z_n \in \Omega$ have correlation functions

$$\left\langle F_1(z_1) \cdots F_n(z_n) \right\rangle_{\Omega}^{\text{D/N}} \quad (\text{the precise definition will be given in Section 7})$$

in this free bosonic CFT. Indeed a field theory should fundamentally consist of the above kind of data: a space of local fields, and correlation functions assigned to any local fields at any finite number of distinct points in any domain. This data should of course also satisfy suitable axioms of a field theory, notably concerning the existence of operator product expansions (OPE), i.e., short-distance expansions of correlation functions of local fields in terms of other local fields contained in the theory³.

As we allow only polynomials in the gradient of φ as local fields, we are excluding not just the field $\varphi(z)$ itself, but also for example the “vertex operators” $e^{i\alpha\varphi(z)}$. Such vertex operators would be in many ways very interesting local fields. Still, reasons to exclude them are again similar — we want local fields that make sense with more general boundary conditions (including Neumann), and we prefer to focus on a simple well-behaved CFT. In the cases where free field is defined without additive constant ambiguities (for example with Dirichlet boundary conditions), extending our calculations to cases involving vertex operators seems feasible. But that would be a distraction from our primary objective, which is to provide a first complete CFT scaling limit result for a lattice model.

²The chiral part \mathcal{F} of this state space is exactly the Heisenberg vertex operator algebra (VOA), see e.g. [Kac98, FrBZ04, LeLi04]. General CFTs have more modules (for the appropriate VOA) in their state space — the bosonic CFT of interest here is particularly simple in this respect.

³In typical conformal field theories, there are particularly important local fields called *primary fields*, but they alone do not form a full conformal field theory for example for the reason that their OPEs involve other local fields (*descendants*) besides just the primary fields themselves.

1.4 Local fields in the lattice model. The discrete Gaussian Free Field (DGFF) is the most straightforward discretization of the GFF to finite subgraphs $\Omega_\delta \subset \delta\mathbb{Z}^2$ of the square lattice with mesh size $\delta > 0$ [She07]. It is a Gaussian measure associated with the discrete analogue

$$S(\phi) = \frac{1}{8\pi} \sum (\phi(\mathbf{z}) - \phi(\mathbf{w}))^2$$

of (1.1), where the sum ranges over pairs of nearest neighbor vertices $\mathbf{z}, \mathbf{w} \in \Omega_\delta$ in the discrete domain (the precise definition will be given in Section 4). Again, we consider ϕ as defined only up to an additive constant, i.e., our Gaussian process is just the discrete gradient of ϕ . This makes sense with both Dirichlet and Neumann boundary conditions (D/N) in arbitrary discrete domains Ω_δ .




Figure 1.1: A local field for a lattice model encodes a rule to construct concrete random variables at arbitrary points \mathbf{z} in arbitrary discrete domains Ω_δ of any mesh size $\delta > 0$ and for any choice of boundary conditions. These random variables take into account some fixed finite set of lattice sites around their points of insertion.

In a lattice model, the notion of a local field obviously should not refer to infinitesimal neighborhoods of a point — rather any finite lattice distance is to be considered microscopic, and we should allow local fields to “see” finite lattice regions around their positions of insertion. An example is provided by the following discretization of the gradient squared of the DGFF. Given a discrete domain $\Omega_\delta \subset \delta\mathbb{Z}^2$, a choice of boundary conditions (Dirichlet or Neumann), and a point $\mathbf{z} \in \Omega_\delta$ (such that all of its four neighbors also belong to Ω_δ), the discrete gradient squared is the random variable

$$\left(\frac{\phi(\mathbf{z} + \delta) - \phi(\mathbf{z} - \delta)}{2} \right)^2 + \left(\frac{\phi(\mathbf{z} + i\delta) - \phi(\mathbf{z} - i\delta)}{2} \right)^2 \quad (1.2)$$

on the probability space of the DGFF in Ω_δ . For different choices of mesh size δ , discrete domain, boundary conditions, and insertion point, we will want to view the correspondingly obtained random variable as an incarnation of the same abstract local field. So, following [GHP19] and [HKV22], the idea of a local field of the lattice model is a fixed rule, as

illustrated schematically in Figure 1.1, to construct random variables in a natural translation invariant fashion from the model's random configuration restricted to a finite lattice neighborhood. The same rule is to be applied in all discrete domains, with any boundary conditions, and at any point — provided just that the domain is large enough around the point so that the given finite neighborhood fits in it, which in scaling limit considerations is guaranteed for all small enough lattice mesh sizes δ .

For specificity and in accordance with the notion of local fields in our chosen free boson CFT, we require the rule defining a DGFF local field moreover be given by polynomials in the differences of the DGFF values. The space of such local *field polynomials* is denoted by \mathcal{P}_∇ (the precise definition will be given in Section 4). A crucial subtlety, emphasized and treated in [HKV22], is that among such field polynomials, there may be different ones which represent the same observable quantity, and should therefore be identified as local fields. Namely, it is possible that for two different field polynomials $P_1, P_2 \in \mathcal{P}_\nabla$, in all sufficiently large discrete domains Ω_δ , with all allowed boundary conditions, we have the coincidence of all expected values involving the random variables associated with P_1 and P_2 at a point \mathbf{z} and other random variables supported at least some microscopic distance $\mathcal{O}(\delta)$ away from that point. In this case P_1 and P_2 were called *correlation equivalent* in [HKV22], and the space $\mathcal{N} \subset \mathcal{P}_\nabla$ of field polynomials which are correlation equivalent to zero was called *null fields*. The appropriate definition of the space of *local fields*, corresponding to distinct observable quantities, is then the quotient

$$\mathcal{F}_\nabla := \mathcal{P}_\nabla / \mathcal{N}. \quad (1.3)$$

In [HKV22] it was proven that the space \mathcal{F}_∇ carries representations of the Heisenberg algebra and the Virasoro algebra: the Heisenberg algebra generators act by discrete Laurent-monomial weighted discrete contour integrations of discrete holomorphic currents, and the Virasoro generators are obtained from them by a Sugawara construction. By repeating the same with antiholomorphic variants, one also obtains second copies of Heisenberg and Virasoro algebras acting on the space \mathcal{F}_∇ , which commute⁴ with the original copies.

1.5 Main results. As described above, both the space \mathcal{F}_∇ of lattice model local fields and the space $\mathcal{F} \otimes \overline{\mathcal{F}}$ of local fields of the free boson CFT carry representations of two commuting copies of Heisenberg and Virasoro algebras. Our first main result is the following.

Theorem (Theorem 6.4, informally stated).

The Fock space $\mathcal{F} \otimes \overline{\mathcal{F}}$ of local fields of the free boson CFT and the space \mathcal{F}_∇ of local fields of the gradient of the discrete Gaussian Free Field are isomorphic,

$$\mathcal{F}_\nabla \cong \mathcal{F} \otimes \overline{\mathcal{F}},$$

⁴To obtain the commutation of these holomorphic and antiholomorphic chiral algebras acting on lattice model local fields, however, it is crucial to fix one convention about the discrete Laurent monomials from the published version of [HKV22]. The fixed convention will be given in our Proposition 3.4, and with this, we provide the details of the commutation in Proposition 4.2.

as representations of two commuting copies of the Heisenberg algebra and as representations of two commuting copies of the Virasoro algebra with central charge $c = 1$.

To spell out the meaning in the most concrete terms, the isomorphism of Theorem 6.4 provides a one-to-one correspondence

$$\left\{ \begin{array}{c} \text{lattice model} \\ \text{local fields} \end{array} \right\} \xleftrightarrow{1\text{-to-}1} \left\{ \begin{array}{c} \text{conformal field theory} \\ \text{local fields} \end{array} \right\}$$

whose two directions have the following interpretations:

- Any random variable, which is a polynomial in differences of the DGFF values at finitely many lattice neighbors of a point, determines by a translation invariant rule an abstract field polynomial $P \in \mathcal{P}_\nabla$, and then by the above isomorphism also a unique associated local field $F \in \mathcal{F} \otimes \overline{\mathcal{F}}$ of the free boson CFT. The CFT local field F is zero in the Fock space if and only if the field polynomial is null⁵, $P \in \mathcal{N}$.
- To any Fock space local field $F \in \mathcal{F} \otimes \overline{\mathcal{F}}$ of the free boson CFT, one can associate a corresponding lattice model local field and a representative abstract field polynomial $P \in \mathcal{P}_\nabla$, and therefore corresponding random variables in the DGFF at any point in any sufficiently large discrete domain. These random variables are polynomials in differences of the DGFF values at finitely many lattice neighbors of the point, given by a translation invariant formula. The field polynomial P associated to F is unique modulo null fields.⁶

The raison d'être of local fields is the formation of correlation functions. In turn, lattice discretizations of quantum field theories are meant to recover such correlation functions as appropriately renormalized scaling limits. Our second main result concerns the scaling limits of probabilistic correlations of local fields in the lattice model when conceptually correct renormalization is applied, and it identifies these scaling limits as CFT correlations. By general principles, the right renormalization of a local field should be determined via the structure of the Virasoro representations, in terms of its $L_0 + \bar{L}_0$ eigenvalue, and our result does precisely this.

Let us concretely illustrate scaling limit considerations with the field polynomial (1.2) from our earlier example. Perhaps contrary to a naive expectation for a squared discrete gradient, this local field has nonvanishing limits for its correlation functions without any renormalization. These limits, however, are trivial in the sense that they are constant as functions of the position where the field is inserted — they in fact accidentally capture a component corresponding to the CFT identity field! A field polynomial which actually corresponds to

⁵In practical terms, being null means that the DGFF random variables given by that formula in general discrete domains and with general boundary conditions have vanishing correlations with anything else at least some microscopic distance away.

⁶In practical terms again, this means uniqueness up to the addition of random variables that do not affect correlations with anything else at least some microscopic distance away.

the regularized gradient squared field in the CFT via the isomorphism $\mathcal{F} \otimes \overline{\mathcal{F}} \cong \mathcal{F}_\nabla$ of our first main result, is obtained by subtracting from (1.2) a suitable constant, $4\pi - 8$. After thus getting rid of the unwanted identity field component, we can apply the renormalization anticipated for the gradient squared, i.e., divide by the square of the lattice mesh δ . The random variables

$$\delta^{-2} \left(\left(\frac{\phi(\mathbf{z} + \delta) - \phi(\mathbf{z} - \delta)}{2} \right)^2 + \left(\frac{\phi(\mathbf{z} + \mathbf{i}\delta) - \phi(\mathbf{z} - \mathbf{i}\delta)}{2} \right)^2 - 4\pi + 8 \right).$$

then indeed have nontrivial scaling limits for their correlation functions. For more general field polynomials, the right procedure of subtracting accidental components and applying the right renormalization is not a priori obvious. Serendipitously, the isomorphism of our first main result directly gives the right counterparts to any CFT fields. Our second main theorem, given below, then identifies the right renormalization and scaling limits of correlations. It states, in particular, that the homogeneous components for renormalization are exactly the eigenspaces of $\mathbf{L}_0 + \overline{\mathbf{L}}_0$ in our space \mathcal{F}_∇ of lattice model local fields. Any Fock space field can be written as a sum of such homogeneous components.

Theorem (Theorem 7.2, informally stated).

Let $F_1, \dots, F_n \in \mathcal{F} \otimes \overline{\mathcal{F}}$ be n eigenvectors of $\mathbf{L}_0 + \overline{\mathbf{L}}_0$ with respective eigenvalues D_1, \dots, D_n . Let $\Omega \subset \mathbb{C}$ be an open simply-connected proper subset of the plane, and let $z_1, \dots, z_n \in \Omega$ be n distinct points in it. Let $(\Omega_\delta; \mathbf{z}_1^\delta, \dots, \mathbf{z}_n^\delta)$ be discrete domains $\Omega_\delta \subset \delta\mathbb{Z}^2$ with n marked vertices approximating $(\Omega; z_1, \dots, z_n)$ in Carathéodory sense as $\delta \rightarrow 0$. Fix a choice of boundary conditions, Dirichlet or Neumann (\mathbb{D}/\mathbb{N}). Then we have the following scaling limit of discrete GFF expected values

$$\frac{1}{\delta^{D_1 + \dots + D_n}} \mathbb{E}_{\Omega_\delta}^{\mathbb{D}/\mathbb{N}} \left[F_1^{\Omega_\delta}(\mathbf{z}_1^\delta) \cdots F_n^{\Omega_\delta}(\mathbf{z}_n^\delta) \right] \xrightarrow{\delta \rightarrow 0} \left\langle F_1(z_1) \cdots F_n(z_n) \right\rangle_\Omega^{\mathbb{D}/\mathbb{N}}$$

where each $F_j^{\Omega_\delta}(\mathbf{z}_j^\delta)$, is a random variable at \mathbf{z}_j^δ associated with the CFT field F_j via the isomorphism $\mathcal{F} \otimes \overline{\mathcal{F}} \cong \mathcal{F}_\nabla$, and the right hand side is the CFT correlation of the Fock space fields F_1, \dots, F_n in Ω with the chosen boundary conditions.

We view the combination of the two main results as a complete realization of the free boson CFT as the scaling limit of the discrete Gaussian Free Field.

1.6 Organization of the article and outlines of the proofs. Let us briefly outline the structure and ideas of the proofs of the two main results, and simultaneously describe the organization of the article.

There are two sections about the necessary preliminaries. Section 2 contains definitions and background related to the Heisenberg algebra, Virasoro algebra, and bosonic Fock spaces. Section 3 contains definitions, conventions and background about discrete complex analysis. Many of the details in these sections could just be consulted when they become relevant to the main arguments.

The gradient of the discrete Gaussian Free Field and its local fields are defined in Section 4, and the constructions of the two commuting Heisenberg and Virasoro algebra representations on the space of these lattice model local fields are given. This mostly amounts to recalling results from [HKV22] but with slight modifications to the exact setup (gradient of DGFF instead of GFF) and one minor change which is necessary to ensure the commutation of the two copies of both algebras.

The proof of the first main result, giving the isomorphism $\mathcal{F} \otimes \overline{\mathcal{F}} \cong \mathcal{F}_\nabla$, is carried out in Sections 5 – 6. The starting point (Corollary 4.4) is to note that an embedding $\mathcal{F} \otimes \overline{\mathcal{F}} \subset \mathcal{F}_\nabla$ is obtained just using the universal property and irreducibility of the Fock space $\mathcal{F} \otimes \overline{\mathcal{F}}$, as soon as a few very simple properties about the Heisenberg algebra actions on the constant field polynomial $1 \in \mathcal{P}_\nabla$ are verified. Therefore the essence of the proof is to get the opposite inclusion $\mathcal{F}_\nabla \subset \mathcal{F} \otimes \overline{\mathcal{F}}$. The rough idea is to show that dimensions of our lattice local fields are bounded from above by the corresponding dimensions in the embedded Fock space, which then rules out the existence of any superfluous local fields in the lattice model. Both spaces are actually infinite-dimensional, so such dimension bounds must be obtained in some appropriate finite-dimensional subspaces instead.

The first step, undertaken in Section 5, is to consider only those lattice local fields which correspond to linear polynomials in the gradient of the DGFF. The space $\mathcal{F}_\nabla^{\text{lin}} \subset \mathcal{F}_\nabla$ of such linear local fields of the lattice model is still infinite-dimensional, but it admits a filtration by finite-dimensional subspaces $\mathcal{F}_\nabla^{\text{lin}}(r)$ of linear local fields which have representative field polynomials supported in a lattice ball of size $r \in \mathbb{N}$. On the one hand, a version of the domain Markov property of the DGFF can be used to find essentially canonical representatives supported on the boundary of the lattice ball. This gives dimension upper bounds $\dim(\mathcal{F}_\nabla^{\text{lin}}(r)) \leq 4r - 1$ in the filtration (Lemma 5.9). On the other hand, for particular linear local fields in the embedded Fock space $\mathcal{F} \otimes \overline{\mathcal{F}} \subset \mathcal{F}_\nabla$ we perform explicit calculations with discrete contour integration and discrete Laurent monomials to get representatives of small enough radius of support, which allow us to conclude that such local fields already saturate the obtained dimension upper bound for $\mathcal{F}_\nabla^{\text{lin}}(r)$ (Lemma 5.4). Therefore the linear local fields in the embedded Fock space exhaust all linear local fields of the DGFF (Theorem 5.1). As an important by-product, we simultaneously obtain an explicit characterization of all linear null fields (Corollary 5.2).⁷

The remaining task of treating higher degree lattice local fields is then undertaken in Section 6. The key tool here is a suitable version of normal ordering on the lattice local fields. The combinatorics of the appropriate normal ordering is identical to the usual Wick products, but it is important to use infinite square grid quantities in the contractions.⁸ The fundamental observation about normal ordering is that field polynomials with linear null field factors behave like an ideal with respect to it: any linear null factor in a normal ordered

⁷The result is that the linear null fields are essentially just the discrete Laplacians of the DGFF at all possible locations. This is not unexpected: these vanishing Laplacians are the “equations of motion of the corresponding (classical) field theory”.

⁸The interpretation is that the infinite square grid plays a role similar to the specification of a local coordinate at the position of the field insertion. Conceptually, local field correlations in CFT require such local coordinate specifications; see, e.g., [FrBZ04] and [KaMa13].

product renders the full result null (Lemma 6.2). This makes normal ordering well-defined for local fields $F \in \mathcal{F}_\nabla = \mathcal{P}_\nabla/\mathcal{N}$, not merely for field polynomials $P \in \mathcal{P}_\nabla$. The proof of this well-definedness relies on the full classification of linear null fields from the previous section. Finally, one needs to relate normal ordering to the Heisenberg algebra actions on local fields (Lemma 6.3). In combination with the earlier result that all linear local fields are in the embedded Fock space, this yields the proof of the nontrivial inclusion $\mathcal{F}_\nabla \subset \mathcal{F} \otimes \overline{\mathcal{F}}$, and finishes the proof of the first main result: $\mathcal{F} \otimes \overline{\mathcal{F}} \cong \mathcal{F}_\nabla$ (Theorem 6.4).

The starting point of the proof of the second main result, about scaling limits, is to write formulas for the CFT correlations of Fock space fields as multiple contour integrals (Proposition 7.1 and Equation (7.11)), since the corresponding lattice local fields are by construction given by discrete contour integrals. The straightforward idea, then, is to expand expected values of DGFF local fields by Wick's formula and interpret the discrete contour integrals appearing in them as Riemann sum approximations of the integrals which give the CFT correlations. The discrete integrals involve derivatives of discrete Green's functions with Dirichlet or Neumann boundary conditions as well as discrete Laurent monomials. Both converge in the scaling limit to their continuum counterparts, with suitable scaling factors extracted from the Laurent monomials. In view of this, the convergence of the discrete integrals is entirely unsurprising, and the remaining part of the proof amounts to taking care of essentially combinatorial details from the specific discretizations. The scaling limit result (Theorem 7.2) then follows.

1.7 Scaling limit results in the literature. We finish the introduction by discussing prior results on conformally invariant scaling limits of correlation functions obtained for probabilistic lattice models. This is a major research topic with a long history, so it would be impossible to give a comprehensive account even of results on a single model. We simply aim to highlight some notable results for comparisons to the present work.

The two-dimensional critical Ising model is a quintessential lattice model which is believed to converge to a CFT in its scaling limit, so it provides a fruitful point of first comparisons.

A landmark result on the correlation functions of the Ising model was the proof by Chelkak, Hongler, and Izyurov of the existence of conformally covariant scaling limits for all spin correlation functions [CHI15]. That result had been preceded by the breakthrough work of Smirnov and others on conformally covariant scaling limits for fermionic observables and their generalizations [Smi10, ChSm12, Izy11] (which was in particular a key input to conformally invariant random geometry descriptions) and of energy correlations [Hon10, HoSm13]. Later the result was generalized to the proof of conformally covariant scaling limits of any mixed correlations of spins, disorders, fermions, and energies, in arbitrary finitely connected domains, with the most general boundary conditions allowed in the corresponding CFT [CHI21], see also [BIVW23]. This is essentially the full extent to which one can hope to understand the scaling limits of primary field correlation functions in the critical Ising model in planar domains. Despite the spectacular success, we argue that this does not yet establish a full CFT as the scaling limit of the Ising model, nor does it fully describe the

scaling limits of all locally defined random variables in the Ising model.

Regarding the former point, note that besides the finitely many primary fields, a full CFT contains a vast amount of other local fields, including infinite-dimensional spaces of descendant fields to every primary field. This is crucial, since it is this infinite-dimensional space of local fields that carries the algebraic structures that are the hallmarks of conformal field theory. Given the algebraic structure, one should furthermore proceed to verify that the scaling limits satisfy the characterizing conditions of CFT correlation functions.

Regarding the latter point, note that there are locally defined random variables in the Ising model more general than the spin at one point, the energy (the product of spins at two lattice neighbors), and the other specific ones mentioned above. Such general local random variables and their scaling limits, “pattern probabilities”, have been studied in [GHP19]. The scaling limits there were, however, taken with the generic renormalization for spin-flip even and spin-flip odd patterns separately, and the limits with these renormalizations therefore only retain the projections to the corresponding CFT primary fields. While impressive, the result does not, therefore, describe the probabilities of arbitrary (finely-tuned to cancel the leading scaling components) local patterns in their true asymptotic scale as a function of the lattice mesh, and it does not correspond to the renormalization of local fields by their scaling dimensions as required for field theory.

Complementarily, the articles [HKV22, Ada23] do identify the algebraic structures of CFTs acting on the infinite-dimensional spaces of local fields for three lattice models: the Ising model, discrete Gaussian Free Field, and discrete symplectic fermions (within the double-dimer model). On the other hand, in these works it is not proven that the space of local fields corresponds exactly to the CFT state space, and scaling limits of correlations of local fields are not treated.

Let us then compare with scaling limit results of discrete models believed to converge to the Gaussian Free Field.

Discrete-valued random height functions on a (discrete) lattice are natural higher-dimensional analogues of random walks, just like the GFF is a higher-dimensional generalization of the Brownian motion. It is natural to conjecture “functional central limit theorems” stating their convergence to GFF in the scaling limit, for at least suitable exact models or under suitable assumptions. Recent progress has been made, e.g., in [DHLRR15, CPST21, DKMO20], but scaling limits for these discretizations remain challenging.

Dimer configurations on suitable planar graphs can be encoded, following Thurston, into discrete-valued discrete height functions (modulo an additive constant) which bear resemblance with the random walk analogues but can be studied by more powerful combinatorial tools related to discrete complex analysis. In a celebrated result, Kenyon proved the conformal invariance of uniform domino tilings (dimers on the square grid) in the scaling limit, and showed that dimer height functions on Temperleyan domains tend to the GFF [Ken00, Ken01]. In more general domains, the same results still hold [Rus18], and, on non-simply connected surfaces, such height functions converge in the scaling limit to compactified free fields [Dub15, Bas23]. Moreover, double dimers, i.e., the superposition of two independent dimer covers, form loops which have been shown to converge in the scaling

limit to the conformal loop ensemble CLE_4 [Ken14, Dub18, BaCh21, BaWa23], which is naturally coupled with the GFF, and the loop scaling limit result can be seen as a convergence result of appropriate height level lines. Many of these results are stated in terms of convergence in law of globally defined random configurations, not in terms of local fields correlations. These and related works contain, however, also scaling limit results for correlations of dimer height gradients [Ken00] and for monomer correlations and the correlations of so-called electric operators (dimer analogues of vertex operators) [Dub15]. From the point of view of CFT, these should again be viewed as scaling limit results of primary field correlations.

There are also recent results on scaling limits of correlations in percolation [Cam24, CaFe24] and in the abelian sandpile model [PoRu17, CCR23]. The scaling limits of these models are believed to be logarithmic conformal field theories: they have non-diagonalizable Virasoro generators L_0 and \bar{L}_0 , and correspondingly local fields with much more intricate scaling behavior. Gaining full CFT scaling limit results for such lattice models could be particularly valuable, as it would shed light on a class of CFTs that remains extremely poorly understood while containing important examples for statistical physics.

Acknowledgments: DAC, DB, and KK were supported by the Research Council of Finland (project 346309: Finnish Centre of Excellence in Randomness and Structures, “FiRST”). The authors would like to thank Mikhail Basok, Nathanaël Berestycki, Dmitry Chelkak, Clément Hongler, Konstantin Izyurov, Richard Kenyon, Antti Kupiainen, and Wioletta Ruszel for valuable discussions.

2 Preliminaries: the Fock space

A central object in the present work is a Fock space, which should be interpreted as the space of polynomial local fields in the gradient and higher order derivatives of the bosonic free field. This interpretation will eventually be made concrete in Section 7.3. By way of preliminaries, we start in this section by explicitly defining the Fock space as a vector space and as a representation of the key algebraic structures present in the free boson conformal field theory (CFT).

In Section 2.1, we fix notation and conventions related to the Heisenberg algebra and its chargeless Fock representation, and we recall some essential properties of these. The Heisenberg algebra plays the role of a chiral symmetry algebra of the free boson CFT. In the full CFT with holomorphic and antiholomorphic chiralities, one has two commuting copies of the Heisenberg algebra, and the full Fock space of interest to us is built from Fock representations for these two chiral parts. A textbook reference for (a more general version of) these topics is [LeLi04, Ch. 6.2 – 6.3].

In Section 2.2 we recall how the Fock space becomes a representation of two commuting copies of Virasoro algebra via the Segal-Sugawara construction. Eigenvalues of particular Virasoro generators give rise to gradings of the Fock space by conformal dimensions, or equivalently by the scaling dimension and spin of the local fields. These scaling dimensions

will later feature crucially in the renormalization of fields in our scaling limit result.

To finish this background section, we make explicit observations in Section 2.3 about Virasoro primary fields in the Fock space.

2.1 Heisenberg algebra and Fock space. The **Heisenberg algebra** is the \mathbb{C} -vector space

$$\mathfrak{hei} = \left(\bigoplus_{k \in \mathbb{Z}} \mathbb{C} j_k \right) \oplus \mathbb{C} k, \quad (2.1)$$

equipped with the Lie algebra structure uniquely determined by the Lie brackets

$$[j_k, j_\ell] = k \delta_{k+\ell} k, \quad [k, j_\ell] = 0, \quad (2.2)$$

where we use the Kronecker delta with notation

$$\delta_n = \begin{cases} 1 & \text{if } n = 0 \\ 0 & \text{if } n \neq 0. \end{cases}$$

As is common, we will only consider representations of \mathfrak{hei} where the central element k acts as the identity, so the reader should feel free to think “ $k = 1$ ”. Somewhat more formally, we consider the associative algebra

$$\text{Hei} = \mathcal{U}(\mathfrak{hei}) / (k - 1) \quad (2.3)$$

obtained as the quotient of the universal enveloping algebra $\mathcal{U}(\mathfrak{hei})$ by the two-sided ideal generated by $k - 1$. A Poincaré-Birkhoff-Witt (PBW) basis of Hei consists of words

$$\cdots j_{-2}^{n_{-2}} j_{-1}^{n_{-1}} j_0^{n_0} j_1^{n_1} j_2^{n_2} \cdots$$

with only finitely many of the exponents $n_k \in \mathbb{Z}_{\geq 0}$ nonzero.

The most fundamental representation of the Heisenberg algebra \mathfrak{hei} is the (chargeless) **Fock representation**

$$\mathcal{F} = \text{Hei} / (j_k : k \geq 0) \quad (2.4)$$

obtained as a quotient of the Hei -module Hei by the left ideal generated by the j_k with nonnegative indices k . The vector $[1] \in \mathcal{F}$, i.e., the equivalence class of the algebra unit $1 \in \text{Hei}$ in the quotient (2.4), is called the **vacuum** vector in the Fock representation \mathcal{F} . From the PBW basis of Hei , one gets a basis of \mathcal{F} consisting of the vectors

$$j_{-k_m} \cdots j_{-k_2} j_{-k_1} [1] \quad \text{with} \quad m \in \mathbb{Z}_{\geq 0}, \quad 0 < k_1 \leq k_2 \leq \cdots \leq k_m. \quad (2.5)$$

Note that the zero-mode j_0 is in the centre of \mathfrak{hei} and acts as zero on the whole chargeless Fock space \mathcal{F} . Let us also mention, although this fact will not be directly used, that the

(chargeless) Fock space \mathcal{F} admits the structure of a *vertex operator algebra* (VOA) [LeLi04, Theorem 6.3.2]; it plays the role of the chiral symmetry algebra of the free boson conformal field theory (CFT).

As usual in bulk conformal field theory, we will in fact need two chiralities: the holomorphic and the antiholomorphic. Algebraically both are exactly the same, but we distinguish the latter by overline in the notation. The full (two-chiral) Lie algebra is the direct sum

$$\mathfrak{hei} \oplus \overline{\mathfrak{hei}}$$

of two commuting copies of the Heisenberg algebra (2.1), and the appropriate associative algebra is the tensor product

$$\text{Hei} \otimes \overline{\text{Hei}}$$

of two commuting copies of the associative algebra (2.3). The **(full) Fock space**

$$\mathcal{F} \otimes \overline{\mathcal{F}}$$

is also the tensor product of two copies of the Fock representation (2.4), with Hei acting on the first tensor factor and $\overline{\text{Hei}}$ acting on the second. Let us denote by $\mathbb{1} = [1] \otimes [1] \in \mathcal{F} \otimes \overline{\mathcal{F}}$ the vector obtained as the tensor product of the vacuum vectors of the two chiral Fock representations. A natural basis of the full Fock space is obtained by tensoring the bases of the form (2.5) in both chiral Fock representations; explicitly, denoting the generators of the two commuting copies by j_k and \bar{j}_k , that basis consists of vectors

$$\begin{aligned} j_{-k_m} \cdots j_{-k_2} j_{-k_1} \bar{j}_{-k'_m} \cdots \bar{j}_{-k'_2} \bar{j}_{-k'_1} \mathbb{1} &\in \mathcal{F} \otimes \overline{\mathcal{F}} \quad \text{with} \\ m, m' \in \mathbb{Z}_{\geq 0}, \quad 0 < k_1 \leq k_2 \leq \cdots \leq k_m, \quad 0 < k'_1 \leq k'_2 \leq \cdots \leq k'_{m'}. \end{aligned} \quad (2.6)$$

We note the following simple fact, which will be used for the easy half of our first main result.

Lemma 2.1. *Suppose that V is a representation of $\text{Hei} \otimes \overline{\text{Hei}}$, and $v \in V \setminus \{0\}$ is a nonzero vector such that $j_k v = 0$ and $\bar{j}_k v = 0$ for all $k \geq 0$. Then there exists a unique map $\mathcal{F} \otimes \overline{\mathcal{F}} \rightarrow V$ of representations of $\text{Hei} \otimes \overline{\text{Hei}}$ such that $\mathbb{1} \mapsto v$. This map is injective and onto the subrepresentation $(\text{Hei} \otimes \overline{\text{Hei}})v \subset V$. In particular if V is irreducible, then the map gives an isomorphism $\mathcal{F} \otimes \overline{\mathcal{F}} \cong V$ of representations of $\text{Hei} \otimes \overline{\text{Hei}}$.*

Proof. The condition $\mathbb{1} \mapsto v$ and the requirement of being a map of $\text{Hei} \otimes \overline{\text{Hei}}$ representations immediately fixes the values of the map on basis vectors (2.6),

$$j_{-k_m} \cdots j_{-k_2} j_{-k_1} \bar{j}_{-k'_m} \cdots \bar{j}_{-k'_2} \bar{j}_{-k'_1} \mathbb{1} \mapsto j_{-k_m} \cdots j_{-k_2} j_{-k_1} \bar{j}_{-k'_m} \cdots \bar{j}_{-k'_2} \bar{j}_{-k'_1} v,$$

so the uniqueness of such a map $\mathcal{F} \otimes \overline{\mathcal{F}} \rightarrow V$ is clear. The existence follows from a universal property (note that the properties assumed of the vector v correspond exactly to the generators of the left ideal factored out in the construction (2.4) of the Fock representation).

It is well-known and easy to check that \mathcal{F} is an irreducible representation of Hei , so $\mathcal{F} \otimes \overline{\mathcal{F}}$ is an irreducible representation of $\text{Hei} \otimes \overline{\text{Hei}}$. Therefore the kernel of the nonzero map $\mathcal{F} \otimes \overline{\mathcal{F}} \rightarrow V$ is zero, giving injectivity. By construction the map is onto the subrepresentation $(\text{Hei} \otimes \overline{\text{Hei}})_v \subset V$. If V is irreducible, this nonzero subrepresentation must be the whole space $(\text{Hei} \otimes \overline{\text{Hei}})_v = V$, so we get an isomorphism. \square

2.2 Virasoro algebra and the Sugawara construction. By construction, the Fock representation \mathcal{F} is a representation of the Heisenberg algebra. It also becomes a representation of the Virasoro algebra by the Sugawara construction, outlined below.

The **Virasoro algebra** is the \mathbb{C} -vector space

$$\mathbf{Vir} = \left(\bigoplus_{n \in \mathbb{Z}} \mathbb{C} L_n \right) \oplus \mathbb{C} C, \quad (2.7)$$

equipped with the Lie algebra structure uniquely determined by the Lie brackets

$$[L_n, L_m] = (n - m) L_{n+m} + \delta_{n+m} \frac{n^3 - n}{12} C, \quad [C, L_n] = 0, \quad (2.8)$$

where the Kronecker delta notation convention is as in (2.2). In CFT, one considers representations where the central element C acts as a fixed scalar multiple of identity, and the value c of the scalar is called the **central charge** of the CFT. For the free boson CFT of interest to us, the central charge is $c = 1$.

The following is the most basic variant of Sugawara constructions.

Lemma 2.2 (Sugawara construction). *On the Fock representation \mathcal{F} of (2.4), the operators defined by the formulas*

$$L_n = \frac{1}{2} \left(\sum_{k \geq 0} j_{n-k} j_k + \sum_{k < 0} j_k j_{n-k} \right) \quad \text{and} \quad C = \text{id}_{\mathcal{F}} \quad (2.9)$$

are well-defined (their action on any vector has only finitely many nonzero terms) and they make \mathcal{F} a representation of the Virasoro algebra with central charge $c = 1$.

Proof. This is well-known; see, e.g., [Kac98, Section 5.7] or [LeLi04, Theorem 6.2.16] for proofs in more general vertex operator algebra setting, or [HKV22, Theorem 4.10] for a similar calculation in a specific setting directly related to that of the present article. \square

The eigenvalues of L_0 give an important grading on the Fock representation \mathcal{F} . A direct calculation shows that the basis vectors (2.1) are L_0 -eigenvectors with eigenvalues $\Delta = k_1 + \dots + k_m$. We thus make \mathcal{F} an $\mathbb{Z}_{\geq 0}$ -graded vector space by

$$\mathcal{F} = \bigoplus_{\Delta \in \mathbb{Z}_{\geq 0}} \mathcal{F}_{\Delta}, \quad \text{where} \quad (2.10)$$

$$\mathcal{F}_{\Delta} = \text{span} \left\{ j_{-k_m} \cdots j_{-k_2} j_{-k_1} [1] \mid m \in \mathbb{Z}_{\geq 0}, 0 < k_1 \leq \dots \leq k_m, \sum_{j=1}^m k_j = \Delta \right\},$$

and then we have

$$L_0 v = \Delta v \quad \text{for any } v \in \mathcal{F}_\Delta.$$

From the Virasoro commutation relations (2.8) it also follows easily that $L_n \mathcal{F}_\Delta \subset \mathcal{F}_{\Delta-n}$, so L_n is a homogeneous operator of degree $-n$ with respect to the grading (2.10).

On the full (two-chiral) Fock space $\mathcal{F} \otimes \overline{\mathcal{F}}$, we similarly obtain two commuting actions of Virasoro algebra, both with central charge $c = 1$. We denote the generators of these by L_n and \bar{L}_n (we don't introduce notation for the central elements, since they simply act as identity on $\mathcal{F} \otimes \overline{\mathcal{F}}$). On the full Fock space $\mathcal{F} \otimes \overline{\mathcal{F}}$ we then have gradings by L_0 and \bar{L}_0 -eigenvalues,

$$\mathcal{F} \otimes \overline{\mathcal{F}} = \bigoplus_{\Delta, \bar{\Delta} \in \mathbb{Z}_{\geq 0}} \mathcal{F}_\Delta \otimes \overline{\mathcal{F}}_{\bar{\Delta}}. \quad (2.11)$$

The above bigrading by the pairs of L_0 and \bar{L}_0 -eigenvalues is said to be according to **conformal dimensions**. The $\mathbb{Z}_{\geq 0}$ -grading by eigenvalues of the sum $L_0 + \bar{L}_0$ is an $\mathbb{Z}_{\geq 0}$ -grading by **scaling dimensions**, and it will be important for us to determine the appropriate renormalization of lattice local fields in the scaling limit. The \mathbb{Z} -grading by the eigenvalues of the difference $L_0 - \bar{L}_0$ is also physically significant, interpreted as giving the *spins* of the corresponding local fields, but this last \mathbb{Z} -grading will not be needed in the present work.

2.3 Primary fields in the Fock space. Let us finish with a comment about primary fields, as these are generally of interest in CFTs. The Fock space $\mathcal{F} \otimes \overline{\mathcal{F}}$ is irreducible as a representation of $\text{Hei} \otimes \overline{\text{Hei}}$, and has the vacuum $\mathbb{1}$ as the unique (up to scalars) **Heisenberg primary state**, i.e., an eigenvector for j_0 and \bar{j}_0 annihilated by j_k and \bar{j}_k for $k > 0$. The situation is different for **Virasoro primary states**, i.e., eigenvectors for L_0 and \bar{L}_0 which are annihilated by L_n and \bar{L}_n for $n > 0$. The Heisenberg primary state $\mathbb{1}$ is also Virasoro primary, but in addition there is a countably infinite number of linearly independent Virasoro primaries in the Fock space, as made explicit in the following.

Remark 2.1. For every $a, b \in \mathbb{Z}_{\geq 0}$ there exists a Virasoro primary state in the Fock space $\mathcal{F} \otimes \overline{\mathcal{F}}$ with conformal weights $\Delta = a^2$, $\bar{\Delta} = b^2$, and this state is unique up to a multiplicative constant. \diamond

The above is a known fact and not logically used in our main results, so we give only a brief outline of the argument leading to it.

Sketch of proof of Remark 2.1. Due to the tensor product form of the Fock space $\mathcal{F} \otimes \overline{\mathcal{F}}$, it is enough to show that the (chiral) Fock representation \mathcal{F} has unique (up to scalars) Virasoro highest weight states of all highest weights $\Delta = a^2$ for $a \in \mathbb{Z}_{\geq 0}$. The Fock representation \mathcal{F} has an invariant inner product, with respect to which $L_n^\dagger = L_{-n}$ for $n \in \mathbb{Z}$. Consequently, any Virasoro submodule in \mathcal{F} has a complementary submodule, and any Virasoro submodule which is a highest weight module must be irreducible. The

argument from here on relies on dimension counting for the L_0 -eigenspaces in \mathcal{F} . On the one hand, the basis (2.5) readily shows that the dimension of $\text{Ker}(L_0 - \Delta) \subset \mathcal{F}$ is $p(\Delta)$, the number of integer partitions of Δ . On the other hand, the irreducible Virasoro highest weight modules of central charge $c = 1$ and highest weight $h = a^2$, with $a \in \mathbb{Z}_{\geq 0}$, has dimension $p(\Delta - a^2) - p(\Delta - (a+1)^2)$ for the corresponding eigenspace (this is the only needed fact whose proof is not completely elementary). Starting from the fact that the vacuum $[1] \in \mathcal{F}$ is a highest weight vector with highest weight $h = 0$, and inductively looking for the complementary subspaces to the subspace already found, and comparing dimensions of L_0 -eigenspaces, one sees that also (up to scalars unique) highest weight vectors of highest weights $1, 4, 9, 25, 36, \dots$ (and only these highest weights) can be found in \mathcal{F} . \square

Formulas for any of these Virasoro primary states in the Fock space $\mathcal{F} \otimes \overline{\mathcal{F}}$ can be obtained by straightforward computation: examples include

$$\mathbb{1} \quad \text{primary with conformal weights } \Delta = 0, \bar{\Delta} = 0 \quad (2.12)$$

$$j_{-1} \mathbb{1} \quad \text{primary with conformal weights } \Delta = 1, \bar{\Delta} = 0 \quad (2.13)$$

$$\bar{j}_{-1} \mathbb{1} \quad \text{primary with conformal weights } \Delta = 0, \bar{\Delta} = 1 \quad (2.14)$$

$$j_{-1} \bar{j}_{-1} \mathbb{1} \quad \text{primary with conformal weights } \Delta = 1, \bar{\Delta} = 1 \quad (2.15)$$

$$j_{-1} \left(\bar{j}_{-1}^4 + \frac{3}{2} j_{-2}^2 - 2 \bar{j}_{-3} \bar{j}_{-1} \right) \mathbb{1} \quad \text{primary with conformal weights } \Delta = 1, \bar{\Delta} = 4 \quad (2.16)$$

$$\begin{aligned} & \left(j_{-1}^9 + 9 j_{-2}^2 j_{-1}^5 - \frac{135}{4} j_{-2}^4 j_{-1} - 12 j_{-3} j_{-1}^6 + 90 j_{-3} j_{-2}^2 j_{-1}^2 + 40 j_{-3}^3 - 90 j_{-4} j_{-2} j_{-1}^3 \right. \\ & \quad \left. - 90 j_{-4} j_{-3} j_{-2} + \frac{135}{2} j_{-4}^2 j_{-1} + 36 j_{-5} j_{-1}^4 + 54 j_{-5} j_{-2}^2 - 72 j_{-5} j_{-3} j_{-1} \right) \mathbb{1} \\ & \quad \text{primary with conformal weights } \Delta = 9, \bar{\Delta} = 0. \end{aligned} \quad (2.17)$$

A few of these will appear throughout the present work. The vacuum (2.12) will correspond to the identity field of the CFT, as made concrete in Section 7.3. The fields corresponding to (2.13) and (2.14), will in various guises play central roles both in the discrete and in the continuum; these fields are called the holomorphic and antiholomorphic currents. The infinitely many other Virasoro primaries will not directly show up in our construction, but we find it worthwhile to remark that those primary fields are nevertheless present in our space of local fields.

3 Preliminaries: discrete complex analysis

A key feature of the lattice models that enabled equipping the spaces of their local fields with representations of Virasoro algebra in [HKV22] is a form of exact solvability expressed in terms of discrete complex analysis. This second background section collects notions and results about discrete complex analysis that we will build on.

Section 3.1 contains the definitions of the sublattices of square grids that we use as well as our conventions about discretizations of differential operators. In Section 3.2, we introduce discrete domains and discrete Green's functions with both Dirichlet and Neumann boundary conditions. The necessary scaling limit results for these discrete Green's functions are addressed in Section 3.3. The final notions of discrete complex analysis needed are discrete contour integration and discrete Laurent monomials, and we give the precise conventions and results about these in Section 3.4.

3.1 Square grids and discrete differential operators. Our discrete setup is based on the square grid \mathbb{Z}^2 and a few related graphs. The classification of lattice model local fields can be formulated on the unit-mesh square grid \mathbb{Z}^2 , but for the scaling limit results of Section 7, we need a general **mesh-size** $\delta > 0$ — the scaling limit amounts to letting δ tend to zero.

All of our square-grid graphs are embedded in the complex plane \mathbb{C} . Throughout the article, without further comments, we often identify vertices with complex numbers via the embedding, and edges and faces (square plaquettes) with complex numbers corresponding to the midpoints of the embedded edges or square plaquettes.

The infinite **square lattice** of mesh size $\delta > 0$ is the set

$$\delta\mathbb{Z}^2 := \{n\delta + im\delta \mid n, m \in \mathbb{Z}\} \subset \mathbb{C}. \quad (3.1)$$

We view $\delta\mathbb{Z}^2$ as a nearest-neighbor graph: an edge connects $\mathbf{z}, \mathbf{w} \in \delta\mathbb{Z}^2$ when $|\mathbf{z} - \mathbf{w}| = \delta$, and this adjacency relation is denoted by $\mathbf{z} \sim \mathbf{w}$. Since for discrete complex analysis we use also various other related grids, we for clarity we occasionally denote $\delta\mathbb{Z}_{\bullet}^2 := \delta\mathbb{Z}^2$ and call this the **primal lattice** to distinguish it from the variants. Two related grids are

$$\begin{aligned} \delta\mathbb{Z}_*^2 &:= \left\{ k\delta + i\ell\delta \mid k, \ell \in \mathbb{Z} + \frac{1}{2} \right\} && \text{(dual lattice)} \\ \delta\mathbb{Z}_m^2 &:= \left\{ \frac{\mathbf{z} + \mathbf{w}}{2} \mid \mathbf{z}, \mathbf{w} \in \delta\mathbb{Z}^2, \mathbf{z} \sim \mathbf{w} \right\} && \text{(medial lattice)} \end{aligned}$$

and we refer to the elements of $\delta\mathbb{Z}_*^2$ as dual vertices or plaquette centers, and to the elements of $\delta\mathbb{Z}_m^2$ as medial vertices or edge midpoints — see Figure 3.1 for an illustration. There is a bipartition $\delta\mathbb{Z}_m^2 = (\delta(\mathbb{Z} + \frac{1}{2}) \times \delta\mathbb{Z}) \cup (\delta\mathbb{Z} \times \delta(\mathbb{Z} + \frac{1}{2}))$ of the medial lattice to midpoints of **horizontal and vertical edges**. Finally, the bipartite grid

$$\delta\mathbb{Z}_{\diamond}^2 := \delta\mathbb{Z}_{\bullet}^2 \cup \delta\mathbb{Z}_*^2 \quad \text{(diamond lattice)}$$

is essentially dual to the medial lattice $\delta\mathbb{Z}_m^2$, and it is a natural domain of definition of some of our discrete functions. The elements of $\delta\mathbb{Z}_{\diamond}^2$ are referred to as diamond vertices.

Finite-difference operators are analogues of differential operators in the discrete setup, i.e., on the above lattices. For concreteness, let us only write the defining formulas of these finite-difference operators acting on complex-valued functions, although the action by




Figure 3.1: The δ -mesh square grid with the sublattices $\delta\mathbb{Z}_{\bullet}^2$, $\delta\mathbb{Z}_{*}^2$, and $\delta\mathbb{Z}_{\text{m}}^2$.

the same formulas will also be used on functions with values in other vector spaces. The (combinatorially normalized) discrete **holomorphic** and **antiholomorphic derivatives** of a function $f: \delta\mathbb{Z}_{\diamond}^2 \rightarrow \mathbb{C}$ on the diamond lattice are functions $\partial_{\sharp}f, \bar{\partial}_{\sharp}f: \delta\mathbb{Z}_{\text{m}}^2 \rightarrow \mathbb{C}$ on the medial lattice given by the formulas

$$\begin{aligned}\partial_{\sharp}f(\mathbf{z}) &:= \frac{f(\mathbf{z} + \frac{\delta}{2}) - f(\mathbf{z} - \frac{\delta}{2})}{2} - \mathfrak{i} \frac{f(\mathbf{z} + \frac{\mathfrak{i}\delta}{2}) - f(\mathbf{z} - \frac{\mathfrak{i}\delta}{2})}{2}, \\ \bar{\partial}_{\sharp}f(\mathbf{z}) &:= \frac{f(\mathbf{z} + \frac{\delta}{2}) - f(\mathbf{z} - \frac{\delta}{2})}{2} + \mathfrak{i} \frac{f(\mathbf{z} + \frac{\mathfrak{i}\delta}{2}) - f(\mathbf{z} - \frac{\mathfrak{i}\delta}{2})}{2}.\end{aligned}\tag{3.2}$$

Similarly, for a function $f: \delta\mathbb{Z}_{\text{m}}^2 \rightarrow \mathbb{C}$ on the medial lattice, $\partial_{\sharp}f, \bar{\partial}_{\sharp}f: \delta\mathbb{Z}_{\diamond}^2 \rightarrow \mathbb{C}$ are functions on the diamond lattice defined by exactly the same formulas as above. If a function f defined either on the diamond or the medial lattice satisfies $\bar{\partial}_{\sharp}f(\mathbf{z}) = 0$ at a medial or diamond vertex \mathbf{z} , it is said to be **discrete holomorphic** at \mathbf{z} . If it satisfies $\partial_{\sharp}f(\mathbf{z}) = 0$, then it is said to be **discrete antiholomorphic** at \mathbf{z} .

The discrete **Laplacian** of a function f on any of the lattices ($\delta\mathbb{Z}_{\bullet}^2$, $\delta\mathbb{Z}_{*}^2$, $\delta\mathbb{Z}_{\text{m}}^2$, or $\delta\mathbb{Z}_{\diamond}^2$) is the function $\Delta_{\sharp}f$ on the same lattice defined by

$$\Delta_{\sharp}f(\mathbf{z}) := f(\mathbf{z} + \delta) + f(\mathbf{z} + \mathfrak{i}\delta) + f(\mathbf{z} - \delta) + f(\mathbf{z} - \mathfrak{i}\delta) - 4f(\mathbf{z}).\tag{3.3}$$

Note that discrete Laplacian admits the following factorization:

$$\Delta_{\sharp} = 4\partial_{\sharp}\bar{\partial}_{\sharp} = 4\bar{\partial}_{\sharp}\partial_{\sharp}.$$

At times we furthermore need versions of the holomorphic and antiholomorphic derivative operators acting on functions defined only on the primal graph. For $f: \delta\mathbb{Z}_{\bullet}^2 \rightarrow \mathbb{C}$ we define $\partial_{\sharp}^{\bullet}f: \delta\mathbb{Z}_{\text{m}}^2 \rightarrow \mathbb{C}$ by

$$\partial_{\sharp}^{\bullet}f(\mathbf{z}) := f\left(\mathbf{z} + \frac{\delta}{2}\right) - f\left(\mathbf{z} - \frac{\delta}{2}\right) \quad \text{and} \quad \partial_{\sharp}^{\bullet}f(\mathbf{z}) := -\mathfrak{i}\left(f\left(\mathbf{z} + \frac{\mathfrak{i}\delta}{2}\right) - f\left(\mathbf{z} - \frac{\mathfrak{i}\delta}{2}\right)\right)\tag{3.4}$$

when \mathbf{z} is, respectively, a horizontal and a vertical edge. Similarly, we define $\bar{\partial}_{\#}^{\bullet} f: \delta\mathbb{Z}_{\mathbf{m}}^2 \rightarrow \mathbb{C}$ by

$$\bar{\partial}_{\#}^{\bullet} f(\mathbf{z}) := f\left(\mathbf{z} + \frac{\delta}{2}\right) - f\left(\mathbf{z} - \frac{\delta}{2}\right) \quad \text{and} \quad \bar{\partial}_{\#}^{\bullet} f(\mathbf{z}) := \mathbf{i} \left(f\left(\mathbf{z} + \frac{\mathbf{i}\delta}{2}\right) - f\left(\mathbf{z} - \frac{\mathbf{i}\delta}{2}\right) \right) \quad (3.5)$$

when \mathbf{z} is, respectively, a horizontal and a vertical edge. For a function $f: \mathbb{Z}_{\diamond}^2 \rightarrow \mathbb{C}$, we then have the following variants of factorizations of the discrete Laplacian

$$\partial_{\#} \bar{\partial}_{\#}^{\bullet} f(\mathbf{z}) = \bar{\partial}_{\#} \partial_{\#}^{\bullet} f(\mathbf{z}) = \begin{cases} \frac{1}{2} \Delta_{\#} f(\mathbf{z}) & \mathbf{z} \in \mathbb{Z}_{\bullet}^2 \\ 0 & \mathbf{z} \in \mathbb{Z}_{*}^2. \end{cases} \quad (3.6)$$

3.2 Discrete domains and Green's functions. By a **discrete domain** with mesh size $\delta > 0$ we mean a planar region bounded by a polygonal Jordan curve made of the edges of the δ -mesh square grid $\delta\mathbb{Z}^2$, such that the induced subgraph of $\delta\mathbb{Z}^2$ consisting of the primal vertices inside the Jordan curve is connected. We write $\Omega_{\delta} \subset \delta\mathbb{Z}^2$ for the set of primal vertices in the closure of the Jordan domain, and $\partial\Omega_{\delta}$ for the set of primal vertices on the Jordan curve. Vertices in $\Omega_{\delta} \setminus \partial\Omega_{\delta}$ are called interior vertices. By a mild abuse of terminology, we refer to Ω_{δ} as the discrete domain.

Discrete differential operators act naturally on functions defined on discrete domains, too, once we specify some boundary conditions.

The **discrete Dirichlet Laplacian** on Ω_{δ} is the operator $\Delta_{\Omega_{\delta}}^{\mathbf{D}}$ acting by the formula (3.3) on functions $f: \Omega_{\delta} \setminus \partial\Omega_{\delta} \rightarrow \mathbb{R}$ defined on the interior primal vertices, with the interpretation that the values of the function are zero outside the interior (in particular on the boundary $\partial\Omega_{\delta}$). This operator is negative definite, and its inverse is up to a sign the **discrete Dirichlet Green's function** in Ω_{δ} , defined as

$$G_{\Omega_{\delta}}^{\mathbf{D}} = -(\Delta_{\Omega_{\delta}}^{\mathbf{D}})^{-1}: \Omega_{\delta} \times \Omega_{\delta} \rightarrow \mathbb{R}, \quad (3.7)$$

setting again the values to zero if either of the arguments is not an interior vertex. Concretely, the discrete Dirichlet Green's function is determined by

$$\begin{cases} \Delta_{\#} G_{\Omega_{\delta}}^{\mathbf{D}}(\cdot, \mathbf{w}) = -\delta_{\mathbf{w}}(\cdot) & \text{on } \Omega_{\delta} \setminus \partial\Omega_{\delta} \\ G_{\Omega_{\delta}}^{\mathbf{D}}(\mathbf{z}, \mathbf{w}) = 0 & \text{if either } \mathbf{z} \in \partial\Omega_{\delta} \text{ or } \mathbf{w} \in \partial\Omega_{\delta}. \end{cases} \quad (3.8)$$

Put differently, the discrete Green's function is a kernel for the Dirichlet problem for the discrete Laplacian: for any $\psi: \Omega_{\delta} \setminus \partial\Omega_{\delta} \rightarrow \mathbb{R}$, the function $f(\mathbf{z}) = \sum_{\mathbf{w} \in \Omega_{\delta} \setminus \partial\Omega_{\delta}} G_{\Omega_{\delta}}^{\mathbf{D}}(\mathbf{z}, \mathbf{w}) \psi(\mathbf{w})$ is the unique solution to $\Delta_{\Omega_{\delta}}^{\mathbf{D}} f = -\psi$ with $f|_{\partial\Omega_{\delta}} \equiv 0$.

The **discrete Neumann Laplacian** is the operator $\Delta_{\Omega_{\delta}}^{\mathbf{N}}$ acting on functions $f: \Omega_{\delta} \rightarrow \mathbb{C}$ by the formula

$$(\Delta_{\Omega_{\delta}}^{\mathbf{N}} f)(\mathbf{z}) := \sum_{\substack{\mathbf{w} \in \Omega_{\delta} \\ |\mathbf{z} - \mathbf{w}| = \delta}} (f(\mathbf{w}) - f(\mathbf{z})). \quad (3.9)$$

The crucial difference to the Dirichlet case is that the coefficient of the “diagonal” term $f(\mathbf{z})$ on the right is proportional to the number of neighbors of \mathbf{z} in the discrete domain Ω_δ for the discrete Neumann Laplacian. The operator $\Delta_{\Omega_\delta}^N$ is negative semidefinite but not invertible, it has a one-dimensional kernel consisting of constant functions (the connectedness of the discrete domain is essential here). The degeneracy can be remedied by restricting to the space

$$\text{Fun}_\nabla(\Omega_\delta) := \left\{ f: \Omega_\delta \rightarrow \mathbb{R} \mid \sum_{\mathbf{z} \in \Omega_\delta} f(\mathbf{z}) = 0 \right\} \quad (3.10)$$

of **zero-average functions**: the Neumann Laplacian $\Delta_{\Omega_\delta}^N$ is injective on $\text{Fun}_\nabla(\Omega_\delta)$ and its range lies in $\text{Fun}_\nabla(\Omega_\delta)$, so

$$\Delta_{\Omega_\delta}^N: \text{Fun}_\nabla(\Omega_\delta) \rightarrow \text{Fun}_\nabla(\Omega_\delta)$$

is invertible. By fixing any probability mass function on the boundary, i.e., $b: \partial\Omega_\delta \rightarrow [0, 1]$ such that $\sum_{x \in \partial\Omega_\delta} b(x) = 1$, we can still define a (choice of the) **discrete Neumann Green’s function**

$$G_{\Omega_\delta}^N: \Omega_\delta \times \Omega_\delta \rightarrow \mathbb{R} \quad \text{by} \quad G_{\Omega_\delta}^N(\cdot, \mathbf{w}) := (\Delta_{\Omega_\delta}^N)^{-1}(b - \delta_{\mathbf{w}}). \quad (3.11)$$

We then have $\Delta_\# G_{\Omega_\delta}^N(\cdot, \mathbf{w}) = -\delta_{\mathbf{w}}(\cdot)$ in $\Omega_\delta \setminus \partial\Omega_\delta$, and for any $\psi \in \text{Fun}_\nabla(\Omega_\delta)$ the function $f(\mathbf{z}) := \sum_{\mathbf{w}} G_{\Omega_\delta}^N(\mathbf{z}, \mathbf{w})\psi(\mathbf{w})$ is the unique zero-average solution to $\Delta_{\Omega_\delta}^N f = -\psi$. Note also that all of the discrete derivatives of Section 3.1 are zero-average linear combinations of values, so the discrete derivatives of $G_{\Omega_\delta}^N$ with respect to its second argument are well-defined (independent of the choice of b).

3.3 Scaling limits of discrete Green’s functions. An ingredient of the proof of our scaling limit result for correlations of local fields is the convergence of discrete double derivatives of the discrete Green’s functions to their continuum counterparts. Below we introduce these continuum objects, and then state the convergence in the form that it will be used.

Let $\Omega \subset \mathbb{C}$ be a nonempty open simply-connected proper subset of the complex plane. Both the Dirichlet and Neumann Green’s functions in such domains Ω can be defined making use of conformal invariance. Choose a conformal map

$$\varphi: \Omega \rightarrow \mathbb{D}$$

to the unit disk $\mathbb{D} = \{z \in \mathbb{C} \mid |z| < 1\}$. The Dirichlet Green’s function in the unit disk is

$$G_{\mathbb{D}}^D(z, w) := \frac{-1}{2\pi} \log \left| \frac{z - w}{1 - z\bar{w}} \right| \quad \text{for } z, w \in \mathbb{D}, \ z \neq w,$$

and (a choice of) the Neumann Green's function in the unit disk is

$$G_{\mathbb{D}}^{\mathbf{N}}(z, w) := \frac{-1}{2\pi} \log \left| (z - w)(1 - z\bar{w}) \right| \quad \text{for } z, w \in \mathbb{D}, z \neq w.$$

We may then define the **Dirichlet Green's function** in Ω by

$$G_{\Omega}^{\mathbf{D}}(z, w) := G_{\mathbb{D}}^{\mathbf{D}}(\varphi(z), \varphi(w)) \quad \text{for } z, w \in \Omega, z \neq w,$$

since it is easy to check that the result does not depend on the chosen φ . For the **Neumann Green's function**, we can use

$$G_{\Omega}^{\mathbf{N}}(z, w) := G_{\mathbb{D}}^{\mathbf{N}}(\varphi(z), \varphi(w)) \quad \text{for } z, w \in \Omega, z \neq w,$$

which depends on the chosen conformal map φ , but the difference of any two choices is of the form $h_1(z) + h_2(w)$, with $h_1, h_2: \Omega \rightarrow \mathbb{R}$ harmonic.

These Green's functions are real-analytic functions on the configuration space

$$\text{Conf}_2(\Omega) = \{(z, w) \in \Omega \times \Omega \mid z \neq w\}.$$

They are Green's functions for the Laplacian $\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$ (where we write $z = x + iy$ with $x, y \in \mathbb{R}$) in the sense that

$$\begin{aligned} \Delta G_{\Omega}^{\mathbf{D/N}}(\cdot, w) &= 0 \quad \text{on } \Omega \setminus \{w\} \\ \text{and} \quad G_{\Omega}^{\mathbf{D/N}}(z, w) &= \frac{1}{2\pi} \log \frac{1}{|z - w|} + g_{\Omega}^{\mathbf{D/N}}(z, w). \end{aligned} \quad (3.12)$$

where $g_{\Omega}^{\mathbf{D/N}}: \Omega \times \Omega \rightarrow \mathbb{R}$ is real-analytic, and harmonic in both variables separately. By direct inspection, we also observe symmetricity of the Green's functions

$$G_{\Omega}^{\mathbf{D/N}}(z, w) = G_{\Omega}^{\mathbf{D/N}}(w, z) \quad \text{for } z, w \in \Omega, z \neq w.$$

We need to consider directional derivatives of discrete and continuum Green's functions with respect to both variables. The (combinatorially normalized) discrete directional derivative of a function $f: \Omega_{\delta} \rightarrow \mathbb{C}$ in a direction $\mu \in \{+1, \mathbf{i}, -1, -\mathbf{i}\}$ is

$$\nabla_{\#}^{\mu} f(\mathbf{z}) = \frac{f(\mathbf{z} + \frac{1}{2}\mu\delta) - f(\mathbf{z} - \frac{1}{2}\mu\delta)}{\mu}. \quad (3.13)$$

In continuum, the directional derivative of a function $f: \Omega \rightarrow \mathbb{C}$ in the direction of a complex number $\mu \in \mathbb{C}$ of unit modulus $|\mu| = 1$, is

$$\nabla^{\mu} f(z) = \left. \frac{d}{dt} f(z + \mu t) \right|_{t=0}. \quad (3.14)$$

When differentiating functions of several variables, we indicate the index or label of the variable acted on by an extra subscript in a hopefully self-explanatory way.

The following convergence results of discrete Dirichlet Green's functions are well-known.

Lemma 3.1. *When discrete domains $(\Omega_\delta)_{\delta>0}$ form an approximation to Ω in Carathéodory sense and $\mathbf{z}^\delta, \mathbf{w}^\delta \in \Omega_\delta$ are the closest points in the discrete domain Ω_δ to given points $z, w \in \Omega$, the discrete Dirichlet Green's functions converge*

$$G_{\Omega_\delta}^D(\mathbf{z}^\delta, \mathbf{w}^\delta) = G_\Omega^D(z, w) + o(1),$$

and the error term $o(1)$ is uniformly small for (z, w) on compact subsets of $\text{Conf}_2(\Omega)$.

Corollary 3.2. *When discrete domains $(\Omega_\delta)_{\delta>0}$ form an approximation to Ω in Carathéodory sense and $\mathbf{z}_m^\delta, \mathbf{w}_m^\delta$ are the closest edges of directions $\mu, \nu \in \{\pm 1, \pm i\}$ in the discrete domain Ω_δ to given points $z, w \in \Omega$, the discrete double derivatives of the Dirichlet Green's functions converge*

$$\delta^{-2}(\nabla_\#^\mu)_z(\nabla_\#^\nu)_w G_{\Omega_\delta}^D(\mathbf{z}_m^\delta, \mathbf{w}_m^\delta) = \nabla_z^\mu \nabla_w^\nu G_\Omega^D(z, w) + o(1), \quad (3.15)$$

and the error term $o(1)$ is uniformly small for (z, w) on compact subsets of $\text{Conf}_2(\Omega)$.

We also need a counterpart of the result (3.15) for Neumann boundary conditions. This is less well-known, so we sketch the proof below.

Lemma 3.3. *When discrete domains $(\Omega_\delta)_{\delta>0}$ form an approximation to Ω in Carathéodory sense and $\mathbf{z}_m^\delta, \mathbf{w}_m^\delta$ are the closest edges of directions $\mu, \nu \in \{\pm 1, \pm i\}$ in the discrete domain Ω_δ to given points $z, w \in \Omega$, the discrete double derivatives of the Neumann Green's functions converge*

$$\delta^{-2}(\nabla_\#^\mu)_z(\nabla_\#^\nu)_w G_{\Omega_\delta}^N(\mathbf{z}_m^\delta, \mathbf{w}_m^\delta) = \nabla_z^\mu \nabla_w^\nu G_\Omega^N(z, w) + o(1), \quad (3.16)$$

and the error term $o(1)$ is uniformly small for (z, w) on compact subsets of $\text{Conf}_2(\Omega)$.

Proof. Fix $w \in \Omega$ and a direction $\nu \in \{\pm 1, \pm i\}$, and let \mathbf{w}_m^δ be the edge of direction ν in Ω_δ that is nearest to w . Consider the discrete Dirichlet Green's function $G_{\Omega_\delta^*}^D$ on the dual graph $\Omega_\delta^* \subset \delta\mathbb{Z}_*^2$, and particularly its discrete derivative in the second variable at the medial vertex \mathbf{w}_m^δ in the direction $-i\nu$,

$$F(\mathbf{z}_*^\delta) := (\nabla_\#^{-i\nu})_w G_{\Omega_\delta^*}^D(\mathbf{z}_*^\delta, \mathbf{w}_m^\delta).$$

By construction, the function $F: \Omega_\delta^* \rightarrow \mathbb{R}$ is discrete harmonic except at $\mathbf{z}_*^\delta = \mathbf{w}_m^\delta \pm i\frac{\delta}{2}\nu$, where its discrete Laplacian is

$$\Delta_{\Omega_\delta^*}^D F(\mathbf{w}_m^\delta \pm i\frac{\delta}{2}\nu) = \pm 1.$$

In particular, we can define a function H (a discrete harmonic conjugate of F), by requiring

$$\nabla_\#^\mu H(\mathbf{z}_m^\delta) = \nabla_\#^{-i\mu} F(\mathbf{z}_m^\delta)$$

for all edges $\mathbf{z}_m^\delta \neq \mathbf{w}_m^\delta$; the single-valuedness of H around \mathbf{w}_m^δ relies on the opposite values of the Laplacian of F at the two adjacent dual vertices $\mathbf{w}_m^\delta \pm \mathbf{i}\frac{\delta}{2}$. From the construction, routine combinatorial considerations yield

$$\Delta_{\Omega_\delta}^N H(\mathbf{z}^\delta) = 0 \quad \text{for } \mathbf{z}^\delta \neq \mathbf{w}_m^\delta \pm \frac{\delta}{2}\nu \quad \text{and} \quad \Delta_{\Omega_\delta}^N H(\mathbf{w}_m^\delta \pm \frac{\delta}{2}\nu) = \pm 1,$$

where the Dirichlet boundary conditions for F are used for the Neumann Laplacian harmonicity of H on the boundary. As a consequence, fixing the additive constant in the harmonic conjugate so that H it is zero-average for definiteness, we find

$$H(\mathbf{z}^\delta) = -(\nabla_\#^\nu)_\mathbf{w} G_{\Omega_\delta}^N(\mathbf{z}^\delta, \mathbf{w}_m^\delta).$$

We can therefore write the double derivative of the discrete Neumann Green's function in the following form

$$\begin{aligned} \delta^{-2} (\nabla_\#^\mu)_\mathbf{z} (\nabla_\#^\nu)_\mathbf{w} G_{\Omega_\delta}^N(\mathbf{z}_m^\delta, \mathbf{w}_m^\delta) &= -\delta^{-2} \nabla_\#^\mu H(\mathbf{z}_m^\delta) \\ &= -\delta^{-2} \nabla_\#^{-\mathbf{i}\mu} F(\mathbf{z}_m^\delta) \\ &= -\delta^{-2} (\nabla_\#^{-\mathbf{i}\mu})_\mathbf{z} (\nabla_\#^{-\mathbf{i}\nu})_\mathbf{w} G_{\Omega_\delta}^D(\mathbf{z}_m^\delta, \mathbf{w}_m^\delta) \\ &= -\nabla_z^{-\mathbf{i}\mu} \nabla_w^{-\mathbf{i}\nu} G_\Omega^D(z, w) + o(1), \end{aligned}$$

where the last step uses (3.15). This shows that the double derivative has a scaling limit, given in terms of different (90° rotated) directional derivatives of the Dirichlet Green's function. It remains to observe the relationship

$$-\nabla_z^{-\mathbf{i}\mu} \nabla_w^{-\mathbf{i}\nu} G_\Omega^D(z, w) = \nabla_z^\mu \nabla_w^\nu G_\Omega^N(z, w).$$

of directional double derivatives of continuum Green's functions, which can for example be directly verified from the defining expressions of $G_\Omega^{D/N}$. \square

3.4 Discrete contour integration and discrete monomial functions. Two final notions of discrete complex analysis are employed in constructing the representations of Heisenberg and Virasoro algebras on the space of local fields of the discrete model as in [HKV22]: discrete analogues of contour integration and of Laurent monomials. These notions are only used as such on the square grid of unit-mesh $\delta = 1$. As a notational distinction, we typically denote vertices of the unit-mesh square grid and the associated diamond and medial lattices by $\mathbf{u}, \mathbf{v} \in \mathbb{Z}^2$, $\mathbf{u}_\diamond, \mathbf{v}_\diamond \in \mathbb{Z}_\diamond^2$, and $\mathbf{u}_m, \mathbf{v}_m \in \mathbb{Z}_m^2$. For the appropriate notion of discrete contour integration, we need yet one more lattice,

$$\mathbb{Z}_c^2 := \left\{ r + \mathbf{i}s \mid r, s \in \frac{1}{2}\mathbb{Z} + \frac{1}{4} \right\} \quad (\text{corner lattice})$$

whose vertices are referred to as corners. A **corner path** is a finite sequence $\gamma = (c_0, \dots, c_\ell)$ of consecutively nearest corners, i.e., $c_j \in \mathbb{Z}_c^2$ for $0 \leq j \leq \ell$ such that $|c_j - c_{j-1}| = \frac{1}{2}$ for

$0 < j \leq \ell$. Then, given two functions $f: \mathbb{Z}_\diamond^2 \rightarrow \mathbb{C}$ and $g: \mathbb{Z}_\text{m}^2 \rightarrow \mathbb{C}$ we define

$$\int_\gamma^\# f(\mathbf{u}_\diamond) g(\mathbf{u}_\text{m}) d^\# \mathbf{u} := \sum_{j=1}^\ell (c_j - c_{j-1}) f(\mathbf{u}_j^\diamond) g(\mathbf{u}_j^\text{m}), \quad \text{and} \quad (3.17)$$

$$\int_\gamma^\# f(\mathbf{u}_\diamond) g(\mathbf{u}_\text{m}) d^\# \bar{\mathbf{u}} := \sum_{j=1}^\ell \overline{(c_j - c_{j-1})} \overline{f(\mathbf{u}_j^\diamond) g(\mathbf{u}_j^\text{m})},$$

where $\mathbf{u}_j^\diamond \in \mathbb{Z}_\diamond^2$ and $\mathbf{u}_j^\text{m} \in \mathbb{Z}_\text{m}^2$ are the unique diamond vertex and medial vertex that have both c_j and c_{j-1} among their nearest corner vertices — see Figure 3.2. Note, furthermore, that bilinear discrete integration may be defined by the same formulas also when one of the functions f, g is complex-valued and the other one takes values in a complex vector space.




Figure 3.2: A discrete contour on the corner lattice. Each step (c_{j-1}, c_j) of it separates a medial vertex $\mathbf{u}_j^\text{m} \in \mathbb{Z}_\text{m}^2$ from a diamond vertex $\mathbf{u}_j^\diamond \in \mathbb{Z}_\diamond^2$.

A corner path $\gamma = (c_0, \dots, c_\ell)$ is said to be closed if $c_\ell = c_0$. If, moreover, the corners c_1, \dots, c_ℓ are distinct, we call γ a **corner contour**. A corner contour γ is essentially a Jordan curve on the lattice; it surrounds a bounded region. The set $\text{int}^\# \gamma \subset \mathbb{Z}_\diamond^2 \cup \mathbb{Z}_\text{m}^2$ of surrounded diamond and medial vertices is called the **interior** of γ , and we also denote $\text{int}_\diamond^\# \gamma := \text{int}^\# \gamma \cap \mathbb{Z}_\diamond^2$ and $\text{int}_\text{m}^\# \gamma := \text{int}^\# \gamma \cap \mathbb{Z}_\text{m}^2$ for the two types of surrounded vertices separately — see Figure 3.2. A corner contour $\gamma = (c_0, \dots, c_\ell)$ is said to be **positively oriented**, if the corners c_1, \dots, c_ℓ appear counterclockwise along γ seen as a Jordan curve. The symbol $\oint_\gamma^\#$ is used for discrete integrations (3.17) along positively oriented corner contours γ .

For calculations with discrete integration, the most important properties are the following analogues of Stokes' formula, contour deformation, and integration by parts.

Let γ be a positively-oriented corner contour. For any two functions $f: \mathbb{Z}_\diamond^2 \rightarrow \mathbb{C}$ and

$g: \mathbb{Z}_m^2 \rightarrow \mathbb{C}$ we have the **discrete Stokes' formulas**

$$\oint_{\gamma}^{\#} f(\mathbf{u}_{\diamond}) g(\mathbf{u}_m) d^{\#} \mathbf{u} = \mathfrak{i} \sum_{\mathbf{v}_{\diamond} \in \text{int}_{\diamond}^{\#} \gamma} f(\mathbf{v}_{\diamond}) \bar{\partial}_{\#} g(\mathbf{v}_{\diamond}) + \mathfrak{i} \sum_{\mathbf{v}_m \in \text{int}_m^{\#} \gamma} \bar{\partial}_{\#} f(\mathbf{v}_m) g(\mathbf{v}_m), \quad (3.18)$$

$$\oint_{\gamma}^{\#} f(\mathbf{u}_{\diamond}) g(\mathbf{u}_m) d^{\#} \bar{\mathbf{u}} = -\mathfrak{i} \sum_{\mathbf{v}_{\diamond} \in \text{int}_{\diamond}^{\#} \gamma} f(\mathbf{v}_{\diamond}) \partial_{\#} g(\mathbf{v}_{\diamond}) - \mathfrak{i} \sum_{\mathbf{v}_m \in \text{int}_m^{\#} \gamma} \partial_{\#} f(\mathbf{v}_m) g(\mathbf{v}_m). \quad (3.19)$$

In particular if the two functions f and g are discrete holomorphic on the symmetric differences of the appropriate subsets of the interior of two positively-oriented corner contours γ_1 and γ_2 (namely, $\bar{\partial}_{\#} f \equiv 0$ on $(\text{int}_m^{\#} \gamma_1 \setminus \text{int}_m^{\#} \gamma_2) \cup (\text{int}_m^{\#} \gamma_2 \setminus \text{int}_m^{\#} \gamma_1)$ and $\bar{\partial}_{\#} g \equiv 0$ on $(\text{int}_{\diamond}^{\#} \gamma_1 \setminus \text{int}_{\diamond}^{\#} \gamma_2) \cup (\text{int}_{\diamond}^{\#} \gamma_2 \setminus \text{int}_{\diamond}^{\#} \gamma_1)$), then one has the **contour deformation** equality

$$\oint_{\gamma_1}^{\#} f(\mathbf{u}_{\diamond}) g(\mathbf{u}_m) d^{\#} \mathbf{u} = \oint_{\gamma_2}^{\#} f(\mathbf{u}_{\diamond}) g(\mathbf{u}_m) d^{\#} \mathbf{u}.$$

A similar contour deformation equality holds for discrete antiholomorphic functions and the integrals $\oint_{\gamma_i}^{\#} \cdots d^{\#} \bar{\mathbf{u}}$.

The **discrete integration by parts** equalities


$$\begin{aligned} \oint_{\gamma}^{\#} f(\mathbf{u}_{\diamond}) \partial_{\#} h(\mathbf{u}_m) d^{\#} \mathbf{u} &= - \oint_{\gamma}^{\#} h(\mathbf{u}_{\diamond}) \partial_{\#} f(\mathbf{u}_m) d^{\#} \mathbf{u}, \quad \text{and} \\ \oint_{\gamma}^{\#} f(\mathbf{u}_{\diamond}) \bar{\partial}_{\#} h(\mathbf{u}_m) d^{\#} \bar{\mathbf{u}} &= - \oint_{\gamma}^{\#} h(\mathbf{u}_{\diamond}) \bar{\partial}_{\#} f(\mathbf{u}_m) d^{\#} \bar{\mathbf{u}} \end{aligned} \quad (3.20)$$

hold whenever $f, h: \mathbb{Z}_{\diamond}^2 \rightarrow \mathbb{C}$ are two discrete holomorphic functions on a discrete neighborhood of a corner contour γ .


The final necessary ingredient is discrete analogues of the Laurent monomials $z \mapsto z^n$ for $n \in \mathbb{Z}$. Such discrete Laurent monomials were constructed in [HKV22], but we crucially need a minor modification here — without this, the exact correspondence between the CFT local fields and the lattice model local fields simply does not work. We state the result of the modified construction here. The modification is in the exact coefficients used in property 5 below. We still refer to the original proof, as it remains in all essential ways similar.

Proposition 3.4 ([HKV22, Proposition 2.1]). *There exists a unique family of \mathbb{C} -valued functions $\{\mathbf{u} \mapsto \mathbf{u}^{[n]}\}_{n \in \mathbb{Z}}$ on $\mathbb{Z}_{\diamond}^2 \cup \mathbb{Z}_m^2$ that satisfies the following properties:*

1. *For all $n \in \mathbb{Z}$, the function $\mathbf{u} \mapsto \mathbf{u}^{[n]}$ has the same square-grid symmetries as the Laurent monomial $z \mapsto z^n$, i.e., $(\mathfrak{i}\mathbf{u})^{[n]} = \mathfrak{i}^n \mathbf{u}^{[n]}$ and $\bar{\mathbf{u}}^{[n]} = \overline{\mathbf{u}^{[n]}}$ for all $\mathbf{u} \in \mathbb{Z}_{\diamond}^2 \cup \mathbb{Z}_m^2$.*
2. *For all $\mathbf{u} \in \mathbb{Z}_{\diamond}^2 \cup \mathbb{Z}_m^2$, $\mathbf{u}^{[0]} = 1$ and, for all $n \in \mathbb{Z}$, $\partial_{\#} \mathbf{u}^{[n]} = n \mathbf{u}^{[n-1]}$.*
3. *For each $\mathbf{u} \in \mathbb{Z}_{\diamond}^2 \cup \mathbb{Z}_m^2$, there exists an $N \in \mathbb{N}$ such that $\mathbf{u}^{[n]} = 0$ for all $n \geq N$.*
4. *For $n < 0$, we have $\mathbf{u}^{[n]} \rightarrow 0$ as $|\mathbf{u}| \rightarrow \infty$.*




(a) The values of $\mathbf{u}^{[1]}$ are simply the complex points where the diamond and medial vertices \mathbf{u} are embedded.




(b) The values of $\mathbf{u}^{[3]}$ vanish in a finite neighborhood of the origin. By property 3 of Proposition 3.4, such vanishing neighborhoods grow with the order of the monomial.

Figure 3.3: The values of the positive Laurent monomials $\mathbf{u}^{[1]}$ and $\mathbf{u}^{[3]}$.



(a) The values of $\mathbf{u}^{[-1]}$.



(b) The values of $\mathbf{u}^{[-3]}$.

Figure 3.4: The values of the negative Laurent monomials $\mathbf{u}^{[-1]}$ and $\mathbf{u}^{[-3]}$.

5. The first negative-power monomial has the following explicit failure of discrete holomorphicity near the origin

$$\frac{1}{2\pi} \bar{\partial}_{\sharp} \mathbf{u}^{[-1]} = \frac{1}{2} \delta_{\mathbf{u},0} + \frac{1}{4} \sum_{|\mathbf{v}|=\frac{1}{2}} \delta_{\mathbf{u},\mathbf{v}} + \frac{1}{8} \sum_{\mathbf{v}=\frac{\pm 1 \pm i}{2}} \delta_{\mathbf{u},\mathbf{v}}.$$

6. For any $n \geq 0$ and all $\mathbf{u} \in \mathbb{Z}_{\diamond}^2 \cup \mathbb{Z}_{\mathbf{m}}^2$ we have $\bar{\partial}_{\sharp} \mathbf{u}^{[n]} = 0$. For any $n < 0$, we have $\bar{\partial}_{\sharp} \mathbf{u}^{[n]} = 0$ except at finitely many points⁹ $\mathbf{u} \in \mathbb{Z}_{\diamond}^2 \cup \mathbb{Z}_{\mathbf{m}}^2$.

7. For any $n, m \in \mathbb{Z}$, we have the discrete residue formula

$$\frac{1}{2\pi i} \oint_{\gamma}^{\sharp} \mathbf{u}_{\diamond}^{[n]} \mathbf{u}_{\mathbf{m}}^{[m]} d^{\sharp} \mathbf{u} = \delta_{n+m+1},$$

for any large enough positively-oriented corner contour γ that encircles the origin.

8. For any $n \in \mathbb{Z}$, as $|\mathbf{u}| \rightarrow \infty$, the discrete monomial has the asymptotics

$$\mathbf{u}^{[n]} = \mathbf{u}^n + o(|\mathbf{u}|^n). \quad (3.21)$$

Example values of some positive monomials are illustrated in Figure 3.3. These values remain identical to [HKV22]. Our modification only affects the values of negative monomials, such as those illustrated in Figure 3.4.

According to property 6 above, for each $k > 0$, there is a finite set

$$S^{[-k]} := \left\{ \mathbf{u} \in \mathbb{Z}_{\diamond}^2 \cup \mathbb{Z}_{\mathbf{m}}^2 \mid \bar{\partial}_{\sharp} \mathbf{u}^{[k]} \neq 0 \right\} \quad (3.22)$$

where the discrete holomorphicity of $\mathbf{u} \mapsto \mathbf{u}^{[-k]}$ fails. Let us also denote $S_{\mathbf{m}}^{[-k]} := S^{[-k]} \cap \mathbb{Z}_{\mathbf{m}}^2$. The values of $\bar{\partial}_{\sharp} \mathbf{u}^{[-3]}$ in Figure 3.5(b) display the shape of $S^{[-3]}$.

In our analysis, we will need the exact growth of $S_{\mathbf{m}}^{[-k]}$ with k . For that purpose, define discrete balls in $\mathbb{Z}_{\diamond}^2 \cup \mathbb{Z}_{\mathbf{m}}^2$ as

$$\mathbf{B}^{\sharp}(r) := \left\{ \mathbf{u} \in \mathbb{Z}_{\diamond}^2 \cup \mathbb{Z}_{\mathbf{m}}^2 \mid \|\mathbf{u}\|_1 \leq r \right\}, \quad (3.23)$$


where $\|\mathbf{u}\|_1 := \operatorname{Re}(\mathbf{u}) + \operatorname{Im}(\mathbf{u})$, and denote by $\mathbf{B}_{\mathbf{m}}^{\sharp}(r) := \mathbf{B}^{\sharp}(r) \cap \mathbb{Z}_{\mathbf{m}}^2$ the set of its medial vertices. We record the following facts for later reference.

Remark 3.1. From property 5 of Proposition 3.4 it follows that $S^{[-1]} \subset \mathbf{B}^{\sharp}(1)$ and that $S_{\mathbf{m}}^{[-1]} \subset \mathbf{B}_{\mathbf{m}}^{\sharp}(\frac{1}{2})$; see also Figure 3.5(a). For $k > 0$, we have


$$S^{[-k]} \subset \mathbf{B}^{\sharp}\left(\frac{k+1}{2}\right) \quad \text{and} \quad S_{\mathbf{m}}^{[-k]} \subset \mathbf{B}_{\mathbf{m}}^{\sharp}\left(\left\lfloor \frac{k}{2} \right\rfloor + \frac{1}{2}\right), \quad (3.24)$$

by virtue of the formula $\mathbf{u}^{[-k]} = \frac{(-1)^{k-1}}{(k-1)!} \partial_{\sharp}^{k-1} \mathbf{u}^{[-1]}$ from property 2 and the form (3.2) of the finite difference operator ∂_{\sharp} . \diamond

⁹This finite set of failure of discrete holomorphicity is made more explicit in the considerations below the proposition.



(a) The values of $\bar{\partial}_{\sharp} \mathbf{u}^{[-1]}$ are prescribed in property 5 of Proposition 3.4, and they are the single crucial modification to the monomial definition that we made compared to [HKV22].



(b) The values of $\bar{\partial}_{\sharp} \mathbf{u}^{[-3]}$ illustrate the shape of the finite set $S^{[-3]} \subset \mathbb{Z}_{\diamond}^2 \cup \mathbb{Z}_{\mathbf{m}}^2$ of failure of discrete holomorphicity of $\mathbf{u}^{[-3]}$.

Figure 3.5: The values of $\bar{\partial}_{\sharp} \mathbf{u}^{[-1]}$ and $\bar{\partial}_{\sharp} \mathbf{u}^{[-3]}$.

4 Discrete Gaussian Free Field and its local fields

In this section we define the discrete Gaussian free field (DGFF), its local fields, and the current modes that provide the representation of two commuting copies of the Heisenberg algebra on the space of these local fields. This essentially amounts to recalling results from [HKV22] — but two differences are worth pointing out. First of all, it is necessary to use the slightly modified discrete Laurent monomials of Section 3.4, to get the commutation of the two chiralities.¹⁰ Second, we allow for both Dirichlet and Neumann boundary conditions for the DGFF, and we correspondingly focus on local fields built from the discrete gradient of the DGFF, as this is exactly what corresponds to the (full) Fock space of local fields of the free boson CFT (see Section 7).

Our probabilistic models, the DGFF with Dirichlet and Neumann boundary conditions, are defined in Section 4.1. In Section 4.2 we introduce the discrete holomorphic and antiholomorphic currents of the DGFF, which are the quantities that enable the subsequent discrete complex analysis approach. Section 4.3 details what we mean by using abstract polynomials to specify fields, and how these become concrete random variables in the model in any fixed domain and either Dirichlet or Neumann boundary conditions. The final notion of local fields is derived from such field polynomials in Section 4.4, by forming equivalence classes of field polynomials which have indistinguishable correlations at macroscopic distances. The construction of the fundamental algebraic structure of local fields is given in Section 4.5: there are two commuting Heisenberg algebra actions on the space of local fields given by Laurent-monomial weighted discrete contour integrals of the discrete currents of the DGFF. Finally, the two commuting Virasoro actions on the space of local fields of the DGFF are obtained by Sugawara constructions. Section 4.6 recalls how the Sugawara construction applies in this discrete setup, and draws attention in particular to the grading of local fields by their scaling dimensions, i.e., eigenvalues of $L_0 + \bar{L}_0$.

4.1 DGFF with Dirichlet and Neumann boundary conditions. Let Ω_δ be a discrete domain on the δ -mesh square grid as in Section 3.2.

The discrete Gaussian free field with Dirichlet boundary conditions on Ω_δ could be defined as the centered Gaussian process $(\phi(\mathbf{z}))_{\mathbf{z} \in \Omega_\delta}$ indexed by the vertices \mathbf{z} of the discrete domain, whose covariance is

$$\mathbb{E}_{\Omega_\delta}^{\mathbf{D}} [\phi(\mathbf{z}) \phi(\mathbf{w})] = 4\pi G_{\Omega_\delta}^{\mathbf{D}}(\mathbf{z}, \mathbf{w})$$

where $G_{\Omega_\delta}^{\mathbf{D}} : \Omega_\delta \times \Omega_\delta \rightarrow [0, \infty)$ is the discrete Dirichlet Green's function (3.8). Equivalently,

¹⁰In [HKV22] only one chiral Heisenberg (and Virasoro) algebra action was written down. It is straightforward to define another chiral copy analogously. But the two chiral copies would not commute if one were to use the discrete Laurent monomials exactly as defined originally.

the probability density of the vector $(\phi(\mathbf{z}))_{\mathbf{z} \in \Omega_\delta \setminus \partial\Omega_\delta}$ is

$$\frac{1}{\mathcal{Z}_{\Omega_\delta}} \exp \left(-\frac{1}{8\pi} \sum_{\substack{\{\mathbf{z}, \mathbf{w}\} \subset \Omega_\delta \\ |\mathbf{z} - \mathbf{w}| = \delta}} (\phi(\mathbf{z}) - \phi(\mathbf{w}))^2 \right) \prod_{\mathbf{z} \in \Omega_\delta \setminus \partial\Omega_\delta} d\phi(\mathbf{z}),$$

where $\mathcal{Z}_{\Omega_\delta}$ is a normalization constant and $\phi(\mathbf{z})$ is interpreted as 0 if $\mathbf{z} \in \partial\Omega_\delta$. The quadratic form in the exponential above is a constant multiple of the discrete Dirichlet energy of ϕ .

Imagining the DGFF concretely as a pointwise defined random field is convenient, but for the purposes of this article we make two adjustments. First, for a closer parallel with the continuum Gaussian free field, it is more appropriate to consider the “mollified” values $\langle \Phi, f \rangle := \delta^2 \sum_{\mathbf{z} \in \Omega_\delta} \phi(\mathbf{z}) f(\mathbf{z})$, and view the DGFF as a process indexed by “test functions” f . Furthermore, the principal object for us is the gradient of the field, so we define the **discrete Gaussian free field with Dirichlet boundary conditions** on Ω_δ (Dirichlet DGFF for short) to be the centered Gaussian process

$$(\langle \Phi, f \rangle)_{f \in \text{Fun}_\nabla(\Omega_\delta)},$$

indexed by zero-average test functions $f \in \text{Fun}_\nabla(\Omega_\delta)$ as defined in (3.10), whose covariance is given by

$$\mathbb{E}_{\Omega_\delta}^{\text{D}} [\langle \Phi, f_1 \rangle \langle \Phi, f_2 \rangle] = 4\pi\delta^4 \sum_{\mathbf{z}_1, \mathbf{z}_2 \in \Omega_\delta} f_1(\mathbf{z}_1) G_{\Omega_\delta}^{\text{D}}(\mathbf{z}_1, \mathbf{z}_2) f_2(\mathbf{z}_2).$$

Discrete derivatives are recovered by suitable zero-average mollifiers, so they are well-defined random variables which we for convenience still denote by

$$\partial_\#^\bullet \phi(\mathbf{z}) = \delta^{-2} \langle \Phi, -\partial_\#^\bullet \delta_\mathbf{z} \rangle, \quad \bar{\partial}_\#^\bullet \phi(\mathbf{z}) = \delta^{-2} \langle \Phi, -\bar{\partial}_\#^\bullet \delta_\mathbf{z} \rangle, \quad \Delta_\# \phi(\mathbf{z}) = \delta^{-2} \langle \Phi, \Delta_\# \delta_\mathbf{z} \rangle, \quad \text{etc.}$$

In principle, the pointwise values $(\phi(\mathbf{z}))_{\mathbf{z} \in \Omega_\delta}$ and the test-function indexed field $(\langle \Phi, f \rangle)_{f \in \text{Fun}_\nabla(\Omega_\delta)}$ contain exactly the same information; $(\langle \Phi, f \rangle)_{f \in \text{Fun}_\nabla(\Omega_\delta)}$ straightforwardly determines the discrete derivatives, and one can “integrate the discrete derivatives” to recover values of ϕ anywhere in Ω_δ starting from the boundary $\partial\Omega_\delta$ where the values are zero. However, it is more appropriate to think of $\langle \Phi, \cdot \rangle$ as a description of the gradient of the free field only, since we will only ever probe the free field $(\langle \Phi, f \rangle)_{f \in \text{Fun}_\nabla(\Omega_\delta)}$ with finitely many finitely supported test functions f (see also Section 7.1 for the analogue in the continuum).

For Neumann boundary conditions, there is an inherent ambiguity about an additive constant in the free field, and in this case only the gradient of the field is meaningful. We define the **discrete Gaussian free field with Neumann boundary conditions** on Ω_δ (Neumann DGFF for short) to be the centered Gaussian process

$$(\langle \Phi, f \rangle)_{f \in \text{Fun}_\nabla(\Omega_\delta)},$$

indexed by zero-average test functions $f \in \text{Fun}_\nabla(\Omega_\delta)$, whose covariance is given by

$$\mathbb{E}_{\Omega_\delta}^{\text{N}} [\langle \Phi, f_1 \rangle \langle \Phi, f_2 \rangle] = -4\pi\delta^2 \sum_{\mathbf{z} \in \Omega_\delta} f_1(\mathbf{z}) ((\Delta_{\Omega_\delta}^{\text{N}})^{-1} f_2)(\mathbf{z}).$$

Note that if $G_{\Omega_\delta}^N$ is any choice of a discrete Neumann Green's function, this covariance may be alternatively written in a form where $G_{\Omega_\delta}^N$ is an approximate integral kernel,

$$\mathbb{E}_{\Omega_\delta}^N [\langle \Phi, f_1 \rangle \langle \Phi, f_2 \rangle] = 4\pi\delta^4 \sum_{\mathbf{z}_1, \mathbf{z}_2 \in \Omega_\delta} f_1(\mathbf{z}_1) G_{\Omega_\delta}^N(\mathbf{z}_1, \mathbf{z}_2) f_2(\mathbf{z}_2).$$

Although the pointwise values of the Neumann DGFF are not defined, the discrete derivatives are recovered by zero-average mollifiers, and by a mild abuse of notation we still denote the corresponding random variables by $\partial_\#^\bullet \phi(\mathbf{z})$, $\bar{\partial}_\#^\bullet \phi(\mathbf{z})$, $\Delta_\# \phi(\mathbf{z})$ as above.

Wick's formula applies to any centered Gaussians, in particular to both the Dirichlet and Neumann DGFF. Concretely, for any $f_1, \dots, f_n \in \text{Fun}_\nabla(\Omega_\delta)$, **Wick's formula** gives

$$\begin{aligned} \mathbb{E}_{\Omega_\delta}^{\text{D/N}} \left[\prod_{i=1}^n \langle \Phi, f_i \rangle \right] &= \sum_{P \in \text{Pair}(n)} \prod_{\{i,j\} \in P} \mathbb{E}_{\Omega_\delta}^{\text{D/N}} [\langle \Phi, f_i \rangle, \langle \Phi, f_j \rangle] \\ &= \delta^{2n} \sum_{\mathbf{z}_1, \dots, \mathbf{z}_n \in \Omega_\delta} f_1(\mathbf{z}_1) \cdots f_n(\mathbf{z}_n) \left(\sum_{P \in \text{Pair}(n)} \prod_{\{i,j\} \in P} 4\pi G_{\Omega_\delta}^{\text{D/N}}(\mathbf{z}_i, \mathbf{z}_j) \right), \end{aligned} \quad (4.1)$$

where the sums are over the set $\text{Pair}(n)$ of **pairings** P of the index set $\{1, 2, \dots, n\}$, i.e., partitions of the index set into subsets of size two each (for n odd, there are no such pairings and the empty sum is zero). It is convenient to keep in mind the more concise formal version

$$\text{“} \mathbb{E}_{\Omega_\delta}^{\text{D/N}} [\phi(\mathbf{z}_1) \cdots \phi(\mathbf{z}_n)] = \sum_{P \in \text{Pair}(n)} \prod_{\{i,j\} \in P} 4\pi G_{\Omega_\delta}^{\text{D/N}}(\mathbf{z}_i, \mathbf{z}_j) \text{”}$$

of (4.1), which is meaningful for the pointwise defined Dirichlet DGFF and can also be used for calculations with the Neumann DGFF if one ensures that only zero-average linear combinations are considered.

Remark 4.1. Let Ω_δ be a discrete domain and let $\mathbf{z} \in \Omega_\delta \setminus \partial\Omega_\delta$. Recall that the discrete Laplacian of the DGFF at \mathbf{z} is the random variable $\Delta_\# \phi(\mathbf{z}) = \delta^{-2} \langle \Phi, \Delta_\# \delta_\mathbf{z} \rangle$. For any $f \in \text{Fun}_\nabla(\Omega_\delta)$ the covariance of $\Delta_\# \phi(\mathbf{z})$ and $\langle \Phi, f \rangle$ simplifies to the following frequently useful formula

$$\begin{aligned} \mathbb{E}_{\Omega_\delta}^{\text{D/N}} [\Delta_\# \phi(\mathbf{z}) \langle \Phi, f \rangle] &= -4\pi\delta^2 f(\mathbf{z}) \\ \text{or formally} \quad \mathbb{E}_{\Omega_\delta}^{\text{D/N}} [\Delta_\# \phi(\mathbf{z}) \phi(\mathbf{w})] &= -4\pi \delta_{\mathbf{z}, \mathbf{w}}. \end{aligned} \quad (4.2)$$

Taking furthermore into account Wick's formula (4.1), a particularly simple consequence of (4.2) is that the correlation functions of $\Delta_\# \phi(\mathbf{z})$ of the form $\mathbb{E}_{\Omega_\delta}^{\text{D/N}} [\Delta_\# \phi(\mathbf{z}) \prod_{i=1}^n \langle \Phi, f_i \rangle]$ vanish when none of the supports of f_1, \dots, f_n contain \mathbf{z} . \diamond

4.2 Discrete holomorphic and antiholomorphic currents. Using the primal lattice discrete holomorphic and antiholomorphic derivatives $\partial_\#^\bullet$ and $\bar{\partial}_\#^\bullet$ defined in (3.4) and (3.5),

we define the **discrete holomorphic current** of the DGFF (either Dirichlet or Neumann) at an edge midpoint \mathbf{z} of Ω_δ as

$$J(\mathbf{z}) := \mathfrak{i} \partial_\#^\bullet \phi(\mathbf{z}) = \frac{\mathfrak{i}\delta}{2} \sum_{\substack{\mathbf{w} \in \Omega_\delta \\ |\mathbf{w} - \mathbf{z}| = \delta/2}} \frac{\phi(\mathbf{w})}{\mathbf{w} - \mathbf{z}}, \quad (4.3)$$

where, as usual, the zero-average linear combination of DGFF values on the right should properly be interpreted as $\langle \Phi, \dots \rangle$ with the suitable zero-average function inserted. Similarly, we define the **discrete antiholomorphic current** at \mathbf{z} as

$$\bar{J}(\mathbf{z}) := -\mathfrak{i} \bar{\partial}_\#^\bullet \phi(\mathbf{z}) = \frac{-\mathfrak{i}\delta}{2} \sum_{\substack{\mathbf{w} \in \Omega_\delta \\ |\mathbf{w} - \mathbf{z}| = \delta/2}} \frac{\phi(\mathbf{w})}{\bar{\mathbf{w}} - \bar{\mathbf{z}}}. \quad (4.4)$$

The following discrete holomorphicity/antiholomorphicity of correlations of the currents is a consequence of the factorizations (3.6) of the Laplacian and the vanishing of correlations of the discrete Laplacian of the free field (see Remark 4.1).

Lemma 4.1 ([HKV22, Lemma 3.5]¹¹). *Let $\Omega_\delta \subset \delta\mathbb{Z}^2$ be a discrete domain and let $f \in \text{Fun}_\nabla(\Omega_\delta)$. Then the functions $\mathbf{z} \mapsto \mathbb{E}_{\Omega_\delta}^{\text{D/N}}[J(\mathbf{z}) \langle \Phi, f \rangle]$ and $\mathbf{z} \mapsto \mathbb{E}_{\Omega_\delta}^{\text{D/N}}[\bar{J}(\mathbf{z}) \langle \Phi, f \rangle]$ are, respectively, discrete holomorphic and discrete antiholomorphic on the set*

$$(\Omega_\delta \cap \delta\mathbb{Z}_\diamond^2) \setminus (\partial\Omega_\delta \cup \{\mathbf{w} \in \Omega_\delta \mid f(\mathbf{w}) \neq 0\})$$

of interior diamond vertices excluding the support of f .

4.3 Field polynomials. Recall the rough idea of a local field in a field theory: a quantity that is determined by the values of the basic fields of the theory in a microscopic neighborhood of its point of insertion, in a manner that does not depend on the domain, boundary conditions, or other details. In the lattice model context, an abstract local field is meant to encode a rule to construct concrete random variables from the basic degrees of freedom in a finite set of lattice sites around any point of any discrete domain — see Figure 1.1 again for illustration.

We define a **field polynomial** of the (gradient of the) DGFF to be a polynomial in indeterminates $X(\mathbf{u})$ indexed by the points $\mathbf{u} \in \mathbb{Z}^2$ of the unit-mesh square grid, i.e., an element of the polynomial ring

$$\mathcal{P}_\nabla := \mathbb{C}[X(\mathbf{u}) : \mathbf{u} \in \mathbb{Z}^2]. \quad (4.5)$$

Any such polynomial is “local” in the sense that there are only finitely many terms in the polynomial. We define the **support** of a field polynomial $F \in \mathcal{P}_\nabla$ to be the minimally chosen (finite) subset of those $\mathbf{u} \in \mathbb{Z}^2$ such that $X(\mathbf{u})$ appears in F ,

$$\text{Supp } F := \bigcap \left\{ S \subset \mathbb{Z}^2 \mid F \in \mathbb{C}[X(\mathbf{u}) : \mathbf{u} \in S] \right\}. \quad (4.6)$$

¹¹In the article [HKV22] there is a factor two mistake in this statement and the proof has an error which is simply fixed by correctly using the factorization (3.6).

When a discrete domain $\Omega_\delta \subset \delta\mathbb{Z}^2$, a choice of boundary conditions, and a point $\mathbf{z} \in \Omega_\delta$ are given, we define an **evaluation** of field polynomials

$$\text{ev}_{\mathbf{z}}^{\Omega_\delta} : \mathcal{P}_\nabla \rightarrow \{\text{random variables for the DGFF in } \Omega_\delta\}.$$

The evaluation of general field polynomials will be determined by linear extension of an evaluation of monomials $X(\mathbf{u}_1) \cdots X(\mathbf{u}_n)$. When $\mathbf{z} + \delta\mathbf{u}_1, \dots, \mathbf{z} + \delta\mathbf{u}_n \in \Omega_\delta$, such a monomial is evaluated to

$$\text{ev}_{\mathbf{z}}^{\Omega_\delta} : X(\mathbf{u}_1) \cdots X(\mathbf{u}_n) \mapsto \left(\phi(\mathbf{z} + \delta\mathbf{u}_1) - \phi(\mathbf{z}) \right) \cdots \left(\phi(\mathbf{z} + \delta\mathbf{u}_n) - \phi(\mathbf{z}) \right).$$

This is the only case we actually care about, because when $(\Omega_\delta)_{\delta>0}$ are discrete approximations to a continuum domain, i.e., an open set $\Omega \subset \mathbb{C}$, then we indeed have $\mathbf{z} + \delta\mathbf{u}_j \in \Omega_\delta$ for any small enough lattice mesh $\delta > 0$. For completeness of the definition, monomials $X(\mathbf{u}_1) \cdots X(\mathbf{u}_n)$ such that $\mathbf{z} + \delta\mathbf{u}_j \notin \Omega_\delta$ for some j are evaluated to zero (this is somewhat arbitrary, but the exact choice in this irrelevant case does not matter).

Example 4.1. For any edge midpoint $\mathbf{u}_m \in \mathbb{Z}_m^2$ of the unit-mesh square lattice, the linear combinations

$$\begin{aligned} \mathbf{j}(\mathbf{u}_m) &:= \mathbf{i} \partial_{\#}^\bullet X(\mathbf{u}_m) = \begin{cases} X(\mathbf{z} + \frac{1}{2}) - X(\mathbf{z} - \frac{1}{2}) & \text{if } \mathbf{u}_m \text{ is on a horizontal edge} \\ -\mathbf{i} X(\mathbf{z} + \frac{\mathbf{i}}{2}) + \mathbf{i} X(\mathbf{z} - \frac{\mathbf{i}}{2}) & \text{if } \mathbf{u}_m \text{ is on a vertical edge} \end{cases} \\ \bar{\mathbf{j}}(\mathbf{u}_m) &:= -\mathbf{i} \bar{\partial}_{\#}^\bullet X(\mathbf{u}_m) = \begin{cases} X(\mathbf{z} + \frac{1}{2}) - X(\mathbf{z} - \frac{1}{2}) & \text{if } \mathbf{u}_m \text{ is on a horizontal edge} \\ \mathbf{i} X(\mathbf{z} + \frac{\mathbf{i}}{2}) - \mathbf{i} X(\mathbf{z} - \frac{\mathbf{i}}{2}) & \text{if } \mathbf{u}_m \text{ is on a vertical edge} \end{cases} \end{aligned}$$

are field polynomials. When evaluated at a point $\mathbf{z} \in \Omega_\delta$ of a discrete domain, they give rise to the random variables

$$\text{ev}_{\mathbf{z}}^{\Omega_\delta}(\mathbf{j}(\mathbf{u}_m)) = \mathbf{J}(\mathbf{z} + \delta\mathbf{u}_m) \quad \text{and} \quad \text{ev}_{\mathbf{z}}^{\Omega_\delta}(\bar{\mathbf{j}}(\mathbf{u}_m)) = \bar{\mathbf{J}}(\mathbf{z} + \delta\mathbf{u}_m)$$

which are the values of the discrete holomorphic and discrete antiholomorphic currents, (4.3) and (4.4), on an edge at a fixed finite number of lattice steps away from the point \mathbf{z} . In particular the field polynomials $\mathbf{j}(\frac{\mathbf{i}}{2})$, $\mathbf{j}(\frac{1}{2})$, $\mathbf{j}(\frac{-\mathbf{i}}{2})$ and $\mathbf{j}(\frac{-1}{2})$ evaluate to the discrete holomorphic currents on the edge to the north, east, south, and west of \mathbf{z} , respectively. Trusting that no confusion arises, the field polynomial valued functions $\mathbf{j} : \mathbb{Z}_m^2 \rightarrow \mathcal{P}_\nabla$ and $\bar{\mathbf{j}} : \mathbb{Z}_m^2 \rightarrow \mathcal{P}_\nabla$ on the unit-mesh medial lattice will still be referred to as the discrete holomorphic and antiholomorphic currents. \diamond

4.4 Local fields. It can happen that two different field polynomials produce random variables which have indistinguishable correlation functions with anything at a macroscopic distance away from their insertion. Such field polynomials do not then really represent different observable quantities in a field theory, so we want to identify them. Following [HKV22], we now define precisely the equivalence relation, and then define local fields as the equivalence classes. These local fields are going to be the main object of our interest.

We say that a field polynomial $F \in \mathcal{P}_\nabla$ is a **null field** if

$$\begin{aligned}
& \text{for} \quad \text{all discrete domains } \Omega_\delta \subset \delta\mathbb{Z}^2, \\
& \quad \text{both choices of boundary conditions (Dirichlet or Neumann)} \\
& \quad \text{any point } \mathbf{z} \in \Omega_\delta \\
& \quad \text{any } n \in \mathbb{N} \text{ and all test functions } f_1, \dots, f_n \in \text{Fun}_\nabla(\Omega_\delta) \\
& \text{such that} \quad \mathbf{z} + \delta \text{Supp } F \subset \Omega_\delta \setminus \bigcup_{j=1}^n \text{supp } f_j \\
& \text{we have} \quad \mathbb{E}_{\Omega_\delta}^{\text{D/N}}[(\text{ev}_{\mathbf{z}}^{\Omega_\delta} F) \langle \Phi, f_1 \rangle \cdots \langle \Phi, f_n \rangle] = 0.
\end{aligned} \tag{4.7}$$

We let $\mathcal{N} \subset \mathcal{P}_\nabla$ denote the set of null fields.

Example 4.2. Any monomial $X(\mathbf{u}_1) \cdots X(\mathbf{u}_n) \in \mathcal{P}_\nabla$ with $\mathbf{u}_j = 0$ for some j is trivially null, since in its evaluation as a random variable, the factor $X(\mathbf{u}_j)$ becomes $\phi(\mathbf{z} + \delta\mathbf{u}_j) - \phi(\mathbf{z}) = \phi(\mathbf{z}) - \phi(\mathbf{z}) = 0$. \diamond

Example 4.3. The discrete Laplacians of the basic field X are null fields: it follows easily from (4.2) and Wick's formula (4.1) that for any $\mathbf{u} \in \mathbb{Z}^2$ we have

$$\Delta_\# X(\mathbf{u}) = \sum_{\substack{\mathbf{v} \in \mathbb{Z}^2 \\ \mathbf{v} \sim \mathbf{u}}} (X(\mathbf{v}) - X(\mathbf{u})) \in \mathcal{N}. \tag{4.8}$$

In view of the factorization (3.6), as a consequence we get that for any $\mathbf{u}_\diamond \in \mathbb{Z}_\diamond^2$

$$\bar{\partial}_\# \mathbf{j}(\mathbf{u}_\diamond) \in \mathcal{N} \quad \text{and} \quad \partial_\# \bar{\mathbf{j}}(\mathbf{u}_\diamond) \in \mathcal{N}. \tag{4.9}$$

This observation is closely related to Lemma 4.1; it is a discrete holomorphicity (resp. antiholomorphicity) property of the current, now viewed as a field polynomial. \diamond

The null fields (4.8) are closely related to the equations of motion of the theory: the minimizers of the discrete Dirichlet energy are discrete harmonic functions. It is natural to anticipate that the null fields $\Delta_\# X(\mathbf{u})$ will play a particularly important role.

We record one more explicit form of null fields that will appear in later calculations.

Example 4.4. For any $\mathbf{u}, \mathbf{v} \in \mathbb{Z}^2$, the local field

$$(\Delta_\# X(\mathbf{u}))X(\mathbf{v}) + 4\pi(\delta_{\mathbf{u},\mathbf{v}} - \delta_{\mathbf{u},0})$$

is null again by (4.2) and Wick's formula (4.1). \diamond

The space of **local fields** of (the gradient of) the DGFF is now defined as the quotient

$$\mathcal{F}_\nabla := \mathcal{P}_\nabla / \mathcal{N}. \tag{4.10}$$

In other words, we view two field polynomials $F_1, F_2 \in \mathcal{P}_\nabla$ as correlation equivalent if their difference $F_1 - F_2$ is null. Since working with concrete field polynomial representatives is

still often convenient, we write $F + \text{Null}$ for the local field that is the equivalence class of $F \in \mathcal{P}_\nabla$ in \mathcal{F}_∇ .

Note that while $\mathcal{N} \subset \mathcal{P}_\nabla$ is evidently a vector subspace, Examples 4.4 and 4.3 show that it is not an ideal in $\mathcal{P}_\nabla = \mathbb{C}[X(\mathbf{u}) : \mathbf{u} \in \mathbb{Z}^2]$ with the usual polynomial ring structure. In particular, the space of local fields (4.10) does not inherit any obvious multiplication from the polynomial ring.

Different representatives of the same equivalence class in \mathcal{F}_∇ may of course have different supports (4.6). A meaningful and useful notion, however, is the **minimal radius of support** of a local field $F + \text{Null}$, defined as the smallest r such that a representative with a support in a ball (3.23) of r lattice units exists, i.e.,

$$R_{\text{Supp}}(F + \text{Null}) := \min \left\{ r \in \mathbb{Z}_{\geq 0} \mid \exists \tilde{F} \in \mathcal{P}_\nabla : \tilde{F} - F \in \mathcal{N} \text{ and } \text{Supp } \tilde{F} \subset \mathbf{B}^\sharp(r) \right\}.$$

Since exactly determining the minimal radius of support for interesting fields is not straightforward (requires controlling all choices of representatives), we postpone examples for later.

Understanding the space (4.10) of local fields is at the heart of the present work. Let us pause to comment on why this is nontrivial. The space $\mathcal{F}_\nabla = \mathcal{P}_\nabla / \mathcal{N}$ is formed as a quotient by null fields, which are defined by a philosophically motivated but very implicit condition. No apparent tractable procedure exists to decide whether two polynomials $F_1, F_2 \in \mathcal{P}_\nabla$ differ by a null field: by definition this would involve inspecting correlation functions in all discrete domains, with all boundary conditions, at all points, and with all possible other fields. With enough algebraic structure on the space \mathcal{F}_∇ and some concretely verifiable dimension bounds, we will, however, ultimately arrive at a fully explicit description of the quotient $\mathcal{F}_\nabla = \mathcal{P}_\nabla / \mathcal{N}$.




Figure 4.1: A visualization of the definition (4.11) for $k = -2$: In purple the support of a representative F , in red the set $S^{[-2]}$ of nonholomorphicity of the monomial $\mathbf{u} \mapsto \mathbf{u}^{[-2]}$, and in light blue a choice of positively oriented corner contour γ ; all of them laid on the infinite square grid \mathbb{Z}^2 and its sublattices \mathbb{Z}_\diamond^2 and \mathbb{Z}_m^2 .

4.5 Current modes. On the space (4.10) of correlation equivalence classes of local fields there are operators closely analogous to the Laurent modes of the currents as operators in conformal field theory.

The **holomorphic current modes** J_k and **antiholomorphic current modes** \bar{J}_k , for $k \in \mathbb{Z}$, are the linear operators on the space of fields \mathcal{F}_∇ defined by

$$\begin{aligned} J_k(F + \text{Null}) &:= \frac{1}{2\pi i} \int_\gamma^\# \mathbf{u}_\diamond^{[k]} j(\mathbf{u}_m) F d^\# \mathbf{u} + \text{Null} \\ \bar{J}_k(F + \text{Null}) &:= -\frac{1}{2\pi i} \oint_\gamma^\# \bar{\mathbf{u}}_\diamond^{[k]} \bar{j}(\mathbf{u}_m) F d^\# \bar{\mathbf{u}} + \text{Null}, \end{aligned} \quad (4.11)$$

where $j: \mathbb{Z}_m^2 \rightarrow \mathcal{P}_\nabla$ and $\bar{j}: \mathbb{Z}_m^2 \rightarrow \mathcal{P}_\nabla$ are the holomorphic and antiholomorphic currents from Example 4.1, and the discrete integration in the sense (3.17) is performed over any positively oriented corner contour γ that surrounds both the support of the field polynomial $F \in \mathcal{P}_\nabla$ and the set (3.22) of nonholomorphicity of the Laurent monomial $\mathbf{u} \mapsto \mathbf{u}^{[k]}$; i.e., γ is required to satisfy $S^{[k]} \cup \text{Supp } F \subset \text{int}^\# \gamma$. The setup is illustrated in Figure 4.1. To see that the operators $J_k, \bar{J}_k: \mathcal{F}_\nabla \rightarrow \mathcal{F}_\nabla$ are indeed well-defined by (4.11), one checks that the discrete integral on the right hand side is in the same correlation equivalence class for any representative F and for any allowed choice of γ , see [HKV22, Lemma 4.2].

Example 4.5. The **identity field** is defined as

$$I := 1 + \text{Null} \in \mathcal{F}_\nabla.$$

We note that

$$I \neq 0,$$

or equivalently $1 \notin \mathcal{N}$, because the random variable $\text{ev}_{\mathbf{z}}^{\Omega_\delta}(1) = 1$ (the evaluation of $1 \in \mathcal{P}_\nabla$) has a nonvanishing expected value. This may sound like an utterly trivial observation, but let us emphasize that 1 is the *only* field polynomial we need to explicitly verify not to be null; current modes allow us to generate new local fields starting from I , and algebraic considerations will imply that we get infinitely many linearly independent ones. Let us first consider the action of a single current mode on the identity field. For $k \in \mathbb{Z}$, using Stokes' formula (3.18) and the definition $j = i\partial_\#^\bullet X$, we find

$$\begin{aligned} J_k I &= \frac{1}{2\pi i} \oint_\gamma^\# \mathbf{u}_\diamond^{[k]} j(\mathbf{u}_m) d^\# \mathbf{u} + \text{Null} \\ &= \frac{i}{2\pi} \sum_{\mathbf{u}_m \in \text{int}_m^\# \gamma} \bar{\partial} \mathbf{u}_m^{[k]} \partial_\#^\bullet X(\mathbf{u}_m) + \frac{i}{2\pi} \sum_{\mathbf{u}_\diamond \in \text{int}_\diamond^\# \gamma} \mathbf{u}_\diamond^{[k]} \bar{\partial} \partial_\#^\bullet X(\mathbf{u}_\diamond) + \text{Null}. \end{aligned}$$

The second term is null for any $k \in \mathbb{Z}$, by the factorization of the Laplacian (3.6) and the null fields in Example 4.3. For $k \geq 0$, the first term is identically 0 by the discrete holomorphicity of the monomials — see Proposition 3.4. We conclude

$$J_k I = 0 \in \mathcal{F}_\nabla \quad \text{for all } k \in \mathbb{Z}_{\geq 0}. \quad (4.12)$$

A formula that is also valid for $k < 0$ is obtained by keeping the first term,

$$J_k I = J_k + \text{Null} \quad \text{where} \quad J_k := \frac{i}{2\pi} \sum_{\mathbf{u}_m \in S_m^{[k]}} \bar{\partial} \mathbf{u}_m^{[k]} \partial_{\sharp}^{\bullet} X(\mathbf{u}_m). \quad (4.13)$$

Note that the formula for the representative $J_k \in \mathcal{P}_{\nabla}$ here is manifestly independent of the choice of the contour γ . The support of the representative is also transparently related to the nonholomorphicity set (3.22) of the Laurent monomial. Specifically, using Remark 3.1 we obtain that

$$\text{Supp } J_{-k} \subset \mathbf{B}_{\bullet}^{\sharp} \left(\left\lfloor \frac{k}{2} \right\rfloor + 1 \right) \quad \text{and thus} \quad R_{\text{Supp}}(J_{-k} I) \leq \left\lfloor \frac{k}{2} \right\rfloor + 1, \quad \text{for } k \in \mathbb{Z}_{>0}.$$

Similarly, $\bar{J}_{-k} I = \bar{J}_{-k} + \text{Null}$ with $\bar{J}_k := \frac{-i}{2\pi} \sum_{\mathbf{u}_m \in S_m^{[k]}} \partial \bar{\mathbf{u}}_m^{[k]} \bar{\partial}_{\sharp}^{\bullet} X(\mathbf{u}_m)$ and the same arguments yield $R_{\text{Supp}}(\bar{J}_{-k} I) \leq \lfloor k/2 \rfloor + 1$. \diamond

The next proposition states that the space of correlation-equivalent local fields \mathcal{F}_{∇} can be equipped with a representation of two commuting copies of $\mathfrak{h}\mathfrak{c}\mathfrak{i}$; one for the holomorphic and one for the antiholomorphic chirality. The construction of the holomorphic representation was the key content in [HKV22], and the antiholomorphic one essentially repeats the same — but it is here that the small convention differences become important: with the exact conventions of [HKV22], the two chiralities would fail to commute! With our slightly modified Laurent monomials, the desired commutation property is recovered.

Proposition 4.2. *The current modes satisfy, for all $k, \ell \in \mathbb{Z}$,*

$$[J_k, J_{\ell}] = [\bar{J}_k, \bar{J}_{\ell}] = k \delta_{k+\ell} \text{id}_{\mathcal{F}_{\nabla}} \quad \text{and} \quad [J_k, \bar{J}_{\ell}] = 0.$$

The proof is presented after an auxiliary result regarding discrete integration. In that auxiliary result, we use the function

$$\epsilon: \mathbb{Z}_{\diamond}^2 \cup \mathbb{Z}_m^2 \rightarrow \{+1, -1\} \quad \text{given by} \quad \epsilon(\mathbf{u}) := (-1)^{2\text{Im}(\mathbf{u})},$$

i.e., the function which takes the value $+1$ on primal vertices and on horizontal edges and that takes the value -1 on dual vertices and vertical edges.

Lemma 4.3. *For any $\ell, k \in \mathbb{Z}$, we have*

$$\oint_{\gamma}^{\sharp} \bar{\mathbf{u}}_{\diamond}^{[\ell]} \epsilon(\mathbf{u}_m) \mathbf{u}_m^{[k]} d^{\sharp} \bar{\mathbf{u}} = 0,$$

where γ is any positively oriented corner contour surrounding the support of the discrete poles of both discrete monomials, i.e., $S^{[k]} \cup S^{[\ell]} \subset \text{int}^{\sharp} \gamma$.

Proof. Note, first of all, that, for any function f on $\mathbb{Z}_{\diamond}^2 \cup \mathbb{Z}_m^2$, we have

$$\partial(\epsilon \cdot f) = \epsilon \cdot \bar{\partial} f \quad \text{and} \quad \bar{\partial}(\epsilon \cdot f) = \epsilon \cdot \partial f$$

where the dot \cdot stands for pointwise multiplication of functions. By Stokes formula (3.18) combined with this observation, the integral can be written as

$$\begin{aligned} \oint_{\gamma}^{\#} \bar{\mathbf{u}}_{\diamond}^{[\ell]} \epsilon(\mathbf{u}_m) \mathbf{u}_m^{[k]} d^{\#} \bar{\mathbf{u}} &= -i \sum_{\mathbf{v}_{\diamond} \in \text{int}_{\diamond}^{\#} \gamma} \bar{\mathbf{v}}_{\diamond}^{[\ell]} \epsilon(\mathbf{v}_{\diamond}) (\bar{\partial}_{\#} \mathbf{v}_{\diamond}^{[k]}) - i \sum_{\mathbf{v}_m \in \text{int}_m^{\#} \gamma} (\partial_{\#} \bar{\mathbf{v}}_m^{[\ell]}) \epsilon(\mathbf{v}_m) \mathbf{v}_m^{[k]} \\ &= -i \sum_{\mathbf{v}_{\diamond} \in S_{\diamond}^{[k]}} \bar{\mathbf{v}}_{\diamond}^{[\ell]} \epsilon(\mathbf{v}_{\diamond}) (\bar{\partial}_{\#} \mathbf{v}_{\diamond}^{[k]}) - i \sum_{\mathbf{v}_m \in S_m^{[\ell]}} \overline{(\bar{\partial}_{\#} \mathbf{v}_m^{[\ell]})} \epsilon(\mathbf{v}_m) \mathbf{v}_m^{[k]}. \end{aligned}$$

If $k, \ell \geq 0$ then there are no discrete poles that contribute, $S^{[\ell]} = S^{[k]} = \emptyset$, so the integral vanishes as asserted.

If $k + \ell \leq -2$, then using the fact that the integral does not depend on γ , we can argue as follows. Taking γ to be a symmetric square path at distance r from the origin, the integrand is $\mathcal{O}(r^{k+\ell})$ and the length of the integration contour is $\mathcal{O}(r)$, so the integral is $\mathcal{O}(r^{1+k+\ell})$, and taking $r \rightarrow \infty$ shows that it must vanish.

It remains to consider the case $k < 0$ and $\ell \geq -1 - k \geq 0$, and the case $\ell < 0$ and $k \geq -1 - \ell \geq 0$. By repeated integration by parts, these can be reduced to cases when the negative exponent is -1 .

For example if $\ell < 0$ then with (3.20) we can rewrite the integral as

$$\begin{aligned} \oint_{\gamma}^{\#} \bar{\mathbf{u}}_{\diamond}^{[\ell]} \epsilon(\mathbf{u}_m) \mathbf{u}_m^{[k]} d^{\#} \bar{\mathbf{u}} &= \frac{1}{\ell + 1} \oint_{\gamma}^{\#} (\bar{\partial}_{\#} \bar{\mathbf{u}}_{\diamond}^{[\ell+1]}) \epsilon(\mathbf{u}_m) \mathbf{u}_m^{[k]} d^{\#} \bar{\mathbf{u}} \\ &= -\frac{1}{\ell + 1} \oint_{\gamma}^{\#} \bar{\mathbf{u}}_m^{[\ell+1]} \epsilon(\mathbf{u}_{\diamond}) (\partial_{\#} \mathbf{u}_{\diamond}^{[k]}) d^{\#} \bar{\mathbf{u}} \\ &= -\frac{k}{\ell + 1} \oint_{\gamma}^{\#} \bar{\mathbf{u}}_m^{[\ell+1]} \epsilon(\mathbf{u}_{\diamond}) \mathbf{u}_{\diamond}^{[k-1]} d^{\#} \bar{\mathbf{u}}. \end{aligned}$$

Applying this recursively, the integral is seen to be proportional to either

$$\oint_{\gamma}^{\#} \bar{\mathbf{u}}_m^{[-1]} \epsilon(\mathbf{u}_{\diamond}) \mathbf{u}_{\diamond}^{[k+\ell+1]} d^{\#} \bar{\mathbf{u}} \quad \text{or} \quad \oint_{\gamma}^{\#} \bar{\mathbf{u}}_{\diamond}^{[-1]} \epsilon(\mathbf{u}_m) \mathbf{u}_m^{[k+\ell+1]} d^{\#} \bar{\mathbf{u}},$$

depending on the parity of $\ell < 0$. These integrals are then evaluated by Stokes' formula (3.18), as above. Note that we have $\bar{\partial}_{\#} \mathbf{v}^{[k+\ell+1]} \equiv 0$ since $k + \ell + 1 \geq 0$, so one of the terms in Stokes' formula does not contribute. The integrals above thus become, up to multiplicative constants,

$$\sum_{|\mathbf{v}_{\diamond}| \leq \frac{1}{\sqrt{2}}} \overline{(\bar{\partial}_{\#} \mathbf{v}_{\diamond}^{[-1]})} \epsilon(\mathbf{v}_{\diamond}) \mathbf{v}_{\diamond}^{[k+\ell+1]} \quad \text{or} \quad \sum_{|\mathbf{v}_m| = \frac{1}{2}} \overline{(\bar{\partial}_{\#} \mathbf{v}_m^{[-1]})} \epsilon(\mathbf{v}_m) \mathbf{v}_m^{[k+\ell+1]}.$$

For $k + \ell + 1 \geq 3$, the monomial $\mathbf{v}^{[k+\ell+1]}$ vanishes on the support of $\bar{\partial}_{\#} \mathbf{v}^{[-1]}$, so all terms in these sums are zero, and the original integral vanishes again as asserted. Only in the cases

$k + \ell + 1 \in \{0, 1, 2\}$ the sums above have nonzero terms. In the case $k + \ell + 1 = 1$ the summands are odd, and they therefore cancel. In the case $k + \ell + 1 = 0$ there are equal contributions with both signs of ϵ , and they therefore cancel. In the case $k + \ell + 1 = 2$ on the medial lattice the relevant values of the monomial $\mathbf{v}_m^{[2]}$ are again zero, and on the diamond lattice the value at the origin is zero and the other four values change sign under 90° -rotations, by virtue of symmetries of the monomial $\mathbf{v}_\diamond^{[2]}$, leading to cancellations again.

Similarly if $k < 0$, with repeated integrations by parts one reduces to two cases with a first order pole, both of which are evaluated by Stokes' formula, and both of which can be explicitly seen to vanish — case by case according to the value of $k + \ell + 1 \geq 0$. \square




Figure 4.2: An example of the sets and contours involved in the computation of $[J_k, \bar{J}_\ell]$. In red the set $S^{[k]} \cup S^{[\ell]}$ of nonholomorphicity of the relevant monomials, in green the support of a representative F , in purple a choice of corner contours γ_- (inner) and γ_+ (outer), and in light blue a choice of corner contour γ ; all of them laid on the infinite square grid \mathbb{Z}^2 and its sublattices \mathbb{Z}_\diamond^2 and \mathbb{Z}_m^2 .

Proof of Proposition 4.2. For the proof of $[J_k, J_\ell] = [\bar{J}_k, \bar{J}_\ell] = k \delta_{k+\ell} \text{id}_{\mathcal{F}_\nabla}$ we refer the reader to [HKV22, Proposition 4.5]; the minor differences in our conventions do not affect the essence of this part of the proof. It remains to prove that $[J_k, \bar{J}_\ell] = 0$. Let $F \in \mathcal{P}_\nabla$ be a local field. Take three disjoint corner contours γ_- , γ , and γ_+ satisfying

$$\text{Supp } F \cup S^{[k]} \cup S^{[\ell]} \subset \text{int}^\# \gamma_- \subset \text{int}^\# \gamma \subset \text{int}^\# \gamma_+,$$

i.e., the contours must be nested and all surround the support of F as well as the sets (3.22) of failure of the discrete (anti)holomorphicity of the monomials of order k and ℓ ; see Figure 4.2. Then, by the discrete Stokes' formula (3.18) and the factorization (3.6) of the

discrete Laplacian we calculate

$$\begin{aligned}
[J_k, \bar{J}_\ell](F + \text{Null}) &= \left(\frac{1}{2\pi i} \oint_{\gamma_+}^\# \mathbf{u}_\diamond^{[k]} j(\mathbf{u}_m) d^\# \mathbf{u} \right) \left(\frac{-1}{2\pi i} \oint_\gamma^\# \bar{\mathbf{v}}_\diamond^{[\ell]} \bar{j}(\mathbf{v}_m) d^\# \bar{\mathbf{v}} \right) F \\
&\quad - \left(\frac{-1}{2\pi i} \oint_\gamma^\# \bar{\mathbf{v}}_\diamond^{[\ell]} \bar{j}(\mathbf{v}_m) d^\# \bar{\mathbf{v}} \right) \left(\frac{1}{2\pi i} \oint_{\gamma_-}^\# \mathbf{u}_\diamond^{[k]} j(\mathbf{u}_m) d^\# \mathbf{u} \right) F + \text{Null} \\
&= \frac{1}{4\pi^2} \oint_\gamma^\# d^\# \bar{\mathbf{v}} \bar{\mathbf{v}}_\diamond^{[\ell]} \left[\oint_{\gamma_+}^\# - \oint_{\gamma_-}^\# \right] d^\# \mathbf{u} \mathbf{u}_\diamond^{[k]} \bar{\partial}_\#^\bullet X(\mathbf{v}_m) \partial_\#^\bullet X(\mathbf{u}_m) F + \text{Null} \\
&= \frac{i}{4\pi^2} \oint_\gamma^\# d^\# \bar{\mathbf{v}} \bar{\mathbf{v}}_\diamond^{[\ell]} \sum_{\mathbf{u}_\diamond \in \text{int}_\diamond^\# \gamma_+ \setminus \text{int}_\diamond^\# \gamma_-} \mathbf{u}_\diamond^{[k]} \bar{\partial}_\#^\bullet X(\mathbf{v}_m) \bar{\partial} \partial_\#^\bullet X(\mathbf{u}_\diamond) F + \text{Null} \\
&= \frac{i}{8\pi^2} \oint_\gamma^\# d^\# \bar{\mathbf{v}} \bar{\mathbf{v}}_\diamond^{[\ell]} \sum_{\mathbf{u}_\bullet \in \text{int}_\bullet^\# \gamma_+ \setminus \text{int}_\bullet^\# \gamma_-} \mathbf{u}_\bullet^{[k]} \bar{\partial}_\#^\bullet X(\mathbf{v}_m) \Delta_\# X(\mathbf{u}_\bullet) F + \text{Null}.
\end{aligned}$$

Now recall the quadratic null fields of Example 4.4, which yield in particular that

$$\bar{\partial}_\#^\bullet X(\mathbf{v}_m) \Delta_\# X(\mathbf{u}_\bullet) = 4\pi (\bar{\partial}_\#^\bullet)_{\mathbf{v}_m} \delta_{\mathbf{u}_\bullet, \mathbf{v}_m} + \text{Null}.$$

We may therefore simplify the earlier calculation to

$$\begin{aligned}
[J_k, \bar{J}_\ell](F + \text{Null}) &= \frac{i}{2\pi} \oint_\gamma^\# d^\# \bar{\mathbf{v}} \bar{\mathbf{v}}_\diamond^{[\ell]} \sum_{\substack{\mathbf{u}_\bullet \in \text{int}_\bullet^\# \gamma_+ \\ \mathbf{u}_\bullet \notin \text{int}_\bullet^\# \gamma_-}} \mathbf{u}_\bullet^{[k]} (\bar{\partial}_\#^\bullet)_{\mathbf{v}_m} (\delta_{\mathbf{u}_\bullet, \mathbf{v}_m}) F + \text{Null} \\
&= \frac{i}{2\pi} \oint_\gamma^\# d^\# \bar{\mathbf{v}} \bar{\mathbf{v}}_\diamond^{[\ell]} \bar{\partial}_\#^\bullet \mathbf{v}_m^{[k]} F + \text{Null}.
\end{aligned}$$

Now note that if f is a function on $\mathbb{Z}_\diamond^2 \cup \mathbb{Z}_m^2$, then at any edge midpoint $\mathbf{v}_m \in \mathbb{Z}_m^2$ we have

$$\bar{\partial}_\#^\bullet f(\mathbf{v}_m) = \bar{\partial}_\# f(\mathbf{v}_m) + \epsilon(\mathbf{v}_m) \partial_\# f(\mathbf{v}_m).$$

In particular if f is discrete holomorphic we have $\bar{\partial}_\#^\bullet f(\mathbf{v}_m) = \epsilon(\mathbf{v}_m) \partial_\# f(\mathbf{v}_m)$. The function $\mathbf{v}_m \mapsto \mathbf{v}_m^{[k]}$ is discrete holomorphic on γ , because the contour γ surrounds the poles of the monomials by assumption, so on γ we get $\bar{\partial}_\#^\bullet \mathbf{v}_m^{[k]} = k \epsilon(\mathbf{v}_m) \mathbf{v}_m^{[k-1]}$, using also the derivative property of the discrete monomials from Proposition 3.4. The earlier calculation thus simplifies to

$$[J_k, \bar{J}_\ell](F + \text{Null}) = \frac{i k}{2\pi} \oint_\gamma^\# d^\# \bar{\mathbf{v}} \bar{\mathbf{v}}_\diamond^{[\ell]} \epsilon(\mathbf{v}_m) \mathbf{v}_m^{[k-1]} F + \text{Null},$$

which vanishes by Lemma 4.3. □

By Proposition 4.2, the space \mathcal{F}_∇ of local fields carries representations of two commuting Heisenberg algebras. Let us again denote correspondingly by Hei and $\overline{\text{Hei}}$ the two

commuting associative algebras (2.3), whose representations on \mathcal{F}_∇ are determined by formulas (4.11) for their generators, i.e., $j_k \mapsto J_k$ and $\bar{j}_k \mapsto \bar{J}_k$, respectively.

Recall that our first main goal is to put the space \mathcal{F}_∇ of local fields of the (gradient of the) discrete Gaussian Free Field into a one-to-one correspondence with the two-chiral Fock space $\mathcal{F} \otimes \bar{\mathcal{F}}$, which serves as the space of local fields for the bosonic CFT of the (gradient of the) continuum Gaussian Free Field. The Fock space $\mathcal{F} \otimes \bar{\mathcal{F}}$ carries representations of two commuting copies of the Heisenberg algebra by construction, and we want our correspondence to respect this structure, i.e., to be a map of representations of $\text{Hei} \otimes \bar{\text{Hei}}$. By mostly abstract nonsense, one direction of the desired one-to-one correspondence now becomes very easy: an isomorphic copy of the Fock space is found inside the space \mathcal{F}_∇ of local fields of the DGFF as follows:

Corollary 4.4. *The $\text{Hei} \otimes \bar{\text{Hei}}$ subrepresentation in \mathcal{F}_∇ generated by the identity field I (Example 4.5) is isomorphic to the full Fock space,*

$$\mathcal{F} \otimes \bar{\mathcal{F}} \cong (\text{Hei} \otimes \bar{\text{Hei}})I \subset \mathcal{F}_\nabla.$$

Proof. In (4.12) we saw that $J_k I = 0$ and $\bar{J}_k I = 0$ for all $k \geq 0$, and we also noted that $I \neq 0 \in \mathcal{F}_\nabla$. The asserted isomorphism therefore follows immediately from Lemma 2.1. \square

4.6 Sugawara construction and homogeneous local fields. In Section 2.2, we recalled how the Fock space representation of the Heisenberg algebra can be rendered a Virasoro representation via the Sugawara construction. A key observation in [HKV22] was that the same construction can be applied in the space \mathcal{F}_∇ of correlation equivalence classes of local fields of the discrete GFF, thanks to the following truncation property.

Lemma 4.5. *For any $F \in \mathcal{P}_\nabla$, there exists $K \in \mathbb{Z}_{>0}$ such that*

$$J_k(F + \text{Null}) = \bar{J}_k(F + \text{Null}) = 0 + \text{Null}$$

for all $k \geq K$.

Proof. See [HKV22, Lemma 4.4] for the holomorphic sector. The argument is identical for the antiholomorphic sector. \square

The important conclusion about the space \mathcal{F}_∇ of correlation equivalence classes of local fields is the following.

Corollary 4.6. *The formulas*

$$\mathsf{L}_n = \frac{1}{2} \left(\sum_{k \geq 0} J_{n-k} J_k + \sum_{k < 0} J_k J_{n-k} \right) \quad \text{and} \quad \bar{\mathsf{L}}_n = \frac{1}{2} \left(\sum_{k \geq 0} \bar{J}_{n-k} \bar{J}_k + \sum_{k < 0} \bar{J}_k \bar{J}_{n-k} \right)$$

equip the space \mathcal{F}_∇ with two commuting Virasoro representations with central charge $c = 1$.

Proof. The Virasoro commutation relations with $c = 1$ for both \mathbf{L}_n and $\bar{\mathbf{L}}_n$ are shown as in [HKV22, Theorem 4.10], using the truncation property of Lemma 4.5. The mutual commutation, $[\mathbf{L}_n, \bar{\mathbf{L}}_m] = 0$, follows from the mutual commutation of the corresponding Heisenberg modes, $[\mathbf{J}_k, \bar{\mathbf{J}}_\ell] = 0$, proven in Proposition 4.2. \square

A particular role is played by the holomorphic and antiholomorphic Virasoro generators with index $n = 0$,

$$\mathbf{L}_0 = \frac{1}{2} \mathbf{J}_0 \mathbf{J}_0 + \sum_{k=1}^{\infty} \mathbf{J}_{-k} \mathbf{J}_k \quad \text{and} \quad \bar{\mathbf{L}}_0 = \frac{1}{2} \bar{\mathbf{J}}_0 \bar{\mathbf{J}}_0 + \sum_{k=1}^{\infty} \bar{\mathbf{J}}_{-k} \bar{\mathbf{J}}_k.$$

In CFT the sum $\mathbf{L}_0 + \bar{\mathbf{L}}_0$ is the Hamiltonian (energy) operator, which in radial quantization serves as the infinitesimal generator of scalings. The difference $\mathbf{L}_0 - \bar{\mathbf{L}}_0$ is the spin operator, which serves as the infinitesimal generator of rotations. For determining the needed renormalization of fields in the scaling limit, the eigenvalues of $\mathbf{L}_0 + \bar{\mathbf{L}}_0$ will be crucial — these are called the scaling dimensions of the fields. The pair of eigenvalues for both \mathbf{L}_0 and $\bar{\mathbf{L}}_0$ carries the information on both scaling dimension and spin; we define the space of **homogeneous** local fields of **conformal dimensions** $\Delta, \bar{\Delta} \in \mathbb{C}$ as the joint eigenspace

$$(\mathcal{F}_{\nabla})_{\Delta, \bar{\Delta}} := \ker(\mathbf{L}_0 - \Delta) \cap \ker(\bar{\mathbf{L}}_0 - \bar{\Delta}) \subset \mathcal{F}_{\nabla}. \quad (4.14)$$

While it was very easy to see that the Fock space $\mathcal{F} \otimes \bar{\mathcal{F}}$ has a grading (2.11) by conformal dimensions, at this stage we have not yet established the same conclusion about the space \mathcal{F}_{∇} : diagonalizability of \mathbf{L}_0 and $\bar{\mathbf{L}}_0$ and finite-dimensionality of the joint eigenspaces still need to be proven in order for the homogeneous components (4.14) to be complete and usable decomposition of correlation equivalence classes of local fields of the discrete GFF. We can, however, already give some examples of homogeneous fields, because by Corollary 4.4, the space \mathcal{F}_{∇} contains a subspace $(\text{Hei} \otimes \overline{\text{Hei}})I$ isomorphic to the Fock space.

Example 4.6. The basis vectors (2.6) of the Fock space are eigenvectors, and correspondingly we have fields

$$\mathbf{J}_{-k_m} \cdots \mathbf{J}_{-k_2} \mathbf{J}_{-k_1} \bar{\mathbf{J}}_{-k'_{m'}} \cdots \bar{\mathbf{J}}_{-k'_2} \bar{\mathbf{J}}_{-k'_1} I \in (\mathcal{F}_{\nabla})_{\Delta, \bar{\Delta}} \quad (4.15)$$

with conformal dimensions $\Delta = \sum_{i=1}^m k_i$ and $\bar{\Delta} = \sum_{j=1}^{m'} k'_j$.

The most obvious special case is the identity field I : the eigenvalue properties $\mathbf{L}_0 I = 0$ and $\bar{\mathbf{L}}_0 I = 0$ in fact also follow easily from (4.12) and we indeed have $I \in (\mathcal{F}_{\nabla})_{0,0}$. \diamond

5 Linear local fields of the DGFF

Our first main goal is to fully work out the structure of the space $\mathcal{F}_{\nabla} := \mathcal{P}_{\nabla}/\mathcal{N}$ of local fields of the DGFF. We seek to show that it has the same structure as the space of local

fields of a CFT, i.e., that it is isomorphic to the Fock space $\mathcal{F} \otimes \overline{\mathcal{F}}$. A priori, the difficulty stems from the fact that $\mathcal{F}_\nabla = \mathcal{P}_\nabla / \mathcal{N}$ involves a quotient by null fields, which are defined by an implicit condition that cannot be decided by a straightforward method.

Recall, however, that one inclusion, $\mathcal{F} \otimes \overline{\mathcal{F}} \cong (\text{Hei} \otimes \overline{\text{Hei}})I \subset \mathcal{F}_\nabla$, was already obtained in Corollary 4.4 by virtue of the Heisenberg algebra actions of Proposition 4.2. Establishing the remaining opposite inclusion $\mathcal{F}_\nabla \subset (\text{Hei} \otimes \overline{\text{Hei}})I$ now amounts to showing that the whole space \mathcal{F}_∇ is exhausted by linear combinations of those fields that can be obtained from the identity field I by repeated actions of the Heisenberg generators (4.11). A natural strategy for doing that is to exhibit concrete upper bounds for the dimensions of some suitably chosen subspaces of \mathcal{F}_∇ , and showing that the upper bounds are already saturated within the subspace $(\text{Hei} \otimes \overline{\text{Hei}})I \subset \mathcal{F}_\nabla$. We will carry out such a strategy in two steps in this section and the next. The present section achieves dimension upper bounds for *linear* local fields, i.e., those corresponding to homogeneous field polynomials of degree one. The task in Section 6 will then be to reduce the case of higher degree fields to such linear factors.

To achieve useful dimension bounds for linear local fields, we must still refine to further subspaces: the space of all linear local fields (even modulo null fields) remains infinite dimensional, so counting arguments without refinement would be doomed. What turns out to work is to construct a filtration of linear local fields by the finite-dimensional subspaces with at most a given radius of support.

We start in Section 5.1 by defining linear local fields and stating the result (Theorem 5.1) which gives an explicit basis for them. Then, in Section 5.2, we introduce the filtration with finite-dimensional subspaces in which dimension counting is to be performed. Here we also already record the dimension lower bounds, which follow from the earlier observation $(\text{Hei} \otimes \overline{\text{Hei}})I \subset \mathcal{F}_\nabla$ and some observations about the radii of supports of some explicit linear field polynomials. The main task of proving the matching upper bounds for the dimensions is done in Section 5.3, and once this is done, we give the proof of the basis theorem for linear local fields (Theorem 5.1). We conclude in Section 5.4 with simple remarks on what the basis theorem says about homogeneous linear local fields.

5.1 Linear local fields. By definition (4.5), the space of field polynomials

$$\mathcal{P}_\nabla = \mathbb{C}[X(\mathbf{u}) : \mathbf{u} \in \mathbb{Z}^2]$$

is the free commutative (polynomial) algebra generated by the symbols $X(\mathbf{u})$. The space of **linear field polynomials** is now defined to be the subspace *spanned* by these symbols,

$$\mathcal{P}_\nabla^{\text{lin}} := \text{span}_{\mathbb{C}} \{ X(\mathbf{u}) : \mathbf{u} \in \mathbb{Z}^2 \} \subset \mathcal{P}_\nabla. \quad (5.1)$$

The space of **linear local fields** is then again defined by identifying field polynomials which differ by a null field,

$$\mathcal{F}_\nabla^{\text{lin}} := \mathcal{P}_\nabla^{\text{lin}} / \mathcal{N} \subset \mathcal{F}_\nabla. \quad (5.2)$$

Example 5.1. Recall from Example 4.5 that, for $k \in \mathbb{Z}_{>0}$, we can write $J_{-k}I = J_{-k} + \text{Null}$ and $\bar{J}_{-k}I = \bar{J}_{-k} + \text{Null}$ with linear field polynomials $J_{-k}, \bar{J}_{-k} \in \mathcal{P}_{\nabla}^{\text{lin}}$ given explicitly as in (4.13). Therefore,

$$J_{-k}I, \bar{J}_{-k}I \in \mathcal{F}_{\nabla}^{\text{lin}}$$

are linear local fields. \diamond

Corollary 4.4, together with the fact that (2.6) is a basis of the Fock space, implies that the linear local fields given in Example 5.1 are linearly independent. The main goal of this section is to prove that they in fact form a basis.

Theorem 5.1. *The set*

$$\{J_{-k}I \mid k \in \mathbb{Z}_{>0}\} \cup \{\bar{J}_{-k}I \mid k \in \mathbb{Z}_{>0}\} \quad (5.3)$$

is a basis of the space $\mathcal{F}_{\nabla}^{\text{lin}}$ of linear local fields.

We moreover obtain a complete characterization of linear null fields, which will also be used in Section 6. Among linear null fields, the explicit ones given in Examples 4.3 and 4.2 are all there is.

Corollary 5.2. *The set*

$$\{\Delta_{\sharp}X(\mathbf{u}) \mid \mathbf{u} \in \mathbb{Z}^2\} \cup \{X(0)\}$$

spans the subspace $\mathcal{P}_{\nabla}^{\text{lin}} \cap \mathcal{N}$ of null linear local fields.

5.2 Filtration by radius. We now present the filtration which enables a dimension counting argument that is the key to proving Theorem 5.1.

Consider the **discrete balls** of radii $r \in \mathbb{Z}_{>0}$

$$\mathbf{B}_{\bullet}^{\sharp}(r) := \left\{ \mathbf{u} \in \mathbb{Z}^2 \mid \|\mathbf{u}\|_1 \leq r \right\}, \quad (5.4)$$

with respect to the Manhattan norm $\|z\|_1 := \text{Re}(z) + \text{Im}(z)$. In the space (5.1) of linear field polynomials, the subspace of those fields whose support is in $\mathbf{B}_{\bullet}^{\sharp}(r)$ is

$$\mathcal{P}_{\nabla}^{\text{lin}}(r) := \text{span}_{\mathbb{C}} \left\{ X(\mathbf{u}) : \mathbf{u} \in \mathbf{B}_{\bullet}^{\sharp}(r) \right\} \subset \mathcal{P}_{\nabla}^{\text{lin}}. \quad (5.5)$$

These subspaces are evidently finite-dimensional since the subset $\mathbf{B}_{\bullet}^{\sharp}(r) \subset \mathbb{Z}$ is finite,

$$\dim(\mathcal{P}_{\nabla}^{\text{lin}}(r)) \leq |\mathbf{B}_{\bullet}^{\sharp}(r)| = 2r^2 + 2r + 1 \quad \text{for } r \in \mathbb{Z}_{>0}.$$

Denote the corresponding subspace of the quotient (5.2) by

$$\mathcal{F}_{\nabla}^{\text{lin}}(r) := \mathcal{P}_{\nabla}^{\text{lin}}(r)/\mathcal{N} \subset \mathcal{F}_{\nabla}^{\text{lin}}. \quad (5.6)$$

These subspaces form a filtration of $\mathcal{F}_{\nabla}^{\text{lin}}$,

$$\mathcal{F}_{\nabla}^{\text{lin}}(1) \subset \mathcal{F}_{\nabla}^{\text{lin}}(2) \subset \mathcal{F}_{\nabla}^{\text{lin}}(3) \subset \cdots \subset \mathcal{F}_{\nabla}^{\text{lin}} \quad \text{and} \quad \sum_{r \in \mathbb{Z}_{>0}} \mathcal{F}_{\nabla}^{\text{lin}}(r) = \mathcal{F}_{\nabla}^{\text{lin}},$$

because the discrete balls form an increasing sequence of (finite) subsets that exhaust the square grid \mathbb{Z}^2 ,

$$\mathbf{B}_{\bullet}^{\sharp}(1) \subset \mathbf{B}_{\bullet}^{\sharp}(2) \subset \mathbf{B}_{\bullet}^{\sharp}(3) \subset \cdots \subset \mathbb{Z}^2 \quad \text{and} \quad \bigcup_{r \in \mathbb{Z}_{>0}} \mathbf{B}_{\bullet}^{\sharp}(r) = \mathbb{Z}^2.$$

Example 5.2. Recall from Example 4.5 that, for $k \in \mathbb{Z}_{>0}$, the linear field polynomials J_{-k} and \bar{J}_{-k} are supported in $\mathbf{B}_{\bullet}^{\sharp}(r_k)$ with $r_k := \lfloor k/2 \rfloor + 1$. By Example 5.1, we therefore see that $J_{-k}I, \bar{J}_{-k}I \in \mathcal{F}_{\nabla}^{\text{lin}}(r_k)$. \diamond

Let us give one slightly more subtle example in the form of a lemma. In this example the precise form (slightly different from [HKV22]) of our monomials defined in Section 3.4 again becomes important.

Lemma 5.3. *For $r \in \mathbb{Z}_{>0}$, we have*

$$J_{-2r}I - \bar{J}_{-2r}I \in \mathcal{F}_{\nabla}^{\text{lin}}(r).$$

Proof. Take a corner contour γ sufficiently large for the action of J_{-2r} . Using integration by parts (3.20) $2r - 1$ times, the properties of the discrete monomials from Proposition 3.4, Stokes' formula (3.18), and the null fields of Example 4.3, we calculate

$$\begin{aligned} J_{-2r}I &= \frac{1}{2\pi} \oint_{\gamma}^{\sharp} \mathbf{u}_{\diamond}^{[-2r]} \partial_{\sharp}^{\bullet} X(\mathbf{u}_{\text{m}}) d^{\sharp} \mathbf{u} + \text{Null} \\ &= \frac{1}{2\pi(2r-1)!} \oint_{\gamma}^{\sharp} \mathbf{u}_{\text{m}}^{[-1]} \partial_{\sharp}^{2r-1} \partial_{\sharp}^{\bullet} X(\mathbf{u}_{\diamond}) d^{\sharp} \mathbf{u} + \text{Null} \\ &= \frac{\mathfrak{i}}{2\pi(2r-1)!} \sum_{\mathbf{u}_{\diamond} \in \text{int}_{\diamond}^{\sharp} \gamma} \bar{\partial}_{\sharp} \mathbf{u}_{\diamond}^{[-1]} \partial_{\sharp}^{2r-1} \partial_{\sharp}^{\bullet} X(\mathbf{u}_{\diamond}) + \text{Null} \\ &= \frac{\mathfrak{i}}{(2r-1)!} \left[\frac{1}{2} \partial_{\sharp}^{2r-1} \partial_{\sharp}^{\bullet} X(0) + \frac{1}{8} \sum_{\mathbf{v}=\frac{\pm 1 \pm \mathfrak{i}}{2}} \partial_{\sharp}^{2r-1} \partial_{\sharp}^{\bullet} X(\mathbf{v}) \right] + \text{Null}. \end{aligned}$$

Similarly we get

$$\bar{J}_{-2r}I = \frac{-\mathfrak{i}}{(2r-1)!} \left[\frac{1}{2} \bar{\partial}_{\sharp}^{2r-1} \bar{\partial}_{\sharp}^{\bullet} X(0) + \frac{1}{8} \sum_{\mathbf{v}=\frac{\pm 1 \pm \mathfrak{i}}{2}} \bar{\partial}_{\sharp}^{2r-1} \bar{\partial}_{\sharp}^{\bullet} X(\mathbf{v}) \right] + \text{Null}.$$

We now make some observations about the compositions $\partial_{\sharp}^{2r-1} \partial_{\sharp}^{\bullet}$ and $\bar{\partial}_{\sharp}^{2r-1} \bar{\partial}_{\sharp}^{\bullet}$ of the finite difference operators appearing in the above formulas. The expression $\partial_{\sharp}^{2r-1} \partial_{\sharp}^{\bullet} X(\mathbf{v})$ is a

linear combination of $X(\mathbf{u})$, with contributions from paths from $\mathbf{v} \in \mathbb{Z}_\diamond^2$ to $\mathbf{u} \in \mathbb{Z}^2$ which use $2r - 1$ arbitrary half lattice unit steps and then one half lattice unit step along a half-edge leading to \mathbf{u} , with coefficients that are products of the weights in the finite difference operators (3.2), (3.4) and (3.5) corresponding to the steps. If $\mathbf{v} \in \mathbb{Z}_\bullet^2$ is a primal vertex, then necessarily an even number of vertical half-steps is used by the path, and an even number of weight factors are imaginary, so the coefficient is real. By contrast, if $\mathbf{v} \in \mathbb{Z}_*^2$ is a dual vertex, then an odd number of vertical half-steps is used, and the coefficient is imaginary. Moreover, the only difference in the weights between $\partial_\#^{2r-1} \partial_\#^\bullet$ and $\bar{\partial}_\#^{2r-1} \bar{\partial}_\#^\bullet$ is a different sign for the vertical steps, so for $\mathbf{v} \in \mathbb{Z}_\bullet^2$ we have $\bar{\partial}_\#^{2r-1} \bar{\partial}_\#^\bullet X(\mathbf{v}) = \partial_\#^{2r-1} \partial_\#^\bullet X(\mathbf{v})$, and for $\mathbf{v} \in \mathbb{Z}_*^2$ we have $\bar{\partial}_\#^{2r-1} \bar{\partial}_\#^\bullet X(\mathbf{v}) = -\partial_\#^{2r-1} \partial_\#^\bullet X(\mathbf{v})$.

The above considerations of coefficients in compositions of finite difference operators show that in $J_{-2r}I - \bar{J}_{-2r}I$ there is a cancellation of the terms corresponding to $\mathbf{v} = \frac{\pm 1 \pm i}{2} \in \mathbb{Z}_*^2$. More precisely, $J_{-2r}I - \bar{J}_{-2r}I$ has a representative

$$\frac{i}{(2r-1)!} \partial_\#^{2r-1} \partial_\#^\bullet X(0) \in \mathcal{P}_\nabla^{\text{lin}}(r),$$

where the sufficiency of the radius of support r is a consequence of using exactly $2r$ half-steps starting from the origin $0 \in \mathbb{Z}_\bullet^2$. This proves $J_{-2r}I - \bar{J}_{-2r}I \in \mathcal{F}_\nabla^{\text{lin}}(r)$. \square

From the explicit examples above, we get lower bounds for the dimension of the subspaces in the filtration.

Lemma 5.4. *For every $r \in \mathbb{Z}_{>0}$, we have*


$$\dim(\mathcal{F}_\nabla^{\text{lin}}(r)) \geq 4r - 1.$$

Proof. Fix $r \in \mathbb{Z}_{>0}$. For $1 \leq k \leq 2r - 1$ we have $J_{-k}I, \bar{J}_{-k}I \in \mathcal{F}_\nabla^{\text{lin}}(r)$ by Example 5.2. This gives $2(2r - 1) = 4r - 2$ fields in $\mathcal{F}_\nabla^{\text{lin}}(r)$. Lemma 5.3 gives one more, $J_{-2r}I - \bar{J}_{-2r}I \in \mathcal{F}_\nabla^{\text{lin}}(r)$. Corollary 4.4 implies that these $4r - 1$ fields are linearly independent. \square


5.3 Dimension upper bounds. We now need dimension upper bounds for the subspaces in the filtration, which match the lower bounds of Lemma 5.4. Starting from the obvious spanning set $\{X(\mathbf{u}) + \text{Null} \mid \mathbf{u} \in \mathbf{B}_\bullet^\sharp(r)\}$ for $\mathcal{F}_\nabla^{\text{lin}}(r)$, which has $|\mathbf{B}_\bullet^\sharp(r)| = 2r^2 + 2r + 1 \gg 4r - 1$ elements, we use a method based on discrete harmonic measures and divergence theorem, which yields a smaller set of canonical representatives.

In order to describe the harmonic measures literally in terms of the discrete Laplacian operators and Green's functions introduced in Section 3.1, we start with choosing an appropriate discrete domain, which is just a slight modification of the discrete ball $\mathbf{B}_\bullet^\sharp(r)$. For $r \in \mathbb{Z}_{>0}$, consider the discrete domain $\mathbf{D}(r) \subset \mathbb{Z}^2$ as illustrated in Figure 5.1(a), with interior

$$\mathbf{D}^\circ(r) = \left\{ \mathbf{u} \in \mathbb{Z}^2 \mid \|\mathbf{u}\|_1 < r \right\}. \quad (5.7)$$



(a) The discrete domain $\mathbf{D}(r) \subset \mathbb{Z}^2$ with $r = 4$. The interior vertices $\mathbf{D}^\circ(r) \subset \mathbf{D}(r)$ are colored gray. The subset $\mathbf{C}(r) \subset \partial\mathbf{D}(r)$ of boundary vertices where harmonic measures are supported are colored white: the other boundary vertices are black.



(b) The discrete harmonic measure $\mathbf{v} \mapsto H_{\mathbf{b}}(\mathbf{v})$ at a given boundary point $\mathbf{b} \in \mathbf{C}(r)$ represents the weight with which the field $X(\mathbf{v})$ at $\mathbf{v} \in \mathbf{D}^\circ(r)$ contributes to the coefficient of $X(\mathbf{b})$ in the linear local field representative which has its support on $\mathbf{C}(r)$. In this figure $r = 30$ and $\mathbf{b} = 19 - 11i$ and the magnitudes of the values of $H_{\mathbf{b}}(\cdot)$ are indicated by colors.

Figure 5.1: The discrete ball-domains and Poisson kernels in them.

The boundary $\partial\mathbf{D}(r)$ contains the subset

$$\mathbf{C}(r) = \left\{ \mathbf{u} \in \mathbb{Z}^2 \mid \|\mathbf{u}\|_1 = r \right\}, \quad (5.8)$$

which will carry all harmonic measure seen from the interior points, and which is for that reason going to be the important part of the boundary for our purposes. In Figure 5.1(a), the subset $\mathbf{C}(r) \subset \partial\mathbf{D}(r)$ and its complement $\partial\mathbf{D}(r) \setminus \mathbf{C}(r)$ are colored white and black, respectively. Note that the discrete ball is the union of the interior and the subset which supports the harmonic measures,

$$\mathbf{B}_\bullet^\sharp(r) = \mathbf{D}^\circ(r) \cup \mathbf{C}(r).$$

A general graph-theoretic fact about Neumann Laplacians gives the following “divergence theorem”.

Lemma 5.5. *Let $\Delta_\sharp^{\mathbb{N};\mathbf{D}(r)}$ denote the discrete Neumann Laplacian in the domain $\mathbf{D}(r)$. Then for any two functions f, g on $\mathbf{D}(r)$ we have the equality*

$$\sum_{\mathbf{u} \in \mathbf{D}(r)} f(\mathbf{u}) ((\Delta_\sharp^{\mathbb{N};\mathbf{D}(r)} g)(\mathbf{u})) = \sum_{\mathbf{u} \in \mathbf{D}(r)} ((\Delta_\sharp^{\mathbb{N};\mathbf{D}(r)} f)(\mathbf{u})) g(\mathbf{u}).$$

Proof. Separate the contribution to each edge in both sums and observe that both sides are equal to $-\sum (f(\mathbf{v}) - g(\mathbf{u}))^2$, where the sum is over the edges $\{\mathbf{u}, \mathbf{v}\}$ of the graph $\mathbf{D}(r)$. \square

Let us now define the harmonic measures in a manner that facilitates applying the divergence theorem. Fix an interior vertex $\mathbf{v} \in \mathbf{D}^\circ(r)$ and consider the Dirichlet Green's function $\mathbf{u} \mapsto G_{\mathbf{D}(r)}^{\mathbf{D}}(\mathbf{u}, \mathbf{v})$ in $\mathbf{D}(r)$. Note that $\Delta_{\sharp}^{\mathbf{N};\mathbf{D}(r)} G_{\mathbf{D}(r)}^{\mathbf{D}}(\cdot, \mathbf{v})$ vanishes outside $\partial\mathbf{D}(r) \cup \{\mathbf{v}\}$, because the Neumann Laplacian $\Delta_{\sharp}^{\mathbf{N};\mathbf{D}(r)}$ and the Dirichlet Laplacian $\Delta_{\sharp}^{\mathbf{D};\mathbf{D}(r)}$ agree at all interior vertices and $\Delta_{\sharp}^{\mathbf{D};\mathbf{D}(r)} G_{\mathbf{D}(r)}^{\mathbf{D}}(\cdot, \mathbf{v}) = -\delta_{\mathbf{v}}(\cdot)$. In fact, $\Delta_{\sharp}^{\mathbf{N};\mathbf{D}(r)} G_{\mathbf{D}(r)}^{\mathbf{D}}(\cdot, \mathbf{v})$ also vanishes on $\partial\mathbf{D}(r) \setminus \mathbf{C}(r)$, i.e., on the black boundary points in Figure 5.1(a), because all neighbors of such a boundary point are themselves boundary points and the Dirichlet Green's function vanishes on the boundary. We therefore define the harmonic measure $H_{\mathbf{b}}(\mathbf{v})$ of $\mathbf{b} \in \mathbf{C}(r)$ seen from $\mathbf{v} \in \mathbf{D}^\circ(r)$ by the formula

$$H_{\mathbf{b}}(\mathbf{v}) = \Delta_{\sharp}^{\mathbf{N};\mathbf{D}(r)} G_{\mathbf{D}(r)}^{\mathbf{D}}(\mathbf{b}, \mathbf{v}),$$

so that the Neumann Laplacian of the Dirichlet Green's function takes the following form.

Lemma 5.6. *For $\mathbf{v} \in \mathbf{D}^\circ(r)$ we have*

$$\Delta_{\sharp}^{\mathbf{N};\mathbf{D}(r)} G_{\mathbf{D}(r)}^{\mathbf{D}}(\cdot, \mathbf{v}) = -\delta_{\mathbf{v}}(\cdot) + \sum_{\mathbf{b} \in \mathbf{C}(r)} H_{\mathbf{b}}(\mathbf{v}) \delta_{\mathbf{b}}(\cdot).$$

Proof. This follows by construction of $H_{\mathbf{b}}(\mathbf{v})$ and the observations preceding it. \square

Observe also that the Neumann Laplacian always yields a zero-average function, so the harmonic measure terms above must compensate the term $-\delta_{\mathbf{v}}(\cdot)$, i.e., we have

$$\sum_{\mathbf{b} \in \mathbf{C}(r)} H_{\mathbf{b}}(\mathbf{v}) = 1. \quad (5.9)$$

With Lemma 5.6, we are ready to give the (almost) canonical representatives for linear local fields supported on the discrete ball $\mathbf{B}_{\bullet}^{\sharp}(r) = \mathbf{D}^\circ(r) \cup \mathbf{C}(r)$.

Proposition 5.7. *For any $\mathbf{v} \in \mathbf{D}^\circ(r)$, we have*

$$X(\mathbf{v}) + \text{Null} = \sum_{\mathbf{b} \in \mathbf{C}(r)} H_{\mathbf{b}}(\mathbf{v}) X(\mathbf{b}) + \text{Null}.$$

Proof. Use the divergence theorem, Lemma 5.5, with $f = G_{\mathbf{D}(r)}^{\mathbf{D}}(\cdot, \mathbf{v})$ and $g = X$ to get

$$\sum_{\mathbf{u} \in \mathbf{D}^\circ(r)} G_{\mathbf{D}(r)}^{\mathbf{D}}(\mathbf{u}, \mathbf{v}) ((\Delta_{\sharp}^{\mathbf{N};\mathbf{D}(r)} X)(\mathbf{u})) = -X(\mathbf{v}) + \sum_{\mathbf{b} \in \mathbf{C}(r)} H_{\mathbf{b}}(\mathbf{v}) X(\mathbf{b}),$$

where on the left-hand-side we used the vanishing of the Dirichlet Green's function f on the boundary to omit the boundary terms, and on the right-hand-side we used the formula of

Lemma 5.6 for the Neumann Laplacian of the Dirichlet Green's function f to omit all terms except $\mathbf{u} = \mathbf{v}$ and $\mathbf{u} \in \mathbf{C}(r)$. On the left hand side, since \mathbf{u} is an interior point of $\mathbf{D}(r)$, the Neumann Laplacian coincides with the square grid Laplacian, so $(\Delta_{\sharp}^{\mathbf{N}; \mathbf{D}(r)} X)(\mathbf{u}) = \Delta_{\sharp} X(\mathbf{u})$, which is null by Example 4.3. As a linear combination of null fields, the entire left hand side is therefore null. This proves the asserted equality modulo null fields, since the right hand side is exactly the difference of the nonnull terms on the two sides of the asserted equality. \square

We thus obtain representatives which are supported just on the boundary.

Lemma 5.8. *For every $r \in \mathbb{Z}_{>0}$, we have*

$$\mathcal{F}_{\nabla}^{\text{lin}}(r) = \text{span}_{\mathbb{C}} \left\{ X(\mathbf{b}) + \text{Null} \mid \mathbf{b} \in \mathbf{C}(r) \right\}.$$

Proof. By definition, the space $\mathcal{P}_{\nabla}^{\text{lin}}(r)$ is spanned by $X(\mathbf{v})$ for $\mathbf{v} \in \mathbf{B}_{\sharp}^{\circ}(r) = \mathbf{D}^{\circ}(r) \cup \mathbf{C}(r)$, and $\mathcal{F}_{\nabla}^{\text{lin}}(r)$ is correspondingly spanned by the equivalence classes modulo null fields. For any $\mathbf{v} \in \mathbf{D}^{\circ}(r)$, Proposition 5.7 expresses $X(\mathbf{v}) + \text{Null}$ as a linear combination of fields of the form $X(\mathbf{b}) + \text{Null}$ with $\mathbf{b} \in \mathbf{C}(r)$, so these suffice to span $\mathcal{F}_{\nabla}^{\text{lin}}(r)$. \square

This also leads to the desired dimension upper bounds.

Lemma 5.9. *For every $r \in \mathbb{Z}_{>0}$, we have*

$$\dim(\mathcal{F}_{\nabla}^{\text{lin}}(r)) = 4r - 1.$$

A basis of $\mathcal{F}_{\nabla}^{\text{lin}}(r)$ is given by $\mathbf{J}_{-k}I$ and $\bar{\mathbf{J}}_{-k}I$, for $1 \leq k \leq 2r - 1$, and $\mathbf{J}_{-2r}I - \bar{\mathbf{J}}_{-2r}I$.

Proof. Recall that by Lemma 5.4, we have $\dim(\mathcal{F}_{\nabla}^{\text{lin}}(r)) \geq 4r - 1$ and the given elements are linearly independent in $\mathcal{F}_{\nabla}^{\text{lin}}(r)$. It therefore only remains to show that $\dim(\mathcal{F}_{\nabla}^{\text{lin}}(r)) \leq 4r - 1$. The spanning set given in Lemma 5.8 is indexed by $\mathbf{C}(r)$, which has $|\mathbf{C}(r)| = 4r$ points; one more than the asserted dimension. However, applying Proposition 5.7 with $\mathbf{v} = 0$ we see that

$$X(0) + \text{Null} = \sum_{\mathbf{b} \in \mathbf{C}(r)} H_{\mathbf{b}}(0) X(\mathbf{b}) + \text{Null}.$$

Since the left-hand-side $X(0)$ is null by Example 4.2, the right-hand-side gives one linear relation in the spanning set, provided that the coefficients $H_{\mathbf{b}}(0)$ are not all zero. Equation (5.9) shows that they are not. \square

We are now ready to finish the proofs of the two main statements of the section.

Proof of Theorem 5.1. We want to show that the fields $\mathbf{J}_{-k}I$ and $\bar{\mathbf{J}}_{-k}I$, for $k \in \mathbb{Z}_{>0}$, form a basis of the space $\mathcal{F}_{\nabla}^{\text{lin}}$ of linear local fields. We already observed, based on Corollary 4.4, that these fields are linearly independent. It remains to show that they span. Using the filtration by radius, we see that any field $F \in \mathcal{F}_{\nabla}^{\text{lin}}$ belongs to $\mathcal{F}_{\nabla}^{\text{lin}}(r)$ for some $r \in \mathbb{Z}_{>0}$. Then by Lemma 5.9, F is indeed a linear combination of the basis elements of $\mathcal{F}_{\nabla}^{\text{lin}}(r)$, which are manifestly already linear combinations of elements of the asserted basis (5.3) of $\mathcal{F}_{\nabla}^{\text{lin}}$. \square

Proof of Corollary 5.2. By the filtrations by radii, it is again enough to focus on $\mathcal{P}_{\nabla}^{\text{lin}}(r)$ for a fixed $r \in \mathbb{Z}_{>0}$. The field polynomials $X(\mathbf{u})$ with $\mathbf{u} \in \mathbf{B}_{\bullet}^{\sharp}(r)$ are a basis of $\mathcal{P}_{\nabla}^{\text{lin}}(r)$ by construction, so the dimension is $\dim(\mathcal{P}_{\nabla}^{\text{lin}}(r)) = |\mathbf{B}_{\bullet}^{\sharp}(r)| = 2r^2 + 2r + 1$. In Lemma 5.9 we found that the dimension of the quotient $\mathcal{F}_{\nabla}^{\text{lin}}(r) = \mathcal{P}_{\nabla}^{\text{lin}}(r)/(\mathcal{N} \cap \mathcal{P}_{\nabla}^{\text{lin}}(r))$ equals $4r - 1$, so the dimension of $\mathcal{N} \cap \mathcal{P}_{\nabla}^{\text{lin}}(r)$ is the difference

$$\begin{aligned} \dim(\mathcal{N} \cap \mathcal{P}_{\nabla}^{\text{lin}}(r)) &= 2r^2 + 2r + 1 - (4r - 1) = 2r^2 - 2r + 2 \\ &= 2(r - 1)^2 + 2(r - 1) + 2 \\ &= |\mathbf{D}^{\circ}(r)| + 1. \end{aligned}$$

The known null fields $\Delta_{\sharp}X(\mathbf{u})$, for $\mathbf{u} \in \mathbf{D}^{\circ}(r)$, and $X(0)$ are all linearly independent in $\mathcal{N} \cap \mathcal{P}_{\nabla}^{\text{lin}}(r)$ (by standard considerations with the discrete Laplacians) and their number matches the above dimension of $\mathcal{N} \cap \mathcal{P}_{\nabla}^{\text{lin}}(r)$ calculated above, so they form a basis. \square

5.4 Homogeneous linear local fields. Recall that homogeneous local fields, i.e., eigenvectors of \mathbf{L}_0 and $\bar{\mathbf{L}}_0$, are important for the reason that their scaling limits are to be formed with definite renormalization related to the eigenvalues. With our complete understanding of the space of linear local fields from Theorem 5.1, let us see the consequences about homogeneous linear local fields.

In the same fashion as in (4.14), define the **homogeneous linear local fields** of conformal dimensions $\Delta, \bar{\Delta} \in \mathbb{C}$ as the subspace

$$(\mathcal{F}_{\nabla}^{\text{lin}})_{\Delta, \bar{\Delta}} := \ker \left(\mathbf{L}_0|_{\mathcal{F}_{\nabla}^{\text{lin}}} - \Delta \text{id}_{\mathcal{F}_{\nabla}^{\text{lin}}} \right) \cap \ker \left(\bar{\mathbf{L}}_0|_{\mathcal{F}_{\nabla}^{\text{lin}}} - \bar{\Delta} \text{id}_{\mathcal{F}_{\nabla}^{\text{lin}}} \right) \subset \mathcal{F}_{\nabla}^{\text{lin}}.$$

Proposition 5.10. *The Virasoro modes \mathbf{L}_0 and $\bar{\mathbf{L}}_0$ are simultaneously diagonalizable on $\mathcal{F}_{\nabla}^{\text{lin}}$, i.e., we have the vector space direct sum $\mathcal{F}_{\nabla}^{\text{lin}} = \bigoplus_{\Delta, \bar{\Delta}} (\mathcal{F}_{\nabla}^{\text{lin}})_{\Delta, \bar{\Delta}}$. Moreover, the non-trivial joint eigenspaces are one-dimensional; explicitly*

$$(\mathcal{F}_{\nabla}^{\text{lin}})_{\Delta, \bar{\Delta}} = \begin{cases} \mathbb{C} \mathbf{J}_{-k} I & \text{if } (\Delta, \bar{\Delta}) = (k, 0) \text{ with } k \in \mathbb{Z}_{>0} \\ \mathbb{C} \bar{\mathbf{J}}_{-k} I & \text{if } (\Delta, \bar{\Delta}) = (0, k) \text{ with } k \in \mathbb{Z}_{>0} \\ \{0\} & \text{otherwise.} \end{cases}$$

Proof. Theorem 5.1 says that the fields $\mathbf{J}_{-k} I$ and $\bar{\mathbf{J}}_{-k} I$ for $k \in \mathbb{Z}_{>0}$ constitute a basis of $\mathcal{F}_{\nabla}^{\text{lin}}$, and for the basis vectors we have

$$\mathbf{L}_0(\mathbf{J}_{-k} I) = k \mathbf{J}_{-k} I, \quad \bar{\mathbf{L}}_0(\bar{\mathbf{J}}_{-k} I) = k \bar{\mathbf{J}}_{-k} I \quad \text{and} \quad \bar{\mathbf{L}}_0(\mathbf{J}_{-k} I) = \mathbf{L}_0(\bar{\mathbf{J}}_{-k} I) = 0 + \text{Null}$$

by the same calculation as in the Fock space $\mathcal{F} \otimes \bar{\mathcal{F}}$. \square

Remark 5.1. Exactly two Virasoro primary states among the ones listed in (2.12) – (2.17) correspond to linear local fields: the counterparts of (2.13) and (2.14) are

$$\begin{aligned} \mathbf{J}_{-1} I & \quad \text{with a representative} \quad J_{-1} = \mathfrak{i} \left(\frac{1}{2} \frac{X(1) - X(-1)}{2} - \frac{\mathfrak{i}}{2} \frac{X(\mathfrak{i}) - X(-\mathfrak{i})}{2} \right) \\ \bar{\mathbf{J}}_{-1} I & \quad \text{with a representative} \quad \bar{J}_{-1} = -\mathfrak{i} \left(\frac{1}{2} \frac{X(1) - X(-1)}{2} + \frac{\mathfrak{i}}{2} \frac{X(\mathfrak{i}) - X(-\mathfrak{i})}{2} \right), \end{aligned}$$

where the representatives were obtained from (4.13). Note that although (2.13) and (2.14) are as CFT fields interpreted as the holomorphic and antiholomorphic currents, in our discrete current mode constructions (4.11) we needed slightly different discretizations of the holomorphic and antiholomorphic currents. The representatives of $J_{-1}I$ and $\bar{J}_{-1}I$ above are, respectively, the averages of $j(\mathbf{u}_m)$ and $\bar{j}(\mathbf{u}_m)$ over the four midpoints $\mathbf{u}_m \in \{\frac{\pm 1}{2}, \frac{\pm i}{2}\}$ of edges adjacent to the origin. \diamond

6 Local fields of higher degree

In the previous section, we fully characterized linear local fields of the DGFF, so the remaining task for proving the isomorphism $\mathcal{F}_\nabla \cong \mathcal{F} \otimes \overline{\mathcal{F}}$ is to characterize local fields of higher degree. By definition (4.5), the space \mathcal{P}_∇ of field polynomials is a polynomial algebra, i.e., a symmetric tensor algebra, which is naturally graded

$$\mathcal{P}_\nabla = \bigoplus_{d \in \mathbb{Z}_{\geq 0}} S^d \mathcal{P}_\nabla^{\text{lin}}$$

by the degrees of the polynomials, i.e., so that the degree d component is the d th symmetric tensor power of the space (5.1) of linear field polynomials,

$$S^d \mathcal{P}_\nabla^{\text{lin}} = \text{span}_{\mathbb{C}} \left\{ \prod_{i=1}^d X(\mathbf{u}_i) \mid \mathbf{u}_1, \dots, \mathbf{u}_d \in \mathbb{Z}^2 \right\}.$$

The space (4.10) of local fields inherits a grading

$$\mathcal{F}_\nabla = \bigoplus_{d \in \mathbb{Z}_{\geq 0}} \mathcal{F}_\nabla^{(d)}$$

where $\mathcal{F}_\nabla^{(d)} = S^d \mathcal{P}_\nabla^{\text{lin}} / \mathcal{N}$ is the quotient of the space of degree d field polynomials by null fields. Recall, however, that while $\mathcal{P}_\nabla = \bigoplus_{d \in \mathbb{Z}_{\geq 0}} S^d \mathcal{P}_\nabla^{\text{lin}}$ is a graded algebra, the space $\mathcal{F}_\nabla = \mathcal{P}_\nabla / \mathcal{N}$ of local fields does not even inherit a multiplication from the polynomials: the null fields do not form an ideal, let alone a graded ideal. The space $\mathcal{F}_\nabla = \bigoplus_{d \in \mathbb{Z}_{\geq 0}} \mathcal{F}_\nabla^{(d)}$ of local fields is merely a graded vector space.

In order to fully characterize higher degree fields and to finish the proof of isomorphism $\mathcal{F}_\nabla \cong \mathcal{F} \otimes \overline{\mathcal{F}}$, we will introduce normal ordering in \mathcal{F}_∇ , which partially rectifies the lack of an algebra structure and allows us to reduce the analysis of higher degree local fields to that of the linear local fields. Let us also note that the normal ordering will be combinatorially similar with Wick products with respect to a Gaussian measure, but it is not a Wick product for the DGFF measure in any discrete domain (with any boundary conditions). Indeed, we are operating at the level of abstract local fields, which can be evaluated in an arbitrary discrete domain and with either Dirichlet or Neumann boundary conditions. Following the idea from CFT that only a local coordinate specification should affect the normal ordering procedure, we use infinite square lattice quantities in the normal ordering contractions.

The normal ordering is defined in Section 6.1. Besides essentially combinatorial lemmas, we prove Lemma 6.2, which expresses a crucial property that could be interpreted as null fields being an ideal with respect to normal ordering. In Section 6.2 we are then ready to state and prove our main result, Theorem 6.4, by combining this lemma with the full classification of linear local fields from the previous section. The proof also involves making a connection between the normal ordering and the algebraic structure provided by the current modes, and we illustrate the concrete computational nature of this procedure by giving a few examples of higher degree in Section 6.3.

6.1 Normal ordering. Let G denote the Green's function on the infinite square lattice \mathbb{Z}^2 [LaLi10], i.e., the unique function

$$G: \mathbb{Z}^2 \rightarrow \mathbb{R} \quad \text{satisfying} \quad \begin{cases} \Delta_{\sharp} G(\mathbf{u}) = -\delta_0(\mathbf{u}) \\ G(0) = 0 \\ G(\mathbf{u}) = -\frac{1}{2\pi} \log |\mathbf{u}| + C + \mathcal{O}(|\mathbf{u}|^{-2}) \text{ as } |\mathbf{u}| \rightarrow \infty, \end{cases} \quad (6.1)$$

where $C = -\frac{1}{2\pi}(\gamma + \frac{3}{2} \log 2)$ and γ is the Euler–Mascheroni constant. The infinite lattice Green's function is symmetric in the sense that $G(-\mathbf{u}) = G(\mathbf{u})$ for any $\mathbf{u} \in \mathbb{Z}^2$.

If $G(\mathbf{u})$ could be informally thought of as “ $\mathbb{E}[\phi(\mathbf{u}_0 + \mathbf{u})\phi(\mathbf{u}_0)]$ ” (for any $\mathbf{u}_0 \in \mathbb{Z}^2$ by virtue of translation invariance), then an informal counterpart two-point function appropriate for the evaluations $X(\mathbf{u}) \mapsto \phi(\mathbf{u}) - \phi(0)$ would be “ $\mathbb{E}[(\phi(\mathbf{u}) - \phi(0))(\phi(\mathbf{v}) - \phi(0))]$ ”. This motivates the definition of

$$G_{\nabla}: \mathbb{Z}^2 \times \mathbb{Z}^2 \rightarrow \mathbb{R} \quad \text{as} \quad G_{\nabla}(\mathbf{u}, \mathbf{v}) := G(\mathbf{u} - \mathbf{v}) - G(\mathbf{u}) - G(\mathbf{v}), \quad (6.2)$$

which has the symmetricity property $G_{\nabla}(\mathbf{u}, \mathbf{v}) = G_{\nabla}(\mathbf{v}, \mathbf{u})$. We use G_{∇} in normal ordering contractions as follows. We define **contractions** of linear field polynomials

$$\begin{aligned} \mathcal{P}_{\nabla}^{\text{lin}} \otimes \mathcal{P}_{\nabla}^{\text{lin}} &\rightarrow \mathbb{C} \\ L_1 \otimes L_2 &\mapsto \overline{L_1} L_2 \end{aligned}$$

by bilinear extension from the contractions of the linear basis monomials given by the formula

$$\overline{X(\mathbf{u})} X(\mathbf{v}) := 4\pi G_{\nabla}(\mathbf{u}, \mathbf{v}) \quad \text{for } \mathbf{u}, \mathbf{v} \in \mathbb{Z}^2.$$

The contractions are symmetric, $\overline{L_1} L_2 = \overline{L_2} L_1$, and thus well-defined on $S^2 \mathcal{P}_{\nabla}^{\text{lin}}$.

Example 6.1. For $\mathbf{u}, \mathbf{v} \in \mathbb{Z}^2$, we have $(\Delta_{\sharp} \overline{X(\mathbf{u})}) X(\mathbf{v}) = -4\pi \delta_{\mathbf{v}}(\mathbf{u}) + 4\pi \delta_0(\mathbf{u})$. \diamond

The **normal ordering** is a linear operation

$$\circ \cdots \circ: \mathcal{P}_{\nabla} \rightarrow \mathcal{P}_{\nabla}$$

on the space $\mathcal{P}_\nabla = \bigoplus_{d \in \mathbb{Z}_{\geq 0}} S^d \mathcal{P}_\nabla^{\text{lin}}$ of field polynomials, defined on the degree d subspace $S^d \mathcal{P}_\nabla^{\text{lin}}$ by

$$\circ \prod_{i=1}^d L_i \circ := \sum_{P' \in \text{PartPair}(d)} (-1)^{|P'|} \prod_{j \notin \bigcup P'} L_j \prod_{\{k, \ell\} \in P'} \overline{L_k L_\ell}, \quad (6.3)$$

where $\text{PartPair}(d)$ denotes the set of **partial pairings** of the d indices, i.e., collections P' of disjoint two-element subsets of $\{1, \dots, d\}$. One can alternatively view (6.3) as a sum of partitions ϖ of the index set $\{1, \dots, d\}$ to subsets of size at most two; then the number of pairs reads $|P'| = d - |\varpi|$, and the first and second products above range over the parts of size one and two, respectively.

Let us record two combinatorial formulas for later calculations.

Lemma 6.1. *Let $L_1, \dots, L_d \in \mathcal{P}_\nabla^{\text{lin}}$. Then we have*

$$\circ \prod_{i=1}^d L_i \circ = \prod_{i=1}^d L_i + R \quad \text{where} \quad R \in \bigoplus_{d' \leq d-2} S^{d'} \mathcal{P}_\nabla^{\text{lin}} \quad (6.4)$$

and

$$L_1 \circ \prod_{i=2}^d L_i \circ = \circ L_1 \prod_{i=2}^d L_i \circ + \sum_{i=2}^d \overline{L_1 L_i} \circ \prod_{\substack{2 \leq j \leq d \\ j \neq i}} L_j \circ \quad (6.5)$$

In particular, (6.4) shows that $\circ \cdots \circ : \mathcal{P}_\nabla \rightarrow \mathcal{P}_\nabla$ is bijective.

Proof. Formula (6.4) is obtained by separating in (6.3) the summand corresponding to the empty partial pairing $P' = \emptyset$, which gives the term $\prod_{i=1}^d L_i$, and the rest, which include at least one pair and therefore give terms of degrees $d' = d - 2|P'| \leq d - 2$. This “upper triangularity” formula allows one to uniquely solve $\prod_{i=1}^d L_i$ in terms of normally ordered products, inductively on the degree d , so it shows bijectivity of $\circ \cdots \circ : \mathcal{P}_\nabla \rightarrow \mathcal{P}_\nabla$.

Formula (6.5) is obtained by grouping the terms in $\circ \prod_{i=1}^d L_i \circ$ to those where the index 1 has no pair in the partial pairing P' (these produce the left hand side) and those where 1 has a pair i (these produce, up to a sign, the i th summand on the right hand side). \square

Example 6.2. For $\mathbf{u}, \mathbf{v} \in \mathbb{Z}^2$, with the formula from Example 6.1, we get

$$\begin{aligned} \circ (\triangle_\# X(\mathbf{u})) X(\mathbf{v}) \circ &= (\triangle_\# X(\mathbf{u})) X(\mathbf{v}) - \overline{(\triangle_\# X(\mathbf{u}))} X(\mathbf{v}) \\ &= (\triangle_\# X(\mathbf{u})) X(\mathbf{v}) + 4\pi \delta_{\mathbf{v}}(\mathbf{u}) - 4\pi \delta_0(\mathbf{u}). \end{aligned}$$

Observe that this field polynomial is null by Example 4.4. \diamond

The observation in the example above generalizes.

Lemma 6.2. *Let $L_1, \dots, L_d \in \mathcal{P}_\nabla^{\text{lin}}$. If $L_j \in \mathcal{N}$ for some j , then also $\circ \prod_{i=1}^d L_i \circ \in \mathcal{N}$.*

We give the proof shortly, but we first note the important consequence. Lemma 6.2 implies that the normally ordered product of d linear field polynomials $\circ \cdots \circ : (\mathcal{P}_\nabla^{\text{lin}})^{\otimes d} \rightarrow \mathcal{P}_\nabla$ factorizes through the quotient by null fields in each tensorand, giving rise to well-defined normally ordered products $\bullet \cdots \bullet : (\mathcal{F}_\nabla^{\text{lin}})^{\otimes d} \rightarrow \mathcal{F}_\nabla$ of d local fields by

$$\bullet (L_1 + \text{Null}) \otimes \cdots \otimes (L_d + \text{Null}) \bullet := \circ \prod_{i=1}^d L_i \circ + \text{Null} \quad (6.6)$$

(we write the tensor products explicitly on the left hand side in order to avoid a false impression of a ring structure on \mathcal{F}_∇). Extending linearly to all degrees d and noting symmetricity in the arguments, we get a normal ordering operation

$$\bullet \cdots \bullet : \mathcal{F}_\nabla \rightarrow \mathcal{F}_\nabla$$

on the space of local fields of the DGFF. Moreover, it follows from Lemma 6.1 that this normal ordering is surjective.

Proof of Lemma 6.2. Without loss of generality, assume $L_1 \in \mathcal{N}$. By definition, we must check (4.7) for $F = \circ \prod_{i=1}^d L_i \circ$. Let $\Omega_\delta \subset \delta\mathbb{Z}^2$ be a discrete domain, $(f_a)_{a \in A}$ a finite collection of zero-average test functions $f_a \in \text{Fun}_\nabla(\Omega_\delta)$, and $\mathbf{z} \in \Omega_\delta$ a point, such that the supports are meaningful and distinct as required in the condition (4.7). We must show the vanishing of

$$\begin{aligned} & \mathbb{E}_{\Omega_\delta}^{\text{D/N}} \left[\left(\text{ev}_{\mathbf{z}}^{\Omega_\delta} \circ \prod_{i=1}^d L_i \circ \right) \prod_{a \in A} \langle \Phi, f_a \rangle \right] \\ &= \sum_{P'} (-1)^{|P'|} \left(\prod_{\{k, \ell\} \in P'} \overline{L_k L_\ell} \right) \mathbb{E}_{\Omega_\delta}^{\text{D/N}} \left[\prod_{j \notin P'} (\text{ev}_{\mathbf{z}}^{\Omega_\delta} L_j) \prod_{a \in A} \langle \Phi, f_a \rangle \right]. \end{aligned}$$

Use Wick's formula to calculate the expected value on the second line as a sum over pairings P of $A \sqcup (\{1, \dots, d\} \setminus P')$, schematically (avoiding cumbersome notation that would be needed to explicitly spell out three possible forms of the pairs in P)

$$\begin{aligned} & \mathbb{E}_{\Omega_\delta}^{\text{D/N}} \left[\left(\text{ev}_{\mathbf{z}}^{\Omega_\delta} \circ \prod_{i=1}^d L_i \circ \right) \prod_{a \in A} \langle \Phi, f_a \rangle \right] \\ &= \sum_{P'} (-1)^{|P'|} \left(\prod_{\{k, \ell\} \in P'} \overline{L_k L_\ell} \right) \sum_P \left(\prod_{\text{pairs} \in P} \mathbb{E}_{\Omega_\delta}^{\text{D/N}} [(\text{pair product})] \right). \end{aligned}$$

Note that for each (P', P) , the index 1 of the linear null field L_1 appears either in one pair in P' (leading to a normal ordering contraction factor) or in one of the pairs in P (leading to a suitable DGFF two-point function factor), and it is in fact useful to consider three separate cases of which exactly one occurs:

- $\{1, a\} \in P$ for some $a \in A$
- $\{1, m\} \in P$ for some $m \in \{2, \dots, d\}$
- $\{1, m\} \in P'$ for some $m \in \{2, \dots, d\}$.

In the first case, there is a factor

$$\mathbb{E}_{\Omega_\delta}^{\text{D/N}} \left[(\text{ev}_{\mathbf{z}}^{\Omega_\delta} L_1) \langle \Phi, f_a \rangle \right],$$

which vanishes since L_1 is a null field and the supports of L_1 and f_a satisfy the disjointness and well-definedness conditions. Therefore only the last two cases contribute to our quantity of interest. Moreover, the contributions of those can be naturally matched pairwise. Namely, fixing $m \in \{2, \dots, d\}$, in the second case we may remove the pair $\{1, m\}$ from P to get a pairing \hat{P} of $(\{1, \dots, d\} \setminus \{1, m\}) \sqcup A$, and in the third case we may remove the pair $\{1, m\}$ from P' to get a partial pairing \hat{P}' of $\{1, \dots, d\} \setminus \{1, m\}$, and then the sum of all contributions with the fixed m combine to take the form

$$\sum_{\hat{P}'} \sum_{\hat{P}} \left((-1)^{|\hat{P}'|+1} \overline{L_1} L_m + (-1)^{|\hat{P}'|} \mathbb{E}_{\Omega_\delta}^{\text{D/N}} \left[(\text{ev}_{\mathbf{z}}^{\Omega_\delta} L_1) (\text{ev}_{\mathbf{z}}^{\Omega_\delta} L_m) \right] \right) A(\hat{P}') B(\hat{P}),$$

where $A(\hat{P}')$ and $B(\hat{P})$ are products of common contraction factors and common DGFF two-point function factors, respectively (writing them explicitly is possible but will not be needed). We will show the vanishing of the first factor, which up to a sign equals

$$-\overline{L_1} L_m + \mathbb{E}_{\Omega_\delta}^{\text{D/N}} \left[(\text{ev}_{\mathbf{z}}^{\Omega_\delta} L_1) (\text{ev}_{\mathbf{z}}^{\Omega_\delta} L_m) \right], \quad (6.7)$$

and this will conclude the proof.

The key to prove the vanishing of (6.7) is the complete classification of linear null fields from the previous section, Corollary 5.2. The classification shows that $L_1 \in \mathcal{P}_{\nabla}^{\text{lin}} \cap \mathcal{N}$ is a linear combination of field polynomials of the forms $\Delta_{\#} X(\mathbf{u})$, for $\mathbf{u} \in \mathbb{Z}^2$, and $X(0)$. By linearity in L_1 , we may assume L_1 is exactly one of these. The latter case is trivial, so we will assume $L_1 = \Delta_{\#} X(\mathbf{u})$ for some $\mathbf{u} \in \mathbb{Z}^2$. By linearity in L_m we may also assume $L_m = X(\mathbf{v})$ for some $\mathbf{v} \in \mathbb{Z}^2$. In this case (6.7) reads

$$-(\Delta_{\#} \overline{X(\mathbf{u})}) X(\mathbf{v}) + \mathbb{E}_{\Omega_\delta}^{\text{D/N}} \left[(\Delta_{\#} \phi(\mathbf{z} + \delta \mathbf{u})) (\phi(\mathbf{z} + \delta \mathbf{v}) - \phi(\mathbf{z})) \right].$$

The first term here was calculated in Example 6.1; it is $4\pi\delta_{\mathbf{v}}(\mathbf{u}) - 4\pi\delta_0(\mathbf{u})$. Directly by the defining covariance of the DGFF, the second term equals

$$\Delta_{\#} G_{\Omega_\delta}^{\text{D/N}}(\mathbf{w}, \mathbf{z} + \delta \mathbf{v}) \big|_{\mathbf{w}=\mathbf{z}+\delta \mathbf{u}} - \Delta_{\#} G_{\Omega_\delta}^{\text{D/N}}(\mathbf{w}, \mathbf{z}) \big|_{\mathbf{w}=\mathbf{z}+\delta \mathbf{u}}.$$

Since $\mathbf{z} + \delta \mathbf{u} \in \Omega_\delta \setminus \partial\Omega_\delta$ by the requirements (4.7) of supports, this second term simplifies to $-4\pi\delta_{\mathbf{v}}(\mathbf{u}) + 4\pi\delta_0(\mathbf{u})$. Therefore it cancels the first term, completing the proof. \square

Let us emphasize again that Lemma 6.2 is key to us, since it allows us to define normal ordering on the level of local fields of the DGFF, not just field polynomials. The key ingredients enabling its proof, in turn, were the full classification of linear null fields from Section 5, and the fact that the infinite lattice Green's functions used in our domain-independent normal orderings had the exact same form of local singularities as the DGFF two-point functions in any discrete domain.

6.2 Local fields form a Fock space. In this section we will prove that the space \mathcal{F}_∇ of local fields of the DGFF is a Fock space. The local fields $\bar{J}_{-k'_m} \cdots \bar{J}_{-k'_1} J_{-k_m} \cdots J_{-k_1} I$ obtained by repeated current mode actions on the identity field correspond to the Fock space basis vectors. In order to show that these fields span \mathcal{F}_∇ , we first want to express them in terms of explicit normal ordered products of linear local fields.

Lemma 6.3. *For any $k_1, \dots, k_m, k'_1, \dots, k'_m \in \mathbb{Z}_{>0}$, we have the following equality in \mathcal{F}_∇ :*

$$\bullet \bar{J}_{-k_m} I \otimes \cdots \otimes \bar{J}_{-k_1} I \otimes J_{-k_m} I \otimes \cdots \otimes J_{-k_1} I \bullet = \bar{J}_{-k'_m} \cdots \bar{J}_{-k'_1} J_{-k_m} \cdots J_{-k_1} I.$$

This expression is the final ingredient of the proof of our first main result, so before giving the proof of the lemma, let us show how we conclude with it.

Theorem 6.4. *The space \mathcal{F}_∇ of local fields of the discrete Gaussian free field and the Fock space $\mathcal{F} \otimes \overline{\mathcal{F}}$ are isomorphic,*

$$\mathcal{F}_\nabla \cong \mathcal{F} \otimes \overline{\mathcal{F}},$$

as representations of two commuting Heisenberg algebras, and as representations of two commuting Virasoro algebras.

Proof. On both \mathcal{F}_∇ and $\mathcal{F} \otimes \overline{\mathcal{F}}$, the Virasoro representations are obtained from the Heisenberg representations by the Sugawara construction, so it suffices to prove isomorphism as representations of Heisenberg algebras. In Corollary 4.4 we already established $\mathcal{F} \otimes \overline{\mathcal{F}} \cong (\text{Hei} \otimes \overline{\text{Hei}})I \subset \mathcal{F}_\nabla$ as representations of Heisenberg algebras, so to complete the proof we must show $\mathcal{F}_\nabla \subset (\text{Hei} \otimes \overline{\text{Hei}})I$.

So let $F + \text{Null} \in \mathcal{F}_\nabla$ be an arbitrary local field, with $F \in \mathcal{P}_\nabla$ denoting a representative field polynomial. By the surjectivity of the normal ordering that was observed in Lemma 6.1, we can find some $\tilde{F} \in \mathcal{P}_\nabla$ such that $\circ \tilde{F} \circ = F$. Split the field polynomial \tilde{F} to homogeneous pieces, i.e., write $\tilde{F} = \sum_{d=0}^D \tilde{F}_d$ with $\tilde{F}_d \in S^d \mathcal{P}_\nabla^{\text{lin}}$. By linearity, it suffices to show that $\circ \tilde{F}_d \circ + \text{Null} \in (\text{Hei} \otimes \overline{\text{Hei}})I$ for each d . Factorize the homogeneous field polynomial \tilde{F}_d into linear factors, i.e., write $\tilde{F}_d = \prod_{i=1}^d L_i$ for some $L_1, \dots, L_d \in \mathcal{P}_\nabla^{\text{lin}}$, and note that by definition (6.6) we then have

$$\circ \tilde{F}_d \circ + \text{Null} = \circ L_1 \cdots L_d \circ + \text{Null} = \bullet (L_1 + \text{Null}) \otimes \cdots \otimes (L_d + \text{Null}) \bullet.$$

Theorem 5.1 gives a basis for linear local fields, and it in particular guarantees that for each $i = 1, \dots, d$, we can write $L_i + \text{Null}$ as a linear combination of $J_{-k}I$ and $\bar{J}_{-k}I$, $k \in \mathbb{Z}_{>0}$.

Lemma 6.3 explicitly shows that normal ordering applied to such linear factors is expressible in terms of local fields of the form $\bar{J}_{-k'_m'} \cdots \bar{J}_{-k'_1} J_{-k_m} \cdots J_{-k_1} I$, which are manifestly in $(\text{Hei} \otimes \overline{\text{Hei}})I$. We conclude that $F + \text{Null} \in (\text{Hei} \otimes \overline{\text{Hei}})I$ as desired. \square

For the proof of Lemma 6.3, we still need one auxiliary result. In it, and in the proof of Lemma 6.3 itself, we will use the explicit expressions given in Example 4.5 for the field polynomials $J_{-k}, \bar{J}_{-k} \in \mathcal{P}_{\nabla}^{\text{lin}}$ which are representatives of the linear local fields $J_{-k}I, \bar{J}_{-k}I \in \mathcal{F}_{\nabla}^{\text{lin}}$, respectively. Recall that the explicit expressions are

$$J_{-k} := \frac{i}{2\pi} \sum_{\mathbf{u}_m \in S_m^{[-k]}} \bar{\partial}_{\sharp} \mathbf{u}_m^{[-k]} \partial_{\sharp}^{\bullet} X(\mathbf{u}_m) \quad \text{and} \quad \bar{J}_{-k} := \frac{-i}{2\pi} \sum_{\mathbf{u}_m \in S_m^{[-k]}} \partial_{\sharp} \bar{\mathbf{u}}_m^{[-k]} \bar{\partial}_{\sharp}^{\bullet} X(\mathbf{u}_m),$$

where as in (3.22), we denote by $S^{[-k]} \subset \mathbb{Z}_{\diamond}^2 \cup \mathbb{Z}_m^2$ the neighborhood of the origin where the function $\mathbf{u} \mapsto \mathbf{u}^{[-k]}$ is not discrete holomorphic, and $S_m^{[-k]} = S^{[-k]} \cap \mathbb{Z}_m^2$.

Lemma 6.5. *For any $k, \ell \in \mathbb{Z}_{>0}$ and any positively oriented corner contour γ satisfying $S^{[k]} \cup S^{[\ell]} \subset \text{int}^{\sharp} \gamma$, we have*

$$\begin{aligned} \oint_{\gamma}^{\sharp} \mathbf{u}_{\diamond}^{[-k]} (\overline{\partial_{\sharp}^{\bullet} X(\mathbf{u}_m)}) J_{-\ell} d^{\sharp} \mathbf{u} &= \oint_{\gamma}^{\sharp} \mathbf{u}_{\diamond}^{[-k]} (\overline{\partial_{\sharp}^{\bullet} X(\mathbf{u}_m)}) \bar{J}_{-\ell} d^{\sharp} \mathbf{u} = 0, \\ \oint_{\gamma}^{\sharp} \bar{\mathbf{u}}_{\diamond}^{[-k]} (\overline{\bar{\partial}_{\sharp}^{\bullet} X(\mathbf{u}_m)}) J_{-\ell} d^{\sharp} \bar{\mathbf{u}} &= \oint_{\gamma}^{\sharp} \bar{\mathbf{u}}_{\diamond}^{[-k]} (\overline{\bar{\partial}_{\sharp}^{\bullet} X(\mathbf{u}_m)}) \bar{J}_{-\ell} d^{\sharp} \bar{\mathbf{u}} = 0. \end{aligned}$$

Proof. The proofs of vanishing of all four integrals are similar, so let us just look at the top left one. Using the explicit expression of $J_{-\ell}$ recalled above and the definition of contractions, we find

$$\begin{aligned} \oint_{\gamma}^{\sharp} \mathbf{u}_{\diamond}^{[-k]} (\overline{\partial_{\sharp}^{\bullet} X(\mathbf{u}_m)}) J_{-\ell} d^{\sharp} \mathbf{u} &= \frac{i}{2\pi} \sum_{\mathbf{v}_m \in S_m^{[\ell]}} \bar{\partial}_{\sharp} \mathbf{v}_m^{[-\ell]} \oint_{\gamma}^{\sharp} \mathbf{u}_{\diamond}^{[-k]} (\overline{\partial_{\sharp}^{\bullet} X(\mathbf{u}_m)}) (\partial_{\sharp}^{\bullet} X(\mathbf{v}_m)) d^{\sharp} \mathbf{u} \\ &= \frac{i}{2\pi} \sum_{\mathbf{v}_m \in S_m^{[\ell]}} \bar{\partial}_{\sharp} \mathbf{v}_m^{[-\ell]} \oint_{\gamma}^{\sharp} \mathbf{u}_{\diamond}^{[-k]} (\partial_{\sharp}^{\bullet})_{\mathbf{u}} (\partial_{\sharp}^{\bullet})_{\mathbf{v}} G_{\nabla}(\mathbf{u}_m, \mathbf{v}_m) d^{\sharp} \mathbf{u}, \end{aligned}$$

Note moreover, that the function $\mathbf{u}_m \mapsto (\partial_{\sharp}^{\bullet})_{\mathbf{u}} (\partial_{\sharp}^{\bullet})_{\mathbf{v}} G_{\nabla}(\mathbf{u}_m, \mathbf{v}_m)$ is discrete holomorphic away from the neighbors of \mathbf{v}_m by the factorization (3.6) of the Laplacian and the discrete harmonicity of $G_{\nabla}(\cdot, \mathbf{v})$ away from 0 and \mathbf{v} . The integrand in the last discrete contour integral has the asymptotic behaviour $\mathcal{O}(|\mathbf{u}|^{-k-2})$ as $|\mathbf{u}| \rightarrow \infty$. By discrete holomorphicity of the integrand, the contour γ can be taken arbitrarily large without changing the result. On a square contour at distance R from the origin, the integral is $\mathcal{O}(R^{-k-1})$, so taking $R \rightarrow \infty$ shows that it vanishes for $k > 0$. \square

We are then ready to provide the proof that we postponed earlier.

Proof of Lemma 6.3. We use the representative field polynomials $J_{-k}, \bar{J}_{-k} \in \mathcal{P}_{\nabla}^{\text{lin}}$ given in Example 4.5 for the linear local fields $J_{-k}I = J_{-k} + \text{Null}$ and $\bar{J}_{-k}I = \bar{J}_{-k} + \text{Null}$. In particular, the left hand side of the asserted equality becomes

$$\bullet \bar{J}_{-k_m} I \otimes \cdots \otimes \bar{J}_{-k_1} I \otimes J_{-k_n} I \otimes \cdots \otimes J_{-k_1} I \bullet = \circ \prod_{i=1}^m J_{-k_i} \prod_{i'=1}^{m'} \bar{J}_{-k_{i'}} \circ + \text{Null}.$$

To compare this with the right hand side in the assertion,

$$\bar{J}_{-k'_{m'}} \cdots \bar{J}_{-k'_1} J_{-k_m} \cdots J_{-k_1} I,$$

we argue by induction on the degree $m + m'$. In order to simplify notation, we spell out the details only in the case of $m' = 0$; the proof of the general case would just consist of splitting the induction step into two similarly handled cases and would involve decorating some of the symbols with bars and primes ($\bar{J}_{-k'_{i'}}$, $\bar{J}_{-k'_{i'}}$, $\bar{\partial}_{\sharp}^{\bullet}$, etc.).

So focusing on $m' = 0$, we seek to prove the equality of

$$\circ \prod_{i=1}^m J_{-k_i} \circ + \text{Null} \quad \text{and} \quad J_{-k_m} \cdots J_{-k_1} I.$$

Normal ordering does nothing to polynomials of degree at most one (there are no nontrivial partial pairings of a set with fewer than two elements), so the cases $m = 0$ and $m = 1$ are clear: for $m = 0$ the equality is just the definition of the identity local field, and for $m = 1$ it is our preferred representative choice $J_{-k_1} + \text{Null} = J_{-k_1} I$.

For brevity, from here on, denote

$$F = \prod_{1 \leq i \leq m} J_{-k_i} \quad \text{and} \quad \widehat{F}_j = \prod_{\substack{1 \leq i \leq m \\ i \neq j}} J_{-k_i} \quad \text{for } j = 1, \dots, m.$$

Inductively, we assume $\circ F \circ + \text{Null} = J_{-k_m} \cdots J_{-k_1} I$, and we must prove the equality of

$$\circ F J_{-k_{m+1}} \circ + \text{Null} \quad \text{and} \quad J_{-k_{m+1}} J_{-k_m} \cdots J_{-k_1} I = J_{-k_{m+1}} (\circ F \circ + \text{Null}).$$

We will use (6.5) to rewrite both sides, and we then compare the results. On the left hand side, rewriting directly gives

$$\circ F J_{-k_{m+1}} \circ = J_{-k_{m+1}} \circ F \circ - \sum_{j=1}^m J_{-k_{m+1}} \overline{J_{-k_j}} \circ \widehat{F}_j \circ. \quad (6.8)$$

On the right hand side we first take a suitable corner contour γ for the action of $J_{-k_{m+1}}$ and unravel the definition of the action of the current mode

$$J_{-k_{m+1}} (\circ F \circ + \text{Null}) = \frac{1}{2\pi} \oint_{\gamma}^{\sharp} \mathbf{u}_{\diamond}^{[-k_{m+1}]} \partial_{\sharp}^{\bullet} X(\mathbf{u}_m) \circ F \circ + \text{Null}.$$

Calculating the discrete contour integral using discrete Stokes' formula (3.18) and recalling the explicit expression for $J_{-k_{m+1}}$, we get

$$J_{-k_{m+1}}(\circ F \circ + \text{Null}) = \left(J_{-k_{m+1}} + \frac{i}{2\pi} \sum_{\mathbf{u}_\diamond \in \text{int}^\# \gamma} \mathbf{u}_\diamond^{[-k_{m+1}]} \bar{\partial}_\# \partial_\#^\bullet X(\mathbf{u}_\diamond) \right) \circ F \circ + \text{Null}.$$

In the second term, we then recall the factorization (3.6) of the discrete Laplacian and rewrite using (6.5) to see that

$$\begin{aligned} & \frac{i}{2\pi} \sum_{\mathbf{u}_\diamond \in \text{int}^\# \gamma} \mathbf{u}_\diamond^{[-k_{m+1}]} \bar{\partial}_\# \partial_\#^\bullet X(\mathbf{u}_\diamond) \circ F \circ \\ &= \frac{i}{4\pi} \sum_{\mathbf{u}_\diamond \in \text{int}^\# \gamma} \mathbf{u}_\diamond^{[-k_{m+1}]} \left(\circ \Delta_\# X(\mathbf{u}_\diamond) F \circ + \sum_{j=1}^m (\Delta_\# \overline{X(\mathbf{u}_\diamond)}) J_{-k_j} \circ \widehat{F}_j \circ \right). \end{aligned}$$

The linear factor $\Delta_\# X(\mathbf{u}_\diamond)$ is null in the fully normal ordered term, so by Lemma 6.2 the corresponding normal ordered product is also null. Combining, we have found that

$$\begin{aligned} & J_{-k_{m+1}}(\circ F \circ + \text{Null}) \\ &= J_{-k_{m+1}} \circ F \circ + \frac{i}{4\pi} \sum_{j=1}^m \sum_{\mathbf{u}_\diamond \in \text{int}^\# \gamma} \mathbf{u}_\diamond^{[-k_{m+1}]} (\Delta_\# \overline{X(\mathbf{u}_\diamond)}) J_{-k_j} \circ \widehat{F}_j \circ + \text{Null}. \end{aligned} \quad (6.9)$$

Comparing (6.8) and (6.9), we see that it suffices to prove that for each $j = 1, \dots, m$, the difference

$$J_{-k_{m+1}} \overline{J_{-k_j}} + \frac{i}{4\pi} \sum_{\mathbf{u}_\diamond \in \text{int}^\# \gamma} \mathbf{u}_\diamond^{[-k_{m+1}]} (\Delta_\# \overline{X(\mathbf{u}_\diamond)}) J_{-k_j}$$

of the explicitly written terms is null. Again writing out the definition of the representative $J_{-k_{m+1}}$ and using the same factorization of the discrete Laplacian and Stokes' formula, this difference becomes

$$\begin{aligned} & \frac{i}{2\pi} \sum_{\mathbf{u}_m \in S_m^{[k_{m+1}]}} \bar{\partial}_\# \mathbf{u}_m^{[-k_{m+1}]} \partial_\#^\bullet \overline{X(\mathbf{u}_m) J_{-k_j}} + \frac{i}{2\pi} \sum_{\mathbf{u}_\diamond \in \text{int}^\# \gamma} \mathbf{u}_\diamond^{[-k_{m+1}]} (\bar{\partial}_\# \partial_\#^\bullet \overline{X(\mathbf{u}_\diamond)}) J_{-k_j} \\ &= \frac{1}{2\pi} \oint_\gamma^\# \mathbf{u}_m^{[-k_{m+1}]} (\partial_\#^\bullet \overline{X(\mathbf{u}_m)}) J_{-k_j} d^\# \mathbf{u}. \end{aligned}$$

This discrete contour integral indeed vanishes by Lemma 6.5, and thus the proof of the induction step is complete, too. \square

6.3 Concrete examples of higher degree local fields. Lemma 6.3 was necessary for our proof of Theorem 6.4, but it can also be applied directly to give concrete representatives of local fields. Let us give two examples.

In our first example, we elaborate on the details of the discretization of the gradient squared of the DGFF which was discussed in the introduction. For this, we start by admitting an observation about our conformal field theory (this fact would properly make sense only in Section 7): the CFT field $\frac{1}{4}\|\nabla\varphi\|^2$, suitably regularized, is the Virasoro primary with conformal weights $(1,1)$ seen in (2.15), i.e., $j_{-1}\bar{j}_{-1}\mathbb{1}$. The local field of the DGFF corresponding to it is

$$J_{-1}\bar{J}_{-1}I = \bullet J_{-1}I \otimes \bar{J}_{-1}I \bullet = \circ J_{-1}\bar{J}_{-1} \circ + \text{Null}.$$

A concrete representative for this quadratic primary field is therefore obtained with formulas of Example 4.5 and some simplification

$$\begin{aligned} \circ J_{-1}\bar{J}_{-1} \circ &= J_{-1}\bar{J}_{-1} - \overline{J_{-1}\bar{J}_{-1}} \\ &= \frac{1}{4} \left(\left(\frac{X(1) - X(-1)}{2} \right)^2 + \left(\frac{X(\mathfrak{i}) - X(-\mathfrak{i})}{2} \right)^2 \right) - (\pi - 2). \end{aligned}$$

This representative of the discrete local field contains a clear discrete analogue of $\frac{1}{4}\|\nabla\varphi\|^2$ and an explicit additive constant that a priori comes from normal ordering. The constant is exactly what is needed to make the corresponding random variable (obtained by evaluation) have asymptotically zero expected value in the scaling limit: the two expected squares of gaussians are cancelled by the negative additive constant. This subtraction is *necessary* before it is meaningful to renormalize by the diverging prefactor δ^{-2} , which corresponds to the scaling dimension 2 of this local field.

Giving concrete representatives also for the other Virasoro primary fields (2.12) – (2.17) etc. would be possible, but the expressions become long and not as enlightening.

As the other explicit example, we choose the holomorphic stress tensor T of the CFT, which in terms of Heisenberg generators is $\frac{1}{2}j_{-1}j_{-1}\mathbb{1}$. The local field of the DGFF corresponding to the holomorphic stress tensor is

$$\frac{1}{2}J_{-1}J_{-1}I = \frac{1}{2}\bullet J_{-1}I \otimes J_{-1}I \bullet = \frac{1}{2}\circ J_{-1}J_{-1} \circ + \text{Null}.$$

A concrete representative is

$$\begin{aligned} \frac{1}{2}\circ J_{-1}J_{-1} \circ &= \frac{1}{2}J_{-1}J_{-1} - \frac{1}{2}\overline{J_{-1}J_{-1}} \\ &= -\frac{1}{2} \left(\frac{1}{2} \frac{X(1) - X(-1)}{2} - \frac{\mathfrak{i}}{2} \frac{X(\mathfrak{i}) - X(-\mathfrak{i})}{2} \right)^2. \end{aligned}$$

7 Scaling limits of local field correlations

The conformal field theory (CFT) of interest to us is known in physics as the massless free boson. The constructive quantum field theory approach to it amounts to studying

the continuum probabilistic model called the Gaussian Free Field (GFF) [Gaw99, KaMa13, WePo21]. Analogously to our conventions for the discrete GFF, we will consider the gradient of the GFF with either Dirichlet or Neumann boundary conditions in simply connected planar domains of general shape. Changing the domain and/or boundary conditions results in a different probabilistic object.¹² Nevertheless, our space of local fields of the CFT will always be the same: the full Fock space $\mathcal{F} \otimes \overline{\mathcal{F}}$ defined in Section 2.1.

Throughout this section, $\Omega \subset \mathbb{C}$ is taken to be a nonempty open simply connected proper subset of the complex plane.

Section 7.1 contains the definition of (the gradient of) the GFF, and a discussion of the sense in which this basic field has pointwise defined correlation functions. In Section 7.2, we then describe correlation functions of something akin to linear local fields of the GFF, which include, most notably, the holomorphic and antiholomorphic currents J and \bar{J} . From these correlation kernels of currents, a standard construction of a bosonic CFT is given in Section 7.3. More precisely, using the currents and operator product expansions as building blocks, we characterize and construct the n -point correlation functions of general Fock space fields, i.e., for a general domain Ω , boundary conditions (D or N), and number $n \in \mathbb{N}$ of points, we describe a map

$$\langle \cdots \rangle_{\Omega}^{\text{D/N}} : (\mathcal{F} \otimes \overline{\mathcal{F}})^{\otimes n} \rightarrow C^{\omega}(\text{Conf}_n(\Omega), \mathbb{C}),$$

assigning in a symmetric multilinear way to an n -tuple of Fock space field a complex-valued real-analytic correlation function defined on the **configuration space**

$$\text{Conf}_n(\Omega) := \Omega^n \setminus \bigcup_{1 \leq i < j \leq n} \{(z_1, \dots, z_n) \in \Omega^n \mid z_i = z_j\} \quad (7.1)$$

of (ordered) n -tuples of distinct points in Ω .

The following main result will be precisely stated and proven in Section 7.4. If the domain Ω is suitably approximated by discrete domains Ω_{δ} as $\delta \rightarrow 0$, then the correlations of local fields of the DGFF on Ω_{δ} , renormalized by the lattice mesh δ to the power of the scaling dimensions of the fields, converge to the corresponding correlation functions of Fock space fields in the bosonic CFT built from the GFF in Ω .

7.1 Gaussian Free Fields. Analogously to the discrete GFF, one would like to think of the continuum GFF in Ω as a random real-valued function φ on Ω with a centered Gaussian distribution and covariance $\mathbb{E}[\varphi(z)\varphi(w)]$ equal to 4π times a continuum Green's function $G_{\Omega}(z, w)$. However, due to the logarithmic singularity (3.12) of the Green's function

¹²In the continuum, conformal invariance relates the free fields with the same boundary conditions in different simply connected domains, but nevertheless defining and viewing these as different probabilistic models is a good perspective, since in the presence of nontrivial conformal moduli (e.g., for multiply connected domains or for domains with marked points) no such reduction by simple coordinate changes could be used. Moreover, Dirichlet and Neumann boundary condition free fields are rather evidently different probabilistic models.

when approaching $z = w$, one cannot actually assign pointwise values to the GFF; rather the GFF in Ω should be constructed as a random generalized function, i.e., a random distribution. Moreover, in the case of Neumann boundary conditions, there is an ambiguity of an additive constant (and in fact no unique choice of the Green's function). We thus seek to define only the gradient of the GFF

$$\nabla\varphi = \left(\frac{\partial\varphi}{\partial x}, \frac{\partial\varphi}{\partial y} \right)$$

as a two-component distribution-valued random variable. Gradients are not arbitrary two-component vector fields — they are necessarily curl-free — so a convenient equivalent perspective is to view the field φ itself as defined up to additive constants. Let us furthermore note already that it will soon be convenient to repackage the two components of the gradient in \mathbb{C} -linear combinations: the Wirtinger derivatives

$$\frac{\partial}{\partial z} = \frac{1}{2} \frac{\partial}{\partial x} - \frac{\mathbf{i}}{2} \frac{\partial}{\partial y} \quad \text{and} \quad \frac{\partial}{\partial \bar{z}} = \frac{1}{2} \frac{\partial}{\partial x} + \frac{\mathbf{i}}{2} \frac{\partial}{\partial y}$$

contain exactly the same information as the partial derivatives $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$. With suitable normalization constants, $\partial\varphi := \frac{\partial\varphi}{\partial z}$ and $\bar{\partial}\varphi := \frac{\partial\varphi}{\partial \bar{z}}$ will be the currents in our CFT.

As ordinary **test functions**, we use compactly supported C^∞ -smooth functions $g: \Omega \rightarrow \mathbb{R}$. The set of such test functions is denoted by $\mathcal{D}(\Omega)$, and is equipped with the topology of uniform convergence on compact subsets for all derivatives up to an arbitrary order. The space of **distributions** is denoted by $\mathcal{D}'(\Omega)$; it is the dual of the space of test functions, consisting of all continuous linear maps $\mathcal{D}(\Omega) \rightarrow \mathbb{R}$. The appropriate topology on $\mathcal{D}'(\Omega)$ is the weak-* topology. We denote the duality pairing of a distribution $\phi \in \mathcal{D}'(\Omega)$ with a test function $g \in \mathcal{D}(\Omega)$ by $\langle \phi, g \rangle \in \mathbb{R}$.

The constant distribution $1' \in \mathcal{D}'(\Omega)$ is defined by

$$\langle 1', g \rangle = \int_{\Omega} g(z) \, d^2z \quad \text{for } g \in \mathcal{D}(\Omega).$$

The subspace of **zero-average test functions** is

$$\mathcal{D}_{\nabla}(\Omega) := \text{Ker } 1' = \left\{ f \in \mathcal{D}(\Omega) \mid \int_{\Omega} f(z) \, d^2z = 0 \right\} \subset \mathcal{D}(\Omega).$$

Note that two distributions $\phi, \psi \in \mathcal{D}'(\Omega)$ differ by a multiple of the constant distribution $1'$ if and only if $\langle \phi, f \rangle = \langle \psi, f \rangle$ for all $f \in \mathcal{D}_{\nabla}(\Omega)$. The quotient space $\mathcal{D}'(\Omega) / \mathbb{R}1'$ of **distributions up to additive constants** can therefore be identified with the space $\mathcal{D}'_{\nabla}(\Omega)$ of continuous linear functionals on $\mathcal{D}_{\nabla}(\Omega)$.

We define the **GFF up to additive constants**, in a domain Ω , with Dirichlet (D) or Neumann (N) boundary conditions, as the $\mathcal{D}'_{\nabla}(\Omega)$ -valued random variable φ whose characteristic function is

$$\mathbb{E}_{\Omega}^{\text{D/N}}[e^{\mathbf{i}\langle \varphi, f \rangle}] = \exp \left(-2\pi \iint_{\Omega \times \Omega} f(z) G_{\Omega}^{\text{D/N}}(z, w) f(w) \, d^2w \, d^2z \right) \quad \text{for } f \in \mathcal{D}_{\nabla}(\Omega). \quad (7.2)$$

Note that even in the Neumann case, the double integral above does not depend on the choice of the Green's function. This characteristic function uniquely determines the law of φ , and an explicit construction of such a process φ can be found, e.g., in [BePo23].

Observe that for any $g \in \mathcal{D}(\Omega)$, the partial derivatives are zero-average, $\frac{\partial g}{\partial x}, \frac{\partial g}{\partial y} \in \mathcal{D}_\nabla(\Omega)$. In particular the distributional derivatives $\frac{\partial \varphi}{\partial x}, \frac{\partial \varphi}{\partial y}$ of the GFF up to additive constants are random ordinary distributions defined by

$$\left\langle \frac{\partial \varphi}{\partial x}, g \right\rangle = -\left\langle \varphi, \frac{\partial g}{\partial x} \right\rangle \quad \text{and} \quad \left\langle \frac{\partial \varphi}{\partial y}, g \right\rangle = -\left\langle \varphi, \frac{\partial g}{\partial y} \right\rangle \quad \text{for } g \in \mathcal{D}(\Omega),$$

and from the $\mathcal{D}'_\nabla(\Omega)$ -valued random variable φ we thus obtain the **gradient of the Gaussian free field (GFF)** $\nabla \varphi = \left(\frac{\partial \varphi}{\partial x}, \frac{\partial \varphi}{\partial y} \right)$ as a two-component $\mathcal{D}'(\Omega)$ -valued random variable.

From the characteristic function (7.2) one gets, in particular, the finite-dimensional marginals, i.e., the joint distributions of $(\langle \varphi, f_i \rangle)_{i=1}^n$ for any finite number n of zero-average test functions $f_1, \dots, f_n \in \mathcal{D}_\nabla(\Omega)$. These marginals are n -dimensional centered Gaussians with covariances

$$C_{ij} = 4\pi \iint_{\Omega \times \Omega} f_i(z) G_\Omega^{\text{D/N}}(z, w) f_j(w) \, d^2 w \, d^2 z, \quad i, j = 1, \dots, n.$$

The directional derivatives ∇^μ of (3.14) are convenient for concisely writing down the covariances of the components of $\nabla \varphi$. For any test functions $g_1, \dots, g_n \in \mathcal{D}(\Omega)$, the joint distribution of $(\langle \nabla^{\mu_i} \varphi, g_i \rangle)_{i=1}^n$ is, by construction, the centered n -dimensional Gaussian with covariances

$$\begin{aligned} C_{ij}^{\mu_i \mu_j} &= 4\pi \iint_{\Omega \times \Omega} (\nabla_i^{\mu_i} g_i)(z_i) G_\Omega^{\text{D/N}}(z_i, z_j) (\nabla_j^{\mu_j} g_j)(z_j) \, d^2 z_i \, d^2 z_j \\ &= 4\pi \iint_{\Omega \times \Omega} g_i(z_i) (\nabla_i^{\mu_i} \nabla_j^{\mu_j} G_\Omega^{\text{D/N}}(z_i, z_j)) g_j(z_j) \, d^2 z_i \, d^2 z_j \quad i, j = 1, \dots, n, \end{aligned}$$

where on the second line the double-derivative of the Green's function is in the distributional sense (delta-like terms do appear on the diagonal $\{z_i = z_j\}$, but if the supports of g_i, g_j are disjoint, then even the second line only involves ordinary integration of smooth functions). Note that even in the Neumann case, the double derivative of the Green's function on the last line above does not depend on the precise choice of the Green's function. As for any centered Gaussians, **Wick's formula** applies, and in this case gives

$$\begin{aligned} &\mathbb{E}_\Omega^{\text{D/N}} \left[\prod_{i=1}^n \langle \nabla^{\mu_i} \varphi, g_i \rangle \right] \tag{7.3} \\ &= \sum_{P \in \text{Pair}(n)} \prod_{\{i,j\} \in P} 4\pi \iint_{\Omega \times \Omega} g_i(z_i) (\nabla_i^{\mu_i} \nabla_j^{\mu_j} G_\Omega^{\text{D/N}}(z_i, z_j)) g_j(z_j) \, d^2 z_i \, d^2 z_j \\ &= \int \cdots \int_{\Omega^n} g_1(z_1) \cdots g_n(z_n) \left((4\pi)^{n/2} \sum_{P \in \text{Pair}(n)} \prod_{\{i,j\} \in P} \nabla_i^{\mu_i} \nabla_j^{\mu_j} G_\Omega^{\text{D/N}}(z_i, z_j) \right) \, d^2 z_1 \cdots d^2 z_n, \end{aligned}$$

where the sum is over the set $\text{Pair}(n)$ of pairings P of the index set $\{1, 2, \dots, n\}$. In this sense of integral kernels (distributional in general, but literal integration of smooth functions when the supports of g_1, \dots, g_n are disjoint), we may thus interpret the n -point **correlation function** of the components of the gradient GFF $\nabla\varphi$ at $(z_1, \dots, z_n) \in \text{Conf}_n(\Omega)$ as

$$“\mathbb{E}_\Omega^{\text{D/N}}[(\nabla^{\mu_1}\varphi)(z_1) \cdots (\nabla^{\mu_n}\varphi)(z_n)]” := (4\pi)^{n/2} \sum_{P \in \text{Pair}(n)} \prod_{\{i,j\} \in P} \nabla_i^{\mu_i} \nabla_j^{\mu_j} G_\Omega^{\text{D/N}}(z_i, z_j). \quad (7.4)$$

One can also view (7.4) as the limit of (7.3) with mollifiers taken as approximate delta-functions ($“g_i^{(\varepsilon)} \xrightarrow{\varepsilon \rightarrow 0} \delta(\cdot - z_i)”$) at distinct points $z_1, \dots, z_n \in \Omega$.

7.2 Current correlation kernels of the GFF. Before addressing correlation functions of general local fields in the CFT, we slightly generalize the above integral kernels for correlations of the gradient of the GFF, and we transform them to the most convenient form for the purposes of constructing the free boson CFT. Namely, consider derivatives of arbitrary order of $\varphi(z)$ — they serve as the natural counterparts of linear local fields in the CFT. It is convenient to change basis from the horizontal and vertical partial derivatives $\frac{\partial\varphi}{\partial x}$ and $\frac{\partial\varphi}{\partial y}$ to the holomorphic $\partial\varphi = \frac{1}{2}\frac{\partial\varphi}{\partial x} - \frac{i}{2}\frac{\partial\varphi}{\partial y}$ and antiholomorphic $\bar{\partial}\varphi = \frac{1}{2}\frac{\partial\varphi}{\partial x} + \frac{i}{2}\frac{\partial\varphi}{\partial y}$ Wirtinger derivatives, and similarly in higher order.

In the same sense as (7.4), the correlation functions of the fields $\partial^a \bar{\partial}^b \varphi$, for $a, b \in \mathbb{N}$ with $a + b > 0$, are defined by

$$\begin{aligned} & “\mathbb{E}_\Omega^{\text{D/N}}[(\partial^{a_1} \bar{\partial}^{b_1} \varphi)(z_1) \cdots (\partial^{a_n} \bar{\partial}^{b_n} \varphi)(z_n)]” \\ & := (4\pi)^{n/2} \frac{\partial^{a_1}}{\partial z_1^{a_1}} \frac{\partial^{b_1}}{\partial \bar{z}_1^{b_1}} \cdots \frac{\partial^{a_n}}{\partial z_n^{a_n}} \frac{\partial^{b_n}}{\partial \bar{z}_n^{b_n}} \sum_{P \in \text{Pair}(n)} \prod_{\{i,j\} \in P} G_\Omega^{\text{D/N}}(z_i, z_j); \end{aligned} \quad (7.5)$$

these correlation functions $\text{Conf}_n(\Omega) \rightarrow \mathbb{C}$ are the integral kernels for correlations of the distributional Wirtinger derivatives of the GFF φ mollified by test functions, analogously to (7.3). Observing the factorization of the Laplacian into Wirtinger derivatives, $\Delta = 4\frac{\partial}{\partial z}\frac{\partial}{\partial \bar{z}}$, and noting that (7.5) only contains sums of products of the Green’s functions, we see that the correlation (7.5) is zero if for some j we have both $a_j > 0$ and $b_j > 0$. This is the continuum counterpart of a null field property for $\Delta\varphi$ and its further derivatives. In particular, the only derivative-fields with interesting correlations are the holomorphic derivatives $\partial^a \varphi$ of order $a \in \mathbb{Z}_{>0}$ and the antiholomorphic derivatives $\bar{\partial}^b \varphi$ of order $b \in \mathbb{Z}_{>0}$.

We define the holomorphic and antiholomorphic **currents** by $J := i\partial\varphi$ and $\bar{J} := -i\bar{\partial}\varphi$, respectively. The integral kernel for m holomorphic and m' antiholomorphic currents will play a key role in the construction of the CFT, and we therefore use the notation

$$K_\Omega^{\text{D/N}} \in C^\omega(\text{Conf}_{m+m'}(\Omega), \mathbb{C}) \quad (7.6)$$

for it. This kernel has the informal interpretation

$$K_\Omega^{\text{D/N}}(z_1, \dots, z_m; w_1, \dots, w_{m'}) = “\mathbb{E}_\Omega^{\text{D/N}}[J(z_1) \cdots J(z_m) \bar{J}(w_1) \cdots \bar{J}(w_{m'})]” ;$$

the explicit formula would be notationally cumbersome, but it is naturally just $i^{m-m'}$ times (7.5) with $n = m + m'$, $(a_i, b_i) = (1, 0)$ for $i = 1, \dots, m$, and $(a_{m+j}, b_{m+j}) = (0, 1)$ for $j = 1, \dots, m'$, together with the relabeling $w_j := z_{m+j}$ for $j = 1, \dots, m'$. Derivatives of any nonzero order of the GFF can be written in terms of the currents: $\partial^a \varphi = -i \partial^{a-1} J$ and $\bar{\partial}^b \varphi = i \partial^{b-1} \bar{J}$, and the integral kernels for their correlations are obtained as further derivatives of $K_\Omega^{\text{D/N}}$.

The Green's functions (3.12) differentiated in both variables have the forms

$$\begin{aligned} \frac{\partial}{\partial z} \frac{\partial}{\partial w} G_\Omega^{\text{D/N}}(z, w) &= \frac{-1/4\pi}{(z-w)^2} + \frac{\partial}{\partial z} \frac{\partial}{\partial w} g_\Omega^{\text{D/N}}(z, w), & \frac{\partial}{\partial z} \frac{\partial}{\partial \bar{w}} G_\Omega^{\text{D/N}}(z, w) &= \frac{\partial}{\partial z} \frac{\partial}{\partial \bar{w}} g_\Omega^{\text{D/N}}(z, w), \\ \frac{\partial}{\partial \bar{z}} \frac{\partial}{\partial w} G_\Omega^{\text{D/N}}(z, w) &= \frac{-1/4\pi}{(\bar{z}-\bar{w})^2} + \frac{\partial}{\partial \bar{z}} \frac{\partial}{\partial w} g_\Omega^{\text{D/N}}(z, w), & \frac{\partial}{\partial \bar{z}} \frac{\partial}{\partial \bar{w}} G_\Omega^{\text{D/N}}(z, w) &= \frac{\partial}{\partial \bar{z}} \frac{\partial}{\partial \bar{w}} g_\Omega^{\text{D/N}}(z, w) \end{aligned}$$

where $g_\Omega^{\text{D/N}}: \Omega \times \Omega \rightarrow \mathbb{R}$ is real-analytic, and harmonic in both variables separately. By virtue of the factorization $\Delta = 4 \frac{\partial}{\partial z} \frac{\partial}{\partial \bar{z}}$ of the Laplacian, the double Wirtinger derivatives of $g_\Omega^{\text{D/N}}$ are then real-analytic functions $\Omega \times \Omega \rightarrow \mathbb{C}$ which are holomorphic or antiholomorphic separately in the two variables. We obtain, in particular, the following explicit formula for the two-point function of the holomorphic current

$$“\mathbb{E}_\Omega^{\text{D/N}}[\partial\varphi(z_1)\partial\varphi(z_2)]” = K_\Omega^{\text{D/N}}(z_1, z_2;) = \frac{1}{(z_1 - z_2)^2} - 4\pi \underbrace{\frac{\partial}{\partial z_1} \frac{\partial}{\partial z_2} g_\Omega^{\text{D/N}}(z_1, z_2)}_{\text{regular as } z_1 \rightarrow z_2}.$$

This shows that $z_1 \mapsto K_\Omega^{\text{D/N}}(z_1, z_2;) - \frac{1}{(z_1 - z_2)^2}$ is holomorphic in the whole Ω . Similarly, $w_1 \mapsto K_\Omega^{\text{D/N}}(; w_1, w_2) - \frac{1}{(\bar{w}_1 - \bar{w}_2)^2}$ is antiholomorphic in the whole Ω . In the mixed two-point functions, there are no singularities: just by themselves, $z \mapsto K_\Omega^{\text{D/N}}(z; w)$ is holomorphic and $w \mapsto K_\Omega^{\text{D/N}}(z; w)$ is antiholomorphic in the whole Ω . Furthermore, since multi-point correlations are obtained by Wick's formula, $K_\Omega^{\text{D/N}}(z_1, \dots, z_m; w_1, \dots, w_{m'})$ is meromorphic in each of the variables z_i with poles of order 2 at $z_1, \dots, z_{i-1}, z_{i+1}, \dots, z_m$ only, and antimeromorphic in each of the variables w_i with (anti)poles of order 2 at $w_1, \dots, w_{i-1}, w_{i+1}, \dots, w_{m'}$ only.

7.3 Correlation functions of CFT fields from current OPEs. One possible way to construct higher degree fields would be by forming normally ordered products of the linear fields, see, e.g., [KaMa13, Section 1.2]. Making sense of these as distribution-valued random variables and defining pointwise correlation functions as integral kernels is genuinely more complicated than for the linear fields, see [KaMa13, Section 2.2]. There is, however, an even more serious drawback: the (probabilistic) normal ordering is not intrinsic to the fields themselves. Indeed, the probabilistic normal ordering depends on the Gaussian measure (the law of φ), which depends on the domain Ω and boundary conditions, so, from a CFT point of view, normal ordering is *not* the right definition of an abstract higher degree field whose correlation functions can be evaluated in all domains and with any reasonable boundary conditions.

The intrinsic (domain-agnostic and boundary-condition-agnostic) way to construct higher degree local fields is by recursively extracting operator-product expansion (OPE) coefficients from lower order fields, as discussed, e.g., in [KaMa13, Section 3]. Starting from the currents $J(z)$ and $\bar{J}(w)$, general Fock space fields are generated via such OPEs. The holomorphic current $J(z)$ has a purely meromorphic OPE with any Fock space field, so an easy way to extract the OPE coefficients is by suitable weighted contour integrals. Similarly the anti-holomorphic current $\bar{J}(w)$ has a purely antimeromorphic OPE with any Fock space field, and again OPE coefficients can be extracted by suitable contour integrals. This is, roughly speaking, the route we take below to define arbitrary n -point correlation functions

$$\begin{aligned} \langle \cdots \rangle_{\Omega}^{\mathbb{D}/\mathbb{N}} : (\mathcal{F} \otimes \overline{\mathcal{F}})^{\otimes n} &\rightarrow C^{\omega}(\text{Conf}_n(\Omega), \mathbb{C}). \\ F_1 \otimes \cdots \otimes F_n &\mapsto \left((z_1, \dots, z_n) \mapsto \left\langle F_1(z_1) \cdots F_n(z_n) \right\rangle_{\Omega}^{\mathbb{D}/\mathbb{N}} \right) \end{aligned} \quad (7.7)$$

for a bosonic CFT with the Fock space $\mathcal{F} \otimes \overline{\mathcal{F}}$ as its space of local fields.

Proposition 7.1. *Consider the domain Ω and boundary conditions \mathbb{D}/\mathbb{N} fixed. Then there exists a unique collection of linear assignments of type (7.7), indexed by $n \in \mathbb{N}$, such that the following conditions hold:*

(BOS) *For any permutation $\sigma \in \mathfrak{S}_n$, we have*

$$\left\langle F_1(z_1) \cdots F_n(z_n) \right\rangle_{\Omega}^{\mathbb{D}/\mathbb{N}} = \left\langle F_{\sigma(1)}(z_{\sigma(1)}) \cdots F_{\sigma(n)}(z_{\sigma(n)}) \right\rangle_{\Omega}^{\mathbb{D}/\mathbb{N}}.$$

(CUR) *For $(z_1, \dots, z_m, w_1, \dots, w_{m'}) \in \text{Conf}_{m+m'}(\Omega)$, we have*

$$\begin{aligned} &\left\langle (j_{-1}\mathbb{1})(z_1) \cdots (j_{-1}\mathbb{1})(z_m) (\bar{j}_{-1}\mathbb{1})(w_1) \cdots (\bar{j}_{-1}\mathbb{1})(w_{m'}) \right\rangle_{\Omega}^{\mathbb{D}/\mathbb{N}} \\ &= K_{\Omega}^{\mathbb{D}/\mathbb{N}}(z_1, \dots, z_m; w_1, \dots, w_{m'}) \end{aligned}$$

where the right-hand side is the integral kernel (7.6) of GFF current correlations in Ω with the chosen boundary conditions \mathbb{D}/\mathbb{N} .

(J-OPE) *For any $F_1, \dots, F_n \in \mathcal{F} \otimes \overline{\mathcal{F}}$ and any $(z_1, \dots, z_n) \in \text{Conf}_n(\Omega)$, we have the following Laurent series expansion of the $n+1$ -point function*

$$\begin{aligned} &\left\langle F_1(z_1) \cdots F_j(z_j) \cdots F_n(z_n) (j_{-1}\mathbb{1})(z) \right\rangle_{\Omega}^{\mathbb{D}/\mathbb{N}} \\ &= \sum_{k \in \mathbb{Z}} (z - z_j)^{-1-k} \left\langle F_1(z_1) \cdots (j_k F_j)(z_j) \cdots F_n(z_n) \right\rangle_{\Omega}^{\mathbb{D}/\mathbb{N}}, \end{aligned} \quad (7.8)$$

in the region $0 < |z - z_j| < \min \{ \min_{i \neq j} |z_j - z_i|, \text{dist}(z_j, \partial\Omega) \}$.

($\bar{\text{J}}$ -OPE) For any $F_1, \dots, F_n \in \mathcal{F} \otimes \overline{\mathcal{F}}$ and any $(z_1, \dots, z_n) \in \text{Conf}_n(\Omega)$, we have the following anti-Laurent series expansion of the $n+1$ -point function

$$\begin{aligned} & \left\langle F_1(z_1) \cdots F_j(z_j) \cdots F_n(z_n) (\bar{\text{J}}_{-1} \mathbb{1})(w) \right\rangle_{\Omega}^{\text{D/N}} \\ &= \sum_{k \in \mathbb{Z}} (\bar{w} - \bar{z}_j)^{-1-k} \left\langle F_1(z_1) \cdots (\bar{\text{J}}_k F_j)(z_j) \cdots F_n(z_n) \right\rangle_{\Omega}^{\text{D/N}}, \end{aligned} \quad (7.9)$$

in the region $0 < |w - z_j| < \min \{ \min_{i \neq j} |z_j - z_i|, \text{dist}(z_j, \partial\Omega) \}$.

Proof. The uniqueness of such $\langle \cdots \rangle_{\Omega}^{\text{D/N}}$ is straightforward by a recursive argument: (CUR) provides the base case, and with repeated coefficient extraction using (J-OPE) and ($\bar{\text{J}}$ -OPE) one obtains expressions for the general correlation functions. The notation introduced in the existence proof will make the explicit expression clear.

To prove the existence of $\langle \cdots \rangle_{\Omega}^{\text{D/N}}$ as in (7.7), we start by constructing correlation functions for n formal expressions

$$F_i := \text{j}_{k_{i;m_i}} \cdots \text{j}_{k_{i;2}} \text{j}_{k_{i;1}} \bar{\text{j}}_{k'_{i;m'_i}} \cdots \bar{\text{j}}_{k'_{i;2}} \bar{\text{j}}_{k'_{i;1}} \mathbb{1}, \quad (7.10)$$

$$\text{with } m_i, m'_i \in \mathbb{Z}_{\geq 0}, \quad k_{i;1}, \dots, k_{i;m_i}, k'_{i;1}, \dots, k'_{i;m'_i} \in \mathbb{Z}.$$

indexed by $i = 1, \dots, n$. To then finally define (7.7), we extend from the formal expressions n -multilinearly, and check that the correlation functions become well-defined, i.e., in each tensorand, they factor through a quotient which defines the Fock space $\mathcal{F} \otimes \overline{\mathcal{F}}$. The quotient has to account for both the Heisenberg commutation relations and the quotient construction (2.4) of the Fock representation in both chiralities.

The n -point correlation function

$$\left\langle F_1(z_1) \cdots F_n(z_n) \right\rangle_{\Omega}^{\text{D/N}}$$

of formal expressions (7.10) at $(z_1, \dots, z_n) \in \text{Conf}_n(\Omega)$ is constructed starting from the kernel (7.6) for current correlations

$$K_{\Omega}^{\text{D/N}} \left(\underbrace{\zeta_{1;1}, \dots, \zeta_{1;m_1}, \dots, \zeta_{n;1}, \dots, \zeta_{n;m_n}}_{\substack{n \text{ groups of variables} \\ m_1 \text{ variables} \quad m_n \text{ variables}}}, \underbrace{\xi_{1;1}, \dots, \xi_{1;m'_1}, \dots, \xi_{n;1}, \dots, \xi_{n;m'_n}}_{\substack{n \text{ groups of variables} \\ m'_1 \text{ variables} \quad m'_n \text{ variables}}} \right)$$

with $\sum_i m_i$ holomorphic currents and $\sum_i m'_i$ antiholomorphic currents, by integrating each of the variables $\zeta_{i;s_i}$ and $\xi_{i;t_i}$ around z_i along suitable radially ordered contours with the appropriate weight depending on $k_{i;s_i}$ or $k'_{i;t_i}$. Specifically, for each $i = 1, \dots, n$, choose radii

$$0 < r'_{i;1} < r'_{i;2} < \cdots < r'_{i;m'_i} < r_{i;1} < r_{i;2} < \cdots < r_{i;m_i} < R,$$

where R is small enough so that the disks $\mathbf{D}_R(z_i)$, $i = 1, \dots, n$, are disjoint and contained in the domain Ω . Then the defining formula for the correlation function of the expressions (7.10) is

$$\begin{aligned} & \left\langle F_1(z_1) \cdots F_n(z_n) \right\rangle_{\Omega}^{\mathbf{D}/\mathbf{N}} \\ &:= \oint \cdots \oint K_{\Omega}^{\mathbf{D}/\mathbf{N}}(\zeta_{1;1}, \dots; \dots, \xi_{n;m'_n}) \prod_{i=1}^n \left(\prod_{s=1}^{m_i} (\zeta_{i;s} - z_i)^{k_{i;s}} \frac{d\zeta_{i;s}}{2\pi i} \prod_{t=1}^{m'_i} (\bar{\xi}_{i;s} - \bar{z}_i)^{k'_{i;t}} \frac{d\bar{\xi}_{i;t}}{(-2\pi i)} \right), \end{aligned} \quad (7.11)$$

where $\zeta_{i;s_i}$ is integrated over the positively oriented circle $\partial\mathbf{D}_{r_{i;s_i}}(z_i)$ and $\xi_{i;t_i}$ over the positively oriented circle $\partial\mathbf{D}_{r'_{i;t_i}}(z_i)$. By holomorphicity and antiholomorphicity of the correlation kernel $K_{\Omega}^{\mathbf{D}/\mathbf{N}}$ of the currents \mathbf{J} and $\bar{\mathbf{J}}$, the value of (7.11) only depends on the homotopy class of these integrations, and in particular any choice of radii with the prescribed ordering yields the same result. Moreover, since there are no singularities at $\zeta = \xi$ of \mathbf{J} - $\bar{\mathbf{J}}$ -correlations $K_{\Omega}^{\mathbf{D}/\mathbf{N}}(\dots, \zeta, \dots; \dots, \xi, \dots)$ the homotopies can be used to move holomorphic current integrations past antiholomorphic current integrations, so the radial ordering convention can be relaxed to separately requiring

$$0 < r'_{i;1} < r'_{i;2} < \cdots < r'_{i;m'_i} < R \quad \text{and} \quad 0 < r_{i;1} < r_{i;2} < \cdots < r_{i;m_i} < R.$$

We must then check that the multi-linear extension of (7.11) gives rise to well-defined correlation functions

$$\langle \cdots \rangle_{\Omega}^{\mathbf{D}/\mathbf{N}} : (\mathcal{F} \otimes \overline{\mathcal{F}})^{\otimes n} \rightarrow C^{\omega}(\text{Conf}_n(\Omega), \mathbb{C}).$$

The first thing to check is that the correlation functions (7.11) of the formal expressions (7.10) satisfy equations corresponding to the commutation relations (2.2) of the two commuting Heisenberg algebras in each of the n tensorands. Concretely, for

$$G_i := \mathbf{j}_k \mathbf{j}_{\ell} F_i \quad \text{and} \quad H_i := \mathbf{j}_{\ell} \mathbf{j}_k F_i,$$

by standard satellite integral arguments and the explicit poles of the current correlation functions (7.6), one can prove equalities of correlation functions (7.11) of the form

$$\left\langle \cdots G_i(z_i) \cdots \right\rangle_{\Omega}^{\mathbf{D}/\mathbf{N}} - \left\langle \cdots H_i(z_i) \cdots \right\rangle_{\Omega}^{\mathbf{D}/\mathbf{N}} = k \delta_{k+\ell} \left\langle \cdots F_i(z_i) \cdots \right\rangle_{\Omega}^{\mathbf{D}/\mathbf{N}}.$$

Similar equations hold relating correlations of $\bar{\mathbf{j}}_k \bar{\mathbf{j}}_{\ell} F_i$, $\bar{\mathbf{j}}_{\ell} \bar{\mathbf{j}}_k F_i$ and F_i . From such commutation equations it follows that the correlation functions of formal linear combinations of symbols (7.10) at least factor through $\text{Hei} \otimes \overline{\text{Hei}}$ in each tensorand, and (7.11) gives rise to a map

$$(\text{Hei} \otimes \overline{\text{Hei}})^{\otimes n} \rightarrow C^{\omega}(\text{Conf}_n(\Omega), \mathbb{C}).$$

For well-definedness on the Fock space $\mathcal{F} \otimes \overline{\mathcal{F}}$, it therefore remains to check equations corresponding to the quotient construction (2.4) of the Fock representations in the two chiralities.

If $F_i = j_k \mathbb{1}$ for some i , then there is exactly one integration around z_i in (7.11), with $\zeta_{i;1}$ as the integration variable and $(\zeta_{i;1} - z_i)^k$ as the weight. But the correlation kernel (7.6) of the currents is holomorphic in $\zeta_{i;1}$, so this integration (alone) picks the coefficient of $(\zeta_{i;1} - z_i)^{-1-k}$ in the Taylor series expansion of the holomorphic function $\zeta_{i;1} \mapsto K_\Omega^{\text{D/N}}(\zeta_{1;1}, \dots; \dots, \xi_{n;m'_n})$ at z_i , which is zero for $k \geq 0$. This shows factorization through $\mathcal{F} = \text{Hei}/(j_k : k \geq 0)$ in the holomorphic chirality of the i th tensorand. For the factorization

$$(\mathcal{F} \otimes \overline{\mathcal{F}})^{\otimes n} \rightarrow C^\omega(\text{Conf}_n(\Omega), \mathbb{C})$$

through the full Fock space in each factor, one repeats an entirely similar argument with the antiholomorphic current.

We thus obtain a collection of well-defined maps (7.7) indexed by n . It remains to check the asserted properties (BOS), (CUR), (J-OPE), and ($\bar{\text{J}}$ -OPE).

Symmetry (BOS) is evident from the symmetry of the kernel $K_\Omega^{\text{D/N}}$ and of the integral (7.11).

The property (CUR) is evident by choosing $m_1 = \dots = m_m = 1$, $m'_1 = \dots = m'_{m'} = 1$, and $k_{1;1} = \dots = k_{m;1} = -1$, $k'_{1;1} = \dots = k'_{m';1} = -1$, and noticing that the current kernel $K_\Omega^{\text{D/N}}(z_1, \dots, z_m; w_1, \dots, w_{m'})$ is then obtained as the residue of the defining integrals (7.11).

The proofs of (J-OPE) and ($\bar{\text{J}}$ -OPE) are similar, so let us only comment on the former. Consider a correlation function of the form (7.11) with $n+1$ fields of the form (7.10), where the $(n+1)$ st one is taken to be simply $F_{n+1} = j_{-1} \mathbb{1}$. This correlation function is meromorphic in the variable z_{n+1} (as shown by induction on the total degree $\sum_i (m_i + m'_i)$). Its Laurent series expansion at z_i is of the form

$$\left\langle F_1(z_1) \cdots F_n(z_n) (j_{-1} \mathbb{1})(z_{n+1}) \right\rangle_\Omega^{\text{D/N}} = \sum_{p=-p_0}^{\infty} (z_{n+1} - z_i)^p C_p(z_1, \dots, z_n).$$

The Laurent series coefficients admit contour integral expressions,

$$C_p(z_1, \dots, z_n) = \frac{1}{2\pi i} \oint_{\partial \mathbf{D}_R(z_i)} \left\langle F_1(z_1) \cdots F_n(z_n) (j_{-1} \mathbb{1})(z_{n+1}) \right\rangle_\Omega^{\text{D/N}} (z_{n+1} - z_i)^{-1-p} dz_{n+1}.$$

Writing out the definition of the correlation function and extracting the residue in its single integration around z_{n+1} as in (CUR), we obtain an expression for the coefficient C_p which coincides, by the definition (7.11), with the n -point correlation function

$$C_p(z_1, \dots, z_n) = \left\langle F_1(z_1) \cdots (j_{-k} F_i)(z_i) \cdots F_n(z_n) \right\rangle_\Omega^{\text{D/N}} \quad \text{with } k = -1 - p$$

involving the higher degree field $j_{-k} F_i$. □

Finally, let us make a few further remarks on the interpretations of the fields in the correlation functions (7.11).

Identity as a Fock space field: If for some i we have $F_i = \mathbb{1}$, i.e., $m_i = 0 = m'_i$, then there are no integrations around z_i and no z_i -dependent weights in (7.11). The resulting correlation function does not depend on z_i , and its value is the same as the correlation obtained by omitting $F_i(z_i)$. Such an F_i is interpreted as the “identity field”.

Holomorphic and antiholomorphic currents and their derivatives as Fock space fields: The fields $j_{-1}\mathbb{1}$ and $\bar{j}_{-1}\mathbb{1}$ should be interpreted as the holomorphic and antiholomorphic current fields in the bosonic CFT. Correlation functions which only involve these are exactly the probabilistic GFF current correlation kernels,

$$\begin{aligned} \left\langle (j_{-1}\mathbb{1})(z_1) \cdots (\bar{j}_{-1}\mathbb{1})(w_{m'}) \right\rangle_{\Omega}^{\text{D/N}} &= K_{\Omega}^{\text{D/N}}(z_1, \dots, z_m; w_1, \dots, w_{m'}) \\ &= “\mathbb{E}_{\Omega}^{\text{D/N}}[J(z_1) \cdots J(z_m) \bar{J}(w_1) \cdots \bar{J}(w_{m'})]”, \end{aligned}$$

according to the property (CUR). The CFT fields $j_{-1}\mathbb{1}$ and $\bar{j}_{-1}\mathbb{1}$, however, have correlation functions not only among themselves, but more generally with any other Fock space fields. In view of this generalization, we denote, from here on,

$$J := j_{-1}\mathbb{1} \in \mathcal{F} \otimes \overline{\mathcal{F}} \quad \text{and} \quad \bar{J} := \bar{j}_{-1}\mathbb{1} \in \mathcal{F} \otimes \overline{\mathcal{F}}.$$

Similarly, for $p \geq 0$, the Fock space fields $j_{-1-p}\mathbb{1}$ and $\bar{j}_{-1-p}\mathbb{1}$ have the interpretations as the higher order derivative linear fields $\frac{1}{p!}\partial^p J$ and $\frac{1}{p!}\bar{\partial}^p \bar{J}$, respectively. Indeed, as in the proof of (CUR) above, the construction of the correlation functions of these involves picking a residue which recovers the corresponding derivative of the GFF correlation kernel.

7.4 Renormalized limits of correlations of lattice local fields. In the scaling limit statements, we let the lattice mesh $\delta > 0$ tend to zero. Continuum domains $\Omega \subset \mathbb{C}$ are approximated by discrete domains $\Omega_{\delta} \subset \delta\mathbb{Z}^2$ on the δ -mesh square grid. We will show that expected values $\mathbb{E}_{\Omega_{\delta}}^{\text{D/N}}[\cdots]$ of products of lattice local fields w.r.t. the DGFF measures of Section 4.1 converge, when suitably renormalized, to the CFT correlation functions $\langle \cdots \rangle_{\Omega}^{\text{D/N}}$ given by Proposition 7.1.

The correspondence between CFT fields and lattice local fields, used in the expected values and CFT correlations, respectively, is provided by the isomorphism

$$\mathcal{F}_{\nabla} \cong \mathcal{F} \otimes \overline{\mathcal{F}}$$

of Theorem 6.4. We (ab)use this one-to-one correspondence to interpret a given lattice local field $F = P + \text{Null} \in \mathcal{F}_{\nabla}$ as a Fock space field $F \in \mathcal{F} \otimes \overline{\mathcal{F}}$. The former has well-defined correlations $\mathbb{E}_{\Omega_{\delta}}^{\text{D/N}}[\text{ev}_{\mathbb{Z}^{\delta}}^{\Omega_{\delta}}(P) \cdots]$ independent of the chosen representative P in cases relevant to the scaling limit, and the latter has meaningful CFT correlations $\langle F(z) \cdots \rangle_{\Omega}^{\text{D/N}}$. To facilitate stating the scaling limit as locally uniform convergence on the configuration spaces $\text{Conf}_n(\Omega)$, we extend the definition of evaluation of field polynomials P to points $z \in \Omega$ in

the continuum domain, by setting $\text{ev}_z^{\Omega_\delta}(P) := \text{ev}_{\mathbf{z}}^{\Omega_\delta}(P)$ with $\mathbf{z} \in \Omega_\delta$ a closest lattice point to $z \in \Omega$. The lattice local fields need to be renormalized in the scaling limit according to their $L_0 + \bar{L}_0$ eigenvalue, so we phrase the statement in terms of homogeneous fields $F \in (\mathcal{F}_\nabla)_{\Delta, \bar{\Delta}}$ defined in (4.14). Note that by virtue of the isomorphism $\mathcal{F}_\nabla \cong \mathcal{F} \otimes \bar{\mathcal{F}}$, any local field can be written as a finite linear combination of such homogeneous local fields.

Theorem 7.2. *Fix boundary conditions D/N and an approximation $(\Omega_\delta)_{\delta>0}$ of a domain Ω in the Carathéodory sense by discrete domains $\Omega_\delta \subset \delta\mathbb{Z}^2$. Let $F_1, \dots, F_n \in \mathcal{F}_\nabla \cong \mathcal{F} \otimes \bar{\mathcal{F}}$ be local fields satisfying $F_i \in (\mathcal{F}_\nabla)_{\Delta_i, \bar{\Delta}_i}$, and fix representative field polynomials $P_i \in \mathcal{P}_\nabla$ so that $F_i = P_i + \text{Null}$. Then, as $\delta \rightarrow 0$, we have*

$$\delta^{-\sum_j(\Delta_j + \bar{\Delta}_j)} \mathbb{E}_{\Omega_\delta}^{D/N} \left[\text{ev}_{z_1}^{\Omega_\delta}(P_1) \cdots \text{ev}_{z_n}^{\Omega_\delta}(P_n) \right] \longrightarrow \left\langle F_1(z_1) \cdots F_n(z_n) \right\rangle_\Omega^{D/N} \quad (7.12)$$

uniformly for (z_1, \dots, z_n) in compact subsets of $\text{Conf}_n(\Omega)$.

The proof strategy should already naturally suggest itself. If F_i are basis vectors of the Fock space of fields, then the right hand side of (7.12) is explicitly given by the multiple contour integral (7.11). The left hand side expected value is given by analogous discrete contour integrations, with discrete monomial weights. The discrete integration contours can be chosen to approximate rectangular continuum contours. The DGFF expected values inside the discrete integrations involve only discrete currents, so they are simply written in terms of discrete double derivatives of discrete Green's functions, which converge to their continuum counterparts, and in the limit they reproduce the integral kernel $K_\Omega^{D/N}$ of GFF currents. The discrete monomial weights also converge to the monomial weights in (7.11), when an appropriate scaling by lattice mesh is taken into account. Together, the scaling factors from the discrete monomials account for the renormalization factor $\delta^{-\sum_j(\Delta_j + \bar{\Delta}_j)}$. Finally, the discrete integrations can be viewed as Riemann sum approximations, so the locally uniform convergence of the integrands implies the locally uniform convergence (7.12). The proof below provides concrete details following this natural strategy.

Proof. By linearity of expected values, it suffices to prove the statement for basis vectors of the joint eigenspaces $(\mathcal{F}_\nabla)_{\Delta_i, \bar{\Delta}_i}$ of L_0 and \bar{L}_0 with eigenvalues Δ_i and $\bar{\Delta}_i$. Fields of the form $F_i = j_{-k_i; m_i} \cdots j_{-k_{i,1}; 1} j_{-k'_{i,1}; 1} \cdots j_{-k'_{i, m'_i}; 1} \mathbb{1}$, with $\sum_{s=1}^{m_i} k_{i,s} = \Delta_i$ and $\sum_{s=1}^{m'_i} k'_{i,s} = \bar{\Delta}_i$ form the convenient basis for us. The essence of the proof becomes clear already from the case $F_i = j_{-k_i; m_i} \cdots j_{-k_{i,1}; 1} \mathbb{1}$, for $i = 1, \dots, n$ so for notational simplicity let us assume this.

The CFT correlation on the right-hand side of the assertion is defined by (7.11), and for

fields of the form $F_i = j_{-k_i; m_i} \cdots j_{-k_i; 1} \mathbb{1}$ it simplifies to

$$\begin{aligned}
& \left\langle F_1(z_1) \cdots F_n(z_n) \right\rangle_{\Omega}^{\text{D/N}} \\
&:= \oint \cdots \oint K_{\Omega}^{\text{D/N}}(\zeta_{1;1}, \dots, \zeta_{n; m_n};) \prod_{i=1}^n \prod_{s=1}^{m_i} (\zeta_{i;s} - z_i)^{-k_{i;s}} \frac{d\zeta_{i;s}}{2\pi i} \\
&= \sum_P \prod_{\{(i,s), (j,t)\} \in P} 4\pi \oint \oint \frac{\partial^2 G_{\Omega}^{\text{D/N}}(\zeta_{i;s}, \zeta_{j;t})}{\partial \zeta_{i;s} \partial \zeta_{j;t}} (\zeta_{i;s} - z_i)^{-k_{i;s}} (\zeta_{j;t} - z_j)^{-k_{j;t}} \frac{d\zeta_{i;s}}{2\pi} \frac{d\zeta_{j;t}}{2\pi},
\end{aligned}$$

where P is summed over the pairings of the set $\{(i, s) \mid 1 \leq i \leq n, 1 \leq s \leq m_i\}$ and each $\zeta_{i;s}$ is integrated in positive direction around z_i along radially ordered circular contours as described below (7.11). We can, however, deform the integrations to radially-ordered concentric square contours $\Gamma_{i;s}$ contained in Ω , centered at z_i — see Figure 7.1. We fix a choice of such contours for the rest of the proof. We will not write full details about the local uniformity of the convergence below, but locally uniform error estimates are routine to obtain if one ensures here that the chosen $\Gamma_{i;s}$ is only changed slightly for nearby values of z_i .

To write down the expected value on the left-hand side of the assertion for a given small mesh size $\delta > 0$, we choose the unit mesh corner contours $\gamma_{i;s}^{(\delta)}$ that are closest from inside to the blown-up and shifted continuum square contours $\delta^{-1}(\Gamma_{i;s} - z_i)$ such that each of their sides has an even number of corner steps — see Figure 7.1. Then, for small $\delta > 0$, the expected value involving evaluations of specific representatives P_i of F_i can by discrete contour deformation (without changing the value of the expectation) be written in the form

$$\begin{aligned}
& \mathbb{E}_{\Omega_{\delta}}^{\text{D/N}} \left[\text{ev}_{z_1}^{\Omega_{\delta}}(P_1) \cdots \text{ev}_{z_n}^{\Omega_{\delta}}(P_n) \right] \tag{7.13} \\
&= \oint^{\#} \cdots \oint^{\#} \mathbb{E}_{\Omega_{\delta}}^{\text{D/N}} \left[\prod_{i=1}^n \prod_{s=1}^{m_i} J(\mathbf{z}_i^{\delta} + \delta \mathbf{u}_m^{i;s}) \right] \prod_{i=1}^n \prod_{s=1}^{m_i} (\mathbf{u}_{\diamond}^{i;s})^{[-k_{i;s}]} \frac{d^{\#} \mathbf{u}^{i;s}}{2\pi i} \\
&= \sum_P \prod_{\{(i,s), (j,t)\} \in P} 4\pi \oint_{\gamma_{i;s}^{(\delta)}}^{\#} \oint_{\gamma_{j;t}^{(\delta)}}^{\#} (\partial_{\#}^{\bullet})_{\mathbf{u}^{i;s}} (\partial_{\#}^{\bullet})_{\mathbf{u}^{j;t}} G_{\Omega_{\delta}}^{\text{D/N}}(\mathbf{z}_i^{\delta} + \delta \mathbf{u}_m^{i;s}, \mathbf{z}_j^{\delta} + \delta \mathbf{u}_m^{j;t}) \\
&\quad (\mathbf{u}_{\diamond}^{i;s})^{[-k_{i;s}]} (\mathbf{u}_{\diamond}^{j;t})^{[-k_{j;t}]} \frac{d^{\#} \mathbf{u}^{i;s}}{2\pi} \frac{d^{\#} \mathbf{u}^{j;t}}{2\pi},
\end{aligned}$$

where $\mathbf{z}_i^{\delta} \in \Omega_{\delta}$ is the closest vertex to $z_i \in \Omega$.

Both the CFT correlation and the discrete expectation were written above as sums of products of double-integral factors, so it suffices to handle the convergence of each such factor. More precisely, remembering the form of the renormalizing prefactor

$$\delta^{-\sum_i \Delta_i} = \delta^{-\sum_i \sum_{s=1}^{m_i} k_{i;s}},$$

it suffices to prove, with $k = k_{i;s}$ and $\ell = k_{j;t}$, that as $\delta \rightarrow 0$ we have

$$\delta^{-k-\ell} \oint_{\gamma_{i;s}^{(\delta)}}^{\#} \oint_{\gamma_{j;t}^{(\delta)}}^{\#} (\partial_{\#}^{\bullet})_{\mathbf{u}} (\partial_{\#}^{\bullet})_{\mathbf{v}} G_{\Omega_{\delta}}^{\mathbf{D}/\mathbf{N}}(\mathbf{z}_i^{\delta} + \delta \mathbf{u}_m, \mathbf{z}_j^{\delta} + \delta \mathbf{v}_m) \mathbf{u}_{\diamond}^{[-k]} \mathbf{v}_{\diamond}^{[-\ell]} d^{\#} \mathbf{u} d^{\#} \mathbf{v} \quad (7.14)$$

$$\longrightarrow \oint_{\Gamma_{i;s}} \oint_{\Gamma_{j;t}} \frac{\partial^2 G_{\Omega}^{\mathbf{D}/\mathbf{N}}(\zeta, \xi)}{\partial \zeta \partial \xi} (\zeta - z_i)^{-k} (\xi - z_j)^{-\ell} d\zeta d\xi, \quad (7.15)$$

with error estimates of the form that ensure local uniformity of the convergence.

It is natural to split the discrete and continuous integrals in (7.14) and (7.15) over rectangle contours to the four sides of the rectangles; with 16 terms in total from the double integrations. Let us for concreteness focus on the term when both integration variables are on the bottom side of their corresponding rectangles.

The remaining slight complication in interpreting (7.14) as a Riemann sum approximation is related to the combinatorics of the corner contour integrations and the primal lattice discrete Wirtinger derivatives $\partial_{\#}^{\bullet}$ of (3.4). Namely, every other step of a single corner contour integral contributes with a horizontal discrete derivative of the Green's function, whereas the subsequent step contributes with a vertical discrete derivative, and one of these comes with a real and the other with an imaginary prefactor. For this reason, in the discrete integrations with even numbers of steps, we group together the four terms of two consecutive steps in each of the two variables, and interpret the sum as δ^{-2} times a double integral of a function $Q_{k,\ell}^{(\delta)}$, to be described below, which is constant on $\delta \times \delta$ squares. We then estimate the error by triangle inequality for integrals

$$\begin{aligned} & \left| (7.14) - (7.15) \right| \\ & \leq \iint \left| \delta^{-k-\ell-2} Q_{k,\ell}^{(\delta)}(\zeta, \xi) - \frac{\partial^2 G_{\Omega}^{\mathbf{D}/\mathbf{N}}(\zeta, \xi)}{\partial \zeta \partial \xi} (\zeta - z_i)^{-k} (\xi - z_j)^{-\ell} \right| |d\zeta| |d\xi| + \mathcal{O}(\delta), \end{aligned}$$

where the last error term is from the $\mathcal{O}(\delta)$ discrepancy of lengths of the rectangle side length and its discretization. The terms combined in the piecewise constant function $Q_{k,\ell}^{(\delta)}$ are the contributions of the following form. Among two consecutive corner lattice steps along the bottom side of the rectangle contour $\gamma_{i;s}^{\delta}$, one separates a horizontal edge $\mathbf{u}_h^{\delta} \in \mathbb{Z}_m^2$ from a dual vertex $\mathbf{u}_*^{\delta} \in \mathbb{Z}_*^2$, and the other separates a vertical edge $\mathbf{u}_v^{\delta} \in \mathbb{Z}_m^2$ from a primal vertex $\mathbf{u}_{\bullet}^{\delta} \in \mathbb{Z}_{\bullet}^2$ — see Figure 7.1. Similarly two consecutive steps of along the bottom side of the other rectangle $\gamma_{j;t}^{\delta}$, involve $\mathbf{v}_h^{\delta} \in \mathbb{Z}_m^2$ separated from $\mathbf{v}_*^{\delta} \in \mathbb{Z}_*^2$, and $\mathbf{v}_v^{\delta} \in \mathbb{Z}_m^2$ separated from $\mathbf{v}_{\bullet}^{\delta} \in \mathbb{Z}_{\bullet}^2$. These lattice points depend (in a piecewise constant way) on the variables ζ and ξ , respectively, but we omit writing the explicit dependence $\mathbf{u}_h^{\delta} = \mathbf{u}_h^{\delta}(\zeta), \dots, \mathbf{v}_{\bullet}^{\delta} = \mathbf{v}_{\bullet}^{\delta}(\xi)$ below. The value of $Q_{k,\ell}^{(\delta)}$ is, then, the combined contribution to the discrete integration




Figure 7.1: The top right figure illustrates a domain Ω with marked points z_1, \dots, z_n and concentric radially-ordered square contours $\Gamma_{i;s}$ around z_i for the integrals in (7.11). The top left figure illustrates a corner lattice discrete integration contour $\gamma_{i;s}^{(\delta)}$ with even side lengths approximating a shifted and rescaled version of $\Gamma_{i;s}$ from the inside. The bottom figure shows a part of bottom side discrete and continuous integration contours, and illustrates the lattice points \mathbf{u}_* , \mathbf{u}_h^δ , $\mathbf{u}_\bullet^\delta$, and \mathbf{u}_v^δ involved in two consecutive steps of discrete integration. The piecewise constant function $Q_{k,l}^{(\delta)}$ is defined in terms of these.

from the two pairs of steps,

$$\begin{aligned}
 Q_{k,\ell}^{(\delta)}(\zeta, \xi) := & \frac{1}{4} \left((\mathbf{u}_\bullet^\delta)^{[-k]} (\mathbf{v}_\bullet^\delta)^{[-\ell]} (\partial_\#^\bullet \mathbf{u}) (\partial_\#^\bullet \mathbf{v}) G_{\Omega_\delta}^{\text{D/N}}(\mathbf{z}_i^\delta + \delta \mathbf{u}_v^\delta, \mathbf{z}_j^\delta + \delta \mathbf{v}_v^\delta) \right. \\
 & + (\mathbf{u}_*^\delta)^{[-k]} (\mathbf{v}_\bullet^\delta)^{[-\ell]} (\partial_\#^\bullet \mathbf{u}) (\partial_\#^\bullet \mathbf{v}) G_{\Omega_\delta}^{\text{D/N}}(\mathbf{z}_i^\delta + \delta \mathbf{u}_h^\delta, \mathbf{z}_j^\delta + \delta \mathbf{v}_v^\delta) \\
 & + (\mathbf{u}_\bullet^\delta)^{[-k]} (\mathbf{v}_*^\delta)^{[-\ell]} (\partial_\#^\bullet \mathbf{u}) (\partial_\#^\bullet \mathbf{v}) G_{\Omega_\delta}^{\text{D/N}}(\mathbf{z}_i^\delta + \delta \mathbf{u}_v^\delta, \mathbf{z}_j^\delta + \delta \mathbf{v}_h^\delta) \\
 & \left. + (\mathbf{u}_*^\delta)^{[-k]} (\mathbf{v}_*^\delta)^{[-\ell]} (\partial_\#^\bullet \mathbf{u}) (\partial_\#^\bullet \mathbf{v}) G_{\Omega_\delta}^{\text{D/N}}(\mathbf{z}_i^\delta + \delta \mathbf{u}_h^\delta, \mathbf{z}_j^\delta + \delta \mathbf{v}_h^\delta) \right).
 \end{aligned} \tag{7.16}$$

The discrete monomial asymptotics (3.21) from Proposition 3.4 yield the following locally uniform convergence as $\delta \rightarrow 0$

$$\delta^{-k}(\mathbf{u}_{*/\bullet}^\delta)^{[-k]} = (\zeta - z_i)^{-k} + o(1) \quad \text{and} \quad \delta^{-\ell}(\mathbf{v}_{*/\bullet}^\delta)^{[-\ell]} = (\xi - z_j)^{-\ell} + o(1).$$

By locally uniform convergence of derivatives of discrete Green's functions, Lemmas 3.1 and 3.3, we also have

$$\begin{aligned} \delta^{-2}(\partial_\#^\bullet)_\mathbf{u}(\partial_\#^\bullet)_\mathbf{v} G_{\Omega_\delta}^{\mathbf{D}/\mathbf{N}}(\mathbf{z}_i^\delta + \delta \mathbf{u}_\mathbf{h}, \mathbf{z}_j^\delta + \delta \mathbf{v}_\mathbf{v}) &= (\partial_1^x)(-\mathfrak{i}\partial_2^y) G_\Omega^{\mathbf{D}/\mathbf{N}}(\zeta, \xi) + o(1), \\ \delta^{-2}(\partial_\#^\bullet)_\mathbf{u}(\partial_\#^\bullet)_\mathbf{v} G_{\Omega_\delta}^{\mathbf{D}/\mathbf{N}}(\mathbf{z}_i^\delta + \delta \mathbf{u}_\mathbf{v}, \mathbf{z}_j^\delta + \delta \mathbf{v}_\mathbf{h}) &= (-\mathfrak{i}\partial_1^y)(\partial_2^x) G_\Omega^{\mathbf{D}/\mathbf{N}}(\zeta, \xi) + o(1), \\ \delta^{-2}(\partial_\#^\bullet)_\mathbf{u}(\partial_\#^\bullet)_\mathbf{v} G_{\Omega_\delta}^{\mathbf{D}/\mathbf{N}}(\mathbf{z}_i^\delta + \delta \mathbf{u}_\mathbf{h}, \mathbf{z}_j^\delta + \delta \mathbf{v}_\mathbf{h}) &= (\partial_1^x)(\partial_2^x) G_\Omega^{\mathbf{D}/\mathbf{N}}(\zeta, \xi) + o(1), \\ \delta^{-2}(\partial_\#^\bullet)_\mathbf{u}(\partial_\#^\bullet)_\mathbf{v} G_{\Omega_\delta}^{\mathbf{D}/\mathbf{N}}(\mathbf{z}_i^\delta + \delta \mathbf{u}_\mathbf{v}, \mathbf{z}_j^\delta + \delta \mathbf{v}_\mathbf{v}) &= (-\mathfrak{i}\partial_1^y)(-\mathfrak{i}\partial_2^y) G_\Omega^{\mathbf{D}/\mathbf{N}}(\zeta, \xi) + o(1). \end{aligned}$$

Regrouping these derivatives into Wirtinger derivatives, we obtain the desired estimate for the error integrand,

$$\left| \delta^{-k-\ell-2} Q_{k,\ell}^{(\delta)}(\zeta, \xi) - \frac{\partial^2 G_\Omega^{\mathbf{D}/\mathbf{N}}(\zeta, \xi)}{\partial \zeta \partial \xi} (\zeta - z_i)^{-k} (\xi - z_j)^{-\ell} \right| = o(1)$$

when $\delta \rightarrow 0$. This error integrand is integrated over a fixed finite pair of rectangle sides, so the total error is $|(7.14) - (7.15)| = o(1)$. This completes the proof. \square

References

- [Ada23] DAVID ADAME-CARRILLO. *Discrete symplectic fermions on double dimers and their Virasoro representation*. Preprint: [arXiv:2304.08163](#) (2023).
- [ArPo21] JUHAN ARU AND ELLEN POWELL. *A characterisation of the continuum Gaussian free field in $d \geq 2$ dimensions*. Preprint: [arXiv:2103.07273](#) (2021).
- [ASW19] JUHAN ARU, AVELIO SEPULVEDA, AND WENDELIN WERNER. *On bounded-type thin local sets of the two-dimensional Gaussian free field*. J. Inst. Math. Jussieu **18**(3):591–618 (2019).
- [BaWa23] TIANYI BAI AND YIJUN WAN. *On the Crossing Estimates for Simple Conformal Loop Ensembles*. Int. Math. Res. Notices **2023**(13):11645–11683 (2023).
- [Bas23] MIKHAIL BASOK. *Dimers on Riemann surfaces and compactified free field*. Preprint: [arXiv:2309.14522](#) (2023).
- [BaCh21] MIKHAIL BASOK AND DMITRY CHELKAK. *Tau-functions à la Dubédat and probabilities of cylindrical events for double-dimers and $CLE(4)$* . J. Eur. Math. Soc. **23**(8):2787–2832 (2021).
- [BIVW23] BARAN BAYRAKTAROGU, KONSTANTIN IZYUROV, TUOMAS VIRTANEN, AND CHRISTIAN WEBB. *Bosonization of primary fields for the critical Ising model on multiply connected planar domains*. Preprint: [arXiv:2312.02960](#) (2023).
- [BePo23] NATHANAËL BERESTYCKI AND ELLEN POWELL. *Gaussian free field and Liouville quantum gravity*. Book draft: <https://homepage.univie.ac.at/nathanael.berestycki/wp-content/uploads/2023/09/master.pdf> (2023).
- [BPR20] NATHANAËL BERESTYCKI, ELLEN POWELL, AND GOURAB RAY. *A characterisation of the Gaussian free field*. Probab. Theory Rel. Fields **176** (2020).
- [BPR21] NATHANAËL BERESTYCKI, ELLEN POWELL, AND GOURAB RAY. *$(1 + \varepsilon)$ moments suffice to characterise the GFF*. European J. Probability **26** (2021).
- [Cam24] FEDERICO CAMIA. *Conformal Covariance of Connection Probabilities and Fields in 2D Critical Percolation*. Communications on Pure and Applied Mathematics **77**(3):2138–2176 (2024).
- [CaFe24] FEDERICO CAMIA AND YU FENG. *Logarithmic singularity in the density four-point function of two-dimensional critical percolation in the bulk*. Preprint: [arXiv:2403.18576](#) (2024).
- [CPST21] NISHANT CHANDGOTIA, RON PELED, SCOTT SHEFFIELD, AND MARTIN TASSY. *Delocalization of Uniform Graph Homomorphisms from \mathbb{Z}^2 to \mathbb{Z}* . Commun. Math. Phys. **387**:621–647 (2021).

- [CHI15] DMITRY CHELKAK, CLÉMENT HONGLER, AND KONSTANTIN IZYUROV. *Conformal invariance of spin correlations in the planar Ising model*. Annals of Mathematics **181**(3):1087–1138 (2015).
- [CHI21] DMITRY CHELKAK, CLÉMENT HONGLER, AND KONSTANTIN IZYUROV. *Correlations of primary fields in the critical Ising model*. Preprint: [arXiv:2103.10263](#) (2021).
- [ChSm12] DMITRY CHELKAK AND STANISLAV SMIRNOV. *Universality in the 2D Ising model and conformal invariance of fermionic observables*. Invent. math. **189**:515–580 (2012).
- [CCRR23] LEANDRO CHIARINI, ALESSANDRA CIPRIANI, ALAN RAPOPORT, AND WIOLETTA RUSZEL. *Fermionic Gaussian free field structure in the Abelian sandpile model and uniform spanning tree*. Preprint: [arXiv:2309.08349](#) (2023).
- [DKRV16] FRANÇOIS DAVID, ANTTI KUPIAINEN, RÉMI RHODES, AND VINCENT VARGAS. *Liouville Quantum Gravity on the Riemann sphere*. Comm. Math. Phys. **342**(3):869–907 (2016).
- [DMS97] PHILIPPE DI FRANCESCO, PIERRE MATHIEU, AND DAVID SÉNÉCHAL. *Conformal Field Theory*. Graduate texts in contemporary physics. Springer-Verlag New York (1997).
- [Dub09] JULIEN DUBÉDAT. *SLE and the free field: partition functions and couplings*. J. Amer. Math. Soc. **22**:995–1054 (2009).
- [Dub15] JULIEN DUBÉDAT. *Dimers and families of Cauchy-Riemann operators I*. J. Amer. Math. Soc. **28**(4):1063–1167 (2015).
- [Dub18] JULIEN DUBÉDAT. *Double dimers, conformal loop ensembles and isomonodromic deformations*. J. Eur. Math. Soc. **21**(1):1–54 (2018).
- [DHLRR15] HUGO DUMINIL-COPIN, MATAN HAREL, BENOIT LASLIER, ARAN RAOUFI, AND GOURAB RAY. *Logarithmic Variance for the Height Function of Square-Ice*. Commun. Math. Phys. **396**:867–902 (2022).
- [DKMO20] HUGO DUMINIL-COPIN, ALEX KARRILA, IOAN MANOLESCU, AND MENDES OULAMARA. *Delocalization of the height function of the six-vertex model*. Preprint: [arXiv:2012.13750](#) (2020).
- [FrBZ04] EDWARD FRENKEL AND DAVID BEN-ZVI. *Vertex Algebras and Algebraic Curves. 2nd ed.* Mathematical Surveys and Monographs, vol. 88.R. Amer. Math. Soc. (2004).
- [Gaw99] KRZYSZTOF GAWĘDZKI. *Lectures on conformal field theory*. Quantum fields and strings: a course for mathematicians, Vol. 1, 2 (Princeton, NJ, 1996/1997), 727–805, Amer. Math. Soc., Providence, RI (1999).

- [GHP19] REZA GHEISSARI, CLÉMENT HONGLER, AND S.C. PARK. *Ising Model: Local Spin Correlations and Conformal Invariance*. Comm. Math. Phys. **367**:771–833 (2019).
- [GlJa81] JAMES GLIMM AND ARTHUR JAFFE *Quantum physics: a functional integral point of view* Springer-Verlag (1981).
- [GKR24] COLIN GUILLARMOU, ANTTI KUPIAINEN, AND RÉMI RHODES. *Review on the probabilistic construction and Conformal bootstrap in Liouville Theory*. Preprint: arXiv:2403.12780 (2024).
- [GKRV21] COLIN GUILLARMOU, ANTTI KUPIAINEN, RÉMI RHODES., AND VINCENT VARGAS. *Segal’s axioms and bootstrap for Liouville Theory*. Preprint: arXiv:2112.14859 (2021).
- [Hon10] CLÉMENT HONGLER. *Conformal Invariance of Ising Model Correlations*, Ph.D. thesis, Université de Genève, <https://archive-ouverte.unige.ch/unige:18163> (2010).
- [HKV22] CLÉMENT HONGLER, KALLE KYTÖLÄ, AND FREKRIK VIKLUND. *Conformal Field Theory at the lattice level: Discrete complex analysis and Virasoro structure*. Comm. Math. Phys. **395**:1–58 (2022).
- [HoSm13] CLÉMENT HONGLER AND STANISLAV SMIRNOV. *The energy density in the planar Ising model*. Acta Math **211**:191–225 (2013).
- [Izy11] KONSTANTIN IZYUROV. *Holomorphic spinor observables and interfaces in the critical Ising model*. PhD thesis, Université de Genève (2011).
- [IzKy13] KONSTANTIN IZYUROV AND KALLE KYTÖLÄ. *Hadamard’s formula and couplings of SLEs with free field*. Probab. Th. Rel. Fields **155**:35–69 (2013).
- [Kac98] VICTOR G. KAC. *Vertex algebras for beginners*. No. 10. American Mathematical Soc. (1998).
- [KaMa13] NAM-GYU KANG AND NIKOLAI G. MAKAROV. *Gaussian free field and conformal field theory*. Astérisque **353** (2013).
- [Ken00] RICHARD KENYON. *Conformal invariance of domino tiling*. Ann. Probab. **28**(2):759–795 (2000).
- [Ken01] RICHARD KENYON. *Dominos and the Gaussian Free Field*. Ann. Probab. **29**(3):1128–1137 (2001).
- [Ken14] RICHARD KENYON. *Conformal Invariance of Loops in the Double-Dimer Model*. Commun. Math. Phys. **326**:477–497 (2014).

- [LaLi10] GREGORY F. LAWLER AND VLADA LIMIC. *Random walk: A modern introduction*. Cambridge University Press (2010).
- [LeLi04] JAMES LEPOWSKY AND HAISHENG LI. *Introduction to vertex operator algebras and their representations*. Vol. 227. Springer Science & Business Media (2004).
- [MiSh16] JASON MILLER AND SCOTT SHEFFIELD. *Imaginary Geometry I: Interacting SLEs*. Probab. Th. Rel. Fields **164**:553–705 (2016).
- [QiWe18] WEI QIAN AND WENDELIN WERNER. *Coupling the Gaussian free fields with free and with zero boundary conditions via common level lines*. Comm. Math. Phys. **361**:53–80 (2018).
- [PeWu19] EVELIINA PELTOLA AND HAO WU. *Global and Local Multiple SLEs for $\kappa \leq 4$ and Connection Probabilities for Level Lines of GFF*. Commun. Math. Phys. **366**:469–536 (2019).
- [PoRu17] ADRIEN PONCELET AND PHILIPPE RUELLE. *Multipoint correlators in the Abelian sandpile model*. J. Stat. Mech.: Theory Exp. **2017** (2017).
- [Rus18] MARIANNA RUSSIKH. *Dominos in hedgehog domains*. Annales de l’Institut Henri Poincaré D **8**(1):1-33, (2018).
- [ScSh09] ODED SCHRAMM AND SCOTT SHEFFIELD. *Contour lines of the two-dimensional discrete Gaussian free field*. Acta Math. **202** (2009).
- [ScSh13] ODED SCHRAMM AND SCOTT SHEFFIELD. *A contour line of the continuum Gaussian free field*. Probab. Th. Rel. Fields **157** (2013).
- [She07] SCOTT SHEFFIELD. *Gaussian free fields for mathematicians*. Probab. Th. Rel. Fields **139**:521-541 (2007).
- [Smi10] STANISLAV SMIRNOV. *Conformal invariance in random cluster models. I. Holomorphic fermions in the Ising model*. Annals of Mathematics **172**(2):1435–1467 (2010).
- [WePo21] WENDELIN WERNER AND ELLEN POWELL. *Lecture notes on the Gaussian free field*. Cours Spécialisés, 28. Société Mathématique de France, Paris (2021).