Sampling from Spherical Spin Glasses in Total Variation via Algorithmic Stochastic Localization

Brice Huang*

Andrea Montanari[†]

Huy Tuan Pham[‡]

April 25, 2024

Abstract

We consider the problem of algorithmically sampling from the Gibbs measure of a mixed p-spin spherical spin glass. We give a polynomial-time algorithm that samples from the Gibbs measure up to vanishing total variation error, for any model whose mixture satisfies

$$\xi''(s) < \frac{1}{(1-s)^2}, \qquad \forall s \in [0,1).$$

This includes the pure *p*-spin glasses above a critical temperature that is within an absolute (*p*-independent) constant of the so-called shattering phase transition.

Our algorithm follows the algorithmic stochastic localization approach introduced in [AMS22]. A key step of this approach is to estimate the mean of a sequence of tilted measures. We produce an improved estimator for this task by identifying a suitable correction to the TAP fixed point selected by approximate message passing (AMP). As a consequence, we improve the algorithm's guarantee over previous work, from normalized Wasserstein to total variation error. In particular, the new algorithm and analysis opens the way to perform inference about one-dimensional projections of the measure.

Contents

1	Intro	oduction 2		
	1.1	Background and related work		
	1.2	Notations		
2	Main result			
	2.1	Mean estimation of tilted measure		
	2.2	Stochastic localization sampling		
	2.3	The correction $\Delta(m)$		
	2.4	Fundamental limits of algorithmic SL, replica symmetry breaking, and shattering 9		
3	Prel	iminaries 10		
	3.1	Stochastic localization		
	3.2	Planted model and contiguity		
	3.3	Basic regularity estimate		
	3.4	Conditioning lemma 13		

*Department of Electrical Engineering and Computer Science, Massachusetts Institute of Technology

[†]Department of Statistics and Department of Mathematics, Stanford University

[‡]Department of Mathematics, Stanford University

4	Analysis of mean computation algorithm	14
5	Analysis of AMP iteration: proof of Proposition 4.35.1State evolution limit	17 17 19 23
6	Description of TAP fixed point: proof of Proposition 4.46.1Existence and uniqueness of TAP fixed point6.2Characterization of Riemannian critical points: proof of Proposition 6.26.3Approximate stationarity and local concavity of \mathcal{F}_{TAP} : proof of Proposition 6.96.4Proof of conditioning bound6.5Determinant concentration and estimate of Kac–Rice integral6.6Algorithmic guarantees and Lipschitz continuity of correction	26 29 32 38 40 44
7	Local computation of magnetization: proof of Proposition 4.67.1Conditional magnetization per band7.1.1Quadratic Hamiltonians7.1.2Estimates of restricted partition functions7.1.3Proof of Lemma 7.87.1.4Magnetization in the band: Proof of Lemma 7.27.2Integrating over bands	48 49 50 57 64 69 76
8	Lognormal fluctuations of partition function	92
9	Completing the proof of Theorem 2.19.1TV-closeness of Euler discretization: Proof of Proposition 9.19.2Log-concavity of late measures	93 94 96
10	Failure of stochastic localization in complementary regime	99

1 Introduction

Let $\gamma_2, \gamma_3, \ldots \ge 0$ satisfy $\sum_{p \ge 2} 2^p \gamma_p^2 < \infty$. The mixed *p*-spin glass Hamiltonian $H_N : \mathbb{R}^N \to \mathbb{R}$ is

$$H_N(\boldsymbol{\sigma}) = \sum_{p \ge 2} \frac{\gamma_p}{N^{(p-1)/2}} \sum_{i_1, \dots, i_p=1}^N G_{i_1, \dots, i_p} \sigma_{i_1} \cdots \sigma_{i_p}, \qquad G_{i_1, \dots, i_p} \stackrel{i.i.d.}{\sim} \mathcal{N}(0, 1).$$
(1.1)

Define the mixture function $\xi(s) = \sum_{p \ge 2} \gamma_p^2 s^p$, so that H_N is the Gaussian process with covariance

$$\mathbb{E} H_N(\boldsymbol{\sigma}^1) H_N(\boldsymbol{\sigma}^2) = N\xi(\langle \boldsymbol{\sigma}^1, \boldsymbol{\sigma}^2 \rangle / N).$$

The Gibbs measure of this model is the probability measure over the sphere $S_N = \{ \boldsymbol{x} \in \mathbb{R}^N : \|\boldsymbol{x}\|_2^2 = N \}$ given by

$$\mu_{H_N}(\mathsf{d}\boldsymbol{\sigma}) = \frac{1}{Z_N} \exp(H_N(\boldsymbol{\sigma})) \ \mu_0(\mathsf{d}\boldsymbol{\sigma}), \qquad Z_N = \int_{S_N} \exp(H_N(\boldsymbol{\sigma})) \ \mu_0(\mathsf{d}\boldsymbol{\sigma}) \,. \tag{1.2}$$

Here and below, μ_0 denotes the uniform probability measure on S_N . We will denote by $\mathbf{G} = (G_{i_1,...,i_p})_{p \ge 2, i_{\ell \le N}}$ the vector of couplings that defines the Hamiltonian.

In this paper, we consider the problem of efficiently sampling from this Gibbs measure. For dist a distance on $\mathcal{P}(\mathbb{R}^N)$ (the set of probability measures over \mathbb{R}^N), we seek a computationally efficient algorithm that generates σ^{alg} whose law μ^{alg} satisfies dist $(\mu^{alg}, \mu_{H_N}) = o_N(1)$, with high probability over H_N .

We follow the algorithmic stochastic localization approach introduced by [AMS22], which is in turn motivated by the stochastic localization process [Eld20], and closely related to the denoising diffusions method in machine learning [SDWMG15, HJA20, SSDK+21] (see [Mon23] for a discussion of the connection). The basic idea is to generate (an approximation of) a sample path from the following Ito diffusion on \mathbb{R}^N :

$$d\boldsymbol{y}_t = \boldsymbol{m}(\boldsymbol{y}_t, t) dt + d\boldsymbol{B}_t, \quad \boldsymbol{y}_0 = \boldsymbol{0}, \qquad (1.3)$$

where $(B_t)_{t\geq 0}$ is a standard Brownian motion and $m(y,t) = \mathbb{E}[\sigma|t\sigma + \sqrt{t}g = y]$ (conditioning over G is implicit here), with the conditional expectation being taken with respect to $(\sigma, g) \sim \mu_{H_N} \otimes \mathcal{N}(0, I_N)$. The key remark (see Section 3.1) is that y_t thus defined has the same distribution at $t\sigma + B'_t$ (with B'_t a different Brownian motion) and therefore y_t/t converges to a sample from the desired measure. Of course, constructing an actual algorithm requires to discretize time and — crucially — to define an efficient algorithm that approximates the conditional mean $m(\cdot, t)$ well enough.

The analysis of [AMS22] establishes that this approach samples from the Gibbs measure of the Sherrington-Kirkpatrick model on the (more difficult) cube $\Sigma_N = \{-1, 1\}^N$, up to vanishing normalized Wasserstein error. That is, with probability $1 - o_N(1)$ over the Sherrington-Kirkpatrick Hamiltonian H_N , there is a coupling of μ_{H_N} and μ^{alg} such that for $(\boldsymbol{\sigma}, \boldsymbol{\sigma}^{alg})$ drawn from this coupling,

$$\frac{1}{N} \mathop{\mathbb{E}}_{\boldsymbol{\sigma}, \boldsymbol{\sigma}^{\mathsf{alg}}} \|\boldsymbol{\sigma} - \boldsymbol{\sigma}^{\mathsf{alg}}\|_2^2 = o_N(1).$$
(1.4)

Our main result is an improved version of this general sampling scheme that samples from μ_{H_N} , in the stronger sense of vanishing **total variation** error, for any spherical spin glass whose mixture satisfies

$$\xi''(s) < \frac{1}{(1-s)^2}, \quad \forall s \in [0,1).$$
 (1.5)

Remark 1.1. For the special case of pure models $\xi(s) = \beta^2 s^p$, (1.5) holds for all $\beta < \beta_{SL}(p)$, where we defined the stochastic localization inverse temperature as

$$\beta_{\mathrm{SL}}(p) := \frac{1}{2} \sqrt{\left(\frac{p}{p-1}\right) \left(\frac{p}{p-2}\right)^{p-2}}.$$

For large p we have $\beta_{SL}(p) = e/2 + O(1/p)$.

As mentioned above, the key challenge in implementing the algorithmic stochastic localization approach is the construction of an efficient algorithm to approximate the mean of the measure $\mu_{H_N}(d\sigma)$, as well as its conditional mean given Gaussian observations. The latter corresponds to the mean of a exponential tilt $\mu_{H_N,y}(d\sigma) \propto \exp(\langle y, \sigma \rangle) \mu_{H_N}(d\sigma)$. Approximating m(y) was achieved in [AMS22] by a variational approach that requires minimizing the so called Thouless-Anderson-Palmer (TAP) free energy [TAP77]. The same paper established that the resulting estimate satisfies (with high probability) $||m(y) - m^{TAP}(y)||^2 = o(N)$. (For the case of a measure supported over S_N , the function $m(\cdot)$ does not depend on t, and we will therefore omit this argument.) Note that $\|\boldsymbol{m}(\boldsymbol{y})\|^2 = \Theta(N)$, and therefore [AMS22] establishes the weakest non-trivial upper bound on $\|\boldsymbol{m}(\boldsymbol{y}) - \boldsymbol{m}^{TAP}(\boldsymbol{y})\|^2$. However, in order to obtain a sampling algorithm with guarantees in total variation distance, it is necessary to construct an efficient estimator $\widehat{\boldsymbol{m}}(\boldsymbol{y})$ satisfying $\|\boldsymbol{m}(\boldsymbol{y}) - \widehat{\boldsymbol{m}}(\boldsymbol{y})\|^2 = o(1)$. The construction and analysis of such an estimator is the main problem solved in the present paper.

In fact we prove the following:

- 1. The TAP estimator is significantly more accurate than what could be hoped from the analysis of [AMS22, AMS23a]. Namely, we prove that $||m(y) m^{TAP}(y)||^2 = O(1)$.
- 2. We design a correction $\Delta(y)$ to the TAP estimator that can be computed efficiently and such that, letting $\widehat{m}(y) = m^{\text{TAP}}(y) + \Delta(y)$, we achieve the desired accuracy $||m(y) \widehat{m}(y)||^2 = o(1)$.

1.1 Background and related work

A substantial line of work in probability theory studies Langevin dynamics for the Gibbs measure (1.2). This is defined as the following diffusion on S_N

$$d\boldsymbol{\sigma}_{t} = \left(\mathsf{P}_{\boldsymbol{\sigma}_{t}}^{\perp} \nabla H_{N}(\boldsymbol{\sigma}_{t}) - \frac{N-1}{2N} \boldsymbol{\sigma}_{t}\right) \mathsf{d}t + \sqrt{2} \mathsf{P}_{\boldsymbol{\sigma}_{t}}^{\perp} \mathsf{d}\boldsymbol{B}_{t}, \qquad (1.6)$$

where B_t is a standard *N*-dimensional Brownian motion, and $\mathsf{P}_{\sigma_t}^{\perp}$ is the projector orthogonal to σ_t . Langevin dynamics is a Markov process reversible for the measure μ_{H_N} of Eq. (1.2). Therefore, suitable discretizations of Langevin dynamics can be used to sample from μ_{H_N} .

An asymptotically exact characterization of Langevin dynamics on short times horizons t = O(1), in the high-dimensional limit $N \to \infty$, is provided by the so-called Cugliandolo-Kurchan equations. These were studied first in physics [CHS93, CK93] and subsequently established rigorously in probability theory [BADG06]. Unfortunately, this approach does not give access to mixing times. On top of that, the Cugliandolo-Kurchan equations proved difficult to analyze rigorously except at sufficiently 'high temperature' (i.e. when $\xi(s) = \beta^2 \xi_1(s)$, for a fixed ξ_1 and β small enough) [DGM07].

Based on a postulated asymptotic form of the Cugliandolo-Kurchan equations, as well as on thermodynamic calculations, physicists conjecture a phase transition in the mixing time of Langevin dynamics, when initialized uniformly at random [CHS93, CK93]. Namely, they expect the mixing time to be polynomial in N for

$$\xi'(q) < \frac{q}{1-q}, \qquad \forall q \in (0,1).$$
 (1.7)

and exponentially large in the opposite case, and more precisely when $\sup_{q \in (0,1)} (1-q)\xi'(q)/q > 1$. This is commonly referred to as the 'dynamical phase transition,' and corresponds to a phase transition in the geometry of the Gibbs measure, known as 'shattering phase transition.' In the homogeneous case $\xi(t) = \beta^2 t^p$, the above formula implies that the dynamical/shattering phase transition takes place at $\beta = \beta_{sh}(p)$ given by

$$\beta_{\rm sh}(p) = \sqrt{\frac{(p-1)^{p-1}}{p(p-2)^{p-2}}}\,.$$
(1.8)

For large p, $\beta_{sh}(p) = \sqrt{e} + O(1/p)$. We also recall that a second phase transition ('condensation' or 'static' or 'replica symmetry breaking') takes place at a lower temperature

$$\beta_c^2(p) = \inf_{s \in [0,1]} \left(\frac{1}{s^p} \log\left(\frac{1}{1-s}\right) - \frac{1}{s^{p-1}} \right) \,. \tag{1.9}$$

This corresponds to a non-analiticity of the free energy, and to the temperature at which the overlap stops concentrating [Che13]. For large p, we have $\beta_c(p) = \sqrt{\log p}(1 + o_p(1))$.

Towards the goal of proving the dynamical phase transition phenomenon, Ben Arous and Jagannath [AJ24] established that — for the homogeneous model — shattering takes place in a non-empty temperature interval, implying in particular $\beta_{sh}(p) < \beta_c(p)$ strictly. A order-optimal bound was proven in [AMS23b], who proved $\beta_{sh}(p) \leq C$ for a *p*-independent constant *C*.

A bolder version of the dynamical phase transition conjecture postulates that not only Langevin dynamics is slow, but indeed sampling is fundamentally hard beyond the shattering phase transition. Rigorous evidence was provided in [AMS23b], which proves that 'stable algorithms' fail to sample from μ_{H_N} under shattering.

In the positive direction Gheissari and Jagannath [GJ19] proved that there exists $\underline{\beta}(p) > 0$ such that Langevin dynamics mixes rapidly for $\beta < \underline{\beta}(p)$. These authors also note that their proof technique extends to mixed models.

A closely related model is the Ising version of model (1.2), whereby the uniform measure μ_0 over the sphere S_N is replaced by the uniform measure over the hypercube $\{+1, -1\}^N$. A dynamical/shattering phase transition was conjectured in that setting as well [KT87], although at a different temperature. In this context, shattering for a non-empty interval of temperatures was proven in [GJK23], while mixing of Glauber dynamics at high temperature was proven in [ABXY22, AJK⁺23]. As for the spherical case, positive and negative results are separated by a large gap, indeed diverging with p.

The algorithmic stochastic localization approach was applied to Ising mixed p-spin spin classes in [AMS23a], which established the Wasserstein guarantee (1.4).

1.2 Notations

Throughout this paper, $\|\boldsymbol{\sigma}\|_N = \|\boldsymbol{\sigma}\|/\sqrt{N} = \sqrt{\boldsymbol{\sigma}^\top \boldsymbol{\sigma}/N}$ is the norm corresponding to the inner product $\langle \boldsymbol{\sigma}_1, \boldsymbol{\sigma}_2 \rangle_N = \langle \boldsymbol{\sigma}_1, \boldsymbol{\sigma}_2 \rangle/N = \boldsymbol{\sigma}_1^\top \boldsymbol{\sigma}_2/N$. There will be no confusion with the ℓ_p norm, which will not appear. Given a matrix \boldsymbol{A} , we denote by $\|\boldsymbol{A}\|_F$ its Frobenius norm. For $\boldsymbol{m} \in \mathbb{R}^N$, measurable $I \subseteq \mathbb{R}$, and $\rho > 0$, we define

$$Band(\boldsymbol{m}, I) := \{\boldsymbol{\sigma} \in S_N : \langle \boldsymbol{m}, \boldsymbol{\sigma} \rangle_N \in I\} ,$$

$$B_N(\boldsymbol{m}, \rho) := \{\boldsymbol{x} \in \mathbb{R}^N : \|\boldsymbol{x} - \boldsymbol{m}\|_N \leq \rho\} .$$

We will occasionally abuse notations and write, for $q \in \mathbb{R}$, Band(m, q) instead of $Band(m, \{q\})$.

We will often state that certain events occur with probability $1 - e^{-cN}$. When we do, c > 0 is an unspecified constant, which may change from line to line and may depend on all parameters other than N. We use p-lim to denote limit in probability.

We write $G \sim \text{GOE}(N)$ if G is a symmetric matrix with independent centered Gaussian entries on or above the diagonal with $G_{ii} \sim \mathcal{N}(0, 2/N)$ and $G_{ij} \sim \mathcal{N}(0, 1/N)$ for i < j.

Throughout the paper, the mixture ξ is fixed and various constants can depend on ξ but we will track this dependence. If ι is a small constant, we write $\iota' = o_{\iota}(1)$ if $|\iota'| \leq h(\iota)$ where h is a function independent of N, such that $\lim_{\iota \to 0} h(\iota) = 0$.

2 Main result

In this section we describe the sampling algorithm and state our main result. Throughout, we assume the model ξ satisfies (1.5).

2.1 Mean estimation of tilted measure

We first describe the main subroutine of our algorithm, which estimates the mean of the following exponentially tilted version of μ_{H_N} . For $\boldsymbol{y} \in \mathbb{R}^N$, define

$$\mu_{H_N,\boldsymbol{y}}(\mathsf{d}\boldsymbol{\sigma}) = \frac{1}{Z(\boldsymbol{y})} \exp\left\{H_N(\boldsymbol{\sigma}) + \langle \boldsymbol{y}, \boldsymbol{\sigma} \rangle\right\} \ \mu_0(\mathsf{d}\boldsymbol{\sigma}) \,. \tag{2.1}$$

The tilt y will be generated by the outer loop of the algorithm described in Subsection 2.2, which implements a discretized version of the stochastic localization process. The outer loop also provides a time t > 0, which this subroutine will take as input. The algorithm consists of three steps as outlined below. We defer the description of the correction $\Delta(m)$ to Section 2.3.

(1) Let $\xi_t(s) = \xi(s) + ts$, and define the sequence $\{q_k : k \ge 0\}$ by $q_0 = 0$ and

$$q_{k+1} = \frac{\xi'_t(q_k)}{1 + \xi'_t(q_k)}.$$
(2.2)

Starting from initialization $m^{-1} = w^0 = 0$, run the approximate message passing (AMP) iteration

$$\boldsymbol{m}^{k} = (1 - q_{k})\boldsymbol{w}^{k}, \qquad \boldsymbol{w}^{k+1} = \nabla H_{N}(\boldsymbol{m}^{k}) + \boldsymbol{y} - (1 - q_{k})\xi''(q_{k})\boldsymbol{m}^{k-1}, \qquad (2.3)$$

for K_{AMP} iterations. Let $m^{\text{AMP}} = m^{K_{\text{AMP}}}$.

(2) Define

$$\theta(s) = \xi(1) - \xi(s) - (1 - s)\xi'(s) \tag{2.4}$$

and the TAP free energy

$$\mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m};\boldsymbol{y}) = H_N(\boldsymbol{m}) + \langle \boldsymbol{y}, \boldsymbol{m} \rangle + \frac{N}{2}\theta(\|\boldsymbol{m}\|_N^2) + \frac{N}{2}\log(1 - \|\boldsymbol{m}\|_N^2).$$
(2.5)

Starting from $\boldsymbol{m}^{\text{AMP}}$, run gradient ascent on $\mathcal{F}_{\text{TAP}}(\cdot; \boldsymbol{y})$ for $K_{\text{GD}}(N) := [K_{\text{GD}}^* \log N]$ iterations, and let the resulting point be $\boldsymbol{m}^{\text{GD}}$.

(3) Output $m^{alg} := m^{GD} + \Delta(m^{GD})$, with $\Delta(m)$ defined as in Section 2.3.

Pseudocode for the computation of m^{alg} is provided in Algorithm 1.

Algorithm 1: APPROXIMATE MEAN COMPUTATION

Input: H_N , $\boldsymbol{y} \in \mathbb{R}^N$, t > 0. Parameters: K_{AMP} , $K_{GD}(N)$, $\eta > 0$ 1 $\boldsymbol{m}^{-1} = \boldsymbol{w}^0 = \boldsymbol{0}$, 2 For $k = 0, \dots, K_{AMP}$, run iteration (2.3) 3 Let $\boldsymbol{u}^0 = \boldsymbol{m}^{AMP} = \boldsymbol{m}^{K_{AMP}}$ 4 for $k = 0, \dots, K_{GD}(N) - 1$ do 5 $| \boldsymbol{u}^{k+1} = \boldsymbol{u}^k - \eta \nabla \mathcal{F}_{TAP}(\boldsymbol{u}^k; \boldsymbol{y})$ 6 end 7 Let $\boldsymbol{m}^{GD} = \boldsymbol{u}^{K_{GD}(N)}$ 8 return $\boldsymbol{m}^{alg}(H_N, \boldsymbol{y}, t) = \boldsymbol{m}^{GD} + \boldsymbol{\Delta}(\boldsymbol{m}^{GD})$

2.2 Stochastic localization sampling

We are now in position to describe the sampling algorithm, which uses Algorithm 1 as a subroutine. The main idea is to truncate the diffusion process (1.3) to the interval [0, T], and to replace it by its Euler discretization (see Step 6 in Algorithm 2 below).

We will prove that, for T a sufficiently large constant, the tilted measure of Eq. (2.1), with $y = y_T$ is well approximated by a strongly log-concave measure. As a consequence, we can sample from it in total variation using standard approaches such as the Metropolis-adjusted Langevin algorithm, or MALA (see [CLA⁺21] and references therein). Formally, define

where $U \in \mathbb{R}^{N \times (N-1)}$ is an orthonormal basis of the orthogonal complement of y, and

$$H_{N,\boldsymbol{y}}^{\mathsf{proj}}(\boldsymbol{\rho}) = H_{N,\boldsymbol{y}}(\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})) - \frac{N}{2}\log(1 + \|\boldsymbol{\rho}\|_{N}^{2}).$$
(2.7)

Note that σ_y is the inverse of the stereographic projection T_y from $S_N \cap \{\sigma : \langle \sigma, y \rangle > 0\}$ to the affine plane $\{\hat{y} + U\rho : \rho \in \mathbb{R}^{N-1}\}$. We will see (Lemma 9.5) that the push-forward of $\mu_{H_N,y}(\cdot | \langle \sigma, y \rangle > 0)$ under T_y is precisely

$$\nu_{H_N,\boldsymbol{y}}^{\mathsf{proj}}(\mathsf{d}\boldsymbol{\rho}) = \frac{1}{\widehat{Z}(\boldsymbol{y})} \exp H_{N,\boldsymbol{y}}^{\mathsf{proj}}(\boldsymbol{\rho}) \, \mathsf{d}\boldsymbol{\rho}.$$
(2.8)

Let $\varepsilon_0 = 0.1$ and $\varphi : [0, +\infty) \rightarrow [0, +\infty)$ be a twice continuously differentiable function satisfying $\varphi(x) = 0$ for $x \in [0, \varepsilon_0]$ and

$$\frac{1}{(1+x)^{3/2}} + \varphi'(x) \ge \varepsilon_0, \qquad \frac{1-2x}{(1+x)^{5/2}} + \varphi'(x) + 2x\varphi''(x) \ge \varepsilon_0$$
(2.9)

for all $x \ge 0$. (Existence of such a function is shown in Fact 9.9.) Define the following measure on \mathbb{R}^{N-1} :

$$\widetilde{\nu}_{H_{N},\boldsymbol{y}}^{\mathsf{proj}}(\mathsf{d}\boldsymbol{\rho}) = \frac{1}{\widetilde{Z}(\boldsymbol{y})} \exp \widetilde{H}_{N,\boldsymbol{y}}^{\mathsf{proj}}(\boldsymbol{\rho}) \, \mathsf{d}\boldsymbol{\rho}, \qquad \widetilde{H}_{N,\boldsymbol{y}}^{\mathsf{proj}}(\boldsymbol{\rho}) = H_{N,\boldsymbol{y}}^{\mathsf{proj}}(\boldsymbol{\rho}) - \frac{TN}{2}\varphi(\|\boldsymbol{\rho}\|_{N}^{2}). \tag{2.10}$$

We will show that for sufficiently large T, $\tilde{\nu}_{H_N,y}^{\text{proj}}$ is strongly log-concave (Proposition 9.8) and approximates $\nu_{H_N,y}^{\text{proj}}$ in total variation (Corollary 9.7). Thus, we may sample from it using MALA, and produce samples from $\mu_{H_N,y}$ by pushing forward through σ_y .

Theorem 2.1. Suppose ξ satisfies (1.5). There exist constants K_{AMP} , K^*_{GD} , η , T depending on ε and ξ such that running Algorithm 2 with parameters K_{AMP} , $K_{GD}(N) = K^*_{GD} \log N$, η , T, the following holds. With probability $1 - o_N(1)$ over H_N , $\mu^{alg} = \mathcal{L}(\sigma^{alg})$ satisfies

$$\mathsf{TV}(\mu^{\mathsf{alg}}, \mu_{H_N}) \leq o_N(1).$$

Further the complexity of the algorithm is upper bounded by $CN^4 (N + \chi_{\nabla H}) \log N + \chi_{\text{log-conc}}$, where $\chi_{\nabla H}$ is the complexity of evaluating $\nabla H_N(\mathbf{m})$ at a point \mathbf{m} with $\|\mathbf{m}\|_N \leq 1$, and $\chi_{\text{log-conc}}$ is the complexity of sampling from a 1-strongly log-concave measure in N dimension using MALA to accuracy 1/N in total variation.

Remark 2.2. The main result of [CLA⁺21] implies that, for a 'warm start' initialization $\chi_{\text{log-conc}}$ is of order $N^{3/2} \log N$. In the present case we do not have a good warm start, and obtain $\chi_{\text{log-conc}} \leq C \cdot N^{5/2}$. We believe this bound is suboptimal, but made no attempt at improving it.

Algorithm 2: SAMPLING

Input: H_N . Parameters: K_{AMP} , $K_{GD}(N)$, η , T > 0, where T is a multiple of N^{-4} 1 Set $\delta = N^{-4}$, $L = T/\delta$ 2 Set $y^0 = 0$ 3 for $\ell = 0, ..., L - 1$ do 4 Let $m^{\ell} = m^{alg}(H_N, y^{\ell}, \ell\delta)$ be the output of Algorithm 1 on input $(H_N, y^{\ell}, \ell\delta, K_{AMP}(N), K_{GD}, \eta)$ 5 Draw $w^{\ell} \sim \mathcal{N}(0, I_N)$ independent of everything else 6 Set $y^{\ell+1} = y^{\ell} + \delta m^{\ell} + \sqrt{\delta} w^{\ell}$ 7 end 8 Let $\tilde{\nu}_{H_N, y^L}^{\text{proj}}$ be defined according to Eq. (2.10) 9 Use MALA to sample from $\rho^{MALA} \sim \nu^{MALA}$, to accuracy $\text{TV}(\nu^{MALA}, \tilde{\nu}_{H_N, y^L}^{\text{proj}}) \leq 1/N$ 10 return $\sigma_{y^L}(\rho^{MALA})$

2.3 The correction $\Delta(m)$

We now describe the computation of the correction $\Delta(m)$. Let T_m be the (N-1)-dimensional subspace orthogonal to m and define $H_N(\cdot; m) : \mathsf{T}_m \to \mathbb{R}$ via $H_N(x; m) := H_N(m + x)$. We then define the tensors

$$\boldsymbol{A}^{(2)}(\boldsymbol{m}) := \nabla_{\boldsymbol{x}}^2 H_N(\boldsymbol{0}; \boldsymbol{m}), \quad \boldsymbol{A}^{(3)}(\boldsymbol{m}) := \nabla_{\boldsymbol{x}}^3 H_N(\boldsymbol{0}; \boldsymbol{m}).$$
(2.11)

These should be interpreted as tensors $A^{(i)}(m) \in \mathsf{T}_{m}^{\otimes i}$. Let $\gamma_{*,N}(m)$ be the unique solution of

$$\begin{cases} \mathsf{Tr}\big((\gamma_{*,N}\boldsymbol{I}_{N-1} - \boldsymbol{A}^{(2)}(\boldsymbol{m}))^{-1}\big) = N \cdot \left(1 - \|\boldsymbol{m}\|^2 / N\right) &, \\ \gamma_{*,N} > \lambda_{\max}(\boldsymbol{A}^{(2)}(\boldsymbol{m})) &. \end{cases}$$
(2.12)

Here I_{N-1} denotes the identity matrix acting on T_m , and the inverse is over quadratic forms on T_m .

Then we define

$$\Delta_{i}(\boldsymbol{m}) = \frac{1}{2} \langle \boldsymbol{A}^{(3)}(\boldsymbol{m}), \boldsymbol{Q}(\boldsymbol{m}) \otimes \boldsymbol{Q}(\boldsymbol{m})_{i,\cdot} \rangle = \frac{1}{2} \sum_{a,b,c=1}^{N} A^{(3)}_{abc}(\boldsymbol{m}) Q_{ia}(\boldsymbol{m}) Q_{bc}(\boldsymbol{m}), \qquad (2.13)$$

$$Q(m) := \left(\gamma_{*,N}(m)I_{N-1} - A^{(2)}(m)\right)^{-1}.$$
(2.14)

It is useful to make two additional remarks about the evaluation of $\Delta(m)$:

- 1. For any fixed \boldsymbol{m} , $\boldsymbol{A}^{(2)}(\boldsymbol{m}) \stackrel{d}{=} \sqrt{\xi''(\|\boldsymbol{m}\|_N^2) \cdot \frac{N-1}{N}} \boldsymbol{W}$, for $\boldsymbol{W} \sim \text{GOE}(N-1)$. It turns out that, although $\boldsymbol{m}^{\text{TAP}}$ is itself random, this nonetheless gives the correct asymptotics for $\gamma_{*,N}(\boldsymbol{m}^{\text{TAP}})$. Let $q_* = q_*(t)$ be the solution to $\frac{q_*}{1-q_*} = \xi'_t(q_*)$, existence and uniqueness of which is shown in Fact 4.2. We will show (see Proposition 4.4) that typically $\|\boldsymbol{m}^{\text{TAP}}\|_N^2 = q_* + o_N(1)$, and (see Lemma 6.22) $\gamma_{*,N}(\boldsymbol{m}^{\text{TAP}}) = \gamma_* + o_N(1)$, for $\gamma_* = (1-q_*)^{-1} + (1-q_*)\xi''(q_*)$. For the computation of $\boldsymbol{\Delta}$, we can replace $\gamma_{*,N}$ by γ_* with negligible error.
- 2. The tensors $A^{(2)}(m)$ and $A^{(3)}(m)$ can be written as explicit linear functions of the couplings g, and hence can be computed efficiently without need to take any numerical derivative.

2.4 Fundamental limits of algorithmic SL, replica symmetry breaking, and shattering

It is useful to compare condition (1.5) with the condition for (absence of) shattering, and replica symmetry breaking:

• As mentioned above (cf. Eq. (1.7)), it is conjectured [CHS93, CS95, BCKM98] that shattering is absent if and only if

$$\xi'(q) < \frac{q}{1-q}, \quad \forall q \in (0,1).$$
 (2.15)

This is implied by the condition under which our algorithm succeeds, namely Eq. (1.5), by integrating once.

• The tight condition for replica symmetry was identified in [Tal06, Proposition 2.3].

$$\xi(q) + q + \log(1 - q) \le 0, \quad \forall q \in [0, 1)$$
(2.16)

Note that this holds under (2.15) by integrating once, and hence under (1.5).

In this section, we prove that the condition (1.5) is necessary not only for Algorithm 2 to succeed, but indeed for a broader class of stochastic localization schemes that we next introduce. This points at a fundamental gap between such schemes and the possible computational limit for sampling, a fact that was suggested in [GDKZ23] and, in a related context, in [MRTS07].

By the key remark below (1.3), the process y_t generated by (1.3) consists of observations of some $\sigma \sim \mu_{H_N}$ through a progressively less noisy Gaussian channel. A natural generalization of this process outputs observations of $\sigma, \sigma^{\otimes 2}, \sigma^{\otimes 3}, \ldots$ through Gaussian channels of varying signal strengths, and can similarly be converted to a sampling algorithm.

Consider any $J \in \mathbb{N}$ and continuously differentiable, coordinate-wise increasing $\tau : [0, +\infty) \rightarrow [0, +\infty)^J$, normalized to $\|\tau(t)\|_1 = t$ for all $t \in [0, +\infty)$, and such that $\lim_{t\to\infty} \tau_j(t) = \infty$ for at least one odd $j \leq J$. For each $j \leq J$, let $(\boldsymbol{B}_t^j)_{t\geq 0}$ be a standard Brownian motion in $(\mathbb{R}^N)^{\otimes j}$. Let $(\vec{\boldsymbol{y}}_t)_{t\geq 0} = (\boldsymbol{y}_t^1, \ldots, \boldsymbol{y}_t^J) \in \mathbb{R}^N \times \cdots \times (\mathbb{R}^N)^{\otimes J}$ be given by the Ito diffusion

$$d\boldsymbol{y}_{t}^{j} = \tau_{j}'(t)\boldsymbol{m}_{j}(\boldsymbol{\vec{y}}_{t},t) dt + \tau_{j}'(t)^{1/2} d\boldsymbol{B}_{t}^{j}, \qquad \boldsymbol{\vec{y}}_{0} = \boldsymbol{0},$$
(2.17)

where, with expectation over $\boldsymbol{\sigma} \sim \mu_{H_N}$ and $\boldsymbol{G}^j \sim \mathcal{N}(0, \boldsymbol{I}_N^{\otimes j})$,

$$\boldsymbol{m}_{j}(\boldsymbol{\vec{y}}_{t},t) = \mathbb{E}[\boldsymbol{\sigma}^{\otimes j}|\boldsymbol{\tau}_{i}(t)\boldsymbol{\sigma}^{\otimes i} + \boldsymbol{\tau}_{i}(t)^{1/2}\boldsymbol{G}^{i} = \boldsymbol{y}_{t}^{i}, \forall 1 \leq i \leq J].$$
(2.18)

The process (1.3) corresponds to the case J = 1. As in that case, a sampling algorithm can be constructed from Eq. (2.17) by discretizing time and approximating the calculation of $m_i(\vec{y}_t, t)$ (see Remark 2.5 below).

For $\mathbf{A} \in (\mathbb{R}^N)^{\otimes j}$ and $1 \leq \ell \leq j$, let $\mathbf{A}^{(\ell)}$ be the tensor obtained by rotating coordinates by $i \pmod{j}$, that is

$$A_{i_1,...,i_j}^{(\ell)} = A_{i_{\ell+1},...,i_j,i_1,...,i_\ell}.$$

Then, for $\boldsymbol{B} \in (\mathbb{R}^N)^{\otimes j-1}$, let $(\boldsymbol{A}, \boldsymbol{B})_{\mathrm{sym}} \in \mathbb{R}^N$ be the vector satisfying

$$\langle \boldsymbol{v}, (\boldsymbol{A}, \boldsymbol{B})_{\mathrm{sym}} \rangle = \sum_{\ell=1}^{j} \langle \boldsymbol{B} \otimes \boldsymbol{v}, \boldsymbol{A}^{(\ell)} \rangle$$

for all $\boldsymbol{v} \in \mathbb{R}^N$. Let

$$\check{\xi}_t(s) = \xi(s) + \sum_{j=1}^J \tau_j(t) s^j$$

and define sequence $\{\check{q}_k : k \ge 0\}$ by $q_0 = 0$ and

$$\check{q}_{k+1} = \frac{\check{\xi}'_t(q_k)}{1 + \check{\xi}'_t(q_k)}$$
(2.19)

Finally define an AMP iteration analogous to (2.3) by

$$\widetilde{\boldsymbol{m}}^{k} = (1 - \widetilde{q}_{j})\widetilde{\boldsymbol{w}}^{k}, \quad \widetilde{\boldsymbol{w}}^{k+1} = \nabla H_{N}(\widetilde{\boldsymbol{m}}^{k}) + \sum_{j=1}^{J} \frac{1}{N^{j-1}} ((\widetilde{\boldsymbol{m}}^{k})^{\otimes j-1}, \boldsymbol{y}_{t}^{j})_{\text{sym}} - (1 - \widetilde{q}_{k})\widetilde{\boldsymbol{\xi}}''(\widetilde{q}_{k})\widetilde{\boldsymbol{m}}^{k-1}.$$
(2.20)

The next theorem is proved in Section 10, under the following condition which is a strict form of (2.16).

 $\xi''(0) < 1, \qquad \xi(q) + q + \log(1-q) < 0, \quad \forall q \in (0,1).$ (2.21)

Theorem 2.3. Suppose that (2.21) holds and that there exists $q \in [0,1)$ such that $\xi''(q) > \frac{1}{(1-q)^2}$. There exists a positive measure set $\mathcal{I} \subseteq [0, +\infty)$ such that for all $t \in \mathcal{I}$ the following holds. There exists $1 \leq j \leq J$ such that $\tau'_i(t) > 0$ and, for \vec{y}_t generated from (2.17),

$$\lim_{k\to\infty} \liminf_{N\to\infty} \mathbb{E}\frac{1}{N^j} \left\| (\widetilde{\boldsymbol{m}}^k)^{\otimes j} - \boldsymbol{m}_j(\vec{\boldsymbol{y}}_t, t) \right\|_2^2 > 0.$$

Remark 2.4. In this theorem we assume Eq. (2.21) to hold, but note that this an artifact of our proof technique. Indeed efficient sampling is believed to be impossible beyond the threshold (2.21). Indeed [AMS23b] implies that 'stable' algorithms fail under replica symmetry breaking.

Remark 2.5. As alluded to above, we can define a natural analog of Algorithm 1 for this generalized setting, which computes an estimator \widetilde{m}^{alg} for $m_1(\vec{y}_t, t)$. For some $K_{AMP} \in \mathbb{N}$, the point $\widetilde{m}^{K_{AMP}}$ is the result of the first phase of this algorithm. The output \widetilde{m}^{alg} of this algorithm satisfies $\|\widetilde{m}^{K_{AMP}} - \widetilde{m}^{alg}\|_N \to 0$ as $K_{AMP} \to \infty$; see Theorem 4.1 and Proposition 4.4 below, which show this for Algorithm 1 when (1.5) holds.

The analog of Algorithm 2 simulates the SDE (2.17) via an Euler discretization, estimating each $m_j(\vec{y}_t, t)$ with $(\widetilde{m}^{alg})^{\otimes j}$. Theorem 2.3 shows that for a interval of t of positive measure, this algorithm fails for a tensor order j relevant to the Euler discretization.

3 Preliminaries

In this section we provide further background. The contents of Subsections 3.1 and 3.2 are known and we often refer to [AMS22, Sections 3 and 4.1] for proofs. Subsection 3.4 introduces a lemma about conditioning a Gaussian process on a random vector: this is a fairly standard but crucial technical tool.

3.1 Stochastic localization

Fix a realization of H_N . The stochastic localization process is defined by the SDE (1.3), which has unique strong solutions provided $\mathbf{y} \mapsto \mathbf{m}(\mathbf{y}, t)$ is Lipschitz continuous. Note that, for μ_{H_N, \mathbf{y}_t} as in (2.1), \mathbf{m} is the mean

$$oldsymbol{m}(oldsymbol{y},t) = \int oldsymbol{\sigma} \ \mu_{H_N,oldsymbol{y}}(\mathrm{d}oldsymbol{\sigma}).$$

Therefore Lipschitz continuity is implied by $\sup_{\boldsymbol{y}} \|\operatorname{Cov}(\mu_{H_N,\boldsymbol{y}})\|_{op} < \infty$ which always holds since $\mu_{H_n,\boldsymbol{y}}$ is supported on a compact set.

As already mentioned in the introduction, we have the following facts (see for instance [AM22]).

Proposition 3.1. Let $(\mathbf{y}_t)_{t\geq 0}$ be the unique solution of the SDE (1.3). Then there exists a standard Brownian motion \mathbf{B}'_t independent of $\boldsymbol{\sigma} \sim \mu_{H_N}$, such that, for all t, $\mathbf{y}_t = t\boldsymbol{\sigma} + \mathbf{B}'_t$.

Further, $\mathbb{E} \operatorname{Cov}(\mu_{H_N, y_t}) \leq I_N / t$. In particular $\mu_{H_N, y_t} \Rightarrow \delta_{\sigma}$ almost surely as $t \to \infty$.

3.2 Planted model and contiguity

Recall that μ_0 denotes the uniform probability measure on S_N . Further, let \mathscr{H}_N be the space of Hamiltonians H_N (i.e. continuous functions $H_N : S_N \to \mathbb{R}$ endowed with the uniform convergence topology and the induced Borel sigma-algebra) and $\mu_{\text{null}} \in \mathcal{P}(\mathscr{H}_N)$ be the law induced on H_N by Eq. (1.1). Define the planted measure $\mu_{\text{pl}} \in \mathcal{P}(S_N \times \mathscr{H}_N)$ by

$$\mu_{\mathsf{pl}}(\mathsf{d}\boldsymbol{x},\mathsf{d}H_N) := \frac{1}{Z_{\mathsf{pl}}} \exp\left\{H_N(\boldsymbol{x})\right\} \mathsf{d}\mu_0(\boldsymbol{x}) \mathsf{d}\mu_{\mathsf{null}}(H_N).$$

For $H_N \in \mathscr{H}_N$, define the partition function

$$Z(H_N) := \int \exp \left\{ H_N(\boldsymbol{\sigma}) \right\} \mu_0(\mathsf{d}\boldsymbol{\sigma}) \,.$$

Lemma 3.2 (Proved in Section 8). Suppose ξ satisfies (2.21). Let $W \sim \mathcal{N}(-\frac{1}{2}\sigma^2, \sigma^2)$, where $\sigma^2 = -\frac{1}{2}\log(1-\xi''(0))$. As $N \to \infty$, for $H_N \sim \mu_{\text{null}}$, the Radon-Nykodym derivative of μ_{pl} with respect to μ_{null} is

$$\frac{\mathrm{d}\mu_{\mathsf{pl}}}{\mathrm{d}\mu_{\mathsf{null}}}(H_N) = \frac{Z(H_N)}{\mathbb{E}\,Z(H_N)} \xrightarrow{d} \exp(W).$$

Remark 3.3. In most of this paper, we are interested in ξ satisfying the condition (1.5), which implies (2.21) by integrating twice. However, the proof of Theorem 2.3 in Section 10 only assumes ξ satisfies (2.21), so we state this lemma with the more general condition.

For any T > 0, let $\mathbb{P}, \mathbb{Q} \in \mathcal{P}(S_N \times \mathscr{H}_N \times C([0,T], \mathbb{R}^N))$ be the laws of $(\boldsymbol{\sigma}, H_N, (\boldsymbol{y}_t)_{t \in [0,T]})$, generated as follows.

• Under Q:

$$H_N \sim \mu_{\text{null}}, \quad \boldsymbol{\sigma} \sim \mu_{H_N}, \quad \boldsymbol{y}_t = t\boldsymbol{\sigma} + \boldsymbol{B}_t, \quad (3.1)$$

for B_t a standard Brownian motion independent of σ , H_N . By Proposition 3.1, an equivalent description of this distribution is: $H_N \sim \mu_{\text{null}}$, $(y_t)_{t \ge 0}$ given by the SDE (1.3) and $\sigma = \lim_{t \to \infty} y_t/t$.

• Under \mathbb{P} :

$$(H_N, \boldsymbol{\sigma}) \sim \mu_{\mathsf{pl}}, \qquad \boldsymbol{y}_t = t\boldsymbol{\sigma} + \boldsymbol{B}_t, \qquad (3.2)$$

for B_t a standard Brownian motion independent of σ , H_N . As before, we can equivalently generate first H_N , then $(y_t)_{t\geq 0}$ given by the SDE (1.3) and finally σ .

The joint distribution of $(H_N, \sigma) \sim \mu_{pl}$ can be described in two equivalent ways. In the first one, we generate first H_N and then σ conditional on H_N :

$$H_N \sim \mu_{\mathsf{pl}}(\mathsf{d}H_N) = \frac{Z(H_N)}{\mathbb{E}Z(H_N)} \mu_{\mathsf{null}}(\mathsf{d}H_N), \qquad \boldsymbol{\sigma} \sim \mu_{H_N}.$$
(3.3)

In the second, we generate first σ and then H_N :

$$\boldsymbol{\sigma} \sim \mu_0, \qquad H_N \sim \mu_{\mathsf{pl}}(\mathsf{d}H_N|\boldsymbol{\sigma}) \propto e^{H_N(\boldsymbol{\sigma})} \mu_{\mathsf{null}}(\mathsf{d}H_N).$$
 (3.4)

A short calculation shows that $H_N \sim \mu_{pl}(\cdot | \boldsymbol{x})$ is given by

$$H_N(\boldsymbol{\sigma}) = N\xi(\langle \boldsymbol{x}, \boldsymbol{\sigma} \rangle_N) + \tilde{H}_N(\boldsymbol{\sigma}), \qquad (3.5)$$

where $\widetilde{H}_N \sim \mu_{\text{null}}$. The above definition has the following immediate consequence.

Proposition 3.4 ([AMS22, Proposition 4.2]). For all $T \ge 0$,

$$\frac{\mathsf{d}\mathbb{P}}{\mathsf{d}\mathbb{Q}}(\boldsymbol{\sigma}, H_N, (\boldsymbol{y}_t)_{t\in[0,T]}) = \frac{Z(H_N)}{\mathbb{E}\,Z(H_N)}.$$

As a consequence of Lemma 3.2 and Proposition 3.4, Le Cam's first lemma implies the following.

Corollary 3.5. The measures \mathbb{P} and \mathbb{Q} are mutually contiguous. That is, for any sequence of events \mathcal{E}_N , $\mathbb{P}(\mathcal{E}_N) \to 0$ if and only if $\mathbb{Q}(\mathcal{E}_N) \to 0$.

Thus it suffices to analyze our algorithm under the planted distribution \mathbb{P} .

3.3 Basic regularity estimate

For a tensor $\boldsymbol{A} \in (\mathbb{R}^N)^{\otimes k}$, define the operator norm

$$\|\boldsymbol{A}\|_{\mathrm{op},N} = rac{1}{N} \sup_{\|\boldsymbol{\sigma}^1\|_N,...,\|\boldsymbol{\sigma}^k\|_N \leqslant 1} |\langle \boldsymbol{A}, \boldsymbol{\sigma}^1 \otimes \cdots \otimes \boldsymbol{\sigma}^k
angle|.$$

Notice that this normalization is different from the standard injective norm $\|\cdot\|_{ini}$ in that $\|A\|_{op,N} = N^{(k-2)/2} \|A\|_{ini}$.

Proposition 3.6 ([HS22, Proposition 2.3]). There exists a sequence of constants $(C_k)_{k\geq 0}$ independent of N for which the following holds. Define the event

$$K_N := \left\{ \sup_{\|\boldsymbol{\sigma}\|_N \leqslant 1} \|\nabla^k H_N(\boldsymbol{\sigma})\|_{\text{op},N} \leqslant C_k \ \forall k \ge 0 \right\}$$

Then $\mathbb{P}(K_N) \ge 1 - e^{-cN}$.

3.4 Conditioning lemma

Lemma 3.7. Let $D \subseteq \mathbb{R}^N$ be an open set and $\mathcal{F} : D \to \mathbb{R}$ be a (not necessarily centered) C^2 Gaussian process on a probability space $(\Omega, \Sigma, \mathbb{P})$. Let X be a random variable on (Ω, Σ) taking values in [0, 1], and \mathbf{m}_0 be a random vector on the same space taking values in \mathbb{R}^N . For ε , c_{spec} , $c_{\text{op}} > 0$ satisfying $\varepsilon \leq c_{\text{spec}}^2/10c_{\text{op}}$, define $U_{\mathbf{m}_0} := \mathsf{B}_N(\mathbf{m}_0, 5\varepsilon/c_{\text{spec}})$ and the events

$$\begin{split} \mathcal{G}(\varepsilon, c_{\mathsf{spec}}) &:= \left\{ \|\nabla \mathcal{F}(\boldsymbol{m}_0)\|_N \leqslant \varepsilon \,, \quad \nabla^2 \mathcal{F}(\boldsymbol{m}_0) \leq -c_{\mathsf{spec}} \boldsymbol{I}_n \right\}, \\ \mathcal{H}(c_{\mathsf{op}}) &:= \left\{ \sup_{\boldsymbol{m} \in D} \|\nabla^2 \mathcal{F}(\boldsymbol{m})\|_{\mathsf{op},N} \leqslant c_{\mathsf{op}}, \quad \sup_{\boldsymbol{m} \in D} \|\nabla^3 \mathcal{F}(\boldsymbol{m})\|_{\mathsf{op},N} \leqslant c_{\mathsf{op}} \right\}, \\ \mathcal{E}_{\mathsf{cond}} &:= \mathcal{G}(\varepsilon, c_{\mathsf{spec}}) \cap \mathcal{H}(c_{\mathsf{op}}) \cap \{\|\boldsymbol{m}_0\|_N \leqslant 1\} \cap \{U_{\boldsymbol{m}_0} \subseteq D\} \,. \end{split}$$

Finally, assume $\mathbf{m} \mapsto \mathbb{E} \nabla \mathcal{F}(\mathbf{m})$ is continuous and $\lambda_{\min}(\operatorname{Cov}(\nabla \mathcal{F}(\mathbf{m})))$ is bounded away from 0 uniformly over $\mathbf{m} \in D$. Then, with $\varphi_{\nabla \mathcal{F}(\mathbf{m})}$ the probability density of $\nabla \mathcal{F}(\mathbf{m})$ w.r.t. Lebesgue measure on \mathbb{R}^N and d^N denoting integration against this measure,

$$\mathbb{E}(X\mathbf{1}\{\mathcal{E}_{\mathsf{cond}}\}) = \int_D \mathbb{E}\left[|\det \nabla^2 \mathcal{F}(\boldsymbol{m})| X\mathbf{1}\{\mathcal{E}_{\mathsf{cond}} \cap \{\boldsymbol{m} \in U_{\boldsymbol{m}_0}\}\} \middle| \nabla \mathcal{F}(\boldsymbol{m}) = \mathbf{0} \right] \varphi_{\nabla \mathcal{F}(\boldsymbol{m})}(\mathbf{0}) \, \mathsf{d}^N \boldsymbol{m}.$$

Proof. On event \mathcal{E}_{cond} , for all $m \in U_{m_0}$ we have

$$\lambda_{\max}(\nabla^{2}\mathcal{F}(\boldsymbol{m})) \leq \lambda_{\max}(\nabla^{2}\mathcal{F}(\boldsymbol{m}_{0})) + c_{\text{op}} \|\boldsymbol{m} - \boldsymbol{m}_{0}\|_{N} \leq -c_{\text{spec}} + \frac{5\varepsilon c_{\text{op}}}{c_{\text{spec}}} \leq -\frac{1}{2}c_{\text{spec}}.$$
 (3.6)

Since $\|\nabla \mathcal{F}(\boldsymbol{m}_0)\|_N \leq \varepsilon$, there is exactly one solution to $\nabla \mathcal{F}(\boldsymbol{m}_*) = \mathbf{0}$ in $U_{\boldsymbol{m}_0}$, which is measurable on (Ω, Σ) and furthermore lies in $B_N(\boldsymbol{m}_0, 4\varepsilon/c_{spec})$. The strong concavity (3.6) implies that $\nabla \mathcal{F}$ is injective on $U_{\boldsymbol{m}_0}$ and its image contains a neighborhood of **0**. By the area formula, for sufficiently small $\iota > 0$,

$$1 = \frac{1}{|\mathsf{B}_N(\mathbf{0},\iota)|} \int_{U_{\boldsymbol{m}_0}} |\det \nabla^2 \mathcal{F}(\boldsymbol{m})| \mathbf{1} \{ \|\nabla \mathcal{F}(\boldsymbol{m})\|_N \leqslant \iota \} \, \mathsf{d}^N \boldsymbol{m}$$

Multiplying by $X1{\mathcal{E}_{cond}}$ and taking expectations of both sides by Fubini yields

$$\begin{split} & \mathbb{E}(X\mathbf{1}\{\mathcal{E}_{\mathsf{cond}}\}) \\ &= \frac{1}{|\mathsf{B}_N(\mathbf{0},\iota)|} \int_D \mathbb{E}\left[|\det \nabla^2 \mathcal{F}(\boldsymbol{m})| X\mathbf{1}\{\mathcal{E}_{\mathsf{cond}} \cap \{\boldsymbol{m} \in U_{\boldsymbol{m}_0}\} \cap \{\|\nabla \mathcal{F}(\boldsymbol{m})\|_N \leqslant \iota\}\} \right] \, \mathsf{d}^N \boldsymbol{m} \\ &= \int_D \mathbb{E}\left[|\det \nabla^2 \mathcal{F}(\boldsymbol{m})| X\mathbf{1}\{\mathcal{E}_{\mathsf{cond}} \cap \{\boldsymbol{m} \in U_{\boldsymbol{m}_0}\}\} \right] \|\nabla \mathcal{F}(\boldsymbol{m})\|_N \leqslant \iota \right] \frac{\mathbb{P}(\|\nabla \mathcal{F}(\boldsymbol{m})\|_N \leqslant \iota)}{|\mathsf{B}_N(\mathbf{0},\iota)|} \, \mathsf{d}^N \boldsymbol{m} \end{split}$$

Note that on \mathcal{E}_{cond} , $|\det \nabla^2 \mathcal{F}(\boldsymbol{m})| \leq c_{op}^N$. Since \mathcal{E}_{cond} is contained in the event $\|\boldsymbol{m}_0\|_N \leq 1$, $\{\boldsymbol{m} \in U_{\boldsymbol{m}_0}\}$ can only occur for \boldsymbol{m} on a bounded set. Since $\lambda_{\min}(\operatorname{Cov}(\nabla \mathcal{F}(\boldsymbol{m})))$ is bounded away from 0, $\varphi_{\nabla \mathcal{F}(\boldsymbol{m})}$ is bounded, and thus so is $\mathbb{P}(\|\nabla \mathcal{F}(\boldsymbol{m})\|_N \leq \iota)/|B_N(\boldsymbol{0},\iota)|$. Therefore the integral in the last display is dominated by a bounded integrable function. Continuity of $\mathbb{E} \nabla \mathcal{F}(\boldsymbol{m})$ implies that $\varphi_{\nabla \mathcal{F}(\boldsymbol{m})}(\boldsymbol{z})$ is continuous in \boldsymbol{z} in a neighborhood of $\boldsymbol{0}$. We take the $\iota \to 0$ limit of the last display by dominated convergence to conclude.

4 Analysis of mean computation algorithm

The next several sections are devoted to the analysis of Algorithm 1. We fix $t \in [0, T]$ and consider $(\boldsymbol{x}, H_N, (\boldsymbol{y}_t)_{t \ge 0}) \in S_N \times \mathscr{H}_N \times C([0, T], \mathbb{R}^N)$ distributed according to the planted law \mathbb{P} defined in Eq. (3.2). Define

$$H_{N,t}(\boldsymbol{\sigma}) = H_N(\boldsymbol{\sigma}) + \langle \boldsymbol{y}_t, \boldsymbol{\sigma} \rangle$$

$$= N\xi(\langle \boldsymbol{x}, \boldsymbol{\sigma} \rangle_N) + \widetilde{H}_N(\boldsymbol{\sigma}) + \langle \boldsymbol{y}_t, \boldsymbol{\sigma} \rangle.$$
(4.1)

where we recall $\widetilde{H}_N(\boldsymbol{\sigma}) \sim \mu_{\text{null}}$. The tilted measure $\mu_t = \mu_{H_N, \boldsymbol{y}_t}$ defined in (2.1) has the form

$$\mu_t(\mathsf{d}\boldsymbol{\sigma}) = \frac{1}{Z} \exp H_{N,t}(\boldsymbol{\sigma}) \ \mu_0(\mathsf{d}\boldsymbol{\sigma}).$$

Let m_t be the mean of μ_t . The main result of our analysis is the following.

Theorem 4.1. Under condition (1.5), there exist parameters $(K_{AMP}, K_{GD}^*, \eta)$ depending only on (ξ, t) such that the point \mathbf{m}^{alg} output by Algorithm 1 on input (H_N, \mathbf{y}_t) , with parameters K_{AMP} , $K_{GD}(N) = K_{GD}^* \log N$, η satisfies

$$\mathbb{E} \|\boldsymbol{m}^{\mathsf{alg}} - \boldsymbol{m}_t\|_N^2 = o(N^{-1}).$$

Recall that we defined $\xi_t(q) = \xi(q) + tq$.

Fact 4.2. For any $t \in [0, \infty)$, there is a unique solution $q_* = q_*(t) \in [0, 1)$ to

$$\xi_t'(q) = \frac{q}{1-q}.$$
(4.2)

Proof. Define $f(q) = \xi'_t(q) - \frac{q}{1-q}$. Since f(0) = t > 0 and $\lim_{q \to 1^-} f(q) = -\infty$, there is at least one solution. As

$$\frac{\mathsf{d}}{\mathsf{d}q}\left(\xi_t'(q) - \frac{q}{1-q}\right) = \xi''(q) - \frac{1}{(1-q)^2} \stackrel{(1.5)}{<} 0,$$

this solution is unique.

Henceforth let q_* denote this solution. It will also be useful to rewrite (4.1) as

$$H_{N,t}(\boldsymbol{\sigma}) = N\xi_t(\langle \boldsymbol{x}, \boldsymbol{\sigma} \rangle_N) + H_{N,t}(\boldsymbol{\sigma}), \qquad (4.3)$$

where

$$\widetilde{H}_{N,t}(\boldsymbol{\sigma}) = \widetilde{H}_N(\boldsymbol{\sigma}) + \langle \boldsymbol{B}_t, \boldsymbol{\sigma} \rangle$$
(4.4)

is a spin glass with mixture ξ_t . In the proofs below, we will switch between these two representations of $H_{N,t}$ as convenient.

The first step of our analysis characterizes the limiting performance of the AMP iteration (2.3), on (H_N, y_t) generated from the planted process (3.2). Recall the TAP free energy \mathcal{F}_{TAP} introduced in (2.5). With the notation (4.3), we can write

$$\mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = N\xi_t(\langle \boldsymbol{x}, \boldsymbol{m} \rangle_N) + \widetilde{H}_{N,t}(\boldsymbol{m}) + \frac{N}{2}\theta(\|\boldsymbol{m}\|_N^2) + \frac{N}{2}\log(1 - \|\boldsymbol{m}\|_N^2).$$

Proposition 4.3. For any $\iota > 0$, there exists $k_0 \in \mathbb{N}$, depending only on (ξ, t, ι) , such that for any fixed k, $k \ge k_0$ the following holds with probability $1 - e^{-cN}$. The AMP iterate \mathbf{m}^k satisfies

$$|\langle \boldsymbol{x}, \boldsymbol{m}^k \rangle_N - q_*|, |\langle \boldsymbol{m}^k, \boldsymbol{m}^k \rangle_N - q_*| \leq \iota$$
 (4.5)

and

$$\left\|\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}^{k})\right\|_{N}, \left\|\nabla \widetilde{H}_{N,t}(\boldsymbol{m}^{k}) + \xi_{t}'(q_{*})\boldsymbol{x} - \left((1-q_{*})\xi''(q_{*}) + \frac{1}{1-q_{*}}\right)\boldsymbol{m}^{k}\right\|_{N} \leqslant \iota.$$
(4.6)

Moreover, with $I = I(\iota) = [q_* - \iota, q_* + \iota]$,

$$\mu_t(\mathsf{Band}(\boldsymbol{m}^k, I) \cap \mathsf{Band}(\boldsymbol{x}, I)) \ge 1 - e^{-cN}.$$
(4.7)

The proof of this proposition is presented in Section 5. For $\iota > 0$, define

$$\mathcal{S}_{\iota} := \left\{ \boldsymbol{m} \in \mathbb{R}^{N} : |\langle \boldsymbol{m}, \boldsymbol{x} \rangle_{N} - q_{*}|, |\langle \boldsymbol{m}, \boldsymbol{m} \rangle_{N} - q_{*}| \leq \iota \right\}.$$
(4.8)

Proposition 4.4. There exist $C_{\max}^{\text{spec}} > C_{\min}^{\text{spec}} > 0$ and L > 0 such that, for any sufficiently small $\iota > 0$, there is an event \mathcal{E}_0 with probability $1 - e^{-cN}$, on which the following holds.

- (a) The event K_N from Proposition 3.6 holds.
- (b) \mathcal{F}_{TAP} has a unique critical point m^{TAP} in \mathcal{S}_{ι} , which further satisfies

$$\operatorname{spec}(\nabla^2 \mathcal{F}_{\operatorname{TAP}}(\boldsymbol{m}^{\operatorname{TAP}})) \subseteq [-C_{\max}^{\operatorname{spec}}, -C_{\min}^{\operatorname{spec}}].$$
 (4.9)

(c) For K_{AMP} large enough (depending on ι), we have $\boldsymbol{m}^{\text{AMP}} \in S_{\iota/2}$ and $\|\boldsymbol{m}^{\text{AMP}} - \boldsymbol{m}^{\text{TAP}}\|_N \leq \iota/2$.

Note that under (a), there exists c_{op} such that $\|\nabla^2 \mathcal{F}_{TAP}(\boldsymbol{m})\|_{op,N}$, $\|\nabla^3 \mathcal{F}_{TAP}(\boldsymbol{m})\|_{op,N} \leq c_{op}$ uniformly over $\boldsymbol{m} \in \mathcal{S}_{\iota}$, for all sufficiently small $\iota > 0$. Let

$$\varepsilon = \min\left(\frac{\iota c_{\rm op}}{10}, \frac{(C_{\rm min}^{\rm spec})^2}{40c_{\rm op}}\right).$$
(4.10)

Let $\mathcal{E} = \mathcal{E}_0 \cap \{ \| \nabla \mathcal{F}_{TAP}(\boldsymbol{m}^{AMP}) \|_N \leq \varepsilon \}$. (For K_{AMP} large enough, this holds with probability $1 - e^{-cN}$ by Proposition 4.3.) We further have:

(d) For any $\delta > 0$ there exists $C_{\delta} > 0$ such that the following holds. For any random variable X with $0 \leq X \leq 1$ almost surely,

$$\mathbb{E}[X\mathbf{1}\{\mathcal{E}\}] \leqslant C_{\delta} \sup_{\boldsymbol{m}\in\mathcal{S}_{\iota}} \mathbb{E}\left[X^{1+\delta}\mathbf{1}\{\mathcal{E}\} \middle| \nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \mathbf{0}\right]^{1/(1+\delta)}.$$

Proposition 4.5. For sufficiently small $\iota > 0$, with probability $1 - e^{-cN}$, the event \mathcal{E} from Proposition 4.4 holds and:

(a) For $I = I(\iota)$ as above, we have

$$\mu_t(\mathsf{Band}(oldsymbol{m}^{ t{TAP}},I)\cap\mathsf{Band}(oldsymbol{x},I))\geqslant 1-e^{-cN}.$$

(b) For η small enough and K^*_{GD} large enough, we have $\|\boldsymbol{m}^{\mathsf{GD}} - \boldsymbol{m}^{\mathsf{TAP}}\|_N \leq N^{-10}$.

(c) For any $\boldsymbol{m}_1, \boldsymbol{m}_2 \in \mathsf{B}_N(\boldsymbol{m}^{\mathsf{TAP}}, \iota)$, we have $\|\boldsymbol{\Delta}(\boldsymbol{m}_1) - \boldsymbol{\Delta}(\boldsymbol{m}_2)\|_N \leq \frac{L}{N} \|\boldsymbol{m}_1 - \boldsymbol{m}_2\|_N$.

The proofs of the last two propositions are given in Section 6. For $\iota > 0$, define the truncated magnetization

$$\widetilde{\boldsymbol{m}}_{\iota}(\boldsymbol{m}) = \frac{\int_{\mathsf{Band}(\boldsymbol{m},I(\iota)) \cap \mathsf{Band}(\boldsymbol{x},I(\iota))} \boldsymbol{\sigma} \exp(H_{N,t}(\boldsymbol{\sigma})) \ \mu_0(\mathsf{d}\boldsymbol{\sigma})}{\int_{\mathsf{Band}(\boldsymbol{m},I(\iota)) \cap \mathsf{Band}(\boldsymbol{x},I(\iota))} \exp(H_{N,t}(\boldsymbol{\sigma})) \ \mu_0(\mathsf{d}\boldsymbol{\sigma})}.$$

Proposition 4.6. Let $\Delta(\cdot)$ be defined as in Section 2.3. Then, for sufficiently small $\iota, \delta > 0$, we have

$$\sup_{\boldsymbol{m}\in\mathcal{S}_{\iota}}\mathbb{E}\left[\|\boldsymbol{m}+\boldsymbol{\Delta}(\boldsymbol{m})-\widetilde{\boldsymbol{m}}_{2\iota}(\boldsymbol{m})\|_{N}^{2+\delta}\big|\nabla\mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})=\boldsymbol{0}\right]\leqslant N^{-(1+\delta)}$$

The proof of this proposition is given in Section 7.

Proof of Theorem 4.1. Let \mathcal{E}_1 be the intersection of \mathcal{E} from Proposition 4.4 and the event in Proposition 4.5. On \mathcal{E}_1 , the point m^{TAP} is well-defined and we can write

$$egin{aligned} m^{\mathsf{alg}} - m_t &= m^{\mathsf{GD}} + \Delta(m^{\mathsf{GD}}) - m_t \ &= (m^{\mathsf{GD}} - m^{\mathsf{TAP}}) + (\Delta(m^{\mathsf{GD}}) - \Delta(m^{\mathsf{TAP}})) + (\widetilde{m}_{2\iota}(m^{\mathsf{TAP}}) - m_t) \ &+ (m^{\mathsf{TAP}} + \Delta(m^{\mathsf{TAP}}) - \widetilde{m}_{2\iota}(m^{\mathsf{TAP}})) \,, \end{aligned}$$

whence

$$\begin{split} \|\boldsymbol{m}^{\mathsf{alg}} - \boldsymbol{m}_t\|_N^2 &\leqslant 4 \|\boldsymbol{m}^{\mathsf{GD}} - \boldsymbol{m}^{\mathsf{TAP}}\|_N^2 + 4 \|\boldsymbol{\Delta}(\boldsymbol{m}^{\mathsf{GD}}) - \boldsymbol{\Delta}(\boldsymbol{m}^{\mathsf{TAP}})\|_N^2 + 4 \|\widetilde{\boldsymbol{m}}_{2\iota}(\boldsymbol{m}^{\mathsf{TAP}}) - \boldsymbol{m}_t\|_N^2 \\ &+ 4 \|\boldsymbol{m}^{\mathsf{TAP}} + \boldsymbol{\Delta}(\boldsymbol{m}^{\mathsf{TAP}}) - \widetilde{\boldsymbol{m}}_{2\iota}(\boldsymbol{m}^{\mathsf{TAP}})\|_N^2 \end{split}$$

The following also holds on \mathcal{E}_1 . By Proposition 4.5(b) and 4.5(c), for some constant C (changing from line to line below),

$$\|\boldsymbol{m}^{\mathsf{GD}} - \boldsymbol{m}^{\mathsf{TAP}}\|_{N}^{2}, \|\boldsymbol{\Delta}(\boldsymbol{m}^{\mathsf{GD}}) - \boldsymbol{\Delta}(\boldsymbol{m}^{\mathsf{TAP}})\|_{N}^{2} \leqslant CN^{-20}$$

By Proposition 4.4(a), the complement of $Band(m^{TAP}, I) \cap Band(x, I)$ accounts for a e^{-cN} fraction of the Gibbs measure. Because the spins σ are bounded, this implies

$$\|\widetilde{\boldsymbol{m}}_{2\iota}(\boldsymbol{m}^{\mathrm{TAP}}) - \boldsymbol{m}_t\|_N^2 \leqslant e^{-cN}$$

Therefore, on \mathcal{E}_1 , for all sufficiently large N

$$\|\boldsymbol{m}^{\mathsf{alg}} - \boldsymbol{m}_t\|_N^2 \leq CN^{-20} + 4\|\boldsymbol{m}^{\mathsf{TAP}} + \boldsymbol{\Delta}(\boldsymbol{m}^{\mathsf{TAP}}) - \widetilde{\boldsymbol{m}}_{2\iota}(\boldsymbol{m}^{\mathsf{TAP}})\|_N^2.$$

Thus

$$\begin{split} \mathbb{E}[\|\boldsymbol{m}^{\mathsf{alg}} - \boldsymbol{m}_t\|_N^2] &\leq \mathbb{P}(\mathcal{E}_1^c) + \mathbb{E}[\|\boldsymbol{m}^{\mathsf{alg}} - \boldsymbol{m}_t\|_N^2 \mathbf{1}\{\mathcal{E}_1\}] \\ &\leq CN^{-20} + 4 \mathbb{E}\left[\|\boldsymbol{m}^{\mathsf{TAP}} + \boldsymbol{\Delta}(\boldsymbol{m}^{\mathsf{TAP}}) - \widetilde{\boldsymbol{m}}_{2\iota}(\boldsymbol{m}^{\mathsf{TAP}})\|_N^2 \mathbf{1}\{\mathcal{E}_1\}\right] \\ &\leq CN^{-20} + 4C_{\delta/2} \sup_{\boldsymbol{m}\in\mathcal{S}_\iota} \mathbb{E}\left[\|\boldsymbol{m} + \boldsymbol{\Delta}(\boldsymbol{m}) - \widetilde{\boldsymbol{m}}_{2\iota}(\boldsymbol{m})\|_N^{2+\delta} \mathbf{1}\{\mathcal{E}_1\} \middle| \nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \mathbf{0} \right]^{1/(1+\delta/2)} \\ &\leq CN^{-20} + 4C_{\delta/2}N^{-(1+\delta)/(1+\delta/2)} = o(N^{-1}). \end{split}$$

In the second-last line, we applied Proposition 4.4(d), noting that on \mathcal{E}_1 and conditioned on $\nabla \mathcal{F}_{TAP}(m) = 0$, we have $m^{TAP} = m$ almost surely. The last line is Proposition 4.6.

5 Analysis of AMP iteration: proof of Proposition 4.3

5.1 State evolution limit

We first prove (4.5) and (4.6) using the state evolution result of [Bol14, BM11, JM13]. Recalling the change of notation (4.3), the AMP iteration (2.3) can be rewritten as $m^{-1} = w^0 = 0$,

$$\boldsymbol{m}^{k} = (1 - q_{k})\boldsymbol{w}^{k},$$

$$\boldsymbol{w}^{k+1} = \nabla H_{N,t}(\boldsymbol{m}^{k}) - (1 - q_{k})\boldsymbol{\xi}''(q_{k})\boldsymbol{m}^{k-1}$$

$$= \nabla \widetilde{H}_{N,t}(\boldsymbol{m}^{k}) + \boldsymbol{\xi}'_{t}(\langle \boldsymbol{x}, \boldsymbol{m}^{k} \rangle_{N})\boldsymbol{x} - (1 - q_{k})\boldsymbol{\xi}''(q_{k})\boldsymbol{m}^{k-1}.$$
(5.1)

Here and below, the sequence $(q_k)_{k\geq 0}$ is defined as per Eq. (2.2).

Set $\gamma_0 = \Sigma_{0,i} = \Sigma_{i,0} = 0$ for all $i \ge 0$, and define the following recurrence. Sample $X \sim \mathcal{N}(0,1)$ and, for $k \ge 0$,

$$(G_1, \ldots, G_k) \sim \mathcal{N}(0, \Sigma_{\leq k}), \qquad W_i = G_i + \gamma_i X.$$

Then, let

$$\gamma_{k+1} = \xi_t'((1 - q_k)\gamma_k)$$
(5.2)

$$\Sigma_{k+1,j+1} = \xi'_t \left((1 - q_k)(1 - q_j) \mathbb{E}[W_k W_j] \right).$$
(5.3)

The following proposition is an immediate consequence of [EAMS21, Proposition 3.1], which generalizes to the tensor case [BM11, Theorem 1].

Proposition 5.1. For any $k \ge 0$, the empirical distribution of the AMP iterates' coordinates converges in W_2 in probability:

$$\frac{1}{N}\sum_{i=1}^N \delta_{x_i,w_i^1,\dots,w_i^k} \xrightarrow{W_2} \mathcal{L}(X,W_1,\dots,W_k).$$

(In words, the left-hand side is the probability distribution on \mathbb{R}^{k+1} that puts mass 1/N on each point $(x_i, w_i^1, \ldots, w_i^k)$, for $i \in [N]$.)

Lemma 5.2. For all $k, j \ge 0$, we have $\Sigma_{k,j} = \gamma_{k \land j} = \frac{q_{k \land j}}{1 - q_{k \land j}}$.

Proof. We first prove by induction that $\gamma_k = \frac{q_k}{1-q_k}$. For k = 0 this is clear, and then by induction

$$\gamma_{k+1} = \xi'_t(q_k) = \frac{q_{k+1}}{1 - q_{k+1}}.$$

Similarly, by induction

$$(1 - q_k)(1 - q_j) \mathbb{E}[W_k W_j] = (1 - q_k)(1 - q_j) (\Sigma_{k,j} + \gamma_k \gamma_j) = (1 - q_{k \lor j})q_{k \land j} + q_k q_j = q_{k \land j},$$

and thus

$$\Sigma_{k+1,j+1} = \xi'_t(q_{k\wedge j}) = \frac{q_{k\wedge j+1}}{1 - q_{k\wedge j+1}}.$$

Lemma 5.3. As $k \to \infty$, we have $q_k \to q_*$.

Proof. Since the function $f(q) = \frac{\xi'_t(q)}{1+\xi'_t(q)}$ is increasing, with f(0) > 0, f(1) < 1, q_k must converge to a solution of q = f(q). This rearranges to $\xi'_t(q) = \frac{q}{1-q}$, which has unique solution q_* by Fact 4.2.

Proposition 5.4. With probability $1 - e^{-cN}$, (4.5) and (4.6) hold for all $k \ge k_0$.

Proof. Let \simeq denote equality up to an additive error $o_{P,N}(1)$ (a term vanishing in probability as $N \to \infty$). By Proposition 5.1,

$$\langle \boldsymbol{x}, \boldsymbol{m}^k \rangle_N = (1 - q_k) \langle \boldsymbol{x}, \boldsymbol{w}^k \rangle_N \simeq (1 - q_k) \gamma_k = q_k.$$
 (5.4)

Moreover,

$$\langle \boldsymbol{m}^{k}, \boldsymbol{m}^{k} \rangle_{N} = (1 - q_{k})^{2} \langle \boldsymbol{w}^{k}, \boldsymbol{w}^{k} \rangle_{N} \simeq (1 - q_{k})^{2} \left(\Sigma_{k,k} + \gamma_{k}^{2} \right) = q_{k}.$$
 (5.5)

By Lemma 5.3, for all k large enough we have $|q_k - q_*| \le \iota/3$, whence (4.5) holds with high probability. Rearranging the AMP iteration gives

$$\nabla \widetilde{H}_{N,t}(\boldsymbol{m}^{k}) = -\xi_{t}'(\langle \boldsymbol{x}, \boldsymbol{m}^{k} \rangle_{N})\boldsymbol{x} + \boldsymbol{w}^{k+1} + (1 - q_{k})\xi''(q_{k})\boldsymbol{m}^{k-1}$$
$$= -\xi_{t}'(\langle \boldsymbol{x}, \boldsymbol{m}^{k} \rangle_{N})\boldsymbol{x} + \frac{1}{1 - q_{k+1}}\boldsymbol{m}^{k+1} + (1 - q_{k})\xi''(q_{k})\boldsymbol{m}^{k-1},$$
(5.6)

By Proposition 5.1, Lemma 5.2, and Lemma 5.3, we have

$$\lim_{k \to \infty} \operatorname{p-lim}_{N \to \infty} \|\boldsymbol{m}^{k+1} - \boldsymbol{m}^{k}\| = 0, \qquad (5.7)$$

$$\lim_{k \to \infty} \operatorname{p-lim}_{N \to \infty} \| \boldsymbol{w}^{k+1} - \boldsymbol{w}^k \| = 0.$$
(5.8)

and therefore, by Eq. (5.6),

$$\lim_{k \to \infty} \operatorname{p-lim}_{N \to \infty} \left\| \nabla \widetilde{H}_{N,t}(\boldsymbol{m}^k) + \xi'_t(q_*) \boldsymbol{x} + \left(\frac{1}{1 - q_*} + (1 - q_*) \xi''(q_*) \right) \boldsymbol{m}^k \right\|_N = 0.$$

As

$$\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \nabla \widetilde{H}_{N,t}(\boldsymbol{m}) + \xi_t'(\langle \boldsymbol{x}, \boldsymbol{m} \rangle_N)\boldsymbol{x} + \left(\frac{1}{1 - \|\boldsymbol{m}\|_N^2} + (1 - \|\boldsymbol{m}\|_N^2)\xi''(\|\boldsymbol{m}\|_N^2)\right)\boldsymbol{m},$$
(5.9)

equations (5.4), (5.5) further imply

$$\lim_{k\to\infty} \operatorname{p-lim}_{N\to\infty} \|\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}^k)\|_N = 0.$$

Thus, for large enough k, (4.6) holds with high probability.

To improve these assertions to $1 - e^{-cN}$ probability, note that by [HS22, Section 8], the AMP iterate m^k is, on an event \mathcal{E}_{Lip} with probability $1 - e^{-cN}$, a O(1)-Lipschitz function of the disorder Gaussians in $\widetilde{H}_{N,t}$. By Kirszbraun's extension theorem, there is a measurable, O(1)-Lipschitz function \widetilde{m}^k of the disorder which agrees with m^k on \mathcal{E}_{Lip} . Thus $\langle x, \widetilde{m}^k \rangle_N$ and $\langle \widetilde{m}^k, \widetilde{m}^k \rangle_N$ are $O(N^{-1/2})$ -Lipschitz in the disorder. By Gaussian concentration of measure

$$|\langle \boldsymbol{x}, \widetilde{\boldsymbol{m}}^k \rangle_N - \mathbb{E} \langle \boldsymbol{x}, \widetilde{\boldsymbol{m}}^k \rangle_N |, |\langle \widetilde{\boldsymbol{m}}^k, \widetilde{\boldsymbol{m}}^k \rangle_N - \mathbb{E} \langle \widetilde{\boldsymbol{m}}^k, \widetilde{\boldsymbol{m}}^k \rangle_N | \leq \iota/3$$

with probability $1 - e^{-cN}$. Since $\mathbf{m}^k = \widetilde{\mathbf{m}}^k$ on \mathcal{E}_{Lip} , (4.5) holds with probability $1 - e^{-cN}$.

By Proposition 3.6, $\mathbf{m} \mapsto \nabla \widetilde{H}_{N,t}(\mathbf{m})$ is also O(1)-Lipschitz over $\|\mathbf{m}\|_N \leq 1$ with probability $1 - e^{-cN}$. A similar argument shows that (4.6) holds with probability $1 - e^{-cN}$.

5.2 Overlap with AMP iterates

The following proposition constitutes the first half of the proof of Eq. (4.7).

Proposition 5.5. Let $\iota > 0$ and $I = I(\iota)$. With probability $1 - e^{-cN}$, for all $k \ge k_0$ (with k_0 a sufficiently large constant depending on (ξ, t, ι)),

$$\mu_t(\mathsf{Band}(\boldsymbol{m}^k, I)) \ge 1 - e^{-cN}$$

To prove Proposition 5.5, we will combine Lemma 5.7 below, which identifies a band on which the Gibbs measure μ_t concentrates, with a self-reduction argument. We return to the earlier representation (4.1) of $H_{N,t}$, which we reproduce below.

$$\begin{split} H_{N,t}(\boldsymbol{\sigma}) &= H_N(\boldsymbol{\sigma}) + \langle \boldsymbol{y}_t, \boldsymbol{\sigma} \rangle, \quad \text{where} \\ H_N(\boldsymbol{\sigma}) &= N\xi(\langle \boldsymbol{x}, \boldsymbol{\sigma} \rangle_N) + \widetilde{H}_N(\boldsymbol{\sigma}), \\ \boldsymbol{y}_t &= t\boldsymbol{x} + \sqrt{t}\boldsymbol{g}, \quad \boldsymbol{g} \sim \mathcal{N}(0, \boldsymbol{I}_N) \end{split}$$

Let $\langle \cdot \rangle$ denote average with respect to $\sigma \sim \mu_t$. The following fact is a restatement of Bayes theorem: sampling x and then y_t is equivalent to sampling y_t and then x from the posterior. In the context of statistical physics, this is known as 'Nishimori's property.'

Fact 5.6. For any bounded measurable $f, \mathbb{E} f(\boldsymbol{x}, \boldsymbol{y}_t) = \mathbb{E} \langle f(\boldsymbol{\sigma}, \boldsymbol{y}_t) \rangle$.

Lemma 5.7. Let $\iota > 0$ be arbitrary. With probability $1 - e^{-cN}$,

$$\left|\left|\boldsymbol{y}_{t}\right|\right|_{N}^{2}-t^{2}-t\right|\leqslant\iota,\quad\left|\langle\boldsymbol{x},\boldsymbol{y}_{t}\rangle_{N}-t\right|\leqslant\iota,\quad\mu_{t}(\mathsf{Band}(\boldsymbol{y}_{t},[t-\iota,t+\iota]))\geqslant1-e^{-cN}.$$

Proof. Clearly $\|\boldsymbol{y}_t\|_N^2 \simeq t^2 + t$ and $\langle \boldsymbol{x}, \boldsymbol{y}_t \rangle_N \simeq t$, so the first two conclusions follow by standard concentration arguments. By Fact 5.6,

$$\mathbb{E} \langle \mathbf{1} \{ \langle \boldsymbol{\sigma}, \boldsymbol{y}_t \rangle_N \notin [t - \iota, t + \iota] \} \rangle = \mathbb{P} \left(\langle \boldsymbol{x}, \boldsymbol{y}_t \rangle_N \notin [t - \iota, t + \iota] \right) \leqslant e^{-cN}.$$

By Markov's inequality,

$$\mathbb{P}\left\{ \langle \mathbf{1}\left\{ \langle \boldsymbol{\sigma}, \boldsymbol{y}_t \rangle_N \notin [t-\iota, t+\iota] \right\} \rangle \ge e^{-cN/2} \right\} \leqslant e^{-cN/2}.$$

This implies the final conclusion after adjusting c.

We next introduce a self-reduction property of models obtained by restriction to a certain band. Define

$$U = \left\{ oldsymbol{\sigma} \in \mathbb{R}^N : \langle oldsymbol{\sigma}, oldsymbol{y}_t
angle_N = 0
ight\}.$$

Recall that $(q_k)_{k \ge 0}$ is defined by Eq. (2.2), and in particular $q_1 = t/(1+t)$. Let $\hat{y}_t = y_t/||y_t||_N$ and $r = \sqrt{q_1}$. Consider the Hamiltonian on $\rho \in U$ defined by

$$\hat{H}(\boldsymbol{\rho}) = H_N(r\hat{\boldsymbol{y}}_t + \sqrt{1 - r^2}\boldsymbol{\rho}) - H_N(r\hat{\boldsymbol{y}}_t).$$

Further define

$$\xi_{(1)}(s) = \xi(q_1 + (1 - q_1)s) - \xi(q_1)$$

Let $r_1 = \langle \boldsymbol{x}, \hat{\boldsymbol{y}}_t \rangle_N$ and define $\boldsymbol{x}^{\perp} \in U$ by $\boldsymbol{x} = r_1 \hat{\boldsymbol{y}}_t + \sqrt{1 - r_1^2} \boldsymbol{x}^{\perp}$. Note that conditionally on $(\boldsymbol{y}_t, r_1), \boldsymbol{x}^{\perp}$ is a uniformly random vector in $U \cap S_N$. Also define the Hamiltonian

$$\widehat{H}'(\boldsymbol{\rho}) = N\xi_{(1)}(\langle \boldsymbol{x}^{\perp}, \boldsymbol{\rho} \rangle_N) + \widetilde{H}'(\boldsymbol{\rho}),$$

where \widetilde{H}' is a Gaussian process on U with covariance

$$\mathbb{E} \widetilde{H}'(\boldsymbol{\rho}^1) \widetilde{H}'(\boldsymbol{\rho}^2) = N\xi_{(1)}(\langle \boldsymbol{\rho}^1, \boldsymbol{\rho}^2 \rangle_N).$$

Note that \hat{H}' is of the form (4.3), with one fewer dimension and $\xi_{(1)}$ in place of ξ_t .

Proposition 5.8 (Self-reduction). There exists a constant C such that the following holds. Let $\iota > 0$. Let S be the (\boldsymbol{y}_t, r_1) -measurable event

$$\left| \|\boldsymbol{y}_t\|_N - \sqrt{t(1+t)} \right|, |\langle \boldsymbol{x}, \hat{\boldsymbol{y}}_t \rangle_N - \sqrt{q_1}| \leq \iota.$$
(5.10)

Then $\mathbb{P}(S) \ge 1 - e^{-cN}$ and for any $(\boldsymbol{y}_t, r_1) \in S$ the following holds. There is a coupling \mathcal{C} of $\mathcal{L}(\hat{H}|\boldsymbol{y}_t, r_1)$ and $\mathcal{L}(\hat{H}')$ such that almost surely,

$$\frac{1}{N} \sup_{\boldsymbol{\rho} \in U \cap S_N} |\hat{H}(\boldsymbol{\rho}) - \hat{H}'(\boldsymbol{\rho})| \leq C\iota,$$

$$\sup_{\boldsymbol{\rho} \in U \cap S_N} \left\| \nabla_U \hat{H}(\boldsymbol{\rho}) - \nabla_U \hat{H}'(\boldsymbol{\rho}) \right\|_N \leq C\iota.$$
(5.11)

Proof. Suppose the event in Lemma 5.7 holds. Then, using $q_1 = t/(1+t)$,

$$r_1 = \frac{\langle \boldsymbol{x}, \boldsymbol{y}_t \rangle_N}{\|\boldsymbol{y}_t\|_N} = \frac{t + O(\iota)}{\sqrt{t(1+t)} + O(\iota)} = \sqrt{q_1} + O(\iota).$$

This proves $\mathbb{P}(S) \ge 1 - e^{-cN}$, after adjusting ι by a constant factor. Now suppose $(\boldsymbol{y}_t, r_1) \in S$. We have $\hat{H}(\boldsymbol{\rho}) = \hat{H}_1(\boldsymbol{\rho}) + \hat{H}_2(\boldsymbol{\rho})$, where

$$\hat{H}_1(\boldsymbol{\rho}) = N \left\{ \xi \left(R \left(r \hat{\boldsymbol{y}}_t + \sqrt{1 - r^2} \boldsymbol{\rho}, r_1 \hat{\boldsymbol{y}}_t + \sqrt{1 - r_1^2} \boldsymbol{x}^\perp \right) \right) - \xi \left(R \left(r \hat{\boldsymbol{y}}_t, r_1 \hat{\boldsymbol{y}}_t + \sqrt{1 - r_1^2} \boldsymbol{x}^\perp \right) \right) \right\},$$

$$\hat{H}_2(\boldsymbol{\rho}) = \left\{ \tilde{H}_N \left(r \hat{\boldsymbol{y}}_t + \sqrt{1 - r^2} \boldsymbol{\rho} \right) - \tilde{H}_N(r \hat{\boldsymbol{y}}_t) \right\}.$$

The first summand simplifies as

$$\widehat{H}_{1}(\boldsymbol{\rho}) = N\left\{\xi\left(rr_{1} + \sqrt{(1-r^{2})(1-r_{1}^{2})}\langle\boldsymbol{\rho},\boldsymbol{x}^{\perp}\rangle_{N}\right) - \xi(rr_{1})\right\} = N\xi_{(1)}(\langle\boldsymbol{\rho},\boldsymbol{x}^{\perp}\rangle_{N}) + N \cdot O(\iota).$$

The second summand is a Gaussian process on U with covariance

$$\mathbb{E}\widehat{H}_{2}(\boldsymbol{\rho}^{1})\widehat{H}_{2}(\boldsymbol{\rho}^{2}) = N\left(\xi(r^{2} + (1 - r^{2})\langle \boldsymbol{\rho}^{1}, \boldsymbol{\rho}^{2}\rangle_{N}) - \xi(r^{2})\right) = N\xi_{(1)}(\langle \boldsymbol{\rho}^{1}, \boldsymbol{\rho}^{2}\rangle_{N}).$$

Thus we can couple \hat{H}_2 and \tilde{H}' so that $\hat{H}_2 = \tilde{H}'$ almost surely.

Define $\hat{q}_0 = 0$ and, similarly to (2.2),

$$\widehat{q}_{k+1} = \frac{\xi'_{(1)}(\widehat{q}_k)}{1 + \xi'_{(1)}(\widehat{q}_k)}.$$

Lemma 5.9. For all $k \ge 0$, we have $q_1 + (1 - q_1)\hat{q}_k = q_{k+1}$.

Proof. We induct on k. The base case k = 0 is trivial. Recalling $q_1 = \frac{t}{1+t}$, the inductive step follows from

$$q_{1} + (1 - q_{1})\widehat{q}_{k+1} = q_{1} + (1 - q_{1})\frac{\xi_{(1)}'(\widehat{q}_{k})}{1 + \xi_{(1)}'(\widehat{q}_{k})} = 1 - (1 - q_{1})\left(1 - \frac{\xi_{(1)}'(\widehat{q}_{k})}{1 + \xi_{(1)}'(\widehat{q}_{k})}\right)$$
$$= 1 - \frac{1 - q_{1}}{1 + (1 - q_{1})\xi'(q_{k+1})} = 1 - \frac{1}{1 + t + \xi'(q_{k+1})}$$
$$= \frac{\xi_{t}'(q_{k+1})}{1 + \xi_{t}'(q_{k+1})} = q_{k+2}.$$

Define the AMP iteration, analogous to (5.1), on the reduced model \hat{H}' , by $\hat{m}^{-1} = \hat{w}^0 = 0$ and

$$\widehat{\boldsymbol{m}}^{k} = (1 - \widehat{q}_{k})\widehat{\boldsymbol{w}}^{k}, \qquad \widehat{\boldsymbol{w}}^{k+1} = \nabla_{U}\widehat{H}'(\widehat{\boldsymbol{m}}^{k}) - (1 - \widehat{q}_{k})\xi_{(1)}''(\widehat{q}_{k})\widehat{\boldsymbol{m}}^{k-1}$$

Note that $\widehat{\boldsymbol{m}}^k, \widehat{\boldsymbol{w}}^k \in U$.

Proposition 5.10 (Self-reduction of AMP iterates). Let $\iota > 0$. Suppose $(\boldsymbol{y}_t, r_1) \in S$ for S as in Proposition 5.8, and couple $\mathcal{L}(\hat{H}|\boldsymbol{y}_t, r_1)$ and \hat{H}' as in that proposition. Then (conditionally on \boldsymbol{y}_t, r_1) with probability $1 - e^{-cN}$, for all $1 \leq k \leq O(1)$,

$$\|\boldsymbol{m}^{k} - \widetilde{\boldsymbol{m}}^{k}\|_{N} \leq O(\iota), \quad \text{where} \quad \widetilde{\boldsymbol{m}}^{k+1} = \sqrt{q_{1}}\widehat{\boldsymbol{y}}_{t} + \sqrt{1 - q_{1}}\widehat{\boldsymbol{m}}^{k}.$$
 (5.12)

Proof. We induct on the claim that (5.12) holds for all $1 \le k \le K$. First, we have

$$\boldsymbol{m}^{1} = (1 - q_{1})\boldsymbol{y}_{t} = \frac{\boldsymbol{y}_{t}}{1 + t}, \qquad \widetilde{\boldsymbol{m}}^{1} = \sqrt{q_{1}}\widehat{\boldsymbol{y}}_{t} = \sqrt{\frac{t}{1 + t}}\widehat{\boldsymbol{y}}_{t}.$$
(5.13)

For $(\boldsymbol{y}_t, r_1) \in S$, we have $|\|\boldsymbol{y}_t\|_N - \sqrt{t(1+t)}| \leq \iota$, and thus

$$\|\boldsymbol{m}^1 - \widetilde{\boldsymbol{m}}^1\|_N = \left|\frac{\sqrt{\|\boldsymbol{y}_t\|_N}}{1+t} - \sqrt{\frac{t}{1+t}}\right| \leq \frac{\iota}{1+t}.$$

This proves the base case K = 1. Suppose (5.12) holds for $1 \le k \le K$. By Proposition 5.1, for all $1 \le j, k \le K + 1$,

$$\langle \boldsymbol{m}^j, \boldsymbol{m}^k \rangle_N \to_p q_{j \wedge k}, \qquad \langle \widehat{\boldsymbol{m}}^j, \widehat{\boldsymbol{m}}^k \rangle_N \to_p \widehat{q}_{j \wedge k},$$

and thus, by Lemma 5.9,

$$\langle \widetilde{\boldsymbol{m}}^j, \widetilde{\boldsymbol{m}}^k \rangle_N \to_p q_1 + (1-q_1) \widehat{q}_{(j-1)\wedge(k-1)} = q_k.$$

Because AMP iterates are Lipschitz in the disorder (see the proof of Proposition 5.4), on an event with probability $1 - e^{-cN}$,

$$\langle \boldsymbol{m}^{j}, \boldsymbol{m}^{k} \rangle_{N}, \langle \widetilde{\boldsymbol{m}}^{j}, \widetilde{\boldsymbol{m}}^{k} \rangle_{N} \in [q_{j \wedge k} - \iota, q_{j \wedge k} + \iota]$$
(5.14)

for all $1 \leq j,k \leq K + 1$. Since m^1 is a multiple of $y_t = \nabla H_{N,t}(\mathbf{0})$,

$$\boldsymbol{m}^{K+1} \in \operatorname{span}(\boldsymbol{m}^1,\ldots,\boldsymbol{m}^K, \nabla H_{N,t}(\boldsymbol{m}^K)) = \operatorname{span}(\boldsymbol{m}^1,\ldots,\boldsymbol{m}^K, \nabla_U H_N(\boldsymbol{m}^K)).$$

$$\widehat{\boldsymbol{m}}^{K} \in \operatorname{span}(\widehat{\boldsymbol{m}}^{1}, \dots, \widehat{\boldsymbol{m}}^{K-1}, \nabla_{U}\widehat{H}'(\widehat{\boldsymbol{m}}^{K-1})),$$

we have

$$\widetilde{\boldsymbol{m}}^{K+1} \in \operatorname{span}(\widetilde{\boldsymbol{m}}^1, \dots, \widetilde{\boldsymbol{m}}^K, \nabla_U \widehat{H}'(\widehat{\boldsymbol{m}}^{K-1})).$$

Note that $\sqrt{1-q_1}\nabla_U H_N(\widetilde{\boldsymbol{m}}^K) = \nabla_U \hat{H}(\widehat{\boldsymbol{m}}^{K-1})$. Thus (on an event where ∇H_N is O(1)-Lipschitz, and the event in Proposition 5.8, both of which are probability $1 - e^{-cN}$)

$$\begin{split} \left\| \sqrt{1 - q_1} \nabla_U H_N(\boldsymbol{m}^K) - \nabla_U \widehat{H}'(\widehat{\boldsymbol{m}}^{K-1}) \right\|_N &\leq \sqrt{1 - q_1} \left\| \nabla_U H_N(\boldsymbol{m}^K) - \nabla_U H_N(\widetilde{\boldsymbol{m}}^K) \right\|_N \\ &+ \left\| \nabla_U \widehat{H}(\widehat{\boldsymbol{m}}^{K-1}) - \nabla_U \widehat{H}'(\widehat{\boldsymbol{m}}^{K-1}) \right\|_N = O(\iota). \end{split}$$

This and (5.14) imply $\|\boldsymbol{m}^{K+1} - \widetilde{\boldsymbol{m}}^{K+1}\|_N = O(\iota)$, completing the induction.

Proposition 5.11. For all $\iota > 0$ and $k \ge 1$ fixed, the following holds. Let

$$V_k(\iota) = \left\{ \boldsymbol{\sigma} \in S_N : |\langle \boldsymbol{\sigma}, \boldsymbol{m}^j \rangle_N - q_j| \leq \iota, \quad \forall 1 \leq j \leq k \right\}.$$

Then, with probability $1 - e^{-cN}$,

$$\mu_t(V_k(\iota)) \ge 1 - e^{-cN}$$

Proof. We induct on k. By Lemma 5.7, with probability $1 - e^{-cN}$,

$$\mu_t(\mathsf{Band}(\boldsymbol{y}_t, [t-\iota, t+\iota])) \ge 1 - e^{-cN}.$$
(5.15)

As calculated in (5.13), $m^1 = y_t/(1+t)$, so $\sigma \in \text{Band}(y_t, [t-\iota, t+\iota])$ if and only if

$$\langle \boldsymbol{\sigma}, \boldsymbol{m}^1 \rangle_N = \frac{t}{1+t} + O(\iota) = q_1 + O(\iota).$$

This proves the base case k = 1 after adjusting ι by a constant factor.

For the inductive step, let ι_1 be suitably small in ι . Let S_1 be the event (5.10) with right-hand side ι_1 . By Proposition 5.8, $(\boldsymbol{y}_t, r_1) \in S_1$ with probability $1 - e^{-cN}$. Condition on any such (\boldsymbol{y}_t, r_1) . Along with (5.15), this implies

$$\mu_t(\mathsf{Band}(\widehat{\boldsymbol{y}}_t, [\sqrt{q_1} - C\iota_1, \sqrt{q_1} + C\iota_1])) \ge 1 - e^{-cN}$$

for suitable C. For $r_2 \in [\sqrt{q_1} - C\iota_1, \sqrt{q_1} + C\iota_1]$, let $\hat{\mu}_t^{r_2}$ be the Gibbs measure on $U \cap S_N$ given by

$$\hat{\mu}_t^{r_2} = Q_{\#} \mu_t(\cdot | \langle \boldsymbol{\sigma}, \hat{\boldsymbol{y}}_t \rangle_N = r_2), \quad \text{where} \quad Q(\boldsymbol{\sigma}) = \frac{P_{\hat{\boldsymbol{y}}_t}^{\perp}(\boldsymbol{\sigma})}{\|P_{\hat{\boldsymbol{y}}_t}^{\perp}(\boldsymbol{\sigma})\|_N}$$

Note that $\hat{\mu}_t^{\sqrt{q_1}}$ is the Gibbs measure on $U \cap S_N$ corresponding to Hamiltonian \hat{H} . Couple \hat{H} and \hat{H}' as in Proposition 5.8, and let $\hat{\mu}_t'$ be the Gibbs measure on $U \cap S_N$ corresponding to Hamiltonian \hat{H}' .

By the inductive hypothesis **applied to Hamiltonian** \hat{H}' and mixture $\xi_{(1)}$, with probability $1 - e^{-cN}$, $\hat{\mu}'_t(\hat{V}_k(\iota)) \ge 1 - e^{-cN}$, where

$$\widehat{V}_k(\iota) = \left\{ \boldsymbol{\rho} \in U \cap S_N : |\langle \boldsymbol{\rho}, \widehat{\boldsymbol{m}}^j \rangle_N - \widehat{q}_j| \leq \iota, \quad \forall 1 \leq j \leq k \right\}.$$

By Proposition 5.8,

$$\frac{1}{N} \sup_{\boldsymbol{\rho} \in U \cap S_N} |\hat{H}(\boldsymbol{\rho}) - \hat{H}'(\boldsymbol{\rho})| \leq \iota_1.$$

For ι_1 small enough in ι , this implies

$$\hat{\mu}_t^{\sqrt{q_1}}(\hat{V}_k(2\iota)) \ge 1 - e^{-cN}$$

By Lipschitz continuity of $H_{N,t}$, for ι_1 small enough in ι , we have

$$\hat{\mu}_t^{r_2}(\hat{V}_k(3\iota)) \ge 1 - e^{-cN}, \quad \forall r_2 \in [\sqrt{q_1} - C\iota_1, \sqrt{q_1} + C\iota_1].$$

This implies $\mu_t(\widetilde{V}_{k+1}(4\iota)) \ge 1 - e^{-cN}$, where

$$\widetilde{V}_k(\iota) = \left\{ \boldsymbol{\sigma} \in S_N : |\langle \boldsymbol{\sigma}, \widetilde{\boldsymbol{m}}^j \rangle_N - q_j| \leq \iota, \quad \forall 1 \leq j \leq k \right\}.$$

However, by Proposition 5.10, with probability $1 - e^{-cN}$, $\|\boldsymbol{m}^j - \widetilde{\boldsymbol{m}}^j\|_N \leq \iota$ for all $1 \leq j \leq k + 1$. On this event, $\widetilde{V}_{k+1}(4\iota) \subseteq V_k(5\iota)$. Thus $\mu_t(V_k(5\iota)) \geq 1 - e^{-cN}$. This completes the induction, upon adjusting ι .

Proof of Proposition 5.5. Let

$$V_k^+(\iota) = \left\{ \boldsymbol{\sigma} \in S_N : |\langle \boldsymbol{\sigma}, \boldsymbol{m}^k \rangle_N - q_k| \leq \iota \right\},\$$

so clearly $V_k^+(\iota) \supseteq V_k(\iota)$. By Proposition 5.4, for all $k \ge k_0$ we have $|q_k - q_*| \le \iota$. Thus

$$\mathsf{Band}(\boldsymbol{m}^k, [q_* - 2\iota, q_* + 2\iota]) \supseteq V_k^+(\iota).$$

By Proposition 5.11, with probability $1 - e^{-cN}$,

$$\mu_t(\mathsf{Band}(\boldsymbol{m}^k, [q_* - 2\iota, q_* + 2\iota])) \ge 1 - e^{-cN}.$$

The result follows by adjusting ι .

5.3 Overlap with planted signal

The following proposition completes the proof of (4.7).

Proposition 5.12. Let $\iota > 0$ and $I = I(\iota)$. With probability $1 - e^{-cN}$,

$$\mu_t(\mathsf{Band}(\boldsymbol{x},I)) \geqslant 1 - e^{-cN}$$

Lemma 5.13. The function

$$f(q) = \xi_t(q) + q + \log(1-q)$$

is maximized over [0, 1] uniquely at $q = q_*$.

Proof. We calculate

$$f'(q) = \xi'_t(q) - \frac{q}{1-q}, \qquad f''(q) = \xi''(q) - \frac{1}{(1-q)^2}.$$

By (4.2), f is stationary at q_* . By (1.5), it is concave on [0, 1).

We will use the following replica-symmetric upper bound on the free energy. Let \hat{H}_N be the Hamiltonian a spherical spin glass with mixture $\hat{\xi}$, which may contain a degree-1 term (i.e., possibly $\hat{\xi}'(0) > 0$).

Define the partition function

$$\widehat{Z}_N = \int_{S_N} \exp\left\{\widehat{H}_N(\boldsymbol{\sigma})\right\} \mu_0(\mathsf{d}\boldsymbol{\sigma}).$$
(5.16)

Proposition 5.14. For any $u \in [0, 1)$, we have

$$p-\lim_{N \to \infty} \frac{1}{N} \log \hat{Z}_N \leqslant \frac{1}{2} \left(\hat{\xi}(1) - \hat{\xi}(u) + \frac{u}{1-u} + \log(1-u) \right).$$
 (5.17)

Furthermore, equality holds if

$$g(s) = \int_0^s \left(\hat{\xi}'(r) - \frac{r}{(1-u)^2}\right) \,\mathrm{d}r \tag{5.18}$$

is maximized over $s \in [0, u]$ at s = u, and $\hat{\xi}_u(s) = \hat{\xi}(u + (1 - u)s) - \hat{\xi}(u) - (1 - u)\hat{\xi}'(u)s$ satisfies

$$\hat{\xi}_u(s) + s + \log(1-s) \leqslant 0 \tag{5.19}$$

for all $s \in [0, 1)$.

Proof. The bound (5.17) is the spherical Parisi formula [Tal06, Theorem 1.1] with order parameter δ_u . The equality condition follows from the extremality condition [Tal06, Proposition 2.1].

Let $H_{N,t}$ be as in (4.3). Let ψ_N denote the probability density of z_1 , where z is a sample from the uniform Haar measure on the unit sphere \mathbb{S}^{N-1} . It is known that

$$\psi_N(q) = \frac{1}{Z_{N,\psi}} (1 - q^2)^{(N-3)/2}, \qquad q \in [-1, 1]$$
 (5.20)

for some normalizing constant $Z_{N,\psi}$. For $q \in [-1, 1]$, define

$$Z(q) = \int_{\mathsf{Band}(\boldsymbol{x},q)} \exp\left\{H_{N,t}(\boldsymbol{\sigma})\right\} d\mu_{(q)}(\boldsymbol{\sigma}), \qquad (5.21)$$

where $\mu_{(q)}$ is the uniform measure on Band(x,q), normalized to $\mu_{(q)}(\text{Band}(x,q)) = \psi_N(q)$. Note that

$$\int_{-1}^{1} Z(q) \, \mathrm{d}q = \int_{S_N} \exp\left\{H_{N,t}(\boldsymbol{\sigma})\right\} \, \mathrm{d}\mu_0(\boldsymbol{\sigma}).$$

Proposition 5.15. For any fixed $q \in (-1, 1)$,

$$\lim_{N \to \infty} \frac{1}{N} \log Z(q) \leq \frac{1}{2} \Big(\xi_t(1) + \xi_t(|q|) + |q| + \log(1 - |q|) \Big).$$
(5.22)

Equality holds for $q = q_*$, and does not hold for any q < 0.

Proof. Consider first $q \in [0,1]$. On $\mathsf{Band}(x,q)$, if we write $\sigma = qx + \sqrt{1-q^2}\rho$, where $\langle x, \rho \rangle = 0$, then the random part

$$\widehat{H}_{N,q}(\boldsymbol{\rho}) := \widetilde{H}_{N,t}(\boldsymbol{\sigma}) - \widetilde{H}_{N,t}(q\boldsymbol{x}) = \widetilde{H}_{N,t}(q\boldsymbol{x} + \sqrt{1 - q^2\boldsymbol{\rho}})$$

is a spin glass with one fewer dimension and mixture ξ replaced by

$$\hat{\xi}(s) = \xi_t(q^2 + (1 - q^2)s) - \xi_t(q^2).$$

Then,

$$\underset{N \to \infty}{\text{p-lim}} \frac{1}{N} \log Z(q) = \xi_t(q) + \frac{1}{2} \log(1 - q^2) + \underset{N \to \infty}{\text{p-lim}} \frac{1}{N} \log \widehat{Z}_{N,q},$$
 (5.23)

where $\hat{Z}_{N,q}$ is the free energy of the spin glass with Hamiltonian $\hat{H}_{N,q}$. By Proposition 5.14 with $u = \frac{q}{1+q}$,

$$p-\lim_{N \to \infty} \frac{1}{N} \log \widehat{Z}_{N,q} \leq \frac{1}{2} \left(\xi_t(1) - \xi_t(q) + q - \log(1+q) \right).$$
 (5.24)

Combining with (5.23) proves (5.22). For q < 0, (5.23) still holds. Since $\xi_t(q) < \xi_t(|q|)$, and the remaining terms on the right-hand side of (5.23) depend on q only through |q|, (5.22) holds with strict inequality.

To show that equality holds in (5.22) for $q = q_*$, we will verify that (5.24) holds with equality. Let $u_* = \frac{q_*}{1+q_*}$. Then

$$\frac{d\xi}{du}(u_*) = (1 - q_*^2)\xi_t'(q_*) \stackrel{(4.2)}{=} q_*(1 + q_*) = \frac{u_*}{(1 - u_*)^2},$$

while

$$\frac{\mathsf{d}^2\hat{\xi}}{\mathsf{d}u^2}(u_*) = (1-q_*^2)^2 \xi''(q_*) \stackrel{(1.5)}{<} (1+q_*)^2 = \frac{1}{(1-u_*)^2}$$

Thus, for g in (5.18), $g'(u_*) = 0$ and $g''(u_*) < 0$. However, over $s \in [0, u_*]$,

$$g'(s) = \hat{\xi}'(s) - \frac{1}{(1-u_*)^2}$$

is convex because $\hat{\xi}'$ is convex. So, g''(s) < 0 for all $s \in [0, u_*]$, which implies $g'(s) \ge 0$ for all $s \in [0, u_*]$. It follows that g(s) is maximized over $s \in [0, u_*]$ at u_* , verifying (5.18). Since

$$\hat{\xi}_{u_*}(s) = \xi_t(q_* + (1 - q_*)s) - \xi_t(q_*) - (1 - q_*)\xi'_t(q_*)s$$

$$\stackrel{(4.2)}{=} \xi_t(q_* + (1 - q_*)s) - \xi_t(q_*) - q_*s,$$

we have

$$\widehat{\xi}_{u_*}(s) + s + \log(1-s) = \left\{ \xi_t(q_* + (1-q_*)s) + (q_* + (1-q_*)s) + \log\left[1 - (q_* + (1-q_*)s)\right] \right\} - \left\{ \xi_t(q_*) + q_* + \log(1-q_*) \right\} \le 0,$$

where the final inequality is by Lemma 5.13. This verifies (5.19) and completes the proof.

Proof of Proposition 5.12. Fix $\iota > 0$ arbitrarily (independent of N). We will choose $\upsilon = \upsilon(\iota)$ a sufficiently small constant to verify the derivations below. Let

$$q_k^+ = q_* + \iota + k\upsilon, \qquad q_k^- = q_* - \iota - k\upsilon,$$

and let k^+ (resp. k^-) be the largest integer such that $q_{k^+}^+ \leq 1$ (resp. $q_{k^-}^- \geq -1$). Let

$$J = \{q_{k^-}^-, \dots, q_1^-, q_1^+, \dots, q_{k^+}^+\}.$$

Define $h(q) = \frac{1}{2}(\xi_t(1) + \xi_t(|q|) + |q| + \log(1 - |q|))$ to be the right-hand side of (5.22). Consider the event:

- K_N from Proposition 3.6 holds,
- $\frac{1}{N}\log Z(q_*) \ge h(q_*) v$,
- $\frac{1}{N}\log Z(q) \leq h(q) + v$ for all $q \in J$.

This holds with probability $1 - e^{-cN}$ by concentration properties of Z(q). Further let

$$Z_{0} = \int_{S_{N}} \mathbf{1}\{\langle \boldsymbol{\sigma}, \boldsymbol{x} \rangle_{N} \in [q_{*} - \upsilon, q_{*} + \upsilon]\} \{ \exp H_{N,t}(\boldsymbol{\sigma}) \} d\mu_{0}(\boldsymbol{\sigma}) = \int_{q_{*} - \upsilon}^{q_{*} + \upsilon} Z(q) dq,$$

$$Z_{k}^{+} = \int_{S_{N}} \mathbf{1}\{\langle \boldsymbol{\sigma}, \boldsymbol{x} \rangle_{N} \in [q_{k}^{+}, q_{k}^{+} + \upsilon]\} \{ \exp H_{N,t}(\boldsymbol{\sigma}) \} d\mu_{0}(\boldsymbol{\sigma}) = \int_{q_{k}^{+}}^{q_{k}^{+} + \upsilon} Z(q) dq,$$

$$Z_{k}^{-} = \int_{S_{N}} \mathbf{1}\{\langle \boldsymbol{\sigma}, \boldsymbol{x} \rangle_{N} \in [q_{k}^{-} - \upsilon, q_{k}^{-}]\} \{ \exp H_{N,t}(\boldsymbol{\sigma}) \} d\mu_{0}(\boldsymbol{\sigma}) = \int_{q_{k}^{-} - \upsilon}^{q_{k}^{-} - \upsilon} Z(q) dq.$$

Since K_N holds, $H_{N,t}(\boldsymbol{\sigma})$ is O(1)-Lipschitz, and thus

$$Z_0 \ge Z(q_*)e^{-o_{\upsilon}(1)N}, \qquad Z_k^+ \le Z(q_k^+)e^{o_{\upsilon}(1)N}, \qquad Z_k^- \le Z(q_k^-)e^{o_{\upsilon}(1)N}.$$

Here and below, $o_{\upsilon}(1)$ denotes a term independent of N that vanishes as $\upsilon \to 0$. So

$$\frac{1}{N}\log\int_{S_N}\mathbf{1}\{\langle\boldsymbol{\sigma},\boldsymbol{x}\rangle_N\in[q_*-\iota,q_*+\iota]\}\exp H_{N,t}(\boldsymbol{\sigma})\mathsf{d}\mu_0(\boldsymbol{\sigma})\geq\frac{1}{N}\log Z_0\geq h(q_*)-o_{\upsilon}(1)$$

while

$$\frac{1}{N}\log\int_{S_N} \mathbf{1}\{\langle \boldsymbol{\sigma}, \boldsymbol{x} \rangle_N \notin [q_* - \iota, q_* + \iota]\} \leqslant \frac{1}{N}\log\left(\sum_{k=0}^{k^+} Z_k^+ + \sum_{k=0}^{k^-} Z_k^-\right)$$
$$\leqslant \max_{q \in J} h(q) + o_{\upsilon}(1).$$

By Lemma 5.13, for v small enough,

$$h(q_*) - o_{\upsilon}(1) > \max_{q \in J} h(q) + o_{\upsilon}(1)$$

and thus $\mu_t([q_* - \iota, q_* + \iota]) \ge 1 - e^{-cN}$.

Proof of Proposition 4.3. Follows from Propositions 5.4, 5.5, and 5.12.

6 Description of TAP fixed point: proof of Proposition 4.4

6.1 Existence and uniqueness of TAP fixed point

We say that m is a ι -approximate critical point of \mathcal{F}_{TAP} if $\|\nabla \mathcal{F}_{TAP}(m)\|_N \leq \iota$. In this subsection we show the following result.

Proposition 6.1. There exist $C_{\text{max}}^{\text{spec}} > C_{\min}^{\text{spec}} > 0$ such that, for sufficiently small $\iota > 0$, the following holds with probability $1 - e^{-cN}$.

- (a) \mathcal{F}_{TAP} has a unique critical point m^{TAP} in S_{ι} , which further satisfies (4.9).
- (b) There exists $\iota' = o_{\iota}(1)$ such that any ι -approximate critical point $\mathbf{m} \in S_{\iota}$ of $\mathcal{F}_{\mathsf{TAP}}$ satisfies $\|\mathbf{m} \mathbf{m}^{\mathsf{TAP}}\|_{N} \leq \iota'$.

The proof of this proposition depends on an understanding of the landscape of $\tilde{H}_{N,t}$ restricted to S_0 , given in Proposition 6.2 below (recall that $\tilde{H}_{N,t}$ is the centered version of the Hamiltonian $H_{N,t}$, cf. Eqs. (4.3) and (4.4)). Note that S_0 is an affine transformation of the sphere S_{N-2} ; we will view it as a Riemannian manifold. We first recall notions of Riemannian gradient and Hessian. For $m \in S_0$, let

$$oldsymbol{m}^{\perp} = rac{oldsymbol{m} - q_*oldsymbol{x}}{\sqrt{q_*(1-q_*)}},$$

so that $\langle \boldsymbol{x}, \boldsymbol{m}^{\perp} \rangle_N = 0$ and $\|\boldsymbol{m}^{\perp}\|_N = 1$. The Riemannian gradient and radial derivative of $\widetilde{H}_{N,t}$ are

$$\nabla_{\mathsf{sp}} \widetilde{H}_{N,t}(\boldsymbol{m}) = P_{\mathsf{span}(\boldsymbol{m},\boldsymbol{x})}^{\perp} \nabla \widetilde{H}_{N,t}(\boldsymbol{m}), \qquad \partial_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m}) = \langle \boldsymbol{m}^{\perp}, \nabla \widetilde{H}_{N,t}(\boldsymbol{m}) \rangle / \sqrt{N}.$$

In the below calculations, it will be convenient to work with the following rescaled radial derivative, whose typical maximum is O(1):

$$\widetilde{\partial}_{\rm rad} \widetilde{H}_{N,t}(\boldsymbol{m}) = \partial_{\rm rad} \widetilde{H}_{N,t}(\boldsymbol{m}) / \sqrt{N} = \langle \boldsymbol{m}^{\perp}, \nabla \widetilde{H}_{N,t}(\boldsymbol{m}) \rangle_N$$

Similarly to above, we say $m \in S_0$ is a **Riemannian critical point** of $\tilde{H}_{N,t}$ if $\nabla_{sp}\tilde{H}_{N,t}(m) = 0$, and an *i*-approximate **Riemannian critical point** if $\|\nabla_{sp}\tilde{H}_{N,t}(m)\|_N \leq \iota$. Further define the tangential and Riemannian Hessian (these will be used in the next subsection)

$$\begin{split} \nabla_{\tan}^{2} \widetilde{H}_{N,t}(\boldsymbol{m}) &= P_{\mathsf{span}(\boldsymbol{m},\boldsymbol{x})}^{\perp} \nabla^{2} \widetilde{H}_{N,t}(\boldsymbol{m}) P_{\mathsf{span}(\boldsymbol{m},\boldsymbol{x})}^{\perp}, \\ \nabla_{\mathsf{sp}}^{2} \widetilde{H}_{N,t}(\boldsymbol{m}) &= \nabla_{\tan}^{2} \widetilde{H}_{N,t}(\boldsymbol{m}) - \frac{\widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m})}{\sqrt{q_{*}(1-q_{*})}} P_{\mathsf{span}(\boldsymbol{m},\boldsymbol{x})}^{\perp}. \end{split}$$

Proposition 6.2. There exist $C_{\text{max}}^{\text{spec}} > C_{\min}^{\text{spec}} > 0$ such that for any $\iota > 0$, the following holds with probability $1 - e^{-cN}$.

(a) $\widetilde{H}_{N,t}$ has exactly two Riemannian critical points m_{\pm} on S_0 , and their (rescaled) radial derivatives satisfy

$$\left|\widetilde{\partial}_{\mathsf{rad}}\widetilde{H}_{N,t}(\boldsymbol{m}_{\pm}) \mp \sqrt{\frac{q_*}{1-q_*}} \left(1 + (1-q_*)^2 \xi''(q_*)\right)\right| \leq \iota.$$
(6.1)

Moreover, there exists $\iota' = o_{\iota}(1)$ such that all ι -approximate Riemannian critical points m on S_0 satisfy $||m - m_+||_N \leq \iota'$ for some choice of sign \pm .

- (b) The point m_+ is an ι -approximate critical point of \mathcal{F}_{TAP} (i.e. $\|\nabla \mathcal{F}_{TAP}(m)\|_N \leq \iota$).
- (c) The point m_+ satisfies

$$\operatorname{spec}(\nabla^2 \mathcal{F}_{\operatorname{TAP}}(\boldsymbol{m}_+)) \subseteq [-C^{\operatorname{spec}}_{\max}, -C^{\operatorname{spec}}_{\min}]$$

We will prove this proposition in Subsection 6.2. We first show Proposition 6.1 given Proposition 6.2.

Lemma 6.3. For sufficiently small $\iota > 0$, with probability $1 - e^{-cN}$, \mathcal{F}_{TAP} has a unique critical point m in the region $||m - m_+||_N \leq \iota$, which further satisfies (4.9).

Proof. Throughout this proof, assume the event K_N from Proposition 3.6 holds, which occurs with probability $1 - e^{-cN}$. By Proposition 6.2(c), with probability $1 - e^{-cN}$, m_+ is well-defined and

$$\mathsf{spec}(
abla^2\mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}_+)) \subseteq [-C^{\mathsf{spec}}_{\max}, -C^{\mathsf{spec}}_{\min}].$$

On K_N , the maps $\boldsymbol{m} \mapsto \lambda_{\max}(\nabla^2 \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}))$ and $\boldsymbol{m} \mapsto \lambda_{\min}(\nabla^2 \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}_+))$ are O(1)-Lipschitz (over $\|\boldsymbol{m}\|_N \leq 1-\varepsilon$, for any $\varepsilon > 0$). Thus, for suitably small ι ,

$$\operatorname{spec}(\nabla^{2}\mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})) \subseteq \left[-2C_{\max}^{\operatorname{spec}}, -\frac{1}{2}C_{\min}^{\operatorname{spec}}\right] \qquad \forall \, \|\boldsymbol{m} - \boldsymbol{m}_{+}\|_{N} \leqslant \iota.$$
(6.2)

Let v be suitably small in ι . By Proposition 6.2(b), with probability $1 - e^{-cN}$, $\|\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}_+)\|_N \leq v$. Combined with (6.2), this implies $\mathcal{F}_{\mathsf{TAP}}$ has a unique critical point \boldsymbol{m} in the region $\|\boldsymbol{m} - \boldsymbol{m}_+\|_N \leq \iota$. By (6.2), this critical point also satisfies (4.9), upon adjusting the constants $C_{\min}^{\mathsf{spec}}, C_{\max}^{\mathsf{spec}}$.

Lemma 6.4. For any sufficiently small $\iota > 0$, there exists $\iota' = o_{\iota}(1)$ such that with probability $1 - e^{-cN}$, all ι -approximate critical points $\mathbf{m} \in S_{\iota}$ of $\mathcal{F}_{\mathsf{TAP}}$ satisfy $\|\mathbf{m} - \mathbf{m}_{+}\|_{N} \leq \iota'$.

Proof. Suppose K_N holds. Let $\boldsymbol{m} \in S_{\iota}$ be an ι -approximate critical point of \mathcal{F}_{TAP} , and let $\widetilde{\boldsymbol{m}}$ be the nearest point in S_0 to \boldsymbol{m} , so that $\|\boldsymbol{m} - \widetilde{\boldsymbol{m}}\|_N \leq 2\iota$. On K_N , the map $\boldsymbol{m} \mapsto \nabla \mathcal{F}_{TAP}(\boldsymbol{m})$ is O(1)-Lipschitz. Thus $\widetilde{\boldsymbol{m}}$ is a $O(\iota)$ -approximate critical point of $\mathcal{F}_{TAP}(\boldsymbol{m})$, i.e.

$$\left\|\nabla \widetilde{H}_{N,t}(\widetilde{\boldsymbol{m}}) + \xi_t'(q_*)\boldsymbol{x} - \left((1-q_*)\xi''(q_*) + \frac{1}{1-q_*}\right)\widetilde{\boldsymbol{m}}\right\|_N \leqslant O(\iota).$$
(6.3)

Thus $\|\nabla_{sp} \widetilde{H}_{N,t}(\widetilde{\boldsymbol{m}})\|_N \leq O(\iota)$. By Proposition 6.2(a), there exists $\iota' = o_\iota(1)$ such that on an event with probability $1 - e^{-cN}$, $\|\widetilde{\boldsymbol{m}} - \boldsymbol{m}_{\pm}\|_N \leq \iota'/2$ for some choice of sign \pm . We now show the sign must be +. By (6.3),

$$\widetilde{\partial}_{\mathsf{rad}}\widetilde{H}_{N,t}(\widetilde{\boldsymbol{m}}) = \frac{1}{\sqrt{q_*(1-q_*)}} R\left(\widetilde{\boldsymbol{m}} - q_*\boldsymbol{x}, -\xi_t'(q_*)\boldsymbol{x} + \left((1-q_*)\xi''(q_*) + \frac{1}{1-q_*}\right)\widetilde{\boldsymbol{m}}\right) + O(\iota) \\ = \sqrt{\frac{q_*}{1-q_*}} \left(1 + (1-q_*)^2\xi''(q_*)\right) + O(\iota).$$
(6.4)

If we had $\|\widetilde{m} - m_{-}\|_{N} \leq \iota'/2$, then Eq. (6.1) and Lipschitzness of $m \mapsto \nabla \mathcal{F}_{TAP}(m)$ would imply

$$\widetilde{\partial}_{\mathsf{rad}}\widetilde{H}_{N,t}(\widetilde{\boldsymbol{m}}) = -\sqrt{\frac{q_*}{1-q_*}} \left(1 + (1-q_*)^2 \xi''(q_*)\right) + O(\iota'),$$

which contradicts (6.4) for small enough ι . Thus $\|\widetilde{\boldsymbol{m}} - \boldsymbol{m}_+\|_N \leq \iota'/2$. Recalling $\|\boldsymbol{m} - \widetilde{\boldsymbol{m}}\|_N \leq 2\iota$ implies the conclusion.

Proof of Proposition 6.1. By Lemma 6.3, $(m_+ \text{ is well-defined and})$ there is a unique critical point of \mathcal{F}_{TAP} in the region $||m - m_+||_N \leq \iota$, which also satisfies (4.9). Let m^{TAP} denote this point.

Let $\iota' = o_{\iota}(1)$ be given by Lemma 6.4. For ι sufficiently small, Lemma 6.3 also implies that m^{TAP} is the unique critical point of \mathcal{F}_{TAP} in the region $||m - m_{+}||_{N} \leq \iota'$.

By Lemma 6.4, all ι -approximate critical points $m \in S_{\iota}$ of \mathcal{F}_{TAP} satisfy $||m - m_+||_N \leq \iota'$. In particular all critical points are in this region, and thus m^{TAP} is the unique critical point. This proves part (a). Furthermore, for ι -approximate critical points $m \in S_{\iota}$,

$$\|\boldsymbol{m}-\boldsymbol{m}^{\mathsf{TAP}}\|_{N} \leqslant \|\boldsymbol{m}-\boldsymbol{m}_{+}\|_{N} + \|\boldsymbol{m}^{\mathsf{TAP}}-\boldsymbol{m}_{+}\|_{N} \leqslant 2\iota'.$$

This proves part (b) upon adjusting ι' .

28

6.2 Characterization of Riemannian critical points: proof of Proposition 6.2

The proof builds on a sequence of recent results on **topological trivialization** in spherical spin glasses [FLD14, Fyo15, BČNS22, HS23b].

Proof of Proposition 6.2(a). For $m \in S_0$, we may write $m = q_* x + \sqrt{q_*(1-q_*)}\tau$, where $\langle x, \tau \rangle_N = 0$ and $\|\tau\|_N = 1$. Let

$$\widehat{H}(\boldsymbol{\tau}) = \widetilde{H}_{N,t}(q_*\boldsymbol{x} + \sqrt{q_*(1-q_*)}\boldsymbol{\tau}) - \widetilde{H}_{N,t}(q_*\boldsymbol{x})$$

This is a spin glass (in 1 fewer dimension) with mixture

$$\widetilde{\xi}(s) = \xi_t(q_*^2 + q_*(1 - q_*)s) - \xi_t(q_*^2).$$
(6.5)

Note that

$$\widetilde{\xi}'(1) = q_*(1-q_*)\xi_t'(q_*) \stackrel{(4.2)}{=} q_*^2, \qquad \qquad \widetilde{\xi}''(1) = q_*^2(1-q_*)^2\xi''(q_*) \stackrel{(1.5)}{<} q_*^2.$$

Thus $\tilde{\xi}'(1) > \tilde{\xi}''(1)$, which is the condition for topological trivialization identified in [Fyo15, Equation 64], see also [BČNS22, Theorem 1.1]. Thus, with high probability, \hat{H} has exactly two critical points τ_{\pm} , which have radial derivative

$$\widetilde{\partial}_{\mathsf{rad}}\widehat{H}(\boldsymbol{\tau}_{\pm}) = \pm \left(\sqrt{\widetilde{\xi}'(1)} + \frac{\widetilde{\xi}''(1)}{\sqrt{\widetilde{\xi}'(1)}}\right) + O(\iota) = \pm q_* \left(1 + (1 - q_*)^2 \xi''(1)\right) + O(\iota).$$

By [HS23b, Theorem 1.6], this actually holds with probability $1 - e^{-cN}$. On this event, $\tilde{H}_{N,t}$ has exactly two Riemannian critical points m_{\pm} on S_0 , which have radial derivative

$$\widetilde{\partial}_{\mathsf{rad}}\widetilde{H}_{N,t}(\boldsymbol{m}_{\pm}) = \frac{1}{\sqrt{q_{*}(1-q_{*})}} \cdot \widetilde{\partial}_{\mathsf{rad}}\widehat{H}(\boldsymbol{\tau}^{\pm}) = \pm \sqrt{\frac{q_{*}}{1-q_{*}}} \left(1 + (1-q_{*})^{2} \boldsymbol{\xi}''(1)\right) + O(\iota).$$

The estimate (6.1) holds by adjusting ι . The claim about approximate critical points also follows from [HS23b, Theorem 1.6], which shows that all approximate critical points are close to exact critical points.

We will prove parts (b) and (c) by slightly modifying the calculation in [Fyo15, BČNS22]. This calculation is based on the Kac–Rice formula, which we now recall. Let Crt denote the set of Riemannian critical points of $\tilde{H}_{N,t}$ on S_0 and μ_{S_0} denote the (N-2)-dimensional Hausdorff measure on S_0 . The Kac–Rice Formula [Ric44, Kac48] (see [AT07] for a textbook treatment), applied to $\nabla \tilde{H}_{N,t}$ on the Riemannian manifold S_0 , states that for any (random) measurable set $\mathcal{T} \subseteq S_0$,

$$\mathbb{E}\left|\operatorname{Crt}\cap\mathcal{T}\right| = \int_{\mathcal{S}_0} \mathbb{E}\left[\left|\det\nabla_{\operatorname{sp}}^2 \widetilde{H}_{N,t}(\boldsymbol{m})|\mathbf{1}\{\boldsymbol{m}\in\mathcal{T}\}\middle|\nabla_{\operatorname{sp}}\widetilde{H}_{N,t}(\boldsymbol{m})=\mathbf{0}\right]\varphi_{\nabla_{\operatorname{sp}}\widetilde{H}_{N,t}(\boldsymbol{m})}(\mathbf{0})\,\mathrm{d}\mu_{\mathcal{S}_0}(\boldsymbol{m}).$$
(6.6)

Here φ_X denotes the probability density of the random variable X, and $\nabla^2_{sp} \widetilde{H}_{N,t}(\boldsymbol{m})$ is understood as a $(N-2) \times (N-2)$ matrix. The following fact is standard, see, e.g., [AA13, Lemma 1].

Fact 6.5. For any $m \in S_0$, the random variables $\partial_{\mathsf{rad}} \widetilde{H}_{N,t}(m)$, $\nabla_{\mathsf{sp}} \widetilde{H}_{N,t}(m)$, and $\nabla_{\mathsf{tan}}^2 \widetilde{H}_{N,t}(m)$ are independent and Gaussian. Moreover, with $G \sim \mathsf{GOE}(N-2)$, we have

$$abla_{\mathsf{tan}}^2 \widetilde{H}_{N,t}(\boldsymbol{m}) \stackrel{d}{=} \sqrt{\xi''(q_*) \frac{N-2}{N}} \boldsymbol{G}$$

We defer the proof of the following lemma to Subsection 6.5.

Lemma 6.6. Let $G \sim \text{GOE}(N)$. For any $t \ge 1$, r > 2, there exists $C_{r,t} > 0$, uniform for r in compact subsets of $(2, +\infty)$, such that

$$\mathbb{E}\left[\left|\det(r\boldsymbol{I}-\boldsymbol{G})\right|^{t}\right]^{1/t} \leq C_{r,t} \mathbb{E}\left[\left|\det(r\boldsymbol{I}-\boldsymbol{G})\right|\right].$$

Proposition 6.7. We have $\mathbb{E} |\mathsf{Crt}| = 2 + o_N(1)$.

Proof. As shown in the proof of Proposition 6.2(a) above, after reparametrizing S_0 to a sphere of radius \sqrt{N} , the restriction of $\tilde{H}_{N,t}$ to S_0 is a spherical spin glass in one fewer dimension with mixture $\tilde{\xi}$ (6.5), which satisfies $\tilde{\xi}'(1) > \tilde{\xi}''(1)$. The claim follows from [Fyo15, Equation 64] or [BČNS22, Theorem 1.2].

We will use (6.6) through the following lemma. Let

$$r_* = \sqrt{\frac{q_*}{1 - q_*}} \left(1 + (1 - q_*)^2 \xi''(q_*) \right).$$
(6.7)

Lemma 6.8. Let $\iota > 0$ be sufficiently small, $I_{\iota} = [r_* - \iota, r_* + \iota]$, and

$$\mathcal{T}_{\iota} = \left\{ \boldsymbol{m} \in \mathcal{S}_0 : \widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m}) \in I_{\iota} \right\}.$$

There exists a constant C > 0 (independent of ι) such that for any measurable $\mathcal{T} \subseteq \mathcal{T}_{\iota}$,

$$\mathbb{E} \left| \mathsf{Crt} \cap \mathcal{T} \right| \leqslant C \sup_{\boldsymbol{m} \in \mathcal{S}_0} \sup_{r \in I_{\iota}} \mathbb{P} \left[\boldsymbol{m} \in \mathcal{T} \middle| \nabla_{\mathsf{sp}} \widetilde{H}_{N,t}(\boldsymbol{m}) = \mathbf{0}, \widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m}) = r \right]^{1/2}.$$

Proof. By Fact 6.5, $\tilde{\partial}_{\mathsf{rad}} \tilde{H}_{N,t}(\boldsymbol{m})$ is independent of $\nabla_{\mathsf{sp}} \tilde{H}_{N,t}(\boldsymbol{m})$. Explicitly integrating $\tilde{\partial}_{\mathsf{rad}} \tilde{H}_{N,t}(\boldsymbol{m})$ in (6.6) gives

$$\begin{split} \mathbb{E} \left| \mathsf{Crt} \cap \mathcal{T} \right| &= \int_{\mathcal{S}_0} \int_{I_{\iota}} \mathbb{E} \left[\left| \det \nabla_{\mathsf{sp}}^2 \widetilde{H}_{N,t}(\boldsymbol{m}) | \mathbf{1} \{ \boldsymbol{m} \in \mathcal{T} \} \right| \nabla_{\mathsf{sp}} \widetilde{H}_{N,t}(\boldsymbol{m}) = \mathbf{0}, \widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m}) = r \right] \\ &\times \varphi_{\widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m})}(r) \varphi_{\nabla_{\mathsf{sp}} \widetilde{H}_{N,t}(\boldsymbol{m})}(\mathbf{0}) \, \mathrm{d}r \, \mathrm{d}\mu_{\mathcal{S}_0}(\boldsymbol{m}). \end{split}$$

By Cauchy-Schwarz,

$$\mathbb{E}\left[|\det \nabla_{\mathsf{sp}}^{2} \widetilde{H}_{N,t}(\boldsymbol{m})|\mathbf{1}\{\boldsymbol{m} \in \mathcal{T}\} \middle| \nabla_{\mathsf{sp}} \widetilde{H}_{N,t}(\boldsymbol{m}) = \mathbf{0}, \widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m}) = r\right]$$

$$\leq \mathbb{E}\left[|\det \nabla_{\mathsf{sp}}^{2} \widetilde{H}_{N,t}(\boldsymbol{m})|^{2} \middle| \nabla_{\mathsf{sp}} \widetilde{H}_{N,t}(\boldsymbol{m}) = \mathbf{0}, \widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m}) = r\right]^{1/2}$$

$$\times \mathbb{P}\left[\boldsymbol{m} \in \mathcal{T} \middle| \nabla_{\mathsf{sp}} \widetilde{H}_{N,t}(\boldsymbol{m}) = \mathbf{0}, \widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m}) = r\right]^{1/2}.$$

By Fact 6.5, conditional on $\nabla_{sp} \widetilde{H}_{N,t}(\boldsymbol{m}) = \boldsymbol{0}, \, \widetilde{\partial}_{rad} \widetilde{H}_{N,t}(\boldsymbol{m}) = r$,

$$\nabla_{\mathsf{sp}}^{2} \widetilde{H}_{N,t}(\boldsymbol{m}) \stackrel{d}{=} \sqrt{\xi''(q_{*}) \frac{N-2}{N}} \boldsymbol{G} - \frac{r}{\sqrt{q_{*}(1-q_{*})}} \boldsymbol{I}$$
$$= \sqrt{\xi''(q_{*}) \frac{N-2}{N}} \left(\boldsymbol{G} - \sqrt{\frac{N}{N-2}} \frac{r}{\sqrt{q_{*}(1-q_{*})\xi''(q_{*})}} \boldsymbol{I} \right).$$

In light of (1.5),

$$\frac{r_*}{\sqrt{q_*(1-q_*)\xi''(q_*)}} = (1-q_*)\xi''(q_*)^{1/2} + \frac{1}{(1-q_*)\xi''(q_*)^{1/2}} > 2,$$

and thus, for $r \in I_{\iota}$ and ι suitably small,

$$\sqrt{\frac{N}{N-2}} \frac{r}{\sqrt{q_*(1-q_*)\xi''(q_*)}} > 2$$

By Lemma 6.6, for some C > 0,

$$\mathbb{E}\left[\left|\det \nabla_{\mathsf{sp}}^{2}\widetilde{H}_{N,t}(\boldsymbol{m})\right|^{2}\left|\nabla_{\mathsf{sp}}\widetilde{H}_{N,t}(\boldsymbol{m})=\boldsymbol{0},\widetilde{\partial}_{\mathsf{rad}}\widetilde{H}_{N,t}(\boldsymbol{m})=r\right]^{1/2}\right]^{1/2}$$

$$=\mathbb{E}\left[\left|\det \sqrt{\xi''(q_{*})}\frac{N-2}{N}\left(\boldsymbol{G}-\sqrt{\frac{N}{N-2}}\frac{r}{\sqrt{q_{*}(1-q_{*})\xi''(q_{*})}}\boldsymbol{I}\right)\right|^{2}\right]^{1/2}$$

$$\leq C\mathbb{E}\left[\left|\det \sqrt{\xi''(q_{*})}\frac{N-2}{N}\left(\boldsymbol{G}-\sqrt{\frac{N}{N-2}}\frac{r}{\sqrt{q_{*}(1-q_{*})\xi''(q_{*})}}\boldsymbol{I}\right)\right|\right]$$

$$=C\mathbb{E}\left[\left|\det \nabla_{\mathsf{sp}}^{2}\widetilde{H}_{N,t}(\boldsymbol{m})\right|\left|\nabla_{\mathsf{sp}}\widetilde{H}_{N,t}(\boldsymbol{m})=\boldsymbol{0},\widetilde{\partial}_{\mathsf{rad}}\widetilde{H}_{N,t}(\boldsymbol{m})=r\right].$$
(6.8)

Combining, we find

$$\begin{split} \mathbb{E} \left| \mathsf{Crt} \cap \mathcal{T} \right| &\leq C \sup_{\boldsymbol{m} \in \mathcal{S}_0} \sup_{r \in I_{\iota}} \mathbb{P} \left[\boldsymbol{m} \in \mathcal{T} \left| \nabla_{\mathsf{sp}} \widetilde{H}_{N,t}(\boldsymbol{m}) = \boldsymbol{0}, \widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m}) = r \right]^{1/2} \\ &\times \int_{\mathcal{S}_0} \int_{I_{\iota}} \mathbb{E} \left[|\det \nabla_{\mathsf{sp}}^2 \widetilde{H}_{N,t}(\boldsymbol{m})| \left| \nabla_{\mathsf{sp}} \widetilde{H}_{N,t}(\boldsymbol{m}) = \boldsymbol{0}, \widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m}) = r \right] \\ &\times \varphi_{\widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m})}(r) \varphi_{\nabla_{\mathsf{sp}} \widetilde{H}_{N,t}(\boldsymbol{m})}(\boldsymbol{0}) \, \mathrm{d}r \, \mathrm{d}\mu_{\mathcal{S}_0}(\boldsymbol{m}). \end{split}$$

By the Kac–Rice formula, the last integral is the expected number of Riemannian critical points m of $\tilde{H}_{N,t}$ with radial derivative $\tilde{\partial}_{rad} \tilde{H}_{N,t}(m) \in I_{\iota}$. This is upper bounded by $\mathbb{E} |Crt| = 2 + o_N(1)$, by Proposition 6.7.

Proposition 6.9. There exist $C_{\max}^{\text{spec}} > C_{\min}^{\text{spec}} > 0$ such that for all sufficiently small $\iota > 0$, there exists $\iota' = h(\iota) = o_{\iota}(1)$ such that the following holds. For any $m \in S_0$ define the events

$$E_1(\boldsymbol{m},\iota') := \left\{ \|\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})\|_N \leqslant \iota' \right\}, \qquad E_2(\boldsymbol{m}) := \left\{ \mathsf{spec}(\nabla^2 \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})) \subseteq \left[-C_{\max}^{\mathsf{spec}}, -C_{\min}^{\mathsf{spec}} \right] \right\}.$$

Then,

$$\inf_{r\in I_{\iota}} \mathbb{P}\left[E_1(\boldsymbol{m}, \iota') \cap E_2(\boldsymbol{m}) \middle| \nabla_{sp} \widetilde{H}_{N,t}(\boldsymbol{m}) = \mathbf{0}, \widetilde{\partial}_{rad} \widetilde{H}_{N,t}(\boldsymbol{m}) = r\right] \ge 1 - e^{-cN}$$

Here the constant c *is uniform over* $m \in S_0$ *.*

We prove this proposition in the next subsection. Assuming it, we first complete the proof of Proposition 6.2.

Proof of Proposition 6.2(b)(c). Let v be small enough that $\max(v, h(v)) \leq \iota$, for the h from Proposition 6.9. Also let $C_{\max}^{\text{spec}}, C_{\min}^{\text{spec}}$ be given by this proposition. Let $\mathcal{T} \subseteq S_0$ be the set of points m such that

- $\tilde{\partial}_{\mathsf{rad}} \tilde{H}_{N,t}(\boldsymbol{m}) \in I_{\upsilon}$, and
- $E_1(\boldsymbol{m},\iota) \cap E_2(\boldsymbol{m})$ does not hold.

Thus $\mathcal{T} \subseteq \mathcal{T}_{v}$. By Lemma 6.8 and Proposition 6.9 (with v for ι)

$$\mathbb{E} \left| \mathsf{Crt} \cap \mathcal{T} \right| \leq C \sup_{\boldsymbol{m} \in \mathcal{S}_0} \sup_{r \in I_v} \mathbb{P} \left[(E_1(\boldsymbol{m}, \iota) \cap E_2(\boldsymbol{m}))^c \middle| \nabla_{\mathsf{sp}} \widetilde{H}_{N,t}(\boldsymbol{m}) = \mathbf{0}, \widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m}) = r \right]^{1/2} \leq e^{-cN}.$$

Thus, with probability $1 - e^{-cN}$, there do not exist points $\boldsymbol{m} \in S_0$ such that $\tilde{\partial}_{\mathsf{rad}} \tilde{H}_{N,t}(\boldsymbol{m}) \in I_v$ and $E_1(\boldsymbol{m}, \iota) \cap E_2(\boldsymbol{m})$ does not hold.

However, by Proposition 6.2(a) with v in place of ι , $\tilde{\partial}_{\mathsf{rad}} \tilde{H}_{N,t}(\boldsymbol{m}_+) \in I_v$ with probability $1 - e^{-cN}$. Thus $E_1(\boldsymbol{m}_+, \iota) \cap E_2(\boldsymbol{m}_+)$ holds, completing the proof.

6.3 Approximate stationarity and local concavity of \mathcal{F}_{TAP} : proof of Proposition 6.9

Lemma 6.10. Let $\mathbf{m} \in S_0$ and $r \in I_{\iota}$. Conditional on $\nabla_{sp} \widetilde{H}_{N,t}(\mathbf{m}) = \mathbf{0}$ and $\widetilde{\partial}_{rad} \widetilde{H}_{N,t}(\mathbf{m}) = r$, $\langle \mathbf{x}, \nabla \widetilde{H}_{N,t}(\mathbf{m}) \rangle$ is Gaussian with mean $q_*(1-q_*)\xi''(q_*) + O(\iota)$ and variance $O(N^{-1})$.

Proof. All the random variables considered are jointly Gaussian, so it suffices to compute the conditional mean and variance. A short linear-algebraic calculation shows

$$\mathbb{E}\left[\langle \boldsymbol{x}, \nabla \widetilde{H}_{N,t}(\boldsymbol{m}) \rangle_{N} \big| \nabla_{\mathsf{sp}} \widetilde{H}_{N,t}(\boldsymbol{m}), \widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m}) \right] = \frac{q_*^{3/2} (1-q_*)^{1/2} \xi''(q_*)}{\xi'_t(q_*) + q_* (1-q_*) \xi''(q_*)} \widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m}).$$

Thus

$$\begin{split} & \mathbb{E}\left[\langle \boldsymbol{x}, \nabla \widetilde{H}_{N,t}(\boldsymbol{m}) \rangle_{N} \middle| \nabla_{\mathsf{sp}} \widetilde{H}_{N,t}(\boldsymbol{m}) = \boldsymbol{0}, \widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m}) = r\right] \\ &= \frac{q_{*}^{3/2} (1 - q_{*})^{1/2} \xi''(q_{*})}{\xi'_{t}(q_{*}) + q_{*}(1 - q_{*}) \xi''(q_{*})} r_{*} + O(\iota) \\ & \stackrel{(4.2),(6.7)}{=} \frac{q_{*}^{3/2} (1 - q_{*})^{1/2} \xi''(q_{*})}{\frac{q_{*}}{1 - q_{*}} + q_{*}(1 - q_{*}) \xi''(q_{*})} \cdot \sqrt{\frac{q_{*}}{1 - q_{*}}} \left(1 + (1 - q_{*})^{2} \xi''(q_{*})\right) + O(\iota) \\ &= q_{*}(1 - q_{*}) \xi''(q_{*}) + O(\iota). \end{split}$$

Before any conditioning, $\langle \boldsymbol{x}, \nabla \tilde{H}_{N,t}(\boldsymbol{m}) \rangle_N$ is Gaussian with variance $O(N^{-1})$, and conditioning only reduces variance.

Proposition 6.11. Let $\mathbf{m} \in S_0$ and $r \in I_{\iota}$. Conditional on $\nabla_{sp} \widetilde{H}_{N,t}(\mathbf{m}) = \mathbf{0}$ and $\widetilde{\partial}_{rad} \widetilde{H}_{N,t}(\mathbf{m}) = r$, $E_1(\mathbf{m}, \iota')$ holds with probability $1 - e^{-cN}$, for some $\iota' = o_{\iota}(1)$.

Proof. By Lemma 6.10, with conditional probability $1 - e^{-cN}$,

$$|\langle \boldsymbol{x}, \nabla H_{N,t}(\boldsymbol{m}) \rangle_N - q_*(1-q_*)\xi''(q_*)| \leq O(\iota).$$

Suppose this event holds. Since $\nabla_{sp} \widetilde{H}_{N,t}(\boldsymbol{m}) = \mathbf{0}$,

$$\begin{split} \nabla \widetilde{H}_{N,t}(\boldsymbol{m}) &= \widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m}) \frac{\boldsymbol{m} - q_* \boldsymbol{x}}{\sqrt{q_*(1 - q_*)}} + \langle \boldsymbol{x}, \nabla \widetilde{H}_{N,t}(\boldsymbol{m}) \rangle \boldsymbol{x} \\ &= \sqrt{\frac{q_*}{1 - q_*}} \left(1 + (1 - q_*)^2 \xi''(q_*) \right) \frac{\boldsymbol{m} - q_* \boldsymbol{x}}{\sqrt{q_*(1 - q_*)}} + q_*(1 - q_*) \xi''(q_*) \boldsymbol{x} + O(\iota) \boldsymbol{x} + O(\iota) \boldsymbol{x} \\ &= -\xi_t'(q_*) \boldsymbol{x} + \left(\frac{1}{1 - q_*} + (1 - q_*) \xi''(q_*) \right) \boldsymbol{m} + O(\iota) \boldsymbol{x} + O(\iota) \boldsymbol{m}. \end{split}$$

Since

$$\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \nabla \widetilde{H}_{N,t}(\boldsymbol{m}) + \xi'_t(q_*)\boldsymbol{x} - \left(\frac{1}{1-q_*} + (1-q_*)\xi''(q_*)\right)\boldsymbol{m},$$

it follows that $\|\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})\|_N \leq O(\iota)$.

The next lemma is a linear-algebraic calculation of the conditional law given $\nabla \tilde{H}_{N,t}(\boldsymbol{m})$ of $\nabla^2 \tilde{H}_{N,t}(\boldsymbol{m})$, now as a Hessian in \mathbb{R}^N rather than a Riemannian Hessian in S_0 . While $\boldsymbol{m} \in S_0$ for the proofs in the current subsection, we will not assume this for use in Fact 6.18 below.

Lemma 6.12. Let
$$\boldsymbol{m} \in \mathbb{R}^N$$
 with $\|\boldsymbol{m}\|_N^2 = q_{\boldsymbol{m}} < 1$. Conditional on $\nabla \widetilde{H}_{N,t}(\boldsymbol{m}) = \boldsymbol{z}$, we have

$$\nabla^2 \widetilde{H}_{N,t}(\boldsymbol{m}) \stackrel{d}{=} \frac{\xi''(q_{\boldsymbol{m}})}{\xi'_t(q_{\boldsymbol{m}})} \cdot \frac{\boldsymbol{m} \boldsymbol{z}^\top + \boldsymbol{z} \boldsymbol{m}^\top}{N} + \frac{\langle \boldsymbol{m}, \boldsymbol{z} \rangle_N}{\xi'_t(q_{\boldsymbol{m}}) + q_{\boldsymbol{m}} \xi''(q_{\boldsymbol{m}})} \left(\xi^{(3)}(q_{\boldsymbol{m}}) - \frac{2\xi''(q_{\boldsymbol{m}})^2}{\xi'_t(q_{\boldsymbol{m}})}\right) \frac{\boldsymbol{m} \boldsymbol{m}^\top}{N} + \boldsymbol{M},$$

where M is the following symmetric random matrix. Let (e_1, \ldots, e_N) be an orthonormal basis of \mathbb{R}^N with $e_1 = m/||m||_2$, and to reduce notation let $M(i, j) = \langle Me_i, e_j \rangle$. Then the random variables $\{M(i, j) : 1 \leq i \leq j \leq N\}$ are independent centered Gaussians with variance

$$\mathbb{E} \mathbf{M}(i,j)^{2} = N^{-1} \times \begin{cases} (irrelevant \ O(1)) & 1 = i = j \\ \xi''(q_{\mathbf{m}}) + q_{\mathbf{m}}\xi^{(3)}(q_{\mathbf{m}}) - \frac{q_{\mathbf{m}}\xi''(q_{\mathbf{m}})^{2}}{\xi'_{t}(q_{\mathbf{m}})} & 1 = i < j \\ 2\xi''(q_{\mathbf{m}}) & 1 < i = j \\ \xi''(q_{\mathbf{m}}) & 1 < i < j \end{cases}$$
(6.9)

Remark 6.13. The covariance calculation in the proof of Lemma 6.12 implies $\xi''(q_m) + q_m \xi^{(3)}(q_m) - \frac{q_m \xi''(q_m)^2}{\xi'_t(q_m)} \ge 0$, but this can also be seen directly by Cauchy-Schwarz:

$$\begin{split} \left(\xi''(q_{m}) + q_{m}\xi^{(3)}(q_{m})\right)\xi_{t}'(q_{m}) &\geq q_{m}\left(\sum_{p\geq 2}p(p-1)^{2}\gamma_{p}^{2}(q_{m})^{p-2}\right)\left(\sum_{p\geq 2}p\gamma_{p}^{2}(q_{m})^{p-2}\right) \\ &\geq q_{m}\left(\sum_{p\geq 2}p(p-1)\gamma_{p}^{2}(q_{m})^{p-2}\right)^{2} = q_{m}\xi''(q_{m})^{2}. \end{split}$$

Proof. It suffices to compute the conditional mean and covariance. Let $u^1, u^2 \in \mathbb{S}^{N-1}$. Then

$$\mathbb{E}\left[\langle \nabla^2 \widetilde{H}_{N,t}(\boldsymbol{m}) \boldsymbol{u}^1, \boldsymbol{u}^2 \rangle \big| \nabla \widetilde{H}_{N,t}(\boldsymbol{m}) \right] = \langle \boldsymbol{v}, \nabla \widetilde{H}_{N,t}(\boldsymbol{m}) \rangle$$

for $\boldsymbol{v} = \boldsymbol{v}(\boldsymbol{u}^1, \boldsymbol{u}^2, \boldsymbol{m})$ such that for all $\boldsymbol{w} \in \mathbb{R}^N$,

$$\langle
abla^2 \widetilde{H}_{N,t}(oldsymbol{m})oldsymbol{u}^1,oldsymbol{u}^2
angle - \langle oldsymbol{v},
abla \widetilde{H}_{N,t}(oldsymbol{m})
angle \mathbb{L}\langle oldsymbol{w},
abla \widetilde{H}_{N,t}(oldsymbol{m})
angle.$$

We calculate that

$$\begin{split} \mathbb{E} \langle \nabla^{2} \widetilde{H}_{N,t}(\boldsymbol{m}) \boldsymbol{u}^{1}, \boldsymbol{u}^{2} \rangle \langle \boldsymbol{w}, \nabla \widetilde{H}_{N,t}(\boldsymbol{m}) \rangle &= N^{-1} \left(\langle \boldsymbol{u}^{1}, \boldsymbol{w} \rangle \langle \boldsymbol{u}^{2}, \boldsymbol{m} \rangle + \langle \boldsymbol{u}^{1}, \boldsymbol{m} \rangle \langle \boldsymbol{u}^{2}, \boldsymbol{w} \rangle \right) \xi''(q_{\boldsymbol{m}}) \\ &+ N^{-2} \langle \boldsymbol{u}^{1}, \boldsymbol{m} \rangle \langle \boldsymbol{u}^{2}, \boldsymbol{m} \rangle \langle \boldsymbol{w}, \boldsymbol{m} \rangle \xi^{(3)}(q_{\boldsymbol{m}}), \\ \mathbb{E} \langle \boldsymbol{v}, \nabla \widetilde{H}_{N,t}(\boldsymbol{m}) \rangle \langle \boldsymbol{w}, \nabla \widetilde{H}_{N,t}(\boldsymbol{m}) \rangle &= \langle \boldsymbol{v}, \boldsymbol{w} \rangle \xi'_{t}(q_{\boldsymbol{m}}) + N^{-1} \langle \boldsymbol{v}, \boldsymbol{m} \rangle \langle \boldsymbol{w}, \boldsymbol{m} \rangle \xi''(q_{\boldsymbol{m}}). \end{split}$$

Thus, *v* must satisfy

$$N^{-1} \left(\langle \boldsymbol{u}^2, \boldsymbol{m} \rangle \boldsymbol{u}^1 + \langle \boldsymbol{u}^1, \boldsymbol{m} \rangle \boldsymbol{u}^2 \right) \xi''(q_{\boldsymbol{m}}) + N^{-2} \langle \boldsymbol{u}^1, \boldsymbol{m} \rangle \langle \boldsymbol{u}^2, \boldsymbol{m} \rangle \xi^{(3)}(q_{\boldsymbol{m}}) \boldsymbol{m} \\ = \xi'_t(q_{\boldsymbol{m}}) \boldsymbol{v} + N^{-1} \langle \boldsymbol{v}, \boldsymbol{m} \rangle \xi''(q_{\boldsymbol{m}}) \boldsymbol{m}.$$

This has solution $\boldsymbol{v} = a_1 \boldsymbol{u}^1 + a_2 \boldsymbol{u}^2 + a_3 \boldsymbol{m}$, where

$$a_{1} = \frac{\xi''(q_{m})}{N\xi'_{t}(q_{m})} \langle \boldsymbol{u}^{2}, \boldsymbol{m} \rangle, \qquad a_{2} = \frac{\xi''(q_{m})}{N\xi'_{t}(q_{m})} \langle \boldsymbol{u}^{1}, \boldsymbol{m} \rangle,$$
$$a_{3} = \frac{\langle \boldsymbol{u}^{1}, \boldsymbol{m} \rangle \langle \boldsymbol{u}^{2}, \boldsymbol{m} \rangle}{N^{2}(\xi'_{t}(q_{m}) + q_{m}\xi''(q_{m}))} \left(\xi^{(3)}(q_{m}) - \frac{2\xi''(q_{m})^{2}}{\xi'_{t}(q_{m})}\right)$$

.

Thus

$$\mathbb{E}\left[\langle \nabla^2 \widetilde{H}_{N,t}(\boldsymbol{m})\boldsymbol{u}^1, \boldsymbol{u}^2 \rangle \big| \nabla \widetilde{H}_{N,t}(\boldsymbol{m}) \right] = a_1 \langle \boldsymbol{u}^1, \nabla \widetilde{H}_{N,t}(\boldsymbol{m}) \rangle + a_2 \langle \boldsymbol{u}^2, \nabla \widetilde{H}_{N,t}(\boldsymbol{m}) \rangle + a_3 \langle \boldsymbol{m}, \nabla \widetilde{H}_{N,t}(\boldsymbol{m}) \rangle,$$

which implies

$$\begin{split} \mathbb{E}\left[\nabla^{2}\widetilde{H}_{N,t}(\boldsymbol{m})\big|\nabla\widetilde{H}_{N,t}(\boldsymbol{m})\right] &= \frac{\xi''(q_{\boldsymbol{m}})}{N\xi'_{t}(q_{\boldsymbol{m}})}(\boldsymbol{m}\nabla\widetilde{H}_{N,t}(\boldsymbol{m})^{\top} + \nabla\widetilde{H}_{N,t}(\boldsymbol{m})\boldsymbol{m}^{\top}) \\ &+ \frac{\langle \boldsymbol{m},\nabla\widetilde{H}_{N,t}(\boldsymbol{m})\rangle_{N}}{N(\xi'_{t}(q_{\boldsymbol{m}}) + q_{\boldsymbol{m}}\xi''(q_{\boldsymbol{m}}))}\left(\xi^{(3)}(q_{\boldsymbol{m}}) - \frac{2\xi''(q_{\boldsymbol{m}})^{2}}{\xi'_{t}(q_{\boldsymbol{m}})}\right)\boldsymbol{m}\boldsymbol{m}^{\top}, \end{split}$$

as desired. The conditionally random part of $\nabla^2 \widetilde{H}_{N,t}$ is thus

$$oldsymbol{M} =
abla^2 \widetilde{H}_{N,t}(oldsymbol{m}) - \mathbb{E}\left[
abla^2 \widetilde{H}_{N,t}(oldsymbol{m}) \Big|
abla \widetilde{H}_{N,t}(oldsymbol{m})
ight].$$

Direct evaluation of covariances $\mathbb{E} M(i_1, j_1) M(i_2, j_2)$ gives the covariance structure (6.9). The calculation is greatly simplified by the fact that $\langle e_i, m \rangle = 0$ for all $i \neq 1$, which implies e.g. that $M(i, j) = \langle \nabla^2 \widetilde{H}_{N,t}(m) e_i, e_j \rangle$ for all $i, j \neq 1$.

Corollary 6.14. Let $\iota > 0$ be sufficiently small. Let $m, z \in \mathbb{R}^N$ with $||m||_N^2 = q_m$ and $\langle m, x \rangle_N = q_x$, such that $|q_m - q_*|, |q_x - q_*|, ||z||_N \leq \iota$. Conditional on $\nabla \mathcal{F}_{\mathsf{TAP}}(m) = z$,

$$\nabla^{2} \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) \stackrel{d}{=} \left(\frac{2+q_{*}}{q_{*}} \xi''(q_{*}) - (1-q_{*})\xi^{(3)}(q_{*}) - \frac{2}{(1-q_{*})^{2}} \right) \frac{\boldsymbol{m}\boldsymbol{m}^{\top}}{N} + \xi''(q_{*}) \frac{(\boldsymbol{x}-\boldsymbol{m})(\boldsymbol{x}-\boldsymbol{m})^{\top}}{N} - \left((1-q_{m})\xi''(q_{m}) + \frac{1}{1-q_{m}} \right) \boldsymbol{I} + \boldsymbol{M} + \boldsymbol{E}.$$
(6.10)

Here, M *is as in* (6.9)*, and* E *is a* (x, m, z)*-measurable symmetric matrix satisfying* $||E||_{op} \leq o_{\iota}(1)$ *, whose kernel contains* span $(x, m, z)^{\perp}$.

Proof. In the below calculations, E is an error term satisfying the above, which may change from line to line. Conditioning on $\nabla \mathcal{F}_{TAP}(m) = z$ is equivalent to conditioning on

$$\nabla \widetilde{H}_{N,t}(\boldsymbol{m}) = \widetilde{\boldsymbol{z}} \equiv \boldsymbol{z} - \xi'_t(q_{\boldsymbol{x}})\boldsymbol{x} + \left((1 - q_{\boldsymbol{m}})\xi''(q_{\boldsymbol{m}}) + \frac{1}{1 - q_{\boldsymbol{m}}}\right)\boldsymbol{m}.$$

By Lemma 6.12,

$$\begin{split} \nabla^{2} \widetilde{H}_{N,t}(\boldsymbol{m}) &= \frac{\xi''(q_{\boldsymbol{m}})}{\xi'_{t}(q_{\boldsymbol{m}})} \cdot \frac{\boldsymbol{m} \widetilde{\boldsymbol{z}}^{\top} + \widetilde{\boldsymbol{z}} \boldsymbol{m}^{\top}}{N} + \frac{\langle \boldsymbol{m}, \widetilde{\boldsymbol{z}} \rangle_{N}}{\xi'_{t}(q_{\boldsymbol{m}}) + q_{\boldsymbol{m}} \xi''(q_{\boldsymbol{m}})} \left(\xi^{(3)}(q_{\boldsymbol{m}}) - \frac{2\xi''(q_{\boldsymbol{m}})^{2}}{\xi'_{t}(q_{\boldsymbol{m}})} \right) \frac{\boldsymbol{m} \boldsymbol{m}^{\top}}{N} + \boldsymbol{M} \\ &= -\xi''(q_{*}) \frac{\boldsymbol{m} \boldsymbol{x}^{\top} + \boldsymbol{x} \boldsymbol{m}^{\top}}{N} + \frac{2\xi''(q_{*})}{\xi'_{t}(q_{*})} \left((1 - q_{*})\xi''(q_{*}) + \frac{1}{1 - q_{*}} \right) \frac{\boldsymbol{m} \boldsymbol{m}^{\top}}{N} \\ &+ \frac{q_{*}(-\xi'_{t}(q_{*}) + (1 - q_{*})\xi''(q_{*}) + \frac{1}{1 - q_{*}})}{\xi'_{t}(q_{*}) + q_{*}\xi''(q_{*})} \left(\xi^{(3)}(q_{*}) - \frac{2\xi''(q_{*})^{2}}{\xi'_{t}(q_{*})} \right) \frac{\boldsymbol{m} \boldsymbol{m}^{\top}}{N} + \boldsymbol{M} + \boldsymbol{E} \\ &\stackrel{(4.2)}{=} -\xi''(q_{*}) \frac{\boldsymbol{m} \boldsymbol{x}^{\top} + \boldsymbol{x} \boldsymbol{m}^{\top}}{N} + \left(\frac{2\xi''(q_{*})}{q_{*}} + (1 - q_{*})\xi^{(3)}(q_{*}) \right) \frac{\boldsymbol{m} \boldsymbol{m}^{\top}}{N} + \boldsymbol{M} + \boldsymbol{E}. \end{split}$$

Then

$$\nabla^{2} \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \nabla^{2} \widetilde{H}_{N,t}(\boldsymbol{m}) + \xi''(q_{*}) \frac{\boldsymbol{x}\boldsymbol{x}^{\top}}{N} - \left((1-q_{*})\xi^{(3)}(q_{*}) - \xi''(q_{*}) + \frac{1}{(1-q_{*})^{2}} \right) \frac{2\boldsymbol{m}\boldsymbol{m}^{\top}}{N} \\ - \left((1-q_{m})\xi''(q_{m}) + \frac{1}{1-q_{m}} \right) \boldsymbol{I} + \boldsymbol{E}.$$

Combining gives the conclusion.

Lemma 6.15. Let $\iota > 0$ be sufficiently small and $|q_m - q_*| \leq \iota$. Fix an orthonormal basis e_1, \ldots, e_N of \mathbb{R}^N as discussed above (6.9). Let M be as in (6.9). Let M_* be sampled from the same law, except with q_m replaced by q_* , and with

$$\boldsymbol{M}_*(i,j) = 0, \qquad \forall i,j \in \{1,2\}.$$

There is a coupling of M, M_* such that with probability $1 - e^{-cN}, \|M - M_*\|_{op} \leq o_{\iota}(1)$.

Proof. Let M' be the matrix with M'(i, j) = 0 for all $i, j \in \{1, 2\}$, and otherwise M'(i, j) = M(i, j). Since the M(i, j) have variance $O(N^{-1})$, with probability $1 - e^{-cN}$, $||M - M'||_{op} \leq \iota$.

For all $(i, j) \notin \{1, 2\}^2$, $|\mathbb{E} M(i, j)^2 - \mathbb{E} M_*(i, j)^2| \leq O(\iota)/N$. We couple M and M_* as follows. If $\mathbb{E} M(i, j)^2 \leq \mathbb{E} M_*(i, j)^2$, we first sample M(i, j) from its law, and then sample

$$\boldsymbol{M}_{*}(i,j) = \boldsymbol{M}(i,j) + \upsilon_{i,j}g_{i,j},$$

for $g_{i,j} \sim \mathcal{N}(0, 1/N)$ and suitable $v_{i,j} = O(\iota^{1/2})$. If $\mathbb{E} \mathbf{M}(i, j)^2 \ge \mathbb{E} \mathbf{M}_*(i, j)^2$, we follow a similar procedure, sampling $\mathbf{M}_*(i, j)$ first. Let $\mathbf{E} = \mathbf{M}' - \mathbf{M}_*$. Then

$$\boldsymbol{E}(i,j) = (\varepsilon_{i,j} \upsilon_{i,j} g_{i,j})_{i,j \in [N]}$$

for some (deterministic) signs $\varepsilon_{i,j} \in \{\pm 1\}$. Let $v = \max_{i,j}(v_{i,j})$. There exists a random symmetric Gaussian matrix E', independent of E, such that $E + E' =_d vG$, where $G \sim \text{GOE}(N)$. Define

$$\mathcal{K} = \left\{ \boldsymbol{A} \in \mathbb{R}^{N \times N} \text{ symmetric} : \|\boldsymbol{A}\|_{\text{op}} \leq 3v \right\},\$$

Note that

$$\mathbb{P}(\boldsymbol{E} + \boldsymbol{E}' \notin \mathcal{K}) = \mathbb{P}(\|\boldsymbol{G}\|_{\text{op}} > 3) \leqslant e^{-cN}$$

while by convexity of \mathcal{K} and symmetry of E',

$$\mathbb{P}(\boldsymbol{E} + \boldsymbol{E}' \notin \mathcal{K} | \boldsymbol{E} \notin \mathcal{K}) \geq \frac{1}{2}$$

It follows that $\mathbb{P}(\boldsymbol{E} \notin \mathcal{K}) \leq 2e^{-cN}$, concluding the proof.

Lemma 6.16. Let $G \sim \text{GOE}(N)$ and $g \sim \mathcal{N}(0, I_N/N)$. For any $a, b \in \mathbb{R}$, $\iota > 0$,

$$\sup_{\boldsymbol{v}\in\mathbb{S}^{N-1}}\left\{a\langle\boldsymbol{G}\boldsymbol{v},\boldsymbol{v}\rangle+2b\langle\boldsymbol{g},\boldsymbol{v}\rangle\right\}-2\sqrt{a^2+b^2}\right|\leqslant\iota$$

with probability $1 - e^{-cN}$.

Proof. By [CS17, Proposition 1.1],

$$\underset{N \to \infty}{\text{p-lim}} \sup_{\|\boldsymbol{v}\|_2 = 1} a \langle \boldsymbol{G} \boldsymbol{v}, \boldsymbol{v} \rangle + 2b \langle \boldsymbol{g}, \boldsymbol{v} \rangle = 2\sqrt{a^2 + b^2}.$$

For each fixed $v \in \mathbb{S}^{N-1}$, $a\langle Gv, v \rangle + 2b\langle g, v \rangle$ has variance $O(N^{-1})$. The result follows from Borell-TIS.

Proposition 6.17. Let $\nabla^2 \mathcal{F}_{TAP}(m)$ be as in Eq. (6.10). There exist $C_{\max}^{\text{spec}} > C_{\min}^{\text{spec}} > 0$ such that for sufficiently small $\iota > 0$,

$$\mathsf{spec}(
abla^2\mathcal{F}_{\mathsf{TAP}}(m{m})) \subseteq [-C^{\mathsf{spec}}_{\max}, -C^{\mathsf{spec}}_{\min}]$$

with probability $1 - e^{-cN}$.

Proof. Let $\widetilde{\boldsymbol{m}} = \boldsymbol{m}/\|\boldsymbol{m}\|_2$, $\widetilde{\boldsymbol{x}} = P_{\boldsymbol{m}}^{\perp}\boldsymbol{x}/\|P_{\boldsymbol{m}}^{\perp}\boldsymbol{x}\|_2$. Throughout this proof, we will denote by \boldsymbol{E} , \boldsymbol{E}_1 , \boldsymbol{E}_2 , and so on error terms with the same meaning as in Corollary 6.14, namely $(\boldsymbol{x}, \boldsymbol{m}, \boldsymbol{z})$ -measurable symmetric matrices satisfying $\|\boldsymbol{E}\|_{op} \leq o_{\iota}(1)$, whose kernel contains span $(\boldsymbol{x}, \boldsymbol{m}, \boldsymbol{z})^{\perp}$. In particular

$$q_*\widetilde{\boldsymbol{m}}\widetilde{\boldsymbol{m}}^\top - \frac{\boldsymbol{m}\boldsymbol{m}^\top}{N} =: \boldsymbol{E}_1, \ (1-q_*)\widetilde{\boldsymbol{x}}\widetilde{\boldsymbol{x}}^\top - \frac{(\boldsymbol{x}-\boldsymbol{m})(\boldsymbol{x}-\boldsymbol{m})^\top}{N} =: \boldsymbol{E}_2$$

Let e_1, \ldots, e_N be an orthonormal basis of \mathbb{R}^N with $e_1 = \widetilde{m}$, $e_2 = \widetilde{x}$. Let M_* be defined in Lemma 6.15, coupled to M so that $\|M - M_*\|_{op} \leq o_{\iota}(1)$ with probability $1 - e^{-cN}$. Taking ι small, it suffices to show

$$-C_{\max}^{\text{spec}} \boldsymbol{I} \le \boldsymbol{A} \le -C_{\min}^{\text{spec}} \boldsymbol{I}, \tag{6.11}$$

for

$$\begin{aligned} \boldsymbol{A} &= \left((2+q_*)\xi''(q_*) - q_*(1-q_*)\xi^{(3)}(q_*) - \frac{2q_*}{(1-q_*)^2} \right) \widetilde{\boldsymbol{m}}\widetilde{\boldsymbol{m}}^\top + (1-q_*)\xi''(q_*)\widetilde{\boldsymbol{x}}\widetilde{\boldsymbol{x}}^\top \\ &- \left((1-q_*)\xi''(q_*) + \frac{1}{1-q_*} \right) \boldsymbol{I} + \boldsymbol{M}_*. \end{aligned}$$

By comparing M_* to a large constant multiple of a GOE, identically to the proof of Lemma 6.15, we can show $\|M_*\|_{op} = O(1)$ with probability $1 - e^{-cN}$. On this event, all terms in A have bounded operator norm, and thus $-C_{\max}^{\text{spec}} I \leq A$. For the upper bound in (6.11), let

$$\psi = \xi''(q_*) + q_*\xi^{(3)}(q_*) - \frac{q_*\xi''(q_*)^2}{\xi_t'(q_*)}.$$
which (recall Remark 6.13) is nonnegative. Then

$$\begin{aligned} \boldsymbol{A} &= \left(-(1-q_*)\psi + q_* \left(\xi''(q_*) - \frac{1}{(1-q_*)^2} \right) - (1-q_*)^2 \left(\xi''(q_*) - \frac{1}{(1-q_*)^2} \right)^2 \right) \boldsymbol{e}_1 \boldsymbol{e}_1^\top \\ &- \frac{1}{1-q_*} \boldsymbol{e}_2 \boldsymbol{e}_2^\top - \left((1-q_*)\xi''(q_*) + \frac{1}{1-q_*} \right) \sum_{i=3}^N \boldsymbol{e}_i \boldsymbol{e}_i^\top + \boldsymbol{M}_* \end{aligned}$$

By (1.5), there exists $c_0 > 0$ depending only on ξ such that

$$\begin{split} \mathbf{A} &\leq \mathbf{A}' - c_0(\mathbf{e}_1 \mathbf{e}_1^\top + \mathbf{e}_2 \mathbf{e}_2^\top), \quad \text{where} \\ \mathbf{A}' &= -(1 - q_*)\psi \mathbf{e}_1 \mathbf{e}_1^\top - (1 - q_*)\xi''(q_*)\mathbf{e}_2 \mathbf{e}_2^\top - \left((1 - q_*)\xi''(q_*) + \frac{1}{1 - q_*}\right)\sum_{i=3}^N \mathbf{e}_i \mathbf{e}_i^\top + \mathbf{M}_* \\ &= -(1 - q_*)\psi \mathbf{e}_1 \mathbf{e}_1^\top - (1 - q_*)\xi''(q_*)\sum_{i=2}^N \mathbf{e}_i \mathbf{e}_i^\top - \frac{1}{1 - q_*}\sum_{i=3}^N \mathbf{e}_i \mathbf{e}_i^\top + \mathbf{M}_*. \end{split}$$

By (6.9), (with $\boldsymbol{M}_{*}(i,j)$ having the same meaning as above)

$$(\boldsymbol{M}_{*}(1,i):3 \leq i \leq N) \stackrel{d}{=} \sqrt{\psi} \boldsymbol{g}^{1}, \qquad \boldsymbol{g}^{1} \sim \mathcal{N}(0, \boldsymbol{I}_{N-2}/N),$$
$$(\boldsymbol{M}_{*}(2,i):3 \leq i \leq N) \stackrel{d}{=} \sqrt{\xi''(q_{*})} \boldsymbol{g}^{2}, \qquad \boldsymbol{g}^{2} \sim \mathcal{N}(0, \boldsymbol{I}_{N-2}/N),$$
$$(\boldsymbol{M}_{*}(i,j):3 \leq i,j \leq N) \stackrel{d}{=} \sqrt{\xi''(q_{*})} \cdot \frac{N-2}{N} \boldsymbol{G} \qquad \boldsymbol{G} \sim \mathsf{GOE}(N-2),$$

and g^1, g^2, G are independent. Fix a, b with $a^2 + b^2 \le 1$ and consider temporarily the restricted set

$$\mathbb{S}_{a,b}^{N-1} = \left\{ \boldsymbol{v} \in \mathbb{S}^{N-1} : \langle \boldsymbol{v}, \boldsymbol{e}_1 \rangle = a, \langle \boldsymbol{v}, \boldsymbol{e}_2 \rangle = b \right\}.$$

For any $\boldsymbol{v} \in \mathbb{S}_{a,b}^{N-1}$ we can write

$$\boldsymbol{v} = a\boldsymbol{e}_1 + b\boldsymbol{e}_2 + \sqrt{1 - a^2 - b^2}\boldsymbol{w},$$

where $\boldsymbol{w} \in \mathbb{S}_{0,0}^{N-1}$. Because we defined $\boldsymbol{M}_{*}(i,j) = 0$ for all $i, j \in \{1,2\}$,

$$\begin{split} \langle \boldsymbol{M}_{*}\boldsymbol{v},\boldsymbol{v}\rangle &= 2a\sqrt{(1-a^{2}-b^{2})\psi}\langle \boldsymbol{g}^{1},\boldsymbol{w}\rangle + 2b\sqrt{(1-a^{2}-b^{2})\xi''(q_{*})}\langle \boldsymbol{g}^{2},\boldsymbol{w}\rangle \\ &+ (1-a^{2}-b^{2})\sqrt{\xi''(q_{*})\cdot\frac{N-2}{N}}\langle \boldsymbol{G}\boldsymbol{w},\boldsymbol{w}\rangle. \end{split}$$

By Lemma 6.16, with probability $1 - e^{-cN}$,

$$\left|\sup_{\boldsymbol{v}\in\mathbb{S}^{N-1}_{a,b}}\langle \boldsymbol{M}_{*}\boldsymbol{v},\boldsymbol{v}\rangle-2\sqrt{f(a,b)}\right|\leqslant\iota,$$

where

$$\begin{aligned} f(a,b) &= (1-a^2-b^2)^2 \xi''(q_*) + a^2(1-a^2-b^2)\psi + b^2(1-a^2-b^2)\xi''(q_*) \\ &= (1-a^2-b^2)\left((1-a^2)\xi''(q_*) + a^2\psi\right). \end{aligned}$$

On this event, for all $\boldsymbol{v} \in \mathbb{S}_{a,b}^{N-1}$,

$$\begin{split} \langle \boldsymbol{A}\boldsymbol{v},\boldsymbol{v}\rangle \leqslant -(1-q_{*})\psi a^{2} - (1-q_{*})\xi''(q_{*})(1-a^{2}) - \frac{1-a^{2}-b^{2}}{1-q_{*}} + 2\sqrt{f(a,b)} - c_{0}(a^{2}+b^{2}) + \iota \\ &= -\left(\sqrt{(1-q_{*})\left((1-a^{2})\xi''(q_{*}) + a^{2}\psi\right)} - \sqrt{\frac{1-a^{2}-b^{2}}{1-q_{*}}}\right)^{2} - c_{0}(a^{2}+b^{2}) + \iota. \end{split}$$

At a = b = 0, the first term is strictly negative by (1.5). So, there exists $c_1 > 0$, depending only on ξ , such that for all $a^2 + b^2 \leq 1$,

$$-\left(\sqrt{(1-q_*)\left((1-a^2)\xi''(q_*)+a^2\psi\right)}-\sqrt{\frac{1-a^2-b^2}{1-q_*}}\right)^2-c_0(a^2+b^2)\leqslant -c_1.$$

We have thus shown that, for fixed a, b, with probability $1 - e^{-cN}$,

$$\sup_{\boldsymbol{v}\in\mathbb{S}^{N-1}_{a,b}}\langle \boldsymbol{A}\boldsymbol{v},\boldsymbol{v}\rangle\leqslant -c_1+\iota.$$
(6.12)

Recall that $\| \boldsymbol{A} \|_{\text{op}} = O(1)$ with probability $1 - e^{-cN}$. So, the map

$$(a,b)\mapsto \sup_{\boldsymbol{v}\in\mathbb{S}^{N-1}_{a,b}}\langle \boldsymbol{A}\boldsymbol{v},\boldsymbol{v}
angle$$

is O(1)-Lipschitz. By a union bound, with proability $1 - e^{-cN}$ (6.12) holds for all (a, b) in a ι -net of $a^2 + b^2 \leq 1$. On this event,

$$\sup_{\boldsymbol{v}\in\mathbb{S}^{N-1}}\langle \boldsymbol{A}\boldsymbol{v},\boldsymbol{v}\rangle\leqslant -c_1+O(\iota)$$

Taking $C_{\min}^{\text{spec}} = c_1/2$ and ι small enough completes the proof.

Proof of Proposition 6.9. Let ι' be given by Proposition 6.11. By this proposition, for any $m \in S_0, r \in I_{\iota}$,

$$\mathbb{P}\left(E_1(\boldsymbol{m},\boldsymbol{\iota}')^c \middle| \nabla_{\mathsf{sp}} \widetilde{H}_{N,t}(\boldsymbol{m}) = \mathbf{0}, \widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m}) = r\right) \leqslant e^{-cN}$$

Since $\|\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})\|_N \leq \iota'$ on $E_1(\boldsymbol{m},\iota')$, and $\nabla_{\mathsf{sp}} \widetilde{H}_{N,t}(\boldsymbol{m}), \widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m})$ are $\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})$ -measurable,

$$\mathbb{P}\left(E_1(\boldsymbol{m},\iota') \cap E_2(\boldsymbol{m})^c \big| \nabla_{sp} \widetilde{H}_{N,t}(\boldsymbol{m}) = \boldsymbol{0}, \widetilde{\partial}_{\mathsf{rad}} \widetilde{H}_{N,t}(\boldsymbol{m}) = r\right) \leqslant \sup_{\|\boldsymbol{z}\|_N \leqslant \iota'} \mathbb{P}\left(E_2(\boldsymbol{m})^c \big| \nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \boldsymbol{z}\right).$$

By Corollary 6.14 and Proposition 6.17, this last probability is $\leq e^{-cN}$. This completes the proof.

6.4 **Proof of conditioning bound**

Propositions 3.6 and 6.1 directly imply parts (a) and (b) of Proposition 4.4. We now prove the remainder of this proposition.

Proof of Proposition 4.4(c). Set v > 0 small enough that $\max(v, \iota'(v)) \leq \iota/2$, for the function ι' from Proposition 6.1. Suppose K_N holds and the events in Propositions 4.3 and 6.1 hold with tolerance v. This occurs with probability $1 - e^{-cN}$.

By (4.5) and (4.6), for suitably large K_{AMP} , $\boldsymbol{m}^{AMP} \in S_{\upsilon} \subseteq S_{\iota/2}$ and \boldsymbol{m}^{AMP} is an υ -approximate critical point of \mathcal{F}_{TAP} . By Proposition 6.1(b), this implies $\|\boldsymbol{m}^{AMP} - \boldsymbol{m}^{TAP}\|_N \leq \iota/2$.

We now turn to the proof of part (d). Define

$$K(\boldsymbol{m}) = P_{\mathsf{span}(\boldsymbol{x},\boldsymbol{m})}^{\perp} \nabla^2 \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) P_{\mathsf{span}(\boldsymbol{x},\boldsymbol{m})}^{\perp}.$$

We will treat this as a $(N-2) \times (N-2)$ matrix, after a suitable change of coordinates. The following fact is a consequence of Corollary 6.14.

Fact 6.18. Let $||m||_N^2 = q_m < 1$. Conditional on $\nabla \mathcal{F}_{TAP}(m) = 0$,

$$K(\boldsymbol{m}) \stackrel{d}{=} -\left((1-q_{\boldsymbol{m}})\xi''(q_{\boldsymbol{m}}) + \frac{1}{1-q_{\boldsymbol{m}}}\right)\boldsymbol{I} + \sqrt{\xi''(q_{\boldsymbol{m}})\frac{N-2}{N}}\boldsymbol{G}, \qquad \boldsymbol{G} \sim \mathsf{GOE}(N-2).$$

The next fact is verified by direct calculation.

Fact 6.19. For any m, $\nabla \tilde{H}_{N,t}(m)$ is Gaussian, with variance $\xi'_t(q_m) + q_m \xi''(q_m)$ in the direction of m and $\xi'_t(q_m)$ in all directions orthogonal to m.

We will need the following technical lemma, which we prove in Subsection 6.5.

Lemma 6.20. For all $\iota > 0$ sufficiently small, there exists a constant C > 0 such that

$$\int_{\mathcal{S}_{\iota}} \mathbb{E}\left[|\det K(\boldsymbol{m})| \big| \nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \boldsymbol{0} \right] \varphi_{\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})}(\boldsymbol{0}) \, \mathsf{d}^{N}(\boldsymbol{m}) \leqslant C.$$

Proof of Proposition 4.4(d). We will apply Lemma 3.7 with \mathcal{F}_{TAP} for \mathcal{F} , m^{AMP} for m_0 , the interior of \mathcal{S}_{ι} for D, and $C_{\min}^{spec}/2$ for c_{spec} . Note that (4.10) implies $\varepsilon \leq c_{spec}^2/10c_{op}$. We next verify that the event \mathcal{E} is contained in the event \mathcal{E}_{cond} defined in Lemma 3.7. Suppose \mathcal{E} holds. Then event $\mathcal{H}(c_{op})$ holds by part (a). $\|\nabla \mathcal{F}_{TAP}(m^{AMP})\|_N \leq \varepsilon$ by definition, and by parts (a), (b), and (c),

$$\lambda_{\max}(\nabla^2 \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}^{\mathsf{AMP}})) \leqslant \lambda_{\max}(\nabla^2 \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}^{\mathsf{TAP}})) + c_{\mathsf{op}} \left\| \boldsymbol{m}^{\mathsf{AMP}} - \boldsymbol{m}^{\mathsf{TAP}} \right\|_{\mathsf{op},N} \leqslant -C_{\min}^{\mathsf{spec}} + c_{\mathsf{op}}\iota/2 \leqslant -C_{\min}^{\mathsf{spec}}/2$$

for small enough ι . Thus $\mathcal{G}(\varepsilon, c_{\text{spec}})$ holds. We have $\|\boldsymbol{m}^{\text{AMP}}\|_N \leq 1$ because $\boldsymbol{m}^{\text{AMP}} \in \mathcal{S}_{\iota/2}$, by part (c). Also, (4.10) implies $\frac{5\varepsilon}{c_{\text{spec}}} \leq \frac{\iota}{2}$, so $U = \mathsf{B}^N(\boldsymbol{m}^{\text{AMP}}, 5\varepsilon/c_{\text{spec}}) \subseteq \mathcal{S}_{\iota}$.

By (5.9), $\mathbb{E} \nabla \mathcal{F}_{TAP}(\boldsymbol{m})$ is continuous in \boldsymbol{m} , and by Fact 6.19, $Cov(\nabla \mathcal{F}_{TAP}(\boldsymbol{m}))$ is uniformly lower bounded for all $\boldsymbol{m} \in S_{\iota}$. This verifies the regularity condition in Lemma 3.7. By this lemma,

$$\mathbb{E}[X\mathbf{1}\{\mathcal{E}\}] \leqslant \int_{S_{\iota}} \mathbb{E}\left[|\det \nabla^2 \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})| X\mathbf{1}\{\mathcal{E}\} \big| \nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \mathbf{0} \right] \varphi_{\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})}(\mathbf{0}) \, \mathsf{d}^N(\boldsymbol{m})$$

By Hölder's inequality,

$$\mathbb{E}\left[|\det \nabla^{2} \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})|X\mathbf{1}\{\mathcal{E}\}|\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \mathbf{0}\right]$$

$$\leq \mathbb{E}\left[|\det \nabla^{2} \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})|^{1+\delta^{-1}}\mathbf{1}\{\mathcal{E}\}|\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \mathbf{0}\right]^{\delta/(1+\delta)} \mathbb{E}\left[X^{1+\delta}\mathbf{1}\{\mathcal{E}\}|\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \mathbf{0}\right]^{1/(1+\delta)}$$

On event \mathcal{E} , the eigenvalues of $\nabla^2 \mathcal{F}_{TAP}(\boldsymbol{m})$ lie in $[-C_{\max}^{spec}, -C_{\min}^{spec}]$ and interlace those of $K(\boldsymbol{m})$. So,

$$|\det \nabla^2 \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})| \leq (C_{\max}^{\mathsf{spec}})^2 |\det K(\boldsymbol{m})|.$$

Thus,

$$\begin{split} & \mathbb{E}\left[\left|\det\nabla^{2}\mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})\right|^{1+\delta^{-1}}\mathbf{1}\{\mathcal{E}\}\Big|\nabla\mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})=\mathbf{0}\right]^{\delta/(1+\delta)}\\ &\leqslant (C_{\max}^{\mathsf{spec}})^{2}\,\mathbb{E}\left[\left|\det K(\boldsymbol{m})\right|^{1+\delta^{-1}}\Big|\nabla\mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})=\mathbf{0}\right]^{\delta/(1+\delta)}\\ &\leqslant (C_{\max}^{\mathsf{spec}})^{2}C_{\delta}'\,\mathbb{E}\left[\left|\det K(\boldsymbol{m})\right|\Big|\nabla\mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})=\mathbf{0}\right]. \end{split}$$

for some $C'_{\delta} > 0$. The last estimate is by Fact 6.18, (1.5), and Lemma 6.6, similarly to (6.8). Combining,

$$\begin{split} \mathbb{E}[X\mathbf{1}\{\mathcal{E}\}] &\leqslant (C_{\max}^{\mathsf{spec}})^2 C_{\delta}' \int_{\mathcal{S}_{\iota}} \mathbb{E}\left[|\det K(\boldsymbol{m})| \big| \nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \mathbf{0} \right] \varphi_{\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})}(\mathbf{0}) \, \mathsf{d}^{N}(\boldsymbol{m}) \\ &\times \sup_{\boldsymbol{m} \in \mathcal{S}_{\iota}} \mathbb{E}\left[X^{1+\delta} \mathbf{1}\{\mathcal{E}\} \big| \nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \mathbf{0} \right]^{1/(1+\delta)}. \end{split}$$

Finally, by Lemma 6.20, this integral is bounded by a constant C > 0. Thus the result holds with $C_{\delta} = (C_{\max}^{\text{spec}})^2 C'_{\delta} C$.

6.5 Determinant concentration and estimate of Kac-Rice integral

In this subsection, we provide the deferred proofs of Lemmas 6.6 and 6.20. These are the final ingredients to the proof of Proposition 4.4.

Proof of Lemma 6.6. For any compact $K \subseteq (2, +\infty)$, we may pick $\varepsilon > 0$ such small enough that $r \ge 2+2\varepsilon$ for all $r \in K$. Let $\mathcal{E}_{\varepsilon}$ be the event that $\|\mathbf{G}\|_{op} \le 2 + \varepsilon$. It is classical that $\mathbb{P}(\mathcal{E}_{\varepsilon}) \ge 1 - e^{-cN}$. For $r \in K$, let

$$f(x) = \log \max(|r - x|, \varepsilon),$$

which is ε^{-1} -Lipschitz. Let $\lambda_1, \ldots, \lambda_N$ be the eigenvalues of G and define

$$\operatorname{Tr} f(\boldsymbol{G}) = \sum_{i=1}^{N} f(\lambda_i).$$

By [GZ00, Theorem 1.1(b)], for all $s \ge 0$,

$$\mathbb{P}\left(|\mathsf{Tr}f(\boldsymbol{G}) - \mathbb{E}\,\mathsf{Tr}f(\boldsymbol{G})| \ge s\right) \le 2\exp(-\varepsilon^2 s^2/8).\tag{6.13}$$

Note that $|\det(rI - G)| \leq \exp(\operatorname{Tr} f(G))$, and equality holds if $G \in \mathcal{E}_{\varepsilon}$. Thus,

$$\mathbb{E}\left[|\det(r\boldsymbol{I}-\boldsymbol{G})|^t\right] \leqslant \mathbb{E}\left[\exp(t\mathsf{Tr}f(\boldsymbol{G}))\right], \qquad \mathbb{E}\left[|\det(r\boldsymbol{I}-\boldsymbol{G})|\right] \geqslant \mathbb{E}\left[\exp(\mathsf{Tr}f(\boldsymbol{G}))\mathbf{1}\{\mathcal{E}_{\varepsilon}\}\right].$$

By (6.13), there exists $C_{\varepsilon,t}$ depending on ε, t such that

$$\mathbb{E}\left[\exp(t\operatorname{Tr} f(\boldsymbol{G}))\right] \leqslant C_{\varepsilon,t}\exp(t\mathbb{E}\operatorname{Tr} f(\boldsymbol{G})).$$

By Cauchy-Schwarz,

$$\mathbb{E}\left[\exp(\mathrm{Tr} f(\boldsymbol{G}))\mathbf{1}\{\mathcal{E}_{\varepsilon}^{c}\}\right] \leqslant \mathbb{E}\left[\exp(2\mathrm{Tr} f(\boldsymbol{G}))\right]^{1/2} \mathbb{P}(\mathcal{E}_{\varepsilon}^{c})^{1/2} \leqslant C_{\varepsilon,2}^{1/2} e^{-cN/2} \exp(\mathbb{E}\operatorname{Tr} f(\boldsymbol{G})),$$

which implies

$$\mathbb{E}\left[\exp(\mathsf{Tr}f(\boldsymbol{G}))\mathbf{1}\{\mathcal{E}_{\varepsilon}\}\right] \ge \mathbb{E}\left[\exp(\mathsf{Tr}f(\boldsymbol{G}))\right] - C_{\varepsilon,2}^{1/2}e^{-cN/2}\exp(\mathbb{E}\operatorname{Tr}f(\boldsymbol{G}))$$
$$\ge (1 - C_{\varepsilon,2}^{1/2}e^{-cN/2})\exp(\mathbb{E}\operatorname{Tr}f(\boldsymbol{G})).$$

Thus,

$$\frac{\mathbb{E}\left[|\det(r\boldsymbol{I}-\boldsymbol{G})|^{t}\right]^{1/t}}{\mathbb{E}\left[|\det(r\boldsymbol{I}-\boldsymbol{G})|\right]} \leqslant \frac{C_{\varepsilon,t}^{1/t}\exp(\mathbb{E}\operatorname{Tr} f(\boldsymbol{G}))}{(1-C_{\varepsilon,2}^{1/2}e^{-cN/2})\exp(\mathbb{E}\operatorname{Tr} f(\boldsymbol{G}))}$$

is bounded by a constant depending only on ε , t.

Lemma 6.21. Let $G \sim \text{GOE}(N)$. For all r in any compact subset of $(2, +\infty)$, there exists C > 0 such that

$$\mathbb{E}[|\det(r\boldsymbol{I} - \boldsymbol{G})|] \leq C \exp(N\Phi(r)),$$

where

$$\Phi(r) = \frac{1}{4}r^2 - \frac{1}{2} - \frac{1}{4}r\sqrt{r^2 - 4} + \log\frac{r + \sqrt{r^2 - 4}}{2}.$$

Proof. Follows from [**BČNS22**, Lemma 2.1 and 2.2(i)] with N+1 for N and $\sqrt{\frac{N}{2(N+1)}}r$ for x. Note that the matrix $GOE_{N-1}(N^{-1})$ therein is defined with typical spectral radius $\sqrt{2}$, while our GOE(N) has spectral radius 2.

Proof of Lemma 6.20. Throughout this proof, C > 0 is a constant, uniform over $m \in S_{\iota}$, which may change from line to line. Let $||m||_{N}^{2} = q_{m}$ and $\langle x, m \rangle_{N} = q_{x}$, so that $q_{m}, q_{x} \in [q_{*} - \iota, q_{*} + \iota]$. By Fact 6.18, for $G \sim \text{GOE}(N-2)$,

$$\begin{split} \mathbb{E}\left[\left|\det K(\boldsymbol{m})\right| \middle| \nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \boldsymbol{0}\right] &= \left(\xi''(q_{\boldsymbol{m}}) \cdot \frac{N-2}{N}\right)^{(N-2)/2} \mathbb{E}\left[\left|\det\left(\sqrt{\frac{N}{N-2}}r_{\boldsymbol{m}}\boldsymbol{I} - \boldsymbol{G}\right)\right|\right]\\ \text{where} \quad r_{\boldsymbol{m}} &= (1-q_{\boldsymbol{m}})\xi''(q_{\boldsymbol{m}})^{1/2} + \frac{1}{(1-q_{\boldsymbol{m}})\xi''(q_{\boldsymbol{m}})^{1/2}}. \end{split}$$

By (1.5), for $q_m \in [q_* - \iota, q_* + \iota]$, r_m takes values in a compact subset of $(2, +\infty)$. By Lemma 6.21,

$$\mathbb{E}\left[\left|\det K(\boldsymbol{m})\right| \middle| \nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \boldsymbol{0}\right] \leqslant C \exp(Nf_1(q_{\boldsymbol{m}})), \qquad f_1(q_{\boldsymbol{m}}) := \frac{1}{2}\log \xi''(q_{\boldsymbol{m}}) + \Phi(r_{\boldsymbol{m}}).$$

By (1.5),

$$\sqrt{r_{\boldsymbol{m}}^2 - 4} = \frac{1}{(1 - q_{\boldsymbol{m}})\xi''(q_{\boldsymbol{m}})^{1/2}} - (1 - q_{\boldsymbol{m}})\xi''(q_{\boldsymbol{m}})^{1/2}.$$

So, f_1 simplifies to

$$f_1(q_m) = \frac{1}{2}(1 - q_m)^2 \xi''(q_m) - \log(1 - q_m)$$

On the other hand, by (5.9),

$$\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \nabla \widetilde{H}_{N,t}(\boldsymbol{m}) + \xi'_t(q_{\boldsymbol{x}}) \left(\boldsymbol{x} - \frac{q_{\boldsymbol{x}}}{q_{\boldsymbol{m}}} \boldsymbol{m} \right) - \left((1 - q_{\boldsymbol{m}}) \xi''(q_{\boldsymbol{m}}) + \frac{1}{1 - q_{\boldsymbol{m}}} - \frac{q_{\boldsymbol{x}} \xi'_t(q_{\boldsymbol{x}})}{q_{\boldsymbol{m}}} \right) \boldsymbol{m}.$$

Since $x - \frac{q_x}{q_m}m$ is orthogonal to m, Fact 6.19 yields

$$\varphi_{\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})}(\boldsymbol{0}) \leqslant C \exp(N f_2(q_{\boldsymbol{m}}, q_{\boldsymbol{x}}))$$

where

$$f_2(q_m, q_x) = -\frac{1}{2} \log(2\pi\xi'_t(q_m)) - \frac{\xi'_t(q_x)^2}{2\xi'_t(q_m)} \left(1 - \frac{q_x^2}{q_m}\right) - \frac{q_m}{2(\xi'_t(q_m) + q_m\xi''(q_m))} \left((1 - q_m)\xi''(q_m) + \frac{1}{1 - q_m} - \frac{q_x\xi'_t(q_x)}{q_m}\right)^2.$$

Combining the above,

$$\int_{\mathcal{S}_{\iota}} \mathbb{E}\left[|\det K(\boldsymbol{m})| |\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \boldsymbol{0} \right] \varphi_{\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})}(\boldsymbol{0}) \, \mathsf{d}^{N}(\boldsymbol{m}) \\ \leq CN \int_{q_{*}-\iota}^{q_{*}+\iota} \int_{q_{*}-\iota}^{q_{*}+\iota} \exp\left(N(f_{1}(q_{\boldsymbol{m}}) + f_{2}(q_{\boldsymbol{m}}, q_{\boldsymbol{x}}) + f_{3}(q_{\boldsymbol{m}}, q_{\boldsymbol{x}}))\right) \, \mathsf{d}q_{\boldsymbol{x}} \mathsf{d}q_{\boldsymbol{m}}.$$
(6.14)

Here $CN \exp(Nf_3(q_m, q_x))$ is a volumetric factor and

$$f_3(q_m, q_x) = \frac{1}{2} + \frac{1}{2}\log(2\pi(q_m - q_x^2)).$$

Let

$$\begin{split} F(q_{m},q_{x}) &= f_{1}(q_{m}) + f_{2}(q_{m},q_{x}) + f_{3}(q_{m},q_{x}) \\ &= \frac{1}{2} + \frac{1}{2} \log \frac{q_{m} - q_{x}^{2}}{\xi_{t}'(q_{m})(1 - q_{m})^{2}} + \frac{1}{2}(1 - q_{m})^{2}\xi''(q_{m}) - \frac{\xi_{t}'(q_{x})^{2}}{2\xi_{t}'(q_{m})} \left(1 - \frac{q_{x}^{2}}{q_{m}}\right) \\ &- \frac{q_{m}}{2(\xi_{t}'(q_{m}) + q_{m}\xi''(q_{m}))} \left((1 - q_{m})\xi''(q_{m}) + \frac{1}{1 - q_{m}} - \frac{q_{x}\xi_{t}'(q_{x})}{q_{m}}\right)^{2}. \end{split}$$

To conclude, we will verify that $F(q_*, q_*) = 0$, $\nabla F(q_*, q_*) = 0$, and F is $\Omega(1)$ -strongly concave over $q_m, q_x \in [q_* - \iota, q_* + \iota]$. This will imply that the integral in (6.14) is $O(N^{-1})$ and finish the proof. Recall from (4.2) that $\xi'_t(q_*) = \frac{q_*}{1-q_*}$. The following identity will be used repeatedly in the calculations below to simplify the final term in F and its derivatives:

$$\frac{1}{\xi_t'(q_*) + q_*\xi''(q_*)} \left((1 - q_*)\xi''(q_*) + \frac{1}{1 - q_*} - \xi_t'(q_*) \right) = \frac{1 - q_*}{q_*}$$

Using this, we verify that

$$F(q_*, q_*) = \frac{1}{2} + \frac{1}{2}(1 - q_*)^2 \xi''(q_*) - \frac{q_*}{2} - \frac{q_*}{2(\frac{q_*}{1 - q_*} + q_*\xi''(q_*))} \left((1 - q_*)\xi''(q_*) + 1\right)^2$$
$$= \frac{1}{2}(1 - q_*)^2 \xi''(q_*) + \frac{1 - q_*}{2} - \frac{1 - q_*}{2} \left((1 - q_*)\xi''(q_*) + 1\right) = 0.$$

We also calculate

$$\begin{split} \frac{\partial F}{\partial q_{m}}(q_{m},q_{x}) &= \frac{1}{2(q_{m}-q_{x}^{2})} - \frac{\xi''(q_{m})}{2\xi'_{t}(q_{m})} + \frac{1}{1-q_{m}} - (1-q_{m})\xi''(q_{m}) + \frac{1}{2}(1-q_{m})^{2}\xi^{(3)}(q_{m}) \\ &+ \frac{\xi'_{t}(q_{x})^{2}\xi''(q_{m})}{2\xi'_{t}(q_{m})^{2}} \left(1 - \frac{q_{x}^{2}}{q_{m}}\right) - \frac{\xi'_{t}(q_{x})^{2}}{2\xi'_{t}(q_{m})}\frac{q_{x}^{2}}{q_{m}^{2}} \\ &- \frac{\xi'_{t}(q_{m}) - q_{m}\xi''(q_{m}) - q_{m}^{2}\xi^{(3)}(q_{m})}{2(\xi'_{t}(q_{m}) + q_{m}\xi''(q_{m}))^{2}} \left((1 - q_{m})\xi''(q_{m}) + \frac{1}{1-q_{m}} - \frac{q_{x}\xi'_{t}(q_{x})}{q_{m}}\right)^{2} \\ &- \frac{q_{m}}{\xi'_{t}(q_{m}) + q_{m}\xi''(q_{m})} \left((1 - q_{m})\xi''(q_{m}) + \frac{1}{1-q_{m}} - \frac{q_{x}\xi'_{t}(q_{x})}{q_{m}}\right) \\ &\times \left(-\xi''(q_{m}) + (1 - q_{m})\xi^{(3)}(q_{m}) + \frac{1}{(1-q_{m})^{2}} + \frac{q_{x}\xi'_{t}(q_{x})}{q_{m}^{2}}\right), \\ \frac{\partial F}{\partial q_{x}}(q_{m},q_{x}) &= -\frac{q_{x}}{q_{m} - q_{x}^{2}} - \frac{\xi'_{t}(q_{x})\xi''(q_{x})}{\xi'_{t}(q_{m})} \left(1 - \frac{q_{x}^{2}}{q_{m}}\right) + \frac{q_{x}\xi'_{t}(q_{x})^{2}}{q_{m}\xi'_{t}(q_{m})} \\ &+ \frac{\xi'_{t}(q_{x}) + q_{x}\xi''(q_{x})}{\xi'_{t}(q_{m})} \left((1 - q_{m})\xi''(q_{m}) + \frac{1}{1-q_{m}} - \frac{q_{x}\xi'_{t}(q_{x})}{q_{m}}\right). \end{split}$$

Thus

$$\begin{aligned} \frac{\partial F}{\partial q_m}(q_*,q_*) &= \frac{1}{2q_*(1-q_*)} - \frac{(1-q_*)\xi''(q_*)}{2q_*} + \frac{1}{1-q_*} - (1-q_*)\xi''(q_*) + \frac{1}{2}(1-q_*)^2\xi^{(3)}(q_*) \\ &+ \frac{(1-q_*)\xi''(q_*)}{2} - \frac{q_*}{2(1-q_*)} - \frac{(1-q_*)^2}{2q_*^2} \left(\frac{q_*}{1-q_*} - q_*\xi''(q_*) - q_*^2\xi^{(3)}(q_*)\right) \\ &- (1-q_*)\left(-\xi''(q_*) + (1-q_*)\xi^{(3)}(q_*) + \frac{1}{(1-q_*)^2} + \frac{1}{1-q_*}\right) = 0,\end{aligned}$$

and

$$\frac{\partial F}{\partial q_{\boldsymbol{x}}}(q_{\ast},q_{\ast}) = -\frac{1}{1-q_{\ast}} - (1-q_{\ast})\xi''(q_{\ast}) + \xi_{t}'(q_{\ast}) + \left((1-q_{\ast})\xi''(q_{\ast}) + \frac{1}{1-q_{\ast}} - \xi_{t}'(q_{\ast})\right) = 0.$$

By similar calculations, we find the following formulas for the second derivative. Let

$$\Delta_0 = \xi''(q_*) - \frac{1}{(1-q_*)^2} \stackrel{(1.5)}{<} 0$$

and

$$\Delta_1 = \frac{(1-q_*)^3 \Delta_0^3 - q_*(1-q_*) \Delta_0^2}{2q_*^2 (1+(1-q_*)\xi''(q_*))}, \qquad \Delta_2 = -\frac{(1-q_*)^2}{q_*} \Delta_0^2 + \Delta_0.$$

Then

$$\frac{\partial^2 F}{\partial q_{\boldsymbol{m}}^2}(q_*,q_*) = \Delta_1 + \Delta_2, \qquad \frac{\partial^2 F}{\partial q_{\boldsymbol{m}} \partial q_{\boldsymbol{x}}}(q_*,q_*) = -\Delta_2, \qquad \frac{\partial^2 F}{\partial q_{\boldsymbol{x}}^2}(q_*,q_*) = \Delta_2.$$

It follows that

$$\nabla^2 F(q_*, q_*) = \Delta_1(1, 0)^{\otimes 2} + \Delta_2(1, -1)^{\otimes 2} \le -CI_2$$

for some C > 0 depending only on ξ . Since $\nabla^2 F$ is clearly locally Lipschitz around $(q_*, q_*), \nabla^2 F(q_m, q_x) \leq -CI_2/2$ for all $q_m, q_x \in [q_* - \iota, q_* + \iota]$ for suitably small ι . This concludes the proof.

6.6 Algorithmic guarantees and Lipschitz continuity of correction

In this subsection, we prove Proposition 4.5.

Proof of Proposition 4.5(a). By (4.7),

$$\mu_t(\mathsf{Band}(\boldsymbol{m}^{\mathsf{AMP}}, [q_* - \iota/2, q_* + \iota/2])) \geqslant 1 - e^{-cN}.$$

Since $\|\boldsymbol{m}^{\mathsf{AMP}} - \boldsymbol{m}^{\mathsf{TAP}}\|_N \leqslant \iota/2$, we have

$$\mathsf{Band}(\boldsymbol{m}^{\mathsf{TAP}}, [q_* - \iota, q_* + \iota]) \supseteq \mathsf{Band}(\boldsymbol{m}^{\mathsf{AMP}}, [q_* - \iota/2, q_* + \iota/2]).$$

Proof of Proposition 4.5(b). On K_N , the maps $\boldsymbol{m} \mapsto \lambda_{\max}(\nabla^2 \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}))$ and $\boldsymbol{m} \mapsto \lambda_{\min}(\nabla^2 \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}_+))$ are O(1)-Lipschitz (over $\|\boldsymbol{m}\|_N \leq 1 - \varepsilon$, for any $\varepsilon > 0$). Combined with (4.9), this implies

$$\operatorname{spec}(\nabla^2 \mathcal{F}_{\operatorname{TAP}}(\boldsymbol{m})) \subseteq \left[-2C_{\max}^{\operatorname{spec}}, -\frac{1}{2}C_{\min}^{\operatorname{spec}}\right], \qquad \forall \|\boldsymbol{m} - \boldsymbol{m}^{\operatorname{TAP}}\|_N \leqslant \iota.$$

Thus $\nabla^2 \mathcal{F}_{TAP}$ is strongly concave and well-conditioned in the convex region $\|\boldsymbol{m} - \boldsymbol{m}^{TAP}\|_N \leq \iota$. It is classical (see e.g. [Nes03]) that for suitable $\eta > 0$, gradient descent

$$oldsymbol{u}^{k+1} = oldsymbol{u}^k - \eta
abla \mathcal{F}_{ extsf{TAP}}(oldsymbol{u}^k)$$

initialized from \boldsymbol{u}^0 in this region satisfies

$$\|\boldsymbol{u}^k-\boldsymbol{m}^{\mathrm{TAP}}\|_N\leqslant (1-\varepsilon)^k\|\boldsymbol{u}^0-\boldsymbol{m}^{\mathrm{TAP}}\|_N\leqslant \iota(1-\varepsilon)^k.$$

for some $\varepsilon > 0$. In particular $u^0 = m^{\text{AMP}}$ is in this region. Recalling $m^{\text{GD}} = u^{K_{\text{GD}}(N)}$ and $K_{\text{GD}}(N) = [K^*_{\text{GD}} \log N]$, we conclude

$$\|\boldsymbol{m}^{\mathsf{GD}}-\boldsymbol{m}^{\mathsf{TAP}}\|_N\leqslant\iota(1-\varepsilon)^{K_{\mathsf{GD}}(N)}\leqslant N^{-10}$$

for suitably large K_{GD}^* . This implies part (b).

We now turn to the proof of part (c). Recall from below (2.12) that I_{N-1} denotes the identity operator on T_m ; we sometimes write this as I_{N-1}^m to emphasize the dependence on m.

Lemma 6.22. Let $\gamma_* = (1-q_*)^{-1} + (1-q_*)\xi''(q_*)$. Let $\iota, \mathbf{m}^{TAP} \in S_\iota$ be as in Proposition 4.4. There exists $\iota' = o_\iota(1)$ such that with probability $1 - e^{-cN}$, (\mathbf{m}^{TAP} is defined and)

spec
$$A^{(2)}(m^{\text{TAP}}) \subseteq [-(2+\iota')\sqrt{\xi''(q_*)}, (2+\iota')\sqrt{\xi''(q_*)}]$$
 (6.15)

and

$$\left|\frac{1}{N}\operatorname{Tr}\left(\left(\gamma_{*}\boldsymbol{I}_{N-1}-\boldsymbol{A}^{(2)}(\boldsymbol{m}^{\mathsf{TAP}})\right)^{-1}\right)-(1-q_{*})\right|\leq\iota'.$$
(6.16)

Proof. Let \mathcal{E}_{spec} be the event that (6.15), (6.16) both hold, and let \mathcal{E} be as in Proposition 4.4. By Proposition 4.4(d) with $\delta = 1/2$,

$$\begin{split} \mathbb{P}(\mathcal{E}_{\mathsf{spec}}^c) &\leqslant \mathbb{P}(\mathcal{E}^c) + \mathbb{P}(\mathcal{E}_{\mathsf{spec}}^c \cap \mathcal{E}) \\ &\leqslant e^{-cN} + C_{1/2} \sup_{\boldsymbol{m} \in \mathcal{S}_{\iota}} \mathbb{P}\left[\mathcal{E}_{\mathsf{spec}}^c \cap \mathcal{E} \middle| \nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \mathbf{0}\right]^{1/2}. \end{split}$$

We will show that this probability is e^{-cN} , uniformly in $m \in S_{\iota}$. Note that on \mathcal{E} , we have deterministically $m^{TAP} = m$.

Let $q_{\boldsymbol{m}} = \|\boldsymbol{m}\|_{N}^{2} = [q_{*} - \iota, q_{*} + \iota]$. One checks analogously to Fact 6.5 that conditional on $\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \mathbf{0}$, we have $\boldsymbol{A}^{(2)}(\boldsymbol{m}) =_{d} \sqrt{\xi''(q_{\boldsymbol{m}})\frac{N-1}{N}}\boldsymbol{G}, \boldsymbol{G} \sim \mathsf{GOE}(N-1)$. It is classical that $\mathsf{spec}(\boldsymbol{G}) \subseteq [-2 - \iota, 2 + \iota]$ with probability $1 - e^{-cN}$, so (6.15) holds with conditional probability $1 - e^{-cN}$. Note that by (1.5),

$$\gamma_* - 2\sqrt{\xi''(q_*)} = \frac{1}{1 - q_*} \left(1 - (1 - q_*)\xi''(q_*)^{1/2} \right)^2 > 0.$$
(6.17)

So, for small enough ι , when (6.15) holds the matrix $\gamma_* I_{N-1} - A^{(2)}(m)$ is positive semidefinite with smallest eigenvalue bounded away from 0. Recall the semicircle measure

$$\rho_{\rm sc}(\lambda) = \frac{1}{2\pi} \sqrt{4 - \lambda^2} \, \mathrm{d}\lambda. \tag{6.18}$$

Applying [GZ00, Theorem 1.1(b)] as in the proof of Lemma 6.6 shows that with probability $1 - e^{-cN}$,

$$\left|\frac{1}{N}\operatorname{Tr}\left((\gamma_*\boldsymbol{I}_{N-1}-\boldsymbol{A}^{(2)}(\boldsymbol{m}))^{-1}\right)-\int\frac{\rho_{\operatorname{sc}}(\mathrm{d}\lambda)}{\gamma_*-\sqrt{\xi''(q_{\boldsymbol{m}})}\lambda}\right|\leqslant\iota.$$

This integral evaluates as

$$\int \frac{\rho_{\mathsf{sc}}(\mathsf{d}\lambda)}{\gamma_* - \sqrt{\xi''(q_*)\lambda}} + o_\iota(1) = 1 - q_* + o_\iota(1).$$

Thus, for suitable ι' , (6.16) holds with conditional probability $1 - e^{-cN}$, as desired.

Lemma 6.23. Suppose the event K_N in Proposition 3.6 holds. For any $\delta > 0$, there exists L such that for all $\delta \leq \|\boldsymbol{m}_1\|_N$, $\|\boldsymbol{m}_2\|_N \leq 1$, (treating $\boldsymbol{A}^{(2)}(\boldsymbol{m}_i)$ as a matrix in $\mathbb{R}^{N \times N}$, and $\boldsymbol{A}^{(3)}(\boldsymbol{m}_i)$ as a tensor in $(\mathbb{R}^N)^{\otimes 3}$)

$$egin{aligned} \|m{A}^{(2)}(m{m}_1) - m{A}^{(2)}(m{m}_2)\|_{ ext{op},N} \leqslant L \|m{m}_1 - m{m}_2\|_N, \ \|m{A}^{(3)}(m{m}_1) - m{A}^{(3)}(m{m}_2)\|_{ ext{op},N} \leqslant L \|m{m}_1 - m{m}_2\|_N. \end{aligned}$$

Proof. Let $\mathsf{P}_{\boldsymbol{m}}^{\perp}$ denote the projection operator to the orthogonal complement of \boldsymbol{m} . Then $\boldsymbol{A}^{(2)}(\boldsymbol{m}) = P_{\boldsymbol{m}}^{\perp} \nabla^2 H_N(\boldsymbol{m}) P_{\boldsymbol{m}}^{\perp}$. So,

$$\begin{split} \|\boldsymbol{A}^{(2)}(\boldsymbol{m}_{1}) - \boldsymbol{A}^{(2)}(\boldsymbol{m}_{2})\|_{\text{op},N} &\leq \|P_{\boldsymbol{m}_{1}}^{\perp} \nabla^{2} H_{N}(\boldsymbol{m}_{1}) P_{\boldsymbol{m}_{1}}^{\perp} - P_{\boldsymbol{m}_{1}}^{\perp} \nabla^{2} H_{N}(\boldsymbol{m}_{1}) P_{\boldsymbol{m}_{2}}^{\perp}\|_{\text{op},N} \\ &+ \|P_{\boldsymbol{m}_{1}}^{\perp} \nabla^{2} H_{N}(\boldsymbol{m}_{1}) P_{\boldsymbol{m}_{2}}^{\perp} - P_{\boldsymbol{m}_{2}}^{\perp} \nabla^{2} H_{N}(\boldsymbol{m}_{1}) P_{\boldsymbol{m}_{2}}^{\perp}\|_{\text{op},N} \\ &+ \|P_{\boldsymbol{m}_{2}}^{\perp} \nabla^{2} H_{N}(\boldsymbol{m}_{1}) P_{\boldsymbol{m}_{2}}^{\perp} - P_{\boldsymbol{m}_{2}}^{\perp} \nabla^{2} H_{N}(\boldsymbol{m}_{2}) P_{\boldsymbol{m}_{2}}^{\perp}\|_{\text{op},N} \\ &\leq 2 \|P_{\boldsymbol{m}_{1}}^{\perp} - P_{\boldsymbol{m}_{2}}^{\perp}\|_{\text{op},N} \max(\|\nabla^{2} H_{N}(\boldsymbol{m}_{1})\|_{\text{op},N}, \|\nabla^{2} H_{N}(\boldsymbol{m}_{2})\|_{\text{op},N}) \\ &+ \|\nabla^{2} H_{N}(\boldsymbol{m}_{1}) - \nabla^{2} H_{N}(\boldsymbol{m}_{2})\|_{\text{op},N} \end{split}$$

On event K_N ,

$$\begin{aligned} \|\nabla^2 H_N(\boldsymbol{m}_1)\|_{\text{op},N}, \|\nabla^2 H_N(\boldsymbol{m}_2)\|_{\text{op},N} &\leq C_2, \\ \|\nabla^2 H_N(\boldsymbol{m}_1) - \nabla^2 H_N(\boldsymbol{m}_2)\|_{\text{op},N} &\leq C_3 \|\boldsymbol{m}_1 - \boldsymbol{m}_2\|_N. \end{aligned}$$

Finally, for a constant C_{δ} depending on δ ,

$$\begin{split} \|P_{\boldsymbol{m}_{1}}^{\perp} - P_{\boldsymbol{m}_{2}}^{\perp}\|_{\text{op},N} &= \left\|\frac{\boldsymbol{m}_{1}\boldsymbol{m}_{1}^{\top}}{\|\boldsymbol{m}_{1}\|^{2}} - \frac{\boldsymbol{m}_{2}\boldsymbol{m}_{2}^{\top}}{\|\boldsymbol{m}_{2}\|^{2}}\right\|_{\text{op},N} \\ &\leqslant \left\|\frac{\boldsymbol{m}_{1}\boldsymbol{m}_{1}^{\top}}{\|\boldsymbol{m}_{1}\|^{2}} - \frac{\boldsymbol{m}_{1}\boldsymbol{m}_{1}^{\top}}{\|\boldsymbol{m}_{2}\|^{2}}\right\|_{\text{op},N} + \left\|\frac{\boldsymbol{m}_{1}\boldsymbol{m}_{1}^{\top}}{\|\boldsymbol{m}_{2}\|^{2}} - \frac{\boldsymbol{m}_{2}\boldsymbol{m}_{2}^{\top}}{\|\boldsymbol{m}_{2}\|^{2}}\right\|_{\text{op},N} \leqslant C_{\delta} \|\boldsymbol{m}_{1} - \boldsymbol{m}_{2}\|_{N}. \end{split}$$

This proves the inequality for $A^{(2)}$. The proof for $A^{(3)}$ is analogous.

s

Lemma 6.24. There exists L > 0 such that with probability $1 - e^{-cN}$, for all $m_1, m_2 \in \mathsf{B}_N(m^{\mathsf{TAP}}, \iota)$ (treating $Q(m_i)$ as a matrix in $\mathbb{R}^{N \times N}$)

$$egin{aligned} & \|oldsymbol{Q}(oldsymbol{m}_1)\|_{ ext{op},N} \leqslant L\,, \ & \|oldsymbol{Q}(oldsymbol{m}_1) - oldsymbol{Q}(oldsymbol{m}_2)\|_{ ext{op},N} \leqslant L\, \|oldsymbol{m}_1 - oldsymbol{m}_2\|_N \,\,. \end{aligned}$$

Proof. Suppose K_N holds and (6.15), (6.16) from Lemma 6.22 hold. Then, for some $\iota'' = o_\iota(1)$ and all $m \in \mathsf{B}_N(m^{\mathsf{TAP}}, \iota)$,

pec
$$A^{(2)}(m) \subseteq [-(2+\iota'')\sqrt{\xi''(q_*)}, (2+\iota'')\sqrt{\xi''(q_*)}]$$
 (6.19)

and

$$\left|\frac{1}{N}\operatorname{Tr}\left((\gamma_*\boldsymbol{I}_{N-1} - \boldsymbol{A}^{(2)}(\boldsymbol{m}))^{-1}\right) - (1 - q_*)\right| \leq \iota''.$$
(6.20)

When (6.19) holds, the calculation (6.17) shows γ_* is bounded away from spec $A^{(2)}(m)$. Thus,

$$\gamma \mapsto \frac{1}{N} \operatorname{Tr} \left((\gamma I_{N-1} - A^{(2)}(\boldsymbol{m}))^{-1} \right)$$

has derivative $\Omega(1)$ in a neighborhood of γ_* . It follows from (6.20) that $\gamma_{*,N}(\boldsymbol{m}) = \gamma_* + o_\iota(1)$ uniformly for all $\boldsymbol{m} \in \mathsf{B}_N(\boldsymbol{m}^{\mathsf{TAP}}, \iota)$. This is also bounded away from spec $\boldsymbol{A}^{(2)}(\boldsymbol{m})$, so

$$\|oldsymbol{Q}(oldsymbol{m})\|_{ ext{op},N} = \|(\gamma_{st,N}(oldsymbol{m})oldsymbol{I}_N - oldsymbol{A}^{(2)}(oldsymbol{m}))^{-1}\|_{ ext{op},N}$$

is bounded. Let $\boldsymbol{m}_1, \boldsymbol{m}_2 \in \mathsf{B}_N(\boldsymbol{m}^{\mathsf{TAP}}, \iota)$. There exists a rotation operator \boldsymbol{R} from $\mathsf{T}_{\boldsymbol{m}_1}$ to $\mathsf{T}_{\boldsymbol{m}_2}$ such that $\|\boldsymbol{R} - \boldsymbol{I}_{N-1}^{\boldsymbol{m}_1}\|_{\mathrm{op},N} \leq \|\boldsymbol{m}_1 - \boldsymbol{m}_2\|_N$. Recall $q_{\boldsymbol{m}} = \|\boldsymbol{m}\|_N^2$. The definition of $\gamma_{*,N}(\boldsymbol{m})$ implies

$$\begin{split} q_{m_2} - q_{m_1} &= \frac{1}{N} \mathsf{Tr} \left((\gamma_{*,N}(m_1) I_{N-1}^{m_1} - A^{(2)}(m_1))^{-1} - (\gamma_{*,N}(m_2) I_{N-1}^{m_1} - A^{(2)}(m_2))^{-1} \right) \\ &= \frac{1}{N} \mathsf{Tr} \left(Q(m_1) \left((\gamma_{*,N}(m_2) - \gamma_{*,N}(m_1)) I_N - (A^{(2)}(m_1) - R^{-1} A^{(2)}(m_2) R) \right) \\ & R^{-1} Q(m_2) R \right). \end{split}$$

Thus,

$$(\gamma_{*,N}(\boldsymbol{m}_{1}) - \gamma_{*,N}(\boldsymbol{m}_{2}))\frac{1}{N}\operatorname{Tr}(\boldsymbol{Q}(\boldsymbol{m}_{1})\boldsymbol{R}^{-1}\boldsymbol{Q}(\boldsymbol{m}_{2})\boldsymbol{R})$$

= $q_{\boldsymbol{m}_{2}} - q_{\boldsymbol{m}_{1}} + \frac{1}{N}\operatorname{Tr}\left(\boldsymbol{Q}(\boldsymbol{m}_{1})(\boldsymbol{A}^{(2)}(\boldsymbol{m}_{1}) - \boldsymbol{R}^{-1}\boldsymbol{A}^{(2)}(\boldsymbol{m}_{2})\boldsymbol{R})\boldsymbol{R}^{-1}\boldsymbol{Q}(\boldsymbol{m}_{2})\boldsymbol{R}\right).$ (6.21)

Note that

$$\|\boldsymbol{A}^{(2)}(\boldsymbol{m}_{1}) - \boldsymbol{R}^{-1}\boldsymbol{A}^{(2)\boldsymbol{R}}(\boldsymbol{m}_{2})\|_{\text{op},N} \leqslant \|\boldsymbol{A}^{(2)}(\boldsymbol{m}_{1})\|_{\text{op},N} \|\boldsymbol{I}_{N-1}^{\boldsymbol{m}_{1}} - \boldsymbol{R}\|_{\text{op},N} + \|\boldsymbol{A}^{(2)}(\boldsymbol{m}_{1}) - \boldsymbol{A}^{(2)}(\boldsymbol{m}_{2})\|_{\text{op},N},$$

and thus the absolute value of the right-hand side of (6.21) is upper bounded by

$$|q_{m_2} - q_{m_1}| + \|\boldsymbol{Q}(m_1)\|_{\text{op},N} \|\boldsymbol{Q}(m_2)\|_{\text{op},N} \|\boldsymbol{A}^{(2)}(m_1) - \boldsymbol{R}^{-1} \boldsymbol{A}^{(2)} \boldsymbol{R}(m_2)\|_{\text{op},N} \leq L \|\boldsymbol{m}_1 - \boldsymbol{m}_2\|_N,$$

by Lemma 6.23. As discussed above, $Q(m_1), R^{-1}Q(m_2)R \ge cI_{N-1}^{m_1}$ for some constant c > 0, so

$$\frac{1}{N}\operatorname{Tr}(\boldsymbol{Q}(\boldsymbol{m}_1)\boldsymbol{R}^{-1}\boldsymbol{Q}(\boldsymbol{m}_2)\boldsymbol{R}) \ge \frac{1}{N}\operatorname{Tr}((c\boldsymbol{I}_{N-1}^{\boldsymbol{m}_1})^2) \ge c^2/2$$

is bounded away from 0. It follows that, after adjusting L,

$$|\gamma_{*,N}(\boldsymbol{m}_1) - \gamma_{*,N}(\boldsymbol{m}_2)| \leqslant L \|\boldsymbol{m}_1 - \boldsymbol{m}_2\|_N$$

Finally, (adjusting L again)

$$\begin{split} \| \boldsymbol{Q}(\boldsymbol{m}_{1}) - \boldsymbol{Q}(\boldsymbol{m}_{2}) \|_{\text{op},N} \\ &= \left\| \boldsymbol{Q}(\boldsymbol{m}_{1}) \left((\gamma_{*,N}(\boldsymbol{m}_{1}) - \gamma_{*,N}(\boldsymbol{m}_{2})) \boldsymbol{I}_{N} - (\boldsymbol{A}^{(2)}(\boldsymbol{m}_{1}) - \boldsymbol{A}^{(2)}(\boldsymbol{m}_{2})) \right) \boldsymbol{Q}(\boldsymbol{m}_{2}) \right\|_{\text{op},N} \\ &\leq \| \boldsymbol{Q}(\boldsymbol{m}_{1}) \|_{\text{op},N} \| \boldsymbol{Q}(\boldsymbol{m}_{2}) \|_{\text{op},N} \left(|\gamma_{*,N}(\boldsymbol{m}_{1}) - \gamma_{*,N}(\boldsymbol{m}_{2})| + \| \boldsymbol{A}^{(2)}(\boldsymbol{m}_{1}) - \boldsymbol{A}^{(2)}(\boldsymbol{m}_{2}) \|_{\text{op},N} \right) \\ &\leq L \| \boldsymbol{m}_{1} - \boldsymbol{m}_{2} \|_{N}. \end{split}$$

<i>Proof of Proposition</i> $4.5(c)$. For	$r \text{ any } \ \boldsymbol{v}\ _2 = 1,$
--	--

$$\begin{aligned} 2|\langle \boldsymbol{\Delta}(\boldsymbol{m}_1) - \boldsymbol{\Delta}(\boldsymbol{m}_2), \boldsymbol{v} \rangle| &= |\langle \boldsymbol{A}^{(3)}(\boldsymbol{m}_1) \otimes \boldsymbol{v}, \boldsymbol{Q}(\boldsymbol{m}_1) \otimes \boldsymbol{Q}(\boldsymbol{m}_1) \rangle - \langle \boldsymbol{A}^{(3)}(\boldsymbol{m}_2) \otimes \boldsymbol{v}, \boldsymbol{Q}(\boldsymbol{m}_2) \otimes \boldsymbol{Q}(\boldsymbol{m}_2) \rangle| \\ &\leq |\langle (\boldsymbol{A}^{(3)}(\boldsymbol{m}_1) - \boldsymbol{A}^{(3)}(\boldsymbol{m}_2)) \otimes \boldsymbol{v}, \boldsymbol{Q}(\boldsymbol{m}_1) \otimes \boldsymbol{Q}(\boldsymbol{m}_1) \rangle| \\ &+ |\langle \boldsymbol{A}^{(3)}(\boldsymbol{m}_2) \otimes \boldsymbol{v}, (\boldsymbol{Q}(\boldsymbol{m}_1) - \boldsymbol{Q}(\boldsymbol{m}_2)) \otimes \boldsymbol{Q}(\boldsymbol{m}_1) \rangle| \\ &+ |\langle \boldsymbol{A}^{(3)}(\boldsymbol{m}_2) \otimes \boldsymbol{v}, \boldsymbol{Q}(\boldsymbol{m}_2) \otimes (\boldsymbol{Q}(\boldsymbol{m}_1) - \boldsymbol{Q}(\boldsymbol{m}_2)) \rangle|. \end{aligned}$$

By the previous two lemmas, this is bounded by $\frac{L}{\sqrt{N}} \|\boldsymbol{m}_1 - \boldsymbol{m}_2\|_N$, for some L > 0. Since this holds for all \boldsymbol{v} , we have $\|\boldsymbol{\Delta}(\boldsymbol{m}_1) - \boldsymbol{\Delta}(\boldsymbol{m}_2)\|_2 \leq \frac{L}{2\sqrt{N}}$, and thus $\|\boldsymbol{\Delta}(\boldsymbol{m}_1) - \boldsymbol{\Delta}(\boldsymbol{m}_2)\|_N \leq \frac{L}{2N}$.

7 Local computation of magnetization: proof of Proposition 4.6

Recall that $H_{N,t}(\sigma) = H_N(\sigma) + \langle y_t, \sigma \rangle$ with $y_t = tx + B_t$, and define

$$oldsymbol{x}^{\perp} = oldsymbol{x} - rac{\langle oldsymbol{x},oldsymbol{m}
angle_N}{\|oldsymbol{m}\|_N^2}oldsymbol{m}$$

as well as the bands (for $\|\boldsymbol{m}\|_N^2 = q$)

$$\mathsf{Band}_*(\iota) := \mathsf{Band}(\boldsymbol{m}, I(\iota)) \cap \mathsf{Band}(\boldsymbol{x}, I(\iota)), \quad I(\iota) := [q_* - \iota, q_* + \iota], \tag{7.1}$$

$$D_N(a,b) = \left\{ \boldsymbol{\sigma} \in S_N : \langle \boldsymbol{\sigma}, \boldsymbol{m} \rangle_N = aq, \, \langle \boldsymbol{\sigma}, \boldsymbol{x} \rangle_N = b \right\}.$$
(7.2)

We recall the definition of truncated magnetization from Proposition 4.6:

$$\widetilde{\boldsymbol{m}}_{2\iota}(\boldsymbol{m}) = \frac{\int_{\mathsf{Band}_{\ast}(2\iota)} \boldsymbol{\sigma} \exp(H_{N,t}(\boldsymbol{\sigma})) \,\mu_0(\mathsf{d}\boldsymbol{\sigma})}{\int_{\mathsf{Band}_{\ast}(2\iota)} \exp(H_{N,t}(\boldsymbol{\sigma})) \,\mu_0(\mathsf{d}\boldsymbol{\sigma})}.$$
(7.3)

In Sections 7.1 and 7.2 we will prove Proposition 4.6. For the readers' convenience, we reproduce the statement below.

Proposition 4.6. Define $A_2 := A^{(2)}(m)$, $A_3 := A^{(3)}(m)$ as per Eq. (2.11) and $\gamma_* = \gamma_{*,N}$ as per Eq. (2.12). Surther recall the definition of S_{ι} on Eq. (4.8), namely $S_{\iota} := \{m \in \mathbb{R}^N : |\langle m, x \rangle_N - q_*|, |\langle m, m \rangle_N - q_*| \leq \iota\}$. Then we have, for appropriate constant $\delta, \iota > 0$,

$$\sup_{\boldsymbol{m}\in\mathcal{S}_{\iota}} \mathbb{E}\left[\|\boldsymbol{m}+\boldsymbol{\Delta}(\boldsymbol{m})-\widetilde{\boldsymbol{m}}_{2\iota}(\boldsymbol{m})\|_{N}^{2+\delta} |\nabla\mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})=\boldsymbol{0}\right]$$

$$=\sup_{\boldsymbol{m}\in\mathcal{S}_{\iota}} \mathbb{E}\left[N^{-1-\delta/2}\left(\sum_{i=1}^{N}\left(\left[\widetilde{\boldsymbol{m}}_{2\iota}(\boldsymbol{m})-\boldsymbol{m}\right]_{i}-\left(\frac{1}{2}\langle\boldsymbol{A}_{3},\boldsymbol{Q}_{i,\cdot}\otimes\boldsymbol{Q}\rangle\right)\right)^{2}\right)^{1+\delta/2} \middle| \nabla\mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m})=\boldsymbol{0}\right]$$

$$\leq N^{-1-\delta},$$

$$\boldsymbol{Q}:=(\gamma_{*}\boldsymbol{I}-\boldsymbol{A}_{2})^{-1}.$$
(7.4)

Our approach to proving Proposition 4.6 is based on decomposing

$$\mathsf{Band}_{\ast}(2\iota) = \bigcup_{r,s \in I(2\iota)} \mathsf{Band}(\boldsymbol{m}, \{r\}) \cap \mathsf{Band}(\boldsymbol{x}, \{s\}) = \bigcup_{a,b \in L(2\iota)} D_N(a, b),$$
(7.6)

where, for $q = \|\boldsymbol{m}\|_N^2$, $c = \langle \boldsymbol{x}, \boldsymbol{m} \rangle_N$,

$$L(2\iota) = \{(a, b) : qa \in I(2\iota), b \in I(2\iota)\}$$

Note that for $m \in S_{\iota}$, we have $q, c \in I(\iota)$, and thus $L(2\iota)$ is a neighborhood of (0,0) of radius of order ι . For any $r, s \in I(2\iota)$, we will see that the Hamiltonian restricted to $\text{Band}(m, \{r\}) \cap \text{Band}(x, \{s\})$ (conditional on $\nabla \mathcal{F}_{\text{TAP}}(m) = 0$) is equivalent to that of a mixed *p*-spin model in its replica symmetric phase with a small magnetic field. We will therefore devote Section 7.1 to study this problem. In Section 7.2 we will use this result, and integrate it over a, b to prove Proposition 4.6.

7.1 Conditional magnetization per band

As anticipated, in this section we will compute a good approximation to the magnetization for general spherical models with small external field. While we will apply this result to the effective Hamiltonian in the band, hence in dimension N - 2, throughout this section, we adopt general notations for such a model, cf. Eq. (1.1) and recast N - 2 as N. We write

$$H(\boldsymbol{\sigma}) = \langle \boldsymbol{u}, \boldsymbol{\sigma} \rangle + H_{\geq 2}(\boldsymbol{\sigma}) = \langle \boldsymbol{u}, \boldsymbol{\sigma} \rangle + \sum_{p \geq 2} H_p(\boldsymbol{\sigma}), \qquad (7.7)$$

$$H_p(\boldsymbol{\sigma}) = \frac{1}{N^{(p-1)/2}} \sum_{i_1,\dots,i_p=1}^N g_{i_1,\dots,i_p} \sigma_{i_1}\dots\sigma_{i_p}, \qquad g_{i_1,\dots,i_p} \stackrel{i.i.d.}{\sim} \mathcal{N}(0,\beta_p^2).$$
(7.8)

We will write throughout $H_{\geq i}(\boldsymbol{\sigma}) = \sum_{p \geq i} H_p(\boldsymbol{\sigma})$. We recast the mixture of $H_{\geq 2}$ as $\xi(s) = \sum_{p \geq 2} \beta_p^2 s^p$.

The results of this subsection hold for all models satisfying the replica symmetry condition (2.21), which we reproduce for convenience:

$$\xi''(0) < 1, \qquad \xi(q) + q + \log(1 - q) < 0 \quad \forall q \in (0, 1).$$
(7.9)

This holds under the main condition (1.5) by integrating twice and, as we will see in (7.116), will hold for the effective model on the band $\text{Band}(\boldsymbol{m}, \{r\}) \cap \text{Band}(\boldsymbol{x}, \{s\})$, for all $r, s \in I(2\iota)$. Note that the first inequality in (7.9) implies $\beta_2^2 < 1/2$.

We will always assume $\|\boldsymbol{u}\|_2 \leq c_0 \sqrt{N}$, with c_0 a small constant, and in some lemmas $\|\boldsymbol{u}\|_2 \leq N^{c_0}$. We will denote by $\mu(d\boldsymbol{\sigma}) \propto \exp(H(\boldsymbol{\sigma})) \mu_0(d\boldsymbol{\sigma})$ the corresponding Gibbs measure.

Note that we can view H_2 as a quadratic form with $H_2(\sigma) = \langle \sigma, W^{(2)}\sigma \rangle$ (with entries $W_{ij}^{(2)} = (g_{ij} + g_{ji})/2\sqrt{N}$). Hence $W^{(2)}$ is a GOE matrix scaled by $\beta_2/\sqrt{2}$. We will work in the orthonormal basis diagonalizing $W^{(2)}$ and its the spectrum of be $\Lambda = (\Lambda_i)_{i \leq N}$, with $\Lambda_1 \geq \Lambda_2 \geq \cdots \geq \Lambda_N$. We will occasionally identify Λ with the diagonal matrix with diagonal entries Λ_i . We also write $W^{(3)}$ for the symmetric 3rd-order tensor such that $H_3(\sigma) = \langle W^{(3)}, \sigma^{\otimes 3} \rangle$, written in the basis of eigenvectors of $W^{(2)}$. That is, $W^{(3)}$ is obtained by rotating $\tilde{W}^{(3)}$ with entries $\tilde{W}_{ijk}^{(3)} = (g_{ijk} + \text{permutations})/6N$.

Given a symmetric matrix $\boldsymbol{A} \in \mathbb{R}^{N \times N}$, we define

$$G(\gamma) = G(\gamma; \boldsymbol{A}, \boldsymbol{u}) := \gamma - \frac{1}{2N} \log \det(\gamma \boldsymbol{I} - \boldsymbol{A}) + \frac{1}{4N} \langle \boldsymbol{u}, (\gamma \boldsymbol{I} - \boldsymbol{A})^{-1} \boldsymbol{u} \rangle,$$
(7.10)

$$\gamma_* = \gamma_*(\boldsymbol{A}, \boldsymbol{u}) = \operatorname*{arg\,min}_{\gamma > \lambda_{\max}(\boldsymbol{A})} G(\gamma; \boldsymbol{A}, \boldsymbol{u}) \,. \tag{7.11}$$

Note that G is convex with $\lim_{\gamma \downarrow \lambda_{\max}(A)} G'(\gamma) = -\infty$, $\lim_{\gamma \uparrow +\infty} G'(\gamma) = +\infty$, so γ_* is also the unique solution to $G'(\gamma) = 0$. We will omit the argument A or u whenever clear from the context (in particular, we typically omit u and omit A when $A = \Lambda$ is the diagonal matrix containing the eigenvalues of W).

Lemma 7.1. There exits $c_0 > 0$ such that, for $||u|| \leq N^{c_0}$, and under the additional assumptions above, the following holds. Let $\gamma_* = \gamma_*(\Lambda)$ and define, for $j \leq N$

$$\hat{m}_j := \frac{u_j}{2(\gamma_* - \Lambda_j)} + \frac{1}{2(\gamma_* - \Lambda_j)} \sum_{i=1}^N \frac{W_{jii}^{(3)}}{\gamma_* - \Lambda_i} \,. \tag{7.12}$$

Then, for some c > 0, with probability $1 - N^{-c}$, the following holds for all $i \in [N]$:

$$\int \sigma_i \,\mu(\mathsf{d}\boldsymbol{\sigma}) = (1 + O(N^{-c})) \Big(\hat{m}_i + O(N^{-1/2-c}) \Big).$$

Together with further estimates, we will use Lemma 7.1 to prove the following lemma, which is the main result of the section.

Lemma 7.2. Let $\alpha \ge 2$. There exists $c_0 > 0$ such that, for $||\mathbf{u}|| \le N^{c_0}$, and under the additional assumptions above, we have for some c > 0 that

$$\mathbb{E}\left[\left\|\int \boldsymbol{\sigma}\,\mu(\mathsf{d}\boldsymbol{\sigma}) - \widehat{\boldsymbol{m}}\right\|_{N}^{\alpha}\right] = O(N^{-\alpha/2-c}).$$
(7.13)

The rest of this section is devoted to the proof of Lemma 7.1 and Lemma 7.2.

7.1.1 Quadratic Hamiltonians

We begin by proving several supporting lemmas about quadratic models. The Laplace transform allows us to compute accurately various statistics of quadratic models. We note that the use of Laplace transforms in studying spherical quadratic models has been utilized before, for example for analyzing the fluctuation of the free energy in [BL16]. We will however need accurate control over a number of statistics beyond the free energy.

Lemma 7.3. For $\mathbf{A} \in \mathbb{R}^{N \times N}$ a GOE matrix scaled by $\alpha < 1/2$, and $\mathbf{u} \in \mathbb{R}^N$ such that $\|\mathbf{u}\|^2 \leq \varepsilon N$ for $\varepsilon > 0$ depending only on $1/2 - \alpha$, there exists a constant c > 0 such that, defining and $G(\gamma) = G(\gamma; \mathbf{A})$, $\gamma_* = \gamma_*(\mathbf{A})$, we have that the following claim holds with probability at least $1 - \exp(-cN)$:

$$\int e^{\langle \boldsymbol{\sigma}, \boldsymbol{A}\boldsymbol{\sigma} \rangle + \langle \boldsymbol{u}, \boldsymbol{\sigma} \rangle} \mu_0(\mathsf{d}\boldsymbol{\sigma}) = (1 + O(N^{-c})) \sqrt{\frac{2}{G''(\gamma_*)}} (2e)^{-N/2} e^{NG(\gamma_*)}, \tag{7.14}$$

and, for v_k any eigenvector of A (uniformly over k)

$$\frac{\int \langle \boldsymbol{v}_k, \boldsymbol{\sigma} \rangle e^{\langle \boldsymbol{\sigma}, \boldsymbol{A} \boldsymbol{\sigma} \rangle + \langle \boldsymbol{u}, \boldsymbol{\sigma} \rangle} \mu_0(\mathsf{d} \boldsymbol{\sigma})}{\int e^{\langle \boldsymbol{\sigma}, \boldsymbol{A} \boldsymbol{\sigma} \rangle + \langle \boldsymbol{u}, \boldsymbol{\sigma} \rangle} \mu_0(\mathsf{d} \boldsymbol{\sigma})} = (1 + O(N^{-c})) \frac{\langle \boldsymbol{v}_k, \boldsymbol{u} \rangle}{2(\gamma_* - \lambda_k(\boldsymbol{A}))} \,.$$
(7.15)

Proof. By a change of basis, we can assume that $A = \Lambda$ is diagonal (and its entries ordered). Let

$$E(\ell) := \ell^{N/2-1} \int \exp\left(\langle \boldsymbol{u}, \boldsymbol{\sigma} \rangle \sqrt{\ell} + \langle \boldsymbol{\Lambda}, \boldsymbol{\sigma}^{\otimes 2} \rangle \ell\right) \mu_0(\mathsf{d}\boldsymbol{\sigma}) \,. \tag{7.16}$$

Then the Laplace transform of E is given by

$$F(t) = \int_0^\infty e^{-t\ell} E(\ell) \,\mathrm{d}\ell,$$

and one has (for $\Re(\gamma) > \Lambda_1 = \max_{i \leq N} \Lambda_i$)

$$E(\ell) = \frac{1}{2\pi i} \int_{N\gamma - i\infty}^{N\gamma + i\infty} e^{t\ell} F(t) \,\mathrm{d}t$$

We evaluate, for $\Re(t) > \Lambda_1$,

$$\begin{split} F(Nt) &= \int_0^\infty \int e^{-Nt\ell} \exp\left(\langle \boldsymbol{u}, \boldsymbol{\sigma} \rangle \sqrt{\ell} + \langle \boldsymbol{\Lambda}, \boldsymbol{\sigma}^{\otimes 2} \rangle \ell\right) \ell^{N/2-1} \mu_0(\mathsf{d}\boldsymbol{\sigma}) \,\mathsf{d}\ell \\ &= \frac{\Gamma(N/2)}{(N\pi)^{N/2}} \int_{\mathbb{R}^N} \exp\left(-t \|\boldsymbol{y}\|^2 + \langle \boldsymbol{u}, \boldsymbol{y} \rangle + \langle \boldsymbol{\Lambda}, \boldsymbol{y}^{\otimes 2} \rangle\right) \mathsf{d}\boldsymbol{y} \\ &= \frac{\Gamma(N/2)}{(N\pi)^{N/2}} \int_{\mathbb{R}^N} \exp\left\{-\sum_{i=1}^N (t - \Lambda_i) y_i^2 + \sum_{i=1}^N u_i y_i\right\} \mathsf{d}\boldsymbol{y} \\ &= \frac{\Gamma(N/2)}{N^{N/2}} \exp\left\{-\frac{1}{2} \sum_{i=1}^N \log(t - \Lambda_i) + \sum_{i=1}^N \frac{u_i^2}{4(t - \Lambda_i)}\right\} \,. \end{split}$$

Hence, by the inverse Laplace transform, for all $\gamma \in \mathbb{R}$, $\gamma > \max_{i \leq N} \Lambda_i$,

$$E(1) = \frac{N}{2\pi i} \int_{\gamma-i\infty}^{\gamma+i\infty} e^{Nt} F(Nt) dt$$

$$= \frac{\Gamma(N/2)}{2\pi N^{N/2-1}} \int_{-\infty}^{\infty} \exp\left\{N(\gamma+iz) - \frac{1}{2} \sum_{i=1}^{N} \log(\gamma+iz-\Lambda_i) + \sum_{i=1}^{N} \frac{u_i^2}{4(\gamma+iz-\Lambda_i)}\right\} dz$$

$$= \frac{\Gamma(N/2)}{2\pi N^{N/2-1}} \int_{-\infty}^{\infty} \exp\left(NG(\gamma+iz)\right) dz,$$
(7.17)

where $G(x) = G(x; \mathbf{\Lambda})$ is defined as per Eq. (7.10).

Let γ_* be defined as per Eq. (7.11). Per the discussion below (7.11), γ_* is the unique solution to $G'(\gamma_*) = 0$. Explicitly,

$$N - \frac{1}{2} \sum_{i=1}^{N} \frac{1}{\gamma_* - \Lambda_i} - \sum_{i=1}^{N} \frac{u_i^2}{4(\gamma_* - \Lambda_i)^2} = 0.$$

Our assumption on α and $\|\boldsymbol{u}\|$ implies that $\gamma_* - \max_i \boldsymbol{\Lambda}_i > \delta$ for an appropriate δ depending only on $1/2 - \alpha$. We will set $\gamma = \gamma_*$ in Eq. (7.17). Note that

$$\Re(G(\gamma_* + iz) - G(\gamma_*)) = -\frac{1}{4N} \sum_{i=1}^N \log(1 + z^2/(\gamma_* - \Lambda_i)^2) - \frac{1}{4N} \sum_{i=1}^N \frac{u_i^2 z^2}{(\gamma_* - \Lambda_i)(z^2 + (\gamma_* - \Lambda_i)^2)}.$$

For $|z| \in ((\log N)/\sqrt{N}, 1)$, we have $\Re(G(\gamma_* + iz) - G(\gamma_*)) < -cz^2$, and for $|z| \ge 1$, we have $\Re(G(\gamma_* + iz) - G(\gamma_*)) < -c\log(1 + cz^2)$. This implies that

$$\int_{|z| > \frac{\log N}{\sqrt{N}}} \exp\left(NG(\gamma_* + iz) - NG(\gamma_*)\right) \mathrm{d}z < e^{-c(\log N)^2} \,.$$

On the other hand, for $|z| \leq (\log N)/\sqrt{N}$ we use the Taylor expansion:

$$G(\gamma_* + iz) = G(\gamma_*) + \sum_{j=1}^k \frac{(iz)^j}{j!} G^{(j)}(\gamma_*) + \operatorname{Err}_{N,k+1},$$

$$\operatorname{Err}_{N,k+1} \leq \frac{C}{k!} \left(\frac{\log N}{\sqrt{N}}\right)^k \sup_{|z| \leq (\log N)/\sqrt{N}} \left| G^{(k)}(\gamma_* + iz) \right|.$$

We have that

$$G^{(1)}(z) = 1 - \frac{1}{2N} \sum_{i=1}^{N} \frac{1}{z - \Lambda_i} - \frac{1}{4N} \sum_{i=1}^{N} \frac{u_i^2}{(z - \Lambda_i)^2},$$
$$G^{(j)}(z) = \frac{(-1)^j (j - 1)!}{2N} \sum_{i=1}^{N} \frac{1}{(z - \Lambda_i)^j} + \frac{(-1)^j j!}{4N} \sum_{i=1}^{N} \frac{u_i^2}{(z - \Lambda_i)^{j+1}}.$$

In particular, with probability $1 - \exp(-cN)$ over \boldsymbol{A} , $\sup_{|z| \leq (\log N)/\sqrt{N}} |G^{(j)}(\gamma_* + iz)| \leq j!C^j$ for a finite constant C > 0 as long as $\|\boldsymbol{u}\|^2 \leq N$. Hence, we have (for $\ell_N := (\log N)/\sqrt{N}$ and $J_N := [-\ell_N, \ell_N]$)

$$\begin{split} \int_{J_N} \exp\left(NG(\gamma_* + iz) - NG(\gamma_*)\right) \mathrm{d}z &= \int_{J_N} e^{-NG^{(2)}(\gamma_*)z^2/2} \, \exp\left(O(N\ell_N^3)\right) \, \mathrm{d}z \\ &= \sqrt{\frac{2\pi}{NG^{(2)}(\gamma_*)}} \left(1 + O(N^{-1/2+\varepsilon})\right). \end{split}$$

Together with Eq. (7.17), we get

$$E(1) = \frac{\Gamma(N/2)}{2\pi N^{N/2-1}} e^{NG(\gamma_*)} \left\{ \sqrt{\frac{2\pi}{NG^{(2)}(\gamma_*)}} + O(N^{-1+\varepsilon}) \right\} ,$$

which yields (7.14) by Stirling's formula.

By a similar argument, we obtain the integral of the spin $\sigma_k = \langle v_k, \sigma \rangle$ (recall we are working in the basis in which A is diagonal). Let

$$E_k(\ell) := \ell^{N/2-1} \int \sqrt{\ell} \sigma_k \exp\left(\langle \boldsymbol{u}, \boldsymbol{\sigma} \rangle \sqrt{\ell} + \langle \boldsymbol{\Lambda}, \boldsymbol{\sigma}^{\otimes 2} \rangle \ell\right) \mu_0(\mathsf{d}\boldsymbol{\sigma}).$$

Then the Laplace transform can be evaluated as

$$\begin{split} F_k(Nt) &= \int_0^\infty \int e^{-Nt\ell} \sqrt{\ell} \sigma_k \exp\left(\langle \boldsymbol{u}, \boldsymbol{\sigma} \rangle \sqrt{\ell} + \langle \boldsymbol{\Lambda}, \boldsymbol{\sigma}^{\otimes 2} \rangle \ell\right) \ell^{N/2-1} \mu_0(\mathsf{d}\boldsymbol{\sigma}) \, \mathsf{d}\ell \\ &= \frac{\Gamma(N/2)}{(N\pi)^{N/2}} \int_{\mathbb{R}^N} y_k \exp\left(-t \|\boldsymbol{y}\|^2 + \langle \boldsymbol{u}, \boldsymbol{y} \rangle + \langle \boldsymbol{\Lambda}, \boldsymbol{y}^{\otimes 2} \rangle\right) \, \mathsf{d}\boldsymbol{y} \\ &= \frac{\Gamma(N/2)}{N^{N/2}} \frac{u_k}{2(t-\Lambda_k)} \exp\left\{-\frac{1}{2} \sum_{i=1}^N \log(t-\Lambda_i) + \sum_{i=1}^N \frac{u_i^2}{4(t-\Lambda_i)}\right\}. \end{split}$$

Then we apply the same strategy as for computing E(1). By inverse Laplace transform:

$$E_k(1) = \frac{\Gamma(N/2)}{2\pi N^{N/2-1}} u_k \int_{-\infty}^{\infty} \frac{1}{2(\gamma_* + iz - \Lambda_k)} \exp\left(NG(\gamma_* + iz)\right) dz,$$
(7.18)

We make a negligible error in restricting to $J_N := [-\ell_N, \ell_N]$ (for $\ell_N := (\log N)/\sqrt{N}$)

$$E_k(1) = \frac{\Gamma(N/2)}{2\pi N^{N/2-1}} u_k \left\{ \int_{J_N} \frac{1}{2(\gamma_* + iz - \Lambda_k)} \exp\left(NG(\gamma_* + iz)\right) dz + O(e^{-c(\log N)^2}) \right\}$$
$$= \frac{\Gamma(N/2)}{2\pi N^{N/2-1}} u_k \left\{ \sqrt{\frac{2\pi}{NG^{(2)}(\gamma_*)}} \frac{1}{2(\gamma_* - \Lambda_k)} + O(N^{-1+\varepsilon}) \right\}.$$

Comparing with the above, we get

$$E_k(1) = (1 + O(N^{-c})) \frac{u_k}{2(\gamma_* - \Lambda_k)} \int e^{\langle \boldsymbol{\sigma}, \boldsymbol{A}\boldsymbol{\sigma} \rangle + \langle \boldsymbol{u}, \boldsymbol{\sigma} \rangle} \mu_0(\mathsf{d}\boldsymbol{\sigma}).$$

This gives (7.15).

Lemma 7.4. Let $\mathbf{A} \in \mathbb{R}^{N \times N}$ be a GOE matrix scaled by $\alpha < 1/2$. Assume that $\mathbf{u} \in \mathbb{R}^N$ is such that $\|\mathbf{u}\| \leq N^{c_0}$. For $\ell \in [L]$, consider a collection of pairs of indices (i_ℓ, j_ℓ) with $i_\ell \neq j_\ell \in [2k]$ and integers $r_\ell \geq 3$. Let $R = \sum_{\ell=1}^L r_\ell$. We have that the following claim holds with probability at least $1 - \exp(-cN)$: Uniformly in $h \in [N]$,

$$\frac{\int \prod_{i=1}^{2k} \boldsymbol{\sigma}_{h}^{i} \prod_{\ell=1}^{L} \langle \boldsymbol{\sigma}^{i_{\ell}}, \boldsymbol{\sigma}^{j_{\ell}} \rangle^{r_{\ell}} \exp\left(\sum_{i=1}^{2k} \langle \boldsymbol{u}, \boldsymbol{\sigma}^{i} \rangle + \langle \boldsymbol{\sigma}^{i}, \boldsymbol{\Lambda} \boldsymbol{\sigma}^{i} \rangle\right) \mu_{0}^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma})}{\int \exp\left(\sum_{i=1}^{2k} \langle \boldsymbol{u}, \boldsymbol{\sigma}^{i} \rangle + \langle \boldsymbol{\sigma}^{i}, \boldsymbol{\Lambda} \boldsymbol{\sigma}^{i} \rangle\right) \mu_{0}^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma})} = O_{k,L} \left(|u_{h}|^{2k-2\min(k,L)} N^{(R-\min(k,L))/2} (1 + ||\boldsymbol{u}||)^{2R} \right).$$
(7.19)

Proof. As before, we perform a change of basis and assume that $A = \Lambda$ is diagonal. Consider

$$E(z_1,\ldots,z_{2k})$$

$$:=(\prod_{i=1}^{2k}z_i)^{N/2-1}\int\prod_{i=1}^{2k}(\boldsymbol{\sigma}_h^i\sqrt{z_i})\prod_{\ell=1}^L\langle\boldsymbol{\sigma}^{i_\ell}\sqrt{z_{i_\ell}},\boldsymbol{\sigma}^{j_\ell}\sqrt{z_{j_\ell}}\rangle^{r_\ell}\exp\left(\sum_{i=1}^{2k}\langle\boldsymbol{u},\boldsymbol{\sigma}^i\rangle\sqrt{z_i}+\langle\boldsymbol{\sigma}^i,\boldsymbol{\Lambda}\boldsymbol{\sigma}^i\rangle z_i\right)\mu_0^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma}).$$

Then the multivariate Laplace transform of E is

$$\begin{split} F(N(t_1,\ldots,t_{2k})) &= \frac{\Gamma(N/2)^{2k}}{(N\pi)^{kN}} \int_{\mathbb{R}^N} \prod_{i=1}^{2k} y_h^i \prod_{\ell=1}^L \langle \boldsymbol{y}^{i_\ell}, \boldsymbol{y}^{j_\ell} \rangle^{r_\ell} \exp\left(-\sum_{i=1}^{2k} \left(t_i \|\boldsymbol{y}^i\|^2 - \langle \boldsymbol{u}, \boldsymbol{y}^i \rangle - \langle \boldsymbol{\Lambda}, (\boldsymbol{y}^i)^{\otimes 2} \rangle\right)\right) \mathrm{d}\boldsymbol{y} \\ &= \frac{\Gamma(N/2)^{2k}}{N^{kN}} \exp\left\{-\sum_{i=1}^{2k} \left(\frac{1}{2} \sum_{h'=1}^N \log(t_i - \Lambda_{h'}) + \sum_{h'=1}^N \frac{u_{h'}^2}{4(t_i - \Lambda_{h'})}\right)\right\} \cdot \mathfrak{G}, \end{split}$$

where, for $y^i = \frac{1}{2}(t_i - \Lambda)^{-1}u + w^i$ and w^i independently distributed according $\mathcal{N}\left(0, \frac{1}{2}(t_i - \Lambda)^{-1}\right)$,

$$\mathfrak{G} := \mathbb{E}\left[\prod_{i=1}^{2k} y_h^i \prod_{\ell=1}^{L} \langle \boldsymbol{y}^{i_{\ell}}, \boldsymbol{y}^{j_{\ell}} \rangle^{r_{\ell}}\right]$$

$$= \mathbb{E}\left[\prod_{i=1}^{2k} \left(\frac{u_h}{2(t_i - \Lambda_h)} + w_h^i\right)\right]$$

$$\prod_{\ell=1}^{L} \left(\langle \boldsymbol{w}^{i_{\ell}}, \boldsymbol{w}^{j_{\ell}} \rangle + \frac{1}{2} \langle \boldsymbol{w}^{i_{\ell}}, (t_{j_{\ell}} - \boldsymbol{\Lambda})^{-1} \boldsymbol{u} \rangle + \frac{1}{2} \langle \boldsymbol{w}^{j_{\ell}}, (t_{i_{\ell}} - \boldsymbol{\Lambda})^{-1} \boldsymbol{u} \rangle + \frac{1}{4} \langle (t_{j_{\ell}} - \boldsymbol{\Lambda})^{-1} \boldsymbol{u}, (t_{i_{\ell}} - \boldsymbol{\Lambda})^{-1} \boldsymbol{u} \rangle\right]$$
(7.20)

Assume that $\min_{i \in [2k]} \Re(t_i - \max_{h'} \Lambda_{h'}) > c > 0$, and recall that $\|\boldsymbol{u}\| \leq N^{c_0}$. Define \mathcal{R} a tuple of sets of length 2k + R, where, for $1 \leq a \leq 2k$, \mathcal{R}_a is a subset of $\{a\}$, and for a > 2k, \mathcal{R}_a is a subset of the pair of indices $\{i_\ell, j_\ell\}$ in the corresponding term. As such, each tuple \mathcal{R} represents a term in the expansion of (7.25). If there exists an index in [2k] that appears an odd number of times among the sets in \mathcal{R} , then the contribution of the corresponding term to (7.25) is 0. Consider the tuples \mathcal{R} where each index appears an even number of

times. Let $B(\mathcal{R})$ be the collection of indices $a \leq 2k$ where $\mathcal{R}_a = \{a\}$, and let $b(\mathcal{R}) = |B(\mathcal{R})|$. The indices $B(\mathcal{R})$ must appear an odd number of times among the remaining sets $(\mathcal{R}_j)_{j=2k+1}^{2k+R}$. In each possible way to pick out terms among $(\mathcal{R}_j)_{j=2k+1}^{2k+R}$ so that each index in $B(\mathcal{R})$ appears at least once, let $d(\mathcal{R})$ denote the number of sets $|\mathcal{R}_j| = 1$ among these terms. Among the remaining terms, each of the index in $B(\mathcal{R})$ not covered by the $d(\mathcal{R})$ sets can be matched to terms among $(\mathcal{R}_j)_{j=2k+1}^{2k+R}$. Consider an arbitrary way to pair up all remaining indices appearing in the terms into pairs; let $f(\mathcal{R}) \leq R - d(\mathcal{R}) - (b(\mathcal{R}) - d(\mathcal{R}))/2$ be the number of such pairs. For each such term and fixed pairing, we can upper bound its contribution to (7.25) by

$$O\left(|u_h|^{2k-b(\mathcal{R})+d(\mathcal{R})}N^{f(\mathcal{R})/2}(1+\|\boldsymbol{u}\|^2)^R\right) \leq O\left(\max_{a \leq \min(k,L)} |u_h|^{2k-2a}N^{(R-a)/2}(1+\|\boldsymbol{u}\|)^{2R}\right),$$

noting the constraints $0 \leq d(\mathcal{R}) \leq b(\mathcal{R})$, $f(\mathcal{R}) \leq R - d(\mathcal{R}) - (b(\mathcal{R}) - d(\mathcal{R}))/2$. Thus, we have

$$\mathfrak{G} = O_{k,L} \left(\max_{a \leq \min(k,L)} |u_h|^{2k-2a} N^{(R-a)/2} (1 + ||\boldsymbol{u}||)^{2R} \right).$$

Hence, for $R = \sum_{\ell} r_{\ell} \ge 3L$,

$$\mathfrak{G} = O_{k,L} \left(|u_h|^{2k - 2\min(k,L)} N^{(R-\min(k,L))/2} (1 + ||\boldsymbol{u}||)^{2R} \right).$$

Taking the inverse Laplace transform and integrating on $t_i = \gamma_* + ix_i$, for γ_* defined in Eq. (7.11), noting that, similar to Lemma 7.3, we can restrict the integration to the range $x_i \in [-\ell_N, \ell_N]$ for $\ell_N = (\log N)/\sqrt{N}$, we obtain that

$$\frac{\int \prod_{i=1}^{2k} \boldsymbol{\sigma}_{h}^{i} \prod_{\ell=1}^{L} \langle \boldsymbol{\sigma}^{i_{\ell}}, \boldsymbol{\sigma}^{j_{\ell}} \rangle^{r_{\ell}} \exp\left(\sum_{i=1}^{2k} \langle \boldsymbol{u}, \boldsymbol{\sigma}^{i} \rangle + \langle \boldsymbol{\sigma}^{i}, \boldsymbol{\Lambda} \boldsymbol{\sigma}^{i} \rangle\right) \mu_{0}^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma})}{\int \exp\left(\sum_{i=1}^{2k} \langle \boldsymbol{u}, \boldsymbol{\sigma}^{i} \rangle + \langle \boldsymbol{\sigma}^{i}, \boldsymbol{\Lambda} \boldsymbol{\sigma}^{i} \rangle\right) \mu_{0}^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma})} = O_{k,L} \left(|u_{h}|^{2k-2\min(k,L)} N^{(R-\min(k,L))/2} (1 + \|\boldsymbol{u}\|)^{2R} \right).$$

The next lemma states that, under a purely quadratic Hamiltonian, and for small field, the overlap concentrates near zero.

Lemma 7.5 (Overlap concentration in quadratic models). Define

$$A_2(t)^c := \{(\boldsymbol{\sigma}^1, \boldsymbol{\sigma}^2) \in S_N \times S_N : |\langle \boldsymbol{\sigma}^1, \boldsymbol{\sigma}^2 \rangle_N | \ge t \}.$$

Assuming that $\|\boldsymbol{u}\|^2 \leq \delta N$ for δ sufficiently small, we have for some constant c > 0 that, with probability $1 - e^{-cN}$,

$$\frac{\int_{A_2(t)^c} \exp(H_{\leq 2}(\boldsymbol{\sigma}^1) + H_{\leq 2}(\boldsymbol{\sigma}^2)) \mu_0^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma})}{\int \exp(H_{\leq 2}(\boldsymbol{\sigma}^1) + H_{\leq 2}(\boldsymbol{\sigma}^2)) \mu_0^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma})} \leq \exp\Big\{-cN\big(t - \|\boldsymbol{u}\|_N\big)_+^2\Big\}.$$
(7.21)

Proof. Consider the Hamiltonian $H(\sigma^1, \sigma^2) = H_{\leq 2}(\sigma^1) + H_{\leq 2}(\sigma^2) + 2\theta \langle \sigma^1, \sigma^2 \rangle$. Let Λ_i be the eigenvalues of the quadratic component A of H. Using the Laplace transform as in Lemma 7.3,

$$\int \exp\left(\langle \boldsymbol{u}, \boldsymbol{\sigma}^{1} + \boldsymbol{\sigma}^{2} \rangle + \langle \boldsymbol{A}, (\boldsymbol{\sigma}^{1})^{\otimes 2} + (\boldsymbol{\sigma}^{2})^{\otimes 2} \rangle\right) \exp\left(2\theta \langle \boldsymbol{\sigma}^{1}, \boldsymbol{\sigma}^{2} \rangle\right) \mu_{0}^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma})$$
$$= \left(\frac{\Gamma(N/2)}{(2\pi)N^{N/2-1}}\right)^{2} \int_{-\infty}^{\infty} \exp\left(2N\tilde{G}_{\theta}(\gamma_{1} + iz_{1}, \gamma_{2} + iz_{2})\right) \mathsf{d}z_{1}\mathsf{d}z_{2},$$

where

$$\tilde{G}_{\theta}(z_1, z_2) = \frac{z_1 + z_2}{2} - \frac{1}{4N} \sum_{i=1}^N \log((z_1 - \Lambda_i)(z_2 - \Lambda_i) - \theta^2) + \frac{1}{8N} \sum_{i=1}^N \frac{u_i^2(z_1 + z_2 - 2\Lambda_i + 2\theta)}{(z_1 - \Lambda_i)(z_2 - \Lambda_i) - \theta^2}$$

We also denote

$$G_{\theta}(z) = G_{\theta}(z, z) = z - \frac{1}{4N} \sum_{i=1}^{N} \log((z - \Lambda_i)^2 - \theta^2) + \frac{1}{4N} \sum_{i=1}^{N} \frac{u_i^2(z - \Lambda_i + \theta)}{(z - \Lambda_i)^2 - \theta^2}.$$

Let $\gamma_*(\theta)$ be a stationary point of G_{θ} on \mathbb{R} so that $\gamma_*(\theta) > \max \Lambda_i + \theta$ (there exists a unique such point), and $\gamma_* = \gamma_*(0)$.

As in Lemma 7.3, noting that

$$\exp\left(2\Re\left(\log((\gamma_{*}(\theta) - \Lambda_{i} + iz_{1})(\gamma_{*}(\theta) - \Lambda_{i} + iz_{2}) - \theta^{2}\right) - \log((\gamma_{*}(\theta) - \Lambda_{i})^{2} - \theta^{2})\right)\right)$$

$$= \left(\frac{(\gamma_{*}(\theta) - \Lambda_{i})^{2} - \theta^{2} - z_{1}z_{2}}{(\gamma_{*}(\theta) - \Lambda_{i})^{2} - \theta^{2}}\right)^{2} + \left(\frac{(\gamma_{*}(\theta) - \Lambda_{i})(z_{1} + z_{2})}{(\gamma_{*}(\theta) - \Lambda_{i})^{2} - \theta^{2}}\right)^{2}$$

$$= \frac{((\gamma_{*}(\theta) - \Lambda_{i})^{2} - \theta^{2})^{2} + (z_{1}z_{2})^{2} + 2\theta^{2}z_{1}z_{2} + (z_{1}^{2} + z_{2}^{2})(\gamma_{*}(\theta) - \Lambda_{i})^{2}}{((\gamma_{*}(\theta) - \Lambda_{i})^{2} - \theta^{2})^{2}}$$

$$\geq 1 + \frac{z_{1}^{2} + z_{2}^{2}}{(\gamma_{*}(\theta) - \Lambda_{i})^{2} - \theta^{2}}.$$
(7.22)

Furthermore,

$$\Re\left(\frac{(\gamma_{*}(\theta) - \Lambda_{i} + \theta) + i(z_{1} + z_{2})/2}{N((\gamma_{*}(\theta) - \Lambda_{i})^{2} - \theta^{2} - z_{1}z_{2} + i(\gamma_{*}(\theta) - \Lambda_{i})(z_{1} + z_{2}))} - \frac{1}{N(\gamma_{*}(\theta) - \Lambda_{i} - \theta)}\right) \\
= \frac{-(z_{1}z_{2})^{2} - \theta(\gamma_{*}(\theta) - \Lambda_{i} + \theta)(z_{1} + z_{2})^{2}/2 - ((\gamma_{*}(\theta) - \Lambda_{i})^{2} - \theta^{2})(z_{1}^{2} + z_{2}^{2})/2}{N(\gamma_{*}(\theta) - \Lambda_{i} - \theta)(((\gamma_{*}(\theta) - \Lambda_{i})^{2} - \theta^{2} - z_{1}z_{2})^{2} + ((\gamma_{*}(\theta) - \Lambda_{i})(z_{1} + z_{2}))^{2})} \\
\leqslant 0.$$
(7.23)

Given (7.22) and (7.23), we can proceed as in Lemma 7.3 to restrict the integral over z_1 and z_2 to the range $|z_1|, |z_2| < (\log N)/\sqrt{N}$, incurring an error $e^{-c(\log N)^2}$. Then by similarly expanding around $(\gamma_*(\theta), \gamma_*(\theta))$, we obtain

$$\int \exp\left(\langle \boldsymbol{u}, \boldsymbol{\sigma}^{1} + \boldsymbol{\sigma}^{2} \rangle + \langle \boldsymbol{A}, (\boldsymbol{\sigma}^{1})^{\otimes 2} + (\boldsymbol{\sigma}^{2})^{\otimes 2} \rangle\right) \exp\left(2\theta \langle \boldsymbol{\sigma}^{1}, \boldsymbol{\sigma}^{2} \rangle\right) \mu_{0}^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma})$$
$$= \left(\frac{\Gamma(N/2)}{(2\pi)N^{N/2-1}}\right)^{2} e^{2NG_{\theta}(\gamma_{*}(\theta))} \left\{\frac{2\pi}{N \det(\nabla^{2}\tilde{G}_{\theta}(\gamma_{*}(\theta), \gamma_{*}(\theta)))^{1/2}} + O(N^{-3/2+\epsilon})\right\}$$

When $\|\boldsymbol{u}\|^2 \leq \delta N$, we have for G as in (7.10) that

$$G_{\theta}(\gamma_*(\theta)) - G(\gamma_*) = O(\theta^2) + O(\theta \|\boldsymbol{u}\|^2 / N).$$

On the other hand, by Lemma 7.3,

$$\int \exp(\langle \boldsymbol{u}, \boldsymbol{\sigma}^{1} + \boldsymbol{\sigma}^{2} \rangle + \langle \boldsymbol{A}, (\boldsymbol{\sigma}^{1})^{\otimes 2} + (\boldsymbol{\sigma}^{2})^{\otimes 2} \rangle) \mu_{0}^{\otimes 2}(\mathrm{d}\boldsymbol{\sigma})$$
$$= (1 + O(N^{-c})) \left(\frac{\Gamma(N/2)}{(2\pi)N^{N/2-1}}\right)^{2} \left(\sqrt{\frac{2\pi}{NG''(\gamma_{*})}} + O(N^{-3/2+\varepsilon})\right)^{2}.$$

In particular,

$$\frac{\int_{t \leq |\langle \boldsymbol{\sigma}^{1}, \boldsymbol{\sigma}^{2} \rangle_{N}|} \exp(H_{\leq 2}(\boldsymbol{\sigma}^{1}) + H_{\leq 2}(\boldsymbol{\sigma}^{2}))\mu_{0}^{\otimes 2}(\mathrm{d}\boldsymbol{\sigma})}{\int_{\boldsymbol{\sigma}^{1}, \boldsymbol{\sigma}^{2}} \exp(H_{\leq 2}(\boldsymbol{\sigma}^{1}) + H_{\leq 2}(\boldsymbol{\sigma}^{2}))\mu_{0}^{\otimes 2}(\mathrm{d}\boldsymbol{\sigma})} \\
\leq \exp(-2N\theta t) \frac{\int_{\boldsymbol{\sigma}^{1}, \boldsymbol{\sigma}^{2}} \exp(\langle \boldsymbol{u}, \boldsymbol{\sigma}^{1} + \boldsymbol{\sigma}^{2} \rangle + N\langle \boldsymbol{A}, (\boldsymbol{\sigma}^{1})^{\otimes 2} + (\boldsymbol{\sigma}^{2})^{\otimes 2} \rangle) \exp\left(2\theta \langle \boldsymbol{\sigma}^{1}, \boldsymbol{\sigma}^{2} \rangle\right) \mu_{0}^{\otimes 2}(\mathrm{d}\boldsymbol{\sigma})}{\int_{\boldsymbol{\sigma}^{1}, \boldsymbol{\sigma}^{2}} \exp(\langle \boldsymbol{u}, \boldsymbol{\sigma}^{1} + \boldsymbol{\sigma}^{2} \rangle + N\langle \boldsymbol{A}, (\boldsymbol{\sigma}^{1})^{\otimes 2} + (\boldsymbol{\sigma}^{2})^{\otimes 2} \rangle) \mu_{0}^{\otimes 2}(\mathrm{d}\boldsymbol{\sigma})} \\
\leq \exp\left(-2N\theta t + O(N\theta^{2}) + O(\theta \|\boldsymbol{u}\|^{2})\right).$$
imizing over θ , we obtain Eq. (7.21).

Optimizing over θ , we obtain Eq. (7.21).

We will also need the following lemma, giving an accurate expansion of moments of overlaps in perturbations of quadratic Hamiltonians.

Lemma 7.6. Let $\mathbf{A} \in \mathbb{R}^{N \times N}$ be a GOE matrix scaled by $\alpha < 1/2$ with eigenvalues given by $\mathbf{\Lambda}$. Let $\Delta \in \mathbb{R}^{N \times N}$ be an independent GOE matrix scaled by $\beta > 0$ and $|\zeta_1|, |\zeta_2| \leq C(\log N)/\sqrt{N}$. For i = 1, 2, let $\tilde{\Lambda}_i = \Lambda + \zeta_i \Delta$. Assume that $u \in \mathbb{R}^N$ is such that $||u|| \leq N^{c_0}$. Let $r \geq 0$ and L > 0. We have that the following claim holds with probability at least $1 - \exp(-cN)$: There exist $C_{i,j} = O_{r,L}(\|\boldsymbol{u}\|^{2r} + N^{\lfloor r/2 \rfloor})$ for $i, j \leq L$ such that

$$\frac{\int \langle \boldsymbol{\sigma}^{1}, \boldsymbol{\sigma}^{2} \rangle^{r} \exp\left(\sum_{i=1}^{2} \langle \boldsymbol{u}, \boldsymbol{\sigma}^{i} \rangle + \langle \boldsymbol{\sigma}^{i}, \tilde{\boldsymbol{\Lambda}}_{i} \boldsymbol{\sigma}^{i} \rangle\right) \mu_{0}^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma})}{\int \exp\left(\sum_{i=1}^{2} \langle \boldsymbol{u}, \boldsymbol{\sigma}^{i} \rangle + \langle \boldsymbol{\sigma}^{i}, \boldsymbol{\Lambda} \boldsymbol{\sigma}^{i} \rangle\right) \mu_{0}^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma})} = C_{0,0} + \sum_{i,j=0, (i,j)\neq (0,0)}^{L} C_{i,j} \zeta_{1}^{i} \zeta_{2}^{j} + O_{L}(N^{-L/2} + e^{-N^{c}}).$$
(7.24)

Proof. Consider

$$E(z_1, z_2) := (\prod_{i=1}^2 z_i)^{N/2-1} \int \langle \boldsymbol{\sigma}^1 \sqrt{z_1}, \boldsymbol{\sigma}^2 \sqrt{z_2} \rangle^r \exp\left(\sum_{i=1}^2 \langle \boldsymbol{u}, \boldsymbol{\sigma}^i \rangle \sqrt{z_i} + \langle \boldsymbol{\sigma}^i, \tilde{\boldsymbol{\Lambda}}_i \boldsymbol{\sigma}^i \rangle z_i\right) \mu_0^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma})$$

Then the multivariate Laplace transform of E is

$$F(N(t_1, t_2)) = \frac{\Gamma(N/2)^2}{(N\pi)^N} \int_{\mathbb{R}^N} \langle \boldsymbol{y}^1, \boldsymbol{y}^2 \rangle^r \exp\left(-\sum_{i=1}^2 \left(t_i \|\boldsymbol{y}^i\|^2 - \langle \boldsymbol{u}, \boldsymbol{y}^i \rangle - \langle \tilde{\boldsymbol{\Lambda}}_i, (\boldsymbol{y}^i)^{\otimes 2} \rangle\right)\right) d\boldsymbol{y}$$
$$= \frac{\Gamma(N/2)^2}{N^N} \exp\left\{-\sum_{i=1}^2 \left(\frac{1}{2} \log \det(t_i \boldsymbol{I}_N - \tilde{\boldsymbol{\Lambda}}_i) + \sum_{h'=1}^N \frac{1}{4N} \langle (t_i \boldsymbol{I}_N - \tilde{\boldsymbol{\Lambda}}_i)^{-1}, \boldsymbol{u} \boldsymbol{u}^T \rangle\right)\right\} \cdot \mathfrak{G}(\sigma)$$

where, for $Z^i = (t_i I_N - \tilde{\Lambda}_i)^{-1}$, $y^i = \frac{1}{2} Z^i u + w^i$ and w^i independently distributed according $\mathcal{N}\left(0, \frac{1}{2} Z^i\right)$,

$$\mathfrak{G}(t_1, t_2; \sigma_1, \sigma_2) := \mathbb{E}\left[\langle \boldsymbol{y}^1, \boldsymbol{y}^2 \rangle^r\right] \\= \mathbb{E}\left[\left(\langle \boldsymbol{w}^1, \boldsymbol{w}^2 \rangle + \frac{1}{2} \langle \boldsymbol{w}^1, \boldsymbol{Z}^2 \boldsymbol{u} \rangle + \frac{1}{2} \langle \boldsymbol{w}^2, \boldsymbol{Z}^1 \boldsymbol{u} \rangle + \frac{1}{4} \langle \boldsymbol{Z}^1 \boldsymbol{u}, \boldsymbol{Z}^2 \boldsymbol{u} \rangle\right)^r\right].$$
(7.25)

Let $\mathfrak{G}_0(t_1, t_2) = \mathfrak{G}(t_1, t_2; 0, 0)$. Note that \mathfrak{G} is a rational function of t_i , and hence extends to complex values of t_i . We next consider the Taylor expansion in ζ_1, ζ_2 of \mathfrak{G} . Write $w^i = (\frac{1}{2}Z^i)^{1/2} \tilde{w}^i$ for $\tilde{w}^i \sim \mathcal{N}(0, I_N)$. Note that

$$\|\partial^i_{\zeta_1} Z^1\|_{\mathrm{op}} = O_i(\beta).$$

We can thus bound the derivatives of $\mathfrak{G}(t_1, t_2; \zeta_1, \zeta_2)$ for $|t_1 - \gamma_*|, |t_2 - \gamma_*| \leq C(\log N)/\sqrt{N}$ as

$$\left|\partial_{\zeta_{1}}^{i}\partial_{\zeta_{2}}^{j}\mathfrak{G}(t_{1},t_{2};\zeta_{1},\zeta_{2})\right| \leq O_{r,i+j}\left(\|\boldsymbol{u}\|^{2r}+N^{r/2}\right)$$
(7.26)

for r even, and

$$\left|\partial_{\zeta_{1}}^{i}\partial_{\zeta_{2}}^{j}\mathfrak{G}(t_{1},t_{2};\zeta_{1},\zeta_{2})\right| \leq O_{r,i+j}\left(\|\boldsymbol{u}\|^{2r} + \|\boldsymbol{u}\|^{2}N^{(r-1)/2}\right)$$
(7.27)

for r odd. We can thus write

$$\mathfrak{G}(t_1, t_2; \zeta_1, \zeta_2) = \mathfrak{G}_0(t_1, t_2) + \sum_{i,j \leq L, (i,j) \neq (0,0)} C_{i,j} \zeta_1^i \zeta_2^j + O(\max(\zeta_1, \zeta_2)^{L+1}),$$

where $|C_{i,j}| = O_{r,i+j} (\|u\|^{2r} + N^{\lfloor r/2 \rfloor}).$ Let

$$F(\zeta_1, \zeta_2) := \frac{\int \langle \boldsymbol{\sigma}^1, \boldsymbol{\sigma}^2 \rangle^r \exp\left(\sum_{i=1}^2 \langle \boldsymbol{u}, \boldsymbol{\sigma}^i \rangle + \langle \boldsymbol{\sigma}^i, \tilde{\boldsymbol{\Lambda}}_i \boldsymbol{\sigma}^i \rangle\right) \mu_0^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma})}{\int \exp\left(\sum_{i=1}^2 \langle \boldsymbol{u}, \boldsymbol{\sigma}^i \rangle + \langle \boldsymbol{\sigma}^i, \boldsymbol{\Lambda} \boldsymbol{\sigma}^i \rangle\right) \mu_0^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma})}.$$
(7.28)

Next, we take the inverse Laplace transform and integrate on $t_i = \gamma_* + ix_i$, for γ_* defined in Eq. (7.11). We note that, for $G(\gamma) = G(\gamma; \mathbf{\Lambda}, \mathbf{u})$ and $\tilde{G}_i(\gamma) = G(\gamma; \mathbf{\Lambda}_i, \mathbf{u})$,

$$\begin{split} \tilde{G}'_i(z) &= G'(z) + \frac{1}{2N} \mathsf{Tr}((z-\mathbf{\Lambda})^{-1}(\mathbf{I} - (z-\mathbf{\Lambda})(z-\tilde{\mathbf{\Lambda}}_i)^{-1})) \\ &+ \frac{1}{4N} \langle \mathbf{u}, (z-\mathbf{\Lambda})^{-1}(\mathbf{I} - (z-\mathbf{\Lambda})(z-\tilde{\mathbf{\Lambda}}_i)^{-2}(z-\mathbf{\Lambda}))(z-\mathbf{\Lambda})^{-1}\mathbf{u} \rangle. \end{split}$$

Moreover, $(z - \Lambda)(z - \tilde{\Lambda})^{-1} = (I - \zeta_i \Delta (z - \Lambda)^{-1})^{-1}$, and $(z - \Lambda)(z - \tilde{\Lambda}_i)^{-2}(z - \Lambda) = (I - \zeta_i \Delta (z - \Lambda)^{-1})^{-1}(I - (z - \Lambda)^{-1}\zeta_i \Delta)^{-1}$. Expanding in $\zeta_i \Delta$, we can show that for $|\zeta_i| \leq C(\log N)/\sqrt{N}$, $|\tilde{G}'_i(\gamma_*)| \leq N^{-1+o(1)}$. Hence, by an argument similar to Lemma 7.3, we can restrict the integration on $t_i = \gamma_* + ix_i$ to the range $x_i \in [-\ell_N, \ell_N]$ for $\ell_N = (\log N)/\sqrt{N}$, and obtain that

$$F(\zeta_1, \zeta_2) = F(0, 0) + \sum_{i,j \leq L, (i,j) \neq (0,0)} C_{i,j} \zeta_1^i \zeta_2^j + O_L(N^{-L/2} + e^{-N^c}).$$

7.1.2 Estimates of restricted partition functions

In this section we estimate modified partition functions that are obtained by suitable restrictions of the integral over σ , always under the assumption (7.9). Namely, for any Borel set $U \subseteq (S_N)^{\otimes m}$,

$$Z_m(U) := \int_U e^{\sum_{i=1}^m H(\boldsymbol{\sigma}^i)} \mu_0^{\otimes m}(\mathsf{d}\boldsymbol{\sigma}), \qquad (7.29)$$

with subscript omitted if m = 1. If $U = S_N$, we write simply $Z = Z(S_N)$. We also denote by $Z_{\leq 2,m}(U)$ the same integral whereby $H(\boldsymbol{\sigma})$ is replaced by $H_{\leq 2}(\boldsymbol{\sigma})$:

$$Z_m(U) := \int_U e^{\sum_{i=1}^m H_{\leq 2}(\boldsymbol{\sigma}^i)} \mu_0^{\otimes m}(\mathsf{d}\boldsymbol{\sigma}), \qquad (7.30)$$

We will occasionally omit the subscript m when the dimension of U is clear from the context.

Throughout this section, we follow the notations $\langle x, y \rangle_N = \langle x, y \rangle/N$, so $\langle x, x \rangle_N = ||x||_N^2$. As for the restrictions, an important role is played by the typical set:

$$T(\delta) = \left\{ \boldsymbol{\sigma} \in S_N : \int_{\boldsymbol{\sigma}': |\langle \boldsymbol{\sigma}', \boldsymbol{\sigma} \rangle_N| > \delta} e^{H(\boldsymbol{\sigma}')} \mu_0(\mathsf{d}\boldsymbol{\sigma}') < e^{-c_1(\delta)N} \min\left(\int e^{H(\boldsymbol{\sigma})} \mu_0(\mathsf{d}\boldsymbol{\sigma}); e^{N\xi(1)/2}\right) \right\}.$$
(7.31)

We further define $A_m(\delta) \subseteq (S_N)^m$ to be the set of *m*-uples of vectors which are nearly orthogonal. Namely:

$$A_m(\delta) := \left\{ (\boldsymbol{\sigma}^i)_{i \leqslant m} : \, \boldsymbol{\sigma}^i \in S_N, |\langle \boldsymbol{\sigma}^i, \boldsymbol{\sigma}^j \rangle_N| \leqslant \delta \,\,\forall i \neq j \right\}.$$
(7.32)

Finally, we consider the set of *m*-uples in $T = T(\delta)$ that are nearly orthogonal:

$$A_m(T,\delta) := \left\{ (\boldsymbol{\sigma}^i)_{i \leqslant m} : \, \boldsymbol{\sigma}^i \in T, |\langle \boldsymbol{\sigma}^i, \boldsymbol{\sigma}^j \rangle_N| \leqslant \delta \,\,\forall i \neq j \right\}.$$
(7.33)

In particular $A_m(T, \delta) = T^m \cap A_m(\delta)$.

Our first lemma establishes that, under the Gibbs measure, non-typical points are exponentially rare.

Lemma 7.7 (Most points are typical). For any $\delta > 0$, there exists $u(\delta), c_1(\delta), c_2(\delta) > 0$ such that the following holds. Let $H(\sigma)$ be defined as per Eq. (7.7) and suppose $\|\mathbf{u}\| \leq u(\delta)\sqrt{N}$. Let $T(\delta)$ be defined as per Eq. (7.31).

Then, with probability at least $1 - \exp(-c_2(\delta)N)$,

$$Z(T(\delta)) \ge (1 - e^{-Nc_2(\delta)}) \cdot Z.$$
(7.34)

Furthermore, there is $c_3(\delta) > 0$ such that, with probability at least $1 - \exp(-c_3(\delta)N)$,

$$Z_{\leq 2}(T(\delta)^c) \leq e^{-c_3(\delta)N} Z_{\leq 2}.$$
(7.35)

Finally

$$\mathbb{E}\left[\int_{T(\delta)^c} e^{H_{\geq 2}(\boldsymbol{\sigma})} \mu_0(\mathsf{d}\boldsymbol{\sigma})\right] \leqslant e^{-c_1(\delta)N} \mathbb{E} Z_{\geq 2}.$$
(7.36)

Proof. The second inequality in (7.9) is termed "strictly RS" in [HS23a], see Eq. (2.7) therein. By Proposition 3.1 of that paper,

$$\mathbb{E}\int_{T(\delta)} e^{H_{\geq 2}(\boldsymbol{\sigma})} \ \mu_0(\mathsf{d}\boldsymbol{\sigma}) \ge (1 - e^{-c_1(\delta)N}) e^{N\xi(1)/2}.$$

(While this proposition states a bound of $(1 - o(1)) \exp(N\xi(1)/2)$, its proof shows the 1 - o(1) is in fact $1 - e^{-c_1(\delta)N}$.) As $\mathbb{E} Z_{\geq 2} = \exp(N\xi(1)/2)$, for $Z_{\geq 2} := \int_{S_N} \exp H_{\geq 2}(\sigma) \mu_0(\mathsf{d}\sigma)$, this implies Eq. (7.36).

By Markov's inequality, with probability $1 - e^{-c_1(\delta)N/5}$,

$$\int_{T(\delta)^c} e^{H_{\geq 2}(\boldsymbol{\sigma})} \mu_0(\mathsf{d}\boldsymbol{\sigma}) \leqslant e^{-4c_1(\delta)N/5} \mathbb{E} Z_{\geq 2}.$$

By [Tal06, Proposition 2.3], (7.9) implies that $\frac{1}{N} \log Z_{\geq 2} \rightarrow_p \xi(1)/2$. By standard concentration properties of $\frac{1}{N} \log Z_{\geq 2}$, with probability $1 - e^{-c_2(\delta)N}$,

$$Z_{\geq 2} \geq e^{-c_1(\delta)N/5} \mathbb{E} Z_{\geq 2}.$$

On the intersection of these events,

$$\int_{T(\delta)^c} e^{H_{\geq 2}(\boldsymbol{\sigma})} \mu_0(\mathrm{d}\boldsymbol{\sigma}) \leqslant e^{-3c_1(\delta)N/5} Z_{\geq 2}.$$

Finally, set $u(\delta) = c_1(\delta)/5$, so that for all $\sigma \in S_N$,

$$|H(\boldsymbol{\sigma}) - H_{\geq 2}(\boldsymbol{\sigma})| = |\langle \boldsymbol{u}, \boldsymbol{\sigma} \rangle| \leq c_1(\delta)N/5.$$

Thus

$$\int_{T(\delta)^c} e^{H(\boldsymbol{\sigma})} \ \mu_0(\mathrm{d}\boldsymbol{\sigma}) \leqslant e^{-c_1(\delta)N/5} \int_{S_N} e^{H(\boldsymbol{\sigma})} \ \mu_0(\mathrm{d}\boldsymbol{\sigma}).$$

The conclusion (7.34) follows with $c(\delta) = \min(c_1(\delta)/6, c_2(\delta)/2)$.

Finally, from Markov's inequality, we have with probability $1 - e^{-c_3(\delta)N}$ that

$$Z_{\leq 2}(T(\delta)^c) \leq e^{-c_3(\delta)N} e^{N\xi_{\leq 2}(1)/2}$$

Then (7.35) follows from standard concentration properties.

The next lemma states that we can anneal over terms of degree higher than 2 in the Hamiltonian. This will be the most important technical result of the section.

Lemma 7.8. Let $H(\sigma)$ be defined as per Eq. (7.7) and define $T = T(\delta)$ as in Eq. (7.31). Assume that $\|u\| \leq N^{c_0}$ for c_0 sufficiently small given ξ . Under assumption (7.9), for all L, k > 0 and $\varepsilon > 0$, there exist C = C(L,k) > 0 such that the following holds with probability at least $1 - \exp(-N/C)$

$$\mathbb{E}_{\geq 3}\left\{\left(Z(T) - \mathbb{E}_{\geq 3}Z(T)\right)^{2k}\right\} \leqslant C N^{-L/2} \left(\mathbb{E}_{\geq 3}Z(T)\right)^{2k},\tag{7.37}$$

and further

$$\mathbb{P}\left\{\left|Z - \mathbb{E}_{\geq 3}Z(T)\right| > \varepsilon \mathbb{E}_{\geq 3}Z(T)\right\} \leq C\varepsilon^{-2L}N^{-L/2} + e^{-N/C}.$$
(7.38)

We also have, with probability at least $1 - \exp(-N/C)$

$$\mathbb{E}_{\geq 3}Z(T) = (1 + O(e^{-N/C}))\mathbb{E}_{\geq 3}Z.$$
(7.39)

Further, letting $(\boldsymbol{v}_k)_{k \leq N}$ be the basis of eigenvectors of \boldsymbol{W}_2 , for each $i \in [N]$,

$$\mathbb{P}\left(\int_{T} \langle \boldsymbol{v}_{i}, \boldsymbol{\sigma} \rangle e^{H(\boldsymbol{\sigma})} \mu_{0}(\mathsf{d}\boldsymbol{\sigma}) \geqslant N^{\varepsilon} \|\boldsymbol{u}\|^{Ck} (|\langle \boldsymbol{v}_{i}, \boldsymbol{u} \rangle| + C N^{-1/2}) \mathbb{E}_{\geqslant 3} \int e^{H(\boldsymbol{\sigma})} \mu_{0}(\mathsf{d}\boldsymbol{\sigma}) \right) \leqslant C \left(N^{-2\varepsilon k} + e^{-N/C}\right)$$
(7.40)

Before proving this lemma, we state and prove a number of key estimates.

Our first lemma establishes that (in expectation) the partition function in $A_{2k}(\delta)$ is dominated by the subset $A_{2k}(\delta, T)$.

Lemma 7.9 (Orthogonal frames are mostly typical). Define $T = T(\delta)$ as in Eq. (7.31). We have for $\delta > 0$ sufficiently small and appropriate c, c' > 0 that, if $\|u\| \leq c'\sqrt{N}$,

$$\mathbb{E} Z_{2k} \left(\left\{ (\boldsymbol{\sigma}^i)_{i \leq 2k} \in A_{2k}(\delta) : \boldsymbol{\sigma}^1 \in T^c \right\} \right) \leq e^{-cN} \mathbb{E} Z_{2k} \left(A_{2k}(\delta) \right).$$
(7.41)

As a consequence,

$$\mathbb{E} Z_{2k} \left(A_{2k}(\delta, T) \right) \ge (1 - e^{-cN}) \mathbb{E} Z_{2k} \left(A_{2k}(\delta) \right).$$
(7.42)

Proof. We have

$$\mathbb{E}\left\{H_{\geq 2}(\boldsymbol{\rho})\Big|\sum_{i=1}^{2k}H_{\geq 2}(\boldsymbol{\sigma}^{i})\right\} = \frac{\mathbb{E}H_{\geq 2}(\boldsymbol{\rho})\sum_{i=1}^{2k}H_{\geq 2}(\boldsymbol{\sigma}^{i})}{\mathbb{E}(\sum_{i=1}^{2k}H_{\geq 2}(\boldsymbol{\sigma}^{i}))^{2}}\sum_{i=1}^{2k}H_{\geq 2}(\boldsymbol{\sigma}^{i})$$
$$= \frac{\sum_{i=1}^{2k}\xi(\langle \boldsymbol{\rho}, \boldsymbol{\sigma}^{i}\rangle_{N})}{\sum_{i,j\in[2k]}\xi(\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j}\rangle_{N})}\sum_{i=1}^{2k}H_{\geq 2}(\boldsymbol{\sigma}^{i}),$$

and for $\hat{H}(\boldsymbol{\rho}) = H_{\geq 2}(\boldsymbol{\rho}) - \mathbb{E}[H_{\geq 2}(\boldsymbol{\rho})|\sum_{i=1}^{2k} H_{\geq 2}(\boldsymbol{\sigma}^{i})],$ $\mathbb{E}[\hat{H}(\boldsymbol{\rho}^{1})\hat{H}(\boldsymbol{\rho}^{2})] = \xi(\langle \boldsymbol{\rho}^{1}, \boldsymbol{\rho}^{2} \rangle_{N}) - \frac{(\sum_{i=1}^{2k} \xi(\langle \boldsymbol{\rho}^{1}, \boldsymbol{\sigma}^{i} \rangle_{N})(\sum_{i=1}^{2k} \xi(\langle \boldsymbol{\rho}^{2}, \boldsymbol{\sigma}^{i} \rangle_{N}))}{\sum_{i,j \in [2k]} \xi(\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j} \rangle_{N})}.$

For each $|q_1| \ge \delta$, and $q_2, \ldots, q_{2k} \in [0, 1]$, consider the band $\text{Band}_*(\{\sigma^i\})$ of vectors ρ with $\langle \rho, \sigma^i \rangle = q_i$ for all $i \in [2k]$. Write $\rho = x + \sqrt{1 - \tilde{q}^2 \tau}$ where $x \in \text{span}(\sigma^1, \ldots, \sigma^{2k})$ and $||\tau||^2 = N, \tau \perp \text{span}(\sigma^1, \ldots, \sigma^{2k})$. Define the process $\overline{H}(\tau) = \hat{H}(\rho)$, which is a *p*-spin model with corresponding mixture $\tilde{\xi}(t) = \tilde{\xi}(t; q, (\sigma^i)_{i=1}^{2k})$ given by

$$\widetilde{\xi}(t;\boldsymbol{q},(\boldsymbol{\sigma}^{i})_{i=1}^{2k}) = \xi(\widetilde{q}^{2} + (1 - \widetilde{q}^{2})t) - \frac{\left(\sum_{i=1}^{2k} \xi(q_{i})\right)^{2}}{\sum_{i,j \in [2k]} \xi(\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j} \rangle_{N})}$$

We define the free energy

$$\Phi(\boldsymbol{q}; (\boldsymbol{\sigma}^i)_{i=1}^{2k}) := \frac{1}{N} \log \int_{\mathsf{Band}_*(\{\boldsymbol{\sigma}^i\})} e^{H_{\geqslant 2}(\boldsymbol{\rho})} \mu_0(\mathsf{d}\boldsymbol{\rho})$$

Following the proof of Lemma 3.3 of [HS23a], the replica-symmetric bound implies that the following holds with high probability:

$$\Phi(\boldsymbol{q};(\boldsymbol{\sigma}^{i})_{i=1}^{2k}) \leq \frac{\sum_{i=1}^{2k} \xi(q_{i})}{\sum_{i,j \in [2k]} \xi(\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j} \rangle_{N})} \sum_{i=1}^{2k} H_{\geq 2}(\boldsymbol{\sigma}^{i}) + \frac{1}{2}\xi(1) - \frac{1}{2}\xi(\tilde{q}) + \frac{1}{2}\tilde{q} + \frac{1}{2}\log(1-\tilde{q}) + o_{N}(1).$$
(7.43)

By the generalized Bessel inequality, we have

$$\sum_{i=1}^{2k} \langle \boldsymbol{x}, \boldsymbol{\sigma}^i \rangle_N^2 \leqslant \| \boldsymbol{x} \|_N^2 (2k)^{-1} \sum_{i,j \in [2k]} \langle \boldsymbol{\sigma}^i, \boldsymbol{\sigma}^j \rangle_N^2 = \| \boldsymbol{x} \|_N^2 (2k)^{-1} (2k + (2k)^2 \delta^2).$$

Hence,

$$\tilde{q}^2 = \|\boldsymbol{x}\|_N^2 \ge \frac{1}{1+2k\delta^2} \sum_{i=1}^{2k} q_i^2,$$

and since $\xi(0) = \xi'(0) = 0$, this implies

$$\sum_{i=1}^{2k} \xi(q_i) \le \xi \left((1 + 2k\delta^2)^{1/2} \tilde{q} \right).$$

We pick δ sufficiently small in c and k, and η small in δ . Given $\sum_{i=1}^{2k} H_{\geq 2}(\sigma^i) = EN$ where $E \leq \sum_{i,j\in[2k]} \xi(\langle \sigma^i, \sigma^j \rangle_N) + \eta$, whenever $q_1 \geq \delta$, we have by assumption (7.9), with high probability

$$\Phi(\boldsymbol{q}; (\boldsymbol{\sigma}^i)_{i=1}^{2k}) \leqslant \frac{1}{2}\xi(1) - 10\eta.$$

Integrating over the $(q_i)_{i \leq 2k}$ and using Gaussian concentration, we deduce that for $E \leq \sum_{i,j \in [2k]} \xi(\langle \sigma^i, \sigma^j \rangle_N) + \eta$, we have

$$\mathbb{P}\left\{ \left. \int_{\boldsymbol{\rho}:\langle \boldsymbol{\rho},\boldsymbol{\sigma}_1 \rangle_N > \delta} e^{H_{\geqslant 2}(\boldsymbol{\rho})} \mu_0(\mathsf{d}\boldsymbol{\rho}) \leqslant e^{N(\xi(1)/2 - 9\eta)} \right| \sum_{i=1}^{2k} H_{\geqslant 2}(\boldsymbol{\sigma}^i) = EN \right\} \geqslant 1 - e^{-c(\eta)N}.$$

Up until now we worked with the Hamiltonian $H_{\geq 2}(\sigma)$, which does not include the term linear in σ . Recall that $H(\sigma) = \langle u, \sigma \rangle + H_{\geq 2}(\sigma)$ and $||u|| \leq c'\sqrt{N}$ so $|H(\sigma) - H_{\geq 2}(\sigma)| \leq |\langle u, \sigma \rangle| \leq c'N$, assuming that $c' < \eta$, we have

$$\mathbb{P}\left\{\left.\int_{\boldsymbol{\rho}:\langle\boldsymbol{\rho},\boldsymbol{\sigma}_1\rangle_N>\delta}e^{H(\boldsymbol{\rho})}\mu_0(\mathsf{d}\boldsymbol{\rho})\leqslant e^{N(\xi(1)/2-8\eta)}\right|\sum_{i=1}^{2k}H_{\geqslant 2}(\boldsymbol{\sigma}^i)=EN\right\}\geqslant 1-e^{-c(\eta)N}.$$

Hence, under the same conditions

$$\mathbb{P}\left\{\boldsymbol{\sigma}^{1} \in T^{c} \left| \sum_{i=1}^{2k} H_{\geq 2}(\boldsymbol{\sigma}^{i}) = EN \right. \right\} \leq e^{-c(\eta)N}$$

Define the event

$$\mathcal{E}(\{\boldsymbol{\sigma}^i\}) := \left\{ \sum_{i=1}^{2k} H_{\geq 2}(\boldsymbol{\sigma}^i) \geq N\left(\sum_{i,j\in[2k]} \xi(\langle \boldsymbol{\sigma}^i, \boldsymbol{\sigma}^j \rangle_N) + \eta\right) \right\}.$$

Thus, since $|H(\boldsymbol{\sigma}) - H_{\geq 2}(\boldsymbol{\sigma})| \leq c'N$, we can then conclude that

$$\begin{split} & \mathbb{E}\left\{\int_{A_{2k}(\delta):\boldsymbol{\sigma}^{1}\in T^{c}}e^{\sum_{i=1}^{2k}H(\boldsymbol{\sigma}^{i})}\mu_{0}^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma})\right\} = \mathbb{E}\left\{\int_{A_{2k}(\delta)}\mathbf{1}_{\boldsymbol{\sigma}^{1}\in T^{c}}e^{\sum_{i=1}^{2k}H(\boldsymbol{\sigma}^{i})}\mu_{0}^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma})\right\} \\ &= \mathbb{E}\left\{\int_{A_{2k}(\delta)}\mathbb{P}\left\{\boldsymbol{\sigma}^{1}\in T^{c}\Big|\sum_{i=1}^{2k}H_{\geqslant 2}(\boldsymbol{\sigma}^{i})\right\}e^{\sum_{i=1}^{2k}H(\boldsymbol{\sigma}^{i})}\mu_{0}^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma})\right\} \\ &\leq e^{-c(\eta)N+c'N}\mathbb{E}\int_{A_{2k}(\delta)}e^{\sum_{i=1}^{2k}H_{\geqslant 2}(\boldsymbol{\sigma}^{i})}\mu_{0}^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma}) + e^{c'N}\mathbb{E}\int_{A_{2k}(\delta)}e^{\sum_{i=1}^{2k}H_{\geqslant 2}(\boldsymbol{\sigma}^{i})}\mathbf{1}_{\mathcal{E}(\{\boldsymbol{\sigma}^{i}\})}\mu_{0}^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma}) \\ &\leq e^{-cN}\mathbb{E}\int_{A_{2k}(\delta)}e^{\sum_{i=1}^{2k}H(\boldsymbol{\sigma}^{i})}\mu_{0}^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma}). \end{split}$$

Here we assume $c' < c(\eta)/4$ and $c = c(\eta)/4$, and in the last step we used, for $U(\{\sigma^i\}) := \sum_{i,j \in [2k]} \xi(\langle \sigma^i, \sigma^j \rangle_N)$,

$$\mathbb{E}\int_{A_{2k}(\delta)} e^{\sum_{i=1}^{2k} H_{\geqslant 2}(\boldsymbol{\sigma}^{i})} \mathbf{1}_{\mathcal{E}(\{\boldsymbol{\sigma}^{i}\})} \mu_{0}^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma}) \leqslant \int_{A_{2k}(\delta)} \exp\left\{N(1-s+s^{2})U(\{\boldsymbol{\sigma}^{i}\})-Ns\eta\right\} \mu_{0}^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma}),$$

and chose δ , s suitably small.

The next lemma shows that integrals of S_N^{2k} with the product Gibbs measure are very precisely approximated by integral over tuples that are very close to orthogonal.

Lemma 7.10 (Near-orthogonal tuples dominate). For $\delta > 0$ sufficiently small and appropriate $c, c_0 > 0$, if $\|\mathbf{u}\| \leq N^{c_0/2}$, then with probability $1 - e^{-cN}$ over $\mathbf{W}^{(2)}$, the following holds:

1. For quadratic Hamiltonians, the unrestricted partition function of 2k replicas is dominated by its restriction to $A_{2k}(N^{-1/2+c})$:

$$Z_{\leq 2,2k} \left(A_{2k} (N^{-1/2+c}) \right) \ge (1 - e^{-N^c}) \cdot \left(Z_{\leq 2} \right)^{2k}.$$
(7.44)

2. The contribution of $A_{2k}(\delta) \setminus A_{2k}(N^{-1/2+c}) = \{(\boldsymbol{\sigma}^i)_{i \leq 2k} : \max_{i \neq j} |\langle \boldsymbol{\sigma}^i, \boldsymbol{\sigma}^j \rangle_N| \in [N^{-1/2+c}, \delta]\}$ is small:

$$\mathbb{E}_{\geq 3} Z_{2k} \big(A_{2k}(\delta) \setminus A_{2k}(N^{-1/2+c}) \big) \leq e^{-N^c + Nk\xi_{\geq 3}(1)} \big(Z_{\leq 2} \big)^{2k}.$$
(7.45)

3. Annealing the restricted partition function over $H_{\geq 3}$ is roughly equivalent to complete annealing:

$$\mathbb{E}_{\geq 3} Z_{2k} \left(A_{2k}(\delta) \right) \geq e^{-4k(\|\boldsymbol{u}\|+1)\sqrt{N}} \mathbb{E} Z_{2k} \left(A_{2k}(\delta) \right).$$
(7.46)

Proof. Proof of 1. By Lemma 7.5, for some constants $c_1, C_1 > 0$ that, with probability $1 - e^{-cN}$ over $W^{(2)}$,

$$Z_{\leq 2,2k} \left(A_{2k} (N^{-1/2+c}) \right) \leq \sum_{i \neq j} \int_{S_N^{2k}} \mathbf{1}_{|\langle \boldsymbol{\sigma}^i, \boldsymbol{\sigma}^j \rangle_N| > N^{-1/2+c}} e^{\sum_{i=1}^{2k} H_{\leq 2}(\boldsymbol{\sigma}^i)} \mu_0^{\otimes 2k} (\mathsf{d}\boldsymbol{\sigma})$$
$$\leq e^{-N^c} \int_{S_N^{2k}} e^{\sum_{i=1}^{2k} H_{\leq 2}(\boldsymbol{\sigma}^i)} \mu_0^{\otimes 2k} (\mathsf{d}\boldsymbol{\sigma}),$$

yielding (7.44).

Proof of 2. By a direct calculation, for any set $U \subseteq (S_N)^{2k}$:

$$\mathbb{E}_{\geq 3} Z_{2k}(U) = e^{Nk\xi_{\geq 3}(1)} \int_{U} e^{\sum_{i=1}^{2k} H_{\leq 2}(\boldsymbol{\sigma}^{i})} \exp\left(N \sum_{i < j < 2k} \xi_{\geq 3}(\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j} \rangle_{N})\right) \mu_{0}^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma}).$$

Applying Lemma 7.5, we have for t > 0 and $\varepsilon_N = N^{-1/2+c}$ that, with probability $1 - e^{-cN}$ over $W^{(2)}$,

$$\frac{1}{(Z_{\leq 2})^{2k}}e^{-Nk\xi_{\geq 3}(1)}\mathbb{E}_{\geq 3}Z_{2k}\left(A_{2k}(t+\varepsilon_N)\backslash A_{2k}(t)\right) \leq$$
(7.47)

$$\frac{1}{(Z_{\leq 2})^{2k}} \int_{\max_{i \neq j} |\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j} \rangle_{N}| \in [t, t+\varepsilon_{N}]} \exp\left(\sum_{i=1}^{2k} H_{\leq 2}(\boldsymbol{\sigma}^{i}) + N \sum_{i \neq j \in [2k]} \xi_{\geq 3}(\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j} \rangle_{N})\right) \mu_{0}^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma}) \leq \\ \leq \exp\left\{-cN\left(t - \|\boldsymbol{u}\|_{N}^{2}\right)_{+}^{2} + N(2k)^{2}\xi_{\geq 3}(t+\varepsilon_{N})\right\}.$$
(7.48)

Under the assumption $\|\boldsymbol{u}\|_N^2 \leq N^{c_0-1}$, $c_0 < c + 1/2$, summing over the range $N^{-1/2+c} < |t| \leq \delta$, we obtain the following with probability $1 - e^{-cN}$ over $\boldsymbol{W}^{(2)}$,

$$\begin{split} &\int_{\max_{i\neq j}|\langle\boldsymbol{\sigma}^{i},\boldsymbol{\sigma}^{j}\rangle_{N}|\in[N^{-1/2+c},\delta]} \exp\left(\sum_{i=1}^{2k} H_{\leqslant 2}(\boldsymbol{\sigma}^{i}) + N \sum_{i\neq j\in[2k]} \xi_{\geqslant 3}(\langle\boldsymbol{\sigma}^{i},\boldsymbol{\sigma}^{j}\rangle_{N})\right) \mu_{0}^{\otimes 2k}(\mathrm{d}\boldsymbol{\sigma}) \\ &\leqslant \exp(-N^{c}) \int_{S_{N}^{2k}} \exp\left(\sum_{i=1}^{2k} H_{\leqslant 2}(\boldsymbol{\sigma}^{i})\right) \mu_{0}^{\otimes 2k}(\mathrm{d}\boldsymbol{\sigma}). \end{split}$$

This gives (7.45). **Proof of 3.** Note that

$$e^{-Nk\xi_{\geq 3}(1)} \mathbb{E}_{\geq 3} Z_{2k} \left(A_{2k}(N^{-1/2+c}) \right) = \\ = \int_{A_{2k}(N^{-1/2+c})} \exp\left(\sum_{i=1}^{2k} H_{\leq 2}(\boldsymbol{\sigma}^{i}) + N \sum_{i \neq j \in [2k]} \xi_{\geq 3}(\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j} \rangle_{N}) \right) \mu_{0}^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma}) \\ = (1 + O(N^{-1/2+3c})) \cdot Z_{\leq 2,2k} \left(A_{2k}(N^{-1/2+c}) \right).$$
(7.49)

Therefore, using Eq. (7.44), we get

$$\mathbb{E}_{\geq 3} Z_{2k} \left(A_{2k} (N^{-1/2+c}) \right) = (1 + O(N^{-1/2+3c})) e^{Nk\xi_{\geq 3}(1)} \left(Z_{\leq 2} \right)^{2k}.$$
(7.50)

Also,

$$\mathbb{E}Z_{\leq 2,2k}\Big(A_{2k}(N^{-1/2+c})\Big) \leq e^{2k\|\boldsymbol{u}\|\sqrt{N}}\exp\left(k\beta_2^2N + (2k)^2\beta_2^2N^{2c}\right).$$
(7.51)

On the other hand, Lemma 7.3 readily implies that with probability at least $1 - e^{-cN}$,

$$(Z_{\leq 2})^{2k} \ge e^{-o(\sqrt{N})} \exp(k\beta_2^2 N).$$
 (7.52)

Combining Eqs. (7.51) and (7.51), we get

$$\mathbb{E}Z_{\leq 2,2k}\left(A_{2k}(N^{-1/2+c})\right) \leq e^{2k(1+\|\boldsymbol{u}\|)\sqrt{N}}(Z_{\leq 2})^{2k}.$$
(7.53)

Finally, using Eq. (7.50) together with the last display, we get

$$\mathbb{E}_{\geq 3} Z_{2k} \left(A_{2k}(N^{-1/2+c}) \right) \geq e^{-3k(\|\boldsymbol{u}\|+1)\sqrt{N}} \mathbb{E} Z_{2k} \left(A_{2k}(N^{-1/2+c}) \right).$$
(7.54)

Combining this with Eq. (7.45) yields the claim.

Lemma 7.11. For any $m \ge 2$, there exists a constant c > 0 such that, for $T = T(\delta)$,

$$Z_m(A_m(T,\delta)) \le Z(T)^m \le (1+e^{-cN}) \cdot Z_m(A_m(\delta)) + e^{-cN + Nm\xi(1)/2}.$$
(7.55)

Proof. The left hand inequality is obvious since $A_m(T, \delta) \subseteq T^{\otimes m}$. For the right inequality consider first the case m = 2. Then we have

$$Z(T)^{2} \leq Z_{2}(A_{2}(T,\delta)) + \int_{T \times T} \mathbf{1}_{|\langle \boldsymbol{\sigma}^{1}, \boldsymbol{\sigma}^{2} \rangle_{N}| \geq \delta} e^{H(\boldsymbol{\sigma}^{1}) + H(\boldsymbol{\sigma}^{2})} \mu_{0}^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma})$$

$$(7.56)$$

$$\leq Z_2(A_2(T,\delta)) + \int_T e^{H(\boldsymbol{\sigma}^1)} \left[\int_T \mathbf{1}_{|\langle \boldsymbol{\sigma}^1, \boldsymbol{\sigma}^2 \rangle_N| \ge \delta} e^{H(\boldsymbol{\sigma}^2)} \mu_0(\mathsf{d}\boldsymbol{\sigma}^2) \right] \mu_0(\mathsf{d}\boldsymbol{\sigma}^1) \tag{7.57}$$

$$\leq Z_2(A_2(T,\delta)) + \int_T e^{H(\boldsymbol{\sigma}^1)} e^{-\delta N + N\xi(1)/2} \mu_0(\mathsf{d}\boldsymbol{\sigma}^1)$$
(7.58)

$$\leq Z_2(A_2(T,\delta)) + e^{-N\delta + N\xi(1)/2}Z(T)$$
(7.59)

$$\leq Z_2(A_2(T,\delta)) + e^{-N\delta + N\xi(1)} + e^{-N\delta}Z(T)^2.$$
 (7.60)

where in the last step we used the AM-GM inequality. Solving this inequality for $Z(T)^2$, we get:

$$Z(T)^{2} \leq (1 + e^{-cN}) Z_{2} (A_{2}(T, \delta)) + 2e^{-\delta N + N\xi(1)}.$$
(7.61)

which proves the claim for m = 2.

Consider now $m \ge 2$. Note that

$$\begin{split} &\int_{T(\delta)^m} e^{\sum_{i=1}^m H(\boldsymbol{\sigma}^i)} \mu_0^{\otimes m}(\mathrm{d}\boldsymbol{\sigma}) - \int_{A_m(T(\delta),\delta)} e^{\sum_{i=1}^m H(\boldsymbol{\sigma}^i)} \mu_0^{\otimes 2m}(\mathrm{d}\boldsymbol{\sigma}) \\ &\leqslant \sum_{i\neq j} \left(\int_{T(\delta)} e^{H(\boldsymbol{\sigma})} \mu_0(\mathrm{d}\boldsymbol{\sigma}) \right)^{m-2} \int_{\boldsymbol{\sigma}^i, \boldsymbol{\sigma}^j \in T(\delta) : |\langle \boldsymbol{\sigma}^i, \boldsymbol{\sigma}^j \rangle_N| > \delta} e^{H(\boldsymbol{\sigma}^i) + H(\boldsymbol{\sigma}^j)} \mu_0(\mathrm{d}\boldsymbol{\sigma}^i) \mu_0(\mathrm{d}\boldsymbol{\sigma}^j) \end{split}$$

whence

$$Z(T)^m - Z_m (A_m(T,\delta)) \leq m^2 Z_2(T^{\otimes 2} \setminus A_2(T,\delta)) \cdot Z(T)^{m-2}$$
$$\leq m^2 \cdot Z(T)^{m-1} \cdot e^{-N\delta + N\xi(1)/2},$$

where in the last inequality we used Eq. (7.59). Using again the AM-GM inequality, we get

$$Z(T)^{m} - Z_{m}(A_{m}(T,\delta)) \leq m^{2} e^{-N\delta} Z(T)^{m} + m^{2} e^{-N\delta + Nm\xi(1)/2},$$

which yields the claim.

7.1.3 Proof of Lemma 7.8

We next prove Lemma 7.8. In the proof, we let c denote small absolute constants that can change from line to line. We will first prove the partition function estimate, Eq. (7.38) and then the magnetization estimate, Eq. (7.40).

Estimating the partition function, Eq. (7.38). By Eq. (7.45) in Lemma 7.10, with probability $1 - e^{-cN}$ over $W^{(2)}$,

$$\int_{\max_{i\neq j} |\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j} \rangle_{N}| \in [N^{-1/2+c}, \delta]} \exp\left(\sum_{i=1}^{2k} H_{\leq 2}(\boldsymbol{\sigma}^{i}) + N \sum_{i < j \leq 2k} \xi_{\geq 3}(\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j} \rangle_{N})\right) \mu_{0}^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma})
\leq \exp(-N^{c}) \int_{S_{N}^{2k}} \exp\left(\sum_{i=1}^{2k} H_{\leq 2}(\boldsymbol{\sigma}^{i})\right) \mu_{0}^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma}).$$
(7.62)

On $A_{2k}(N^{-1/2+c})) = \{ |\langle \boldsymbol{\sigma}^i, \boldsymbol{\sigma}^j \rangle_N | \leqslant N^{-1/2+c} \} \ \forall i \neq j \}$, we can expand

$$\exp\left(N\sum_{i< j}\xi_{\geq 3}(\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j}\rangle_{N})\right) = \sum_{\ell=0}^{L-1}\frac{1}{\ell!}\left(N\sum_{i< j}\xi_{\geq 3}(\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j}\rangle_{N})\right)^{\ell} + O(N^{-L/2+3cL}).$$

Thus, for $T = T(\delta)$, the following holds with probability at least $1 - e^{-cN}$ over W_2 ,

$$\mathbb{E}_{\geq 3} \left\{ \left(Z(T) - \mathbb{E}_{\geq 3} Z(T) \right)^{2k} \right\}$$
(7.63)
$$\stackrel{(a)}{\leq} \sum_{r \leq 2k} \binom{2k}{2k - r} (-1)^r \left(\mathbb{E}_{\geq 3} Z(T) \right)^{2k - r} \cdot \mathbb{E}_{\geq 3} Z \left(A_r(T, \delta) \right) \\
+ e^{-Nc} \sum_{r \leq 2k} \binom{2k}{2k - r} \left(\mathbb{E}_{\geq 3} Z(T) \right)^{2k - r} \cdot \left(\mathbb{E}_{\geq 3} Z \left(A_r(T, \delta) \right) + e^{Nr\xi(1)/2} \right) \\
\stackrel{(b)}{\leq} \sum_{r \leq 2k} \binom{2k}{2k - r} (-1)^r \left(\mathbb{E}_{\geq 3} Z(T) \right)^{2k - r} \cdot \mathbb{E}_{\geq 3} Z \left(A_r(T, \delta) \right) \\
+ e^{-Nc} \max_{r \leq 2k} e^{N(2k - r)\xi(1)/2} \cdot \left(\mathbb{E}_{\geq 3} Z \left(A_r(T, \delta) \right) + e^{Nr\xi(1)/2} \right),$$
(7.64)

where (a) follows from Lemma 7.10, (b) holds because $\mathbb{E}_{\geq 3}Z(T) \leq e^{c'N}\mathbb{E}Z$ with the claimed probability by Markov inequality.

We define the error terms

$$\mathsf{Err}_{1} := e^{-cN + Nk\xi(1)} + e^{-cN} \max_{1 \le r \le 2k} \left(\mathbb{E}_{\ge 3} Z(A_{r}(T, \delta)) \right)^{2k/r} + \mathbb{E}_{\ge 3} Z(A_{2k}(\delta) \cap \{ \sigma^{1} \in T^{c} \}), \quad (7.66)$$

$$\operatorname{Err}_{2} := N^{-L/2} e^{Nk\xi_{\geq 3}(1)} Z_{\leq 2,2k}(A_{2k}(\delta)), \qquad (7.67)$$

so that the bound (7.65) implies

$$\mathbb{E}_{\geq 3}\left\{\left(Z(T) - \mathbb{E}_{\geq 3}Z(T)\right)^{2k}\right\} \leq \sum_{r \leq 2k} \binom{2k}{2k-r} (-1)^r \left(\mathbb{E}_{\geq 3}Z(T)\right)^{2k-r} \cdot \mathbb{E}_{\geq 3}Z\left(A_r(T,\delta)\right) + O_k(\mathsf{Err}_1).$$
(7.68)

Next note that

$$\begin{split} \left(\mathbb{E}_{\geqslant 3}Z(T)\right)^{2k-r} \cdot \mathbb{E}_{\geqslant 3}Z\left(A_{r}(T,\delta)\right) \\ &= e^{Nk\xi_{\geqslant 3}(1)} \left(\int_{T(\delta)} e^{H_{\leqslant 2}(\sigma)}\mu_{0}(\mathrm{d}\boldsymbol{\sigma})\right)^{2k-r} \cdot \\ &\quad \cdot \int_{A_{r}(\delta)} e^{\sum_{i=1}^{r} H_{\leqslant 2}(\sigma^{i})} \exp\left(N\sum_{i < j}\xi_{\geqslant 3}(\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j} \rangle_{N})\right) \mu_{0}^{\otimes r}(\mathrm{d}\boldsymbol{\sigma}) + O_{k}(\mathrm{Err}_{1}) \\ &= e^{Nk\xi_{\geqslant 3}(1)} \int_{A_{2k}(\delta)} \exp\left(\sum_{i'=1}^{2k-r} H_{\leqslant 2}((\boldsymbol{\sigma}')^{i'}) + \sum_{i=1}^{r} H_{\leqslant 2}(\boldsymbol{\sigma}^{i})\right) \cdot \\ &\quad \cdot \left\{\sum_{\ell=0}^{L-1} \frac{1}{\ell!} \left(N\sum_{i < j \leqslant r} \xi_{\geqslant 3}(\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j} \rangle_{N})\right)^{\ell}\right\} \mu_{0}^{\otimes r}(\mathrm{d}\boldsymbol{\sigma}) \mu_{0}^{\otimes(2k-r)}(\mathrm{d}\boldsymbol{\sigma}') + O_{k}(\mathrm{Err}_{1} + \mathrm{Err}_{2}), \end{split}$$

where the last inequality holds with probability $1 - e^{-cN}$ over $W^{(2)}$ by Eq. (7.45).

Substituting in Eq. (7.68), we get

$$\mathbb{E}_{\geq 3}\left\{\left(Z(T) - \mathbb{E}_{\geq 3}Z(T)\right)^{2k}\right\} \\
\leqslant e^{Nk\xi_{\geq 3}(1)} \int_{A_{2k}(\delta)} e^{\sum_{i=1}^{2k} H_{\leq 2}(\sigma^{i})} \sum_{\ell \leqslant L} \frac{1}{\ell!} \sum_{r \leqslant 2k} (-1)^{r} \sum_{S \subseteq [2k]:|S|=r} \left(N \sum_{i < j \in S} \xi_{\geq 3}(\langle \sigma^{i}, \sigma^{j} \rangle_{N})\right)^{\ell} \mu_{0}^{\otimes 2k}(\mathsf{d}\sigma) \\
+ O_{k}(\mathsf{Err}_{1} + \mathsf{Err}_{2}),$$
(7.69)

We can expand the ℓ -th power in (7.69), thus getting a sum indexed by sets of pairs $S = \{(i_t, j_t) : t \leq \ell\} \subseteq {\binom{[2k]}{2}}$. Denoting by n(S) the number of distinct elements of [2k] appearing in S, the coefficient of such therm is its coefficient is, for n(S) < 2k,

$$\sum_{\ell \leqslant r \leqslant 2k} (-1)^r \binom{2k - n(S)}{r - n(S)} = 0$$

for $|\{i_t, j_t : t \leq \ell\}| < 2k$. Hence, taking L < k, we have

$$\mathbb{E}_{\geq 3}\left\{\left(Z(T) - \mathbb{E}_{\geq 3}Z(T)\right)^{2k}\right\} = O_k(\mathsf{Err}_1 + \mathsf{Err}_2).$$
(7.70)

We now estimate the error terms.

Error term Err₂. Using Lemma 7.7, we have

$$\left(\mathbb{E}_{\geq 3}Z(T)\right)^{2k} = \left(\mathbb{E}_{\geq 3}Z - \mathbb{E}_{\geq 3}Z(T^{c})\right)^{2k}$$
$$\geq (1 - e^{-cN/8}) \left(\mathbb{E}_{\geq 3}Z\right)^{2k}$$
$$\geq c e^{Nk\xi_{\geq 3}(1)} \left(Z_{\leq 2}\right)^{2k}$$
$$\geq c e^{Nk\xi_{\geq 3}(1)} Z_{\leq 2} \left(A_{2k}(\delta)\right).$$

From this estimate, we obtain with probability at least $1 - \exp(-cN/8)$ over $W^{(2)}$ that

$$\mathsf{Err}_2 \leqslant C \cdot N^{-L/2} \cdot \left(\mathbb{E}_{\geqslant 3} Z(T)\right)^{2k}.$$
(7.71)

Error term Err₁. Using Lemma 7.9 by Markov inequality, with probability $1 - \exp(-cN/2)$ over $W^{(2)}$,

$$\mathbb{E}_{\geq 3}Z(A_{2k}(\delta) \cap \{\boldsymbol{\sigma}^1 \in T^c\}) \leq e^{-cN/2} \mathbb{E}Z(A_{2k}(\delta)).$$
(7.72)

Further using Eq. (7.46) in Lemma 7.10, and using the assumption on $\|\boldsymbol{u}\|_2$, with probability $1 - \exp(-cN/4)$ over $\boldsymbol{W}^{(2)}$,

$$\mathbb{E}_{\geq 3}Z(A_{2k}(\delta) \cap \{\boldsymbol{\sigma}^1 \in T^c\}) \leq e^{-cN/2} \mathbb{E}_{\geq 3}Z(A_{2k}(\delta)).$$
(7.73)

Hence, with probability at least $1 - \exp(-cN/8)$ over $\boldsymbol{W}^{(2)}$,

$$\mathsf{Err}_{1} \leqslant e^{-cN + Nk\xi(1)} + e^{-cN} \max_{1 \leqslant r \leqslant 2k} \left(\mathbb{E}_{\geqslant 3} Z(A_{r}(\delta)) \right)^{2k/r}$$
(7.74)

Further, with probability at least $1 - \exp(-cN/8)$ over $oldsymbol{W}^{(2)}$,

$$\mathbb{E}_{\geq 3}Z(A_r(\delta)) = \mathbb{E}_{\geq 3}Z(A_r(N^{-1/2+c})) + \mathbb{E}_{\geq 3}Z(A_r(\delta) \setminus A_r(N^{-1/2+c}))$$

$$\leq 2e^{Nr\xi_{\geq 3}(1)/2} Z_{\leq 2}(A_r(N^{-1/2+c})) + e^{-N^c + Nr\xi_{\geq 3}(1)/2} (Z_{\leq 2})^r, \qquad (7.75)$$

where in the last line we used Eq. (7.45), and the fact that

$$\begin{split} \mathbb{E}_{\geq 3} Z \big(A_r(N^{-1/2+c}) \big) &= \int_{A_r(N^{-1/2+c})} e^{\sum_{i=1}^r H_{\leq 2}(\boldsymbol{\sigma}^i)} \exp\left(\frac{N}{2} \sum_{i,j \leq r} \xi_{\geq 3}(\langle \boldsymbol{\sigma}^i, \boldsymbol{\sigma}^j \rangle_N) \right) \mu_0^{\otimes r}(\mathsf{d}\boldsymbol{\sigma}) \\ &\leq \left(1 + O(N^{-1/2+3c})\right) e^{Nk\xi_{\geq 3}(1)} Z_{\leq 2} \big(A_r(N^{-1/2+c}) \big) \,. \end{split}$$

Using Eq. (7.14) in Eq. (7.75), we get

$$\mathbb{E}_{\geq 3}Z(A_r(\delta)) \leqslant N^C e^{Nr\xi(1)/2}, \qquad (7.76)$$

whence Eq. (7.74) simplifies to

$$\operatorname{Err}_{1} \leqslant e^{-cN + Nr\xi(1)} \,. \tag{7.77}$$

On the other hand, by Lemma 7.7 and Markov inequality, with probability $1 - \exp(-cN/4)$ over $W^{(2)}$,

$$\mathbb{E}_{\geq 3} \int_{T(\delta)^c} e^{H(\boldsymbol{\sigma})} \mu_0(\mathsf{d}\boldsymbol{\sigma}) \leqslant e^{-cN/4} e^{N\xi(1)/2}.$$

Using Lemma 7.3, we obtain that, with probability at least $1 - \exp(-cN/8)$ over $W^{(2)}$,

$$\mathbb{E}_{\geq 3} Z = \int e^{N\xi_{\geq 3}(1)/2} e^{H_{\leq 2}(\boldsymbol{\sigma})} \mu_0(\mathsf{d}\boldsymbol{\sigma})$$
$$\geq e^{N\xi(1)/2 - cN/10},$$

whence Eq. (7.77) yields

$$\operatorname{Err}_{1} \leqslant e^{-cN/16} \left(\mathbb{E}_{\geqslant 3} Z(T) \right)^{2k}.$$
(7.78)

We also note here the estimate

$$\mathbb{E}_{\geq 3}Z(T) = \mathbb{E}_{\geq 3}Z - \mathbb{E}_{\geq 3}Z(T^c) \ge (1 - e^{-cN/10})\mathbb{E}_{\geq 3}Z,$$
(7.79)

which holds with probability at least $1 - \exp(-cN/8)$ over $W^{(2)}$, as claimed in Eq. (7.39).

Combining the error estimates (7.78), (7.71) in the moment bound (7.70), we get, with probability at least $1 - \exp(-Nc)$ with respect to W_2 ,

$$\mathbb{E}_{\geq 3}\left\{\left(Z(T) - \mathbb{E}_{\geq 3}Z(T)\right)^{2k}\right\} \leq C N^{-L/2} \left(\mathbb{E}_{\geq 3}Z(T)\right)^{2k}.$$
(7.80)

Adjusting c, we have

.

$$\mathbb{P}\left(|Z - \mathbb{E}_{\geq 3}Z(T(\delta))| > \varepsilon \mathbb{E}_{\geq 3}Z(T(\delta))\right) \leq \varepsilon^{-2L} N^{-L/2} + e^{-cN}$$

Estimating the magnetization, Eq. (7.40). We next apply the same argument to the magnetization. First, we note that

$$\mathbb{E}_{\geq 3}\left\{\left(\int_{T(\delta)} \sigma_1 e^{H(\boldsymbol{\sigma})} \mu_0(\mathsf{d}\boldsymbol{\sigma})\right)^{2k}\right\} = \mathbb{E}_{\geq 3} \int_{T(\delta)^{2k}} \prod_{i=1}^{2k} \sigma_1^i e^{\sum_{i=1}^{2k} H(\boldsymbol{\sigma}^i)} \mu_0^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma})$$
(7.81)

$$= \mathbb{E}_{\geq 3} \int_{A_{2k}(\delta)} \prod_{i=1}^{2k} \sigma_1^i e^{\sum_{i=1}^{2k} H(\boldsymbol{\sigma}^i)} \mu^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma}) + \mathsf{Err}_3, \qquad (7.82)$$

where

$$\operatorname{Err}_{3} := \mathbb{E}_{\geq 3} \int_{T(\delta)^{2k}} \prod_{i=1}^{2k} \sigma_{1}^{i} e^{\sum_{i=1}^{2k} H(\boldsymbol{\sigma}^{i})} \mu^{\otimes 2k} (\mathrm{d}\boldsymbol{\sigma}) - \mathbb{E}_{\geq 3} \int_{A_{2k}(\delta)} \prod_{i=1}^{2k} \sigma_{1}^{i} e^{\sum_{i=1}^{2k} H(\boldsymbol{\sigma}^{i})} \mu^{\otimes 2k} (\mathrm{d}\boldsymbol{\sigma}).$$
(7.83)

We have

$$\begin{aligned} \left| \int_{T(\delta)^{2k}} \prod_{i=1}^{2k} \sigma_{1}^{i} e^{\sum_{i=1}^{2k} H(\sigma^{i})} \mu_{0}^{\otimes 2k} (\mathrm{d}\sigma) - \int_{A_{2k}(T(\delta),\delta)} \prod_{i=1}^{2k} \sigma_{1}^{i} e^{\sum_{i=1}^{2k} H(\sigma^{i})} \mu_{0}^{\otimes 2k} (\mathrm{d}\sigma) \right| \\ &= \left| \int_{T(\delta)^{2k}:\max_{i\neq j} |\langle \sigma^{i}, \sigma^{j} \rangle_{N}| > \delta} \prod_{i=1}^{2k} \sigma_{1}^{i} e^{\sum_{i=1}^{2k} H(\sigma^{i})} \mu_{0}^{\otimes 2k} (\mathrm{d}\sigma) \right| \\ \stackrel{(a)}{\leq} N^{k} (2k)^{2} e^{-cN + N\xi(1)/2} Z(T(\delta))^{2k-1} \\ \stackrel{(b)}{\leqslant} e^{-cN/(2k) + Nk\xi(1)} + e^{-cN/(2k)} Z(T(\delta))^{2k} \\ \stackrel{(c)}{\leqslant} e^{-cN/(2k) + Nk\xi(1)} + e^{-cN/(2k)} \left(\mathbb{E}_{\geq 3} Z(T(\delta)) \right)^{2k} , \end{aligned}$$
(7.84)

where in (a) we used Lemma 7.7, in (b) the AM-GM inequality, and (c) holds with probability at least $1 - \exp(-cN)$ by Eq. (7.37).

Using Eq. (7.84) and Lemma 7.3 we obtain that, with probability at least $1 - e^{-cN}$ over $W^{(2)}$,

$$|\mathsf{Err}_3| \leqslant e^{-cN} \left(\mathbb{E}_{\geqslant 3} Z(T(\delta)) \right)^{2k}.$$
(7.85)

Turning to the main term in Eq. (7.82),

$$\mathbb{E}_{\geq 3} \int_{A_{2k}(\delta)} \prod_{i=1}^{2k} \sigma_1^i e^{\sum_{i=1}^{2k} H(\boldsymbol{\sigma}^i)} \mu^{\otimes 2k} (\mathsf{d}\boldsymbol{\sigma}) \\ = e^{Nk\xi_{\geq 3}(1)} \int_{A_{2k}(\delta)} \prod_{i=1}^{2k} \sigma_1^i \exp\left\{\sum_{i=1}^{2k} H_{\leq 2}(\boldsymbol{\sigma}^i) + \frac{N}{2} \sum_{i\neq j} \xi_{\geq 3}(\langle \boldsymbol{\sigma}^i, \boldsymbol{\sigma}^j \rangle_N)\right\} \mu_0^{\otimes 2k} (\mathsf{d}\boldsymbol{\sigma})$$

By Eqs. (7.44) and (7.45) in Lemma 7.10, we can bound

$$\begin{split} \mathbb{E}_{\geqslant 3} & \int_{A_{2k}(\delta)} \prod_{i=1}^{2k} \sigma_1^i \, e^{\sum_{i=1}^{2k} H(\sigma^i)} \mu^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma}) \\ &= e^{Nk\xi_{\geqslant 3}(1)} \int_{A_{2k}(N^{-1/2+c})} \prod_{i=1}^{2k} \sigma_1^i \exp\left\{\sum_{i=1}^{2k} H_{\leqslant 2}(\boldsymbol{\sigma}^i) + \frac{N}{2} \sum_{i \neq j} \xi_{\geqslant 3}(\langle \boldsymbol{\sigma}^i, \boldsymbol{\sigma}^j \rangle_N)\right\} \mu_0^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma}) \\ &+ O\left(N^k e^{-N^c + Nk\xi_{\geqslant 3}(1)} Z_{\leqslant 2, 2k}(A_{2k}(\delta))\right). \end{split}$$

To bound the first term, using Lemma 7.5,

$$\begin{split} &\int_{A_{2k}(\delta)} \prod_{i=1}^{2k} \sigma_1^i \exp\left\{\sum_{i=1}^{2k} H_{\leq 2}(\boldsymbol{\sigma}^i)\right\} \mu_0^{\otimes 2k}(\mathrm{d}\boldsymbol{\sigma}) \\ &= \int_{S_N^{2k}} \prod_{i=1}^{2k} \sigma_1^i \exp\left\{\sum_{i=1}^{2k} H_{\leq 2}(\boldsymbol{\sigma}^i)\right\} \mu_0^{\otimes 2k}(\mathrm{d}\boldsymbol{\sigma}) + O_k\left(N^k e^{-c\delta^2 N} (Z_{\leq 2})^{2k}\right) \\ &= \left(\int_{S_N} \sigma_1 \exp\left\{H_{\leq 2}(\boldsymbol{\sigma})\right\} \mu_0(\mathrm{d}\boldsymbol{\sigma})\right)^{2k} + O_k\left(N^k e^{-c\delta^2 N} (Z_{\leq 2})^{2k}\right). \end{split}$$

By Lemma 7.3, we then obtain

$$\int_{A_{2k}(\delta)} \prod_{i=1}^{2k} \sigma_1^i \exp\left\{\sum_{i=1}^{2k} H_{\leq 2}(\boldsymbol{\sigma}^i)\right\} \mu_0^{\otimes 2k}(\mathsf{d}\boldsymbol{\sigma}) \leq C_k \left(|u_1|^{2k} + N^k e^{-c\delta^2 N}\right) (Z_{\leq 2})^{2k}.$$
(7.86)

On the other hand, by taking the Taylor expansion of $\exp\left\{\frac{N}{2}\sum_{i\neq j}\xi_{\geq 3}(\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j}\rangle_{N})\right\}$ up to terms of order L = Ck for C > 2, we obtain that, for $\xi_{\geq 3, \leq \ell}(s) = \sum_{3 \leq p \leq \ell} \beta_{p}^{2} s^{p}$,

$$\frac{1}{(Z_{\leq 2})^{2k}} \int_{A_{2k}(N^{-1/2+c})} \prod_{i=1}^{2k} \sigma_{1}^{i} \left(\exp\left(N \sum_{i < j} \xi_{\geq 3}(\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j} \rangle_{N})\right) - 1 \right) e^{\sum_{i=1}^{2k} H_{\leq 2}(\boldsymbol{\sigma}^{i})} \mu_{0}^{\otimes 2k}(\mathrm{d}\boldsymbol{\sigma}) \\
= O(N^{-k}) + \frac{1}{(Z_{\leq 2})^{2k}} \sum_{\ell \leq L} \frac{N^{\ell}}{\ell!} \int_{A_{2k}(N^{-1/2+c})} \prod_{i=1}^{2k} \sigma_{1}^{i} \left(\sum_{i < j} \xi_{\geq 3}(\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j} \rangle_{N})\right)^{\ell} e^{\sum_{i=1}^{2k} H_{\leq 2}(\boldsymbol{\sigma}^{i})} \mu_{0}^{\otimes 2k}(\mathrm{d}\boldsymbol{\sigma}) \\
= O(N^{-k}) + \frac{1}{(Z_{\leq 2})^{2k}} \sum_{\ell \leq L} \frac{N^{\ell}}{\ell!} \int_{A_{2k}(N^{-1/2+c})} \prod_{i=1}^{2k} \sigma_{1}^{i} \left(\sum_{i < j} \xi_{\geq 3, \leq 4k}(\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j} \rangle_{N})\right)^{\ell} e^{\sum_{i=1}^{2k} H_{\leq 2}(\boldsymbol{\sigma}^{i})} \mu_{0}^{\otimes 2k}(\mathrm{d}\boldsymbol{\sigma}) \\
\stackrel{(a)}{=} O(N^{-k} + e^{-N^{c}}) + \frac{1}{(Z_{\leq 2})^{2k}} \sum_{\ell \leq L} \frac{N^{\ell}}{\ell!} \int_{S_{N}^{2k}} \prod_{i=1}^{2k} \sigma_{1}^{i} \left(\sum_{i < j} \xi_{\geq 3, \leq 2k}(\langle \boldsymbol{\sigma}^{i}, \boldsymbol{\sigma}^{j} \rangle_{N})\right)^{\ell} e^{\sum_{i=1}^{2k} H_{\leq 2}(\boldsymbol{\sigma}^{i})} \mu_{0}^{\otimes 2k}(\mathrm{d}\boldsymbol{\sigma}) \\
\stackrel{(b)}{=} O(N^{-k} + e^{-N^{c}}) + (1 + \|\boldsymbol{u}\|)^{O(k^{2})} O_{k} \left(\sum_{\ell < k} |\boldsymbol{u}|^{2k-2\ell} N^{-\ell} + \sum_{k < \ell < L} N^{-\ell/2-k/2}\right) \tag{7.87}$$

$$= (1 + \|\boldsymbol{u}\|)^{O(k^2)} O(N^{-k} + e^{-N^c} + |u_1|^{2k}),$$
(7.88)

where in (a) we used again Lemma 7.10 and in (b) Lemma 7.4.

We thus have from Eqs. (7.85), (7.86), (7.88),

$$\mathbb{E}_{\geq 3} \left\{ \left(\int_{T(\delta)} \sigma_1 e^{H(\boldsymbol{\sigma})} \mu_0(\mathsf{d}\boldsymbol{\sigma}) \right)^{2k} \right\}$$

$$\leq C_k (1 + \|\boldsymbol{u}\|)^{Ck^2} \left(|u_1|^{2k} + N^{-k} + N^k e^{-N^c} + e^{-cN} \right) \left(\mathbb{E}_{\geq 3} Z(T(\delta)) \right)^{2k}$$

The desired claim (7.40) follows from Markov Inequality upon adjusting the constant c.

7.1.4 Magnetization in the band: Proof of Lemma 7.2

In the remaining of this section, we denote by μ the Gibbs measure associated to $H(\boldsymbol{\sigma})$, i.e.

$$\mu(\mathsf{d}\boldsymbol{\sigma}) \propto \exp(H(\boldsymbol{\sigma})) \,\mu_0(\mathsf{d}\boldsymbol{\sigma})$$

In the following we estimate the components of $\langle \boldsymbol{\sigma} \rangle = (\langle \sigma_1 \rangle, \dots, \langle \sigma_N \rangle)$ in the basis of eigenvectors of the quadratic part of the Hamiltonian $\boldsymbol{W}^{(2)}$. For simplicity of notation, we consider the component $\langle \sigma_1 \rangle$ but we emphasize that this does not necessarily correspond to the largest (or smallest) eigenvalue of $\boldsymbol{W}^{(2)}$. Defining $\boldsymbol{\sigma}_{-1} = (\sigma_2, \dots, \sigma_N)$, we have

$$\int \sigma_1 e^{H(\boldsymbol{\sigma})} \mu_0(\mathsf{d}\boldsymbol{\sigma}) = \frac{1}{Z} \int \sigma_1 e^{\sigma_1 u_1 + \Lambda_1 \sigma_1^2} \hat{E}(\sigma_1) \mathsf{d}\sigma_1.$$
(7.89)

where we defined

$$\hat{E}(\sigma_{1}) = C_{N}(1 - \sigma_{1}^{2}/N)^{(N-3)/2} \int \exp\left(\frac{\sigma_{1}}{N} \sum_{i,j=2}^{N} \tilde{g}_{1ij}\sigma_{i}\sigma_{j}\right) e^{H_{\sigma_{1}}(\boldsymbol{\sigma}_{-1})} \mu_{0,\sqrt{N-\sigma_{1}^{2}}}(\mathsf{d}\boldsymbol{\sigma}_{-1}),$$

 $\mu_{0,\rho}$ denotes the uniform measure over the sphere of radius ρ ,

$$C_N := \frac{\Gamma(N-1)}{\Gamma((N-1)/2)^2 2^{N-2} \sqrt{N}} = \frac{1}{\sqrt{2\pi}} + O(N^{-1}),$$

and

$$H_{\sigma_1}(\boldsymbol{\sigma}_{-1}) := \sum_{i=2}^N (\sigma_i u_i + \Lambda_i \sigma_i^2) + N^{-1} \sum_{i,j,k>1} g_{ijk}^{(3)} \sigma_i \sigma_j \sigma_k + \sum_{p \ge 4} H_p(\boldsymbol{\sigma}).$$

Here \tilde{g}_{1ij} is the sum of g over permutations of (1, i, j). In particular $\tilde{g}_{1ij} = \tilde{g}_{1ji}$

$$(\widetilde{g}_{1ij})_{1 < i < j} \sim_{iid} \mathcal{N}(0, 3\beta_3^2/2), \quad (\widetilde{g}_{1ii})_{1 < i} \sim_{iid} \mathcal{N}(0, 3\beta_3^2).$$
 (7.90)

We set $\hat{E}(\sigma_1) = 0$ for $|\sigma_1| > \sqrt{N}$.

By Lemma 7.3 and Lemma 7.8, with probability $1 - e^{-cN} - N^{-C}$,

$$Z = (1 + O(N^{-c})) \sqrt{\frac{2}{G''(\gamma_*)}} \cdot \exp\left\{N\left[\xi_{\geq 3}(1) - \frac{1}{2}\log(2e) + G(\gamma_*)\right]\right\},\$$

where $G(\gamma)$ and γ_* where defined in Eqs. (7.10) and (7.11).

In estimating $\langle \sigma_1 \rangle$, we first anneal over $g_{\geq 4}$ and $g_{3-} := (g_{ijk} : 1 < i < j < k)$. We have

$$E(\sigma_1) := \mathbb{E}_{\boldsymbol{g}_{3-}, \boldsymbol{g}_{\geq 4}}[\hat{E}(\sigma_1)] = C_N \left(1 - \frac{\sigma_1^2}{N}\right)^{(N-3)/2} \int \exp\left(\frac{\sigma_1}{N} \sum_{i,j=2}^N \widetilde{g}_{1ij} \sigma_i \sigma_j\right) \\ \exp\left\{H_{\leq 2}(\boldsymbol{\sigma}_{-1}) + N\xi_{\geq 4}(1)/2 + N\beta_3^2 (1 - \sigma_1^2/N)^3/2\right\} \mu_{0,\sqrt{N-\sigma_1^2}}(\mathsf{d}\boldsymbol{\sigma}_{-1}).$$

The next lemma show that this expectation is an accurate approximation of $\hat{E}(\sigma_1)$.

Lemma 7.12. We have for an appropriate $c \in (0, 1/8)$ that, with probability $1 - N^{-c}$,

$$\int \sigma_1 e^{u_1 \sigma_1 + \Lambda_1 \sigma_1^2} \hat{E}(\sigma_1) \, \mathrm{d}\sigma_1 =$$

$$= \int \sigma_1 e^{u_1 \sigma_1 + \Lambda_1 \sigma_1^2} E(\sigma_1) \, \mathrm{d}\sigma_1 + O\left(N^{-1/2+c}(|u_1| + N^{-1/2}) \int e^{u_1 \sigma_1 + \Lambda_1 \sigma_1^2} E(\sigma_1) \, \mathrm{d}\sigma_1\right).$$

Before proving Lemma 7.12, we use it to prove Lemma 7.1.

Proof of Lemma 7.1. For $U(\sigma_1) := N\xi_{\geq 4}(1)/2 + N\beta_3^2(1 - \sigma_1^2/N)^3/2$, we have

$$E(\sigma_{1}) = C_{N} \left(1 - \frac{\sigma_{1}^{2}}{N}\right)^{(N-3)/2} \int \exp\left(\frac{\sigma_{1}}{N} \sum_{i,j=2}^{N} \widetilde{g}_{1ij} \sigma_{i} \sigma_{j}\right) e^{H_{\leq 2}(\boldsymbol{\sigma}_{-1}) + U(\sigma_{1})} \mu_{0,\sqrt{N-\sigma_{1}^{2}}}(\mathsf{d}\boldsymbol{\sigma}_{-1}).$$

Again by Lemma 7.3, for $V = V(\sigma_1) := \Lambda_{-1} + \Delta$, where Λ_{-1} is the diagonal matrix with entries corresponding to the spectrum of $W^{(2)}$, with Λ_1 replaced by 0, and $\Delta := \sigma_1 N^{-1} \tilde{G}$ with $\tilde{G}_{ij} = \tilde{g}_{1ij}$,

$$E(\sigma_1) = (1 + O(N^{-1})) \frac{1}{(2e)^{(N-1)/2} \sqrt{2\pi}} (1 - \sigma_1^2/N)^{-1} \sqrt{\frac{2}{G_{\sigma_1}''(\gamma_*(\sigma_1))}} \exp\left(U(\sigma_1) + NG_{\sigma_1}(\gamma_*(\sigma_1))\right),$$
(7.91)

where we defined

$$G_{\sigma_1}(\gamma) := (1 - \sigma_1^2/N)\gamma - \frac{1}{2N}\log\det(\gamma \boldsymbol{I}_{N-1} - \boldsymbol{V}) + \frac{1}{4N}\langle \boldsymbol{u}, (\gamma \boldsymbol{I}_{N-1} - \boldsymbol{V})^{-1}\boldsymbol{u}\rangle,$$
(7.92)

$$\gamma_*(\sigma_1) = \arg\max G_{\sigma_1}(\gamma) \,. \tag{7.93}$$

By Lemma 7.12, we have

$$\int \sigma_1 \,\mu(\mathsf{d}\boldsymbol{\sigma}) = \frac{\int \sigma_1 e^{u_1 \sigma_1 + \Lambda_1 \sigma_1^2} E(\sigma_1) \mathsf{d}\sigma_1}{\int e^{u_1 \sigma_1 + \Lambda_1 \sigma_1^2} E(\sigma_1) \mathsf{d}\sigma_1} + O\left(N^{-1/2+c}(|u_1| + N^{-1/2})\right) \,. \tag{7.94}$$

We next estimate these integrals by approximating their argument for small σ_1 . Note that by Lemma 7.5 and Lemma 7.7, we can restrict these integrals to $|\sigma_1| \leq C \log N$ making a negligible error.

It is easy to see that, for $\sigma_1 = 0$, we recover $G_{\sigma_1}(\gamma) = G_0(\gamma)$, where $G_0(\gamma)$ is the same function defined in Eq. (7.10), with N replaced by N - 1. To leading order, we can expand

$$\begin{aligned} G_{\sigma_1}(\gamma) &= \\ &= (1 - \sigma_1^2/N)\gamma - \frac{1}{2N}\log\det(\gamma \boldsymbol{I} - \boldsymbol{V}) + \frac{1}{4N}\langle \boldsymbol{u}, (\gamma \boldsymbol{I} - \boldsymbol{V})^{-1}\boldsymbol{u} \rangle \\ &= (1 - \sigma_1^2/N)\gamma - \frac{1}{2N}\log\det(\gamma \boldsymbol{I} - \boldsymbol{V}) + \frac{1}{4N}\langle \boldsymbol{u}, (\boldsymbol{I} + (\gamma \boldsymbol{I} - \boldsymbol{\Lambda}_{-1})^{-1}\boldsymbol{\Delta} + \boldsymbol{E}_N)(\gamma \boldsymbol{I} - \boldsymbol{\Lambda}_{-1})^{-1}\boldsymbol{u} \rangle. \end{aligned}$$

where $\|\boldsymbol{E}_N\|_{\text{op}} = O(N^{-1})$ with probability $1 - \exp(-cN)$ over $\boldsymbol{W}^{(3)}$. Therefore

$$G_{\sigma_{1}}(\gamma) - G_{0}(\gamma) = -\frac{\gamma \sigma_{1}^{2}}{N} + \frac{1}{2N} \log(\gamma - \Lambda_{1}) - \frac{1}{2N} \log \det \left(\boldsymbol{I} - (\gamma \boldsymbol{I} - \boldsymbol{\Lambda}_{-1})^{-1/2} \boldsymbol{\Delta} (\gamma \boldsymbol{I} - \boldsymbol{\Lambda}_{-1})^{-1/2} \right) + \frac{1}{4N} \langle \boldsymbol{u}, (\gamma \boldsymbol{I} - \boldsymbol{\Lambda}_{-1})^{-1} \boldsymbol{\Delta} (\gamma \boldsymbol{I} - \boldsymbol{\Lambda}_{-1})^{-1} \boldsymbol{u} \rangle + O(\|\boldsymbol{u}\|^{2}/N^{2}).$$
(7.95)

on $\gamma > \max_i \Lambda_i + \varepsilon$. Since the above difference (and its derivative with respect to λ) is of order σ_1/\sqrt{N} and G is strongly convex in a neighborhood of γ_* , it follows that $\gamma_*(\sigma_1) = \gamma_* + O(\sigma_1^2/N)$. We will therefore restrict ourselves to $|\gamma - \gamma_*| \leq CN^{-1}(\log N)^2$.

We next expand the log-determinant term in the difference. Defining

$$D_{2} := \sum_{i,j=1}^{N} \left(N^{-1} (\gamma \boldsymbol{I} - \boldsymbol{\Lambda}_{-1})^{-1/2} \tilde{\boldsymbol{G}} (\gamma \boldsymbol{I} - \boldsymbol{\Lambda}_{-1})^{-1/2} \right)_{ij}^{2},$$
(7.96)

we have

$$\mathsf{Tr}\Big((\gamma I - \Lambda_{-1})^{-1/2} \Delta(\gamma I - \Lambda_{-1})^{-1/2}) = \frac{\sigma_1}{N} \sum_{i \neq 1} (\gamma - \Lambda_i)^{-1} \tilde{g}_{1ii} + O(N^{-1}),$$
(7.97)

$$\mathsf{Tr}\Big(\big(\gamma \boldsymbol{I} - \boldsymbol{\Lambda}_{-1}\big)^{-1/2} \boldsymbol{\Delta} (\gamma \boldsymbol{I} - \boldsymbol{\Lambda}_{-1})^{-1/2} \Big)^2 \Big) = D_2 \sigma_1^2, \tag{7.98}$$

$$\operatorname{Tr}\left(\left((\gamma \boldsymbol{I} - \boldsymbol{\Lambda}_{-1})^{-1/2} \boldsymbol{\Delta}(\gamma \boldsymbol{I} - \boldsymbol{\Lambda}_{-1})^{-1/2}\right)^k\right) = O(N^{-1}) \quad \text{for } k \ge 3.$$
(7.99)

Thus, with high probability,

$$\frac{1}{2N}\log\det\left(\boldsymbol{I} - (\gamma \boldsymbol{I} - \boldsymbol{\Lambda}_{-1})^{-1/2}\boldsymbol{\Delta}(\gamma \boldsymbol{I} - \boldsymbol{\Lambda}_{-1})^{-1/2}\right) = -\frac{\sigma_1}{2N^2}\sum_{i\neq 1}(\gamma - \Lambda_i)^{-1}\widetilde{g}_{1ii} - \frac{D_2}{4N}\sigma_1^2 + O(N^{-2}).$$

For $\gamma = \gamma_*(\sigma_1) = \gamma_* + O(\sigma_1^2/N)$, we can compute

$$\mathbb{E}D_2 = \frac{3\beta_3^2}{2N^2} \Big(\sum_{i \neq 1} (\gamma_* - \Lambda_i)^{-1}\Big)^2 + O(N^{-1})$$

and

$$\operatorname{Var}(D_2) = \frac{\beta_3^4}{N^4} O\left(\left(\sum_{i \neq 1} \left(\gamma_* - \Lambda_i \right)^{-2} \right)^2 \right) = O(N^{-2}).$$

Furthermore, recalling the stationarity condition $G'(\gamma_*) = 0$, which yields

$$\frac{1}{2N}\sum_{i=1}^{N}\frac{1}{\gamma_{*}-\Lambda_{i}} = 1 + \frac{1}{4N}\sum_{i=1}^{N}\frac{u_{i}^{2}}{(\gamma_{*}-\Lambda_{i})^{2}}$$

which yields (for $\|\boldsymbol{u}\| \leq N^{c_0} \sum_{i \geq 1} (\gamma_* - \Lambda_i)^{-1} = 2N + O(N^{2c_0})$, and therefore

$$\mathbb{E}D_2 = 6\beta_3^2 + O(N^{-1}).$$
(7.100)

Substituting the above estimates in Eq. (7.95) the following holds with probability at least $1 - \exp(-N^c)$, for $|\sigma_1| \leq C \log N$,

$$\min_{\gamma} G_{\sigma_1}(\gamma) - G_0(\gamma_*) = -\frac{\gamma_* \sigma_1^2}{N} + \frac{1}{2N} \log(\gamma_* - \Lambda_1) + \frac{\sigma_1}{2N^2} \sum_{i=1}^N (\gamma_* - \Lambda_i)^{-1} \widetilde{g}_{1ii} + \frac{D_2}{4N} \sigma_1^2 + O(N^{-2+3c_0}).$$

Letting $a_N := C \log N$, and using Eq. (7.91),

$$\begin{split} \int \sigma_{1} e^{\sigma_{1}u_{1}+\Lambda_{1}\sigma_{1}^{2}} E(\sigma_{1}) \mathrm{d}\sigma_{1} \\ &= \frac{1}{(2e)^{(N-1)/2}\sqrt{2\pi}} \sqrt{\frac{2}{G''(\gamma_{*})}} \int_{[-a_{N},a_{N}]} (1-\sigma_{1}^{2}/N)^{-1}\sigma_{1} \\ &\quad \exp\left\{NG_{0}(\gamma_{*}) + \frac{1}{2}\log(\gamma_{*}-\Lambda_{1}) + U(\sigma_{1}) + \sigma_{1}\left(u_{1} + \frac{1}{2N}\sum_{i}(\gamma-\Lambda_{i})^{-1}\widetilde{g}_{1ii}\right) \right. \\ &\quad - \left(-\Lambda_{1} - \frac{1}{4}D_{2} + \gamma_{*}\right)\sigma_{1}^{2} + O(N^{-1+3c_{0}})\right\} \mathrm{d}\sigma_{1} + \delta_{N} \\ \stackrel{(a)}{=} (2e)^{-(N-1)/2} \sqrt{\frac{1}{\pi G''(\gamma_{*})}} \int_{[-a_{N},a_{N}]} \sigma_{1} \exp\left\{NG_{0}(\gamma_{*}) + \frac{1}{2}\log(\gamma_{*}-\Lambda_{1}) \right. \\ &\quad + U(\sigma_{1}) + \sigma_{1}\left(u_{1} + \frac{1}{2N}\sum_{i}(\gamma-\Lambda_{i})^{-1}\widetilde{g}_{1ii}\right) - \left(-\Lambda_{1} - \frac{3}{2}\beta_{3}^{2} + \gamma_{*}\right)\sigma_{1}^{2} + O(N^{-1+3c_{0}})\right\} \mathrm{d}\sigma_{1} + \delta_{N} \\ &= (1 + O(N^{-1}))(2e)^{-(N-1)/2} \sqrt{\frac{1}{\pi G''(\gamma_{*})}} \int_{[-a_{N},a_{N}]} \sigma_{1} \exp\left\{NG_{0}(\gamma_{*}) + \frac{1}{2}\log(\gamma_{*}-\Lambda_{1}) \right. \\ &\quad + \frac{N}{2}(\xi_{\geq 4}(1) + \beta_{3}^{2})/2 + \sigma_{1}\left(u_{1} + \frac{1}{2N}\sum_{i}(\gamma-\Lambda_{i})^{-1}\widetilde{g}_{1ii}\right) - (-\Lambda_{1} + \gamma_{*})\sigma_{1}^{2} + O(N^{-1+3c_{0}})\right\} \mathrm{d}\sigma_{1} + \delta_{N} , \end{split}$$
where in (a) we used Eq. (7.100), and

$$|\delta_N| \leqslant N^{-1} \int e^{\sigma_1 u_1 + \Lambda_1 \sigma_1^2} E(\sigma_1) \mathsf{d}\sigma_1 \,. \tag{7.101}$$

Therefore, we obtain

$$\frac{\int \sigma_1 \exp(\sigma_1 u_1 + \Lambda_1 \sigma_1^2) E(\sigma_1) \mathsf{d}\sigma_1}{\int \exp(\sigma_1 u_1 + \Lambda_1 \sigma_1^2) E(\sigma_1) \mathsf{d}\sigma_1} = \frac{u_1 + N^{-1} \sum_i (\gamma_* - \Lambda_i)^{-1} \widetilde{g}_{1ii}}{2(\gamma_* - \Lambda_1)} + O(N^{-1}).$$
(7.102)

which completes the proof using Eq. (7.94).

Finally, we prove Lemma 7.12. The main idea is that the error in annealing can be controlled by accurate estimates of certain quantities involving overlap over the quadratic model on σ_{-1} , which follows from Laplace transform and expansion of the dependence on σ_1 .

Proof of Lemma 7.12. Define

$$W(\boldsymbol{\sigma}^{1}, \boldsymbol{\sigma}^{2}) := \frac{1}{N} \mathbb{E} \Big\{ \Big(H_{3}(\boldsymbol{\sigma}_{-1}^{1}) + H_{\geq 4}(\boldsymbol{\sigma}^{1}) \Big) \Big(H_{3}(\boldsymbol{\sigma}_{-1}^{2}) + H_{\geq 4}(\boldsymbol{\sigma}^{2}) \Big) \Big\} \\ = \beta_{3}^{2} \langle \boldsymbol{\sigma}_{-1}^{1}, \boldsymbol{\sigma}_{-1}^{2} \rangle_{N}^{3} + \xi_{\geq 4} \big(\langle \boldsymbol{\sigma}_{-1}^{1}, \boldsymbol{\sigma}_{-1}^{2} \rangle_{N} + \sigma_{1}^{1} \sigma_{1}^{2} / N \big) \,,$$

where, with an abuse of notation, $H_3(\boldsymbol{\sigma}_{-1}^a) := N^{-1} \sum_{i,j,k=2}^N \sigma_i^a \sigma_j^a \sigma_k^a$ (and a similar notation will be used for $H_{\leq 2}(\boldsymbol{\sigma}_{-1}^a)$ below). Note that $W(\boldsymbol{\sigma}^1, \boldsymbol{\sigma}^1) = \xi_{\geq 4}(1) + \beta_3^2(1 - (\sigma_1^1)^2/N)^3$. For a Borel set $U \subseteq S_N^2$, define

$$\begin{split} Q(U) &:= \int_{U} \sigma_{1}^{1} \sigma_{1}^{2} e^{u_{1}(\sigma_{1}^{1} + \sigma_{1}^{2}) + \Lambda_{1}((\sigma_{1}^{1})^{2} + (\sigma_{1}^{2})^{2})} \cdot \\ & \quad \cdot \exp\left\{ N^{-1} \Big(\sigma_{1}^{1} \sum_{i,j=2}^{N} \widetilde{g}_{1ij} \sigma_{i}^{1} \sigma_{j}^{1} + \sigma_{1}^{2} \sum_{i,j=2}^{N} \widetilde{g}_{1ij} \sigma_{i}^{2} \sigma_{j}^{2} \Big) + H_{\leq 2}(\boldsymbol{\sigma}_{-1}^{1}) + H_{\leq 2}(\boldsymbol{\sigma}_{-1}^{2}) \right\} \\ & \quad e^{N[W(\boldsymbol{\sigma}^{1}, \boldsymbol{\sigma}^{1}) + W(\boldsymbol{\sigma}^{2}, \boldsymbol{\sigma}^{2})]/2} \big\{ \exp[NW(\boldsymbol{\sigma}^{1}, \boldsymbol{\sigma}^{2})] - 1 \big\} \mu_{0}^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma}) \, . \end{split}$$

Expanding the square and taking expectation, we obtain

$$\mathbb{E}_{\boldsymbol{g}_{\geq 4},\boldsymbol{g}_{3-}}\left[\left\{\int_{T(\delta)}\sigma_1\left(e^{H(\boldsymbol{\sigma})}-\mathbb{E}_{\boldsymbol{g}_{\geq 4},\boldsymbol{g}_{3-}}e^{H(\boldsymbol{\sigma})}\right)\,\mu_0(\mathsf{d}\boldsymbol{\sigma})\right\}^2\right]=Q(T(\delta)\times T(\delta))\,.$$

Further, writing $T = T(\delta)$, and $A_2 = A_2(N^{-1/2+c})$, we obtain that, with probability at least $1 - \exp(-N^c)$,

$$\begin{aligned} \left|Q(A_{2}) - Q(T \times T)\right| &= N \cdot Q\left(A_{2} \setminus T \times T\right) + N \cdot Q\left(T \times T \setminus A_{2}\right) \\ &\stackrel{(a)}{\leqslant} N \cdot Z_{\leqslant 2,2}\left(A_{2} \setminus T \times T\right) e^{N\xi_{\geqslant 3}(1)} + N \cdot Z_{\leqslant 2,2}\left(T \times T \setminus A_{2}\right) e^{N\xi_{\geqslant 3}(1)} \\ &\stackrel{(b)}{\leqslant} e^{-cN}(Z_{\leqslant 2})^{2} e^{N\xi_{\geqslant 3}(1)} + e^{-N^{c}}(Z_{\leqslant 2})^{2} e^{N\xi_{\geqslant 3}(1)} \end{aligned}$$

where in (a) we used the fact that $|\sigma_1^1 \sigma_1^2| \leq N$, and in (b) the first term was bounded by using $Z_{\leq 2,2}((T \times T)^c) \leq 2Z_{\leq 2}(T^c)Z_{\leq 2}$ and applying Lemma 7.7, see Eq. (7.35), and the second by $Z_{\leq 2,2}(T \times T \setminus A_2) \leq C_{\leq 2,2}(T \times T \setminus A_2)$

 $Z_{\leqslant 2,2}(A_2^c)$ and using Lemma 7.10, Eq. (7.44). Hence we conclude that

$$\mathbb{E}_{\boldsymbol{g}_{\geq 4},\boldsymbol{g}_{3-}}\left[\left\{\int_{T(\delta)}\sigma_1\left(e^{H(\boldsymbol{\sigma})} - \mathbb{E}_{\boldsymbol{g}_{\geq 4},\boldsymbol{g}_{3-}}e^{H(\boldsymbol{\sigma})}\right)\,\mu_0(\mathsf{d}\boldsymbol{\sigma})\right\}^2\right]$$
(7.103)

$$= Q(A_2(N^{-1/2+c})) + O\left(e^{-N^c + N\xi_{\ge 1}(1)}(Z_{\le 2})^2\right).$$
(7.104)

By Taylor expansion, always using the shorthand $A_2 = A_2(N^{-1/2+c})$,

$$\begin{split} Q(A_2) &= \int_{A_2} \sigma_1^1 \sigma_1^2 \exp \left\{ \sum_{i=1}^2 \left(\left(u_1 \sigma_1^i + \Lambda_1(\sigma_1^i)^2 \right) + H_{\leqslant 2}(\boldsymbol{\sigma}_{-1}^i) + NW(\boldsymbol{\sigma}^i, \boldsymbol{\sigma}^i) \right) \right\} \\ & \exp \left\{ \frac{\sigma_1^1}{N} \sum_{i,j=2}^N \widetilde{g}_{1ij} \sigma_i^1 \sigma_j^1 + \frac{\sigma_1^2}{N} \sum_{i,j=2}^N \widetilde{g}_{1ij} \sigma_i^2 \sigma_j^2 \right\} \cdot \left\{ \sum_{\ell=1}^L \frac{1}{\ell!} (NW(\boldsymbol{\sigma}^1, \boldsymbol{\sigma}^2))^\ell + O(N^{-L/2+c}) \right\} \mu_0^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma}) \,. \end{split}$$

We estimate each term

$$T_{\ell}(a,b) := \int_{A_2} \sigma_1^1 \sigma_1^2 \exp\left\{\sum_{i=1}^2 \left(\left(u_1 \sigma_1^i + \Lambda_1(\sigma_1^i)^2\right) + H_{\leqslant 2}(\boldsymbol{\sigma}_{-1}^i) + NW(\boldsymbol{\sigma}^i, \boldsymbol{\sigma}^i) \right) \right\}$$
(7.105)
$$\exp\left\{\frac{\sigma_1^1}{N} \sum_{1 < i < j} \widetilde{g}_{1ij} \sigma_i^1 \sigma_j^1 + \frac{\sigma_1^2}{N} \sum_{i,j=2}^N \widetilde{g}_{1ij} \sigma_i^2 \sigma_j^2 \right\} \cdot N^{\ell} \langle \boldsymbol{\sigma}_{-1}^1, \boldsymbol{\sigma}_{-1}^2 \rangle_N^a (\sigma_1^1 \sigma_1^2/N)^b \, \mu_0^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma}) \, .$$

We can restrict ourselves to terms with $a \ge 3\ell$ and b = 0, or $a + b \ge 3\ell + 1$, since these are the terms that can arise in $Q(A_2)$. Let

$$\begin{split} \widehat{T}_{\ell}(a,b) &:= \int_{S_N^2} \sigma_1^1 \sigma_1^2 \, \exp\Big\{ \sum_{i=1}^2 \Big(\Big(u_1 \sigma_1^i + \Lambda_1(\sigma_1^i)^2 \Big) + H_{\leqslant 2}(\boldsymbol{\sigma}_{-1}^i) + NW(\boldsymbol{\sigma}^i, \boldsymbol{\sigma}^i) \Big) \Big\} \\ &\quad \exp\Big\{ \frac{\sigma_1^1}{N} \sum_{i,j=2}^N \widetilde{g}_{1ij} \sigma_i^1 \sigma_j^1 + \frac{\sigma_1^2}{N} \sum_{i,j=2}^N \widetilde{g}_{1ij} \sigma_i^2 \sigma_j^2 \Big\} \cdot N^{\ell} \langle \boldsymbol{\sigma}_{-1}^1, \boldsymbol{\sigma}_{-1}^2 \rangle_N^a (\sigma_1^1 \sigma_1^2/N)^b \, \mu_0^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma}) \,. \end{split}$$

By Lemma 7.10, Eq. (7.44), we have

$$|T_{\ell}(a,b) - \widehat{T}_{\ell}(a,b)| \leqslant e^{-N^c}.$$

Applying Lemma 7.6, we have, for appropriate $C_{i,j} = O(\|\boldsymbol{u}\|^{2a} + N^{\lfloor a/2 \rfloor})$,

$$\begin{split} |\hat{T}_{\ell}(a,b)| &\leqslant N^{\ell-b-a} \int_{S_N^2} (\sigma_1^1 \sigma_1^2)^{b+1} \exp \Big\{ \sum_{i=1}^2 \left(\left(u_1 \sigma_1^i + \Lambda_1(\sigma_1^i)^2 \right) + H_{\leqslant 2}(\boldsymbol{\sigma}_{-1}^i) + NW(\boldsymbol{\sigma}^i, \boldsymbol{\sigma}^i) \right) \Big\} \\ & \left\{ C_{0,0} + \sum_{i,j=0,(i,j) \neq (0,0)}^L C_{i,j} N^{-(i+j)/2} (\sigma_1^1)^i (\sigma_1^2)^j + O_L(N^{-L/2}) \right\} \, \mu_0^{\otimes 2}(\mathrm{d}\boldsymbol{\sigma}). \end{split}$$

Note that when b + 1 + i or b + 1 + j is odd,

$$\frac{\int_{S_N^2} (\sigma_1^1 \sigma_1^2)^{b+1} \exp\left\{\sum_{i=1}^2 \left(\left(u_1 \sigma_1^i + \Lambda_1(\sigma_1^i)^2\right) + H_{\leq 2}(\boldsymbol{\sigma}_{-1}^i) + W(\boldsymbol{\sigma}^i, \boldsymbol{\sigma}^i) \right) \right\} C_{i,j}(\sigma_1^1)^i (\sigma_1^2)^j \mu_0^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma}) \\
\int_{S_N^2} \exp\left\{\sum_{i=1}^2 \left(\left(u_1 \sigma_1^i + \Lambda_1(\sigma_1^i)^2\right) + H_{\leq 2}(\boldsymbol{\sigma}_{-1}^i) + W(\boldsymbol{\sigma}^i, \boldsymbol{\sigma}^i) \right) \right\} \mu_0^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma}) \\
= O_{b+i+j} \left(|u_1|(1+|u_1|)^{2(b+1)+i+j} (\|\boldsymbol{u}\|^{2a} + N^{\lfloor a/2 \rfloor}) \right).$$

When both of them are odd,

$$\frac{\int_{S_N^2} (\sigma_1^1 \sigma_1^2)^{b+1} \exp\left\{\sum_{i=1}^2 \left(\left(u_1 \sigma_1^i + \Lambda_1(\sigma_1^i)^2\right) + H_{\leqslant 2}(\boldsymbol{\sigma}_{-1}^i) + W(\boldsymbol{\sigma}^i, \boldsymbol{\sigma}^i) \right) \right\} C_{i,j}(\sigma_1^1)^i (\sigma_1^2)^j \, \mu_0^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma})}{\int_{S_N^2} \exp\left\{\sum_{i=1}^2 \left(\left(u_1 \sigma_1^i + \Lambda_1(\sigma_1^i)^2\right) + H_{\leqslant 2}(\boldsymbol{\sigma}_{-1}^i) + W(\boldsymbol{\sigma}^i, \boldsymbol{\sigma}^i) \right) \right\} \mu_0^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma})} = O_{b+i+j} \left(|u_1|^2 (1 + |u_1|)^{2(b+1)+i+j} (\|\boldsymbol{u}\|^{2a} + N^{\lfloor a/2 \rfloor}) \right).$$

Otherwise, when b + 1 + i and b + 1 + j are both even,

$$\frac{\left|\int_{S_N^2} (\sigma_1^1 \sigma_1^2)^{b+1} \exp\left\{\sum_{i=1}^2 \left(\left(u_1 \sigma_1^i + \Lambda_1(\sigma_1^i)^2\right) + H_{\leq 2}(\boldsymbol{\sigma}_{-1}^i) + W(\boldsymbol{\sigma}^i, \boldsymbol{\sigma}^i)\right) \right\} C_{i,j}(\sigma_1^1)^i (\sigma_1^2)^j \, \mu_0^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma})}{\int_{S_N^2} \exp\left\{\sum_{i=1}^2 \left(\left(u_1 \sigma_1^i + \Lambda_1(\sigma_1^i)^2\right) + H_{\leq 2}(\boldsymbol{\sigma}_{-1}^i) + W(\boldsymbol{\sigma}^i, \boldsymbol{\sigma}^i)\right) \right\} \mu_0^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma})} \\ \leqslant O_{b+i+j} \left((1+|u_1|)^{2(b+1)+i+j} (\|\boldsymbol{u}\|^{2a} + N^{\lfloor a/2 \rfloor}) \right).$$

Therefore, under the assumption $\|\boldsymbol{u}\| \leq N^{c_0}$, for $\ell \leq L$,

$$\begin{split} \frac{|\hat{T}_{\ell}(a,b)|}{\int_{S_{N}^{2}} \exp\left\{\sum_{i=1}^{2}\left(\left(u_{1}\sigma_{1}^{i}+\Lambda_{1}(\sigma_{1}^{i})^{2}\right)+H_{\leq 2}(\boldsymbol{\sigma}_{-1}^{i})+W(\boldsymbol{\sigma}^{i},\boldsymbol{\sigma}^{i})\right)\right\}\mu_{0}^{\otimes 2}(\mathsf{d}\boldsymbol{\sigma})} \\ & \leq \left(N^{\lfloor a/2 \rfloor}+\|\boldsymbol{u}\|^{2a}\right)\cdot\left[O_{L}\left(|u_{1}|^{2}\sum_{i,j\leqslant L}N^{\ell-b-a-(i+j)/2}\right)+\sum_{\substack{i,j\leqslant L\\i,j=b+1\mod 2}}O_{L}\left(N^{\ell-b-a-(i+j)/2}\right)\right.\\ & \left.+\sum_{\substack{i,j\leqslant L\\i\neq j\mod 2}}O_{L}\left(|u_{1}|N^{\ell-b-a-(i+j)/2}\right)\right] \\ & = O_{L}(N^{-2}+N^{-3/2}|u_{1}|+N^{-1}|u_{1}|^{2})=O_{L}(N^{-1}|u_{1}|^{2}+N^{-2}), \end{split}$$

where in the last step we used the fact that $\ell \ge 1$, and $a \ge 3\ell$ when b = 0, or $a + b \ge 3\ell + 1$, otherwise.

Take L = 4, and combining the terms in Eq. (7.105), we obtain

$$Q(A_2(N^{-1/2+c})) \leq O\left((N^{-2} + N^{-1}|u_1|)(Z_{\leq 2})^2 e^{N\xi_{\geq 3}(1)}\right),$$

and therefore, using Eq. (7.104)

$$\mathbb{E}_{\boldsymbol{g}_{\geq 4},\boldsymbol{g}_{3-}}\left[\left\{\int_{T(\delta)}\sigma_1\left(e^{H(\boldsymbol{\sigma})} - \mathbb{E}_{\boldsymbol{g}_{\geq 4},\boldsymbol{g}_{3-}}e^{H(\boldsymbol{\sigma})}\right)\,\mu_0(\mathsf{d}\boldsymbol{\sigma})\right\}^2\right] = O\left((N^{-2} + N^{-1}|u_1|^2)(Z_{\leq 2})^2 e^{N\xi_{\geq 3}(1)}\right).$$

Thus, with probability at least $1 - N^{-c}$, we have

$$\left| \int_{T(\delta)} \sigma_1 \left(e^{H(\boldsymbol{\sigma})} - \mathbb{E}_{\boldsymbol{g}_{\geq 4}, \boldsymbol{g}_{3-}} e^{H(\boldsymbol{\sigma})} \right) \, \mu_0(\mathsf{d}\boldsymbol{\sigma}) \right| \leq Z_{\leq 2} N^c (N^{-1} + N^{-1/2} |u_1|).$$

This yields the desired claim upon using Lemma 7.7.

We note that (7.40) in Lemma 7.8 immediately gives the following high probability bound on the magnetization.

Lemma 7.13. For any $\varepsilon, C > 0$, there exists $c_0 > 0$ such that, for $||\mathbf{u}|| \le N^{c_0}$, with probability at least $1 - N^{-C}$, we have

$$\left\|\int \boldsymbol{\sigma}\mu(\mathsf{d}\boldsymbol{\sigma})\right\|^2 \leqslant N^{\varepsilon}\,,\tag{7.106}$$

for N sufficiently large.

Proof. We work, as before, in the basis of eigenvectors of the quadratic part W_2 of the Hamiltonian. By (7.40), with $k = 4C/\varepsilon$, with probability at least $1 - N^{-2C}$,

$$\int_{T} \sigma_{i} e^{H(\boldsymbol{\sigma})} \mu_{0}(\mathsf{d}\boldsymbol{\sigma}) \leq N^{\varepsilon/4} \|\boldsymbol{u}\|^{Ck} (|u_{i}| + CN^{-1/2}) \mathbb{E}_{\geq 3} \int e^{H(\boldsymbol{\sigma})} \mu_{0}(\mathsf{d}\boldsymbol{\sigma}).$$

By (7.38) with L = 4C and the union bound over $i \in [N]$, we then have, with probability at least $1 - \varepsilon^{-8C} N^{-C}$, for all $i \in [N]$,

$$\frac{1}{Z} \int \sigma_i e^{H(\boldsymbol{\sigma})} \mu_0(\mathbf{d}\boldsymbol{\sigma}) \leqslant N^{\varepsilon/2} \|\boldsymbol{u}\|^{Ck} (|u_i| + CN^{-1/2}).$$

Assuming that c_0 is chosen so that $c_0 L < \varepsilon/4$, we then obtain (7.106).

Lemma 7.2 now follows.

Proof of Lemma 7.2. Let $\hat{m} = m + \Delta(m)$. From Lemma 7.1 we have, with probability at least $1 - N^{-c}$

$$\|\langle \boldsymbol{\sigma} \rangle - \hat{\boldsymbol{m}} \|^2 \leq O \left(N^{-c} + N^{-c} \| \boldsymbol{u} \|^2 \right).$$

Therefore, using Lemma 7.13 and the trivial bound $\|\langle \sigma \rangle\| \leq \sqrt{N}$, we can pick $\varepsilon > 0$ sufficiently small and k sufficiently large such that, upon adjusting the constant c,

$$\mathbb{E}[\|\langle \boldsymbol{\sigma} \rangle - \hat{\boldsymbol{m}}\|^{\alpha}] = O(N^{-c\alpha} + N^{-c\alpha} \|\boldsymbol{u}\|^{\alpha}) + N^{-C} + O(N^{\varepsilon\alpha} \cdot N^{-c})$$
$$= O(N^{-c/2}).$$

7.2 Integrating over bands

Using the results in the previous section, we will complete the proof of Proposition 4.6. We will assume the setup of Proposition 4.6. We sample $\boldsymbol{x} \sim \mu_{\text{unif}}, \boldsymbol{y} = t\boldsymbol{x} + \boldsymbol{B}_t$, and $\tilde{H}(\cdot) \sim \mu_{\text{null}}$ (the Gaussian process with covariance $\mathbb{E} \tilde{H}(\boldsymbol{\sigma}^1)\tilde{H}(\boldsymbol{\sigma}^2) = N\xi(\langle \boldsymbol{\sigma}^1, \boldsymbol{\sigma}^2 \rangle)$) with $\boldsymbol{x}, \boldsymbol{B}, \tilde{H}$ independent. We define the tilted disorder $H(\boldsymbol{\sigma}) = \tilde{H}(\boldsymbol{\sigma}) + \langle \boldsymbol{y}, \boldsymbol{\sigma} \rangle + N\xi(\langle \boldsymbol{x}, \boldsymbol{\sigma} \rangle_N)$, so that $(\boldsymbol{x}, H, \boldsymbol{y}) \sim \mathbb{P}$ are distributed according to the planted model, cf. Eq. (3.2). (For simplicity of notation, we drop the dependence on t in the notation of H, \boldsymbol{y} in this section compared to the notation in Section 4.) In this section, we will estimate the mean of the Gibbs measure given by H.

Recall that

$$\mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = N\xi(\langle \boldsymbol{x}, \boldsymbol{m}
angle_N) + \widetilde{H}(\boldsymbol{m}) + \langle \boldsymbol{y}, \boldsymbol{m}
angle + rac{N}{2} heta(\|\boldsymbol{m}\|_N^2) + rac{N}{2}\log(1 - \|\boldsymbol{m}\|_N^2),$$

where $\theta(s) = \xi(1) - \xi(s) - (1 - s)\xi'(s)$.

Let $\boldsymbol{m} \in \mathbb{R}^N$ and $q = \|\boldsymbol{m}\|_N^2$. The following lemma follows from standard calculations.

Lemma 7.14. The distribution of $\widetilde{H}(\boldsymbol{\sigma})$ given $\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = 0$ is a Gaussian process with

$$N^{-1}\mathbb{E}[\tilde{H}(\boldsymbol{\sigma}) \mid \nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = 0, \boldsymbol{y}, \boldsymbol{x}] = \frac{\xi'(\langle \boldsymbol{m}, \boldsymbol{\sigma} \rangle_N) \langle \boldsymbol{z}, \boldsymbol{\sigma} \rangle_N}{\xi'(q)} - \frac{\xi''(q) \langle \boldsymbol{m}, \boldsymbol{z} \rangle_N}{\xi'(q) \zeta(q)} \xi'(\langle \boldsymbol{m}, \boldsymbol{\sigma} \rangle_N) \langle \boldsymbol{m}, \boldsymbol{\sigma} \rangle_N,$$
(7.107)

with $\zeta(q) = \xi'(q) + q\xi''(q)$ and $\boldsymbol{z} = -\boldsymbol{y} - \xi'(\langle \boldsymbol{x}, \boldsymbol{m} \rangle_N)\boldsymbol{x} + (1-q)\xi''(q)\boldsymbol{m} + \frac{\boldsymbol{m}}{1-q}$, and covariance

$$N^{-1} \operatorname{Cov}[\tilde{H}(\boldsymbol{\sigma}^{1}), \tilde{H}(\boldsymbol{\sigma}^{2}) | \nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = 0, \boldsymbol{y}, \boldsymbol{x}]$$

$$= \xi(\langle \boldsymbol{\sigma}^{1}, \boldsymbol{\sigma}^{2} \rangle_{N}) - \frac{\xi'(\langle \boldsymbol{m}, \boldsymbol{\sigma}^{1} \rangle_{N})\xi'(\langle \boldsymbol{m}, \boldsymbol{\sigma}^{2} \rangle_{N})}{\xi'(q)} \langle \boldsymbol{\sigma}^{1}, \boldsymbol{\sigma}^{2} \rangle_{N}$$

$$+ \frac{\xi''(q)\xi'(\langle \boldsymbol{m}, \boldsymbol{\sigma}^{1} \rangle_{N})\xi'(\langle \boldsymbol{m}, \boldsymbol{\sigma}^{2} \rangle_{N})\langle \boldsymbol{m}, \boldsymbol{\sigma}^{1} \rangle_{N} \langle \boldsymbol{m}, \boldsymbol{\sigma}^{2} \rangle_{N}}{\xi'(q)}.$$
(7.108)

Let $\sigma^{\perp} = \mathsf{P}_{\{x,m\}^{\perp}}(\sigma)$ be the projection of σ on $\{x,m\}^{\perp}$, and similarly define y^{\perp} , z^{\perp} . Define the band

$$D_N(a,b) := \left\{ \boldsymbol{\sigma} \in S_N : \langle \boldsymbol{\sigma}, \boldsymbol{m} \rangle_N = aq \text{ and } \langle \boldsymbol{\sigma}, \boldsymbol{x} \rangle_N = b \right\},$$
(7.109)

and let $r(a, b) = \|\boldsymbol{\sigma} - \boldsymbol{\sigma}^{\perp}\|_N^2$ for $\boldsymbol{\sigma} \in D_N(a, b)$. Throughout the rest of the section, we will condition on the event $\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = 0$, and on $\boldsymbol{y} - \boldsymbol{y}^{\perp}$ and \boldsymbol{x} . Conditional on $\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = 0, \boldsymbol{y} - \boldsymbol{y}^{\perp}, \boldsymbol{x}$, we can write

$$N^{-1}H(\boldsymbol{\sigma}) = \xi(b) + \frac{\xi'(aq)\langle \boldsymbol{z}, \boldsymbol{\sigma} \rangle_N}{\xi'(q)} - \frac{\xi''(q)\xi'(aq)aq\langle \boldsymbol{m}, \boldsymbol{z} \rangle_N}{\xi'(q)\zeta(q)} + N^{-1}\widehat{H}(\boldsymbol{\sigma}^{\perp}) + \langle \boldsymbol{y}^{\perp}, \boldsymbol{\sigma}^{\perp} \rangle_N + \langle \boldsymbol{y} - \boldsymbol{y}^{\perp}, \boldsymbol{\sigma} - \boldsymbol{\sigma}^{\perp} \rangle_N,$$

where \hat{H} is a centered Gaussian process with covariance

$$N^{-1}\operatorname{Cov}(\widehat{H}(\boldsymbol{\sigma}^{\perp,1}),\widehat{H}(\boldsymbol{\sigma}^{\perp,2})) = \xi\left(r(a,b) + \langle \boldsymbol{\sigma}^{\perp,1}, \boldsymbol{\sigma}^{\perp,2} \rangle_N\right) - \frac{\xi'(aq)^2}{\xi'(q)} \langle \boldsymbol{\sigma}^{\perp,1}, \boldsymbol{\sigma}^{\perp,2} \rangle_N - \frac{\xi'(aq)^2 r(a,b)}{\xi'(q)} + \frac{\xi'(aq)^2 \xi''(q)(aq)^2}{\zeta(q)\xi'(q)}.$$

Let $\tilde{\boldsymbol{\sigma}} = \boldsymbol{\sigma}^{\perp} / \| \boldsymbol{\sigma}^{\perp} \|_N$. We can then write

$$\begin{split} &\int_{D_{N}(a,b)} e^{H(\boldsymbol{\sigma})} \mu_{0}^{a,b}(\mathrm{d}\boldsymbol{\sigma}) \\ &= \exp\left(N\left[\xi(b) + \frac{\xi'(aq)\langle \boldsymbol{z} + \boldsymbol{y}, \boldsymbol{\sigma} \rangle_{N}}{\xi'(q)} - \frac{\xi''(q)\xi'(aq)aq\langle \boldsymbol{m}, \boldsymbol{z} \rangle_{N}}{\xi'(q)\zeta(q)} + \left(1 - \frac{\xi'(aq)}{\xi'(q)}\right)\langle \boldsymbol{y} - \boldsymbol{y}^{\perp}, \boldsymbol{\sigma} - \boldsymbol{\sigma}^{\perp} \rangle_{N}\right]\right) \\ &\int_{S_{N-2}} \exp\left(N\left(1 - \frac{\xi'(aq)}{\xi'(q)}\right)(1 - r(a,b))^{1/2}\langle \boldsymbol{y}^{\perp}, \tilde{\boldsymbol{\sigma}} \rangle_{N} + \underline{\tilde{H}}(\tilde{\boldsymbol{\sigma}}) + \frac{N-3}{2}\log(1 - r(a,b))\right)\mu_{0}(\mathrm{d}\tilde{\boldsymbol{\sigma}}) \\ &= \exp\left(N\Gamma_{N}(\boldsymbol{y}, \boldsymbol{m}; a, b) + \frac{N-3}{2}\log(1 - r(a,b))\right)\int_{S_{N-2}} e^{N^{1/2}g_{a,b} + \underline{H}(\tilde{\boldsymbol{\sigma}})}\mu_{0}(\mathrm{d}\tilde{\boldsymbol{\sigma}}), \end{split}$$
(7.110)

where $\mu_0^{a,b}$ is the measure induced on $D_N(a,b)$ by μ_0 , we defined Γ_N via

$$\Gamma_{N}(\boldsymbol{y},\boldsymbol{m};a,b) := \xi(b) + \frac{\xi'(aq)\langle \boldsymbol{z} + \boldsymbol{y}, \boldsymbol{\sigma} \rangle_{N}}{\xi'(q)} - \frac{\xi''(q)\xi'(aq)aq\langle \boldsymbol{m}, \boldsymbol{z} \rangle_{N}}{\xi'(q)\zeta(q)} + \left(1 - \frac{\xi'(aq)}{\xi'(q)}\right)\langle \boldsymbol{y} - \boldsymbol{y}^{\perp}, \boldsymbol{\sigma} - \boldsymbol{\sigma}^{\perp} \rangle_{N},$$
(7.111)

and <u>*H*</u> is a Hamiltonian on $\tilde{\sigma}$ with mixture $\tilde{\xi}(q) = \sum_{k \ge 1} \tilde{\xi}_k$ given by

$$\tilde{\xi}_1 = (1 - r(a, b)) \left(\xi'(r(a, b)) - \frac{\xi'(aq)^2}{\xi'(q)} + \left(1 - \frac{\xi'(aq)}{\xi'(q)}\right)^2 t \right) =: \tilde{\gamma}_1^2,$$
(7.112)

$$\tilde{\xi}_2 = \frac{1}{2} \xi''(r(a,b))(1-r(a,b))^2 =: \tilde{\gamma}_2^2,$$
(7.113)

$$\tilde{\xi}_p = \frac{1}{p!} \xi^{(p)}(r(a,b)) (1 - r(a,b))^p, \quad p \ge 3.$$
(7.114)

Finally, $g_{a,b}$ is a Gaussian independent of <u>H</u> with standard deviation $\tilde{\gamma}_0$ given by

$$\tilde{\gamma}_0^2 := \xi(r(a,b)) - \frac{\xi'(aq)^2 r(a,b)}{\xi'(q)} + \frac{\xi'(aq)^2 \xi''(q)(aq)^2}{\zeta(q)\xi'(q)}.$$
(7.115)

Note that

$$\tilde{\xi}_{\geq 2}(s) = \sum_{p \geq 2} \frac{1}{p!} \xi^{(p)}(r(a,b)) (1 - r(a,b))^p s^p$$

= $\xi(r(a,b) + (1 - r(a,b))s) - \xi(r(a,b)) - \xi'(r(a,b))(1 - r(a,b))s$

and therefore

$$\tilde{\xi}_{\geqslant 2}''(s) = (1 - r(a, b))^2 \xi''(r(a, b) + (1 - r(a, b))s)$$

$$\stackrel{(1.5)}{<} \frac{(1 - r(a, b))^2}{(1 - (r(a, b) + (1 - r(a, b))s))^2} = \frac{1}{(1 - s)^2}.$$
(7.116)

Integrating twice shows $\tilde{\xi}_{\geq 2}$ satisfies condition (7.9), and thus the results in Subsection 7.1 apply to $\tilde{\xi}_{\geq 2}$. Similarly, note that

$$\tilde{\xi}_{\geq 3}(1) = \xi(1) - \xi(r(a,b)) - \xi'(r(a,b))(1 - r(a,b)) - \frac{1}{2}\xi''(r(a,b))(1 - r(a,b))^2.$$
(7.117)

Following Subsection 7.1, we write the quadratic component of \underline{H} as $\langle \mathbf{A}^{(2)}, \tilde{\boldsymbol{\sigma}}^{\otimes 2} \rangle$ for $\mathbf{A}^{(2)} = \mathbf{A}^{(2)}(a, b)$ a GOE matrix scaled by $\tilde{\gamma}_2/\sqrt{2}$. Recall the definition of $G(\gamma) = G(\gamma; \mathbf{A}, \mathbf{u})$ in Eq. (7.10). We take \mathbf{u} to be the external field $\mathbf{u} = \tilde{\gamma}_1 \mathbf{g}$, and $\mathbf{A} = \mathbf{A}^{(2)}$. Note that \mathbf{u} and $\mathbf{A}^{(2)}$ depend on the parameters a, b. Let $\gamma_{a,b} = \arg \min_{z>z_*} G(z; \mathbf{A}^{(2)}, \mathbf{u}), z_* := \lambda_{\max}(\mathbf{A}^{(2)})$. From Lemma 7.8 Eqs. (7.38) and (7.39) and Lemma 7.3, when

$$\tilde{\gamma}_1^2 = (1 - r(a, b)) \left(\xi'(r(a, b)) - \frac{\xi'(aq)^2}{\xi'(q)} + \left(1 - \frac{\xi'(aq)}{\xi'(q)}\right)^2 t \right) \le N^{c_0 - 1},$$

we have (with probability at least $1 - N^{-c}$, conditional on $\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = 0, \boldsymbol{y} - \boldsymbol{y}^{\perp}, \boldsymbol{x}$) that

$$\begin{split} \int_{D_{N}(a,b)} e^{H(\boldsymbol{\sigma})} \mu_{0}^{a,b}(\mathbf{d}\boldsymbol{\sigma}) \\ &= (1+O(N^{-c}))(2e)^{-(N-2)/2} \sqrt{\frac{2}{NG''(\gamma_{a,b}; \boldsymbol{A}^{(2)}, \boldsymbol{u})}} \\ &\quad \exp\left(N \bigg[N^{-1/2}g_{a,b} + \Gamma_{N}(\boldsymbol{y}, \boldsymbol{m}; a, b) + \frac{N-3}{2N} \log(1-r(a, b)) + \min_{z>z_{*}} G(z; \boldsymbol{A}^{(2)}, \boldsymbol{u}) \right. \\ &\quad + \frac{1}{2} \left(\xi(1) - \xi(r(a, b)) - \xi'(r(a, b))(1-r(a, b)) - \frac{1}{2} \xi''(r(a, b))(1-r(a, b))^{2} \right) \bigg] \bigg), \end{split}$$
(7.118)

where we have simplified using Eq. (7.117). By independence of $\boldsymbol{u}, \tilde{W}^{(2)}$, and the fact that $\tilde{W}^{(2)}$ is a GOE matrix scaled by $\tilde{\gamma}_2/\sqrt{2}$, the following holds with probability at least $1 - \exp(-N^c)$ provided $z > \tilde{\gamma}_2\sqrt{2} + \delta$ for some constant $\delta > 0$

$$G(z; \mathbf{A}^{(2)}, \mathbf{u}) = G_{a,b}(z) + O(1/N),$$
(7.119)

where

$$G_{a,b}(z) := z - \frac{1}{2} \left(\psi(z\sqrt{2}/\tilde{\gamma}_2) + \log(\tilde{\gamma}_2/\sqrt{2}) \right) + \frac{1}{4} \left(\tilde{\gamma}_1^2 + (1 - r(a, b)) \left(1 - \frac{\xi'(aq)}{\xi'(q)} \right)^2 t \right) \phi(z\sqrt{2}/\tilde{\gamma}_2),$$
(7.120)

and, for x > 2,

$$\phi(x) = \frac{1}{2}(x - \sqrt{x^2 - 4}), \qquad \psi(x) = \frac{1}{2}((x - \sqrt{x^2 - 4})/2)^2 - \log((x - \sqrt{x^2 - 4})/2).$$

Note that $\phi(x) = \int (x-u)^{-1} \mu_{sc}(du)$ and $\psi(x) = \int \log(x-u) \mu_{sc}(du)$ where μ_{sc} is the semicircular law. Moreover, $\psi'(x) = \phi(x)$.

Thus,

$$\int_{D_N(a,b)} e^{H(\boldsymbol{\sigma})} \mu_0^{a,b}(\mathsf{d}\boldsymbol{\sigma}) = \sqrt{\frac{2}{NG''(\gamma_{a,b}; \boldsymbol{A}^{(2)}, \boldsymbol{u})}} \exp\left(NE(a,b) + N^{1/2}g_{a,b} + O(1)\right),$$
(7.121)

where we define

$$E(a,b) := -\frac{N-2}{2N}\ln(2e) + \frac{N-3}{2N}\log(1-r(a,b)) + \min_{z>\tilde{\gamma}_2\sqrt{2}}G_{a,b}(z) + \Gamma_N(\boldsymbol{y},\boldsymbol{m};a,b) + \frac{1}{2}\left(\xi(1) - \xi(r(a,b)) - \xi'(r(a,b))(1-r(a,b)) - \frac{1}{2}\xi''(r(a,b))(1-r(a,b))^2\right).$$
 (7.122)

Let $b_* = \langle \boldsymbol{x}, \boldsymbol{m} \rangle_N$. Note that $r(1, b_*) = q$. Furthermore, we have

$$r(a,b) = a^2 q + \frac{(b - a\langle \boldsymbol{x}, \boldsymbol{m} \rangle_N)^2}{1 - \langle \boldsymbol{x}, \boldsymbol{m} \rangle_N^2/q}.$$
(7.123)

We will next verify several properties of E(a, b), starting with the observation that $(a, b) = (1, b_*)$ is a stationary point of E.

Lemma 7.15. We have $\nabla E(a,b)|_{(a,b)=(1,b_*)} = 0$. (Here ∇ denotes gradient with respect to (a,b).)

Proof. We first compute $\nabla \min_z G_{a,b}(z)$. Let $z_*(a,b) = \arg \min_z G_{a,b}(z)$ and $z_* = \arg \min_z G_{1,b_*}(z)$ so

$$\partial_z G_{a,b}(z)|_{z_*(a,b)} = 0 \Leftrightarrow 1 - \frac{1}{\sqrt{2}\tilde{\gamma}_2}\phi(z\sqrt{2}/\tilde{\gamma}_2) = 0.$$
(7.124)

For $\alpha \in \{a, b\}$,

$$\partial_{\alpha} z_{*}(a,b) = (\partial_{z}^{2} G_{a,b}(z))^{-1}|_{(a,b,z_{*}(a,b))} \partial_{\alpha} \partial_{z} G_{a,b}(z)|_{(a,b,z_{*}(a,b))}.$$
(7.125)

A quick calculation shows that $z_* = 1/2 + \tilde{\gamma}_2^2$ when $(a, b) = (1, b_*)$, and for $\alpha \in \{a, b\}$,

$$\partial_{\alpha} \min_{z} G_{a,b}(z)|_{(a,b)} = \partial_{\alpha} G_{a,b}(z_{*}(a,b))|_{(a,b)}.$$

Also note that

$$\nabla \left(\tilde{\gamma}_1^2 + (1 - r(a, b)) \left(1 - \frac{\xi'(aq)}{\xi'(q)} \right)^2 t \right) \bigg|_{(1, b_*)} = 0; \quad \left(\tilde{\gamma}_1^2 + (1 - r(a, b)) \left(1 - \frac{\xi'(aq)}{\xi'(q)} \right)^2 t \right) \bigg|_{(1, b_*)} = 0.$$
(7.126)

From the definition of G and the stationary condition (7.124), we obtain that

$$\nabla \min_{z} G_{a,b}(z)|_{(1,b_{*})} = \frac{1}{2} \left(-\tilde{\gamma}_{2}^{-1} + \sqrt{2}\tilde{\gamma}_{2}^{-2}z_{*}\phi(z_{*}\sqrt{2}/\tilde{\gamma}_{2}) \right) \nabla \tilde{\gamma}_{2} = \tilde{\gamma}_{2}\nabla \tilde{\gamma}_{2} = \frac{1}{2}\nabla(\tilde{\gamma}_{2}^{2}).$$
(7.127)

Furthermore,

$$\nabla \tilde{\gamma}_2^2 = -\xi''(r(a,b))(1-r(a,b))\nabla r(a,b) + \frac{1}{2}\xi'''(r(a,b))(1-r(a,b))^2\nabla r(a,b).$$

We have

$$\nabla\left(\frac{1}{2}\log(1-r(a,b)) + \frac{1}{2}(\xi(1)-\xi(r(a,b))-\xi'(r(a,b))(1-r(a,b)) - \frac{1}{2}\xi''(r(a,b))(1-r(a,b))^2)\right)$$

$$= \frac{-1}{2(1-r(a,b))}\nabla r(a,b) - \frac{1}{4}\xi'''(r(a,b))(1-r(a,b))^2\nabla r(a,b),$$
(7.128)

and furthermore $\partial_a r(a,b)|_{(1,b_*)} = 2q$, $\partial_b r(a,b)|_{(1,b_*)} = 0$ and $r(1,b_*) = q$. Recall

$$\langle \boldsymbol{z} + \boldsymbol{y}, \boldsymbol{\sigma} \rangle_N = -\xi'(\langle \boldsymbol{x}, \boldsymbol{m} \rangle_N)b + aq\left((1-q)\xi''(q) + \frac{1}{1-q}\right).$$

Moreover,

$$\langle \boldsymbol{y} - \boldsymbol{y}^{\perp}, \boldsymbol{\sigma} - \boldsymbol{\sigma}^{\perp}
angle_N = a \langle \boldsymbol{y}, \boldsymbol{m}
angle_N + rac{b - a \langle \boldsymbol{x}, \boldsymbol{m}
angle_N}{1 - \langle \boldsymbol{x}, \boldsymbol{m}
angle_N^2 / q} \left(\langle \boldsymbol{y}, \boldsymbol{x}
angle_N - \langle \boldsymbol{x}, \boldsymbol{m}
angle_N \langle \boldsymbol{y}, \boldsymbol{m}
angle_N / q
ight).$$

Hence,

$$\begin{split} \frac{\xi'(aq)\langle \boldsymbol{y} + \boldsymbol{z}, \boldsymbol{\sigma} \rangle_{N}}{\xi'(q)} &- \frac{\xi''(q)\xi'(aq)aq\langle \boldsymbol{m}, \boldsymbol{z} \rangle_{N}}{\xi'(q)\zeta(q)} + \left(1 - \frac{\xi'(aq)}{\xi'(q)}\right)\langle \boldsymbol{y} - \boldsymbol{y}^{\perp}, \boldsymbol{\sigma} - \boldsymbol{\sigma}^{\perp} \rangle_{N} \\ &= \frac{\xi'(aq)(-\xi'(\langle \boldsymbol{x}, \boldsymbol{m} \rangle_{N})b + aq((1-q)\xi''(q) + \frac{1}{1-q}))}{\xi'(q)} - \frac{\xi''(q)\xi'(aq)aq\langle \boldsymbol{m}, \boldsymbol{z} \rangle_{N}}{\xi'(q)\zeta(q)} \\ &+ \left(1 - \frac{\xi'(aq)}{\xi'(q)}\right)\left(a\langle \boldsymbol{y}, \boldsymbol{m} \rangle_{N} + \frac{b - a\langle \boldsymbol{x}, \boldsymbol{m} \rangle_{N}}{1 - \langle \boldsymbol{x}, \boldsymbol{m} \rangle_{N}^{2}/q}\left(\langle \boldsymbol{y}, \boldsymbol{x} \rangle_{N} - \langle \boldsymbol{x}, \boldsymbol{m} \rangle_{N}\langle \boldsymbol{y}, \boldsymbol{m} \rangle_{N}/q\right)\right) \\ &= \frac{\xi'(aq)(-\xi'(\langle \boldsymbol{x}, \boldsymbol{m} \rangle_{N})b + aq((1-q)\xi''(q) + \frac{1}{1-q}))}{\xi'(q)} \\ &+ \left(1 - \frac{\xi'(aq)}{\xi'(q)}\right)\frac{b - a\langle \boldsymbol{x}, \boldsymbol{m} \rangle_{N}}{1 - \langle \boldsymbol{x}, \boldsymbol{m} \rangle_{N}^{2}/q}\left(\langle \boldsymbol{y}, \boldsymbol{x} \rangle_{N} - \langle \boldsymbol{x}, \boldsymbol{m} \rangle_{N}\langle \boldsymbol{y}, \boldsymbol{m} \rangle_{N}/q\right) \\ &+ \frac{\xi''(q)\xi'(aq)aq\left(\xi'(\langle \boldsymbol{x}, \boldsymbol{m} \rangle_{N})\langle \boldsymbol{x}, \boldsymbol{m} \rangle_{N} - q\xi''(q)(1-q) - \frac{q}{1-q}\right)}{\xi'(q)} \\ &+ \langle \boldsymbol{y}, \boldsymbol{m} \rangle_{N}\left(\frac{\xi''(q)\xi'(aq)aq}{\xi'(q)\zeta(q)} + a\left(1 - \frac{\xi'(aq)}{\xi'(q)}\right)\right). \end{split}$$

Note that

$$\partial_a \left(\left(1 - \frac{\xi'(aq)}{\xi'(q)} \right) \frac{b - a \langle \boldsymbol{x}, \boldsymbol{m} \rangle_N}{1 - \langle \boldsymbol{x}, \boldsymbol{m} \rangle_N^2 / q} \right) \Big|_{(1,b_*)} = \partial_b \left(\left(1 - \frac{\xi'(aq)}{\xi'(q)} \right) \frac{b - a \langle \boldsymbol{x}, \boldsymbol{m} \rangle_N}{1 - \langle \boldsymbol{x}, \boldsymbol{m} \rangle_N^2 / q} \right) \Big|_{(1,b_*)} = 0,$$

and $\partial_a(\xi'(aq)aq)|_{(1,q)}=q\zeta(q)$ so

$$\partial_a \left(\frac{\xi''(q)\xi'(aq)aq}{\xi'(q)\zeta(q)} + a \left(1 - \frac{\xi'(aq)}{\xi'(q)} \right) \right) |_{(1,b_*)} = \partial_b \left(\frac{\xi''(q)\xi'(aq)aq}{\xi'(q)\zeta(q)} + a \left(1 - \frac{\xi'(aq)}{\xi'(q)} \right) \right) |_{(1,b_*)} = 0.$$

Thus, we can compute

$$\partial_{a} \left(\frac{\xi'(aq)\langle \boldsymbol{y} + \boldsymbol{z}, \boldsymbol{\sigma} \rangle_{N}}{\xi'(q)} - \frac{\xi''(q)\xi'(aq)aq\langle \boldsymbol{m}, \boldsymbol{z} \rangle_{N}}{\xi'(q)\zeta(q)} + \left(1 - \frac{\xi'(aq)}{\xi'(q)}\right) \langle \boldsymbol{y} - \boldsymbol{y}^{\perp}, \boldsymbol{\sigma} - \boldsymbol{\sigma}^{\perp} \rangle_{N} \right) |_{(1,b_{*})}$$

$$= q(1-q)\xi''(q) + \frac{q}{1-q}.$$
(7.129)

Similarly,

$$\partial_{b} \left(\frac{\xi'(aq) \langle \boldsymbol{y} + \boldsymbol{z}, \boldsymbol{\sigma} \rangle_{N}}{\xi'(q)} - \frac{\xi''(q) \xi'(aq) aq \langle \boldsymbol{m}, \boldsymbol{z} \rangle_{N}}{\xi'(q) \zeta(q)} + \left(1 - \frac{\xi'(aq)}{\xi'(q)} \right) \langle \boldsymbol{y} - \boldsymbol{y}^{\perp}, \boldsymbol{\sigma} - \boldsymbol{\sigma}^{\perp} \rangle_{N} \right) |_{(1,b_{*})}$$

= $-\xi'(\langle \boldsymbol{x}, \boldsymbol{m} \rangle_{N}).$ (7.130)

Combining (7.127), (7.128), (7.129), (7.130), we obtain the desired claim that $\nabla E(a, b)|_{(a,b)=(1,b_*)} = 0.$

Lemma 7.16. We have

$$\mathbb{E}\left[\int_{D_N(a,b)} e^{H(\boldsymbol{\sigma})} \mu_0^{a,b}(\mathsf{d}\boldsymbol{\sigma}) \middle| \nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = \boldsymbol{0}, \boldsymbol{x}, \boldsymbol{y} - \boldsymbol{y}^{\perp}, g_{a,b}\right] = \exp\left\{N\hat{E}(a,b) - \log(2e) + \sqrt{N}g_{a,b}\right\},\tag{7.131}$$

where

$$\hat{E}(a,b) := \frac{1}{N}\ln(2e) + \frac{N-3}{2N}\log(1-r(a,b)) + \left(\frac{1}{2}\tilde{\gamma}_2^2 + \frac{1}{2}\tilde{\gamma}_1^2\right) + \Gamma_N(\boldsymbol{y},\boldsymbol{m};a,b) \\ + \frac{1}{2}\left(\xi(1) - \xi(r(a,b)) - \xi'(r(a,b))(1-r(a,b)) - \frac{1}{2}\xi''(r(a,b))(1-r(a,b))^2\right).$$
(7.132)

Furthermore, E(a,b) is uniformly upper bounded by $\hat{E}(a,b)$, $E(1,b_*) = \hat{E}(1,b_*)$, and $\nabla E(1,b_*) = \nabla \hat{E}(1,b_*) = 0$.

Proof. Eq (7.131) follows from a direct calculation. For the last claim, let $\tilde{\gamma}_2 = \tilde{\gamma}_2(a, b)$ and $\tilde{\gamma}_1 = \tilde{\gamma}_1(a, b)$. Given a quadratic Hamiltonian $H_{\leq 2}(\sigma) = \tilde{\gamma}_2 \langle \sigma, A\sigma \rangle / \sqrt{2} + \tilde{\gamma}_1 \langle g, \sigma \rangle$ where A is a GOE matrix and $g \sim \mathcal{N}(0, I_N)$,

$$\mathbb{E}\left[\int e^{H_{\leq 2}(\boldsymbol{\sigma})}\mu_0(\mathsf{d}\boldsymbol{\sigma})\right] = e^{N\tilde{\gamma}_2^2/2 + N\tilde{\gamma}_1^2/2}.$$

On the other hand, by Lemma 7.3, we have, for $\tilde{\gamma}_1$ sufficiently small, with probability at least $1 - e^{-cN}$,

$$\int e^{H_{\leq 2}(\boldsymbol{\sigma})} \mu_0(\mathsf{d}\boldsymbol{\sigma}) \ge \exp\Big\{(1-o(1))N\Big(\min_{z>\tilde{\gamma}_2\sqrt{2}} G_{a,b}(z) - \frac{1}{2}\log(2e)\Big)\Big\}.$$

Since this holds for all N, Markov inequality implies

$$\min_{z > \tilde{\gamma}_2 \sqrt{2}} G_{a,b}(z) - \frac{1}{2} \log(2e) \leq \frac{1}{2} \tilde{\gamma}_2^2 + \frac{1}{2} \tilde{\gamma}_1^2$$

The last claim follows immediately upon this observation.

When $\langle \boldsymbol{x}, \boldsymbol{m} \rangle_N = q$, $\langle \boldsymbol{y}, \boldsymbol{m} \rangle_N = t$, $\langle \boldsymbol{y}, \boldsymbol{x} \rangle_N = t$, $\|\boldsymbol{y}\|^2 = t + t^2$, under the constraint $\xi'(q) + t = \frac{q}{1-q}$, we can simplify $\hat{E}(a, b)$ as

$$\tilde{E}(a,b) := \frac{1}{N}\ln(2e) + \frac{N-3}{2N}\log(1-r(a,b)) + \frac{1}{2}\tilde{\gamma}_1^2 + \xi(b) - b\xi'(aq) + \frac{\xi'(aq)aq}{(1-q)\xi'(q)} + at\left(1 - \frac{\xi'(aq)}{\xi'(q)}\right) + \frac{1}{2}\left(\xi(1) - \xi(r(a,b)) - \xi'(r(a,b))(1-r(a,b))\right).$$
(7.133)

Indeed, under these values and constraints,

$$\begin{split} \Gamma_{N}(\boldsymbol{y},\boldsymbol{m};a,b) &= \xi(b) + \frac{\xi'(aq)\langle \boldsymbol{y} + \boldsymbol{z}, \boldsymbol{\sigma} \rangle}{\xi'(q)} - \frac{\xi''(q)\xi'(aq)aq\langle \boldsymbol{m}, \boldsymbol{z} \rangle}{\xi'(q)\theta(q)} + \left(1 - \frac{\xi'(aq)}{\xi'(q)}\right) \langle \boldsymbol{y} - \boldsymbol{y}^{\perp}, \boldsymbol{\sigma} - \boldsymbol{\sigma}^{\perp} \rangle \\ &= \xi(b) - \frac{\xi''(q)\xi'(aq)aq(-t - \xi'(q)q + \frac{q}{1-q} + q(1-q)\xi''(q))}{\xi'(q)\theta(q)} \\ &+ \frac{\xi'(aq)(-\xi'(q)b + aq((1-q)\xi''(q) + \frac{1}{1-q}))}{\xi'(q)} + t\left(1 - \frac{\xi'(aq)}{\xi'(q)}\right) \\ &= \xi(b) - b\xi'(aq) + \frac{\xi'(aq)aq}{(1-q)\xi'(q)} + at\left(1 - \frac{\xi'(aq)}{\xi'(q)}\right). \end{split}$$

Furthermore, in this case, $b_* = q$.

Lemma 7.17. For $\varepsilon > 0$ sufficiently small, there is $\eta > 0$ such that for all (a, b) satisfying $|aq-q|+|b-q| \leq \varepsilon$, we have $\nabla^2 \tilde{E}(a, b) \leq -\eta I_2$.

Proof. We have

$$\hat{\partial}_b^2 \tilde{E}(a,b)|_{(1,q)} = 2\xi''(q) - \left(\frac{1}{2(1-r(a,b))} + \frac{1}{2}\xi''(r(a,b))(1-r(a,b))\right) \hat{\partial}_b^2(r(a,b))|_{(1,q)} = -\frac{1}{(1-q)^2} + \xi''(q),$$
(7.134)

and

$$\partial_b \partial_a \tilde{E}(a,b)|_{(1,q)} = -\left(\frac{1}{2(1-r(a,b))} + \frac{1}{2}\xi''(r(a,b))(1-r(a,b))\right)\partial_{a,b}(r(a,b))|_{(1,q)}$$

= $-q\xi''(q) + \frac{q}{(1-q)^2}.$ (7.135)

Finally, we compute

$$\begin{aligned} \partial_a^2 \tilde{E}(a,b)|_{(1,q)} &= -q^3 \xi'''(q) + q(2\xi''(q) + q\xi'''(q)) + (1-q)(q^2\xi'''(q) + q\xi''(q) - q^2\xi''(q)^2/\xi'(q) + q^2v\xi''(q)^2/\xi'(q)^2) \\ &- \frac{q(2q+1)}{(1-q)^2} - 2q^2(1-q)\xi'''(q) + q(2q-1)\xi''(q) \\ &= -\frac{q(2q+1)}{(1-q)^2} + q(q+2)\xi''(q) + (1-q)(-q^2\xi''(q)^2/\xi'(q) + q^2v\xi''(q)^2/\xi'(q)^2). \end{aligned}$$
(7.136)

Using the constraints $v = t - \frac{t^2(1-q)}{q}$, and that $t = \frac{q}{1-q} - \xi'(q)$, we can simplify

$$\partial_a^2 \tilde{E}(a,b)|_{(1,q)} = -\frac{q(2q+1)}{(1-q)^2} + q(q+2)\xi''(q) - q(1-q)^2\xi''(q)^2.$$
(7.137)

Consider a change of variable $\tilde{a} = aq$ and let $\underline{E}(\tilde{a}, b) = \tilde{E}(\tilde{a}/q, b)$. Combining (7.134), (7.135), (7.137), under the condition $\xi''(q) < \frac{1}{(1-q)^2}$, that $\underline{E}(\tilde{a}, b)$ is strictly concave at $(\tilde{a}, b) = (q, q)$ is equivalent to

$$\frac{1}{q} \left((1-q)^2 \xi''(q)^2 - (q+2)\xi''(q) + \frac{(2q+1)}{(1-q)^2} \right) > \left(\frac{1}{(1-q)^2} - \xi''(q) \right)$$

$$\Leftrightarrow \frac{1}{q} \left((1-q)\xi''(q) - \frac{1}{1-q} \right)^2 + \frac{1}{(1-q)^2} > 0.$$

Notice that, for (a, b) in a neighborhood of $(1, b_*)$, the Hessian of $\hat{E}(a, b)$ is a continuous rational function of q, $\xi(b)$, $\xi(q)$, $\xi'(q)$, $\xi''(q)$, $\xi''(q)$, $\xi''(\langle x, m \rangle_N)$, and $\langle x, y \rangle_N$, $\langle x, m \rangle_N$, $\langle y, m \rangle_N$, $\|y\|_N^2$. Hence, we have the following implication of the previous lemma.

Corollary 7.18. There exist $\varepsilon, \eta > 0$ such that, for $|\langle \boldsymbol{x}, \boldsymbol{m} \rangle_N - q| \leq \varepsilon$, $|\langle \boldsymbol{y}, \boldsymbol{m} \rangle_N - t| \leq \varepsilon$, $|\langle \boldsymbol{y}, \boldsymbol{x} \rangle_N - t| \leq \varepsilon$, and $|||\boldsymbol{y}||_N^2 - t| \leq \varepsilon$, all (a, b) such that $|aq - a_*q| + |b - b_*| \leq \varepsilon$, we have $\nabla^2 \hat{E}(a, b) \leq -\eta \boldsymbol{I}_2$. (Here $(a_*, b_*) = (1, q)$.)

We will next prove several simple preliminary estimates before giving the proof of Proposition 4.6.

Recall that on $D_N(a, b)$, we have defined the Hamiltonian $\underline{H}(\tilde{\sigma})$, which is a spin glass with mixture given by Eqs. (7.112) to (7.114). Let $A^{(p)}(a, b) = \nabla^p \underline{H}(\mathbf{0})$ and $u(a, b) = \nabla \underline{H}(\mathbf{0})$.

By Lemma 7.15, Lemma 7.16, and Corollary 7.18 and the preceding remark, there is a unique local maxima $(a_*, b_*) = (1, b_*)$ of E(a, b) and $\hat{E}(a, b)$ with $|qa_* - q| + |b_* - q| \le \varepsilon$, and $\hat{E}(a, b)$ is strongly concave at (a_*, b_*) . In particular, there is $\eta > 0$ such that, for sufficiently small ε and (a, b) such that $|qa - qa_*| + |b - b_*| \le \varepsilon$, we have

$$E(a,b) \leq E(a_*,b_*) - \eta(|qa - qa_*|^2 + |b - b_*|^2).$$
(7.138)

For each a, b let $\boldsymbol{m}(a, b)$ be the unique point in $V := \operatorname{span}(\boldsymbol{m}, \boldsymbol{x})$ with $\|\boldsymbol{m}(a, b)\|_N^2 = qa$ and $\langle \boldsymbol{m}(a, b), \boldsymbol{x} \rangle_N = b$.

The following lemma follows from standard control on suprema of Gaussian processes (see, e.g. [MS23, Lemma A.3]).

Lemma 7.19. For $\varepsilon > 0$ sufficiently small there exist $c = c(\varepsilon)$, $C = C(\varepsilon) > 0$ depending uniquely on ε such that the following holds with probability at least $1 - e^{-cN}$ conditional on $\nabla \mathcal{F}_{\mathsf{TAP}}(\boldsymbol{m}) = 0$. For (a, b) such that $|qa - qa_*| + |b - b_*| \leq \varepsilon$, we have

$$\nabla H(\boldsymbol{m}(a,b)) = \nabla H(\boldsymbol{m}) + \nabla^2 H(\boldsymbol{m})(\boldsymbol{m}(a,b) - \boldsymbol{m}) + \mathsf{Err},$$

where $\|\operatorname{Err}\| \leq CN^{-1/2} \|\boldsymbol{m}(a,b) - \boldsymbol{m}\|^2$. Furthermore, $\mathsf{P}_V^{\perp}(\nabla H(\boldsymbol{m})) = 0$, and for $\boldsymbol{v} \in V$, $\|\nabla^2 H(\boldsymbol{m})\boldsymbol{v}\|^2 \leq C \|\boldsymbol{v}\|^2$. (Here $V = \operatorname{span}(\boldsymbol{m}, \boldsymbol{x})$.)

As a corollary of Lemma 7.19, we obtain the following control on the effective fields $\boldsymbol{u}(a,b) = (1 - r(a,b))^{1/2} \mathsf{P}_V^{\perp}(\nabla H(\boldsymbol{m})).$

Lemma 7.20. For ε , $\delta > 0$ sufficiently small, the following holds with probability at least $1 - e^{-cN}$. There exists $\mathbf{u}_1, \mathbf{u}_2 \in \mathbb{R}^{N-3}$ with $\|\mathbf{u}_1\|, \|\mathbf{u}_2\| = O(N^{1/2})$ such that, for any (a, b) with $|qa - qa_*| + |b - b_*| \leq \varepsilon$, we have $\|\mathbf{u}(a, b) - (qa - qa_*)\mathbf{u}_1 - (b - b_*)\mathbf{u}_2\| \leq CN^{1/2}(|qa - qa_*|^2 + |b - b_*|^2)$.

Furthermore, for $\gamma > \delta + \mathbb{E}\lambda_{\max}(\mathbf{A}^{(2)}(a_*, b_*))$ and $i, j \in \{1, 2\}$, there is $c = c(\delta) > 0$ such that, with probability $1 - e^{-N^c}$, $\langle \mathbf{u}_i, (\gamma \mathbf{I} - \mathbf{A}^{(2)}(a_*, b_*))^{-1}\mathbf{u}_j \rangle$, concentrates in a window of size $O(N^{1/2+c})$ around its expectation.

Proof. The first part follows from Lemma 7.19, using $\mathsf{P}_V^{\perp}(\nabla H(\boldsymbol{m})) = \mathbf{0}$.

The second part holds by concentration of Lipschitz functions of Gaussian random variables. Indeed note that u_i depend linearly on $A^{(2)}(a_*, b_*)$ as well as on independent Gaussian random variables. Under the high probability event $\mathbb{E}\lambda_{\max}(A^{(2)}(a_*, b_*)) + \delta/2 \leq \lambda_{\max}(A^{(2)}(a_*, b_*)) \leq C$, the quantity $\langle u_i, (\gamma I - A^{(2)}(a_*, b_*))^{-1}u_j \rangle$ is indeed Lipschitz in these Gaussians as well as on $A^{(2)}(a_*, b_*)$.

Let

$$R := \left\{ (a,b) : q|a - a_*| + |b - b_*| \le N^{-1/2+c} \right\}.$$
(7.139)

Recall the random shifts $g_{a,b}$ in Eq. (7.110). We have the following control on $g_{a,b}$, again from standard control on Gaussian processes.

Lemma 7.21. We have that $g_{a,b}$, for $(a,b) \in R$, forms a Gaussian process with $\mathbb{E}[(g_{a,b} - g_{a',b'})^2] = O(||\mathbf{m}(a,b) - \mathbf{m}(a',b')||_N^2)$. Furthermore, with probability at least $1 - e^{-cN}$, we have, for all $(a,b) \in R$ that

$$|g_{a,b} - g_{a_*,b_*}| \leq C(|q(a - a_*)| + |b - b_*|).$$

Proof. The first claim follows from a standard calculation, and the second claim follows Sudakov-Fernique inequality, comparing with the linear process $\langle \boldsymbol{g}, \boldsymbol{m}(a, b) - \boldsymbol{m}(a_*, b_*) \rangle / \sqrt{N}$ for \boldsymbol{g} a standard normal vector.

Lemma 7.22. The scaled GOE matrices $A^{(2)}(a, b)$ for $(a, b) \in R$ form a Gaussian process with metric

$$\mathbb{E}\left\{\left\|\boldsymbol{A}^{(2)}(a,b) - \boldsymbol{A}^{(2)}(a',b')\right\|_{\mathrm{F}}^{2}\right\} \leq CN(|qa - qa'|^{2} + |b - b'|^{2}).$$

Furthermore, for any $\eta > 0$ there exist constants c, C > 0 such that, with probability at least $1 - e^{-cN}$,

$$\|\boldsymbol{A}^{(2)}(a,b) - \boldsymbol{A}^{(2)}(a',b')\|_{\mathrm{F}}^{2} \leqslant CN(|qa - qa'|^{2} + |b - b'|^{2})^{1+\eta}, \quad \forall (a,b), (a',b') \in R,$$
(7.140)

$$\|\boldsymbol{A}^{(2)}(a,b) - \boldsymbol{A}^{(2)}(a',b')\|_{\text{op}}^2 \leq C(|qa - qa_*|^2 + |b - b_*|^2)^{1+\eta}, \quad \forall (a,b), (a',b') \in R.$$
(7.141)

and

$$\sup_{(a,b)\in R} \|\boldsymbol{A}^{(2)}(a,b) - \boldsymbol{A}^{(2)}(a_*,b_*)\|_{\mathrm{F}}^2 \leq CN^{2c},$$
(7.142)

$$\sup_{(a,b)\in R} \|\boldsymbol{A}^{(2)}(a,b) - \boldsymbol{A}^{(2)}(a_*,b_*)\|_{\rm op}^2 \leq CN^{-1+2c} \,.$$
(7.143)

Proof. The bound on the canonical distance of $A^{(2)}$ follows from a straightforward calculation.

The bounds (7.140) and (7.142) follow from chaining on R, together with the standard bound on chisquared random variables

$$\mathbb{P}\left(\|\boldsymbol{A}^{(2)}(a,b) - \boldsymbol{A}^{(2)}(a',b')\|_{\mathrm{F}}^{2} > \kappa \mathbb{E}\|\boldsymbol{A}^{(2)}(a,b) - \boldsymbol{A}^{(2)}(a',b')\|_{\mathrm{F}}^{2}\right) \leq 2e^{-cN^{2}[(\kappa-1)\wedge(\kappa-1)^{2}]}$$

for $\kappa > 1$.

The bound (7.141) and (7.143) follow from a similar chaining argument. Indeed, $A^{(2)}(a,b) - A^{(2)}(a',b')$ is a matrix with independent entries with variance bounded by $C(|qa - qa_*|^2 + |b - b_*|^2)/N$, whence by standard estimates on the norm of Gaussian random matrices, the following holds for all $\kappa > \kappa_0$

$$\mathbb{P}\left(\|\boldsymbol{A}^{(2)}(a,b) - \boldsymbol{A}^{(2)}(a',b')\|_{\text{op}}^{2} > \kappa N^{-1}\mathbb{E}\|\boldsymbol{A}^{(2)}(a,b) - \boldsymbol{A}^{(2)}(a',b')\|_{\text{F}}^{2}\right) \leq 2e^{-cN\kappa}.$$

Recall $G(\gamma; \mathbf{A}, \mathbf{u})$ in (7.10) and $G_{a,b}(\gamma)$ in (7.120). The next lemma gives control over G and $G_{a,b}$ for $(a, b) \in \mathbb{R}$.

Lemma 7.23. Given a compact interval $I \subseteq [M, \infty)$, $M := \varepsilon + \mathbb{E}\lambda_{\max}(\mathbf{A}^{(2)}(a_*, b_*))$, the following holds with probability at least $1 - e^{-N^{\delta}}$ for appropriate $C_0, c, \delta > 0$ depending on $\varepsilon > 0$:

1. We have

$$\sup_{\gamma \in I, (a,b) \in \mathbb{R}} \left| N(G(\gamma; \mathbf{A}^{(2)}(a,b), \mathbf{u}(a,b)) - G_{a,b}(\gamma)) - N(G(\gamma; \mathbf{A}^{(2)}(a_*,b_*), \mathbf{u}(a_*,b_*)) - G_{a_*,b_*}(\gamma)) \right| = O(N^{-1/2+c}).$$
(7.144)

2. We have

$$\sup_{\substack{(a,b),(a',b')\in R:(a,b)+(a',b')=2(a_{*},b_{*})\\\gamma\in I}} \left| NG(\gamma; \boldsymbol{A}^{(2)}(a,b), \boldsymbol{u}(a,b)) - NG(\gamma; \boldsymbol{A}^{(2)}(a',b'), \boldsymbol{u}(a',b')) \right|$$
$$= O(N^{-1/2+c}).$$
(7.145)

3. The event in Lemma 7.3 holds uniformly in $(a, b) \in R$. Namely,

$$Z_{\leq 2}(a,b) = \int e^{\langle \boldsymbol{\sigma}, \boldsymbol{A}^{(2)}(a,b)\boldsymbol{\sigma} \rangle + \langle \boldsymbol{u}(a,b), \boldsymbol{\sigma} \rangle} \mu_0^{a,b}(\mathsf{d}\boldsymbol{\sigma})$$

= $(1 + \mathsf{Err}_{a,b}(N))(2e)^{-(N-2)/2} \sqrt{\frac{2}{G''(\gamma_{a,b}; \boldsymbol{A}^{(2)}(a,b), \boldsymbol{u}(a,b))}} e^{NG(\gamma_{a,b}; \boldsymbol{A}^{(2)}(a,b), \boldsymbol{u}(a,b))},$ (7.146)

where $\sup_{(a,b)\in R} |\mathsf{Err}_{a,b}(N)| \leq C_0, N^{-c}$.

Proof. We can represent $A^{(2)}(a,b) = A^{(2)}(a_*,b_*) + \Delta(a,b)$ where each entry of $\Delta(a,b)$ forms an independent Gaussian process with metric $\mathbb{E}[(\Delta_{i,j}(a,b) - \Delta_{i,j}(a',b'))^2]^{1/2} \leq CN^{-1/2}(q|a-a'|+|b-b'|)$, and $\Delta(a_*,b_*) = 0$.

Letting $Q_*(\gamma) = \gamma I - A^{(2)}(a_*, b_*)$, we can expand

$$G(\gamma; \mathbf{A}^{(2)}(a, b), \mathbf{u}(a, b)) =$$

$$= \gamma - \frac{1}{2N} \log \det(\gamma \mathbf{I} - \mathbf{A}^{(2)}(a, b)) + \frac{1}{4N} \langle \mathbf{u}(a, b), (\gamma \mathbf{I} - \mathbf{A}^{(2)}(a, b))^{-1} \mathbf{u}(a, b) \rangle$$

$$= G(\gamma; \mathbf{A}^{(2)}(a_{*}, b_{*}), \mathbf{u}(a_{*}, b_{*})) - \frac{1}{2N} \log \det \left(\mathbf{I} - \mathbf{Q}_{*}(\gamma)^{-1/2} \mathbf{\Delta}(a, b) \mathbf{Q}_{*}(\gamma)^{-1/2}\right)$$
(7.147)
$$+ \frac{1}{4N} \langle \mathbf{u}(a, b), (\mathbf{Q}_{*}(\gamma) - \mathbf{\Delta}(a, b))^{-1} \mathbf{u}(a, b) \rangle - \frac{1}{4N} \langle \mathbf{u}(a, b), \mathbf{Q}_{*}(\gamma)^{-1} \mathbf{u}(a, b) \rangle.$$

Next, for $k \ge 2$, let

$$X_k(a,b) = \mathsf{Tr}\left(\left(\boldsymbol{Q}_*(\gamma)^{-1/2}\boldsymbol{\Delta}(a,b)\boldsymbol{Q}_*(\gamma)^{-1/2}\right)^k\right).$$
(7.148)

We have

$$|X_k(a,b) - X_k(a',b')| \leq C_k N^{-(k-1)(1/2-c)+1/2} \| \mathbf{A}^{(2)}(a,b) - \mathbf{A}^{(2)}(a',b') \|_{\mathrm{F}},$$

under the event in Lemma 7.22. Recall that this also guarantees

$$\sup_{(a,b)\in R} \left\{ \| \boldsymbol{\Delta}(a,b) \|_{\rm op} \vee \| \boldsymbol{Q}_{*}(\gamma)^{-1/2} \boldsymbol{\Delta}(a,b) \boldsymbol{Q}_{*}(\gamma)^{-1/2} \|_{\rm op} \right\} \leqslant C N^{-1/2+c},$$
(7.149)

Hence, under the event in Lemma 7.22, we have $|X_k(a,b) - X_k(a',b')| \le N^{-(k-1)(1/2-c)-L+1}$ whenever $q|a - a'| + |b - b'| < N^{-L}$.

Let \mathbb{P}_{Δ} and \mathbb{E}_{Δ} denote probability and expectation with respect to $\Delta(a, b)$ only, i.e. conditional on $A^{(2)}(a_*, b_*)$. Also let

$$\boldsymbol{M}(a,b) = \boldsymbol{Q}_{*}(\gamma)^{-1/2} \boldsymbol{\Delta}(a,b) \boldsymbol{Q}_{*}(\gamma)^{-1/2}$$

be the matrix appearing in $X_k(a,b)$. On the $A^{(2)}(a_*,b_*)$ -measurable, probability $1 - e^{-cN}$ event that $Q_*(\gamma)^{-1/2}$ is bounded in operator norm, M(a,b) is (conditional on $A^{(2)}(a_*,b_*)$), in a suitable basis) a random matrix with independent centered gaussian entries, with variances not equal but bounded uniformly by N^{-2+2c} . It is well known, cf. [AGZ10, Chapter 2] that tracial moments of M(a,b) amount to certain (weighted) cycle counts, and from this a routine calculation gives

$$\mathbb{E}_{\mathbf{\Delta}} X_k(a,b) = \begin{cases} 0 & k \text{ odd,} \\ O_k(N^{1-k(1/2-c)}) & k \text{ even,} \end{cases} \quad \text{Var}_{\mathbf{\Delta}} [X_k(a,b)] = O_k(N^{-k(1-2c)}).$$

(The last estimate amounts to computing cycle counts for $\mathbb{E}[\text{Tr}(M(a,b)^k)^2]$ and $\mathbb{E}[\text{Tr}(M(a,b)^k)]^2$, cf. [AGZ10, Proof of Lemma 2.1.7].) For any fixed $(a,b) \in R$, Gaussian hypercontractivity gives

$$\mathbb{P}_{\Delta}\left\{|X_k(a,b) - \mathbb{E}_{\Delta}X_k(a,b)| \ge t\sqrt{\operatorname{Var}_{\Delta}(X_k(a,b))}\right\} \le \exp(-t^{c_k}),$$

because $X_k(a, b)$ is a degree k-polynomial in the entries of M(a, b). By a union bound over a N^{-L} -net of R (of size N^{2L}), with probability $1 - e^{-N^{c_k\delta}}$ over $\Delta(a, b)$ the following holds.

For k even, uniformly in $(a, b) \in R$

$$\operatorname{Tr}\left(\left(\boldsymbol{Q}_{*}(\boldsymbol{\gamma})^{-1/2}\boldsymbol{\Delta}(a,b)\boldsymbol{Q}_{*}(\boldsymbol{\gamma})^{-1/2}\right)^{k}\right) - \mathbb{E}\operatorname{Tr}\left(\left(\boldsymbol{Q}_{*}(\boldsymbol{\gamma})^{-1/2}\boldsymbol{\Delta}(a,b)\boldsymbol{Q}_{*}(\boldsymbol{\gamma})^{-1/2}\right)^{k}\right) = O_{k}(N^{\delta-k(1/2-c)}),$$

and for k odd,

$$\sup_{(a,b)\in R} \operatorname{Tr}\left(\left(\boldsymbol{Q}_{*}(\gamma)^{-1/2}\boldsymbol{\Delta}(a,b)\boldsymbol{Q}_{*}(\gamma)^{-1/2}\right)^{k}\right) = O_{k}(N^{\delta-k(1/2-c)}).$$

Recall that $|\log(1-x) + x + x^2/2| \le |x|^3$ for all $|x| \le 1/4$. Therefore, uniformly in $(a, b) \in R$, for all δ , c small enough

$$\log \det \left(\boldsymbol{I} - \boldsymbol{Q}_{*}(\gamma)^{-1/2} \boldsymbol{\Delta}(a, b) \boldsymbol{Q}_{*}(\gamma)^{-1/2} \right)$$

= $-\sum_{k=1}^{2} \frac{1}{k} \operatorname{Tr} \left(\left(\boldsymbol{Q}_{*}(\gamma)^{-1/2} \boldsymbol{\Delta}(a, b) \boldsymbol{Q}_{*}(\gamma)^{-1/2} \right)^{k} \right) + O(N^{-1+c})$
= $-\mathbb{E} \operatorname{Tr} \left(\left(\boldsymbol{Q}_{*}(\gamma)^{-1/2} \boldsymbol{\Delta}(a, b) \boldsymbol{Q}_{*}(\gamma)^{-1/2} \right)^{2} \right) + O(N^{-1/2+c+\delta}).$ (7.150)

We next turn to the term $\langle \boldsymbol{u}(a,b), (\boldsymbol{Q}_*(\gamma) - \boldsymbol{\Delta}(a,b))^{-1}\boldsymbol{u}(a,b) \rangle$ in Eq. (7.147). From Lemma 7.20, there are $\boldsymbol{u}_1, \boldsymbol{u}_2$ with $\|\boldsymbol{u}_1\|, \|\boldsymbol{u}_2\| = O(N^{1/2})$ such that letting $\boldsymbol{u}_0(a,b) := q(a-a_*)\boldsymbol{u}_1 + (b-b_*)\boldsymbol{u}_2$, we have $\|\boldsymbol{u}(a,b) - \boldsymbol{u}_0(a,b)\| \leq CN^{-1/2+2c}$ and $\|\boldsymbol{u}_0(a,b)\| \leq CN^c$ for any $(a,b) \in R$ Therefore, with probability $1 - e^{-N^{\delta}}$,

$$\begin{aligned} \langle \boldsymbol{u}_0(a,b), (\boldsymbol{Q}_*(\gamma) - \boldsymbol{\Delta}(a,b))^{-1} \boldsymbol{u}_0(a,b) \rangle \\ &= \langle \boldsymbol{u}_0(a,b), \boldsymbol{Q}_*(\gamma)^{-1} \boldsymbol{u}_0(a,b) \rangle + \langle \boldsymbol{u}_0(a,b), \boldsymbol{Q}_*(\gamma)^{-1} \boldsymbol{\Delta}(a,b) \boldsymbol{Q}_*(\gamma)^{-1} \boldsymbol{u}_0(a,b) \rangle + O(N^{-1+3c}) \\ &= \langle \boldsymbol{u}_0(a,b), \boldsymbol{Q}_*(\gamma)^{-1} \boldsymbol{u}_0(a,b) \rangle + O(N^{-1+3c+\delta}). \end{aligned}$$

where the first estimate follows from Eq. (7.149), and the second from independence of $\Delta(a, b)$ and u_1, u_2 , together with the fact that the entries of $\Delta(a, b)$ have variance bounded by N^{-2+2c} . Therefore, we obtain that, with probability $1 - e^{-N^{\delta}}$,

$$\langle \boldsymbol{u}(a,b), (\boldsymbol{Q}_{*}(\gamma) - \boldsymbol{\Delta}(a,b))^{-1} \boldsymbol{u}(a,b) \rangle$$

= $\langle q(a-a_{*})\boldsymbol{u}_{1} + (b-b_{*})\boldsymbol{u}_{2}, \boldsymbol{Q}_{*}(\gamma)^{-1}(q(a-a_{*})\boldsymbol{u}_{1} + (b-b_{*})\boldsymbol{u}_{2}) \rangle + O(N^{-1+4c+\delta}).$ (7.151)

By similarly taking a union bound over a net of R of radius N^{-L} and using the continuity in Lemma 7.22, we can guarantee Eq. (7.151) uniformly in $(a, b) \in R$.

Combining the last conclusion in Lemma 7.20, Eqs. (7.150) and (7.151) and a union bound over γ , over any compact interval of γ , upon changing δ , with probability at least $1 - e^{-N^{\delta}}$, that

$$\sup_{\substack{\gamma,(a,b)\in R}} \left| N(G(\gamma; \mathbf{A}^{(2)}(a,b), \boldsymbol{u}(a,b)) - G_{a,b}(\gamma) - G(\gamma; \mathbf{A}^{(2)}(a_*,b_*), \boldsymbol{u}(a_*,b_*)) + G_{a_*,b_*}(\gamma)) \right| \\ \leq O(N^{-1/2+c}).$$

Thus, we have, with probability at least $1 - e^{-N^{\delta}}$, uniformly in $(a, b) \in R$ and γ , that Eq. (7.144) holds. Given $(a, b), (a', b') \in R$ such that $(a, b) + (a', b') = 2(a_*, b_*)$, we have that

$$\mathbb{E}\mathsf{Tr}\left(\left(\boldsymbol{Q}_{*}(\gamma)^{-1/2}\boldsymbol{\Delta}(a,b)\boldsymbol{Q}_{*}(\gamma)^{-1/2}\right)^{2}\right) = \mathbb{E}\mathsf{Tr}\left(\left(\boldsymbol{Q}_{*}(\gamma)^{-1/2}\boldsymbol{\Delta}(a',b')\boldsymbol{Q}_{*}(\gamma)^{-1/2}\right)^{2}\right).$$

Combining with Eqs. (7.150) and (7.151), we then obtain Eq. (7.145).

Finally, we recall that for each $(a, b) \in R$, the event in Lemma 7.3 holds with probability at least $1 - e^{-cN}$. On the other hand, for appropriate C > C' > 1, using Lemma 7.20 and Lemma 7.22, with probability $1 - e^{-cN}$, uniformly over $(a, b), (a', b') \in R$ with $|q(a - a')| + |b - b'| \leq N^{-C}$, we have $Z_{\leq 2}(a, b) = (1 + O(N^{-C'}))Z_{\leq 2}(a', b')$. Similar continuity estimates hold for the right hand side of Eq. (7.146). Taking a net of radius N^{-C} of R and apply the union bound, we obtain Eq. (7.146) uniformly in $(a, b) \in R$.

Proof of Proposition 4.6. Consider appropriate constants c > c' > 0. Define

$$D_N(\varepsilon) := \left\{ \boldsymbol{\sigma} \in S_N : N^{-1/2+c} \leq |\langle \boldsymbol{\sigma}, \boldsymbol{m} \rangle_N - a_* q| + |\langle \boldsymbol{\sigma}, \boldsymbol{x} \rangle_N - b_*| \leq \varepsilon \right\},$$
(7.152)

$$\hat{D}_N(\varepsilon) := \left\{ (a,b) \in \mathbb{R}^2 : N^{-1/2+c} \le |aq - a_*q| + |b - b_*| \le \varepsilon \right\},\tag{7.153}$$

Using Markov's Inequality and that the annealing upper bound $\hat{E}(a, b)$ is strongly concave for $(a, b) \in \hat{D}_N(\varepsilon)$ (see Lemma 7.16 and Corollary 7.18), we obtain that with probability $1 - \exp(-N^{c'})$, for all sufficiently small $\varepsilon > 0$, there is $\eta > 0$ such that

$$\int_{D_{N}(\varepsilon)} e^{H(\sigma)} \mu_{0}(\mathrm{d}\sigma) = \int_{\hat{D}_{N}(\varepsilon)} \left\{ \int e^{H(\sigma)} \mu_{0}^{a,b}(\mathrm{d}\sigma) \right\} \mathrm{d}a \mathrm{d}b$$

$$\leq e^{-\eta(|qa-qa_{*}|^{2}+|b-b_{*}|^{2})+N\hat{E}(a_{*},b_{*})+\sqrt{N}g_{a,b}}$$

$$= e^{-\eta(|qa-qa_{*}|^{2}+|b-b_{*}|^{2})+NE(a_{*},b_{*})+\sqrt{N}g_{ab}}.$$
(7.154)

Denote by $\langle \cdot \rangle_{a,b}$ the average with respect to the Gibbs measure restricted to band $D_N(a,b)$, namely with respect to $\mu^{a,b}(d\sigma) \propto \exp\{NH(\sigma)\} \mu_0^{a,b}(d\sigma)$. Note that $u(a_*, b_*) = 0$, and for $|qa - qa_*| + |b - b_*| \leq N^{-1/2+c}$, we have $||u(a,b)|| = O(N^{c'})$.

Recall $\gamma_{a,b} = \arg \min_{z > \lambda_{\max}(\boldsymbol{A}^{(2)}(a,b))} G(z; \boldsymbol{A}^{(2)}(a,b), \boldsymbol{u}(a,b))$. Let

$$\Delta(a,b) := \frac{1}{2} (\gamma_{a,b} - \mathbf{A}^{(2)}(a,b))^{-1} \boldsymbol{u}(a,b) + \frac{1}{2} (\gamma_{a,b} - \mathbf{A}^{(2)}(a,b))^{-1} \langle \mathbf{A}^{(3)}(a,b), (\gamma_{a,b} - \mathbf{A}^{(2)}(a,b))^{-1} \rangle.$$

We have $\Delta(a_*, b_*) = \Delta(m) + O(N^{-c})$, where $\Delta(m)$ is defined as per Eq. (2.13). Let

$$Z(a,b) := \int e^{H(\boldsymbol{\sigma})} \mu_0^{a,b}(\boldsymbol{\sigma}),$$

and

$$Z := \int_{\mathsf{Band}_{*}(2\iota)} \exp(H_{N,t}(\boldsymbol{\sigma})) \ \mu_{0}(\mathsf{d}\boldsymbol{\sigma}).$$

Recall the definitions

$$R = \{(a,b) \in \mathbb{R}^2 : |qa - qa_*| + |b - b_*| \leq N^{-1/2+c} \},\$$

$$R_+ = \{(a,b) \in \mathbb{R}^2 : |qa - qa_*| + |b - b_*| \leq N^{-1/2} \}.$$

Using Eq. (7.154), we have that, with probability at least $1 - \exp(-N^{c'})$,

$$\widetilde{\boldsymbol{m}}_{2\iota}(\boldsymbol{m}) = \frac{\int_{\mathsf{Band}_{\ast}(2\iota)} \boldsymbol{\sigma} \exp(H_{N,t}(\boldsymbol{\sigma})) \ \mu_0(\mathsf{d}\boldsymbol{\sigma})}{\int_{\mathsf{Band}_{\ast}(2\iota)} \exp(H_{N,t}(\boldsymbol{\sigma})) \ \mu_0(\mathsf{d}\boldsymbol{\sigma})} \\ = O(e^{-\eta N^{2c}}) + \int_R \frac{Z(a,b)}{Z} \langle \boldsymbol{\sigma} \rangle_{a,b} \mathsf{d}(a,b)$$

Let

$$Z_T(a,b) = \int \mathbf{1}\{\boldsymbol{\sigma} \in T(a,b)\} e^{H(\boldsymbol{\sigma})} \mu_0^{a,b}(\mathsf{d}\boldsymbol{\sigma}), \qquad Z_T = \int_{\mathsf{Band}_*(2\iota)} \mathbf{1}\{\boldsymbol{\sigma} \in T(a,b)\} e^{H(\boldsymbol{\sigma})} \mu_0(\mathsf{d}\boldsymbol{\sigma}),$$

where $T(a,b) \subseteq D_N(a,b)$ is the typical set (7.31) defined for the effective model on $D_N(a,b)$. Recall from Lemma 7.7 that for each $(a,b) \in R$, with probability $1 - e^{-cN}$,

$$Z_T(a,b) \ge (1-e^{-cN})Z(a,b), \qquad \mathbb{E}_{\ge 3}Z_T(a,b) \ge (1-e^{-cN})\mathbb{E}_{\ge 3}Z(a,b).$$
 (7.155)

By a union bound over a $e^{-cN/10}$ -net of R and standard continuity properties of H, with probability $1 - e^{-cN/2}$ this holds simultaneously for all $(a, b) \in R$. By integrating, on this event we also have

$$Z_T \ge (1 - e^{-cN})Z, \qquad \mathbb{E}_{\ge 3}Z_T \ge (1 - e^{-cN})\mathbb{E}_{\ge 3}Z.$$
 (7.156)

Note that, by Eq. (7.37) in Lemma 7.8, for $k > L \ge 1$, the following holds with probability at least $1 - e^{-cN}$,

$$\mathbb{E}_{\geq 3}\left[(Z_T(a,b) - \mathbb{E}_{\geq 3} Z_T(a,b))^{2k} \right] \leq C_L N^{-L} \left(\mathbb{E}_{\geq 3} Z_T(a,b) \right)^{2k},$$
(7.157)

and therefore

$$\mathbb{E}_{\geq 3}\left[Z_T(a,b)^{2k}\right] \leq (1 + C_L N^{-L}) \left(\mathbb{E}_{\geq 3} Z_T(a,b)\right)^{2k}.$$
(7.158)

Again by standard continuity estimates and the union bound over a net of R of radius $e^{-c'N}$, the above estimates hold uniformly in $(a, b) \in R$ with probability at least $1 - e^{-cn/2}$. By Eq. (7.156), the same estimates hold for Z in place of Z_T uniformly in $(a, b) \in R$ with probability at least $1 - e^{-c'N}$.

By Eqs. (7.144), (7.146) of Lemma 7.23, together with Eqs. (7.110) and Lemma 7.21 (which implies that $e^{\sqrt{N}(g_{a,b}-g_{a',b'})} = O(1)$ for all $(a,b), (a',b') \in R_+$), with probability at least $1 - e^{-N^{\delta}}$, uniformly in $(a,b) \in R_+$,

$$\mathbb{E}_{\geq 3}Z(a,b) \geq \Omega\left(\mathbb{E}_{\geq 3}Z(a_*,b_*)\right). \tag{7.159}$$

From strict concavity of E(a, b) at (a_*, b_*) (see (7.138)), and the simple estimate

$$\sup_{a,b} \left(N^{1/2} (|q(a-a_*)| + |b-b_*|) - \eta N (|q(a-a_*)|^2 + |b-b_*|^2) \right)$$

= $O_\eta(1) - \eta N (|q(a-a_*)|^2 + |b-b_*|^2),$ (7.160)

we obtain that, uniformly in $(a, b) \in R$,

$$\mathbb{E}_{\geq 3}Z(a,b) \leq O\left(\mathbb{E}_{\geq 3}Z(a_{*},b_{*}) \cdot e^{-\eta N(|qa-qa_{*}|^{2}+|b-b_{*}|^{2})/2}\right).$$
(7.161)

Further, by Lemma 7.8, we also have

$$\mathbb{P}\left\{\left|Z_T - \mathbb{E}_{\geqslant 3}Z_T\right| > \frac{1}{2} \mathbb{E}_{\geqslant 3}Z_T\right\} \leqslant CN^{-L/2}.$$
(7.162)

Since R_+ has volume $\Theta(N^{-1})$, on the event in (7.159) we have

$$\mathbb{E}_{\geqslant 3}Z \geqslant \int_{R_+} \mathbb{E}_{\geqslant 3}Z(a,b) \operatorname{d}(a,b) = \Omega(N^{-1}\mathbb{E}_{\geqslant 3}Z(a_*,b_*)).$$

Furthermore, when (7.161) holds we have

$$\mathbb{E}_{\geq 3}Z = \int_R \mathbb{E}_{\geq 3}Z(a,b) \operatorname{\mathsf{d}}(a,b) \leqslant O(N^{-1}\mathbb{E}_{\geq 3}Z(a_*,b_*))),$$

where the N^{-1} comes from integrating the exponential in (7.161). Thus, with probability at least $1 - e^{-N^{\delta}}$,

$$\mathbb{E}_{\geq 3}Z = \Theta(N^{-1}\mathbb{E}_{\geq 3}Z(a_*, b_*)).$$
(7.163)

Let \mathcal{E} denote the event that estimates (7.155), (7.157), (7.158), (7.159), (7.161), (7.162), (7.163) all hold. By the above, we have $\mathbb{P}(\mathcal{E}) \ge 1 - CN^{-L/2}$. Further, for $\hat{Z}(a, b) = e^{NE(a, b)}$ and $\hat{Z} = \int_R \hat{Z}(a, b) d(a, b)$,

$$\int_{R} \mathbb{E}\left[\mathbf{1}_{\mathcal{E}}\left(\frac{Z(a,b)}{Z}\right)^{2k}\right] \mathsf{d}(a,b) = O\left(\int_{R} \mathbb{E}\left[\mathbf{1}_{\mathcal{E}}\left(\frac{\hat{Z}(a,b)}{\hat{Z}}\right)^{2k}\right] \mathsf{d}(a,b)\right)$$
(7.164)

Under the event \mathcal{E} , from Eqs. (7.159), (7.161), we thus obtain

$$\int_{R} \mathbb{E}\left[\mathbf{1}_{\mathcal{E}}\left(\frac{Z(a,b)}{Z}\right)^{2}\right]^{1/2} \mathsf{d}(a,b) = O(1), \tag{7.165}$$

By Jensen and Cauchy-Schwarz inequality,

$$\mathbb{E}\left[\mathbf{1}_{\mathcal{E}}\left\|\int_{R}\frac{Z(a,b)}{Z}(\langle\boldsymbol{\sigma}\rangle_{a,b}-\boldsymbol{\Delta}(a,b)-\boldsymbol{m}(a,b))\mathsf{d}(a,b)\right\|^{2+\delta}\right]$$

$$\leq \int_{R}\mathbb{E}\left[\mathbf{1}_{\mathcal{E}}\frac{Z(a,b)}{Z}\left\|(\langle\boldsymbol{\sigma}\rangle_{a,b}-\boldsymbol{\Delta}(a,b)-\boldsymbol{m}(a,b))\right\|^{2+\delta}\right]\mathsf{d}(a,b)$$

$$\leq \int_{R}\mathbb{E}\left[\mathbf{1}_{\mathcal{E}}\left(\frac{Z(a,b)}{Z}\right)^{2}\right]^{1/2}\mathbb{E}\left[\left\|(\langle\boldsymbol{\sigma}\rangle_{a,b}-\boldsymbol{\Delta}(a,b)-\boldsymbol{m}(a,b))\right\|^{2(2+\delta)}\right]^{1/2}\mathsf{d}(a,b).$$

By Lemma 7.2, we have

$$\mathbb{E}\left[\|\langle \boldsymbol{\sigma} \rangle_{a,b} - \Delta(a,b) - \boldsymbol{m}(a,b)\|^{2(2+\delta)}\right] \leq N^{-c}.$$
(7.166)

Combining with Eq. (7.165), we obtain that

$$\mathbb{E}\left[\mathbf{1}_{\mathcal{E}}\left\|\int_{R}\frac{Z(a,b)}{Z}(\langle\boldsymbol{\sigma}\rangle_{a,b}-\Delta(a,b)-\boldsymbol{m}(a,b))\mathsf{d}(a,b)\right\|^{2+\delta}\right] \leq O(N^{-c}).$$
(7.167)

On the other hand, by Lemma 7.20, with probability $1 - e^{-cN}$, there are u_1, u_2 with $||u_1||, ||u_2|| = O(N^{1/2})$ such that, for $|qa - qa_*| + |b - b_*| \leq N^{-1/2+c}$,

$$\|\boldsymbol{u}(a,b) - (q(a-a_*)\boldsymbol{u}_1 + (b-b_*)\boldsymbol{u}_2)\| = O(N^{-1/2+2c}).$$
(7.168)

Using this, letting $\overline{Z} = \mathbb{E}_{\geqslant 3}Z(a_*, b_*)$ and defining $\overline{a} := a - a_*, \overline{b} := b - b_*$, we have

$$\begin{split} &\int_{R} \frac{Z(a,b)}{Z} \boldsymbol{u}(a,b) \mathsf{d}(a,b) \\ &= \int_{R} \frac{Z(a,b) - \overline{Z}}{Z} (q \bar{a} \boldsymbol{u}_{1} + \bar{b} \boldsymbol{u}_{2}) \mathsf{d}(a,b) + \int_{R} \frac{\overline{Z}}{Z} (q \bar{a} \boldsymbol{u}_{1} + \bar{b} \boldsymbol{u}_{2}) \mathsf{d}(a,b) + O(N^{-1/2+2c}) \\ &= \int_{R} \frac{Z(a,b) - \overline{Z}}{Z} (q \bar{a} \boldsymbol{u}_{1} + \bar{b} \boldsymbol{u}_{2}) \mathsf{d}(a,b) + O(N^{-1/2+2c}) \\ &= \int_{R} \frac{Z(a,b) - \mathbb{E}_{\geq 3} Z(a,b)}{Z} (q \bar{a} \boldsymbol{u}_{1} + \bar{b} \boldsymbol{u}_{2}) \mathsf{d}(a,b) \\ &+ \int_{R} \frac{\mathbb{E}_{\geq 3} Z(a,b) - \mathbb{E}_{\geq 3} Z(a_{*},b_{*})}{Z} (q \bar{a} \boldsymbol{u}_{1} + \bar{b} \boldsymbol{u}_{2}) \mathsf{d}(a,b) + O(N^{-1/2+2c}). \end{split}$$

Furthermore, by Hölder's inequality on the measure $\frac{\mathbf{1}\{(a,b)\in R\}\mathbf{d}(a,b)}{\operatorname{Vol}(R)}$,

$$\mathbb{E}\left[\mathbf{1}_{\mathcal{E}}\left\|\int_{R}\frac{Z(a,b)-\mathbb{E}_{\geq 3}Z(a,b)}{Z}(q(a-a_{*})\boldsymbol{u}_{1}+(b-b_{*})\boldsymbol{u}_{2})\mathsf{d}(a,b)\right\|^{2+\delta}\right]$$

$$\leq \int_{R}\mathbb{E}\left[\left(\mathbf{1}_{\mathcal{E}}\left|\frac{Z(a,b)-\mathbb{E}_{\geq 3}Z(a,b)}{Z}\right|\|q(a-a_{*})\boldsymbol{u}_{1}+(b-b_{*})\boldsymbol{u}_{2}\|\operatorname{Vol}(R)\right)^{2+\delta}\right]\frac{\mathsf{d}(a,b)}{\operatorname{Vol}(R)}$$

By (7.161) and (7.163), $Z = \Omega(N^{-1}Z_{\geq 3}(a,b))$. Moreover, we have the estimates $|q(a - a_*)|, |b - b_*| \leq N^{-1/2+c}$ by definition of R, $\operatorname{Vol}(R) \leq N^{-1+2c}$, and $||\mathbf{u}_1||, ||\mathbf{u}_2|| \leq \sqrt{N}$. Combining these estimates, the last display is bounded by

$$O(N^{3c(2+\delta)}) \int_{R} \mathbb{E}\left[\mathbf{1}_{\mathcal{E}} \left| \frac{Z(a,b) - \mathbb{E}_{\geq 3}Z(a,b)}{Z_{\geq 3}(a,b)} \right|^{2+\delta} \right] \frac{\mathsf{d}(a,b)}{\mathsf{Vol}(R)}$$
(7.169)

Finally, since \mathcal{E} contains the event that (7.155), (7.157) holds for (a, b), for any $(a, b) \in \mathbb{R}$ we have the estimate

$$\mathbb{E}\left[\mathbf{1}_{\mathcal{E}} \left| \frac{Z(a,b) - \mathbb{E}_{\geq 3} Z(a,b)}{Z_{\geq 3}(a,b)} \right|^{2+\delta} \right]$$

$$\leq \mathbb{E}\left[\mathbf{1}((7.155), (7.157) \text{ holds for } (a,b))\mathbb{E}_{\geq 3} \left| \frac{Z(a,b) - \mathbb{E}_{\geq 3} Z(a,b)}{Z_{\geq 3}(a,b)} \right|^{2+\delta} \right]$$

$$\leq \mathbb{E}\left[\mathbf{1}((7.155), (7.157) \text{ holds for } (a,b)) \frac{(\mathbb{E}_{\geq 3} |Z_T(a,b) - \mathbb{E}_{\geq 3} Z_T(a,b)|^{2k})^{(2+\delta)/2k}}{|Z_{\geq 3}(a,b)|^{2+\delta}} \right] + e^{-cN},$$

and by (7.157) this is bounded by $N^{-1/2}$. Then, for c small enough, (7.169) is bounded by $O(N^{-c})$.

By Eqs. (7.146) and (7.145) of Lemma 7.23,

$$\left\|\int_{R} \frac{\mathbb{E}_{\geq 3}Z(a,b) - \mathbb{E}_{\geq 3}Z(a_{*},b_{*})}{Z} (q\bar{a}\boldsymbol{u}_{1} + \bar{b}\boldsymbol{u}_{2})\mathsf{d}(a,b)\right\| = O(N^{-c}).$$

Similarly, we have

$$\mathbb{E}\left[\left\|\int_{R}\frac{Z(a,b)}{Z}\boldsymbol{m}(a,b)\mathsf{d}(a,b) - \int_{R}\frac{\mathbb{E}_{\geq 3}Z(a,b)}{Z}\boldsymbol{m}(a,b)\mathsf{d}(a,b)\right\|^{2+\delta}\right] = O(N^{-c}),$$

Again by Eqs. (7.146) and (7.145) of Lemma 7.23, noting that m(a, b) + m(a', b') = 2m if $(a, b) + (a', b') = 2(a_*, b_*)$,

$$\mathbb{E}\left[\left\|\int_{R}\frac{\mathbb{E}_{\geq 3}Z(a,b)}{Z}\boldsymbol{m}(a,b)\mathsf{d}(a,b)-\boldsymbol{m}\right\|^{2+\delta}\right]=O(N^{-c}),$$

Thus, we obtain

$$\mathbb{E}\left[\|\widetilde{m}_{2\iota}(m) - m - \Delta(m)\|^{2+\delta}\right] \leq O(N^{1+\delta/2}e^{-\eta N^c}) + O(N^{-c}) + O(N^{1+\delta/2} \cdot N^{-L}) = O(N^{-c}).$$
(7.170)

Proposition 4.6 then follows.

8 Lognormal fluctuations of partition function

Proof of Lemma 3.2. Recall that $H_{N,2}$ denotes the degree-2 part of H_N , which is of the form

$$H_{N,2}(oldsymbol{\sigma}) = rac{\xi''(0)^{1/2}}{2} \langle oldsymbol{G} oldsymbol{\sigma}, oldsymbol{\sigma}
angle,$$

for $\boldsymbol{G} \sim \mathsf{GOE}(N)$. Let

$$Z_{N,2} = \int_{S_N} \exp H_{N,2}(\boldsymbol{\sigma}) \, \mathrm{d}\mu_0(\boldsymbol{\sigma})$$

It follows from [BL16, Theorem 1.2] (with $w_2 = 2, W_4 = 3$) that, with $\sigma^2 = -\frac{1}{2}\log(1 - \xi''(0))$ and $W \sim \mathcal{N}(-\frac{1}{2}\sigma^2, \sigma^2)$,

$$\frac{Z_{N,2}}{\mathbb{E}Z_{N,2}} = \frac{Z_{N,2}}{\exp(N\xi''(0)/2)} \xrightarrow{d} \exp(W).$$
(8.1)

Recall that the results in Section 7.1 only assume (2.21) rather than (1.5), and thus apply in the present proof. Let $\delta > 0$ be small and $T = T(\delta)$ as in (7.31), and recall the restricted partition function

$$Z_N(T) = \int_T \exp H_N(\boldsymbol{\sigma}) \, \mathrm{d}\mu_0(\boldsymbol{\sigma})$$

By (7.36), in Lemma 7.7, we have

$$\mathbb{E}[Z_N - Z_N(T)] \leqslant e^{-cN} \mathbb{E}[Z_N]$$

By Markov's inequality, applied respectively to the randomness of H_N and $H_{N,2}$, with probability $1 - e^{-cN}$,

$$(Z_N - Z_N(T)) \vee \mathbb{E}_{\geq 3}[Z_N - Z_N(T)] \leq e^{-cN} \mathbb{E}[Z_N].$$

By (7.70) (for k = 2), we also have, with probability 1 - o(1),

$$\mathbb{E}_{\geq 3}\left[\left(Z_N(T) - \mathbb{E}_{\geq 3}[Z_N(T)]\right)^2\right] = o(1) \mathbb{E}[Z_N]^2.$$

Thus with probability 1 - o(1),

$$|Z_N(T) - \mathbb{E}_{\geq 3} Z_N(T)| \leq o(1) \mathbb{E}[Z_N].$$

On the intersection of these events,

$$\left|\frac{Z_N}{\mathbb{E}Z_N} - \frac{\mathbb{E}_{\geq 3}Z_N}{\mathbb{E}Z_N}\right| \leq \frac{|Z_N - Z_N(T)|}{\mathbb{E}Z_N} + \frac{|Z_N(T) - \mathbb{E}_{\geq 3}Z_N(T)|}{\mathbb{E}Z_N} + \frac{|\mathbb{E}_{\geq 3}Z_N - \mathbb{E}_{\geq 3}Z_N(T)|}{\mathbb{E}Z_N} = o(1).$$

Since

$$\frac{\mathbb{E}_{\geq 3} Z_N}{\mathbb{E} Z_N} = \frac{Z_{N,2}}{\mathbb{E} Z_{N,2}}$$

the result follows from (8.1).

9 Completing the proof of Theorem 2.1

The following two propositions are the final ingredients in the proof of Theorem 2.1. Let δ, L be as in Algorithm 2 and $T' = \delta L$.

Proposition 9.1. Let $(H_N, y_{T'})$ be sampled from the marginal of the planted distribution \mathbb{P} (as defined in Eq. (3.2)). Let y^L be generated as in Algorithm 2, run on input H_N . Then,

$$\mathbb{E}_{H_N}\mathsf{TV}\left(\mathcal{L}(\boldsymbol{y}_{T'}|H_N),\mathcal{L}(\boldsymbol{y}^L|H_N)\right) = o_N(1).$$

Proposition 9.2. Let $(H_N, \sigma, y_{T'})$ be sampled from the marginal of the planted distribution \mathbb{P} . Let ρ^{MALA} be the (random) output of MALA run on $\tilde{\nu}_{H_N, y_{T'}}^{\text{proj}}$ (recall Eq. (2.10)) and $\hat{\sigma} = \sigma_{y_{T'}}(\rho^{MALA})$ (recall Eq. (2.6)). Then,

$$\mathbb{E}_{H_N,\boldsymbol{y}_{T'}}\mathsf{TV}\left(\mathcal{L}(\boldsymbol{\sigma}|H_N,\boldsymbol{y}_{T'}),\mathcal{L}(\widehat{\boldsymbol{\sigma}}|H_N,\boldsymbol{y}_{T'})\right) = o_N(1).$$

Proof of Theorem 2.1. Let $\mathcal{K} : \mathscr{H}_N \times \mathbb{R}^N \to \mathbb{R}^N$ be the random map that, given input (H_N, \boldsymbol{y}) , generates $\boldsymbol{\rho}^{\text{MALA}}$ by running MALA on $\nu_{H_N, \boldsymbol{y}}$ and outputs $\hat{\boldsymbol{\sigma}} = \boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho}^{\text{MALA}})$. Let $(H_N, \boldsymbol{\sigma}, \boldsymbol{y}_{T'})$ be sampled from the marginal of \mathbb{P} . Let $\mathbb{P}_{\text{alg}, H_N}$ denote the law of the output of \boldsymbol{y}^L generated by Algorithm 2 on input H_N . Then,

$$\begin{split} \mathbb{E}_{H_{N}\sim\mathbb{P}}\mathsf{TV}(\mu_{H_{N}},\mu^{\mathsf{alg}}) &= \mathbb{E}_{H_{N}\sim\mathbb{P}}\mathsf{TV}\left(\mathbb{E}_{\boldsymbol{y}_{T'}\sim\mathbb{P}(\cdot|H_{N})}\mathcal{L}(\boldsymbol{\sigma}|H_{N},\boldsymbol{y}_{T'}),\mathbb{E}_{\boldsymbol{y}^{L}\sim\mathbb{P}_{\mathsf{alg},H_{N}}}\mathcal{L}(\mathcal{K}(H_{N},\boldsymbol{y}^{L}))\right) \\ &\leq \mathbb{E}_{H_{N}\sim\mathbb{P}}\mathsf{TV}\left(\mathbb{E}_{\boldsymbol{y}_{T'}\sim\mathbb{P}(\cdot|H_{N})}\mathcal{L}(\boldsymbol{\sigma}|H_{N},\boldsymbol{y}_{T'}),\mathbb{E}_{\boldsymbol{y}_{T'}\sim\mathbb{P}(\cdot|H_{N})}\mathcal{L}(\mathcal{K}(H_{N},\boldsymbol{y}_{T'}))\right) \\ &+ \mathbb{E}_{H_{N}\sim\mathbb{P}}\mathsf{TV}\left(\mathbb{E}_{\boldsymbol{y}_{T'}\sim\mathbb{P}(\cdot|H_{N})}\mathcal{L}(\mathcal{K}(H_{N},\boldsymbol{y}_{T'})),\mathbb{E}_{\boldsymbol{y}^{L}\sim\mathbb{P}_{\mathsf{alg},H_{N}}}\mathcal{L}(\mathcal{K}(H_{N},\boldsymbol{y}^{L}))\right) \\ &\leq \mathbb{E}_{(H_{N},\boldsymbol{y}_{T'})\sim\mathbb{P}}\mathsf{TV}\left(\mathcal{L}(\boldsymbol{\sigma}|H_{N},\boldsymbol{y}_{T'}),\mathcal{L}(\mathcal{K}(H_{N},\boldsymbol{y}_{T'}))\right) \\ &+ \mathbb{E}_{H_{N}\sim\mathbb{P}}\mathsf{TV}\left(\mathcal{L}(\boldsymbol{y}_{T'}|H_{N}),\mathcal{L}(\boldsymbol{y}^{L}|H_{N})\right). \end{split}$$

The last inequality is by data processing. By Propositions 9.1 and 9.2, the final bound is $o_N(1)$. Thus, with probability $1 - o_N(1)$ over $H_N \sim \mathbb{P}$, $\mathsf{TV}(\mu_{H_N}, \mu^{\mathsf{alg}}) = o_N(1)$. By Corollary 3.5, the same is true for $H_N \sim \mathbb{Q}$.

9.1 TV-closeness of Euler discretization: Proof of Proposition 9.1

We prove Proposition 9.1 by an application of Girsanov's theorem, an approach introduced [CCL+22] in a related context. For all $0 \le l \le L - 1$, define

$$\widehat{\boldsymbol{m}}(\boldsymbol{y},\ell\delta) = \boldsymbol{m}^{\mathsf{alg}}(H_N,\boldsymbol{y},\ell\delta)$$

to be the output of Algorithm 1 with these inputs. Then, define the process $(\hat{y}_t)_{t \in [0,T]}$ by $\hat{y}_0 = 0$ and, for $t \in [\ell \delta, (\ell + 1)\delta)$,

$$\mathrm{d}\widehat{\boldsymbol{y}}_t = \widehat{\boldsymbol{m}}(\widehat{\boldsymbol{y}}_{\ell\delta}, \ell\delta) \,\mathrm{d}t + \mathrm{d}\boldsymbol{B}_t. \tag{9.1}$$

On each interval $[\ell\delta, (\ell+1)\delta)$, the drift in (9.1) is constant, so this SDE can be integrated directly: conditional on $H_N, \hat{y}_{\ell\delta}$,

$$\widehat{\boldsymbol{y}}_{(\ell+1)\delta} = \widehat{\boldsymbol{y}}_{\ell\delta} + \delta \boldsymbol{m}^{\mathsf{alg}}(H_N, \widehat{\boldsymbol{y}}_{\ell\delta}, \ell\delta) + \boldsymbol{B}_{(\ell+1)\delta} - \boldsymbol{B}_{\ell\delta}$$

Note that $B_{(\ell+1)\delta} - B_{\ell\delta} =_d \sqrt{\delta} w^{\ell}$ for $w^{\ell} \sim \mathcal{N}(0, I_N)$, so this is precisely the Euler discretization in Algorithm 2. It follows that

$$\mathcal{L}(\hat{\boldsymbol{y}}_T|H_N) = \mathcal{L}(\boldsymbol{y}^L|H_N).$$
(9.2)

Lemma 9.3. Given H_N , let $(\boldsymbol{y}_t)_{t \in [0,T]}$ be sampled from (1.3) and $(\hat{\boldsymbol{y}}_t)_{t \in [0,T]}$ be sampled from (9.1). Then,

$$\mathbb{E}_{H_N \sim \mathbb{P}} \mathsf{KL}(\mathcal{L}(\boldsymbol{y}_T | H_N), \mathcal{L}(\widehat{\boldsymbol{y}}_T | H_N)) \leqslant \frac{1}{2} \sum_{\ell=0}^{L-1} \int_{\ell\delta}^{(\ell+1)\delta} \mathbb{E}_{\mathbb{P}} \left\| \widehat{\boldsymbol{m}}(\boldsymbol{y}_{\ell\delta}, \ell\delta) - \boldsymbol{m}(\boldsymbol{y}_t, t) \right\|^2 \, \mathrm{d}t$$

Proof. Fix any realization of H_N . For $0 \le \ell \le L - 1$ and $t \in [\ell \delta, (\ell + 1)\delta)$, define the process

$$\boldsymbol{b}_t = \widehat{\boldsymbol{m}}(\boldsymbol{y}_{\ell\delta}, \ell\delta) - \boldsymbol{m}(\boldsymbol{y}_t, t)$$

Let

$$\mathcal{E}_t = \exp\left(\int_0^t \langle \boldsymbol{b}_s, \mathsf{d} \boldsymbol{B}_s
angle - rac{1}{2}\int_0^t \|\boldsymbol{b}_s\|^2 \,\,\mathsf{d} s
ight).$$

Let Q be the probability measure (conditional on H_N) under which $(\mathbf{B}_t)_{t \in [0,T]}$ is a Brownian motion and let P be the probability measure with $\frac{dP}{dQ} = \mathcal{E}_T$. By Girsanov's theorem [LG16, Theorem 5.22],

$$\boldsymbol{\beta}_t = \boldsymbol{B}_t - \int_0^t \boldsymbol{b}_s \, \mathrm{d}s$$

is a Brownian motion under *P*. (Since $\|\widehat{\boldsymbol{m}}(\boldsymbol{y}_{\ell\delta}, \ell\delta)\|$, $\|\boldsymbol{m}(\boldsymbol{y}_t, t)\| \leq \sqrt{N}$, \boldsymbol{b}_t is a.s. bounded, and thus the conditions of Girsanov's theorem are satisfied.) The SDE (1.3) rearranges as

$$\mathsf{d} \boldsymbol{y}_t = (\boldsymbol{m}(\boldsymbol{y}_t,t) + \boldsymbol{b}_t) \, \mathsf{d} t + \mathsf{d} \boldsymbol{\beta}_t = \widehat{\boldsymbol{m}}(\boldsymbol{y}_{\ell\delta},\ell\delta) \, \mathsf{d} t + \mathsf{d} \boldsymbol{\beta}_t, \qquad t \in [\ell\delta,(\ell+1)\delta).$$

Thus, under P, the law of $(\boldsymbol{y}_t)_{t \in [0,T]}$ is that of $(\hat{\boldsymbol{y}}_t)_{t \in [0,T]}$. By data processing,

$$\mathsf{KL}(\mathcal{L}(\boldsymbol{y}_T|H_N), \mathcal{L}(\widehat{\boldsymbol{y}}_T|H_N)) \leqslant \mathsf{KL}(Q, P) = \mathbb{E}_Q \log \frac{\mathsf{d}Q}{\mathsf{d}P} = \frac{1}{2} \int_0^T \mathbb{E}_Q \|\boldsymbol{b}_t\|^2 \, \mathsf{d}t.$$

The result follows by taking expectation over H_N .

Lemma 9.4. For all $0 \leq \ell \leq L - 1$, $t \in [\ell \delta, (\ell + 1)\delta)$, we have $\mathbb{E}_{\mathbb{P}} \|\widehat{\boldsymbol{m}}(\boldsymbol{y}_{\ell\delta}, \ell\delta) - \boldsymbol{m}(\boldsymbol{y}_t, t)\|^2 = o_N(1)$.

Proof. We first estimate

$$\mathbb{E}_{\mathbb{P}} \left\| \widehat{\boldsymbol{m}}(\boldsymbol{y}_{\ell\delta}, \ell\delta) - \boldsymbol{m}(\boldsymbol{y}_t, t) \right\|^2 \leq 2\mathbb{E}_{\mathbb{P}} \left\| \widehat{\boldsymbol{m}}(\boldsymbol{y}_{\ell\delta}, \ell\delta) - \boldsymbol{m}(\boldsymbol{y}_{\ell\delta}, \ell\delta) \right\|^2 + 2\mathbb{E}_{\mathbb{P}} \left\| \boldsymbol{m}(\boldsymbol{y}_{\ell\delta}, \ell\delta) - \boldsymbol{m}(\boldsymbol{y}_t, t) \right\|^2.$$

The first term on the right-hand side is $o_N(1)$ by Theorem 4.1, so it suffices to bound the second term. Recall that for $(H_N, \boldsymbol{x}, (\boldsymbol{y}_t)_{t \in [0,T]}) \sim \mathbb{P}$, conditional on (H_N, \boldsymbol{y}_t) the posterior law on \boldsymbol{x} is $\mu_t(\boldsymbol{\sigma}) \propto e^{H_{N,t}(\boldsymbol{\sigma})}$, for $H_{N,t}(\boldsymbol{\sigma})$ as in (4.1). Furthermore, for $s = t - \ell \delta, \boldsymbol{g} \sim \mathcal{N}(0, I_N)$,

$$H_{N,t}(\boldsymbol{\sigma}) = H_{N,\ell\delta}(\boldsymbol{\sigma}) + \langle s\boldsymbol{x} + \sqrt{s}\boldsymbol{g}, \boldsymbol{\sigma} \rangle$$

Let $\Delta_{t,\ell\delta}(\boldsymbol{\sigma}) = H_{N,t}(\boldsymbol{\sigma}) - H_{N,\ell\delta}(\boldsymbol{\sigma})$. With probability $1 - e^{-cN}$, $\|\boldsymbol{g}\| \leq 2\sqrt{N}$. Let \mathcal{E} denote this event. On \mathcal{E} ,

$$\sup_{\boldsymbol{\sigma}\in S_N} \|\boldsymbol{\Delta}_{t,\ell\delta}(\boldsymbol{\sigma})\| \leq \delta\sqrt{N} \, \|\boldsymbol{x}\| + \sqrt{\delta N} \, \|\boldsymbol{g}\| \leq 3\sqrt{\delta}N = 3/N.$$
(9.3)

So,

$$\begin{split} \boldsymbol{m}(\boldsymbol{y}_{\ell\delta},\ell\delta) - \boldsymbol{m}(\boldsymbol{y}_{t},t) &= \frac{\int_{S_{N}} \boldsymbol{\sigma} e^{H_{N,\ell\delta}(\boldsymbol{\sigma})}}{\int_{S_{N}} e^{H_{N,\ell\delta}(\boldsymbol{\sigma})}} - \frac{\int_{S_{N}} \boldsymbol{\sigma} e^{H_{N,t}(\boldsymbol{\sigma})}}{\int_{S_{N}} e^{H_{N,t}(\boldsymbol{\sigma})}} \\ &= \frac{\iint \boldsymbol{\sigma}^{1}(e^{H_{N,\ell\delta}(\boldsymbol{\sigma}^{1}) + H_{N,t}(\boldsymbol{\sigma}^{2})} - e^{H_{N,\ell\delta}(\boldsymbol{\sigma}^{2}) + H_{N,t}(\boldsymbol{\sigma}^{1})}) \, \mu_{0}^{\otimes 2}(\mathrm{d}\boldsymbol{\sigma})}{\iint e^{H_{N,\ell\delta}(\boldsymbol{\sigma}^{1}) + H_{N,t}(\boldsymbol{\sigma}^{2})} \, \mu_{0}^{\otimes 2}(\mathrm{d}\boldsymbol{\sigma})} \\ &= \frac{\iint \boldsymbol{\sigma}^{1}(e^{\boldsymbol{\Delta}_{t,\ell\delta}(\boldsymbol{\sigma}^{1})} - e^{\boldsymbol{\Delta}_{t,\ell\delta}(\boldsymbol{\sigma}^{2})}) e^{H_{N,\ell\delta}(\boldsymbol{\sigma}^{1}) + H_{N,\ell\delta}(\boldsymbol{\sigma}^{2})} \, \mu_{0}^{\otimes 2}(\mathrm{d}\boldsymbol{\sigma})}{\iint e^{\boldsymbol{\Delta}_{t,\ell\delta}(\boldsymbol{\sigma}^{2})} e^{H_{N,\ell\delta}(\boldsymbol{\sigma}^{1}) + H_{N,\ell\delta}(\boldsymbol{\sigma}^{2})} \, \mu_{0}^{\otimes 2}(\mathrm{d}\boldsymbol{\sigma})}. \end{split}$$

By (9.3),

$$\left\|\boldsymbol{\sigma}^{1}\right\|\left|e^{\boldsymbol{\Delta}_{t,\ell\delta}(\boldsymbol{\sigma}^{1})}-e^{\boldsymbol{\Delta}_{t,\ell\delta}(\boldsymbol{\sigma}^{2})}\right|=O(N^{-1/2})$$

$$\mathbb{E}_{\mathbb{P}} \|\boldsymbol{m}(\boldsymbol{y}_{\ell\delta}, \ell\delta) - \boldsymbol{m}(\boldsymbol{y}_t, t)\|^2 \leq \mathbb{E}_{\mathbb{P}} \mathbf{1}\{\mathcal{E}\} \|\boldsymbol{m}(\boldsymbol{y}_{\ell\delta}, \ell\delta) - \boldsymbol{m}(\boldsymbol{y}_t, t)\|^2 + \mathbb{E}_{\mathbb{P}} \mathbf{1}\{\mathcal{E}^c\} \|\boldsymbol{m}(\boldsymbol{y}_{\ell\delta}, \ell\delta) - \boldsymbol{m}(\boldsymbol{y}_t, t)\|^2 \\ \leq O(N^{-1/2}) + e^{-cN} \cdot 4N = o_N(1).$$

п	-	

Proof of Proposition 9.1. By (9.2) and Lemmas 9.3 and 9.4,

$$\mathbb{E}_{H_N \sim \mathbb{P}} \mathsf{KL}(\mathcal{L}(\boldsymbol{y}_T | H_N), \mathcal{L}(\boldsymbol{y}^L | H_N)) = o_N(1).$$

The result follows from Pinsker's inequality and Jensen's inequality:

$$\mathbb{E}_{H_N \sim \mathbb{P}} \mathsf{KL}(\mathcal{L}(\boldsymbol{y}_T | H_N), \mathcal{L}(\boldsymbol{y}^L | H_N)) \ge 2\mathbb{E}_{H_N \sim \mathbb{P}} \left[\mathsf{TV}(\mathcal{L}(\boldsymbol{y}_T | H_N), \mathcal{L}(\boldsymbol{y}^L | H_N))^2 \right] \ge 2 \left[\mathbb{E}_{H_N \sim \mathbb{P}} \mathsf{TV}(\mathcal{L}(\boldsymbol{y}_T | H_N), \mathcal{L}(\boldsymbol{y}^L | H_N)) \right]^2.$$

9.2 Log-concavity of late measures

In this subsection, we prove Proposition 9.2. Let e_1, \ldots, e_N be the standard basis. By a change of coordinates, we may assume without loss of generality that $\hat{y} = y/||y||_N = e_N \sqrt{N}$ and $U = (e_1, \ldots, e_{N-1})$.

Lemma 9.5. For any $y \neq 0$, the push-forward of $\mu_{H_N,y}(\cdot |\langle \sigma, y \rangle > 0)$ under the stereographic projection T_y is $\nu_{H_N,y}^{\text{proj}}$, defined in (2.8).

Proof. Note that (denoting by DF the Jacobian of map F):

$$\boldsymbol{D}\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})^{\top} = \frac{[\boldsymbol{I}_{N-1}, \boldsymbol{0}]}{\sqrt{1 + \|\boldsymbol{\rho}\|_{N}^{2}}} - \frac{\boldsymbol{\rho}\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})^{\top}/N}{1 + \|\boldsymbol{\rho}\|_{N}^{2}}$$

Since $[I_{N-1}, \mathbf{0}] \sigma_y(\rho) = \frac{\rho}{\sqrt{1+\|\rho\|_N^2}}$, we have

$$\boldsymbol{D}\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})^{\top}\boldsymbol{D}\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho}) = \frac{\boldsymbol{I}_{N-1}}{1 + \|\boldsymbol{\rho}\|_{N}^{2}} - \frac{\boldsymbol{\rho}\boldsymbol{\rho}^{\top}/N}{(1 + \|\boldsymbol{\rho}\|_{N}^{2})^{2}} = \frac{\boldsymbol{I}_{N-1}}{1 + \|\boldsymbol{\rho}\|_{N}^{2}} \left(\boldsymbol{I}_{N-1} - \frac{\boldsymbol{\rho}\boldsymbol{\rho}^{\top}/N}{1 + \|\boldsymbol{\rho}\|_{N}^{2}}\right)$$

The stereographic projection thus incurs a change of density factor of

$$\det(\boldsymbol{D}\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})^{\top}\boldsymbol{D}\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho}))^{1/2} = (1 + \|\boldsymbol{\rho}\|_{N}^{2})^{-N/2}.$$

This precisely accounts for the term $-\frac{N}{2}\log(1+\|\boldsymbol{\rho}\|_N^2)$ in (2.7).

Lemma 9.6. For sufficiently large T, with probability $1 - o_N(1)$ over (H_N, \boldsymbol{y}_T) as in Proposition 9.2, $\nu_{H_N, \boldsymbol{y}_T}^{\text{proj}}(\|\boldsymbol{\rho}\|_N^2 \leq \varepsilon_0) = 1 - o_N(1)$ and $\mu_{H_N, \boldsymbol{y}_T}(\langle \boldsymbol{\sigma}, \boldsymbol{y}_T \rangle_N \leq 0) = o_N(1)$.

Proof. Let $(H_N, \boldsymbol{x}, \boldsymbol{y}_T)$ be a sample from \mathbb{P} , and let $q_* = q_*(T)$ be as in Fact 4.2. Note that $q_* > 1 - \frac{1}{T}$, as

$$\xi'_T(1-1/T) \ge T + \xi'_T(1-1/T) \ge T > T - 1 = \frac{1-1/T}{1/T}.$$

By Proposition 5.12, with probability $1 - o_N(1)$,

$$\mu_{H_N,\boldsymbol{y}_T}(\langle \boldsymbol{\sigma}, \boldsymbol{x} \rangle_N \ge 1 - 1/T) = 1 - o_N(1).$$

With probability $1 - o_N(1)$, we have $\|\boldsymbol{y}\|_N = \sqrt{T(T+1)} + o_N(1)$, so

$$\langle \boldsymbol{x}, \boldsymbol{\hat{y}} \rangle = \frac{\langle \boldsymbol{x}, \boldsymbol{y} \rangle}{\|\boldsymbol{y}\|_N} = \sqrt{1 - \frac{1}{T+1}} + o_N(1).$$

On this event, $\{ \boldsymbol{\sigma} \in S_N : \langle \boldsymbol{\sigma}, \boldsymbol{x} \rangle_N \ge 1 - 1/T \} \subseteq \{ \boldsymbol{\sigma} \in S_N : \langle \boldsymbol{\sigma}, \hat{\boldsymbol{y}} \rangle_N \ge 1 - 2/T \}$. So, with probability $1 - o_N(1)$,

$$\mu_{H_N,\boldsymbol{y}_T}(\langle \boldsymbol{\sigma}, \boldsymbol{\hat{y}} \rangle_N \ge 1 - 2/T) = 1 - o_N(1).$$

(This of course implies $\mu_{H_N, \boldsymbol{y}_T}(\langle \boldsymbol{\sigma}, \boldsymbol{y}_T \rangle_N \leq 0) = o_N(1)$.) For sufficiently large T, the stereographic projection $T_{\boldsymbol{y}}$ maps $\{\boldsymbol{\sigma} \in S_N : \langle \boldsymbol{\sigma}, \hat{\boldsymbol{y}} \rangle_N \geq 1 - 2/T\}$ into $\{\boldsymbol{\rho} \in \mathbb{R}^{N-1} : \|\boldsymbol{\rho}\|_N^2 \leq \varepsilon_0\}$. The conclusion follows from Lemma 9.5.

Corollary 9.7. Recall definition (2.10) of $\nu_{H_N, \boldsymbol{y}_T}^{\text{proj}}$, $\widetilde{\nu}_{H_N, \boldsymbol{y}_T}^{\text{proj}}$. For sufficiently large T, with probability $1 - o_N(1)$ over (H_N, \boldsymbol{y}_T) , $\mathsf{TV}(\nu_{H_N, \boldsymbol{y}_T}^{\text{proj}}, \widetilde{\nu}_{H_N, \boldsymbol{y}_T}^{\text{proj}}) = o_N(1)$.

Proof. Since $\varphi(x) = 0$ for $x \in [0, \varepsilon_0]$, and $\varphi(x) \ge 0$ for $x > \varepsilon_0$, we have

$$\begin{split} & \int_{\|\boldsymbol{\rho}\|_{N}^{2} \leqslant \varepsilon_{0}} \exp \widetilde{H}_{N,\boldsymbol{y}_{T}}^{\mathsf{proj}}(\boldsymbol{\rho}) \, \mathrm{d}\boldsymbol{\rho} = \int_{\|\boldsymbol{\rho}\|_{N}^{2} \leqslant \varepsilon_{0}} \exp H_{N,\boldsymbol{y}_{T}}^{\mathsf{proj}}(\boldsymbol{\rho}) \, \mathrm{d}\boldsymbol{\rho}, \\ & \int_{\|\boldsymbol{\rho}\|_{N}^{2} > \varepsilon_{0}} \exp \widetilde{H}_{N,\boldsymbol{y}_{T}}^{\mathsf{proj}}(\boldsymbol{\rho}) \, \mathrm{d}\boldsymbol{\rho} \leqslant \int_{\|\boldsymbol{\rho}\|_{N}^{2} > \varepsilon_{0}} \exp H_{N,\boldsymbol{y}_{T}}^{\mathsf{proj}}(\boldsymbol{\rho}) \, \mathrm{d}\boldsymbol{\rho}. \end{split}$$

Combined with Lemma 9.6, it follows that with probability $1 - o_N(1)$,

$$\widetilde{\nu}_{H_N,\boldsymbol{y}_T}^{\text{proj}}(\|\boldsymbol{\rho}\|_N^2 \leqslant \varepsilon_0) \geqslant \nu_{H_N,\boldsymbol{y}_T}^{\text{proj}}(\|\boldsymbol{\rho}\|_N^2 \leqslant \varepsilon_0) \geqslant 1 - o_N(1).$$

Since $\tilde{\nu}_{H_N, \boldsymbol{y}_T}^{\text{proj}}$ and $\nu_{H_N, \boldsymbol{y}_T}^{\text{proj}}$ are furthermore proportional on $\{\|\boldsymbol{\rho}\|_N^2 \leq \varepsilon_0\}$, the conclusion follows.

Proposition 9.8. For sufficiently large T, there exist $C_{\min}, C_{\max} > 0$ (depending on T) such that with probability $1 - o_N(1)$, for all $\rho \in \mathbb{R}^{N-1}$,

$$-C_{\max}\boldsymbol{I}_{N-1} \leq \nabla^2 \widetilde{H}_{N,\boldsymbol{y}_T}^{\mathsf{proj}}(\boldsymbol{\rho}) \leq -C_{\min}\boldsymbol{I}_{N-1}$$

Proof. Let $\boldsymbol{y} = \boldsymbol{y}_T \, \hat{\boldsymbol{y}} = \boldsymbol{y}/\|\boldsymbol{y}\|_N$, and assume without loss of generality $\hat{\boldsymbol{y}} = \sqrt{N} \boldsymbol{e}_N$. Let $\boldsymbol{U}^\top = [\boldsymbol{I}_{N-1}, \boldsymbol{0}] \in \mathbb{R}^{(N-1)\times N}$ be the projection onto the orthogonal complement of $\hat{\boldsymbol{y}}$.

A direct calculation shows

$$\begin{split} \nabla^{2} \widetilde{H}_{N,\boldsymbol{y}}^{\mathsf{proj}}(\boldsymbol{\rho}) &= \frac{\langle \nabla H_{N,\boldsymbol{y}}(\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})), \boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho}) \rangle}{N(1+\|\boldsymbol{\rho}\|_{N}^{2})} \left(-\boldsymbol{I}_{N-1} + \frac{3\boldsymbol{\rho}\boldsymbol{\rho}^{\top}}{N(1+\|\boldsymbol{\rho}\|_{N}^{2})} \right) \\ &+ \frac{\langle \nabla^{2} H_{N,\boldsymbol{y}}(\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})), \boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho}) \otimes^{2} \rangle}{N(1+\|\boldsymbol{\rho}\|_{N}^{2})^{2}} \cdot \frac{\boldsymbol{\rho}\boldsymbol{\rho}^{\top}}{N} + \frac{\boldsymbol{U}^{\top} \nabla^{2} H_{N,\boldsymbol{y}}(\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})) \boldsymbol{U}}{1+\|\boldsymbol{\rho}\|_{N}^{2}} \\ &- \frac{\boldsymbol{\rho}\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})^{\top} \nabla^{2} H_{N,\boldsymbol{y}}(\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})) \boldsymbol{U} + \boldsymbol{U}^{\top} \nabla^{2} H_{N,\boldsymbol{y}}(\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})) \boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho}) \boldsymbol{\rho}^{\top}}{N(1+\|\boldsymbol{\rho}\|_{N}^{2})} \\ &- \frac{\boldsymbol{\rho} \nabla H_{N,\boldsymbol{y}}(\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho}))^{\top} \boldsymbol{U} + \boldsymbol{U}^{\top} \nabla H_{N,\boldsymbol{y}}(\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})) \boldsymbol{\rho}^{\top}}{N(1+\|\boldsymbol{\rho}\|_{N}^{2})^{3/2}} \\ &- \left(T \varphi'(\|\boldsymbol{\rho}\|_{N}^{2}) + \frac{1}{1+\|\boldsymbol{\rho}\|_{N}^{2}}\right) \boldsymbol{I}_{N-1} - \left(T \varphi''(\|\boldsymbol{\rho}\|_{N}^{2}) - \frac{1}{(1+\|\boldsymbol{\rho}\|_{N}^{2})^{2}}\right) \frac{2\boldsymbol{\rho}\boldsymbol{\rho}^{\top}}{N} \end{split}$$

By Proposition 3.6, there exists C > 0 (independent of T) such that with probability $1 - o_N(1)$,

$$\sup_{\boldsymbol{\sigma}\in S_N} \|\nabla H_N(\boldsymbol{\sigma})\|_N, \sup_{\boldsymbol{\sigma}\in S_N} \|\nabla^2 H_N(\boldsymbol{\sigma})\|_{_{\mathrm{op}}} \leqslant C.$$

We will show that on this event,

$$\nabla^{2} \widetilde{H}_{N,\boldsymbol{y}}^{\mathsf{proj}}(\boldsymbol{\rho}) = \frac{\|\boldsymbol{y}\|_{N}}{(1+\|\boldsymbol{\rho}\|_{N}^{2})^{3/2}} \left(-\boldsymbol{I}_{N-1} + \frac{3\boldsymbol{\rho}\boldsymbol{\rho}^{\top}}{N(1+\|\boldsymbol{\rho}\|_{N}^{2})} \right) - T\varphi'(\|\boldsymbol{\rho}\|_{N}^{2})\boldsymbol{I}_{N-1} - T\varphi''(\|\boldsymbol{\rho}\|_{N}^{2}) \cdot \frac{2\boldsymbol{\rho}\boldsymbol{\rho}^{\top}}{N} + O(1),$$
(9.4)

where O(1) denotes a matrix of operator norm O(1), independent of T. Note that

$$\frac{\langle \nabla H_{N,\boldsymbol{y}}(\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})), \boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho}) \rangle}{N(1+\|\boldsymbol{\rho}\|_{N}^{2})} = \frac{\langle \nabla H_{N}(\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})) + \boldsymbol{y}, \boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho}) \rangle}{N(1+\|\boldsymbol{\rho}\|_{N}^{2})} = \frac{\langle \nabla H_{N}(\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})), \boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho}) \rangle}{N(1+\|\boldsymbol{\rho}\|_{N}^{2})} + \frac{\|\boldsymbol{y}\|_{N}}{(1+\|\boldsymbol{\rho}\|_{N}^{2})^{3/2}}.$$

The first term on the right-hand side is bounded independently of T, as

$$\frac{|\langle \nabla H_N(\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})), \boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho}) \rangle|}{N} \leq \|\nabla H_N(\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho}))\|_N \|\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})\|_N$$

Similarly, all other terms in the expansion of $\nabla^2 \widetilde{H}_{N,\boldsymbol{y}}^{\mathsf{proj}}(\boldsymbol{\rho})$ above, aside from $T\varphi'(\|\boldsymbol{\rho}\|_N^2)\boldsymbol{I}_{N-1}$ and $T\varphi''(\|\boldsymbol{\rho}\|_N^2)$. $\frac{2\rho\rho^{\top}}{N}$, are bounded independently of T, due to the following inequalities:

$$\begin{aligned} \left\| \nabla^{2} H_{N,\boldsymbol{y}}(\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})) \right\|_{\text{op}} &= \left\| \nabla^{2} H_{N}(\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})) \right\|_{\text{op}} = O(1), \\ \frac{\left\| \boldsymbol{U}^{\top} \nabla H_{N,\boldsymbol{y}}(\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})) \boldsymbol{\rho}^{\top} \right\|_{\text{op}}}{N} &\leq \left\| \boldsymbol{U}^{\top} \nabla H_{N,\boldsymbol{y}}(\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})) \right\|_{N} \|\boldsymbol{\rho}\|_{N} \\ &= \left\| \boldsymbol{U}^{\top} \nabla H_{N}(\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})) \right\|_{N} \|\boldsymbol{\rho}\|_{N} \leq O(1) \|\boldsymbol{\rho}\|_{N} \end{aligned}$$

and $\|\boldsymbol{\rho}\boldsymbol{\rho}^{\top}\|_{\text{op}}/N = \|\boldsymbol{\rho}\|_{N}^{2}, \|\boldsymbol{\rho}\boldsymbol{\sigma}_{\boldsymbol{y}}(\boldsymbol{\rho})^{\top}\|_{\text{op}}/N = \|\boldsymbol{\rho}\|_{N}$. (Note that each of these terms, each copy of $\|\boldsymbol{\rho}\|_{N}^{2}$ in the resulting bound is compensated by at least one copy of $1 + \|\boldsymbol{\rho}\|_N^2$ in the denominator.) This proves (9.4). With probability $1 - o_N(1)$, we have $\|\boldsymbol{y}\|_N = \sqrt{T(T+1)} + o_N(1)$. On this event, (9.4) yields

$$\nabla^2 \tilde{H}_{N,\boldsymbol{y}}^{\text{proj}}(\boldsymbol{\rho}) = T(-\boldsymbol{M}(\boldsymbol{\rho}) + o_T(1)),$$

where $o_T(1)$ denotes a matrix with operator norm vanishing with T and

$$\boldsymbol{M}(\boldsymbol{\rho}) = \frac{\boldsymbol{I}_{N-1}}{(1+\|\boldsymbol{\rho}\|_N^2)^{3/2}} - \frac{3\boldsymbol{\rho}\boldsymbol{\rho}^\top}{N(1+\|\boldsymbol{\rho}\|_N^2)^{5/2}} + \varphi'(\|\boldsymbol{\rho}\|_N^2)\boldsymbol{I}_{N-1} + \varphi''(\|\boldsymbol{\rho}\|_N^2) \cdot \frac{2\boldsymbol{\rho}\boldsymbol{\rho}^\top}{N}.$$

From this it is clear that $-C_{\max} I_{N-1} \leq \nabla^2 \widetilde{H}_{N, y_T}^{\text{proj}}(\rho)$ for suitable C_{\max} . For the other direction, note that $M(\rho)$ has eigenvalue $\frac{1}{(1+\|\rho\|_N^2)^{3/2}} + \varphi'(\|\rho\|_N^2)$ in all directions orthogonal to ρ , and

$$\frac{1-2\|\boldsymbol{\rho}\|_{N}^{2}}{(1+\|\boldsymbol{\rho}\|_{N}^{2})^{5/2}}+\varphi'(\|\boldsymbol{\rho}\|_{N}^{2})+2\|\boldsymbol{\rho}\|_{N}^{2}\varphi''(\|\boldsymbol{\rho}\|_{N}^{2})$$

in the direction of ρ . By (2.9), $M(\rho) \geq \varepsilon_0 I_{N-1}$, and thus $\nabla^2 \widetilde{H}_{N, y_T}^{\text{proj}}(\rho) \leq -C_{\min} I_{N-1}$ for $C_{\min} =$ $T\varepsilon_0/2.$

Finally, we verify that φ satisfying (2.9) exists.

Fact 9.9. For suitable C > 0, the function

$$\varphi(x) = C\mathbf{1}\{x > \varepsilon_0\} \left(x - \frac{\varepsilon_0^2}{x} - 2\varepsilon_0 \log \frac{x}{\varepsilon_0}\right)$$

is nonnegative, twice continuously differentiable, and satisfies (2.9).

Proof. Note that for $x > \varepsilon_0$,

$$\varphi'(x) = C\left(1 - \frac{\varepsilon_0}{x}\right)^2, \qquad \qquad \varphi''(x) = \frac{2C\varepsilon}{x^2}\left(1 - \frac{\varepsilon_0}{x}\right).$$

Thus $\lim_{x\downarrow \varepsilon_0} \varphi''(x) = 0$, so φ is twice continuously differentiable. Note that $\varphi' \ge 0$, so integrating shows $\varphi \ge 0$. Let

$$C_0 = \min_{x \ge 0} \frac{1 - 2x}{(1 + x)^{5/2}}$$

and set C so that $C_0 + \varphi'(2\varepsilon_0) \ge \varepsilon_0$. Note $\varphi'' \ge 0$, and thus φ' is increasing; thus (2.9) holds for all $x \ge 2\varepsilon_0$. For all $x \in [0, 2\varepsilon_0]$, we verify that

$$\frac{1}{(1+x)^{3/2}} \ge \frac{1-2x}{(1+x)^{5/2}} \ge \frac{1-4\varepsilon_0}{(1+2\varepsilon_0)^{5/2}} \ge \varepsilon_0,$$

so (2.9) holds.

Proof of Proposition 9.2. By Proposition 9.8, $\tilde{\nu}_{H_N, \boldsymbol{y}_T}^{\text{proj}}$ is O(1)-smooth and strongly log-concave. By [CLA+21, Theorem 3], MALA run for time $\chi_{\text{log-conc}} = \text{poly}(N)$ outputs $\boldsymbol{\rho}^{\text{MALA}} \sim \nu^{\text{MALA}}$, where $\text{TV}(\nu^{\text{MALA}}, \tilde{\nu}_{H_N, \boldsymbol{y}_T}^{\text{proj}}) \leq 1/N$. Combined with Corollary 9.7, we find that (with probability $1 - o_N(1)$), $\text{TV}(\nu_{H_N, \boldsymbol{y}_T}^{\text{proj}}, \nu^{\text{MALA}}) = o_N(1)$. Lemma 9.5 completes the proof.

10 Failure of stochastic localization in complementary regime

In this section, we prove Theorem 2.3. Similarly to Subsection 3.2, we may analyze the process (2.17) by passing to a planted model. For any T > 0, let $\tilde{\mathbb{P}}, \tilde{\mathbb{Q}} \in \mathcal{P}(S_N \times \mathscr{H}_N \times C([0,T], \mathbb{R}^N \times \cdots \times (\mathbb{R}^N)^{\otimes J}))$ be the laws of $(\sigma, H_N, (\vec{y}_t)_{t \in [0,T]})$, generated as follows.

• Under $\check{\mathbb{Q}}$,

$$H_N \sim \mu_{\mathsf{null}}, \qquad \boldsymbol{\sigma} \sim \mu_{H_N}, \qquad \boldsymbol{y}_t^j = \tau_j(t) \boldsymbol{\sigma}^{\otimes j} + \boldsymbol{B}_{\tau_j(t)}^j, \quad \forall j = 1, \dots, J$$

for $(\boldsymbol{B}_t^1, \ldots, \boldsymbol{B}_t^J)_{t \ge 0}$ independent of $(H_N, \boldsymbol{\sigma})$. Equivalently, $H_N \sim \mu_{\text{null}}, (\boldsymbol{\vec{y}}_t)_{t \ge 0}$ is given by the SDE (2.17), and for any odd j such that $\lim_{t\to\infty} \tau_j(t) = \infty, \boldsymbol{\sigma}$ is the unique solution to $\boldsymbol{\sigma}^{\otimes j} = \lim_{t\to\infty} \boldsymbol{y}_t^j / \tau_j(t)$.

• Under Ď,

$$(H_N, \boldsymbol{\sigma}) \sim \mu_{\mathsf{pl}}, \qquad \boldsymbol{y}_t^j = \tau_j(t) \boldsymbol{\sigma}^{\otimes j} + \boldsymbol{B}_{\tau_j(t)}^j, \quad \forall j = 1, \dots, J$$

for $(\boldsymbol{B}_t^1, \ldots, \boldsymbol{B}_t^J)_{t \ge 0}$ independent of $(H_N, \boldsymbol{\sigma})$. Equivalently, we can generate first H_N , then $(\vec{\boldsymbol{y}}_t)_{t \ge 0}$ by (2.17), and finally $\boldsymbol{\sigma}$ as above. Furthermore, the law of $(H_N, \boldsymbol{\sigma}) \sim \mu_{\mathsf{pl}}$ can be described by either (3.3) or (3.4).

Analogously to Proposition 3.4, we have

$$\frac{\mathrm{d}\tilde{\mathbb{P}}}{\mathrm{d}\tilde{\mathbb{Q}}}(\boldsymbol{\sigma}, H_N, (\boldsymbol{\vec{y}}_t)_{t \in [0,T]}) = \frac{Z(H_N)}{\mathbb{E}Z(H_N)},$$

and this ratio is tight by Lemma 3.2. Thus $\check{\mathbb{P}}$ and $\check{\mathbb{Q}}$ are mutually contiguous.

Therefore, it suffices to analyze the AMP iteration (2.20) under \mathbb{P} . Similarly to (4.1), we find that conditional on \vec{y}_t , the posterior law of σ under \mathbb{P} is

$$\check{\mu}_t(\mathsf{d}\boldsymbol{\sigma}) = \frac{1}{Z} \exp \check{H}_{N,t}(\boldsymbol{\sigma}) \mu_0(\mathsf{d}\boldsymbol{\sigma})$$

where

$$\check{H}_{N,t}(\boldsymbol{\sigma}) = N\xi(\langle \boldsymbol{x}, \boldsymbol{\sigma} \rangle_N) + \tilde{H}_N(\boldsymbol{\sigma}) + \sum_{j=1}^J \frac{1}{N^{j-1}} \langle \boldsymbol{y}_t^j, \boldsymbol{\sigma}^{\otimes j} \rangle \stackrel{d}{=} N\check{\xi}_t(\langle \boldsymbol{x}, \boldsymbol{\sigma} \rangle_N) + \tilde{H}_{N,t}(\boldsymbol{\sigma}),$$

-		
I		1
I		I
		I

for $\widetilde{H}_{N,t}$ a spin glass with mixture

$$\check{\xi}_t(q) = \xi(q) + \sum_{j=1}^J \tau_j(t)^2 q^j$$
.

Let $q_{\text{AMP}} = q_{\text{AMP}}(t)$ be the smallest solution to $\check{\xi}'_t(q) = \frac{q}{1-q}$. Note that a solution exists because $\check{\xi}'_t(0) \ge 0$ and $\lim_{q \uparrow 1} \frac{q}{1-q} = +\infty$.

Proposition 10.1. We have

$$\lim_{k \to \infty} \operatorname{p-lim}_{N \to \infty} \langle \boldsymbol{x}, \widecheck{\boldsymbol{m}}^k \rangle_N = \lim_{k \to \infty} \operatorname{p-lim}_{N \to \infty} \langle \widecheck{\boldsymbol{m}}^k, \widecheck{\boldsymbol{m}}^k \rangle_N = q_{\mathsf{AMP}}.$$

Consequently, for all $1 \leq j \leq J$ *,*

$$\lim_{k \to \infty} \lim_{N \to \infty} \mathbb{E} \frac{1}{N^j} \left\| \boldsymbol{x}^{\otimes j} - (\widecheck{\boldsymbol{m}}^k)^{\otimes j} \right\|_2^2 = 1 - q_{\text{AMP}}^j.$$

Proof. Since $q \mapsto \frac{\xi'_t(q)}{1+\xi'_t(q)}$ is increasing, the sequence $(\check{q}_k)_{k\geq 0}$ defined in (2.19) is increasing. Furthermore, if $\check{q}_k \leq q_{AMP}$, then

$$\breve{q}_{k+1} = \frac{\breve{\xi}_t'(\breve{q}_k)}{1+\breve{\xi}_t'(\breve{q}_k)} \leqslant \frac{\breve{\xi}_t'(q_{\mathsf{AMP}})}{1+\breve{\xi}_t'(q_{\mathsf{AMP}})} = q_{\mathsf{AMP}},$$

and therefore by induction $(\check{q}_k)_{k\geq 0}$ is bounded above by q_{AMP} . As the limit of $(\check{q}_k)_{k\geq 0}$ must be a fixed point of $q \mapsto \frac{\check{\xi}'_t(q)}{1+\check{\xi}'_t(q)}$, we have $\lim_{k\to\infty} \check{q}_k = q_{AMP}$. By state evolution, similarly to the proof of Proposition 4.3, the first conclusion follows. Since

$$\frac{1}{N^j} \left\| \boldsymbol{x}^{\otimes j} - (\widetilde{\boldsymbol{m}}^k)^{\otimes j} \right\|_2^2 = \langle \boldsymbol{x}, \boldsymbol{x} \rangle_N^j - 2 \langle \boldsymbol{x}, \widetilde{\boldsymbol{m}}^k \rangle_N^j + \langle \widetilde{\boldsymbol{m}}^k, \widetilde{\boldsymbol{m}}^k \rangle_N^j,$$

the second conclusion follows from the first.

Let

$$Q_{\text{bayes}} = Q_{\text{bayes}}(t) = \arg\max_{q \in [0,1)} \left\{ \check{\xi}_t(q) + q + \log(1-q) \right\} \subseteq [0,1)$$
(10.1)

be the set of all maximizers of this quantity, and let

$$q_{\text{bayes}} = q_{\text{bayes}}(t) = \inf Q_{\text{bayes}}(t).$$

Lemma 10.2. For any t, the equation $\check{\xi}'_t(q) = \frac{q}{1-q}$ has finitely many solutions $q \in [0,1)$. Moreover, $Q_{\text{bayes}}(t)$ is a finite set for all t. If $T_1 \subseteq [0, +\infty)$ is the set of t_1 such that $|Q_{\text{bayes}}(t_1)| > 1$, then for each $t_1 \in T_1$, there exists $\delta > 0$ such that $(t_1 - \delta, t_1 + \delta) \cap T_1 = \{t_1\}$.

Proof. Let $f_t(q) = (1-q)\check{\xi}'_t(q) - q$, so any solution to $\check{\xi}'_t(q) = \frac{q}{1-q}$ is a zero of f_t . Note that f_t is not identically zero: if it were, then $\check{\xi}'_t(q) = \frac{q}{1-q}$, contradicting that the coefficients γ_p^2 of ξ satisfy $\sum_{p \ge 2} 2^p \gamma_p^2 < \infty$. Since f_t is complex analytic in the unit disc, its zero set has no limit point, and in particular it has finitely many zeros in [0, 1). This shows that there are finitely many solutions to $\check{\xi}'_t(q) = \frac{q}{1-q}$.

Note that $\frac{d}{dq}(\check{\xi}_t(q) + q + \log(1-q)) = \check{\xi}'_t(q) - \frac{q}{1-q}$. Any interior maximizer of (10.1) must therefore satisfy the stationarity condition $\check{\xi}'_t(q) = \frac{q}{1-q}$. Since $\check{\xi}'_t(0) \ge 0$, 0 can be a maximizer only if it also solves this equation. Thus $Q_{\text{bayes}}(t)$ is finite.

Consider an arbitrary $t_1 \in T_1$ and let $Q = Q_{\text{bayes}}(t_1)$. For each $\tilde{q} \in Q$, let $I_{\tilde{q}} = [\tilde{q} - \varepsilon, \tilde{q} + \varepsilon]$, where $\varepsilon > 0$ is small enough that these intervals do not overlap. By continuity, for sufficiently small δ and all $t \in (t_1 - \delta, t_1 + \delta)$, all maximizers of $\check{\xi}_t(q) + q + \log(1 - q)$ lie in $\bigcup_{\tilde{a} \in Q} I_{\tilde{q}}$. Let

$$m(t,\tilde{q}) = \max_{q \in I_{\tilde{q}}} \left\{ \check{\xi}_t(q) + q + \log(1-q) \right\}, \qquad q(t,\tilde{q}) = \arg\max_{q \in I_{\tilde{q}}} \left\{ \check{\xi}_t(q) + q + \log(1-q) \right\}.$$

Note that $q(t_1, \tilde{q}) = \tilde{q}$ for each $\tilde{q} \in Q$. Since the maximum of $\check{\xi}_{t_1}(q) + q + \log(1-q)$ is attained over $I_{\tilde{q}}$ uniquely at \tilde{q} , by continuity $\lim_{t \to t_1} q(t, \tilde{q}) = \tilde{q}$.

For $\widetilde{q} \in Q$, $t \in (t_1, t_1 + \delta)$, we have

$$\frac{m(t,\tilde{q}) - m(t_1,\tilde{q})}{t - t_1} \ge \frac{\check{\xi}_t(q(t_1,\tilde{q})) - \check{\xi}_{t_1}(q(t_1,\tilde{q}))}{t - t_1} = \sum_{j=1}^J \tau'_j(t_1)q(t_1,\tilde{q})^j + O(t - t_1),$$
$$\frac{m(t,\tilde{q}) - m(t_1,\tilde{q})}{t - t_1} \le \frac{\check{\xi}_t(q(t,\tilde{q})) - \check{\xi}_{t_1}(q(t,\tilde{q}))}{t - t_1} = \sum_{j=1}^J \tau'_j(t_1)q(t,\tilde{q})^j + O(t - t_1).$$

Taking the limit $t \downarrow t_1$ yields

$$\lim_{t \downarrow t_1} \frac{m(t, \widetilde{q}) - m(t_1, \widetilde{q})}{t - t_1} = \sum_{j=1}^J \tau'_j(t_1) \widetilde{q}^j.$$

A similar argument shows the left-derivative is also equal to this. Therefore

$$\frac{\partial}{\partial t}m(t,\widetilde{q})\big|_{t=t_1} = \sum_{j=1}^J \tau'_j(t_1)\widetilde{q}^j$$

This quantity is distinct for different $\tilde{q} \in Q$. Therefore, for all $t \in (t_1 - \delta, t_1 + \delta) \setminus \{t_1\}, |Q_{bayes}(t)| = 1.$

Proposition 10.3. Suppose $t \notin T_1$ satisfies $q_{\text{bayes}}(t) > 0$. Let $\check{\xi}_t(q) = \sum_{p \ge 1} \beta_p^2 q^p$ (where we suppress the dependence of the β_p on t). For any p such that $\beta_p > 0$, we have (recall the definition of $m_p(\vec{y}_t, t)$ in Eq. (2.18)):

$$\lim_{N \to \infty} \mathbb{E} \frac{1}{N^p} \left\| \boldsymbol{x}^{\otimes p} - \boldsymbol{m}_p(\boldsymbol{\vec{y}}_t, t) \right\|_2^2 = 1 - q_{\mathsf{bayes}}^p$$

We first prove a preparatory lemma. In what follows, we let $\tilde{\beta}_{p'} = \beta_{p'}$ be fixed for all $p' \neq p$ and treat $\tilde{\beta}_p$ as a variable. Define $\tilde{\xi}(q) = \sum_{p' \ge 1} \tilde{\beta}_{p'}^2 q^{p'}$; we sometimes emphasize the dependence on $\tilde{\beta}_p$ by writing $\tilde{\xi}^{\tilde{\beta}_p}(q)$. Let \mathcal{P} denote the Parisi functional for spherical spin glasses, see e.g. [Tal06, Equation (1.12)]. (In the proof below we will only need the replica-symmetric case of this functional, which is given in Proposition 5.14.) Further, for $q \in (-1, 1)$, let

$$\widetilde{\xi}_q(s) = \widetilde{\xi}(q^2 + (1 - q^2)s) - \widetilde{\xi}(q^2),$$

and define

$$P(\widetilde{\beta}_p) = \sup_{q \in [0,1)} \left\{ \widetilde{\xi}(q) + \mathcal{P}(\widetilde{\xi}_q) + \frac{1}{2} \log(1-q^2) \right\}$$

Lemma 10.4. Assume the setting of Proposition 10.3. For all $\tilde{\beta}_p$ in a neighborhood of β_p ,

$$P(\tilde{\beta}_p) = \frac{1}{2} \sup_{q \in [0,1)} \left\{ \tilde{\xi}(1) + \tilde{\xi}(q) + q + \log(1-q) \right\}.$$
 (10.2)

Furthermore, P is differentiable at β_p , with

$$P'(\beta_p) = \beta_p (1 + q_{\mathsf{bayes}}^p). \tag{10.3}$$

Proof. By Proposition 5.14 with $u = \frac{q}{1+q}$, for all $q \in [0, 1)$,

$$\mathcal{P}(\widetilde{\xi}_q) \leqslant \frac{1}{2} \left\{ \widetilde{\xi}_q(1) - \widetilde{\xi}_q(u) + \frac{u}{1-u} + \log(1-u) \right\} = \frac{1}{2} \left\{ \widetilde{\xi}(1) - \widetilde{\xi}(q) + q - \log(1+q) \right\}, \quad (10.4)$$

and thus

$$P(\widetilde{\beta}_p) \leq \frac{1}{2} \sup_{q \in [0,1)} \left\{ \widetilde{\xi}(1) + \widetilde{\xi}(q) + q + \log(1-q) \right\}.$$

Since $\lim_{q \uparrow 1} \log(1-q) = -\infty$, the supremum is attained. Let $q(\tilde{\beta}_p)$ denote the maximizer. Arguing identically to the proof of Proposition 5.15, (10.4) is an equality at $q = q(\tilde{\beta}_p)$. This proves (10.2).

Note that $q(\beta_p) = q_{\text{bayes}}$ by definition. Since $t \notin T_1$, the maximum in (10.2) at $\tilde{\beta}_p = \beta_p$ is attained uniquely at q_{bayes} . By continuity, $\lim_{\tilde{\beta}_p \to \beta_p} q(\tilde{\beta}_p) = q_{\text{bayes}}$ as well. Note that for any $\tilde{\beta}_p > \beta_p$,

$$\frac{P(\widetilde{\beta}_p) - P(\beta_p)}{\widetilde{\beta}_p - \beta_p} \ge \frac{\widetilde{\xi}^{\widetilde{\beta}_p}(1) + \widetilde{\xi}^{\widetilde{\beta}_p}(q(\beta_p)) - \widetilde{\xi}^{\beta_p}(1) - \widetilde{\xi}^{\beta_p}(q(\beta_p))}{\widetilde{\beta}_p - \beta_p} = 2\beta_p(1 + q(\beta_p)^p) + O(\widetilde{\beta}_p - \beta_p),$$
$$\frac{P(\widetilde{\beta}_p) - P(\beta_p)}{\widetilde{\beta}_p - \beta_p} \le \frac{\widetilde{\xi}^{\widetilde{\beta}_p}(1) + \widetilde{\xi}^{\widetilde{\beta}_p}(q(\widetilde{\beta}_p)) - \widetilde{\xi}^{\beta_p}(1) - \widetilde{\xi}^{\beta_p}(q(\widetilde{\beta}_p))}{\widetilde{\beta}_p - \beta_p} = 2\beta_p(1 + q(\widetilde{\beta}_p)^p) + O(\widetilde{\beta}_p - \beta_p).$$

Taking the limit $\widetilde{\beta}_p \downarrow \beta_p$ yields

$$\lim_{\widetilde{\beta}_p \downarrow \beta_p} \frac{P(\widetilde{\beta}_p) - P(\beta_p)}{\widetilde{\beta}_p - \beta_p} = 2\beta_p (1 + q_{\mathsf{bayes}}^p).$$

A similar argument shows the left derivative also equals this, proving (10.3).

Proof of Proposition 10.3. Let \widetilde{H}_N be a spin glass Hamiltonian with mixture $\widetilde{\xi}$, and let

$$F_N(\widetilde{\beta}_p) = \frac{1}{N} \mathbb{E} \log \int_{S_N} \exp \left\{ N \widetilde{\xi}(\langle \boldsymbol{x}, \boldsymbol{\sigma} \rangle_N) + \widetilde{H}_N(\boldsymbol{\sigma}) \right\} \, \mathrm{d}\mu_0(\boldsymbol{\sigma}).$$

Since the restriction of \widetilde{H}_N to the band $\langle \boldsymbol{x}, \boldsymbol{\sigma} \rangle_N = q$ is a spin glass with mixture $\widetilde{\xi}_q$, the Parisi formula [Tal06, Theorem 1.1] implies

$$\lim_{N \to \infty} F_N(\widetilde{\beta}_p) = \sup_{q \in (-1,1)} \left\{ \widetilde{\xi}(q) + \mathcal{P}(\widetilde{\xi}_q) + \frac{1}{2} \log(1-q^2) \right\}.$$

This equals $P(\tilde{\beta}_p)$ because the supremum over (-1, 0] is clearly at most the supremum over [0, 1). By Hölder's inequality, $F_N(\tilde{\beta}_p)$ is convex in $\tilde{\beta}_p$. So, for any $\delta > 0$,

$$\frac{F_N(\beta_p) - F_N(\beta_p - \delta)}{\delta} \leqslant F'_N(\beta_p) \leqslant \frac{F_N(\beta_p + \delta) - F_N(\beta_p)}{\delta}.$$

Differentiability of P (by Lemma 10.4) then implies

$$\lim_{N \to \infty} F'_N(\beta_p) = P'(\beta_p) = \beta_p (1 + q^p_{\mathsf{bayes}}).$$
(10.5)

Let $\langle \cdot \rangle$ denote average w.r.t. the Gibbs measure corresponding to Hamiltonian $\check{H}_{N,t}$, which coincides in law with $N \tilde{\xi}(\langle \boldsymbol{x}, \boldsymbol{\sigma} \rangle_N) + \tilde{H}_N(\boldsymbol{\sigma})$ for $\tilde{\beta}_p = \beta_p$. Note that $\boldsymbol{m}_p(\boldsymbol{\vec{y}}_t, t) = \langle \boldsymbol{\sigma}^{\otimes p} \rangle$. We calculate that

$$\begin{split} F'_{N}(\beta_{p}) &= 2\beta_{p}\mathbb{E}\langle\langle \boldsymbol{x}, \boldsymbol{\sigma} \rangle_{N}^{p} \rangle + \beta_{p} \left(1 - \mathbb{E}\langle\langle \boldsymbol{\sigma}, \boldsymbol{\sigma} \rangle_{N}^{p} \rangle\right) \\ &= \beta_{p} \left(1 + 2\mathbb{E}\frac{\langle \boldsymbol{x}^{\otimes p}, \boldsymbol{m}_{p}(\boldsymbol{\vec{y}}_{t}, t) \rangle}{N^{p}} - \mathbb{E}\frac{\langle \boldsymbol{m}_{p}(\boldsymbol{\vec{y}}_{t}, t), \boldsymbol{m}_{p}(\boldsymbol{\vec{y}}_{t}, t) \rangle}{N^{p}}\right). \end{split}$$

Comparing with (10.5) shows

$$\lim_{N \to \infty} \left\{ 2\mathbb{E} \frac{\langle \boldsymbol{x}^{\otimes p}, \boldsymbol{m}_p(\boldsymbol{\vec{y}}_t, t) \rangle}{N^p} - \mathbb{E} \frac{\langle \boldsymbol{m}_p(\boldsymbol{\vec{y}}_t, t), \boldsymbol{m}_p(\boldsymbol{\vec{y}}_t, t) \rangle}{N^p} \right\} = q_{\text{bayes}}^p$$

Since

$$\mathbb{E}\frac{1}{N^p} \left\| \boldsymbol{x}^{\otimes p} - \boldsymbol{m}_p(\boldsymbol{\vec{y}}_t, t) \right\|_2^2 = 1 - 2\mathbb{E}\frac{\langle \boldsymbol{x}^{\otimes p}, \boldsymbol{m}_p(\boldsymbol{\vec{y}}_t, t) \rangle}{N^p} + \mathbb{E}\frac{\langle \boldsymbol{m}_p(\boldsymbol{\vec{y}}_t, t), \boldsymbol{m}_p(\boldsymbol{\vec{y}}_t, t) \rangle}{N^p},$$

blows.

the result follows.

Lemma 10.5. If there exists $q \in [0,1)$ such that $\xi''(q) > \frac{1}{(1-q)^2}$, then there exists $t \ge 0$ such that $\xi'_t(q) = \frac{q}{1-q}$ has more than one solution.

Proof. Let $g_t(q) = \check{\xi}'_t(q) - \frac{q}{1-q}$, so solutions to $\check{\xi}'_t(q) = \frac{q}{1-q}$ are zeros of g_t . Suppose for contradiction that for all $t \ge 0$, g_t has unique zero $q_{\mathsf{AMP}}(t)$. Then, for all $t, g_t > 0$ on $[0, q_{\mathsf{AMP}}(t))$ (this is vacuous if $q_{\mathsf{AMP}}(t) = 0$) and $g_t < 0$ on $(q_{\mathsf{AMP}}(t), 1)$. Note that for each $q, g_t(q)$ is continuous and increasing in t, and thus $q_{\mathsf{AMP}}(t)$ is also continuous and increasing.

Recall that $\|\tau(t)\|_1 = t$ for all t. For each $q \in (0, 1)$,

$$g_t(q) \ge \sum_{j=1}^J j\tau_j(t)q^{j-1} - \frac{q}{1-q} \ge \|\tau(t)\|_1 q^{J-1} - \frac{q}{1-q} = tq^{J-1} - \frac{q}{1-q}.$$
 (10.6)

It follows that $g_t(q) > 0$ for sufficiently large t. Thus $\lim_{t\to+\infty} g_{AMP}(t) = 1$, so $q_{AMP}(t)$ ranges over all of [0,1) as t ranges over $[0,+\infty)$.

Since $\xi''(q) > \frac{1}{(1-q)^2}$ for some $q \in [0, 1)$, the function g_0 is not monotonically decreasing. Let $0 \leq q_1 < q_2 < 1$ be such that $g_0(q_1) < g_0(q_2)$. Note that

$$g_t(q_1) - g_0(q_1) = \sum_{j=1}^J j\tau_j(t)q_1^{j-1} \le \sum_{j=1}^J j\tau_j(t)q_2^{j-1} = g_t(q_2) - g_0(q_2)$$

Thus $g_t(q_1) < g_t(q_2)$. Set t such that $q_1 = q_{AMP}(t)$, so that $g_t(q_1) = 0$. This implies that $g_t(q_2) > 0$, and therefore g_t has another zero in $[q_2, 1)$.

Lemma 10.6. If there exists $t \ge 0$ such that $\xi'_t(q) = \frac{q}{1-q}$ has more than one solution, then there exists a nontrivial interval $I = [t_-, t_+] \subseteq [0, +\infty)$ such that for all $t' \in I$, $q_{\mathsf{AMP}}(t') \ne q_{\mathsf{bayes}}(t')$.

Proof. Let g_t be defined as in the proof of Lemma 10.5 and $q_1 = q_{AMP}(t)$, so that q_1 is the smallest zero of g_t . Let $q_2 > q_1$ be the next smallest zero of g_t . Note that by Lemma 10.2, either $g_t(q) > 0$ for all $q \in (q_1, q_2)$ or $g_t(q) < 0$ for all $q \in (q_1, q_2)$.

Suppose the former case holds. We will show the conclusion holds with $I = [t, t - \delta]$ for small δ . We first show that we must have t > 0, so this is a valid interval. Suppose for contradiction that t = 0; then $q_1 = 0$. So, $g_0(q) = \xi'(q) - \frac{q}{1-q}$ is positive on $(0, q_2)$. This implies that for $q \in (0, q_2]$,

$$\xi(q) + q + \log(1 - q) = \int_0^q g_0(s) \, \mathrm{d}s > 0,$$

contradicting (2.21). Note that

$$\left(\check{\xi}_t(q_2) + q_2 + \log(1 - q_2)\right) - \left(\check{\xi}_t(q_1) + q_1 + \log(1 - q_1)\right) = \int_{q_1}^{q_2} g_t(q) \, \mathrm{d}q > 0.$$

We claim that $q_{AMP}(t')$ is continuous on $t' \in I$, for small enough δ . If $q_1 = 0$, this is clear because $q_{AMP}(t)$ is increasing. Otherwise, since $g_t(0) \ge 0$ and q_1 is the smallest zero of g_t , we have $g_t(q) > 0$ for $q \in [0, q_1)$. Since the $g_t(q)$ are continuous and increasing in t, the claim follows. It follows that for sufficiently small δ , for all $t' \in I$ and $q'_1 = q_{AMP}(t')$,

$$\left(\check{\xi}_{t'}(q_2) + q_2 + \log(1 - q_2)\right) - \left(\check{\xi}_{t'}(q_1') + q_1' + \log(1 - q_1')\right) > 0.$$

Thus $q_{AMP}(t') \neq q_{bayes}(t')$ for all $t' \in I$.

Finally, we consider the case that $g_t(q) < 0$ for all $q \in (q_1, q_2)$. Then, $g_t > 0$ on $[0, q_1)$ (vacuously if $q_1 = 0$) and $g_t < 0$ on (q_1, q_2) . Let t'' be the smallest time such that $\inf_{q \in [q_1, q_2]} g_{t''}(q) \ge 0$; this is finite by the discussion surrounding (10.6). Since $g_t(q)$ is increasing in t, we have $g_{t''} \ge 0$ for $q \in [0, q_2]$, with equality attained at some $q \in [q_1, q_2]$. By definition, $q_{AMP}(t'')$ is the smallest such q. As $f_{t''}(q_2) > g_t(q_2) = 0$, we have $q_{AMP}(t'') < q_2$. The result now follows from the first case.

Proof of Theorem 2.3. By the last two lemmas, there exists a nontrivial interval $I = [t_-, t_+] \subseteq [0, +\infty)$ such that $q_{\mathsf{AMP}}(t) \neq q_{\mathsf{bayes}}(t)$ for all $t \in I$. Since $q_{\mathsf{bayes}}(t)$ is a maximizer of (10.1), it satisfies the stationarity condition $\check{\xi}'_t(q) = \frac{q}{1-q}$, and therefore $q_{\mathsf{AMP}}(t) < q_{\mathsf{bayes}}(t)$. It also follows that $q_{\mathsf{bayes}}(t) > 0$.

Let U(t) be the number of nonzero coefficients of ξ_t of degree at most J. This is an increasing function with at most J discontinuities; let T_0 be the set of these discontinuities.

We will show the theorem holds with $\mathcal{I} = I \setminus (T_0 \cup T_1)$. (Recall the definition of T_1 in Lemma 10.2.) This is a positive measure set by Lemma 10.2. Consider any $t \in \mathcal{I}$. Since $t \notin T_0$, there exists $1 \leq j \leq J$ such that the q^j coefficient of ξ_t is positive and $\tau'_i(t) > 0$. By Propositions 10.1 and 10.3,

$$\lim_{k \to \infty} \lim_{N \to \infty} \mathbb{E} \frac{1}{N^j} \left\| \boldsymbol{x}^{\otimes j} - (\widetilde{\boldsymbol{m}}^k)^{\otimes j} \right\|_2^2 = 1 - q_{\mathsf{AMP}}(t)^j,$$
$$\lim_{N \to \infty} \mathbb{E} \frac{1}{N^j} \| \boldsymbol{x}^{\otimes j} - \boldsymbol{m}_j(\vec{\boldsymbol{y}}_t, t) \|^2 \leq 1 - q_{\mathsf{bayes}}(t)^j.$$

Since $q_{AMP}(t) < q_{bayes}(t)$, the conclusion follows.

References

[AA13] Antonio Auffinger and Gerard Ben Arous. Complexity of random smooth functions on the high-dimensional sphere. *Annals of Probability*, 41(6):4214–4247, 2013.

- [ABXY22] Arka Adhikari, Christian Brennecke, Changji Xu, and Horng-Tzer Yau. Spectral gap estimates for mixed *p*-spin models at high temperature. *arXiv*:2208.07844, 2022.
- [AGZ10] Greg W Anderson, Alice Guionnet, and Ofer Zeitouni. *An introduction to random matrices*. Number 118. Cambridge university press, 2010.
- [AJ24] Gérard Ben Arous and Aukosh Jagannath. Shattering versus metastability in spin glasses. *Communications on Pure and Applied Mathematics*, 77(1):139–176, 2024.
- [AJK⁺23] Nima Anari, Vishesh Jain, Frederic Koehler, Huy Tuan Pham, and Thuy-Duong Vuong. Universality of spectral independence with applications to fast mixing in spin glasses. *arXiv:2307.10466*, 2023.
- [AM22] Ahmed El Alaoui and Andrea Montanari. An information-theoretic view of stochastic localization. *IEEE Transactions on Information Theory*, 68(11):7423–7426, 2022.
- [AMS22] Ahmed El Alaoui, Andrea Montanari, and Mark Sellke. Sampling from the Sherrington-Kirkpatrick Gibbs measure via algorithmic stochastic localization. In *Proc. 63rd FOCS*, pages 323–334, 2022.
- [AMS23a] Ahmed El Alaoui, Andrea Montanari, and Mark Sellke. Sampling from mean-field gibbs measures via diffusion processes. *arXiv:2310.08912*, 2023.
- [AMS23b] Ahmed El Alaoui, Andrea Montanari, and Mark Sellke. Shattering in pure spherical spin glasses. *arXiv:2307.04659*, 2023.
- [AT07] Robert J Adler and Jonathan E Taylor. *Random fields and geometry*, volume 80. Springer, 2007.
- [BADG06] Gérard Ben Arous, Amir Dembo, and Alice Guionnet. Cugliandolo-Kurchan equations for dynamics of spin-glasses. *Probability theory and related fields*, 136(4):619–660, 2006.
- [BCKM98] Jean-Philippe Bouchaud, Leticia F Cugliandolo, Jorge Kurchan, and Marc Mézard. Out of equilibrium dynamics in spin-glasses and other glassy systems. *Spin glasses and random fields*, 12:161, 1998.
- [BČNS22] David Belius, Jiří Černý, Shuta Nakajima, and Marius A Schmidt. Triviality of the geometry of mixed p-spin spherical hamiltonians with external field. *Journal of Statistical Physics*, 186(1):12, 2022.
- [BL16] Jinho Baik and Ji Oon Lee. Fluctuations of the free energy of the spherical sherringtonkirkpatrick model. *Journal of Statistical Physics*, 165:185–224, 2016.
- [BM11] Mohsen Bayati and Andrea Montanari. The dynamics of message passing on dense graphs, with applications to compressed sensing. *IEEE Transactions on Information Theory*, 57(2):764–785, 2011.
- [Bol14] Erwin Bolthausen. An iterative construction of solutions of the tap equations for the sherrington-kirkpatrick model. *Communications in Mathematical Physics*, 325(1):333–366, 2014.

- [CCL⁺22] Sitan Chen, Sinho Chewi, Jerry Li, Yuanzhi Li, Adil Salim, and Anru R Zhang. Sampling is as easy as learning the score: theory for diffusion models with minimal data assumptions. *arXiv:2209.11215*, 2022.
- [Che13] Wei-Kuo Chen. The Aizenman-Sims-Starr scheme and Parisi formula for mixed *p*-spin spherical models. *Electronic Journal of Probability*, 18, 2013.
- [CHS93] Andrea Crisanti, Heinz Horner, and H J Sommers. The spherical p-spin interaction spin-glass model: the dynamics. *Zeitschrift für Physik B Condensed Matter*, 92:257–271, 1993.
- [CK93] Leticia F Cugliandolo and Jorge Kurchan. Analytical solution of the off-equilibrium dynamics of a long-range spin-glass model. *Physical Review Letters*, 71(1):173, 1993.
- [CLA⁺21] Sinho Chewi, Chen Lu, Kwangjun Ahn, Xiang Cheng, Thibaut Le Gouic, and Philippe Rigollet. Optimal dimension dependence of the metropolis-adjusted langevin algorithm. In *Conference on Learning Theory*, pages 1260–1300. PMLR, 2021.
- [CS95] A Crisanti and H-J Sommers. Thouless-anderson-palmer approach to the spherical p-spin spin glass model. *Journal de Physique I*, 5(7):805–813, 1995.
- [CS17] Wei-Kuo Chen and Arnab Sen. Parisi formula, disorder chaos and fluctuation for the ground state energy in the spherical mixed p-spin models. *Communications in Mathematical Physics*, 350:129–173, 2017.
- [DGM07] Amir Dembo, Alice Guionnet, and Christian Mazza. Limiting dynamics for spherical models of spin glasses at high temperature. *Journal of Statistical Physics*, 126:781–815, 2007.
- [EAMS21] Ahmed El Alaoui, Andrea Montanari, and Mark Sellke. Optimization of mean-field spin glasses. *The Annals of Probability*, 49(6):2922–2960, 2021.
- [Eld20] Ronen Eldan. Taming correlations through entropy-efficient measure decompositions with applications to mean-field approximation. *Probability Theory and Related Fields*, 176(3-4):737–755, 2020.
- [FLD14] Yan V Fyodorov and Pierre Le Doussal. Topology trivialization and large deviations for the minimum in the simplest random optimization. *Journal of Statistical Physics*, 154(1-2):466– 490, 2014.
- [Fyo15] YV Fyodorov. High-dimensional random fields and random matrix theory. *Markov Processes and Related Fields*, 21(3):483–518, 2015.
- [GDKZ23] Davide Ghio, Yatin Dandi, Florent Krzakala, and Lenka Zdeborová. Sampling with flows, diffusion and autoregressive neural networks: A spin-glass perspective. *arXiv:2308.14085*, 2023.
- [GJ19] Reza Gheissari and Aukosh Jagannath. On the spectral gap of spherical spin glass dynamics. 55(2):756–776, 2019.
- [GJK23] David Gamarnik, Aukosh Jagannath, and Eren C Kızıldağ. Shattering in the ising pure *p*-spin model. *arXiv*:2307.07461, 2023.

- [GZ00] A Guionnet and O Zeitouni. Concentration of the spectral measure for large matrices. *Electronic Communications in Probability*, 5:119–136, 2000.
- [HJA20] Jonathan Ho, Ajay Jain, and Pieter Abbeel. Denoising diffusion probabilistic models. *Advances in Neural Information Processing Systems*, 33:6840–6851, 2020.
- [HS22] Brice Huang and Mark Sellke. Tight lipschitz hardness for optimizing mean field spin glasses. In 2022 IEEE 63rd Annual Symposium on Foundations of Computer Science (FOCS), pages 312–322. IEEE, 2022.
- [HS23a] Brice Huang and Mark Sellke. A constructive proof of the spherical parisi formula. *arXiv:2311.15495*, 2023.
- [HS23b] Brice Huang and Mark Sellke. Strong topological trivialization of multi-species spherical spin glasses. *arXiv:2308.09677*, 2023.
- [JM13] Adel Javanmard and Andrea Montanari. State evolution for general approximate message passing algorithms, with applications to spatial coupling. *Information and Inference: A Journal of the IMA*, 2(2):115–144, 2013.
- [Kac48] M Kac. On the Average Number of Real Roots of a Random Algebraic Equation (II). *Proceedings of the London Mathematical Society*, 2(1):390–408, 1948.
- [KT87] Theodore R Kirkpatrick and Devarajan Thirumalai. p-spin-interaction spin-glass models: Connections with the structural glass problem. *Physical Review B*, 36(10):5388, 1987.
- [LG16] Jean-François Le Gall. *Brownian motion, martingales, and stochastic calculus*. Springer, 2016.
- [Mon23] Andrea Montanari. Sampling, diffusions, and stochastic localization. *arXiv:2305.10690*, 2023.
- [MRTS07] Andrea Montanari, Federico Ricci-Tersenghi, and Guilhem Semerjian. Solving constraint satisfaction problems through belief propagation-guided decimation. *arXiv:0709.1667*, 2007.
- [MS23] Andrea Montanari and Eliran Subag. Solving overparametrized systems of random equations: I. Model and algorithms for approximate solutions. *arXiv:2306.13326*, 2023.
- [Nes03] Yurii Nesterov. *Introductory lectures on convex optimization: A basic course*, volume 87. Springer Science & Business Media, 2003.
- [Ric44] Stephen O Rice. Mathematical analysis of random noise. *The Bell System Technical Journal*, 23(3):282–332, 1944.
- [SDWMG15] Jascha Sohl-Dickstein, Eric Weiss, Niru Maheswaranathan, and Surya Ganguli. Deep unsupervised learning using nonequilibrium thermodynamics. In *International Conference on Machine Learning*, pages 2256–2265. PMLR, 2015.
- [SSDK⁺21] Yang Song, Jascha Sohl-Dickstein, Diederik P Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. Score-based generative modeling through stochastic differential equations. In *International Conference on Learning Representations (ICLR)*, 2021.

- [Tal06] Michel Talagrand. Free energy of the spherical mean field model. *Probability theory and related fields*, 134(3):339–382, 2006.
- [TAP77] David J. Thouless, Philip W. Anderson, and Richard G. Palmer. Solution of 'solvable model of a spin glass'. *Philosophical Magazine*, 35(3):593–601, 1977.