

Axial Gravitational Perturbations of Slowly-Rotating Compact Objects in General Relativity and Beyond

Xing-Hui Feng^{1*} and Jun Peng^{2†}

¹*Center for Joint Quantum Studies and Department of Physics, School of Science,
Tianjin University, Tianjin 300350, China*

²*Faculty of Science, Beijing University of Technology, Beijing 100124, China*

ABSTRACT

We study the axial gravitational perturbations of slowly-rotating compact objects which are assumed to be supported by anisotropic fluids. We find that the gravitational perturbations decouple from the matter perturbations for axial sectors. We obtain a master wave equation whose potential is fully determined by the metric functions. This equation makes the calculations of gravitational QNMs for rotating compact objects extremely easy in specific background configurations.

* xhfeng@tju.edu.cn † junpeng@bjut.edu.cn

1 Introduction

Black hole (BH) perturbation theory is an important tool to study the stability and quasi-normal modes (QNMs) of BH, the ringdown signal from compact binary coalescences [1, 2, 3]. The metric perturbations of static BHs can well be established according to their spherical symmetry. We can expand the metric perturbations using spherical harmonics and separate them into the axial and polar parts depending on their behavior under parity transformations. The axial perturbations of Schwarzschild BH is studied in [4] and the master wave equation is the famous Regge-Wheeler equation. The polar perturbations of Schwarzschild BH is studied in [5, 6] and the master wave equation is the so-called Zerilli equation.

However we have no well-posed metric perturbations for rotating BH so far. The perturbation of Kerr BH is completed in [7] based on Newmann-Penrose (NP) formalism. The master wave equation is the famous Teukolsky equation, whose solutions are two-dimensional eigenvalue problem. When applying the NP form to the perturbations of Kerr-Newmann (KN) BH, we have no ways to obtain a variables-separated and perturbations-decoupled master wave equation [8]. Anyway the analytical perturbations of non-Kerr and spinning BH in (or beyond) general relativity (GR) are open problems.

Things become possible when we consider the slowly-rotating backgrounds. Since the spin parameter is viewed as a small modification with respect to the static metric, the general decomposition of metric perturbations, as well as the gauge choosing are still valid for slowly-rotating ones. Based on Kojima's elegant work about the angular separation of perturbation equations [9], Pani et al proposed an advanced perturbation method for slowly-rotating spacetime [10]. This advanced method is applied in the perturbations of Kerr BH [10], KN BH [11, 12], BHs in Einstein-dilaton Gauss-Bonnet gravity [13, 14], dynamical Chern-Simons gravity [15, 16], higher-derivative gravity [17], Einstein-Bumblebee gravity [18].

In this paper, we investigate the axial gravitational perturbations of slowly-rotating compact objects in GR and beyond. We assume the general slowly-rotating metric ansatz is supported by the effective stress tensor of an anisotropic fluid, seeing the set up in section 2. We analyse the axial gravitational perturbations in section 3 and derive the master wave equation in section 4. We give a brief discussion in section 5.

2 The set up

Understanding the gravitational field of a rotating body is one of great astrophysical and theoretical significance. Although a number of exact solutions with rotation are known in Einstein gravity, for general modified gravities they are quite difficult to obtain. On the other hand slowly-rotating solutions are somewhat easier to come by. The general ansatz of slowly-rotating metric is

$$\bar{d}s^2 = -A(r)dt^2 + \frac{dr^2}{B(r)} + C(r)(d\theta^2 + \sin^2\theta d\phi^2) - 2aD(r)\sin^2\theta dt d\phi \quad (2.1)$$

where a is the spin parameter. In ab-initio calculation, the metric should solve the equations of motion for certain theory of gravity to leading order in the spin parameter a . From a phenomenological point of view, we can assume the solution is governed by the Einstein equation with an effective stress tensor

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 8\pi T_{\mu\nu} \quad (2.2)$$

Without loss of generality, the stress tensor supporting the metric (2.1) is described by an anisotropic fluid [19]

$$T_{\mu\nu} = \rho u_\mu u_\nu + P_r k_\mu k_\nu + P_t \Pi_{\mu\nu} \quad (2.3)$$

where u_μ is the fluid's 4-velocity and k_μ is the unit spacelike vector orthogonal to u_μ , i.e.

$$-u^\mu u_\mu = k^\mu k_\mu = 1, \quad u^\mu k_\mu = 0 \quad (2.4)$$

and

$$\Pi_{\mu\nu} = g_{\mu\nu} + u_\mu u_\nu - k_\mu k_\nu \quad (2.5)$$

is a projection operator onto a two-surface orthogonal to u^μ and k^μ , i.e.

$$u_\mu \Pi^{\mu\nu} V_\nu = k_\mu \Pi^{\mu\nu} V_\nu = 0 \quad (2.6)$$

for any vector V^μ . We can assume

$$u^\mu = \frac{1}{\sqrt{A}}(1, 0, 0, \Omega), \quad k^\mu = \sqrt{B}(0, 1, 0, 0) \quad (2.7)$$

where Ω is the fluid angular velocity.

3 Axial gravitational perturbations

In order to study the gravitational perturbations, one needs to perturb the gravitational equations as well as the energy momentum tensor. As in the spherically symmetric case,

the metric perturbations $g_{\mu\nu} = \bar{g}_{\mu\nu} + h_{\mu\nu}$ can be decomposed into the axial and polar parts. In Regge-Wheller gauge they reads

$$\begin{aligned} h_{\mu\nu}^- dx^\mu dx^\nu &= 2[h_0^{\ell m} dt + h_1^{\ell m} dr] \times [S_\theta^{\ell m} d\theta + S_\phi^{\ell m} d\phi] \\ h_{\mu\nu}^+ dx^\mu dx^\nu &= [AH_0^{\ell m} dt^2 + 2H_1^{\ell m} dt dr + H_2^{\ell m}/B dr^2 + K^{\ell m} d\Omega_2^2] Y^{\ell m} \end{aligned} \quad (3.1)$$

where $h_0^{\ell m}, h_1^{\ell m}, H_0^{\ell m}, H_1^{\ell m}, H_2^{\ell m}, K^{\ell m}$ are free functions of t, r , and $S_a^{\ell m}$ is the axial vector spherical harmonics

$$(S_\theta^{\ell m}, S_\phi^{\ell m}) = \left(-\frac{Y_{,\phi}^{\ell m}}{\sin \theta}, \sin \theta Y_{,\theta}^{\ell m} \right) \quad (3.2)$$

In the case of a spherically symmetric background, perturbations with different values of (ℓ, m) , as well as perturbations with opposite parity, are decoupled. In a rotating, axially symmetric background, perturbations with different values of m are still decoupled but perturbations with different values of ℓ are not. However in [20, 21], it's proved that if we are only interested in the eigenvalue spectrum to first order in spin a , the polar and axial perturbations—as well as perturbations with different values of the harmonic indices—are decoupled from each other, and can be studied independently. In the following, we only consider axial perturbations and continue in tetrad formalism for simplicity. We define the tetrad basis $e^{\bar{a}} = e_{\mu}^{\bar{a}} dx^\mu$ as

$$\begin{aligned} e^{\bar{0}} &= \sqrt{A} dt - \frac{aD \sin^2 \theta}{\sqrt{A}} d\phi, & e^{\bar{1}} &= \frac{dr}{\sqrt{B}}, \\ e^{\bar{2}} &= \sqrt{C} d\theta + \frac{h_0^{\ell m} S_\theta^{\ell m}}{\sqrt{C}} dt + \frac{h_1^{\ell m} S_\theta^{\ell m}}{\sqrt{C}} dr \\ e^{\bar{3}} &= \sqrt{C} \sin \theta d\phi + \frac{h_0^{\ell m} S_\phi^{\ell m}}{\sqrt{C} \sin \theta} dt + \frac{h_1^{\ell m} S_\phi^{\ell m}}{\sqrt{C} \sin \theta} dr \end{aligned} \quad (3.3)$$

The inverse tetrad $E_{\bar{a}}^\mu$ are given by

$$e_{\mu}^{\bar{a}} E_{\bar{a}}^\nu = \delta_{\mu}^{\nu}, \quad e_{\mu}^{\bar{a}} E_{\bar{b}}^\mu = \delta_{\bar{b}}^{\bar{a}} \quad (3.4)$$

The tensor components in coordinate and tetrad basis are related by

$$\begin{aligned} A_{\mu} &= e_{\mu}^{\bar{a}} A_{\bar{a}}, & A_{\bar{a}} &= E_{\bar{a}}^{\mu} A_{\mu} \\ B_{\mu\nu} &= e_{\mu}^{\bar{a}} e_{\nu}^{\bar{b}} B_{\bar{a}\bar{b}}, & B_{\bar{a}\bar{b}} &= E_{\bar{a}}^{\mu} E_{\bar{b}}^{\nu} B_{\mu\nu} \end{aligned} \quad (3.5)$$

In tetrad formalism, we rewrite the effective Einstein equation as

$$R_{\bar{a}\bar{b}} = 8\pi T_{\bar{a}\bar{b}}^t \quad (3.6)$$

where

$$T_{\bar{a}\bar{b}}^t \equiv (T_{\bar{a}\bar{b}} - \frac{1}{2}T\eta_{\bar{a}\bar{b}}) \quad (3.7)$$

The perturbation equations are given by

$$\mathcal{E}_{\bar{a}\bar{b}} \equiv \delta R_{\bar{a}\bar{b}} - 8\pi\delta T_{\bar{a}\bar{b}}^t = 0 \quad (3.8)$$

The perturbations of Ricci tensor can be calculated based on the metric perturbation (3.1). The perturbations of stress tensor are discussed in appendix A. The axial perturbation equations can be divided into two groups [9, 10]. The first group is

$$\begin{aligned} \mathcal{E}_{\bar{L}\bar{2}} &= a\tilde{\alpha}_{\ell m}^L \cos\theta Y_{,\theta}^{\ell m} - \beta_{\ell m}^L \frac{Y_{,\phi}^{\ell m}}{\sin\theta} + a\eta_{\ell m}^L \sin\theta Y^{\ell m} + a\chi_{\ell m}^L \sin\theta W^{\ell m} \\ \mathcal{E}_{\bar{L}\bar{3}} &= \beta_{\ell m}^L Y_{,\theta}^{\ell m} + a\tilde{\alpha}_{\ell m}^L \cos\theta \frac{Y_{,\phi}^{\ell m}}{\sin\theta} + a\chi_{\ell m}^L X^{\ell m} \end{aligned} \quad (3.9)$$

where $L = 0, 1$. The second group reads

$$\begin{aligned} \mathcal{E}_{2\bar{3}} &= ag_{\ell m} Y_{,\phi}^{\ell m} + t_{\ell m} W^{\ell m} \\ \mathcal{E}_{(-)} &= \mathcal{E}_{\bar{2}\bar{2}} - \mathcal{E}_{\bar{3}\bar{3}} = ag_{\ell m} \sin\theta Y_{,\theta}^{\ell m} - t_{\ell m} \frac{X^{\ell m}}{\sin\theta} \end{aligned} \quad (3.10)$$

Here $\tilde{\alpha}_{\ell m}^L, \beta_{\ell m}^L, \eta_{\ell m}^L, \chi_{\ell m}^L, t_{\ell m}, g_{\ell m}$ consist of axial perturbation variables $h_0^{\ell m}, h_1^{\ell m}, U^{\ell m}, Q^{\ell m}$. Their explicit expressions are given in appendix B, where all perturbation variables are in frequency domain with Fourier transformation $e^{-i\omega t}$.

4 Master wave equation

In order to derive a master wave equation, we need separate the angular dependence. This can be performed using the orthogonality properties of the spherical harmonics [9, 10]

$$\begin{aligned} P_L &= \int d\Omega \left[Y_{,\theta}^{*\ell'm'} \mathcal{E}_{\bar{L}\bar{3}} - \frac{Y_{,\phi}^{*\ell'm'}}{\sin\theta} \mathcal{E}_{\bar{L}\bar{2}} \right] \\ &= J\beta_{\ell m}^L + ima[(J-2)\chi_{\ell m}^L + \tilde{\alpha}_{\ell m}^L + \eta_{\ell m}^L] = 0 \end{aligned} \quad (4.1)$$

$$\begin{aligned} P_2 &= \int d\Omega \frac{1}{J-2} \left[W^{*\ell'm'} \mathcal{E}_{\bar{2}\bar{3}} - \frac{X^{*\ell'm'}}{\sin\theta} \mathcal{E}_{(-)} \right] \\ &= Jt_{\ell m} + imag_{\ell m} = 0 \end{aligned} \quad (4.2)$$

where $d\Omega = \sin\theta d\theta d\phi$ is the volume form of the two-sphere and $J = \ell(\ell+1)$. We have three equations $P_0 = P_1 = P_2 = 0$ for the axial perturbations $h_0^{\ell m}, h_1^{\ell m}$ and $U^{\ell m}$, in which

$$P_1 = J\beta_{\ell m}^1 + ima[(J-2)\chi_{\ell m}^1 + \tilde{\alpha}_{\ell m}^1 + \eta_{\ell m}^1] = 0 \quad (4.3)$$

$$P_2 = Jt_{\ell m} + imag_{\ell m} = 0 \quad (4.4)$$

are two first order equations involved only in $h_0^{\ell m}, h_1^{\ell m}$. We start with the perturbation of static background

$$\begin{aligned}\beta_{\ell m}^1 &= 0 \\ t_{\ell m} &= 0\end{aligned}\tag{4.5}$$

The second equation gives

$$h_0 = a(r)h_1 + b(r)h_1'\tag{4.6}$$

From now on, we omit the upscript ℓm in all perturbation variables. Substituting it into the first equation, we have

$$h_1'' + c(r)h_1' + d(r)h_1 = 0\tag{4.7}$$

where $a(r), b(r), c(r), d(r)$ are certain functions. Finally, we can introduce a master variable

$$\Psi^{(0)} = \sqrt{\frac{AB}{C}}h_1\tag{4.8}$$

Then we obtain a Schrodinger-like wave equation

$$\frac{d^2\Psi^{(0)}}{dr_*^2} + (\omega^2 - V^{(0)})\Psi = 0\tag{4.9}$$

where r_* is the tortoise coordinate defined by $dr_* = dr/\sqrt{AB}$. The potential V is given by

$$V^{(0)} = (J - 2)\frac{A}{C} + \sqrt{ABC}\left(\sqrt{AB}\left(\frac{1}{\sqrt{C}}\right)'\right)'\tag{4.10}$$

Note that the equation $P_0 = 0$ gives the solution of fluid perturbation $U^{\ell m}$

$$U^{\ell m} = -h_0^{\ell m}\tag{4.11}$$

Now we turn to the rotating background. Since we consider only the slowly-rotating case, i.e. the spin parameter a contributes a linear correction to the static case, we can assume the master variable has the following form

$$\Psi = \sqrt{\frac{AB}{C}}(h_1 + maf_{h_1})\tag{4.12}$$

Then we can also obtain a Schrodinger-like wave equation

$$\frac{d^2\Psi}{dr_*^2} + (\omega^2 - V^{(0)} - maV^{(a)})\Psi = 0\tag{4.13}$$

We find that the corrective function f_{h_1} is

$$f_{h_1} = c + \frac{1}{\omega}\frac{D}{C}\tag{4.14}$$

where c is a constant and the corrective potential $V^{(a)}$ is

$$V^{(a)} = 2\omega \frac{D}{C} + \frac{1}{J\omega} \left[\sqrt{AB} \left(\sqrt{AB} \left(\frac{D}{C} \right)' \right)' - AB \frac{C'}{C} \left(\frac{D}{C} \right)' \right] \quad (4.15)$$

We can see that the axial gravitational perturbations decouple from perturbations of the effective stress tensor. A similar result was derived in [22] to parameterize derivations from general relativistic expectations at the level of the action governing the dynamics of BH perturbations. Their result applies to perturbations around BHs with scalar hair. In fact, our parameterized derivations attribute to the phenomenological stress tensor. Thus our result is more applicable, even for theories with vector hairs and high-dimensional operators whose perturbation dynamics is neglected. For KN BH, to first order in spin a , we have

$$A(r) = B(r) = \Delta/r^2, \quad C(r) = r^2, \quad D(r) = \frac{2M}{r} - \frac{Q^2}{r^2} \quad (4.16)$$

where $\Delta = r^2 - 2Mr + Q^2$, the potentials reduce to

$$V^{(0)} = \frac{\Delta}{r^5} \left[Jr - 6M + \frac{4Q^2}{r} \right] \\ V^{(a)} = \frac{8\Delta (3Mr^2(3r - 7M) + 7Q^2r(4M - r) - 9Q^4)}{J\omega r^{10}} + \frac{2\omega (2Mr - Q^2)}{r^4} \quad (4.17)$$

These results suppose to give the gravitational QNMs of slowly-rotating and weakly-charged KN BH (Freezing the perturbations of electromagnetic field).

5 Discussion

In this paper, we study the axial gravitational perturbations of general slowly-rotating compact objects in GR and beyond. We find that the gravitational parts decouple from the effective matter parts for axial sectors. Eventually we obtain a modified Regge-Wheeler equation whose potential is fully determined by the metric functions. This equation makes the calculations of gravitational QNMs for rotating compact objects extremely easy in specific background configurations. We leave the polar perturbations for future's work.

Acknowledgement

X.H.F. is supported by NSFC (National Natural Science Foundation of China) Grant No. 11905157 and No. 11935009.

A Perturbations of stress tensor

The most general perturbations of the stress tensor are

$$\begin{aligned}\delta T^{\mu\nu} = & (\rho + P_t)\delta(u^\mu u^\nu) + (P_r - P_t)\delta(k^\mu k^\nu) + P_t\delta g^{\mu\nu} \\ & + (\delta\rho + \delta P_t)u^\mu u^\nu + (\delta P_r - \delta P_t)k^\mu k^\nu + \delta P_t g^{\mu\nu}\end{aligned}\quad (\text{A.1})$$

The perturbation of the energy density and pressures can be expanded by the scalar spherical harmonics

$$\delta\rho = \delta\rho^{\ell m} Y^{\ell m}, \quad \delta P_r = \delta P_r^{\ell m} Y^{\ell m}, \quad \delta P_t = \delta P_t^{\ell m} Y^{\ell m} \quad (\text{A.2})$$

Similar to the decomposition of metric perturbation, the perturbation of fluid's 4-velocity u^μ are described by three functions $R^{\ell m}, U^{\ell m}, V^{\ell m}$

$$\begin{aligned}\delta u^r &= \frac{1}{\sqrt{AB}} R^{\ell m} Y^{\ell m} \\ \delta u^\theta &= \frac{1}{\sqrt{AC}} (U^{\ell m} S_\theta^{\ell m} + V^{\ell m} Y_{,\theta}^{\ell m}) \\ \delta u^\phi &= \frac{1}{\sqrt{AC} \sin^2 \theta} (U^{\ell m} S_\phi^{\ell m} + V^{\ell m} Y_{,\phi}^{\ell m})\end{aligned}\quad (\text{A.3})$$

The perturbations of k^μ are described by two functions $Q^{\ell m}, P^{\ell m}$

$$\begin{aligned}\delta k^\theta &= \frac{\sqrt{B}}{C} (Q^{\ell m} S_\theta^{\ell m} + P^{\ell m} Y_{,\theta}^{\ell m}) \\ \delta k^\phi &= \frac{\sqrt{B}}{C \sin^2 \theta} (Q^{\ell m} S_\phi^{\ell m} + P^{\ell m} Y_{,\phi}^{\ell m})\end{aligned}\quad (\text{A.4})$$

The other perturbed components are determined by conditions (2.4). According to the classifications, $R^{\ell m}, V^{\ell m}, P^{\ell m}, \delta\rho^{\ell m}, \delta P_r^{\ell m}, \delta P_t^{\ell m}$ belong to the polar perturbations, and $U^{\ell m}, Q^{\ell m}$ belong to axial perturbations. In this paper, we only consider the axial perturbations. Eventually, we obtain the components of perturbed stress tensor in tetrad form

$$\delta T_{\bar{a}\bar{b}}^t = \begin{pmatrix} \frac{2a\Omega\kappa(U^{\ell m} + h_0^{\ell m})S_\phi^{\ell m}}{A} & -\frac{a\Omega\sqrt{B}\sigma(Q^{\ell m} + h_1^{\ell m})S_\phi^{\ell m}}{\sqrt{A}} & -\frac{\sigma(U^{\ell m} + h_0^{\ell m})S_\theta^{\ell m}}{\sqrt{AC}} & -\frac{\sigma(U^{\ell m} + h_0^{\ell m})S_\phi^{\ell m}}{\sqrt{AC}\sin\theta} \\ \text{sym} & 0 & \frac{\sqrt{B}\sigma(Q^{\ell m} + h_1^{\ell m})S_\theta^{\ell m}}{\sqrt{C}} & \frac{\sqrt{B}\sigma(Q^{\ell m} + h_1^{\ell m})S_\phi^{\ell m}}{\sqrt{C}\sin\theta} \\ \text{sym} & \text{sym} & 0 & \frac{a\Omega\kappa(U^{\ell m} + h_0^{\ell m})\sin\theta S_\theta^{\ell m}}{A} \\ \text{sym} & \text{sym} & \text{sym} & \frac{2a\Omega\kappa(U^{\ell m} + h_0^{\ell m})S_\phi^{\ell m}}{A} \end{pmatrix} \quad (\text{A.5})$$

where we have defined $\kappa = \rho + P_t$ and $\sigma = P_r - P_t$. We can see from the perturbed stress tensor A.5 that $Q^{\ell m} + h_1^{\ell m}$ corresponds to the momentum flow. Since the fluid we consider has no friction, we set

$$Q^{\ell m} + h_1^{\ell m} = 0 \quad (\text{A.6})$$

B Axial perturbation variables

$$\begin{aligned}
\beta^0 &= \frac{(-BA'C' + AB'C' + 2A(BC'' + J - 2))h_0}{4(AC)^{3/2}} + \frac{(BA' - AB')h'_0}{4A^{3/2}\sqrt{C}} - \frac{Bh''_0}{2\sqrt{AC}} \\
&\quad - \frac{i\omega(-BCA' + ACB' + 2ABC')h_1}{4(AC)^{3/2}} - \frac{i\omega Bh'_1}{2\sqrt{AC}} + \frac{8\pi\kappa(U + h_0)}{\sqrt{AC}} \\
\beta^1 &= -\frac{i\omega\sqrt{B}C'h_0}{2AC^{3/2}} + \frac{i\omega\sqrt{B}h'_0}{2A\sqrt{C}} + \left(\frac{\sqrt{B}(J - 2)}{2C^{3/2}} - \frac{\sqrt{B}\omega^2}{2A\sqrt{C}} \right) h_1 - \frac{8\pi\sqrt{B}\sigma(Q + h_1)}{\sqrt{C}} \\
\chi^0 &= -\frac{i\omega Dh_0}{2(AC)^{3/2}} + \frac{(ABD' - BDA')h_1}{4(AC)^{3/2}} \\
\chi^1 &= -\frac{\sqrt{B}(CDA' + ADC' - ACD')h_0}{4A^2C^{5/2}} + \frac{\sqrt{B}Dh'_0}{4AC^{3/2}} - \frac{i\omega\sqrt{B}Dh_1}{4AC^{3/2}} \\
\tilde{\alpha}^0 &= \frac{(CDB' - 2BDC' + 3BCD')h_1}{2\sqrt{AC}^{5/2}} + \frac{BDh'_1}{\sqrt{AC}^{3/2}} \\
\tilde{\alpha}^1 &= -\frac{\sqrt{B}(CDA' + 3ADC' - 3ACD')h_0}{2A^2C^{5/2}} + \frac{\sqrt{B}Dh'_0}{2AC^{3/2}} - \frac{i\omega\sqrt{B}Dh_1}{2AC^{3/2}} \\
\eta^0 &= \frac{JB(AD' - DA')h_1}{4(AC)^{3/2}} \\
\eta^1 &= \frac{J\sqrt{B}(CDA' + 3ADC' - ACD')h_0}{4A^2C^{5/2}} - \frac{3J\sqrt{B}Dh'_0}{4AC^{3/2}} - \frac{3Ji\omega\sqrt{B}Dh_1}{4AC^{3/2}} \\
t &= \frac{i\omega h_0}{2AC} + \frac{(BA' + AB')h_1}{4AC} + \frac{Bh'_1}{2C} \\
g &= \frac{(-BCDA'C' + ACD(B'C' + 2BC'' - 4) + 2ABCC'D' - 2ABDC'^2)h_0}{4A^2C^3} \\
&\quad + \frac{(BCDA' - ACDB' + 2ABDC' - 2ABCD')h'_0}{4A^2C^2} - \frac{BDh''_0}{2AC} \\
&\quad - \frac{i\omega(-BDA' + ADB' + 2ABD')h_1}{4A^2C} - \frac{i\omega BDh'_1}{2AC} + \frac{8\pi\Omega\kappa(U + h_0)}{A}
\end{aligned} \tag{B.1}$$

References

- [1] K. D. Kokkotas and B. G. Schmidt, “Quasinormal modes of stars and black holes,” *Living Rev. Rel.* **2**, 2 (1999) [arXiv:gr-qc/9909058 [gr-qc]].
- [2] E. Berti, V. Cardoso and A. O. Starinets, “Quasinormal modes of black holes and black branes,” *Class. Quant. Grav.* **26**, 163001 (2009) [arXiv:0905.2975 [gr-qc]].
- [3] R. A. Konoplya and A. Zhidenko, “Quasinormal modes of black holes: From astrophysics to string theory,” *Rev. Mod. Phys.* **83**, 793-836 (2011) [arXiv:1102.4014 [gr-qc]].

- [4] T. Regge and J. A. Wheeler, “Stability of a Schwarzschild singularity,” *Phys. Rev.* **108**, 1063-1069 (1957)
- [5] F. J. Zerilli, “Gravitational field of a particle falling in a schwarzschild geometry analyzed in tensor harmonics,” *Phys. Rev. D* **2**, 2141-2160 (1970)
- [6] F. J. Zerilli, “Effective potential for even parity Regge-Wheeler gravitational perturbation equations,” *Phys. Rev. Lett.* **24**, 737-738 (1970)
- [7] S. A. Teukolsky, “Perturbations of a rotating black hole. 1. Fundamental equations for gravitational electromagnetic and neutrino field perturbations,” *Astrophys. J.* **185**, 635-647 (1973)
- [8] S. Chandrasekhar, “The mathematical theory of black holes” (Oxford University Press, New York, 1992)
- [9] Y. Kojima, “Equations governing the nonradial oscillations of a slowly rotating relativistic star,” *Phys. Rev. D* **46**, 4289-4303 (1992)
- [10] P. Pani, “Advanced Methods in Black-Hole Perturbation Theory,” *Int. J. Mod. Phys. A* **28**, 1340018 (2013) [arXiv:1305.6759 [gr-qc]].
- [11] P. Pani, E. Berti and L. Gualtieri, “Gravitoelectromagnetic Perturbations of Kerr-Newman Black Holes: Stability and Isospectrality in the Slow-Rotation Limit,” *Phys. Rev. Lett.* **110**, no.24, 241103 (2013) [arXiv:1304.1160 [gr-qc]].
- [12] P. Pani, E. Berti and L. Gualtieri, “Scalar, Electromagnetic and Gravitational Perturbations of Kerr-Newman Black Holes in the Slow-Rotation Limit,” *Phys. Rev. D* **88**, 064048 (2013) [arXiv:1307.7315 [gr-qc]].
- [13] L. Pierini and L. Gualtieri, “Quasi-normal modes of rotating black holes in Einstein-dilaton Gauss-Bonnet gravity: the first order in rotation,” *Phys. Rev. D* **103**, 124017 (2021) [arXiv:2103.09870 [gr-qc]].
- [14] L. Pierini and L. Gualtieri, “Quasinormal modes of rotating black holes in Einstein-dilaton Gauss-Bonnet gravity: The second order in rotation,” *Phys. Rev. D* **106**, no.10, 104009 (2022) [arXiv:2207.11267 [gr-qc]].
- [15] P. Wagle, N. Yunes and H. O. Silva, “Quasinormal modes of slowly-rotating black holes in dynamical Chern-Simons gravity,” *Phys. Rev. D* **105**, no.12, 124003 (2022) [arXiv:2103.09913 [gr-qc]].

- [16] M. Srivastava, Y. Chen and S. Shankaranarayanan, “Analytical computation of quasinormal modes of slowly rotating black holes in dynamical Chern-Simons gravity,” *Phys. Rev. D* **104**, no.6, 064034 (2021) [arXiv:2106.06209 [gr-qc]].
- [17] P. A. Cano, K. Fransen, T. Hertog and S. Maenaut, “Gravitational ringing of rotating black holes in higher-derivative gravity,” *Phys. Rev. D* **105**, no.2, 024064 (2022) [arXiv:2110.11378 [gr-qc]].
- [18] W. Liu, X. Fang, J. Jing and J. Wang, “QNMs of slowly rotating Einstein–Bumblebee black hole,” *Eur. Phys. J. C* **83**, no.1, 83 (2023) [arXiv:2211.03156 [gr-qc]].
- [19] G. Raposo, P. Pani, M. Bezares, C. Palenzuela and V. Cardoso, “Anisotropic stars as ultracompact objects in General Relativity,” *Phys. Rev. D* **99**, no.10, 104072 (2019) [arXiv:1811.07917 [gr-qc]].
- [20] Y. Kojima, ”Coupled pulsations between polar and axial modes in a slowly rotating relativistic star : astrophysics and relativity,” *Progress of Theoretical Physics* 90, 977 (1993).
- [21] P. Pani, V. Cardoso, L. Gualtieri, E. Berti and A. Ishibashi, “Perturbations of slowly rotating black holes: massive vector fields in the Kerr metric,” *Phys. Rev. D* **86**, 104017 (2012) [arXiv:1209.0773 [gr-qc]].
- [22] L. Hui, A. Podo, L. Santoni and E. Trincherini, “Effective Field Theory for the perturbations of a slowly rotating black hole,” *JHEP* **12**, 183 (2021) [arXiv:2111.02072 [hep-th]].