

Recovering Semiclassical Einstein's Equation Using Generalized Entropy

Naman Kumar*

School of Basic Sciences, Indian Institute of Technology, Bhubaneswar, India,
752050

Abstract

In this letter, we show that the Semiclassical Einstein's Field Equation can be recovered using the generalized entropy S_{gen} . This approach is reminiscent of non-equilibrium thermodynamics. Furthermore, contrary to the entanglement equilibrium approach of deriving the semiclassical Einstein's equation, this approach does not require any such assumptions and still recovers its correct form. Therefore, in a sense, we also show the validity of the semiclassical approximation, a crucial approach for establishing a number of important ideas such as the Hawking effect.

Keywords Semiclassical Gravity; Generalized Entropy; Quantum Expansion; Causal Diamond; Non-equilibrium Thermodynamics.

*namankumar5954@gmail.com

1 Introduction

The connection between gravity and thermodynamics has long been explored. The first hints came from black hole thermodynamics when Hawking[1] showed that the area of a horizon(A) is a non-decreasing function of time.

$$\frac{dA}{dt} \geq 0 \tag{1}$$

Later, Bekenstein[2] proposed the equality of horizon area and entropy as

$$S = \gamma A \tag{2}$$

where γ is a constant of proportionality. Hawking[3] fixed this constant of proportionality by deriving the black hole temperature as¹

$$\gamma = \frac{1}{4G\hbar} \tag{3}$$

where G is the universal gravitational constant and \hbar is the reduced Planck's constant. In 1995, Jacobson[4] derived Einstein's field equation from this proportionality of horizon area and entropy and the fundamental thermodynamic result $dQ = TdS$ under equilibrium conditions. Some works under the non-equilibrium conditions were also presented later[5, 6]. Jacobson took Q as the heat flux across a causal horizon while temperature T was given by Unruh temperature[7] as observed by a local Rindler observer. Using these ingredients, he was able to derive Einstein's field equation in the form

$$R_{ab} - \frac{1}{2}Rg_{ab} + \Lambda g_{ab} = \frac{2\pi}{\hbar\eta}T_{ab} \tag{4}$$

where the constant $\eta = (4G\hbar)^{-1}$ and the constant Λ was identified as the cosmological constant. The connection between gravity and thermodynamics was further explored by Padmanabhan (see [8] for a review) with some closely related follow-up articles from his collaborators[9, 10, 11, 12]. Recently, Verlinde[13] combined thermodynamic arguments and the holographic principle[14, 15] and argued that gravity is an entropic force arising from the underlying microscopic theory to maximize its entropy. Very recently in 2016, Bousso et al.[16] conjectured the Quantum Focusing(QFC) which conjectures that the quantum expansion(Θ) cannot increase along any congruence, even in quantum states that would violate the classical focussing theorem:

$$\frac{d\Theta}{d\lambda} \leq 0 \tag{5}$$

The quantum expansion allows to generalize the classical focusing theorem to the semiclassical case. The QFC has a wide variety of applications. It implies a Quantum Bousso Bound which has already gathered a considerable amount of evidence[17, 18, 19, 20, 21, 16, 22, 23]. It also implies the quantum singularity theorems[24, 25], the generalized second law of causal horizons and holographic screens[26] and new property of nongravitational theories, the Quantum Null Energy Condition(QNEC)[16, 27, 28, 29]. Arvin[30] presented a restricted quantum focusing which he argued is sufficient to derive all known essential implications of the quantum focusing and also proved it on brane-world semiclassical gravity theories. The Semiclassical Einstein's Field equation was derived using the entanglement equilibrium approach[31, 32] which was essentially based on two assumptions: first, the vacuum reduced to a small ball is maximal when we vary it keeping the volume fixed and second, the variation of the entanglement entropy coming from the variation of the matter fields takes the following particular form

$$\delta S_{IR} = 2\pi \frac{\Omega_{d-2}R^d}{d^2 - 1} \left(\delta \langle T_{00} \rangle - \delta \langle X \rangle \right) \tag{6}$$

where X is an operator in the QFT and the remaining symbols have the usual meaning. This second assumption was critically analyzed and was found to be somewhat problematic to an extent

¹In this paper we set $k_B = c = 1$

such that the entanglement equilibrium approach can only reveal linearized Einstein's equation or the full non-linear equation by restricting to some special choices of states(see [33] for details). In this letter, we show that the generalized entropy approach can bypass all such assumptions and we can recover the full non-linear Semiclassical Einstein's Field equation which reflects on the naturalness and robustness of our approach. In the next section, we present our idea elaborately.

2 Recovering the Semiclassical Einstein's Field Equation

Hawking[3] showed that black holes could evaporate, thus their horizon area can decrease via Hawking radiation. One of the key assumptions he made is the validity of semiclassical Einstein's field equation in the semiclassical regime given by

$$G_{\mu\nu} = \frac{8\pi G}{c^4} \langle T_{\mu\nu} \rangle \quad (7)$$

where $G_{\mu\nu}$ is the Einstein tensor and $\langle T_{\mu\nu} \rangle$ is the expectation value of the energy-momentum tensor. We show how to recover this critical equation, thus establishing the validity of the semiclassical approximation under the assumption that the fluctuations in T_{ab} are negligible.

We consider a causal diamond in the large limit in the de-Sitter static patch (see Fig.(1)). In this limit, the boundary of the diamond coincides with the cosmological horizon of the dS space², that is, $r = L$. The conformal killing vector ζ^a then becomes the timelike killing vector and \mathcal{H} the killing horizon[34].




Figure 1: A causal diamond in the de-Sitter static patch for a ball-shaped spacelike surface Σ with $d\Sigma$ as its boundary. p and p' are the future and past vertices of the diamond. ζ is a timelike killing vector in the large limit of the diamond when the boundary of the diamond coincides with the cosmological horizon. The dashed curves are the flow lines of the killing vector ζ . \mathcal{H} is the killing horizon.

The generalized entropy S_{gen} is defined as

$$S_{gen}[\mathcal{H}] = \frac{A[\mathcal{H}]}{4G\hbar} + S_{out}[\mathcal{H}] \quad (8)$$

where $S_{out}[\mathcal{H}]$ is the entropy restricted to one side of \mathcal{H} (here, the matter entropy outside the horizon \mathcal{H} of the causal diamond). The Quantum Expansion (Θ) is defined similarly to the classical expansion θ as the response of S_{gen} to deformations of \mathcal{H} along the generator and the amount of deformation is measured by an affine parameter λ along the generator. This translates to the definition of Θ as

$$\Theta = \lim_{\mathcal{A} \rightarrow 0} \frac{4G\hbar}{\mathcal{A}} \frac{dS_{gen}}{d\lambda} \quad (9)$$

²The metric for a static patch in de-Sitter space is given by $ds^2 = -[1 - (r/L)^2]dt^2 + [1 - (r/L)^2]^{-1}dr^2 + r^2d\Omega_2^2$

Therefore, the variation of the generalized entropy is given by

$$\delta S_{gen} = \eta \int_{\mathcal{H}} \Theta d\lambda d\mathcal{A} \quad (10)$$

where $\eta = (4G\hbar)^{-1}$. The cosmological horizon \mathcal{H} has a temperature given by

$$T = \frac{\hbar\kappa}{2\pi} \quad (11)$$

where κ is the surface gravity. Since the interior of the causal diamond is unobservable as it is causally disconnected from the exterior, the integrated energy of the energy current of the matter given by $\langle T_{ab} \rangle \zeta^a$, is identified as heat[4]. The energy flux across \mathcal{H} can therefore be written as

$$\delta Q = \int_{\mathcal{H}} d\Sigma^a \langle T_{ab} \rangle \zeta^b \quad (12)$$

$$= -2\kappa \int_{\mathcal{H}} \lambda \langle T_{ab} \rangle k^a k^b d\lambda d\mathcal{A} \quad (13)$$

where k^a is the (null) tangent vector to the horizon generators for an affine parameter λ so that $\zeta^a = -2\kappa\lambda k^a$. Note the extra factor of 2 in the equation of heat (13). We recall that the relation between affine parameter λ and killing parameter v on a killing horizon is generally given as $\lambda = -e^{-\kappa v}$. But as we show here, in the semiclassical case, we should use instead $\lambda = -e^{-2\kappa v}$ so as to recover the semiclassical equation. This can be understood as the vanishing of quantum expansion Θ to zeroth order in λ must occur at twice the rate to correctly impose the condition on spacetime curvature. Therefore, we get $\zeta^a = -2\kappa\lambda k^a$, and δQ takes the form as given in (13). In the semiclassical case, the natural generalization to work with the generalized entropy is quantum expansion. The infinitesimal evolution of Θ is given by a linear expansion around its equilibrium value at p , up to the first order in λ , as it moves away from the equilibrium surface at $\lambda = 0$ along the generator

$$\Theta = \Theta_p + \lambda \left. \frac{d\Theta}{d\lambda} \right|_p + \mathcal{O}(\lambda^2) \quad (14)$$

To find Θ in terms of known quantities, we use its definition as[16]

$$\Theta = \theta + \frac{4G\hbar}{\mathcal{A}} S'_{out} \quad (15)$$

where θ is the classical expansion and S_{out} the outside matter entropy. The evolution of Quantum Expansion Θ is governed by

$$\frac{d\Theta}{d\lambda} = \frac{d\theta}{d\lambda} + \frac{d}{d\lambda} \left(\frac{4G\hbar}{\mathcal{A}} S'_{out} \right) \quad (16)$$

$$= -\frac{1}{2}\theta^2 - \xi^2 - R_{ab}k^a k^b + \frac{d}{d\lambda} \left(\frac{4G\hbar}{\mathcal{A}} S'_{out} \right) \quad (17)$$

Therefore (10) becomes

$$\delta S_{gen} = \eta \int_{\mathcal{H}} \left[\Theta_p - \lambda \left(\frac{\theta^2}{2} + \xi^2 + R_{ab}k^a k^b - \frac{d}{d\lambda} \frac{4G\hbar}{\mathcal{A}} S'_{out} \right) \right] \Big|_p d\lambda d\mathcal{A} \quad (18)$$

Since at zero order³ in λ , $\Theta_p = 0$, we have

$$\theta_p = - \left. \frac{4G\hbar}{\mathcal{A}} S'_{out} \right|_p \quad (19)$$

³As can be noted here. In contrast with the classical case, the semiclassical case requires the vanishing of quantum expansion Θ to zeroth order and not the classical expansion θ for the semiclassical Einstein's equation to hold. This can be understood as follows: The quantum fields violate the classical focusing while even in this the quantum focusing ($d\Theta/d\lambda \leq 0$) holds. Therefore, focusing to the past of p must bring the quantum expansion to zero so that the increase in generalized entropy S_{gen} is proportional to the killing energy across it. This imposes a condition on the spacetime curvature that is governed by the evolution equation (17). The same cannot be said for the classical focusing since it does not hold in the semiclassical case.

We therefore get

$$\delta S_{gen} = \eta \int_{\mathcal{H}} \left[-2\lambda \left(\frac{\theta^2}{2} + \xi^2 + R_{ab} k^a k^b \right) \right] d\lambda d\mathcal{A} \quad (20)$$

Before we use the Clausius relation $\delta Q = T\delta S$, let us write δS_{gen} as $\delta S_{gen} = d_i S + d_e S$, where

$$d_i S = -2\eta \int_{\mathcal{H}} \lambda \left(\frac{\theta^2}{2} + \xi^2 \right) d\lambda d\mathcal{A} \quad (21)$$

and

$$d_e S = -2\eta \int_{\mathcal{H}} \lambda R_{ab} k^a k^b d\lambda d\mathcal{A} \quad (22)$$

$d_i S$ is identified as the internal entropy production rate due to some dissipation [6, 35] and it must be positive as required by the second law. Interestingly, this is twice that of the classical one and has contributions from both scalar and tensorial degrees of freedom. To see why $d_e S$ and $d_i S$ take these expressions can be understood by expressing (21) in terms of the Killing parameter v as

$$d_i S = \frac{\eta}{\kappa} \int_{\mathcal{H}} dv d\mathcal{A} \left(\frac{\theta^2}{2} + \xi^2 \right) \geq 0 \quad (23)$$

as required by the second law (since the affine parameter λ and Killing parameter v on the horizon are related by $\lambda = -e^{-2\kappa v}$). $d_e S$ is then identified as the reversible entropy change dS_{rev} . Therefore, by subtracting $d_i S$ from δS_{gen} and using the Clausius relation $\delta Q = TdS_{rev} = Td_e S$, we obtain

$$\langle T_{ab} \rangle k^a k^b = \frac{\hbar\eta}{2\pi} R_{ab} k^a k^b \quad (24)$$

This equation implies

$$\frac{2\pi}{\hbar\eta} \langle T_{ab} \rangle = R_{ab} + f g_{ab} \quad (25)$$

for some function f . Local conservation implies $\langle T_{ab} \rangle$ is divergenceless, therefore, using the Bianchi identity, we recover the semiclassical Einstein's equation

$$R_{ab} - \frac{1}{2} R g_{ab} + \Lambda g_{ab} = \frac{2\pi}{\hbar\eta} \langle T_{ab} \rangle \quad (26)$$

where Λ is identified as the cosmological constant of the de-Sitter space and $\eta = (4G\hbar)^{-1}$. Therefore, the constant Λ obtained in this case also carries a natural meaning.

3 Conclusion

In this work, we showed that the semiclassical Einstein's field equation can be recovered in the semiclassical case using the generalized entropy S_{gen} . Since S_{gen} is a cutoff independent quantity, it reveals information about the full quantum gravity theory and shows that the semiclassical approximation can be trusted as long as the fluctuations in T_{ab} are negligible. Furthermore, we showed that this approach is reminiscent of non-equilibrium thermodynamics. The validity of the semiclassical equation has been previously established using the entanglement equilibrium approach. However, this approach applied to causal diamonds makes a number of assumptions and some of which turned out to be problematic. On the contrary, we need no such assumptions and still, we were successful in recovering the semiclassical equation making this approach natural and robust.

Acknowledgements

We would like to thank the anonymous referee who provided detailed comments and useful suggestions that greatly improved the quality of the manuscript.

Conflict of Interest

The author declares no conflict of interest.

References

- [1] Stephen W Hawking. Gravitational radiation from colliding black holes. *Physical Review Letters*, 26(21):1344, 1971.
- [2] Jacob D. Bekenstein. Black holes and entropy. *Phys. Rev. D*, 7:2333–2346, Apr 1973.
- [3] S. W. Hawking. Particle creation by black holes. *Communications in Mathematical Physics*, 43(3):199–220, August 1975.
- [4] Jacobson. Thermodynamics of spacetime: The einstein equation of state. *Physical review letters*, 75 7:1260–1263, 1995.
- [5] Christopher Eling, Raf Guedens, and Ted Jacobson. Nonequilibrium thermodynamics of spacetime. *Physical Review Letters*, 96(12):121301, 2006.
- [6] Goffredo Chirco and Stefano Liberati. Nonequilibrium thermodynamics of spacetime: The role of gravitational dissipation. *Physical Review D*, 81(2):024016, 2010.
- [7] William G Unruh. Notes on black-hole evaporation. *Physical Review D*, 14(4):870, 1976.
- [8] Thanu Padmanabhan. Thermodynamical aspects of gravity: new insights. *Reports on Progress in Physics*, 73(4):046901, 2010.
- [9] Ayan Mukhopadhyay and T Padmanabhan. Holography of gravitational action functionals. *Physical Review D*, 74(12):124023, 2006.
- [10] Dawood Kothawala, Sudipta Sarkar, and T Padmanabhan. Einstein’s equations as a thermodynamic identity: The cases of stationary axisymmetric horizons and evolving spherically symmetric horizons. *Physics Letters B*, 652(5-6):338–342, 2007.
- [11] Dawood Kothawala and T Padmanabhan. Thermodynamic structure of lanczos-lovelock field equations from near-horizon symmetries. *Physical Review D*, 79(10):104020, 2009.
- [12] Sanved Kolekar and T Padmanabhan. Holography in action. *Physical Review D*, 82(2):024036, 2010.
- [13] Erik Verlinde. On the origin of gravity and the laws of newton. *Journal of High Energy Physics*, 2011(4):1–27, 2011.
- [14] Leonard Susskind. The world as a hologram. *Journal of Mathematical Physics*, 36(11):6377–6396, 1995.
- [15] Gerard’t Hooft. Dimensional reduction in quantum gravity. *arXiv preprint gr-qc/9310026*, 1993.
- [16] Raphael Bousso, Zachary Fisher, Stefan Leichenauer, and Aron C Wall. Quantum focusing conjecture. *Physical Review D*, 93(6):064044, 2016.
- [17] Raphael Bousso. A covariant entropy conjecture. *Journal of High Energy Physics*, 1999(07):004, 1999.
- [18] Raphael Bousso. The holographic principle. *Reviews of Modern Physics*, 74(3):825, 2002.
- [19] Raphael Bousso, Eanna E Flanagan, and Donald Marolf. Simple sufficient conditions for the generalized covariant entropy bound. *Physical Review D*, 68(6):064001, 2003.

- [20] Raphael Bousso, Horacio Casini, Zachary Fisher, and Juan Maldacena. Proof of a quantum bousso bound. *Physical Review D*, 90(4):044002, 2014.
- [21] Raphael Bousso, Horacio Casini, Zachary Fisher, and Juan Maldacena. Entropy on a null surface for interacting quantum field theories and the bousso bound. *Physical Review D*, 91(8):084030, 2015.
- [22] Eanna E Flanagan, Donald Marolf, and Robert M Wald. Proof of classical versions of the bousso entropy bound and of the generalized second law. *Physical Review D*, 62(8):084035, 2000.
- [23] Andrew Strominger and David Thompson. Quantum bousso bound. *Physical Review D*, 70(4):044007, 2004.
- [24] Aron C Wall. The generalized second law implies a quantum singularity theorem. *Classical and Quantum Gravity*, 30(16):165003, 2013.
- [25] Raphael Bousso and Arvin Shahbazi-Moghaddam. Quantum singularities. *arXiv preprint arXiv:2206.07001*, 2022.
- [26] Raphael Bousso and Netta Engelhardt. Generalized second law for cosmology. *Physical Review D*, 93(2):024025, 2016.
- [27] Raphael Bousso, Zachary Fisher, Jason Koeller, Stefan Leichenauer, and Aron C Wall. Proof of the quantum null energy condition. *Physical Review D*, 93(2):024017, 2016.
- [28] Srivatsan Balakrishnan, Thomas Faulkner, Zuhair U Khandker, and Huajia Wang. A general proof of the quantum null energy condition. *Journal of High Energy Physics*, 2019(9):1–86, 2019.
- [29] Fikret Ceyhan and Thomas Faulkner. Recovering the qnec from the anec. *Communications in Mathematical Physics*, 377:999–1045, 2020.
- [30] Arvin Shahbazi-Moghaddam. Restricted quantum focusing. *arXiv preprint arXiv:2212.03881*, 2022.
- [31] Ted Jacobson. Entanglement equilibrium and the einstein equation. *Physical review letters*, 116(20):201101, 2016.
- [32] Pablo Bueno, Vincent S Min, Antony J Speranza, and Manus R Visser. Entanglement equilibrium for higher order gravity. *Physical Review D*, 95(4):046003, 2017.
- [33] Horacio Casini, Damián A Galante, and Robert C Myers. Comments on jacobson’s “entanglement equilibrium and the einstein equation”. *Journal of High Energy Physics*, 2016(3):1–34, 2016.
- [34] Theodore Jacobson and Manus R Visser. Gravitational thermodynamics of causal diamonds in (a) ds. *SciPost Physics*, 7(6):079, 2019.
- [35] Ramit Dey, Stefano Liberati, and Daniele Pranzetti. Spacetime thermodynamics in the presence of torsion. *Physical Review D*, 96(12):124032, 2017.