

A JWST Medium Resolution MIRI Spectrum and Models of the Type Ia supernova 2021aefx at +415 d

C. ASHALL ¹, P. HOEFELICH ², E. BARON ^{3,4}, M. SHAHBANDEH ⁵, J. M. DERKACY ¹, K. MEDLER ¹, B. J. SHAPPEE ⁶,
M. A. TUCKER ^{7,*}, E. FEREIDOUNI ², T. MERA ², J. ANDREWS ⁸, D. BAADE ⁹, K. A. BOSTROEM ^{10,†}, P. J. BROWN ¹¹,
C. R. BURNS ¹², A. BURROW ¹³, A. CIKOTA ¹⁴, T. DE JAEGER ¹⁵, A. DO ¹⁶, Y. DONG ¹⁷, I. DOMINGUEZ ¹⁸, O. FOX ¹⁹,
L. GALBANY ^{19,20}, E. Y. HSIAO ², K. KRISCIUNAS ¹¹, B. KHAGHANI ¹, S. KUMAR ²¹, J. LU ²², J. R. MAUND ²³,
P. MAZZALI ^{24,25}, N. MORRELL ²⁶, F. PATAT ⁹, C. PFEFFER ¹, M. M. PHILLIPS ²⁶, J. SCHMIDT ²⁷, S. STANGL ¹³,
C. P. STEVENS ¹, M. D. STRITZINGER ²⁸, N. B. SUNTZEFF ¹¹, C. M. TELESKO ²⁹, L. WANG ³⁰, AND Y. YANG ^{31,‡}

¹Department of Physics, Virginia Tech, Blacksburg, VA 24061, USA

²Department of Physics, Florida State University, 77 Chieftan Way, Tallahassee, FL 32306, USA

³Planetary Science Institute, 1700 East Fort Lowell Road, Suite 106, Tucson, AZ 85719-2395 USA

⁴Hamburger Sternwarte, Gojenbergsweg 112, D-21029 Hamburg, Germany

⁵Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218-2410, USA

⁶Institute for Astronomy, University of Hawai'i at Manoa, 2680 Woodlawn Dr., Hawai'i, HI 96822, USA

⁷Center for Cosmology and AstroParticle Physics, The Ohio State University, 191 W. Woodruff Ave., Columbus, OH 43210, USA

⁸Gemini Observatory/NSF's NOIRLab, 670 North A'ohoku Place, Hilo, HI 96720-2700, USA

⁹European Organization for Astronomical Research in the Southern Hemisphere (ESO), Karl-Schwarzschild-Str. 2, 85748 Garching b. München, Germany

¹⁰Steward Observatory, University of Arizona, 933 North Cherry Avenue, Tucson, AZ 85721-0065, USA

¹¹George P. and Cynthia Woods Mitchell Institute for Fundamental Physics and Astronomy, Texas A&M University, Department of Physics and Astronomy, College Station, TX 77843, USA

¹²Observatories of the Carnegie Institution for Science, 813 Santa Barbara Street, Pasadena, CA 91101, USA

¹³Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, 440 W. Brooks, Rm 100, Norman, OK 73019-2061, USA

¹⁴Gemini Observatory/NSF's NOIRLab, Casilla 603, La Serena, Chile

¹⁵Sorbonne Université, CNRS/IN2P3, LPNHE, F-75005, Paris, France

¹⁶Institute of Astronomy and Kavli Institute for Cosmology, Madingley Road, Cambridge, CB3 0HA, UK

¹⁷Department of Physics, University of California, 1 Shields Avenue, Davis, CA 95616-5270, USA

¹⁸Universidad de Granada, 18071, Granada, Spain

¹⁹Institute of Space Sciences (ICE, CSIC), Campus UAB, Carrer de Can Magrans, s/n, E-08193 Barcelona, Spain

²⁰Institut d'Estudis Espacials de Catalunya (IEEC), E-08034 Barcelona, Spain

²¹Department of Astronomy, University of Virginia, 530 McCormick Rd, Charlottesville, VA 22904, USA

²²Department of Physics and Astronomy, Michigan State University, East Lansing, MI 48824, USA

²³Department of Physics and Astronomy, University of Sheffield, Hicks Building, Hounsfield Road, Sheffield S3 7RH, U.K.

²⁴Astrophysics Research Institute, Liverpool John Moores University, UK

²⁵Max-Planck Institute for Astrophysics, Garching, Germany

²⁶Las Campanas Observatory, Carnegie Observatories, Casilla 601, La Serena, Chile

²⁷Citizen scientist

²⁸Department of Physics and Astronomy, Aarhus University, Ny Munkegade 120, DK-8000 Aarhus C, Denmark

²⁹Department of Astronomy, University of Florida, Gainesville, FL 32611 USA

³⁰Department of Physics and Astronomy, Texas A&M University, College Station, TX 77843, USA

³¹Department of Astronomy, University of California, Berkeley, CA 94720-3411, USA

(Received xxx; Revised xxx; Accepted xxx)

Submitted to ApJ

ABSTRACT

We present a JWST MIRI/MRS spectrum (5–27 μm) of the Type Ia supernova (SN Ia), SN 2021aefx at +415 days past *B*-band maximum. The spectrum, which was obtained during the iron-dominated nebular phase, has

been analyzed in combination with previous *JWST* observations of SN 2021aefx, to provide the first *JWST* time series analysis of an SN Ia. We find the temporal evolution of the [Co III] 11.888 μm feature directly traces the decay of ^{56}Co . The spectra, line profiles, and their evolution are analyzed with off-center delayed-detonation models. Best fits were obtained with White Dwarf (WD) central densities of $\rho_c = 0.9 - 1.1 \times 10^9 \text{ g cm}^{-3}$, a WD mass of $M_{\text{WD}} = 1.33 - 1.35 M_{\odot}$, a WD magnetic field of $\approx 10^6 \text{ G}$, and an off-center deflagration-to-detonation transition at $\approx 0.5 M_{\odot}$ seen opposite to the line of sight of the observer. The inner electron capture core is dominated by energy deposition from γ -rays whereas a broader region is dominated by positron deposition, placing SN 2021aefx at +415 d in the transitional phase of the evolution to the positron-dominated regime. The formerly “flat-tilted” profile at 9 μm now has significant contribution from [Ni IV], [Fe II], and [Fe III] and less from [Ar III], which alters the shape of the feature as positrons excite mostly the low-velocity Ar. Overall, the strength of the stable Ni features in the spectrum is dominated by positron transport rather than the Ni mass. Based on multi-dimensional models, our analysis strongly supports a single-spot, close-to-central ignition with an indication for a pre-existing turbulent velocity field, and excludes a multiple-spot, off-center ignition.

Keywords: supernovae: general - supernovae: individual (SN 2021aefx), *JWST*

1. INTRODUCTION

Type Ia Supernovae (SNe Ia) originate from the thermonuclear disruption of a carbon-oxygen (C-O) White Dwarf (WD) in a multiple star system (Hoyle & Fowler 1960); yet to date, the exact details of their progenitor scenarios or explosion mechanisms are unknown. Determining the exact origin of SNe Ia is essential if we are to understand the nucleosynthesis of heavy elements and improve the use of SNe Ia as extragalactic distance indicators (see Seitzzahl & Townsley 2017 and Riess 2017 for recent reviews).

Potential SNe Ia progenitor scenarios include: (1) the single degenerate (SD) scenario which consists of a C-O WD and a non-degenerate companion star (Whelan & Iben 1973), (2) the double degenerate (DD) scenario which consists of two WDs (Iben & Tutukov 1984; Webbink 1984), and (3) a triple/quadruple system consisting of at least two C-O WDs (Thompson 2011; Pejcha et al. 2013). There is also a complex interplay between the progenitor scenario and the explosion mechanism, where many explosion mechanisms can theoretically occur within each progenitor scenario. Two of the leading explosion mechanisms include the explosion of a near Chandrasekhar mass (M_{Ch}) WD (Iben & Tutukov 1984; Khokhlov 1991) and the detonation of a sub- M_{Ch} WD (Livne & Arnett 1995). Both of which can occur in SD and DD systems. In the near- M_{Ch} explosion the WD accretes H, He, or C material from a non-degenerate or degenerate companion star until it approaches the M_{Ch} , during which time densities in the center of the star become high enough for a simmering stage to occur and a thermonuclear disruption begins (Khokhlov 1991). The flame can propagate as a deflagration, detonation or both via a deflagration-to-detonation transition (DDT)

(Khokhlov 1991; Hoeflich et al. 1995; Gamezo et al. 2003; Poludnenko et al. 2019). In the sub- M_{Ch} scenario, a surface He layer detonates and produces an inwards shockwave that disrupts the whole WD. This can occur for a variety of core masses (0.6-1.1 M_{\odot}) and He shell (0.01-0.2 M_{\odot}) masses. Although only the higher mass WDs and smaller He shell masses are expected to reproduce the observed properties of SNe Ia (Livne & Arnett 1995; Shen et al. 2018; Boos et al. 2021).

Observations of SNe Ia during the nebular phase reveal their inner layers. Spectra at these epochs can be used to measure the high-density burning regions in the ejecta (e.g. Axelrod 1980; Mazzali et al. 2007; Ashall et al. 2016; Mazzali et al. 2020; Jerkstrand et al. 2020; Maguire et al. 2018; Hoeflich et al. 2021; Kumar et al. 2023). A spectral region of particular interest is at mid-infrared (MIR) wavelengths ($\sim 5\text{--}27 \mu\text{m}$), as it contains lines from critical ions that do not have suitable transitions in the optical or near-infrared (NIR). These ions can be used to distinguish between the leading progenitor and explosion scenarios. Prior to the launch of the *James Webb Space Telescope* (*JWST*) there were four SNe Ia which had MIR ($\lambda > 5 \mu\text{m}$) spectral observations, and there were seven published MIR spectra of SNe Ia in total (see the introduction of DerKacy et al. (2023a) for more details). These data provided new insights into SNe Ia explosions but the interpretation was hampered by low S/N and low spectral resolution.

The launch of *JWST* has opened up a new era of transient astronomy. To date, *JWST* spectra of three nearby SNe Ia have been published. These are: SN 2022xkq (DerKacy et al. 2024a), SN 2022pul (Siebert et al. 2024; Kwok et al. 2023a), and SN 2021aefx (Kwok et al. 2023b; DerKacy et al. 2023a). Of these, SN 2021aefx is the best observed SN Ia. Broad-band NIR+MIR *JWST* imaging was obtained at +255 d after *B*-band maximum, which enabled the late time NIR+MIR decline rates and MIR flux contribution in

* CCAPP Fellow

† LSSTC Catalyst Fellow

‡ Bengier-Winslow-Robertson Postdoctoral Fellow




Figure 1. A stacked composite image of NGC 1566 with photometric data obtained with *JWST*’s MIRI instruments, individual bands are published in (Chen et al. 2023). Here, SN 2021aefx is at a phases of +357 d past *B*-band maximum and is highlighted in the white circle. North is 1.5° counter-clockwise from left. An inset around the SN is shown in the bottom right corner. For the original image see [here](#).

the explosion to be determined (Mayker Chen et al. 2023). Spectra of SN 2021aefx were acquired at +255 d (Kwok et al. 2023b), and +323 d (DerKacy et al. 2023a) past *B*-band maximum. These spectra revealed many unique features including multiple stable Ni lines, which are indicative of high-density burning, a “flat-tilted” [Ar III] $8.991\ \mu\text{m}$ profile, and a strong [Co III] $11.888\ \mu\text{m}$ resonance feature (Kwok et al. 2023b; DerKacy et al. 2023a). Spectral modeling of the *JWST* data indicated that SN 2021aefx is consistent with a delayed detonation near- M_{Ch} explosion of a C-O WD, that had an off-center DDT, and produced $5.9 \times 10^{-2}\ M_\odot$ of ^{58}Ni (DerKacy et al. 2023a; although see Blondin et al. 2023). However, all of the previous spectra of SN 2021aefx were obtained with the low-resolution spectral mode and at wavelengths less than $14\ \mu\text{m}$. The medium resolution spectrometer (MRS) on the Mid-Infrared Instrument (MIRI) has a resolving power of ~ 2700 , a wavelength coverage from $\sim 5\text{--}27\ \mu\text{m}$ and is therefore ideal for obtaining precision velocity measurements as well as observations at long wavelengths.

Here we present the first observed *JWST* MIRI/MRS spectrum of a SN Ia¹. In § 2 we discuss the observations and data reduction. In § 3 we present line identification, followed by spectral comparisons in § 4. In § 5 we discuss the time series spectral evolution. In § 6 we use self-consistent multi-dimensional models to produce synthetic spectra, where we demonstrate how the MIRI/MRS data can be used in conjunction with synthetic spectra and line profiles to understand the underlying explosion physics, and the conditions at the thermonuclear runaway. A summary of our results and our conclusions are presented in § 7.

2. OBSERVATIONS & DATA REDUCTION

SN 2021aefx was discovered on 2021 Nov 11.3 (MJD=59529.5) by the Distance Less Than 40 Mpc Survey (Tartaglia et al. 2018). It was located at $\alpha = 04^{\text{h}}19^{\text{m}}53^{\text{s}}.40$, $\delta = -54^\circ 56' 53''.09$, south-west of the center of its host galaxy NCG 1566 ($z=0.0050$). SN 2021aefx is a normal luminosity

¹ The first published *JWST* MIRI/MRS spectra of a SN Ia was SN 2022xkq (DerKacy et al. 2024a).

Table 1. Log of *JWST* observations. The spectrum was taken at MJD=59964.34 which is rest-frame +415 d relative to *B*-band maximum.

Parameter	Value	Value	Value
MIRI/MRS Spectra			
Sub-band	Short	Medium	Long
Groups per Integration	35	35	36
Integrations per Exp.	3	3	3
Exposures per Dither	1	1	1
Total Dithers	12	12	12
Exp Time [s]	10224.88	10224.88	10511.565
Readout Pattern	SLOWR1	SLOWR1	SLOWR1

SN Ia (Ashall et al. 2022; Hosseinzadeh et al. 2022; Ni et al. 2023). The location of SN 2021aefx in its host galaxy is shown in Fig. 1.

JWST MIRI/MRS spectral observations of SN 2021aefx were triggered through *JWST*-GO-2114 (Ashall et al. 2021). Observations began on 20th January 2023 at 03:34:38 (MJD=59964.15), and ended on 20th January 2023 at 12:47:50 (MJD=59964.53). We take our time of observation as the mid-point, MJD = 59964.34. Throughout this work, we use a time of *B*-band maximum of MJD = 59547.25 (DerKacy et al. 2023a) implying the spectrum was obtained at rest-frame +415 d past *B*-band maximum light.

Spectral data were acquired using the MRS on MIRI with the short, medium, and long gratings and with each channel to produce a continuous spectrum from $\sim 5\text{--}27\ \mu\text{m}$. The details of the instrument setup can be found in Table 1. The total exposure time was 8.6 hours.

The data were reduced using a custom-built pipeline² designed to extract observations of faint point sources that have complex backgrounds in MIRI/MRS data cubes (see Shahbandeh et al. 2024). In short, the pipeline creates a master background based on 20 different positions away from the source within the field of view. This master background is subtracted from the whole data cube. The SN flux is then extracted along the data cube, using the *Extract1dStep* in stage 3 of the *JWST* reduction pipeline. The resulting data cube subtraction is shown in Fig. 2. All the previously unpublished data used in this paper can be found in this Digital Object Identifier (DOI)³.

With this custom-made pipeline the flux level in Channel 4 (which covers wavelengths longer than $17.7\ \mu\text{m}$) is uncertain due to the dominant instrumental background flux which

increases with wavelength. Therefore, for the spectral analysis, we choose to extract the data from Channel 4 separately using a manually-selected background region, to ensure that the continuum level is close to flat, as would be expected from a normal SN Ia in the nebular phase (see Appendix A for more details). We emphasize that this may cause large uncertainties in terms of line strengths and ratios in Channel 4, and the reduction of Channel 4 data may change as the *JWST* MIRI/MRS pipeline improves. The reduction shown here utilized version 1.9.4 of the *JWST* Calibration pipeline (Bushouse et al. 2023) and Calibration Reference Data System files version 11.16.20.

The final spectrum has been smoothed with *Specextractor* (Burrow et al. 2020) channel-by-channel to properly account for the differences in resolution across the full MIRI/MRS wavelength coverage. Throughout this work, the spectra were corrected to rest frame using the recessional velocity of $1500\ \text{km s}^{-1}$, and a rotational galactic velocity of $65 \pm 60\ \text{km s}^{-1}$ at the location of the SN (Elagali et al. 2019).

Comparing the MIRI/low-resolution spectrometer (LRS) and MIRI/MRS spectra of SN 2021aefx is useful for checking both the flux and wavelength calibration. It is known from the comparison between *JWST* MIRI photometry and *JWST* MIRI/LRS spectra that the absolute flux calibration of the MIRI/LRS data of SN 2021aefx is within 2% (Kwok et al. 2023b; Chen et al. 2023). Generally, for MIRI/MRS data it is thought that the flux calibration is accurate to $5.6 \pm 0.7\%$ (Argyriou et al. 2023). This was also confirmed using MIRI/MRS data of SN 2022acko where the spectral flux in Channels 1, 2 and 3 were found to be consistent with simultaneous MIRI broad-band photometry (Shahbandeh et al. 2024). However, it is not yet known how well the pipeline extracts the SN flux in Channel 4, and how successful it is at accurately removing the background from both the instrument and the underlying host galaxy. Overall, the flux of SN 2021aefx MIRI/MRS spectrum at +415 d is lower than that at the previous epoch (see § 5). It is also apparent that the decrease in peak flux is more rapid at earlier epochs compared to later ones. This is consistent with the light curve flattening with time. Although there is no simultaneous photometry at +415 d, this time series behavior suggests that the flux calibration of the MIRI/MRS spectrum is accurate. We also note that, the wavelength calibration of MIRI/MRS spectra is accurate to within $9\ \text{km s}^{-1}$ at $5\ \mu\text{m}$ and to $27\ \text{km s}^{-1}$ at $28\ \mu\text{m}$ (Argyriou et al. 2023).

3. LINE IDENTIFICATIONS

The fully stitched 4-channel MIRI/MRS spectrum is presented in Fig. 3. The strongest lines that contribute to the formation of the spectral features have been labeled. These line identifications are based on radiation hydrodynamical models discussed in section § 6. At short wavelengths ($< 14\ \mu\text{m}$),

² https://github.com/shahbandeh/MIRI_MRS

³ <https://doi.org/10.17909/p6j0-gp74>




Figure 2. *Top panel:* MIRI/MRS cube of SN 2021aefx before background subtraction divided into 4 channels. Each channel is collapsed sum of all its slices. *Bottom panel:* MIRI/MRS cube of SN 2021aefx after background subtraction.




Figure 3. The MIRI/MRS spectrum of SN 2021aefx at +415 d past maximum light. The raw data is plotted in light gray behind, and a spectrum smoothed to the instrumental resolution is overlaid in black. The dominant ions that contribute to each feature are labeled.

the line IDs are consistent with those presented in Kwok et al. (2023b) and DerKacy et al. (2023a). However, the individual line strengths and profiles have evolved over the intervening 92 d between observations — demonstrating that the physical processes driving the spectral formation have profoundly changed (see § 6).

We identify four dominant regions of line formation, which are presented in Fig. 4. The lines contributing to each spectral region are described here, with the exact details of the line formation within each region discussed in the following subsections.

Between 6–7.5 μm the spectrum is dominated by stable Ni lines including [Ni II] 6.636 μm and [Ni I] 7.507 μm , as well as [Ar II] 6.985 μm . Between 7.5–10.0 μm lines of [Ni IV] 8.405, 8.945 μm , and [Ar III] 8.991 μm are the strongest. In the wavelength range of 10.0–12.0 μm , [Ni II] 10.682 μm ; [Co II] 10.523 μm ; and [Co III] 11.888 μm dominate. Finally, past 12.0 μm the strongest features are [Fe II] 17.936, 22.902 μm ; [Fe III] 22.925 μm ; and [Co II] 23.196 μm .

3.1. The 5.0 – 10.0 μm Region

We identify many spectral lines which contribute to the spectral formation in the region from 5.0 – 10.0 μm . In the




Figure 4. The +415 d spectrum of SN 201aeix with the main contributing ions labeled.

complex blend between 5.8–7.8 μm we find [Ni I] 5.893 μm ; [Ni II] 5.953 μm ; [Co II] 6.214 μm ; [Ni II] 6.636, 6.920 μm ; [Ar II] 6.985 μm ; [Co I] 7.045, 7.108 μm ; [Fe III] 7.147 μm ; [Co I] 7.202 μm ; [Ni III] 7.349 μm ; [Ni I] 7.507 μm ; and [Fe III] 7.791 μm . The lines contributing to the region 8.0–9.5 μm are [Fe III] 8.211 μm ; [Fe II] 8.299 μm ; [Ni IV] 8.405 μm ; [Co II] 8.644 μm ; [Fe II] 8.733 μm ; [Ni IV] 8.945 μm ; and [Ar III] 8.991 μm .

3.2. The 10.0 – 15.0 μm Region

In the 10.0 – 15.0 μm region, there are two significant areas of line formation. The first is from 10–11 μm . We identify the main ions contributing to this feature as [Fe II] 10.189 μm ; [Co II] 10.523 μm ; [Ni II] 10.682 μm ; and [Ni IV] 11.130 μm . The second region of line formation is at ~ 12 μm . The main ion contributing to this region is the ground state [Co III] 11.888 μm line, but the red edge of this feature may also have flux from [Ni I] 12.001 μm . Longwards of 12 μm there are several weaker lines primarily coming from [Co I] 12.255, 13.058, 14.356 μm ; [Co II] 14.739 μm ; [Co III] 12.681, 13.820, 14.006 μm ; [Mn II] 12.261 μm ; [Fe II] 12.642 μm ; [Ni I] 14.814 μm ; and [Ni II] 12.729 μm . SN 2021aefx does not have any contribution from [Ne II] 12.81 μm .

3.3. The 15.0 – 20.0 μm Region

In the 15.0 – 20.0 μm region, the ejecta consist of a broad area of emission between 15–17 μm which is formed from a blend of many Co features including [Co I] 16.925 μm ; [Co II] 15.459, 16.299 μm ; [Co III] 16.391 μm ; and a strong emission line profile at 18 μm . The 18 μm profile consists of [Fe II] 17.936 μm , with a redward wing that comes from [Ni II] 18.241 μm ; [Co I] 18.265 μm ; and [Co II] 18.389 μm . Going further to the red, several Co, Ni, and Fe lines contribute to the formation of the spectrum, including [Fe II] 19.007, and 19.056 μm .

3.4. The 20.0 – 26.0 μm Region

The strongest feature in the 20.0 – 27.0 μm region is at 23 μm , and is dominated by [Fe III] 22.925 μm , with some contribution from [Fe II] 22.902 μm . The blue shoulder of this feature is comprised of [Fe I] 22.297 μm . There is also a red shoulder to this feature consisting of [Co II] 23.196 μm , and [Fe III] 23.389 μm . Possible emission at longer wavelengths may come from, [Co II] 25.689 μm ; [Co III] 24.070 μm ; [Fe I] 24.042 μm ; and [Fe II] 24.519, 25.988 μm . It should be noted that these line IDs are tentative, as the background subtraction is uncertain at these wavelengths. Furthermore, the flux calibration at these wavelengths is highly uncertain and the spectrum at wavelengths > 25 μm should not be trusted until a more accurate reduction in Channel 4 is available. Finally, we note that it is not possible to confidently identify all




Figure 5. A selection of MIR SNe Ia spectra. Times relative to B -band maximum are provided next to each observation. All SNe Ia in this plot are normal SNe Ia except for SN 2022xkq which is sub-luminous.

spectral features, as data for many atomic line transitions in these MIR regions are missing. Through spectral modeling we can identify some of the unknown line strengths as well as estimates of the flux contribution from the weaker lines (see § 6.3.1 and Appendix D).

4. SPECTRAL COMPARISON

Figure 5 shows a spectral comparison plot between SN 2021aefx and a sample of published MIR SNe Ia observations. The previous sample covers the evolution of SNe Ia from +39 to +323 d past B -band maximum. All previous MIRI/LRS spectral observations of SN 2021aefx were re-reduced using the method outlined in (DerKacy et al. 2023a), but using pipeline version 1.13.4 and CRDS version 1223.pmap. This version of the pipeline and calibration files fixed previously known issues with the MIRI/LRS wavelength calibration.

All spectra share similarities and consist of broad emission features that are composed of forbidden line transitions from Ar, Co, Fe, and Ni. The strongest of these is the [Co III]




Figure 6. (*Top:*) A spectral comparison between the MIRI/MRS data of SN 2022xkq and SN 2021aefx. The dominant ions contributing to various spectral regions are highlighted with vertical colored bars. We note that SN 2022xkq is a low luminosity SN Ia. (*Bottom:*) Time series comparison of all published MIR spectra of SN 2021aefx.

11.888 μm resonance feature. However, the capabilities and power of *JWST* become apparent with the first two spectra of SN 2021aefx. These spectra were observed using MIRI/LRS ($R \sim 100$) and revealed many spectral features that had not been observed before. In particular, a “flat-tilted” [Ar III] 8.991 μm profile and multiple ionization states of Ni were observed (Kwok et al. 2023b; DerKacy et al. 2023a). The MIRI/MRS spectrum presented in this work is dominated by similar spectral features as earlier epochs, although with the higher resolution ($R \sim 2700$) we can more accurately determine the exact region of the ejecta in which these ions are located as well as examine the data for resolved features. Overall, the MIRI/MRS data allow for detailed line IDs and analysis of ejecta structure at a level not possible with MIRI/LRS data.

To date, the spectrum of SN 2022xkq at 130 d past maximum light is the only published MIRI/MRS spectrum of a SN Ia (DerKacy et al. 2024a). The top panel of Fig. 6 shows a comparison between the low-luminosity SN 2022xkq and the normal-luminosity SN 2021aefx. Although they are taken 285 d apart, many of the same ions form the spectrum, but with different strengths and ratios; reflecting the variations in ionization balances due to phase and sub-type differences. In SN 2021aefx the stable Ni features are stronger than those in

SN 2022xkq relative to both the Ar and Co features. There is also a significant contribution from Fe in the spectral formation of SN 2021aefx which is prevalent in these epochs due to the decay of ^{56}Co . The spectra of SN 2021aefx and SN 2022xkq also vary in the ionization state of Ar where for the low-luminosity SN 2022xkq [Ar II] is the dominant ionization state but for the normal-luminosity SN 2021aefx [Ar III] is the dominant ionization state of Ar. For SN 2021aefx the Ar features have significantly more line blending from Fe-group elements than in SN 2022xkq.

5. TEMPORAL EVOLUTION OF SN 2021aefx

The time evolution of the *JWST* MIR spectra of SN 2021aefx from +215 to +415 d relative to rest-frame *B* band maximum light in absolute flux is presented in the bottom panel of Fig. 6. As the data of SN 2021aefx are the first MIR time series of an SN from *JWST*, they allow us to follow the evolution of features as a function of time. We start by analyzing the [Co III] 11.888 μm feature, as it is the most prominent feature in the MIR spectra, and then proceed to the evolution of other dominant lines.

5.1. [Co III] 11.888 μm

The [Co III] 11.888 μm feature gets weaker as a function of time (see the top left panel of Fig. 7). We quantitatively ex-

amine the evolution of this feature. For simplicity, we assume the feature can be modeled by a single Gaussian function, although there is likely some blending with the [Ni II] 12.001 μm line.

The peak velocity (v_{peak}) of the [Co III] feature is $\sim 500 \text{ km s}^{-1}$ at +255 d and +323 d, and increases to $\sim 750 \text{ km s}^{-1}$ by +415 d. The values in v_{peak} are consistent between the MIRI/LRS and MIRI/MRS data, once the error due to the low spectral resolution of the LRS data is considered.

The measured FWHM of the [Co III] 11.888 μm feature increases from $\sim 11,100 \text{ km s}^{-1}$ at +255 d to $\sim 11,700 \text{ km s}^{-1}$ at +415 d. The cause of this is likely driven by the strength of the [Co III] 11.888 μm line decreasing as a function of time and the [Ni II] 12.001 μm line getting stronger.

The peak flux of the [Co III] feature decreases from $\sim 4.2 \times 10^{-18} \text{ erg cm}^{-2} \text{ s}^{-2} \text{ \AA}^{-1}$ at +255 d to $\sim 1 \times 10^{-18} \text{ erg cm}^{-2} \text{ s}^{-2} \text{ \AA}^{-1}$ at +415 d. The evolution of the feature is well described by an exponential decay, with a half-life of $76.72 \pm 4.61 \text{ d}$. This is consistent with the half-life of ^{56}Co which is 77.27 d, demonstrating that the time evolution of this feature can be used to directly trace the energy deposition of the ^{56}Co in the ejecta. This trend is not found in the optical [Co III] feature at 6200 \AA which has been found to decrease faster than the ^{56}Co cooling rate (McClelland et al. 2013). This highlights the uniqueness of the [Co III] 11.888 μm feature. It is a resonance line and all of the recombination passes through it. It also demonstrates that the [Co III] line dominates this feature throughout, as well as confirming that the flux calibration of spectra is consistent between the MIRI/LRS and MIRI/MRS data. Continued observations of this feature in Cycle 2 and 3 (DerKacy et al. 2023b, 2024b) up to +1150 days past maximum light will probe the region where radioactive ^{57}Co may dominate the heating. If there is little mixing in the inner part of the ejecta this ^{57}Co will be located in the central $\sim 1,000 \text{ km s}^{-1}$ (Hoeftlich et al. 2021). This may cause the [Co III] feature to become stronger and narrower over time. Following the evolution of this feature with future spectral observations will allow for the distribution and strength of the ^{57}Co to be characterized.

5.2. The evolution of other important features

We now turn our attention to the temporal evolution of four other features. These are the [Ni II] 7.349 μm , the [Ni IV] 8.405 μm , the [Co III] 10.523 μm , and the [Ar II] 8.991 μm regions. Due to the more complex structures and blends in these features, we choose to concentrate on the evolution of only the peak flux. Figure 8 shows the evolution of these features in velocity space, as well as their peak flux as a function of time.

In the earlier phases, the [Ni II] 7.349 μm feature appears to be dominated by one transition, but by day +323 a blue wing




Figure 7. (Top Left:) Evolution of the [Co III] 11.888 μm feature as a function of time, for simplicity it is fit with a single Gaussian profile. (Top Right:) The velocity evolution of the peak of the emission profile. (Bottom Left:) The evolution of the feature’s FWHM. (Bottom Right:) The peak flux of the feature as a function of time, along with a line of best-fit with a decaying exponential function. For all panels, if the error bars are not visible they are smaller than the markers. Furthermore, we have only plotted the fit error and not resolution error.

appears in the emission profile. In the MIRI/LRS data, this blue wing is blended with the main [Ni II] 7.349 μm feature, however, it is resolved in the MIRI/MRS data at +415 d, and shows two distinct components with many smaller overlapping emission features in the bluest wavelengths. We identify this blue feature as [Co I] 7.202 μm . It is clear through this comparison that MIRI/MRS data is required to fully resolve profiles, but even with the higher resolution, the emission profiles are a complex blend of many lines, some of which are unknown. The peak flux of this wavelength region as a function of time does not follow the radioactive decay of ^{56}Co .

The [Ni IV] 8.405 μm feature appears to be dominated by one spectral ion throughout the MIRI/LRS time series of data. However, the MIRI/MRS data reveal that there are at least two resolved profiles contributing to this feature, one closer to the rest wavelength of [Ni IV] 8.405 μm and one at $\sim 9000 \text{ km s}^{-1}$. We identify the blue feature to be a blend of [Co I] 8.283 μm and [Fe II] 8.299 μm . The peak flux of this feature does decrease as a function of time, but unlike the [Co III] 11.888 μm feature, it does not follow the half-life of ^{56}Co .

The [Co III] 10.523 μm feature is the second most prominent [Co III] region in the MIR spectrum, the shape of this profile

changes dramatically between the MIRI/LRS and MIRI/MRS data. It is much narrower and peaked within the MIRI/MRS data. The evolution of the peak flux of this feature follows a half-life of 206.50 ± 22.79 d, which is more than twice that of what would be expected of ^{56}Co decay. This demonstrates the uniqueness of the $[\text{Co III}]$ $11.888 \mu\text{m}$ feature, as it is both isolated and a resonance line, so all recombination passes through the transition. It also highlights that an apparently isolated feature, such as the $[\text{Co III}]$ $10.523 \mu\text{m}$, is still heavily blended, and not dominated by one single transition, meaning any velocity extracted from such features will be highly uncertain.

The $[\text{Ar II}]$ $8.991 \mu\text{m}$ feature is unique as it shows a flat top profile. This flat-top profile (at earlier epochs this was referred to as a “flat-tilted” profile) is seen throughout all of the spectra in the time series, and has been interpreted as being caused by a shell of Ar in the ejecta (Kwok et al. 2023b), which may come from an off-center deflagration to detonation transition (DerKacy et al. 2023a). The blue side of this feature is clearly contaminated by emission from the $[\text{Ni IV}]$ $8.405 \mu\text{m}$ transition. We also see that the peak of this feature does not follow the half life of ^{56}Co . Overall, we stress that all of the line profiles shown in this section are highly blended, and only with improved atomic data can the true components that contribute to the MIRI/MRS spectrum of SN 2021aefx be extracted.

5.3. $[\text{Ni II}] / [\text{Co III}]$ ratio as a function of time

Examining the peak flux ratio $[\text{Ni II}]$ $7.349 \mu\text{m}$ / $[\text{Co III}]$ $11.888 \mu\text{m}$ as a function of time is another useful measurable. It allows us to determine how the temporal variation in the energy deposition, heating, and mixing between the ^{56}Ni region and the electron capture element region evolves. As can be seen in Fig. 9, this ratio gradually increases throughout the time series. At later epochs if the decay of ^{57}Co begins to dominate it could be expected that this trend changes direction. However, this may heavily depend on the location of the ^{57}Co with relation to the stable Ni in the ejecta.

5.4. Half Width at Half Maximum vs v_{peak}

There are many overlapping spectral lines that contribute to the formation of the nebular phase spectrum of SN 2021aefx. MIRI/MRS observations allow for more of these lines to be resolved. However, many of the atomic line transitions and strengths are not known. Regardless of this, we attempt to fit the emission profiles of spectral features in order to get a broad understanding of where in the ejecta the emitting regions of certain ions are located. The full analysis is given in Appendix B. We stress that the lack of full line IDs and the assumption that all emitting line profiles can be explained by simple functions (where we inherently assume they are




Figure 8. (Left:) The velocity profile evolution of four important wavelength ranges within MIR spectra of SN 2021aefx. For most regions the MIRI/MRS data can resolve structure within features which appear blended in the MIRI/LRS mode, see also the bottom panel. (Right:) The peak flux of the profiles as a function of time fit with exponential decay curves. (Bottom:) The MIRI/MRS data has been resampled to the spectral resolution of the MIRI/LRS data ($R \approx 70$), demonstrating that many of the isolated features cannot be resolved in the MIRI/LRS mode. We also note here that all MIRI/LRS data is heavily over sampled, and should be resampled to the spectral resolution, as we have done in this panel, before direct comparison to the MIRI/MRS data.




Figure 9. The ratio of the peak flux of the [Ni II] 7.349 μm to [Co III] 11.888 μm features as a function of time demonstrates that emission from stable Ni increasingly dominates the spectral formation. We chose to examine the peak flux here as it follows the bulk of the emission, and the features are too blended to obtain accurate values of HWHM.

symmetrical), and the priors given in the fitting procedure make the results highly uncertain.

Figure 10 shows the Half Width at Half Maximum (HWHM) vs the peak velocity of ions that have been fit in Appendix B. Overall, we see no general trend with any of the ions. This is likely to be caused by the number of Gaussians required to reproduce the observed spectra, along with unknown spectral lines, and their corresponding strengths. Therefore, we emphasize that fitting spectra with multiple Gaussians has very little physical meaning, except in the case of isolated strong features such as the [Co III] 11.888 μm line. Hence, we turn our attention to the models below, to further understand the physics and formation of the spectra. We also emphasize that both the high spectral resolution and the extended wavelength coverage of the MRS mode on MIRI are critical for the physical interpretation described below.

6. MODELS

In the following section, we compare the data to models. The simulations utilized in this work are computed using the HYDrodynamical RADiation code (HYDRA; e.g. Höflich 2003a) that consists of physics-based modules which provide solutions for: the rate equations that determine the nuclear reactions, the statistical equations needed to determine the atomic level populations, the equation-of-state, the matter opacities, the hydrodynamic evolution, and the radiation-




Figure 10. The peak velocity vs HWHM of selected spectral features. As discussed in the text, these values are highly uncertain, and the detailed spectral modeling presented in § 6 is needed to extract further information about the explosion.

transport equations (RTE). Here, the RTE is treated using Variable Eddington Tensor methods, with a Monte-Carlo (MC) scheme providing the necessary closure relation to the momentum equations needed to solve the generalized scattering and non-LTE problem (Höflich 2002; Höflich 2003b; Penney & Höflich 2014; Diamond et al. 2015; Höflich 2017; Hristov et al. 2021). The relevant physics and current limitations of the simulations in the nebular phase are discussed in detail in Höflich et al. (2021). The growth of the WD mass towards M_{Ch} has been simulated following the approach by Sugimoto et al. (1979) and Nomoto (1982). The accretion rate and composition of the accreting material during the final stages has been tuned so that the ignition is triggered at a central density ρ_c (Höflich 2002; Diamond et al. 2015). To trigger a thermonuclear runaway at a ρ_c below $\approx 0.8 - 1.0 \times 10^9 \text{ g cm}^{-3}$ requires accretion rates in excess of $2 \times 10^{-6} M_{\odot}/\text{yr}$. To avoid over-Eddington luminosity of the accreting WD, He or C-rich matter is required, at least during the final stages. This limit depends on the details of the physical processes which occur during the accretion, such as the Urca-cooling by neutrinos (see Gamow 1970; Haensel 1995 and, Sect. 5.7 of Diamond et al. 2015). For a review on a wide variety of progenitor evolutions see Alsabti & Murrin (2017) and references therein.

In this work, the explosion model is based on simulations of off-center delayed-detonation models following Livne (1999). This class of models has been successfully used to reproduce the photospheric and nebular phase flux and polarization spectra of SNe Ia. It has also been used to reproduce

the abundance distributions in SN-remnants and to study the role of SNe Ia as producers of the positrons observed in our Galaxy (Höflich et al. 2006; Fesen et al. 2015; Telesco et al. 2015; Hristov et al. 2021; Höflich et al. 2021; Mera Evans et al. 2022; Penney & Höflich 2014).

Here, the simulations are used with parameters that have been shown to reproduce the evolution of the polarization and flux spectra of the normal-luminosity SN 2019np (Höflich et al. 2023). These model parameters have been successfully applied to SN 2021aefx at +323 d if seen from $\Theta \approx -30^\circ$ (DerKacy et al. 2023a) where Θ is the angle between the equatorial plane defined by the orthogonal vector between the kinematic center and the location of the deflagration-to-detonation transition. Note that a variety of Θ values have been found for different SNe using both line profiles and spectropolarimetry, e.g. SN 2003hv (Höflich et al. 2006; Motohara et al. 2006), SN 2003du (Höflich et al. 2004), SN 2012ke (Patat et al. 2012), SN 2019np (Höflich et al. 2023), and SN 2020qxp (Höflich et al. 2021). For nebular profiles of different SNe, the distribution of Θ is consistent with a random orientation relative to the observer. Line-polarization during the photospheric phase favors angles close to the northern pole (i.e. Θ larger than 30° Höflich et al. 2023). This can be understood by selective line absorption by a large-scale abundance asymmetry, as produced in an off-center DDT. Both effects combined support the notion of a loop-sided asymmetry. For objects with multi-epoch observations, consistent values of Θ have been found, suggesting a large-scale abundance asymmetry. We note that the point of ignition and the deflagration-to-detonation transition are distinct and are related to very different physical processes.

In this work, the basic model parameters (the initial magnetic field of the WD, B ; and the central density ρ_c) have been extended to cover a larger range in values, to take advantage of the increase in spectral resolution of the MIRI/MRS data.

The model parameters are given in Table 2. The base model, Model 25, serves as a reference to a series of simulations with various ρ_c , see Fig. 11. To first order, the central density of the WD at the time of the runaway depends on the accretion rate, and WD magnetic field. Note that the accretion on the WD is governed by the configuration of the progenitor, and the range of possible accretion depends on the composition of the accreted material (Sugimoto et al. 1979; Nomoto 1982). For comprehensive overviews, see the textbook by Alsabti & Murdin (2017). The WD masses of models considered here are close to M_{Ch} .

6.1. Resolution of Simulations

Medium-resolution spectra are crucial for accurately interpreting the physical characteristics of individual observed and synthetic features. For our discussion below, where we compare observations with theory, the spectral resolution sets the

Table 2. Model 25 Parameters (*: base model 25 is used as reference).

Parameter	Mod.25a	Mod.25b	Mod.25	Mod.25c	Mod.25d
$M_{ej} [M_\odot]$	1.30	1.33	1.35*	1.35	1.38
$\rho_c [10^9 \text{ g cm}^{-3}]$	0.5	0.9	1.1*	1.1	4.0
$M_{tr} [M_\odot]$	0.24	0.24	0.24*	0.24	0.24
$M_{DDT} [M_\odot]$	0.5	0.5	0.5*	0.5	0.5
$B(WD) [10^3 \text{ G}]$	1000	1000	1000*	1	1000

limit to resolve small features and their profiles. The spectral resolution of the data varies with wavelength but $R \sim 2700$ corresponds to a velocity resolution of about $\sim 111 \text{ km s}^{-1}$. As in previous works (e.g. Höflich et al. 2023), the model domain is spatially resolved by a equidistant Cartesian grid of 300 points per dimension which covers an expansion velocity of $\approx 26,000 \text{ km s}^{-1}$. Taking into account the second order discretization in spatial coordinates of the simulations, this translates into a spectral resolving power $R \approx 600$. This corresponds to a velocity of $\approx 500 \text{ km s}^{-1}$ for underlying physical features in the full-domain simulation. This resolution is employed for solving for the temperature structure and rate equations because optical depth effects in the UV are important for the ionization balance due to the incomplete Rosseland cycle (Höflich et al. 2021)⁴.

For the emitted synthetic spectra and line profiles, more than 99.9% of the flux in the optical to MIR originates within $15,000 \text{ km s}^{-1}$ because the energy is produced from the radioactive decay of $^{56}\text{Ni}/^{56}\text{Co}$ (see Fig. 11). Therefore, the computed domain has been reduced to boost the resolving power to $R \approx 1200$, which corresponds to a velocity of $\approx 300 \text{ km s}^{-1}$. This allows for a direct comparison with the MIRI/MRS spectra without artificial convolution of the observed data. The model has approximately the same resolving power as the Channel 4 data.

The typical scale over which the density and abundances vary in our models (see Fig. 11), is $\approx 1,000 \text{ km s}^{-1}$. Physically, smaller scales down to 100 km s^{-1} can be expected as a result of downwards cascading in Rayleigh-Taylor instabilities in the outer layers. However, due to the large number of small plumes, a detection of density and chemical inhomogeneities requires a S/N better than 300 in the polarization spectra (Höflich et al. 2023) or in the peak flux of strong emission lines in flux spectra.

6.2. Energy Deposition and Ionization

⁴ Even at 1000+ days, the UV optical depth remains large in the iron-rich region.




Figure 11. (Left:) The angle-averaged composition of our best fit model, Model 25 from [Hoeftlich et al. \(2017\)](#) and [Hoeftlich et al. \(2023\)](#). The abundances are stratified, with asymmetries limited to the outer edge of the quasi-statistical equilibrium and nuclear statistical equilibrium layers in velocity. EC marks the region of electron capture elements with the thin, red line being ^{58}Ni . (Right:) Angle-averaged energy deposition by γ -rays and positrons, E_X , for our reference model with $(\rho_c, B) = (1.1 \times 10^9 \text{ g cm}^{-3}, 10^6 \text{ G})$; red). Also shown are explosion models with a large ρ_c , $(4 \times 10^9 \text{ g cm}^{-3})$ and $B = 10^6 \text{ G}$; cyan), and a lower WD initial magnetic field $(1.1 \times 10^9 \text{ g cm}^{-3}, 10^3 \text{ G})$; blue). Note that the shift of the sharp drop of E_X corresponds to the inner edge of the ^{56}Ni distribution as a function of ρ_c ([Diamond et al. 2015](#)). Positron-transport effects become important for small B . (E.g., on top of E_γ), E_X in the central electron capture-region depends sensitively on the magnetic B -field. As a result, the emission from electron capture elements will change by a factor of 2. Though large ρ_c increases the ^{58}Ni production by a factor of 2, the specific energy input is halved, leading to similar electron capture line strengths in the corresponding features. The degeneracy can be broken by line profiles with a resolution of better than 1000 km s^{-1} , or a time series of data and models that cover the γ -ray to the positron-dominated regime. The excitation in the Ar-region depends sensitively on the magnetic field of the WD, which has consequences for the spectra and a profound impact on the Ar line profiles (see Sect. 6.3).




Figure 12. Same as the left panel from [Figure 11](#) but with the angle-averaged ionization levels of I-IV per particle zoomed in on the spectra-forming region. At day +415, most of the ^{56}Co (blue) has decayed to ^{56}Fe with II-III being the dominant ionization stages where a small amount of I/IV are present. Various isotopes of electron capture elements (Fe/Co/Ni) are found in stages I-III and are seen at low velocities.

The angle-averaged ionization structures at day +415 are shown in [Figure 12](#). ^{56}Fe dominates the nuclear statistical equilibrium region through singly and doubly ionized species, as can be expected for normal-luminosity SNe Ia and under-luminous SNe Ia at earlier times ([Wilk et al. 2018](#); [Shingles](#)

[et al. 2020](#); [Hoeftlich et al. 2021](#)). Because the density increases inwards, and the recombination rate depends on the square of the density, the ionization decreases towards the center. In the models, the ionization rate shifts towards more neutral ions compared to nebular calculation at $\approx 200 - 300 \text{ d}$ ([Wilk et al. 2018](#); [Shingles et al. 2020](#); [DerKacy et al. 2023a](#)). At day +415, we see ~ 5 to 10% of neutral iron group elements. One of the reasons for the lower ionization is that temperatures decrease by $\sim 1000\text{--}1500 \text{ K}$ at day +415 from $\sim 5000 \text{ K}$ at day $200 - 300$ ([Kozma & Fransson 1992](#); [Fransson & Jerkstrand 2015](#)). The temperature structure of SN 2021aefx at +415 d resembles an under-luminous SNe Ia at about 190 days ([Hoeftlich et al. 2021](#)).

The significant difference between the new spectrum of SN 2021aefx and the previous ones is that +415 d marks the transition from the energy input being dominated by hard γ -rays to being dominated by positrons. This will be fully reached at $\sim +500 \text{ d}$ ([Penney & Hoeftlich 2014](#)). Note that this time also corresponds to a shift in optical emission properties in SN 2011fe ([Tucker et al. 2022](#)). Hence the magnetic field of the WD is a critical variable in the nebular simulations ([Penney & Hoeftlich 2014](#)). We also note that in our models between 400–500 days γ -rays are just as important as positrons to the energy position in the most central regions and both have a significant effect on the spectral formation.

Table 3. Line contributions to the *JWST* spectrum at Day +415 from the reference model with $\rho_c = 1.1 \times 10^9 \text{ g cm}^{-3}$ and $B = 10^6 \text{ G}$.

S	$\lambda [\mu\text{m}]$	Ion	S	$\lambda [\mu\text{m}]$	Ion	S	$\lambda [\mu\text{m}]$	Ion	S	$\lambda [\mu\text{m}]$	Ion	S	$\lambda [\mu\text{m}]$	Ion
	4.8603	[Fe III]	**	4.8891	[Fe II]		5.0623	[Co I]		5.1635	[Co I]		5.1796	[Co II]
**	5.1865	[Ni II]		5.2112	[Co I]	**	5.3402	[Fe II]		5.4394	[Co II]		5.4652 [†]	[V I]
**	5.6739	[Fe II]	*	5.6870 [†]	[V I]		5.7044	[Co II]		5.7391	[Fe II]	**	5.8933	[Ni I]
	5.9395	[Co II]	*	5.9527	[Ni II]	*	6.2135	[Co II]		6.273	[Co I]		6.2738	[Co II]
	6.3683	[Ar III]		6.379	[Fe II]	**	6.636	[Ni II]		6.7213	[Fe II]	**	6.9196	[Ni II]
**	6.9853	[Ar II]		7.0454	[Co I]	*	7.103	[Co III]	*	7.1473	[Fe III]		7.2019	[Co I]
*	7.3492	[Ni III]	**	7.5066	[Ni I]	*	7.7906	[Fe III]		8.044	[Co I]	*	8.211	[Fe III]
	8.2825	[Co I]	*	8.2993	[Fe II]	**	8.405	[Ni IV]	*	8.6107	[Fe III]	*	8.6438	[Co II]
*	8.7325	[Fe II]	*	8.9147 [†]	[Ti II]	**	8.945	[Ni IV]	**	8.9914	[Ar III]	*	9.1969 [†]	[Ti II]
	9.279 [†]	[Co IV]	*	9.618	[Ni II]		9.8195	[Co I]	*	10.08	[Ni II]	**	10.1637 [†]	[Ti II]
**	10.189	[Fe II]		10.203	[Fe III]	*	10.5105	[Ti II]	*	10.5105	[S IV]	*	10.523	[Co II]
**	10.682	[Ni II]	**	11.13	[Ni IV]		11.167	[Co II]	*	11.238 [†]	[Ti II]		11.307	[Ni I]
* **	11.888	[Co III]	**	12.001	[Ni I]	*	12.1592 [†]	[Ti II]	**	12.255	[Co I]	*	12.261	[Mn II]
**	12.286 [†]	[Fe II]	**	12.642	[Fe II]		12.681	[Co III]	**	12.729	[Ni II]		12.736	[Ni IV]
	12.811	[Ne II]		13.058	[Co I]		13.82	[Co III]		13.924 [†]	[Co IV]		14.006	[Co III]
	14.356	[Co I]		14.391	[Co I]	*	14.739	[Co II]	*	14.814	[Ni I]	*	14.977	[Co II]
*	15.459	[Co II]		16.299	[Co II]	*	16.391	[Co III]	*	16.925	[Co I]	* **	17.936	[Fe II]
*	18.241	[Ni II]		18.265	[Co I]	*	18.39	[Co II]	**	18.713	[S III]		18.804	[Co II]
	18.985	[Co II]	*	19.007	[Fe II]	**	19.056	[Fe II]		19.138	[Ni II]		19.232	[Fe III]
	20.167	[Fe III]		20.928 [†]	[Fe II]		21.17 [†]	[Fe I]		21.986 [†]	[Fe II]		22.106 [†]	[Ni I]
*	21.481	[Fe II]		20.167	[Fe III]		20.928 [†]	[Fe II]		21.17 [†]	[Fe I]		21.986 [†]	[Fe II]
	22.106 [†]	[Ni I]	*	21.481	[Fe II]	*	21.829	[Ar III]	*	22.297	[Fe I]		22.80 [†]	[Co IV]
*	22.902	[Fe II]	* **	22.925	[Fe III]		23.086	[Ni II]	*	23.196	[Co II]		23.389	[Fe III]
	24.04 [†]	[Co IV]	* **	24.042	[Fe I]	**	24.07	[Co III]	* **	24.519	[Fe II]		24.847	[Co I]
**	25.249	[S I]	*	25.689	[Co II]	**	25.89	[O IV]		25.986	[Co II]	* **	25.988	[Fe II]
	26.1	[Co III]		26.13	[Fe III]		26.601	[Fe II]		27.53	[Co II]		27.55	[Co I]
*	28.466	[Fe I]		29.675	[Mn II]	* **	33.038	[Fe III]		33.481	[S III]		34.66	[Fe II]
*	34.713	[Fe I]	**	34.815	[Si II]	* **	35.349	[Fe II]	*	35.777	[Fe II]	*	38.801	[Fe I]
	39.272	[Co II]	* **	51.301	[Fe II]	* **	51.77	[Fe III]	*	54.311	[Fe I]	*	56.311	[S I]
	60.128	[Fe II]												

NOTE—The relative strengths are indicated by the number of *. For transitions without known lifetimes (marked by [†]), $A_{i,j}$ are assumed from the equivalent iron levels.

6.3. Flux Spectra and Profiles

6.3.1. Atomic data, line identifications, and strengths

The atomic data used in our simulations come from [Diamond et al. \(2015, 2018\)](#); [van Hoof \(2018\)](#), and references therein. In addition, the atomic data have been supplemented by fits of the lifetimes, A_{ij} , of unknown weak features between 5 and 27 μm based on the observed spectrum (see Appendix C, Tables D.1 & D.2). Full lists with line strengths of transitions contributing to the flux spectrum of SN 2021aefx at +415 d from maximum are shown in Tables 3 & C.1. Line IDs are only shown for transitions with measured spontaneous lifetimes, i.e. Einstein A_{ij} values. Our atomic models have been verified and used in previous papers ([DerKacy et al. 2023a, 2024a](#)), and cross-checked with line identifications by [Kwok et al. \(2023b\)](#).

6.3.2. Understanding the overall spectra and evolution

The observed and modeled spectra are presented with identifications of notable features in Fig. 13. Here, we discuss the overall spectrum and underlying physics of our reference

model before optimizing the parameters. The synthetic spectrum agrees reasonably well with the observations.

At first glance, the spectrum at +415 d resembles earlier epochs. However, this apparent similarity masks a physical regime change in the ejecta. For example, at +415 d, most of the radioactive ^{56}Co has decayed to ^{56}Fe (Fig. 11) with only $\sim 2\%$ of the initial ^{56}Co remaining. The observed and model spectra show broad features of [Fe II] and [Fe III] at 18 and 23 μm . In the models the +415 d spectrum shows an equally strong, peaked feature at the location of [Co III] 11.888 μm rather than a rounded profile seen a year earlier. This is also seen in several weaker features which could be attributed to [Co II] (e.g. 10.5 μm) at earlier epochs. Furthermore, the observed spectrum and model at +415 d also show strong features at 7 and 9 μm . This is similar to earlier epochs, but both the profile shapes and dominant lines in the features have changed. At earlier times, the dominant ions have been identified as [Ar II] and [Ar III] ([Gerardy et al. 2007](#); [Telesco et al. 2015](#); [DerKacy et al. 2023a](#); [Kwok et al. 2023b](#)). In fact, at +415 d the 8.991 μm feature is “flat-topped” rather




Figure 15. Same as Fig. 14, but the sensitivity of the initial magnetic field of the WD, B , on the synthetic spectra is shown. The lines become narrower and more peaked with decreasing B . This is very similar to the low ρ_c case. However, the effects can be separated by the overall spectral model.

^{54}Fe . The isotopic shift between ^{56}Co and ^{57}Co is small, and the rest wavelengths of $[\text{Ni I}]$ and $[\text{Co I}]$ are similar to $[\text{Co III}]$. This leads to narrower, additional components boosting the central peak of the $11.888 \mu\text{m}$ feature.

6.4. Sensitivity of the free parameters on the spectra and profiles

In this section, the formation of spectra and specific spectral profiles is considered in more detail, and discussed in the context of variations between models.

6.4.1. Dependence on the WD central density

Figure 14 shows how the central density, ρ_c , of the WD (which determines the size of the inner ^{56}Ni hole and the amount of electron capture elements produced in the explosion) affects the spectral formation. With lower ρ_c , the intrinsic line profiles of ^{56}Fe and ^{56}Co become narrower because an increase of emission at low velocities produces a rise in the

peak (Penney & Hoefflich 2014). This can be seen in the broad $[\text{Fe II}]$ and $[\text{Fe III}]$ dominated features at ≈ 18 and $23 \mu\text{m}$. Note that, though large ρ_c increases the ^{58}Ni production by a factor of 2, the specific energy input is halved, leading to similar electron capture line strengths, but different widths, in the features (Fig. 11), resulting in similar strengths in ^{58}Ni lines (Fig. 14 vs. 15). Furthermore, a high $\rho_c = 4 \times 10^9 \text{ g cm}^{-3}$ can be ruled out because it would produce profiles that are much broader than the data, by a factor of two based on a spherical model series (Diamond et al. 2015; Galbany et al. 2019). The MIR spectrum of such densities are not shown here because our simulation used do not have enough grid-points to resolve the complex radial structure of low Y_e electron capture elements such as Mn and Cr that are important (see Figs. 25 and 26 in Galbany et al. 2019).

Another spectral indicator of the central density, ρ_c , is the trend that the features dominated by electron capture elements show a decreasing flux with decreasing ρ_c . This becomes




Figure 16. Same as Fig. 14 but the evolution over one e-folding time for ^{56}Co decay is shown. The flux at +550 d is scaled up in luminosity to compensate for the ^{56}Co decay. Thus, line profiles and ratios between the models can be compared.

obvious in the 6.7 and 7.4 μm regions. The sensitivity to ρ_c is also seen by narrow mostly unblended weak $[\text{Ni II}]$ and $[\text{Co II}]$ lines at 6 and 6.2 μm , respectively.

Models with $\rho_c = 1.1$ and $0.9 \times 10^9 \text{ g cm}^{-3}$ do equally well at reproducing the observations. The former shows better agreement with the flux level, whereas the latter produces slightly narrower forbidden $[\text{Fe II}]$ and $[\text{Fe III}]$ features at 18 and 23 μm . Note, that for $\rho_c \geq 0.9 \times 10^9 \text{ g cm}^{-3}$, the total emission of Ni features hardly depends on the ^{58}Ni mass produced (in unmixed models) because the smaller electron capture core also increases the specific heating which boosts the Ni emission (Fig. 11). Because heating is produced by γ -rays and positrons, the degeneracy can only be broken by time-series of spectra well beyond +500 d when positrons dominate everywhere. Moreover, at very late times, the variations due to positron transport effects can separate macroscopic from microscopic mixing.

Both the $[\text{Fe II}]$ and $[\text{Fe III}]$ features at 18 and 23 μm may favor the lowest ρ_c ($0.5 \times 10^9 \text{ g cm}^{-3}$). However, this model

is incompatible with the features and profiles at 7, 10.5 and 12 μm due to the low mass of electron capture elements.

The 7 μm feature is complex with a structure dominated by narrow Ni components with a width of $\sim 3000 \text{ km s}^{-1}$, and a broad $[\text{Ar II}]$ component (with a width of $\sim 7000\text{--}10000 \text{ km s}^{-1}$) which ‘fills the emission gap’ between the $[\text{Ni I}]$ and $[\text{Ni II}]$ dominated peaks. Other lines such as $[\text{Fe III}]$ contribute to the complexity.

The emission complex between 8.1 and 9.3 μm shows two components. The blue part of this wavelength region ($\sim 8.4 \mu\text{m}$) is dominated by $[\text{Ni IV}]$ 8.405 μm . Observations and models of SN 2021aefx at +323 d show, that the red part of this region ($\sim 9 \mu\text{m}$) was dominated by $[\text{Ar III}]$ (DerKacy et al. 2023a), which is similar to MIR observations of other SNe Ia (Gerardy et al. 2007; Telesco et al. 2015). At these phases, SN 2021aefx showed a tilted profile, which was interpreted as evidence for off-center DDT (DerKacy et al. 2023a). Such a profile can be tilted if the high-velocity region is asymmetric in the abundance distribution at the outer-edge, e.g. produced

by an off-center DDT (Höflich et al. 2021). However, in the latest spectrum of SN 2021aefx at +415 d, the $9\ \mu\text{m}$ profile is “flat-topped”, not tilted, and is dominated by [Ni iv] $8.945\ \mu\text{m}$ and [Fe ii]/[Fe iii] lines with only a non-dominant contribution of [Ar iii]. This leads to a bump at the red end of the profile. Only about 50% of the red end of the profile comes from Ar and is produced after the transition to the positron-dominated regime. The tilting vanishes because the outer, high-velocity Ar bulge is located outside the layers heated by positrons, resulting in a flat contribution to the profile at the latest epoch of observations.⁵ About 50% of the total emission can be attributed to [Ni iv] and multiple lines of [Fe ii] and [Fe iii].

In the early nebular phase, blending by [Fe ii] and [Fe iii] was suggested as a main contributor to the [Ar iii] feature. However, this leads to an opposite tilt to the observed spectrum (Blondin et al. 2023).⁶ Thus, the tilt and evolution of the feature supports our interpretation as a geometrical effect.

It is well established that the $12\ \mu\text{m}$ feature at earlier times can be attributed to [Co iii] although some weaker additional components are present. In the models by +415 d, [Ni i], [Ti ii], and many weak Fe-lines, contribute about 1/3 of the flux. The significant contribution beyond [Co iii] also becomes evident from Fig. 14 by showing a pointed peak compared to earlier observations.

As mentioned above, in both the models and data most spectral features show a very narrow peak because they are dominated by electron capture elements but still have a broad component. This characteristic is valid for all strong Fe features in our model spectrum because the ^{56}Fe distribution traces ^{56}Co and ^{56}Ni .

We note that microscopic mixing (i.e. mixing of species on atomic scales) over large scales would drastically change the evolution and was previously excluded for SN 2021aefx (DerKacy et al. 2023a).

6.4.2. Dependence on the initial magnetic field of the WD

Fig. 15 shows the sensitivity of the initial magnetic field of the WD on the synthetic spectra. We use a turbulent morphology on scales of the Rayleigh-Taylor instabilities because even large-scale initial dipole fields would be tangled by passive flows during the deflagration phase of burning (Hristov et al. 2021). The changes of the spectra and profiles can be understood in the same way as the previous discussion. Prior to day ≈ 300 , all positrons locally annihilate, regardless of the strength of the initial magnetic field of the WD (Pen-

ney & Höflich 2014). The changes in the specific energy distributions can be seen in Fig. 11.

For weaker magnetic fields, positrons increasingly escape from the ^{56}Ni layers to regions of higher velocities. This leads to narrower mean half-width Fe and Co profiles, similar to changes in ρ_c , by reducing the emission at high Doppler shifts, namely ^{56}Co at high velocities (see Fig. 11). The second effect is a boost of features where quasi-statistical equilibrium elements, such as Si and S, contribute significantly to the line flux, e.g. see the feature at [Si iv] $10.5\ \mu\text{m}$, which is very similar to the potential appearance of the [Si ii] $1.3\ \mu\text{m}$ feature in the NIR (Diamond et al. 2015). The other effect is the shifting of the energy deposition towards the inner, higher-density layers, which results in a boost of the [Fe ii] line at $18\ \mu\text{m}$ compared to the [Fe iii] at $23\ \mu\text{m}$, because some positron escape leads to slightly lower temperatures and, thus, a shift in the flux from the optical to the MIR (Penney & Höflich 2014; Mera Evans et al. 2022). We require initial B-fields in excess of 10^6G , which is at the upper end of previous limits for SNe Ia based on late-time light curves and spectra (Höflich et al. 2004; Diamond et al. 2015; Kumar et al. 2023). For the creation of high B fields, we possibly need a strong dynamo during the smoldering phase, as discussed in the above publications.

6.4.3. Future time evolution of the spectra

In the discussion in § 5, we made use of past observations. From the models, we identified that the main physical driver of the evolution is the abundance change from ^{56}Co to ^{56}Fe , and the change of the mode of energy input, from a γ -ray to the positron-dominated regime.

Here, we consider the future evolution of SN 2021aefx using the same model configuration and parameters presented above. This model has been evolved to one e-folding time of ^{56}Co later (+550 d, see Fig 16), hence the absolute flux is lower. The main change in the evolution of the spectral profiles is a boost in features dominated by intermediate mass elements, such as [Ar ii] at $7\ \mu\text{m}$ and [Ar iii] at $9\ \mu\text{m}$. Furthermore, there is a narrowing of features due to a combination of positron transport effects and a decrease in the contribution from γ -rays by $\approx 60\%$ produced by the geometrical dilution.

At +550 d the spectrum resembles an SN Ia; however, not unexpectedly, it does not match the data from +415 days.

What would one learn if the prediction looks different from future data, or if the spectral evolution is slow? One would have to restore positron trapping by increasing the magnetic field of the WD because, in our analysis, the magnetic field is a lower limit, and decreases with the square of time. In general, positron transport effects on the spectra depend on the size and morphology of both the magnetic field and the 3D abundance distributions in the central region.

⁵ At +323 days, heating was facilitated by γ -rays leading to a “flat-tilted” Ar-dominated profile.

⁶ At +323d, our simulations show blends at the same wavelength but weaker. If those blends dominate the profile, they would cause a tilt opposite to the observation.

6.5. Discussion and Implications for the Underlying Explosion Physics

To prevent duplication of discussions on alternative scenarios, we direct readers to the analyses provided in earlier studies (Hristov et al. 2021; Hoeflich et al. 2023) and, for SN 2021aefx to DerKacy et al. (2023a). Here, we focus on our new findings and their implications.

SN 2021aefx can be understood in the framework of an off-center delayed-detonation model with central WD densities of $\rho_c \approx 0.9 - 1.1 \times 10^9 \text{ g cm}^{-3}$ and with $M_{\text{WD}} \approx 1.33 - 1.35 M_{\odot}$. These parameters can be attributed to a near M_{Ch} mass WD, and place it into a regime of high-density burning with a proton to nucleon ratio $Y_e \approx 0.49$. As discussed at the beginning of Sect. 6 in context of the model construction, the relatively low ρ_c requires a large accretion rate that may be more compatible with He or C rather than H accretors.

For the +415 d spectra, the strength of the ^{58}Ni hardly depends on the total mass of ^{58}Ni which is governed by ρ_c . As is obvious from Fig. 11, the amount of ^{58}Ni may vary by a factor of 2, but this can be compensated by the specific energy input, i.e. the energy input per gram. Time series of data and models are needed that cover the γ -ray dominated nebular regime (~ 200 d) to the positron-dominated regime which starts at ~ 500 days after the explosion.

The result is that the main isotopes in the core are stable electron capture elements e.g. ^{58}Ni , ^{57}Co , and ^{54}Fe . This places SN 2021aefx into a similar physical regime as other normal-luminosity SNe Ia such as SN 2014J (Telesco et al. 2015).

However, in our models not all SNe Ia have the same ρ_c . In fact, they seem to span a wide range between $5 - 50 \times 10^8 \text{ g cm}^{-3}$, where there is evidence for high ρ_c in sub-luminous SNe Ia such as SN 2016hmk, SN 2020qxp, and SN 2022xkq (Höflich et al. 2004; Penney & Hoeflich 2014; Diamond et al. 2015, 2018; Galbany et al. 2019; DerKacy et al. 2024a)⁷.

Probing the transition between the γ -ray and positron dominated regimes is important for establishing the Urca-process, its inner working, its impact on the energy balance during the smoldering phase, and the nature of weak interactions (see Diamond et al. 2018; Tucker et al. 2022, and references therein). Moreover, nuclear cross-sections played a role (e.g. Brachwitz et al. 2000; Höflich et al. 2006) and will continue to play a central role (Thielemann 2019) in our understanding of the explosion mechanism and thermonuclear runaway. For example, improvements in the electron capture rates (Langanke & Martínez-Pinedo 2000) resulted in a drastic change of the [Fe II] at $1.644 \mu\text{m}$ nebular line profile (Höflich 2006) from flat-topped to rounded (Penney & Hoeflich 2014), which

demonstrated the need for high ρ_c WDs close to where an accretion induced collapse would occur (e.g. Gronow et al. 2021).

6.6. Implications for the physics of the thermonuclear runaway

The comparison of synthetic and observed spectral profiles places strong constraints on the thermonuclear runaway in the delayed-detonation scenario, specifically regarding the initiation of thermonuclear explosive burning.

Detailed multi-dimensional simulations of the central single-point runaway show only mixing of the inner electron capture layers during the deflagration phase (see e.g. Figs. 1, 2, and 15 in Khokhlov 2000). Mixing occurs only during the subsonic deflagration phase and not during the subsequent detonation phase of DDT models, which leads to overall structures very similar to spherical DDT models (see Fig. 3 in Gamezo et al. 2005). In contrast, multiple-spot, off-center ignitions mix the electron capture rich material to high velocities (Fink et al. 2014; Pakmor et al. 2024). The differences between these hydro-simulations can be simply understood by the lack of gravitation in the center, which results in close-to-laminar burning for about 1 second of high-density burning before Rayleigh-Taylor instabilities develop. Whereas, in multi-spot far off-center ignitions, the Rayleigh-Taylor-dominated burning phase starts right away. As shown by Khokhlov (2000), the long delay between ignition happens regardless of pre-existing turbulence produced during the smoldering phase (Höflich & Stein 2002) or low C/O-ratios (e.g. Domínguez et al. 2001). Pre-existing turbulence will introduce some mixing by dragging material, but this effect is limited to the innermost slowly expanding layers. As found in the analysis of previous SNe Ia (Galbany et al. 2019; Hoeflich et al. 2021), again the spectra here are consistent with a near-to-central ignition and inconsistent with multi-spot strong-off-center ignition.

Moreover, the narrow width of the lines produced by electron capture elements puts tight limits on the macroscopic mixing, and confines it to the inner $\approx 3000 \text{ km s}^{-1}$ which may be expected from the passive drag of electron capture material by pre-existing turbulent fields produced during the smoldering phase (e.g. Höflich & Stein 2002). Note that high WD magnetic fields ($B \geq 10^6 \text{ G}$) are required based on studies of light curves⁸ and spectra. Such fields may decrease the mixing of electron capture element even further. For detailed discussions, see Höflich et al. (2004); Diamond et al. (2015); Hristov et al. (2021); Hoeflich et al. (2023); DerKacy et al. (2024a). A turbulence field can be expected to develop during the smoldering phase (Höflich & Stein 2002), which

⁷High central densities, close to an Accretion Induced Collapse, found in many SNe Ia do not imply one specific unique explosion/progenitor scenario for all SNe Ia.

⁸To measure B fields of the order of 10^6 G , an accuracy of 0.025 mag is required for late-time bolometric light curves (Hristov et al. 2021)

drags the freshly formed products of high-density burning by passive flow. The result is an inhomogeneous mixture of non-radioactive electron capture elements and ^{56}Ni material which is heated over days by radioactive decay. Pressure equilibrium will compress the electron capture elements and non-radioactive layers into thin sheets (Hoeftlich 2017). Pre-existing turbulence may produce caustic distributions with a wall thickness of $\approx 1000 - 1500 \text{ km s}^{-1}$ but limited to the inner region (Fesen et al. 2015; Hoeftlich 2017). The models used here do not take into account these structures, but their existence has been recently indicated by late-time polarization (Yang et al. 2022).

7. CONCLUSIONS

We present nebular phase *JWST* MIRI/MRS observations of SN 2021aefx at +415 d past maximum. Our work demonstrates how combining MIR medium-resolution data with detailed spectral models allows for the physics of SNe Ia to be understood in a way that was previously not possible. *JWST* promises to transform this area of research over the coming years. The main results of our study can be summarized as follows:

- These new data and models covering $5\text{--}27 \mu\text{m}$ have allowed us to produce an extended list of line identifications during the iron-dominated nebular phase (see, § 2 & Table 3). The higher resolution of the spectrum relative to previous epochs has allowed for many spectral features to be resolved. The spectrum is dominated by iron-group elements, with a strong contribution of stable Ni.
- The *JWST* MIRI/MRS spectrum at +415 d has been analyzed in conjunction with previous MIRI/LRS observations of SN 2021aefx, allowing for a time series analysis from +255 to +415 d to be performed. We find that the peak evolution of the resonance [Co III] $11.888 \mu\text{m}$ feature is consistent with the half-life of radioactive decay of ^{56}Co (see Sect. § 5.1 & Fig. 7).
- The spectrum has been analyzed using a new series of off-center delayed-detonation models. All models had a central point-of-ignition, and a point of the DDT at $0.5 M_{\odot}$, which results in asymmetric abundance distributions (see § 6). The viewing angle Θ is measured between the line-of-sight and the line defined by the point of DDT and the kinematic center. The spectrum at +415 d can be reproduced using the same $\Theta \approx -30^\circ$ that has been found in our previous study of SN 2021aefx at +323 d (DerKacy et al. 2023a). For other SNe Ia, a wide range of values for Θ from -90° to $+90^\circ$ have been found between SNe. This supports our interpretation that the profiles of SN 2021aefx are produced by a geometric effect.
- Variations in the central density and initial magnetic field of the WD in the models were studied. We find that the progenitor of SN 2021aefx had a WD mass of $\approx 1.33 - 1.35 M_{\odot}$, and central density of $\rho_c = 0.9 - 1.1 \times 10^9 \text{ g cm}^{-3}$, and an initial magnetic field of $\geq 10^6 \text{ G}$.
- Comparison between these new data and models has revealed a profound change in the physics of the spectral formation compared to previous epochs. At +415 d the SN made the transition from a ^{56}Co -dominated to a mostly ^{56}Fe -dominated regime. For the energy input, this is the transition phase from a globally γ -ray-dominated regime towards a positron-dominated regime. However, at +415 d γ -rays still dominate the energy input in the central region. The evolution of the line profiles can be understood in terms of the above physics (see § 6.4).
- The fact that we observe a narrow region of electron capture elements means there is very limited mixing in the inner regions of the ejecta. Thus, the point of ignition in SN 2021aefx is consistent with being near-to-central, and is inconsistent with being strongly off center if burning started as a deflagration front (see § 6.6).
- At +415 d, the strength of the Ni features is dominated by the size of the B-field in the WD and is rather insensitive to the amount of Ni produced through high central density burning $\geq 10^9 \text{ g cm}^{-3}$.
- At earlier epochs the spectral features at 7 and $9 \mu\text{m}$ were mostly attributed to [Ar II] and [Ar III]. The profiles were interpreted as being caused by asymmetric abundance distribution produced by the off-center nature of the DDT. This resulted in ‘flat-tilted’ line profile because the entire Ar region was excited by gamma-rays. At +415 d these features have significantly changed. The Ar-region is no longer excited by γ -rays but by positrons, which deposit their energy locally. Hence, the emission is governed by the overlapping region between ^{56}Ni and Ar. The Ar contribution to the 7 and $9 \mu\text{m}$ features is small and flat topped, because the contribution of the asymmetric component from the outer layers is suppressed relative to earlier epochs. Overall, these features are dominated by iron-group elements at this phase (§ 6.4).
- Spectra and line profiles are sensitive to the initial B field of the WD because positron transport alters the energy input relative to the abundances (Penney & Hoeftlich 2014; Diamond et al. 2015). Here, the importance of positron transport and the need for high B fields has

been demonstrated. Where we find the need for B fields which are larger than fields typically found in WDs. These values of B may be produced during the smoldering phase prior to the explosion. Later time-spectra are needed to push the limits to the regime where they heavily modify the properties of nuclear-burning fronts (Remming & Khokhlov 2014; Hristov et al. 2018).

Finally, we want to emphasize the future prospects for our analysis of SN 2021aefx. After about 550 days, the spectra enter the full positron-dominated regime. The positron transport effects have been demonstrated and the lower limit of B has been derived for SN 2021aefx, but spectra taken after day 700 (DerKacy et al. 2023b, 2024b) will be more sensitive to the morphology of the magnetic field of the WD and will provide the measurements of B beyond the current limit of 10^6 G. Current *JWST* observations indicate a low density and high mass WD which had a high magnetic field. This hints both towards a lack of full understanding of the accretion and smoldering phase, and the Urca cooling, which may be affected by large B-fields and WD rotation. Future *JWST* observations may address these problems.

Overall, SN 2021aefx has demonstrated how intensive studies with an extended time-series of data consisting of low, and medium-resolution spectra can revolutionize our understanding of the physical processes governing SNe Ia. However, SN 2021aefx is just one object and future low to medium resolution *JWST* observations of SNe Ia are required to examine the true diversity both within the physical processes in the ejecta, and the explosion scenarios within the universe.

ACKNOWLEDGMENTS

C.A., P.H., E.B., J.D., and K.M. acknowledge support by NASA grants JWST-GO-02114, JWST-GO-02122, JWST-GO-03726, JWST-GO-04436, and JWST-GO-04522. M.S. acknowledges support by NASA grants JWST-GO-03726, JWST-GO-04436, and JWST-GO-04522. Support for programs #2114, #2122, #3726, #4436, and #4522 were provided by NASA through a grant from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-03127. P.H. and E.B. acknowledge support by NASA grant 80NSSC20K0538. P.H. acknowledges support by NSF grant AST-2306395.

L.G. acknowledges financial support from the Spanish Ministerio de Ciencia e Innovación (MCIN) and the Agencia Estatal de Investigación (AEI) 10.13039/501100011033 under the PID2020-115253GA-I00 HOSTFLOWS project, from Centro Superior de Investigaciones Científicas (CSIC) under the PIE project 20215AT016 and the program Unidad de Excelencia María de Maeztu CEX2020-001058-M, and from the Departament de Recerca i Universitats de la Generalitat de Catalunya through the 2021-SGR-01270 grant. I.D. is supported by the project PID2021-123110NB-I00 financed by MCIN/AEI/10.13039/501100011033/ & FEDER A way to make Europe, UE.

Facilities: *JWST* (MIRI/MRS), MAST (*JWST*)

Software: jwst (ver. 1.9.4, Bushouse et al. 2023), jdaviz (ver. 3.2.1, Developers et al. 2023), HYDRA (Höflich 2003b, 2009; Höflich et al. 2017), Astropy (Astropy Collaboration et al. 2013, 2018, 2022), NumPy (Harris et al. 2020), SciPy (Virtanen et al. 2020), Matplotlib (Hunter 2007).

REFERENCES

- Alsabti, A. W., & Murdin, P. 2017, Handbook of Supernovae (Berlin: Springer), doi: [10.1007/978-3-319-21846-5](https://doi.org/10.1007/978-3-319-21846-5)
- Argyriou, I., Glasse, A., Law, D. R., et al. 2023, A&A, 675, A111, doi: [10.1051/0004-6361/202346489](https://doi.org/10.1051/0004-6361/202346489)
- Ashall, C., Mazzali, P. A., Pian, E., & James, P. A. 2016, MNRAS, 463, 1891, doi: [10.1093/mnras/stw2114](https://doi.org/10.1093/mnras/stw2114)
- Ashall, C., Baron, E., Höflich, P. A., et al. 2021, MIR Spectroscopy of Type Ia Supernovae: The Key to Unlocking their Explosions and Element Production, *JWST* Proposal. Cycle 1, ID. #2114
- Ashall, C., Lu, J., Shappee, B. J., et al. 2022, ApJL, 932, L2, doi: [10.3847/2041-8213/ac7235](https://doi.org/10.3847/2041-8213/ac7235)
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, A&A, 558, A33, doi: [10.1051/0004-6361/201322068](https://doi.org/10.1051/0004-6361/201322068)
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, AJ, 156, 123, doi: [10.3847/1538-3881/aabc4f](https://doi.org/10.3847/1538-3881/aabc4f)
- Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., et al. 2022, apj, 935, 167, doi: [10.3847/1538-4357/ac7c74](https://doi.org/10.3847/1538-4357/ac7c74)
- Axelrod, T. S. 1980, PhD thesis, University of California, Santa Cruz
- Berger, M. B., Hubbell, J. H., Seltzer, S., Coursey, J., & Zucker, D. 1998, NBSIR, 87-3597, ["http://physics.nist.gov/PhysRefData/Xcom/Text/XCOM.html"](http://physics.nist.gov/PhysRefData/Xcom/Text/XCOM.html)
- Blondin, S., Dessart, L., Hillier, D. J., Ramsbottom, C. A., & Storey, P. J. 2023, A&A, 678, A170, doi: [10.1051/0004-6361/202347147](https://doi.org/10.1051/0004-6361/202347147)
- Boos, S. J., Townsley, D. M., Shen, K. J., Caldwell, S., & Miles, B. J. 2021, ApJ, 919, 126, doi: [10.3847/1538-4357/ac07a2](https://doi.org/10.3847/1538-4357/ac07a2)

- Brachwitz, F., Dean, D. J., Hix, W. R., et al. 2000, *ApJ*, 536, 934, doi: [10.1086/308968](https://doi.org/10.1086/308968)
- Burrow, A., Baron, E., Ashall, C., et al. 2020, *ApJ*, 901, 154, doi: [10.3847/1538-4357/abafa2](https://doi.org/10.3847/1538-4357/abafa2)
- Bushouse, H., Eisenhamer, J., Dencheva, N., et al. 2023, JWST Calibration Pipeline, 1.9.4, Zenodo, doi: [10.5281/zenodo.7577320](https://doi.org/10.5281/zenodo.7577320)
- Chen, N. M., Tucker, M. A., Hoyer, N., et al. 2023, *ApJL*, 944, L28, doi: [10.3847/2041-8213/acb6d8](https://doi.org/10.3847/2041-8213/acb6d8)
- Childress, M. J., Hillier, D. J., Seitzenzahl, I., et al. 2015, *MNRAS*, 454, 3816, doi: [10.1093/mnras/stv2173](https://doi.org/10.1093/mnras/stv2173)
- DerKacy, J. M., Ashall, C., Hoefflich, P., et al. 2023a, *ApJL*, 945, L2, doi: [10.3847/2041-8213/acb8a8](https://doi.org/10.3847/2041-8213/acb8a8)
- DerKacy, J. M., Baron, E., Hoefflich, P. A., et al. 2023b, Examining the Heart of Type Ia Supernova 2021aefx with Ultra-Late Time Spectra, JWST Proposal. Cycle 2, ID. #3726
- DerKacy, J. M., Ashall, C., Hoefflich, P., et al. 2024a, *ApJ*, 961, 187, doi: [10.3847/1538-4357/ad0b7b](https://doi.org/10.3847/1538-4357/ad0b7b)
- DerKacy, J. M., Baron, E., Hoefflich, P. A., et al. 2024b, Examining the Heart of Type Ia Supernova 2021aefx with Ultra-Late Time Spectra, JWST Proposal. Cycle 3, ID. #6582
- Developers, J., Averbukh, J., Bradley, L., et al. 2023, Jdaviz, 3.2.1, Zenodo, doi: [10.5281/zenodo.7600492](https://doi.org/10.5281/zenodo.7600492)
- Diamond, T. R., Hoefflich, P., & Gerardy, C. L. 2015, *ApJ*, 806, 107, doi: [10.1088/0004-637X/806/1/107](https://doi.org/10.1088/0004-637X/806/1/107)
- Diamond, T. R., Hoefflich, P., Hsiao, E. Y., et al. 2018, *ApJ*, 861, 119, doi: [10.3847/1538-4357/aac434](https://doi.org/10.3847/1538-4357/aac434)
- Domínguez, I., Höflich, P., & Straniero, O. 2001, *ApJ*, 557, 279, doi: [10.1086/321661](https://doi.org/10.1086/321661)
- Elagali, A., Staveley-Smith, L., Rhee, J., et al. 2019, *MNRAS*, 487, 2797, doi: [10.1093/mnras/stz1448](https://doi.org/10.1093/mnras/stz1448)
- Fesen, R. A., Höflich, P. A., & Hamilton, A. J. S. 2015, *ApJ*, 804, 140, doi: [10.1088/0004-637X/804/2/140](https://doi.org/10.1088/0004-637X/804/2/140)
- Fink, M., Kromer, M., Seitzenzahl, I. R., et al. 2014, *MNRAS*, 438, 1762, doi: [10.1093/mnras/stt2315](https://doi.org/10.1093/mnras/stt2315)
- Fransson, C., & Jerkstrand, A. 2015, *ApJL*, 814, L2, doi: [10.1088/2041-8205/814/1/L2](https://doi.org/10.1088/2041-8205/814/1/L2)
- Galbany, L., Ashall, C., Höflich, P., et al. 2019, *A&A*, 630, A76, doi: [10.1051/0004-6361/201935537](https://doi.org/10.1051/0004-6361/201935537)
- Gamezo, V. N., Khokhlov, A. M., & Oran, E. S. 2005, *ApJ*, 623, 337, doi: [10.1086/428767](https://doi.org/10.1086/428767)
- Gamezo, V. N., Khokhlov, A. M., Oran, E. S., Chtchelkanova, A. Y., & Rosenberg, R. O. 2003, *Science*, 299, 77, doi: [10.1126/science.299.5603.77](https://doi.org/10.1126/science.299.5603.77)
- Gamow, G. 1970, *My world line: An informal autobiography*.
- Gerardy, C. L., Meikle, W. P. S., Kotak, R., et al. 2007, *ApJ*, 661, 995, doi: [10.1086/516728](https://doi.org/10.1086/516728)
- Gronow, S., Côté, B., Lach, F., et al. 2021, *A&A*, 656, A94, doi: [10.1051/0004-6361/202140881](https://doi.org/10.1051/0004-6361/202140881)
- Haensel, P. 1995, *SSRv*, 74, 427, doi: [10.1007/BF00751429](https://doi.org/10.1007/BF00751429)
- Harris, C. R., Millman, K. J., van der Walt, S. J., et al. 2020, *Nature*, 585, 357, doi: [10.1038/s41586-020-2649-2](https://doi.org/10.1038/s41586-020-2649-2)
- Hoefflich, P. 2002, arXiv e-prints, astro, doi: [10.48550/arXiv.astro-ph/0207103](https://doi.org/10.48550/arXiv.astro-ph/0207103)
- . 2017, in *Handbook of Supernovae*, ed. A. W. Alsabti & P. Murdin (Berlin: Springer), 1151, doi: [10.1007/978-3-319-21846-5_56](https://doi.org/10.1007/978-3-319-21846-5_56)
- Hoefflich, P., Khokhlov, A. M., & Wheeler, J. C. 1995, *ApJ*, 444, 831, doi: [10.1086/175656](https://doi.org/10.1086/175656)
- Hoefflich, P., Hsiao, E. Y., Ashall, C., et al. 2017, *ApJ*, 846, 58, doi: [10.3847/1538-4357/aa84b2](https://doi.org/10.3847/1538-4357/aa84b2)
- Hoefflich, P., Ashall, C., Bose, S., et al. 2021, *ApJ*, 922, 186, doi: [10.3847/1538-4357/ac250d](https://doi.org/10.3847/1538-4357/ac250d)
- Hoefflich, P., Yang, Y., Baade, D., et al. 2023, *MNRAS*, 520, 560, doi: [10.1093/mnras/stad172](https://doi.org/10.1093/mnras/stad172)
- Höflich, P. 2003a, in *Astronomical Society of the Pacific Conference Series*, Vol. 288, *Stellar Atmosphere Modeling*, ed. I. Hubeny, D. Mihalas, & K. Werner, 371, doi: [10.48550/arXiv.astro-ph/0209420](https://doi.org/10.48550/arXiv.astro-ph/0209420)
- Höflich, P. 2003b, in *Astronomical Society of the Pacific Conference Series*, Vol. 288, *Stellar Atmosphere Modeling*, ed. I. Hubeny, D. Mihalas, & K. Werner, 185
- Höflich, P. 2006, *NuPhA*, 777, 579, doi: [10.1016/j.nuclphysa.2004.12.038](https://doi.org/10.1016/j.nuclphysa.2004.12.038)
- Höflich, P. 2009, in *American Institute of Physics Conference Series*, Vol. 1171, *Recent Directions in Astrophysical Quantitative Spectroscopy and Radiation Hydrodynamics*, ed. I. Hubeny, J. M. Stone, K. MacGregor, & K. Werner, 161–172, doi: [10.1063/1.3250057](https://doi.org/10.1063/1.3250057)
- Höflich, P., Gerardy, C. L., Marion, H., & Quimby, R. 2006, *NewAR*, 50, 470, doi: [10.1016/j.newar.2006.06.074](https://doi.org/10.1016/j.newar.2006.06.074)
- Höflich, P., Gerardy, C. L., Nomoto, K., et al. 2004, *ApJ*, 617, 1258, doi: [10.1086/425571](https://doi.org/10.1086/425571)
- Höflich, P., & Stein, J. 2002, *ApJ*, 568, 779, doi: [10.1086/338981](https://doi.org/10.1086/338981)
- Hosseinzadeh, G., Sand, D. J., Lundqvist, P., et al. 2022, *ApJL*, 933, L45, doi: [10.3847/2041-8213/ac7cef](https://doi.org/10.3847/2041-8213/ac7cef)
- Hoyle, F., & Fowler, W. A. 1960, *ApJ*, 132, 565, doi: [10.1086/146963](https://doi.org/10.1086/146963)
- Hristov, B., Collins, D. C., Hoefflich, P., Weatherford, C. A., & Diamond, T. R. 2018, *ApJ*, 858, 13, doi: [10.3847/1538-4357/aab7f2](https://doi.org/10.3847/1538-4357/aab7f2)
- Hristov, B., Hoefflich, P., & Collins, D. C. 2021, *ApJ*, 923, 210, doi: [10.3847/1538-4357/ac0ef8](https://doi.org/10.3847/1538-4357/ac0ef8)
- Hunter, J. D. 2007, *Computing in Science and Engineering*, 9, 90, doi: [10.1109/MCSE.2007.55](https://doi.org/10.1109/MCSE.2007.55)
- Iben, I., J., & Tutukov, A. V. 1984, *ApJS*, 54, 335, doi: [10.1086/190932](https://doi.org/10.1086/190932)
- Jerkstrand, A., Maeda, K., & Kawabata, K. S. 2020, *Science*, 367, 415, doi: [10.1126/science.aaw1469](https://doi.org/10.1126/science.aaw1469)
- Khokhlov, A. M. 1991, *A&A*, 245, 114

- . 2000, arXiv e-prints, astro, doi: [10.48550/arXiv.astro-ph/0008463](https://doi.org/10.48550/arXiv.astro-ph/0008463)
- Kozma, C., & Fransson, C. 1992, *ApJ*, 390, 602, doi: [10.1086/171311](https://doi.org/10.1086/171311)
- Kumar, S., Hsiao, E. Y., Ashall, C., et al. 2023, *ApJ*, 945, 27, doi: [10.3847/1538-4357/acad73](https://doi.org/10.3847/1538-4357/acad73)
- Kurucz, R. L. 1993, *Physica Scripta Volume T*, 47, 110, doi: [10.1088/0031-8949/1993/T47/017](https://doi.org/10.1088/0031-8949/1993/T47/017)
- . 1995, *Highlights of Astronomy*, 10, 579
- Kwok, L. A., Siebert, M. R., Johansson, J., et al. 2023a, arXiv e-prints, arXiv:2308.12450, doi: [10.48550/arXiv.2308.12450](https://doi.org/10.48550/arXiv.2308.12450)
- Kwok, L. A., Jha, S. W., Temim, T., et al. 2023b, *ApJL*, 944, L3, doi: [10.3847/2041-8213/acb4ec](https://doi.org/10.3847/2041-8213/acb4ec)
- Langanke, K., & Martínez-Pinedo, G. 2000, *NuPhA*, 673, 481, doi: [10.1016/S0375-9474\(00\)00131-7](https://doi.org/10.1016/S0375-9474(00)00131-7)
- Livne, E. 1999, *ApJL*, 527, L97, doi: [10.1086/312405](https://doi.org/10.1086/312405)
- Livne, E., & Arnett, D. 1995, *ApJ*, 452, 62, doi: [10.1086/176279](https://doi.org/10.1086/176279)
- Maguire, K., Sim, S. A., Shingles, L., et al. 2018, *MNRAS*, 477, 3567, doi: [10.1093/mnras/sty280](https://doi.org/10.1093/mnras/sty280)
- Mayker Chen, N., Tucker, M. A., Hoyer, N., et al. 2023, arXiv e-prints, arXiv:2301.05718, doi: [10.48550/arXiv.2301.05718](https://doi.org/10.48550/arXiv.2301.05718)
- Mazzali, P. A., Röpke, F. K., Benetti, S., & Hillebrandt, W. 2007, *Science*, 315, 825, doi: [10.1126/science.1136259](https://doi.org/10.1126/science.1136259)
- Mazzali, P. A., Bikmaev, I., Sunyaev, R., et al. 2020, *MNRAS*, 494, 2809, doi: [10.1093/mnras/staa839](https://doi.org/10.1093/mnras/staa839)
- McClelland, C. M., Garnavich, P. M., Milne, P. A., Shappee, B. J., & Pogge, R. W. 2013, *ApJ*, 767, 119, doi: [10.1088/0004-637X/767/2/119](https://doi.org/10.1088/0004-637X/767/2/119)
- Mera Evans, T. B., Hoeflich, P., & Diehl, R. 2022, *ApJ*, 930, 107, doi: [10.3847/1538-4357/ac5253](https://doi.org/10.3847/1538-4357/ac5253)
- Motohara, K., Maeda, K., Gerardy, C. L., et al. 2006, *ApJL*, 652, L101, doi: [10.1086/509919](https://doi.org/10.1086/509919)
- Ni, Y. Q., Moon, D.-S., Drout, M. R., et al. 2023, arXiv e-prints, arXiv:2304.00625, doi: [10.48550/arXiv.2304.00625](https://doi.org/10.48550/arXiv.2304.00625)
- Nomoto, K. 1982, *ApJ*, 253, 798, doi: [10.1086/159682](https://doi.org/10.1086/159682)
- Pakmor, R., Seitzzahl, I. R., Ruiter, A. J., et al. 2024, arXiv e-prints, arXiv:2402.11010, doi: [10.48550/arXiv.2402.11010](https://doi.org/10.48550/arXiv.2402.11010)
- Patat, F., Höflich, P., Baade, D., et al. 2012, *A&A*, 545, A7, doi: [10.1051/0004-6361/201219146](https://doi.org/10.1051/0004-6361/201219146)
- Pejcha, O., Antognini, J. M., Shappee, B. J., & Thompson, T. A. 2013, *MNRAS*, 435, 943, doi: [10.1093/mnras/stt1281](https://doi.org/10.1093/mnras/stt1281)
- Penney, R., & Hoeflich, P. 2014, *ApJ*, 795, 84, doi: [10.1088/0004-637X/795/1/84](https://doi.org/10.1088/0004-637X/795/1/84)
- Poludnenko, A. Y., Chambers, J., Ahmed, K., Gamezo, V. N., & Taylor, B. D. 2019, *Science*, 366, aau7365, doi: [10.1126/science.aau7365](https://doi.org/10.1126/science.aau7365)
- Remming, I. S., & Khokhlov, A. M. 2014, *ApJ*, 794, 87, doi: [10.1088/0004-637X/794/1/87](https://doi.org/10.1088/0004-637X/794/1/87)
- Riess, A. G. 2017, in *Handbook of Supernovae*, ed. A. W. Alsabti & P. Murdin, 2615, doi: [10.1007/978-3-319-21846-5_105](https://doi.org/10.1007/978-3-319-21846-5_105)
- Seitzzahl, I. R., & Townsley, D. M. 2017, in *Handbook of Supernovae*, ed. A. W. Alsabti & P. Murdin, 1955, doi: [10.1007/978-3-319-21846-5_87](https://doi.org/10.1007/978-3-319-21846-5_87)
- Shahbandeh, M., Ashall, C., Hoeflich, P., et al. 2024, arXiv e-prints, arXiv:2401.14474, <https://arxiv.org/abs/2401.14474>
- Shen, K. J., Kasen, D., Miles, B. J., & Townsley, D. M. 2018, *ApJ*, 854, 52, doi: [10.3847/1538-4357/aaa8de](https://doi.org/10.3847/1538-4357/aaa8de)
- Shingles, L. J., Sim, S. A., Kromer, M., et al. 2020, *MNRAS*, 492, 2029, doi: [10.1093/mnras/stz3412](https://doi.org/10.1093/mnras/stz3412)
- Siebert, M. R., Kwok, L. A., Johansson, J., et al. 2024, *ApJ*, 960, 88, doi: [10.3847/1538-4357/ad0975](https://doi.org/10.3847/1538-4357/ad0975)
- Sugimoto, D., Fujimoto, M. Y., Nariai, K., & Nomoto, K. 1979, in *IAU Colloq. 53: White Dwarfs and Variable Degenerate Stars*, ed. H. M. van Horn, V. Weidemann, & M. P. Savedoff, 280
- Tartaglia, L., Sand, D. J., Valenti, S., et al. 2018, *ApJ*, 853, 62, doi: [10.3847/1538-4357/aaa014](https://doi.org/10.3847/1538-4357/aaa014)
- Telesco, C. M., Höflich, P., Li, D., et al. 2015, *ApJ*, 798, 93, doi: [10.1088/0004-637X/798/2/93](https://doi.org/10.1088/0004-637X/798/2/93)
- Thielemann, F.-K. 2019, in *Nuclei in the Cosmos XV*, Vol. 219, 125–134, doi: [10.1007/978-3-030-13876-9_21](https://doi.org/10.1007/978-3-030-13876-9_21)
- Thompson, T. A. 2011, *ApJ*, 741, 82, doi: [10.1088/0004-637X/741/2/82](https://doi.org/10.1088/0004-637X/741/2/82)
- Tucker, M. A., Ashall, C., Shappee, B. J., et al. 2022, *ApJL*, 926, L25, doi: [10.3847/2041-8213/ac4fbd](https://doi.org/10.3847/2041-8213/ac4fbd)
- van Hoof, P. A. M. 2018, *Galaxies*, 6, 63, doi: [10.3390/galaxies6020063](https://doi.org/10.3390/galaxies6020063)
- Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, *Nature Methods*, 17, 261, doi: [10.1038/s41592-019-0686-2](https://doi.org/10.1038/s41592-019-0686-2)
- Webbink, R. F. 1984, *ApJ*, 277, 355, doi: [10.1086/161701](https://doi.org/10.1086/161701)
- Whelan, J., & Iben, Icko, J. 1973, *ApJ*, 186, 1007, doi: [10.1086/152565](https://doi.org/10.1086/152565)
- Wilk, K. D., Hillier, D. J., & Dessart, L. 2018, *MNRAS*, 474, 3187, doi: [10.1093/mnras/stx2816](https://doi.org/10.1093/mnras/stx2816)
- Yang, Y., Yan, H., Wang, L., et al. 2022, *ApJ*, 939, 18, doi: [10.3847/1538-4357/ac8d5f](https://doi.org/10.3847/1538-4357/ac8d5f)

APPENDIX

A. CHANNEL 4 REDUCTION

Due to the increased noise from the background in Channel 4, we chose to manually extract the SN flux in this region. To do this we use an STScI notebook⁹ designed to perform extraction on a point source in JWST NIRSpec IFU data. We adapt this for use with MIRI/MRS data, using the final s3d data cube obtained from MAST. We extract the spectrum using a linearly expanding circular aperture (cone) because the PSF size increases with wavelength. The spectrum at the position of the SN is obtained by extracting the raw flux using various-size apertures at the location of the SN. After visual inspection, we chose to proceed with an aperture radius of 0.273 arcsec (this is 1 pixel in radius). Although this aperture size is likely to be smaller than the true PSF of the SN, using a larger size dramatically increases the noise in the data. To remove the background contribution, individual background spectra are constructed at 32 locations away from the SN across the field of view. For each spectrum, the background is subtracted from the raw SN flux, see Fig. A.1. There is a large variation in the final flux depending on the location of the background selected. We opt to use a background where the continuum is roughly flat. Our chosen optimum background position is centered around pixels $x=15$, $y=23$.

Figure A.2 shows the final SN spectrum in Channel 4. Due to the small aperture this spectrum has to be scaled by a factor of 6 to match the flux in Channel 3. Overall, we highlight that the flux in Channel 4 is uncertain, but we are confident in the width and location of the features identified. Finally, we note that combining all 32 background spectra before subtracting from the raw SN spectrum does not successfully remove the background flux and leaves a large excess at longer wavelengths.




Figure A.1. 32 spectra extracted using different background positions within the IFU data cube.

⁹ https://spacetelescope.github.io/jdat_notebooks/notebooks/ifu_optimal/ifu_optimal.html

xpixel=15, ypixel=23




Figure A.2. The raw flux at the position of the SN (red), with the flux at the position of the optimal background region centered on pixel $x=15$, $y=23$. The final Channel 4, produced from subtracting the extractions is plotted in black.

B. VELOCITIES

In this section, we fit spectral line profiles of the dominant ions in selected wavelength ranges to determine the region in the ejecta in which they are formed. As the spectral resolution of MIRI/MRS is much higher than the previous MIRI/LRS observations, this analysis has the potential to allow us to determine more accurately the location of the emitting regions within the ejecta. However, as discussed in the main text, many of the atomic line transitions are not known in this region and detailed spectral modeling is required to make significant progress. Despite this, we fit the spectra below.

For the fitting process, the *scipy.optimize* package is used, and the best fits are determined via using the Levenberg-Marquardt algorithm and the least squares statistic option. In each wavelength region, the emission features within the regions are assumed to be composed of Gaussian profiles. Although treating emission line profiles as Gaussians makes assumptions about the distribution of the material in the emitting region, it does provide us with a quantitative way to analyze the data. Priors and bounds are provided to each fit to ensure that they are consistent with the line IDs provided from the models in § 6. If there is a region in which a line ID is not provided from the models, but there is a clear feature we also add an emission profile in this region. The spectral fits can be found in Fig. B.1, and the corresponding values, plotted in Fig. 10, are presented in table B.1. Finally, we choose not to fit the feature between 8-10 μm due to the complex blend around the “flat-topped” Ar region, and we do not fit longwards of 18.6 μm due to the uncertainty in the Channel 4 data, and the fit around the [Co III] 11.888 μm feature is shown in the main body of the text (see § 5.1).

C. OPTICAL TO 4.7 μm LINE LIST FROM MODELS

For completeness, in Table C.1 the line list from our models in the wavelength regime shorter than our observations is provided.

Table B.1. The best fit parameters from the ions used in Fig. 10.

Ion	Wavelength	ν_{peak}	Error ν_{peak}	Amplitude	Error Amplitude	σ	Error σ
	μm	μm	F_λ	F_λ	μm	μm	
[Ni II]	6.636	6.644	0.004	1.699	0.052	0.092	0.008
[Fe II]	6.721	6.771	0.005	1.040	0.109	0.064	0.008
[Ar II]	6.985	6.966	0.008	1.501	0.051	0.086	0.012
[Ni III]	7.349	7.360	0.005	2.564	0.024	0.139	0.013
[Co I]	7.202	7.202	0.004	1.288	0.167	0.054	0.005
[Co III]	7.103	7.093	0.006	1.203	0.152	0.066	0.012
[Co I]	7.507	7.472	0.003	0.489	0.195	0.048	0.011
[Fe II]	17.936	17.926	0.020	1.011	0.245	0.100	0.042
[Co I]	18.265	18.282	0.023	0.587	0.079	0.100	0.036
[Fe II]	10.189	10.212	0.011	0.337	0.019	0.100	0.016
[Co III]	10.523	10.555	0.010	0.784	0.162	0.085	0.023
[Co II]	10.682	10.712	0.027	0.439	0.059	0.100	0.040
[Co III]	11.888	11.917	0.002	1.013	0.008	0.196	0.002

D. MIR TRANSITIONS WITHOUT CROSS-SECTIONS

Weak MIRI/MRS features are used to calibrate the atomic models. One of the main uncertainties in the modeling is the lack of atomic data for many weak transitions (Hoeflich et al. 2021) or, here, the lifetimes or Einstein coefficients A_{lu} for the spontaneous decay.

In principle, we follow the method employed by Kurucz (1993, 1995) who calibrated allowed cross-sections by comparing synthetic spectra with the observed solar spectrum. Similarly, we make use of the MIRI spectrum of SN 2021aefx to estimate Einstein A-values. We do not change the atomic models and only adjust atomic cross-sections between known energy levels but without previously measured values. Because of the low densities and non-thermal excitation, the full rate equations are simulated for the given background.

However, the applicability of the method is limited by the $S/N \approx 5 - 10$ in the continuum and the uncertainty in the absolute calibration between the channels (Sect. 6.3.2). Moreover, the features are smeared out over several 1000 km s^{-1} . Many weak features are present in the quasi-continuum in both the observations and synthetic spectra (see Fig. 13), e.g. between 13 and $23 \mu\text{m}$, but not always: Two features are predicted at $15.9 \mu\text{m}$ and $16.8 \mu\text{m}$, whereas a single, broad feature has been observed without a corresponding transition in Table D.1 to fill the gap. The method can be made more complete by future SNe Ia observations with MIRI/MRS during the early-time nebular phase. For SN 2021aefx at day +415, only a few features are well above the noise level and can be used individually. However, the underlying flux is produced by many weak lines of Fe, Co, and Ni I-IV and, at +415 d, a quasi-continuum of free-free and bound-free and some allowed lines produced in inner layers (Diamond et al. 2015). The A_{lu} values have been found by a Monte-Carlo scheme for some individual transitions by attributing the ‘missing’ continuum flux by a global *default* A_{lu} value, with the latter determined by the residual flux in the continuum around $9.5 \pm 0.4 \mu\text{m}$ and $15 \pm 2 \mu\text{m}$. The values adopted are given in Tabs. D.1 and D.2. The corrections to the flux are on the percent level.




Figure B.1. Spectral fits of three main regions in the data.

Table C.1. Line contributions to the spectra from the optical to 4.7 μm at +415 d from the Reference Model with $\rho_c = 1.1 \times 10^9 \text{ g cm}^{-3}$ and $B = 10^6 \text{ G}$.

S	$\lambda [\mu\text{m}]$	Ion	S	$\lambda [\mu\text{m}]$	Ion	S	$\lambda [\mu\text{m}]$	Ion	S	$\lambda [\mu\text{m}]$	Ion	S	$\lambda [\mu\text{m}]$	Ion
	0.3689	[Co II]		0.4104	[Co II]		0.4116	[Fe II]		0.4178	[Fe II]	** *	0.4245	[Fe II]
*	0.4246	[Fe II]	**	0.4278	[Fe II]	*	0.4289	[Fe II]		0.4307	[Fe II]	*	0.4321	[Fe II]
	0.4327	[Ni II]	*	0.4348	[Fe II]	*	0.4354	[Fe II]		0.436	[Fe II]		0.4361	[Fe II]
*	0.4374	[Fe II]		0.4415	[Fe II]	*	0.4418	[Fe II]		0.4453	[Fe II]	*	0.4459	[Fe II]
	0.4476	[Fe II]		0.449	[Fe II]		0.4494	[Fe II]		0.4501	[Co III]		0.4608	[Fe III]
	0.4624	[Co II]		0.4641	[Fe II]	** *	0.4659	[Fe III]	*	0.4703	[Fe III]	*	0.4729	[Fe II]
*	0.4735	[Fe III]		0.4749	[Co II]	*	0.4756	[Fe III]	*	0.4771	[Fe III]		0.4804	[Co II]
*	0.4882	[Fe III]	*	0.4891	[Fe II]		0.4932	[Fe III]	**	0.5013	[Fe III]		0.5086	[Fe III]
*	0.5113	[Fe II]	**	0.516	[Fe II]		0.5222	[Fe II]	*	0.5263	[Fe II]		0.527	[Co II]
**	0.5272	[Fe III]		0.5298	[Fe II]	*	0.5335	[Fe II]	*	0.5378	[Fe II]		0.5414	[Fe III]
	0.5472	[Co II]		0.5548	[Co II]		0.5562	[Co II]		0.589	[Co III]		0.5976	[Fe I]
	0.6016	[Fe II]		0.6129	[Co III]		0.6197	[Co III]		0.6578	[Co III]		0.6586	[Co I]
	0.6669	[Ni II]		0.6855	[Fe II]		0.6934	[Co II]		0.7138	[Ar III]		0.7155	[Co III]
** *	0.7157	[Fe II]	*	0.7174	[Fe II]		0.7249	[Co I]	** *	0.7293	[Ca II]	**	0.7326	[Ca II]
*	0.738	[Ni II]	*	0.739	[Fe II]	*	0.7414	[Ni II]	**	0.7455	[Fe II]		0.7541	[Co II]
*	0.764	[Fe II]	*	0.7689	[Fe II]		0.7892	[Ni III]		0.803	[Co II]	**	0.8123	[Co II]
	0.8123	[Co II]		0.8303	[Ni II]		0.8336	[Co II]		0.8466	[Co II]		0.8469	[Co II]
	0.8502	[Ni III]		0.8546	[Co I]	*	0.8574	[Co II]		0.8583	[Co II]		0.8597	[Co I]
**	0.8619	[Fe II]	** *	0.8894	[Fe II]	*	0.9036	[Fe II]	**	0.9054	[Fe II]		0.9071	[S III]
**	0.9229	[Fe II]	*	0.927	[Fe II]	*	0.9339	[Co II]	*	0.9345	[Co II]		0.9447	[Fe III]
	0.9474	[Fe II]		0.9533	[S III]		0.9642	[Co II]	*	0.9642	[Co II]		0.9697	[Co I]
	0.9705	[Fe III]		0.9983	[Fe II]	*	0.9483	[Fe II]	** *	1.0191	[Co II]	*	1.0248	[Co II]
	1.0283	[Co II]		1.0283	[Co II]		1.0611	[Fe III]		1.0718	[Ni II]		1.0718	[Ni II]
**	1.0824	[S I]		1.0885	[Fe III]	*	1.0976	[Co II]		1.0994	[Si I]	*	1.1283	[Co II]
*	1.1309	[S I]		1.1616	[Ni II]		1.2122	[Fe II]		1.2525	[Fe II]	** *	1.257	[Fe II]
**	1.2707	[Fe II]	*	1.2791	[Fe II]	*	1.2946	[Fe II]	*	1.2981	[Fe II]	**	1.3209	[Fe II]
*	1.321	[Fe I]	**	1.3281	[Fe II]		1.3422	[Fe I]		1.3556	[Fe I]		1.3676	[Fe I]
*	1.3722	[Fe II]	**	1.3733	[Fe I]		1.3762	[Fe I]		1.4055	[Co II]	**	1.4434	[Fe I]
	1.4972	[Co II]	** *	1.5339	[Fe II]		1.5474	[Co II]		1.5488	[Co III]		1.5694	[Co II]
** *	1.5999	[Fe II]		1.6073	[Si I]		1.6267	[Co II]	*	1.6347	[Co II]	** *	1.644	[Fe II]
*	1.6459	[Si I]	** *	1.6642	[Fe II]	**	1.6773	[Fe II]	*	1.7116	[Fe II]		1.7289	[Co II]
	1.7366	[Co II]		1.7413	[Co III]	**	1.7454	[Fe II]	**	1.7976	[Fe II]	**	1.8005	[Fe II]
*	1.8099	[Fe II]		1.8119	[Fe II]	*	1.904	[Co II]	**	1.9393	[Ni II]		1.9581	[Co III]
	2.0028	[Co III]		2.0073	[Fe II]		2.0418	[Ti II]	**	2.0466	[Fe II]		2.0492	[Ni II]
	2.0979	[Co III]		2.1334	[Fe II]	*	2.1457	[Fe III]		2.1605	[Ti II]	**	2.2187	[Fe III]
**	2.2425	[Fe III]	**	2.2443	[Fe II]		2.3086	[Ni II]	** *	2.3486	[Fe III]		2.3695	[Ni II]
** *	2.4781	[Fe II]	**	2.5255	[Co I]	*	2.6521	[Co I]	**	2.7173	[Fe III]	*	2.8713	[Co I]
**	2.8742	[Fe III]	**	2.9048	[Fe III]	**	2.9114	[Ni II]	*	2.9542	[Co I]	**	2.9610	[Fe II]
	3.0305	[Co I]	**	3.0439	[Fe III]		3.0457	[Co I]	** *	3.12	[Ni I]		3.2294	[Fe III]
	3.3942	[Ni III]		3.4917	[Co III]	*	3.6334	[Co I]		3.7498	[Co I]		3.8023	[Ni III]
**	3.9524	[Ni I]	** *	4.0763	[Fe II]		4.082	[Fe II]	**	4.115	[Fe II]		4.3071	[Co II]
	4.5196	[Ni I]	**	4.6077	[Fe II]									

NOTE—The relative strengths are indicated by the number of *. For transitions without known lifetimes (marked by †), $A_{i,j}$ are assumed from the equivalent iron levels.

Table D.1. Life-times of transitions without experimental values have been calculated based on a combination of observations and the base model.

λ [μm]	Ion	Term	$J_{l,u}$	A_{lu}	$E_{l,u}$ [cm^{-1}]	λ [μm]	Ion	Term	$J_{l,u}$	A_{lu}	$E_{l,u}$ [cm^{-1}]
5.022021	[Fe II]	a2G-a2P	7/2,3/2	1E-3	16369.41-18360.64	5.051034	[Fe II]	b4P-b4F	5/2,7/2	1E-3	20830.55-22810.35
5.022021	[Fe II]	a2G-a2P	7/2,3/2	1E-3	16369.41-18360.64	5.051034	[Fe II]	b4P-b4F	5/2,7/2	1E-3	20830.55-22810.35
5.0623456	[Fe II]	a6D-a4F	3/2,5/2	1E-3	862.61-2837.98	5.105920	[Co II]	b1G-a3H	4,6	1E-3	25147.23-27105.74
5.144614	[Fe II]	a2P-b4P	1/2,5/2	1E-3	18886.77-20830.55	5.149301	[Co II]	a1D-a3P	2,0	1E-3	11651.28-13593.29
5.157755	[Co II]	b3F-a3P	2,2	1E-3	11321.86-13260.69	5.282999	[Fe II]	a2D2-b4P	5/2,1/2	1E-3	20516.95-22409.82
5.301864	[Fe II]	a4G-a2F	11/2,7/2	1E-3	25428.79-27314.92	5.3401693	[Fe II]	a6D-a4F	9/2,9/2	1E-2	0.00-1872.60
5.3736547	[Fe II]	a6D-a4F	1/2,5/2	1E-2	977.05-2837.98	5.396797	[Co II]	a3G-b3P	3,1	1E-3	22414.43-24267.38
5.439462	[Co II]	b3F-a1D	4,2	2E-3	9812.86-11651.28	5.456068	[Fe II]	b2P-a2F	3/2,5/2	1E-3	25787.58-27620.40
5.460209	[Fe II]	a2H-b4F	9/2,9/2	1E-3	20805.76-22637.19	5.477981	[Fe I]	a5P-a3P	2,1	1E-3	17726.99-19552.48
5.509410	[Fe II]	a4G-a2F	9/2,5/2	1E-3	25805.33-27620.40	5.535132	[Fe II]	b4P-b4F	5/2,9/2	1E-3	20830.55-22637.19
5.596278	[Co II]	b3P-c3P	2,0	1E-3	24074.42-25861.32	5.604112	[Co II]	a3D-a3D	3,1	1E-3	27484.37-29268.78
5.6316	[Co III]	a4P-a2G	5/2,9/2	1E-3	15201.90-16977.60	5.6739070	[Fe II]	a6D-a4F	5/2,7/2	1E-2	667.68-2430.14
5.801083	[Co II]	c3P-a1P	0,1	1E-3	25861.32-27585.14	5.802874	[Fe II]	a2D2-b4F	3/2,3/2	1E-3	21308.00-23031.28
5.826	[Co IV]	3H-3F2	6,4	1E-3	23679.50-25396.00	5.845	[Ni IV]	4P-2G	5/2,9/2	1E-3	18118.60-19829.60
5.868	[Co IV]	3H-3F2	5,3	1E-3	24031.80-25735.90	5.893	[Co IV]	3H-3F2	4,2	1E-3	24272.00-25969.00
6.024136	[Co II]	a3G-b3P	3,2	5E-3	22414.43-24074.42	6.102182	[Fe II]	a4G-a2F	7/2,5/2	5E-3	25981.65-27620.40
6.129884	[Fe II]	a2D2-b4F	3/2,5/2	5E-3	21308.00-22939.35	6.134290	[Fe II]	a2P-a2D2	1/2,5/2	5E-3	18886.77-20516.95
6.153483	[Fe I]	a5P-a3P	1,1	5E-3	17927.38-19552.48	6.227663	[Fe II]	a4H-a6S	7/2,5/2	5E-3	21711.90-23317.64
6.260968	[Co II]	a3F-a3F	4,2	3E-2	0.00,1597.20	6.332062	[Fe II]	b4P-b4P	5/2,1/2	5E-3	20830.55-22409.82
6.3794832	[Fe II]	a6D-a4F	3/2,7/2	2E-2	862.61-2430.14	6.389812	[Fe II]	a4G-a2F	5/2,5/2	1E-3	26055.41-27620.40
6.547345	[Fe II]	b2P-a2F	3/2,7/2	1E-3	25787.58-27314.92	6.624310	[Fe II]	a4G-a2F	9/2,7/2	1E-3	25805.33-27314.92
6.626905	[Co II]	b3F-b3F	4,2	2E-3	9812.86-11321.86	6.62998	[Ni I]	3D-3D	3,1	1E-2	204.79-1713.09
6.641916	[Fe II]	b4P-a6S	3/2,5/2	1E-3	21812.05-23317.64	6.656252	[Fe II]	a2D2-b4F	3/2,7/2	1E-3	21308.00-22810.35
6.721277	[Fe II]	a6D-a4F	7/2,9/2	5E-2	384.79-1872.60	6.831	[Co IV]	3H-3F2	4,3	1E-3	24272.00-25735.90
6.890	[Co III]	a4F-a4F	9/2,5/2	1E-2	0.00-1451.30	6.986	[Ni IV]	4F-4F	7/2,3/2	5E-3	1189.70-2621.10
7.201	[Co IV]	3P2-3H	2,4	1E-3	22883.30-24272.00	7.217029	[Fe II]	a4H-b4F	13/2,9/2	1E-3	21251.58-22637.19
7.246432	[Fe II]	a4H-b4F	11/2,7/2	1E-3	21430.36-22810.35	7.290488	[Fe II]	a2H-a4H	11/2,7/2	1E-3	20340.25-21711.90
7.330	[Co IV]	3H-3F2	5,4	1E-3	24031.80-25396.00	7.48092	[Ni I]	3D-3F	2,2	2E-2	879.82-2216.55
7.500338	[Fe II]	a4G-a2F	7/2,7/2	1E-3	25981.65-27314.92	7.579280	[Fe II]	a4H-b4F	7/2,3/2	1E-3	21711.90-23031.28
7.721459	[Fe II]	a2D2-b4P	5/2,3/2	1E-3	20516.95-21812.05	7.888699	[Fe II]	b2H-a2F	9/2,5/2	1E-3	26352.77-27620.40
7.939619	[Fe II]	a4G-a2F	5/2,7/2	1E-3	26055.41-27314.92	8.055622	[Fe II]	a2H-a4H	11/2,9/2	1E-3	20340.25-21581.62
8.066	[Co IV]	3P2-3F2	1,2	1E-3	24729.20-25969.00	8.138479	[Fe II]	a4H-b4F	9/2,7/2	1E-3	21581.62-22810.35
8.146939	[Fe II]	a4H-b4F	7/2,5/2	1E-3	21711.90-22939.35	8.201847	[Fe II]	b4P-b4F	3/2,3/2	1E-3	21812.05-23031.28
8.262448	[Co II]	a5F-a5F	5,3	2E-3	3350.49-4560.79	8.286112	[Fe II]	a4H-b4F	11/2,9/2	1E-3	21430.36-22637.19
8.299328	[Fe II]	a6D-a4F	5/2,9/2	1E-3	667.68-1872.60	8.317945	[Co II]	a5P-a1G	3,4	1E-3	17771.51-18973.73
8.368601	[Fe II]	a2D2-a4H	5/2,7/2	1E-3	20516.95-21711.90	8.37951	[Ni I]	3F-1D	2,2	1E-3	2216.55-3409.94
8.735629	[Fe II]	b2H-a2F	11/2,7/2	1E-3	26170.18-27314.92	8.87015	[Ni I]	3D-3F	3,3	1E-2	204.79-1332.16
8.870707	[Fe II]	b4P-b4F	3/2,5/2	1E-3	21812.05-22939.35	8.897	[Co IV]	3H-3F2	4,4	1E-3	24272.00-25396.00
9.075902	[Fe II]	a2D2-b4P	3/2,1/2	1E-3	21308.00-22409.82	9.103740	[Fe II]	a4H-b4F	7/2,7/2	1E-3	21711.90-22810.35
9.173384	[Fe II]	a2H-a4H	11/2,11/2	1E-3	20340.25-21430.36	9.279	[Co IV]	5D-5D	4,2	1E-2	0.00-1077.70
9.321289	[Co II]	b3P-b1G	2,4	1E-3	24074.42-25147.23	9.392655	[Fe II]	a2D2-a4H	5/2,9/2	1E-3	20516.95-21581.62
9.473466	[Fe II]	a4H-b4F	9/2,9/2	1E-3	21581.62-22637.19	9.522585	[Co II]	b3P-c3P	1,1	1E-3	24267.38-25317.52
9.750	[Co III]	a4F-a4F	7/2,3/2	1E-2	841.20-1866.80	9.933	[Co IV]	3P2-3F2	1,3	1E-3	24729.20-25735.90
10.017024	[Fe II]	b4P-b4F	3/2,7/2	1E-3	21812.05-22810.35	10.077445	[Fe I]	a3F-a3F	4,2	1E-3	11976.24-12968.55
10.257231	[Co II]	c3P-c3P	2,0	1E-3	24886.40-25861.32	10.358606	[Fe II]	a4F-a4F	9/2,5/2	1E-3	1872.60-2837.98
10.393377	[Fe II]	b2H-a2F	9/2,7/2	1E-3	26352.77-27314.92	10.605061	[Co II]	b3F-a1D	3,2	1E-3	10708.33-11651.28
10.612332	[Co II]	a5P-a1G	2,4	1E-3	18031.43-18973.73	10.807321	[Fe II]	a4H-b4F	7/2,9/2	1E-3	21711.90-22637.19
10.822771	[Fe II]	a4G-b2H	11/2,9/2	1E-3	25428.79-26352.77	10.856891	[Co II]	a5F-a5F	4,2	1E-3	4028.99-4950.06
10.972887	[Fe II]	a2H-a4H	11/2,13/2	1E-3	20340.25-21251.58						

NOTE—The relative strengths of individual features are fitted. For many weak lines, a standard value has been assumed based on the overall spectra (see text).

Table D.2. Same as Table 4 but for 11-30 μm .

λ [μm]	Ion	Term	$J_{l,u}$	A_{lu}	$E_{l,u}[\text{cm}^{-1}]$	λ [μm]	Ion	Term	$J_{l,u}$	A_{lu}	$E_{l,u}[\text{cm}^{-1}]$
11.015432	[Fe II]	b4P-a6S	1/2,5/2	1E-3	22409.82-23317.64	11.035906	[Fe II]	a2H-a4H	9/2,7/2	1E-3	20805.76-21711.90
11.037065	[Co II]	b3P-c3P	0,1	1E-3	24411.48-25317.52	11.36601	[Ni I]	3F-3D	4,2	2E-2	0.00-879.82
11.398313	[Fe II]	a4G-b2P	5/2,1/2	1E-3	26055.41-26932.73	11.825	[Co IV]	3G-3G	5,3	1E-3	29021.80-29867.50
11.908	[Ni IV]	4P-4P	5/2,1/2	1E-3	18118.60-18958.40	12.077222	[Fe I]	a5P-a3P	3,2	1E-3	17550.18-18378.19
12.315574	[Co II]	b3P-c3P	2,2	1E-3	24074.42-24886.40	12.503266	[Fe I]	a5F-a5F	5,3	2E-3	6928.27-7728.06
12.55624	[Co II]	a3H-a3H	6,4	1E-3	27105.74-27902.16	12.659782	[Co II]	a3G-a3G	5,3	1E-3	21624.53-22414.43
12.889058	[Fe II]	a2H-a4H	9/2,9/2	1E-3	20805.76-21581.62	13.314485	[Fe II]	b4P-a4H	5/2,9/2	1E-3	20830.55-21581.62
13.48815	[Fe II]	a4G-b2H	11/2,11/2	1E-3	25428.79-26170.18	13.790212	[Fe II]	a4D-a4D	7/2,3/2	1E-3	7955.32-8680.47
13.924	[Co IV]	5D-5D	3,1	1E-2	639.10-1357.30	14.204404	[Fe I]	a5D-a5D	4,2	1E-2	0.00-704.01
14.54189	[Fe II]	b2P-a2F	1/2,5/2	1E-3	26932.73-27620.40	14.548612	[Fe II]	a4F-a4F	7/2,3/2	5E-3	2430.14-3117.49
14.69637	[Fe II]	b4F-a6S	9/2,5/2	1E-3	22637.19-23317.64	14.977170	[Fe II]	a6D-a6D	9/2,5/2	1E-2	0.00-667.68
15.35631	[Fe I]	a5P-a3P	2,2	1E-3	17726.99-18378.19	15.53014	[Co II]	a5F-a5F	3,1	1E-3	4560.79-5204.70
16.01042	[Fe II]	a2H-a4H	9/2,11/2	1E-3	20805.76-21430.36	16.09101	[Fe II]	b4P-b4F	1/2,3/2	1E-3	22409.82-23031.28
16.407	[Co III]	a4P-a4P	5/2,1/2	1E-3	15201.90-15811.40	16.41982	[Fe I]	a5F-a5F	4,2	5E-3	7376.76-7985.78
16.878	[Co IV]	3H-3H	6,4	1E-3	23679.50-24272.00	17.572	[Ni III]	3P-3P	2,0	1E-3	16661.60,17230.70
17.63255	[Co II]	a5P-a5P	3,1	1E-3	17771.51-18338.64	18.08790	[Fe II]	a4G-a4G	11/2,7/2	1E-3	25428.79-25981.65
18.26684	[Fe II]	a4G-b2H	9/2,9/2	1E-3	25805.33-26352.77	18.88455	[Fe II]	b4P-b4F	1/2,5/2	1E-3	22409.82-22939.35
19.71262	[Fe II]	b4F-a6S	7/2,5/2	1E-3	22810.35-23317.64	19.83944	[Fe II]	a2D2-b4P	3/2,3/2	1E-3	21308.00-21812.05
19.8624	[Ni I]	3D-3F	1,2	2E-3	1713.09,2216.55	20.928182	[Fe II]	a6D-a6D	7/2,3/2	3E-3	384.79-862.61
20.94479	[Fe II]	b4P-a2D2	5/2,3/2	3E-3	20830.55-21308.00	21.05613	[Co II]	b3P-c3P	0,2	2E-3	24411.48-24886.40
21.17751	[Fe I]	a5D-a5D	3,1	2E-3	415.93-888.13	22.1069	[Ni I]	3D-3F	2,3	1E-2	879.82-1332.16
22.18259	[Fe I]	a5P-a3P	1,2	1E-3	17927.38-18378.19	22.43057	[Fe II]	a2H-a4H	9/2,13/2	2E-3	20805.76-21251.58
23.23350	[Fe II]	a4P-a4P	5/2,1/2	2E-3	13474.45-13904.86	23.43820	[Fe I]	a5F-a5F	3,1	2E-3	7728.06-8154.71
23.93558	[Co II]	a3D-a3H	3,4	2E-3	27484.37-27902.16	24.04	[Co IV]	5D-5D	2,0	1E-2	1077.70-1493.60
24.5422	[Ni I]	3P-3P	2,0	1E-3	15609.84-16017.31	24.75877	[Fe II]	a2D2-a4H	3/2,7/2	1E-3	21308.00-21711.90
26.2520	[Ni I]	3F-3D	3,1	2E-3	1332.16-1713.09	26.43517	[Fe II]	b4F-a6S	5/2,5/2	1E-3	22939.35-23317.64
26.51106	[Fe I]	a5P-a5P	3,1	1E-3	17550.18-17927.38	26.63	[Co III]	a2D2-a2H	5/2,9/2	1E-3	23058.80-23434.30
26.94532	[Fe II]	a4G-b2H	7/2,9/2	1E-3	25981.65-26352.77	27.40822	[Fe II]	a4G-b2H	9/2,11/2	1E-3	25805.33-26170.18

NOTE—The relative strengths of individual features are fitted. For many weak lines, a standard value has been assumed based on the overall spectra (see text).