

From infinite to infinitesimal: Using the Universe as a dataset to probe Casimir corrections to the vacuum energy from fields inhabiting the dark dimension

Luis A. Anchordoqui,^{1,2,3} Ignatios Antoniadis,^{4,5} Dieter Lüst,^{6,7} Neena T. Noble,^{1,3} and Jorge F. Soriano¹

¹*Department of Physics and Astronomy, Lehman College, City University of New York, NY 10468, USA*

²*Department of Physics, Graduate Center, City University of New York, NY 10016, USA*

³*Department of Astrophysics, American Museum of Natural History, NY 10024, USA*

⁴*High Energy Physics Research Unit, Faculty of Science, Chulalongkorn University, Bangkok 1030, Thailand*

⁵*Laboratoire de Physique Théorique et Hautes Énergies - LPTHE, Sorbonne Université, CNRS, 4 Place Jussieu, 75005 Paris, France*

⁶*Max-Planck-Institut für Physik, Werner-Heisenberg-Institut, 80805 München, Germany*

⁷*Arnold Sommerfeld Center for Theoretical Physics, Ludwig-Maximilians-Universität München, 80333 München, Germany*

(Dated: April 2024)

Promptly after high-resolution experiments harbinger the field of precision cosmology low- and high-redshift observations abruptly gave rise to a tension in the measurement of the present-day expansion rate of the Universe (H_0) and the clustering of matter (S_8). The statistically significant discrepancies between the locally measured values of H_0 and S_8 and the ones inferred from observations of the cosmic microwave background assuming the canonical Λ cold dark matter (CDM) cosmological model have become a new cornerstone of theoretical physics. Λ_s CDM is one of the many beyond Standard Model setups that have been proposed to simultaneously resolve the cosmological tensions. This setup relies on an empirical conjecture, which postulates that Λ switched sign (from negative to positive) at a critical redshift $z_c \sim 2$. We reexamine a stringy model that can describe the transition in the vacuum energy hypothesized in Λ_s CDM. The model makes use of the Casimir forces driven by fields inhabiting the incredible bulk of the dark dimension scenario. Unlike the Λ_s CDM setup the model deviates from Λ CDM in the early universe due to the existence of relativistic neutrino-like species. Using the Boltzmann solver CLASS in combination with MontePython we confront predictions of the stringy model to experimental data (from the Planck mission, Pantheon+ supernova type Ia, BAO, and KiDS-1000). We show that the string-inspired model provides a satisfactory fit to the data and can resolve the cosmological tensions.

I. INTRODUCTION

Just about a century after the expansion of the Universe was established [1], the Hubble constant, which measures its rate ($H_0 \equiv 100h$ km/s/Mpc), continues to encounter challenging shortcomings. Inferring H_0 from the measured spectra of temperature and polarization anisotropies in the cosmic microwave background (CMB) requires a cosmological model to describe the evolution of the baryon-photon fluid at early times (i.e., before the last-scattering surface) as well as the evolution of the Universe at later times (i.e., after the last-scattering surface). Λ cold dark matter (CDM) is the simplest model that can provide a good phenomenological fit to current cosmological data [2], and therefore it is generally adopted as benchmark to determine H_0 . Likewise, Λ CDM is assumed in the CMB determination of the growth of cosmic structure (parameterized by $S_8 \equiv \sigma_8 \sqrt{\Omega_m}/0.3$, where σ_8 describes the matter fluctuations at scales of 8 Mpc/ h and Ω_m is the present day value of the non-relativistic matter density).

The $h = 0.674 \pm 0.005$ and $S_8 = 0.834 \pm 0.016$ inferred from *Planck*'s CMB data assuming Λ CDM [3] are in $\sim 5\sigma$ tension with $h = 0.73 \pm 0.01$ from the SH0ES distance ladder measurement (using Cepheid-calibrated

type-Ia supernovae) [4, 5] and in $\sim 3\sigma$ tension with $S_8 = 0.766^{+0.020}_{-0.014}$ from the cosmic shear data of the Kilo-Degree Survey [6], respectively. As a matter of fact, it has been suggested that the H_0 tension is actually a tension on the type Ia supernovae absolute magnitude M_B [7, 8], because the SH0ES H_0 measurement comes directly from M_B estimates. Counting them all, these statistically significant discrepancies have become a new cornerstone of theoretical physics, and many beyond Standard Model (SM) physics models are rising to the challenge [9–12].

Of particular interest herein, Λ_s CDM is one of the models that have been proposed to simultaneously resolve the H_0 , S_8 , and M_B tensions [13–16]. This model relies on an empirical conjecture which postulates that Λ switched sign (from negative to positive) at critical redshift $z_c \sim 2$;

$$\Lambda \rightarrow \Lambda_s \equiv \Lambda_0 \operatorname{sgn}[z_c - z], \quad (1)$$

where $\Lambda_0 > 0$ and $\operatorname{sgn}[x] = -1, 0, 1$ for $x < 0$, $x = 0$ and $x > 0$, respectively. Apart from resolving the three major cosmological tensions, Λ_s CDM achieves quite a good fit to Lyman- α data provided $z_c \lesssim 2.3$ [13], and it is in agreement with the otherwise puzzling JWST observations [17, 18].

Despite the remarkable success of Λ_s CDM to accommodate current experimental data, the model is theoret-

ically unsatisfactory because it postulates that the Universe experienced a rapid transition from an anti-de Sitter (AdS) vacuum to a de Sitter (dS) vacuum, and this holds out against the AdS swampland distance conjecture, which posits that at zero temperature AdS and dS vacua are an infinite distance apart in metric space [19]. In this work, we reexamine a stringy model developed elsewhere [20], which can justify the AdS \rightarrow dS transition hypothesized in Λ_s CDM. Bearing this in mind, throughout we will refer to the model as Λ_s CDM⁺.

The layout of the paper is as follows. In Sec. II we first review the basic setting of Λ_s CDM⁺ and after that we discuss phenomenological implications of the model. In Sec. III we carry out a numerical analysis to confront Λ CDM⁺ with current astrophysical and cosmological observations. The paper wraps up in Sec. IV with conclusions and some discussion.

II. SWAMPLANDISH COSMOLOGY

A. The Dark Dimension

Low-energy effective field theories (EFTs) are the driving gear in the description of low-energy particle physics, cosmology, and gravitational phenomena. It is common ground that the SM and General Relativity should both be understood as leading terms in an EFT expansion. Fundamental principles of string theory or black hole physics can enforce consistency requirements on the space of consistent low-energy EFTs. Indeed, this is the goal of the Swampland program in the quest to understand which are the “good” low-energy EFTs that can couple to gravity consistently (e.g. the landscape of superstring theory vacua) and distinguish them from the “bad” ones that cannot [21]. In theory space, the frontier discerning the good theories from those downgraded to the swampland is drawn by a family of conjectures classifying the properties that an EFT should call for/avoid to enable a consistent completion into quantum gravity [22–25].

For example, the distance conjecture states that infinite distance limits $\Delta\phi \rightarrow \infty$ in the moduli space of massless scalar fields are accompanied by an infinite tower of exponentially light states $m \sim e^{-\alpha\Delta\phi}$, where distance and masses are measured in Planck units [26]. Connected to the distance conjecture is the AdS distance conjecture, which correlates the dark energy density to the mass scale m characterizing the infinite tower of states, $m \sim |\Lambda|^\alpha$, as the negative AdS vacuum energy $\Lambda \rightarrow 0$, where α is an $\mathcal{O}(1)$ positive constant [19]. In addition, under the hypothesis that this scaling behavior holds in dS (or quasi dS) space, an unbounded number of massless modes also pop up in the limit $\Lambda \rightarrow 0$.

The AdS distance conjecture in dS space provides a pathway, dubbed the dark dimension scenario [27], to clear up the origin of the cosmological hierarchy $\Lambda/M_p^4 \sim 10^{-120}$, because it connects the size of the compact space

R_\perp to the dark energy scale $\Lambda^{-1/4}$ via

$$R_\perp \sim \lambda \Lambda^{-1/4}, \quad (2)$$

where M_p is the reduced Planck mass and the proportionality factor is estimated to be within the range $10^{-1} < \lambda < 10^{-4}$.¹ Actually, (2) derives from constraints by theory and experiment. On the theoretical side, since the Kaluza-Klein (KK) tower contains massive spin-2 bosons, the Higuchi bound [30] places an absolute upper limit to the exponent of Λ^α , whereas explicit string calculations of the vacuum energy (see e.g. [31–34]) set a lower bound on α . Altogether, these constraints yield $1/4 \leq \alpha \leq 1/2$. On the experimental side, constraints on deviations from Newton’s gravitational inverse-square law [35] and neutron star heating [36] give rise to the conclusion encapsulated in (2): *The cosmological hierarchy problem can be addressed if there exists one extra dimension of radius R_\perp in the micron range, and the lower bound for $\alpha = 1/4$ is basically saturated [27]*. Within this picture, the SM must be localized on a D-brane transverse to the dark dimension [37]. A theoretical amendment on the connection between the cosmological and KK mass scales confirms $\alpha = 1/4$ [38].

A point worth noting at this juncture is that the KK tower of the dark dimension opens up at the mass scale $m_{\text{KK}} \sim 1/R_\perp$ in the eV range. As a consequence, in the dark dimension scenario the five-dimensional (5D) Planck scale (or species scale where gravity becomes strong [39–42]) is found to be

$$M_* \sim m_{\text{KK}}^{1/3} M_p^{2/3} \sim 10^9 \text{ GeV}. \quad (3)$$

Alternatively, it was recently speculated that the dark dimension can be understood as a line interval with end-of-the-world 9-branes attached at each end. Of course this is equivalent to a semicircular dimension endowed with S^1/\mathbb{Z}_2 symmetry [43].

Adding to the story, the dark dimension scenario has the potential to explain the smallness of neutrino masses. This proposal envisions the right-handed neutrinos as 5D bulk states with Yukawa couplings to the left-handed lepton and Higgs doublets that are localized states on the SM brane [44–46]. The neutrino masses are then suppressed due to the wave function of the bulk states. To be more specific, this is accomplished by introducing three 5D Dirac fermions Ψ_α , which are singlets under the SM gauge symmetries and interact in our brane with the three active left-handed neutrinos in a way that conserves lepton number. The S^1/\mathbb{Z}_2 symmetry in the 5th dimension coordinate x_5 contains x_5 to $-x_5$, which acts as chirality (γ_5) on spinors. Then, in the Weyl basis each Dirac field can be decomposed into two two-component spinors $\Psi_\alpha \equiv (\psi_{\alpha L}, \psi_{\alpha R})^T$.

¹ Auger data of highest energy cosmic rays favor $\lambda \sim 10^{-3}$ [28, 29].

The generation of neutrino masses originates in 5D bulk-brane interactions of the form

$$\mathcal{L} \supset h_{ij} \bar{L}_i \tilde{H} \psi_{jR}(x_5 = 0), \quad (4)$$

where $\tilde{H} = -i\sigma_2 H^*$, L_i denotes the lepton doublets (localized on the SM brane), ψ_{jR} stands for the three bulk (right-handed) R -neutrinos evaluated at the position of the SM brane, $x_5 = 0$ in the fifth-dimension, and h_{ij} are coupling constants. This gives a coupling with the L -neutrinos of the form $\langle H \rangle \bar{\nu}_{L_i} \psi_{jR}(x_5 = 0)$, where $\langle H \rangle = 175$ GeV is the Higgs vacuum expectation value. Expanding ψ_{jR} into modes canonically normalized leads for each of them to a Yukawa 3×3 matrix suppressed by the square root of the volume of the bulk $\sqrt{\pi R_\perp M_s}$, i.e.,

$$Y_{ij} = \frac{h_{ij}}{\sqrt{\pi R_\perp M_s}} \sim h_{ij} \frac{M_s}{M_p}, \quad (5)$$

where $M_s \lesssim M_*$ is the string scale, and where in the second rendition we have dropped factors of π 's and of the string coupling.

Now, neutrino oscillation data can be well-fitted in terms of two nonzero differences $\Delta m_{ij}^2 = m_i^2 - m_j^2$ between the squares of the masses of the three mass eigenstates; namely, $\Delta m_{21}^2 = (7.53 \pm 0.18) \times 10^{-5}$ eV² and $\Delta m_{32}^2 = (2.453 \pm 0.033) \times 10^{-3}$ eV² or $\Delta m_{32}^2 = -(2.536 \pm 0.034) \times 10^{-3}$ eV² [2]. It is easily seen that to obtain the correct order of magnitude of neutrino masses the coupling h_{ij} should be of order 10^{-4} to 10^{-5} for $10^9 \lesssim M_s/\text{GeV} \lesssim 10^{10}$ [47]. In the presence of bulk masses [48, 49], the mixing of the first KK modes to active neutrinos can be suppressed [50, 51].

The latest chapter in the story contemplates dark matter candidates. The dark dimension provides a colosseum for dark matter contenders. In particular, it was observed in [52–54] that the universal coupling of the SM fields to the massive spin-2 KK excitations of the graviton in the dark dimension provides a dark matter candidate. The “dark-to-dark” decays driving the dynamics of the KK modes provide a specific realization of dynamical dark matter framework [55]. Complementary to the dark gravitons, it was discussed in [56–58] that primordial black holes with Schwarzschild radius smaller than a micron could also be good dark matter candidates, possibly even with an interesting close relation to the dark gravitons [59]. Finally, the dark dimension can also accommodate fuzzy dark matter [60].

All in all, the dark dimension facilitates a structured scenario to describe the cosmological evolution of the dark sector. We now turn to discuss how this scenario could also help to resolve the current discordances in the cosmological parameters.

B. $\Lambda_s \text{CDM}^+$

Inspired by the discussion in the previous section, we consider 5D Einstein-de Sitter gravity compactified on a

circle S^1 endowed with S^1/\mathbb{Z}_2 symmetry, and we assume that the SM is localized on a D-brane transverse to the compact 5th dimension, whereas gravity spills into the dark dimension. The effective 4D potential of the modulus controlling the radius (or radion field) R is found to be

$$V(R) = \frac{2\pi \Lambda_5 r^2}{R} + \left(\frac{r}{R}\right)^2 T_4 + V_C(R), \quad (6)$$

where Λ_5 is the 5D cosmological constant, $r \equiv \langle R \rangle$ is the vacuum expectation value of the radion, T_4 is the total 3-brane tension of the model, and V_C stands for the quantum corrections to the vacuum energy due to Casimir forces [61]. These corrections are expected to become important in the deep infrared region, because the Casimir contribution to the potential falls off exponentially at large R compared to the particle wavelength. Indeed, as R decreases different particle thresholds open up,

$$V_C(R) = \sum_i \frac{\pi r^2}{32\pi^7 R^6} (N_F - N_B) \Theta(R_i - R), \quad (7)$$

where $m_i = R_i^{-1}$ are the masses of the 5D fields, Θ is a step function, and $N_F - N_B$ stands for the difference between the number of light fermionic and bosonic degrees of freedom. At the classical level, i.e. considering only the first two terms in (6), it is straightforward to see that the potential develops a maximum at

$$R_{\text{max}} = -T_4/(\pi\Lambda_5), \quad (8)$$

requiring a negative tension T_4 . Note that if the fermionic degrees of freedom overwhelm the bosonic contribution, they would give rise to possible minima, as long as $R_i < R_{\text{max}}$.




FIG. 1: The potential $V(R)$ for $(\Lambda_5)^{1/5} = 22.6$ meV and $|T_4|^{1/4} = 24.2$ meV considering $N_F - N_B = 6$ (AdS) and $N_F - N_B = 7$ (dS).

We now have to define the 5D mass spectrum. Counting the six 5D Weyl fields we find that neutrinos contribute with $N_F = 12$. On the other hand, the 5D graviton contributes with 5 bosonic degrees of freedom. In addition, we assume that the 5D spectrum contains a real

scalar field φ . We further assume that φ has a potential with two local minima, with very small difference in vacuum energy and bigger curvature (mass) of the lower one. Around $z \sim 2$ the false vacuum “tunnels” to its true vacuum state. After the quantum tunneling φ becomes more massive and its contribution to the Casimir energy becomes exponentially suppressed. Altogether, this implies that for $z \gtrsim 2$, the number of bosonic degrees of freedom is $N_B = 6$, but for $z \lesssim 2$, it reduces to $N_B = 5$. In Fig. 1 we show an illustrative example of the phase transition of AdS \rightarrow dS vacua generated by the dark sector described above.

Worthy of mention, the 5D vacuum transition creates a δV contribution to Λ_5 , where $\delta V = V_{\min} - V_{\text{localmin}}$ corresponding to the vacuum energies of the upper (localmin) and the lower (min) minima. We take $\delta V \ll \Lambda_5$ so that it does not perturb the analysis producing the curves shown in Fig. 1.

Now, the AdS \rightarrow dS transition shown in Fig. 1 slightly deviates from the model analyzed in [16], because the fields characterizing the deep infrared region of the dark sector contribute to the effective number of relativistic neutrino-like species N_{eff} [62]. Using conservation of entropy, fully thermalized relics with g_* degrees of freedom contribute

$$\Delta N_{\text{eff}} = g_* \left(\frac{43}{4g_s} \right)^{4/3} \begin{cases} 4/7 & \text{for bosons} \\ 1/2 & \text{for fermions} \end{cases}, \quad (9)$$

where g_s denotes the effective degrees of freedom for the entropy of the other thermalized relativistic species that are present when they decouple [63]. The 5D graviton has 5 helicities, but the spin-1 helicities do not have zero modes, because we assume the compactification has S^1/\mathbb{Z}_2 symmetry and so the ± 1 helicities are projected out. The spin-0 is the radion and the spin-2 helicities form the massless (zero mode) graviton. This means that for the 5D graviton, $g_* = 3$. The scalar field φ contributes with $g_* = 1$. The (bulk) left-handed neutrinos are odd, but the right-handed neutrinos are even and so each counts as a Weyl neutrino, for a total $g_* = 2 \times 3$. Assuming that the dark sector decouples from the SM sector before the electroweak phase transition we have $g_s = 106.75$. This gives $\Delta N_{\text{eff}} = 0.25$. It is of interest to investigate whether these extra-relativistic degrees of freedom can spoil the $\Lambda_s\text{CDM}$ predictions for H_0 and S_8 .

III. NUMERICAL ANALYSIS

We adopt the CLASS+MontePython code [64–66] to explore the full parameter space of the $\Lambda_s\text{CDM}^+$ model and compare its predictions to ΛCDM and $\Lambda_s\text{CDM}$. We derive constraints on the cosmological parameters of these three models from data-sets and likelihoods given below using the Metropolis-Hastings mode, while ensuring a Gelman-Rubin convergence criterion of $R - 1 < 10^{-2}$ in all the runs [67]. The baseline seven free parameters of the $\Lambda_s\text{CDM}$ and $\Lambda_s\text{CDM}^+$ models are given by

$\mathcal{P} = \{\omega_b, \omega_{cdm}, \theta_s, \tau_{\text{reio}}, n_s, A_s, z_c\}$, where the first six are the common ones with ΛCDM : (i) the baryon density $\omega_b \equiv \Omega_b h^2$, (ii) the CDM density $\omega_{cdm} \equiv \Omega_c h^2$, (iii) the angular size of the sound horizon at recombination θ_s , (iv) the Thomson scattering optical depth due to reionization τ_{reio} , (v) the scalar spectral index n_s , and (vi) the power spectrum amplitude of adiabatic scalar perturbations A_s . For ΛCDM and $\Lambda_s\text{CDM}$, we take $N_{\text{eff}} = 3.044$ [68], whereas for $\Lambda_s\text{CDM}^+$, we take $N_{\text{eff}} = 3.294$.

To constrain the free parameters of the models we make use of a series of astrophysical and cosmological probes:

- **Planck 2018 CMB data:** The CMB temperature and polarization angular power spectra from the *Planck* 2018 legacy release [3, 69].
- **Transversal BAO:** Measurements of 2D baryon acoustic oscillations (BAO), $\theta_{\text{BAO}}(z)$, obtained in a weakly model-dependent approach, and compiled in Table I of [70].
- **Pantheon+:** The 1701 light curves of 1550 distinct supernovae type Ia, which are distributed in the redshift interval $0.001 \leq z \leq 2.26$ [71].
- **Cosmic Shear:** KiDS-1000 data [72, 73], including the weak lensing two-point statistics data for both the auto and cross-correlations across five tomographic redshift bins [74]. We follow the KiDS team analysis and adopt the COSEBIs (Complete Orthogonal Sets of E/B-Integrals) likelihood [75].

In Fig. 2 we show the one-dimensional posterior distributions and joint marginalized contours (at 68% and 95% CL) of the free parameters of the ΛCDM , $\Lambda_s\text{CDM}$, and $\Lambda_s\text{CDM}^+$ models. One can immediately appreciate the qualitative similarities in the predictions of $\Lambda_s\text{CDM}$ and $\Lambda_s\text{CDM}^+$. The best parameter to distinguish the models seem to be ω_{cdm} and n_s has the potential to discriminate between $\Lambda_s\text{CDM}$ and $\Lambda_s\text{CDM}^+$. There is quite a good agreement between the ΛCDM^+ predicted value of the Hubble constant $h \sim 0.74$ and the SH0ES measurement $h = 0.73 \pm 0.01$ [4, 5]. The situation is even more interesting for S_8 : an estimate of structure growth predicted by $\Lambda_s\text{CDM}^+$ follows from $\sigma_8 \simeq 0.835$, $\omega_b \simeq 2.28 \times 10^{-2}$, $\omega_{cdm} \simeq 0.121$, yielding $S_8 \simeq 0.78$. The predicted S_8 is within 1σ of KiDS-1000 measurement, $S_8 = 0.766^{+0.020}_{-0.014}$ [6].

The one-dimensional posterior distribution of the seventh parameter z_c is shown in Fig. 3. We find that $\Lambda_s\text{CDM}^+$ slightly shifts the transition to higher redshifts, when compared with the value of $z_c = 1.72^{+0.09}_{-0.12}$ reported in [16].

IV. CONCLUSIONS

$\Lambda_s\text{CDM}$ is a promising model for solving the H_0 and S_8 cosmological tensions by demanding the cosmological constant Λ to switch sign (from negative to positive) at a critical redshift $z_c \sim 2$ [13–16]. We have reexam-




FIG. 2: One-dimensional posterior distributions and joint marginalized contours (at 68% and 95% CL) of the free parameters of the Λ CDM, Λ_s CDM, and Λ_s CDM⁺ models using data from *Planck*, transversal BAO, Pantheon+, and KiDS-1000.




FIG. 3: One-dimensional distribution of the critical redshift for the Λ_s CDM and Λ_s CDM⁺ models using data from *Planck*, transversal BAO, Pantheon+, and KiDS-1000.

ined a string-inspired model which can describe the re-

quired transition in the vacuum energy making use of the Casimir forces driven by fields inhabiting the incredible bulk of the dark dimension scenario [20]. The model, dubbed Λ_s CDM⁺, deviates from Λ CDM in the early universe due to the existence of relativistic neutrino-like species. We used the Boltzmann solver CLASS in combination with MontePython to confront model prediction to experimental data. We have shown that Λ_s CDM⁺ provides a satisfactory fit to the data and can resolve the cosmological tensions.

We end by reviewing some key aspects of the H_0 conundrum and after that we call attention to a pressing difference between the Λ_s CDM and Λ_s CDM⁺ models, which could be of utmost importance in resolving the cosmological tensions. At the moment, there is a huge set of astrophysical and cosmological observations that (on average) seem to show a discrepant trend, in which “early-time” model dependent measurements prefer a low H_0 ($0.65 < h < 0.70$), whereas “late-time” direct measurements prefer a high H_0 ($h \sim 0.73$) [12]. To understand the origin of this discrepancy it is worthwhile to look at

the interplay between H_0 and the comoving sound horizon at the end of the baryonic-drag epoch r_{drag} , which sets the transverse BAO scale. The rationale here is that BAO data play an important role in defining the H_0 tension and also in illuminating possible paths to resolve it. Let us then focus on current experimental constraints placed on the H_0 - r_{drag} plane. Firstly, requiring Λ CDM to accommodate Planck data we obtain a constraint on the sound horizon $r_{\text{drag}} = 147.18 \pm 0.29$ Mpc and another one on the Hubble constant $h = 0.674 \pm 0.005$ [3]. Secondly, the allowed region of the H_0 - r_s plane by BAO distance measurements and uncalibrated supernovae is a U-shaped hyperbolic area; see Fig. 1 in [76]. Actually, the central values of the allowed region are at the locus of a hyperbola, because these measurements fix the product between the Hubble constant and the sound horizon. Finally, the local H_0 measurements by the SHOES team are independent of r_{drag} and thus the allowed region represents a band which is parallel to the r_{drag} axis. Putting all this together, we can conclude that to resolve the H_0 tension the “new” cosmological model should be able to bring the Planck constraints from Λ CDM to the region where they can agree with both (i) the local SHOES measurements and (ii) the inverse distance ladder measurements from BAO and Hubble flow supernovae. It is obvious that this requires lowering the sound horizon, because if the product $H_0 \times r_{\text{drag}}$ is fixed, to increase H_0 we must decrease r_{drag} . The comoving linear size of the sound horizon is given by

$$r_{\text{drag}}(z) = \int_{z_{\text{drag}}}^{\infty} \frac{c_s(z')}{H(z')} dz', \quad (10)$$

where

$$c_s(z) = \frac{c}{\sqrt{3\{1 + 3\omega_b/[4\omega_\gamma(1+z)]\}}} \quad (11)$$

is the sound speed of the photon-baryon fluid and $H(z)$ is the Hubble parameter describing the expansion rate [77]. To leading order $c_s \sim c/\sqrt{3}$, and so the sound horizon is given by the integral of the inverse of the expansion rate. At face value, this implies that to solve the H_0 tension we need new physics before the end of the baryonic-drag epoch, because this is the only way we can reduce the sound horizon.

We can also look at the H_0 tension from a perspective which is independent of CMB data. To remain totally independent from CMB measurements, we use data from BAO and uncalibrated supernovae. We take BAO data calibrated in the early universe (using the sound horizon as a free parameter) to readjust the absolute value of the expansion rate at late times, and then we use Hubble flow supernovae to control the shape of the expansion history. Combining information from these observations (while remaining totally independent from the CMB data) we can try reconstructing the late time expansion history of the Universe. However, in doing so, one inexorably finds that deviations from Λ CDM are very tightly constrained,

to no more than 5% for $z < 0.6$, which is well below the level required to address the H_0 tension using only new physics at late times [78]. This exercise certainly provides support to our previous finding; namely, that we need new physics before the end of the baryonic-drag epoch to resolve the H_0 tension.²

In summary, the H_0 tension is actually a calibrate tension between the absolute magnitude of supernovae (which controls the local H_0 value) and the sound horizon (which controls the absolute scale set by BAO data). BAO and uncalibrated supernovae alone are very unforgiving, in the sense that they do not allow huge deviation from Λ CDM at late time. All in all, we arrive at a “no go theorem” which states that *to resolve the H_0 tension we need new physics at early times*.

Now, Λ_s CDM can resolve the cosmological tensions, but does not deviate from Λ CDM for $z > z_c$. This implies that Λ_s CDM violates the H_0 “no go theorem.” Such a violation may be ascribed to the fact that Λ_s CDM has a hidden sudden singularity at z_c [80]. Indeed, it is easily seen that the scale factor a is continuous and non-zero at $t = t_c$, but its first derivative \dot{a} is discontinuous, and its second derivative \ddot{a} diverges.³ It is important to stress that there is an inherent difference between the two models: in contrast to Λ_s CDM, the proposed Λ_s CDM⁺ satisfies the H_0 “no go theorem.” It is not clear whether the Λ_s CDM⁺ model described in Sec. II carries with it a sudden singularity. To a first approximation, we have implemented the model into the CLASS Boltzmann solver using a signum function to characterize the tunneling of φ . In the spirit of [82], an investigation to smoothed-out this sharp transition is obviously important to be done.

Acknowledgements

The work of L.A.A. is supported by the U.S. National Science Foundation (NSF Grant PHY-2112527). I.A. is supported by the Second Century Fund (C2F), Chulalongkorn University. The work of D.L. is supported by the Origins Excellence Cluster and by the German-Israel-Project (DIP) on Holography and the Swampland. N.T.N. is supported by the AstroCom NYC program through NSF Grant AST-2219090.

² Arguments have also been given insinuating that early-time new physics alone is not sufficient to solve the H_0 tension [79].

³ It has been shown that the sudden singularity at z_c does not result in the dissociation of bound systems [81].

- [1] E. Hubble, **A relation between distance and radial velocity among extra-galactic nebulae**, Proc. Nat. Acad. Sci. **15**, 168-173 (1929) doi:10.1073/pnas.15.3.168
- [2] R. L. Workman *et al.* [Particle Data Group], **Review of Particle Physics**, PTEP **2022**, 083C01 (2022) doi:10.1093/ptep/ptac097
- [3] N. Aghanim *et al.* [Planck], **Planck 2018 results VI: Cosmological parameters**, Astron. Astrophys. **641**, A6 (2020) [erratum: Astron. Astrophys. **652**, C4 (2021)] doi:10.1051/0004-6361/201833910 [arXiv:1807.06209 [astro-ph.CO]].
- [4] A. G. Riess, W. Yuan, L. M. Macri, D. Scolnic, D. Brout, S. Casertano, D. O. Jones, Y. Murakami, L. Breuval and T. G. Brink, *et al.* **A comprehensive measurement of the local value of the Hubble constant with 1 km s⁻¹ Mpc⁻¹ uncertainty from the Hubble Space Telescope and the SH0ES Team**, Astrophys. J. Lett. **934** (2022) no.1, L7 doi:10.3847/2041-8213/ac5c5b [arXiv:2112.04510 [astro-ph.CO]].
- [5] Y. S. Murakami, A. G. Riess, B. E. Stahl, W. D. Kenworthy, D. M. A. Pluck, A. Macorettta, D. Brout, D. O. Jones, D. M. Scolnic and A. V. Filippenko, **Leveraging SN Ia spectroscopic similarity to improve the measurement of H₀**, JCAP **11**, 046 (2023) doi:10.1088/1475-7516/2023/11/046 [arXiv:2306.00070 [astro-ph.CO]].
- [6] C. Heymans, T. Tröster, M. Asgari, C. Blake, H. Hildebrandt, B. Joachimi, K. Kuijken, C. A. Lin, A. G. Sánchez and J. L. van den Busch, *et al.*, **KiDS-1000 Cosmology: Multi-probe weak gravitational lensing and spectroscopic galaxy clustering constraints**, Astron. Astrophys. **646**, A140 (2021) doi:10.1051/0004-6361/202039063 [arXiv:2007.15632 [astro-ph.CO]].
- [7] G. Efstathiou, **To H₀ or not to H₀?**, Mon. Not. Roy. Astron. Soc. **505**, no.3, 3866-3872 (2021) doi:10.1093/mnras/stab1588 [arXiv:2103.08723 [astro-ph.CO]].
- [8] D. Camarena and V. Marra, **On the use of the local prior on the absolute magnitude of Type Ia supernovae in cosmological inference**, Mon. Not. Roy. Astron. Soc. **504**, 5164-5171 (2021) doi:10.1093/mnras/stab1200 [arXiv:2101.08641 [astro-ph.CO]].
- [9] E. Di Valentino, O. Mena, S. Pan, L. Visinelli, W. Yang, A. Melchiorri, D. F. Mota, A. G. Riess and J. Silk, **In the realm of the Hubble tension—a review of solutions**, Class. Quant. Grav. **38**, no.15, 153001 (2021) doi:10.1088/1361-6382/ac086d [arXiv:2103.01183 [astro-ph.CO]].
- [10] N. Schöneberg, G. Franco Abellán, A. Pérez Sánchez, S. J. Witte, V. Poulin and J. Lesgourgues, **The H0 Olympics: A fair ranking of proposed models**, Phys. Rept. **984**, 1-55 (2022) doi:10.1016/j.physrep.2022.07.001 [arXiv:2107.10291 [astro-ph.CO]].
- [11] L. Perivolaropoulos and F. Skara, **Challenges for Λ CDM: An update**, New Astron. Rev. **95**, 101659 (2022) doi:10.1016/j.newar.2022.101659 [arXiv:2105.05208 [astro-ph.CO]].
- [12] E. Abdalla, G. Franco Abellán, A. Aboubrahim, A. Agnello, Ö. Akarsu, Y. Akrami, G. Alestas, D. Aloni, L. Amendola and L. A. Anchordoqui, *et al.*, **Cosmology intertwined: A review of the particle physics, astrophysics, and cosmology associated with the cosmological tensions and anomalies**, JHEAp **34**, 49-211 (2022) doi:10.1016/j.jheap.2022.04.002 [arXiv:2203.06142 [astro-ph.CO]].
- [13] Ö. Akarsu, J. D. Barrow, L. A. Escamilla and J. A. Vazquez, **Graduated dark energy: Observational hints of a spontaneous sign switch in the cosmological constant**, Phys. Rev. D **101** (2020) no.6, 063528 doi:10.1103/PhysRevD.101.063528 [arXiv:1912.08751 [astro-ph.CO]].
- [14] Ö. Akarsu, S. Kumar, E. Özülker and J. A. Vazquez, **Relaxing cosmological tensions with a sign switching cosmological constant**, Phys. Rev. D **104** (2021) no.12, 123512 doi:10.1103/PhysRevD.104.123512 [arXiv:2108.09239 [astro-ph.CO]].
- [15] Ö. Akarsu, S. Kumar, E. Özülker, J. A. Vazquez and A. Yadav, **Relaxing cosmological tensions with a sign switching cosmological constant: Improved results with Planck, BAO, and Pantheon data**, Phys. Rev. D **108** (2023) no.2, 023513 doi:10.1103/PhysRevD.108.023513 [arXiv:2211.05742 [astro-ph.CO]].
- [16] Ö. Akarsu, E. Di Valentino, S. Kumar, R. C. Nunes, J. A. Vazquez and A. Yadav, **Λ_s CDM model: A promising scenario for alleviation of cosmological tensions**, [arXiv:2307.10899 [astro-ph.CO]].
- [17] S. A. Adil, U. Mukhopadhyay, A. A. Sen and S. Vagnozzi, **Dark energy in light of the early JWST observations: case for a negative cosmological constant?**, JCAP **10** (2023), 072 doi:10.1088/1475-7516/2023/10/072 [arXiv:2307.12763 [astro-ph.CO]].
- [18] N. Menci, S. A. Adil, U. Mukhopadhyay, A. A. Sen and S. Vagnozzi, **Negative cosmological constant in the dark energy sector: tests from JWST photometric and spectroscopic observations of high-redshift galaxies**, [arXiv:2401.12659 [astro-ph.CO]].
- [19] D. Lüst, E. Palti and C. Vafa, **AdS and the Swampland**, Phys. Lett. B **797**, 134867 (2019) doi:10.1016/j.physletb.2019.134867 [arXiv:1906.05225 [hep-th]].
- [20] L. A. Anchordoqui, I. Antoniadis and D. Lüst, **Anti-de Sitter \rightarrow de Sitter transition driven by Casimir forces and mitigating tensions in cosmological parameters**, [arXiv:2312.12352 [hep-th]].
- [21] C. Vafa, **The string landscape and the swampland**, [arXiv:hep-th/0509212 [hep-th]].
- [22] E. Palti, **The swampland: introduction and review**, Fortsch. Phys. **67**, no.6, 1900037 (2019) doi:10.1002/prop.201900037 [arXiv:1903.06239 [hep-th]].
- [23] M. van Beest, J. Calderón-Infante, D. Mirfendereski and I. Valenzuela, **Lectures on the Swampland Program in string compactifications**, Phys. Rept. **989**, 1-50 (2022) doi:10.1016/j.physrep.2022.09.002 [arXiv:2102.01111 [hep-th]].
- [24] M. Graña and A. Herráez, **The Swampland Conjectures: A bridge from quantum gravity to particle physics**, Universe **7**, no.8, 273 (2021) doi:10.3390/universe7080273 [arXiv:2107.00087 [hep-th]].
- [25] N. B. Agmon, A. Bedroya, M. J. Kang and C. Vafa, **Lectures on the string landscape and the swampland**, [arXiv:2212.06187 [hep-th]].
- [26] H. Ooguri and C. Vafa, **On the Geometry of the String**

- Landscape and the Swampland, Nucl. Phys. B **766**, 21-33 (2007) doi:10.1016/j.nuclphysb.2006.10.033 [arXiv:hep-th/0605264 [hep-th]].
- [27] M. Montero, C. Vafa and I. Valenzuela, **The dark dimension and the Swampland**, JHEP **02**, 022 (2023) doi:10.1007/JHEP02(2023)022 [arXiv:2205.12293 [hep-th]].
- [28] L. A. Anchordoqui, **Dark dimension, the swampland, and the origin of cosmic rays beyond the Greisen-Zatsepin-Kuzmin barrier**, Phys. Rev. D **106**, no.11, 116022 (2022) doi:10.1103/PhysRevD.106.116022 [arXiv:2205.13931 [hep-ph]].
- [29] N. T. Noble, J. F. Soriano and L. A. Anchordoqui, **Probing the Dark Dimension with Auger data**, Phys. Dark Univ. **42** (2023), 101278 doi:10.1016/j.dark.2023.101278 [arXiv:2306.03666 [hep-ph]].
- [30] A. Higuchi, **Forbidden mass range for spin-2 field theory in de Sitter space-time**, Nucl. Phys. B **282** (1987), 397-436 doi:10.1016/0550-3213(87)90691-2
- [31] H. Itoyama and T. R. Taylor, **Supersymmetry restoration in the compactified $O(16) \times O(16)$ -prime heterotic string theory**, Phys. Lett. B **186** (1987), 129-133 doi:10.1016/0370-2693(87)90267-X
- [32] H. Itoyama and T. R. Taylor, **Small cosmological constant in string models**, FERMILAB-CONF-87-129-T.
- [33] I. Antoniadis and C. Kounnas, **Superstring phase transition at high temperature**, Phys. Lett. B **261** (1991), 369-378 doi:10.1016/0370-2693(91)90442-S
- [34] Q. Bonnefoy, E. Dudas and S. Lüster, **On the weak gravity conjecture in string theory with broken supersymmetry**, Nucl. Phys. B **947** (2019), 114738 doi:10.1016/j.nuclphysb.2019.114738 [arXiv:1811.11199 [hep-th]].
- [35] J. G. Lee, E. G. Adelberger, T. S. Cook, S. M. Fleischer and B. R. Heckel, **New test of the gravitational $1/r^2$ law at separations down to 52 μm** , Phys. Rev. Lett. **124**, no.10, 101101 (2020) doi:10.1103/PhysRevLett.124.101101 [arXiv:2002.11761 [hep-ex]].
- [36] S. Hannestad and G. G. Raffelt, **Supernova and neutron star limits on large extra dimensions reexamined**, Phys. Rev. D **67**, 125008 (2003) [erratum: Phys. Rev. D **69**, 029901 (2004)] doi:10.1103/PhysRevD.69.029901 [arXiv:hep-ph/0304029 [hep-ph]].
- [37] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, **New dimensions at a millimeter to a Fermi and superstrings at a TeV**, Phys. Lett. B **436** (1998), 257-263 doi:10.1016/S0370-2693(98)00860-0 [arXiv:hep-ph/9804398 [hep-ph]].
- [38] L. A. Anchordoqui, I. Antoniadis, D. Lüster and S. Lüster, **On the cosmological constant, the KK mass scale, and the cut-off dependence in the dark dimension scenario**, Eur. Phys. J. C **83**, no.11, 1016 (2023) doi:10.1140/epjc/s10052-023-12206-2 [arXiv:2309.09330 [hep-th]].
- [39] G. Dvali, **Black holes and large N species solution to the hierarchy problem**, Fortsch. Phys. **58**, 528-536 (2010) doi:10.1002/prop.201000009 [arXiv:0706.2050 [hep-th]].
- [40] G. Dvali and M. Redi, **Black hole bound on the number of species and quantum gravity at LHC**, Phys. Rev. D **77**, 045027 (2008) doi:10.1103/PhysRevD.77.045027 [arXiv:0710.4344 [hep-th]].
- [41] N. Cribiori, D. Lüster and G. Staudt, **Black hole entropy and moduli-dependent species scale**, Phys. Lett. B **844**, 138113 (2023) doi:10.1016/j.physletb.2023.138113 [arXiv:2212.10286 [hep-th]].
- [42] D. van de Heisteeg, C. Vafa, M. Wiesner and D. H. Wu, **Species scale in diverse dimensions**, [arXiv:2310.07213 [hep-th]].
- [43] J. H. Schwarz, **Concerning a hypothetical mesoscopic dark dimension**, [arXiv:2403.12899 [hep-th]].
- [44] K. R. Dienes, E. Dudas and T. Gherghetta, **Neutrino oscillations without neutrino masses or heavy mass scales: A Higher dimensional seesaw mechanism**, Nucl. Phys. B **557**, 25 (1999) doi:10.1016/S0550-3213(99)00377-6 [arXiv:hep-ph/9811428 [hep-ph]].
- [45] N. Arkani-Hamed, S. Dimopoulos, G. R. Dvali and J. March-Russell, **Neutrino masses from large extra dimensions**, Phys. Rev. D **65**, 024032 (2001) doi:10.1103/PhysRevD.65.024032 [arXiv:hep-ph/9811448 [hep-ph]].
- [46] G. R. Dvali and A. Y. Smirnov, **Probing large extra dimensions with neutrinos**, Nucl. Phys. B **563**, 63-81 (1999) doi:10.1016/S0550-3213(99)00574-X [arXiv:hep-ph/9904211 [hep-ph]].
- [47] L. A. Anchordoqui, I. Antoniadis and D. Lüster, **Aspects of the dark dimension in cosmology**, Phys. Rev. D **107**, no.8, 083530 (2023) doi:10.1103/PhysRevD.107.083530 [arXiv:2212.08527 [hep-ph]].
- [48] A. Lukas, P. Ramond, A. Romanino and G. G. Ross, **Solar neutrino oscillation from large extra dimensions**, Phys. Lett. B **495**, 136-146 (2000) doi:10.1016/S0370-2693(00)01206-5 [arXiv:hep-ph/0008049 [hep-ph]].
- [49] A. Lukas, P. Ramond, A. Romanino and G. G. Ross, **Neutrino masses and mixing in brane world theories**, JHEP **04**, 010 (2001) doi:10.1088/1126-6708/2001/04/010 [arXiv:hep-ph/0011295 [hep-ph]].
- [50] M. Carena, Y. Y. Li, C. S. Machado, P. A. N. Machado and C. E. M. Wagner, **Neutrinos in Large Extra Dimensions and Short-Baseline ν_e Appearance**, Phys. Rev. D **96** (2017) no.9, 095014 doi:10.1103/PhysRevD.96.095014 [arXiv:1708.09548 [hep-ph]].
- [51] L. A. Anchordoqui, I. Antoniadis and J. Cunat, **The dark dimension and the standard model landscape**, Phys. Rev. D **109**, no.1, 016028 (2024) doi:10.1103/PhysRevD.109.016028 [arXiv:2306.16491 [hep-ph]].
- [52] E. Gonzalo, M. Montero, G. Obied and C. Vafa, **Dark Dimension Gravitons as Dark Matter**, JHEP **11**, 109 (2023) doi:10.1007/JHEP11(2023)109 [arXiv:2209.09249 [hep-ph]].
- [53] J. A. P. Law-Smith, G. Obied, A. Prabhu and C. Vafa, **Astrophysical Constraints on Decaying Dark Gravitons**, [arXiv:2307.11048 [hep-ph]].
- [54] G. Obied, C. Dvorkin, E. Gonzalo and C. Vafa, **Dark Dimension and Decaying Dark Matter Gravitons**, [arXiv:2311.05318 [astro-ph.CO]].
- [55] K. R. Dienes and B. Thomas, **Dynamical dark matter I: Theoretical overview**, Phys. Rev. D **85**, 083523 (2012) doi:10.1103/PhysRevD.85.083523 [arXiv:1106.4546 [hep-ph]].
- [56] L. A. Anchordoqui, I. Antoniadis and D. Lüster, **Dark dimension, the swampland, and the dark matter fraction composed of primordial black holes**, Phys. Rev. D **106**, no.8, 086001 (2022) doi:10.1103/PhysRevD.106.086001 [arXiv:2206.07071 [hep-th]].
- [57] L. A. Anchordoqui, I. Antoniadis and D. Lüster, **The dark dimension, the swampland, and the dark matter fraction**

- composed of primordial near-extremal black holes, *Phys. Rev. D* (in press) [arXiv:2401.09087 [hep-th]].
- [58] L. A. Anchordoqui, I. Antoniadis and D. Lüst, **More on black holes perceiving the dark dimension**, [arXiv:2403.19604 [hep-th]].
- [59] L. A. Anchordoqui, I. Antoniadis and D. Lüst, **The dark universe: Primordial black hole \rightleftharpoons dark graviton gas connection**, *Phys. Lett. B* **840**, 137844 (2023) doi:10.1016/j.physletb.2023.137844 [arXiv:2210.02475 [hep-th]].
- [60] L. A. Anchordoqui, I. Antoniadis and D. Lüst, **Fuzzy dark matter and the dark dimension**, *Eur. Phys. J. C* **84**, no.3, 273 (2024) doi:10.1140/epjc/s10052-024-12622-y [arXiv:2307.01100 [hep-ph]].
- [61] L. A. Anchordoqui and I. Antoniadis, **Large extra dimensions from higher-dimensional inflation**, *Phys. Rev. D* (in press) [arXiv:2310.20282 [hep-ph]].
- [62] G. Steigman, D. N. Schramm and J. E. Gunn, **Cosmological limits to the number of massive leptons**, *Phys. Lett. B* **66** (1977), 202-204 doi:10.1016/0370-2693(77)90176-9
- [63] L. A. Anchordoqui, I. Antoniadis, D. Lüst, J. F. Soriano and T. R. Taylor, **H_0 tension and the String Swampland**, *Phys. Rev. D* **101** (2020), 083532 doi:10.1103/PhysRevD.101.083532 [arXiv:1912.00242 [hep-th]].
- [64] D. Blas, J. Lesgourgues and T. Tram, **The Cosmic Linear Anisotropy Solving System (CLASS) II: Approximation schemes**, *JCAP* **07**, 034 (2011) doi:10.1088/1475-7516/2011/07/034 [arXiv:1104.2933 [astro-ph.CO]].
- [65] J. Lesgourgues, **The Cosmic Linear Anisotropy Solving System (CLASS) I: Overview**, [arXiv:1104.2932 [astro-ph.IM]].
- [66] T. Brinckmann and J. Lesgourgues, **MontePython 3: boosted MCMC sampler and other features**, *Phys. Dark Univ.* **24**, 100260 (2019) doi:10.1016/j.dark.2018.100260 [arXiv:1804.07261 [astro-ph.CO]].
- [67] A. Gelman and D. B. Rubin, **Inference from iterative simulation using multiple sequences**, *Statist. Sci.* **7**, 457-472 (1992) doi:10.1214/ss/1177011136
- [68] S. Gariazzo, P. F. de Salas and S. Pastor, **Thermalisation of sterile neutrinos in the early Universe in the 3+1 scheme with full mixing matrix**, *JCAP* **07**, 014 (2019) doi:10.1088/1475-7516/2019/07/014 [arXiv:1905.11290 [astro-ph.CO]].
- [69] N. Aghanim *et al.* [Planck], **Planck 2018 results V: CMB power spectra and likelihoods**, *Astron. Astrophys.* **641**, A5 (2020) doi:10.1051/0004-6361/201936386 [arXiv:1907.12875 [astro-ph.CO]].
- [70] R. C. Nunes, S. K. Yadav, J. F. Jesus and A. Bernui, **Cosmological parameter analyses using transversal BAO data**, *Mon. Not. Roy. Astron. Soc.* **497**, no.2, 2133-2141 (2020) doi:10.1093/mnras/staa2036 [arXiv:2002.09293 [astro-ph.CO]].
- [71] D. Brout, D. Scolnic, B. Popovic, A. G. Riess, J. Zuntz, R. Kessler, A. Carr, T. M. Davis, S. Hinton and D. Jones, *et al.* **The Pantheon+ analysis: Cosmological constraints**, *Astrophys. J.* **938**, no.2, 110 (2022) doi:10.3847/1538-4357/ac8e04 [arXiv:2202.04077 [astro-ph.CO]].
- [72] K. Kuijken, C. Heymans, A. Dvornik, H. Hildebrandt, J. T. A. de Jong, A. H. Wright, T. Erben, M. Bilicki, B. Giblin and H. Y. Shan, *et al.* **The fourth data release of the Kilo-Degree Survey: ugri imaging and nine-band optical-IR photometry over 1000 square degrees**, *Astron. Astrophys.* **625**, A2 (2019) doi:10.1051/0004-6361/201834918 [arXiv:1902.11265 [astro-ph.GA]].
- [73] B. Giblin, C. Heymans, M. Asgari, H. Hildebrandt, H. Hoekstra, B. Joachimi, A. Kannawadi, K. Kuijken, C. A. Lin and L. Miller, *et al.* **KiDS-1000 catalogue: Weak gravitational lensing shear measurements**, *Astron. Astrophys.* **645**, A105 (2021) doi:10.1051/0004-6361/202038850 [arXiv:2007.01845 [astro-ph.CO]].
- [74] H. Hildebrandt, J. L. Van Den Busch, A. H. Wright, C. Blake, B. Joachimi, K. Kuijken, T. Tröster, M. Asgari, M. Bilicki and J. T. A. De Jong, *et al.* **KiDS-1000 catalogue: Redshift distributions and their calibration**, *Astron. Astrophys.* **647**, A124 (2021) doi:10.1051/0004-6361/202039018 [arXiv:2007.15635 [astro-ph.CO]].
- [75] M. Asgari *et al.* [KiDS], **KiDS-1000 Cosmology: Cosmic shear constraints and comparison between two point statistics**, *Astron. Astrophys.* **645**, A104 (2021) doi:10.1051/0004-6361/202039070 [arXiv:2007.15633 [astro-ph.CO]].
- [76] L. Knox and M. Millea, **Hubble constant hunter's guide**, *Phys. Rev. D* **101**, no.4, 043533 (2020) doi:10.1103/PhysRevD.101.043533 [arXiv:1908.03663 [astro-ph.CO]].
- [77] W. Hu and M. J. White, **Acoustic signatures in the cosmic microwave background**, *Astrophys. J.* **471**, 30-51 (1996) doi:10.1086/177951 [arXiv:astro-ph/9602019 [astro-ph]].
- [78] J. L. Bernal, L. Verde and A. G. Riess, **The trouble with H_0** , *JCAP* **10**, 019 (2016) doi:10.1088/1475-7516/2016/10/019 [arXiv:1607.05617 [astro-ph.CO]].
- [79] S. Vagnozzi, **Seven hints that early-time new physics alone is not sufficient to solve the Hubble tension**, *Universe* **9**, no.9, 393 (2023) doi:10.3390/universe9090393 [arXiv:2308.16628 [astro-ph.CO]].
- [80] J. D. Barrow, **Sudden future singularities**, *Class. Quant. Grav.* **21**, L79-L82 (2004) doi:10.1088/0264-9381/21/11/L03 [arXiv:gr-qc/0403084 [gr-qc]].
- [81] E. A. Paraskevas, A. Cam, L. Perivolaropoulos and O. Akarsu, **Transition dynamics in the Λ_s CDM model: Implications for bound cosmic structures**, [arXiv:2402.05908 [astro-ph.CO]].
- [82] O. Akarsu, A. De Felice, E. Di Valentino, S. Kumar, R. C. Nunes, E. Ozulker, J. A. Vazquez and A. Yadav, **Λ_s CDM cosmology from a type-II minimally modified gravity**, [arXiv:2402.07716 [astro-ph.CO]].