ON ELLIPTICAL AND INVERSE ELLIPTICAL WISHART DISTRIBUTIONS: REVIEW, NEW RESULTS, AND APPLICATIONS

Imen Ayadi *, Florent Bouchard†, Frederic Pascal‡

Université Paris-Saclay, CNRS, CentraleSupélec, Laboratoire des Signaux et Systèmes 91190, Gif-sur-Yvette, France

ABSTRACT

This paper deals with matrix-variate distributions, from Wishart to Inverse Elliptical Wishart distributions over the set of symmetric definite positive matrices. Similar to the multivariate scenario, (Inverse) Elliptical Wishart distributions form a vast and general family of distributions, encompassing, for instance, Wishart or *t*-Wishart ones. The first objective of this study is to present a unified overview of Wishart, Inverse Wishart, Elliptical Wishart, and Inverse Elliptical Wishart distributions through their fundamental properties. This involves leveraging the stochastic representation of these distributions to establish key statistical properties of the Normalized Wishart distribution. Subsequently, this enables the computation of expectations, variances, and Kronecker moments for Elliptical Wishart and Inverse Elliptical Wishart distributions. As an illustrative application, the practical utility of these generalized Elliptical Wishart distributions is demonstrated using a real electroencephalographic dataset. This showcases their effectiveness in accurately modeling heterogeneous data

Keywords Covariance matrices · EEG · Elliptical Wishart · Kronecker moments · Matrix-variate statistics · Normalized Wishart · Stochastic representation.

1 Introduction

Covariance matrices have always been at the core of multivariate estimation theory and have recently gained interest in Machine Learning applications as key features. They have proven their relevance in biological signals analysis, radar, and image processing [59], etc. In these applications, statistics over the set of symmetric definite positive matrices (SPD), the native space of non-degenerate covariance matrices, are leveraged. In this perspective, the most classical distribution is the Wishart one [62], known as the distribution of the sample covariance matrix of independent Gaussian vectors. It has various uses in multivariate analysis. For instance, in Bayesian inference, the Wishart distribution is the conjugate prior to the inverse covariance matrix of a multivariate normal random vector [31]. Furthermore, it corresponds to the distribution of the Wishart process introduced by Bru [13] as the solution of a stochastic differential equation with many financial applications, essentially in portfolio theory. Closely linked to the Wishart matrix, the distribution of its inverse, known as Inverse Wishart or Inverted Wishart distribution, plays a major role in multivariate statistical analysis as a conjugate prior to multivariate normal sampling. Several statistical properties of the Wishart and Inverse Wishart distributions were derived, including the probability density function, the characteristic function, the cumulative density function (CDF), the joint probability density function of eigenvalues, the invariance property, different types of moments such that expectation, variance, Kronecker moments (up to small orders for Wishart and for arbitrary order for Inverse Wishart) and invariant moments, etc. Readers are referred to [23] and [30] for detailed review with proofs.

Due to its significance, the Wishart distribution has received several expansions and extensions. The first ones considered singular random matrices as discussed in [19]. Furthermore, a non-central matrix parameter was added

^{*}imen.ayadi@centralesupelec.fr

[†]florent.bouchard@centralesupelec.fr

[†]frederic.pascal@centralesupelec.fr

in [3]; the resulting distribution, called non-central Wishart, can be applied in modelling images corrupted by speckle [58]. Sutradhar and Ali [54] generalized the Wishart distribution by studying the sample covariance matrix of multivariate elliptical vectors, while Teng et al. [56] focused on matrix-variate elliptical models in their work. Multiple Wishart-related distributions were introduced: for instance, one can cite the Laplace-Wishart distribution suggested by Wong and Wang in [63] to establish a Cochran-type theorem; the normal quasi-Wishart distribution by Letac and Massam [33], and the hyper-inverse Wishart introduced by [50] for Bayesian inference in graphical models. Bryc [14] introduced the compound Wishart and q-Wishart, a family of matrices with non-commutative entries that generalize the classical real Wishart matrices. A generalization to Wishart-Laguerre ensembles was suggested by Abul-Magd et al. [1]. Furthermore, Adhikari [2] extended the Wishart distribution for probabilistic structural dynamics. At the same time, it was a prominent tool for real normed division algebras in Dáz-Garca and Gutiérrez-Jáimez's work [18]. Additionally, Munilla and Cantet [41] developed a unique structure for the Wishart distribution to apply to the maternal animal model (MAM) using a conjugated Bayesian approach. In Multiple-Input Multiple-Output (MIMO) systems, Kumar et al. [49] employed the eigenvalue distribution of the sum of Wishart random variables, whereas Lopez-Martinez et al. [36] applied the complex Wishart distribution. Bekker et al. [10] introduced the Hypergeometric Elliptical Wishart as a combination of a generalized matrix-variate hypergeometric distribution and the Elliptical Wishart distribution.

Among these various generalizations, this paper focuses on pursuing the work of Teng *et al.* [56] about the generalized Wishart distribution under elliptical models, also called Elliptical Wishart. The motivation behind our interest in this family of distributions over SPDs is its great potential in modelling intrinsic non-Gaussianity in signal processing problems, wherein extremely aberrant recordings and even mislabelling are frequent. Such issues made classical multivariate robust techniques [39] inefficient and thus urged us to consider the outliers at the scale of the covariance matrices and not only at the scale of a single measurement [7]. Accordingly, we leverage robust matrix-variate statistics by generalizing the Wishart modeling into the Elliptical Wishart one. Although they were introduced a long time ago, only a few statistical properties of Elliptical Wishart and Inverse Elliptical Wishart distributions have already been derived, including the probability density function, the characteristic function (only for Elliptical Wishart), and other properties involving only the statistics of eigenvalues such that the joint probability density of eigenvalues, moments of the trace and the determinant [9], the expectation of zonal polynomials and cumulative density function [16].

In this paper, we suggest studying some statistical properties of the Elliptical Wishart and Inverse Elliptical Wishart that are lacking in the literature. More precisely, our main theoretical contributions consist in computing the expectation, variance, and Kronecker moments up to any order of these distributions. For that, the basic tool is the stochastic representation of a random matrix drawn from an Elliptical Wishart distribution and the already-derived results about Wishart and Inverse Wishart distributions. Our contributions are not limited to Elliptical and Inverse Elliptical Wishart distributions: one major contribution is to derive the Kronecker moments of a Wishart matrix up to any order. Prior to this paper, it was only known up to order four [53]. From a practical point of view, we first propose a very computationally efficient way to generate samples from Elliptical and Inverse Elliptical Wishart distributions. Then, the interest of Elliptical Wishart distributions is emphasized on electroencephalography (EEG), which involves recording brain signals, *e.g.*, for medical purposes or assisting people with movement impediments via a brain-computer interface (BCI). Fitting real noisy EEG measurements illustrate the potential of an Elliptical Wishart distribution. For the data we consider, the so-called *t*-Wishart distribution clearly outperforms the Wishart distribution. The code to reproduce numerical experiments is provided in https://github.com/IA3005/jmva_paper_code.git.

The article is organized as follows. Section 2 provides a brief reminder about the basic statistical tools needed to manipulate random vectors and matrices and a background about multivariate elliptical distributions. Section 3 reviews the Wishart and Inverse Wishart distributions and their elliptical extensions. Then, Section 4 derives the paper's main contributions, namely the flagship theorem of the stochastic representation. This theorem is the basis for studying the Normalized Wishart distribution in Section 4.1 and further deriving the expectation, the variance, and the Kronecker moments of Elliptical Wishart and Inverse Elliptical Wishart, respectively, in Section 4.2 and Section 4.3. Some numerical experiments in Section 5 validate the merit of Elliptical Wishart distributions, particularly *t*-Wishart, for BCIs by fitting real EEG signals. Finally, concluding remarks and perspectives are drawn in Section 6.

2 Preliminaries

Before turning to the main contributions of this work, this section aims at introducing the main notations used in the paper as well as presenting existing results on the various distributions of interest.

2.1 Notations

Vectors (respectively matrices) are denoted by bold-faced lowercase letters (respectively uppercase letters). \otimes , $\operatorname{tr}(\cdot)$, $|\cdot|$ and $|\cdot|_2$ correspond to the Kronecker product, trace, determinant, and ℓ_2 norm, respectively. A^{\perp} denotes the transpose of the matrix A. vec(·) is the operator that transforms a matrix $p \times n$ into a vector of length pn, using columnwise concatenation. Moreover, \hat{I} is the identity matrix, and 0 is the matrix of zeros with appropriate dimensions, while K is the commutation matrix, which transforms vec(A) into $\text{vec}(A^{\top})$. Let \mathcal{S}_d and \mathcal{S}_d^{++} denote the set of $d \times d$ real symmetric matrices and the set of $d \times d$ real symmetric definite positive matrices, respectively. We define the unit sphere on $\mathbb{R}^{p \times n}$ as $\mathcal{C}_{p,n} = \{ \boldsymbol{U} \in \mathbb{R}^{p \times n} | \|\boldsymbol{U}\|_2 = 1 \}$. \sim means "distributed as" while $\stackrel{d}{=}$ stands for "shares the same distribution as". $E[\cdot]$ denotes the statistical expectation. $\Gamma(\cdot)$ is the Gamma function and $\Gamma_p(\cdot)$ is the multivariate Gamma function of dimension p. For $\kappa = (k_1, \dots, k_p) \in \mathbb{N}^p$, we denote by $(\cdot)_{\kappa}$ the partitional shifted factorial defined for $a \in \mathbb{R}$ at κ as $(a)_{\kappa} = \prod_{j=1}^{p} \left(a - \frac{j-1}{2}\right)_{k_j}$ where $(a)_{k_j} = a(a+1) \dots (a+k_j-1)$, with the convention $(a)_0 = 1.$

2.2 Statistical properties reminders

This paper focuses on distributions of random matrices over $\mathbb{R}^{p \times n}$. This section provides basic statistical notions and notations needed to manipulate these matrix-variate distributions. As for any distribution, we are interested in the probability density functions, the characteristic functions, and the moments.

The probability density function (PDF) of a random matrix can be defined like the ones of a random uni-variate variable or random vector. If $X \in \mathbb{R}^{p \times n}$ is a random matrix, the matrix-variate PDF of X, denoted as f_X , is a mapping from $\mathbb{R}^{p \times n}$ to \mathbb{R}_+ with some parameters describing some characteristics of that distribution. To be a proper PDF, f_X must

$$\int_{\mathbb{R}^{p\times n}} f_{\mathbf{X}}(\mathbf{X}) d\mathbf{X} = 1,\tag{1}$$

where $dX = \prod_{k \neq l} dX_{k\ell}$ denotes the Lebesgue measure on $\mathbb{R}^{p \times n}$.

To study a distribution, we systematically derive its moments as they are paramount in characterizing the mean, dispersion around the mean, skewness, and tail's heaviness. The definition of moments in multivariate statistics was easily extended to the matrix variate case. In particular, the first moment of a random matrix X is its expectation E[X], which is the $p \times n$ matrix whose elements are

$$E[\mathbf{X}]_{k\ell} = E[\mathbf{X}_{k\ell}] = \int_{\mathbb{D}_{p\times n}} \mathbf{X}_{k\ell} f_{\mathbf{X}}(\mathbf{X}) d\mathbf{X}.$$
 (2)

The second moment is $E[\text{vec}(X) \text{vec}(X)^{\top}]$. Hence, the variance of the random matrix X, denoted var[X], is given by

$$\operatorname{var}[\boldsymbol{X}] = E[\operatorname{vec}(\boldsymbol{X})\operatorname{vec}(\boldsymbol{X})^{\top}] - \operatorname{vec}(E[\boldsymbol{X}])\operatorname{vec}(E[\boldsymbol{X}])^{\top}. \tag{3}$$

More generally, it is possible to define the k^{th} moment (if it exists) for random matrices to improve our understanding of the distribution shape. Its definition can vary from one author to another, but it is more likely given in literature by $E[X_{j_0j_1}...X_{j_{k-1}j_k}]$ [30], the expectation of the product of k elements of X, which is somehow equivalent to the expectation of a Kronecker product of order k. For a random matrix $S \in S_p$, the kth Kronecker moment corresponds to the expectation of k^{th} Kronecker product, i.e., $E[\otimes^k \mathbf{S}] = E[\mathbf{S} \otimes \cdots \otimes \mathbf{S}] \in \mathbb{R}^{p^k \times p^k}$. Thanks to these moments, it is possible to compute other types of moments using some vectorization and Kronecker-product tricks. Examples of such moments are $E[S^k]$ and $E[tr(S)^q S^r]$ where $q, r \in \mathbb{N}$ [60]. One of the available tools to compute Kronecker moments is the differentiation of the characteristic function, which is defined for a random matrix X as the Fourier transform of f_X . The characteristic function $\Psi_X : \mathbb{R}^{p \times n} \to \mathbb{R}$ is defined as

$$\Psi_{\boldsymbol{X}}(\boldsymbol{Z}) = E[\exp(i\operatorname{tr}(\boldsymbol{X}^{\top}\boldsymbol{Z}))] = \int_{\mathbb{R}^{p \times n}} \exp(i\operatorname{tr}(\boldsymbol{X}^{\top}\boldsymbol{Z}))f_{\boldsymbol{X}}(\boldsymbol{X})d\boldsymbol{X}, \tag{4}$$

where $i^2 = -1$. If n = 1, we regain the definition of the characteristic function for the multivariate case. For symmetric random matrices, Kronecker moments $E[\otimes^k S]$ are related to the k^{th} derivative⁴ of the characteristic function Ψ_S with respect to $Z \in \mathcal{S}_p$ at 0 (if it exists):

$$\left. \frac{\partial^k \Psi_{\boldsymbol{S}}(\boldsymbol{Z})}{\partial \boldsymbol{Z}^k} \right|_{\boldsymbol{Z} \to \boldsymbol{0}} = i^k E[\otimes^k \boldsymbol{S}].$$
 (5)

 $[\]frac{\partial^k \Psi_{\boldsymbol{S}}(\boldsymbol{Z})}{\partial \boldsymbol{Z}^k}\bigg|_{\boldsymbol{Z} \to 0} = i^k E[\otimes^k \boldsymbol{S}]. \tag{5}$ $\frac{^4 \text{Let } \boldsymbol{X} \in \mathbb{R}^{p \times p} \text{ symmetric and } \boldsymbol{Y} \in \mathbb{R}^{q \times r}. \text{ We choose the Kroneckerian arrangement for defining our matrix derivative}$ $\frac{\partial \boldsymbol{Y}}{\partial \boldsymbol{X}} = \sum_{m,j,k,\ell} \alpha_{k\ell} \frac{\partial \boldsymbol{Y}_{mj}}{\partial \boldsymbol{X}_{k\ell}} (\boldsymbol{s}_m \boldsymbol{t}_j^\top) \otimes (\boldsymbol{e}_k \boldsymbol{e}_\ell^\top) \text{ with } \alpha_{k\ell} = 1 \text{ if } k = \ell, \text{ otherwise } 1/2; \text{ where } \{\boldsymbol{s}_m\}_{m=1}^q, \{\boldsymbol{t}_j\}_{j=1}^r \text{ and } \{\boldsymbol{e}_k\}_{k=1}^p \text{ denote the canonical bases of } \mathbb{R}^q, \mathbb{R}^r \text{ and } \mathbb{R}^p, \text{ respectively. For more details, see [43, 60]}$

Remark 1. We highlight that the second Kronecker moment, $E[S \otimes S]$, is different from the second moment, $E[\text{vec}(S) \text{vec}(S)^{\top}]$. However, they can be linked to each other by:

$$\operatorname{vec}(E[\boldsymbol{S} \otimes \boldsymbol{S}]) = (\boldsymbol{I}_p \otimes \boldsymbol{K}_{p,p} \otimes \boldsymbol{I}_p) \operatorname{vec}(E[\operatorname{vec}(\boldsymbol{S}) \operatorname{vec}(\boldsymbol{S})^{\top}]), \tag{6}$$

using the formula $\text{vec}(\boldsymbol{A} \otimes \boldsymbol{B}) = (\boldsymbol{I}_{n_1} \otimes \boldsymbol{K}_{n_2,p_1} \otimes \boldsymbol{I}_{p_2}) (\text{vec}(\boldsymbol{A}) \otimes \text{vec}(\boldsymbol{B}))$ for any $\boldsymbol{A} \in \mathbb{R}^{p_1 \times n_1}$ and $\boldsymbol{B} \in \mathbb{R}^{p_2 \times n_2}$ [44].

2.3 Multivariate and matrix-variate elliptical distributions

This subsection provides a short overview of the multivariate and matrix-variate elliptical distributions. These are leveraged to define and study the Elliptical Wishart and Inverse Wishart distributions in the following sections. We first provide a definition of the multivariate elliptical distribution in Definition 1. For a full review, the reader is referred to [46].

Definition 1 (Multivariate elliptical distribution). A random vector $x \in \mathbb{R}^d$ is said to have a centered elliptical distribution, denoted as $\mathcal{E}_d(\mathbf{0}, \mathbf{\Sigma}, h_d)$ if its probability density function has the form

$$f_{\mathbf{x}}(\mathbf{x}) = |\mathbf{\Sigma}|^{-1/2} h_d(\mathbf{x}^{\mathsf{T}} \mathbf{\Sigma}^{-1} \mathbf{x}), \tag{7}$$

where $\Sigma \in \mathcal{S}_d^{++}$ is the covariance matrix and $h_d: \mathbb{R}_+ \to \mathbb{R}$ is the density generator function of the distribution. To ensure that h_d is a proper density generator function, one must have $\frac{\pi^{d/2}}{\Gamma(d/2)} \int_0^{+\infty} h_d(t) t^{\frac{d}{2}-1} dt = 1$.

The elliptical model can be naturally extended to matrices via the vectorization operator. This family of distributions plays a key role in the definition of Elliptical Wishart distributions. The matrix-variate elliptical distribution is defined in Definition 2. For a thorough review, see [24].

Definition 2 (Matrix-variate elliptical distributions). A random matrix $X \in \mathbb{R}^{p \times n}$ is said to be drawn from a centered elliptical distribution, denoted as $\mathcal{E}_{p,n}(\mathbf{0}, \mathbf{\Sigma}, \mathbf{\Psi}, h_{np})$ if $\operatorname{vec}(X) \sim \mathcal{E}_{np}(\mathbf{0}, \mathbf{\Psi} \otimes \mathbf{\Sigma}, h_{np})$ wherein $\mathbf{\Sigma} \in \mathcal{S}_p^{++}$ is the covariance matrix controlling the correlation between X's rows and $\mathbf{\Psi} \in \mathcal{S}_n^{++}$ is the covariance matrix controlling the correlation between X's columns.

Multivariate and matrix-variate elliptical distributions were well studied in the literature. In particular, their statistical properties, such as the expectation and the variance, were derived in both cases; see e.g., [46, 24]. In this paper, we need two tools from the multivariate case: the stochastic representation and moments of an elliptical vector's squared ℓ_2 norm. They indeed play a major role in Section 4. The stochastic representation of $\boldsymbol{x} \sim \mathcal{E}_d(\mathbf{0}, \boldsymbol{\Sigma}, h_d)$ is

$$\boldsymbol{x} \stackrel{d}{=} \sqrt{\mathcal{Q}} \boldsymbol{\Sigma}^{1/2} \boldsymbol{u}, \tag{8}$$

where $\mathcal{Q} = \|\mathbf{\Sigma}^{-1/2}\boldsymbol{x}\|_2^2$ is a non-negative random variable and $\boldsymbol{u} \in \mathbb{R}^d$ is uniformly distributed over the unit sphere $\mathcal{C}_d := \{\boldsymbol{v} \in \mathbb{R}^d | \|\boldsymbol{v}\|_2 = 1\}$. Moreover, \mathcal{Q} and \boldsymbol{u} are independent. The variable \mathcal{Q} is called the second-order modular of \boldsymbol{x} and its probability density function is $t \mapsto \frac{\pi^{d/2}}{\Gamma(d/2)}h_d(t)t^{\frac{d}{2}-1}$. Furthermore, when $\boldsymbol{x} \sim \mathcal{E}_d(0, \boldsymbol{I}_d, h_d)$, the expectation $m_k^{(h_d)}$ of $\|\boldsymbol{x}\|_2^{2k}$ is given by

$$m_k^{(h_d)} = E[\|\boldsymbol{x}\|_2^{2k}] = E[\mathcal{Q}^k] = \frac{\pi^{d/2}}{\Gamma(d/2)} \int_0^{+\infty} h_d(t) t^{\frac{d}{2} + k - 1} dt, \tag{9}$$

for $k \in \mathbb{Z}$. However, this moment does not necessarily exist for all orders. In particular, the convergence of the integral should be checked at 0 and $+\infty$.

3 Review: Wishart, Inverse Wishart, and elliptical extensions

This section aims to provide a detailed review of the Wishart and Inverse Wishart distributions, along with their elliptical extensions. In particular, their definitions, probability density functions, and some of their existing statistical properties – mainly the characteristic functions, the first and second moments, and the Kronecker moments. Notice that in this paper, only non-degenerate distributions are considered, *i.e.*, it is always assumed that the number of samples n is greater than the number of features p.

3.1 Wishart distribution

First, we introduce the Wishart distribution; see *e.g.*, [11, 23]. Wishart random matrices are characterized in Definition 3.

Definition 3 (Wishart distribution). Let n independent and identically distributed (i.i.d.) vectors $\{x_j\}_{1 \leq j \leq n}$ in \mathbb{R}^p drawn from the multivariate Gaussian distribution $\mathcal{N}_p(\mathbf{0}, \mathbf{\Sigma})$ and $\mathbf{S} = \sum_{j=1}^n \mathbf{x}_j \mathbf{x}_j^{\top}$. Then, \mathbf{S} is said to have the Wishart distribution with n degrees of freedom and a center matrix $\mathbf{\Sigma}$, denoted as $\mathcal{W}(n, \mathbf{\Sigma})$.

Remark 2. Notice that if we consider $X \in \mathbb{R}^{p \times n}$ the matrix whose columns are the x_j 's, then X has the matrix-variate Gaussian distribution $\mathcal{N}_{p,n}(\mathbf{0}, \Sigma, I_n)$ [11]. We can write $S = XX^{\top}$, which yields an alternative way to define the Wishart distribution [23]. Such a matrix formulation is at the core of elliptical extensions.

The probability density function of $S \sim \mathcal{W}(n, \Sigma)$ is given by

$$f_{\mathbf{S}}(\mathbf{S}) = \frac{1}{2^{np/2} \Gamma_n(n/2)} |\mathbf{\Sigma}|^{-n/2} |\mathbf{S}|^{\frac{n-p-1}{2}} \exp\left(-\frac{1}{2} \operatorname{tr}(\mathbf{\Sigma}^{-1} \mathbf{S})\right).$$
(10)

Various statistical properties of the Wishart distribution are derived thanks to the independence of the x_j 's and the statistical properties of the multivariate Gaussian distribution. For instance, from [11], the characteristic function of $S \sim W(n, \Sigma)$ is, for $Z \in \mathcal{S}_p$,

$$\Psi_{S}(Z) = |I_{p} - 2iZ\Sigma|^{-\frac{n}{2}}.$$
(11)

The first and second moments of the Wishart distribution are well known [11]. Indeed, its expectation and its variance are

$$E[S] = n\Sigma, \tag{12}$$

and

$$var[S] = n(I_{p^2} + K_{p,p})(\Sigma \otimes \Sigma).$$
(13)

The Kronecker moments were derived in [38, 53] up to order 4 and were obtained through (5), *i.e.*, the differentiation of the characteristic function (11) of the Wishart distribution. In particular, the Kronecker moments of orders 2 and 3 are

$$E[S \otimes S] = n^{2}(\Sigma \otimes \Sigma) + n\left(K_{p,p}(\Sigma \otimes \Sigma) + \operatorname{vec}(\Sigma)\operatorname{vec}(\Sigma)^{\top}\right), \tag{14}$$

$$E[\otimes^{3}S] = n^{3} \otimes^{3} \Sigma + n^{2} \left(P + APA + BAPAB + (A + B + BAB) \otimes^{3} \Sigma \right) + n(CAP + PAC + BAPA + AVAB + (AB + BA) \otimes^{3} \Sigma),$$

$$(15)$$

where $P = (\text{vec}(\Sigma) \text{ vec}(\Sigma)^{\top}) \otimes \Sigma$, $A = I_p \otimes K_{p,p}$ and $B = K_{p,p} \otimes I_p$ and $C = I_{p^3} + B$. The detailed expression for order 4 can be found in [53]. To the best of our knowledge, no explicit formula was derived for Kronecker moments of arbitrary order for the Wishart distributions. However, closely linked to Kronecker moments, $E[S_{12}S_{34}\dots S_{2k-1,2k}]$ were shown to be expressed as weighted generating functions of graphs associated with the Wishart distribution in [45].

In the literature, several other types of moments of random matrices drawn from the Wishart distribution can also be found. For instance, explicit formulas for the moments of arbitrary polynomials in the entries of a Wishart-distributed matrix are obtained in [37]. These rely on MacMahon's master theorem, differentiation of moment generating function, and the representation theory for the symmetric group. In [34, 12], invariant moments of the form E[Q(S)], where Q(S) is a polynomial that only depends on the eigenvalues of S, are computed. Implementations of the formulas derived in [34] are provided in SAGE software [48]. Moreover, [21] derived $E[\operatorname{tr}(S^q)S^r]$ by leveraging the Wishart process and Itô calculus. Recently, moments $E[S^k]$ were computed recursively, with explicit formulas up to order 4 [25]. In the meantime, [42] gave explicit expressions for $E[(\operatorname{tr}(S^q))^r(\operatorname{tr}(S^m))^j]$ for $m+j+q+r\leq 5$. Finally, one of the invariant moments that has gained a lot of interest in literature is the expectation of zonal polynomials of S^5 . We recall that a zonal polynomial of S at any order $\kappa \in \mathbb{N}^p$, denoted as $C_{\kappa}(S)$, where $\kappa = (k_1, \ldots, k_p)$ and $k_1 \geq \cdots \geq k_p$, is a symmetric, homogeneous polynomial of degree $k := \sum_{j=1}^p k_j$ of the eigenvalues of S; see e.g., [40] for a detailed definition. It is known that, for any deterministic $Z \in S_p$, [29]

$$E[C_{\kappa}(\mathbf{Z}\mathbf{S})] = 2^{k} \left(\frac{n}{2}\right)_{\kappa} C_{\kappa}(\mathbf{Z}\mathbf{\Sigma}). \tag{16}$$

⁵In particular, it entitles us with an alternative way to regain the characteristic function of the Elliptical Wishart distribution.

3.2 Inverse Wishart distribution

From the Wishart distribution, it is possible to define another major matrix-variate distribution over S_p^{++} : the Inverse Wishart distribution. It corresponds to the distribution of the inverse of a random matrix drawn from the Wishart distribution. A formal definition is provided in Definition 4.

Definition 4. We say that S has the Inverse Wishart distribution with n degrees of freedom and covariance matrix Σ , denoted $W^{-1}(n, \Sigma)$, if $S^{-1} \sim W(n, \Sigma^{-1})$.

Using the inverse substitution in the integral in (10), the probability density function of $W^{-1}(n, \Sigma)$ is given by

$$f_{S}(S) = \frac{1}{2^{np/2} \Gamma_{p}(n/2)} |\Sigma|^{n/2} |S|^{-\frac{n+p+1}{2}} \exp\left(-\frac{1}{2} tr(\Sigma S^{-1})\right).$$
(17)

Unlike the Wishart distribution, no closed-form expression is found for the characteristic function of the Inverse Wishart distribution. The first and second moments need additional assumptions on n and p to exist [23]. When n > p+1, the expectation of $S \sim W^{-1}(n, \Sigma)$ is

$$E[S] = \frac{1}{n - p - 1} \Sigma,\tag{18}$$

and when n > p + 3, the variance exists and is given by

$$\operatorname{var}[\boldsymbol{S}] = \frac{1}{(n-p)(n-p-1)(n-p-3)} \left[(\boldsymbol{I}_{p^2} + \boldsymbol{K}_{p,p})(\boldsymbol{\Sigma} \otimes \boldsymbol{\Sigma}) + \frac{2}{n-p-1} \operatorname{vec}(\boldsymbol{\Sigma}) \operatorname{vec}(\boldsymbol{\Sigma})^{\top} \right].$$
(19)

Interestingly, in contrast to Wishart, the Kronecker moments of the Inverse Wishart random matrix are well known up to any order [60] thanks to Stokes' theorem. Given k such that n > p + 2k + 1, the (k + 1)th Kronecker moment is given by

$$\operatorname{vec}(E[\otimes^{k+1} \mathbf{S}]) = \frac{1}{n-p-1} \prod_{i=0}^{k-1} (\mathbf{I}_{p^{2i}} \otimes \mathbf{A}(k-i))(\otimes^{k+1} \operatorname{vec}(\mathbf{\Sigma})), \tag{20}$$

where $\boldsymbol{A}(k) = (n-p-1)\boldsymbol{I}_{p^{2k}} - \sum_{j=0}^{k-1}(\boldsymbol{P}_1(j) + \boldsymbol{P}_2(j))$ with $\boldsymbol{P}_1(j) = \left(\otimes^2(\boldsymbol{I}_{p^j}\otimes\boldsymbol{K}_{p,p^{k-1-j}}\otimes\boldsymbol{I}_p)\right)(\boldsymbol{I}_{p^k}\otimes\boldsymbol{K}_{p,p^k}\otimes\boldsymbol{I}_p)(\boldsymbol{I}_{p^{k-1}} + \boldsymbol{K}_{p^{k+2},p})$ and $\boldsymbol{P}_2(j) = \left(\otimes^2(\boldsymbol{I}_{p^j}\otimes\boldsymbol{K}_{p,p^{k-1-j}}\otimes\boldsymbol{I}_p)\right)(\boldsymbol{I}_{p^{2k}} + \boldsymbol{K}_{p,p})$. Further notice that moments $E[\otimes^k \operatorname{vec}(\boldsymbol{S})]$, which are different from our definition of Kronecker moments, were derived in [61] with the help of a factorization theorem, moments for normally distributed variables, and inverse moments for Chi-squared variables.

As for the Wishart distribution, various moments were derived for the Inverse Wishart distribution, such as the expectation of zonal polynomials; see [29]. This allowed to derive the expectation of powers of traces, i.e., $\operatorname{tr}(S)^k$, powers of determinant, i.e., $|S|^k$, and powers of sums, i.e., $\operatorname{tr}(S^k)$ with $S \sim \mathcal{W}^{-1}(n, \Sigma)$ [26]. In addition, [28] computed moments of $S^r P(S)$ where P is a power-sum symmetric function. Finally, [42] obtained $E[(\operatorname{tr}(S^q))^r(\operatorname{tr}(S^i))^j]$ for $i+j+q+r \leq 5$.

3.3 Elliptical extensions

The Wishart distribution can be generalized in various ways. In this work, we are interested in Elliptical Wishart distributions. They generalize the Wishart distribution the same way multivariate elliptical distributions generalize the Gaussian one. These are formally defined in Definition 5.

Definition 5. We say that a random matrix S follows the Elliptical Wishart distribution $\mathcal{EW}(n, \Sigma, h_{np})$, with n degrees of freedom, center Σ , and density generator $h_{np} : \mathbb{R}_+ \to \mathbb{R}$, if $S = XX^{\top}$ such that $X \sim \mathcal{E}_{p,n}(0, \Sigma, I_n, h_{np})$.

Consequently, the same density generators h_{np} as in the multivariate case can be considered to define Elliptical Wishart distributions. For instance, one can leverage density generators from the Student t-, generalized Gaussian, Kotz, Bessel models, etc. Examples are provided in Section 4.4. The probability density function of $\mathcal{EW}(n, \Sigma, h_{np})$, which can be obtained through some integration substitution tricks [4], is

$$f_{\mathbf{S}}(\mathbf{S}) = \frac{\pi^{np/2}}{\Gamma_p(n/2)} |\mathbf{\Sigma}|^{-\frac{n}{2}} |\mathbf{S}|^{\frac{n-p-1}{2}} h_{np} \left(\operatorname{tr}(\mathbf{\Sigma}^{-1} \mathbf{S}) \right).$$
 (21)

Few theoretical results about Elliptical Wishart distributions are already known. One of the few statistical properties that were derived is the characteristic function of $S \sim \mathcal{EW}(n, \Sigma, h_{np})$ when h_{np} admits a Taylor series expansion [9]. It is given, for any $Z \in \mathcal{S}_n$, by

$$\Psi_{\mathbf{S}}(\mathbf{Z}) = \pi^{np/2} \sum_{k=0}^{+\infty} \frac{i^k}{k!} \left(\frac{\int_0^{+\infty} h_{np}(r) r^{\frac{np}{2} + k - 1} dr}{\Gamma\left(\frac{np}{2} + k\right)} \right) \sum_{|\kappa| = k} \left(\frac{n}{2}\right)_{\kappa} C_{\kappa}(\mathbf{Z}\boldsymbol{\Sigma}).$$
(22)

In Section 4.2, a more straightforward way to derive it is provided. Neither the expectation nor the variance of Elliptical Wishart distributions have been computed in the literature. One of the significant contributions of this work is to derive them in Section 4.2. However, some other statistical properties have been established, such as the joint density function of eigenvalues⁶, $E[\operatorname{tr}(S)^k]$ and $E[|S|^k]$; see *e.g.*, [9, 18]. The expectation of zonal polynomials and the cumulative function have also been derived [16]. Notice that, from a more practical point of view, the information geometry along with estimation and classification methods based on Elliptical Wishart distributions are proposed in [6, 7].

Akin to the definition of the Inverse Wishart distributions, it is possible to define the Inverse Elliptical Wishart definitions. A proper definition is provided in Definition 6.

Definition 6. We say that S has the Inverse Elliptical Wishart distribution $\mathcal{EW}^{-1}(n, \Sigma, h_{np})$ with n degrees of freedom, center Σ and density generator $h_{np}: \mathbb{R}_+ \to \mathbb{R}$, if $S^{-1} \sim \mathcal{EW}(n, \Sigma^{-1}, h_{np})$.

We would highlight that Definition 6 holds as long as $n \geq p$ and Σ is non-singular. In fact, in this case, $X \sim \mathcal{E}_{p,n}(\mathbf{0}, \Sigma, \mathbf{I}_n, h_{np})$ has a full rank with probability 1. Consequently, $S = XX^{\top}$ is non-singular with probability 1. From [20], the probability density function of the Inverse Elliptical Wishart distribution is

$$f_{\mathbf{S}}(\mathbf{S}) = \frac{\pi^{np/2}}{\Gamma_p(n/2)} |\mathbf{\Sigma}|^{-n/2} |\mathbf{S}|^{-\frac{n+p+1}{2}} h_{np} \left(\operatorname{tr}(\mathbf{\Sigma} \mathbf{S}^{-1}) \right).$$
 (23)

Remark 3. Considering that the Inverse Elliptical Wishart distribution is the extension of the Inverse Wishart distribution that consists in replacing the Gaussian density generator function in (17) with a general density generator h_{np} , it is obvious that the "Inverse Elliptical Wishart" is the same thing as the "Elliptical Inverse Wishart".

For the Inverse Elliptical Wishart distribution, apart from deriving the probability density function (23) of $S \sim \mathcal{EW}^{-1}(n, \Sigma, h_{np})$, no other statistical properties have been derived yet. In this paper, more precisely in Section 4.3, the expectation, variance and Kronecker moments of an Inverse-Elliptical-Wishart-distributed matrix are derived.

To conclude this section, a summary of all the known statistical properties prior to this paper of the Wishart, Inverse Wishart, Elliptical Wishart and Inverse Elliptical Wishart distributions is provided in Table 1. With the present paper, Table 1 changes quite a lot. Indeed, only the characteristic functions of the Inverse Wishart and Inverse Elliptical Wishart distributions remain unknown. Hence, the expectation, variance, and Kronecker moments are derived from any order of Elliptical Wishart and Inverse Elliptical Wishart distributions. In particular, the Kronecker moments of the Wishart distribution, which were only known up to order four, are derived up to any order.

Distribution	PDF	characteristic function	Expectation	Variance	Kronecker moments
Wishart $\mathcal{W}(n, \mathbf{\Sigma})$	(10)	(11)	(12)	(13)	up to order 4, [53]
Inverse Wishart $\mathcal{W}^{-1}(n, \Sigma)$	(17)	No closed form	(18)	(19)	(20)
Elliptical Wishart $\mathcal{EW}(n, \mathbf{\Sigma}, h_{np})$	(21)	(22)	Not derived yet	Not derived yet	Not derived yet
Inverse Elliptical Wishart $\mathcal{EW}^{-1}(n, \Sigma, h_{np})$	(23)	Not derived yet	Not derived yet	Not derived yet	Not derived yet

Table 1: Recapitulating table of the review about Wishart, Inverse Wishart, Elliptical Wishart, and Inverse Elliptical Wishart distributions

⁶As a matter of fact, [40, Theorem 3.2.17.] provides a formula to compute the joint PDF of eigenvalues of a random SPD matrix knowing its PDF.

4 Main contributions: Elliptical Wishart and Inverse Elliptical Wishart statistical properties

This section contains the main methodological contributions of this paper. In particular, it provides the expectation and variance of the Elliptical Wishart and Inverse Elliptical Wishart distributions. Concerning the characteristic function of the Elliptical Wishart distribution, an alternative (and simpler) proof is obtained as compared to the one given in [9] while the characteristic function of the Inverse Elliptical Wishart remains without a closed form. All these new results stem from the stochastic representation of an Elliptical-Wishart-distributed matrix, given in Theorem 1. As for the multivariate elliptical case (see Section 3), this representation relies on the second-order modular, Q, and a random matrix uniformly distributed on the unit sphere.

Theorem 1 (Stochastic Representation). Let $S \sim \mathcal{EW}(n, \Sigma, h_{np})$ where $\Sigma \in \mathcal{S}_p^{++}$. There exist a non negative random variable Q and a random matrix $U \in \mathbb{R}^{p \times n}$ such that

$$S \stackrel{d}{=} Q \Sigma^{1/2} (UU^{\top}) \Sigma^{1/2}, \tag{24}$$

and

- U is uniformly distributed on the unit sphere $C_{p,n}$.
- The probability density function of Q is $t \mapsto \frac{\pi^{np/2}}{\Gamma(np/2)} h_{np}(t) t^{\frac{np}{2}-1}$.
- *U* and *Q* are independent.

Proof. By definition, there exists $X \in \mathbb{R}^{p \times n}$ such that $S = XX^{\top}$ and $X \sim \mathcal{E}_{p,n}(0, \Sigma, I_n, h_{np})$. Recall that $\operatorname{vec}(X) \sim \mathcal{E}_{pn}(0, I_n \otimes \Sigma, h_{np})$. Hence, $\operatorname{vec}(X) \stackrel{d}{=} \sqrt{\mathcal{Q}}(I_n \otimes \Sigma)^{1/2} \boldsymbol{u}$, where \mathcal{Q} is a non negative random variable and \boldsymbol{u} is uniformly distributed on the np-dimensional unit sphere. Moreover, \boldsymbol{u} and \mathcal{Q} are independent. The density function of \mathcal{Q} , which can be found in [15], is $t \mapsto \frac{\pi^{np/2}}{\Gamma(np/2)}h_{np}(t)t^{\frac{np}{2}-1}$. Let $\boldsymbol{U} \in \mathbb{R}^{p \times n}$ such that $\operatorname{vec}(\boldsymbol{U}) = \boldsymbol{u}$. It is easy to check that $\operatorname{tr}(\boldsymbol{U}\boldsymbol{U}^{\top}) = \boldsymbol{u}^{\top}\boldsymbol{u} = 1$. Thus, $\boldsymbol{U} \in \mathcal{C}_{p,n}$. Furthermore, $(\boldsymbol{I}_n \otimes \Sigma)^{1/2} = \boldsymbol{I}_n \otimes \Sigma^{1/2}$. It follows that $\operatorname{vec}(\boldsymbol{X}) \stackrel{d}{=} \sqrt{\mathcal{Q}}(\boldsymbol{I}_n \otimes \Sigma^{1/2})\operatorname{vec}(\boldsymbol{U}) = \sqrt{\mathcal{Q}}\operatorname{vec}(\Sigma^{1/2}\boldsymbol{U})$. Consequently, $\boldsymbol{X} \stackrel{d}{=} \sqrt{\mathcal{Q}}\Sigma^{1/2}\boldsymbol{U}$ and $\boldsymbol{S} \stackrel{d}{=} \mathcal{Q}\Sigma^{1/2}(\boldsymbol{U}\boldsymbol{U}^{\top})\Sigma^{1/2}$. This concludes the proof.

From the stochastic representation of Theorem 1, the statistical properties of \mathcal{Q} are well-known. Moreover, one can recognize that the random matrix UU^{\top} follows the Normalized Wishart distribution [55]. To obtain the statistical properties of the Elliptical Wishart and Inverse Elliptical Wishart distributions, we first need to obtain those of UU^{\top} . This is achieved in Section 4.1. From there, the main results for the Elliptical Wishart and Inverse Elliptical Wishart distributions are given in Sections 4.2 and 4.3, respectively.

Remark 4. All distributions considered here (Elliptical Wishart and Inverse Elliptical Wishart with I_p as covariance matrix and Normalized Wishart) are rotationally-invariant, i.e., $S \stackrel{d}{=} HSH^{\top}$ for any orthogonal matrix $H \in \mathbb{R}^{p \times p}$. For such distributions, [11, Proposition 13.2] shows that $E[S] = \alpha I_p$ and $\text{var}[S] = \sigma_1(I_{p^2} + K_{p,p}) + \sigma_2 \text{vec}(I_p) \text{vec}(I_p)^{\top}$, where $\alpha \in \mathbb{R}$, $\sigma_1 \geq 0$ and $\sigma_2 \geq \frac{-2\sigma_1}{p}$. This proposition thus provides the general form of the expectation and variance of a rotationally invariant distribution. For the distributions we consider, while it enables us to obtain the expectation easily, it does not allow us to identify σ_1 and σ_2 straightforwardly.

4.1 Normalized Wishart distribution

In this section, the statistical properties of UU^{\top} in the stochastic representation of Theorem 1 are studied. One can observe that, given $S \sim W(n, I_p)^7$, $UU^{\top} \stackrel{d}{=} \frac{S}{\operatorname{tr}(S)}$. Thus, as mentioned above, UU^{\top} follows the so-called Normalized Wishart distribution [55]. In the following, let $V = UU^{\top}$. After properly defining the Normalized Wishart distribution in Definition 7, first and second-order moments of V and V^{-1} are derived in Proposition 1.

Definition 7 (Normalized Wishart distribution). Let $S \sim \mathcal{W}(n, I_p)$ a Wishart matrix. The corresponding Normalized Wishart matrix is given by

$$V = \frac{S}{\operatorname{tr}(S)}.$$
 (25)

⁷As a matter of fact, it works for any Elliptical Wishart distribution, i.e., $S \sim \mathcal{EW}(n, I_p, h_{np})$.

Its distribution is denoted $\mathcal{NW}(n, p)$.

Proposition 1. Let $V \sim \mathcal{NW}(n, p)$. Then,

1.
$$E[V] = \frac{1}{p}I_p$$
,

2.
$$E[\operatorname{vec}(\boldsymbol{V})\operatorname{vec}(\boldsymbol{V})^{\top}] = \frac{1}{np(np+2)} \left(n(\boldsymbol{I}_{p^2} + \boldsymbol{K}_{p,p}) + n^2 \operatorname{vec}(\boldsymbol{I}_p) \operatorname{vec}(\boldsymbol{I}_p)^{\top} \right),$$

3. V is invertible with probability 1,

4.
$$E[V^{-1}] = \frac{np-2}{n-p-1}I_p$$
,

5.
$$E\left[\operatorname{vec}(\mathbf{V}^{-1})\operatorname{vec}(\mathbf{V}^{-1})^{\top}\right] = \frac{(np-2)(np-4)}{(n-p)(n-p-1)(n-p-3)}\left[(\mathbf{I}_{p^2} + \mathbf{K}_{p,p}) + (n-p-2)\operatorname{vec}(\mathbf{I}_p)\operatorname{vec}(\mathbf{I}_p)^{\top}\right].$$

Proof. Let $S \sim \mathcal{W}(n, \mathbf{I}_p)$ such that $V = S/\operatorname{tr}(S)$. From Theorem 1, $S = \mathcal{Q}V$, where $\mathcal{Q} \sim \chi^2_{np}$ and $V \stackrel{d}{=} UU^{\top}$ with U uniformly distributed on the unit sphere $\mathcal{C}_{p,n}$. Moreover, \mathcal{Q} and U are independent (hence, \mathcal{Q} and V are also independent). First, to prove 1., remembering that $E[\mathcal{Q}] = np$ and $E[S] = nI_p$ (from (12)), it is enough to notice that

$$E[S] = nI_p = E[Q]E[V] = npE(V).$$

To prove 2., recall that $E[Q^2] = np(np+2)$ and $E[\text{vec}(\boldsymbol{S}) \text{vec}(\boldsymbol{S})^{\top}] = n(\boldsymbol{I}_{p^2} + \boldsymbol{K}_{p,p}) + n^2 \text{vec}(\boldsymbol{I}_p) \text{vec}(\boldsymbol{I}_p)^{\top}$ (from (13)). It is then enough to compute

$$E[\operatorname{vec}(\boldsymbol{S})\operatorname{vec}(\boldsymbol{S})^{\top}] = n(\boldsymbol{I}_{p^2} + \boldsymbol{K}_{p,p}) + n^2 \operatorname{vec}(\boldsymbol{I}_p) \operatorname{vec}(\boldsymbol{I}_p)^{\top}$$
$$= E[\mathcal{Q}^2] E[\operatorname{vec}(\boldsymbol{V})\operatorname{vec}(\boldsymbol{V})^{\top}] = np(np+2) E[\operatorname{vec}(\boldsymbol{V})\operatorname{vec}(\boldsymbol{V})^{\top}].$$

To prove 3., it is enough to notice that, by construction, S is invertible with probability 1. To prove 4., remember that $E[Q^{-1}] = \frac{1}{np-2}$ and $E[S^{-1}] = \frac{1}{n-p-1}I_p$ (Equation (18)). To conclude, it is then enough to observe

$$E[S^{-1}] = \frac{1}{n-p-1}I_p = E[Q^{-1}]E[V^{-1}] = \frac{1}{np-2}E[V^{-1}].$$
 (26)

Finally, to prove 5., recall $E[Q^{-2}] = \frac{1}{(np-2)(np-4)}$ and, from (19),

$$E[\text{vec}(\boldsymbol{S}^{-1}) \text{ vec}(\boldsymbol{S}^{-1})^{\top}] = \frac{1}{(n-p)(n-p-1)(n-p-3)} [(\boldsymbol{I}_{p^2} + \boldsymbol{K}_{p,p}) + (n-p-2) \text{ vec}(\boldsymbol{I}_p) \text{ vec}(\boldsymbol{I}_p)^{\top}].$$

The result follows immediately from combining these with

$$E[\operatorname{vec}(\boldsymbol{S}^{-1})\operatorname{vec}(\boldsymbol{S}^{-1})^{\top}] = E[\mathcal{Q}^{-2}]E[\operatorname{vec}(\boldsymbol{V}^{-1})\operatorname{vec}(\boldsymbol{V})^{-1})^{\top}].$$

This concludes the proof.

The stochastic representation of Theorem 1 along with Normalized Wishart distribution offers an alternative way to compute the characteristic function of the Elliptical Wishart distribution (22). To obtain it, we thus need the characteristic function of the Normalized Wishart distribution. It is provided in Proposition 2. Before we get to the proposition, the first required thing is the expectation of zonal polynomials of the matrix ZV. This is achieved in Lemma 1.

Lemma 1. Let $V \sim \mathcal{NW}(n,p)$ and let $Z \in \mathcal{S}_p$. The expectation of the κ -order zonal polynomial of ZV is given by

$$E[C_{\kappa}(\mathbf{Z}\mathbf{V})] = \frac{\Gamma\left(\frac{np}{2}\right)}{\Gamma\left(\frac{np}{2} + k\right)} \left(\frac{n}{2}\right)_{\kappa} C_{\kappa}(\mathbf{Z}). \tag{27}$$

Proof. Let $S \sim \mathcal{W}(n, I_p)$ such that $V = S/\operatorname{tr}(S)$. From Theorem 1, $S = \mathcal{Q}V$, where $\mathcal{Q} \sim \chi_{np}^2$ and $V \stackrel{d}{=} UU^{\top}$ with U uniformly distributed on the unit sphere $\mathcal{C}_{p,n}$. Furthermore, \mathcal{Q} and V are independent. Therefore, since $C_{\kappa}(\alpha A) = \alpha^k C_{\kappa}(A)$ for $\alpha \in \mathbb{R}$ and $A \in \mathcal{S}_p$,

$$E[C_{\kappa}(\mathbf{Z}\mathbf{S})] = E[C_{\kappa}(\mathcal{Q}\mathbf{Z}\mathbf{V})] = E[\mathcal{Q}^{k}C_{\kappa}(\mathbf{Z}\mathbf{V})] = E[\mathcal{Q}^{k}]E[C_{\kappa}(\mathbf{Z}\mathbf{V})].$$

From [51], we get $E[Q^k] = 2^k \frac{\Gamma(\frac{np}{2} + k)}{\Gamma(\frac{np}{2})}$. The result then stems from (16).

Proposition 2. Let $V \sim \mathcal{NW}(n,p)$. The characteristic function of V at $Z \in \mathcal{S}_p$ is

$$\Psi_{\mathbf{V}}(\mathbf{Z}) = \sum_{k=0}^{+\infty} \frac{i^k}{k!} \frac{\Gamma\left(\frac{np}{2}\right)}{\Gamma\left(\frac{np}{2} + k\right)} \sum_{|\kappa| = k} \left(\frac{n}{2}\right)_{\kappa} C_{\kappa}(\mathbf{Z}). \tag{28}$$

Proof. Since $V \in \mathcal{S}_p^+$, we have $\operatorname{tr}(ZV) \leq \|Z\| \operatorname{tr}(V) = \|Z\|$ where $\|Z\|$ is the spectral norm of Z, and consequently, $E[\operatorname{tr}(ZV)^k] \leq \|Z\|^k$. Thus, using the dominated convergence theorem, one can write

$$\Psi_{\boldsymbol{V}}(\boldsymbol{Z}) = E[\exp(i\operatorname{tr}(\boldsymbol{Z}\boldsymbol{V}))] = \sum_{k=0}^{+\infty} \frac{i^k}{k!} E[\operatorname{tr}(\boldsymbol{Z}\boldsymbol{V})^k].$$

Moreover, from $\operatorname{tr}(\boldsymbol{Z}\boldsymbol{V})^k = \sum_{|\kappa|=k} C_{\kappa}(\boldsymbol{Z}\boldsymbol{V})$ and Lemma 1, we get $E[\operatorname{tr}(\boldsymbol{Z}\boldsymbol{V})^k] = \frac{\Gamma\left(\frac{np}{2}\right)}{\Gamma\left(\frac{np}{2}+k\right)} \sum_{|\kappa|=k} \left(\frac{n}{2}\right)_{\kappa} C_{\kappa}(\boldsymbol{Z})$. This is enough to conclude.

4.2 Elliptical Wishart distribution

We now have all the necessary tools to derive the statistical properties of the Elliptical Wishart distribution. First, leveraging the stochastic representation from Theorem 1 along with the expectation and variance of the Normalized Wishart distribution from Proposition 1, the expectation and variance of the Elliptical Wishart distribution $\mathcal{EW}(n, \Sigma, h_{np})$ are derived. This is achieved in Propositions 3 and 4.

Proposition 3 (Expectation). Suppose that moment $m_1^{(h_{np})}$ from (9) is finite. The expectation of $S \sim \mathcal{EW}(n, \Sigma, h_{np})$

$$\mathbb{E}[S] = \frac{m_1^{(h_{np})}}{p} \Sigma. \tag{29}$$

Proof. From Theorem 1, $S = \mathcal{Q} \Sigma^{1/2} U U^{\top} \Sigma^{1/2}$. Since U and \mathcal{Q} are independent, $E[S] = E[\mathcal{Q}] \Sigma^{1/2} E[U U^{\top}] \Sigma^{1/2}$. As $UU^{\top} \sim \mathcal{NW}(n, p)$, the point 1. of Proposition 1 yields the result.

Proposition 4 (Variance). Suppose that $m_1^{(h_{np})}$ and $m_2^{(h_{np})}$ from (9) are finite. The variance of $S \sim \mathcal{EW}(n, \Sigma, h_{np})$

$$\operatorname{var}[\mathbf{S}] = n a_{n,p} (\mathbf{I}_{p^2} + \mathbf{K}_{p,p}) (\mathbf{\Sigma} \otimes \mathbf{\Sigma}) + n^2 b_{n,p} \operatorname{vec}(\mathbf{\Sigma}) \operatorname{vec}(\mathbf{\Sigma})^{\top},$$
where $a_{n,p} = \frac{m_2^{(h_{np})}}{np(np+2)}$ and $b_{n,p} = a_{n,p} - \frac{(m_1^{(h_{np})})^2}{(np)^2}$. (30)

Proof. From Theorem 1, $S = \mathcal{Q}\Sigma^{1/2}UU^{\top}\Sigma^{1/2}$. Therefore, since $(C^{\top} \otimes A)\operatorname{vec}(B) = \operatorname{vec}(ABC)$, we have $\operatorname{vec}(S) = \mathcal{Q}(\Sigma^{1/2} \otimes \Sigma^{1/2}) \operatorname{vec}(UU^{\top})$. Since U and \mathcal{Q} are independent,

$$E[\operatorname{vec}(\boldsymbol{S})\operatorname{vec}(\boldsymbol{S})^{\top}] = E[\mathcal{Q}^2](\boldsymbol{\Sigma}^{1/2} \otimes \boldsymbol{\Sigma}^{1/2}) E[\operatorname{vec}(\boldsymbol{U}\boldsymbol{U}^{\top})\operatorname{vec}(\boldsymbol{U}\boldsymbol{U}^{\top})^{\top}](\boldsymbol{\Sigma}^{1/2} \otimes \boldsymbol{\Sigma}^{1/2}).$$

The result is finally obtained by using 2. from Proposition 1.

Remark 5 (Characteristic function). The stochastic representation from Theorem 1 offers an alternative way from [17] to derive the characteristic function (22) of the Elliptical Wishart distribution. It is actually simpler because it does not use the integral equality proved in of [17, Theorem 2.1.1]. Instead, the independence of Q and U from the stochastic representation along with characteristic functions of Q and of the Normalized Wishart distribution from Proposition 2 are leveraged. Indeed, from $S = \mathcal{Q}\Sigma^{1/2}(UU^{\top})\Sigma^{1/2}$, we get

$$\Psi_{\boldsymbol{S}}(\boldsymbol{Z}) = E\left[E\left[\exp\left(i\boldsymbol{\mathcal{Q}}\operatorname{tr}(\boldsymbol{Z}\boldsymbol{\Sigma}^{1/2}\boldsymbol{U}\boldsymbol{U}^{\top}\boldsymbol{\Sigma}^{1/2})\right)|\boldsymbol{\mathcal{Q}}\right]\right] = E\left[\Psi_{\boldsymbol{U}\boldsymbol{U}^{\top}}(\boldsymbol{\mathcal{Q}}\boldsymbol{\Sigma}^{1/2}\boldsymbol{Z}\boldsymbol{\Sigma}^{1/2})\right].$$

Then,
$$C_{\kappa}(\mathbf{\Sigma}^{1/2}\mathbf{Z}\mathbf{\Sigma}^{1/2}) = C_{\kappa}(\mathbf{\Sigma}\mathbf{Z})$$
 and (28) yields $\Psi_{UU^{\top}}(\mathcal{Q}\mathbf{\Sigma}^{1/2}\mathbf{Z}\mathbf{\Sigma}^{1/2}) = \sum_{k=0}^{+\infty}\frac{i^k}{k!}\mathcal{Q}^k\frac{\Gamma\left(\frac{np}{2}\right)}{\Gamma\left(\frac{np}{2}+k\right)}\sum_{|\kappa|=k}\left(\frac{n}{2}\right)_{\kappa}C_{\kappa}(\mathbf{\Sigma}\mathbf{Z}).$ From there, let ρ : $x \in \mathbb{R}$ \mapsto

$$\sum_{k=0}^{+\infty} \sqrt{m_{2k}^{(h_{np})}} \left(\frac{\Gamma\left(\frac{np}{2}\right)}{\Gamma\left(\frac{np}{2}+k\right)}\right)^{1/p} \frac{\Gamma\left(\frac{n}{2}+k\right)}{\Gamma\left(\frac{n}{2}\right)} \frac{x^k}{k!}.$$
 Assuming h_{np} admits a Taylor series expansion, we have

 $\sum_{k=0}^{+\infty} m_k^{(h_{np})} \frac{x^k}{k!} < +\infty. \text{ Therefore, using the D'Alembert criterion, one can prove that } \rho(x) \text{ converges absolutely for any } x. \text{ From [22, Lemma 6.5], } |C_\kappa(\boldsymbol{A})| \leq p^k \|\boldsymbol{A}\|^k \text{ for any } \boldsymbol{A} \in \mathcal{S}_p. \text{ Since } k! \geq k_1! \dots k_p! \text{ (where } k = \sum_{j=1}^p k_j), \text{ we also have}$

$$\sum_{k=0}^{+\infty} \left| m_k^{(h_{np})} \frac{\Gamma\left(\frac{np}{2}\right)}{\Gamma\left(\frac{np}{2}+k\right)} \sum_{|\kappa|=k} \left(\frac{n}{2}\right)_{\kappa} \frac{C_{\kappa}(\mathbf{\Sigma}\mathbf{Z})}{k!} \right| \leq \sum_{k=0}^{+\infty} \sum_{|\kappa|=k} m_k^{(h_{np})} \frac{\Gamma\left(\frac{np}{2}\right)}{\Gamma\left(\frac{np}{2}+k\right)} \left(\frac{n}{2}\right)_{\kappa} \frac{p^k \|\mathbf{Z}\mathbf{\Sigma}\|^k}{k_1! \dots k_p!}.$$

Furthermore, notice that $E[\mathcal{Q}^k] = E[\prod_{j=1}^p \mathcal{Q}^{k_j}] \leq \prod_{j=1}^p \sqrt{E[\mathcal{Q}^{2k_j}]}$. In addition, $\left(\frac{\Gamma\left(\frac{np}{2}+k\right)}{\Gamma\left(\frac{np}{2}\right)}\right)^p \geq \prod_{j=1}^p \frac{\Gamma\left(\frac{np}{2}+k_j\right)}{\Gamma\left(\frac{np}{2}\right)}$ and $\left(\frac{n}{2}\right)_{\kappa} \leq \prod_{j=1}^p \left(\frac{n}{2}\right)_{k_j} = \prod_{j=1}^p \frac{\Gamma\left(\frac{n}{2}+k_j\right)}{\Gamma\left(\frac{n}{2}\right)}$. Hence,

$$\sum_{k=0}^{+\infty} \left| m_k^{(h_{np})} \frac{\Gamma(\frac{np}{2})}{\Gamma(\frac{np}{2}+k)} \sum_{|\kappa|=k} \left(\frac{n}{2}\right)_{\kappa} \frac{C_{\kappa}(\boldsymbol{\Sigma}\boldsymbol{Z})}{k!} \right| \leq \prod_{j=1}^{p} \sum_{k_j=0}^{+\infty} \sqrt{m_{2k_j}^{(h_{np})}} \left(\frac{\Gamma(\frac{np}{2})}{\Gamma(\frac{np}{2}+k_j)}\right)^{1/p} \frac{\Gamma(\frac{n}{2}+k_j)}{\Gamma(\frac{n}{2})} \frac{p^{k_j} \|\boldsymbol{Z}\boldsymbol{\Sigma}\|^{k_j}}{k_j!},$$

which is finite thanks to the absolute convergence of ρ . Finally, the use of the dominated convergence theorem yields the result $\Psi_{\mathbf{Z}}(\mathbf{S}) = \sum_{k=0}^{+\infty} \frac{i^k}{k!} m_k^{(h_{np})} \frac{\Gamma(\frac{np}{2})}{\Gamma(\frac{np}{2}+k)} \sum_{|\kappa|=k} \left(\frac{n}{2}\right)_{\kappa} C_{\kappa}(\mathbf{\Sigma}\mathbf{Z}).$

The last statistical property of the Elliptical Wishart distribution studied in this work is the Kronecker moments at any order. Again, we resort to the stochastic representation from Theorem 1. Hence, to obtain Kronecker moments of the Elliptical Wishart distribution, we rely on Kronecker moments of the Wishart distribution. As mentioned in Section 3.1, Kronecker moments of a Wishart-distributed matrix are derived in [53] up to order 4 by recursively differentiating 4 times the characteristic function of the Wishart distribution. Our first major contribution is to provide a novel recursive way to compute Kronecker moments of the Wishart distribution up to any order, which leads to non-recursive explicit formulas. This is achieved in Proposition 5. From there, Kronecker moments of the Elliptical Wishart distribution are obtained in Proposition 6.

Proposition 5 (Kronecker moments). Let $S \sim W(n, \Sigma)$ where $\Sigma \in \mathcal{S}_p^{++}$. Then, for $k \in \mathbb{N}^*$,

$$E[\otimes^{k+1} \mathbf{S}] = nE[\otimes^k \mathbf{S}] \otimes \mathbf{\Sigma} - 2 \frac{\partial E[\otimes^k \mathbf{S}]}{\partial \mathbf{\Sigma}^{-1}}.$$
(31)

In a non-recursive way,

$$\operatorname{vec}(E[\otimes^{k} \mathbf{S}]) = \left(\otimes^{2k} \mathbf{\Sigma}^{1/2}\right) \prod_{l=0}^{k-1} \left[\mathbf{I}_{p^{2l}} \otimes \mathbf{M}_{(k-1-l)} \right] \left(\otimes^{k} \operatorname{vec}(\mathbf{I}_{p})\right), \tag{32}$$

where

$$\begin{split} \boldsymbol{M}_{(k)} &= \left[n(\boldsymbol{I}_{p^k} \otimes \boldsymbol{K}_{p,p^k} \otimes \boldsymbol{I}_p) + \boldsymbol{J}_{(k)} \right] \boldsymbol{K}_{p^{2k},p^2}, \\ \boldsymbol{J}_{(k)} &= \left[\sum_{l=0}^{k-1} (\boldsymbol{H}_{(k,l)} \otimes \boldsymbol{H}_{(k,l)}) \right] (\boldsymbol{I}_{p^{k-1}} \otimes \boldsymbol{K}_{p,p^{k-1}} \otimes \boldsymbol{I}_p) \left(\boldsymbol{I}_{p^{k-1}} \otimes \boldsymbol{G} \right) (\boldsymbol{I}_{p^{k-1}} \otimes \boldsymbol{K}_{p^{k-1},p^2} \otimes \boldsymbol{I}_{p^2}) (\boldsymbol{I}_{p^k} \otimes \boldsymbol{K}_{p,p^k} \otimes \boldsymbol{I}_p), \\ \boldsymbol{H}_{(k,l)} &= \left(\boldsymbol{I}_{p^l} \otimes \boldsymbol{K}_{p,p^{k-1-l}} \otimes \boldsymbol{I}_p \right), \\ \boldsymbol{G} &= \frac{1}{2} \left[(\boldsymbol{K}_{pp} \otimes \boldsymbol{K}_{pp}) (\boldsymbol{I}_p \otimes \boldsymbol{K}_{pp} \otimes \boldsymbol{I}_p) + \boldsymbol{I}_{p^4} \right] \left[\boldsymbol{K}_{pp} \otimes (\boldsymbol{I}_{p^2} + \boldsymbol{K}_{pp}) \right]. \end{split}$$

Proposition 6 (Kronecker moments). Let $S \sim \mathcal{EW}(n, \Sigma, h_{np})$. Then, for $k \in \mathbb{N}^*$ such that $m_k^{(h_{np})}$ defined in (9) is finite,

$$\operatorname{vec}(E[\otimes^{k} \mathbf{S}]) = \frac{\Gamma(\frac{np}{2})}{2^{k} \Gamma(\frac{np}{2} + k)} m_{k}^{(h_{np})} \left(\otimes^{2k} \mathbf{\Sigma}^{1/2} \right) \prod_{l=0}^{k-1} \left[\mathbf{I}_{p^{2l}} \otimes \mathbf{M}_{(k-1-l)} \right] \left(\otimes^{k} \operatorname{vec}(\mathbf{I}_{p}) \right), \tag{33}$$

where $M_{(k)}$ is defined in Proposition 5.

Proof. From Theorem 1, one has $S = \mathcal{Q}\Sigma^{1/2}V\Sigma^{1/2}$ with $V \stackrel{d}{=} UU^{\top}$. Consequently, $\otimes^k S = \mathcal{Q}^k \otimes^k (\Sigma^{1/2}V\Sigma^{1/2})$ and

$$\operatorname{vec}(E[\otimes^k \mathbf{S}]) = E[\mathcal{Q}^k] \operatorname{vec}(E[\otimes^k (\mathbf{\Sigma}^{1/2} \mathbf{V} \mathbf{\Sigma}^{1/2})]) = m_k^{(h_{np})} \operatorname{vec}(E[\otimes^k (\mathbf{\Sigma}^{1/2} \mathbf{V} \mathbf{\Sigma}^{1/2})]).$$
(34)

It remains to determine $\text{vec}(E[\otimes^k(\mathbf{\Sigma}^{1/2}V\mathbf{\Sigma}^{1/2})])$. To do so, we leverage the fact that $\text{vec}(E[\otimes^k\mathbf{S}])$ is known in the Wishart case. Indeed, in this specific case, using $m_k^{(h_{np})}=2^k\frac{\Gamma(\frac{np}{2}+k)}{\Gamma(\frac{np}{2})}$ and injecting (33) from Proposition 5 in (34) yields

$$\left(\otimes^{2k}\boldsymbol{\Sigma}^{1/2}\right)\prod_{l=0}^{k-1}\left[\boldsymbol{I}_{p^{2l}}\otimes\boldsymbol{M}_{(k-1-l)}\right]\left(\otimes^{k}\operatorname{vec}(\boldsymbol{I}_{p})\right)=2^{k}\frac{\Gamma(\frac{np}{2}+k)}{\Gamma(\frac{np}{2})}\operatorname{vec}(E[\otimes^{k}(\boldsymbol{\Sigma}^{1/2}\boldsymbol{V}\boldsymbol{\Sigma}^{1/2}]).$$

This is enough to conclude.

4.3 Inverse Elliptical Wishart distribution

After studying the Elliptical Wishart distribution, we turn to the Inverse Elliptical Wishart distribution. As in the previous section, the stochastic representation from Theorem 1 along with the statistical properties of the Normalized Wishart distribution from Section 4.1 are exploited. First, Proposition 1 is exploited to derive the expectation and variance of the Inverse Elliptical Wishart distribution $\mathcal{EW}^{-1}(n, \Sigma, h_{nn})$. These are in Propositions 7 and 8.

Proposition 7 (Expectation). Suppose that n > p+1 and moment $m_{-1}^{(h_{np})}$ from (9) is finite. The expectation of $S \sim \mathcal{EW}^{-1}(n, \Sigma, h_{np})$ is

$$E[S] = \frac{np-2}{n-p-1} m_{-1}^{(h_{np})} \Sigma.$$
(35)

Proof. From Theorem 1, $S = \frac{1}{\mathcal{Q}} \Sigma^{1/2} (UU^{\top})^{-1} \Sigma^{1/2}$. Since U and \mathcal{Q} are independent, $E[S] = E\left[\frac{1}{\mathcal{Q}}\right] \Sigma^{1/2} E[(UU^{\top})^{-1}] \Sigma^{1/2}$. The result is then obtained using point 4. from Proposition 1.

Proposition 8 (Variance). Suppose that n > p+3 and moments $m_{-1}^{(h_{np})}$ and $m_{-2}^{(h_{np})}$ are finite. The variance of $S \sim \mathcal{EW}^{-1}(n, \Sigma, h_{np})$ is

$$\operatorname{var}[\mathbf{S}] = c_{n,p}(\mathbf{I}_{p^2} + \mathbf{K}_{p,p})(\mathbf{\Sigma} \otimes \mathbf{\Sigma}) + d_{n,p}\operatorname{vec}(\mathbf{\Sigma})\operatorname{vec}(\mathbf{\Sigma})^{\top}, \tag{36}$$

where
$$c_{n,p} = \frac{(np-2)(np-4)}{(n-p)(n-p-1)(n-p-3)} m_{-2}^{(h_{np})}$$
 and $d_{n,p} = (n-p-2)c_{n,p} - \left(\frac{np-2}{n-p-1}m_{-1}^{(h_{np})}\right)^2$.

Proof. From Theorem 1, $S = \frac{1}{Q} \Sigma^{1/2} (UU^\top)^{-1} \Sigma^{1/2}$. Therefore, $\text{vec}(S) = \frac{1}{Q} (\Sigma^{1/2} \otimes \Sigma^{1/2}) \text{ vec}((UU^\top)^{-1})$. Since U and Q are independent,

$$E[\operatorname{vec}(\boldsymbol{S})\operatorname{vec}(\boldsymbol{S})^{\top}] = E\left[\frac{1}{\mathcal{Q}^{2}}\right] (\boldsymbol{\Sigma}^{1/2} \otimes \boldsymbol{\Sigma}^{1/2}) E[\operatorname{vec}((\boldsymbol{U}\boldsymbol{U}^{\top})^{-1})\operatorname{vec}((\boldsymbol{U}\boldsymbol{U}^{\top})^{-1})^{\top}] (\boldsymbol{\Sigma}^{1/2} \otimes \boldsymbol{\Sigma}^{1/2}). \tag{37}$$

Point 5. of Proposition 1 concludes the proof.

As for the Inverse Wishart distribution, the characteristic function of the Inverse Elliptical Wishart distribution appears quite complicated to obtain and remains unknown. The last statistical property of the Inverse Elliptical Wishart distribution $\mathcal{EW}^{-1}(n, \Sigma, h_{np})$ that is derived is the Kronecker moments. These are provided in Proposition 9.

Proposition 9 (Kronecker moments). Let $S \sim \mathcal{EW}^{-1}(n, \Sigma, h_{np})$. For $k < \frac{np}{2} - 1$ such that $m_{-k-1}^{(h_{np})}$ defined in (9) is finite, we have

$$\operatorname{vec}(E[\otimes^{k+1} \mathbf{S}]) = \frac{2^{k+1} \Gamma(\frac{np}{2})}{(n-p-1) \Gamma(\frac{np}{2}-k-1)} m_{-k-1}^{(h_{np})} \prod_{i=0}^{k-1} (\mathbf{I}_{p^{2i}} \otimes \mathbf{A}(k-i)) (\otimes^{k+1} \operatorname{vec}(\mathbf{\Sigma})), \tag{38}$$

where A(k) is defined in (20).

Proof. From Theorem 1, $S = \frac{1}{\mathcal{Q}} \Sigma^{1/2} V^{-1} \Sigma^{1/2}$ with $V \stackrel{d}{=} U U^{\top}$. Consequently, $\otimes^{k+1} S = \frac{1}{\mathcal{Q}^{k+1}} \otimes^{k+1} (\Sigma^{1/2} V^{-1} \Sigma^{1/2})$ and

$$\operatorname{vec}(E[\otimes^{k+1} \mathbf{S}]) = E[\mathcal{Q}^{-k-1}] \operatorname{vec}(E[\otimes^{k+1} (\mathbf{\Sigma}^{1/2} \mathbf{V}^{-1} \mathbf{\Sigma}^{1/2}]) = m_{-k-1}^{(h_{np})} \operatorname{vec}(E[\otimes^{k+1} (\mathbf{\Sigma}^{1/2} \mathbf{V}^{-1} \mathbf{\Sigma}^{1/2}]).$$
(39)

It remains to obtain $\text{vec}(E[\otimes^{k+1}(\mathbf{\Sigma}^{1/2}\mathbf{V}^{-1}\mathbf{\Sigma}^{1/2})])$. Similarly to the proof of Proposition 6, we exploit the fact that $\text{vec}(E[\otimes^{k+1}\mathbf{S}])$ is known in the Inverse Wishart case. In this specific case, using $E[\mathcal{Q}^{-k-1}] = 2^{-k-1}\frac{\Gamma\left(\frac{np}{2}-k-1\right)}{\Gamma\left(\frac{np}{2}\right)}$ and injecting (20) in (39) yields

$$\frac{1}{n-p-1}\prod_{i=0}^{k-1}(\boldsymbol{I}_{p^{2i}}\otimes\boldsymbol{A}(k-i))(\otimes^{k+1}\operatorname{vec}(\boldsymbol{\Sigma}))=2^{-k-1}\frac{\Gamma\left(\frac{np}{2}-k-1\right)}{\Gamma\left(\frac{np}{2}\right)}\operatorname{vec}(E[\otimes^{k+1}(\boldsymbol{\Sigma}^{1/2}\boldsymbol{V}^{-1}\boldsymbol{\Sigma}^{1/2})]).$$

This is enough to conclude.

4.4 Examples with particular Elliptical Wishart distributions

Finally, this section is dedicated to some special cases of the Elliptical Wishart and Inverse Elliptical Wishart distributions. We deal with the Gaussian, t-, Generalized Gaussian and Kotz density generators. In each case, to be able to apply results from Sections 4.2 and 4.3, explicit values of moments $m_k^{(h_{np})}$ for $k \in \{-2, -1, 1, 2\}$ are provided. For further details on these moments, see e.g., [32]. Here, we use them to compute the expectations and variances from Propositions 3, 4, 7 and 8.

4.4.1 Gaussian density generator

For the usual Wishart and Inverse Wishart distributions, the density generator is $h_{np}: t \mapsto (2\pi)^{-np/2} \exp\left(-\frac{t}{2}\right)$. As mentioned previously, with the notations of Theorem (1), we have $\mathcal{Q} \sim \chi_{np}^2$ and,

$$m_k^{(h_{np})} = 2^k \frac{\Gamma\left(\frac{np}{2} + k\right)}{\Gamma\left(\frac{np}{2}\right)}.$$
 (40)

By plugging these moments for $k \in \{1, 2, -1, -2\}$ in Equations (29), (30), (35) and (36), the expectations (12) and (18), and variances (13) and (19) of the Wishart and Inverse Wishart distributions, respectively.

4.4.2 *t*-density generator

The t- density generator $h_{np}: t\mapsto (\pi\nu)^{-np/2}\frac{\Gamma(\frac{\nu+np}{2})}{\Gamma(\frac{\nu}{2})}\left(1+\frac{t}{\nu}\right)^{-\frac{\nu+np}{2}}$, which is characterized by its degree of freedom $\nu>0$, yields the t-Wishart distribution. With the notations of Theorem 1, we recognize that $\frac{\mathcal{Q}}{np}\sim\mathcal{F}(np,\nu)$, the F-distribution with np and ν degrees of freedom [47]. Recall that the k-th moment of an F- random variable is

$$m_k^{(h_{np})} = \nu^k \frac{\Gamma\left(\frac{np}{2} + k\right)}{\Gamma\left(\frac{np}{2}\right)} \frac{\Gamma\left(\frac{\nu}{2} - k\right)}{\Gamma\left(\frac{\nu}{2}\right)}, \quad -\frac{np}{2} < k < \frac{\nu}{2}. \tag{41}$$

Notice that $m_{-1}^{(h_{np})}$ of the t- distribution is the same as in the Gaussian case. Consequently, the Inverse Wishart and Inverse t-Wishart share the same expectation. Moreover, as expected, when $\nu \to +\infty$, moments $m_k^{(h_{np})}$ of the the t-distribution converges to those of the Normal distribution. This simply arises from the fact that the density generator h_{np} of the t- distribution converges point-wisely to the density generator of the Normal distribution and from the monotone convergence theorem. For the expectations and variances, we obtain, given $S \sim t$ - $W(n, \Sigma, \nu)$,

$$E[S] = \frac{v}{v-2} n \Sigma, \quad \text{for } \nu > 2,$$

$$\operatorname{var}[S] = \frac{v^2}{(\nu-2)(\nu-4)} n (I_{p^2} + K_{p,p}) (\Sigma \otimes \Sigma) + \frac{2\nu^2}{(\nu-2)^2(\nu-4)} n^2 \operatorname{vec}(\Sigma) \operatorname{vec}(\Sigma)^{\top}, \quad \text{for } \nu > 4,$$

$$E[S^{-1}] = \frac{1}{n-p-1} \Sigma^{-1}, \quad \text{for } n > p+1$$

$$\operatorname{var}[S^{-1}] = \frac{1}{(n-p)(n-p-1)(n-p-3)} \left[\left(1 + \frac{2}{\nu} \right) (I_{p^2} + K_{p,p}) (\Sigma^{-1} \otimes \Sigma^{-1}) + \frac{2}{n-p-1} \left(1 + \frac{(n-p-2)(n-p-1)}{\nu} \right) \operatorname{vec}(\Sigma^{-1}) \operatorname{vec}(\Sigma^{-1}) \right], \quad \text{for } n > p+3.$$

$$(42)$$

4.4.3 Generalized Gaussian density generator

The density generator of the Generalized Gaussian distribution, also called Power Exponential distribution, is defined as $h_{np}(t) = \beta \left(2^{1/\beta}\pi\right)^{-np/2} \exp(-t^{\beta}/2)$, characterized by the shape parameter $\beta > 0$. When $0 < \beta < 1$, it features heavier tail regions than the normal distribution and hence can be useful in providing robustness against outliers [35]. With the notations of Theorem (1), $\mathcal{Q}^{\beta} \sim \chi^{2}_{\frac{np}{\beta}}$, the Chi-squared distribution with $\frac{np}{\beta}$ degrees of freedom (not necessarily an integer), which corresponds to the Gamma distribution $\Gamma\left(\frac{np}{2\beta},2\right)$. From [57], the moments $m_k^{(h_{np})}$ are

$$m_k^{(h_{np})} = 2^{\frac{k}{\beta}} \frac{\Gamma\left(\frac{np}{2\beta} + \frac{k}{\beta}\right)}{\Gamma\left(\frac{np}{2\beta}\right)}.$$
 (43)

For the expectations and variances, we obtain, given $S \sim \mathcal{GGW}(n, \Sigma, \beta)$,

$$E[S] = \frac{2^{\frac{1}{\beta}}}{p} \frac{\Gamma\left(\frac{np}{2\beta} + \frac{1}{\beta}\right)}{\Gamma\left(\frac{np}{2\beta}\right)} \Sigma,$$

$$\operatorname{var}[S] = \frac{2^{\frac{2}{\beta}}}{p(np+2)} \frac{\Gamma\left(\frac{np}{2\beta} + \frac{2}{\beta}\right)}{\Gamma\left(\frac{np}{2\beta}\right)} \left[(\boldsymbol{I}_{p^2} + \boldsymbol{K}_{p,p})(\boldsymbol{\Sigma} \otimes \boldsymbol{\Sigma}) + n\left(1 - \left(1 + \frac{2}{np}\right) \frac{\Gamma\left(\frac{np}{2\beta} + \frac{1}{\beta}\right)^2}{\Gamma\left(\frac{np}{2\beta} + \frac{2}{\beta}\right) \Gamma\left(\frac{np}{2\beta}\right)} \right) \operatorname{vec}(\boldsymbol{\Sigma}) \operatorname{vec}(\boldsymbol{\Sigma})^{\top} \right],$$

$$E[S^{-1}] = 2^{-\frac{1}{\beta}} \frac{np-2}{n-p-1} \frac{\Gamma\left(\frac{np}{2\beta} - \frac{1}{\beta}\right)}{\Gamma\left(\frac{np}{2\beta}\right)} \Sigma^{-1},$$

$$\operatorname{var}[S^{-1}] = \frac{2^{-\frac{2}{\beta}}(np-2)(np-4)}{(n-p)(n-p-1)(n-p-3)} \frac{\Gamma\left(\frac{np}{2\beta} - \frac{2}{\beta}\right)}{\Gamma\left(\frac{np}{2\beta}\right)} \left[(\boldsymbol{I}_{p^2} + \boldsymbol{K}_{p,p})(\boldsymbol{\Sigma}^{-1} \otimes \boldsymbol{\Sigma}^{-1}) + \left((n-p-2) - \frac{(n-p)(np-3)(np-2)}{(n-p-1)(np-4)} \frac{\Gamma\left(\frac{np}{2\beta} - \frac{1}{\beta}\right)^2}{\Gamma\left(\frac{np}{2\beta} - \frac{2}{\beta}\right) \Gamma\left(\frac{np}{2\beta}\right)} \right) \operatorname{vec}(\boldsymbol{\Sigma}^{-1}) \operatorname{vec}(\boldsymbol{\Sigma}^{-1})^{\top} \right].$$

4.4.4 Kotz density generator

The Kotz Wishart distribution $\mathcal{KW}(n, \Sigma, \alpha, \beta, R)$ with parameters α, β and R has as density generator

$$h_{np}(t) = \beta \pi^{-np/2} R^{\frac{np+2(\alpha-1)}{2\beta}} \frac{\Gamma(\frac{np}{2})}{\Gamma(\frac{np}{2\beta} + \frac{\alpha-1}{\beta})} t^{\alpha-1} \exp(-Rt^{\beta}),$$

where $\alpha+\frac{np}{2}>1$ and R>0. Kotz model is a generalization of Gaussian and Generalized Gaussian distributions. When $\alpha=1$ and $R=\frac{1}{2}$, we get the Generalized Gaussian with parameter β . If, in addition, $\beta=1$, we obtain the Gaussian distribution. With the notations of Theorem (1), $\mathcal{Q}\sim\Gamma\left(R^{-1/\beta},\alpha-1+\frac{np}{2},\beta\right)$, the generalized Gamma distribution. From [52], it follows that, for $k>-\frac{np}{2}-\alpha+1$,

$$m_k^{(h_{np})} = R^{-\frac{k}{\beta}} \frac{\Gamma(\frac{np}{2\beta} + \frac{\alpha + k - 1}{\beta})}{\Gamma(\frac{np}{2\beta} + \frac{\alpha - 1}{\beta})}.$$
 (45)

We thus obtain for $S \sim \mathcal{KW}(n, \Sigma, \alpha, \beta, R)$

$$E[S] = \frac{R^{\frac{1}{\beta}}}{p} \frac{\Gamma\left(\frac{np}{2\beta} + \frac{\alpha}{\beta}\right)}{\Gamma\left(\frac{np}{2\beta} + \frac{\alpha-1}{\beta}\right)} \mathbf{\Sigma},$$

$$\operatorname{var}[S] = \frac{R^{\frac{2}{\beta}}}{p(np+2)} \frac{\Gamma\left(\frac{np}{2\beta} + \frac{\alpha+1}{\beta}\right)}{\Gamma\left(\frac{np}{2\beta} + \frac{\alpha-1}{\beta}\right)} \left[(\mathbf{I}_{p^2} + \mathbf{K}_{p,p})(\mathbf{\Sigma} \otimes \mathbf{\Sigma}) + n\left(1 - \left(1 + \frac{2}{np}\right) \frac{\Gamma\left(\frac{np}{2\beta} + \frac{\alpha}{\beta}\right)^2}{\Gamma\left(\frac{np}{2\beta} + \frac{\alpha+1}{\beta}\right)\Gamma\left(\frac{np}{2\beta} + \frac{\alpha-1}{\beta}\right)} \right) \operatorname{vec}(\mathbf{\Sigma}) \operatorname{vec}(\mathbf{\Sigma})^{\top} \right],$$

$$E[S^{-1}] = R^{-\frac{1}{\beta}} \frac{np-2}{n-p-1} \frac{\Gamma\left(\frac{np}{2\beta} + \frac{\alpha-2}{\beta}\right)}{\Gamma\left(\frac{np}{2\beta} + \frac{\alpha-1}{\beta}\right)} \mathbf{\Sigma}^{-1},$$

$$\operatorname{var}[S^{-1}] = \frac{R^{-\frac{2}{\beta}}(np-2)(np-4)}{(n-p)(n-p-1)(n-p-3)} \frac{\Gamma\left(\frac{np}{2\beta} + \frac{\alpha-3}{\beta}\right)}{\Gamma\left(\frac{np}{2\beta} + \frac{\alpha-1}{\beta}\right)} \left[(\mathbf{I}_{p^2} + \mathbf{K}_{p,p})(\mathbf{\Sigma}^{-1} \otimes \mathbf{\Sigma}^{-1}) + \left((n-p-2) - \frac{(n-p)(np-3)(np-2)}{(n-p-1)(np-4)} \frac{\Gamma\left(\frac{np}{2\beta} + \frac{\alpha-2}{\beta}\right)^2}{\Gamma\left(\frac{np}{2\beta} + \frac{\alpha-3}{\beta}\right)\Gamma\left(\frac{np}{2\beta} + \frac{\alpha-1}{\beta}\right)} \right) \operatorname{vec}(\mathbf{\Sigma}^{-1}) \operatorname{vec}(\mathbf{\Sigma}^{-1})^{\top} \right].$$
(46)

5 Numerical experiments

In this section, we highlight the merit of Elliptical Wishart distributions on real data compared to the Wishart distribution. Indeed, we assess to which extent some encephalographic signals can be fitted with Elliptical Wishart models. In order to perform such a distribution fitting efficiently, we need first to generate samples from Elliptical Wishart distributions in a numerically practical way. This is achieved in Section 5.1 wherein we also suggest a deft method to draw samples from Inverse Elliptical distributions. This accounts for a contribution of the paper as our new technique of generating samples drawn from Elliptical and Inverse Elliptical Wishart distributions outperforms the existing technique. Thereafter, we exploit this method in Section 5.2 to fit the empirical cumulative density function of real EEG data in order to show that the Elliptical Wishart distribution offers a more relevant fitting than the Wishart distribution, commonly used as the state-of-the-art model based on the Gaussianity hypothesis.

5.1 Samples generation

This first subsection is dedicated to developing an efficient way to draw samples from Elliptical and Inverse Elliptical Wishart distributions. In fact, by definition of the Inverse Elliptical Wishart distribution (Definition 6), it is enough to be able to generate samples from the Elliptical Wishart distribution in order to get random samples in both cases. Again, to achieve this, we heavily rely on the stochastic representation from Theorem 1 and on existing results for the Wishart distribution.

A naive way to draw samples from the Elliptical Wishart distribution $\mathcal{EW}(n, \Sigma, h_{np})$ with center $\Sigma \in \mathcal{S}_p^{++}$ is to exploit Definition 5. Indeed, one can simply draw $x \in \mathbb{R}^{np}$ from the multivariate elliptical distribution $\mathcal{E}_{np}(\mathbf{0}, I_{np}, h_{np})^8$ to construct the random matrix $X \sim \mathcal{E}_{p,n}(\mathbf{0}, I_p, I_n, h_{np})$ such that vec(X) = x (Definition 2). Then, from the Cholesky decomposition $\Sigma = LL^{\top 9}$, a random matrix $S \sim \mathcal{EW}(n, \Sigma, h_{np})$ is obtained by taking $S = LXX^{\top}L^{\top}$. However, this procedure quickly becomes computationally very demanding when increasing the dimension p or degree of freedom n since it implies generating a \mathbb{R}^{np} vector. Since the latter usually corresponds to the amount of available samples, it is common for it to be quite big. In such a case, the method described above is really not satisfying.

⁸The usual way to generate such random vector \boldsymbol{x} is to leverage their stochastic representation (8), i.e., $\boldsymbol{x} = \mathcal{Q}\boldsymbol{u}$. To generate a uniformly distributed vector \boldsymbol{u} on the unit sphere \mathcal{C}_{np} , one can draw a Gaussian vector $\boldsymbol{z} \sim \mathcal{N}_{np}(\boldsymbol{0}, \boldsymbol{I}_{np})$ and then set $\boldsymbol{u} = \frac{\boldsymbol{z}}{\|\boldsymbol{z}\|_2}$. Finally, the second-order modular random variable \mathcal{Q} is generated from its corresponding distribution.

 $^{{}^{9}}L \in \mathbb{R}^{p \times p}$, lower triangular matrix with strictly positive diagonal elements.

To deal with this issue in the Wishart case, the so-called Bartlett decomposition [8] is exploited. It indeed provides a very computationally efficient way to generate random Wishart matrices. First, to build $S \sim \mathcal{W}(n, \Sigma)$, $R \sim \mathcal{W}(n, I_p)$ is constructed. S is then obtained through $S = LRL^{\top}$, where $\Sigma = LL^{\top}$ is the Cholesky decomposition of Σ . To synthesize R, its Cholesky decomposition $R = TT^{\top}$ is also exploited. Indeed, from the Bartlett decomposition, random values $(T_{k\ell})_{1 \leq \ell \leq k \leq p}$ are independent. Moreover, every diagonal element T_{kk} $(1 \leq k \leq p)$ is drawn from the Chi distribution χ_{n-k+1} with n-k+1 degrees of freedom, while the lower triangular elements $T_{k\ell}$ $(1 \leq \ell < k \leq p)$ are drawn from the normal distribution $\mathcal{N}(0,1)$. This yields a simple way to generate T and thus $S \sim \mathcal{W}(n,\Sigma)$.

Similarly to the multivariate case, the stochastic representation can be exploited to simulate random Elliptical Wishart matrices. To get $S \sim \mathcal{EW}(n, \Sigma, h_{np})$, we exploit Theorem 1 and set $S = \mathcal{Q}LVL^{\top}$, where L corresponds to the Cholesky decomposition of Σ , \mathcal{Q} is the so-called second-order modular random variable and V follows the Normalized Wishart distribution $\mathcal{NW}(n,p)$. Randomly generating \mathcal{Q} can be achieved using classic techniques as in the multivariate elliptical case. To construct a proper V, we synthesize $R \sim \mathcal{W}(n, I_p)$ with the Bartlett decomposition approach described above and then set $V = \frac{R}{\operatorname{tr}(R)}$. The proposed method to generate data from the Elliptical Wishart distribution is summarized in Algorithm 1.

Algorithm 1 Random sample generation from the Elliptical Wishart distribution

```
Input: dimension p, degrees of freedom n \geq p, center \Sigma \in \mathcal{S}_p^{++}, density generator h_{np}
```

Output: random matrix $S \sim \mathcal{EW}(n, \Sigma, h_{np})$

Generate
$$\mathcal Q$$
 with PDF $t\mapsto rac{\pi^{np/2}}{\Gamma(np/2)}h_{np}(t)t^{rac{np}{2}-1}$

Generate lower triangular matrix $T \in \mathbb{R}^{p \times p}$ such that: $T_{kk} \stackrel{\text{i.i.d}}{\sim} \chi_{n-k+1}$, $1 \leq k \leq p$ and $T_{k\ell} \stackrel{\text{i.i.d}}{\sim} \mathcal{N}(0,1)$, $1 \leq \ell < k \leq p$, independent from the diagonal entries.

Set R = TT

Let $V=rac{R}{\operatorname{tr}(R)}$

Compute Cholesky decomposition $oldsymbol{\Sigma} = oldsymbol{L} oldsymbol{L}^ op$

Let $S = \mathcal{Q}LVL^{\mathsf{T}}$

5.2 Electroencephalographic data fitting

We perform some distribution fitting over EEG data to assess the interest of Elliptical Wishart distributions in practice. We consider the Steady-State Visually Evoked Potential (SSVEP) dataset [27], which contains recordings from 12 subjects. In the SSVEP paradigm, subjects are exposed to a light blinking at various frequencies. Data are thus divided into several classes corresponding to these various stimuli frequencies. In our case, the dataset contains four classes corresponding to stimuli at 11Hz, 17Hz, and 21Hz, along with a resting state. To keep our experiment simple, we limit ourselves to subject 12, which features the cleanest data. These data contain 160 trials equally partitioned between the four classes. For each trial, we have p=8 electrodes and n=768 samples (3 seconds sampled at 256Hz). For further details on these data, the reader is referred to [27] and to the MOABB platform [5].

One distribution fitting is achieved for each class. For class z, we first estimate the center $\mathbf{\Sigma}^{(z)}$ of the tested distribution from the covariance matrices $\mathbf{S}_i^{(z)} = \mathbf{X}_i^{(z)} \mathbf{X}_i^{(z) \top}$ of the trials $\mathbf{X}_i^{(z)} \in \mathbb{R}^{np}$ belonging to class z. In this work, we consider the following distributions for the testing: the Wishart distribution $\mathcal{W}(n,\mathbf{\Sigma}^{(z)})$ and the t-Wishart distribution t- $\mathcal{W}(n,\mathbf{\Sigma}^{(z)},\nu^{(z)})$ with $\nu^{(z)}$ degrees of freedom for the t- density generator. The latter parameter $\nu^{(z)}$ is chosen through trials and errors in order to get the best fitting results possible. The actual values used in our experiments are given in Table 2. $\mathbf{\Sigma}^{(z)}$ is set as the maximum likelihood estimator of the distribution. For the Wishart distribution, it is simply given by $\mathbf{\Sigma}^{(z)} = \frac{1}{nK_z} \sum_i \mathbf{S}_i^{(z)}$, where K_z corresponds to the number of trials belonging to class z. For the t-Wishart distribution, no closed form exists and we resort to the iterative algorithm derived in [6] to estimate it.

The actual fitting procedure consists in estimating the empirical CDFs of the traces, ℓ_2 -norms and determinants of real EEG covariances $\{S_i^{(z)}\}_{i=1}^{K_z}$, for every class $z \in \{1, 2, 3, 4\}$. We compare these to the CDFs of the traces, ℓ_2 -

ν_1	ν_2	ν_3	ν_4
24	20	28	41

Table 2: Degrees of freedom to generate t-Wishart samples for fitting EEG data.

norms and determinants of the Wishart distribution $\mathcal{W}(n, \Sigma^{(z)})$ and t-Wishart distribution t- $\mathcal{W}(n, \Sigma^{(z)}, \nu^{(z)})$. To avoid the computation of exact CDFs of these quantities for the Wishart and t-Wishart models, we instead generate random samples from these distributions with the procedures introduced in Section 5.1 and then empirically estimate the CDFs. All these CDFs are plotted in Figure 1, 2 and 3. To not only rely on visual inspection of the results, but we also perform goodness-to-git Kolmogorov-Smirnov tests of the traces, ℓ_2 -norms, and determinants of the data at hand. Since we simulate data from the Wishart and t-Wishart distributions, we rely on two-sample tests. The null hypothesis is that compared samples are drawn from the same distribution. The p-values of the Kolmogorov-Smirnov tests of the traces, ℓ_2 -norms, and determinants are provided in Tables 3, 4 and 5, respectively.

From every Figure and Table, the conclusion is very clear. On these real EEG data, the *t*-Wishart distribution better fits the covariances than the Wishart distribution for every class. As a matter of fact, while the correspondence between the Wishart distribution and the data at hand is quite poor, the *t*-Wishart distribution is a pretty good fit. Even though the considered dataset is very specific and quite small, this clearly demonstrates the practical interest of such distribution. For instance, this spurs the design of a Bayesian classifier of EEG signals that exploits *t*-Wishart in a discriminant analysis framework, as presented in [7], wherein the proposed classifier outperforms the state-of-the-art method on the tested datasets.




Fig. 1: Empirical CDF of the traces of the covariance matrices extracted from EEG signals along with the CDF of the trace of Wishart matrix and the CDF of the trace of t-Wishart matrix, obtained as the empirical CDFs of a large number (10^5) of traces of generated samples from Wishart $\mathcal{W}(n, \widehat{\Sigma}_k^{\mathcal{W}})$ and t-Wishart t- $\mathcal{W}(n, \widehat{\Sigma}_k^{t-\mathcal{W}}, \nu_k)$ respectively.

6 Conclusions and perspectives

In this paper, we derived some statistical properties of the Elliptical Wishart and Inverse Elliptical Wishart that are not covered in the literature. More specifically, we computed the expectation, variance, and Kronecker moments. The

	Class 1	Class 2	Class 3	Class 4
Wishart	6.10^{-6}	3.10^{-8}	7.10^{-9}	10^{-5}
t-Wishart	0.7196	0.6006	0.305	0.765

Table 3: p-values of the two-sample Kolmogorov-Smirnov tests on traces




Fig. 2: Empirical CDF of the ell_2 norms of the covariance matrices extracted from EEG signals along with the CDF of the ell_2 norm of Wishart matrix, obtained as the empirical CDFs of a large number (10⁵) of ell_2 norms of generated samples from Wishart $\mathcal{W}(n, \widehat{\Sigma}_k^{\mathcal{W}})$ and t-Wishart t- $\mathcal{W}(n, \widehat{\Sigma}_k^{t-\mathcal{W}}, \nu_k)$ respectively.

				Class 4
Wishart	4.10^{-6}	10^{-6}	5.10^{-6}	10^{-5}
t-Wishart	0.6325	0.5661	0.1311	0.6383

Table 4: p-values of the two-sample Kolmogorov-Smirnov tests on ell_2 norms

	Class 1	Class 2	Class 3	Class 4
Wishart	7.10^{-11}	10^{-15}	10^{-10}	4.10^{-10}
t-Wishart	0.3211	0.044	0.2437	0.1877

Table 5: p-values of the two-sample Kolmogorov-Smirnov tests on determinants




Fig. 3: Empirical CDF of the $-\log_{10}$ determinants of the covariance matrices extracted from EEG signals along with the CDF of the $-\log_{10}$ determinant of Wishart matrix and the CDF of the $-\log_{10}$ determinants of t-Wishart matrix, obtained as the empirical CDFs of a large number (10^5) of $-\log_{10}$ determinants of generated samples from Wishart $\mathcal{W}(n,\widehat{\Sigma}_k^{\mathcal{W}})$ and t-Wishart t- $\mathcal{W}(n,\widehat{\Sigma}_k^{t-\mathcal{W}})$ respectively.

fundamental instrument for these results is the stochastic representation of a random matrix drawn from an Elliptical Wishart distribution. It also simplifies the existing proofs based on tedious integral calculations to compute the already-derived properties, such as the characteristic function of Elliptical Wishart. In order to derive the Kronecker moments for Elliptical Wishart, we first had to explicitly compute the Kronecker moments of a Wishart matrix to any order, which is one of the contributions of this paper. The usefulness of Elliptical Wishart distributions in the field of electroencephalography is underlined thanks to goodness-to-fit statistical tests, which demonstrated the great potential of this family of distributions, particularly *t*-Wishart, in fitting real EEG signals.

It would be interesting to study the asymptotic behavior of Elliptical Wishart matrices in the future. Furthermore, on the one hand, it is worth comparing the Elliptical Wishart model and the sample covariance matrix of i.i.d vectors drawn from a multivariate elliptical distribution. On the other hand, approaching the asymptotic distribution of M-estimators with Elliptical Wishart seems relevant to us. Finally, the question of estimating the parameters of Elliptical Wishart distributions, particularly the center, is very interesting, especially if we want to exploit them for classification. This involves studying the theoretical properties of the developed estimators, such as the existence, uniqueness, estimation bound, asymptotic distribution, *etc*.

Acknowledgments

This work is partially supported by a public grant overseen by the French National Research Agency (ANR) through the program UDOPIA, a project funded by the ANR-20-THIA-0013-01. This research was also supported by DATAIA Convergence Institute as part of the "Programme d'Investissement d'Avenir", (ANR-17-CONV-0003) operated by laboratoire des Signaux et Systèmes.

References

- [1] AY Abul-Magd, Gernot Akemann, and P Vivo. Superstatistical generalizations of wishart–laguerre ensembles of random matrices. *Journal of Physics A: Mathematical and Theoretical*, 42(17):175207, 2009.
- [2] Sondipon Adhikari. Generalized wishart distribution for probabilistic structural dynamics. *Computational Mechanics*, 45:495–511, 2010.
- [3] Theodore W Anderson. The non-central wishart distribution and certain problems of multivariate statistics. *The Annals of Mathematical Statistics*, pages 409–431, 1946.
- [4] Theodore Wilbur Anderson. An introduction to multivariate statistical analysis, volume 2. Wiley New York, 1958.
- [5] Bruno Aristimunha, Igor Carrara, Pierre Guetschel, Sara Sedlar, Pedro Rodrigues, Jan Sosulski, Divyesh Narayanan, Erik Bjareholt, Quentin Barthelemy, Robin Tibor Schirrmeister, Emmanuel Kalunga, Ludovic Darmet, Gregoire Cattan, Ali Abdul Hussain, Ramiro Gatti, Vladislav Goncharenko, Jordy Thielen, Thomas Moreau, Yannick Roy, Vinay Jayaram, Alexandre Barachant, and Sylvain Chevallier. Mother of all BCI Benchmarks, 2023.
- [6] I. Ayadi, F. Bouchard, and F. Pascal. Elliptical Wishart distribution: maximum likelihood estimator from information geometry. *International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, 2023.
- [7] Imen Ayadi, Florent Bouchard, and Frédéric Pascal. t-WDA: A novel Discriminant Analysis applied to EEG classification. In 2023 31st European Signal Processing Conference (EUSIPCO), pages 1793–1797. IEEE, 2023.
- [8] Maurice Stevenson Bartlett. Xx.—on the theory of statistical regression. *Proceedings of the Royal Society of Edinburgh*, 53:260–283, 1934.
- [9] A Bekker, M Arashi, and J Van Niekerk. Wishart generator distribution. arXiv preprint arXiv:1502.07300, 2015.
- [10] Andriëtte Bekker, Janet Van Niekerk, and Mohammad Arashi. Wishart distributions: Advances in theory with bayesian application. *Journal of Multivariate Analysis*, 155:272–283, 2017.
- [11] Martin Bilodeau and David Brenner. *Theory of multivariate statistics*. Springer Science & Business Media, 1999.
- [12] Adrian N Bishop, Pierre Del Moral, Angèle Niclas, et al. An introduction to wishart matrix moments. *Foundations and Trends*® *in Machine Learning*, 11(2):97–218, 2018.
- [13] Marie-France Bru. Wishart processes. *Journal of Theoretical Probability*, 4:725–751, 1991.
- [14] Włodzimierz Bryc. Compound real wishart and q-wishart matrices. *International Mathematics Research Notices*, 2008(9):rnn079–rnn079, 2008.
- [15] Stamatis Cambanis, Steel Huang, and Gordon Simons. On the theory of elliptically contoured distributions. *Journal of Multivariate Analysis*, 11(3):368–385, 1981.
- [16] Francisco J Caro-Lopera, Graciela González Farías, and Narayanaswamy Balakrishnan. Matrix-variate distribution theory under elliptical models-4: Joint distribution of latent roots of covariance matrix and the largest and smallest latent roots. *Journal of Multivariate Analysis*, 145:224–235, 2016.
- [17] Francisco José Caro Lopera. Noncentral elliptical configuration density. 2008.
- [18] José A Di, Ramón Gutiérrez-Jáimez, et al. On wishart distribution: some extensions. *Linear Algebra and its Applications*, 435(6):1296–1310, 2011.
- [19] José A Díaz-García and Ramón Gutiérrez-Jáimez. Distribution of the generalised inverse of a random matrix and its applications. *Journal of statistical planning and inference*, 136(1):183–192, 2006.
- [20] Kai-Tang Fang and Yao-Ting Zhang. Generalized multivariate analysis. (No Title), 1990.
- [21] Piotr Graczyk and Lyudmila Vostrikova. The moments of wishart processes via itô calculus. *Theory of Probability & Its Applications*, 51(4):609–625, 2007.

- [22] Kenneth I Gross and Donald St P Richards. Special functions of matrix argument. i. algebraic induction, zonal polynomials, and hypergeometric functions. *Transactions of the American Mathematical Society*, 301(2):781–811, 1987.
- [23] Arjun K Gupta and Daya K Nagar. Matrix variate distributions. Chapman and Hall/CRC, 2018.
- [24] Arjun K Gupta and Tamas Varga. *Elliptically contoured models in statistics*, volume 240. Springer Science & Business Media, 2012.
- [25] Grant Hillier and Raymond Kan. Moments of a wishart matrix. *Journal of Quantitative Economics*, 19:141–162, 2021.
- [26] Grant Hillier and Raymond Kan. Properties of the inverse of a noncentral wishart matrix. *Econometric Theory*, 38(6):1092–1116, 2022.
- [27] Emmanuel K Kalunga, Sylvain Chevallier, Quentin Barthélemy, Karim Djouani, Eric Monacelli, and Yskandar Hamam. Online SSVEP-based BCI using Riemannian geometry. *Neurocomputing*, 191:55–68, 2016.
- [28] Raymond Kan and Grant Hillier. On the expectations of equivariant matrix-valued functions of wishart and inverse wishart matrices. *Available at SSRN 4096101*, 2022.
- [29] CG Khatri. On certain distribution problems based on positive definite quadratic functions in normal vectors. *The Annals of Mathematical Statistics*, pages 468–479, 1966.
- [30] Tonu Kollo. Advanced multivariate statistics with matrices. Springer, 2005.
- [31] Gary Koop, Dimitris Korobilis, et al. Bayesian multivariate time series methods for empirical macroeconomics. *Foundations and Trends*® *in Econometrics*, 3(4):267–358, 2010.
- [32] Samuel Kotz. Multivariate distributions at a cross road. In A Modern Course on Statistical Distributions in Scientific Work: Volume 1—Models and Structures Proceedings of the NATO Advanced Study Institute held at the University of Calgagry, Calgary, Alberta, Canada July 29–August 10, 1974, pages 247–270. Springer, 1975.
- [33] GÉRARD Letac and H Massam. The normal quasi-wishart distribution. Contemporary Mathematics, 287:231–240, 2001.
- [34] Gérard Letac and Hélène Massam. All invariant moments of the wishart distribution. *Scandinavian Journal of Statistics*, 31(2):295–318, 2004.
- [35] JK Lindsey. Multivariate elliptically contoured distributions for repeated measurements. *Biometrics*, 55(4):1277–1280, 1999.
- [36] F Javier Lopez-Martinez, Eduardo Martos-Naya, Jose F Paris, and Andrea Goldsmith. Eigenvalue dynamics of a central wishart matrix with application to mimo systems. *IEEE Transactions on Information Theory*, 61(5):2693– 2707, 2015.
- [37] I-Li Lu and Donald St P Richards. Macmahon's master theorem, representation theory, and moments of wishart distributions. *Advances in Applied Mathematics*, 27(2-3):531–547, 2001.
- [38] D Maiwald and D Kraus. Calculation of moments of complex wishart and complex inverse wishart distributed matrices. *IEE Proceedings-Radar, Sonar and Navigation*, 147(4):162–168, 2000.
- [39] Ricardo Antonio Maronna. Robust M-estimators of multivariate location and scatter. The annals of statistics, pages 51–67, 1976.
- [40] Robb J Muirhead. Aspects of multivariate statistical theory. John Wiley & Sons, 2009.
- [41] S Munilla and Rodolfo Juan Carlos Cantet. Bayesian conjugate analysis using a generalized inverted wishart distribution accounts for differential uncertainty among the genetic parameters—an application to the maternal animal model. *Journal of Animal Breeding and Genetics*, 129(3):173–187, 2012.
- [42] Daya K Nagar, Alejandro Roldán-Correa, and Saralees Nadarajah. Expected values of scalar-valued functions of a complex wishart matrix. *Mathematics*, 11(9):2162, 2023.
- [43] Heinz Neudecker. On two matrix derivatives by kollo and von rosen. *SORT-Statistics and Operations Research Transactions*, pages 153–163, 2003.
- [44] Heinz Neudecker and Tom Wansbeek. Some results on commutation matrices, with statistical applications. *Canadian Journal of Statistics*, 11(3):221–231, 1983.
- [45] Yasuhide Numata and Satoshi Kuriki. On formulas for moments of the wishart distributions as weighted generating functions of matchings. *Discrete Mathematics & Theoretical Computer Science*, (Proceedings), 2010.
- [46] Esa Ollila, David E Tyler, Visa Koivunen, and H Vincent Poor. Complex elliptically symmetric distributions: Survey, new results and applications. *IEEE Transactions on signal processing*, 60(11):5597–5625, 2012.

- [47] PB Patnaik. The non-central χ 2-and f-distribution and their applications. Biometrika, 36(1/2):202–232, 1949.
- [48] Carlos Antunes Percíncula, Liliana Forzani, and Ricardo Toledano. Invariant moments of the wishart distribution: A sage package and website implementation. *journal of statistical software*, 3 2022.
- [49] Gabriel Fernando Pivaro, Santosh Kumar, Gustavo Fraidenraich, and Claudio Ferreira Dias. On the exact and approximate eigenvalue distribution for sum of wishart matrices. *IEEE Transactions on Vehicular Technology*, 66(11):10537–10541, 2017.
- [50] Alberto Roverato. Hyper inverse wishart distribution for non-decomposable graphs and its application to bayesian inference for gaussian graphical models. *Scandinavian Journal of Statistics*, 29(3):391–411, 2002.
- [51] Marvin Kenneth Simon. Probability distributions involving Gaussian random variables: A handbook for engineers and scientists. Springer, 2002.
- [52] Edney W Stacy. A generalization of the gamma distribution. *The Annals of mathematical statistics*, pages 1187–1192, 1962.
- [53] Shagufta A Sultan and Derrick S Tracy. Moments of wishart distribution. *Stochastic analysis and applications*, 14(2):237s–243, 1996.
- [54] Brajendra C Sutradhar and Mir M Ali. A generalization of the wishart distribution for the elliptical model and its moments for the multivariate t model. *Journal of Multivariate Analysis*, 29(1):155–162, 1989.
- [55] Abigael Taylor, Philippe Forster, Franck Daout, Hélène M Oriot, and Laurent Savy. A generalization of the fixed point estimate for packet-scaled complex covariance matrix estimation. *IEEE Transactions on Signal Processing*, 65(20):5393–5405, 2017.
- [56] Ch Teng, H Fang, and W Deng. The generalized noncentral wishart distribution. In *J. of Mathematical Research and Exposition*, volume 9, pages 479–488, 1989.
- [57] Herbert CS Thom. A note on the gamma distribution. Monthly weather review, 86(4):117–122, 1958.
- [58] J-Y Tourneret, André Ferrari, and Gérard Letac. The noncentral wishart distribution: Properties and application to speckle imaging. In *IEEE/SP 13th Workshop on Statistical Signal Processing*, 2005, pages 924–929. IEEE, 2005.
- [59] Oncel Tuzel, Fatih Porikli, and Peter Meer. Pedestrian detection via classification on riemannian manifolds. *IEEE transactions on pattern analysis and machine intelligence*, 30(10):1713–1727, 2008.
- [60] Dietrich Von Rosen. Moments for the inverted wishart distribution. *Scandinavian Journal of Statistics*, pages 97–109, 1988.
- [61] Dietrich Von Rosen. On moments of the inverted wishart distribution. *Statistics: A Journal of Theoretical and Applied Statistics*, 30(3):259–278, 1997.
- [62] John Wishart. The generalised product moment distribution in samples from a normal multivariate population. *Biometrika*, pages 32–52, 1928.
- [63] Chi Song Wong and Tonghui Wang. Laplace-wishart distributions and cochran theorems. *Sankhyā: The Indian Journal of Statistics, Series A*, pages 342–359, 1995.

A Proof of Proposition 5

The Kronecker moment of $S \sim \mathcal{W}(n, \Sigma)$ of arbitrary order k exits thanks to the dominated convergence theorem. In fact, for $1 \leq i, j \leq p^k$, $|(\otimes^k S)_{ij}| \leq ||S||_2^k \leq \mathcal{Q}^k ||\Sigma||_2^k$ and $E[\mathcal{Q}^k] = \frac{\Gamma(\frac{np}{2} + k)}{\Gamma(\frac{np}{2})} < \infty$.

Let f_S denote the probability density function of the Wishart distribution $\mathcal{W}(n, \Sigma)$. Using the fact that $\frac{\partial f_S(S)}{\partial \Sigma^{-1}} = \left(\frac{n}{2}\Sigma - \frac{1}{2}S\right) f_S(S)$, we get:

$$\frac{\partial E[\otimes^{k} \mathbf{S}]}{\partial \mathbf{\Sigma}^{-1}} = \int_{\mathcal{S}_{p}^{++}} \frac{\partial}{\partial \mathbf{\Sigma}^{-1}} \left((\otimes^{k} \mathbf{S}) \otimes f_{\mathbf{S}}(\mathbf{S}) \right) d\mathbf{S} = \int_{\mathcal{S}_{p}^{++}} (\otimes^{k} \mathbf{S}) \otimes \frac{\partial f_{\mathbf{S}}(\mathbf{S})}{\partial \mathbf{\Sigma}^{-1}} d\mathbf{S}$$

$$= \frac{n}{2} \int_{\mathcal{S}_{p}^{++}} (\otimes^{k} \mathbf{S}) \otimes \mathbf{\Sigma} f_{\mathbf{S}}(\mathbf{S}) d\mathbf{S} - \frac{1}{2} \int_{\mathcal{S}_{p}^{++}} (\otimes^{k+1} \mathbf{S}) f_{\mathbf{S}}(\mathbf{S}) d\mathbf{S}$$

$$= \frac{n}{2} E(\otimes^{k} \mathbf{S}) \otimes \mathbf{\Sigma} - \frac{1}{2} E[\otimes^{k+1} \mathbf{S}] \quad (47)$$

where $dS = \prod_{i \leq j} dS_{ij}$ and the first equality is a result of differentiating under the integral. This leads to the recursion in Equation (31). Since $\otimes^{k+1}S = S \otimes (\otimes^k S)$, then, by applying the formula of the derivative of a Kronecker product:

$$\frac{\partial \otimes^{k+1} \mathbf{S}}{\partial \mathbf{\Sigma}^{-1}} = \mathbf{S} \otimes \frac{\partial \otimes^{k} \mathbf{S}}{\partial \mathbf{\Sigma}^{-1}} + \left(\mathbf{K}_{p,p^{k}} \otimes \mathbf{I}_{p} \right) \left(\otimes^{k} \mathbf{S} \otimes \frac{\partial \mathbf{S}}{\partial \mathbf{\Sigma}^{-1}} \right) \left(\mathbf{K}_{p^{k},p} \otimes \mathbf{I}_{p} \right)$$
(48)

Accordingly, with a simple recursion, it can be shown that:

$$\frac{\partial \otimes^{k} \mathbf{S}}{\partial \mathbf{\Sigma}^{-1}} = \sum_{l=0}^{k-1} \left(\mathbf{I}_{p^{l}} \otimes \mathbf{K}_{p,p^{k-1-l}} \otimes \mathbf{I}_{p} \right) \left(\otimes^{k-1} \mathbf{S} \otimes \frac{\partial \mathbf{S}}{\partial \mathbf{\Sigma}^{-1}} \right) \left(\mathbf{I}_{p^{l}} \otimes \mathbf{K}_{p^{k-1-l},p} \otimes \mathbf{I}_{p} \right), \tag{49}$$

and thus:

$$vec\left(\frac{\partial \otimes^{k} \mathbf{S}}{\partial \mathbf{\Sigma}^{-1}}\right) = \left[\sum_{l=0}^{k-1} (\mathbf{H}_{(k,l)} \otimes \mathbf{H}_{(k,l)})\right] (\mathbf{I}_{p^{k-1}} \otimes \mathbf{K}_{p^{2},p^{k-1}} \otimes \mathbf{I}_{p^{2}}) \left\{vec\left(\otimes^{k-1} \mathbf{S}\right) \otimes vec\left(\frac{\partial \mathbf{S}}{\partial \mathbf{\Sigma}^{-1}}\right)\right\}. \quad (50)$$

Now, let us compute $vec\left(\frac{\partial S}{\partial \Sigma^{-1}}\right)$. For this, we write $S \stackrel{d}{=} \Sigma^{1/2} R \Sigma^{1/2}$ where R does not depend on Σ . Therefore,

$$\frac{\partial \mathbf{S}}{\partial \mathbf{\Sigma}^{-1}} = \frac{\partial \mathbf{\Sigma}^{1/2}}{\partial \mathbf{\Sigma}^{-1}} \left(\mathbf{I}_p \otimes \mathbf{R} \mathbf{\Sigma}^{1/2} \right) + \left(\mathbf{I}_p \otimes \mathbf{\Sigma}^{1/2} \right) \frac{\partial \mathbf{R} \mathbf{\Sigma}^{1/2}}{\partial \mathbf{\Sigma}^{-1}}, \tag{51}$$

$$\frac{\partial \mathbf{R} \mathbf{\Sigma}^{1/2}}{\partial \mathbf{\Sigma}^{-1}} = (\mathbf{I}_p \otimes \mathbf{R}) \frac{\partial \mathbf{\Sigma}^{1/2}}{\partial \mathbf{\Sigma}^{-1}}.$$
 (52)

So,

$$vec\left(\frac{\partial S}{\partial \Sigma^{-1}}\right) = vec\left(\frac{\partial \Sigma^{1/2}}{\partial \Sigma^{-1}}\left(I_{p} \otimes R\Sigma^{1/2}\right) + \left(I_{p} \otimes \Sigma^{1/2}\right)\left(I_{p} \otimes R\right)\frac{\partial \Sigma^{1/2}}{\partial \Sigma^{-1}}\right)$$

$$= vec\left(\frac{\partial \Sigma^{1/2}}{\partial \Sigma^{-1}}\left(I_{p} \otimes R\Sigma^{1/2}\right) + \left(I_{p} \otimes \Sigma^{1/2}R\right)\frac{\partial \Sigma^{1/2}}{\partial \Sigma^{-1}}\right)$$

$$= \left(I_{p^{2}} \otimes I_{p} \otimes \Sigma^{1/2}R + I_{p} \otimes \Sigma^{1/2}R \otimes I_{p^{2}}\right)vec\left(\frac{\partial \Sigma^{1/2}}{\partial \Sigma^{-1}}\right).$$
(53)

Let us compute $vec\left(\frac{\partial \Sigma^{1/2}}{\partial \Sigma^{-1}}\right)$:

$$\begin{array}{ll} (i) & \boldsymbol{\Sigma}^{1/2} \cdot \boldsymbol{\Sigma}^{1/2} \cdot \boldsymbol{\Sigma}^{-1} = \boldsymbol{I}_{p}, \\ (ii) & d\boldsymbol{\Sigma}^{1/2} \cdot \boldsymbol{\Sigma}^{-1/2} + \boldsymbol{\Sigma}^{1/2} \cdot d\boldsymbol{\Sigma}^{1/2} \cdot \boldsymbol{\Sigma}^{-1} + \boldsymbol{\Sigma} \cdot d\boldsymbol{\Sigma}^{-1} = 0, \\ (iii) & \boldsymbol{\Sigma}^{-1} \cdot d\boldsymbol{\Sigma}^{1/2} \cdot \boldsymbol{\Sigma}^{-1/2} + \boldsymbol{\Sigma}^{-1/2} \cdot d\boldsymbol{\Sigma}^{1/2} \cdot \boldsymbol{\Sigma}^{-1} = -d\boldsymbol{\Sigma}^{-1}, \\ (iv) & \left(\boldsymbol{\Sigma}^{-1/2} \otimes \boldsymbol{\Sigma}^{-1} + \boldsymbol{\Sigma}^{-1} \otimes \boldsymbol{\Sigma}^{-1/2}\right) \operatorname{vec}(d\boldsymbol{\Sigma}^{1/2}) = -\operatorname{vec}(d\boldsymbol{\Sigma}^{-1}), \\ (iv) & \left(\boldsymbol{\Sigma}^{-1/2} \otimes \boldsymbol{\Sigma}^{-1} + \boldsymbol{\Sigma}^{-1} \otimes \boldsymbol{\Sigma}^{-1/2}\right) & \operatorname{vec}(d\boldsymbol{\Sigma}^{1/2}) = -\operatorname{vec}(d\boldsymbol{\Sigma}^{-1}), \\ (v) & d\operatorname{vec}(\boldsymbol{\Sigma}^{1/2}) = -\left[\boldsymbol{\Sigma}^{-1/2} \otimes \boldsymbol{\Sigma}^{-1} + \boldsymbol{\Sigma}^{-1} \otimes \boldsymbol{\Sigma}^{-1/2}\right]^{-1} d\operatorname{vec}(\boldsymbol{\Sigma}^{-1}), \end{array}$$

where (ii) is obtained by differentiating (i); (iii) results from the multiplication of (ii) by Σ^{-1} ; vectorizing (iii) leads to (iv) using $\text{vec}(\boldsymbol{A}\boldsymbol{B}\boldsymbol{C}) = (\boldsymbol{C}^{\top} \otimes \boldsymbol{A}) \text{vec}(\boldsymbol{B})$ and finally, since vec is linear, we have $\text{vec}(d\cdot) = d \text{vec}(\cdot)$, which leads to (v). Therefore,

$$\frac{\partial \operatorname{vec}(\mathbf{\Sigma}^{1/2})}{\partial \operatorname{vec}(\mathbf{\Sigma}^{-1})^{\top}} = -\frac{1}{4} (\mathbf{I}_{p^2} + \mathbf{K}_{pp}) \left(\mathbf{\Sigma}^{-1/2} \otimes \mathbf{\Sigma}^{-1} + \mathbf{\Sigma}^{-1} \otimes \mathbf{\Sigma}^{-1/2} \right)^{-1} (\mathbf{I}_{p^2} + \mathbf{K}_{pp})
= -\frac{1}{2} (\mathbf{I}_{p^2} + \mathbf{K}_{pp}) (\mathbf{\Sigma}^{-1} \oplus \mathbf{\Sigma}^{-1/2})^{-1},$$
(54)

with the notation $A \oplus B = A \otimes B + B \otimes A$. Finally, using the fact that:

$$\operatorname{vec}\left(\frac{\partial \mathbf{Y}}{\partial \mathbf{X}}\right) = (\mathbf{K}_{pp} \otimes \mathbf{K}_{pp})(\mathbf{I}_{p} \otimes \mathbf{K}_{pp} \otimes \mathbf{I}_{p}) \operatorname{vec}\left(\frac{\partial \operatorname{vec}(\mathbf{Y})}{\partial \operatorname{vec}(\mathbf{X})^{\top}}\right), \tag{55}$$

we get:

$$\operatorname{vec}\left(\frac{\partial \mathbf{\Sigma}^{1/2}}{\partial \mathbf{\Sigma}^{-1}}\right) = -(\mathbf{K}_{pp} \otimes \mathbf{K}_{pp})(\mathbf{I}_{p} \otimes \mathbf{K}_{pp} \otimes \mathbf{I}_{p}) \operatorname{vec}\left(\frac{1}{2}(\mathbf{I}_{p^{2}} + \mathbf{K}_{pp})(\mathbf{\Sigma}^{-1} \oplus \mathbf{\Sigma}^{-1/2})^{-1}\right).$$
(56)

By injecting Equation (56) into Equation (50) and using some matrix manipulations, we have:

$$\operatorname{vec}\left(\frac{\partial \mathbf{S}}{\partial \mathbf{\Sigma}^{-1}}\right) = -\frac{1}{2}\left[(\mathbf{K}_{pp} \otimes \mathbf{K}_{pp})(\mathbf{I}_{p} \otimes \mathbf{K}_{pp} \otimes \mathbf{I}_{p}) + \mathbf{I}_{p^{4}} \right]$$

$$\left[\mathbf{K}_{pp} \otimes \left((\mathbf{I}_{p^{2}} + \mathbf{K}_{pp})(\mathbf{I}_{p^{2}} + \mathbf{\Sigma}^{1/2} \otimes \mathbf{\Sigma}^{-1/2})^{-1} \mathbf{K}_{pp} \right) \right] \operatorname{vec}\left(\mathbf{S} \otimes \mathbf{\Sigma}\right) \quad (57)$$

Thus,

$$\operatorname{vec}(\otimes^{k-1} \mathbf{S}) \otimes \operatorname{vec}\left(\frac{\partial \mathbf{S}}{\partial \mathbf{\Sigma}^{-1}}\right) = -\frac{1}{2} \left(\mathbf{I}_{p^{2k-2}} \otimes \mathbf{G}_{\mathbf{\Sigma}} \right) \left\{ \operatorname{vec}\left(\otimes^{k-1} \mathbf{S}\right) \otimes \operatorname{vec}\left(\mathbf{S} \otimes \mathbf{\Sigma}\right) \right\}$$
(58)

where $G_{\Sigma} = \left[(\boldsymbol{K}_{pp} \otimes \boldsymbol{K}_{pp}) (\boldsymbol{I}_{p} \otimes \boldsymbol{K}_{pp} \otimes \boldsymbol{I}_{p}) + \boldsymbol{I}_{p^{4}} \right] \left[\boldsymbol{K}_{pp} \otimes \left((\boldsymbol{I}_{p^{2}} + \boldsymbol{K}_{pp}) (\boldsymbol{I}_{p^{2}} + \boldsymbol{\Sigma}^{1/2} \otimes \boldsymbol{\Sigma}^{-1/2})^{-1} \boldsymbol{K}_{pp} \right) \right] \in \mathbb{R}^{p^{4} \times p^{4}}$. Moreover,

$$\operatorname{vec}\left(\otimes^{k-1}\mathbf{S}\right) \otimes \operatorname{vec}\left(\mathbf{S} \otimes \mathbf{\Sigma}\right) = \left(\mathbf{I}_{p^{k-1}} \otimes \mathbf{K}_{p^{k-1}, p^{2}} \otimes \mathbf{I}_{p^{2}}\right) \operatorname{vec}\left(\left(\otimes^{k}\mathbf{S}\right) \otimes \mathbf{\Sigma}\right) \\ = \left(\mathbf{I}_{p^{k-1}} \otimes \mathbf{K}_{p^{k-1}, p^{2}} \otimes \mathbf{I}_{p^{2}}\right) \left(\mathbf{I}_{p^{k}} \otimes \mathbf{K}_{p, p^{k}} \otimes \mathbf{I}_{p}\right) \left\{\operatorname{vec}\left(\otimes^{k}\mathbf{S}\right) \otimes \operatorname{vec}\left(\mathbf{\Sigma}\right)\right\}$$

$$(59)$$

Combining Equation (50), Equation (59) and Equation (58):

$$\operatorname{vec}\left(\frac{\partial E[\otimes^k \mathbf{S}]}{\partial \mathbf{\Sigma}^{-1}}\right) = -\frac{1}{2}J_{(k),\mathbf{\Sigma}}\left\{\operatorname{vec}(E[\otimes^k \mathbf{S}]) \otimes \operatorname{vec}(\mathbf{\Sigma})\right\}$$
(60)

where $m{J}_{(k),\Sigma} = \left[\sum_{l=0}^{k-1} (m{H}_{(k,l)} \otimes m{H}_{(k,l)})\right] (m{I}_{p^{k-1}} \otimes m{K}_{p^2,p^{k-1}} \otimes m{I}_{p^2}) (m{I}_{p^{2k-2}} \otimes m{G}_{\Sigma}) (m{I}_{p^{k-1}} \otimes m{K}_{p^{k-1},p^2} \otimes m{I}_{p^2}) (m{I}_{p^k} \otimes m{K}_{p,p^k} \otimes m{I}_p).$ By injecting Equation (60) in the recursive equation (31),

$$\operatorname{vec}(E[\otimes^{k+1} \mathbf{S}]) = \left[n(\mathbf{I}_{p^k} \otimes \mathbf{K}_{p,p^k} \otimes \mathbf{I}_p) + \mathbf{J}_{(k),\Sigma} \right] \mathbf{K}_{p^{2k},p^2} \left\{ \operatorname{vec}(\mathbf{\Sigma}) \otimes \operatorname{vec}(E[\otimes^k \mathbf{S}]) \right\}, \tag{61}$$

Let $m{M}_{(k), \Sigma} = \left[n(m{I}_{p^k} \otimes m{K}_{p, p^k} \otimes m{I}_p) + m{J}_{(k), \Sigma} \right] m{K}_{p^{2k}, p^2}$. Hence, using the fact that $E[m{S}] = n m{\Sigma}$,

$$\operatorname{vec}(E[\otimes^k \mathbf{S}]) = \prod_{l=0}^{k-1} \left[\mathbf{I}_{p^{2l}} \otimes \mathbf{M}_{(k-1-l), \mathbf{\Sigma}} \right] \left\{ \otimes^k \operatorname{vec}(\mathbf{\Sigma}) \right\}.$$
 (62)

Notice that the resulting formula derives explicitly and non-recursively the k^{th} Kronecker moment for $S \sim \mathcal{W}(n, \Sigma)$. However, the expression of $J_{(k),\Sigma}$ depends on Σ , which is not an elegant way to compute arbitrary orders. To overcome this issue, we use the invariance of the Wishart distribution. In fact, $R = \Sigma^{-1/2} S \Sigma^{-1/2} \sim \mathcal{W}(n, I_p)$ and it can be shown with recursion that $\otimes^k (\Sigma^{1/2} R \Sigma^{1/2}) = (\otimes^k \Sigma^{1/2})(\otimes^k R)(\otimes^k \Sigma^{1/2})$. Thus, it suffices to derive the Kronecker moments for $\Sigma = I_p$. Using Equation (62),

$$\operatorname{vec}(E[\otimes^k \mathbf{R}]) = \prod_{l=0}^{k-1} \left[\mathbf{I}_{p^{2l}} \otimes \mathbf{M}_{(k-1-l)} \right] \left\{ \otimes^k \operatorname{vec}(\mathbf{I}_p) \right\}$$
(63)

where $m{M}_{(k)} = m{M}_{(k), m{I}_p}$ and $m{G}_{m{I}_p} = \frac{1}{2} \left[(m{K}_{pp} \otimes m{K}_{pp}) (m{I}_p \otimes m{K}_{pp} \otimes m{I}_p) + m{I}_{p^4} \right] \left[m{K}_{pp} \otimes (m{I}_{p^2} + m{K}_{pp}) \right]$. Therefore,

$$\operatorname{vec}(E[\otimes^{k} \mathbf{S}]) = \left(\otimes^{2k} \mathbf{\Sigma}^{1/2}\right) \prod_{l=0}^{k-1} \left[\mathbf{I}_{p^{2l}} \otimes \mathbf{M}_{(k-1-l)} \right] \left\{ \otimes^{k} \operatorname{vec}(\mathbf{I}_{p}) \right\}.$$
(64)