BOUNDEDNESS OF COMPOSITION OPERATORS FROM LORENTZ SPACES TO ORLICZ SPACES

NAOYA HATANO, MASAHIRO IKEDA AND RYOTA KAWASUMI

ABSTRACT. The boundedness (continuity) of composition operators from some function space to another one is significant, though there are few results about this problem. Thus, in this study, we provide necessary and sufficient conditions on the boundedness of composition operators from Lorentz spaces to Orlicz spaces. We emphasize that the measure spaces associated with the Lorentz space may be different from those associated with the Orlicz spaces. Moreover we give some examples and counterexamples of the composed mappings in the conditions.

1. INTRODUCTION

Throughout this paper, the simbol $L^0(X,\mu)$ is defined by the space of all measurable functions over the measure space (X,μ) . Let $X = (X,\mu)$ and $Y = (Y,\nu)$ be σ -finite measure spaces.

Definition 1.1. Let $\tau : Y \to X$ be a measurable map, and assume that τ is nonsingular, namely, $\nu(\tau^{-1}(E)) = 0$ for each μ -null set E. The composition operator C_{τ} is defined by

$$C_{\tau}f(y) \equiv f(\tau(y)), \quad y \in Y$$

for all $f \in L^0(X, \mu)$.

Hereafter τ stands for a measurable and nonsingular map from Y to X.

The boundedness of the composition operators on the Lebesgue spaces is classical and well known by Singh [15]. After that many authors gave some generalizations for some function spaces ([1, 5, 4, 10, 6, 7, 13]). Especially, the boundedness on the Orlicz spaces is given by Cui, Hudzik, Kumar and Maligranda in [4]. In this paper, we aim the boundedness from one function space to another one. When 0 , the necessary and sufficient condition $of the boundedness of the composition operator <math>C_{\tau}$ from $L^p(X,\mu) \to L^q(Y,\nu)$ is for all μ measurable sets $E \subset X$,

$$\nu(\tau^{-1}(E))^p \le C\mu(E)^q.$$

Moreover, when $0 < q < p < \infty$, there is a counterexample of τ that the composition operator C_{τ} is bounded from $L^{p}(X,\mu)$ to $L^{q}(Y,\nu)$. As mentioned in [5] in the framework of the Lorentz spaces, we can show the boundedness of $C_{\tau} : L^{p,r}(X,\mu) \to L^{q,r}(Y,\nu)$ without the condition $0 , where <math>L^{p,r}(X,\mu)$ and $L^{q,r}(Y,\nu)$ are Lorentz spaces with $r \in (0,\infty]$.

In this paper, we provide necessary and sufficient conditions on the boundedness of composition operators from Lorentz spaces to Orlicz spaces. We emphasize that the measure spaces associated with the Lorentz space may be different from those associated with the Orlicz spaces.

Here and below, we will mention the main theorems. The mapping $\Phi : [0, \infty) \to [0, \infty)$ is called Young function when it satisfies the following conditions:

²⁰²⁰ Mathematics Subject Classification. Primary 42B35; Secondary 42B25.

Key words and phrases. Composition operator, Boundedness, Orlicz spaces, Lorentz spaces.

- (1) Φ is a positive function on $(0, \infty)$.
- (2) Φ is a convex function.

(3) $\lim_{t \downarrow 0} \Phi(t) = 0.$

Hereafter we assume that Φ is a Young function.

Definition 1.2. Let $\Phi : [0, \infty) \to [0, \infty)$ satisfy that $\Phi((\cdot)^{1/q})$ is a Young function for some $0 < q < \infty$. The Orlicz space $L^{\Phi}(X, \mu)$ is defined by

$$L^{\Phi}(X,\mu) \equiv \left\{ f \in L^{0}(X,\mu) : \int_{X} \Phi(\epsilon |f(x)|) \,\mathrm{d}\mu(x) < \infty, \quad \text{for every} \quad \epsilon > 0 \right\}$$

endowed with the (quasi-)norm

$$||f||_{L^{\Phi}(X)} \equiv \inf \left\{ \lambda > 0 : \int_{X} \Phi\left(\frac{|f(x)|}{\lambda}\right) d\mu(x) \le 1 \right\}.$$

Orlicz spaces are (quasi-)Banach spaces. If $\Phi(t) = t^p$, $0 , then <math>L^{\Phi}(X, \mu)$ is $L^p(X, \mu)$. Moreover, when $\Phi(t) = t \log(3 + t)$ or $e^t - 1$, the function space $L^{\Phi}(X, \mu)$ is an Orlicz space which is given as $L \log L(X, \mu)$ or $\exp L(X, \mu)$.

Remark 1.3. σ -finiteness is not necessary in the above definition.

Definition 1.4. Let $0 and <math>0 < q \le \infty$. The Lorentz space $L^{p,q}(X,\mu)$ is defined by the space of all $f \in L^0(X,\mu)$ with the finite quasi-norm

$$||f||_{L^{p,q}(X)} \equiv \begin{cases} \left(\int_0^\infty \left[t\mu(\{x \in X : |f(x)| > t\})^{\frac{1}{p}} \right]^q \frac{\mathrm{d}t}{t} \right)^{\frac{1}{q}}, & q < \infty, \\ \sup_{t>0} t\mu(\{x \in X : |f(x)| > t\})^{\frac{1}{p}}, & q = \infty. \end{cases}$$

Since $L^{p,p}(X,\mu) = L^p(X,\mu)$, the Lorentz spaces are generalization for the Lebesgue spaces. Additionally, the Lorentz spaces $L^{p,\infty}(X,\mu)$ stand for the weak Lebesgue spaces. When $p = \infty$, the Lorentz space $L^{\infty,q}(X,\mu)$ is equivalent to $L^{\infty}(X,\mu)$, formally. The Lorentz spaces $L^{p,q}(X,\mu)$ are quasi-Banach spaces. But it is well known that p,q > 1 implies the Lorentz spaces $L^{p,q}(X,\mu)$ are normable (cf. [8]).

Theorem 1.5. Let $0 and <math>\Phi : [0, \infty) \to [0, \infty)$ be a Young function. Then, the composition operator C_{τ} is bounded from $L^{p,1}(X,\mu)$ to $L^{\Phi}(Y,\nu)$ if and only if there exists a constant $D \ge 1$ such that for all μ -measurable sets $E \subset X$,

$$\nu(\tau^{-1}(E)) \le \left\{ \Phi\left(\frac{1}{D\mu(E)^{\frac{1}{p}}}\right) \right\}^{-1}.$$

By this theorem and the properties

$$\|f\|_{L^{p,q}(X)} = \||f|^q\|_{L^{p/q,1}(X)}^{\frac{1}{q}}, \quad \|f\|_{L^{\Phi}(X)} = \||f|^q\|_{L^{\Phi}((\cdot)^{1/q})(X)}^{\frac{1}{q}}$$

for each $f \in L^0(X, \mu)$, the following theorem can be obtained immediately.

Theorem 1.6. Let $0 < p, q < \infty$ and $\Phi : [0, \infty) \to [0, \infty)$. If $\Phi((\cdot)^{1/q})$ is a Young function, then, the composition operator C_{τ} is bounded from $L^{p,q}(X,\mu)$ to $L^{\Phi}(Y,\nu)$ if and only if there exists a constant $D \ge 1$ such that for all μ -measurable sets $E \subset X$,

$$\nu(\tau^{-1}(E)) \le \left\{ \Phi\left(\frac{1}{D\mu(E)^{\frac{1}{p}}}\right) \right\}^{-1}.$$

We remark that the boundedness of the composition operators from some Orlicz spaces to the different Orlicz spaces is interesting. Chawziuk et al. [3] (Musielak-Orlicz space) and Labuschagne and Majewski [11] gave the necessary and sufficient condition on τ of the boundedness of the composition operator from an Orlicz space to another Orlicz one under the assumption that τ is surjective.

We will use standard notation for inequalities. We use C to denote a positive constant that may vary from one occurrence to another. If $f \leq Cg$, then we write $f \leq g$ or $g \gtrsim f$, and if $f \leq g \leq f$, then we write $f \sim g$.

We organize the remaining part of the paper as follows: We write the preliminaries for the proof of theorems in Section 2. We prove the main theorem in Section 3. In Section 4, we give examples of τ in the condition of the main theorems.

2. Preliminaries

In this section, we introduce some Lemmas for Orlicz and Lorentz spaces to prove the main theorem.

Lemma 2.1. For a Young function Φ , the function φ satisfying

$$\Phi(t) = \int_0^t \varphi(s) \,\mathrm{d}s$$

is said left derivative of Φ . Then, the following assertions hold:

(1)
$$\int_X \Phi(|f(x)|) d\mu(x) = \int_0^\infty \varphi(t)\mu(\{x \in X : |f(x)| > t\}) dt.$$

(2)
$$\frac{\Phi(t)}{t} \le \varphi(t) \le \frac{\Phi(2t)}{t}.$$

For (1), see [8, Proposition 1.1.4] and for (2), see [14, P.20].

Lemma 2.2. [14, Corllary 7 in P.78] Let Φ be a Young function. Then, for all measurable sets $E \subset X$, the following holds:

$$\|\chi_E\|_{L^{\Phi}(X)} = \left\{ \Phi^{-1}\left(\frac{1}{\mu(E)}\right) \right\}^{-1}$$

Lemma 2.3 ([8, Proposition 1.4.10]). Let $0 and <math>0 < q_1 \le q_2 \le \infty$. Then,

$$L^{p,q_1}(X,\mu) \hookrightarrow L^{p,q_2}(X,\mu).$$

Lemma 2.4 ([8, Example 1.4.8]). Let $0 and <math>0 < q \le \infty$. Then, for all μ -measurable sets $E \subset X$,

$$\|\chi_E\|_{L^{p,q}(X)} = q^{-\frac{1}{q}} \mu(E)^{\frac{1}{p}}.$$

3. Proof of the main theorem

In this section, we will prove Theorem 1.5. In this proof, for a measurable function f, we write the distribution functions as

$$\mu_f(t) \equiv \mu(\{x \in X : |f(x)| > t\}), \quad \nu_f(t) \equiv \nu(\{x \in Y : |f(x)| > t\}), \quad t \ge 0$$

At first, we prove the "if part". Since the embedding $L^{p,1}(X,\mu) \hookrightarrow L^{p,\infty}(X,\mu)$ holds (see Lemma 2.3), we only consider the case $||f||_{L^{p,\infty}(X)} \leq (2D)^{-1}$ by normalization. Using Lemma

2.1(1) and the assumption, we have

$$\int_{Y} \Phi(|C_{\tau}f(y)|) \,\mathrm{d}\nu(y) = \int_{0}^{\infty} \varphi(t)\nu_{C_{\tau}f}(t) \,\mathrm{d}t \le \int_{0}^{\infty} \varphi(t) \left\{ \Phi\left(\frac{1}{D\mu_{f}(t)^{\frac{1}{p}}}\right) \right\}^{-1} \,\mathrm{d}t.$$

Since the mapping $t \mapsto \Phi(t)/t$ is increasing, we deduced that

$$\Phi\left(\frac{1}{D\mu_{f}(t)^{\frac{1}{p}}}\right) = \left\{2D \cdot t\mu_{f}(t)^{\frac{1}{p}}\Phi\left(\frac{2t}{2D \cdot t\mu_{f}(t)^{\frac{1}{p}}}\right)\right\} \cdot \frac{1}{2D \cdot t\mu_{f}(t)^{\frac{1}{p}}} \ge \frac{\Phi(2t)}{2D \cdot t\mu_{f}(t)^{\frac{1}{p}}}.$$
by Lemma 2.1 (2).

Thus, by Lemma 2.1 (2),

$$\int_{Y} \Phi(|C_{\tau}f(y)|) \,\mathrm{d}\nu(y) \le \int_{0}^{\infty} \varphi(t) \left\{ \frac{\Phi(2t)}{2D \cdot t\mu_{f}(t)^{\frac{1}{p}}} \right\}^{-1} \,\mathrm{d}t \le 2D \int_{0}^{\infty} \mu_{f}(t)^{\frac{1}{p}} \,\mathrm{d}t$$
$$= 2D \|f\|_{L^{p,1}(X)}.$$

Next, we prove the "only if part". Fix a μ -measurable set E. Then, assuming the boundedness of the composition operator C_{τ} from $L^{p,1}(X,\mu)$ to $L^{\Phi}(Y,\nu)$ with the operator norm D, we have

$$\|C_{\tau}\chi_E\|_{L^{\Phi}(Y)} \le D\|\chi_E\|_{L^{p,1}(X)}.$$

By $C_{\tau}\chi_E = \chi_{\tau^{-1}(E)}$, using Lemmsa 2.2 and 2.4, we have

$$\Phi^{-1}\left(\frac{1}{\nu(\tau^{-1}(E))}\right) \ge \frac{1}{D\mu(E)^{\frac{1}{p}}},$$

or equivalently,

$$\nu(\tau^{-1}(E)) \le \left\{ \Phi\left(\frac{1}{D\mu(E)^{\frac{1}{p}}}\right) \right\}^{-1},$$

as desired.

4. Examples

Example 4.1. If we set $\tau(x) = x$, the volume estimate

$$|\tau^{-1}(E)| = |E|$$

holds for any measurable set $E \subset \mathbb{R}$. If $p \geq 1$, then the composition operator C_{τ} is bounded from $L^{p,1}(\mathbb{R})$ to $L^p(\mathbb{R}^n)$. But when p < 1, the boundedness $C_{\tau} : L^{p,1}(\mathbb{R}) \to L^p(\mathbb{R}^n)$ fails. In fact, choosing

$$f(x) = (1 + |x|)^{-\frac{1}{p}} (\log(3 + |x|))^{-\frac{1}{p}}, \quad x \in \mathbb{R}$$

we have $f = C_{\tau} f \in L^{p,1}(\mathbb{R}) \setminus L^p(\mathbb{R})$. Here, see [8, Excercise 1.4.8] for details of the example f(x).

Example 4.2. Let $0 < p, q < \infty$, $0 < r \le \infty$, $(X, \mu) = (\mathbb{Z}, \sharp)$ and $(Y, \nu) = (\mathbb{R}, |\cdot|)$, and set $\tau(x) = [|x|^{p/q}]$, where $[\cdot]$ is the Gauss symbol. Note that for each $k \in \{0\} \cup \mathbb{N}$,

$$\tau^{-1}(\{k\}) = \left(-(k+1)^{\frac{q}{p}}, -k^{\frac{q}{p}}\right] \cup \left[k^{\frac{q}{p}}, (k+1)^{\frac{q}{p}}\right).$$

(1) Let $E = \{n, n+1, \dots, m-1\} \subset \{0\} \cup \mathbb{N} \ (n < m),$

$$\sharp E = m - n, \quad |\tau^{-1}(E)|^{\frac{p}{q}} \le \left\{ 2\sum_{j=n}^{m-1} \left[(j+1)^{\frac{q}{p}} - j^{\frac{q}{p}} \right] \right\}^{\frac{p}{q}} = \left\{ 2\left(m^{\frac{q}{p}} - n^{\frac{q}{p}}\right) \right\}^{\frac{p}{q}}$$

Then, $p \ge q$ implies

$$\left(m^{\frac{q}{p}} - n^{\frac{q}{p}}\right)^{\frac{p}{q}} \le m - n,$$

and then

$$|\tau^{-1}(E)|^{\frac{p}{q}} \lesssim \sharp E.$$

But, when p < q, this estimate fails. In fact, when m = n + 1, there exists $\theta \in (0, 1)$ such that

$$\frac{1}{2} \left(\frac{|\tau^{-1}(E)|^{\frac{p}{q}}}{\sharp E} \right)^{\frac{1}{p}} = (n+1)^{\frac{q}{p}} - n^{\frac{q}{p}} = \frac{q}{p} (n+\theta)^{\frac{q}{p}-1},$$

and then,

$$\lim_{n \to \infty} \frac{|\tau^{-1}(E)|^{\frac{p}{q}}}{\sharp E} = \infty.$$

(2) Assume that $p \ge q$, and let E be a general finite subset of Z. Then, by the convexity of $t \mapsto t^{p/q}$,

$$|\tau^{-1}(E)| \le 2|\tau^{-1}(\{0, 1, 2, \dots, k-1\})|$$

holds, where $k = \sharp E$. It follows from (1) as n = 0 and m = k that

a

$$|\tau^{-1}(E)|^{\frac{p}{q}} \lesssim \sharp E$$

can be obtained, and therefore, the composition operator C_{τ} is bounded from $L^{p,r}(\mathbb{Z})$ to $L^{q,r}(\mathbb{R})$. Here, $L^{p,r}(\mathbb{Z})$ stands for the Lorentz sequence $\ell^{p,r}(\mathbb{Z})$.

(3) When p > q, the boundedness from the Lebesgue sequence space $L^p(\mathbb{Z}) = \ell^p(\mathbb{Z})$ to $L^q(\mathbb{R})$ fails. In fact, choosing

$$f(n) = (1 + |n|)^{-\frac{1}{p}} (\log(3 + |n|))^{-\frac{1}{q}}, \quad n \in \mathbb{Z},$$

we have

$$|C_{\tau}f(x)| = \left(1 + \left[|x|^{\frac{p}{q}}\right]\right)^{-\frac{1}{p}} \left(\log\left(3 + \left[|x|^{\frac{p}{q}}\right]\right)\right)^{-\frac{1}{q}}$$
$$\geq (1 + |x|)^{-\frac{1}{q}} (\log(3 + |x|))^{-\frac{1}{q}} \times \left(\frac{p}{q}\right)^{-\frac{1}{q}}.$$

Then, $f \in \ell^p(\mathbb{Z})$ but $C_{\tau}f \notin L^q(\mathbb{R})$. Here, see [8, Excercise 1.4.8] for details of the example f(n).

(4) Let $K = \{1, 2, ..., k\}$. Then, the estimate

$$|\tau^{-1}(E)|^{\frac{p}{q}} \lesssim \sharp E$$

always holds for any $E \subset K$. Especially, since $\ell^p(K) \cong \mathbb{C}^k$ independent of p, the boundedness of the composition operator $C_\tau : \mathbb{C}^k \to L^q(\mathbb{R})$ can be obtained.

Example 4.3. Let $p \ge 1$, $\Phi(t) = t \log(3 + t)$, and let $\tau : \mathbb{R} \to \mathbb{Z}$ be

$$\tau(x) = \left[\left\{ \Phi^{-1}\left(\frac{1}{|x|}\right) \right\}^{-p} \right], \quad x \in \mathbb{R}.$$

Since the volume estimate

$$|\tau^{-1}(E)| \lesssim \left\{ \Phi\left(\frac{1}{(\sharp E)^{\frac{1}{p}}}\right) \right\}^{-1}$$

for each $E \subset \mathbb{Z}$, therefore Theorem 1.5 the composition operator C_{τ} is bounded from $\ell^{p,1}(\mathbb{Z})$ to $L \log L(\mathbb{R})$. Note that for each $k \in \{0\} \cup \mathbb{N}$,

$$\tau^{-1}(\{k\}) = \left(-\left\{ \Phi\left(\frac{1}{(k+1)^{\frac{1}{p}}}\right) \right\}^{-1}, -\left\{ \Phi\left(\frac{1}{k^{\frac{1}{p}}}\right) \right\}^{-1} \right] \cup \left[\left\{ \Phi\left(\frac{1}{k^{\frac{1}{p}}}\right) \right\}^{-1}, \left\{ \Phi\left(\frac{1}{(k+1)^{\frac{1}{p}}}\right) \right\}^{-1} \right) \right]$$

(1) Let $E = \{n, n + 1, ..., m - 1\} \subset \mathbb{N}$ (n < m). Note that

$$\sharp E = m - n,$$

$$\begin{aligned} |\tau^{-1}(E)| &= 2\sum_{j=n}^{m} \left[\left\{ \Phi\left(\frac{1}{(k+1)^{\frac{1}{p}}}\right) \right\}^{-1} - \left\{ \Phi\left(\frac{1}{k^{\frac{1}{p}}}\right) \right\}^{-1} \right] \\ &= 2\left[\left\{ \Phi\left(\frac{1}{m^{\frac{1}{p}}}\right) \right\}^{-1} - \left\{ \Phi\left(\frac{1}{n^{\frac{1}{p}}}\right) \right\}^{-1} \right]. \end{aligned}$$

(2) Let *E* be a general finite subset of \mathbb{Z} . Then, assuming that the mapping $t \mapsto \{\Phi^{-1}(1/t)\}^{-p}$ is equivalent to some convex function at t > 1, we have

$$|\tau^{-1}(E)| \lesssim |\tau^{-1}(\{n, n+1, \dots, m-1\})|$$

holds, where the natural numbers 1 < n < m are taken by $n = \min E$ and $\sharp E = m - n$. It follows from (1) that

$$|\tau^{-1}(E)| \lesssim \left\{ \Phi\left(\frac{1}{(k+1)^{\frac{1}{p}}}\right) \right\}^{-1} \lesssim \left\{ \Phi\left(\frac{1}{(\sharp E)^{\frac{1}{p}}}\right) \right\}^{-1}$$

can be obtained, and therefore, the composition operator C_{τ} is bounded from $\ell^{p,1}(\mathbb{Z})$ to $L \log L(\mathbb{R})$.

(3) Let us confirm that the mapping $t \mapsto \{\Phi^{-1}(1/t)\}^{-p}$ is equivalent to the convex function $t \mapsto t^p$ on $(1, \infty)$, that is, for all t > 1,

$$\left\{\Phi^{-1}\left(\frac{1}{t}\right)\right\}^{-p} \sim t^p$$

holds. It is well known by O'Neil [12] that for all t > 0,

$$t \le \Psi^{-1}(t)\widetilde{\Psi}^{-1}(t) \le 2t,$$

where $\widetilde{\Psi}$ is a complementary function of the Young function Ψ , that is,

$$\Psi(t) \equiv \sup\{st - \Psi(s) : s \ge 0\}, \quad t \ge 0,$$

and Ψ^{-1} is a generalized inverse function of the nondecreasing function, that is,

$$\Psi^{-1}(t) \equiv \inf\{s \ge 0 : \Psi(s) > t\}, \quad t \ge 0.$$

Additionally, the complementary function $\widetilde{\Phi}$ is provided by the first, third authors and Ono [9, Appendix B] as follows:

$$\widetilde{\Phi}(t) \sim \begin{cases} 0, & 0 \le t \le \log 3, \\ (t - \log 3)^2, & \log 3 < t \le 3, \\ e^t, & t > 3. \end{cases}$$

Then,

$$\widetilde{\Phi}^{-1}(t) \sim \begin{cases} \sqrt{t} + \log 3, & 0 < t < (3 - \log 3)^2, \\ \log t, & t \ge (3 - \log 3)^2. \end{cases}$$

It follows that for each t > 1,

$$\left\{\Phi^{-1}\left(\frac{1}{t}\right)\right\}^{-p} \sim \left\{t\widetilde{\Phi}^{-1}\left(\frac{1}{t}\right)\right\}^p \sim (\sqrt{t} + t\log 3)^p \sim t^p,$$

and then t > 1 implies that the mapping $t \mapsto \{\Phi^{-1}(1/t)\}^{-p}$ is equivalent to the convex function $t \mapsto t^p$.

Example 4.4. Let $p \ge 1$, and let $\tau : \mathbb{R} \to \mathbb{Z}$ be

$$\tau(x) = \left[\left\{ \log \left(1 + \frac{1}{|x|} \right) \right\}^{-p} \right], \quad x \in \mathbb{R}.$$

Then, the volume estimate

(4.1)
$$|\tau^{-1}(E)| \lesssim \left\{ \exp\left(\frac{1}{(\sharp E)^{\frac{1}{p}}}\right) - 1 \right\}^{-1}$$

holds for each $E \subset \mathbb{Z}$, and therefore Theorem 1.5 implies that the composition operator C_{τ} is bounded from $\ell^{p,1}(\mathbb{Z})$ to $\exp L(\mathbb{R})$. In fact, since the mapping $t \mapsto \{\log(1+1/t)\}^{-p}$ is equivalent to the convex function $t \mapsto t^p$ on $(1, \infty)$, it follows from the similar reason of the previous example that the equation (4.1) holds. That equivalency follows from the following argument: when t > 1, there exists $\theta = \theta(t) \in (0, 1)$ such that

$$\log\left(1+\frac{1}{t}\right) = \log(t+1) - \log t = \frac{1}{t+\theta} \sim \frac{1}{t},$$

and then,

$$\left\{ \log\left(1+\frac{1}{t}\right) \right\}^{-p} \sim t^p.$$

Acknowledgements. The authors are thankful to Professor Yoshihiro Sawano for his giving very helpful comments and advice. The first-named author (N.H.) was supported by Grant-in-Aid for Research Activity Start-up Grant Number 23K19013. The second author (M.I.) is supported by the Grant-in-Aid for Scientific Research (C) (No.23K03174), Japan Society for the Promotion of Science and JST CREST, No. JPMJCR1913.

References

- S. C. Arora, G. Datt, S. Verma, Composition operators on Lorentz spaces, Bull. Austral. Math. Soc. 76 (2007), no. 2, 205–214.
- [2] S. C. Arora, G. Datt and S. Verma, Operators on Lorentz sequences spaces Math. Bohem. 134 (2009), no. 1, 87–98.
- [3] T. Chawziuk, Y. Cui, Y. Estaremi, H. Hudzik and R. Kaczmarek, Composition and multiplication operators between Orlicz function spaces, J. Inequal. Appl. (2016), Paper No. 52, 18 pp.
- [4] Y. Cui, H. Hudzik, R. Kumar and L. Maligranda, Composition operators in Orlicz spaces, (English summary) J. Aust. Math. Soc. 76 (2004), no. 2, 189–206.
- [5] N. A. Evseev, Bounded composition operator on Lorentz spaces, Math. Notes 102 (2017), no. 5–6, 763–769.
- [6] N. Evseev and A. Menovschikov, Bounded operators on mixed norm Lebesgue spaces, Complex Anal. Oper. Theory 13 (2019), no. 5, 2239–2258.
- [7] N. A. Evseev and A. V. Menovshchikov, The composition operator on mixed-norm Lebesgue spaces, (Russian) Mat. Zametki 105 (2019), no. 6, 816–823; translation in Math. Notes 105 (2019), no. 5–6, 812–817.
- [8] L. Grafakos, Classical Fourier Analysis, Texts in Mathematics, Springer, New York, Third edition 249 (2014).
- [9] N. Hatano, R. Kawasumi and T. Ono, Predual of weak Orlicz spaces and its applications to Fefferman-Stein vector-valued maximal inequality, Tokyo J. Math. 46 (2023), no. 1, 125–160.
- [10] R. Kumar, Composition operators on Orlicz spaces, Integral Equations Operator Theory 29 (1997), no. 1, 17–22.
- [11] L.E. Labuschagne, and W.A. Majewski, Maps on noncommutative Orlicz spaces, Illinois Journal of Mathematics, 55(2011) no.3, 1053-1081.
- [12] R. O'Neil, Fractional Integration in Orlicz Space, Trans. Amer. Math. Soc., 115 (1965), 300–328.
- [13] K. Raj and S. K. Sharma, Composition operators on Musielak-Orlicz spaces of Bochner type, Math. Bohem. 137 (2012), no. 4, 449–457.
- [14] M. M. Rao and Z. D. Ren, Theory of Orlicz Spaces, Marcel Dekker, Inc., New York, Basel and Hong Kong, 1991.
- [15] R. K. Singh, Composition operators induced by rational functions, Proc. Amer. Math. Soc. 59 (1976), no. 2, 329–333.

(Naoya Hatano) Department of Department of Mathematics, Chuo University, 1-13-27, Kasuga, Bunkyo-ku, Tokyo 112-8551, Japan,

(Masahiro Ikeda) Center for Advanced Intelligence Project, RIKEN, Japan/Department of Mathematics, Faculty of Science and Technology, Keio University, 3-14-1 Hiyoshi, Kohoku-ku, Yokohama 223-8522, Japan,

(Ryota Kawasumi) Faculty of Business Administration, Kobe Gakuin University, 1-1-3 Minatojima, Chuo-ku, Kobe, Hyogo, 650-8586, Japan

Email address, Naoya Hatano: n.hatano.chuo@gmail.com

Email address, Masahiro Ikeda: masahiro.ikeda@riken.jp

Email address, Ryota Kawasumi: kawasumi@ba.kobegakuin.ac.jp