ON THE BOSONIZATION OF THE ENVELOPING ALGEBRA OF A FINITE DIMENSIONAL LIE SUPERALGEBRA

NICOLÁS ANDRUSKIEWITSCH AND KEN A. BROWN

ABSTRACT. We exhibit a PI Hopf algebra that is not a finite module over its center. We survey some ring-theoretical properties of the bosonizations of enveloping algebras of Lie superalgebras.

1. INTRODUCTION

It is well-known that a ring which is a finite module over its center satisfies a polynomial identity (it is PI, for short), cf. § 2.3. But the converse is false, even for affine prime Noetherian rings, see §4. Nevertheless, one might hope that the converse is true when one restricts attention to algebras which are not "too pathological" - for example a prime Noetherian PI-algebra which is a maximal order is always finite over its center, see [21, Propositions 13.9.8, 13.9.11].

There are various sorts of quantum groups that are PI because they are finite over their centers. Thus it is natural to consider the following questions, see [7, 12]:

Questions. Let H be a PI Hopf algebra. Under which conditions is H a finite module either

- (a) over its center $\mathcal{Z}(H)$, or equivalently over a central subalgebra?
- (b) Or over its Hopf center $\mathcal{HZ}(H)$, or equivalently over a central Hopf subalgebra?
- (c) Or at least over a normal commutative one?

Here the Hopf center is the largest central Hopf subalgebra. It was shown in [11] that being semiprime Noetherian is not enough to answer question (c) positively:

Theorem 1. [11] The bosonization $H(\mathfrak{gl}(1,1))$ of the enveloping algebra of the Lie superalgebra $\mathfrak{gl}(1,1)$ is a semiprime Noetherian PI Hopf algebra, but is not a finite module over any normal commutative Hopf subalgebra.

²⁰²⁰ Mathematics Subject Classification. Primary 16R20, Secondary 16T20, 17B35.

The work of N.A. was partially supported by CONICET (PIP 11220200102916CO), FONCyT-ANPCyT (PICT-2019-03660), by the Secyt (UNC) and by the International Center of Mathematics, Southern University of Science and Technology, Shenzhen.

The work of K.B. was supported by Leverhulme Emeritus Fellowship EM 2017-081.

Notice that the main result in [11] claims that $H(\mathfrak{gl}(1,1))$ is prime rather than semiprime, but this is not the case as we show in Lemma 5. Our main result answers the question (a) negatively for the same Hopf algebra.

Theorem 2. The bosonization $H(\mathfrak{gl}(1,1))$ is not a finite module over its center.

It is open whether such an example still exists if 'semiprime Noetherian' is strengthened to 'prime Noetherian' or even 'Noetherian domain'.

Clearly (b) implies (a), so we wonder whether the converse is true. This is indeed the case for enveloping algebras in positive characteristic by a classical Theorem of Jacobson. Also, this was asked in [9, Question 3.6] in the setting of Hopf algebras in positive characteristic that are iterated Ore extensions.

The paper is organized as follows: In Section 2 we collect basic facts on bosonizations of enveloping algebras of Lie superalgebras. Section 3 contains the proof of Theorem 2. Finiteness of semiprime Noetherian PI Hopf algebras over their centers is discussed in Section 4. Our reference for Hopf algebras is [24]; recent surveys on infinite-dimensional Hopf algebras are [1,8].

1.1. Notation and definitions. Let \Bbbk be an algebraically closed field of characteristic 0. All vector spaces, algebras, Lie algebras and so on are over \Bbbk .

Let $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$ be a finite dimensional Lie superalgebra with even part \mathfrak{g}_0 , odd part \mathfrak{g}_1 and superbracket [,]. Denote the enveloping algebra of \mathfrak{g} by $U(\mathfrak{g})$ and write $H(\mathfrak{g})$ for the bosonization of $U(\mathfrak{g})$ with the group algebra of the cyclic group C_2 . Thus $H(\mathfrak{g})$ is a smash product $U(\mathfrak{g})\#\Bbbk C_2$ where the generator t of C_2 acts on \mathfrak{g}_0 with eigenvalue 1 and on \mathfrak{g}_1 with eigenvalue -1; $H(\mathfrak{g})$ is a pointed Hopf algebra where the elements of \mathfrak{g}_0 are primitive, those of \mathfrak{g}_1 are (t, 1)-skew primitive and t is grouplike. Below |x| denotes the degree of a homogeneous element x of a C_2 -graded vector space.

1.2. Key example. Let $\mathfrak{g} = \mathfrak{gl}(1,1)$, that is, $\mathfrak{g} = M_2(\Bbbk)$, decomposed so that $\mathfrak{g}_0 = \Bbbk x \oplus \Bbbk y$, $\mathfrak{g}_1 = \Bbbk u \oplus \Bbbk v$, with the superbracket given by \mathfrak{g}_0 being abelian and

 $[u,v] = x, \quad [u,u] = [v,v] = 0, \quad [x,u] = [x,v] = 0, \quad [y,u] = u, \quad [y,v] = -v.$

Hence $U(\mathfrak{gl}(1,1))$ is presented by generators x, y, u, v with defining relations

x is central,
$$yu - uy = u$$
, $yv - vy = -v$,
 $u^2 = 0$, $v^2 = 0$, $uv + vu = x$. (1.1)

Then $H(\mathfrak{gl}(1,1)) = U(\mathfrak{gl}(1,1)) \# \mathbb{k}C_2$ is presented by generators x, y, u, v, t with relations (1.1) and

 $t^2 = 1,$ tx = xt, ty = yt, tu = -ut, tv = -vt.

In general, given $m, n \in \mathbb{N}$, $\mathfrak{gl}(m, n) \coloneqq M_{m+n}(\Bbbk)$, with elements viewed as block matrices $X = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$, where $A \in M_m(\Bbbk)$ and $D \in M_n(\Bbbk)$. Here $\mathfrak{g}_0 = \left\{ \begin{pmatrix} A & 0 \\ 0 & D \end{pmatrix} \right\}$, while $\mathfrak{g}_1 = \left\{ \begin{pmatrix} 0 & B \\ C & 0 \end{pmatrix} \right\}$. Thus in the above presentation of $\mathfrak{gl}(1, 1)$ the generators are $x = \mathrm{id} = e_{11} + e_{22}$, $y = e_{11}$, $u = e_{12}$ and $v = e_{21}$. For $X \in \mathfrak{gl}(m, n)(\Bbbk)$ as above, its supertrace is $\mathrm{str}(X) \coloneqq \mathrm{tr}(A) - \mathrm{tr}(D)$.

2. RING-THEORETICAL PROPERTIES

We review some properties satisfied by $H(\mathfrak{gl}(1,1))$, as consequences of more general statements. In this Section, \mathfrak{g} denotes a finite dimensional Lie superalgebra.

2.1. Noetherianity and Gelfand-Kirillov dimension. By definition, $U(\mathfrak{g})$ is affine (i.e. finitely generated as algebra). Now it follows from the PBW theorem for enveloping algebras of Lie superalgebras [23, 6.1.1] that $U(\mathfrak{g})$ is Noetherian; in particular, by [23, 6.1.3] $U(\mathfrak{g})$ is a finitely generated module over its subalgebra $U(\mathfrak{g}_0)$, which is Noetherian by the classical PBW theorem.

Similarly, $H(\mathfrak{g})$ is a finite right (and left) module over $U(\mathfrak{g})$, and so it is an affine Noetherian pointed Hopf algebra.

By [18, Prop. 5.5], if A is a subalgebra of an algebra B and B is a finitely generated right or left A-module, then $\operatorname{GKdim} A = \operatorname{GKdim} B$. By [18, Ex. 6.9],

$$\operatorname{GKdim} H(\mathfrak{g}) = \operatorname{GKdim} U(\mathfrak{g}) = \operatorname{GKdim} U(\mathfrak{g}_0) = \operatorname{dim} \mathfrak{g}_0.$$

2.2. **Primeness.** We start with a general fact. If y_1, \ldots, y_m is a basis of \mathfrak{g}_1 , set

$$D(\mathfrak{g}) \coloneqq \det([y_i, y_j])_{1 \le i, j \le m} \in S(\mathfrak{g}_0),$$

which does not depend on the basis up to a nonzero scalar multiple.

Theorem 3. [5], [23, 15.4.1]. If $D(g) \neq 0$, then U(g) is prime.

Sketch of the proof. Consider the Clifford filtration of $U(\mathfrak{g})$ where the elements of \mathfrak{g}_1 have degree 1, and those of \mathfrak{g}_0 have degree 2. In the associated graded algebra R, $U(\mathfrak{g}_0)$ is replaced by the symmetric algebra $S(\mathfrak{g}_0)$, which is central in R. Then R is prime provided that $D(\mathfrak{g}) \neq 0$. By a standard argument, so is $U(\mathfrak{g})$.

It is apparently not known whether primeness always fails when $D(\mathfrak{g}) = 0$.

Recall that \mathfrak{g} is *classical simple* if it is simple and \mathfrak{g}_0 acts semisimply on \mathfrak{g}_1 . Below, $\mathfrak{sl}(m,n)$ denotes the subalgebra of $\mathfrak{gl}(m,n)$ with supertrace 0; for the definition of $\mathfrak{d}(n)$, $(n \ge 2)$, see [5, §3.5].

Theorem 4. [5, Theorem 3.6], [23, 15.4.2]. Suppose that \mathfrak{g} is a direct sum of Lie superalgebras each of which either

(i) has zero odd part, or

(ii) is isomorphic to some $\mathfrak{gl}(m,n)$ or $\mathfrak{sl}(m,n)$ or $\mathfrak{d}(n)$, or

(iii) is a classical simple Lie superalgebra which is not of type $\mathfrak{p}(n)$.

Then $D(\mathfrak{g}) \neq 0$ and so $U(\mathfrak{g})$ is prime.

The enveloping algebras $U(\mathfrak{p}(n))$ are *not* prime, but do have a unique minimal prime ideal, see [17]. If \mathfrak{g} is a classical simple Lie superalgebra, then $U(\mathfrak{g})$ is a domain iff $\mathfrak{g} \simeq \mathfrak{osp}(1, 2n)$ [23, p. 349].

However the passage from $U(\mathfrak{g})$ to $H(\mathfrak{g})$ is not guaranteed. We observe that it is claimed in [11, 3.8] that $H(\mathfrak{gl}(1, 1))$ is prime, invoking [13, 2.3iii]. But this last result cannot be applied if the action of the grouplike is inner. In actual fact we have:

Lemma 5. $H(\mathfrak{gl}(1,1))$ is semiprime but not prime.

Proof. First note that $H(\mathfrak{gl}(1,1))$ is semiprime by [10, Theorem 7], since it is a skew group algebra $U(\mathfrak{gl}(1,1)) * \langle t \rangle$, where |t| = 2 and $U(\mathfrak{gl}(1,1))$ has no 2-torsion.

To see that primeness fails, set w := -x + 2uv. Then the following hold:

$$wx = xw, wy = yw, wu = -uw, wv = -vw, wt = tw,$$
 (2.1)

$$wt \in \mathcal{Z}\big(H(\mathfrak{gl}(1,1))\big),\tag{2.2}$$

$$(x - wt)(x + wt) = x^2 - w^2 = 0.$$
(2.3)

We prove (2.1): The first equality is evident since x is central. Now

$$wy = (-x + 2uv)y = -xy + 2u(yv + v) = -yx + 2(yu - u)v + 2uv = yw;$$

$$wu = (-x + 2uv)u = -xu + 2u(x - uv) = ux;$$

$$uw = u(-x + 2uv) = -ux = -wu;$$

$$wv = (-x + 2uv)v = -xv;$$

$$vw = v(-x + 2uv) = -xv + 2(x - uv)v = xv = -wv.$$

The proof of (2.2) and the first equality in (2.3) follow at once. Finally,

$$w^{2} = (-x + 2uv)(-x + 2uv) = x^{2} - 4xuv + 4uvuv$$
$$= x^{2} - 4xuv + 4(x - vu)uv = x^{2}.$$

Set $H = H(\mathfrak{gl}(1,1))$ and consider the left ideals I = H(x - wt), J = H(x + wt). By (2.2), (x - wt) and (x + wt) are central, hence I and J are ideals. Since IJ = 0 by (2.3), H is not prime.

Remark 6. The ideals I and J in the proof can in fact be shown to be prime, and are the only minimal prime ideals of $H = H(\mathfrak{gl}(1,1))$. Indeed, Since His a skew group algebra $U(\mathfrak{gl}(1,1)) * \langle t \rangle$, where |t| = 2, there are at most 2 minimal primes by [19]. 2.3. Polynomial identity. First, if a ring S is a finite module over a subring R which is PI, then S is also PI, see e.g. [21, Corollary 13.4.9] (and vice versa, as a subring of a PI ring is evidently PI). Since commutative algebras are PI, algebras that are finite modules over the center are PI. Second, the enveloping algebra of a Lie algebra \mathfrak{p} satisfies a PI if and only if \mathfrak{p} is abelian [20] (recall that char $\Bbbk = 0$). Indeed we have the following theorem.

Theorem 7. [4] The enveloping algebra $U(\mathfrak{g})$ is PI if and only if \mathfrak{g}_0 is abelian.

Corollary 8. $H(\mathfrak{g})$ is PI if and only if \mathfrak{g}_0 is abelian.

This reduces drastically the chances that $H(\mathfrak{g})$ is prime and PI simultaneously.

3. Centers

3.1. Center of $U(\mathfrak{g})$. We shall use the following result.

Lemma 9. [21, 13.6.14] Let R be a semiprime PI ring. The following are equivalent:

- the center $\mathcal{Z}(R)$ is Noetherian;
- R is right Noetherian and a finitely generated $\mathcal{Z}(R)$ -module.

In other words, if R is a semiprime PI right Noetherian ring, then it is a finitely generated $\mathcal{Z}(R)$ -module iff $\mathcal{Z}(R)$ is Noetherian too.

To proceed with the center of $U(\mathfrak{gl}(1,1))$, we recall the generalization of the Harish-Chandra theorem on centers of enveloping algebras of semisimple Lie algebras to Lie superalgebras, see [23, Chapter 13]. Given a Lie superalgebra \mathfrak{g} , set

$$\mathcal{Z}(\mathfrak{g}) \coloneqq \mathcal{Z}(U(\mathfrak{g})).$$

A classical simple Lie superalgebra \mathfrak{g} is called *basic* if it admits an even nondegenerate \mathfrak{g} -invariant bilinear form, which is necessarily supersymmetric [23, 1.2.4]. Albeit $\mathfrak{gl}(m, n)$ is not simple, it bears a nondegenerate invariant supersymmetric even bilinear form, using the supertrace [23, Exercise 2.7.1].

Let \mathfrak{g} be either a basic classical simple Lie superalgebra or $\mathfrak{gl}(m, n)$ where $m, n \in \mathbb{N}$. Let \mathfrak{h} be a Cartan subalgebra of \mathfrak{g} ; e.g., the space of diagonal matrices when $\mathfrak{g} = \mathfrak{gl}(m, n)$. Let $S(\mathfrak{h})^W$ be subalgebra of the symmetric algebra $S(\mathfrak{h})$ of W-invariant functions on \mathfrak{h}^* where W is the Weyl group.

Now let $I(\mathfrak{h})$ be the subalgebra of $S(\mathfrak{h})^W$ consisting of those ϕ such that if α is an isotropic root and $\lambda \in \mathbb{H}_{\alpha}$, where \mathbb{H}_{α} the hyperplane orthogonal to α , then

$$\phi(\lambda) = \phi(\lambda + \ell \alpha), \qquad \text{for all } \ell \in \mathbb{k}.$$

Here is the super version of the theorem of Harish-Chandra.

Theorem 10. Assume that \mathfrak{g} is either a basic classical simple Lie superalgebra of type different from A or that $\mathfrak{g} = \mathfrak{gl}(m, n)$ for some positive integers m and n.

- (i) [16] There is an algebra monomorphism ψ from $\mathcal{Z}(\mathfrak{g})$ into $S(\mathfrak{h})^W$.
- (ii) [15, 16, 25] The image of the map ψ is $I(\mathfrak{h})$.

A detailed exposition of the proof is given in $[23, \S 13.2]$.

Corollary 11. [22, Theorem 2.8], [23, 13.2.11]. For \mathfrak{g} as in Theorem 10, $\mathcal{Z}(\mathfrak{g})$ is Noetherian if and only if $\mathfrak{g} = \mathfrak{osp}(1, 2n)$ for $n \in \mathbb{N}$.

Combined with Lemma 9, this last result gives:

Corollary 12. $U(\mathfrak{gl}(1,1))$ is not finitely generated as a module over its center.

Proof. The Noetherian algebra $U(\mathfrak{gl}(1,1))$ is prime and PI by Theorems 4 and 7. Since $\mathcal{Z}(\mathfrak{gl}(1,1))$ is not Noetherian by Corollary 11, Lemma 9 applies.

Remark 13. It is well known that $\mathcal{Z}(U(\mathfrak{gl}(1,1)))$ is isomorphic to the subalgebra of the polynomial algebra $\Bbbk[x,y]$ of polynomials constant on the line x = y, which is generated by $x^n - y^n$ for all positive integers n. See [25, 0.6.1] taking into account [23, Cor. 13.3.8].

3.2. Center of $H(\mathfrak{g})$. Let \mathfrak{g} be a Lie superalgebra, not necessarily finite dimensional. Note that $H(\mathfrak{g})$ is C_2 -graded if we set the degree of t to be 0, that is $H(\mathfrak{g}) = H(\mathfrak{g})_0 \oplus H(\mathfrak{g})_1$ where

$$H(\mathfrak{g})_0 = U(\mathfrak{g})_0 \# \mathbb{k}[t], \qquad \qquad H(\mathfrak{g})_1 = U(\mathfrak{g})_1 \# \mathbb{k}[t].$$

Hence the center $\mathcal{Z}(H(\mathfrak{g}))$ of $H(\mathfrak{g})$ is also C_2 -graded,

$$\mathcal{Z}(H(\mathfrak{g})) = \mathcal{Z}(H(\mathfrak{g}))_0 \oplus \mathcal{Z}(H(\mathfrak{g}))_1,$$

where

$$\mathcal{Z}(H(\mathfrak{g}))_0 = \mathcal{Z}(H(\mathfrak{g})) \cap U(\mathfrak{g})_0 \# \Bbbk[t], \quad \mathcal{Z}(H(\mathfrak{g}))_1 = \mathcal{Z}(H(\mathfrak{g})) \cap U(\mathfrak{g})_1 \# \Bbbk[t].$$

To describe the center of $H(\mathfrak{g})$ we need the following definitions. Let M be a graded $U(\mathfrak{g})$ -bimodule. Recall that the adjoint action of \mathfrak{g} on M is given by

$$\mathrm{ad}(u)(m) \coloneqq um - (-1)^{|u||m|} mu,$$

for homogeneous elements $u \in \mathfrak{g}$ and $m \in M$. The submodule of \mathfrak{g} -invariants (with respect to the adjoint action) is

$$M^{\epsilon} \coloneqq \{ m \in M : \mathrm{ad}(u)(m) = 0 \text{ for all } u \in \mathfrak{g} \}.$$

Let $M \mapsto \Pi(M)$ be the parity functor, that interchanges even and odd components. The *twisted adjoint action* ad' of \mathfrak{g} on M is the adjoint action on $\Pi(M)$ [3,14], i.e.,

$$ad'(u)(m) \coloneqq um - (-1)^{|u|(|m|+1)}mu$$

for homogeneous $u \in \mathfrak{g}$ and $m \in M$. Consider now $U(\mathfrak{g})$ as a bimodule over itself.

Definition 14. [14, § 2.1] The anticenter $A(\mathfrak{g}) \coloneqq \Pi(U(\mathfrak{g}))^{\varepsilon}$ is the submodule of \mathfrak{g} -invariants of $U(\mathfrak{g})$ with respect to the twisted adjoint action ad', that is

$$A(\mathfrak{g}) = \{ x \in U(\mathfrak{g}) : \mathrm{ad}'(u)(x) = 0 \text{ for all } u \in \mathfrak{g} \}.$$

Clearly, $A(\mathfrak{g})$ is a graded submodule of $U(\mathfrak{g})$ with respect to ad'.

The following facts are recorded in [14]. Notice that the meaning of $\mathcal{Z}(\mathfrak{g})$ in [14] is different than here: it is the super center, i.e., $U(\mathfrak{g})^{\varepsilon}$ in the notation above, not the center of the associative algebra $U(\mathfrak{g})$ as in the present article. See [14, §3.5].

Lemma 15. (i) $A(\mathfrak{g})_1 = \mathcal{Z}(\mathfrak{g})_1$.

(ii) $x \in U(\mathfrak{g})_0$ belongs to $A(\mathfrak{g})_0$ if and only if

 $xu_0 = u_0 x$ and $xu_1 = -u_1 x$, for all $u_0 \in \mathfrak{g}_0$, $u_1 \in \mathfrak{g}_1$. (3.1)

- (iii) $A(\mathfrak{g})$ is a $\mathcal{Z}(\mathfrak{g})_0$ -submodule of $U(\mathfrak{g})$.
- (iv) If $x, y \in A(\mathfrak{g})_0$, then $xy \in \mathcal{Z}(\mathfrak{g})_0$.
- (v) If dim \mathfrak{g}_1 is even, then $A(\mathfrak{g}) = A(\mathfrak{g})_0$ and $\mathcal{Z}(\mathfrak{g}) = \mathcal{Z}(\mathfrak{g})_0$.

Proof. (i) to (iv) are straightforward. The first equality in (v) is proved in [14, Cor. 3.1.3]; this says that $0 = A(\mathfrak{g})_1 = \mathcal{Z}(\mathfrak{g})_1$ by(i), and so $\mathcal{Z}(\mathfrak{g}) = \mathcal{Z}(\mathfrak{g})_0$.

Now we move on to consider the bosonization $H(\mathfrak{g})$ of $U(\mathfrak{g})$.

Lemma 16. We have $\mathcal{Z}(H(\mathfrak{g}))_1 = 0$, and hence

$$\mathcal{Z}(H(\mathfrak{g})) = \mathcal{Z}(\mathfrak{g})_0 \oplus A(\mathfrak{g})_0 t.$$
(3.2)

Proof. Let $x \in \mathcal{Z}(H(\mathfrak{g}))_1$. Then tx = xt by centrality but tx = -xt, so x = 0. Let $x = a + bt \in H(\mathfrak{g})_0$, where $a, b \in U(\mathfrak{g})_0$. Given $u_0 \in \mathfrak{g}_0$ and $u_1 \in \mathfrak{g}_1$, we have

$$u_0x = xu_0 \iff u_0(a+bt) = (a+bt)u_0 \iff u_0a = au_0 \text{ and } u_0b = bu_0;$$

 $u_1x = xu_1 \iff u_1(a+bt) = (a+bt)u_1 \iff u_1a = au_1 \text{ and } u_1b = -bu_1.$
Since x commutes with t, we see that $x \in \mathcal{Z}(H(\mathfrak{g}))$ iff $a \in \mathcal{Z}(\mathfrak{g})_0$ and $b \in A(\mathfrak{g})_0.$

We can now deduce that the centers of the bosonizations of many enveloping algebras of Lie superalgebras are not Noetherian:

Proposition 17. Let \mathfrak{g} be as in Theorem 10, but $\mathfrak{g} \not\simeq \mathfrak{osp}(1, 2n)$ for any $n \in \mathbb{N}$. Assume that dim \mathfrak{g}_1 is even. Then $\mathcal{Z}(H(\mathfrak{g}))$ is not Noetherian.

Proof. By Lemmas 15 (v) and 16, as we assume that $\dim \mathfrak{g}_1$ is even, we have

$$\mathcal{Z}(H(\mathfrak{g})) = \mathcal{Z}(\mathfrak{g})_0 \oplus A(\mathfrak{g})_0 t = \mathcal{Z}(\mathfrak{g}) \oplus A(\mathfrak{g}) t, \qquad (3.3)$$

which is a decomposition of $\mathcal{Z}(\mathfrak{g})$ -modules by Lemma 15 (iii). Also $A(\mathfrak{g}) t A(\mathfrak{g}) t \subset \mathcal{Z}(\mathfrak{g})$ by Lemma 15 (iv). Since $\mathcal{Z}(\mathfrak{g})$ is not Noetherian by Corollary 11, we conclude that $\mathcal{Z}(H(\mathfrak{g}))$ is not Noetherian by Lemma 18 below.

For completeness, we give a proof of the following result.

Lemma 18. Let S be a subring of a Noetherian commutative ring R that admits a S-submodule T such that $R = S \oplus T$ and $T \cdot T \subset S$. Then S is Noetherian.

Proof. If J is an ideal in S, then $I = J \oplus JT$ is an ideal of R:

$$RI = (S+T)(J+JT) = SJ + SJT + JT + TJT \subset J + JT.$$

Clearly $I \cap S = J$. Given an ascending chain \mathfrak{C} of ideals in S, we get in this way an ascending chain of ideals in R that stabilizes because R is Noetherian. Intersecting with S, we see that \mathfrak{C} stabilizes, so S is Noetherian.

We can now prove Theorem 2, namely that $H(\mathfrak{gl}(1,1))$ is not finitely generated as a module over its center.

PROOF OF THEOREM 2. The Noetherian algebra $H(\mathfrak{gl}(1,1))$ is semiprime by Lemma 5 and PI by Corollary 8. Since the center $\mathcal{Z}(H(\mathfrak{gl}(1,1)))$ is not Noetherian by Proposition 17, Lemma 9 applies.

In fact, an explicit description of $\mathcal{Z}(H(\mathfrak{gl}(1,1)))$ follows from Proposition 17:

Theorem 19. (i) $\mathcal{Z}(H(\mathfrak{gl}(1,1))) = \mathcal{Z}(\mathfrak{gl}(1,1)) \oplus A(\mathfrak{gl}(1,1)) t$.

(ii) Let τ be the algebra automorphism of $\Bbbk[x, y]$ defined by $\tau(x) = x$ and $\tau(y) = y - 1$. Then

 $A(\mathfrak{gl}(1,1)) = \{ x\omega - (\omega + \tau(\omega))uv : \omega \in \Bbbk[x,y] \}.$

Proof. (i) Since dim $\mathfrak{gl}(1,1)$ is even, this follows from (3.3).

(ii) Let $\alpha \in A(\mathfrak{gl}(1,1)) = A(\mathfrak{gl}(1,1))_0 \subseteq U(\mathfrak{gl}(1,1))_0 = \Bbbk[x,y] \oplus \Bbbk[x,y]uv$, so that $\alpha = r(x,y) + s(x,y)uv$ for unique elements r = r(x,y), $s = s(x,y) \in \Bbbk[x,y]$. Since $U(\mathfrak{gl}(1,1))_0$ is commutative we see from (3.1) that

$$\alpha \in A(\mathfrak{gl}(1,1)) \iff \alpha u = -u\alpha \ and \ \alpha v = -v\alpha. \tag{3.4}$$

By (1.1), we have that yu = u(y+1), yv = v(y-1), hence

 $f(x,y)u = uf(x,y+1), \quad f(x,y)v = vf(x,y-1), \text{ for any } f(x,y) \in \mathbb{k}[x,y].$ Therefore, since uvu = ux and vuv = vx, we see that

$$\begin{aligned} &\alpha u = (r(x,y) + s(x,y)uv) \, u = u \left(r(x,y+1) + xs(x,y+1) \right); \\ &u\alpha = u \left(r(x,y) + s(x,y)uv \right) = ur(x,y); \\ &\alpha v = (r(x,y) + s(x,y)uv) \, v = vr(x,y-1); \\ &v\alpha = v \left(r(x,y) + xs(x,y) \right). \end{aligned}$$

Hence both equalities in the right hand side of 3.4 translate to the same condition r(x, y) + xs(x, y) = -r(x, y - 1), that is

$$\alpha \in A(\mathfrak{gl}(1,1)) \iff r + \tau(r) = -xs. \tag{3.5}$$

Writing $r = \sum_{0 \le j \le m} r_j x^j$ for unique $r_j \in \mathbb{k}[y]$ and substituting in (3.5), we find that

$$r_0 + \tau(r_0) = 0; (3.6)$$

considering the highest degree terms in y in (3.6), we see that $r_0 = 0$. Equivalently, $r = x\omega$ for $\omega \in \mathbb{k}[x, y]$. Substituting in (3.5) we get that $s = -\omega - \tau(\omega)$ for $\omega \in \mathbb{k}[x, y]$. Since all the steps can be reversed, the proof is complete.

4. ON NOETHERIAN PI HOPF ALGEBRAS FINITE OVER THEIR CENTERS

As is known, examples of Noetherian prime affine PI algebras that are *not* finite over their center are scarce. Here is an example: it is shown in [21, §13.10.2] that a subalgebra S of the algebra R introduced in [21, §5.3.7 (iii)] is affine Noetherian prime PI, but $\mathcal{Z}(S)$ is not Noetherian. Thus S is not finite over $\mathcal{Z}(S)$ by Lemma 9.

On the positive side, here is a criterion, obtained by combining a couple of deep results, that apparently has not been recorded elsewhere.

Theorem 20. Let H be an affine Noetherian Hopf algebra satisfying a polynomial identity. Suppose that the global homological dimension of H is finite. Then H is a finite direct sum of prime rings, and is a finite module over its center.

Proof. By [27, Theorems 0.1, 0.2], H is Auslander-Gorenstein and AS-Gorenstein. Thus, if gldim $H < \infty$, it is Auslander-regular and AS-regular. By [26, 1.4], H is a finite direct sum of prime rings and is integral over its center. One easily sees that an affine ring R which is integral over $\mathcal{Z}(R)$ is a finite module over $\mathcal{Z}(R)$.

Remark 21. The results quoted from [27] are valid over all fields \Bbbk , and the hypothesis that H is affine can be weakened to "all simple H-modules are finite dimensional over \Bbbk ".

Remark 22. Note that it follows immediately from Theorems 2 and 20 that

$$\operatorname{gldim} H(\mathfrak{gl}(1,1)) = \infty. \tag{4.1}$$

But in fact it is easy to see this directly: for consider the Hopf subalgebra $S := \Bbbk \langle u, t \rangle$ of $H(\mathfrak{gl}(1, 1))$, which is isomorphic to the Sweedler Hopf algebra. Clearly gldim $S = \infty$ and $H(\mathfrak{gl}(1, 1))$ is free over S by the PBW theorem. By a standard argument, see e.g. [2, Lemma 4.25], (4.1) follows. Notice that S could be replaced in the above argument by the subalgebra $R := \Bbbk \langle u \rangle$.

Remark 23. Returning to an issue raised in §1, namely whether an affine Noetherian Hopf algebra H which is a domain and satisfies a PI has to be finite over $\mathcal{Z}(H)$, we note that this cannot simply be proved by showing that gldim H is finite and applying Theorem 20. For the affine Hopf domains $B(n, \{p_i\}_{i=1}^s, q, \{\alpha_i\}_{i=1}^s)$ constructed in [28] are Noetherian, finite modules over their centers but have finite global dimension only for some very special values of the parameters.

Remark 24. Taking further a comment made in the opening paragraph of §1, it would be interesting to look for conditions ensuring that a prime Noetherian Hopf algebra is a maximal order. For example, filtered Noetherian domains whose associated graded rings are maximal orders inherit the same property [21, Theorem 5.1.6]. It may also be more than a coincidence that, in the characterisation [6, Theorem F] of those group algebras of polycyclicby-finite groups which are maximal orders, the prime 2 is key, as is the case in the Lie superalgebra setting.

Acknowledgements. We thank Ken Goodearl, Ed Letzter, Ian Musson and Vera Serganova for providing useful information. N.A. thanks Efim Zelmanov and Slava Futorny for their warm hospitality durin his visit to the SUSTech.

References

- N. Andruskiewitsch, On infinite-dimensional Hopf algebras, Preprint, arXiv:2308.13120 [math.QA] (2023); to appear in the Proceedings of the ICRA 2022.
- [2] N. Andruskiewitsch, S. Natale, and B. Torrecillas, A class of finite-by-cocommutative Hopf algebras, Annals of Representation Theory 1, pp. 73-94 (2024).
- [3] D. Arnaudon, M. Bauer, and L. Frappat, On Casimir's ghost, Commun. Math. Phys. 187, 429–439 (1997).
- [4] Yu. A. Bakhturin, Identities of a universal enveloping algebra for a Lie superalgebra, Mat. Sb., Nov. Ser. 127(169), No. 3(7), 384–397 (1985).
- [5] A. D. Bell, A criterion for primeness of enveloping algebras of Lie superalgebras, J. Pure Appl. Algebra 69, No. 2, 111–120 (1990)
- [6] K. A. Brown, *Height one primes of polycyclic group rings*, J. London Math. Soc. (2) 32, 426–438 (1985).
- [7] K. A. Brown. Representation theory of Noetherian Hopf algebras satisfying a polynomial identity, Contemp. Math. 229, pp. 49–79 (1998).
- [8] K. A. Brown and J. J. Zhang, Survey on Hopf algebras of GK-dimension 1 and 2, Contemp. Math. 771, pp. 43–62 (2021).
- K. A. Brown and J. J. Zhang, Iterated Hopf Ore extensions in positive characteristic, J. Noncommut. Geom. 16, 787–837 (2022).
- [10] J.W. Fisher and S. Montgomery, Semiprime skew group rings, J.Algebra 52, 241-247 (1978).
- [11] S. Gelaki and E. S. Letzter, An affine PI Hopf algebra not finite over a normal commutative Hopf subalgebra, Proc. Amer. Math. Soc. 131, 2673–2679 (2003).
- [12] K. R. Goodearl, Noetherian Hopf algebras, Glasg. Math. J. 55A, 75–87 (2013).
- [13] K. R. Goodearl and E. S. Letzter, Prime ideals in skew and q-skew polynomial rings, Mem. Am. Math. Soc. 521, 106 p. (1994). Providence, RI: American Mathematical Society (AMS).

- [14] M. Gorelik. On the ghost centre of Lie superalgebras, Ann. Inst. Fourier 50, 1745– 1764 (2000).
- [15] M. Gorelik, The Kac construction of the centre of U(g) for Lie superalgebras, J. Nonlinear Math. Phys. 11, 325–349 (2004).
- [16] V. G. Kac, Laplace operators of infinite-dimensional Lie algebras and theta functions, Proc. Natl. Acad. Sci. USA 81, 645–647 (1984).
- [17] E. Kirkman and J. Kuzmanovich, Minimal prime ideals in enveloping algebras of Lie superalgebras, Proc. Am. Math. Soc. 124, 1693–1702 (1996).
- [18] G. R. Krause and T. H. Lenagan, Growth of algebras and Gelfand-Kirillov dimension, revised edn., Graduate Studies in Mathematics 22, Amer. Math. Soc., Providence, RI, 2000.
- M. Lorenz and D. S. Passman, Prime ideals in crossed products of finite groups, Isr. J. Math. 33, 89–132 (1979).
- [20] V. N. Latyshev, Two remarks on PI-algebras, Sib. Mat. Zh. 4, 1120–1121 (1963).
- [21] J. C. McConnell and J. C. Robson, *Noncommutative Noetherian rings*. Revised edn., Graduate Studies in Mathematics 30, American Mathematical Society, Providence, RI, 2001.
- [22] I. M. Musson. On the center of the enveloping algebra of a classical simple Lie superalgebra, J. Algebra 193, 75–101 (1997).
- [23] I. M. Musson. Lie superalgebras and enveloping algebras. Providence, RI: American Mathematical Society (AMS) (2012).
- [24] D. E. Radford. Hopf algebras. Series on Knots and Everything 49. Hackensack, NJ: World Scientific. xxii, 559 p. (2012).
- [25] A. Sergeev, The invariant polynomials on simple Lie superalgebras, Represent. Theory 3, 250–280 (1999).
- [26] J. T. Stafford and J. J. Zhang. Homological properties of (graded) Noetherian PI rings, J. Algebra 168, 988–1026 (1994).
- [27] Q.-S. Wu and J. J. Zhang, Noetherian PI Hopf algebras are Gorenstein, Trans. Amer. Math. Soc. 355,1043–1066 (2003).
- [28] D.-G. Wang, J.J. Zhang and G. Zhuang, Hopf algebras of GK-dimension two with vanishing Ext-group. J. Algebra 388 (2013), 219–247.

Facultad de Matemática, Astronomía y Física, Universidad Nacional de Córdoba. CIEM – CONICET. Medina Allende s/n (5000) Ciudad Universitaria, Córdoba, Argentina

 $Email \ address:$ nicolas.andruskiewitsch@unc.edu.ar

School of Mathematics and Statistics, University of Glasgow, Glasgow G12 8QW, Scotland

Email address: ken.brown@glasgow.ac.uk