

Degeneration in discriminantal arrangements

Takuya Saito*

May 6, 2024

Abstract

Discriminantal arrangements are hyperplane arrangements, which are generalized braid ones. They are constructed from given hyperplane arrangements, but their combinatorics are not invariant under combinatorial equivalence. However, it is known that the combinatorics of the discriminantal arrangement are constant on a Zariski open set of the space of hyperplane arrangements. In the present paper, we introduce non-very generic varieties in the space of hyperplane arrangements to classify discriminantal arrangements and show that the Zariski open set is the complement of non-very generic varieties. We study their basic properties and construction and provide examples, including infinite families of non-very generic varieties. In particular, the construction we call degeneration is a powerful tool for constructing non-very generic varieties. As an application, we provide lists of non-very generic varieties for spaces of small line arrangements.

1 Introduction

The discriminantal arrangement $\mathcal{B}(\mathcal{A})$ is a collection of hyperplanes in the space of parallel translations of a given arrangement \mathcal{A} . Manin and Schechtman introduced this as a generalized braid arrangement ([MS89]). The defined motivations for the discriminantal arrangement are the higher Bruhat order and higher braid groups. Originally, it was defined for generic arrangements, but Bayer–Brandt defined it for arbitrary arrangements, and hyperplanes in $\mathcal{B}(\mathcal{A})$ correspond to circuits of \mathcal{A} . It is also known as an arrangement that spans points at general positions, and from this point of view, the characteristic polynomial is calculated in [KNT12, NT12]. Applications of discriminantal arrangements include braided monoidal 2-categories [KV94], vanishing cohomology of toric varieties [Per11], and the maximum likelihood degree of very affine varieties of point configurations [ABF⁺23]. For a general theory of hyperplane arrangements, refer to [OT92].

Each point in the discriminantal arrangement $\mathcal{B}(\mathcal{A})$ corresponds to a parallel translation \mathcal{A}^t of an original arrangement \mathcal{A} . Several invariants of parallel translations in a discriminantal arrangement have been studied in [CFW21, FW23]. From the perspective of parallel translations, there is a toric version, namely, a discriminantal toric arrangement [Pag19].

The concept of discriminantal arrangements is equivalent to a circuit basis for a binary matroid by Longyear [Lon80], concurrence geometry, the geometry of circuits by Crapo [Cra84, Cra85], derived matroid by Oxley and Wang [OW19], derived code introduced to determine extended coset weight enumerators by Jurrius and Pellikaan [JP15]. A similar concept has been pointed out as secondary polytopes, or more generally, a fiber zonotope [BB97, ER96]. All of them are defined for representations of matroids. Here, representations of matroids are almost equivalent to hyperplane arrangements. Some similar combinatorial constructions are adjoint [Che74], which generalize these to geometric lattices without representations, and combinatorial derived matroids, which provide a purely combinatorial definition by [FHJK23]. However, these are not matroid invariants except for combinatorial derived matroid. In other words, they are constructions that cannot be uniquely determined from the given matroid. Alternatively, in terms of hyperplane arrangements, the constant combinatorics of discriminantal arrangement are not obtained even if the arrangements have the same combinatorics. In the context of discriminantal arrangement, Falk was the first to point this fact out in [Fal94]. Another fundamental example is by Crapo [Cra84] (see Example 2.1).

*Affiliations: Department of Mathematics, Faculty of Science, Hokkaido University, Sapporo 060-0810, Japan/ Department of Economics and Statistics, University of Turin, 10124 Turin, Italy. e-mail: takuya.saito@unito.it

2020 Mathematics Subject Classification: primary; 52C35, secondary; 52C30, 14N20, 05B35
Key words: hyperplane arrangement, intersection lattice, discriminantal arrangement

Therefore, one of the problems in the theory of discriminantal arrangements is determining the possible discriminantal arrangements $\mathcal{B}(\mathcal{A})$ for an arrangement \mathcal{A} with given combinatorics. Let $\mathbf{Arr}_{\mathbb{K}}(n, k)$ denote the set of arrangements consisting of n hyperplanes in \mathbb{K}^k . As is well known, this is equivalent to the quotient of the Grassmannian $\mathrm{Gr}_k(\mathbb{K}^n)$ by the maximal torus $(\mathbb{K}^\times)^n$. We focus on the subset $\mathbf{Arr}^\circ(n, k)$, consisting of generic arrangements in $\mathbf{Arr}(n, k)$. It is known that an arrangement on a Zariski open set in the realization space of a generic arrangement yields a discriminantal arrangement with constant combinatorics [Ath99]. Arrangements on such Zariski open set are called *very generic*, and arrangements that are not are called *non-very generic*.

Athanasiadis gives the combinatorial sufficient condition for parallel translation to be non-very generic, but there are examples in [SY22a] where the converse does not hold. Some other sufficient conditions for non-very genericity have been given in [SY22b, SY22a]. The set of non-very generic arrangements is an algebraic set; thus, it defines a stratification of $\mathbf{Arr}_{\mathbb{K}}(n, k)$ (or the Grassmannian $\mathrm{Gr}_k(\mathbb{K}^n)$), which is a refinement of matroid stratification (see [Fal94]). The Pappus hypersurfaces in $\mathrm{Gr}_3(\mathbb{K}^n)$ were introduced in [SSY17] to study and classify Falk's example. As a generalization of the Pappus hypersurface, we introduce the *non-very generic variety* $V_{(\mathbb{T}, r)}$ in $\mathbf{Arr}(n, k)$. We show that this definition is appropriate in Theorem 3.9 and Theorem 3.10; the set of non-very generic arrangements is the union of non-very generic varieties. In order to examine the stratification of $\mathbf{Arr}(n, k)$, which is defined by non-very generic varieties, we must classify and determine them. Some known simple descriptions of the stratification using an algebraic set in $\mathbf{Arr}^\circ(n, k)$ are in [SSY19], [SS22]. However, simple descriptions of the defining equations are generally unknown. We construct $2n$ -wheels as an infinite family of non-very generic variety and give their explicit equation in Theorem 4.1. Furthermore, we examine the behavior of non-very generic varieties for deleted/restricted arrangements (Theorem 5.6). We show that the very generic arrangements are closed under the operations deletion and restriction. In particular, deleting parallel elements, which we call *degeneration*, is a powerful tool for finding non-very generic arrangements. By applying this to $2n$ -wheels, we obtain another infinite family of non-very generic varieties, which we call $2n+2$ -ladders (Proposition 6.1). It is possible to determine all non-very generic varieties of 8 or fewer lines (Theorem 6.3). Other studies on non-very generic arrangements are [LS18, Yam23].

This paper is organized as follows. Section 2 includes the basic definitions of discriminantal arrangements and presents some examples of discriminantal arrangements with different combinatorics constructed from arrangements with the same combinatorics. In Section 3, we revisit the definition of non-very generic arrangements, translations, and intersections and introduce non-very generic varieties. We discuss some fundamental properties related to necessary and sufficient conditions for non-very generic arrangements. In Section 4, we construct an infinite family of non-very generic line arrangements called $2n$ -wheels and provide the defining equations of their non-very generic varieties. Moreover, we give a sufficient condition for the existence of (degenerated) $2n$ -wheels. In Section 5, we review the discriminantal arrangement for multiarrangements. Then, we discuss the deletion and restriction of arrangements, mainly focusing on the case of deleting parallel hyperplanes, which we refer to as degeneration. In Section 6, by applying the construction from Section 5 to $2n$ -wheels, we construct $2n+2$ -ladders. We will determine all the defining equations for a minimal non-very generic line arrangement with eight lines. Additionally, we discuss non-very generic line arrangements with nine lines.

Acknowledgment

This work was supported by JSPS KAKENHI Grant Number JP23KJ0031. The author wants to thank Mutsumi Saito, Simona Settepanella and So Yamagata for helpful comments.

2 Discriminantal arrangements

In this section, we give the definitions of parallel translations of the arrangement, discriminantal arrangements, and examples.

2.1 Notations and basics on hyperplane arrangements

Throughout this paper, \mathbb{K} is a commutative field, and W is a k -dimensional vector space over \mathbb{K} . We sometimes abbreviate the set $\{a, b, \dots, c\}$ as $ab\dots c$.

First, we recall the notions of hyperplane arrangements. Let $\mathcal{A} = \{H_1, \dots, H_n\}$ be a central hyperplane arrangement in W with normals $\alpha_i \in W^*$. The *combinatorics* of \mathcal{A} form a graded lattice of the intersections of hyperplanes in \mathcal{A} , ordered by reverse inclusion and denoted by $\mathcal{L}(\mathcal{A})$. The rank of $\mathcal{L}(\mathcal{A})$ is defined as $\mathrm{rank} X = k - \dim X$. A central arrangement \mathcal{A} is *generic* if $\mathrm{rank}(\bigcap_{i \in S} H_i) = \min\{|S|, k\}$ for each $S \subset [n] = \{1, \dots, n\}$. Two arrangements are *combinatorially equivalent* if their combinatorics are lattice isomorphic and *linearly equivalent* if a linear map exists between ambient spaces, inducing a bijection between these arrangements. For convenience, we require \mathcal{A} to be an *essential* arrangement, that is, $\dim \bigcap_{H \in \mathcal{A}} H = 0$. Until Section 4, we assume that \mathcal{A} is generic.

2.2 The space $\mathbb{S}(\mathcal{A})$ of parallel translations of \mathcal{A} and its subspaces

We define the space of parallel translations of an arrangement. There is a natural isomorphism $\alpha_i^{-1} : \mathbb{K} \rightarrow W/H_i; t_i \mapsto \alpha_i^{-1}(t_i)$, which induces the isomorphism from \mathbb{K}^n to the space $\mathbb{S}(\mathcal{A}) = \bigoplus_{i=1}^n W/H_i$ of parallel translations of \mathcal{A} . Define \mathcal{A}^t to be the image of $t = (t_i)_{i=1}^n \in \mathbb{K}^n$. Since $H_i^t = \alpha_i^{-1}(t_i)$ is a hyperplane in W , \mathcal{A}^t can be regarded as a hyperplane arrangement $\{H_i^t \mid i = 1, \dots, n\}$, which is a *parallel translation* of \mathcal{A} by t .

For each subset S of $[n] = \{1, \dots, n\}$, define the subspace $D_S = D_S^{\mathcal{A}}$ in $\mathbb{S}(\mathcal{A})$ as the set of translations \mathcal{A}^t whose intersection of hyperplanes indexed by S is not empty, i.e.,

$$D_S = \{\mathcal{A}^t \in \mathbb{S}(\mathcal{A}) \mid \bigcap_{i \in S} H_i^t \neq \emptyset\}.$$

The dimension of D_S is $\dim D_S = n - |S| + \text{rank}(\bigcap_{i \in S} H_i) = \min\{n, n+k - |S|\}$. Thus, D_S is a hyperplane in $\mathbb{S}(\mathcal{A})$ if and only if $|S| = k+1$. Let $\mathcal{C}(\mathcal{A})$ denote $\{C \subset [n] \mid |C| = k+1\}$, then D_C forms a hyperplane in $\mathbb{S}(\mathcal{A})$ with the normal $\alpha_C = \det \begin{pmatrix} e_{i_1}^* & \cdots & e_{i_{k+1}}^* \\ \alpha_{i_1} & \cdots & \alpha_{i_{k+1}} \end{pmatrix}$ for each $C = \{i_1, \dots, i_{k+1}\} \in \mathcal{C}(\mathcal{A})$. Here e_i 's, in which each e_i spans W/H_i for each $i \in [n]$, are the canonical basis of $\mathbb{S}(\mathcal{A})$, and e_i^* are the dual basis.

The *discriminantal arrangement* associated to \mathcal{A} is the hyperplane arrangement

$$\mathcal{B}(\mathcal{A}) = \{D_C \subset \mathbb{S}(\mathcal{A}) \mid C \in \mathcal{C}(\mathcal{A})\}.$$

If the original arrangements are linearly equivalent, then the discriminantal arrangements associated to them are also linearly equivalent. When $k = 1$, it is equivalent to the braid arrangement, which consists of $D_{ij} = \{(t_1, \dots, t_n) \in \mathbb{K}^n \mid t_i = t_j\}$ for all $1 \leq i < j \leq n$. The normals consist of non-zero scalars; thus, $\mathcal{C}(\mathcal{A}) = \binom{[n]}{2}$. The hyperplanes are $D_{ij} = \{\mathcal{A}^t \in \mathbb{S}(\mathcal{A}) \mid t_i \cap t_j \neq \emptyset\} = \{\mathcal{A}^t \in \mathbb{S}(\mathcal{A}) \mid t_i = t_j\}$.

Let us define a representative of an intersection $X \in \mathcal{L}(\mathcal{B}(\mathcal{A}))$. If $\mathcal{A}^{t_1}, \mathcal{A}^{t_2}$ are parallel translations of \mathcal{A} such that $\mathcal{L}(\mathcal{A}^{t_1}) \cong \mathcal{L}(\mathcal{A}^{t_2})$, then we have for any $S \in 2^{[n]}$, $\mathcal{A}^{t_1} \in D_S$ if and only if $\mathcal{A}^{t_2} \in D_S$. We call an element \mathcal{A}^t of X *representative* of X , and write $[\mathcal{A}^t]$ as X if and only if $\mathcal{A}^t \in X \setminus (\bigcup_{Y \subsetneq X} Y)$. If $\mathcal{A}^{t_1}, \mathcal{A}^{t_2}$ are both representatives of X then $\mathcal{L}(\mathcal{A}^{t_1}) \cong \mathcal{L}(\mathcal{A}^{t_2})$. We call D_S a *component* of X if $|S| \geq k+1$ and $X \not\subset D_{S \cup \{i\}}$ for all $i \in [n] \setminus S$. Define the *canonical presentation* of $X = [\mathcal{A}^t]$, denoted by $\mathbb{T}(X) = \mathbb{T}([\mathcal{A}^t])$, as the set consisting of all the components of X , i.e.

$$\mathbb{T}(X) = \{S \subset [n] \mid D_S \text{ is a component of } X\}$$

The following example is fundamental.

Example 2.1 (Crapo [Cra84]). Let $n = 6$ and $k = 2$, and let the coefficient field be the complex field \mathbb{C} . Take $\lambda \in \mathbb{C} \setminus \{0, 1/2, 1, 2\}$ and take the vectors

$$(\alpha_1 \alpha_2 \alpha_3 \alpha_4 \alpha_5 \alpha_6) = \begin{pmatrix} 1 & 2 & 1 & 1 & 0 & \lambda \\ 0 & 1 & 1 & 2 & 1 & 1 \end{pmatrix}.$$

Let \mathcal{A} be a generic arrangement with the normals $\{\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6\}$. We have $\mathcal{C}(\mathcal{A}) = \binom{[6]}{3} = \{S \subset [6] \mid |S| = 3\}$. Since \mathcal{A} is generic, $\mathcal{L}(\mathcal{A})$ is independent of a choice of λ . Now we consider the intersection of hyperplanes $D_{123}, D_{156}, D_{246}, D_{345}$ in $\mathcal{B}(\mathcal{A})$. Let X be $D_{123} \cap D_{156} \cap D_{246} \cap D_{345}$. We have

$$\begin{aligned} \alpha_{123} &= e_1^* - e_2^* + e_3^*, & \alpha_{156} &= -\lambda e_1^* - e_5^* + e_6^*, \\ \alpha_{246} &= (1-2\lambda)e_2^* + (\lambda-2)e_4^* + 3e_6^*, & \alpha_{345} &= e_3^* - e_4^* + e_5^*. \end{aligned}$$

Then

$$\text{rank } X = 6 - \dim \langle \alpha_{123}, \alpha_{156}, \alpha_{246}, \alpha_{345} \rangle = \begin{cases} 4 & \text{if } \lambda \neq -1, \\ 3 & \text{if } \lambda = -1. \end{cases}$$

Therefore, we get the representatives as in Figure 3 and the canonical presentations

$$\mathbb{T}(X) = \begin{cases} \{123456\} & \text{if } \lambda \neq -1, \\ \{123, 156, 246, 345\} & \text{if } \lambda = -1. \end{cases}$$

And we have a representative of X is $\mathcal{A}^{(0,0,0,0,0,0)} = \mathcal{A}^0$ if $\lambda \neq -1$, or $\mathcal{A}^{(0,1,1,1,0,0)}$ if $\lambda = -1$.




Figure 1: $\mathcal{A} = \mathcal{A}^0$ in Example 2.1




Figure 2: a representative of $D_{123} \cap D_{345}$




Figure 3: a representative of $D_{123} \cap D_{156} \cap D_{345}$

Geometrically, one of the representatives of $D_{123} \cap D_{345}$ is $\mathcal{A}^{(0,1,1,1,0,t_6)}$ for a generic t_6 (see Figure 2). Then we obtain

$$H_1^t \cap H_2^t \cap H_3^t = \{(0, 1)\}, H_3^t \cap H_4^t \cap H_5^t = \{(1, 0)\}, H_1^t \cap H_5^t = \{(0, 0)\}.$$

Move H_6^t through the intersection of H_1^t and H_5^t , then it should be $t_6 = 0$. This means that $\mathcal{A}^{(0,1,1,1,0,0)}$ is in D_{156} . Now $H_2^t \cap H_4^t = \{(1/2, 1/2)\}$, and H_6^t through the intersection if and only if $\lambda = -1$. In other words, $D_{123} \cap D_{156} \cap D_{345} \subset D_{246}$ if $\lambda = -1$. If $\lambda \neq -1$, then we have $D_{123} \cap D_{156} \cap D_{345} \not\subset D_{246}$. Then, we can get the rank of X . In some cases, it is possible to know the rank of the intersection by observing the translation of \mathcal{A} in this way.

More generally, in any commutative coefficient fields, a discriminantal arrangement associated to a line arrangement \mathcal{A} , consisting of 6 lines, have an intersection $D_{123} \cap D_{156} \cap D_{246} \cap D_{345}$ of rank 3 if and only if

$$\Delta_{16}\Delta_{24}\Delta_{35} - \Delta_{15}\Delta_{26}\Delta_{34} = 0 \quad (1)$$

where we use the symbol $\Delta_{ij} = \det(\alpha_i, \alpha_j)$. This equation is shown in [Cra84],[SS22], and we call a representative of a non-very generic intersection Crapo configuration, whose canonical presentation is \mathbb{T} or its permutation $\sigma.\mathbb{T} = \{\{\sigma.i_1, \sigma.i_2, \sigma.i_3\} \mid i_1 i_2 i_3 \in \mathbb{T}\}$ for $\sigma \in \mathfrak{S}_n$. Notice that the canonical representation combinatorially determines this condition. This example is one example of discriminantal arrangements not invariant under the combinatorial equivalence relation.

A phenomenon where intersections of hyperplanes in $\mathcal{B}(\mathcal{A})$ with the same indices have different ranks also occurs in other parameters n, k . The following example differs from the previous example in that the intersections of hyperplanes in $\mathcal{B}(\mathcal{A})$ with the same indices have different ranks, but the canonical presentation is the same. The next section will explain the main difference between Example 2.1 and Example 2.2.

Example 2.2 (Settepanella, Yamagata [SY22a, Example 4.3.]). Let $n = 10, k = 3$, and let the coefficient field be the complex field \mathbb{C} . And let $C_1 = \{1, 2, 3, 4\}, C_2 = \{1, 5, 6, 7\}, C_3 = \{2, 5, 8, 9\}, C_4 = \{3, 6, 8, 10\}$ and $C_5 = \{4, 7, 9, 10\}$ For a sufficiently generic arrangement \mathcal{A} , we get $\text{rank} \bigcap_{i=1}^5 D_{C_i} = 5$. However if we take normals α_i of H_i ($i = 1, \dots, 10$) as follows;

$$(\alpha_1, \dots, \alpha_{10}) = \begin{pmatrix} 0 & 20 & 2 & 3 & 0 & 1 & 1 & 4 & 314 & 139 \\ 10 & 0 & -3 & 1 & 0 & -1 & 2 & -1 & -40 & 30 \\ 3 & -9 & 0 & 0 & 1 & 1 & 2 & -3 & -197 & -43 \end{pmatrix},$$

then $\text{rank} \bigcap_{i=1}^5 D_{C_i} = 4$. The difference from the previous example is that the canonical presentation of intersection $X = \bigcap_{i=1}^5 D_{C_i}$ is the same $\{C_1, C_2, C_3, C_4, C_5\}$ in both cases of $\text{rank} X = 4$ or 5. Hence, this condition is not combinatorially determined.

3 Non-very generic arrangements

As we saw in the previous section, discriminantal arrangements with different combinatorics can be obtained from arrangements with the same combinatorics. We define classes of original arrangements based on this problem. First, Bayer–Brandt defined very generic arrangements for the real case. An arrangement \mathcal{A} consisting of n hyperplanes

in \mathbb{R}^k is *very generic* if the cardinality of $\mathcal{L}(\mathcal{B}(\mathcal{A}))$ is the largest possible for a discriminantal arrangement associated to an essential arrangement with n hyperplanes in \mathbb{R}^k .

If \mathcal{A} is generic, then each k element subset of $[n]$ is contained in at most one member of $\mathbb{T}([\mathcal{A}^t])$ for all translation \mathcal{A}^t of \mathcal{A} . In other words, if a family \mathbb{T} of subsets of $[n]$ is a canonical presentation of some intersection of a discriminantal arrangement, then \mathbb{T} satisfies the following conditions;

(Q0) $S_1 \not\subset S_2$ if $S_1, S_2 \in \mathbb{T}$,

(Q1) each element of \mathbb{T} has at least $k + 1$ element,

(Q2) each k element subset of $[n]$ is contained in at most one member of \mathbb{T} .

"In order to consider all possible combinatorics of parallel translations of the generic arrangement, we define the set $Q(n, k) = Q([n], k)$ to be a set \mathbb{T} consisting of all elements satisfying the above two conditions (Q0), (Q1) and (Q2). The order on $Q(n, k)$ is defined as follows: for \mathbb{T}_1 and \mathbb{T}_2 , $\mathbb{T}_1 \leq \mathbb{T}_2$ if for all $S_1 \in \mathbb{T}_1$, there exists $S_2 \in \mathbb{T}_2$ such that $S_1 \subset S_2$. From now on, we always assume that a set \mathbb{T} satisfies (Q0).

Following the works of Bayer–Brandt and Athanasiadis, we define a map $\nu : Q(n, k) \rightarrow \mathbb{N}$ and subposet $P(n, k)$ of $Q(n, k)$.

Definition 3.1 (Bayer–Brandt [BB97], Athanasiadis [Ath99]). Let \mathbb{T} be an element in $Q(n, k)$ and define

$$\nu(\mathbb{T}) = \sum_{S \in \mathbb{T}} (|S| - k)$$

for each $\mathbb{T} \in Q(n, k)$. The family \mathbb{T} satisfies the *Bayer-Brandt-Athanasiadis condition (the BBA-condition)* if

1. $|S| > k$ for all $S \in \mathbb{T}$, and
2. $|\mathbb{T}'| > 1$ then $\nu(\{\bigcup_{S \in \mathbb{T}'} S\}) > \nu(\mathbb{T}')$ for all $\mathbb{T}' \subset \mathbb{T}$.

The *Bayer-Brandt-Athanasiadis poset (BBA-poset)* $P(n, k)$ is a subposet of the poset graded by ν of $Q(n, k)$. It consists of the set of all $\mathbb{T} \in Q(n, k)$ satisfies the BBA-condition.

The following theorem is conjectured by Bayer–Brandt, and proven by Athanasiadis.

Theorem 3.2 (Bayer–Brandt [BB97], Athanasiadis [Ath99]). Let $n \geq k + 1 \geq 2$. There exists a real arrangement $\mathcal{A} \in \mathbf{Arr}^\circ(n, k)$, which is very generic and $\mathcal{L}(\mathcal{B}(\mathcal{A}))$ isomorphic to $P(n, k)$. The isomorphism is defined by the canonical presentation $\mathbb{T} : \mathcal{L}(\mathcal{B}(\mathcal{A})) \rightarrow P(n, k)$.

We define very generic arrangements over general fields.

Definition 3.3. Let \mathcal{A} be a generic arrangement with n hyperplanes in k dimension space W , and let X be an intersection in $\mathcal{L}(\mathcal{B}(\mathcal{A}))$.

- an arrangement \mathcal{A} is called *very generic* if $\mathcal{L}(\mathcal{B}(\mathcal{A})) \cong P(n, k)$;
- an intersection $X \in \mathcal{L}(\mathcal{B}(\mathcal{A}))$ is *very generic* if $\mathbb{T}(X) \in P(n, k)$ and $\text{rank } X = \nu(\mathbb{T}(X))$;
- a translation \mathcal{A}^t is *very generic* if it is representative of some very generic intersection.

Otherwise, they are called *non-very generic*. A non-very generic intersection X is *minimal* if Y is very generic whenever $X \subsetneq Y \in \mathcal{L}(\mathcal{B}(\mathcal{A}))$.

Notice that (non-)very generic intersections are first defined in [SS22]; however, since it is unclear, we will use the above definition. The condition of very generic intersections means that if $X = \bigcap_{S \in \mathbb{T}(X)} D_S$ is very generic then $D_S (S \in \mathbb{T}(X))$ intersect transversally, i.e. $\text{rank } X = \sum_{S \in \mathbb{T}(X)} \text{rank } D_S$. Also, note that if $\mathbb{T}(X)$ fails the BBA-condition, then X is non-very generic. However, the converse does not hold. Such cases are $\mathbb{T}(X) \in P(n, k)$ and $\text{rank } X < \nu(\mathbb{T}(X))$; one of the examples is in Example 2.2.

Proposition 3.4. Let $X \in \mathcal{L}(\mathcal{B}(\mathcal{A}))$. Then $\text{rank } X \leq \nu(\mathbb{T}(X))$.

Proof. By induction on the size of $\mathbb{T} = \mathbb{T}(X)$. If $\mathbb{T} = 1$ then $\text{rank } X = \nu(\mathbb{T}(X))$. Assume that $\text{rank } X' \leq \nu(\mathbb{T}(X'))$ holds for every X' such that $|\mathbb{T}(X')| = m - 1$. If $\mathbb{T} = \{S_1, \dots, S_m\}$, then $X = \bigcap_{i=1}^{m-1} D_{S_i} \cap D_{S_m}$. By the semimodularity of the rank function,

$$\begin{aligned} \text{rank } X &\leq \text{rank } X + \text{rank} \left(\left(\bigcap_{i=1}^{m-1} D_{S_i} \right) + D_{S_m} \right) \\ &\leq \text{rank} \bigcap_{i=1}^{m-1} D_{S_i} + \text{rank } D_{S_m} \leq \nu \left(\mathbb{T} \left(\bigcap_{i=1}^{m-1} D_{S_i} \right) \right) + |S_m| - k = \nu(\mathbb{T}(X)). \end{aligned}$$

□

Let $\mathbf{Arr}(n, k)$ denote the set of linear isomorphism class of essential central multiarrangements consisting of n hyperplanes in $W = \mathbb{K}^k$, and let $\mathbf{Arr}^\circ(n, k) \subset \mathbf{Arr}(n, k)$ denote the set of generic arrangements. If \mathcal{A} is a generic arrangement, then their normals $\alpha_1, \dots, \alpha_n$ can be written as follows, by appropriate coordinate transformations and scalar products;

$$(\alpha_1, \dots, \alpha_n) = \begin{pmatrix} 1 & 0 & \cdots & 0 & 0 & 1 & a_{1,k+2} & \cdots & a_{1,n} \\ 0 & 1 & & 0 & 0 & 1 & a_{2,k+2} & \cdots & a_{2,n} \\ \vdots & & \ddots & & \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & & 1 & 0 & 1 & a_{k-1,k+2} & \cdots & a_{k-1,n} \\ 0 & 0 & \cdots & 0 & 1 & 1 & 1 & \cdots & 1 \end{pmatrix}.$$

Therefore, $\mathbf{Arr}^\circ(n, k)$ is the isomorphic to a Zariski open set of a $(k-1) \times (n-k-1)$ dimensional affine space. Note that all representative of the isomorphism class of arrangements is written as $(\alpha_1, \dots, \alpha_n) \in (W^*)^n$, and all lists of normals that are transformed by the automorphism of W and the scaling of each vector are included in the same equivalence class. We have $\mathbf{Arr}(n, k) \subset \mathbb{K}^{k \times n} / \text{GL}(W) \times (\mathbb{K}^\times)^n \cong \text{Gr}_k(\mathbb{K}^n) / (\mathbb{K}^\times)^n$. Therefore, we can treat $\mathbf{Arr}(n, k)$ as a quotient of the Grassmannian $\text{Gr}_k(\mathbb{K}^n)$ and will use Plücker coordinates for $\mathbf{Arr}(n, k)$.

Definition 3.5. Let \mathbb{T} be an element in $Q(n, k)$, and let r be a non-negative integer. Define

$$\begin{aligned} V_{(\mathbb{T}, r)} &= \left\{ \mathcal{A} \in \mathbf{Arr}(n, k) \mid \text{rank} \bigcap_{S \in \mathbb{T}} D_S^{\mathcal{A}} \leq r \right\}, \\ V_{(\mathbb{T}, r)}^\circ &= V_{(\mathbb{T}, r)} \cap \mathbf{Arr}^\circ(n, k). \end{aligned}$$

We call $V_{(\mathbb{T}, r)}$ the *non-very generic variety* of (\mathbb{T}, r) if $V_{(\mathbb{T}, r)} \neq \emptyset$ and $V_{(\mathbb{T}, r)} \neq \mathbf{Arr}(n, k)$.

Note that $V_{(\mathbb{T}, r)}$ is defined by vanishing all the $r \times r$ minor of the matrix $(\alpha_C)_{C \subset S \in \mathbb{T}}$, so it is an algebraic set in $\mathbf{Arr}(n, k)$. However, it is not necessarily irreducible.

We know $\mathcal{A} \in V_{(\mathbb{T}_0, r)}$ if $\mathbb{T}([\mathcal{A}^t]) = \mathbb{T}_0$ for some $t \in \mathbb{K}^n$. Therefore, if a coefficient field is the real or the complex numbers and $\mathbb{T} \in P(n, k)$, then we have $V_{(\mathbb{T}, \nu(\mathbb{T}))} = \mathbf{Arr}(n, k)$. Also, by definition, the following proposition holds.

Proposition 3.6. Let $\mathbb{T}, \mathbb{T}' \in Q(n, k)$ and let r and r' be non-negative integers. If $r \leq r'$ and $\mathbb{T} \geq \mathbb{T}'$, then $V_{(\mathbb{T}, r)} \subset V_{(\mathbb{T}', r')}$.

Proof. If $\mathbb{T} \geq \mathbb{T}'$, then it holds that $\bigcap_{S \in \mathbb{T}} D_S \subset \bigcap_{S \in \mathbb{T}'} D_S$ and so $\text{rank} \bigcap_{S \in \mathbb{T}} D_S \geq \text{rank} \bigcap_{S \in \mathbb{T}'} D_S$. Hence, we have

$$\left\{ \mathcal{A} \in \mathbf{Arr}(n, k) \mid \text{rank} \bigcap_{S \in \mathbb{T}} D_S^{\mathcal{A}} \leq r \right\} \subset \left\{ \mathcal{A} \in \mathbf{Arr}(n, k) \mid \text{rank} \bigcap_{S \in \mathbb{T}'} D_S^{\mathcal{A}} \leq r' \right\}.$$

□

The rank decreasing surjection $P(n, k) \rightarrow \mathcal{L}(\mathcal{B}(\mathcal{A})); \mathbb{T} \mapsto \bigcap_{S \in \mathbb{T}} D_S$ exists. Hence, we have $\nu(\mathbb{T}) \geq \text{rank} \bigcap_{S \in \mathbb{T}} D_S$ for $\mathbb{T} \in P(n, k)$. Therefore, for an intersection X in $\mathcal{L}(\mathcal{B}(\mathcal{A}))$,

$$\text{rank } X < \min \{ \nu(\mathbb{T}') \mid \mathbb{T}' \in P(n, k), \mathbb{T}(X) < \mathbb{T}' \}$$

holds. In particular, if X is very generic, then $\min \{ \nu(\mathbb{T}') \mid \mathbb{T}' \in P(n, k), \mathbb{T}(X) < \mathbb{T}' \} = \text{rank } X + 1$. And we have $V_{(\mathbb{T}, r)}^\circ = \mathbf{Arr}^\circ(n, k)$ if $r \geq \min \{ \nu(\mathbb{T}') \mid \mathbb{T}' \in P(n, k), \mathbb{T} < \mathbb{T}' \}$.

For simplicity, we write $V_{\mathbb{T}}$ and $V_{\mathbb{T}}^\circ$ as $V_{(\mathbb{T}, r)}$ and $V_{(\mathbb{T}, r)}^\circ$ if $\min \{ \nu(\mathbb{T}') \mid \mathbb{T}' \in P(n, k), \mathbb{T} < \mathbb{T}' \} = r+1$. Then we have if $\mathcal{A} \in V_{\mathbb{T}}$ for some $\mathbb{T} \in Q(n, k) \setminus P(n, k)$, then \mathcal{A} is non-very generic. An example of a case where a non-very generic variety is not equal to $V_{\mathbb{T}}$ is in Example 2.2; take \mathcal{A} and X as in Example 2.2, then $\min \{ \nu(\mathbb{T}') \mid \mathbb{T}' \in P(n, k), \mathbb{T} < \mathbb{T}' \} = 6 = \text{rank } X + 2$ holds.

Example 3.7. Consider the same intersection as in example 2.1. Take $\mathbb{T} = \{123, 156, 246, 345\}$ For $n = 6, k = 2$, $\mathbb{T} = \{123, 156, 246, 345\}$ is in $Q(6, 2)$ but not in $P(6, 2)$. The element larger than \mathbb{T} in $P(6, 2)$ is only $\{123456\}$, hence $\min\{\nu(\mathbb{T}') \mid \mathbb{T}' \in P(n, k), \mathbb{T} < \mathbb{T}'\} = 6 - 2 = 4$. Then

$$V_{(\mathbb{T}, r)} = \{\mathcal{A} \in \mathbf{Arr}(6, 2) \mid \text{rank } D_{123}^{\mathcal{A}} \cap D_{156}^{\mathcal{A}} \cap D_{246}^{\mathcal{A}} \cap D_{345}^{\mathcal{A}} \leq r\}$$

If $\min\{\nu(\mathbb{T}') \mid \mathbb{T}' \in P(n, k), \mathbb{T} < \mathbb{T}'\} = r + 1$, then we have $V_{\mathbb{T}} = V_{(\mathbb{T}, 3)} = \{\mathcal{A} \in \mathbf{Arr}(6, 2) \mid \Delta_{16}\Delta_{24}\Delta_{35} - \Delta_{15}\Delta_{26}\Delta_{34} = 0\}$ where α_i is a normal of H_i and $\Delta_{ij} = \det(\alpha_i \alpha_j)$.

Similarly, for $n = 7, k = 2$, take $\mathbb{T} = \{123, 147, 156, 246, 357\}$ in $Q(7, 2)$ but not in $P(7, 2)$. Then $\min\{\nu(\mathbb{T}') \mid \mathbb{T}' \in P(n, k), \mathbb{T} < \mathbb{T}'\} = 7 - 2 = 5$. We have $V_{\mathbb{T}} = V_{(\mathbb{T}, 4)}$ and the defining equation of $V_{\mathbb{T}}$ is

$$\Delta_{14}\Delta_{16}\Delta_{27}\Delta_{35} - \Delta_{17}\Delta_{15}\Delta_{26}\Delta_{34} = 0. \quad (2)$$

The author shows such the defining equation in [SS22].

Let us justify that it is fundamental to consider $V_{\mathbb{T}}$; that is, the BBA-condition is a necessary and sufficient condition for determining very generic arrangements.

Lemma 3.8. Let $\mathcal{A} \in \mathbf{Arr}^\circ(n, k)$.

1. Let X be a non-very generic intersection in $\mathcal{L}(\mathcal{B}(\mathcal{A}))$. If X satisfies the BBA-condition, then there exists an intersection $Y \in \mathcal{L}(\mathcal{B}(\mathcal{A}))$ which fails the BBA-condition with the condition $X < Y$.
2. A generic arrangement \mathcal{A} is non-very generic if and only if there is a non-very generic translation \mathcal{A}^t .

Proof. 1. If there is no such intersection Y , then the length of every maximal chain contains X in $\mathcal{L}(\mathcal{B}(\mathcal{A}))$ is at most $n - k$. It is contradictory that the length of a maximal chain of a geometric lattice of rank $n - k$ is $n - k + 1$.

2. Suppose that \mathcal{A} is non-very generic, but every intersection in $\mathcal{L}(\mathcal{B}(\mathcal{A}))$ is very generic. Then there is a proper subposet P' of $P(n, k)$ with the same grading ν such that $P' \cong \mathcal{L}(\mathcal{B}(\mathcal{A}))$. In other words, there exists \mathbb{T}_0 in $P(n, k) \setminus P'$. Let \mathbb{T}_1 be a canonical presentation of $\bigcap_{S \in \mathbb{T}_0} D_S$, then it holds that $\nu(\mathbb{T}_1) = \text{rank} \bigcap_{S \in \mathbb{T}_0} D_S < \nu(\mathbb{T}_0)$. Since \mathbb{T}_0 is not canonical presentation of $X = \bigcap_{S \in \mathbb{T}_0} D_S$, there exists canonical presentation \mathbb{T}_1 of X satisfying the BBA-condition with a property $\mathbb{T}_1 > \mathbb{T}_0$. It leads $\text{rank } X < \nu(\mathbb{T}_0) < \nu(\mathbb{T}_1)$. However it contradicts to $\text{rank } X = \nu(\mathbb{T}_1)$. □

The above lemma and Proposition 3.6 yield the following theorems.

Theorem 3.9. Let X be a non-very generic intersection of the rank r in $\mathcal{L}(\mathcal{B}(\mathcal{A}))$. There exists $\mathbb{T}_0 \in Q(n, k)$ such that $\mathbb{T}_0 \notin P(n, k)$, $\mathbb{T}(X) < \mathbb{T}_0$, $\mathcal{A} \in V_{(\mathbb{T}(X), r)} \subset V_{\mathbb{T}_0}$.

Proof. By Lemma 3.8, there exists an intersection Y such that $X < Y$ and $\mathbb{T}(Y)$ fails the BBA-condition. Take $\mathbb{T}_0 = \mathbb{T}(Y)$, then $\mathbb{T}_0 \notin P(n, k)$ and $\mathbb{T}(X) < \mathbb{T}_0$ hold. Finally, by Proposition 3.6, $V_{(\mathbb{T}(X), r)} \subset V_{(\mathbb{T}_0, \text{rank } Y)} \subset V_{\mathbb{T}_0}$. □

Theorem 3.10. The set of generic but non-very generic arrangements in W with n hyperplanes is

$$\bigcup_{\mathbb{T} \in Q(n, k) \setminus P(n, k)} V_{\mathbb{T}}^\circ.$$

4 Generalized Crapo configuration; wheels

In this section, we discuss non-very generic line arrangements, generalized Crapo configuration, and the corresponding non-very generic variety. In this section, we assume $n \geq 3$.

We define a couple of families of subsets of $[2n]$ as follows:

$$\begin{aligned} \mathbb{W}_{2n} &= \{\{2i - 1, 2i, 2i + 1\} \subset [2n] \mid i \in [n]\} \cup \{\{2, 4, \dots, 2n\}\}, \\ \mathbb{W}_{2n}^\dagger &= \{\{2i, 2i + 1, 2i + 2\} \subset [2n] \mid i \in [n]\} \cup \{\{1, 3, \dots, 2n - 1\}\}. \end{aligned}$$

Here, we identify $[2n]$ with the cyclic group $\mathbb{Z}/2n\mathbb{Z}$. We call \mathbb{W}_{2n} and a permutations $\sigma \cdot \mathbb{W}_{2n}$ of \mathbb{W}_{2n} $2n$ -wheel and call $\sigma \cdot \mathbb{W}_{2n}^\dagger$ a twin of $\sigma \cdot \mathbb{W}_{2n}$ where $\sigma \cdot \mathbb{T}$ denotes $\{\sigma \cdot S = \{\sigma \cdot i \mid i \in S\} \mid S \in \mathbb{T}\}$ for subcollection $\mathbb{T} \subset 2^{[n]}$ and a permutation $\sigma \in \mathfrak{S}_n$.

Also, a $2n$ -wheel arrangement defined as an affine line arrangement $\{H_1, \dots, H_{2n}\}$ with the condition: $\bigcap_{i \in S} H_i \neq \emptyset$ and $H_j \cap \bigcap_{i \in S} H_i = \emptyset$ for all $j \notin S \in \mathbb{W}_{2n}$ cf. Figure 4. This name comes from the fact that the intersection graph of \mathbb{W}_{2n} is a wheel graph. It is easy to check that $\mathbb{W}_{2n} \in Q(2n, 2)$ and $\mathbb{W}_{2n} \notin P(2n, 2)$. Since $\mathbb{W}_{2n} \setminus \{123\}$ satisfies the BBA-condition, we obtain

$$2n - 3 = \nu(\mathbb{W}_{2n} \setminus \{123\}) < \min\{\nu(\mathbb{T}) \mid \mathbb{T} \in P(2n, 2), \mathbb{W}_{2n} \leq \mathbb{T}\} \leq 2n - 2.$$

Hence we obtain $\min\{\nu(\mathbb{T}') \mid \mathbb{T}' \in P(n, k), \mathbb{W}_{2n} < \mathbb{T}'\} - 1 = 4$ and $V_{\mathbb{W}_{2n}} = V_{(\mathbb{W}_{2n}, 2n-3)}$.




Figure 4: $2n$ -wheel arrangement




Figure 5: add H_* to $2n+2$ -wheel

Theorem 4.1. Let $\mathcal{A} = \{H_1, \dots, H_{2n}\}$ be a generic line arrangement consisting of $2n$ lines in \mathbb{K}^2 . The arrangement \mathcal{A} admits a translation into the $2n$ -wheel arrangement if and only if satisfying the equation;

$$\prod_{i=1}^n \Delta_{2i, 2i-1} - \prod_{i=1}^n \Delta_{2i, 2i+1} = 0. \quad (3)$$

In other words, the defining equation of $V_{\mathbb{W}_{2n}}^\circ$ is the above equation 3.

When $n = 3$, it looks different form from the equation (1), but it follows that these equations are consistent with the three terms Plücker relation; $\Delta_{ab}\Delta_{cd} - \Delta_{ac}\Delta_{bd} + \Delta_{ad}\Delta_{bc} = 0$. The equivalence between that \mathcal{A} satisfying the equation (1) and the existence of a non-very generic intersection whose canonical presentation is $\{123, 156, 246, 345\}$ in $\mathcal{L}(\mathcal{B}(\mathcal{A}))$ holds even if $H_1 = H_4$, $H_2 = H_5$ or $H_3 = H_6$. It can be proved in the same way as the case \mathcal{A} generic.

Proof. We prove this by induction on n . The case $n = 3$ has already been shown. Suppose that the claim holds for n .

First, suppose that there exists translation \mathcal{A}^t admitting a $2n+2$ -wheel arrangement. Then, let H_* be the line through $H_1^t \cap H_2^t \cap H_3^t$ and $H_{2n-1}^t \cap H_{2n}^t \cap H_{2n+1}^t$, and let α_* be the normal vector of H_* . Now, $\{\{\star, 1, 2\}, \{\star, 2n, 2n+1\}, \{1, 2n+1, 2n+2\}, \{2, 2n, 2n+2\}\}$ is a 6-wheel and $\mathbb{W}_{2n+2} \setminus \{\{1, 2, 3\}, \{2n-1, 2n, 2n+1\}, \{1, 2n+1, 2n+2\}, \{2, 4, \dots, 2n+2\}\} \cup \{\{\star, 2, 3\}, \{2n-1, 2n, \star\}, \{2, 4, \dots, 2n\}\}$ is a $2n$ -wheel (c.f. Figure 5).

Then, by the assumption,

$$\frac{\det(\alpha_2 \alpha_\star)}{\Delta_{2,3}} \frac{\Delta_{2n, 2n-1}}{\det(\alpha_{2n} \alpha_\star)} \prod_{i=2}^{n-1} \frac{\Delta_{2i, 2i-1}}{\Delta_{2i, 2i+1}} = 1, \quad (4)$$

and

$$\det(\alpha_2 \alpha_\star) \Delta_{2n, 2n+1} \Delta_{2n+2, 1} - \Delta_{2, 1} \det(\alpha_{2n} \alpha_\star) \Delta_{2n+2, 2n+1} = 0$$

is equivalent to

$$\frac{\det(\alpha_2 \alpha_\star)}{\det(\alpha_{2n} \alpha_\star)} = \frac{\Delta_{2, 1} \Delta_{2n+2, 2n+1}}{\Delta_{2n, 2n+1} \Delta_{2n+2, 1}} \quad (5)$$

hold. Combining them yields the given equation (3).

Conversely, suppose that the equation (3) holds. Taking α_* such that the equation (5) is satisfied, then from the assumption that the equation (4) is also satisfied. The equation (4) provides an isomorphic arrangement to the n -wheel, so removing H_* gets us the arrangement \mathcal{W}_{n+1} . \square

Corollary 4.2. The non-very generic variety of a $2n$ -wheel and the twin of it are same, i.e., $V_{\mathbb{W}_{2n}} = V_{\mathbb{W}^\dagger_{2n}}$.

Definition 4.3. We call a family of subset of \mathbb{T} a *degenerated wheel* (resp. a *degenerated wheel minus a hab*) if \mathbb{T} is formed by $\{\{i_l, i_{l+1}, j_l\} \mid l = 1, \dots, n\} \cup \{\{j_1, \dots, j_n\}\}$ (resp. $\{\{i_l, i_{l+1}, j_l\} \mid l = 1, \dots, n\}$) with the conditions;

- For each $l \in [n]$, i_l does not belong elements of \mathbb{T} without $\{i_{l-1}, i_l, j_{l-1}\}, \{i_l, i_{l+1}, j_l\}$ (allow $j_l = j_{l'}$ for some $l, l' \in [n]$), and
- $i_l = i_{l'}$ if and only if $l - l' \in n\mathbb{Z}$.

Proposition 4.4. Let \mathcal{A} be a generic line arrangement and let X be a minimal non-very generic intersection in $\mathcal{L}(\mathcal{B}(\mathcal{A}))$. If there exists a subcollection of $\mathbb{T}(X)$ formed by $\{\{i_l, i_{l+1}, j_l\} \mid l = 1, \dots, n\}$ which is a degenerated wheel minus a hub, then the normals of \mathcal{A} satisfy the following equation;

$$\prod_{l=1}^n \Delta_{j_l i_l} - \prod_{l=1}^n \Delta_{j_l i_{l+1}} = 0.$$

Proof. Without loss in general, we assume $i_1 < \dots < i_n < j_l$ for $l \in [n]$. For each component D_S of X and $l \in \{3, \dots, |S|\}$, $\alpha_{S,\iota}$ denotes $\alpha_{p_1 p_2 p_\iota}$ where $S = \{p_1 < \dots < p_{|S|}\}$. The orthogonal space $D_S^\perp \subset \mathbb{S}(\mathcal{A})^*$ of D_S is spanned by $\{\alpha_{S,3}, \dots, \alpha_{S,|S|}\}$. Hence the orthogonal space of X is spanned by $\bigcup_{S \in \mathbb{T}(X)} \{\alpha_{S,3}, \dots, \alpha_{S,|S|}\}$ whose cardinality $\nu(\mathbb{T}(X)) = \sum_{S \in \mathbb{T}(X)} \text{rank } D_S$. Since X is non-very generic, $\dim D_S^\perp = \text{rank } X < \nu(\mathbb{T}(X))$. Thus $\bigcup_{S \in \mathbb{T}(X)} \{\alpha_{S,3}, \dots, \alpha_{S,|S|}\}$ is linearly dependent. In particular, it is minimally dependent because X is minimal non-very generic. In other words, there exists non-zero scalars $c_{S,\iota}$ ($S \in \mathbb{T}(X), \iota \in \{3, \dots, |S|\}$) such that $\sum_{S \in \mathbb{T}(X)} \sum_{\iota=3}^{|S|} c_{S,\iota} \alpha_{S,\iota} = 0$. Fix such $c_{S,\iota}$'s. The above equation also holds for each coefficient of e_i^* , i.e.,

$$\sum_{S \in \mathbb{T}(X)} \sum_{\iota=3}^{|S|} c_{S,\iota} a_{S,\iota,i} e_i^* = 0$$

where $a_{S,\iota} = \sum_{i=1}^{|A|} a_{S,\iota,i} e_i^*$.

When $|S| = 3$, we write c_S instead of $c_{S,3}$. The normal of $D_{i_1 i_n j_n}$ is $\alpha_{i_1 i_n j_n} = \Delta_{i_n j_n} e_{i_1}^* - \Delta_{i_1 j_n} e_{i_n}^* + \Delta_{i_1 i_n} e_{j_n}^*$. And the normal of $D_{i_l i_{l+1} j_l}$ is $\alpha_{i_l i_{l+1} j_l} = \Delta_{i_{l+1} j_l} e_{i_l}^* - \Delta_{i_l j_l} e_{i_{l+1}}^* + \Delta_{i_l i_{l+1}} e_{j_l}^*$ for $l = 2, \dots, n$. By the assumption, $a_{S,\iota,i} \neq 0$ if and only if $S = \{i_{l-1}, i_l, j_{l-1}\}, \{i_l, i_{l+1}, j_l\}$. Hence we get

$$\begin{aligned} c_{i_1 i_2 j_1} \Delta_{i_2 j_1} + c_{i_1 i_n j_n} \Delta_{i_n j_n} &= 0, \\ -c_{i_{l-1} i_l j_{l-1}} \Delta_{i_{l-1} j_{l-1}} + c_{i_l i_{l+1} j_l} \Delta_{i_{l+1} j_l} &= 0 \text{ for } l = 2, \dots, n-1, \\ -c_{i_{n-1} i_n j_{n-1}} \Delta_{i_{n-1} j_{n-1}} - c_{i_1 i_n j_n} \Delta_{i_1 j_n} &= 0. \end{aligned}$$

Therefore

$$1 = \frac{c_{i_1 i_2 j_1}}{c_{i_1 i_n j_n}} \left(\prod_{l=2}^{n-1} \frac{c_{i_l i_{l+1} j_l}}{c_{i_{l-1} i_l j_{l-1}}} \right) \frac{c_{i_1 i_n j_n}}{c_{i_{n-1} i_n j_{n-1}}} = \frac{\Delta_{i_n j_n}}{\Delta_{i_2 j_1}} \left(\prod_{l=2}^{n-1} \frac{\Delta_{i_{l+1} j_l}}{\Delta_{i_{l-1} j_{l-1}}} \right) \frac{\Delta_{i_{n-1} j_{n-1}}}{\Delta_{i_1 j_n}} = \prod_{i=1}^n \frac{\Delta_{j_i i_i}}{\Delta_{j_i i_{i+1}}}.$$

□

5 Deletion, contraction, degeneration

Recall the definition of discriminantal arrangement in the general case (see [Cra84, Cra85, BB97, OW19, CFW21, FW23]). Let $\mathcal{A} = \{H_i = \text{Ker } \alpha_i \mid i = 1, \dots, n\}$ be a multiarrangement with normals $\{\alpha_1, \dots, \alpha_n\}$ where we allow the α_i 's to be parallel. (Multiarrangements are usually defined as a pair of an arrangement and a weight function, but here, they are defined as a multiset indexed by $[n]$. Hence, we allow $\alpha_i = \alpha_j$ for some $i, j \in [n]$.) As in the case of generic arrangement, we define $\mathbb{S}(\mathcal{A}) = \bigoplus_{i \in [n]} W/H_i$ and $D_S = \{\mathcal{A}^t \in \mathbb{S}(\mathcal{A}) \mid \bigcap_{i \in S} H_i^t \neq \emptyset\}$.

Let $f_S : V \rightarrow \mathbb{S}(\mathcal{A}); v \mapsto \sum_{i \in S} H_i^{\alpha_i(v)}$ for each $S \subset [n]$. Then, we obtain $D_S = \text{Im } f_S \oplus \mathbb{S}(\mathcal{A} \setminus S)$, and so $\text{codim } D_S = |S| - \dim \text{Im}((\alpha_i)_{i \in S} : W \rightarrow \mathbb{K}^S) = |S| - \text{rank}(\bigcap_{i \in S} H_i)$. Hence $D_S \neq \mathbb{S}(\mathcal{A})$ if and only if $\{\alpha_i \mid i \in S\}$ is linearly dependent, and D_S is hyperplane if $\text{rank}(\bigcap_{i \in S} H_i) = |S| - 1$. A subset C of $[n]$ is a *circuit* if $\{\alpha_i \mid i \in S\}$ is minimally linearly dependent, and $\mathcal{C} = \mathcal{C}(\mathcal{A})$ denotes the set of circuits in \mathcal{A} . Then D_C is a hyperplane in $\mathbb{S}(\mathcal{A})$ for a circuit $C \in \mathcal{C}$, and $D_C \neq D_{C'}$ whenever C and C' in \mathcal{C} are distinct. The discriminantal arrangement associated to \mathcal{A} is a hyperplane arrangement consisting of all D_C for all circuits $C \in \mathcal{C}$. If the cardinality of the circuit is all

$k + 1$, it is a generic arrangement. Thus, this is a generalization of what was defined in Section 2. Since we are interested in generic arrangements, we will focus on the primary case where the cardinality of all circuits is 2 or $k + 1$. In other words, they are generic arrangements with some parallel hyperplanes added. In other words, they are generic arrangements with some parallel hyperplanes added. If $C = \{i, j\}$ then $\alpha_C = c_i e_i^* - c_j e_j^*$ where c_i, c_j satisfy $c_i \alpha_i = c_j \alpha_j$. Here e_i 's, in which each e_i spans W/H_i for each $i \in [n]$, are the canonical basis of $\mathbb{S}(\mathcal{A})$, and e_i^* are the dual basis.

Example 5.1. Let the coefficient field be real numbers. Take the vectors

$$(\alpha_1 \alpha_2 \alpha_3 \alpha_4) = \begin{pmatrix} 1 & 2 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix}.$$

Then the circuits of the arrangement \mathcal{A} whose normals are $\alpha_1, \alpha_2, \alpha_3, \alpha_4$ are $\mathcal{C}(\mathcal{A}) = \{12, 134, 234\}$. Hence $\mathcal{B}(\mathcal{A})$ consists of 3 hyperplanes in 4 dimensional space $\mathbb{S}(\mathcal{A})$, and the normals are

$$\alpha_{12} = -2e_1^* + e_2^*, \quad \alpha_{134} = -e_1^* - e_3^* - e_4^*, \quad \alpha_{234} = -e_2^* - 2e_3^* - 2e_4^*.$$

As is well known, deletion and restriction are basic constructions of arrangements. For a (multi)arrangement \mathcal{A} in W and $H_i \in \mathcal{A}$, define the deletion $\mathcal{A} \setminus H_i = \mathcal{A} \setminus \{H_i\}$ and the restriction $\mathcal{A}^{H_i} = \{H_i \cap H \mid H \in \mathcal{A} \setminus H_i\}$. Then $\mathcal{C}(\mathcal{A} \setminus H_i)$ consists of all circuits of \mathcal{A} not containing i , is $\{C \mid i \notin C \in \mathcal{C}(\mathcal{A})\}$. And $\mathcal{C}(\mathcal{A}^{H_i})$ consists all minimal elements of $\{C \setminus i \mid i \in C \in \mathcal{C}(\mathcal{A})\}$. The discriminantal arrangement associated to restriction \mathcal{A}^{H_i} is discussed in [LS18], but we mention it here again.

Proposition 5.2. Let \mathcal{A} be an arrangement and let $H_i \in \mathcal{A}$.

- i) The discriminantal arrangement $\mathcal{B}(\mathcal{A} \setminus H_i)$ associated to $\mathcal{A} \setminus H_i$ is linearly equivalent to $\{D_C \cap \mathbb{S}(\mathcal{A} \setminus H_i) \mid i \notin C \in \mathcal{C}(\mathcal{A})\}$. In particular $H_i = H_j$ for some $j \neq i$, $\mathcal{B}(\mathcal{A} \setminus H_i)$ is linearly equivalent to $\mathcal{B}(\mathcal{A})^{D_{ij}}$.
- ii) The discriminantal arrangement $\mathcal{B}(\mathcal{A}^{H_i})$ associated to \mathcal{A}^{H_i} is linearly equivalent to

$$\begin{aligned} & \{D_C \cap \mathbb{S}(\mathcal{A} \setminus H_i) \mid i \in C \in \mathcal{C}(\mathcal{A})\} \\ & \cup \{D_{C \cup \{i\}} \cap \mathbb{S}(\mathcal{A} \setminus H_i) \mid i \notin C \in \mathcal{C}(\mathcal{A}), C \cup \{i\} \text{ contains no circuits of } \mathcal{A} \text{ except } C\}. \end{aligned}$$

In particular, if \mathcal{A} is a generic arrangement, then $\mathcal{B}(\mathcal{A}^{H_i}) \cong \{D_C \cap \mathbb{S}(\mathcal{A} \setminus H_i) \mid i \in C \in \mathcal{C}(\mathcal{A})\}$.

Proof. i) The first claim follows from $\mathcal{C}(\mathcal{A} \setminus H_i) = \{C \in \mathcal{C}(\mathcal{A}) \mid i \notin C\}$. If \mathcal{A} has $j \in [n]$ such that $H_i = H_j$, then $D_{ij} = \{\mathcal{A}^t \mid H_i^t = H_j^t\}$. Hence the isomorphism between D_{ij} and $\mathbb{S}(\mathcal{A} \setminus H_i)$ by $\mathcal{A}^t \mapsto \mathcal{A}^t \setminus H_i^t$ provides the linear equivalence $\mathcal{B}(\mathcal{A})^{D_{ij}} \cong \mathcal{B}(\mathcal{A} \setminus H_i)$.

- ii) Fix a natural isomorphism between $\mathbb{S}(\mathcal{A}^{H_i}) = \bigoplus_{H \in \mathcal{A} \setminus H_i} H_i / (H_i \cap H)$ and $\mathbb{S}(\mathcal{A} \setminus H_i) = \bigoplus_{H \in \mathcal{A} \setminus H_i} W/H$. Let $(\mathcal{A} \setminus H_i)^t \in D_S(\mathcal{A} \setminus H_i)$, then $\bigcap_{i \in S} (H_i^t \cap H_i) = H_i \cap \bigcap_{i \in S} H_i^t \neq \emptyset$. Thus if $0 \in C \in \mathcal{C}(\mathcal{A})$, then $C \setminus i$ is a circuit of \mathcal{A} and $D_{C \setminus i}(\mathcal{A}^{H_i}) = D_C(\mathcal{A}) \cap \mathbb{S}(\mathcal{A} \setminus H_i)$. Assume $i \notin C \in \mathcal{C}(\mathcal{A})$, then $D_C(\mathcal{A}^{H_i}) \cong D_{C \cup \{i\}}(\mathcal{A}) \cap \mathbb{S}(\mathcal{A} \setminus H_i)$. If $C \cup \{i\}$ contains circuits C' not C of \mathcal{A} , then $D_{C \cup \{i\}} \subsetneq D_{C'}$. In this case, $D_{C \cup \{i\}}$ is not hyperplane in $\mathbb{S}(\mathcal{A})$. Conversely, $C \cup \{i\}$ contains no circuits except C , then $D_{C \cup \{i\}}$ is a hyperplane in $\mathbb{S}(\mathcal{A})$. □

Therefore, the very generic arrangements are closed under the operations deletion and restriction.

Corollary 5.3. Let \mathcal{A} be a generic arrangement. If $\mathcal{A} \setminus H$ or \mathcal{A}^H is a non-very generic arrangement for some hyperplane $H \in \mathcal{A}$, then \mathcal{A} is also non-very generic.

In particular, we call the deletion of parallel elements degeneration.

Definition 5.4. Let $i, j \in [n]$ and let $\mathbb{T} \in Q([n], k)$. Define

$$\tilde{\mathbb{T}}_{i \rightarrow j} = \{S \cup \{j\} \setminus \{i\} \mid i \in S \in \mathbb{T}\} \cup \{S \mid i \notin S \in \mathbb{T}\} \setminus \binom{[n]}{k}$$

and $\mathbb{T}_{i \rightarrow j}$ to be a minimal element of $Q([n] \setminus i, k)$ equal to or larger than $\tilde{\mathbb{T}}_{i \rightarrow j}$. The family $\mathbb{T}_{i \rightarrow j}$ of subset of $[n] \setminus i$ is said to be a *degeneration* of \mathbb{T} from i to j . We write $\gamma = \gamma(\mathbb{T}; i, j)$ as the cardinality of $\{S \in \mathbb{T} \mid \{i, j\} \subset S\}$.




Figure 6: representatives of the minimal non-very generic intersection with 6 or 7 lines

Example 5.5. Let \mathbb{T} be $\{123, 147, 156, 246, 357\}$. The degeneration of \mathbb{T} from 7 to 4 is $\mathbb{T}_{7 \rightarrow 4} = \{123, 156, 246, 345\}$. Refer to the two non-very generic varieties in Example 3.7, these defined by equation (1) and (2), i.e.

$$V_{\mathbb{T}} = \{\mathcal{A} \in \mathbf{Arr}(7, 2) \mid \Delta_{14}\Delta_{16}\Delta_{27}\Delta_{35} - \Delta_{17}\Delta_{15}\Delta_{26}\Delta_{34} = 0\}$$

$$V_{\mathbb{T}_{7 \rightarrow 4}} = \{\mathcal{A} \in \mathbf{Arr}(6, 2) \mid \Delta_{16}\Delta_{24}\Delta_{35} - \Delta_{15}\Delta_{26}\Delta_{34} = 0\}$$

where α_i is a normal of H_i and $\Delta_{ij} = \det(\alpha_i \alpha_j)$. For real or complex coefficients, degeneration of \mathbb{T} from 4 to 7 corresponds to continuously deforming the representative and taking the limit $H_7^t \rightarrow H_4^t$ while maintaining the canonical presentation, cf. Figure 6. Equation (1) is obtained by substituting α_4 for α_7 in equation (2). It says that; if $\mathcal{A} = \{H_1, \dots, H_7\} \in V_{\mathbb{T}}$ and $H_4 = H_7$, then $\mathcal{A} \setminus H_7 \in V_{\mathbb{T}_{7 \rightarrow 4}}$.

Generalizing the above example, we obtain the following theorem.

Theorem 5.6. Let $\mathbb{T} \in Q(n, k)$ and let $\mathbb{T}_{n \rightarrow j}$ be the degeneration of \mathbb{T} from n to j .

1. If $\left| \bigcup_{S \in \mathbb{T}_{n \rightarrow j}} S \right| = \left| \bigcup_{S \in \mathbb{T}} S \right| - 1$, $\mathbb{T} \notin P(n-1, k)$, $\gamma(\mathbb{T}; i, j) = 1$ and $\tilde{\mathbb{T}}_{n \rightarrow j} \in Q(n-1, k)$, then $\mathbb{T}_{n \rightarrow j} \notin P(n, k)$.
2. If $\gamma(\mathbb{T}; n, j) > 0$, $\mathcal{A} = \{H_1, \dots, H_n\} \in V_{(\mathbb{T}, r)}$ and $H_n = H_j$, then $\mathcal{A} \setminus H_n \in V_{(\mathbb{T}_{n \rightarrow j}, r-1)}$.

Proof. 1. Since $\mathbb{T} \notin P(n, k)$, there exists $\mathbb{T}' \subset \mathbb{T}$ such that $\nu(\mathbb{T}') \geq \left| \bigcup_{S \in \mathbb{T}'} S \right| - k$. Let us show that $\mathbb{T}'_{n \rightarrow j} \notin P(n-1, k)$. By the assumption, $\tilde{\mathbb{T}}_{n \rightarrow j} = \mathbb{T}_{n \rightarrow j}$ for each $\mathbb{T}' \subset \mathbb{T}$. We obtain $\nu(\mathbb{T}'_{n \rightarrow j}) = \nu(\tilde{\mathbb{T}}'_{n \rightarrow j}) = \nu(\mathbb{T}') - \gamma(\mathbb{T}'; i, j)$. Because $\gamma(\mathbb{T}'; i, j) \leq \gamma(\mathbb{T}; i, j)$ for $\mathbb{T}' \subset \mathbb{T}$, $0 \leq \gamma(\mathbb{T}'; i, j) \leq 1$. If $\gamma(\mathbb{T}'; i, j) = 1$, then $\left| \bigcup_{S \in \mathbb{T}'_{n \rightarrow j}} S \right| = \left| \bigcup_{S \in \mathbb{T}'} S \setminus \{n\} \right| = \left| \bigcup_{S \in \mathbb{T}'} S \right| - 1$, hence we have $\nu(\mathbb{T}'_{n \rightarrow j}) \geq \left| \bigcup_{S \in \mathbb{T}'_{n \rightarrow j}} S \right| - k$. If $\gamma(\mathbb{T}'; i, j) = 0$, then we have $\nu(\mathbb{T}'_{n \rightarrow j}) = \nu(\mathbb{T}') \geq \left| \bigcup_{S \in \mathbb{T}'} S \right| - k \geq \left| \bigcup_{S \in \mathbb{T}'_{n \rightarrow j}} S \right| - k$.

2. We identify $\mathcal{B}(\mathcal{A} \setminus H_n)$ and $\mathcal{B}(\mathcal{A})^{D_{nj}}$ via Proposition 5.2. By the assumption, $D_{nj} \supset \bigcap_{S \in \mathbb{T}} D_S$ and $\text{rank}_{\mathcal{B}(\mathcal{A})} \bigcap_{S \in \mathbb{T}} D_S \leq r$. Hence

$$\bigcap_{S \in \mathbb{T}} D_S = D_{nj} \cap \bigcap_{S \in \mathbb{T}} D_S = D_{nj} \cap \bigcap_{S \in \mathbb{T}_{n \rightarrow j}} D_S$$

and

$$\text{rank}_{\mathcal{B}(\mathcal{A})^{D_{nj}}} D_{nj} \cap \bigcap_{S \in \mathbb{T}_{n \rightarrow j}} D_S = \text{rank}_{\mathcal{B}(\mathcal{A})} D_{nj} \cap \bigcap_{S \in \mathbb{T}_{n \rightarrow j}} D_S - 1 = \text{rank}_{\mathcal{B}(\mathcal{A})} \bigcap_{S \in \mathbb{T}} D_S - 1 \leq r - 1.$$

□

6 Applications

In this section, we provide some applications of the degeneration construction and wheels. The $2n$ -wheel was defined for a generic arrangement consisting of $2n$ hyperplanes. However it can be extended to a general arrangement by adding the following assumption: $H_i \neq H_{i+1}$ and $H_i \neq H_{i+2}$ for each $i \in [2n]$. Also, even under this assumption, Theorem 4.1 holds, i.e. the non-very generic variety $V_{\mathbb{W}_{2n}}$ is defined by the equation (3). Define $\mathbb{L}_{2n+2} = \{\{2i-1, 2i, 2n+1\} \mid i \in [n]\} \cup \{\{2i, 2i+1, 2n+2\} \mid i \in [n-1]\} \cup \{\{1, 2n, 2n+2\}\}$, and we call it a $2n+2$ -ladder. Then $\mathbb{L}_{2n+2} \in Q(2n+2, 2) \setminus P(2n+2, 2)$.

Proposition 6.1. The defining equation of $V_{\mathbb{L}_{2n+2}}$ is

$$\prod_{i=1}^n \Delta_{2i,2n+2} \Delta_{2i-1,2n+1} - \prod_{i=1}^n \Delta_{2i,2n+1} \Delta_{2i-1,2n+2} = 0.$$

Proof. Because \mathbb{L}_{2n+2} can be obtained by degenerate of \mathbb{W}_{4n} from $4i-2 (i=2, \dots, n)$ to 2 and from $4i (i=2, \dots, n)$ to 4 and permutation of indices. Then it follows from Theorem 5.6 and Theorem 4.1 \square

Also, by applying Theorem 5.6 to Theorem 4.1 in the same way as the proof above and combining Proposition 4.4, we obtain the following proposition.

Proposition 6.2. Let \mathcal{A} be a generic line arrangement and let X be a minimal non-very generic intersection in $\mathcal{L}(\mathcal{B}(\mathcal{A}))$. If there exists a subcollection of $\mathbb{T}(X)$ formed by $\{\{i_l, i_{l+1}, j_l\} \mid l=1, \dots, n\}$ which is a degenerated wheel minus a hub, then there exists a non-very generic intersection Y in $\mathcal{L}(\mathcal{B}(\mathcal{A}))$, whose canonical presentation is a degenerated wheel $\{\{i_l, i_{l+1}, j_l\} \mid l=1, \dots, n\} \cup \{\{j_1, \dots, j_n\}\}$.

Let us list non-very generic varieties corresponding to minimal non-very generic intersections in **Arr**(8, 2). Assume a line arrangement \mathcal{A} with 8 lines has a minimal non-very generic intersection X in $\mathcal{L}(\mathcal{B}(\mathcal{A}))$. We will denote n' as $|\bigcup_{S \in \mathbb{T}(X)} S|$. Because X minimal non-very generic, $\text{rank } X + 1 = \nu(\mathbb{T}(X)) = \sum_{S \in \mathbb{T}(X)} (|S| - 2) = \sum_{S \in \mathbb{T}(X)} |S| - 2|\mathbb{T}(X)|$ and each line in $\bigcup_{S \in \mathbb{T}(X)} S$ is contained by at least two subset $S \in \mathbb{T}(X)$. Hence, $2n' \leq \sum_{S \in \mathbb{T}(X)} |S|$. Subtracting $2\nu(\mathbb{T}(X)) \geq 2|\mathbb{T}(X)|$ from both sides of this inequality, $2(n' - \nu(\mathbb{T}(X))) \leq 2(n' - |\mathbb{T}(X)|) \leq \nu(\mathbb{T}(X))$, then $2n'/3 \leq \nu(\mathbb{T}(X)) = \text{rank}(X) + 1$. Since $n' \geq 6$, we should consider the cases $n' = 6, 7, 8$. In the case $n' = 6, 7$, since $2n'/3 - 1 \leq \text{rank } X \leq n' - 3$, X is either as in Example 3.7 or these permutation. In the case $n' = 8$, since $13/3 \leq \text{rank } X \leq 5$, we obtain $\text{rank } X = 5$ and $\nu(\mathbb{T}(X)) = 6$. If $|S| = 3$ for all $S \in \mathbb{T}(X)$, then there exist two lines H_{i_1}, H_{i_2} in \mathcal{A} such that $|\{S \in \mathbb{T}(X) \mid i_l \in S\}| = 3$ for $l = 1, 2$. Then $\mathbb{T}(X)$ is a 8-ladder or degenerated 10-wheel. If $|S| = 3$ for some $S \in \mathbb{T}(X)$, then $\mathbb{T}(X)$ is a permutation of the 8-wheel. Then, the equations follow from Theorem 5.6 and Theorem 4.1. Hence, the following proposition holds.

Theorem 6.3. Let \mathcal{A} be a generic arrangement with 8 lines in a plane. The canonical representation of minimal non-very generic intersection in $\mathcal{L}(\mathcal{B}(\mathcal{A}))$ is one of the following 5 families or permutations of them: 6-wheel \mathbb{W}_6 , degenerated 8-wheel $(\mathbb{W}_8)_{8 \rightarrow 4}$, 8-wheel \mathbb{W}_8 , degeneration of indices 8 and 9 swapped 10-wheel $((89).\mathbb{W}_{10})_{10 \rightarrow 6} 9 \rightarrow 4$. The defining equations of these non-very generic varieties are as follows:

$$\begin{aligned} \mathbb{W}_6 &: \Delta_{21} \Delta_{43} \Delta_{65} - \Delta_{23} \Delta_{45} \Delta_{61} = 0, \\ (\mathbb{W}_8)_{8 \rightarrow 4} &: \Delta_{21} \Delta_{43} \Delta_{65} \Delta_{47} - \Delta_{23} \Delta_{45} \Delta_{67} \Delta_{41} = 0, \\ \mathbb{W}_8 &: \Delta_{21} \Delta_{43} \Delta_{65} \Delta_{87} - \Delta_{23} \Delta_{45} \Delta_{67} \Delta_{81} = 0, \\ \mathbb{L}_8 &: \Delta_{17} \Delta_{37} \Delta_{57} \Delta_{28} \Delta_{48} \Delta_{68} - \Delta_{18} \Delta_{38} \Delta_{58} \Delta_{27} \Delta_{47} \Delta_{67} = 0, \\ ((89).\mathbb{W}_{10})_{10 \rightarrow 6} 9 \rightarrow 4 &: \Delta_{21} \Delta_{43} \Delta_{65} \Delta_{47} \Delta_{68} - \Delta_{23} \Delta_{45} \Delta_{67} \Delta_{48} \Delta_{61} = 0. \end{aligned}$$

What happens if there are 9 lines or more? Since $2n'/3 - 1 \leq \text{rank } X \leq n' - 2$, the rank of X cannot be uniquely determined by n' . For example, let $n' = 9$, then $\text{rank } X = 5$ or 6. We give an example of the case $\text{rank } X = 5$.

Example 6.4. Let \mathcal{A} be a line arrangement with normals

$$(\alpha_1 \dots \alpha_9) = \begin{pmatrix} -4 & -3 & -2 & -1 & 0 & 1 & 2 & 3 & 4 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{pmatrix}.$$

And let $\mathbb{T}_0 = \{123, 456, 789, 147, 258, 369\}$. Consider the intersection X of $D_S (S \in \mathbb{T}_0)$ in $\mathcal{L}(\mathcal{B}(\mathcal{A}))$. We obtain

$$\begin{pmatrix} \alpha_{123} \\ \alpha_{456} \\ \alpha_{789} \\ \alpha_{147} \\ \alpha_{258} \\ \alpha_{369} \end{pmatrix} = \begin{pmatrix} -1 & 2 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 2 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 2 & -1 \\ -3 & 0 & 0 & 6 & 0 & 0 & -3 & 0 & 0 \\ 0 & -3 & 0 & 0 & 6 & 0 & 0 & -3 & 0 \\ 0 & 0 & -3 & 0 & 0 & 6 & 0 & 0 & -3 \end{pmatrix}.$$

The non-very generic intersection X of rank 5 is minimal because $\nu(\mathbb{T}_0 \setminus i) = \bigcap_{C \in \mathbb{T}_0 \setminus i} D_C = 4$ where $\mathbb{T}_0 \setminus i = \{C \in \mathbb{T}_0 \mid i \notin C\}$ for $i = 1, \dots, 9$. We have $\min \{\nu(\mathbb{T}') \mid \mathbb{T}' \in P(n, k), \mathbb{T} < \mathbb{T}'\} = \text{rank } X + 2$. It is easy to check that \mathbb{T}_0 satisfies the BBA-condition. Therefore, there exists an intersection Y which is greater than X in $\mathcal{L}(\mathcal{B}(\mathcal{A}))$




Figure 7: a representative of X in Example 6.4

by Theorem 3.9. By Proposition 6.2, \mathcal{A} belongs non-very generic varieties of nine 8-wheels $\mathbb{T}_1, \dots, \mathbb{T}_9$ and six degenerated 12-wheels $\mathbb{T}_{10}, \dots, \mathbb{T}_{15}$ as follows;

$$\begin{aligned} \mathbb{T}_1 &= \{123, 456, 147, 258, 3678\}, & \mathbb{T}_2 &= \{123, 789, 147, 258, 3459\}, & \mathbb{T}_3 &= \{123, 456, 147, 369, 2579\}, \\ \mathbb{T}_4 &= \{123, 789, 147, 369, 2468\}, & \mathbb{T}_5 &= \{123, 456, 258, 369, 1489\}, & \mathbb{T}_6 &= \{123, 789, 258, 369, 1567\}, \\ \mathbb{T}_7 &= \{456, 789, 147, 258, 1269\}, & \mathbb{T}_8 &= \{456, 789, 147, 369, 1358\}, & \mathbb{T}_9 &= \{456, 789, 258, 369, 2347\}, \\ \mathbb{T}_{10} &= \{123, 456, 789, 147, 258, 369, 159\}, & \mathbb{T}_{11} &= \{123, 456, 789, 147, 258, 369, 168\}, \\ \mathbb{T}_{12} &= \{123, 456, 789, 147, 258, 369, 249\}, & \mathbb{T}_{13} &= \{123, 456, 789, 147, 258, 369, 267\}, \\ \mathbb{T}_{14} &= \{123, 456, 789, 147, 258, 369, 348\}, & \mathbb{T}_{15} &= \{123, 456, 789, 147, 258, 369, 357\}. \end{aligned}$$

Then we obtain $V_{\mathbb{T}_0} \subset V_{\mathbb{T}_i}$ for $i \in [15]$. Note that, $V_{\mathbb{T}_1}, \dots, V_{\mathbb{T}_{15}}$ do not intersect transversal. Of course, no intersection is larger than X whose canonical representation is 8-wheel. However consider $X \cap \bigcap_{S \in \mathbb{T}_1} D_S$, then its rank is 6 and the canonical presentation is $\{123, 456, 147, 258, 36789\}$. Also, the intersections corresponding to these degenerated 12-wheels are the first example of intersections that are not minimal non-very generic intersections, but all proper subsets of canonical presentation that satisfy the BBA-condition.

References

- [ABF⁺23] Daniele Agostini, Taylor Brysiewicz, Claudia Fevola, Lukas Kühne, Bernd Sturmfels, and Simon Telen. Likelihood degenerations. *Adv. Math.*, 414:Paper No. 108863, 39, 2023. With an appendix by Thomas Lam.
- [Ath99] Christos A Athanasiadis. The largest intersection lattice of a discriminantal arrangement. *Beitr. Algebra Geom*, 40(2):283–289, 1999.
- [BB97] Margaret M Bayer and Keith A Brandt. Discriminantal arrangements, fiber polytopes and formality. *Journal of Algebraic Combinatorics*, 6(3):229–246, 1997.
- [CFW21] Beifang Chen, Houshan Fu, and Suijie Wang. Parallel translates of represented matroids. *Advances in Applied Mathematics*, 127:102176, 2021.
- [Che74] Alan L. C. Cheung. Adjoints of a geometry. *Canadian Mathematical Bulletin*, 17(3):363–365, 1974.
- [Cra84] Henry Crapo. Concurrence geometries. *Advances in Mathematics*, 54(3):278–301, dec 1984.

- [Cra85] Henry Crapo. The combinatorial theory of structures. In *Matroid Theory*, volume 40 of *Colloquia mathematica Societatis János Bolyai*, pages 107–213. North-Holland, 1985.
- [ER96] P. H. Edelman and V. Reiner. Free arrangements and rhombic tilings. *Discrete Comput. Geom.*, 15(3):307–340, 1996.
- [Fal94] Michael Falk. A note on discriminantal arrangements. *Proceedings of the American Mathematical Society*, 122(4):1221–1227, 1994.
- [FHJK23] Ragnar Freij-Hollanti, Relinde Jurrius, and Olga Kuznetsova. Combinatorial derived matroids. *Electron. J. Combin.*, 30(2):Paper No. 2.8, 28, 2023.
- [FW23] Houshan Fu and Suijie Wang. Modifications of hyperplane arrangements. *J. Combin. Theory Ser. A*, 200:Paper No. 105797, 34, 2023.
- [JP15] Relinde Jurrius and Ruud Pellikaan. The coset leader and list weight enumerator. In *Topics in finite fields*, volume 632 of *Contemp. Math.*, pages 229–251. Amer. Math. Soc., Providence, RI, 2015.
- [KNT12] Hiroshi Koizumi, Yasuhide Numata, and Akimichi Takemura. On intersection lattices of hyperplane arrangements generated by generic points. *ANNALS OF COMBINATORICS*, 16(4):789–813, 12 2012.
- [KV94] M. Kapranov and V. Voevodsky. Braided monoidal 2-categories and Manin-Schechtman higher braid groups. *J. Pure Appl. Algebra*, 92(3):241–267, 1994.
- [Lon80] Judith Q Longyear. The circuit basis in binary matroids. *Journal of Number Theory*, 12(1):71–76, 1980.
- [LS18] Anatoly Libgober and Simona Settepanella. Strata of discriminantal arrangements. *Journal of Singularities*, 18:440–454, 2018.
- [MS89] Yu. I. Manin and V. V. Schechtman. Arrangements of hyperplanes, higher braid groups and higher Bruhat orders. In *Algebraic number theory*, volume 17 of *Adv. Stud. Pure Math.*, pages 289–308. Academic Press, Boston, MA, 1989.
- [NT12] Yasuhide Numata and Akimichi Takemura. On computation of the characteristic polynomials of the discriminantal arrangements and the arrangements generated by generic points. *Harmony of Grobner Bases and the Modern Industrial Society, (Takayuki Hibi, editor)*, World Scientific, pages 228–252, 2012.
- [OT92] Peter Orlik and Hiroaki Terao. *Arrangements of hyperplanes*, volume 300 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1992.
- [OW19] James Oxley and Suijie Wang. Dependencies among dependencies in matroids. *The Electronic Journal of Combinatorics*, pages P3–46, 2019.
- [Pag19] Roberto Pagaria. Combinatorics of toric arrangements. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.*, 30(2):317–349, 2019.
- [Per11] Markus Perling. Divisorial cohomology vanishing on toric varieties. *Documenta Mathematica*, 16:209–251, 2011.
- [SS22] Takuya Saito and Simona Settepanella. Non-very generic arrangements in low dimension. *arXiv preprint arXiv:2202.04794*, 2022.
- [SSY17] Sumire Sawada, Simona Settepanella, and So Yamagata. Discriminantal arrangement, 3×3 minors of plücker matrix and hypersurfaces in grassmannian $Gr(3, \mathbf{C}^n)$. *Comptes Rendus Mathématique*, 355(11):1111–1120, 2017.
- [SSY19] Sumire Sawada, Simona Settepanella, and So Yamagata. Pappus’s theorem in grassmannian $Gr(3, \mathbf{C}^n)$. *Ars Mathematica Contemporanea*, 16(1):257–276, 2019.
- [SY22a] Simona Settepanella and So Yamagata. A linear condition for non-very generic discriminantal arrangements. *arXiv preprint arXiv:2205.04664*, 2022.
- [SY22b] Simona Settepanella and So Yamagata. On the non-very generic intersections in discriminantal arrangements. *C. R. Math. Acad. Sci. Paris*, 360:1027–1038, 2022.
- [Yam23] So Yamagata. A classification of combinatorial types of discriminantal arrangements. *J. Algebraic Combin.*, 57(4):1007–1032, 2023.