

Stochastic dynamics of two-compartment models with regulatory mechanisms for hematopoiesis

Ren-Yi Wang¹, Marek Kimmel², and Guodong Pang³

^{1,2}Department of Statistics, Rice University, Houston, TX, 77005, USA

³Department of Computational Applied Mathematics and Operations Research,
Rice University, Houston, TX, 77005, USA

^{1,2,3}`rw47,kimmel,gdpang@rice.edu`

May 7, 2024

Abstract

We present an asymptotic analysis of a stochastic two-compartmental cell proliferation system with regulatory mechanisms. We model the system as a state-dependent birth and death process. Proliferation of hematopoietic stem cells (HSCs) is regulated by population density of HSC-derived clones and differentiation of HSC is regulated by population density of HSCs. By scaling up the initial population, we show the density of dynamics converges in distribution to the solution of a system of ordinary differential equations (ODEs). The system of ODEs has a unique non-trivial equilibrium that is globally stable. Furthermore, we show the scaled fluctuation of the population converges in law to a linear diffusion with time-dependent coefficients. With initial data being Gaussian, the limit is a Gauss-Markov process, and it behaves like the FCLT limit under equilibrium with constant coefficients at large times. This is proved by establishing exponential convergence in the 2-Wasserstein metric for the associated Gaussian measures in a \mathcal{L}_2 Hilbert space. We apply our results to analyze and compare two regulatory mechanisms in the hematopoietic system. Simulations are conducted

to verify our large-scale and long-time approximation of the dynamics. We demonstrate some regulatory mechanisms are efficient (converge to steady state rapidly) but not effective (have large fluctuation around the steady state).

Keywords: Hematopoiesis; Regulatory mechanism; Stochastic two-compartment model; Functional limit theorems; Stability; Large-time asymptotic behavior

1 Introduction

Hematopoiesis is the blood cell production process. A hematopoietic stem cell (HSC) can self-renew, differentiate, or die. When a HSC differentiates, it becomes a multi-potent progenitor cell. A multi-potent progenitor cell will further differentiate until mature cells are produced, leading to a clone that will eventually become extinct. To maintain homeostasis (stability of the cell production system), regulatory mechanisms are required. In Duncan et al. (2005) and Blank et al. (2008), the authors point out Notch and Wnt signaling pathways play crucial roles in the stability of hematopoietic system. Specifically, Notch signaling regulates differentiation and Wnt signaling regulates self-renewal. In this paper, we model Notch signaling by a short-range feedback and Wnt signaling by a long-range feedback. Kopan (2012) and Bigas and Espinosa (2012) contain justifications for modeling Notch signaling as a short-range feedback. Mechanisms that allow Wnt to achieve long-range signaling are detailed in Nusse et al. (2008) and Buechling and Boutros (2011).

Deterministic models for cell production systems with regulatory mechanisms are well-studied in the literature. Getto et al. (2013) studied global stability of two-compartment models with regulatory mechanisms and Marciniak-Czochra et al. (2009) conducted simulations to compare efficiencies among various feedback mechanisms in multicompartment models. Differential equation models for fluctuation (oscillation) of hematopoietic system are studied in Knauer et al. (2020) and Bonnet et al. (2021). Bonnet and Méléard (2021) proposed a stochastic model without regulatory mechanisms to explain unexpected fluctuations of the mature blood cell number. In this paper, we introduce a framework to analyze efficiency and effectiveness of regulatory mechanisms in stochastic two-compartment models.

We view hematopoiesis as a two-compartment model as in Catlin et al. (2011) with first compartment $N_0^{(r)}$ containing HSCs and the second compartment $N_1^{(r)}$ containing HSC-derived clones. Initial conditions are $N_0^{(r)}(0) = n_0^{(r)}$ and $N_1^{(r)}(0) = n_1^{(r)}$, where r is a scaling parameter. HSCs self-renew at rate α and differentiate at rate γ . α depends on the density of derived clones ($\bar{N}_1^{(r)} = N_1^{(r)}/r$) and γ depends on the density of HSCs ($\bar{N}_0^{(r)} = N_0^{(r)}/r$). We are interested in the dynamics when initial population sizes are large (i.e. $n^{(r)} \rightarrow \infty$ as $r \rightarrow \infty$). With conditions for α and γ in Section 2.1, we derive a functional law of large numbers (FLLN) as $r \rightarrow \infty$ in Section 3.1. The FLLN says the density of cell populations $\bar{\mathbf{N}}^{(r)}$ converges weakly to a deterministic function $\bar{\mathbf{N}}$, satisfying a system of ordinary differential equations (ODEs) with a globally stable non-trivial equilibrium. In Section 3.2, we consider the difference between the density of cell dynamics and FLLN limit, and then scale up the difference by \sqrt{r} , denoted as $\hat{\mathbf{N}}^{(r)}$, to derive a functional central limit theorem (FCLT), which has a linear diffusion limit process $\hat{\mathbf{N}}$ with time-dependent coefficients. This diffusion describes the fluctuations of the dynamics around its FLLN limit function in the large-scale system. We show that the time-dependent coefficients of the limiting diffusion $\hat{\mathbf{N}}$ become constant when the FLLN limit starts at its equilibrium in Section 4.1, resulting in a stationary linear diffusion process \mathbf{G}^* . An immediate consequence is that the covariance function of the limiting diffusion under the FLLN equilibrium converges to a constant, which can be solved by a linear equation when we specify a regulatory mechanism (the functional form of α and γ). Next, we show that the limiting diffusion $\hat{\mathbf{N}}$ resembles the linear diffusion \mathbf{G}^* under the FLLN equilibrium at large times. Specifically, we show that the process $\mathbf{G}_T(t) = \hat{\mathbf{N}}(T+t)$ converges to $\mathbf{G}^*(t)$ over $[0, M]$ for arbitrary $M > 0$ as $T \rightarrow \infty$, where we cast these processes to Gaussian measures in an \mathcal{L}_2 Hilbert space and use the 2-Wasserstein metric. We derive the explicit 2-Wasserstein metric for the Gaussian measures corresponding to $\mathbf{G}_T(t)$ and \mathbf{G}^* and show that the convergence is exponentially fast.

Finally, in Section 5, we apply our results to two specific regulatory mechanisms and conduct simulations with parameter values assumed by Catlin et al. (2011) to validate our large-scale and long-time approximation of the dynamics. We demonstrate some regulatory mechanisms are efficient (converge to steady state rapidly) but not effective (have large limiting variance).

2 Model and Assumptions

Denote $N_0^{(r)}(t)$ as the stem cell (type 0 individual) population with initial condition $n_0^{(r)}$ and $N_1^{(r)}(t)$ as the number of HSC-derived clones (type 1 individuals) with initial condition $n_1^{(r)}$. Define the density by $\bar{N}_i^{(r)} = N_i^{(r)}/r$, $i \in \{0, 1\}$. A stem cell can either proliferate (denote this action by $(0, b)$) with rate α and increase the stem cell population by 1 or differentiate (denoted by $(0, di)$) with rate γ to initiate a clone. A HSC-derived clone exhausts with rate δ (denoted by $(1, d)$). Let $\mathcal{A} = \{(0, b), (0, di), (1, d)\}$ denote the set of actions for type 0 and type 1 individuals. Notice that we neglect the action for stem cell death since death rate for the stem cell is small (cf. Domen et al. (2000) and Catlin et al. (2011)). Hence, if the birth rate for stem cell is α and death rate is $\beta \ll \alpha$, we can absorb the death rate into the birth rate ($\tilde{\alpha} = \alpha - \beta$; $\tilde{\beta} = 0$) to approximate the dynamics. Our model of the hematopoietic system with regulatory feedback is defined by the following transitions,

$$\begin{aligned} (N_0^{(r)}, N_1^{(r)}) &\rightarrow (N_0^{(r)} + 1, N_1^{(r)}) && \text{at rate } \alpha(\bar{N}_1^{(r)})N_0^{(r)} \\ (N_0^{(r)}, N_1^{(r)}) &\rightarrow (N_0^{(r)} - 1, N_1^{(r)} + 1) && \text{at rate } \gamma(\bar{N}_0^{(r)})N_0^{(r)} \\ (N_0^{(r)}, N_1^{(r)}) &\rightarrow (N_0^{(r)}, N_1^{(r)} - 1) && \text{at rate } \delta N_1^{(r)}. \end{aligned}$$

2.1 Construction of Dynamics

In this section, we develop a general framework to analyze the stochastic two-compartment model with regulatory feedback influencing rates of self-renewal α and differentiation γ . We introduce three sequences of independent Poisson random measures $\mathcal{N}_a^{(r)}$, $a \in \mathcal{A}$ on \mathbb{R}_+^2 with mean measure being Lebesgue. The natural filtration is

$$\mathcal{F}_t^{(r)} = \sigma\{\mathcal{N}_a^{(r)}([0, s], B) \mid s \leq t, B \in \mathcal{B}(\mathbb{R}_+)\}. \quad (1)$$

Following the notation in Bonnet and Méléard (2021), we set up our dynamics as follows. For all

$t \geq 0$,

$$\begin{aligned}
N_0^{(r)}(t) &= n_0^{(r)} - \int_0^t \int_{\mathbb{R}_+} \mathbf{1}_{\{u \leq \gamma(\bar{N}_0^{(r)}(s-))N_0^{(r)}(s-)\}} \mathcal{N}_{(0,di)}^{(r)}(du, ds) \\
&\quad + \int_0^t \int_{\mathbb{R}_+} \mathbf{1}_{\{u \leq \alpha(\bar{N}_1^{(r)}(s-))N_0^{(r)}(s-)\}} \mathcal{N}_{(0,b)}^{(r)}(du, ds) \\
N_1^{(r)}(t) &= n_1^{(r)} - \int_0^t \int_{\mathbb{R}_+} \mathbf{1}_{\{u \leq \delta N_1^{(r)}(s-)\}} \mathcal{N}_{(1,d)}^{(r)}(du, ds) \\
&\quad + \int_0^t \int_{\mathbb{R}_+} \mathbf{1}_{\{u \leq \gamma(\bar{N}_0^{(r)}(s-))N_0^{(r)}(s-)\}} \mathcal{N}_{(0,di)}^{(r)}(du, ds).
\end{aligned}$$

We write our model in a more compact notation as in Pang et al. (2007). Let $\{P_a^{(r)} \mid r \in \mathbb{N}, a \in \mathcal{A}\}$ be a set of independent rate 1 Poisson processes. Then we can rewrite the dynamics as

$$\begin{aligned}
N_0^{(r)}(t) &= n_0^{(r)} - P_{(0,di)}^{(r)} \left(\int_0^t \gamma(\bar{N}_0^{(r)}(s))N_0^{(r)}(s)ds \right) + P_{(0,b)}^{(r)} \left(\int_0^t \alpha(\bar{N}_1^{(r)}(s))N_0^{(r)}(s)ds \right) \\
N_1^{(r)}(t) &= n_1^{(r)} - P_{(1,d)}^{(r)} \left(\delta \int_0^t N_1^{(r)}(s)ds \right) + P_{(0,di)}^{(r)} \left(\int_0^t \gamma(\bar{N}_0^{(r)}(s))N_0^{(r)}(s)ds \right).
\end{aligned}$$

We assume $\alpha, \gamma \in \mathcal{C}^2(\mathbb{R}_+; \mathbb{R}_+)$ and for all x, y , $\alpha'(y) \leq 0$; $\gamma'(x) \geq 0$. Furthermore, we assume $\alpha(0) > \gamma(0) > 0$ and

$$\lim_{y \rightarrow \infty} \alpha(y) < \gamma(0) \text{ or } \alpha(0) < \lim_{x \rightarrow \infty} \gamma(x). \tag{2}$$

The assumption $\alpha(0) > \gamma(0)$ implies that type 0 population is growing when regulatory mechanisms are absent (for instance, in precancerous or cancer state) and the subsequent assumption provides stability to the system as we will see in the next section.

3 Large-scale Approximations

In this section, we scale up initial population sizes to obtain limiting dynamics for density (FLLN) and fluctuation around equilibrium (FCLT). More specifically, we show global stability for the FLLN and study asymptotic behaviors for the FCLT under equilibrium.

3.1 Functional Law of Large Numbers

Assume $\frac{\mathbf{n}^{(r)}}{r} \rightarrow \bar{\mathbf{n}} > 0$ as $r \rightarrow \infty$ and denote the scaled process as $\bar{\mathbf{N}}^{(r)}$ where

$$\bar{N}_0^{(r)} = \frac{N_0^{(r)}}{r}; \quad \bar{N}_1^{(r)} = \frac{N_1^{(r)}}{r}.$$

The scaled dynamics are

$$\begin{aligned} \bar{N}_0^{(r)}(t) &= \bar{n}_0^{(r)} - \frac{1}{r} P_{(0,di)}^{(r)} \left(r \int_0^t \gamma(\bar{N}_0^{(r)}(s)) \bar{N}_0^{(r)}(s) ds \right) + \frac{1}{r} P_{(0,b)}^{(r)} \left(r \int_0^t \alpha(\bar{N}_1^{(r)}(s)) \bar{N}_0^{(r)}(s) ds \right) \\ \bar{N}_1^{(r)}(t) &= \bar{n}_1^{(r)} - \frac{1}{r} P_{(1,d)}^{(r)} \left(r \int_0^t \delta \bar{N}_1^{(r)}(s) ds \right) + \frac{1}{r} P_{(0,di)}^{(r)} \left(r \int_0^t \gamma(\bar{N}_0^{(r)}(s)) \bar{N}_0^{(r)}(s) ds \right). \end{aligned}$$

Denote the centered and scaled Poisson processes by

$$\begin{aligned} M_{(0,b)}^{(r)} &= \frac{1}{r} P_{(0,b)}^{(r)} \left(r \int_0^t \alpha(\bar{N}_1^{(r)}(s)) \bar{N}_0^{(r)}(s) ds \right) - \int_0^t \alpha(\bar{N}_1^{(r)}(s)) \bar{N}_0^{(r)}(s) ds \\ M_{(0,di)}^{(r)} &= \frac{1}{r} P_{(0,di)}^{(r)} \left(r \int_0^t \gamma(\bar{N}_0^{(r)}(s)) \bar{N}_0^{(r)}(s) ds \right) - \int_0^t \gamma(\bar{N}_0^{(r)}(s)) \bar{N}_0^{(r)}(s) ds \\ M_{(1,d)}^{(r)} &= \frac{1}{r} P_{(1,d)}^{(r)} \left(r \int_0^t \delta \bar{N}_1^{(r)}(s) ds \right) - \int_0^t \delta \bar{N}_1^{(r)}(s) ds. \end{aligned}$$

In Proposition 1 we show $M_{(0,b)}^{(r)}$, $M_{(0,di)}^{(r)}$, and $M_{(1,d)}^{(r)}$ are all L^2 martingales. In Proposition 2, we show for all $a \in \mathcal{A}$, $M_a^{(r)} \Rightarrow 0$. Denote $\bar{\mathbf{N}}^{(r)} = (\bar{N}_0^{(r)}, \bar{N}_1^{(r)})^\top$, $\bar{\mathbf{n}}^{(r)} = (\bar{n}_0^{(r)}, \bar{n}_1^{(r)})^\top$ and rewrite the scaled dynamics as follows,

$$\bar{\mathbf{N}}^{(r)} = I(\bar{\mathbf{n}}^{(r)}, \mathbf{M}^{(r)}); \quad \bar{\mathbf{N}}^{(r)}(t) = \bar{\mathbf{n}}^{(r)} + \mathbf{M}^{(r)}(t) + \int_0^t \mathbf{F}(\bar{\mathbf{N}}^{(r)}), \quad (3)$$

where $\mathbf{M}^{(r)}$ includes all the martingale terms and \mathbf{F} is defined by

$$\mathbf{F}((x_0, x_1)^\top) = \begin{pmatrix} \alpha(x_1)x_0 - \gamma(x_0)x_0 \\ \gamma(x_0)x_0 - \delta x_1 \end{pmatrix}.$$

In Lemma 1, we show the operator I defined by Eq. (3) is continuous at $(\mathbf{b}, \mathbf{0})$ for all $\mathbf{b} > \mathbf{0}$.

Since $\bar{\mathbf{n}}^{(r)} \rightarrow \bar{\mathbf{n}} > 0$ and $\mathbf{M}^{(r)} \Rightarrow \mathbf{0}$, we have $\bar{\mathbf{N}}^{(r)} = I(\bar{\mathbf{n}}^{(r)}, \mathbf{M}^{(r)}) \Rightarrow I(\bar{\mathbf{n}}, \mathbf{0})$ by Theorem 1. Denote the limit by $\bar{\mathbf{N}}$, then it is a solution to the following IVP,

$$\begin{aligned}\bar{N}'_0(t) &= \alpha(\bar{N}_1(t))\bar{N}_0(t) - \gamma(\bar{N}_0(t))\bar{N}_0(t) \\ \bar{N}'_1(t) &= \gamma(\bar{N}_0(t))\bar{N}_0(t) - \delta\bar{N}_1(t) \\ \bar{N}_0(0) &= \bar{n}_0; \bar{N}_1(0) = \bar{n}_1.\end{aligned}$$

Notice that since $\bar{\mathbf{N}}$ is deterministic, $\bar{\mathbf{N}}^{(r)} \rightarrow \bar{\mathbf{N}}$ in probability as well. By setting derivatives to zeros and denote the non-trivial equilibrium by $\bar{\mathbf{N}}^*$, we have

$$\begin{aligned}\alpha(\bar{N}_1^*) &= \gamma(\bar{N}_0^*) \\ \delta\bar{N}_1^* &= \gamma(\bar{N}_0^*)\bar{N}_0^*.\end{aligned}$$

From the second relation, we have

$$\begin{aligned}\bar{N}_1^* &= \frac{\gamma(\bar{N}_0^*)\bar{N}_0^*}{\delta} \\ \implies \alpha\left(\frac{\gamma(\bar{N}_0^*)\bar{N}_0^*}{\delta}\right) &= \gamma(\bar{N}_0^*).\end{aligned}$$

Since $\alpha(0) > \gamma(0)$, the expression has at least a solution. By monotonicity of α and γ , we deduce the non-trivial steady state $\bar{\mathbf{N}}^*$ is unique. The Jacobian of \mathbf{F} evaluated at the equilibrium is

$$A^* = \begin{pmatrix} -\gamma'(\bar{N}_0^*)\bar{N}_0^* & \alpha'(\bar{N}_1^*)\bar{N}_0^* \\ \gamma'(\bar{N}_0^*)\bar{N}_0^* + \gamma(\bar{N}_0^*) & -\delta \end{pmatrix}.$$

Since $\bar{N}_0^* > 0$ and $\bar{N}_1^* > 0$, we have

$$\begin{aligned}Tr(A^*) &= -\gamma'(\bar{N}_0^*)\bar{N}_0^* - \delta < 0 \\ \det(A^*) &= \delta\gamma'(\bar{N}_0^*)\bar{N}_0^* - \alpha'(\bar{N}_1^*)\bar{N}_0^*[\gamma'(\bar{N}_0^*)\bar{N}_0^* + \gamma(\bar{N}_0^*)] > 0.\end{aligned}$$

Hence, the system is locally asymptotically stable. Moreover, define $\lambda = \max\{\Re(\sigma(A^*))\}$, then for every $\eta \in (\lambda, 0)$, there is a neighborhood \mathcal{V}_η of $\bar{\mathbf{N}}^*$ such that for all initial data $\bar{\mathbf{n}} \in \mathcal{V}_\eta$,

$$\|\bar{\mathbf{N}}(t) - \bar{\mathbf{N}}^*\| = O(\exp(\eta t)) \text{ as } t \rightarrow \infty. \quad (4)$$

In Theorem 3, we show the system is also globally asymptotically stable. As a consequence, for any fixed $\eta \in (\lambda, 0)$ and initial data $\bar{\mathbf{n}}$,

$$\|\bar{\mathbf{N}}(t) - \bar{\mathbf{N}}^*\| = o(\exp(\eta t)) \text{ as } t \rightarrow \infty. \quad (5)$$

3.2 Functional Central Limit Theorem

Now we further assume $\hat{\mathbf{n}}^{(r)} = \sqrt{r}(\bar{\mathbf{n}}^{(r)} - \bar{\mathbf{n}}) \Rightarrow \hat{\mathbf{n}} \sim N(\mathbf{u}, U)$ and define $\hat{\mathbf{N}}^{(r)} = \sqrt{r}(\bar{\mathbf{N}}^{(r)} - \bar{\mathbf{N}})$. The scaled dynamics become

$$\begin{aligned} \hat{N}_0^{(r)}(t) &= \hat{n}_0 + \sqrt{r}M_{(0,b)}^{(r)}(t) - \sqrt{r}M_{(0,di)}^{(r)}(t) \\ &\quad + \sqrt{r} \int_0^t \alpha(\bar{N}_1^{(r)}(s))\bar{N}_0^{(r)}(s)ds - \alpha(\bar{N}_1(s))\bar{N}_0(s) - \sqrt{r} \int_0^t \gamma(\bar{N}_0^{(r)}(s))\bar{N}_0^{(r)}(s) - \gamma(\bar{N}_0(s))\bar{N}_0(s)ds \\ \hat{N}_1^{(r)}(t) &= \hat{n}_1 + \sqrt{r}M_{(0,di)}^{(r)}(t) - \sqrt{r}M_{(1,d)}^{(r)}(t) \\ &\quad - \sqrt{r} \int_0^t \gamma(\bar{N}_0^{(r)}(s))\bar{N}_0^{(r)}(s) - \gamma(\bar{N}_0(s))\bar{N}_0(s)ds + \sqrt{r} \int_0^t \delta\bar{N}_1^{(r)}(s) - \delta\bar{N}_1(s)ds. \end{aligned}$$

We rewrite the scaled dynamics as

$$\hat{\mathbf{N}}^{(r)} = \hat{\mathbf{n}}^{(r)} + \hat{\mathbf{M}}^{(r)} + \sqrt{r} \int_0^t \mathbf{F}(\bar{\mathbf{N}}^{(r)}(s)) - \mathbf{F}(\bar{\mathbf{N}}(s))ds. \quad (6)$$

In Lemma 2, we show $\{\hat{\mathbf{N}}^{(r)}\}$ is stochastically bounded and $\hat{\mathbf{M}}^{(r)} \Rightarrow \hat{\mathbf{M}}$, where

$$\hat{\mathbf{M}}(t) = \begin{pmatrix} B_{(0,b)}(\int_0^t \alpha(\bar{N}_1(s))\bar{N}_0(s)ds) - B_{(0,di)}(\int_0^t \gamma(\bar{N}_0(s))\bar{N}_0(s)ds) \\ B_{(0,di)}(\int_0^t \gamma(\bar{N}_0(s))\bar{N}_0(s)ds) - B_{(1,d)}(\int_0^t \delta\bar{N}_1(s)ds) \end{pmatrix}.$$

$\{B_a \mid a \in \mathcal{A}\}$ is a set of IID standard Brownian motions. In Theorem 2, we show $\hat{\mathbf{N}}^{(r)} \Rightarrow \hat{\mathbf{N}}$ with

$$\hat{\mathbf{N}}(t) = \hat{\mathbf{n}} + \hat{\mathbf{M}}(t) + \int_0^t \nabla \mathbf{F}(\bar{\mathbf{N}}(s)) \hat{\mathbf{N}}(s) ds.$$

The FCLT limit process $\hat{\mathbf{N}}$ is a linear diffusion with time-dependent parameters

$$d\hat{\mathbf{N}}(t) = A(t)\hat{\mathbf{N}}(t)dt + \sigma(t)d\mathbf{B}(t),$$

where

$$A(t) = \nabla \mathbf{F}(\bar{\mathbf{N}}(t)); \quad \sigma(t) = \begin{pmatrix} \sqrt{\alpha(\bar{N}_1(t))\bar{N}_0(t)} & -\sqrt{\gamma(\bar{N}_0(t))\bar{N}_0(t)} & 0 \\ 0 & \sqrt{\gamma(\bar{N}_0(t))\bar{N}_0(t)} & -\sqrt{\delta\bar{N}_1(t)} \end{pmatrix}. \quad (7)$$

Since the initial condition $\hat{\mathbf{n}}$ is Gaussian, $\hat{\mathbf{N}}$ is a Gauss-Markov process and it can be characterized by its mean function and autocovariance function as in Section 5.6 of Karatzas and Shreve (2012).

3.3 FCLT Under Equilibrium

If $\bar{\mathbf{n}} = \bar{\mathbf{N}}^*$, the time-dependent coefficients of the limiting linear diffusion become constant as follows:

$$A^* = \begin{pmatrix} -\gamma'(\bar{N}_0^*)\bar{N}_0^* & \alpha'(\bar{N}_1^*)\bar{N}_0^* \\ \gamma'(\bar{N}_0^*)\bar{N}_0^* + \gamma(\bar{N}_0^*) & -\delta \end{pmatrix}; \quad \sigma^* = \sqrt{\gamma(\bar{N}_0^*)\bar{N}_0^*} \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \end{pmatrix}.$$

We henceforth assume A^* has distinct eigenvalues. This assumption can be easily achieved by adding small perturbations to α or γ . The fundamental matrix $\Phi(t)$ for the mean function satisfies the matrix differential equation

$$\Phi'(t) = A^*\Phi(t); \Phi(0) = I_2.$$

Hence, the mean function is $\mathbf{m}(t) = \exp(A^*t)\mathbf{u} = O(e^{\lambda t})$, which converges to 0 exponentially fast.

The variance function $V(t)$ satisfies

$$V'(t) = A^*V(t) + [A^*V(t)]^\top + \sigma^*[\sigma^*]^\top; \quad V(0) = U.$$

Define $\Sigma^* = \sigma^*[\sigma^*]^\top$ and

$$W(t) = \begin{pmatrix} V_{1,1}(t) \\ V_{1,2}(t) \\ V_{2,2}(t) \end{pmatrix}; \quad B^* = \begin{pmatrix} 2A_{1,1}^* & 2A_{1,2}^* & 0 \\ A_{2,1}^* & A_{1,1}^* + A_{2,2}^* & A_{1,2}^* \\ 0 & 2A_{2,1}^* & 2A_{2,2}^* \end{pmatrix}; \quad S^* = \begin{pmatrix} \Sigma_{1,1}^* \\ \Sigma_{1,2}^* \\ \Sigma_{2,2}^* \end{pmatrix}.$$

We write the matrix differential equation as a vector differential equation

$$W'(t) = B^*W(t) + S^*.$$

Since the spectrum of B^* is $\sigma(B^*) = \sigma(A^*) \cup \{Tr(A^*)\}$, $\max\{\Re(\sigma(B^*))\} = \lambda < 0$. Hence,

$$W(t) = \exp(B^*t)W(0) + \int_0^t \exp(B^*(t-s))S^*ds \rightarrow -(B^*)^{-1}S^* \text{ as } t \rightarrow \infty.$$

Define $W^* = -(B^*)^{-1}S^*$. Then,

$$\begin{aligned} (W(t) - W^*)' &= B^*(W(t) - W^*) \\ \implies \|W(t) - W^*\| &= O(\exp(\lambda t)). \end{aligned}$$

From W^* , we deduce the limit V^* for the variance function satisfies

$$A^*V^* + [A^*V^*]^\top = -\Sigma^*. \tag{8}$$

From this relation between A^* and V^* , it seems reasonable that a smaller λ implies smaller limiting variances. However, this is not the case as we will see in Section 5. It is possible for regulatory mechanism 1 to converge to steady state faster than regulatory mechanism 2 (more

efficient) but feedback 1 has larger limiting variances (less effective).

4 Asymptotic Analysis

4.1 Large-time behavior of FCLT limit process

In this section, we discard the assumption $\bar{\mathbf{n}} = \bar{\mathbf{N}}^*$ and study the long-time behavior of the process $\hat{\mathbf{N}}$. Since \mathbf{F} is \mathcal{C}^2 , $\nabla\mathbf{F}$ is locally Lipschitz. Eq. (5) in Section 3.1 suggests the trajectory of \bar{N} lies in a compact set. Hence, there exists a constant C such that for all $\eta \in (\lambda, 0)$,

$$\|A(t) - A^*\| = \|\nabla\mathbf{F}(\bar{\mathbf{N}}(t)) - \nabla\mathbf{F}(\bar{\mathbf{N}}^*)\| \leq C\|\bar{\mathbf{N}}(t) - \bar{\mathbf{N}}^*\| = o(\exp(\eta t)). \quad (9)$$

Define $R(t) = A(t) - A^*$, we have $\int_0^\infty \|R(t)\| dt < \infty$. Therefore, the fundamental matrix for the mean function satisfies

$$\Phi'(t) = [A^* + R(t)]\Phi(t); \quad \Phi(0) = I_2.$$

By assumption in Section 3.3, A^* has distinct eigenvalues, so it is diagonalizable. We apply Theorem 2.7 (Levinson's Fundamental Theorem) in Bodine et al. (2015) to conclude as $t \rightarrow \infty$,

$$\Phi(t) = [1 + o(1)] \exp(A^* t).$$

Hence, $\mathbf{m}(t) = [1 + o(1)] \exp(A^* t) \mathbf{u} = O(\exp(\lambda t))$.

Define $\Sigma(t) = \sigma(t)\sigma^\top(t)$ and

$$W(t) = \begin{pmatrix} V_{1,1}(t) \\ V_{1,2}(t) \\ V_{2,2}(t) \end{pmatrix}; B(t) = \begin{pmatrix} 2A_{1,1}(t) & 2A_{1,2}(t) & 0 \\ A_{2,1}(t) & A_{1,1}(t) + A_{2,2}(t) & A_{1,2}(t) \\ 0 & 2A_{2,1}(t) & 2A_{2,2}(t) \end{pmatrix}; S(t) = \begin{pmatrix} \Sigma_{1,1}(t) \\ \Sigma_{1,2}(t) \\ \Sigma_{2,2}(t) \end{pmatrix}.$$

The variance function satisfies $W'(t) = B(t)W(t) + S(t)$. By Eq. (5) in Section 3.1, we have $\|B(t) - B^*\| = o(\exp(\eta t))$ and $\|S(t) - S^*\| = o(\exp(\eta t))$ for all $\eta \in (\lambda, 0)$. It is evident that $W(t) \rightarrow W^*$ as $t \rightarrow \infty$. To analyze the rate of convergence, we write the system in terms of the difference

$$(W(t) - W^*)' = (B^* + B(t) - B^*)(W(t) - W^*) + B(t)W^* + S(t).$$

By exponential stability, $\|B(t)W^* + S(t)\| = o(\exp(\eta t))$. In addition, since $\|W(t) - W^*\| \rightarrow 0$, $\|(B(t) - B^*)(W(t) - W^*)\| = o(\exp(\eta t))$ for all $\eta \in (\lambda, 0)$ and we may write the system as

$$(W(t) - W^*)' = B^*(W(t) - W^*) + Err(t),$$

where $\|Err(t)\| = o(\exp(\eta t))$ for all $\eta \in (\lambda, 0)$. The solution to this IVP is

$$W(t) - W^* = \exp(B^*t)(W(0) - W^*) + \int_0^t \exp(B^*(t-s))Err(s)ds.$$

The first term is $O(\exp(\lambda t))$. For the second term, for a fixed $\eta \in (\lambda, 0)$, there exists a constant $C > 0$ such that

$$\begin{aligned} \left\| \int_0^t \exp(B^*(t-s))Err(s)ds \right\| &\leq C \int_0^t \exp(\lambda(t-s)) \exp(\eta s) ds \\ &= \frac{C}{\eta - \lambda} (\exp(\eta t) - \exp(\lambda t)) = O(\exp(\eta t)). \end{aligned}$$

We conclude

$$\|W(t) - W^*\| = o(\exp(\eta t)), \quad \forall \eta \in (\lambda, 0). \quad (10)$$

4.2 Convergence in Wasserstein Metric

In this section, we show that the limiting diffusion of the FCLT at large times resembles a Gaussian process \mathbf{G}^* with mean function $\mathbf{m}^*(t) = \mathbf{0}$ and autocovariance function $K^*(s, t) = V^* \exp((t - s)[A^*]^\top)$ for $s \leq t$. Let $T \in \mathbb{N}$ and define $\mathbf{G}_T(t) = \hat{\mathbf{N}}(T + t)$ with mean function \mathbf{m}_T and

autocovariance function K_T , where

$$\mathbf{m}_T(t) = \exp\left(\int_0^t A(s+T)ds\right)\mathbf{u}; \quad K_T(s, t) = V(s+T) \exp\left(\int_s^t A^\top(u+T)du\right), \quad s \leq t. \quad (11)$$

We will show \mathbf{G}_T converges to \mathbf{G}^* as $T \rightarrow \infty$. To introduce the notion of convergence, we follow Minh (2023) to establish a one-to-one correspondence between the space of Gaussian processes and the space of Gaussian measures on $\mathcal{H} = \mathcal{L}_2([0, M]^2, \mathcal{B}([0, M]^2), Leb)$, where $M > 0$ is fixed. \mathcal{H} is a separable Hilbert space. For a Gaussian process with mean function $\mathbf{m} \in \mathcal{H}$ and covariance function $K(s, t)$ being continuous, symmetric, and positive-definite, we invoke Mercer's theorem to induce a Gaussian measure $\mathcal{N}(\mathbf{m}, C_K)$. $C_K : \mathcal{H} \rightarrow \mathcal{H}$ is the covariance operator defined by

$$(C_K)f(s) = \int_{[0, M]} K(s, t)f(t)dt.$$

The covariance operator C_K is of trace class, self-adjoint, and positive.

Since mean functions $\mathbf{m}_T, \mathbf{m}^* \in \mathcal{H}$ and the autocovariance functions K_T, K_* are continuous, symmetric, and positive-definite, we induce a sequence of Gaussian measures $\mathcal{G}_T \sim \mathcal{N}(\mathbf{m}_T, C_T)$ and $\mathcal{G}^* \sim \mathcal{N}(\mathbf{0}, C_*)$. The square of the 2-Wasserstein distance is

$$W_2^2(\mathcal{G}_T, \mathcal{G}_*) = \|\mathbf{m}_T\|_2^2 + Tr(C_T + C_* - 2(C_*^{1/2}C_TC_*^{1/2})^{1/2}).$$

In Theorem 4, we show for any fixed M , $W_2(\mathcal{G}_T, \mathcal{G}_*) = o(\exp(\frac{\eta}{2}T))$ for all $\eta \in (\lambda, 0)$. To interpret this result, we focus on $\hat{\mathbf{N}}$ in a moving interval $[T, T+M]$. As we increase $T \rightarrow \infty$, the dynamics of $\hat{\mathbf{N}}$ will become more and more similar to that of \mathbf{G}^* . The length of the interval M can be arbitrarily large.

5 Applications

We apply our framework to study two regulatory mechanisms. The purpose of this section is to demonstrate high efficiency (fast convergence of FLLN dynamics) does not guarantee high

effectiveness (small limiting variance for FCLT dynamics) as claimed in Section 3.3. Let $\alpha > \gamma > 0$ be constants. We set up two regulatory mechanisms

$$\textbf{Model 1: } \alpha(y) = \frac{\alpha}{1 + \kappa_y y}; \gamma(x) = \gamma + \kappa_x x^2;$$

$$\textbf{Model 2: } \alpha(y) = \max(\alpha - \kappa_y y^2, 0); \gamma(x) = \gamma + \kappa_x x.$$

Notice that in model 2, $\alpha(\cdot)$ is not in \mathcal{C}^2 and this issue can be easily rectified by convolution with a mollifier ϕ_ϵ . By selecting ϵ small enough, our theory will approximate the dynamics of model 2.

We take estimated parameters in Catlin et al. (2011) as our base parameters for simulation. That is, for a human hematopoietic system, we have $\alpha = \frac{1}{40} - \frac{1}{285.7}$ per week, $\gamma = \frac{1}{56.1}$ per week, and $\delta = \frac{1}{6.7}$ per week. We simulate the dynamics under a pathological state with parameter values in Table 1 (faster self-renewal and slower differentiation). Since our approximation requires $r \rightarrow \infty$, we set r to be a large number $r = R = 2000$ in the simulation. To compare two regulation mechanisms, we set κ_x, κ_y so that $\bar{N}_0^* = \frac{11000}{R} = 5.5$ and $\bar{N}_0^* = \frac{1275}{R} = 0.6375$ for all simulations. The numbers 11000 and 1275 are estimated steady-states for HSCs and derived clones in Catlin et al. (2011). We set initial conditions to be $\bar{n}_0 = 11$ and $\bar{n}_1 = \frac{\gamma \bar{n}_0}{\alpha - \gamma + \delta}$. This initial condition corresponds to letting the pathological system, defined as an exponentially growing population without regulation, progress until the system reaches $\bar{n}_0 = 11$. The relation between \bar{n}_0 and \bar{n}_1 is based on the formula in Catlin et al. (2011) on page 4463.

Model Parameters						
	α	γ	δ	κ_x	κ_y	λ
Model 1	$\frac{1.5}{40} - \frac{1.5}{285.7}$	$\frac{1}{1.5 \cdot 56.1}$	$\frac{1}{6.7}$	0.000179052	1.355547131	-0.02671738
Model 2	$\frac{1.5}{40} - \frac{1.5}{285.7}$	$\frac{1}{1.5 \cdot 56.1}$	$\frac{1}{6.7}$	0.000984786	0.036785540	-0.07733503

Table 1: Parameters used in simulations for model 1 and model 2. λ is the real part of the dominating eigenvalue for matrix A^* , representing the efficiency of the regulatory mechanism.

As we see in Table 1, λ for model 2 is smaller than that of model 1, suggesting that the regulatory mechanism in model 2 is more efficient. This is indeed the case since the FLLN scaled dynamics for model 2 converges to steady-state more rapidly in Figure 1. In Table 2, the limiting variances for

the derived clones are similar for both models. However, model 2 has greater variance for HSCs, meaning the regulatory mechanism in model 2 is less effective.




Figure 1: Simulations for the two-compartment cell dynamics for model 1 (top) and 2 (bottom). Initial conditions are $\bar{n}_0 = 11$, $\bar{n}_1 = \frac{\gamma \bar{n}_0}{\alpha - \gamma + \delta}$, $\mathbf{u} = \mathbf{0}$, and $U = 1.5 \cdot V^*$. For visual clarity, we display 10 simulations for the simulated FLLN and FCLT dynamics (first two columns). In total, 100 simulations are conducted and boxplots (third column) use all simulations.

Theoretical and Estimated Limiting Variance				
		$V_{1,1}^*$	$V_{2,2}^*$	$V_{1,2}^*$
Model 1	Theory	6.8311833	0.6951164	0.3056770
	Estimation	6.1895931	0.7504925	0.3406715
Model 2	Theory	12.3610797	0.6541369	0.1093108
	Estimation	12.9098659	0.7213062	0.0638457

Table 2: For each parameter specification, we compare the theoretical limiting covariance with estimated covariance from simulations. “Theory” refers to expression in Eq. (8) in Section 3.3. “Estimation” refers to average of the sample variance of the last 5 time points in the boxplots in Figure 1.

6 Discussions

In this paper, we develop a framework to analyze the efficiency and effectiveness of a given regulatory mechanism on the two-compartment cell proliferation model. We first scale up initial population sizes to study the large-scale approximation (FLLN and FCLT) of the dynamics. Then, we pass with time to infinity to study the long-time asymptotic behavior of FLLN and FCLT. We study the stability property of the FLLN and FCLT dynamics. More specifically, we show global stability for the non-trivial steady state and study the rate of convergence to the steady state. For the FCLT dynamics, we show exponential convergence in 2–Wasserstein metric to a Gauss-Markov process with zero mean function and a covariance function that will eventually stabilize. To interpret the results, we view the rate of convergence of FLLN to the non-trivial steady state as the efficiency of the regulatory mechanism. The FCLT limit at large times represents fluctuation around equilibrium and we associate the limiting variance of fluctuation with effectiveness of the regulatory mechanism. For a regulatory mechanism to be effective, the limiting variance of the FCLT should be small. We show by example that efficiency and effectiveness are not necessarily coupled. There are efficient regulatory mechanisms that are not effective, and vice versa.

In hematologic malignancies, non-canonical activation or dysregulation of Notch and Wnt signaling pathways are common (Zhou et al. (2022); Liu et al. (2022)). Specifically, Zhou et al. (2022) point out more than 50% of the T-ALL patients have NOTCH1 somatic activating mutations and Liu et al. (2022) state that deletion of GSK3B leaves hematopoietic stem cells in a precancerous state. Furthermore, both papers indicate activation of Notch or Wnt signaling endow cancer cell with therapeutic drug resistance for various cancer types. Therapies targeting Notch or Wnt signaling pathways for various cancer types are reviewed in Li et al. (2023) and Zhang and Wang (2020). One application of our model in the context of targeted therapy can be summarized as follows. Suppose dysregulation of signaling pathways occurred in the hematopoietic system and the system in this pathological state progresses until treatment (time 0 in our model). The regulatory mechanism in our model can be interpreted as target therapies aiming to restore or modify various signaling pathways. With the functional forms of $\alpha(\cdot)$ and $\gamma(\cdot)$ estimated from clinical data, one can study

the efficiency and effectiveness of the therapy according to our framework.

We conclude by mentioning possible extensions for our framework. In the present paper, we adopt the assumptions from Model 1 in Arino and Kimmel (1986) that there is a short-range feedback regulating differentiation and a long-range feedback regulating self-renewal. It is conceivable that our procedure of deriving large-scale approximations (FLLN and FCLT) will also work under assumptions from Model 2 (differentiation and self-renewal both regulated by long-range signals) and 3 (differentiation regulated by a long-range signal and self-renewal regulated by a short-range signal) from Arino and Kimmel (1986). Naturally, one needs to modify Eq. (2) in Section 2.1 to obtain stability for FLLN and FCLT dynamics. Notice that conditions for $\alpha(\cdot), \gamma(\cdot)$ in Section 2.1 are imposed to guarantee stability of FLLN and FCLT dynamics. If we discard Eq. (2), FLLN and FCLT can still be derived. However, FLLN might not admit a non-trivial steady state. In this case, the covariance function will not stabilize. Depending on the biological context, the instability of FLLN and FCLT dynamics might be of interest. Finally, extension to multi-compartment models is conceptually straightforward, but complication levels up as the number of compartment increases (chaos might appear in dimension ≥ 3).

7 Acknowledgements

Ren-Yi Wang was supported by the Department of Statistics at Rice University and by the NIH grant R01 268380 to Wenyi Wang. Marek Kimmel was supported in part by the NSF DMS RAPID COLLABORATION grant 2030577 to Marek Kimmel and Simon Tavaré, and by the National Science Center (Poland) grant number 2021/41/B/NZ2/04134.

A Appendix

A.1 FLLN

In this section, we establish the FLLN to the density dynamics $\{\bar{\mathbf{N}}^{(r)}\}$ (Theorem 1). We follow closely the martingale and continuous mapping approach in Pang et al. (2007). The challenge here is that \mathbf{F} in the integral representation in Lemma 1 is not a Lipschitz function. Hence, we cannot show continuity of the integral operator I on its entire domain. Instead, we show continuity at the subset of domain that is of interest in Lemma 1.

Proposition 1. *For each action $a \in \mathcal{A}$, as defined in Section 2, $M_a^{(r)}$ is a L^2 martingale with respect to the natural filtration $\{\mathcal{F}_t^{(r)} \mid t \geq 0\}$ defined by Eq. (1) in Section 2.1.*

Proof. Define $E_i^{(r)}(t) = \mathbb{E}(\bar{N}_0^{(r)}(t))$. Since $\alpha(y) \leq \alpha(0)$ for all $y \geq 0$, we have

$$\begin{aligned} E_0^{(r)}(t) &\leq \bar{n}_0^{(r)} + \int_0^t \alpha(0) E_0^{(r)}(s) ds \\ \implies E_0^{(r)}(t) &\leq \bar{n}_0^{(r)} \exp(\alpha(0)t), \text{ by Grönwall's inequality.} \end{aligned}$$

This implies

$$\begin{aligned} \mathbb{E} \int_0^t \alpha(\bar{N}_1^{(r)}(s)) \bar{N}_0^{(r)}(s) ds &\leq \int_0^t \alpha(0) E_0^{(r)}(s) ds \\ &\leq \int_0^t \alpha(0) \bar{n}_0^{(r)} \exp(\alpha(0)s) ds \\ &= \bar{n}_0^{(r)} \exp(\alpha(0)t) - \bar{n}_0^{(r)} < \infty; \\ \mathbb{E} \int_0^t \gamma(\bar{N}_0^{(r)}(s)) \bar{N}_0^{(r)}(s) ds &\leq \int_0^t \alpha(0) E_0^{(r)}(s) ds + \bar{n}_0^{(r)} \\ &\leq \bar{n}_0^{(r)} \exp(\alpha(0)t) < \infty. \end{aligned}$$

In addition,

$$E_0^{(r)}(t) + E_1^{(r)}(t) \leq \bar{n}_0^{(r)} + \bar{n}_1^{(r)} + \int_0^t \alpha(0) E_0^{(r)}(s) ds$$

$$\begin{aligned}
&\implies E_1^{(r)}(t) \leq \bar{n}_1^{(r)} + \bar{n}_0^{(r)} \exp(\alpha(0)t) \\
&\implies \mathbb{E} \int_0^t \delta \bar{N}_1^{(r)}(s) ds \leq \delta \bar{n}_1^{(r)} t + \delta \bar{n}_0^{(r)} \frac{\exp(\alpha(0)t) - 1}{\alpha(0)} < \infty.
\end{aligned}$$

Therefore, by Pang et al. (2007) Lemma 3.2, we deduce $M_{(0,b)}^{(r)}$, $M_{(0,di)}^{(r)}$, and $M_{(1,d)}^{(r)}$ are all L^2 martingales with predictable quadratic variations (PQV) being

$$\begin{aligned}
\langle M_{(0,b)}^{(r)} \rangle(t) &= \frac{1}{r} \int_0^t \alpha(\bar{N}_1^{(r)}(s)) \bar{N}_0^{(r)}(s) ds \\
\langle M_{(0,di)}^{(r)} \rangle(t) &= \frac{1}{r} \int_0^t \gamma(\bar{N}_0^{(r)}(s)) \bar{N}_0^{(r)}(s) ds \\
\langle M_{(1,d)}^{(r)} \rangle(t) &= \frac{1}{r} \int_0^t \delta \bar{N}_1^{(r)}(s) ds.
\end{aligned}$$

□

Proposition 2. *For all $a \in \mathcal{A}$, $\{\sqrt{r}M_a^{(r)}\}$ is stochastically bounded and $M_a^{(r)} \Rightarrow 0$ as $r \rightarrow \infty$.*

Proof. By Lemma 5.6 in Pang et al. (2007), it suffices to show for any fixed $T > 0$, there exists $K > 0$ such that

$$\sup_r \mathbb{E}((\sqrt{r}M_a^{(r)}(T))^2) = \sup_r \mathbb{E}(r \langle M_a^{(r)} \rangle(T)) \leq K.$$

In Proposition 1, we derived $\langle M_a^{(r)} \rangle$ for all $a \in \mathcal{A}$. Fix $T > 0$, since $\bar{n}_0^{(r)} \rightarrow \bar{n}_0$ as $r \rightarrow \infty$, there exists $K_{(0,b)} > 0$ such that for all r ,

$$\begin{aligned}
\mathbb{E}(r \langle M_{(0,b)}^{(r)} \rangle(T)) &= \mathbb{E} \int_0^T \alpha(\bar{N}_1^{(r)}(s)) \bar{N}_0^{(r)}(s) ds \\
&\leq \int_0^T \alpha(0) E_0^{(r)}(s) ds \\
&\leq \sup_r \{\bar{n}_0^{(r)} \exp(\alpha(0)T) - \bar{n}_0^{(r)}\} \leq K_{(0,b)}.
\end{aligned}$$

Analogously, there exist $K_{(0,di)} > 0$ and $K_{(1,d)} > 0$ such that for all r ,

$$\begin{aligned}\mathbb{E}(r\langle M_{(0,di)}^{(r)} \rangle(T)) &= \mathbb{E} \int_0^T \gamma(\bar{N}_0^{(r)}(s)) \bar{N}_0^{(r)}(s) ds \leq \sup_r \{\bar{n}_0^{(r)} \exp(\alpha(0)T)\} < K_{(0,di)} \\ \mathbb{E}(r\langle M_{(1,d)}^{(r)} \rangle(T)) &= \mathbb{E} \int_0^T \delta \bar{N}_1^{(r)}(s) ds \leq \sup_r \{\delta \bar{n}_1^{(r)} T + \delta \bar{n}_0^{(r)} \frac{\exp(\alpha(0)T) - 1}{\alpha(0)}\} < K_{(1,d)}.\end{aligned}$$

Therefore, by Lemma 5.9 of Pang et al. (2007), we have for all $a \in \mathcal{A}$, $M_a^{(r)} \Rightarrow 0$ in (\mathcal{D}, J_1) , Skorokhod space equipped with J_1 topology. \square

Lemma 1. Let $\mathbf{b} \in \mathbb{R}^2$ and $\mathbf{g} \in \mathcal{D}^2$ and define the operator $I : \mathbb{R}^2 \times \mathcal{D}^2 \rightarrow \mathcal{D}^2$ and $I(\mathbf{b}, \mathbf{g}) = \mathbf{x}$ such that

$$\mathbf{x}(t) = \mathbf{b} + \mathbf{g}(t) + \int_0^t \mathbf{F}(\mathbf{x}(s)) ds.$$

Then the operator I is continuous at $(\mathbf{b}, \mathbf{0})$ for $\mathbf{b} > 0$.

Proof. Endow \mathbb{R}^2 with the max norm $\|\cdot\|$ and \mathcal{D}^2 with SJ_1 (strong J_1) topology generated by truncated Skorokhod metrics $\{d_T \mid T > 0\}$. For topologies on \mathcal{D}^2 , we refer the readers to page 83 of Whitt (2002). We will show for all $\mathbf{b}^{(n)} \rightarrow \mathbf{b} > 0$ in $(\mathbb{R}^2, \|\cdot\|)$ and $\mathbf{g}^{(n)} \rightarrow \mathbf{0}$ in (\mathcal{D}^2, d_T) that we have $I(\mathbf{b}^{(n)}, \mathbf{g}^{(n)}) \rightarrow I(\mathbf{b}, \mathbf{0})$ in (\mathcal{D}^2, d_T) for all $T \geq 0$. Since $\mathbf{0}$ and $I(\mathbf{b}, \mathbf{0})$ are both continuous functions, convergence in d_T is equivalent to convergence in the truncated sup norm $\|\cdot\|_T$. Since $\mathbf{b} > 0$ and $\|\mathbf{g}^{(n)}\|_T \rightarrow 0$, we may without loss of generality take sequences such that $b_0^{(n)} > \|g_0^{(n)}\|_T$ and $b_1^{(n)} > \|g_1^{(n)}\|_T$. If $I(\mathbf{b}^{(n)}, \mathbf{g}^{(n)}) = (x_0^{(n)}, x_1^{(n)})^\top$, we have for all $t \in [0, T]$, $x_0^{(n)}(t) \geq 0$, $x_1^{(n)}(t) \geq 0$ and

$$\begin{aligned}x_0^{(n)}(t) &\leq b_0^{(n)} + \|g_0^{(n)}\|_T + \int_0^t \alpha(0)x_0^{(n)}(s) ds \\ \implies \|x_0^{(n)}\|_T &\leq (b_0^{(n)} + \|g_0^{(n)}\|_T) \exp(\alpha(0)T).\end{aligned}$$

Consequently,

$$\begin{aligned}
x_1^{(n)}(t) &\leq b_0^{(n)} + \|g_0^{(n)}\|_T + b_1^{(n)} + \|g_1^{(n)}\|_T + \int_0^t \alpha(0)x_0^{(n)}(s)ds \\
\implies \|x_1^{(n)}\|_T &\leq b_0^{(n)} + \|g_0^{(n)}\|_T + b_1^{(n)} + \|g_1^{(n)}\|_T + T\alpha(0)(b_0^{(n)} + \|g_0^{(n)}\|_T) \exp(\alpha(0)T).
\end{aligned}$$

Since both upper bounds converge as $n \rightarrow \infty$, there exists N such that for all $n \geq N$ and $t \in [0, T]$, $(x_0^{(n)}(t), x_1^{(n)}(t))^\top$ lies in a compact set K_T . Therefore, \mathbf{F} is Lipschitz with Lipschitz constant L_T on K_T .

Denote $I(\mathbf{b}, \mathbf{0}) = \mathbf{x}$. Then for all $t \in [0, T]$,

$$\begin{aligned}
\|\mathbf{x}^{(n)}(t) - \mathbf{x}(t)\| &\leq \|\mathbf{b}^{(n)} - \mathbf{b}\| + \|\mathbf{g}^{(n)}\|_T + \int_0^t L_T \|\mathbf{x}^{(n)}(s) - \mathbf{x}(s)\| ds \\
\implies \|\mathbf{x}^{(n)} - \mathbf{x}\|_T &\leq (\|\mathbf{b}^{(n)} - \mathbf{b}\| + \|\mathbf{g}^{(n)}\|_T) \exp(TL_T).
\end{aligned}$$

The last inequality follows from Grönwall's inequality. Fix $\epsilon > 0$, we select N such that for all $n \geq N$, we have $\|\mathbf{b}^{(n)} - \mathbf{b}\| + \|\mathbf{g}^{(n)}\|_T \leq \epsilon \exp(-TL_T)$. Continuity then follows. \square

Theorem 1. $\bar{\mathbf{N}}^{(r)} \rightarrow \bar{\mathbf{N}}$ in probability as $r \rightarrow \infty$, where $\bar{\mathbf{N}}' = \mathbf{F}(\bar{\mathbf{N}})$ and $\bar{\mathbf{N}}(0) = \bar{\mathbf{n}}$.

Proof. By Proposition 2, we have $(\bar{\mathbf{n}}^{(r)}, \mathbf{M}^{(r)}) \Rightarrow (\bar{\mathbf{n}}, \mathbf{0})$ with $\bar{\mathbf{n}} > 0$. By Lemma 1, we have

$$\bar{\mathbf{N}}^{(r)} = I(\bar{\mathbf{n}}^{(r)}, \mathbf{M}^{(r)}) \Rightarrow I(\bar{\mathbf{n}}, \mathbf{0}) \triangleq \bar{\mathbf{N}}.$$

Therefore,

$$\begin{aligned}
\bar{\mathbf{N}}(t) &= \bar{\mathbf{n}} + \int_0^t \mathbf{F}(\bar{\mathbf{N}}(s)) ds \\
\implies \bar{\mathbf{N}}'(t) &= \mathbf{F}(\bar{\mathbf{N}}(t)); \quad \bar{\mathbf{N}}(0) = \bar{\mathbf{n}}.
\end{aligned}$$

Since $\bar{\mathbf{N}}$ is deterministic, we strengthen the notion of convergence to $\bar{\mathbf{N}}^{(r)} \rightarrow \bar{\mathbf{N}}$ in probability. \square

A.2 FCLT

In this section, we establish weak convergence for the fluctuation dynamics $\hat{\mathbf{N}}^{(r)}$ as $r \rightarrow \infty$ by applying the continuous mapping theorem. To show the FCLT, we take a different integral operator J , and show convergence for the argument of the integral operator (Theorem 2). The convergence of the argument of the integral operator relies on stochastic boundedness of $\{\hat{\mathbf{N}}^{(r)}\}$, which is shown in Lemma 2.

Lemma 2. $\{\hat{\mathbf{N}}^{(r)}\}$ is stochastically bounded. That is, the set of random variables $\{|\hat{\mathbf{N}}^{(r)}|_T\}$ is tight for all $T \geq 0$.

Proof. Recall

$$\hat{\mathbf{N}}^{(r)} = \hat{\mathbf{n}}^{(r)} + \hat{\mathbf{M}}^{(r)} + \sqrt{r} \int_0^t [\mathbf{F}(\bar{\mathbf{N}}^{(r)}(s)) - \mathbf{F}(\bar{\mathbf{N}}(s))] ds.$$

By Skorohod representation theorem, we may assume $\hat{\mathbf{n}}^{(r)} \rightarrow \hat{\mathbf{n}}$ almost surely and show tightness under this assumption since the law of each $\hat{\mathbf{N}}^{(r)}$ remains intact. Define

$$\mathbf{Y}^{(r)}(\mathbf{t}) = \begin{pmatrix} \sqrt{r}(\frac{1}{r}P_{(0,b)}(rt_1) - t_1) \\ \sqrt{r}(\frac{1}{r}P_{(0,di)}(rt_2) - t_2) \\ \sqrt{r}(\frac{1}{r}P_{(1,d)}(rt_3) - t_3) \end{pmatrix}; \quad \mathbf{B}(\mathbf{t}) = \begin{pmatrix} B_{(0,b)}(t_1) \\ B_{(0,di)}(t_2) \\ B_{(1,d)}(t_3) \end{pmatrix}.$$

By the FCLT for Poisson process (Theorem 4.2 in Pang et al. (2007)), we have $\mathbf{Y}^{(r)} \Rightarrow \mathbf{B}$, a vector of three independent standard Brownian motions. We invoke Skorohod representation theorem again and assume $\mathbf{Y}^{(r)} \rightarrow \mathbf{B}$ almost surely.

Since $\bar{\mathbf{N}}^{(r)} \rightarrow \bar{\mathbf{N}}$ in probability, $\mathbf{I}^{(r)} \rightarrow \mathbf{I}$ in probability, where we now denote

$$\mathbf{I}^{(r)}(t) = \begin{pmatrix} \int_0^t \alpha(\bar{N}_1^{(r)}(s)) ds \\ \int_0^t \gamma(\bar{N}_0^{(r)}(s)) ds \\ \int_0^t \delta \bar{N}_1^{(r)}(s) ds \end{pmatrix}; \quad \mathbf{I}(t) = \begin{pmatrix} \int_0^t \alpha(\bar{N}_1(s)) ds \\ \int_0^t \gamma(\bar{N}_0(s)) ds \\ \int_0^t \delta \bar{N}_1(s) ds \end{pmatrix}.$$

Consequently, we have joint convergence in probability,

$$(\mathbf{Y}^{(r)}, \mathbf{I}^{(r)})^\top \rightarrow (\mathbf{B}, \mathbf{I})^\top.$$

Since a standard Brownian motion is locally $\frac{1}{4}$ Hölder, we have

$$\begin{aligned} \|\mathbf{Y}^{(r)} \circ \mathbf{I}^{(r)} - \mathbf{B} \circ \mathbf{I}\|_T &\leq \|\mathbf{Y}^{(r)} \circ \mathbf{I}^{(r)} - \mathbf{B} \circ \mathbf{I}^{(r)}\|_T + \|\mathbf{B} \circ \mathbf{I}^{(r)} - \mathbf{B} \circ \mathbf{I}\|_T \\ &\leq \|\mathbf{Y}^{(r)} \circ \mathbf{I}^{(r)} - \mathbf{B} \circ \mathbf{I}^{(r)}\|_T + C\|\mathbf{I}^{(r)} - \mathbf{I}\|_T^{\frac{1}{4}}. \end{aligned}$$

Since $\|\mathbf{I}^{(r)} - \mathbf{I}\|_T \rightarrow 0$ in probability as $r \rightarrow \infty$, we have $\|\mathbf{I}^{(r)} - \mathbf{I}\|_T^{\frac{1}{4}} \rightarrow 0$ in probability and $\|\mathbf{I}^{(r)}\|_T \rightarrow \|\mathbf{I}\|_T$ in probability. Hence, for any fixed $\epsilon > 0$, we have

$$\begin{aligned} \limsup_{r \rightarrow \infty} \mathbb{P}(\|\mathbf{Y}^{(r)} \circ \mathbf{I}^{(r)} - \mathbf{B} \circ \mathbf{I}^{(r)}\|_T \geq \epsilon) &\leq \limsup_{r \rightarrow \infty} [\mathbb{P}(\|\mathbf{Y}^{(r)} - \mathbf{B}\|_K \geq \epsilon) + \mathbb{P}(\|\mathbf{I}^{(r)}\|_T > K)] \\ &= \mathbb{P}(\|\mathbf{I}\|_T > K). \end{aligned}$$

Since the inequality holds for all K , the limit is zero. Therefore, we conclude $\mathbf{Y}^{(r)} \circ \mathbf{I}^{(r)} \rightarrow \mathbf{B} \circ \mathbf{I}$ in probability. As a consequence, $\hat{\mathbf{M}}^{(r)} \rightarrow \hat{\mathbf{M}}$ in probability, where

$$\begin{aligned} \hat{M}_0(t) &= B_{(0,b)} \left(\int_0^t \alpha(\bar{N}_1(s)) \bar{N}_0(s) ds \right) - B_{(0,di)} \left(\int_0^t \gamma(\bar{N}_0(s)) \bar{N}_0(s) ds \right) \\ \hat{M}_1(t) &= B_{(0,di)} \left(\int_0^t \gamma(\bar{N}_0(s)) \bar{N}_0(s) ds \right) - B_{(1,d)} \left(\int_0^t \delta \bar{N}_1(s) ds \right). \end{aligned}$$

Using multivariate Taylor's theorem, there are error terms $H^{(r)} \rightarrow 0_{2 \times 2}$ in probability and

$$\hat{\mathbf{N}}^{(r)}(t) = \hat{\mathbf{n}}^{(r)} + \hat{\mathbf{M}}^{(r)}(t) + \int_0^t [\nabla \mathbf{F}(\bar{\mathbf{N}}(s)) + H^{(r)}(s)] \hat{\mathbf{N}}^{(r)}(s) ds.$$

Since $\{\hat{\mathbf{n}}^{(r)}\}$, $\{\hat{\mathbf{M}}^{(r)}\}$, and $\{\nabla \mathbf{F}(\bar{\mathbf{N}}(s)) + H^{(r)}(s)\}$ are stochastically bounded, for any fixed $1 - \epsilon$ and T , there exist K, M such that for all $t \in [0, T]$ and $r \in \mathbb{N}$,

$$\mathbb{P}(\|\hat{\mathbf{N}}^{(r)}(t)\| \leq K + \int_0^t M \|\hat{\mathbf{N}}^{(r)}(s)\| ds) \geq 1 - \epsilon.$$

By Grönwall's inequality, we have

$$\sup_r \mathbb{P}(\|\hat{\mathbf{N}}^{(r)}\|_T \leq K \exp(MT)) \geq 1 - \epsilon.$$

This concludes stochastic boundedness of $\{\hat{\mathbf{N}}^{(r)}\}$. \square

Theorem 2. $\hat{\mathbf{N}}^{(r)} \Rightarrow \hat{\mathbf{N}}$, where

$$\hat{\mathbf{N}}(t) = \hat{\mathbf{n}} + \hat{\mathbf{M}}(t) + \int_0^t \nabla \mathbf{F}(\bar{\mathbf{N}}(s)) \hat{\mathbf{N}}(s) ds.$$

Proof. Define an operator $J : \mathcal{D}^2 \rightarrow \mathcal{D}^2$ such that $J(\mathbf{b}) = \mathbf{x}$ with

$$\mathbf{x}(t) = \mathbf{b}(t) + \int_0^t \nabla \mathbf{F}(\bar{\mathbf{N}}(s)) \mathbf{x}(s) ds.$$

It is easy to check the operation J is continuous in the SJ_1 topology.

Arguing by Grönwall's inequality, we have $\mathbf{b}_n \rightarrow \mathbf{b} \in \mathcal{C}^2$ implies $J(\mathbf{b}_n) \rightarrow J(\mathbf{b})$. We write

$$\hat{\mathbf{N}}^{(r)} = J(\hat{\mathbf{n}} + \hat{\mathbf{M}}^{(r)} + \int_0^\cdot H^{(r)}(s) \hat{\mathbf{N}}^{(r)}(s) ds).$$

Without invoking Skorohod representation theorem as in Theorem 3, we only have joint convergence in distribution by Slutsky's theorem

$$(\mathbf{Y}^{(r)}, \mathbf{I}^{(r)}) \Rightarrow (\mathbf{B}, \mathbf{I}).$$

This is sufficient for us to conclude $\mathbf{Y}^{(r)} \circ \mathbf{I}^{(r)} \Rightarrow \mathbf{B} \circ \mathbf{I}$. Hence, $\hat{\mathbf{M}}^{(r)} \Rightarrow \hat{\mathbf{M}}$. By Lemma 2, $\{\hat{\mathbf{N}}^{(r)}\}$ is stochastically bounded. Combining with the statement $H^{(r)} \rightarrow 0_{2 \times 2}$ in probability, we have $\int_0^\cdot H^{(r)}(s) \hat{\mathbf{N}}^{(r)}(s) ds \Rightarrow \mathbf{0}$. Hence,

$$\hat{\mathbf{n}} + \hat{\mathbf{M}}^{(r)} + \int_0^\cdot H^{(r)}(s) \hat{\mathbf{N}}^{(r)}(s) ds \Rightarrow \hat{\mathbf{n}} + \hat{\mathbf{M}}.$$

Since $\hat{\mathbf{n}} + \hat{\mathbf{M}}$ has continuous sample paths, $\hat{\mathbf{N}}^{(r)} \Rightarrow \hat{\mathbf{N}}$ by continuity of J . □

A.3 Stability of the FLLN Dynamics

Since the FLLN dynamics is a two-dimensional autonomous system of ODEs, global stability can be established by Poincaré–Bendixson theorem. We first show the forward orbit through the initial data $\bar{\mathbf{n}}$ lies in a compact set (Proposition 3) and rule out undesired cases (Theorem 3).

Proposition 3. *For any fixed initial condition $\bar{\mathbf{n}}$, the trajectory of the FLLN limit $\{\bar{\mathbf{N}}(t) \mid t \geq 0\}$ lies in a compact subset of \mathbb{R}^2 .*

Proof. The FLLN limit is defined by $\bar{\mathbf{N}}(0) = \bar{\mathbf{n}}$ and

$$\begin{aligned}\bar{N}'_0(t) &= [\alpha(\bar{N}_1(t)) - \gamma(\bar{N}_0(t))]\bar{N}_0(t) \\ \bar{N}'_1(t) &= -\delta\bar{N}_1(t) + \gamma(\bar{N}_0(t))\bar{N}_0(t).\end{aligned}$$

To show the trajectory lies in a compact set, it suffices to show $\bar{N}_0(t)$ is uniformly bounded for all $t \geq 0$.

We consider two cases in Eq. (2) from Section 2.1. For the first case, we assume $\alpha(0) < \lim_{x \rightarrow \infty} \gamma(x)$ and define an IVP with $\tilde{\mathbf{N}}(0) = \bar{\mathbf{n}}$ and

$$\begin{aligned}\tilde{N}'_0(t) &= [\alpha(0) - \gamma(\tilde{N}_0(t))]\tilde{N}_0(t) \\ \tilde{N}'_1(t) &= -\delta\tilde{N}_1(t) + \gamma(\tilde{N}_0(t))\tilde{N}_0(t).\end{aligned}$$

This IVP corresponds to deletion of the regulation on α . Since the first equation is autonomous, $\tilde{N}_0(t)$ is monotonic. Furthermore, since $\alpha(0) < \lim_{x \rightarrow \infty} \gamma(x)$, $\tilde{N}_0(t)$ is bounded. Since $\alpha(y) \leq \alpha(0)$, $\tilde{N}_0(t) \geq \bar{N}_0(t)$ for all $t \geq 0$, which implies $\bar{N}_0(t)$ is also bounded. As a consequence, $\bar{N}_1(t)$ is bounded for all t .

For the second case, $\lim_{y \rightarrow \infty} \alpha(y) < \gamma(0)$, we set up another IVP with initial condition $\tilde{\mathbf{N}}(0) = \bar{\mathbf{n}}$

and

$$\begin{aligned}\tilde{N}'_0(t) &= [\alpha(\tilde{N}_1(t)) - \gamma(0)]\tilde{N}_0(t) \\ \tilde{N}'_1(t) &= -\delta\tilde{N}_1(t) + \gamma(0)\tilde{N}_0(t).\end{aligned}$$

This IVP corresponds to deletion of regulation on γ . Since $\gamma(x) \geq \gamma(0)$ for all $x \geq 0$, we have $\tilde{N}_0(t) \geq \bar{N}_0(t)$ for all $t \geq 0$. We show there is a compact trapping region \mathcal{K} that contains $\bar{\mathbf{n}}$ by Theorem 4.2.3 in Schaeffer and Cain (2018). To see this, let y^* denote the unique solution to $\alpha(y) = \gamma(0)$, then for $y \geq y^*$, we have $\alpha(y) - \gamma(0) \leq 0$. Therefore, for $y \geq y^*$,

$$\begin{pmatrix} (\alpha(y) - \gamma(0))x \\ -\delta y + \gamma(0)x \end{pmatrix} \cdot \begin{pmatrix} -1 \\ \frac{\alpha(y) - \gamma(0)}{\gamma(0)} \end{pmatrix} \geq 0.$$

The curve with normal vector $(-1, \frac{\alpha(y) - \gamma(0)}{\gamma(0)})^\top$ has the form

$$x(y) = K + \int_{y^*}^y \frac{\alpha(t) - \gamma(0)}{\gamma(0)} dt.$$

Here K is a positive constant that will be specified later. To complete the boundary for the trapping region with $y < y^*$, suppose the normal vector is of the form $(-1, C)^\top$ for some positive C , then we must have for all $y < y^*$,

$$\begin{pmatrix} (\alpha(y) - \gamma(0))x \\ -\delta y + \gamma(0)x \end{pmatrix} \cdot \begin{pmatrix} -1 \\ C \end{pmatrix} = [(1 + C)\gamma(0) - \alpha(y)]x - C\delta y \geq 0.$$

The curve with normal vector $(-1, C)^\top$ has the form $x(y) = K - C(y^* - y)$, so the desired condition is equivalent to the following: for all $y < y^*$,

$$[(1 + C)\gamma(0) - \alpha(y)][K - C(y^* - y)] - C\delta y \geq 0.$$

We may select C^* large enough such that $(1 + C^*)\gamma(0) > \alpha(0) \geq \alpha(y)$, then select K^* large enough such that $[(1 + C^*)\gamma(0) - \alpha(y)][K^* - C^*(y^* - y)] - C^*\delta y \geq 0$ and $\bar{\mathbf{n}} \in \mathcal{K}$, where

$$\begin{aligned} \mathcal{K} = & \left\{ (x, y) \mid y \geq y^*; 0 \leq x \leq K^* + \int_{y^*}^y \frac{\alpha(t) - \gamma(0)}{\gamma(0)} dt \right\} \\ & \cup \{(x, y) \mid 0 \leq y < y^*; 0 \leq x \leq K^* - C^*(y^* - y)\}. \end{aligned}$$

The set \mathcal{K} is a compact trapping region with piecewise \mathcal{C}^1 boundaries. Hence, $\tilde{N}_0(t)$ is bounded, which implies that both $\bar{N}_0(t)$ and $\bar{N}_1(t)$ are bounded. \square

Theorem 3. $\bar{\mathbf{N}}(t) \rightarrow \bar{\mathbf{N}}^*$ as $t \rightarrow \infty$ with any fixed initial condition $\bar{\mathbf{n}}$. Moreover, for any fixed $\eta \in (\lambda, 0)$ and initial data $\bar{\mathbf{n}} > 0$, $\|\bar{\mathbf{N}}(t) - \bar{\mathbf{N}}^*\| = o(\exp(\eta t))$.

Proof. By Proposition 3, the trajectory of FLLN lies in a compact set \mathcal{K} . To rule out periodic orbits, we define $\mathcal{U} = \{(x, y) \mid x > 0, y > 0\}$. Then by Dulac's theorem (Theorem 7.2.5 in Schaeffer and Cain (2018)), we have

$$\nabla \cdot \left[\frac{-1}{x} \mathbf{F}(x, y) \right] = \gamma'(x) + \frac{\delta}{x} > 0 \text{ on } \mathcal{U}.$$

Therefore, there are no periodic orbits in \mathcal{U} . In addition, the forward orbit through $\bar{\mathbf{n}}$ lies in a compact subset of \mathcal{U} and \mathcal{U} contains exactly one point. By Eq. (4) in Section 3.1, $\bar{\mathbf{N}}^*$ is locally stable. We invoke Strong Poincaré–Bendixson Theorem (Theorem 7.2.5 in Schaeffer and Cain (2018)) to conclude $\bar{\mathbf{N}}^*$ is globally stable. Consequently, for any fixed initial data $\bar{\mathbf{n}} > 0$ and $\eta \in (\lambda, 0)$, we have

$$\|\bar{\mathbf{N}}(t) - \bar{\mathbf{N}}^*\| = O(\exp(\eta t)) \text{ as } t \rightarrow \infty.$$

Since $(\lambda, 0)$ is open, we can strengthen the statement. For any fixed $\bar{\mathbf{n}}$ and $\eta \in (\lambda, 0)$,

$$\|\bar{\mathbf{N}}(t) - \bar{\mathbf{N}}^*\| = o(\exp(\eta t)) \text{ as } t \rightarrow \infty.$$

\square

A.4 Large-time Asymptotics of the FCLT Dynamics

To study large-time behaviors of the FCLT dynamics $\hat{\mathbf{N}}$, we fix $M > 0$ and focus on the dynamics in time interval $[T, T + M]$. Recall we define $\mathbf{G}_T(t) = \hat{\mathbf{N}}(T + t)$ and the Gaussian process \mathbf{G}^* with mean function $\mathbf{m}^*(t) = \mathbf{0}$ and autocovariance function $K^*(s, t) = V^* \exp((t - s)[A^*]^\top)$ for $s \leq t$. As we increase T to infinity, the FCLT dynamics $\hat{\mathbf{N}}$ in the moving interval $[T, T + M]$ will behave like the Gaussian process \mathbf{G}^* . Also recall that we cast the processes to the space of Gaussian measures on $\mathcal{H} = \mathcal{L}_2([0, M]^2, \mathcal{B}([0, M]^2), Leb)$, as \mathcal{G}_T 's and \mathcal{G}_* . We consider the 2-Wasserstein distance of these two Gaussian measures in the space \mathcal{H} . We prove the following result.

Theorem 4. $W(\mathcal{G}_T, \mathcal{G}_*) = o(\exp(\frac{\eta}{2}T))$, for all $\eta \in (\lambda, 0)$.

Proof. Let $\|\cdot\|_{Tr}$ denote the trace norm and $\|\cdot\|_{HS}$ denote the Hilbert-Schmidt norm.

$$\begin{aligned} W_2^2(\mathcal{G}_T, \mathcal{G}_*) &= \|\mathbf{m}_T\|_2^2 + Tr(C_T + C_* - 2(C_*^{1/2}C_T C_*^{1/2})^{1/2}) \\ &\leq \|\mathbf{m}_T\|_2^2 + \|C_T + C_* - 2(C_*^{1/2}C_T C_*^{1/2})^{1/2}\|_{Tr}. \end{aligned}$$

Since all covariance operators are positive, by operator monotonicity of the square root function (page 2 of Phillips (1987)),

$$\|C_T + C_* - 2(C_*^{1/2}C_T C_*^{1/2})^{1/2}\|_{Tr} \leq \|(C_T + C_*)^2 - 4(C_*^{1/2}C_T C_*^{1/2})\|_{Tr}^{\frac{1}{2}}.$$

Since C_* is positive, self-adjoint, and of trace class, $C_*^{\frac{1}{2}}$ is positive, self-adjoint, and Hilbert-Schmidt. This implies $(C_T + C_*)^2 - 4(C_*^{1/2}C_T C_*^{1/2})$ is a positive operator. To see this, let (e_i) be an orthonormal eigenbasis for $C_*^{\frac{1}{2}}$ corresponding to eigenvalues (λ_i) . Then, we have

$$\begin{aligned} \langle C_*^{1/2}C_T C_*^{1/2} e_i, e_i \rangle &= \langle C_T C_*^{1/2} e_i, C_*^{1/2} e_i \rangle \\ &= \lambda_i^2 \langle C_T e_i, e_i \rangle \\ &= \langle C_* C_T e_i, e_i \rangle = \langle C_T C_* e_i, e_i \rangle. \end{aligned}$$

Therefore,

$$\langle (C_T + C_*)^2 - 4(C_*^{1/2}C_TC_*^{1/2})e_i, e_i \rangle = \langle (C_T - C_*)^2 e_i, e_i \rangle \geq 0.$$

Since the trace norm of a positive operator is equal to its trace, we have

$$\begin{aligned} \|(C_T + C_*)^2 - 4(C_*^{1/2}C_TC_*^{1/2})\|_{Tr}^{\frac{1}{2}} &= Tr((C_T + C_*)^2 - 4(C_*^{1/2}C_TC_*^{1/2}))^{\frac{1}{2}} \\ &= Tr((C_T - C_*)^2)^{\frac{1}{2}} \\ &= \|C_T - C_*\|_{HS}. \end{aligned}$$

By Proposition 7.10.23 in Bogachev and Smolyanov (2020), we have

$$\begin{aligned} \|C_T - C_*\|_{HS}^2 &= \|K_T - K_*\|_2^2 \\ &= 2 \int_0^M \int_s^M \|V(s+T) \exp(\int_s^t A^\top(u+T)du) - V^* \exp(A^*(t-s))^\top\|^2 dt ds \\ &\leq 2 \int_0^M \int_s^M (\|V(s+T) \exp(\int_s^t A^\top(u+T)du) - V^* \exp(\int_s^t A^\top(u+T)du)\| \\ &\quad + \|V^* \exp(\int_s^t A^\top(u+T)du) - V^* \exp(A^*(t-s))^\top\|)^2 dt ds, \end{aligned}$$

where K_T and K_* are defined by Eq. (11) in Section 4.2. To show $\|C_T - C_*\|_{HS}^2 = o(\exp(2\eta T))$, it suffices to show both

$$\begin{aligned} \int_0^M \int_s^M \|[V(s+T) - V_*] \exp(\int_s^t A^\top(u+T)du)\|^2 dt ds &= o(\exp(2\eta T)); \\ \int_0^M \int_s^M \|V^*[\exp(\int_s^t A^\top(u+T)du) - \exp(A^*(t-s))^\top]\|^2 dt ds &= o(\exp(2\eta T)). \end{aligned}$$

For the first term, by Eq. 10 in Section 4.1, there exist T large enough and a constant C such that

$$\int_0^M \int_s^M \|[V(s+T) - V_*] \exp(\int_s^t A^\top(u+T)du)\|^2 dt ds$$

$$\begin{aligned}
&\leq \int_0^M C \exp(2\eta(s+T)) \int_s^M \|\exp(\int_s^t A^\top(u+T)du)\|^2 dt ds \\
&= C \exp(2\eta T) \int_0^M \exp(2\eta s) \int_s^M \|\exp(\int_s^t A^\top(u+T)du)\|^2 dt ds.
\end{aligned}$$

Since $A(\cdot + T) \rightarrow A^*$ as $T \rightarrow \infty$ (see Eq. 9), the integral term is bounded as $T \rightarrow \infty$. Hence,

$$\begin{aligned}
&\int_0^M \int_s^M \|[V(s+T) - V^*] \exp(\int_s^t A^\top(u+T)du)\|^2 dt ds = O(\exp(2\eta T)), \forall \eta \in (\lambda, 0) \\
\implies &\int_0^M \int_s^M \|[V(s+T) - V^*] \exp(\int_s^t A^\top(u+T)du)\|^2 dt ds = o(\exp(2\eta T)), \forall \eta \in (\lambda, 0).
\end{aligned}$$

For the second term, since $A(\cdot + T) \rightarrow A^*$ as $T \rightarrow \infty$ (see Eq. 9), by the mean value inequality in Banach space (Theorem 44 in Guirao et al. (2022)), there exists a constant $D > 0$ (independent of T) such that

$$\begin{aligned}
&\int_0^M \int_s^M \|V^* [\exp(\int_s^t A^\top(u+T)du) - \exp(A^*(t-s)^\top)]\|^2 dt ds \\
&\leq \|V^*\|^2 \int_0^M \int_s^M D \|\int_s^t A^\top(u+T)du - [A^*]^\top(t-s)\|^2 dt ds \\
&= \|V^*\|^2 \int_0^M \int_s^M D \|\int_s^t A^\top(u+T) - [A^*]^\top du\|^2 dt ds \\
&\leq \tilde{D} \cdot \int_0^M \int_s^M \exp(2\eta T) (\exp(\eta s) - \exp(\eta t)) dt ds, \text{ for some } \tilde{D} > 0.
\end{aligned}$$

Therefore,

$$\begin{aligned}
&\int_0^M \int_s^M \|V^* [\exp(\int_s^t A^\top(u+T)du) - \exp(A^*(t-s)^\top)]\|^2 dt ds = O(\exp(2\eta T)), \forall \eta \in (\lambda, 0) \\
\implies &\int_0^M \int_s^M \|V^* [\exp(\int_s^t A^\top(u+T)du) - \exp(A^*(t-s)^\top)]\|^2 dt ds = o(\exp(2\eta T)), \forall \eta \in (\lambda, 0).
\end{aligned}$$

Since $\|\mathbf{m}\|_T = O(\exp(2\lambda T))$, we conclude

$$\begin{aligned}
W_2^2(\mathcal{G}_T, \mathcal{G}_*) &\leq \|\mathbf{m}_T\|_2^2 + \|C_T - C_*\|_{HS} \\
&= o(\exp(\eta T)).
\end{aligned}$$

Therefore, $W_2(\mathcal{G}_T, \mathcal{G}) = o(\exp(\frac{\eta}{2}T))$ for all $\eta \in (\lambda, 0)$ and this result holds for all M . \square

References

- Andrew W Duncan, Frédérique M Rattis, Leah N DiMascio, Kendra L Congdon, Gregory Pazianos, Chen Zhao, Keejung Yoon, J Michael Cook, Karl Willert, Nicholas Gaiano, et al. Integration of Notch and Wnt signaling in hematopoietic stem cell maintenance. *Nature Immunology*, 6(3): 314–322, 2005.
- Ulrika Blank, Göran Karlsson, and Stefan Karlsson. Signaling pathways governing stem-cell fate. *Blood, The Journal of the American Society of Hematology*, 111(2):492–503, 2008.
- Raphael Kopan. Notch signaling. *Cold Spring Harbor Perspectives in Biology*, 4(10):a011213, 2012.
- Anna Bigas and Lluís Espinosa. Hematopoietic stem cells: to be or Notch to be. *Blood, The Journal of the American Society of Hematology*, 119(14):3226–3235, 2012.
- Roel Nusse, C Fuerer, W Ching, K Harnish, C Logan, A Zeng, D Ten Berge, and Y Kalani. Wnt signaling and stem cell control. In *Cold Spring Harbor Symposia on Quantitative Biology*, volume 73, pages 59–66. Cold Spring Harbor Laboratory Press, 2008.
- Tina Buechling and Michael Boutros. Wnt signaling: signaling at and above the receptor level. *Current Topics in Developmental Biology*, 97:21–53, 2011.
- Philipp Getto, Anna Marciniak-Czochra, Yukihiro Nakata, et al. Global dynamics of two-compartment models for cell production systems with regulatory mechanisms. *Mathematical Biosciences*, 245(2):258–268, 2013.
- Anna Marciniak-Czochra, Thomas Stiehl, Anthony D Ho, Willi Jäger, and Wolfgang Wagner. Modeling of asymmetric cell division in hematopoietic stem cells—regulation of self-renewal is essential for efficient repopulation. *Stem Cells and Development*, 18(3):377–386, 2009.

- Franziska Knauer, Thomas Stiehl, and Anna Marciniak-Czochra. Oscillations in a white blood cell production model with multiple differentiation stages. *Journal of Mathematical Biology*, 80: 575–600, 2020.
- Céline Bonnet, Panhong Gou, Simon Girel, Vincent Bansaye, Catherine Lacout, Karine Bailly, Marie-Hélène Schlagetter, Evelyne Lauret, Sylvie Méléard, and Stéphane Giraudier. Combined biological and modeling approach of hematopoiesis: from native to stressed erythropoiesis. *Available at SSRN 3777468*, 2021.
- Céline Bonnet and Sylvie Méléard. Large fluctuations in multi-scale modeling for rest hematopoiesis. *Journal of Mathematical Biology*, 82(6):58, 2021.
- Sandra N Catlin, Lambert Busque, Rosemary E Gale, Peter Gutter, and Janis L Abkowitz. The replication rate of human hematopoietic stem cells in vivo. *Blood, The Journal of the American Society of Hematology*, 117(17):4460–4466, 2011.
- Jos Domen, Samuel H Cheshier, and Irving L Weissman. The role of apoptosis in the regulation of hematopoietic stem cells: Overexpression of Bcl-2 increases both their number and repopulation potential. *The Journal of Experimental Medicine*, 191(2):253–264, 2000.
- Guodong Pang, Rishi Talreja, and Ward Whitt. Martingale proofs of many-server heavy-traffic limits for Markovian queues. *Probability Surveys*, 4:193–267, 2007.
- Ioannis Karatzas and Steven Shreve. *Brownian motion and stochastic calculus*. Springer Science & Business Media, 2012.
- Sigrun Bodine, Donald A Lutz, et al. *Asymptotic integration of differential and difference equations*. Springer, 2015.
- Ha Quang Minh. Entropic regularization of Wasserstein distance between infinite-dimensional Gaussian measures and Gaussian processes. *Journal of Theoretical Probability*, 36(1):201–296, 2023.

- Binghan Zhou, Wanling Lin, Yaling Long, Yunkai Yang, Huan Zhang, Kongming Wu, and Qian Chu. Notch signaling pathway: Architecture, disease, and therapeutics. *Signal Transduction and Targeted therapy*, 7(1):95, 2022.
- Jiaqi Liu, Qing Xiao, Jiani Xiao, Chenxi Niu, Yuanyuan Li, Xiaojun Zhang, Zhengwei Zhou, Guang Shu, and Gang Yin. Wnt/ β -catenin signalling: function, biological mechanisms, and therapeutic opportunities. *Signal Transduction and Targeted Therapy*, 7(1):3, 2022.
- Xinxin Li, Xianchun Yan, Yufeng Wang, Balveen Kaur, Hua Han, and Jianhua Yu. The Notch signaling pathway: a potential target for cancer immunotherapy. *Journal of Hematology & Oncology*, 16(1):45, 2023.
- Ya Zhang and Xin Wang. Targeting the Wnt/ β -catenin signaling pathway in cancer. *Journal of Hematology & Oncology*, 13:1–16, 2020.
- Ovide Arino and Marek Kimmel. Stability analysis of models of cell production systems. *Mathematical Modelling*, 7(9-12):1269–1300, 1986.
- Ward Whitt. Stochastic-process limits: An introduction to stochastic-process limits and their application to queues. 2002.
- Schaeffer and Cain. *Ordinary differential equations: Basics and beyond*. Springer, 2018.
- John Phillips. On the uniform continuity of operator functions and generalized Powers-Stormer inequalities. Technical report, 1987.
- Vladimir I Bogachev and Oleg G Smolyanov. *Real and functional analysis*. Springer, 2020.
- Antonio José Guirao, Vicente Montesinos, and Václav Zizler. *Renormings in Banach Spaces: A Toolbox*. Springer Nature, 2022.