

Monadic aspects of the ideal lattice functor on the category of distributive lattices

Ando Razafindrakoto^{1*}

¹Department of Mathematics and Applied Mathematics, University of the Western Cape, Robert Sobukwe Rd, Bellville, 7535, Cape Town, South Africa.

Corresponding author(s). E-mail(s): arazafindrakoto@uwc.ac.za;

Abstract

It is known that the construction of the frame of ideals from a distributive lattice induces a monad whose algebras are precisely the frames and frame homomorphisms. Using the Fakir construction of an idempotent approximation of a monad, we extend B. Jacobs' results on lax idempotent monads and show that the sequence of monads and comonads generated by successive iterations of this ideal functor on its algebras and coalgebras do not strictly lead to a new category. We further extend this result and provide a new proof of the equivalence between distributive lattices and coherent frames by showing that when the first inductive step in the Fakir construction is the identity monad, then the ambient category is equivalent to the free algebras.

Keywords: Monad, algebras, distributive lattices, frames, ideal functor, continuous lattices, Stone duality, split equalizer

MSC Classification: 06A11 , 06B35 , 06D22 , 18C15 , 18C20 , 18F60 , 18F70 , 54A99 , 54B30

1 Introduction

The importance and pervasiveness of the ideal lattice functor, or simply the ideal functor, in the study and history of Pointfree Topology are well-known, and its presence can be appreciated from the fact that it is to distributive lattices what the downset functor is to meet-semilattices ([1, Remark 6.6]). In particular, it is a central tool in certain fundamental results: on one hand it provides a richer structure in

the construction of compactifications of frames ([2–4]) and completions of uniform frames ([5, 6]), and on the other it plays a pivotal role in the Stone representation of distributive lattices ([7]) as well as in the theory of dualities¹ as exemplified by the dual adjunction that exists between distributive lattices and topological spaces via a schizophrenic object ([7–9]). In general, functorial constructions from the ideal lattice produce either a monad or a coreflector, i.e. an idempotent comonad, in different parts of the existing literature. For instance, while P. Johnstone and H. Simmons consider the monad generated by the dual adjunction between spaces and lattices in [7] and [9] respectively, B. Banaschewski describes various reflective and coreflective subcategories with the help of the ideal lattice in [2]. These seemingly contradictory descriptions are not only a consequence of the change in the domain and codomain categories, but more precisely from a much deeper reality: they arise as successive iterations of the functor. This qualitative transformation between opposites can be understood as a dialectical process as explained by W. Lawvere in the examples provided in the paper [10]. Here, the cylindrical model ([10, Section I]) of two sections united by a retraction (Cf. Lemma 2) qualitatively evolves into its opposite after another (quantitative) iteration of the functor. We therefore concentrate our investigation on the monadic aspects of the ideal functor in relation to this iteration. Generally, a monad $\mathbb{T} = (T, m, e)$ on a category \mathbf{C} induces an alternating chain of monads and comonads ([11]) on the various subcategories of algebras and coalgebras:

$$\dots \mathbf{Alg}(\mathbb{T}_n) \rightarrow \mathbf{Coalg}(\mathbb{K}_n) \rightarrow \dots \rightarrow \mathbf{Alg}(\mathbb{T}_1) \rightarrow \mathbf{Coalg}(\mathbb{K}) \rightarrow \mathbf{Alg}(\mathbb{T}) \rightarrow \mathbf{C}.$$

The question naturally arises as to whether this sequence stops in a meaningful way or continues indefinitely. M. Barr has provided a complete answer in [11] for the category of sets, pointed sets and vector spaces over a field. In the category of distributive lattices endowed with the ideal functor, it actually follows from B. Jacobs’ results ([12, Theorem 4.5]) on Kock-Zöberlein monads, also called lax idempotent monads, that $\mathbf{Alg}(\mathbb{T})$ and $\mathbf{Alg}(\mathbb{T}_1)$ are equivalent.

In this paper, we show that it is sufficient for \mathbf{C} to have either equalizers or coequalizers, for $\mathbf{Alg}(\mathbb{T})$ and $\mathbf{Alg}(\mathbb{T}_1)$ to be equivalent in the sense that $\mathbf{Alg}(\mathbb{T})$ is monadic over $\mathbf{Coalg}(\mathbb{K})$. Thus we reproduce B. Jacobs’ result, but free of any condition on the monad \mathbb{T} . With an additional mild condition on \mathbb{T} , we can lift this equivalence to the level of \mathbf{C} and the category of free \mathbb{T} -algebras, hence generalising the Stone representation theorem for distributive lattices.

The structure of the paper is very simple. We first introduce the necessary background in Section 2, and then give a brief representation of algebras and coalgebras through split coequalizers and split equalizers in Section 3. This representation is necessary to compensate for the absence of the chain of Galois connections used for lax idempotent monads ([12–16]). The main results are then shown and discussed in Section 4 together with a few illustrations. One key tool that we use is the Fakir construction of an idempotent approximation of a monad [17, 18]. Its main idea stems from the observation that there is a natural and intimate relationship between

¹“Just as in ring theory, the concepts of elements and ideals are of utmost importance in lattice theory.” ([8])

\mathbb{T} -algebras and equalizers of pairs (Te_X, e_{TX}) . The proofs then rely almost entirely on the manipulation of split equalizers and split coequalizers. In the case of the equivalence in Theorem 12, a split coequalizer is traced around the chain diagram described above. We reproduce here a step of B. Jacobs' proof of [12, Theorem 4.5], and duly reference it, in order to properly separate it from the context of a lax idempotent monad and for the presentation to be self-contained. We end the paper with a note on projectives in the category $\mathbf{Alg}(\mathbb{T})$; this is one categorical synthesis of the characterisations given in the papers [19–22].

2 Preliminaries and background

Distributive lattices and frames.

A distributive lattice ([7, 9]) is a partially ordered set² set admitting finite joins and finite meets³, and for which the identity

$$a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$$

and its order-dual hold. A homomorphism between distributive lattices is understood as a function that preserves the operations \wedge and \vee , including the top element $\wedge \emptyset = 1$ and the bottom element $\vee \emptyset = 0$. These form the category \mathbf{DLat} .

A frame ([7, 23]) L is a complete lattice with the equational identity

$$a \wedge \left(\bigvee S \right) = \bigvee \{a \wedge s \mid s \in S\}$$

for all $a \in L$ and any $S \subseteq L$. Frame homomorphisms or frame maps are functions $f : L \rightarrow M$ that preserve finite meets and arbitrary joins. Frames and their homomorphisms form a category denoted by \mathbf{Frm} . There is forgetful functor $\mathbf{Frm} \rightarrow \mathbf{DLat}$.

Definition 1. A subset $J \subseteq L$ is called an ideal if J is a downset, i.e. $b \in J$ provided there is $a \in J$ with $b \leq a$, and if it is closed under the formation of finite joins.

The set of all ideals on a frame L is denoted by $\mathfrak{I}L$. $\mathfrak{I}L$ is a frame with the following operations: meets are given by set-intersections and joins given by

$$\bigvee \mathcal{J} = \bigcup \{I_1 \vee I_2 \vee \cdots \vee I_n \mid I_1, I_2, \dots, I_n \in \mathcal{J} \text{ and } n \in \mathbb{N}\},$$

where $I_1 \vee I_2 \vee \cdots \vee I_n = \{i_1 \vee i_2 \vee \cdots \vee i_n \mid i_k \in I_k \text{ for } 1 \leq k \leq n\}$. Thus, if \mathcal{J} is a directed set, then $\bigvee \mathcal{J} = \bigcup \mathcal{J}$. If $f : L \rightarrow M$ is a homomorphism between distributive lattices, then the function $\mathfrak{I}f : \mathfrak{I}L \rightarrow \mathfrak{I}M$ given by

$$\mathfrak{I}f(I) = \{b \mid b \leq f(a) \text{ for some } a \in I\},$$

²In the context of the paper, "set" means small set.

³The lattice is then bounded.

is a frame homomorphism. In particular \mathfrak{J} is an endofunctor on **DLat**. The frames of the form $\mathfrak{J}D$, where D is a distributive lattice, are called *coherent frames* ([7, 24]). We are interested in the way-below relation \ll which is defined by

$$a \ll b \text{ if for any } S \subseteq L, \text{ if } b \leq \bigvee S \text{ then } a \leq \bigvee F \text{ for some finite } F \subseteq S.$$

A frame L is said to be *stably compact* ([9, 25, 26]) if \ll is a sublattice of $L \times L$ and if for all $a \in L, a = \bigvee \{b \mid b \ll a\}$. The category of stably compact frames and frame homomorphisms that preserve⁴ \ll is denoted by **StKFrm**. The category of coherent frames and frame homomorphisms preserving *compact elements*, i.e. elements satisfying $a \ll a$, is denoted by **CohFrm**. We have inclusion functors **CohFrm** \rightarrow **StKFrm** \rightarrow **Frm**.


Notations for natural transformations.

Given two pairs of functors $F, G : \mathbf{C} \rightarrow \mathbf{D}$ and $H, K : \mathbf{D} \rightarrow \mathbf{E}$ and two natural transformations $\alpha : F \rightarrow G$ and $\beta : H \rightarrow K$, the horizontal composition $\beta \circ \alpha : HF \rightarrow KG$ is defined by

$$(\beta \circ \alpha)_X = \beta_{GX}H(\alpha_X) = K(\alpha_X)\beta_{FX}$$

for each X in \mathbf{C} , for which we simply write $\beta \circ \alpha = \beta G \cdot H\alpha = K\alpha \cdot \beta F$. If $L : \mathbf{C} \rightarrow \mathbf{D}$ is another functor and $\delta : G \rightarrow L$ another natural transformation, then the vertical composition $\delta \cdot \alpha : F \rightarrow L$ is simply defined as $\delta_X \cdot \alpha_X$ for each $X \in \mathbf{C}$.

Lemma 1. (*Middle-Interchange Law*) *Given natural transformations*



we have $(\beta' \circ \alpha') \cdot (\beta \circ \alpha) = (\beta' \cdot \beta) \circ (\alpha' \cdot \alpha)$. ([27, 28].)

Monads.

Recall that a *monad* ([27–30]) \mathbb{T} on a category \mathbf{C} is a triple (T, m, e) , where $m : TT \rightarrow T$ and $e : 1 \rightarrow T$ are natural transformations satisfying the identities

$$m \cdot Tm = m \cdot mT \text{ and } m \cdot eT = m \cdot Te = 1_T.$$

Example 1. *The ideal functor \mathfrak{J} , together with the natural transformations provided by the set-union⁵ $\bigcup : \mathfrak{J}\mathfrak{J} \rightarrow \mathfrak{J}$ and the principal ideal map $\downarrow : 1 \rightarrow \mathfrak{J}$, is a monad (Cf. [9, Section 2] and [7, Exercise 4.6]).*

⁴These are called proper maps or perfect maps. The condition is equivalent to the right adjoint of the frame map preserving directed joins.

⁵An ideal on $\mathfrak{J}L$ is by definition directed.

A \mathbb{T} -algebra (or an *Eilenberg-Moore algebra*) is a pair (X, a) , where $X \in \mathbf{C}$ and $a : TX \rightarrow X$ a morphism such that

$$a \cdot Ta = a \cdot m_X \text{ and } a \cdot e_X = 1_X.$$

If (X, a) and (Y, b) are \mathbb{T} -algebras, then a \mathbb{T} -algebra homomorphism $f : (X, a) \rightarrow (Y, b)$ is a morphism $f : X \rightarrow Y$ in \mathbf{C} such that $f \cdot a = b \cdot Tf$. The category of \mathbb{T} -algebras and \mathbb{T} -algebra homomorphisms is denoted by $\mathbf{C}^{\mathbb{T}}$ or by $\mathbf{Alg}(\mathbb{T})$. The forgetful functor $G^{\mathbb{T}} : \mathbf{C}^{\mathbb{T}} \rightarrow \mathbf{C} : (X, a) \mapsto X$ admits a left adjoint $F^{\mathbb{T}} : \mathbf{C} \rightarrow \mathbf{C}^{\mathbb{T}}$ defined by $F^{\mathbb{T}}(f : X \rightarrow Y) = Tf : (TX, m_X) \rightarrow (TY, m_Y)$. The unit of this adjunction is given by $e_X : X \rightarrow G^{\mathbb{T}}F^{\mathbb{T}}X$ and the co-unit is provided by the algebra morphisms $\varepsilon_{(X, a)} = a : F^{\mathbb{T}}G^{\mathbb{T}}(X, a) \rightarrow (X, a)$. It is important to note in addition that $G^{\mathbb{T}}$ reflects isomorphisms.

Definition 2. ([27, 30]) *The diagram*


$$\begin{array}{ccc} A & \xrightarrow{f} & B & \xrightarrow{q} & C \\ & \searrow g & & \swarrow s & \\ & & t & & s \end{array}$$

is said to be a *split coequalizer* if $q \cdot f = q \cdot g$, $q \cdot s = 1_C$, $f \cdot t = 1_B$ and $s \cdot q = g \cdot t$.

Split coequalizers are absolute, i.e. they are preserved by any functor.

Any adjunction $(F \dashv G, \eta, \epsilon)$ where $G : \mathbf{A} \rightarrow \mathbf{C}$ and $F : \mathbf{C} \rightarrow \mathbf{A}$ induces a monad (GF, GeF, η) on \mathbf{C} ([29]). If the adjunction $F \dashv G$ induces the same monad \mathbb{T} on \mathbf{C} , that is $GF = T$, $GeF = m$ and $\eta = e$, then there is a unique functor $C : \mathbf{A} \rightarrow \mathbf{C}^{\mathbb{T}}$ - called *comparison functor*, such that $CF = F^{\mathbb{T}}$ and $G^{\mathbb{T}}C = G$. We say that \mathbf{A} is monadic (resp. strictly monadic) over \mathbf{C} if C is an equivalence (resp. isomorphism). Beck's monadicity criterion ([27, 30, 31]) implies in this case that G creates coequalizers of G -split pairs: i.e. if $f = Gu$ and $g = Gv$ in Definition 2, then u and v admit a coequalizer r in $\mathbf{Alg}(\mathbb{T})$ such that the G -image of the diagram $r \cdot u = r \cdot v$ is isomorphic to the split coequalizer. This relation of isomorphism is strict for $G^{\mathbb{T}}$.

The adjunction $F^{\mathbb{T}} \dashv G^{\mathbb{T}}$ induces a comonad $\mathbb{K} = (T, F^{\mathbb{T}}eG^{\mathbb{T}}, \epsilon)$ on the category of algebras $\mathbf{Alg}(\mathbb{T})$, where $F^{\mathbb{T}}eG^{\mathbb{T}} = Te : (T, m) \rightarrow (TT, mT)$. This generates an alternating sequence of monads and comonads as follows:



We have sequences of comparison functors $F_n^{\mathbb{T}} : \mathbf{Alg}(\mathbb{T}_{n-1}) \rightarrow \mathbf{Alg}(\mathbb{T}_n)$ and $C_n^{\mathbb{K}} : \mathbf{Coalg}(\mathbb{K}_{n-1}) \rightarrow \mathbf{Coalg}(\mathbb{K}_n)$ for $n \geq 1$, where $\mathbb{T}_0 = \mathbb{T}$ and $\mathbb{K}_0 = \mathbb{K}$.

Definition 3. A category \mathbf{C} is said to be order-enriched if each hom-sets $\text{Hom}(X; Y)$ is a partially ordered set for any pair $X, Y \in \mathbf{C}$ and the composition of morphisms preserves the order. The monad \mathbb{T} is said to be order-enriched if the underlying functor T preserves the order.

Definition 4. ([27, 32]) An order-enriched monad \mathbb{T} is said to be lax idempotent or a Kock-Zöberlein monad if $Te_X \leq e_{TX}$ for any $X \in \mathbf{C}$.

We will omit mentioning that \mathbb{T} is order-enriched each time we consider a lax idempotent monad, as this is clear from the context⁶.

Lemma 2. ([27, Section II.4.9]) For a monad \mathbb{T} , we have the following equivalences for any $X \in \mathbf{C}$:

$$Te_X \leq e_{TX} \iff Te_X \dashv m_X \iff m_X \dashv e_{TX}.$$

In such a case, any morphism $a : TX \rightarrow X$ with $a \cdot e_X = 1_X$ necessarily satisfies $a \dashv e_X$ and this makes (X, a) a \mathbb{T} -algebra.

We mention the following contribution by B. Jacobs.

Theorem 3. ([12]) For a lax idempotent monad \mathbb{T} on a category \mathbf{C} with equalisers, one has $\mathbf{Alg}(\mathbb{T}) \simeq \mathbf{Alg}(\mathbb{T}_1)$.

3 Presentation of algebras and coalgebras

A \mathbb{T} -algebra (X, a) can be presented as in the following split coequalizer ([27, 30]):

$$TTX \begin{array}{c} \xrightarrow{Ta} \\ \xrightarrow{m_X} \\ \xleftarrow{e^T} \end{array} TX \begin{array}{c} \xrightarrow{a} \\ \xleftarrow{e} \end{array} X.$$

⁶Unlike in [27], we consider the order to be separated, i.e. $x \leq y$ and $y \leq x$ imply $x = y$.

A \mathbb{K} -coalgebra structure on (X, a) is then a \mathbb{T} -algebra morphism $c : (X, a) \rightarrow (TX, m_X)$ that is part of the following split equalizer:

$$(X, a) \begin{array}{c} \xrightarrow{c} \\ \xleftarrow{a} \end{array} (TX, m_X) \begin{array}{c} \xrightarrow{Tc} \\ \xleftarrow{Te} \\ \xrightarrow{m_X} \end{array} (TTX, m_{TX}).$$

We shall denote this \mathbb{K} -coalgebra as an ordered triple (X, a, c) . Thus a \mathbb{T}_1 -algebra structure on (X, a, c) is a \mathbb{K} -coalgebra morphism $b : (TX, m_X, Te_X) \rightarrow (X, a, c)$ that is a split coequalizer:

$$(TTX, m_{TX}, Te_{TX}) \begin{array}{c} \xrightarrow{Tb} \\ \xleftarrow{Ta} \\ \xrightarrow{Te} \end{array} (TX, m_X, Te_X) \begin{array}{c} \xrightarrow{b} \\ \xleftarrow{c} \end{array} (X, a, c).$$

A \mathbb{T}_1 -algebra as described above is denoted by (X, a, c, b) . Thus free \mathbb{K} -algebras and free \mathbb{T}_1 -algebras are given by (TX, m_X, Te_X) and (TX, m_X, Te_X, Ta) respectively so that the functor $F^{\mathbb{T}}$ restricts itself to the comparison functor $F_1^{\mathbb{T}} : \mathbf{Alg}(\mathbb{T}) \rightarrow \mathbf{Alg}(\mathbb{T}_1)$ between the adjunctions that induce the monad \mathbb{T}_1 . It is a restriction in a sense that

$$(G^{\mathbb{K}}G^{\mathbb{T}_1}) \cdot F_1^{\mathbb{T}} = K = F^{\mathbb{T}} \cdot G^{\mathbb{T}}.$$

For a lax idempotent monad, the description of the algebras and coalgebras can be greatly simplified. In the table below, we list the identities characterising the algebras and coalgebras for a lax idempotent monad.

(co)Monads	(co)Algebras
$\mathbb{T} = (T, m, e)$	$a \dashv e_X$ and $a \cdot e_X = 1$
$\mathbb{K} = (T, Te, a)$	$c \dashv a$ and $a \cdot c = 1$
$\mathbb{T}_1 = (T, Ta, c)$	$b \dashv c$ and $b \cdot c = 1$
...	...

The simplification in the table is due to various work that can be traced back to the work of R.-E. Hoffman on continuous posets [13], the papers by A. Kock ([32]) and V. Zöberlein ([33]) on lax idempotent monads. These characterisations are also described by B. Jacobs in [12].

Remark 1. For a lax idempotent monad \mathbb{T} , the fact that $c : X \rightarrow TX$ is as well a \mathbb{T} -algebra morphism is due to its being a left adjoint of a . This is given by [32, Proposition 2.5] and [16, Proposition 3.8]. Split algebras (Cf. [16, Definition 3.6] and [14]) are then characterised as coalgebras of the comonad \mathbb{K} .

We mention the following:

- Theorem 4.** 1. A distributive lattice D is a frame if and only if $\downarrow: D \rightarrow \mathfrak{J}D$ admits a left adjoint a such that $a \cdot \downarrow = 1_D$. In this case, the left adjoint is given by the join \bigvee .
2. A frame L is stably compact if and only if the join $\bigvee: \mathfrak{J}L \rightarrow L$ admits a left adjoint c such that $\bigvee \cdot c = 1_D$. Here $c(x) = \{y \in L \mid y \ll x\}$ ⁷.

With the help of Lemma 2, the above characterisation can be deduced from the facts that **Frm** is the category of algebras for the ideal monad ([7, Exercise 4.6]), and that **StKFrm** is the category of coalgebras of the comonad $(\mathfrak{J}, \mathfrak{J}(\downarrow), \bigvee)$ as shown in [25, Section 3]. It is known since [32, Theorem 4.2] and [13] that the way below relation \ll on a poset is directly linked to the condition of a certain join map admitting a left adjoint. We refer the reader to the compendium [34], as well as the seminal paper [35] by D. Scott and the paper [36] by Alan Day for a brief background on continuous lattices. We note that D. Scott was the first to introduce and systematically study continuous lattices. The category of algebras of the induced monad $(\mathfrak{J}, \mathfrak{J}(\bigvee), c)$ is also familiar to us:

Theorem 5. The algebras of the monad $(\mathfrak{J}, \mathfrak{J}(\bigvee), c)$ are characterised by those stably compact L for which $c_L: L \rightarrow \mathfrak{J}L$ admits a left adjoint $b: \mathfrak{J}L \rightarrow L$ such that $b \cdot c_L = 1_L$.

Proof. This can be obtained by following an argument similar to [15, Theorem 3.4, Lemma 3.6] and [13, Theorem 2.6]. \square

We observe that b , being a left adjoint to c , is given by:

$$b(J) = \bigvee \{y \in J \mid y \ll y\}.$$

L then becomes a coherent frame, that is $L = \mathfrak{J}D$ where $D = \{y \in L \mid y \ll y\}$. However, not all coherent frames satisfy Theorem 5. The following result is at the heart of the equivalence in Theorem 3.

Proposition 6. ([12]) Let \mathbb{T} be a lax idempotent monad and let us assume that e_X is the equalizer of the pair (e_{TX}, Te_X) for a fixed X . It follows that X admits a \mathbb{T} -algebra structure if and only if TX admits a \mathbb{T}_1 -algebra structure.

Proof. The necessary condition is obvious. Conversely, suppose that $b \cdot Te_X = 1_{TX}$ exists. As shown in [12, Theorem 4.5], the sequence $b \dashv Te_X \dashv m_X \dashv e_{TX}$ makes e_X a split monomorphism: we have

$$\begin{aligned} Te_X \cdot b \cdot e_{TX} &= Tb \cdot Te_{TX} \cdot e_{TX} \text{ (} b \text{ is a coalgebra map)} \\ &= Tb \cdot e_{TTX} \cdot e_{TX} \\ &= e_{TX} \cdot b \cdot e_{TX}. \end{aligned}$$

There is a unique morphism $r: TX \rightarrow X$ such that $e_X \cdot r = b \cdot e_{TX}$. Composing with e_X from the right gives $e_X \cdot r \cdot e_X = b \cdot Te_X \cdot e_X = e_X$. One has $r \cdot e_X = 1$ ([12]). \square

⁷This expression is forced by the identities $\bigvee \cdot c = 1_D$ and $c \cdot \bigvee \leq 1_{\mathfrak{J}D}$.

Remark 2. For a distributive lattice D , since \mathbf{DLat} is monadic over \mathbf{Set} ([7, Proposition 3.7]), $\downarrow: D \rightarrow \mathfrak{J}D$ is an equalizer of $(\downarrow_{\mathfrak{J}D}, \mathfrak{J}(\downarrow_D))$. Consequently, a coherent frame $\mathfrak{J}D$ is an algebra of the monad $(\mathfrak{J}, \mathfrak{J}(\vee), c)$ if and only if D is a frame. A similar situation is described in [13, Corollary 2.7] for algebraic lattices, and subsequently discussed in [14, Section 8].

4 Natural equivalences

In order to show that under some mild conditions $F_1^{\mathbb{T}}$ is an equivalence, we shall introduce a functor $\Phi: F^{\mathbb{T}}(\mathbf{C}) \rightarrow \mathbf{C}$ that is provided by the Fakir construction of idempotent approximation of a monad ([17]). For a given monad $\mathbb{T} = (T, m, e)$, consider the equalizer

$$T^\varphi \xrightarrow{\varphi} T \xrightleftharpoons[eT]{Te} TT.$$

Theorem 7. (Fakir [17]) The functor T^φ underlies a monad $\mathbb{I}_T = (T^\varphi, m^\varphi, e^\varphi)$ where m^φ is unique such that $\varphi \cdot m^\varphi = m \cdot (\varphi \circ \varphi)$ and e^φ is unique such that $\varphi \cdot e^\varphi = e$.

We mention the following results which are of interest to us.

Proposition 8. ([17, Proposition 3]) $e^\varphi T: T \rightarrow T^\varphi T$ is an isomorphism, and the following are equivalent:

- \mathbb{I}_T is idempotent;
- $Te^\varphi: T \rightarrow TT^\varphi$ is an isomorphism;

Now consider the category $F^{\mathbb{T}}(\mathbf{C})$ of free \mathbb{T} -algebras and morphisms $F^{\mathbb{T}}f$ where f is in \mathbf{C} . We consider the functor $\theta^{\mathbb{T}}: \mathbf{C} \rightarrow F^{\mathbb{T}}(\mathbf{C})$ so that if $E^{\mathbb{T}}$ is the inclusion functor $F^{\mathbb{T}}(\mathbf{C}) \rightarrow \mathbf{Alg}(\mathbb{T})$, then $E^{\mathbb{T}} \cdot \theta^{\mathbb{T}} = F^{\mathbb{T}}$. In a similar manner, we consider $\theta_1^{\mathbb{T}}: \mathbf{Alg}(\mathbb{T}) \rightarrow F_1^{\mathbb{T}}(\mathbf{Alg}(\mathbb{T}))$, where $F_1^{\mathbb{T}}(\mathbf{Alg}(\mathbb{T}))$ is the category of free \mathbb{T}_1 -algebras with morphisms of the form $F_1^{\mathbb{T}}f$, as well as the inclusion $E_1^{\mathbb{T}}: F^{\mathbb{T}}(\mathbf{C}) \rightarrow \mathbf{Alg}(\mathbb{T})$. We have

$$\begin{array}{ccccc} \mathbf{C} & \xrightarrow{\theta^{\mathbb{T}}} & F^{\mathbb{T}}(\mathbf{C}) & \xrightarrow{E^{\mathbb{T}}} & \mathbf{Alg}(\mathbb{T}) \\ \uparrow & & \uparrow & & \uparrow \\ \mathbf{Alg}(\mathbb{T}) & \xrightarrow{\theta_1^{\mathbb{T}}} & F_1^{\mathbb{T}}(\mathbf{Alg}(\mathbb{T})) & \xrightarrow{E_1^{\mathbb{T}}} & \mathbf{Alg}(\mathbb{T}_1) \end{array}$$

where vertical functors are forgetful.

Definition 5. We define $\Phi: F^{\mathbb{T}}(\mathbf{C}) \rightarrow \mathbf{C}$ by $\Phi(F^{\mathbb{T}}f) = T^\varphi f$ for all f in \mathbf{C} . In particular $\Phi \cdot \theta^{\mathbb{T}} = T^\varphi$. This gives two natural transformations $e^\varphi: 1 \rightarrow \Phi \cdot \theta^{\mathbb{T}}$ and $\theta^{\mathbb{T}} e^\varphi: 1 \rightarrow \theta^{\mathbb{T}} \cdot \Phi$.

Theorem 9. If \mathbb{I}_T is an idempotent monad, then $\theta^{\mathbb{T}}(e^\varphi)$ is a natural isomorphism. In particular if \mathbb{I}_T is the identity monad, then we have an equivalence between the category of free \mathbb{T} -algebras and the ambient category \mathbf{C} .

Proof. We note that $\theta^{\mathbb{T}}e^\varphi = Te^\varphi : (T, m) \rightarrow (TT^\varphi, mT^\varphi)$. Since $G^{\mathbb{T}}$ reflects isomorphisms, Te^φ is an isomorphism if and only if $\theta^{\mathbb{T}}e^\varphi$ is. The result easily follows from Proposition 8. \square

We shall restrict Φ to $F_1^{\mathbb{T}}(\mathbf{Alg}(\mathbb{T})) \rightarrow \mathbf{Alg}(\mathbb{T})$. We first give a strengthening of the fact that $e^\varphi T$ is an isomorphism.

Lemma 10. *For a \mathbb{T} -algebra (X, a) , e_X^φ is an isomorphism.*

Proof. If (X, a) is a \mathbb{T} -algebra, then we have a split equalizer

$$\begin{array}{ccccc} X & \xrightarrow{e} & TX & \xrightleftharpoons[eT]{Te} & TTX, \\ & \searrow a & \swarrow & \searrow eT & \swarrow Ta \end{array}$$

which makes e_X^φ an isomorphism. \square

This allows us to define the restriction $\Phi_1 : F_1^{\mathbb{T}}(\mathbf{Alg}(\mathbb{T})) \rightarrow \mathbf{Alg}(\mathbb{T})$ of Φ by $\Phi_1(TX, m_X, Te, Ta) = (X, a)$ and $\Phi_1(F_1^{\mathbb{T}}f) = f$.

Lemma 11. *Suppose that \mathbf{C} has either equalizers or coequalizers. If (X, a, c, b) is a \mathbb{T}_1 -algebra, then we have the following split coequalizer and split equalizer*

$$\begin{array}{ccccccc} TX & \xrightleftharpoons[a]{b} & X & \xrightarrow{r} & X_c & \xrightarrow{\kappa} & X & \xrightleftharpoons[e]{c} & TX \\ & \searrow a & \swarrow & \searrow \kappa & \swarrow r & \searrow e & \swarrow c & \swarrow b \end{array}$$

Proof. The identities $a \cdot e = 1$ and $b \cdot c = 1$ hold as (X, a, b, c) is a \mathbb{T}_1 -algebra. Suppose that \mathbf{C} has equalizers⁸ and denote by κ the equalizer of (c, e) . As shown in the proof of [12, Theorem 4.5]: $c \cdot b \cdot e = Tb \cdot Te \cdot e = Tb \cdot eT \cdot e = e \cdot b \cdot e$; there is a unique r such that $\kappa \cdot r = b \cdot e$. One shows that $\kappa \cdot r \cdot \kappa = b \cdot c \cdot \kappa = \kappa$ and so $r \cdot \kappa = 1$. We have a split equalizer on the right. It remains to show that $r \cdot a = r \cdot b$ to have a split coequalizer on the left. We have

$$\begin{aligned} \kappa \cdot r \cdot b &= b \cdot e \cdot b \\ &= b \cdot Tb \cdot eT \\ &= b \cdot Ta \cdot eT \text{ (} b \text{ coequalizes } Ta \text{ and } Tb) \\ &= b \cdot e \cdot a \\ &= \kappa \cdot r \cdot a. \end{aligned}$$

Since κ is a monomorphism, the identity $r \cdot a = r \cdot b$ follows. \square

Theorem 12. *If \mathbf{C} admits either coequalizers for \mathbb{T} -algebra morphisms or equalizers for any parallel pair of morphisms, then $F_1^{\mathbb{T}}$ is an equivalence.*

⁸The case where it has coequalizers is treated analogously.

Proof. First, since the restriction of \mathbb{I}_T on $\mathbf{Alg}(\mathbb{T})$ is essentially the identity monad, we have $\Phi_1 \cdot \theta_1^\mathbb{T} \cong 1$ and $\theta_1^\mathbb{T} \cdot \Phi_1 \cong 1$. Now, let (X, a, c, b) be a \mathbb{T}_1 -algebra and consider the split coequalizer from Lemma 11:

$$\begin{array}{ccc} TX & \begin{array}{c} \xrightarrow{b} \\ \xrightarrow{a} \end{array} & X & \xrightarrow{r} & X_c \\ & \searrow e & & \swarrow \kappa & \\ & & & & \end{array}$$

As an absolute coequalizer, this is preserved by the comparison functor from \mathbf{C} to $\mathbf{Coalg}(\mathbb{K})$ which arises from the adjunctions inducing the comonad \mathbb{K} on $\mathbf{Alg}(\mathbb{T})$. Thus we have a split coequalizer

$$\begin{array}{ccccc} (TTX, m_{TX}, Te_{TX}) & \begin{array}{c} \xrightarrow{Tb} \\ \xrightarrow{Ta} \end{array} & (TX, m_X, Te_X) & \xrightarrow{Tr} & (TX_c, m_{X_c}, Te_c) \\ & \searrow Te & & \swarrow T\kappa & \\ & & & & \end{array}$$

in $\mathbf{Coalg}(\mathbb{K})$. Now, both Ta and Tb are morphisms of (free) \mathbb{T}_1 -algebras. By Beck's monadicity theorem, since $\mathbf{Alg}(\mathbb{T}_1)$ is monadic over $\mathbf{Coalg}(\mathbb{K})$ in the strict sense, there is a \mathbb{T}_1 -algebra structure γ on (TX_c, m_{X_c}, Te_c) and Tr becomes a \mathbb{T}_1 -algebra morphism. The universality of the coequalizer b in $\mathbf{Alg}(\mathbb{T}_1)$ shows that

$$(TX_c, m_{X_c}, Te_c, \gamma) \cong (X, a, c, b) \text{ and } Tr \cong b \cong \gamma.$$

Finally since $\mathbf{Alg}(\mathbb{T})$ is monadic over \mathbf{C} , X_c admits a \mathbb{T} -algebra structure a_c , as both a and b are \mathbb{T} -algebra morphisms. The isomorphism $TX_c \cong X$ implies that $r \cong a_c$ as morphisms of \mathbb{T} -algebras. If $f : (X, a, c, b) \rightarrow (X', a', c', b')$ is a \mathbb{T}_1 -algebra morphism, then we form the (split) equalizers κ, κ' and (split) coequalizers r, r' . There is a unique morphism f_c such that $f_c \cdot r = r' \cdot f$ and $f \cdot \kappa = \kappa' \cdot f_c$. Thus f_c is a \mathbb{T} -algebra morphism and $\theta_1^\mathbb{T} f_c \cong f$. It follows that the inclusion $E_1^\mathbb{T}$ is an equivalence. \square

Corollary 13. *With the assumption of Theorem 12, $F^\mathbb{K}$ is monadic, $F^{\mathbb{T}_1}$ is comonadic and the action of \mathbb{K}_1 on $\mathbf{Alg}(\mathbb{T}_1)$ is equivalent to that of \mathbb{K} on $\mathbf{Alg}(\mathbb{T})$.*

We note that B. Jacobs' result in [12] requires that certain equalizers exist in \mathbf{C} in order to construct the object X_c . The identification of the algebra (X, a, c, b) with $(TX_c, m_{X_c}, Te_c, Ta_c)$ proceeds from the fact that for a lax idempotent monad, the adjoints $a \dashv c \dashv b$ coincide with $m_{X_c} \dashv Te_c \dashv Ta$ due to the property of universality of Galois connections. The verification of the equivalence in Theorem 3 then requires one to check that the relevant diagrams commute. The absence of the Galois connections has been remedied here by the use of the characterisation of algebras through split coequalizers, reaffirming their importance in the theory of monads.

Example 2. *We are mostly interested in the ideal functor and the down set functor. Similar examples that are treated by Bart Jacobs in [15] and [12] are assumed to be known. These*

functors have the common property that they are lax idempotent, and so the results in [15] apply. Here, we augment these results with the equivalence from Theorem 9.

- (i) In the category **DLat**, the first inductive step $(\mathfrak{J}^\varphi, \mathfrak{V}^\varphi, \mathfrak{D}^\varphi)$ of Fakir construction is essentially the identity monad (Cf. Remark 2). The equivalence between **DLat** and the category **CohFrm** then follows from Theorem 9. This is part of the Stone representation for distributive lattices ([7, Section 3.3], [24]). We emphasize the fact that **DLat** is complete and so Theorem 12 applies as well. A case that is analogous to this is that of the category $\sigma\mathbf{Frm}$ ([37]) of σ -frames with the monad \mathcal{H} taking a σ -frame L to the frame $\mathcal{H}L$ of its σ -ideals. This establishes an equivalence between $\sigma\mathbf{Frm}$ and the category $\sigma\mathbf{CohFrm}$ of σ -coherent frames via Theorem 9. The functor Φ (Definition 5) is given by the functor Lind that extracts the Lindelöf elements from the free algebras $\mathcal{H}L$.
- (ii) Consider the category **MLat** of meet-semilattices and meet-semilattice homomorphisms with the downset functor $\mathfrak{D} : \mathbf{MLat} \rightarrow \mathbf{MLat}$. This form a monad \mathbb{T} with the natural transformations given by set-theoretic unions $\cup : \mathcal{D}\mathcal{D} \rightarrow \mathcal{D}$ and downset maps $\downarrow : 1 \rightarrow \mathcal{D}$. The category of algebras $\mathbf{Alg}(\mathbb{T})$ is **Frm** and the category **Coalg**(\mathbb{K}) is the category of supercontinuous frames with their corresponding homomorphisms ([1, 20]). Here $(\mathfrak{D}^\varphi, \mathfrak{V}^\varphi, \mathfrak{D}^\varphi)$ is essentially the identity. Theorem 9 shows that **MLat** is equivalent to the category **SCohFrm** of supercontinuous frames (Cf. [20, Remark 3] and [1, Proposition 6.2]).
- (iii) In [21], Z. Dhongsheng considers the functor $Z : \mathbf{MLat} \rightarrow \mathbf{MLat}$ which is a submonad \mathbb{T} of \mathfrak{D} from the previous example⁹. From this follow three subcategories: the category of Z -frames, the category of stably Z -continuous semilattices, and the category of coherent Z -frames. These are respectively the category $\mathbf{Alg}(\mathbb{T})$, the category **Coalg**(\mathbb{K}) and the category of free \mathbb{T} -algebras. Theorem 9 applies here as well.

Example 3. The following classical examples trivially satisfy the condition of Theorem 12: the category of topological spaces and continuous maps with the ultrafilter monad or with the prime open filter monad ([9, 38]), the category of T_0 -spaces and continuous maps with the open filter monad ([16, 39, 40]). For those which are lax idempotent, B. Jacobs' result applies to these instances. Theorem 12 allows for the extension of the results to arbitrary monads. However, for those classical instances we do not know if the first inductive step of the Fakir idempotent approximation reduces to the identity monad.

Example 4. M. Barr considers the following examples in [11]: **Set**, the category $(1, \mathbf{Set})$ of pointed sets, and the category \mathbf{V}_K of vector spaces over a field K and looks at an arbitrary monad \mathbb{T} . It is shown that - up to constants - **C** is equivalent to **Coalg**(\mathbb{K}) through the comparison functor ([11, Theorem 11]). Since **Set** is complete, Theorem 12 applies, confirming the general result that the sequence stabilises. However, in this case our result is weaker than the one obtained in [11] by M. Barr. Indeed he showed that $F^{\mathbb{T}}$ is essentially comonadic as well, whereas Theorem 12 does not necessarily imply this, as illustrated by Example 2. At the same time, Example 2 indicates that it is not easy, if not impossible, to

⁹That this is a submonad is straightforward, given the requirements of Definition 1.1 in [21]

borrow the assumptions and method used by M. Barr and adapt them for categories enriched over partially ordered sets.

5 Projectives in the category of algebras

One of the most interesting aspects of Example 2 is the characterisations of projectives in the ambient category as shown in [20, Proposition 5], [22, Proposition 2.2] and [21, Theorem 3.2]. These characterisations actually depend on the pair of forgetful-free functors as a parameter that renders the ambient category a category of algebras. However this is not explicitly mentioned. Indeed, J. Madden writes in [22] with regard to the characterisations of stably compact frames as retracts of coherent frames ([19]) that *I am sure Johnstone himself has long been aware of this interpretation of his result, but it seems to have escaped many others.* We give here a definition of projectives that varies according to the monad \mathbb{T} and that specialises into the specific results obtained in the above-mentioned sources. This definition is an easy dualisation of the notion of injective spaces in the sense of Escardó ([39, 40]).

Definition 6. *Let \mathbb{T} be a monad on a category \mathbf{C} . An algebra (X, a) is projective in $\mathbf{Alg}(\mathbb{T})$ or simply projective if (X, a) is projective with respect to the (co)free functor $F^{\mathbb{K}}$.*

The above definition subsumes Definition 2.1 of [22], the definition of E -projective Z -frames in [21] and the definition of projective frames in [20]. With regard to lax idempotent monads, we have the following equivalences.

Theorem 14. ([16, 39, 40]) *Let \mathbb{T} be a lax idempotent monad on a category \mathbf{C} . The following are equivalent:*

1. (X, a) is projective with respect to $\{\alpha : (TA, m_A) \rightarrow (A, \alpha) \mid (A, \alpha) \in \mathbf{Alg}(\mathbb{T})\}$.
2. (X, a) has a structure of a \mathbb{K} -coalgebra.
3. (X, a) is a retract in $\mathbf{Alg}(\mathbb{T})$ of free \mathbb{T} -algebras.
4. (X, a) is a \mathbb{T} -split algebra. (Cf. [16, Definition 3.6])

\mathbb{K} -coalgebras are therefore projectives. In the absence of the condition of lax idempotency, we only have the following characterisation.

Theorem 15. *Projectives are precisely the retracts in $\mathbf{Alg}(\mathbb{T})$ of free \mathbb{T} -algebras.*

Example 5. *The projectives in the particular instances of Example 2 are respectively: the stably compact frames ([19, 22, 24]) and the stably continuous σ -frames ([22, 37]), the stably supercontinuous frames ([20, Proposition 5]), the stably Z -continuous semilattices ([21, Theorem 3.2]).*

Example 6. (Gleason, 1958) [7, 29, 41] *If \mathbb{T} is the monad generated by the forgetful functor from the category \mathbf{KHaus} of compact Hausdorff spaces to \mathbf{Set} and the free functor $\beta : \mathbf{Set} \rightarrow \mathbf{KHaus}$ - which is the Stone-Čech compactification of a discrete space, then the*

*projectives with respect to \mathbb{T} are exactly the extremally disconnected spaces in **KHaus**. Indeed, these are precisely the retracts of spaces of the form βX .*

Acknowledgements. I thank B. Jacobs for brief and useful communication with regard to Theorem 4.5 of his paper [12].

Declarations

- No funding was obtained for this study.
- The author declares that there is no conflict of interests.
- Availability of data and materials not available.

References

- [1] Banaschewski, B., Pultr, A.: Adjointness aspects of the down-set functor. *Appl. Categ. Structures* **9**(4), 419–436 (2001)
- [2] Banaschewski, B.: Compactification of frames. *Math. Nachr.* **149**, 105–116 (1990)
- [3] Bezhnashvili, G., Harding, J.: Stable compactifications. *Cahiers de Topologie et Géométrie Différentielle Catégoriques* **LV**(1), 37–65 (2014)
- [4] Frith, J., Schauerte, A.: Compactifications of partial frames via strongly regular ideals. *Mathematica Slovaca* **68**(2), 285–298 (2018)
- [5] Frith, J.L.: The category of uniform frames. *Cahiers de Topologie et Géométrie Différentielle Catégoriques* **31**(4), 305–313 (1990)
- [6] Banaschewski, B., Pultr, A.: Samuel compactification and completion of uniform frames. *Mathematical Proceedings of the Cambridge Philosophical Society* **108**, 63–78 (1990)
- [7] Johnstone, P.T.: *Stone Spaces*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge (1982)
- [8] Hofmann, K.H., Mislove, M., Stralka, A.: Application of duality to lattice theory, In: *The Pontryagin Duality of Compact O-Dimensional Semilattices and its Applications*, pp. 53–91. Springer, Berlin, Heidelberg (1974)
- [9] Simmons, H.: A couple of triples. *Topology and its Applications* **13**(2), 201–223 (1982)
- [10] Lawvere, F.: Unity and identity of opposites in calculus and physics. *Applied Categorical Structures* **4**(2-3), 167–174 (1996)
- [11] Barr, M.: Coalgebras in a category of algebras. In: Hilton, P.J. (ed.) *Category Theory, Homology Theory and Their Applications I*, pp. 1–12. Springer, Berlin, Heidelberg (1969)

- [12] Jacobs, B.: Bases as coalgebras. In: Corradini, A., Klin, B., Cîrstea, C. (eds.) *Algebra and Coalgebra in Computer Science*, pp. 237–252. Springer, Berlin, Heidelberg (2011)
- [13] Hoffmann, R.-E.: Continuous posets and adjoint sequences. *Semigroup Forum* **18**(2), 173–188 (1979)
- [14] Rosebrugh, R., Wood, R.E.: Split structures. *Theories and Applications of Categories* **13**(12), 172–183 (2004)
- [15] Jacobs, B.: Coalgebras and approximation. In: Nerode, A., Matiyasevich, Y.V. (eds.) *Logical Foundations of Computer Science*, pp. 173–183. Springer, Berlin, Heidelberg (1994)
- [16] Hofmann, D., Sousa, L.: Aspects of algebraic algebras. *Logical Methods in Computer Science* **13**, 1–25 (2017)
- [17] Fakir, S.: Monade idempotente associée à une monade. *Comptes Rendus de l'Académie des Sciences de Paris* **270**, 99–101 (1970)
- [18] Lambek, J., Rattray, B.A.: Localization at injectives in complete categories. *Proceedings of the American Mathematical Society* **41**(1), 1–9 (1973)
- [19] Johnstone, P.T.: The Gleason cover of a topos. II. *J. Pure Appl. Algebra* **22**(3), 229–247 (1981)
- [20] Banaschewski, B., Niefield, S.B.: Projective and supercoherent frames. In: *Proceedings of the Conference on Locales and Topological Groups (Curaçao, 1989)*, vol. 70, pp. 45–51 (1991)
- [21] Dongsheng, Z.: On projective z-frames. *Canadian Mathematical Bulletin* **40**(1), 39–46 (1997)
- [22] Madden, J.: κ -frames. *Journal of Pure and Applied Algebra* **70**(1), 107–127 (1991)
- [23] Picado, J., Pultr, A.: *Frames and Locales: Topology Without Points*. *Frontiers in Mathematics*, Birkhäuser Basel (2012)
- [24] Banaschewski, B.: Coherent frames. In: Banaschewski, B., Hoffmann, R.-E. (eds.) *Continuous Lattices*. *Proceedings of the Conference Held in Berlin, Univ. Bremen, November 9 - 11, 1979*, vol. 42, pp. 1–11. Springer, Germany (1981)
- [25] Banaschewski, B., Brümmer, G.C.L.: Stably continuous frames. *Mathematical Proceedings of the Cambridge Philosophical Society* **104**(1), 7–19 (1988)
- [26] Jung, A.: Stably compact spaces and the probabilistic powerspace construction. *Electronic Notes in Theoretical Computer Science* **87**, 5–20 (2004)

- [27] Hofmann, D., Seal, G.J., Tholen, W. (eds.): Monoidal Topology: A Categorical Approach to Order, Metric, and Topology. Encyclopedia of Mathematics and its Applications, p. 492. Cambridge University Press, Cambridge (2014)
- [28] MacLane, S.: Categories for the Working Mathematician. Graduate Texts in Mathematics. Springer, New York (1995)
- [29] Eilenberg, S., Moore, J.C.: Adjoint functors and triples. Illinois Journal of Mathematics **9**(3), 381–398 (1965)
- [30] Riehl, E.: Category Theory in Context. Dover Publications, Cambridge (2017)
- [31] Beck, J.: Triples, algebras and cohomology. Reprints in Theory and Applications of Categories **2**, 1–59 (2003)
- [32] Kock, A.: Monads for which structures are adjoint to units. Journal of Pure and Applied Algebra **104**(1), 41–59 (1995)
- [33] Zöberlein, V.: Doctrines on 2-categories. Mathematische Zeitschrift **148**(3), 267–279 (1976)
- [34] Gierz, G., *et al.*: A Compendium of Continuous Lattices. Springer, Heidelberg (1980)
- [35] Scott, D.: Continuous lattices. In: Lawvere, F.W. (ed.) Toposes, Algebraic Geometry and Logic, pp. 97–136. Springer, Berlin, Heidelberg (1972)
- [36] Day, A.: Filter monads, continuous lattices and closure systems. Canadian Journal of Mathematics **27**(1), 50–59 (1975)
- [37] Banaschewski, B., Matutu, P.: Remarks on the frame envelope of a σ -frame. Journal of Pure and Applied Algebra **177**, 231–236 (2003)
- [38] Razafindrakoto, A.: Separated and prime compactifications. Topology and its Applications **309**, 107911 (2022)
- [39] Escardó, M.: Properly injective spaces and function spaces. Topology and its Applications **89**(1), 75–120 (1998)
- [40] Escardó, M.: Injective locales over perfect embeddings and algebras of the upper powerlocale monad. Applied General Topology **4**, 193–200 (2002)
- [41] Gleason, A.M.: Projective topological spaces. Illinois Journal of Mathematics **2**, 482–489 (1958)