

CENTRAL LIMIT THEOREM FOR TENSOR PRODUCTS OF FREE VARIABLES

CÉCILIA LANCIEN, PATRICK OLIVEIRA SANTOS, AND PIERRE YOUSSEF

ABSTRACT. We establish a central limit theorem for tensor product random variables $c_k := a_k \otimes a_k$, where $(a_k)_{k \in \mathbb{N}}$ is a free family of variables. We show that if the variables a_k are centered, the limiting law is the semi-circle. Otherwise, the limiting law depends on the mean and variance of the variables a_k and corresponds to a free interpolation between the semi-circle law and the classical convolution of two semi-circle laws.

1. INTRODUCTION

The Free Central Limit Theorem serves as a foundational principle in free probability [15], [11, Lecture 8]. It asserts that as the number of freely independent operators summed together approaches infinity, the distribution of the normalized sum tends towards an asymptotically semi-circular shape. This mirrors the classical Central Limit Theorem but with independence conditions replaced by free independence (also known as freeness) and the Gaussian limit substituted with a semi-circular limit. More precisely, let (\mathcal{A}, τ) be a unital noncommutative probability space equipped with a faithful tracial state τ [11, Lecture 1]. We say that subalgebras $\mathcal{A}_1, \dots, \mathcal{A}_d \subset \mathcal{A}$ are free if

$$\tau(a_1 \dots a_p) = 0,$$

whenever $p \geq 1$, $a_i \in \mathcal{A}_{j_i}$, $\tau(a_i) = 0$ for all $i \in [p]$ and $j_1 \neq j_2 \neq \dots \neq j_p$. We say that random variables $a_1, \dots, a_d \in \mathcal{A}$ are free if their generated algebras are free. We say that a sequence of (self-adjoint) variables $a_n \in (\mathcal{A}_n, \tau_n)$ converges in distribution to a variable $a \in (\mathcal{A}, \tau)$ if

$$\tau_n(a_n^p) \rightarrow \tau(a^p),$$

for all integers $p \geq 0$ and we denote it $a_n \Rightarrow a$. We denote $a - \lambda := a - \lambda \mathbf{1}$, where $\mathbf{1} \in \mathcal{A}$ is the unit in the algebra. As usual, $\tau(a)$ is the mean of a and the variance is given by

$$\text{var}(a) = \tau((a - \tau(a))^2).$$

The Free Central Limit Theorem states that if $a_1, \dots, a_n \in (\mathcal{A}, \tau)$ are free self-adjoint i.i.d random variables with mean λ and variance σ^2 , then

$$\frac{1}{\sigma\sqrt{n}} \sum_{k \in [n]} (a_k - \lambda) \Rightarrow \mu_{sc},$$

where μ_{sc} denotes the semi-circle distribution whose density is given by

$$f_{sc}(x) = \frac{1}{2\pi} \sqrt{4 - x^2} \mathbf{1}_{|x| \leq 2}.$$

The goal of this paper is to establish a central limit theorem for the tensor product of free random variables. Concretely, given $a_1, \dots, a_n \in (\mathcal{A}, \tau)$ free self-adjoint i.i.d random variables, we aim at studying the convergence of the normalized sequence

$$\frac{1}{\sqrt{n}} \sum_{k \in [n]} (a_k \otimes a_k - \tau \otimes \tau(a_k \otimes a_k)), \quad (1)$$

in the product space $(\mathcal{A} \otimes \mathcal{A}, \tau \otimes \tau)$.

Just as free probability captures the limiting behavior of random matrices, the above expression appears naturally as the limiting object corresponding to several models of random quantum channels [7]. Indeed, given $M_1, \dots, M_n \in \mathcal{M}_d(\mathbb{C})$ independent random self-adjoint matrices, it was shown in [7] that the empirical spectral distribution (ESD) of the quantum channel

$$\Delta_{d,n} := \frac{1}{\sqrt{n}} \sum_{k \in [n]} (M_k \otimes M_k - \mathbb{E}[M_k \otimes M_k]),$$

having the M_k 's as random Kraus operators and with fixed Kraus rank n , converges as $d \rightarrow \infty$ to the expression in (1) with the a_k 's being the corresponding limits of the ESD of the M_k 's. Moreover, it was in particular shown that if the random matrices M_k are centered, then the ESD of $\Delta_{d,n}$ converges as $n, d \rightarrow \infty$ to the semi-circle distribution. These two statements combined suggest that, in the case where the a_k 's are centered, an analogue of the free central limit theorem should hold for the $a_k \otimes a_k$'s. Whereas these heuristics indicate that the semi-circle distribution should appear as the limit of (1) when the a_k 's are centered, the convergence and the explicit limit are not clear in the general case. The goal of this paper is to address this by establishing the convergence of the expression in (1) and identifying the limiting object. The latter, as we show, depends on the mean and variance of the variables a_k 's and represents a free interpolation between a semi-circle distribution and the classical convolution of two semi-circle distributions.

Random matrix models of the form

$$M = \sum_{k \in [n]} M_k \otimes M_k, \quad (2)$$

for $M_1, \dots, M_n \in \mathcal{M}_d(\mathbb{C})$ independent random self-adjoint matrices, are in fact useful in other areas of Quantum Information Theory. When the M_k 's are positive semidefinite matrices, normalizing M by its trace produces a model for a random separable quantum state. Little is known about the typical asymptotic spectrum of separable states, contrary to that of entangled ones [1]. Moreover, a random matrix M of the form (2) appears naturally when performing a so-called realignment operation on a quantum state. Understanding the spectrum of the realignment of a state is important as it gives information on the entanglement of the state. In [2], this was done in the particular case where the M_k 's are Gaussian matrices (corresponding to the case where the state is a normalized Wishart matrix). The results and techniques we develop here (combined with those in [7]) could be useful in addressing the questions mentioned above.

Given a measure μ , we denote

$$(t\mu)(A) := \mu(t^{-1}A),$$

its dilation by $t \neq 0$, where A is any Borel set in \mathbb{R} . Equivalently, if a is a random variable with distribution μ , then ta has distribution $t\mu$.

The following is our main theorem.

Theorem 1.1. *Let $a \in (\mathcal{A}, \tau)$ be a self-adjoint random variable with mean $\tau(a) = \lambda$ and variance $\text{var}(a) = \sigma^2 \neq 0$. Denote*

$$\delta^2 := \text{var}(a \otimes a) = \sigma^2(\sigma^2 + 2\lambda^2),$$

and

$$q := \frac{2\lambda^2}{\sigma^2 + 2\lambda^2} \in [0, 1].$$

Given $(a_k)_{k \in \mathbb{N}}$ a sequence of free copies of a , the normalized sum

$$S_n := \frac{1}{\delta \sqrt{n}} \sum_{k \in [n]} (a_k \otimes a_k - \lambda^2)$$

converges in distribution as $n \rightarrow \infty$ to

$$\mu_q := \sqrt{q} \left(\frac{1}{\sqrt{2}} \mu_{sc} + \frac{1}{\sqrt{2}} \mu_{sc} \right) \boxplus \sqrt{1-q} \mu_{sc}, \quad (3)$$

where $+$ denotes the classical convolution and \boxplus denotes the free convolution.

The difficulty in analyzing S_n stems from the complicated dependence structure exhibited by tensors, combining classical independence (between the two legs of the tensor) and freeness (between the variables across tensors). In the centered case, similar computations were made for semi-circle random variables [10, 5]. It would be of interest to design a general notion of independence corresponding to the tensor case, analyze its properties, derive the corresponding limit theorems, and characterize the corresponding universal objects. One particular generalization is by replacing the tensor product with the product of ε -independent random variables [14, 9, 13]. A direct consequence of Theorem 1.1 is that such a notion cannot, in general, reduce to freeness.

Corollary 1.2. *Let $a_1, \dots, a_n \in (\mathcal{A}, \tau)$ be self-adjoint free i.i.d noncentered random variables. Then $\{a_k \otimes a_k : k \in [n]\}$ are not free.*

The above corollary trivially follows from Theorem 1.1, since if the $a_k \otimes a_k$'s were free, the limit of their normalized sum would be the semi-circle distribution contradicting the conclusion of Theorem 1.1 (when $\lambda \neq 0$). This fact was originally proved in [4] where, more generally, the freeness of tensors of free variables was characterized.

This paper is organized as follows. In Section 2, we recall some definitions and notations. In Section 3, we provide some properties of the limiting measure appearing in Theorem 1.1. Section 4 establishes the existence of the limit, while Section 5 is dedicated to the proof of Theorem 1.1.

Acknowledgments. The last named author would like to thank Guillaume Cébron and Roland Speicher for helpful discussions. The second named author also thanks Philippe Biane for helpful discussions. Part of this work was initiated during a stay of the second named author at New York University in Abu Dhabi, partly funded by a doctoral mobility grant delivered by Université Gustave Eiffel; he would like to thank both institutions for their support and the excellent working assumptions. The first named author was supported by the ANR projects ESQuisses (grant number ANR-20-CE47-0014-01), STARS (grant number ANR-20-CE40-0008), and QTraj (grant number ANR-20-CE40-0024-01).

2. PRELIMINARIES AND NOTATIONS

Given $p \in \mathbb{N}$, a partition $\pi = \{V_1, \dots, V_k\}$ of $[p]$ is a collection of disjoint sets V_1, \dots, V_k called blocks such that

$$V_1 \cup \dots \cup V_k = [p].$$

We denote by $P(p)$ the set of partitions of $[p]$. We say that a partition $\pi \in P(p)$ is *connected* (also referred to as a linked diagram in [12]) if no proper subinterval of $[p]$ can be written as the union of blocks of π . A partition $\pi \in P(p)$ has a crossing $i < k < j < l$ if there exist two disjoint blocks $V_1, V_2 \in \pi$ such that $\{i, j\} \subset V_1$ and $\{k, l\} \subset V_2$. In this case, we say that V_1 crosses V_2 . A block $V \in \pi$ is crossing if there exists another $V' \in \pi$ such that V' crosses V , and noncrossing if it does not cross any other block. We say that

a partition $\pi \in P(p)$ is a *noncrossing* partition if all its blocks are noncrossing. We denote by $P^{\text{con}}(p)$ (resp. $NC(p)$) the set of all connected (resp. noncrossing) partitions of $[p]$; see Figure 1. We note that the cardinal $|NC(p)|$ is equal to C_p , the p -th Catalan number.




FIGURE 1. Examples of partitions.

Finally, for a partition $\pi \in P(p)$, we denote by $G(\pi)$ its *intersection graph*. It is the graph over the blocks of π such that two blocks are connected if they cross, under some arbitrary labeling. We say that a partition π is a bipartite partition if its intersection graph is bipartite and denote it $\pi \in P^{\text{bi}}(p)$; see Figure 2 for the partitions in Figure 1.




FIGURE 2. Examples of intersection graphs.

Given $\pi \in P(p)$, we denote $|\pi|$ its number of blocks and $\mathbf{cr}(\pi)$ its number of crossing blocks. Therefore, the number of noncrossing blocks of π is

$$\mathbf{ncr}(\pi) := |\pi| - \mathbf{cr}(\pi).$$

Given $\pi \in P(p)$, we denote $\mathbf{cc}(\pi)$ its number of connected components. We denote $P_2(p)$, $P_2^{\text{bi}}(p)$, $P_2^{\text{con}}(p)$, $P_2^{\text{bicon}}(p)$ and $NC_2(p)$ the set of pair partitions, bipartite pair partitions, connected pair partitions, bipartite connected pair partitions, and noncrossing pair partitions, respectively, that is, those partitions such that all of their blocks have cardinality two.

A pair partition $\pi \in P_2(p)$ can be decomposed into its crossing connected components, namely, let $\hat{\pi} \in P(p)$ be the choice of connected components and, for each block $T \in \hat{\pi}$, draw a connected pair partition $\pi_T \in P_2^{\text{con}}(T)$. By definition, $\hat{\pi} \in NC(p)$ as otherwise two disjoint components would meet ($\hat{\pi}$ is called the noncrossing closure of π in [8]). The mapping

$$\Phi : \pi \mapsto (\hat{\pi}, (\pi_T)_{T \in \hat{\pi}}) \tag{4}$$

is a bijection that will be used throughout the proof of Theorem 1.1; see Figure 3.




FIGURE 3. A partition π and its image $\Phi(\pi)$.

Note, for instance, that $|\hat{\pi}| = \mathbf{cc}(\pi)$. We denote

$$\text{Proj}(\hat{\pi}) := \{((\pi_T)_{T \in \hat{\pi}}) : \pi_T \in P_2^{\text{con}}(T), \forall T \in \hat{\pi}\}.$$

Given $a_1, \dots, a_n \in \mathcal{A}$, we denote by $\kappa_n^{\text{free}}(a_1, \dots, a_n)$ their free cumulants, namely, for any $i \in [n]^k$, we have

$$\begin{aligned} \tau(a_{i_1} \cdots a_{i_k}) &= \sum_{\pi \in NC(k)} \kappa_{\pi}^{\text{free}}(a_{i_1}, \dots, a_{i_k}), \\ \kappa_{\pi}^{\text{free}}(a_{i_1}, \dots, a_{i_k}) &= \prod_{V=\{v_1 < \cdots < v_l\} \in \pi} \kappa_{|V|}^{\text{free}}(a_{i_{v_1}}, \dots, a_{i_{v_l}}). \end{aligned}$$

This is known as the moment-cumulant formula [11, Notation 11.5]. Here and throughout the paper, we denote $V = \{v_1 < \cdots < v_l\}$ a block $V = \{v_1, \dots, v_l\}$ such that $v_1 < \cdots < v_l$; see [11, Lecture 11]. Note also that if the variables a_1, \dots, a_n are free, the free mixed cumulants vanish [11, Proposition 11.15]. We denote $\kappa_n^{\text{free}}(a)$ the free cumulants of a random variable a . The moment-cumulant formula implies the following, which is going to be used extensively in Subsection 5.3; see [11, Lecture 5, Equation 5.6].

Lemma 2.1. *Let a, c_1, c_2 be variables such that a is free from $\{c_1, c_2\}$. Then*

$$\tau(ac_1ac_2) = \text{var}(a)\tau(c_1)\tau(c_2) + \tau^2(a)\tau(c_1c_2).$$

We will equivalently denote $\kappa_n^{\text{free}}(\mu)$ the free cumulants of a random variable a with distribution μ . Given two measures μ_a and μ_b , the free convolution $\mu_a \boxplus \mu_b$ denotes the distribution of $a + b$, where a and b are free random variables with distribution μ_a and μ_b , respectively. The classical convolution $\mu_a + \mu_b$ denotes the distribution of $a + b$, where now a and b are classical independent random variables with distribution μ_a and μ_b , respectively.

3. PROPERTIES OF THE LIMITING MEASURE

In this section, we summarize some of the properties of the measure μ_q appearing in (3). We start by computing the free cumulants and the moments of μ_1 .

Proposition 3.1. *Let*

$$\mu_1 = \frac{1}{\sqrt{2}}\mu_{sc} + \frac{1}{\sqrt{2}}\mu_{sc}.$$

Then the following hold.

(1) *Its odd moments vanish and, for every $p \in \mathbb{N}$, its $2p$ -th moment is given by*

$$\int x^{2p} d\mu_1 = 2^{-p} \sum_{l=0}^p \binom{2p}{2l} C_l C_{p-l} = 2^{-p} \sum_{\pi \in P_2^{\text{bi}}(2p)} 2^{\text{cc}(\pi)},$$

where we recall that C_l denotes the l -th Catalan number.

(2) *Its odd free cumulants vanish, $\kappa_2^{\text{free}}(\mu_1) = 1$ and for any even integer $n \geq 4$, we have*

$$\kappa_n^{\text{free}}(\mu_1) = 2 \left(\frac{1}{2}\right)^{n/2} |P_2^{\text{bicon}}(n)|.$$

Proof. Let x_1, x_2 be two classical i.i.d semi-circle random variables. Then

$$\int x^{2p} d\mu_1 = 2^{-p} \tau((x_1 + x_2)^{2p}).$$

Since the variables are classical independent, and in particular, they commute, we get

$$\tau((x_1 + x_2)^{2p}) = 2^{-p} \sum_{l=0}^{2p} \binom{2p}{l} \tau(x_1^l) \tau(x_2^{2p-l}).$$

The odd moments of x_1 vanish and, for any $l \geq 1$, $\tau(x_1^{2l}) = C_l$, hence

$$\int x^{2p} d\mu_1 = 2^{-p} \sum_{l=0}^p \binom{2p}{2l} C_l C_{p-l}.$$

Now consider the sequence m_p given by

$$m_p := 2^{-p} \sum_{\pi \in P_2^{\text{bi}}(2p)} 2^{\text{cc}(\pi)}.$$

For every bipartite pair partition π , we can decompose it into its bipartite sets. Namely, let $\mathcal{V}_1 = \{V_1, \dots, V_m\}$ and $\mathcal{V}_2 = \{V_{m+1}, \dots, V_p\}$ be the bipartition of its blocks in two disjoint families of vertices. Then, the blocks V_1, \dots, V_m are noncrossing from one another, and so are V_{m+1}, \dots, V_p . Let

$$I = \bigcup_{i=1}^m V_i \subseteq [2p].$$

Then

$$\begin{aligned} \pi_1 &:= \{V_1, \dots, V_m\} \in NC_2(I); \\ \pi_2 &:= \{V_{m+1}, \dots, V_p\} \in NC_2(I^c). \end{aligned}$$

We denote (π_1, π_2, I, I^c) a left-right noncrossing representation of π . We note that if (π_1, π_2, I, I^c) is a left-right noncrossing representation of π , so is (π_2, π_1, I^c, I) . Let $R(\pi)$ be the set of all left-right noncrossing representations of π . We note that $|R(\pi)|$ corresponds to the number of ways the vertices of $G(\pi)$ can be split into two independent families of vertices. Therefore, it is clear that

$$|R(\pi)| = 2^{\text{cc}(\pi)}.$$

Hence

$$\sum_{\pi \in P_2^{\text{bi}}(2p)} 2^{\text{cc}(\pi)} = \sum_{\pi \in P_2^{\text{bi}}(2p)} |R(\pi)|. \quad (5)$$

Since $R(\pi)$ is the set of all left-right noncrossing representations of π and we sum over all $\pi \in P_2^{\text{bi}}(2p)$, we get

$$\sum_{\pi \in P_2^{\text{bi}}(2p)} |R(\pi)| = |\{(\pi_1, \pi_2, I, I^c) : I \subseteq [2p], \pi_1 \in NC_2(I), \pi_2 \in NC_2(I^c)\}|.$$

Note that for fixed $I \subseteq [2p]$, each $(\pi_1, \pi_2) \in NC_2(I) \times NC_2(I^c)$ will correspond to a unique partition $\pi \in P_2^{\text{bi}}(2p)$ such that (π_1, π_2, I, I^c) is a left-right noncrossing representation of π . Hence we can first sum over $I \subseteq [2p]$ so that

$$\sum_{\pi \in P_2^{\text{bi}}(2p)} |R(\pi)| = \sum_{I \subseteq [2p]} |\{(\pi_1, \pi_2) : \pi_1 \in NC_2(I), \pi_2 \in NC_2(I^c)\}|.$$

Therefore, by (5), we have

$$\sum_{\pi \in P_2^{\text{bi}}(2p)} 2^{\text{cc}(\pi)} = \sum_{I \subseteq [2p]} |\{(\pi_1, \pi_2) : \pi_1 \in NC_2(I), \pi_2 \in NC_2(I^c)\}|.$$

The latter cardinality can be written as a product over the cardinals and only depends on the length of I . More precisely, we have

$$\sum_{\pi \in P_2^{\text{bi}}(2p)} 2^{\text{cc}(\pi)} = \sum_{l=0}^p \binom{2p}{2l} C_l C_{p-l},$$

where the cardinal $|NC_2(2l)|$ is equal to C_l . Hence

$$\int x^{2p} d\mu_1 = m_p,$$

and the first statement follows. For the second, we use the bijection Φ in (4) to write $\Phi(\pi) = (\hat{\pi}, (\pi_T)_{T \in \hat{\pi}})$. We note that

$$\begin{aligned} \sum_{\substack{T \in \hat{\pi} \\ |T| \geq 4}} 1 &= \mathbf{cc}(\pi) - \mathbf{ncr}(\pi), \\ \sum_{\substack{T \in \hat{\pi} \\ |T| \geq 4}} \binom{|T|}{2} &= \mathbf{cr}(\pi). \end{aligned}$$

Hence

$$\begin{aligned} \sum_{\pi \in P_2^{\text{bi}}(2p)} 2^{\mathbf{cc}(\pi)-p} &= \sum_{\pi \in P_2^{\text{bi}}(2p)} 2^{\mathbf{cc}(\pi)-\mathbf{ncr}(\pi)} \left(\frac{1}{2}\right)^{\mathbf{cr}(\pi)} \\ &= \sum_{\hat{\pi} \in NC(2p)} \sum_{\substack{T \in \hat{\pi} \\ \pi_T \in P_2^{\text{bicon}}(T)}} \prod_{\substack{T \in \hat{\pi} \\ |T| \geq 4}} 2 \prod_{\substack{T \in \hat{\pi} \\ |T| \geq 4}} \left(\frac{1}{2}\right)^{|T|/2}, \end{aligned}$$

where we use that $\mathbf{ncr}(\pi) + \mathbf{cr}(\pi) = p$ in the first equality. We thus have

$$\sum_{\pi \in P_2^{\text{bi}}(2p)} 2^{\mathbf{cc}(\pi)-p} = \sum_{\hat{\pi} \in NC(2p)} \prod_{\substack{T \in \hat{\pi} \\ |T| \geq 4}} \left\{ 2 \left(\frac{1}{2}\right)^{|T|/2} |P_2^{\text{bicon}}(|T|)| \right\}.$$

Since the free cumulants $\kappa_n^{\text{free}}(\mu_1)$ are uniquely characterized by the moments, it follows that for any odd integer n , we have $\kappa_n^{\text{free}}(\mu_1) = 0$, $\kappa_2^{\text{free}}(\mu_1) = 1$ and for any even integer $n \geq 4$, we have

$$\kappa_n^{\text{free}}(\mu_1) = 2 \left(\frac{1}{2}\right)^{n/2} |P_2^{\text{bicon}}(n)|.$$

□

Remark 3.2. For any $p \geq 1$, it was shown in [6] that

$$\sum_{l=0}^p \binom{2p}{2l} C_l C_{p-l} = C_p C_{p+1}.$$

Therefore, the $2p$ -th moment of μ_1 is also characterized by $2^{-p} C_p C_{p+1}$. The sequence $(C_p C_{p+1})_{p \geq 1}$ is A005568 in Sloane's encyclopedia (<https://oeis.org/A005568>), where several combinatorial objects counted by it are shown.

We are ready to compute the moments and free cumulants of μ_q .

Proposition 3.3. *Given $q \in [0, 1]$, let*

$$\mu_q := \sqrt{q} \left(\frac{1}{\sqrt{2}} \mu_{sc} + \frac{1}{\sqrt{2}} \mu_{sc} \right) \boxplus \sqrt{1-q} \mu_{sc}.$$

Then the following hold.

- (1) *The odd free cumulants of μ_q vanish, $\kappa_2^{\text{free}}(\mu_q) = 1$ and for any even integer $n \geq 4$, we have*

$$\kappa_n^{\text{free}}(\mu_q) = 2 \left(\frac{q}{2}\right)^{n/2} |P_2^{\text{bicon}}(n)|.$$

(2) The odd moments of μ_q vanish and, for every $p \in \mathbb{N}$, its $2p$ -th moment is given by

$$\sum_{\pi \in P_2^{\text{bi}}(2p)} 2^{\mathbf{cc}(\pi) - p} q^{\mathbf{cr}(\pi)}.$$

Proof. We will use the notion of R -transform; see [11, Lecture 16]. For a random variable $a \in \mathcal{A}$ with distribution μ , let

$$R_a(z) = R_\mu(z) = \sum_{n \geq 1} \kappa_n^{\text{free}}(a) z^n,$$

be its R -transform, defined as a formal series. The R -transform of a standard semi-circle law is given by

$$R_{\mu_{sc}}(z) = z^2,$$

whereas Proposition 3.1 shows that the R -transform of μ_1 is given by

$$R_{\mu_1}(z) = z^2 + 2 \sum_{n \geq 2} \left(\frac{1}{2}\right)^n |P_2^{\text{bicon}}(2n)| z^{2n}.$$

Since the free cumulants linearize the free convolution, we deduce that

$$\begin{aligned} R_{\mu_q}(z) &= R_{\sqrt{q}\mu_1}(z) + R_{\sqrt{1-q}\mu_{sc}}(z) \\ &= R_{\mu_1}(\sqrt{q}z) + R_{\mu_{sc}}(\sqrt{1-q}z) \\ &= z^2 + 2 \sum_{n \geq 2} \left(\frac{q}{2}\right)^n |P_2^{\text{bicon}}(2n)| z^{2n}. \end{aligned}$$

This proves the first part of the proposition. To prove the second part, we use the moment-cumulant formula to deduce that the odd moments vanish, while the $2p$ -th moment can be expressed as

$$\begin{aligned} \sum_{\hat{\pi} \in NC(2p)} \prod_{T \in \hat{\pi}} \kappa_{|T|}^{\text{free}}(\mu_q) &= \sum_{\hat{\pi} \in NC(2p)} \prod_{\substack{T \in \hat{\pi} \\ |T| \geq 4}} \left\{ 2 \left(\frac{q}{2}\right)^{|T|/2} |P_2^{\text{bicon}}(T)| \right\} \\ &= \sum_{\hat{\pi} \in NC(2p)} \sum_{(\pi_T)_{T \in \hat{\pi}}} \prod_{\substack{T \in \hat{\pi} \\ |T| \geq 4}} \left\{ 2 \left(\frac{q}{2}\right)^{|T|/2} \right\}, \end{aligned}$$

where the second summation is over $(\pi_T)_{T \in \hat{\pi}} \in \text{Proj}(\hat{\pi})$ such that $\pi_T \in P_2^{\text{bicon}}(T)$ for $T \in \hat{\pi}$. Now noting that

$$|\{T \in \hat{\pi} : |T| \geq 4\}| = \mathbf{cc}(\pi) - \mathbf{ncr}(\pi),$$

and

$$\sum_{\substack{T \in \hat{\pi} \\ |T| \geq 4}} \frac{|T|}{2} = \mathbf{cr}(\pi),$$

we finish the proof after using the bijection Φ from (4) to rewrite the above expression. \square

4. EXISTENCE OF THE LIMIT

The goal of this section is to show that the expression in (1) admits a limit that depends only on the first and second moments of the variables at hand. Let us first prove the following centering lemma.

Lemma 4.1. *Let (\mathcal{A}, τ) be a unital faithful tracial noncommutative probability space. Let $\{a_k : k \in [n]\}, \{d_k : k \in [n]\} \subset \mathcal{A}$ be two collections of free random variables and let*

$$c_k := a_k \otimes d_k - \tau \otimes \tau(a_k \otimes d_k),$$

for every $k \in [n]$. Then for any $m \geq 1$ and $i_1, \dots, i_m \in [n]$ such that there exists an index $l^* \in [m]$ satisfying $i_{l^*} \neq i_j$ for all $j \neq l^*$, we have

$$\tau \otimes \tau(c_{i_1} \cdots c_{i_m}) = 0.$$

In particular, we have

$$\tau \otimes \tau(c_{i_1} \cdots c_{i_{l^*-1}} a_{i_{l^*}} \otimes d_{i_{l^*}} c_{i_{l^*+1}} \cdots c_{i_m}) = \tau(a_{i_{l^*}}) \tau(d_{i_{l^*}}) \tau \otimes \tau(c_{i_1} \cdots c_{i_{l^*-1}} c_{i_{l^*+1}} \cdots c_{i_m}).$$

Proof. By cyclicity of the trace, we can assume that $l^* = 1$. We write

$$\tau \otimes \tau(c_{i_1} \cdots c_{i_m}) = \tau \otimes \tau(a_{i_1} \otimes d_{i_1} c_{i_2} \cdots c_{i_m}) - \tau \otimes \tau(a_{i_1} \otimes d_{i_1}) \tau \otimes \tau(c_{i_2} \cdots c_{i_m}). \quad (6)$$

For the first term, we open the expressions for c_{i_j} for $j \geq 2$ to get that

$$\begin{aligned} & \tau \otimes \tau(a_{i_1} \otimes d_{i_1} c_{i_2} \cdots c_{i_m}) \\ &= \sum_{I \subseteq \{2, \dots, m\}} (-1)^{|I|} \prod_{j \in I} (\tau \otimes \tau(a_{i_j} \otimes d_{i_j})) \tau \otimes \tau \left(a_{i_1} \otimes d_{i_1} \prod_{j \in J^c}^{\rightarrow} a_{i_j} \otimes d_{i_j} \right), \end{aligned}$$

where \prod^{\rightarrow} denotes the product respecting the ordering. Since a_{i_1}, d_{i_1} are free from a_{i_j}, d_{i_j} for all $j \geq 2$, freeness implies that

$$\begin{aligned} & \tau \otimes \tau(a_{i_1} \otimes d_{i_1} c_{i_2} \cdots c_{i_m}) \\ &= \tau \otimes \tau(a_{i_1} \otimes d_{i_1}) \sum_{I \subseteq \{2, \dots, m\}} (-1)^{|I|} \prod_{j \in I} (\tau \otimes \tau(a_{i_j} \otimes d_{i_j})) \tau \otimes \tau \left(\prod_{j \in J^c}^{\rightarrow} a_{i_j} \otimes d_{i_j} \right). \end{aligned}$$

The summation can be easily written as $\tau \otimes \tau(c_{i_2} \cdots c_{i_m})$, hence

$$\tau \otimes \tau(a_{i_1} \otimes d_{i_1} c_{i_2} \cdots c_{i_m}) = \tau \otimes \tau(a_{i_1} \otimes d_{i_1}) \tau \otimes \tau(c_{i_2} \cdots c_{i_m}).$$

By (6), we then have

$$\tau \otimes \tau(c_{i_1} \cdots c_{i_m}) = 0.$$

Finally, since $\tau \otimes \tau(a_{i_1} \otimes d_{i_1}) = \tau(a_{i_1}) \tau(d_{i_1})$, we get that

$$\begin{aligned} \tau \otimes \tau(a_{i_1} \otimes d_{i_1} c_{i_2} \cdots c_{i_m}) &= \tau \otimes \tau(a_{i_1} \otimes d_{i_1}) \tau \otimes \tau(c_{i_2} \cdots c_{i_m}) \\ &= \tau(a_{i_1}) \tau(d_{i_1}) \tau \otimes \tau(c_{i_2} \cdots c_{i_m}). \end{aligned}$$

□

We are now ready to prove the existence of the limit and that it only depends on the first and second moments of the a_k 's.

Proposition 4.2 (Existence). *Let $(a_n)_{n \in \mathbb{N}} \in (\mathcal{A}, \tau)$ be self-adjoint free i.i.d random variables with mean λ , variance σ^2 , and denote $\delta^2 := \text{var}(a_1 \otimes a_1) = \sigma^2(\sigma^2 + 2\lambda^2)$. For every $k \in \mathbb{N}$, denote*

$$b_k := \frac{1}{\delta} (a_k \otimes a_k - \tau \otimes \tau(a_k \otimes a_k)),$$

and for every $n \in \mathbb{N}$, denote

$$S_n := \frac{1}{\sqrt{n}} \sum_{k \in [n]} b_k.$$

Then there exists a random variable $\mathbf{S} \in (\mathcal{A}', \tau')$ such that $S_n \Rightarrow \mathbf{S}$. Moreover, the law of \mathbf{S} depends only on λ and σ , its odd moments vanish and, for every $p \in \mathbb{N}$,

$$\tau'(\mathbf{S}^{2p}) = \sum_{\pi \in P_2(2p)} \tau \otimes \tau(b_{i_1} \cdots b_{i_{2p}}),$$

where $i \in [p]^{2p}$ is any sequence such that $i_j = i_k$ if and only if $\{j, k\} \in \pi$.

Proof. We begin by writing

$$\tau \otimes \tau(S_n^p) = \frac{1}{n^{p/2}} \sum_{i \in [n]^p} \tau \otimes \tau(b_{i_1} \cdots b_{i_p}).$$

Since b_1, \dots, b_n are identically distributed, the expression

$$\tau \otimes \tau(b_{i_1} \cdots b_{i_p}) \tag{7}$$

depends only on the partition $\pi = \pi(i) \in P(p)$ given by $l \sim_\pi k$ (that is, l, k belong to the same block of π) if and only if $i_l = i_k$. Denote the common value of (7) by $\tau \otimes \tau(\pi)$. Then we have

$$\tau \otimes \tau(S_n^p) = \frac{1}{n^{p/2}} \sum_{\pi \in P(p)} \tau \otimes \tau(\pi) |\{i \in [n]^p : \pi(i) = \pi\}|.$$

To count the cardinality, we choose an index for each block. Therefore, we have

$$|\{i \in [n]^p : \pi(i) = \pi\}| = n(n-1) \cdots (n - |\pi| + 1) \sim n^{|\pi|}.$$

By Lemma 4.1, if π has a block of size 1, $\tau \otimes \tau(\pi) = 0$. Thus, we have

$$\tau \otimes \tau(S_n^p) = \sum_{\substack{\pi \in P(p) \\ |V| \geq 2; \forall V \in \pi}} \tau \otimes \tau(\pi) \frac{n(n-1) \cdots (n - |\pi| + 1)}{n^{p/2}}.$$

Since $|V| \geq 2$ for all blocks $V \in \pi$, we have $|\pi| \leq p/2$. If there exists a block $V \in \pi$ such that $|V| \geq 3$, we immediately have $|\pi| < p/2$, and its contribution is negligible. In particular, this implies that the odd moments of S_n are asymptotically vanishing. We deduce that

$$\lim_{n \rightarrow \infty} \tau \otimes \tau(S_n^p) = \lim_{n \rightarrow \infty} \sum_{\substack{\pi \in P(p) \\ |V|=2; \forall V \in \pi}} \tau \otimes \tau(\pi) \frac{n(n-1) \cdots (n - |\pi| + 1)}{n^{p/2}}.$$

In this case, π is a pair partition and $|\pi| = p/2$, hence we deduce the formula

$$\lim_{n \rightarrow \infty} \tau \otimes \tau(S_n^p) = \sum_{\pi \in P_2(p)} \tau \otimes \tau(\pi),$$

which shows that S_n converges. To prove that the limit depends only on λ and σ , we write

$$\tau \otimes \tau(\pi) = \tau \otimes \tau(b_{i_1} \cdots b_{i_p}) = \frac{1}{\delta^p} \sum_{I \subseteq [p]} (-1)^{|I|} \lambda^{2|I|} \tau^2 \left(\prod_{l \in I^c}^{\rightarrow} a_{i_l} \right),$$

where $\pi(i) = \pi$. By the moment-cumulant formula, we have

$$\tau \left(\prod_{l \in I^c}^{\rightarrow} a_{i_l} \right) = \sum_{\sigma \in NC(I^c)} \kappa_{\sigma}((a_{i_l})_{l \in I^c}).$$

Since the mixed cumulants of free variables vanish, the only partitions $\sigma \in NC(I^c)$ that contribute are those such that every block $V \in \sigma$ has cardinality at most two. We then have

$$\kappa_{\sigma}((a_{i_l})_{l \in I^c}) = \prod_{\substack{V \in \sigma \\ |V|=2}} \kappa_{|V|}(a, a) \prod_{\substack{V \in \sigma \\ |V|=1}} \kappa_{|V|}(a) = \sigma^{2\#\{V \in \sigma: |V|=2\}} \lambda^{\#\{V \in \sigma: |V|=1\}}.$$

This concludes the proof. \square

5. PROOF OF THEOREM 1.1

After proving the existence of the limit in the previous section, the goal here is to identify this limit as stated in Theorem 1.1. In all this section, $(a_n)_{n \in \mathbb{N}}$ denote free copies of a random variable a with mean λ and variance σ^2 . Moreover, the common law of the normalized tensors will be denoted by

$$b = \frac{1}{\delta} (a \otimes a - \lambda^2), \quad (8)$$

where $\delta^2 = \text{var}(a \otimes a) = \sigma^2(\sigma^2 + 2\lambda^2)$.

Throughout the proof, we will assume p is an even integer. Following Proposition 4.2, we denote \mathbf{S} the limit of S_n and note that

$$\tau'(\mathbf{S}^p) = \sum_{\pi \in P_2(p)} \tau \otimes \tau(\pi),$$

where $\tau \otimes \tau(\pi) = \tau \otimes \tau(b_{i_1} \cdots b_{i_p})$ with $\pi(i) = \pi$.

5.1. Contribution of noncrossing blocks. We begin by removing interval blocks.

Lemma 5.1. *Let $\pi \in P_2(p)$ and suppose that there exists $l \in [p]$ such that $\{l, l+1\} \in \pi$ (with the convention that $p+1 := 1$). Then*

$$\tau \otimes \tau(\pi) = \tau \otimes \tau(\pi \setminus \{l, l+1\}).$$

Proof. By cyclicity, we can assume $l = p-1$. We then have

$$\begin{aligned} \delta^2 \tau \otimes \tau(\pi) &= \tau \otimes \tau \left(b_{i_1} \cdots b_{i_{p-2}} \cdot \left(a_{i_p}^2 \otimes a_{i_p}^2 + \lambda^4 1 \otimes 1 - 2\lambda^2 a_{i_p} \otimes a_{i_p} \right) \right) \\ &= \tau \otimes \tau \left(b_{i_1} \cdots b_{i_{p-2}} \cdot a_{i_p}^2 \otimes a_{i_p}^2 \right) + \lambda^4 \tau \otimes \tau (b_{i_1} \cdots b_{i_{p-2}}) \\ &\quad - 2\lambda^2 \tau \otimes \tau (b_{i_1} \cdots b_{i_{p-2}} a_{i_p} \otimes a_{i_p}) \\ &=: I + II + III. \end{aligned}$$

We immediately recognize

$$II = \lambda^4 \tau \otimes \tau(\pi \setminus \{p-1, p\}).$$

By Lemma 4.1, we have

$$\begin{aligned} I &= \tau^2(a^2) \tau \otimes \tau(\pi \setminus \{p-1, p\}); \\ III &= -2\lambda^4 \tau \otimes \tau(\pi \setminus \{p-1, p\}). \end{aligned}$$

Hence

$$\delta^2 \tau \otimes \tau(\pi) = (\tau^2(a^2) - \lambda^4) \tau \otimes \tau(\pi \setminus \{p-1, p\}).$$

To conclude, we note that $\tau^2(a^2) - \lambda^4 = \delta^2$ and finish the proof. \square

Recall that a noncrossing pair partition $\pi \in NC_2(p)$ always has an interval block $V = \{l, l+1\} \in \pi$ such that $\pi \setminus V$ is a noncrossing pair partition. In particular, by induction, Lemma 5.1 implies the following.

Corollary 5.2. *For any $\pi \in NC_2(p)$, we have $\tau \otimes \tau(\pi) = 1$.*

5.2. Decomposition of pair partitions. In order to capture the contribution of crossing partitions, we need to decompose a partition $\pi \in P_2(p) \setminus NC_2(p)$ using smaller partitions. We denote $\pi = \pi_1 \oplus \cdots \oplus \pi_k$ if $[p]$ can be decomposed into k intervals I_1, \dots, I_k such that $\pi_k \in P_2(I_k)$ and $V \in \pi$ if $V \in \pi_l$ for some $1 \leq l \leq k$. For $I \subseteq [p]$, let

$$a_I := \prod_{l \in I}^{\rightarrow} a_{i_l}, \quad (9)$$

and $a_\emptyset := 1$.

Lemma 5.3. *Let $\pi \in P_2(p)$. Then, the following hold.*

(5.3.i) *If $\pi = \pi_1 \oplus \cdots \oplus \pi_l$, then*

$$\tau \otimes \tau(\pi) = \tau \otimes \tau(\pi_1) \cdots \tau \otimes \tau(\pi_l).$$

(5.3.ii) *If $\pi = \{1, p\} \cup \pi_1$, where $\pi_1 \in P_2(\{2, \dots, p-1\})$, then*

$$\tau \otimes \tau(\pi) = \tau \otimes \tau(\pi_1).$$

Moreover, if there exists an interval $I \subseteq [p]$ such that $\pi|_I$ is a pair partition, then

$$\tau \otimes \tau(\pi) = \tau \otimes \tau(\pi|_I) \tau \otimes \tau(\pi|_{I^c}).$$

(5.3.iii) *If π has a block $V = \{r, s\}$ such that for any block $U = \{l, k\} \in \pi$ with $r < l < s$, we have $r < k < s$ (i.e., every point inside V matches another one inside V), we have*

$$\tau \otimes \tau(\pi) = \tau \otimes \tau(\pi|_{V_-}) \tau \otimes \tau(\pi|_{V_+}),$$

where $\pi|_{V_-}$ is the restriction of π to inside of V and $\pi|_{V_+}$ is the restriction of π to outside of V .

Proof. (5.3.i) By induction, it suffices to prove the case $\pi = \pi_1 \oplus \pi_2$. Let I_1, I_2 be the disjoint decomposition of $[p]$ given by π_1 and π_2 . Then, we can write

$$\tau \otimes \tau(\pi) = \frac{1}{\delta^p} \sum_{\substack{J_1 \subseteq I_1 \\ J_2 \subseteq I_2}} (-1)^{|J_1|+|J_2|} \lambda^{2(|J_1|+|J_2|)} \tau^2 \left(\prod_{l \in J_1^c} a_{i_l} \prod_{l \in J_2^c} a_{i_l} \right).$$

Since π is the direct sum of π_1, π_2 , the variables $a_{J_1^c}, a_{J_2^c}$ are free and we can write

$$\tau \otimes \tau(\pi) = \frac{1}{\delta^p} \sum_{\substack{J_1 \subseteq I_1 \\ J_2 \subseteq I_2}} (-1)^{|J_1|+|J_2|} \lambda^{2(|J_1|+|J_2|)} \tau^2 \left(\prod_{l \in J_1^c} a_{i_l} \right) \tau^2 \left(\prod_{l \in J_2^c} a_{i_l} \right).$$

It is immediate to check that the right-hand-side is equal to $\tau \otimes \tau(\pi_1) \tau \otimes \tau(\pi_2)$.

(5.3.ii) By cyclicity, we have

$$\tau \otimes \tau(\pi) = \tau \otimes \tau(b_{i_2} \cdots b_{i_{p-1}} b_{i_p} b_{i_1}).$$

Since $\{1, p\} \in \pi$, Lemma 5.1 implies that

$$\tau \otimes \tau(\pi) = \tau \otimes \tau(b_{i_2} \cdots b_{i_{p-1}}) = \tau \otimes \tau(\pi_1).$$

The second part follows again by cyclicity as we can assume $I = \{1, \dots, k\}$ for some $k \in [p]$, and the variables are free.

(5.3.iii) By cyclicity, we can assume that $V = \{1, k\}$ for some $k \in [p]$. Note that V creates a direct sum $\pi = (V \cup \pi|_{V_-}) \oplus \pi|_{V_+}$. The result follows by (5.3.i) and (5.3.ii). \square

Note that any block $V \in \pi$ that does not cross any other block of π is either an interval block or a block that satisfies (5.3.iii). In particular, its removal does not affect the value of $\tau \otimes \tau(\pi)$. It is clear then that $\tau \otimes \tau(\pi)$ is a multiplicative function [3] over the connected components of π . Using the mapping Φ defined in (4), we can write

$$\tau \otimes \tau(\pi) = \prod_{T \in \hat{\pi}} \tau \otimes \tau(\pi_T), \quad (10)$$

where $(\hat{\pi}, (\pi_T)_{T \in \hat{\pi}}) = \Phi(\pi)$. In view of this, we will now focus on the case where $\pi \in P_2^{\text{con}}(p)$, for $p \geq 4$, as the case $p = 2$ corresponds to noncrossing blocks.

5.3. Contribution of connected partitions. Given $\pi \in P_2^{\text{con}}(p)$, for an even integer $p \geq 4$, we recall that $\pi \in P_2^{\text{bicon}}(p)$ if its intersection graph $G(\pi)$ is a bipartite connected graph.

The following is the main proposition of this subsection.

Proposition 5.4. *Let $p \geq 4$ be an even integer and $\pi \in P_2^{\text{con}}(p)$. Then the following hold.*

- (1) *If $\pi \notin P_2^{\text{bicon}}(p)$, then $\tau \otimes \tau(\pi) = 0$;*
- (2) *If $\pi \in P_2^{\text{bicon}}(p)$, then*

$$\tau \otimes \tau(\pi) = 2 \left(\frac{q}{2} \right)^{\frac{p}{2}},$$

$$\text{where } q = \frac{2\lambda^2}{\sigma^2 + 2\lambda^2}.$$

To prove Proposition 5.4, our goal is to remove the blocks $V \in \pi$ one at a time and track its influence on the rest of the partition $\pi \setminus V$. To this end, we will define two main quantities associated with V . First, we will define a coloring $w_V \in \{0, 1\}$ of V that encodes how the removal of V affects the other blocks. Secondly, we will define a binary vector $\theta_V \in \{0, 1\}^4$ satisfying

$$\sum_{j \in [4]} \theta_j = 1,$$

that is, only one coordinate of θ_V is equal to 1. It encodes how we remove the block V from π , that is, its weight to $\tau \otimes \tau(\pi)$. Let us begin this description now.

Let $\pi \in P_2^{\text{con}}(p)$ and $i \in [p]^p$ such that $\pi(i) = \pi$. Given an integer $t \geq 0$, blocks $V_1, \dots, V_t, V_{t+1} \in \pi$, and a coloring $w \in \{0, 1\}^{t+1}$, we denote $V_j^{(w_j)}$ the block V_j under the color w_j . We then define the iterated joint law of b_{i_j}

$$B_k := B_k \left(V_1^{(w_1)}, \dots, V_k^{(w_k)} \right) = (b_{i_j, k})_{j \in [p]}$$

for $k \leq t + 1$ as follows. Let $(\tilde{a}_j)_{j \in \mathbb{N}}$ be a family of free independent copies of a , free from $(a_j)_{j \in \mathbb{N}}$. For $k = 0$, we define $B_0 = (b_{i_j})_{j \in [p]}$ and for any $1 \leq k \leq t + 1$, we have

$$B_k = \frac{1}{\delta} \left(a_{i_j}^{(l)} \otimes a_{i_j}^{(r)} - \lambda^2 \right)_{j \in [p]},$$

where

$$a_{i_j}^{(l)}, a_{i_j}^{(r)} \in \{a_{i_j}, \tilde{a}_{i_j}\}, \quad (11)$$

for any $j \in [p]$. We then recursively define the choices in (11) as follows. Given

$$B_k = \frac{1}{\delta} \left(a_{i_j}^{(l)} \otimes a_{i_j}^{(r)} - \lambda^2 \right)_{j \in [p]},$$

we consider the block $V_{k+1} = \{s_1 < s_2\}$ and its color w_{k+1} , for $1 \leq k \leq t$. If $w_{k+1} = 0$, for all blocks $V = \{t_1, t_2\} \in \pi$ that cross V_{k+1} , we replace the left legs of $(b_{i_{t_1}, k}, b_{i_{t_2}, k})$ by a pair of free variables. Concretely, we define

$$\begin{aligned} b_{i_{t_1}, k+1} &= \frac{1}{\delta} \left(\tilde{a}_{i_{t_1}} \otimes a_{i_{t_1}}^{(r)} - \lambda^2 \right); \\ b_{i_{t_2}, k+1} &= \frac{1}{\delta} \left(a_{i_{t_1}} \otimes a_{i_{t_2}}^{(r)} - \lambda^2 \right). \end{aligned}$$

Otherwise, if $w_{k+1} = 1$, we replace the right legs of $(b_{i_{t_1}, k}, b_{i_{t_2}, k})$ by a pair of free variables instead,

$$\begin{aligned} b_{i_{t_1}, k+1} &= \frac{1}{\delta} \left(a_{i_{t_1}}^{(l)} \otimes \tilde{a}_{i_{t_1}} - \lambda^2 \right); \\ b_{i_{t_2}, k+1} &= \frac{1}{\delta} \left(a_{i_{t_2}}^{(l)} \otimes a_{i_{t_1}} - \lambda^2 \right). \end{aligned}$$

All the other blocks remain unchanged. Recall that for a vector B and a set $I \subseteq [p]$, we set as in (9)

$$B_I = \prod_{l \in I}^{\rightarrow} B_{i_l},$$

and $B_{\emptyset} = 1$. We then define

$$\begin{aligned} \tau \otimes \tau \left(\pi, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right) &:= \tau \otimes \tau \left((B_k)_{[p]} \right) \\ \tau \otimes \tau \left(\pi \setminus \{U_1, \dots, U_m\}, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right) &:= \tau \otimes \tau \left((B_k)_{[p] \setminus (U_1 \cup \dots \cup U_m)} \right), \end{aligned}$$

where $U_1, \dots, U_m, V_1, \dots, V_k \in \pi$ and

$$B_k = B_k \left(V_1^{(w_1)}, \dots, V_k^{(w_k)} \right).$$

Note first that at each step k , the distribution of $(b_{i_{v_1}, k}, b_{i_{v_2}, k})$ for a block $V = \{v_1 < v_2\} \in \pi$ can be written as one of the following four cases.

(1) The blocks V_1, \dots, V_k do not cross V , and we have

$$\begin{aligned} b_{i_{v_1}, k} &= \frac{1}{\delta} \left(a_{i_{v_1}} \otimes a_{i_{v_1}} - \lambda^2 \right); \\ b_{i_{v_2}, k} &= \frac{1}{\delta} \left(a_{i_{v_1}} \otimes a_{i_{v_1}} - \lambda^2 \right), \end{aligned}$$

where we recall that $i_{v_1} = i_{v_2}$.

(2) All blocks V_j that cross V have the same color $w_j = 0$, and we have

$$\begin{aligned} b_{i_{v_1}, k} &= \frac{1}{\delta} \left(\tilde{a}_{i_{v_1}} \otimes a_{i_{v_1}} - \lambda^2 \right); \\ b_{i_{v_2}, k} &= \frac{1}{\delta} \left(a_{i_{v_1}} \otimes a_{i_{v_1}} - \lambda^2 \right). \end{aligned}$$

(3) All blocks V_j that cross V have the same color $w_j = 1$, and we have

$$\begin{aligned} b_{i_{v_1}, k} &= \frac{1}{\delta} \left(a_{i_{v_1}} \otimes \tilde{a}_{i_{v_1}} - \lambda^2 \right); \\ b_{i_{v_2}, k} &= \frac{1}{\delta} \left(a_{i_{v_1}} \otimes a_{i_{v_1}} - \lambda^2 \right). \end{aligned}$$

(4) Otherwise, there exist two blocks V_{j_1}, V_{j_2} that cross V such that $w_{j_1} \neq w_{j_2}$, that is, they have different colors, and then

$$\begin{aligned} b_{i_{v_1}, k} &= \frac{1}{\delta} (\tilde{a}_{i_{v_1}} \otimes \tilde{a}_{i_{v_1}} - \lambda^2); \\ b_{i_{v_2}, k} &= \frac{1}{\delta} (a_{i_{v_1}} \otimes a_{i_{v_1}} - \lambda^2). \end{aligned}$$

We remark that even if the choices of replacements do not always happen for $b_{i_{v_1}}$, the joint distribution of $(b_{i_{v_1}, k}, b_{i_{v_2}, k})$ can always be written as one of those four cases since the variables are free i.i.d. For instance, we have

$$(a \otimes a, a \otimes \tilde{a}) \stackrel{d}{=} (a \otimes \tilde{a}, a \otimes a) \stackrel{d}{=} (\tilde{a} \otimes a, \tilde{a} \otimes \tilde{a}),$$

and so on. Let

$$\begin{aligned} \theta &\in \{0, 1\}^4, \\ \sum_{j \in [4]} \theta_j &= 1, \end{aligned}$$

and denote

$$\begin{aligned} b_{i_{v_1}}(\theta) &= \frac{1}{\delta} (\theta_1 a_{i_{v_1}} \otimes a_{i_{v_1}} + \theta_2 \tilde{a}_{i_{v_1}} \otimes a_{i_{v_1}} + \theta_3 a_{i_{v_1}} \otimes \tilde{a}_{i_{v_1}} + \theta_4 \tilde{a}_{i_{v_1}} \otimes \tilde{a}_{i_{v_1}} - \lambda^2); \\ b_{i_{v_2}}(\theta) &= \frac{1}{\delta} (a_{i_{v_1}} \otimes a_{i_{v_1}} - \lambda^2). \end{aligned}$$

The choice of $\theta := \theta(V, V_1^{(w_1)}, \dots, V_k^{(w_k)})$ corresponds to the case in which we are, that is, we define θ as the binary vector such that

$$(b_{i_{v_1}}(\theta), b_{i_{v_2}}(\theta)) \stackrel{d}{=} (b_{i_{v_1}, k}, b_{i_{v_2}, k}).$$

Let $V = \{v_1 < v_2\} \in \pi \setminus \{V_1, \dots, V_k\}$ and

$$\begin{aligned} I_1 &:= \{1, \dots, v_1 - 1\} \setminus \left(\bigcup_{j \in [k]} V_j \right); \\ I_2 &:= \{v_1 + 1, \dots, v_2 - 1\} \setminus \left(\bigcup_{j \in [k]} V_j \right); \\ I_3 &:= \{v_2 + 1, \dots, p\} \setminus \left(\bigcup_{j \in [k]} V_j \right). \end{aligned}$$

Our first goal is to obtain the contribution and coloring of V to

$$\tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k\}, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right) = \tau \otimes \tau \left((B_k)_{I_1} b_{i_{v_1}, k} (B_k)_{I_2} b_{i_{v_2}, k} (B_k)_{I_3} \right),$$

according to the value of $\theta = \theta(V, V_1^{(w_1)}, \dots, V_k^{(w_k)}) \in \{0, 1\}^4$, where

$$B_k = B_k(V_1^{(w_1)}, \dots, V_k^{(w_k)}).$$

We recall that

$$\begin{aligned} \tau \otimes \tau \left((B_k)_{I_1} (B_k)_{I_2} (B_k)_{I_3} \right) &= \tau \otimes \tau \left((B_k)_{[p] \setminus (V_1 \cup \dots \cup V_k \cup V)} \right) \\ &= \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right) \end{aligned}$$

To simplify the notation, we denote

$$h := |I_1| + |I_2| + |I_3| = p - 2(k + 1). \quad (12)$$

We divide the four cases into the next four lemmas.

Lemma 5.5 (Case $\theta_1 = 1$). *Let $p \geq 4$ be an even integer, $\pi \in P_2^{\text{con}}(p)$. Let $V_1, \dots, V_k \in \pi$ be blocks, $w \in \{0, 1\}^k$ be a coloring of V_1, \dots, V_k and*

$$B_k = B_k \left(V_1^{(w_1)}, \dots, V_k^{(w_k)} \right) = (b_{i_j, k})_{j \in [p]}.$$

Let $V = \{v_1 < v_2\} \in \pi \setminus \{V_1, \dots, V_k\}$ and the joint law

$$(b_{i_{v_1}, k}, b_{i_{v_2}, k}) \stackrel{d}{=} (b_{i_{v_1}}(\theta), b_{i_{v_2}}(\theta)).$$

If $\theta_1 = 1$, we have

$$\begin{aligned} & \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k\}, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right) \\ &= \frac{q}{2} \sum_{w=0,1} \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)}, V^{(w)} \right). \end{aligned}$$

Proof. To simplify the notation, let

$$T := \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k\}, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right).$$

Then, we must prove that

$$T = \frac{q}{2} \sum_{w=0,1} \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)}, V^{(w)} \right).$$

As $\theta_1 = 1$, we can write

$$\begin{aligned} b_{i_{v_1}, k} &= \frac{1}{\delta} (a \otimes a - \lambda^2); \\ b_{i_{v_2}, k} &= \frac{1}{\delta} (a \otimes a - \lambda^2), \end{aligned}$$

where $a := a_{i_{v_1}}$. We get by Lemma 4.1 that

$$\begin{aligned} T &= \frac{1}{\delta^2} \tau \otimes \tau \left\{ (B_k)_{I_1} a \otimes a (B_k)_{I_2} a \otimes a (B_k)_{I_3} \right\} \\ &\quad - \frac{\lambda^4}{\delta^2} \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right). \end{aligned} \quad (13)$$

Let us write

$$B_k = \frac{1}{\delta} \left(a_{i_j}^{(l)} \otimes a_{i_j}^{(r)} - \lambda^2 \right)_{j \in [p]}.$$

Then

$$\begin{aligned} & \tau \otimes \tau \left\{ (B_k)_{I_1} a \otimes a (B_k)_{I_2} a \otimes a (B_k)_{I_3} \right\} \\ &= \frac{1}{\delta^h} \sum_{\substack{J_1 \subseteq I_1 \\ J_2 \subseteq I_2 \\ J_3 \subseteq I_3}} (-1)^{|J_1|+|J_2|+|J_3|} \lambda^{2(|J_1|+|J_2|+|J_3|)} \tau \left(a_{J_1^c}^{(l)} a_{J_2^c}^{(l)} a_{J_3^c}^{(l)} \right) \tau \left(a_{J_1^c}^{(r)} a_{J_2^c}^{(r)} a_{J_3^c}^{(r)} \right), \end{aligned}$$

where h is defined as in (12). We use Lemma 2.1 and compute

$$\begin{aligned} R(J_1, J_2, J_3) &:= \tau \left(a_{J_1^c}^{(l)} a_{J_2^c}^{(l)} a_{J_3^c}^{(l)} \right) \tau \left(a_{J_1^c}^{(r)} a_{J_2^c}^{(r)} a_{J_3^c}^{(r)} \right) \\ &= \left(\sigma^2 \tau(a_{J_1^c}^{(l)} a_{J_2^c}^{(l)}) \tau(a_{J_3^c}^{(l)}) + \lambda^2 \tau(a_{J_1^c}^{(l)} a_{J_2^c}^{(l)} a_{J_3^c}^{(l)}) \right) \left(\sigma^2 \tau(a_{J_1^c}^{(r)} a_{J_2^c}^{(r)}) \tau(a_{J_3^c}^{(r)}) + \lambda^2 \tau(a_{J_1^c}^{(r)} a_{J_2^c}^{(r)} a_{J_3^c}^{(r)}) \right). \end{aligned}$$

After the expansion, we have

$$R(J_1, J_2, J_3) =: \sigma^4 R_1(J_1, J_2, J_3) + \sigma^2 \lambda^2 (R_2(J_1, J_2, J_3) + R_3(J_1, J_2, J_3)) + \lambda^4 R_4(J_1, J_2, J_3),$$

where we choose R_2 to be the term with factor $\tau(a_{J_1^c}^{(l)} a_{J_2^c}^{(l)} a_{J_3^c}^{(l)})$. For R_1 , we have

$$\begin{aligned} &\frac{1}{\delta^h} \sum_{\substack{J_1 \subseteq I_1 \\ J_2 \subseteq I_2 \\ J_3 \subseteq I_3}} (-1)^{|J_1|+|J_2|+|J_3|} \lambda^{2(|J_1|+|J_2|+|J_3|)} R_1(J_1, J_2, J_3) \\ &= \tau \otimes \tau \left((B_k)_{I_1} (B_k)_{I_3} \right) \tau \otimes \tau \left((B_k)_{I_2} \right). \end{aligned}$$

Since π is connected, V crosses at least one block $U = \{j, j'\} \in \pi$. Therefore, we assume $j \in I_2$ whose matching symbol $j' \in I_1 \cup I_3$. By Lemma 4.1, we have

$$\tau \otimes \tau \left((B_k)_{I_2} \right) = 0,$$

and the contribution of R_1 is zero. For R_4 , we have

$$\begin{aligned} &\frac{1}{\delta^h} \sum_{\substack{J_1 \subseteq I_1 \\ J_2 \subseteq I_2 \\ J_3 \subseteq I_3}} (-1)^{|J_1|+|J_2|+|J_3|} \lambda^{2(|J_1|+|J_2|+|J_3|)} R_4(J_1, J_2, J_3) \\ &= \tau \otimes \tau \left((B_k)_{I_1} (B_k)_{I_2} (B_k)_{I_3} \right). \end{aligned}$$

Since

$$\tau \otimes \tau \left((B_k)_{I_1} (B_k)_{I_2} (B_k)_{I_3} \right) = \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right),$$

its contribution cancels out with $\frac{\lambda^4}{\delta^2} \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right)$ in (13).

For R_2 , let us first denote

$$B_{k+1} = B_{k+1} \left(V_1^{(w_1)}, \dots, V_k^{(w_k)}, V^{(1)} \right) =: \frac{1}{\delta} \left(\bar{a}_{i_j}^{(l)} \otimes \bar{a}_{i_j}^{(r)} - \lambda^2 \right)_{j \in [p]}.$$

We then have

$$\begin{aligned} &\tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)}, V^{(1)} \right) \\ &= \frac{1}{\delta^h} \sum_{\substack{J_1 \subseteq I_1 \\ J_2 \subseteq I_2 \\ J_3 \subseteq I_3}} (-1)^{|J_1|+|J_2|+|J_3|} \lambda^{2(|J_1|+|J_2|+|J_3|)} \tau \left(\bar{a}_{J_1^c}^{(l)} \bar{a}_{J_2^c}^{(l)} \bar{a}_{J_3^c}^{(l)} \right) \tau \left(\bar{a}_{J_1^c}^{(r)} \bar{a}_{J_2^c}^{(r)} \bar{a}_{J_3^c}^{(r)} \right). \end{aligned}$$

By definition, the color $w_{k+1} = 1$ does not affect the left legs, hence

$$\bar{a}_{i_j}^{(l)} = a_{i_j}^{(l)}.$$

For the right legs and a block $U = \{t_1 < t_2\}$, let us divide it into two cases.

(1) If U does not cross V , we have

$$\begin{aligned} \bar{a}_{i_{t_1}}^{(r)} &= a_{i_{t_1}}^{(r)}; \\ \bar{a}_{i_{t_2}}^{(r)} &= a_{i_{t_2}}^{(r)}. \end{aligned}$$

(2) Otherwise, either $t_1 \in I_2$ and $t_2 \in I_1 \cup I_3$ or $t_1 \in I_1 \cup I_3$ and $t_2 \in I_2$. By the definition of the color $w_{k+1} = 1$, we have

$$\begin{aligned}\bar{a}_{i_{t_1}}^{(r)} &= \tilde{a}_{i_{t_1}}; \\ \bar{a}_{i_{t_2}}^{(r)} &= a_{i_{t_1}}.\end{aligned}$$

Since $(a_k)_{k \in \mathbb{N}}$ and $(\tilde{a}_k)_{k \in \mathbb{N}}$ are free i.i.d copies of a , the following holds. First, let $t \in I_2$. Consider t' its matching symbol, that is, $\{t, t'\} \in \pi$ and $i_t = i_{t'} \neq i_j$, for all $j \notin \{t, t'\}$. Recall that, by the definition in (11), we have

$$\bar{a}_{i_t}^{(r)}, \bar{a}_{i_{t'}}^{(r)} \in \{a_{i_t}, \tilde{a}_{i_t}\}.$$

If $t' \in I_2$, Case (1) implies that the law of $(\bar{a}_{i_t}^{(r)}, \bar{a}_{i_{t'}}^{(r)})$ is equal to the law of $(a_{i_t}^{(r)}, a_{i_{t'}}^{(r)})$. If $t' \in I_1 \cup I_3$, as a_{i_t}, \tilde{a}_{i_t} are free i.i.d and free from $(a_k)_{k \neq i_t}, (\tilde{a}_k)_{k \neq i_t}$, we have

$$\left((\bar{a}_{i_j}^{(r)})_{j \in I_2 \setminus \{t\}}, \bar{a}_{i_t}^{(r)} \right) \stackrel{d}{=} \left((\bar{a}_{i_j}^{(r)})_{j \in I_2 \setminus \{t\}}, a_{i_t} \right) \stackrel{d}{=} \left((\bar{a}_{i_j}^{(r)})_{j \in I_2 \setminus \{t\}}, a_{i_t}^{(r)} \right). \quad (14)$$

Applying the above equalities in distribution for all $t \in I_2$, namely, Case (1) for $t' \in I_2$ and (14) for $t' \notin I_2$, we get

$$\left(\bar{a}_{i_j}^{(r)} \right)_{j \in I_2} \stackrel{d}{=} \left(a_{i_j}^{(r)} \right)_{j \in I_2}.$$

By symmetry, we also have

$$\left(\bar{a}_{i_j}^{(r)} \right)_{j \in I_1 \cup I_3} \stackrel{d}{=} \left(a_{i_j}^{(r)} \right)_{j \in I_1 \cup I_3}.$$

Additionally, $\left(\bar{a}_{i_j}^{(r)} \right)_{j \in I_2}$ is now free from $\left(\bar{a}_{i_j}^{(r)} \right)_{j \in I_1 \cup I_3}$. Indeed, let us show that each variable $\bar{a}_{i_t}^{(r)}$ is free from $\left(\bar{a}_{i_j}^{(r)} \right)_{j \in I_1 \cup I_3}$, for all $t \in I_2$. Consider t' its matching symbol.

First, if $t' \in I_2$, the fact that $(a_k)_{k \in \mathbb{N}}$ and $(\tilde{a}_k)_{k \in \mathbb{N}}$ are free i.i.d collections implies that $\bar{a}_{i_t}^{(r)}$ (and $\bar{a}_{i_{t'}}^{(r)}$) is free from $\left(\bar{a}_{i_j}^{(r)} \right)_{j \in I_1 \cup I_3}$. If $t' \in I_1 \cup I_3$, Case (2) implies that $\bar{a}_{i_t}^{(r)}$ is free from $\bar{a}_{i_{t'}}^{(r)}$ and then again $\bar{a}_{i_t}^{(r)}$ is free from $\left(\bar{a}_{i_j}^{(r)} \right)_{j \in I_1 \cup I_3}$.

In summary, by freeness, we have

$$\begin{aligned}\tau \left(\bar{a}_{J_1^c}^{(l)} \bar{a}_{J_2^c}^{(l)} \bar{a}_{J_3^c}^{(l)} \right) \tau \left(\bar{a}_{J_1^c}^{(r)} \bar{a}_{J_2^c}^{(r)} \bar{a}_{J_3^c}^{(r)} \right) &= \tau \left(a_{J_1^c}^{(l)} a_{J_2^c}^{(l)} a_{J_3^c}^{(l)} \right) \tau \left(a_{J_1^c}^{(r)} a_{J_2^c}^{(r)} \right) \tau \left(a_{J_2^c}^{(r)} \right) \\ &= R_2(J_1, J_2, J_3).\end{aligned}$$

Therefore

$$\begin{aligned}& \frac{1}{\delta^h} \sum_{\substack{J_1 \subseteq I_1 \\ J_2 \subseteq I_2 \\ J_3 \subseteq I_3}} (-1)^{|J_1|+|J_2|+|J_3|} \lambda^{2(|J_1|+|J_2|+|J_3|)} R_2(J_1, J_2, J_3) \\ &= \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)}, V^{(1)} \right),\end{aligned}$$

Following the same reasoning, it follows that

$$\begin{aligned}& \frac{1}{\delta^h} \sum_{\substack{J_1 \subseteq I_1 \\ J_2 \subseteq I_2 \\ J_3 \subseteq I_3}} (-1)^{|J_1|+|J_2|+|J_3|} \lambda^{2(|J_1|+|J_2|+|J_3|)} R_3(J_1, J_2, J_3) \\ &= \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)}, V^{(0)} \right).\end{aligned}$$

Combining the above relations in T , we get

$$T = \frac{\sigma^2 \lambda^2}{\delta^2} \sum_{w=0,1} \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)}, V^{(w)} \right).$$

It remains to note that

$$\frac{\sigma^2 \lambda^2}{\delta^2} = \frac{q}{2},$$

to finish the proof. \square

Lemma 5.6 (Case $\theta_2 = 1$). *Let $p \geq 4$ be an even integer, $\pi \in P_2^{\text{con}}(p)$. Let $V_1, \dots, V_k \in \pi$ be blocks, $w \in \{0, 1\}^k$ be a coloring of V_1, \dots, V_k and*

$$B_k = B_k \left(V_1^{(w_1)}, \dots, V_k^{(w_k)} \right) = (b_{i_j, k})_{j \in [p]}.$$

Let $V = \{v_1 < v_2\} \in \pi \setminus \{V_1, \dots, V_k\}$ and the joint law

$$(b_{i_{v_1}, k}, b_{i_{v_2}, k}) \stackrel{d}{=} (b_{i_{v_1}}(\theta), b_{i_{v_2}}(\theta)).$$

If $\theta_2 = 1$, we have

$$\begin{aligned} & \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k\}, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right) \\ &= \frac{q}{2} \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)}, V^{(1)} \right). \end{aligned}$$

Proof. We must prove that

$$\begin{aligned} T &:= \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k\}, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right) \\ &= \frac{q}{2} \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)}, V^{(1)} \right). \end{aligned}$$

As $\theta_2 = 1$, we can write

$$\begin{aligned} b_{i_{v_1}, k} &= \frac{1}{\delta} (\tilde{a} \otimes a - \lambda^2); \\ b_{i_{v_2}, k} &= \frac{1}{\delta} (a \otimes a - \lambda^2), \end{aligned}$$

where $a := a_{i_{v_1}}$. We have

$$\begin{aligned} T &= \frac{1}{\delta^2} \tau \otimes \tau \left\{ (B_k)_{I_1} \tilde{a} \otimes a (B_k)_{I_2} a \otimes a (B_k)_{I_3} \right\} \\ &\quad - \frac{\lambda^4}{\delta^2} \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right). \end{aligned} \tag{15}$$

We write again

$$B_k = \frac{1}{\delta} \left(a_{i_j}^{(l)} \otimes a_{i_j}^{(r)} - \lambda^2 \right)_{j \in [p]}.$$

Then

$$\begin{aligned} & \tau \otimes \tau \left\{ (B_k)_{I_1} \tilde{a} \otimes a (B_k)_{I_2} a \otimes a (B_k)_{I_3} \right\} \\ &= \frac{1}{\delta^h} \sum_{\substack{J_1 \subseteq I_1 \\ J_2 \subseteq I_2 \\ J_3 \subseteq I_3}} (-1)^{|J_1|+|J_2|+|J_3|} \lambda^{2(|J_1|+|J_2|+|J_3|)} \tau \left(a_{J_1^c}^{(l)} \tilde{a} a_{J_2^c}^{(l)} a a_{J_3^c}^{(l)} \right) \tau \left(a_{J_1^c}^{(r)} a a_{J_2^c}^{(r)} a a_{J_3^c}^{(r)} \right), \end{aligned}$$

where h is defined in (12). We use Lemmas 2.1 and 4.1 to compute

$$\begin{aligned} R(J_1, J_2, J_3) &:= \tau \left(a_{J_1^c}^{(l)} \tilde{a} a_{J_2^c}^{(l)} a_{J_3^c}^{(l)} \right) \tau \left(a_{J_1^c}^{(r)} a_{J_2^c}^{(r)} a_{J_3^c}^{(r)} \right) \\ &= \lambda^2 \tau \left(a_{J_1^c}^{(l)} a_{J_2^c}^{(l)} a_{J_3^c}^{(l)} \right) \left(\sigma^2 \tau(a_{J_1^c}^{(r)} a_{J_2^c}^{(r)}) \tau(a_{J_2^c}^{(r)}) + \lambda^2 \tau(a_{J_1^c}^{(r)} a_{J_2^c}^{(r)} a_{J_3^c}^{(r)}) \right) \end{aligned}$$

Hence

$$\begin{aligned} R(J_1, J_2, J_3) &= \lambda^2 \sigma^2 \tau \left(a_{J_1^c}^{(l)} a_{J_2^c}^{(l)} a_{J_3^c}^{(l)} \right) \tau \left(a_{J_1^c}^{(r)} a_{J_2^c}^{(r)} \right) \tau \left(a_{J_2^c}^{(r)} \right) \\ &\quad + \lambda^4 \tau \left(a_{J_1^c}^{(l)} a_{J_2^c}^{(l)} a_{J_3^c}^{(l)} \right) \tau \left(a_{J_1^c}^{(r)} a_{J_2^c}^{(r)} a_{J_3^c}^{(r)} \right). \end{aligned}$$

Both terms already appeared in the proof of Lemma 5.5. The second term cancels out with $\frac{\lambda^4}{\delta^2} \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right)$ in (15), and the contribution of the first is equal to

$$\begin{aligned} &\frac{1}{\delta^h} \sum_{\substack{J_1 \subseteq I_1 \\ J_2 \subseteq I_2 \\ J_3 \subseteq I_3}} (-1)^{|J_1|+|J_2|+|J_3|} \lambda^{2(|J_1|+|J_2|+|J_3|)} \tau \left(a_{J_1^c}^{(l)} a_{J_2^c}^{(l)} a_{J_3^c}^{(l)} \right) \tau \left(a_{J_1^c}^{(r)} a_{J_2^c}^{(r)} \right) \tau \left(a_{J_2^c}^{(r)} \right) \\ &= \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)}, V^{(1)} \right). \end{aligned}$$

We then have

$$T = \frac{\sigma^2 \lambda^2}{\delta^2} \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)}, V^{(1)} \right),$$

and the result follows by the definition of q . \square

The case $\theta_3 = 1$ is symmetric to the case $\theta_2 = 1$ in Lemma 5.6, and we omit the proof.

Lemma 5.7 (Case $\theta_3 = 1$). *Let $p \geq 4$ be an even integer, $\pi \in P_2^{\text{con}}(p)$. Let $V_1, \dots, V_k \in \pi$ be blocks, $w \in \{0, 1\}^k$ be a coloring of V_1, \dots, V_k and*

$$B_k = B_k \left(V_1^{(w_1)}, \dots, V_k^{(w_k)} \right) = (b_{i_j, k})_{j \in [p]}.$$

Let $V = \{v_1 < v_2\} \in \pi \setminus \{V_1, \dots, V_k\}$ and the joint law

$$(b_{i_{v_1}, k}, b_{i_{v_2}, k}) \stackrel{d}{=} (b_{i_{v_1}}(\theta), b_{i_{v_2}}(\theta)).$$

If $\theta_3 = 1$, we have

$$\begin{aligned} &\tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k\}, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right) \\ &= \frac{q}{2} \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)}, V^{(0)} \right). \end{aligned}$$

Finally, the case $\theta_4 = 1$ has a null contribution.

Lemma 5.8 (Case $\theta_4 = 1$). *Let $p \geq 4$ be an even integer, $\pi \in P_2^{\text{con}}(p)$. Let $V_1, \dots, V_k \in \pi$ be blocks, $w \in \{0, 1\}^k$ be a coloring of V_1, \dots, V_k and*

$$B_k = B_k \left(V_1^{(w_1)}, \dots, V_k^{(w_k)} \right) = (b_{i_j, k})_{j \in [p]}.$$

Let $V = \{v_1 < v_2\} \in \pi \setminus \{V_1, \dots, V_k\}$ and the joint law

$$(b_{i_{v_1}, k}, b_{i_{v_2}, k}) \stackrel{d}{=} (b_{i_{v_1}}(\theta), b_{i_{v_2}}(\theta)).$$

If $\theta_4 = 1$, we have

$$\tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k\}, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right) = 0.$$

Proof. Similarly to the previous proofs, we must show that

$$T := \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k\}, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right) = 0.$$

As $\theta_4 = 1$, we can write

$$\begin{aligned} b_{i_{v_1}, k} &= \frac{1}{\delta} (\tilde{a} \otimes \tilde{a} - \lambda^2); \\ b_{i_{v_2}, k} &= \frac{1}{\delta} (a \otimes a - \lambda^2), \end{aligned}$$

where $a := a_{i_{v_1}}$. First note that

$$\begin{aligned} T &= \frac{1}{\delta^2} \tau \otimes \tau \left\{ (B_k)_{I_1} \tilde{a} \otimes \tilde{a} (B_k)_{I_2} a \otimes a (B_k)_{I_3} \right\} \\ &\quad - \frac{\lambda^4}{\delta^2} \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right). \end{aligned}$$

Applying Lemma 4.1 to both tensors, we get that

$$\tau \otimes \tau \left\{ (B_k)_{I_1} \tilde{a} \otimes \tilde{a} (B_k)_{I_2} a \otimes a (B_k)_{I_3} \right\} = \lambda^4 \tau \otimes \tau \left\{ (B_k)_{I_1} (B_k)_{I_2} (B_k)_{I_3} \right\}.$$

The result follows by recalling that

$$\begin{aligned} \tau \otimes \tau \left\{ (B_k)_{I_1} (B_k)_{I_2} (B_k)_{I_3} \right\} &= \tau \otimes \tau \left((B_k)_{[p] \setminus (V_1 \cup \dots \cup V_k \cup V)} \right) \\ &= \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_k, V\}, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right). \end{aligned}$$

□

Proof of Proposition 5.4. Let $\pi \notin P_2^{\text{bicon}}(p)$, and let $2l^* + 1$ be the length of the smallest odd cycle in the intersection graph of π . We denote $V_1, \dots, V_{2l^*+1} \in \pi$ (any) ordered sequence of blocks in this smallest odd cycle, where V_j crosses V_{j+1} for each $j = 1, \dots, 2l^* + 1$ and $V_{2l^*+2} := V_1$. We first remove the first block V_1 , and by Lemma 5.5 we get that

$$\tau \otimes \tau(\pi) = \frac{q}{2} \sum_{w=0,1} \tau \otimes \tau \left(\pi \setminus V_1, V_1^{(w)} \right).$$

However, as it is the first block we remove, the distribution of $B_0 = (b_{ij})_{j \in [p]}$ is symmetric in both legs. In particular, we have

$$B_1(V_1^{(0)}) \stackrel{d}{=} B_1(V_1^{(1)}),$$

and

$$\tau \otimes \tau \left(\pi \setminus V_1, V_1^{(0)} \right) = \tau \otimes \tau \left(\pi \setminus V_1, V_1^{(1)} \right).$$

Hence

$$\tau \otimes \tau(\pi) = q \tau \otimes \tau \left(\pi \setminus V_1, V_1^{(0)} \right).$$

We then fix the color of V_1 to be $w_1 = 0$. We aim to show by induction that the colors $w_1, w_2, \dots, w_{2l^*}$ of V_1, \dots, V_{2l^*} are deterministic and alternating,

$$w = (w_1, \dots, w_{2l^*}) = (0, 1, 0, \dots, 1),$$

and so is $\theta^{(k+1)} := \theta \left(V_k, V_1^{(w_1)}, \dots, V_k^{(w_k)} \right)$ for $1 \leq k \leq 2l^*$ in the sense that

$$\left(\theta_2^{(2)}, \theta_3^{(3)}, \theta_2^{(4)}, \dots, \theta_2^{(2l^*)} \right) = (1, 1, 1, \dots, 1).$$

First, as V_2 crosses V_1 and $w_1 = 0$, we have $\theta_2^{(2)} = 1$. By Lemma 5.6, we have $w_2 = 1$. Suppose then that the result holds for some $1 < k < 2l^*$. Let us prove that it also holds for $k + 1$. To simplify the reading, let us divide into two cases whether k is even or odd.

Case k is even. The induction hypothesis implies that

$$(w_2, w_3, \dots, w_k) = (1, 0, \dots, 1);$$

$$\left(\theta_2^{(2)}, \theta_3^{(3)}, \theta_2^{(4)}, \dots, \theta_2^{(k)}\right) = (1, 1, 1, \dots, 1).$$

Since V_{k+1} crosses V_k , we automatically have that $\theta_1^{(k+1)} = 0$. Moreover, as $w_k = 1$, the definition of θ implies that

$$\theta_2^{(k+1)} = 0.$$

Hence either $\theta_4^{(k+1)} = 1$ or $\theta_3^{(k+1)} = 1$. Now consider all blocks V_{j_t} that might potentially cross V_{k+1} , for $j_t \leq k$. Since $2l^* + 1$ is the length of the smallest odd cycle, the cycles $(V_{j_t}, \dots, V_k, V_{k+1})$ are of even length. By a parity check, j_t must be even, and hence

$$w_{j_t} = 1.$$

This implies that all crossing blocks V_{j_t} of V_{k+1} for $j_t \leq k$ have the same color $w_{j_t} = 1$. In particular, we have

$$\theta_3^{(k+1)} = 1.$$

We then apply Lemma 5.7 to get that $w_{k+1} = 0$.

Case k is odd. This follows similarly to the even case. We first have that $\theta_1^{(k+1)} = 0$ and as $w_k = 0$, we have

$$\theta_3^{(k+1)} = 0.$$

By the parity check on the even cycles V_{k+1} might belong to, all crossing blocks V_{j_t} of V_{k+1} for $j_t \leq k$ have the same color $w_{j_t} = 0$. Hence

$$\theta_2^{(k+1)} = 1,$$

and Lemma 5.6 implies that $w_{k+1} = 1$. This finishes the induction.

To conclude the proof, the block V_{2l^*+1} crosses both V_1 of color $w_1 = 0$ and V_{2l^*} of color $w_{2l^*} = 1$. Hence

$$\theta_4^{(2l^*+1)} = 1.$$

Lemma 5.8 applied to $V = V_{2l^*+1}$ implies that

$$\tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_{2l^*}\}, V_1^{(0)}, \dots, V_{2l^*}^{(1)} \right) = 0.$$

Applying Lemma 5.6 for blocks $V = V_k$ when k is even and Lemma 5.7 for $V = V_k$ when k is odd, we get that

$$\begin{aligned} \tau \otimes \tau(\pi) &= q\tau \otimes \tau \left(\pi \setminus V_1, V_1^{(0)} \right) \\ &= q \cdot \frac{q}{2} \tau \otimes \tau \left(\pi \setminus \{V_1, V_2\}, V_1^{(0)}, V_2^{(1)} \right) \\ &= \dots \\ &= q \left(\frac{q}{2} \right)^{2l^*-1} \tau \otimes \tau \left(\pi \setminus \{V_1, \dots, V_{2l^*}\}, V_1^{(0)}, \dots, V_{2l^*}^{(1)} \right) \\ &= 0. \end{aligned}$$

Therefore, for all $\pi \notin F_2^{\text{bicon}}(p)$, we have $\tau \otimes \tau(\pi) = 0$. This proves the first part of the proposition.

Assume now that $\pi \in P_2^{\text{bicon}}(p)$. Let $V_1, \dots, V_{p/2}$ be an ordering of the blocks of π such that V_k crosses at least one V_j for $j < k$ and all k . This can always be done as π is connected. Let $\mathcal{V}_1, \mathcal{V}_2$ be the bipartite sets of vertices of the intersection graph of π ,

$$\begin{aligned}\mathcal{V}_1 &= \{V_1, V_{j_1}, \dots, V_{j_m}\}; \\ \mathcal{V}_2 &= \{V_2, V_{h_1}, \dots, V_{h_n}\}.\end{aligned}$$

Since π is bipartite, if $V_k \in \mathcal{V}_1$, all crossing blocks V_j of V_k belong to \mathcal{V}_2 , and similarly if $V_k \in \mathcal{V}_2$. We assume again that the color of V_1 is $w_1 = 0$, since

$$\tau \otimes \tau(\pi) = q\tau \otimes \tau(\pi \setminus V_1, V_1^{(0)}).$$

We will prove that both the color w_k and the binary vector $\theta^{(k)}$ of V_k depend only on which bipartite set V_k belongs to, namely, for all $k \geq 2$, the following hold.

- (1) If $V_k \in \mathcal{V}_1$, then $w_k = 0$ and $\theta_3^{(k)} = 1$;
- (2) Otherwise, $V_k \in \mathcal{V}_2$, $w_k = 1$ and $\theta_2^{(k)} = 1$.

Indeed, note first that $V_2 \in \mathcal{V}_2$, $\theta_2^{(2)} = 1$ as it crosses V_1 and $w_1 = 0$. Then, Lemma 5.6 implies that $w_2 = 1$. Assume then that the result holds for some l and for all $1 \leq k \leq l$. Let us prove that it holds for V_{l+1} as well. Indeed, assume without loss of generality that $V_{l+1} \in \mathcal{V}_1$. Then V_{l+1} only crosses blocks V_{j_t} such that $V_{j_t} \in \mathcal{V}_2$, for $j_t \leq l$. Since all colors $w_{j_t} = 1$, we deduce that

$$\theta_3^{(l+1)} = 1.$$

Then, Lemma 5.7 implies that $w_{l+1} = 0$ and induction is proved. We then apply Lemma 5.5 for V_1 , Lemma 5.6 for all blocks $V_k \in \mathcal{V}_2$ and Lemma 5.7 for all blocks $V_k \in \mathcal{V}_1$ for $k \geq 2$, so that

$$\begin{aligned}\tau \otimes \tau(\pi) &= q\tau \otimes \tau(\pi \setminus V_1, V_1^{(0)}) \\ &= q \cdot \frac{q}{2}\tau \otimes \tau(\pi \setminus \{V_1, V_2\}, V_1^{(0)}, V_2^{(1)}) \\ &= \dots \\ &= q \left(\frac{q}{2}\right)^{p/2-1}.\end{aligned}$$

It follows then that

$$\tau \otimes \tau(\pi) = q \left(\frac{q}{2}\right)^{p/2-1} = 2 \left(\frac{q}{2}\right)^{p/2},$$

and the result is proved. \square

5.4. Proof of Theorem 1.1. We are now ready to prove the main theorem. Recall that

$$\tau'(\mathbf{S}^p) = \sum_{\pi \in P_2(p)} \tau \otimes \tau(\pi) = \sum_{\hat{\pi}, (\pi_T)_{T \in \hat{\pi}}} \prod_{T \in \hat{\pi}} \tau \otimes \tau(\pi_T),$$

where the second summation runs over $(\hat{\pi}, (\pi_T)_{T \in \hat{\pi}}) \in \Phi(P_2(p))$, using the bijection Φ defined in (4). Note that if $|T| = 2$, then by Corollary 5.2 we have $\tau \otimes \tau(\pi_T) = 1$. Therefore, we deduce that

$$\tau'(\mathbf{S}^p) = \sum_{\hat{\pi}, (\pi_T)_{T \in \hat{\pi}}} \prod_{\substack{T \in \hat{\pi} \\ |T| \geq 4}} \tau \otimes \tau(\pi_T).$$

Using Proposition 5.4, we get

$$\tau'(\mathbf{S}^p) = \sum_{\hat{\pi} \in \mathcal{NC}(p)} \sum_{(\pi_T)_{T \in \hat{\pi}}} \prod_{\substack{T \in \hat{\pi} \\ |T| \geq 4}} 2 \left(\frac{q}{2}\right)^{\frac{|T|}{2}},$$

where the second summation runs over bipartite connected pair partitions π_T , for $T \in \hat{\pi}$. Finally, note that the number of size-two blocks is precisely $\mathbf{ncr}(\pi)$ and thus

$$|\{T \in \hat{\pi} : |T| \geq 4\}| = \mathbf{cc}(\pi) - \mathbf{ncr}(\pi).$$

Since

$$\sum_{\substack{T \in \hat{\pi} \\ |T| \geq 4}} \frac{|T|}{2} = \mathbf{cr}(\pi),$$

using again the bijection Φ , we deduce that

$$\tau'(\mathbf{S}^p) = \sum_{\pi \in \mathcal{P}_2^{\text{bi}}(p)} 2^{\mathbf{cc}(\pi) - \mathbf{ncr}(\pi)} \left(\frac{q}{2}\right)^{\mathbf{cr}(\pi)}.$$

This finishes the proof in view of Proposition 3.3 and of the fact that

$$\mathbf{ncr}(\pi) + \mathbf{cr}(\pi) = |\pi| = \frac{p}{2}.$$

REFERENCES

- [1] A. Ambainis, A. W. Harrow, and M. B. Hastings. Random tensor theory: Extending random matrix theory to mixtures of random product states. *Communications in Mathematical Physics*, 310(1):25–74, 2012. doi: 10.1007/s00220-011-1411-x.
- [2] G. Aubrun and I. Nechita. Realigning random states. *J. Math. Phys.*, 53(10):102210, 2012.
- [3] M. Bożejko and R. Speicher. Interpolations between bosonic and fermionic relations given by generalized Brownian motions. *Math. Z.*, 222(1):135–159, 1996. ISSN 0025-5874,1432-1823. doi: 10.1007/PL00004258. URL <https://doi.org/10.1007/PL00004258>.
- [4] B. Collins and P. Y. G. Lamarre. *-freeness in finite tensor products. *Advances in Applied Mathematics*, 83:47–80, 2017.
- [5] P. Dehornoy and P. Biane. Dual garside structure of braids and free cumulants of products. *Séminaire Lotharingien de Combinatoire*, 72:15, 2014.
- [6] D. Gouyou-Beauchamps. *Chemins sous-diagonaux et tableaux de Young*, page 112–125. Springer Berlin Heidelberg, 1986. ISBN 9783540474029. doi: 10.1007/bfb0072513. URL <http://dx.doi.org/10.1007/BFb0072513>.
- [7] C. Lancien, P. O. Santos, and P. Youssef. Limiting spectral distribution of random self-adjoint quantum channels, 2023.
- [8] F. Lehner. Free cumulants and enumeration of connected partitions. *European J. Combin.*, 23(8):1025–1031, 2002. ISSN 0195-6698,1095-9971. doi: 10.1006/eujc.2002.0619. URL <https://doi.org/10.1006/eujc.2002.0619>.
- [9] W. Młotkowski. Λ -free probability. *Infinite Dimensional Analysis, Quantum Probability and Related Topics*, 07(01):27–41, Mar. 2004. ISSN 1793-6306. doi: 10.1142/s0219025704001517. URL <http://dx.doi.org/10.1142/S0219025704001517>.
- [10] A. Nica. Free probability aspect of irreducible meandric systems, and some related observations about meanders. *Infinite Dimensional Analysis, Quantum Probability and Related Topics*, 19(02):1650011, June 2016. ISSN 1793-6306. doi: 10.1142/s0219025716500119. URL <http://dx.doi.org/10.1142/S0219025716500119>.

- [11] A. Nica and R. Speicher. *Lectures on the combinatorics of free probability*, volume 13. Cambridge University Press, 2006.
- [12] A. Nijenhuis and H. S. Wilf. The enumeration of connected graphs and linked diagrams. *J. Combin. Theory Ser. A*, 27(3):356–359, 1979. ISSN 0097-3165,1096-0899. doi: 10.1016/0097-3165(79)90023-2. URL [https://doi.org/10.1016/0097-3165\(79\)90023-2](https://doi.org/10.1016/0097-3165(79)90023-2).
- [13] R. Speicher and M. Weber. Quantum groups with partial commutation relations. *Indiana University Mathematics Journal*, 68(6):1849–1883, 2019.
- [14] R. Speicher and J. Wysoczański. Mixtures of classical and free independence. *Archiv der Mathematik*, 107(4):445–453, Sept. 2016. ISSN 1420-8938. doi: 10.1007/s00013-016-0955-6. URL <http://dx.doi.org/10.1007/s00013-016-0955-6>.
- [15] D. Voiculescu. *Symmetries of some reduced free product C^* -algebras*, page 556–588. Springer Berlin Heidelberg, 1985. ISBN 9783540395140. doi: 10.1007/bfb0074909. URL <http://dx.doi.org/10.1007/BFb0074909>.

CÉCILIA LANCIEN. CNRS & INSTITUT FOURIER UMR 5582, UNIVERSITÉ GRENOBLE ALPES, GRENOBLE, FRANCE.

Email address: cecilia.lancien@univ-grenoble-alpes.fr

PATRICK OLIVEIRA SANTOS. LABORATOIRE D'ANALYSE ET DE MATHÉMATIQUES APPLIQUÉES UMR 8050, UNIVERSITÉ GUSTAVE EIFFEL, UNIVERSITÉ PARIS EST CRÉTEIL, MARNE-LA-VALLÉE, FRANCE.

Email address: patrick.oliveirasantos@u-pem.fr

PIERRE YOUSSEF. DIVISION OF SCIENCE, NYU ABU DHABI, ABU DHABI, UAE & COURANT INSTITUTE OF MATHEMATICAL SCIENCES, NEW YORK UNIVERSITY, NEW YORK, USA.

Email address: yp27@nyu.edu