

IR-FIXED EUCLIDEAN VACUUM FOR LINEARIZED GRAVITY ON DE SITTER SPACE

CHRISTIAN GÉRARD AND MICHAŁ WROCHNA

ABSTRACT. We consider the Euclidean vacuum for linearized gravity on the global de Sitter space, obtained from the Euclidean Green's function on the 4-sphere. We use the notion of Calderón projectors to recover a quantum state for the Lorentzian theory on de Sitter space. We show that while the state is gauge invariant and Hadamard, it is not positive on the whole of the phase space. We show however that a suitable modification at low energies yields a well-defined Hadamard state on global de Sitter space.

1. INTRODUCTION AND MAIN RESULT

1.1. Introduction. The quantization of linearized gravity poses significant mathematical challenges as compared to scalar fields and Dirac fermions. Building on earlier advances [18, 7, 3, 14] (see e.g. [10, 8, 23, 6, 9, 15, 30, 24] for results for Maxwell, Proca, Yang–Mills and Teukolsky fields), the existence of a well-defined state satisfying the Hadamard condition has only recently been shown on spacetimes with a compact Cauchy surface [13]. Even so, requiring in addition that the state is invariant under spacetime symmetries could lead to further issues [22, 27, 26, 17].

A particularly important case in which one expects various simplifications is gravity linearized around the global de Sitter solution. In coordinates, de Sitter space is $dS^4 = \mathbb{R}_t \times \mathbb{S}^3$, equipped with the metric

$$g = -dt^2 + \cosh^2(t)h,$$

where h is the canonical metric on $\Sigma = \mathbb{S}^3$. Wick rotation $t \mapsto is$ yields the metric

$$\tilde{g} = ds^2 + \cos^2(s)h,$$

which is just the standard metric on the sphere \mathbb{S}^4 away from the two poles. The de Donder gauge-fixed linearized Einstein operator on dS^4 (denoted in the sequel by D_2) Wick rotates to an elliptic operator \tilde{D}_2 on \mathbb{S}^4 . It is invertible, so the Euclidean Green's function obtained from \tilde{D}_2^{-1} provides a natural candidate for the two-point function of a state after Wick rotating back to Lorentzian signature. The so-obtained “state”, if well-defined, is called *Euclidean* or *Bunch–Davies vacuum* in the physics literature. An explicit expression for the Green's function was derived by Allen [1], and further properties, formulas in different gauges, symmetries considerations

Date: May 2024.

2020 Mathematics Subject Classification. 81T20, 83C05, 58J47, 58J45, 58J32.

Key words and phrases. linearized Einstein equations, microlocal analysis, Quantum Field Theory on curved spacetimes, de Sitter space, Hadamard states, elliptic boundary value problems.

and computations of physical quantities have been obtained [1, 2, 21, 11], paralleled by results in the conformally flat chart [22, 27, 28, 26].

In the present paper we demonstrate that the Euclidean Green's function on \mathbb{S}^4 does actually *not* yield a state on de Sitter space due to positivity issues. We then propose a modification that resolves the problem, at the cost of breaking part of the de Sitter symmetries.

To explain the main issue, let us first recall the relevant preliminaries on linearized gravity. Let P be the Einstein operator linearized around the de Sitter metric g , and let K be the operator that generates linearized gauge transformations $u \mapsto u + Kw$. We will actually use a convention in which all operators are composed with a trace reversal $(Iu)_{ab} = u_{ab} - \frac{1}{2}\text{tr}_g(u)g_{ab}$, see Sect. 2, in particular $(Kw)_{ab} = \nabla_{(a}w_{b)} - \frac{1}{2}g_{ab}\nabla^c w_c$. The pre-symplectic space of smooth solutions modulo gauge is then the quotient space

$$\text{Ker } P / \text{Ran } K,$$

equipped with the canonical pre-symplectic form (or Hermitian form, as we prefer to use the complex formalism). In quantization it is often more convenient to work on the dual level with an isomorphic space of compactly supported smooth tensors

$$(1.1) \quad \text{Ker}_c K^* / \text{Ran}_c P,$$

where K^* is the formal adjoint of K for a non-positive Hermitian form involving I , explicitly $(K^*u)_b = -2\nabla^a u_{ab}$ in our convention. Then, in order to be interpreted as bi-solutions of P , two-point functions need to be defined as Hermitian forms on the quotient space (1.1). As we are working with the complex formalism, to define a state one actually needs a pair of two-point functions Λ_2^\pm (called *covariances* in the main part of the text to distinguish them from the associated operators and Schwartz kernels), see Subsect. 2.7.

The *de Donder gauge* or *harmonic gauge* consists in considering the *gauge-fixed* operator

$$D_2 = P + KK^*,$$

which in contrast to P is hyperbolic. Then, solutions of $Pu = 0$ are obtained by solving $D_2u = 0$ with the harmonic gauge condition $K^*u = 0$. Concerning quantization, let us stress that if we have candidates for two-point functions Λ_2^\pm which are bi-solutions of D_2 , they do not necessarily induce sesquilinear forms on the physical space (1.1). To ensure that they do, Λ_2^\pm need to satisfy the *weak gauge-invariance* condition¹

$$(1.2) \quad \Lambda_2^\pm = 0 \text{ on } \text{Ker}_c K^* \times \text{Ran}_c KK^*.$$

Furthermore, candidates for two-point functions need to satisfy a *positivity* condition to define a state. In order to ensure that the UV behaviour of fields is as good as possible, it is in addition highly desirable that Λ_2^\pm is defined by a pair of distributions which satisfies the *Hadamard condition*.

As already mentioned, the Green's function of \tilde{D}_2 can be Wick rotated to yield a candidate for a two-point function in Lorentzian signature. To

¹In fact, $\bar{u} \cdot \Lambda_2^+ P v = -\bar{u} \cdot \Lambda_2^+ K K^* v = 0$ for all $u \in \text{Ker}_c K^*$ and all test tensors v iff (1.2) holds. If Λ_2^+ is Hermitian then one gets automatically $\overline{Pv} \cdot \Lambda_2^+ u = 0$ as well.

have that under control it is best to work with Cauchy data at $\Sigma = \{s = 0\} = \{t = 0\}$. The Euclidean vacuum construction is then equivalent to the following.

Definition 1.1. *The Calderón projectors \tilde{c}_2^\pm of \tilde{D}_2 are the projections to the subspace of Cauchy data of L^2 solutions of $\tilde{D}_2 u = 0$ in $\pm s > 0$.*

Next, we define a new pair of operators c_2^\pm by multiplying tensor components of \tilde{c}_2^\pm by suitable powers of i . The operators c_2^\pm play then the role of Cauchy data of two-point function candidates on the Lorentzian side.

To carry out our analysis we use the *TT-gauge*, i.e. we use the residual gauge freedom to work with traceless tensors. Without going into details, this means in practice that the physical phase space is isomorphic to a quotient

$$\mathcal{E}_{\text{TT}}/\mathcal{F}_{\text{TT}},$$

equipped with a Hermitian form $\mathbf{q}_{I,2}$, where \mathcal{E}_{TT} is the space of Cauchy data of smooth traceless solutions D_2 satisfying the harmonic gauge, \mathcal{F}_{TT} describes the remaining gauge freedom, and $\mathbf{q}_{I,2}$ can be computed explicitly. The list of sufficient conditions for c_2^\pm to define a state becomes:

$$(1.3) \quad \begin{aligned} 1) \quad & c_2^+ + c_2^- = \mathbb{1} \text{ on } \mathcal{E}_{\text{TT}}, \\ 2) \quad & c_2^{\pm*} \mathbf{q}_{I,2} = \mathbf{q}_{I,2} c_2^\pm \text{ on } \mathcal{E}_{\text{TT}}, \\ 3) \quad & c_2^\pm K_{21\Sigma} f \in \text{Ran } K_{21\Sigma} + \text{Ran } K_{20\Sigma}, \quad f \in \text{Ran } K_{21\Sigma}^\dagger, \\ 4) \quad & \bar{f} \cdot \mathbf{q}_{I,2} c_2^\pm f \geq 0, \quad f \in \mathcal{E}_{\text{TT}}, \end{aligned}$$

where $K_{21\Sigma}$ is the Cauchy data version of the operator K mapping 1-tensors to 2-tensors, $K_{21\Sigma}^\dagger$ is the Cauchy data version of K^* , and $\text{Ran } K_{20\Sigma}$ is spanned by Cauchy data of the de Sitter metric g . Condition 3) corresponds to gauge invariance, and 4) is positivity. The Hadamard condition can also be expressed in terms of c_2^\pm , see Prop. 2.16. Our first result can be formulated as follows.

Theorem 1.2 (cf. Prop. 7.2–7.5). *The Euclidean Green's function on \mathbb{S}^4 does not define a state for linearized gravity on dS^4 by Wick rotation. More precisely, c_2^\pm satisfy properties 1)–3) and the Hadamard condition, but do not satisfy 4).*

The obstructions to positivity are low energy, finite-dimensional issues (and appear to be unrelated with the IR divergences in mode expansions in the conformally flat chart [22, 27, 26]).

The reason why 4) is an extremely delicate condition is that positivity of the Wick rotated operator \tilde{D}_2 does hold true (on traceless tensors, to be more precise) with respect to a *non-physical* Euclidean scalar product, different from the physical non-positive Hermitian form $\mathbf{q}_{I,2}$. One can try to find a gauge transformation which reduces the positivity condition to a subspace where the two Hermitian forms coincide. But the existence of such gauge transformation is far from being guaranteed; it is in particular highly sensitive to low-energy aspects. In fact we show that the physical space of

Cauchy data decomposes as

$$\mathcal{E}_{\text{TT}} = \mathcal{E}_{\text{TT,reg}} \oplus \mathcal{E}_{\text{TT},3} \oplus \mathcal{E}_{\text{TT},4},$$

where $\mathcal{E}_{\text{TT},3} \oplus \mathcal{E}_{\text{TT},4}$ are explicit problematic finite dimensional subspaces, and $\mathcal{E}_{\text{TT,reg}}$ is a “good” subspace on which the gauge transformation exists indeed. As a result, positivity holds true on $\mathcal{E}_{\text{TT,reg}}$, but on the other hand we also find by explicit computations that positivity is violated on a subspace of $\mathcal{E}_{\text{TT},3} \oplus \mathcal{E}_{\text{TT},4}$.

To deal with the non-positivity, we construct a projection of the form

$$\pi = \mathbb{1} + K_{21\Sigma} R : \mathcal{E}_{\text{TT}} \rightarrow \mathcal{E}_{\text{TT,reg}}$$

for some smoothing operator R . Then, we define a new state by composition with π . The special form of π means that the composition can be viewed as a further gauge transformation.

Theorem 1.3 (cf. Thm. 8.2). *The modified Euclidean vacuum ω_{mod} defined by composition with π is a well-defined Hadamard state for linearized gravity on global de Sitter space. It is invariant under the action of $O(4)$.*

The price to pay is that the state is not invariant under the action of the full group of de Sitter isometries.

One can also ask about the interpretation of the modes responsible for the positivity issues (as we show, they are also responsible for the fact that the Hermitian form $\mathbf{q}_{I,2}$ is degenerate on $\mathcal{E}_{\text{TT}}/\mathcal{F}_{\text{TT}}$). It turns out that a special role is played by the space of Killing one-forms \mathcal{K} . This becomes clearer once if we introduce the auxiliary hyperbolic operator $D_1 = K^*K$ acting on $(0,1)$ -tensors, which is related with the gauge-fixed operator D_2 by $D_2K = KD_1$. Its Wick rotated counterpart \tilde{D}_1 is elliptic but *not* invertible, with kernel isomorphic to \mathcal{K} or equivalently, to a corresponding subspace of Cauchy data \mathcal{K}_Σ . It turns out that c_2^\pm satisfies

$$(1.4) \quad c_2^\pm K_{21\Sigma} f \in \text{Ran } K_{21\Sigma} + \text{Ran } K_{20\Sigma} \text{ iff } f \in \mathcal{K}_\Sigma^{\mathbf{q}_1},$$

where $\mathcal{K}_\Sigma^{\mathbf{q}_1}$ is the orthogonal of \mathcal{K}_Σ with respect to the canonical conserved Hermitian form \mathbf{q}_1 associated to D_1 . We show that $\text{Ran } K_\Sigma^\dagger \subset \mathcal{K}_\Sigma^{\mathbf{q}_1}$, so (1.4) implies the weak gauge-invariance property 3) discussed above. On the other hand, the other implication in (1.4) means that c_2^\pm is *not* invariant with respect to transformations generated by elements outside of $\mathcal{K}_\Sigma^{\mathbf{q}_1}$. These kind of subtle phenomena involving the space of Killing 1-forms \mathcal{K} are ultimately responsible for the finite-dimensional issues with positivity.

We remark that in the literature [29, 19, 20], the space of infinitesimal symmetries \mathcal{K} of de Sitter space has entered the discussion of perturbative quantum gravity for an a priori different reason: it coincides with the space of *linearization instabilities*, which represents linear perturbations that are not relevant for non-linear Einstein equations. In our construction, thanks to modifying the Euclidean vacuum state with the projection π , we do not have to remove any special subspace from the phase space of linearized gravity (for instance through imposing non-linear constraints).

Finally, we show that the obstructions to positivity and the construction of the modified Euclidean vacuum ω_{mod} are not specific to linearized gravity.

In fact we carry out a similar analysis in the case of Maxwell fields, and obtain analogous conclusions.

1.2. Plan of the paper. In Subsect. 1.3 we introduce notation used throughout the paper.

Sect. 2 explains our conventions for linearized gravity and introduces preliminaries on quantization.

Sect. 3 briefly discusses Wick rotation of tensors.

Sect. 4 introduces Calderón projectors for second order differential operators on compact manifolds with an interface. The main novelty are results in the case when the operator is not invertible (as in the case of the operator \tilde{D}_1).

Sect. 5 focuses on Wick rotation of global de Sitter space. In particular it introduces key results on the spectral theory of the Wick-rotated operators \tilde{D}_2, \tilde{D}_1 .

Sect. 6 studies at length the TT gauge physical phase space $\mathcal{E}_{\text{TT}}/\mathcal{F}_{\text{TT}}$ and its decomposition, and constructs the projection $\pi : \mathcal{E}_{\text{TT}} \rightarrow \mathcal{E}_{\text{TT,reg}}$.

Sect. 7 shows crucial properties of the Euclidean vacuum pseudo state, including positivity on $\mathcal{E}_{\text{TT,reg}}$ and invariance under de Sitter symmetries, and proves Theorem 1.2.

Sect. 8 carries out the construction of the modified Euclidean vacuum state ω_{vac} and proves Theorem 1.3.

Appendix A contains various auxiliary results on Euclidean linearized gravity on \mathbb{S}^4 and the relevant elliptic boundary value problems.

Appendix B repeats the analysis in the case of Maxwell fields.

1.3. Notation. We now collect various notation used throughout the paper.

1.3.1. Isomorphisms of vector spaces. If E, F are vector spaces and $A \in L(E, F)$ we write $A : E \xrightarrow{\sim} F$ if A is an isomorphism. If E, F are topological vector spaces, we use the same notation if A is a homeomorphism.

1.3.2. Sesquilinear forms. If E is a complex vector space, its antidual is denoted by E^* . A sesquilinear form \mathbf{A} on E is an element of $L(E, E^*)$ and its action on elements of E is denoted by $\bar{u} \cdot \mathbf{A}v$. Often E is equipped with a reference Hilbertian scalar product $(\cdot|\cdot)$ and we associate to \mathbf{A} the operator $A \in L(E)$ by $\bar{u} \cdot \mathbf{A}v =: (v|Au)$.

1.3.3. Operators on quotient spaces. Let $F_i \subset E_i, i = 1, 2$ be vector spaces and let $A \in L(E_1, E_2)$. Then the induced map

$$[A] \in L(E_1/F_1, E_2/F_2),$$

defined in the obvious way, is

- 1) well-defined if $AE_1 \subset E_2$ and $AF_1 \subset F_2$,
 - 2) injective iff $A^{-1}F_2 = F_1$,
 - 3) surjective iff $E_2 = AE_1 + F_2$.
- (1.5)

1.3.4. *Sesquilinear forms on quotients.* Let now $F \subset E$ be vector spaces and let $\mathbf{A} \in L(E, E^*)$. We denote by $F^\circ \subset E^*$ the annihilator of F . Then the induced map

$$[\mathbf{A}] \in L(E/F, (E/F)^*),$$

defined as before, is

- (1.6) 1) well-defined if $\mathbf{A}E \subset F^\circ$, $F \subset \text{Ker } \mathbf{A}$,
 2) non-degenerate iff $F = \text{Ker } \mathbf{A}$.

If \mathbf{A} is hermitian or anti-hermitian then the condition $F \subset \text{Ker } \mathbf{A}$ implies the other one $\mathbf{A}E \subset F^\circ$ (and vice versa).

2. LINEARIZED GRAVITY

2.1. **Notation and background.** We start by fixing notation. Let (M, g) be a 4-dimensional Lorentzian manifold.

2.1.1. *Convention for the Riemann tensor.* We use the same convention as in e.g. [31, 7, 3, 14] for the sign of the Riemann tensor $R_{abcd} = R_{abc}{}^e g_{ed}$, i.e.

$$(\nabla_a \nabla_b - \nabla_b \nabla_a)u_c = R_{abc}{}^d u_d$$

on $(0, 1)$ -tensors in terms of the Levi-Civita connection ∇_a on (M, g) . Let us recall that the Ricci tensor is the symmetric tensor

$$\text{Ric}_{ab} = R_{acb}{}^c = R^c{}_{acb},$$

and the scalar curvature is $R = g^{ab} \text{Ric}_{ab}$. If $\dim M = 4$ then Einstein equations with cosmological constant Λ , i.e. $\text{Ric} - \frac{1}{2}gR + \Lambda g = 0$, are equivalent to

$$(2.1) \quad \text{Ric} = \Lambda g,$$

and as a consequence of (2.1) one gets $R = 4\Lambda$.

2.1.2. *Hermitian forms on tensors.* We denote by

$$V_k := \mathbb{C} \otimes_{\mathbb{S}}^k T^*M$$

the complex bundle of symmetric $(0, k)$ -tensors. We will only need the cases $k = 0, 1, 2$. V_k is equipped with the non-degenerate Hermitian form

$$(2.2) \quad (u|u)_{V_k} := \bar{u} \cdot k!(g^{\otimes k})^{-1}u.$$

In abstract index notation,

$$(u|u)_{V_k} = k! g^{a_1 b_1} \dots g^{a_k b_k} \bar{u}_{a_1 \dots a_k} u_{b_1 \dots b_k}.$$

For example for $k = 2$ we have

$$(2.3) \quad (u|u)_{V_2} = 2\text{tr}(u^* g^{-1} u g^{-1}).$$

The $k!$ normalization differs from the most common convention, it has however the advantage that various expressions involving adjoints have a more symmetric appearance.

For $U \subset M$ open, the Hermitian form (2.2) on fibers induces a Hermitian form on compactly supported smooth sections,

$$(2.4) \quad (u|v)_{V_k(U)} = \int_U (u(x)|v(x))_{V_k} d\text{vol}_g, \quad u, v \in C_0^\infty(U; V_k).$$

The adjoint of $A : C_0^\infty(M; V_k) \rightarrow C^\infty(M; V_l)$ for those Hermitian forms will be denoted by A^* .

If $\Sigma \subset M$ is a Cauchy surface, we set

$$(u|v)_{V_k(\Sigma)} = \int_{\Sigma} (u(x)|v(x))_{V_k} d\text{vol}_h, \quad u, v \in C_0^\infty(\Sigma; V_k),$$

where $d\text{vol}_h$ is the induced density on Σ .

2.1.3. Decomposition of tensors. Let us assume that $M = I \times \Sigma$ where $I \subset \mathbb{R}$ is an open interval, Σ a smooth manifold with variables (t, x) and

$$g = -dt^2 + h(t, x)dx^2,$$

where $h \in C^\infty(M, \otimes_s^2 T^* \Sigma)$ is a smooth t -dependent Riemannian metric on Σ . We set

$$V_{k\Sigma} = \mathbb{C} \otimes_s^k T^* \Sigma.$$

2.1.4. Decomposition of $(0, 1)$ -tensors. We identify

$$(2.5) \quad \begin{aligned} C^\infty(M; V_1) &\xrightarrow{\sim} C^\infty(I; C^\infty(\Sigma; V_{0\Sigma})) \oplus C^\infty(I; C^\infty(\Sigma; V_{1\Sigma})) \text{ by} \\ w &\mapsto (w_t, w_\Sigma), \\ w &=: w_t dt + w_\Sigma. \end{aligned}$$

The scalar product $(\cdot|\cdot)_{V_1}$ reads then

$$(w|w)_{V_1} = -|w_t|^2 + (w_\Sigma|w_\Sigma)_{V_{1\Sigma}} = -|w_t|^2 + (w_\Sigma|h^{-1}w_\Sigma).$$

2.1.5. Decomposition of $(0, 2)$ -tensors. Similarly we identify

$$C^\infty(M; V_2) \xrightarrow{\sim} C^\infty(I; C^\infty(\Sigma; V_{0\Sigma})) \oplus C^\infty(I; C^\infty(\Sigma; V_{2\Sigma})) \oplus C^\infty(I; C^\infty(\Sigma; V_{2\Sigma}))$$

by

$$(2.6) \quad \begin{aligned} u &\mapsto (u_{tt}, u_{t\Sigma}, u_{\Sigma\Sigma}), \\ u &=: u_{tt} dt \otimes dt + u_{t\Sigma} \otimes dt + dt \otimes u_{t\Sigma} + u_{\Sigma\Sigma}. \end{aligned}$$

The scalar product $(\cdot|\cdot)_{V_2}$ reads:

$$(2.7) \quad (u|u)_{V_2} = 2|u_{tt}|^2 - 4(u_{t\Sigma}|u_{t\Sigma})_{V_{1\Sigma}} + (u_{\Sigma\Sigma}|u_{\Sigma\Sigma})_{V_{2\Sigma}}.$$

2.2. The differential and its adjoint. Let

$$d : \begin{aligned} C^\infty(M; V_k) &\rightarrow C^\infty(M; V_{k+1}) \\ (du)_{a_1, \dots, a_{k+1}} &= \nabla_{(a_1} u_{a_2, \dots, a_{k+1})}, \end{aligned}$$

where $u_{(a_1 \dots a_k)}$ is the symmetrization of $u_{a_1 \dots a_k}$, and

$$\delta : \begin{aligned} C^\infty(M; V_k) &\rightarrow C^\infty(M; V_{k-1}) \\ (\delta u)_{a_1, \dots, a_{k-1}} &= -k \nabla^a u_{aa_1 \dots a_{k-1}}. \end{aligned}$$

With these (non-standard) conventions, we have $d^* = \delta$ w.r.t. the Hermitian form (2.4).

2.3. Operators on tensors.

2.3.1. *Trace reversal.* The operator of *trace reversal* I is given by

$$I := \mathbb{1} - \frac{1}{4}|g\rangle(g|,$$

where $\mathbb{1}$ is the identity and

$$(g| : u_2 \mapsto (g|u_2)_{V_2}, \quad |g\rangle : u_0 \mapsto u_0 g,$$

i.e. I is the orthogonal symmetry w.r.t. the line $\mathbb{C}g$. Equivalently,

$$(Iu)_{ab} = u_{ab} - \frac{1}{2}\mathrm{tr}_g(u)g_{ab}, \quad \mathrm{tr}_g(u) := g^{ab}u_{ab} = \frac{1}{2}(g|u)_{V_2}.$$

It satisfies

$$(2.8) \quad I^2 = \mathbb{1}, \quad I = I^* \text{ on } C^\infty(M; V_2).$$

2.3.2. *Ricci operator.* The *Ricci operator* is

$$\mathrm{Riem}(u)_{ab} := R_a^c{}^d{}_b u_{cd} = R_{ab}^c{}^d u_{cd}, \quad u \in C^\infty(M; V_2).$$

The fact that Riem preserves symmetric $(0, 2)$ -tensors follows from the symmetries of the Riemann tensor.

Lemma 2.1. *The Ricci operator satisfies:*

$$(2.9) \quad \begin{aligned} i) & \quad \mathrm{Riem} g = -\mathrm{Ric}, \\ ii) & \quad \mathrm{Riem} \circ I = I \circ \mathrm{Riem} \text{ if } g \text{ is Einstein,} \\ iii) & \quad \mathrm{Riem} = \mathrm{Riem}^*. \end{aligned}$$

2.4. **Lichnerowicz operators.** Let $-\square_i$ be the rough d'Alembertian acting on sections of V_k , i.e.

$$-\square_i u_i = -g^{ab}\nabla_{e^a, e^b}^2 u_i,$$

where $(e_a)_{0 \leq a \leq d}$ is a local frame and $\nabla_{u,v}^2 = \nabla_u \circ \nabla_v - \nabla_{\nabla_u v}$ for vector fields u, v on M .

The *Lichnerowicz operators* acting on sections of V_k , see [25], are defined by:

$$(2.10) \quad \begin{aligned} D_{0,L} &= -\square_0, \\ D_{1,L} &= -\square_1 + \mathrm{Ric} \circ g^{-1}, \\ D_{2,L} &= -\square_2 + \mathrm{Ric} \circ g^{-1} \circ \cdot + \cdot \circ g^{-1} \circ \mathrm{Ric} + 2\mathrm{Riem}. \end{aligned}$$

They satisfy

$$D_{i,L} = D_{i,L}^*.$$

The proofs of the next facts can be found for instance in [4]. The Lichnerowicz operators satisfy

$$(2.11) \quad \begin{aligned} D_{1,L} &= \delta_a \circ d_a + d_a \circ \delta_a = \delta \circ d - d \circ \delta + 2\mathrm{Ric} \circ g^{-1}, \\ D_{2,L} &= \delta \circ d - d \circ \delta + 2(\mathrm{Ric} \circ g^{-1} \circ \cdot + \cdot \circ g^{-1} \circ \mathrm{Ric}) + 4\mathrm{Riem}, \end{aligned}$$

where d_a, δ_a are the anti-symmetric differential and codifferential.

Proposition 2.2. *If (M, g) is Einstein then for $i = 0, 1$:*

$$\begin{aligned} D_{i+1,L} \circ d &= d \circ D_{i,L}, \quad \delta \circ D_{i+1,L} = D_{i,L} \circ \delta, \\ (g| \circ D_{2,L} &= D_{0,L} \circ (g|, \quad D_{2,L} \circ |g\rangle = |g\rangle \circ D_{0,L}. \end{aligned}$$

2.5. Linearized gravity as a gauge theory. Let (M, g) be a globally hyperbolic spacetime of dimension 4. We assume that (M, g) is Einstein. Let us introduce the differential operators

$$(2.12) \quad \begin{aligned} P &:= -\square_2 - I \circ d \circ \delta + 2 \text{Riem} \in \text{Diff}^2(M; V_2), \\ K &:= I \circ d \in \text{Diff}^1(M; V_1, V_2). \end{aligned}$$

$Pu = 0$ is the *linearized Einstein equation* (in the trace-reversed form). The condition $K^*u = 0$, where K^* is defined below, is the linearized *de Donder* or *harmonic gauge*.

2.5.1. Physical Hermitian form. We consider V_k , $k = 0, 1, 2$ as Hermitian bundles, where the Hermitian forms on fibers is now

$$(2.13) \quad (u|u)_{I, V_k} := (u|u)_{V_k}, \quad k = 0, 1, \quad (u|u)_{I, V_2} := (u|Iu)_{V_2}.$$

The corresponding Hermitian form on smooth sections of V_k , $k = 1, 2$ is

$$(2.14) \quad (u|u)_{I, V_k(U)} = \int_U (u(x)|u(x))_{I, V_k} d\text{vol}_g, \quad u, v \in C_0^\infty(U; V_k).$$

We denote by A^* the corresponding formal adjoint of A for $(\cdot|\cdot)_{I, V_k(M)}$ to distinguish it from the formal adjoint A^* for $(\cdot|\cdot)_{V_k(M)}$. The two are related as follows:

$$(2.15) \quad \begin{aligned} A^* &= IA^*I \text{ if } A : C^\infty(M; V_2) \rightarrow C^\infty(M; V_2), \\ A^* &= A^*I \text{ if } A : C^\infty(M; V_k) \rightarrow C^\infty(M; V_2), \quad k = 0, 1, \\ A^* &= IA^* \text{ if } A : C^\infty(M; V_2) \rightarrow C^\infty(M; V_k), \quad k = 0, 1, \\ A^* &= A^* \text{ if } A : C^\infty(M; V_i) \rightarrow C^\infty(M; V_j), \quad i, j \neq 2. \end{aligned}$$

In particular,

$$(2.16) \quad K^* = K^*I = \delta \circ I \circ I = \delta.$$

2.5.2. Operators in linearized gravity. Let us set:

$$D_k := D_{k,L} - 2\Lambda, \quad k = 0, 1, 2.$$

Then

$$(2.17) \quad \begin{aligned} K^*K &= D_1 = -\square_1 - \Lambda = \delta \circ d - d \circ \delta, \\ P + KK^* &= D_2 = -\square_2 + 2\text{Riem} = \delta \circ d - d \circ \delta + 4\text{Riem} + 2\Lambda. \end{aligned}$$

The operator D_0 is useful in connection with the *traceless gauge*.

2.5.3. Cauchy problem. Let $\Sigma \subset M$ a smooth space-like Cauchy surface. For $k = 0, 1, 2$ we set

$$\varrho_k u = \begin{pmatrix} u|_\Sigma \\ i^{-1} \nabla_\nu u|_\Sigma \end{pmatrix} = \begin{pmatrix} f_0 \\ f_1 \end{pmatrix}, \quad u \in C_{\text{sc}}^\infty(M; V_k),$$

where ν is the future directed unit normal to Σ .

We denote by U_k the operator which assigns to a set of Cauchy data the corresponding solution in $\text{Ker}_{\text{sc}} D_k$, i.e. U_k is the inverse of $\varrho_k : \text{Ker}_{\text{sc}} D_k \rightarrow C_0^\infty(\Sigma; V_k \otimes \mathbb{C}^2)$. In other terms,

$$(2.18) \quad \begin{cases} D_k U_k = 0, \\ \varrho_k U_k = \mathbb{1}. \end{cases}$$

2.5.4. *Conserved charges.* Let G_k be the causal propagator of D_k , i.e. the difference of the retarded and advanced propagator. There exist a unique Hermitian form $\mathbf{q}_k : C_0^\infty(\Sigma; V_k \otimes \mathbb{C}^2) \rightarrow C_0^\infty(\Sigma; V_k \otimes \mathbb{C}^2)^*$ called the *charge* of D_k , such that

$$(\phi_k | iG_k \phi_k)_{V_k(M)} = \overline{\varrho_k u_k} \cdot \mathbf{q}_k \varrho_k u_k$$

for $\phi_k \in C_0^\infty(M; V_k)$ and $u_k = G_k \phi_k \in \text{Ker}_{\text{sc}} D_k$. By [14, Lemma 2.2], one can compute \mathbf{q}_k using the identity

$$\begin{aligned} & (u_k | D_k v_k)_{V_k(J_\pm(\Sigma))} - (D_k u_k | v_k)_{V_k(J_\pm(\Sigma))} \\ &= \pm i^{-1} \overline{\varrho_k u_k} \cdot \mathbf{q}_k \varrho_k u_k, \quad u_k, v_k \in C_0^\infty(M; V_k). \end{aligned}$$

2.5.5. *Operators on Cauchy data and physical charge.* We follow here [15, Subsect. 2.4].

To the operator K we associate an operator K_Σ acting on Cauchy data by setting

$$(2.19) \quad K_\Sigma := \varrho_2 K U_1.$$

Similarly, since $[I, D_2] = 0$ we can define

$$(2.20) \quad I_\Sigma := \varrho_2 I U_2 = I \otimes \mathbb{C}^2.$$

We obtain that $\mathbf{q}_2 I_\Sigma = I_\Sigma^* \mathbf{q}_2$ and as in 2.5.1 we define the Hermitian form

$$\mathbf{q}_{I,2} := \mathbf{q}_2 \circ I_\Sigma,$$

called the *physical charge* for D_2 .

We denote by K_Σ^\dagger the adjoint of K_Σ for the Hermitian forms $\mathbf{q}_1, \mathbf{q}_{I,2}$, i.e.

$$(2.21) \quad \overline{K_\Sigma^\dagger f_2} \cdot \mathbf{q}_1 f_1 = \overline{f_2} \cdot \mathbf{q}_{I,2} K_\Sigma f_1, \quad f_k \in C_0^\infty(\Sigma, V_k \otimes \mathbb{C}^2).$$

We have:

$$K_\Sigma^\dagger = \varrho_1 K^* U_2.$$

Lemma 2.3. *We have:*

- (1) $K \circ U_1 = U_2 \circ K_\Sigma$, $K^* \circ U_2 = U_1 \circ K_\Sigma^\dagger$;
- (2) $\varrho_2 \circ K = K_\Sigma \circ \varrho_1$ on $\text{Ker}_{\text{sc}} D_1$, $\varrho_1 \circ K^* = K_\Sigma^\dagger \circ \varrho_2$ on $\text{Ker}_{\text{sc}} D_2$;
- (3) $K_\Sigma^\dagger \circ K_\Sigma = 0$.

2.6. **Phase spaces.** We use the notation Ker_c and Ran_c (resp. Ker_{sc} and Ran_{sc}) to denote the kernel and range of differential operators acting on compactly supported (resp. space-compact) smooth tensors.

Proposition 2.4. *The maps*

$$\begin{aligned} [G_2] &: \frac{\text{Ker}_c K^*}{\text{Ran}_c P} \longrightarrow \frac{\text{Ker}_{\text{sc}} P}{\text{Ran}_{\text{sc}} K}, \\ [\iota] &: \frac{\text{Ker}_{\text{sc}} D_2 \cap \text{Ker}_{\text{sc}} K^*}{K \text{Ker}_{\text{sc}} D_1} \longrightarrow \frac{\text{Ker}_{\text{sc}} P}{\text{Ran}_{\text{sc}} K}, \end{aligned}$$

where $\iota : \text{Ker}_{\text{sc}} D_2 \cap \text{Ker}_{\text{sc}} K^* \rightarrow \text{Ker}_{\text{sc}} P$ is the canonical injection, are well defined and bijective.

Let us define the Hermitian forms \mathbf{Q}_k on $C_0^\infty(M; V_k)$:

$$\overline{u}_k \cdot \mathbf{Q}_k u_k := i(u_k | G_k u)_{V_k(M)},$$

and the *physical charge*

$$\mathbf{Q}_{I,2} := \mathbf{Q}_2 \circ I = I^* \circ \mathbf{Q}_2.$$

Definition 2.5. *The physical phase space is the Hermitian space $(\mathcal{V}_P, \mathbf{Q}_P)$, where:*

$$\mathcal{V}_P = \frac{\text{Ker}_c K^*}{\text{Ran}_c P}, \quad [\overline{u}] \cdot \mathbf{Q}_P [v] = \overline{u} \cdot \mathbf{Q}_{I,2} v, \quad [u], [v] \in \frac{\text{Ker}_c K^*}{\text{Ran}_c P}.$$

\mathbf{Q}_P is a well-defined Hermitian form on \mathcal{V}_P .

2.6.1. *Phase space of Cauchy data.*

Proposition 2.6. *The induced map*

$$[\varrho_2] : \frac{\text{Ker}_{\text{sc}} D_2 \cap \text{Ker}_{\text{sc}} K^*}{K \text{Ker}_{\text{sc}} D_1} \longrightarrow \frac{\text{Ker}_c K_\Sigma^\dagger}{\text{Ran}_c K_\Sigma}$$

is well defined and bijective.

Proposition 2.7. *The map*

$$[\varrho_2 G_2] : \left(\frac{\text{Ker}_c K^*}{\text{Ran}_c P}, \mathbf{Q}_P \right) \longrightarrow \left(\frac{\text{Ker}_c K_\Sigma^\dagger}{\text{Ran}_c K_\Sigma}, \mathbf{q}_{I,2} \right)$$

is an isomorphism of Hermitian spaces.

2.7. Quantization. The algebraic quantization of linear gauge theories is discussed in detail in [15, Sect. 3]. The algebraic framework reduces the quantization problem to showing the existence of physically relevant quantum states on the CCR $*$ -algebra $\text{CCR}(\mathcal{V}_P, \mathbf{Q}_P)$ associated to the Hermitian space $(\mathcal{V}_P, \mathbf{Q}_P)$ defined in Subsect. 2.6. The notions of quasi-free states and covariances (or two-point functions) are explained in [15, Sect. 3] and references therein.

2.7.1. *Covariances.* In the complex formalism, a quasi-free state on the CCR $*$ -algebra $\text{CCR}(\mathcal{V}_P, \mathbf{Q}_P)$ is determined by a pair $\mathbf{\Lambda}_P^\pm$ of *covariances*, i.e. of Hermitian forms on \mathcal{V}_P such that

$$\begin{cases} \mathbf{\Lambda}_P^\pm \geq 0, \\ \mathbf{\Lambda}_P^+ - \mathbf{\Lambda}_P^- = \mathbf{Q}_P. \end{cases}$$

We will consider quasi-free states ω on $\text{CCR}(\mathcal{V}_P, \mathbf{Q}_P)$ with covariances obtained from a pair of continuous Hermitian forms $\mathbf{\Lambda}_2^\pm$ on $C_0^\infty(M; V_2)$ (called the *spacetime covariances* of ω) by:

$$(2.22) \quad [\overline{u}] \cdot \mathbf{\Lambda}_P^\pm [u] = \overline{u} \cdot \mathbf{\Lambda}_2^\pm u, \quad [u] \in \frac{\text{Ker}_c K^*}{\text{Ran}_c P}.$$

Lemma 2.8. *Suppose that $\mathbf{\Lambda}_2^\pm \in L(C_0^\infty(M; V_2), C_0^\infty(M; V_2)^*)$ are such that:*

$$(2.23) \quad \begin{aligned} & i) \quad D_2^* \circ \mathbf{\Lambda}_2^\pm = \mathbf{\Lambda}_2^\pm \circ D_2 = 0, \\ & ii) \quad \mathbf{\Lambda}_2^+ - \mathbf{\Lambda}_2^- = \mathbf{Q}_{I,2} \text{ on } \text{Ker}_c K^*, \\ & iii) \quad \mathbf{\Lambda}_2^\pm = 0 \text{ on } \text{Ker}_c K^* \times \text{Ran}_c K K^*, \\ & v) \quad \mathbf{\Lambda}_2^\pm = \mathbf{\Lambda}_2^{\pm*}, \quad \mathbf{\Lambda}_2^\pm \geq 0 \text{ on } \text{Ker}_c K^*. \end{aligned}$$

Then Λ_2^\pm are the covariances of a quasi-free state on $\text{CCR}(\mathcal{V}_P, \mathbf{Q}_P)$.

Lemma 2.8 (its proof is straightforward) has appeared in a very similar form in [15, 14, 13], but with *iii*) replaced by the stronger condition

$$\Lambda_2^\pm = 0 \text{ on } \text{Ker}_c K^* \times \text{Ran}_c K.$$

The weak gauge-invariance condition *iii*) is actually sufficient, see the discussion around (1.2) in Subsect. 1.1.

2.7.2. Hadamard condition. We use the following definition of Hadamard states [32], cf. [15, Subsect. 3.4] and references therein. The general consensus is that only states satisfying the *Hadamard condition* (the *Hadamard states*) are physical. We recall that

$$\mathcal{N} = \{(x, \xi) \in T^*M \setminus o : \xi \cdot g^{-1}(x)\xi = 0\}$$

is the characteristic set of the wave operator on (M, g) , and

$$\mathcal{N}^\pm = \mathcal{N} \cap \{(x, \xi) : \pm v \cdot \xi > 0 \ \forall v \in T_x M \text{ future-directed time-like}\}$$

are its two connected components.

To formulate the Hadamard condition, we need to identify the Hermitian forms Λ_2^\pm with distributional kernels $\Lambda_2^\pm(\cdot, \cdot) \in \mathcal{D}'(M \times M; L(V_2))$, called *two-point functions*.

This identification is equivalent to embedding $C_0^\infty(M; V_2)$ in $\mathcal{D}'(M; V_2)$ and is defined by the identity

$$(2.24) \quad \bar{u} \cdot \Lambda_2^\pm v =: \int_{M \times M} (u(x) | \Lambda_2^\pm(x, y) v(y))_{V_2} d\text{vol}_g(x) d\text{vol}_g(y)$$

for all $u, v \in C_0^\infty(M; V_2)$. One can of course use other Hermitian forms on the fibers of V_2 to do this identification, like for example $(\cdot | \cdot)_{I, V_2}$. This change amounts to composing $\Lambda_2^\pm(x, y)$ by smooth linear operators acting on the fibers of V_2 over x and y and it does not change the Hadamard condition (2.25) below.

Note that most of the literature on QFT on curved spacetimes uses the real formalism. The commonly used *real two-point functions* $\omega_2(x, y)$ are obtained by the formula $\omega_2 = \text{Re}(\Lambda_2^\pm \pm \frac{1}{2} Q_{I,2})$, where $Q_{I,2}$ is the distribution obtained from $\mathbf{Q}_{I,2}$ using the above identification, see e.g. [12, Sect. 4].

Definition 2.9. A quasi-free state ω on $\text{CCR}(\mathcal{V}_P, \mathbf{Q}_P)$ given by covariances Λ_2^\pm as in Lemma 2.8 is Hadamard if in addition to (2.23) it satisfies:

$$(2.25) \quad \text{WF}(\Lambda_2^\pm)' \subset \mathcal{N}^\pm \times \mathcal{N}^\pm.$$

2.7.3. Hadamard condition on a Cauchy surface. One can equivalently consider Hermitian forms $\lambda_{2\Sigma}^\pm$ on the space of Cauchy data $C_0^\infty(\Sigma; V_2 \otimes \mathbb{C}^2)$ called *Cauchy surface covariances*. Namely if we have a pair of Hermitian forms

$$\lambda_{2\Sigma}^\pm \in L(C_0^\infty(\Sigma; V_2 \otimes \mathbb{C}^2), C_0^\infty(\Sigma; V_2 \otimes \mathbb{C}^2)^*),$$

then we set

$$(2.26) \quad \Lambda_2^\pm = (\varrho_2 G_2)^* \lambda_{2\Sigma}^\pm (\varrho_2 G_2) \in L(C_0^\infty(M; V_2), C_0^\infty(M; V_2)^*).$$

Note that $\lambda_{2\Sigma}^\pm$ generate covariances $[\lambda_{2\Sigma}^\pm]$ on $(\text{Ker}_c K_\Sigma^\dagger / \text{Ran}_c K_\Sigma, \mathbf{q}_{I,2})$ iff

$$(2.27) \quad \begin{aligned} i) \quad & \lambda_{2\Sigma}^+ - \lambda_{2\Sigma}^- = \mathbf{q}_{I,2} \text{ on } \text{Ker}_c K_\Sigma^\dagger, \\ ii) \quad & \lambda_{2\Sigma}^\pm = 0 \text{ on } \text{Ker}_c K_\Sigma^\dagger \times \text{Ran}_c K_\Sigma, \\ iii) \quad & \lambda_{2\Sigma}^\pm = \lambda_{2\Sigma}^{\pm*}, \lambda_{2\Sigma}^\pm \geq 0 \text{ on } \text{Ker}_c K_\Sigma^\dagger. \end{aligned}$$

We will call property (2.27) *ii*) the *strong gauge invariance* property.

However, in order for the spacetime covariances Λ_2^\pm given by (2.26) to satisfy (2.23) the following weaker conditions are sufficient:

$$(2.28) \quad \begin{aligned} i) \quad & \lambda_{2\Sigma}^+ - \lambda_{2\Sigma}^- = \mathbf{q}_{I,2} \text{ on } \text{Ker}_c K_\Sigma^\dagger, \\ ii) \quad & \lambda_{2\Sigma}^\pm = 0 \text{ on } \text{Ker}_c K_\Sigma^\dagger \times \text{Ran}_c K_\Sigma K_\Sigma^\dagger, \\ iii) \quad & \lambda_{2\Sigma}^\pm = \lambda_{2\Sigma}^{\pm*}, \lambda_{2\Sigma}^\pm \geq 0 \text{ on } \text{Ker}_c K_\Sigma^\dagger. \end{aligned}$$

We call property (2.28) *ii*) the *weak gauge invariance* property.

Since $\mathbf{q}_{I,2}$ is non-degenerate, we can also define

$$(2.29) \quad \lambda_{2\Sigma}^\pm =: \pm \mathbf{q}_{I,2} \circ c_2^\pm.$$

Proposition 2.10. *Suppose $c_2^\pm : C_0^\infty(\Sigma; V_2 \otimes \mathbb{C}^2) \rightarrow C^\infty(\Sigma; V_2 \otimes \mathbb{C}^2)$ is a pair of operators such that:*

$$(2.30) \quad \begin{aligned} i) \quad & c_2^+ + c_2^- = \mathbb{1}, \\ ii) \quad & c_2^\pm : \text{Ran}_c K_\Sigma K_\Sigma^\dagger \rightarrow \text{Ran } K_\Sigma, \\ iii) \quad & \mathbf{q}_{I,2} \circ c_2^\pm = c_2^{\pm*} \circ \mathbf{q}_{I,2}, \quad \pm \mathbf{q}_{I,2} \circ c_2^\pm \geq 0 \text{ on } \text{Ker}_c K_\Sigma^\dagger. \end{aligned}$$

Then Λ_2^\pm given by (2.26) and (2.29) are the covariances of a quasi-free state on $\text{CCR}(\mathcal{V}_P, \mathbf{Q}_P)$. Furthermore if for some neighborhood \mathcal{U} of Σ in M we have:

$$(2.31) \quad iv) \quad \text{WF}(U_2 \circ c_2^\pm)' \subset (\mathcal{N}^\pm \cup \mathcal{F}) \times T^*\Sigma$$

over $\mathcal{U} \times \Sigma$, where $\mathcal{F} \subset T^*M$ is a conic set with $\mathcal{F} \cap \mathcal{N} = \emptyset$, then the associated state is Hadamard.

The proof of (2.30) is analogous to the one found in [15, Sect. 3.4]. The proof of the statement on the Hadamard condition can be found in [12, Sect. 11.1].

Remark 2.11. *Condition (2.30) *iii*) on c_2^\pm is equivalent to the weak gauge invariance property (2.28) *iii*). The corresponding condition for the strong gauge invariance property (2.27) *iii*) is*

$$c_2^\pm : \text{Ran}_c K_\Sigma \rightarrow \text{Ran } K_\Sigma.$$

Remark 2.12. *If $[c_2^\pm, I_\Sigma] = 0$ then we can replace the first condition in (2.30) *iii*) by the simpler*

$$\mathbf{q}_2 \circ c_2^\pm = c_2^{\pm*} \circ \mathbf{q}_2.$$

Conversely if c_2^\pm satisfy the conditions in Prop. 2.10 then setting

$$\hat{c}_2^\pm = \frac{1}{2}(c_2^\pm + I_\Sigma \circ c_2^\pm \circ I_\Sigma),$$

we obtain that \hat{c}_2^\pm also satisfies the conditions in Prop. 2.10 and $[\hat{c}_2^\pm, I_\Sigma] = 0$. The only point deserving some attention is the microlocal condition iv), which follows from the fact that $\text{WF}(I)'$ and $\text{WF}(I_\Sigma)'$ are included in the diagonal of $T^*M \times T^*M$ and $T^*\Sigma \times T^*\Sigma$ respectively.

2.8. TT gauge condition. We now explain how to add the traceless condition $(g|u_2)_{V_2} = 0$ to the harmonic gauge condition $\delta u_2 = 0$, following [7]. These two conditions together are called the *TT gauge condition*.

It is a natural gauge fixing condition, as it does not single out a particular Cauchy surface and is preserved by all isometries of the spacetime (M, g) .

In the rest of the paper we will set

$$\begin{aligned} K_{21} &= K = I \circ d : C^\infty(M; V_1) \rightarrow C^\infty(M; V_2), \\ K_{20} &= |g| : C^\infty(M; V_0) \rightarrow C^\infty(M; V_2), \\ K_{10} &= d : C^\infty(M; V_0) \rightarrow C^\infty(M; V_1). \end{aligned}$$

The adjoints defined as in (2.15), are

$$K_{21}^* = \delta, \quad K_{20}^* = -(g|, \quad K_{10}^* = \delta,$$

and we have

$$(2.32) \quad D_i K_{ij} = K_{ij} D_j, \quad K_{20}^* K_{21} = -2K_{10}^*,$$

as well as the adjoint identities, using that $D_k = D_k^*$.

2.8.1. Operators on Cauchy data. In addition to $K_{21\Sigma}$, $K_{21\Sigma}^\dagger$ we can define more generally $K_{ij\Sigma}$, $K_{ij\Sigma}^\dagger$, by

$$K_{ij\Sigma} := \varrho_i K_{ij} U_j, \quad K_{ij\Sigma}^\dagger := \varrho_j K_{ij}^* U_i.$$

We then have:

$$\begin{aligned} \overline{f}_2 \cdot \mathbf{q}_{I,2} K_{20\Sigma} f_0 &= \overline{K_{20\Sigma}^\dagger} f_2 \cdot \mathbf{q}_0 f_0, \\ \overline{f}_1 \cdot \mathbf{q}_1 K_{10\Sigma} f_0 &= \overline{K_{10\Sigma}^\dagger} f_1 \cdot \mathbf{q}_0 f_0, \end{aligned}$$

for $f_i \in C_c^\infty(\Sigma; V_i \otimes \mathbb{C}^2)$.

2.8.2. Imposing the TT gauge. The TT gauge is imposed by intersecting the corresponding spaces with $\text{Ker}_{\text{sc}} K_{20}^*$.

We assume in the rest of this subsection that $\Lambda \neq 0$ and we set $S_0 := -(4\Lambda)^{-1} K_{10} K_{20}^*$. We denote by $\iota : \text{Ker}_{\text{sc}} D_2 \cap \text{Ker}_{\text{sc}} K_{21}^* \cap \text{Ker}_{\text{sc}} K_{20}^* \rightarrow \text{Ker}_{\text{sc}} D_2 \cap \text{Ker}_{\text{sc}} K_{21}^*$ the canonical injection.

Proposition 2.13. *Assume that $\Lambda \neq 0$. Then the quotient map $[\iota]$:*

$$\left(\frac{\text{Ker}_{\text{sc}} D_2 \cap \text{Ker}_{\text{sc}} K_{21}^* \cap \text{Ker}_{\text{sc}} K_{20}^*}{K_{21} \text{Ker}_{\text{sc}} D_1 \cap \text{Ker}_{\text{sc}} K_{20}^*}, \mathbf{q}_{I,2} \right) \longrightarrow \left(\frac{\text{Ker}_{\text{sc}} D_2 \cap \text{Ker}_{\text{sc}} K_{21}^*}{K_{21} \text{Ker}_{\text{sc}} D_1}, \mathbf{q}_{I,2} \right)$$

is an isomorphism of Hermitian spaces with inverse

$$[\mathbb{1} - K_{21} S_0] : \frac{\text{Ker}_{\text{sc}} D_2 \cap \text{Ker}_{\text{sc}} K_{21}^*}{K_{21} \text{Ker}_{\text{sc}} D_1} \longrightarrow \frac{\text{Ker}_{\text{sc}} D_2 \cap \text{Ker}_{\text{sc}} K_{21}^* \cap \text{Ker}_{\text{sc}} K_{20}^*}{K_{21} \text{Ker}_{\text{sc}} D_1 \cap \text{Ker}_{\text{sc}} K_{20}^*}.$$

Remark 2.14. *Since $K_{20}^* K_{21} = -2K_{10}^*$ we have*

$$K_{21} \text{Ker}_{\text{sc}} D_1 \cap \text{Ker}_{\text{sc}} K_{20}^* = K_{21} (\text{Ker}_{\text{sc}} D_1 \cap \text{Ker}_{\text{sc}} K_{10}^*).$$

Proof of Prop. 2.13. Note the identities

$$(2.33) \quad D_1 S_0 = S_0 D_2, \quad K_{20}^* K_{21} S_0 = K_{20}^* + (2\Lambda)^{-1} K_{20}^* D_2,$$

obtained using (2.32) and the fact that $D_0 = K_{10}^* K_{10} - 2\Lambda$. They imply that $[\iota]$ and $[\mathbb{1} - K_{21} S_0]$ are well defined. Clearly $(\mathbb{1} - K_{21} S_0) \circ \iota = \mathbb{1}$. Moreover using (2.33) one obtains also that $\iota \circ (\mathbb{1} - K_{21} S_0) = \mathbb{1}$ modulo $K_{21} \text{Ker}_{\text{sc}} D_1$. Therefore $[\iota]$ and $[\mathbb{1} - K_{21} S_0]$ are each other's inverses. \square

2.8.3. *Phase space of Cauchy data.* We have seen in Prop. 2.6 that

$$(2.34) \quad [\varrho_2] : \left(\frac{\text{Ker}_{\text{sc}} D_2 \cap \text{Ker}_{\text{sc}} K_{21}^\dagger}{K_{21} \text{Ker}_{\text{sc}} D_1}, \mathbf{q}_{I,2} \right) \rightarrow \left(\frac{\text{Ker}_c K_{21\Sigma}^\dagger}{\text{Ran}_c K_{21\Sigma}}, \mathbf{q}_{I,2} \right)$$

is an isomorphism of Hermitian spaces. The same argument shows that

$$(2.35) \quad [\varrho_2] : \left(\frac{\text{Ker}_{\text{sc}} D_2 \cap \text{Ker}_{\text{sc}} K_{21}^* \cap \text{Ker}_{\text{sc}} K_{20}^*}{K_{21} \text{Ker}_{\text{sc}} D_1 \cap \text{Ker}_{\text{sc}} K_{20}^*}, \mathbf{q}_{I,2} \right) \\ \rightarrow \left(\frac{\text{Ker}_c K_{21\Sigma}^\dagger \cap \text{Ker}_c K_{20\Sigma}^\dagger}{\text{Ran}_c K_{21\Sigma} \cap \text{Ker}_c K_{20\Sigma}^\dagger}, \mathbf{q}_{I,2} \right)$$

is an isomorphism of Hermitian spaces.

Note that $\rho_2 S_0 U_2 := S_{0\Sigma} = (-4\Lambda)^{-1} K_{10\Sigma} K_{20\Sigma}^\dagger$. The Cauchy surface version of Prop. 2.13 is then the statement that

$$[\iota] : \left(\frac{\text{Ker}_c K_{21\Sigma}^\dagger \cap \text{Ker}_c K_{20\Sigma}^\dagger}{\text{Ran}_c K_{21\Sigma} \cap \text{Ker}_c K_{20\Sigma}^\dagger}, \mathbf{q}_{I,2} \right) \rightarrow \left(\frac{\text{Ker}_c K_{21\Sigma}^\dagger}{\text{Ran}_c K_{21\Sigma}}, \mathbf{q}_{I,2} \right)$$

is an isomorphism of Hermitian spaces with inverse

$$[\mathbb{1} - K_{21\Sigma} S_{0\Sigma}] : \frac{\text{Ker}_c K_{21\Sigma}^\dagger}{\text{Ran}_c K_{21\Sigma}} \rightarrow \frac{\text{Ker}_c K_{21\Sigma}^\dagger \cap \text{Ker}_c K_{20\Sigma}^\dagger}{\text{Ran}_c K_{21\Sigma} \cap \text{Ker}_c K_{20\Sigma}^\dagger}.$$

Definition 2.15. We set

$$\begin{aligned} \mathcal{E}_{\text{TT}} &:= \text{Ker}_c K_{21\Sigma}^\dagger \cap \text{Ker}_c K_{20\Sigma}^\dagger, \\ \mathcal{F}_{\text{TT}} &:= \text{Ran}_c K_{21\Sigma} \cap \text{Ker}_c K_{20\Sigma}^\dagger, \\ \mathcal{F}_{\text{TT,w}} &:= \text{Ran}_c K_{21\Sigma} K_{21\Sigma}^\dagger \cap \text{Ker}_c K_{20\Sigma}^\dagger, \end{aligned}$$

By the previous discussion, the physical phase space is isomorphic to $(\mathcal{E}_{\text{TT}}/\mathcal{F}_{\text{TT}}, \mathbf{q}_{I,2})$. Note that $\mathcal{F}_{\text{TT}} = K_{21\Sigma} \text{Ker}_c K_{10\Sigma}^\dagger$ by (2.32), and $\mathbf{q}_{I,2} = \mathbf{q}_2$ on \mathcal{E}_{TT} . The subspace $\mathcal{F}_{\text{TT,w}}$ represents the restricted gauge transformations relevant for the weak gauge invariance condition expressed in different forms in (2.23), (2.28) and (2.30).

2.8.4. *Hadamard states in the TT gauge.* Assume as in 2.7.3 that we have a pair of Hermitian forms $\lambda_{2\Sigma}^\pm \in L(C_0^\infty(\Sigma; V_2 \otimes \mathbb{C}^2), C_0^\infty(\Sigma; V_2 \otimes \mathbb{C}^2)^*)$.

They induce covariances on $(\mathcal{E}_{\text{TT}}/\mathcal{F}_{\text{TT}}, \mathbf{q}_{I,2})$ if

$$(2.36) \quad \begin{aligned} i) \quad & \lambda_{2\Sigma}^+ - \lambda_{2\Sigma}^- = \mathbf{q}_2 \text{ on } \mathcal{E}_{\text{TT}}, \\ ii) \quad & \lambda_{2\Sigma}^\pm = 0 \text{ on } \mathcal{E}_{\text{TT}} \times \mathcal{F}_{\text{TT}}, \\ iii) \quad & \lambda_{2\Sigma}^\pm = \lambda_{2\Sigma}^{\pm*}, \lambda_{2\Sigma}^\pm \geq 0 \text{ on } \mathcal{E}_{\text{TT}}. \end{aligned}$$

With c_2^\pm defined in (2.29) the conditions in (2.36) become:

$$(2.37) \quad \begin{aligned} & i) \quad c_2^+ + c_2^- = \mathbb{1} \text{ on } \mathcal{E}_{\text{TT}}, \\ & ii) \quad c_2^\pm : \mathcal{F}_{\text{TT}} \rightarrow \text{Ran } K_{21\Sigma} + \text{Ran } K_{20\Sigma}, \\ & iii) \quad c_2^\pm \mathbf{q}_{I,2} = \mathbf{q}_{I,2} c_2^\pm, \pm \bar{f} \cdot \mathbf{q}_{I,2} c_2^\pm f \geq 0 \text{ on } \mathcal{E}_{\text{TT}}. \end{aligned}$$

If $[c_2^\pm, I_\Sigma] = 0$ then we can replace (2.37) *iii*) by

$$(2.38) \quad iii) \quad c_2^\pm \mathbf{q}_2 = \mathbf{q}_2 c_2^\pm, \pm \bar{f} \cdot \mathbf{q}_2 c_2^\pm f \geq 0 \text{ on } \mathcal{E}_{\text{TT}},$$

since $I_\Sigma = \mathbb{1}$ on $\text{Ker}_c K_{20\Sigma}^\dagger$.

Similarly as before will call property (2.36) *ii*) or equivalently (2.37) *ii*) the *strong TT gauge invariance property*.

2.8.5. *Spacetime covariances.* We now set

$$(2.39) \quad \Lambda_{2,\text{TT}}^\pm = (\varrho_2(\mathbb{1} - K_{21}S_0)G_2)^* \lambda_{2\Sigma}^\pm (\varrho_2(\mathbb{1} - K_{21}S_0)G_2).$$

Proposition 2.16. *Assume that*

$$(2.40) \quad \begin{aligned} & i) \quad c_2^+ + c_2^- = \mathbb{1} \text{ on } \mathcal{E}_{\text{TT}}, \\ & ii) \quad c_2^\pm : \mathcal{F}_{\text{TT,w}} \rightarrow \text{Ran } K_{21\Sigma} + \text{Ran } K_{20\Sigma}, \\ & iii) \quad c_2^\pm \mathbf{q}_{I,2} = \mathbf{q}_{I,2} c_2^\pm, \pm \bar{f} \cdot \mathbf{q}_{I,2} c_2^\pm f \geq 0 \text{ on } \mathcal{E}_{\text{TT}}. \end{aligned}$$

Then $\Lambda_{2,\text{TT}}^\pm$ given by (2.39) are the covariances of a quasi-free state on $\text{CCR}(\mathcal{V}_P, \mathbf{Q}_P)$. Furthermore, if for some neighborhood \mathcal{U} of Σ in M we have:

$$(2.41) \quad iv) \quad \text{WF}(U_2 \circ c_2^\pm)' \subset (\mathcal{N}^\pm \cup \mathcal{F}) \times T^*\Sigma,$$

over $\mathcal{U} \times \Sigma$, where $\mathcal{F} \subset T^*M$ is a conic set with $\mathcal{F} \cap \mathcal{N} = \emptyset$, then the associated state is Hadamard.

We call property (2.40) *ii*) the *weak TT gauge invariance property*.

Proof. To prove the first statement of the proposition, we need to check that $\Lambda_{2,\text{TT}}^\pm$ vanish on $\text{Ker}_c K_{21}^* \times \text{Ran}_c P$, or equivalently (since $G_2 D_2 = 0$) on $\text{Ker}_c K_{21}^* \times \text{Ran}_c K_{21} K_{21}^*$. Using property (2.40) *ii*), this will hold if $(\mathbb{1} - K_{21}S_0)$ maps $K_{21} K_{21}^* \text{Ker}_{\text{sc}} D_2$ into itself.

We use (2.32) and its adjoint identity to obtain:

$$\begin{aligned} S_0 K_{21} &= (2\Lambda)^{-1} K_{10} K_{10}^* = (-4\Lambda)^{-1} K_{21}^* K_{20} K_{10}^*, \\ K_{21} S_0 K_{21} &= (-4\Lambda)^{-1} K_{21} K_{21}^* K_{20} K_{10}^*, \end{aligned}$$

which shows that $(\mathbb{1} - K_{21}S_0)$ preserves $K_{21} K_{21}^* \text{Ker}_{\text{sc}} D_2$.

To prove the second property we write

$$\Lambda_{2,\text{TT}}^\pm = (\varrho_2 G_2)^* \lambda_{2\Sigma,\text{TT}}^\pm (\varrho_2 G_2),$$

for

$$\lambda_{2\Sigma,\text{TT}}^\pm = (\mathbb{1} - K_{21\Sigma} S_{0\Sigma})^* \lambda_{2\Sigma}^\pm (\mathbb{1} - K_{21\Sigma} S_{0\Sigma}).$$

Setting $\lambda_{2\Sigma,\text{TT}}^\pm =: \pm \mathbf{q}_{I,2}^{-1} c_{2,\text{TT}}^\pm$, we have

$$c_{2,\text{TT}}^\pm = (\mathbb{1} - K_{21\Sigma} S_{0\Sigma}) \circ c_2^\pm \circ (\mathbb{1} - K_{21\Sigma} S_{0\Sigma}).$$

The operator $(\mathbb{1} - K_{21\Sigma}S_{0\Sigma})$ is a matrix of differential operators on Σ , so does not enlarge the wavefront set. On the other hand we have

$$U_2 \circ (\mathbb{1} - K_{21\Sigma}S_{0\Sigma}) = (\mathbb{1} - K_{21}S_0) \circ U_2.$$

Since $\mathbb{1} - K_{21}S_0$ is a differential operator on M , we have finally

$$\text{WF}(U_2 \circ c_{2,\text{TT}}^\pm)' \subset \text{WF}(U_2 \circ c_2^\pm)',$$

so the Hadamard condition is satisfied by $c_{2,\text{TT}}^\pm$. This completes the proof of the proposition. \square

If $[c_2^\pm, I_\Sigma] = 0$ then we can replace $\mathbf{q}_{I,2}$ by \mathbf{q}_2 in the previous proposition.

3. WICK ROTATION

3.1. Wick rotation. Let us assume that $M = I \times \Sigma$ where $I \subset \mathbb{R}$ is some open interval with $0 \in I$, Σ a smooth manifold, and denote the points of M by (t, \mathbf{x}) . We assume that g is real analytic in t and extends holomorphically to a strip $I \times i\tilde{I}$, where \tilde{I} is another open interval with $0 \in \tilde{I}$. We set $\tilde{M} = \tilde{I} \times \Sigma$, with variables (s, \mathbf{x}) .

3.1.1. Wick rotation of tensors. If $a \in \{0, \dots, d\}$ we set $\epsilon(a) = \delta_a^0$. If $A = (a_1, \dots, a_n)$ we set $\epsilon(A) = \sum_i \epsilon(a_i)$.

If $T = T^A_B(t)$ is a tensor, real analytic in t , we define the *Wick rotation* of T ,

$$\mathcal{W}T = \tilde{T},$$

by

$$\tilde{T}^A_B(s) := i^{\epsilon(B) - \epsilon(A)} T^A_B(is).$$

Wick rotation consists in replacing t by is , dt by ids , ∂_t by $-i\partial_s$, keeping the \mathbf{x} variables and the associated tensor coordinates fixed.

3.1.2. Wick rotation of metric. We set $\tilde{g} = \mathcal{W}g$. In general \tilde{g} is only a symmetric complex metric. Taking \tilde{I} small enough we can assume that \tilde{g} is non-degenerate. We denote by $\tilde{\nabla}$ the associated covariant derivative. Then, computing the Christoffel symbols, one easily checks that

$$\tilde{\nabla}_{\tilde{X}} \tilde{T} = \widetilde{\nabla_X T}.$$

Similarly we have

$$\text{R}(\tilde{g}) = \widetilde{\text{R}(g)},$$

where $\text{R}(g), \text{R}(\tilde{g})$ denotes the Riemann tensor for g, \tilde{g} .

The Wick rotation \mathcal{W} commutes with tensor multiplication, action of tensors on tensors, raising and lowering indices, summation over repeated indices, etc.

We set for $k = 0, 1, 2$:

$$\tilde{V}_k = \mathbb{C} \otimes_s^k T^* \tilde{M},$$

and use the sesquilinear forms $(\cdot | \cdot)_{\tilde{V}_k}$ on the fibers of \tilde{V}_k defined with \tilde{g} instead of g . In general $(\cdot | \cdot)_{\tilde{V}_k}$ are not Hermitian.

We have

$$\mathcal{W}(g|u)_{V_2} = (\tilde{g}|\mathcal{W}u)_{\tilde{V}_2},$$

hence

$$\mathcal{W} \circ I = \tilde{I} \circ \mathcal{W}, \text{ for } \tilde{I}\tilde{u} = \tilde{u} - \frac{1}{4}(\tilde{g}|_{\tilde{V}_2})\tilde{g}.$$

3.1.3. Wick rotation of differential operators. If D is a differential operator acting on tensors, with coefficients analytic in t , we define its Wick rotation \tilde{D} by

$$\tilde{D}\tilde{T} := \widetilde{DT}.$$

3.2. Traces on Σ . Let us assume additionally that $g = -dt^2 + h(t, \mathbf{x})dx^2$ where $t \mapsto h(t, \cdot)dx^2$ is analytic with values in Riemannian metrics on Σ , so $\tilde{g} = ds^2 + h(is, \mathbf{x})dx^2$. We identify Σ with $\{0\} \times \Sigma$.

We will use the decomposition of $(0, k)$ -tensors on M explained in 2.1.3. Using this decomposition the trace operators ϱ_k become

$$\varrho_k u_k = \begin{pmatrix} u_k|_{\Sigma} \\ \mathbf{i}^{-1}\partial_t u_k|_{\Sigma} \end{pmatrix}, \quad u_k \in C^\infty(M; V_k).$$

We use the same decomposition for $(0, k)$ -tensors on \tilde{M} and define for $\tilde{u} \in C^\infty(\tilde{M}; \tilde{V}_i)$:

$$\tilde{\varrho}_k \tilde{u}_k = \begin{pmatrix} \tilde{u}_k|_{\Sigma} \\ -\partial_s \tilde{u}_k|_{\Sigma} \end{pmatrix}, \quad \tilde{u}_k \in C^\infty(\tilde{M}; \tilde{V}_k).$$

If u_k is real analytic in t then

$$\tilde{\varrho}_k \mathcal{W} u_k = \mathcal{F}_{k\Sigma} \varrho_k u_k,$$

where

$$(3.1) \quad \mathcal{F}_{k\Sigma} := \begin{pmatrix} \mathcal{F}_k & 0 \\ 0 & \mathcal{F}_k \end{pmatrix}, \quad \mathcal{F}_0 = 1, \quad \mathcal{F}_1 = \begin{pmatrix} \mathbf{i} & 0 \\ 0 & \mathbb{1} \end{pmatrix}, \quad \mathcal{F}_2 = \begin{pmatrix} -1 & 0 & 0 \\ 0 & \mathbf{i} & 0 \\ 0 & 0 & \mathbb{1} \end{pmatrix}.$$

If $A_{ij\Sigma}$ is a linear operator from $C^\infty(\Sigma; V_j \otimes \mathbb{C}^2)$ to $C^\infty(\Sigma; V_i \otimes \mathbb{C}^2)$ we set

$$(3.2) \quad \tilde{A}_{ij\Sigma} := \mathcal{F}_{i\Sigma} \circ A_{ij\Sigma} \circ \mathcal{F}_{j\Sigma}^{-1}.$$

If we define \tilde{I}_Σ by $\tilde{I}_\Sigma \tilde{\varrho}_2 \tilde{u}_2 := \tilde{\varrho}_2 \tilde{I} \tilde{u}_2$, then this notation is coherent, i.e.

$$\tilde{I}_\Sigma = \mathcal{F}_{2\Sigma} \circ I_\Sigma \circ \mathcal{F}_{2\Sigma}^{-1}.$$

4. CALDERÓN PROJECTORS

In this section we study Calderón projectors for elliptic operators on a compact manifold.

4.1. Background.

4.1.1. *Extendible distributions.* Let \widetilde{M} be a compact manifold, and $\widetilde{V} \xrightarrow{\pi} \widetilde{M}$ a finite rank Hilbertian vector bundle. We equip \widetilde{M} with a smooth density. If \widetilde{M} is equipped with a Riemannian metric \widetilde{g} , one chooses the induced density on \widetilde{M} . We denote by $C^\infty(\widetilde{M}; \widetilde{V})$ the smooth sections of \widetilde{V} and by $(\cdot|\cdot)_{\widetilde{M}}$ the scalar product on $C^\infty(\widetilde{M}; \widetilde{V})$ obtained from the fiber scalar product $(\cdot|\cdot)_{\widetilde{V}}$ and the density on \widetilde{M} . Concretely we have

$$(u|u)_{\widetilde{M}} = \int_{\widetilde{M}} (u(x)|u(x))_{\widetilde{V}} dx, \quad u \in C^\infty(\widetilde{M}; \widetilde{V}).$$

If $\Omega \subset \widetilde{M}$ is an open set with smooth boundary, we denote by $(\cdot|\cdot)_\Omega$ the analog scalar products on $C_0^\infty(\Omega; \widetilde{V})$.

If $\mathcal{F}(\widetilde{M}; \widetilde{V}) \subset \mathcal{D}'(\widetilde{M}; \widetilde{V})$ we denote by $\overline{\mathcal{F}}(\Omega; \widetilde{V}) \subset \mathcal{D}'(\Omega; \widetilde{V})$ the image of $\mathcal{F}(\widetilde{M}; \widetilde{V})$ by the restriction map $\mathcal{D}'(\widetilde{M}; \widetilde{V}) \rightarrow \mathcal{D}'(\Omega; \widetilde{V})$. For example $\overline{\mathcal{D}'}(\Omega; \widetilde{V})$ is the space of extendible distributions on Ω , while $\overline{C^\infty}(\Omega; \widetilde{V})$ is the space of smooth sections of \widetilde{V} over Ω bounded with all derivatives.

It is known that any element of $\overline{\mathcal{D}'}(\Omega; \widetilde{V})$ can be extended by 0 in $\widetilde{M} \setminus \Omega^{\text{cl}}$, i.e. is the restriction to Ω of a distribution in $\mathcal{D}'(\widetilde{M}; \widetilde{V})$ supported in Ω^{cl} .

4.1.2. *Hypersurfaces.* Let $\Sigma \subset \widetilde{M}$ be a smooth compact hypersurface such that $\widetilde{M} \setminus \Sigma$ is the union of two disjoint open sets Ω^+ and Ω^- . We equip Σ with a smooth density dx , equal to the induced Riemannian density if \widetilde{M} is equipped with a Riemannian metric \widetilde{g} .

We can consider the Hilbertian vector bundle $\widetilde{V} \otimes \mathbb{C}^2 \xrightarrow{\pi} \Sigma$ and denote by $(\cdot|\cdot)_\Sigma$ the associated scalar product on $C^\infty(\Sigma; \widetilde{V} \otimes \mathbb{C}^2)$.

We embed $C^\infty(\widetilde{M}; \widetilde{V})$ into $\mathcal{D}'(\widetilde{M}; \widetilde{V})$ and $C^\infty(\Sigma; \widetilde{V} \otimes \mathbb{C}^2)$ into $\mathcal{D}'(\Sigma; \widetilde{V} \otimes \mathbb{C}^2)$ using the above scalar products.

We denote by $H^s(\widetilde{M}; \widetilde{V}), H^s(\Sigma; \widetilde{V} \otimes \mathbb{C}^2)$ the canonical Sobolev spaces.

4.1.3. *Trace operators.* We fix a vector field ν transverse to Σ pointing out of Ω^+ and set for $u \in C^\infty(\widetilde{M}; \widetilde{V})$:

$$\tilde{\varrho}u := \begin{pmatrix} u|_\Sigma \\ \nu u|_\Sigma \end{pmatrix},$$

and for $u \in \overline{C^\infty}(\Omega^\pm; \widetilde{V})$:

$$\tilde{\varrho}^\pm u := \begin{pmatrix} u|_\Sigma \\ \nu u|_\Sigma \end{pmatrix}.$$

If $(\widetilde{M}, \widetilde{g})$ is Riemannian, we take ν equal to the exterior unit normal vector field to Ω^+ .

We denote by $\tilde{\varrho}^* : \mathcal{D}'(\Sigma; \widetilde{V} \otimes \mathbb{C}^2) \rightarrow \mathcal{D}'(\widetilde{M}; \widetilde{V})$ the adjoint of the operator $\tilde{\varrho} : C^\infty(\widetilde{M}; \widetilde{V}) \rightarrow C^\infty(\Sigma; \widetilde{V} \otimes \mathbb{C}^2)$ for the above scalar products.

4.1.4. *Green's formulas.* Let $\tilde{D} = \tilde{D}(x, \partial_x)$ be a second order differential operator acting on $C^\infty(\widetilde{M}; \widetilde{V})$. For simplicity we assume that its principal symbol $\tilde{p}(x, \xi)$ is scalar.

We can associate to \tilde{D} a sesquilinear form $\mathbf{Q}_{\widetilde{M}}$ on $C^\infty(\widetilde{M}; \widetilde{V})$, bounded on $H^1(\widetilde{M}; \widetilde{V})$ such that $(u|\tilde{D}v)_{\widetilde{M}} = \mathbf{Q}(u, v)_{\widetilde{M}}$ for $u, v \in C^\infty(\widetilde{M}; \widetilde{V})$. In local

coordinates we have

$$(4.1) \quad Q(u, v)_{\widetilde{M}} = \int_{\widetilde{M}} \sum_{|\alpha|, |\beta| \leq 1} (a_{\alpha, \beta}(x) \partial_x^\alpha u | \partial_x^\beta v)_{\widetilde{V}} dx,$$

for $a_{\alpha, \beta} \in C^\infty(\widetilde{M}; L(\widetilde{V}))$. For $u, v \in \overline{C^\infty}(\Omega^\pm; \widetilde{V})$ we denote by $Q(u, v)_{\Omega^\pm}$ the analogous sesquilinear forms obtained by integrating only on Ω^\pm .

Lemma 4.1. (1) *There exists $\tilde{\sigma} \in C^\infty(\Sigma; L(\widetilde{V} \otimes \mathbb{C}^2))$ such that*

$$(u | \tilde{D}v)_{\Omega^\pm} - (\tilde{D}^*u | v)_{\Omega^\pm} = \pm (\tilde{\rho}^\pm u | \tilde{\sigma} \tilde{\rho}^\pm v)_\Sigma, \quad u, v \in \overline{C^\infty}(\Omega^\pm; \widetilde{V}).$$

(2) *There exists $\tilde{q} \in C^\infty(\Sigma; L_h(\widetilde{V} \otimes \mathbb{C}^2))$ such that for all $u, v \in \overline{C^\infty}(\Omega^\pm; \widetilde{V})$,*

$$(u | \tilde{D}v)_{\Omega^\pm} + (\tilde{D}u | v)_{\Omega^\pm} = Q(u, v)_{\Omega^\pm} + \overline{Q}(v, u)_{\Omega^\pm} \mp (\tilde{\rho}^\pm u | \tilde{q} \tilde{\rho}^\pm v)_\Sigma.$$

(3) *If $[\tilde{D} - \tilde{D}^*, \mathbb{1}_{\Omega^+}] = 0$, then $\tilde{\sigma} \in C^\infty(\Sigma; L_a(\widetilde{V} \otimes \mathbb{C}^2))$.*

(4) *If Σ is non characteristic for \tilde{D} , then $\tilde{\sigma}$ is non-degenerate.*

Proof. Let us fix two extension maps $e_{\Omega^\pm} : \overline{C^\infty}(\Omega^\pm; \widetilde{V}) \rightarrow C^\infty(\widetilde{M}; \widetilde{V})$. Let us first prove (1). Since \tilde{D} is a local operator, we have:

$$(4.2) \quad \begin{aligned} (u | \tilde{D}v)_{\Omega^\pm} - (\tilde{D}^*u | v)_{\Omega^\pm} &= (e_{\Omega^\pm} u | \mathbb{1}_{\Omega^\pm} \tilde{D} e_{\Omega^\pm} v)_{\widetilde{M}} - (\tilde{D}^* e_{\Omega^\pm} u | \mathbb{1}_{\Omega^\pm} e_{\Omega^\pm} v)_{\widetilde{M}} \\ &= (e_{\Omega^\pm} u | [\mathbb{1}_{\Omega^\pm}, \tilde{D}] e_{\Omega^\pm} v)_{\widetilde{M}}. \end{aligned}$$

Computing in local coordinates near Σ we obtain that

$$(4.3) \quad (e_{\Omega^\pm} u | [\mathbb{1}_{\Omega^\pm}, \tilde{D}] e_{\Omega^\pm} v)_{\widetilde{M}} = (\tilde{\rho}^\pm u | \tilde{\sigma}^\pm \tilde{\rho}^\pm v)_\Sigma,$$

for some $\tilde{\sigma}^\pm \in L(C^\infty(\Sigma; \widetilde{V} \otimes \mathbb{C}^2))$. If $f, g \in C^\infty(\Sigma; \widetilde{V} \otimes \mathbb{C}^2)$ we pick $u, v \in C^\infty(\widetilde{M}; \widetilde{V})$ such that $\tilde{\rho}^\pm u = f$ and $\tilde{\rho}^\pm v = g$. Since

$$\begin{aligned} (u | \tilde{D}v)_{\Omega^+} - (\tilde{D}^*u | v)_{\Omega^+} + (u | \tilde{D}v)_{\Omega^-} - (\tilde{D}^*u | v)_{\Omega^-} \\ = (u | \tilde{D}v)_{\widetilde{M}} - (\tilde{D}^*u | v)_{\widetilde{M}} = 0, \end{aligned}$$

we obtain that $(f | (\tilde{\sigma}^+ + \tilde{\sigma}^-)g)_\Sigma = 0$, which proves (1). The fact that $\tilde{\sigma} := \tilde{\sigma}^+$ is anti-selfadjoint if $[\mathbb{1}_{\Omega^+}, \tilde{D} - \tilde{D}^*] = 0$ is immediate, using (4.2). This proves (3).

Let us now prove (2). We have

$$\begin{aligned} (u | \tilde{D}v)_{\Omega^\pm} + (\tilde{D}u | v)_{\Omega^\pm} &= (\mathbb{1}_{\Omega^\pm} e_{\Omega^\pm} u | \tilde{D} e_{\Omega^\pm} v)_{\widetilde{M}} + (\tilde{D} e_{\Omega^\pm} u | \mathbb{1}_{\Omega^\pm} e_{\Omega^\pm} v)_{\widetilde{M}} \\ &= Q(\mathbb{1}_{\Omega^\pm} u, v)_{\widetilde{M}} + \overline{Q}(\mathbb{1}_{\Omega^\pm} v, u)_{\widetilde{M}}. \end{aligned}$$

Computing in local coordinates using (4.1) we obtain that

$$Q(\mathbb{1}_{\Omega^\pm} u, v)_{\widetilde{M}} + \overline{Q}(\mathbb{1}_{\Omega^\pm} v, u)_{\widetilde{M}} = Q(u, v)_{\Omega^\pm} + \overline{Q}(v, u)_{\Omega^\pm} + (\tilde{\rho}^\pm u | \tilde{q}^\pm \tilde{\rho}^\pm v)_\Sigma$$

for some $\tilde{q}^\pm \in C^\infty(\Sigma; L(\widetilde{V} \otimes \mathbb{C}^2))$. For $v = u$ all the terms above are real hence \tilde{q}^\pm are Hermitian. To show that $\tilde{q}^- = -\tilde{q}^+$ we argue as in the proof of (1).

To prove (4) it suffices to work in local coordinates (x^0, x') in which $\Omega^+ = \{x^0 > 0\}$. One can choose $\nu = -\partial_{x^0}$ and check that, modulo a non-vanishing factor coming from the densities on \widetilde{M} and Σ , $\tilde{\sigma}$ is invertible if $p(0, x', dx^0) \neq 0$, where $p(x, \xi) \mathbb{1}_{\widetilde{V}}$ is the (scalar) principal symbol of \tilde{D} . \square

4.2. Calderón projectors. The definition of Calderón projectors for \tilde{D} is standard if \tilde{D} is invertible, we are however interested in generalizations when \tilde{D} is merely Fredholm. We first start with the former case, and discuss the more general case subsequently.

4.2.1. Calderón projectors. Let us assume from now on that \tilde{D} is elliptic. This is in particular the case if $p(x, \xi) = \xi \cdot \tilde{g}^{-1}(x) \xi$. Then $\tilde{D} + \lambda$ is invertible for $\lambda \gg 1$ hence \tilde{D} is Fredholm of index 0.

Definition 4.2. Assume that $\tilde{D} : H^2(\tilde{M}; \tilde{V}) \rightarrow L^2(\tilde{M}; \tilde{V})$ is invertible. By ellipticity $\tilde{D} : H^s(\tilde{M}; \tilde{V}) \rightarrow H^{s-2}(\tilde{M}; \tilde{V})$ is invertible for any $s \in \mathbb{R}$. One defines the Calderón projectors \tilde{c}^\pm by

$$\tilde{c}^\pm f := \mp \tilde{\varrho}^\pm \tilde{D}^{-1} \tilde{\varrho}^* \tilde{\sigma} f, \quad f \in C^\infty(\Sigma; \tilde{V} \otimes \mathbb{C}^2).$$

Proposition 4.3. The Calderón projectors \tilde{c}^\pm satisfy:

- i) $\tilde{c}^+ + \tilde{c}^- = \mathbb{1}$,
- ii) $(\tilde{c}^\pm)^2 = \tilde{c}^\pm$,
- iii) for $u \in \mathcal{D}'(\tilde{M}; \tilde{V})$, $\tilde{D}u = 0$ in $\Omega^\pm \Rightarrow \tilde{\varrho}^\pm u = \tilde{c}^\pm \tilde{\varrho}^\pm u$.

We include the proof for reference.

Proof. Let $f, g \in C^\infty(\Sigma; \tilde{V} \otimes \mathbb{C}^2)$. We fix $v \in C^\infty(\tilde{M}; \tilde{V})$ such that $\tilde{\varrho}v = g$ and set $w = \tilde{D}^{-1} \tilde{\varrho}^* \tilde{\sigma} f$, $u^\pm = \mp w|_{\Omega^\pm}$. Then by Lemma 4.1:

$$\begin{aligned} (g|\tilde{\sigma}(\tilde{c}^+ + \tilde{c}^-)f)_\Sigma &= (\tilde{\varrho}^+ v|\tilde{\sigma}\tilde{\varrho}^+ u^+)_\Sigma + (\tilde{\varrho}^- v|\tilde{\sigma}\tilde{\varrho}^- u^-)_\Sigma \\ &= (v|\tilde{D}u^+)_\Omega^+ - (\tilde{D}^*v|u^+)_\Omega^+ - (v|\tilde{D}u^-)_\Omega^- + (\tilde{D}^*v|u^-)_\Omega^- \\ &= (\tilde{D}^*v|w)_{\tilde{M}} = (v|\tilde{\varrho}^* \tilde{\sigma} f)_{\tilde{M}} = (\tilde{\varrho}v|\tilde{\sigma} f)_\Sigma = (g|\tilde{\sigma} f)_\Sigma. \end{aligned}$$

Since $\tilde{\sigma}$ is non-degenerate, this proves *i*). Let us now prove *ii*). For $f \in C^\infty(\Sigma; \tilde{V} \otimes \mathbb{C}^2)$ we set $v = \tilde{D}^{-1} \tilde{\varrho}^* \tilde{\sigma} f$, so that $\mp \tilde{\varrho}^\pm v = \tilde{c}^\pm f$. For $u \in C^\infty(\tilde{M}; \tilde{V})$ we have

$$(u|\tilde{D}\mathbb{1}_{\Omega^\pm}v)_{\tilde{M}} = (u|[\tilde{D}, \mathbb{1}_{\Omega^\pm}]v)_{\tilde{M}} = \mp (u|\tilde{\varrho}^* \tilde{\sigma} \tilde{\varrho}^\pm v)_{\tilde{M}},$$

using (4.3) and the fact that $\tilde{D}v = 0$ in Ω^\pm . Therefore $\tilde{D}\mathbb{1}_{\Omega^\pm}v = \mp \tilde{\varrho}^* \tilde{\sigma} \tilde{\varrho}^\pm v = \tilde{\varrho}^* \tilde{\sigma} \tilde{c}^\pm f$. Since \tilde{D} is invertible this implies that $\mathbb{1}_{\Omega^\pm}v = \tilde{D}^{-1} \tilde{\varrho}^* \tilde{\sigma} \tilde{c}^\pm f$ hence $\tilde{c}^\pm f = \tilde{c}^\pm \circ \tilde{c}^\pm f$ by applying $\tilde{\varrho}^\pm$ to the above identity.

Finally $\tilde{D}\mathbb{1}_{\Omega^\pm}u = \mp \tilde{\varrho}^* \tilde{\sigma} \tilde{\varrho}^\pm u$, hence $\mathbb{1}_{\Omega^\pm}u = \mp \tilde{D}^{-1} \tilde{\varrho}^* \tilde{\sigma} \tilde{\varrho}^\pm u$, which implies that $\tilde{c}^\pm \tilde{\varrho}^\pm u = \tilde{\varrho}^\pm u$ and proves *iii*). \square

4.2.2. Selfadjointness of Calderón projectors. Let us now assume that there exist a smooth involution $\chi : \tilde{M} \rightarrow \tilde{M}$ with

$$\chi : \Omega^\pm \rightarrow \Omega^\mp, \quad \chi = \mathbb{1} \text{ on } \Sigma,$$

Without loss of generality we can assume that the density dx is invariant under χ .

We interpret χ as a reflection in Σ and we assume that χ lifts to a bundle involution $\kappa : \tilde{V} \rightarrow \tilde{V}$ preserving the fiber scalar product. Since $\chi = \mathbb{1}$ over Σ , κ preserves the fibers of \tilde{V} over Σ ; we denote by κ_Σ its restriction to

\tilde{V} over Σ . To define the trace operator \tilde{q} we can choose a vector field ν such that $\chi^*\nu = -\nu$. It follows that

$$(4.4) \quad \tilde{q}^\pm \kappa u = \begin{pmatrix} \kappa_\Sigma & 0 \\ 0 & -\kappa_\Sigma \end{pmatrix} \tilde{q}^\mp u.$$

Proposition 4.4. *Assume that \tilde{D} satisfies the following reflection property*

$$(4.5) \quad \tilde{D}^* = \kappa \tilde{D} \kappa.$$

Then

- (1) $q := \tilde{\sigma} \circ \begin{pmatrix} \kappa_\Sigma & 0 \\ 0 & -\kappa_\Sigma \end{pmatrix}$ satisfies $q = q^*$;
- (2) one has $\tilde{c}^{\pm*} q = q \tilde{c}^\pm$.

In concrete applications, when \tilde{D} is obtained by a Wick rotation, the operator q corresponds to the Lorentzian charge, see 4.4.2 below.

Proof. Let $u, v \in C^\infty(\tilde{M}; \tilde{V})$. Then using Lemma 4.1 and the fact that $\tilde{\sigma} = -\tilde{\sigma}^*$ we obtain

$$\begin{aligned} (u|\kappa \tilde{D} \kappa v)_{\Omega^+} - (\kappa \tilde{D}^* \kappa u|v)_{\Omega^+} &= (\tilde{q}^+ u|\tilde{\sigma} \tilde{q}^+ v)_\Sigma \\ &= (\kappa u|\tilde{D} \kappa v)_{\Omega^-} - (\tilde{D}^* \kappa u|\kappa v)_{\Omega^-} = -(\tilde{q}^- \kappa u|\tilde{\sigma} \tilde{q}^- \kappa u)_\Sigma, \end{aligned}$$

which implies that $q = q^*$ since $\tilde{q}^\pm = \tilde{q}$ on $C^\infty(\tilde{M}; \tilde{V})$.

To prove 2) we repeat the proof in [14, Prop 6.2] for the reader's convenience. For $f, g \in C^\infty(\Sigma; \tilde{V} \otimes \mathbb{C}^2)$ we set

$$u = -(\tilde{D}^{-1} \tilde{q}^* \tilde{\sigma} f)|_{\Omega^+}, \quad v = -(\kappa \tilde{D}^{-1} \tilde{q}^* \tilde{\sigma} g)|_{\Omega^+}.$$

Since $\tilde{D}u = \tilde{D}^*v = 0$ in Ω^+ , we deduce from Lemma 4.1 that $(\tilde{q}^+ v|\tilde{\sigma} \tilde{q}^+ u)_\Sigma = 0$.

Using (4.4) we obtain

$$\tilde{q}^+ u = \tilde{c}^+ f, \quad \tilde{q}^+ v = \begin{pmatrix} \kappa_\Sigma & 0 \\ 0 & -\kappa_\Sigma \end{pmatrix} \tilde{c}^- g.$$

Then $(\tilde{q}^+ v|\tilde{\sigma} \tilde{q}^+ u)_\Sigma = 0$ implies that $c^{-*} q \tilde{c}^+ = 0$ hence $c^{+*} q = q \tilde{c}^+$ as claimed. \square

4.2.3. Calderón projectors in the non invertible case I. Let us assume now that $\text{Ker } \tilde{D} \neq \{0\}$. Since \tilde{D} is Fredholm, $\text{Ran } \tilde{D} = (\text{Ker } \tilde{D}^*)^\perp$, and we can define

$$\tilde{c}^\pm f := \mp \tilde{q}^\pm \tilde{D}^{-1} \tilde{q}^* \tilde{\sigma} f$$

if $\tilde{\sigma} f \in (\tilde{q}^* \text{Ker } \tilde{D}^*)^\perp$, where $\tilde{D}^{-1} : \text{Ran } \tilde{D} \rightarrow L^2(\tilde{M}; \tilde{V}) \ominus \text{Ker } \tilde{D}$.

Assume that there exist a reflection κ on \tilde{V} such that (4.5) holds. Then it is easy to check that

$$(4.6) \quad \tilde{q}^* \tilde{\sigma} f \in (\text{Ker } \tilde{D}^*)^\perp \text{ iff } f \in (\tilde{q} \text{Ker } \tilde{D})^{\mathbf{q}},$$

where $F^{\mathbf{q}}$ denotes the \mathbf{q} -orthogonal of F and the Hermitian form \mathbf{q} is

$$\bar{f} \cdot \mathbf{q} f = (u|qu)_{\tilde{V}(\Sigma) \otimes \mathbb{C}^2}.$$

Note that since $\text{Ker } \tilde{D} \subset C^\infty(\tilde{M}; \tilde{V})$ we have $\tilde{q} = \tilde{q}^\pm$ on $\text{Ker } \tilde{D}$.

Proposition 4.5. *We have:*

- (1) $\tilde{\varrho} \text{Ker } \tilde{D} \subset (\tilde{\varrho} \text{Ker } \tilde{D})^{\mathfrak{q}}$ and $[\mathfrak{q}]$ is non-degenerate on $\frac{(\tilde{\varrho} \text{Ker } \tilde{D})^{\mathfrak{q}}}{\tilde{\varrho} \text{Ker } \tilde{D}}$,
- (2) the maps

$$[\tilde{c}^{\pm}] : \frac{(\tilde{\varrho} \text{Ker } \tilde{D})^{\mathfrak{q}}}{\tilde{\varrho} \text{Ker } \tilde{D}} \rightarrow \frac{(\tilde{\varrho} \text{Ker } \tilde{D})^{\mathfrak{q}}}{\tilde{\varrho} \text{Ker } \tilde{D}}$$

are well defined,

- (3) $[\tilde{c}^+] + [\tilde{c}^-] = \mathbb{1}$ and $[\tilde{c}^{\pm}]^2 = [\tilde{c}^{\pm}]$.

Proof. Let $u \in \text{Ker } \tilde{D}$, $f = \tilde{\varrho}u$. Then $\tilde{D}\mathbb{1}_{\Omega^{\pm}}u = \mp \tilde{\varrho}^* \tilde{\sigma} f$, hence $\tilde{\varrho}^* \tilde{\sigma} f \in (\text{Ker } \tilde{D}^*)^{\perp}$ and thus $f \in (\text{Ker } \tilde{D})^{\mathfrak{q}}$. Since \tilde{D} is elliptic, $\tilde{\sigma}$ and hence \mathfrak{q} are non-degenerate. Since $\tilde{\varrho} \text{Ker } \tilde{D}$ is finite-dimensional we have $((\tilde{\varrho} \text{Ker } \tilde{D})^{\mathfrak{q}})^{\mathfrak{q}} = \tilde{\varrho} \text{Ker } \tilde{D}$, so $[\mathfrak{q}]$ is non-degenerate.

Moreover $\mathbb{1}_{\Omega^{\pm}}u = \mp \tilde{D}^{-1} \tilde{\varrho}^* \tilde{\sigma} f + v^{\pm}$ for $v^{\pm} \in \text{Ker } \tilde{D}$ hence $f = \tilde{c}^{\pm} f + \tilde{\varrho}v^{\pm}$, so \tilde{c}^{\pm} preserves $\tilde{\varrho} \text{Ker } \tilde{D}$. To complete the proof of (2) we need to show that \tilde{c}^{\pm} preserves $(\tilde{\varrho} \text{Ker } \tilde{D})^{\mathfrak{q}}$. If $f \in (\tilde{\varrho} \text{Ker } \tilde{D})^{\mathfrak{q}}$ and $v = \tilde{D}^{-1} \tilde{\varrho}^* \tilde{\sigma} f$, $u \in \text{Ker } \tilde{D}$ then $\tilde{D}v = \tilde{D}^* \kappa u = 0$ in Ω^{\pm} so by Green's formula we have $(\tilde{\varrho}^{\mp} \kappa u | \tilde{\sigma} \tilde{\varrho}^{\pm} v)_{\Sigma} = \mp (\tilde{\varrho}u | \tilde{\varrho} \tilde{c}^{\pm} f)_{\Sigma} = 0$, so \tilde{c}^{\pm} preserves $(\tilde{\varrho} \text{Ker } \tilde{D})^{\mathfrak{q}}$.

Let us now prove (3). We have $\tilde{c}^+ + \tilde{c}^- = \mathbb{1}$ on $(\tilde{\varrho} \text{Ker } \tilde{D})^{\mathfrak{q}}$ by the same proof as in the invertible case. This proves the first statement of (3). The second statement is proved as before, except at the end: from $\tilde{D}\mathbb{1}_{\Omega^{\pm}}v = \tilde{\varrho}^* \tilde{\sigma} \tilde{c}^{\pm} f$ we obtain $\mathbb{1}_{\Omega^{\pm}}v = \tilde{D}^{-1} \tilde{\varrho}^* \tilde{\sigma} \tilde{c}^{\pm} f$ modulo $\text{Ker } \tilde{D}$ hence $\tilde{c}^{\pm} f = (\tilde{c}^{\pm})^2 f$ modulo $\tilde{\varrho} \text{Ker } \tilde{D}$. \square

4.2.4. *Subspaces of traces.* Let us assume the hypotheses in 4.2.3 and let

$$E^{\pm} = \{\tilde{\varrho}^{\pm} u : u \in \overline{C^{\infty}(\Omega^{\pm}; \tilde{V})}, \tilde{D}u = 0 \text{ in } \Omega^{\pm}\}.$$

Lemma 4.6. *We have:*

- (1) $E^+ \cap E^- = (\tilde{\varrho} \text{Ker } \tilde{D})$,
- (2) $E^+ + E^- = (\tilde{\varrho} \text{Ker } \tilde{D})^{\mathfrak{q}}$.

Proof. If $f = \tilde{\varrho}^{\pm} u^{\pm}$ with $\tilde{D}u^{\pm} = 0$ in Ω^{\pm} and $u = \mathbb{1}_{\Omega^+} u^+ + \mathbb{1}_{\Omega^-} u^-$ then $\tilde{D}u = 0$ and $f = \tilde{\varrho}u$, which proves (1).

Next let $f \in E^+$ with $f = \tilde{\varrho}^+ u$ and $\tilde{D}u = 0$ in Ω^+ . If $e_{\Omega^-} u$ is the extension of u by 0 in Ω^- we have $\tilde{D}e_{\Omega^-} u = -\tilde{\varrho}^* \tilde{\sigma} f$ hence $f \in (\tilde{\varrho} \text{Ker } \tilde{D})^{\mathfrak{q}}$, which shows that $E^+ \subset (\tilde{\varrho} \text{Ker } \tilde{D})^{\mathfrak{q}}$. The same argument shows that $E^- \subset (\tilde{\varrho} \text{Ker } \tilde{D})^{\mathfrak{q}}$. Finally if $f \in (\tilde{\varrho} \text{Ker } \tilde{D})^{\mathfrak{q}}$ we have $\tilde{c}^{\pm} f = \mp \tilde{\varrho}^{\pm} \tilde{D}^{-1} \tilde{\varrho}^* \tilde{\sigma} f$ so $\tilde{c}^{\pm} f \in E^{\pm}$. Since $\tilde{c}^+ + \tilde{c}^- = \mathbb{1}$ on $(\tilde{\varrho} \text{Ker } \tilde{D})^{\mathfrak{q}}$ this proves (2). \square

4.2.5. *Calderón projectors in the non-invertible case II.* Let us give another generalization of Calderón projectors in the non-invertible case. We fix a smoothing operator $r_{-\infty} : \mathcal{D}'(\tilde{M}; \tilde{V}) \rightarrow C^{\infty}(\tilde{M}; \tilde{V})$ such that $\tilde{D} + r_{-\infty} : H^s(\tilde{M}; \tilde{V}) \rightarrow H^{s-2}(\tilde{M}; \tilde{V})$ is invertible. If \tilde{D} is selfadjoint, we can take $r_{-\infty} = \mathbb{1}_{\{0\}}(\tilde{D})$. We set

$$(4.7) \quad \tilde{c}^{\pm} f := \mp \tilde{\varrho}^{\pm} (\tilde{D} + r_{-\infty})^{-1} \tilde{\varrho}^* \tilde{\sigma} f, \quad f \in C^{\infty}(\Sigma; \tilde{V} \otimes \mathbb{C}^2).$$

Proposition 4.7. *One has*

$$\hat{c}^+ + \hat{c}^- = \mathbb{1}.$$

Note that \hat{c}^\pm are not projections anymore, one merely has that $(\hat{c}^\pm)^2 - \hat{c}^\pm$ is smoothing.

Proof. We use the same notation as in the proof of Prop. 4.3 and obtain that

$$(g|\tilde{\sigma}(\hat{c}^+ + \hat{c}^-)f)_\Sigma = (v|\tilde{D}u^+)_{\Omega^+} - (v|\tilde{D}u^-)_{\Omega^-} + (v|\tilde{D}w)_{\tilde{M}}.$$

Since \tilde{D} is local we have

$$\begin{aligned} \tilde{D}u^\pm &= \mp \tilde{D}w|_{\Omega^\pm} = \mp(\tilde{\varrho}^*\tilde{\sigma}f - r_{-\infty}(\tilde{D} + r_{-\infty})^{-1}\tilde{\varrho}^*\tilde{\sigma}f)|_{\Omega^\pm} \\ &= \pm r_{-\infty}(\tilde{D} + r_{-\infty})^{-1}\tilde{\varrho}^*\tilde{\sigma}f|_{\Omega^\pm}. \end{aligned}$$

Since

$$\tilde{D}w = \tilde{\varrho}^*\tilde{\sigma}f - r_{-\infty}(\tilde{D} + r_{-\infty})^{-1}\tilde{\varrho}^*\tilde{\sigma}f,$$

we obtain:

$$(g|\tilde{\sigma}(\hat{c}^+ + \hat{c}^-)f)_\Sigma = (v|\tilde{\varrho}^*\tilde{\sigma}f)_{\tilde{M}} = (g|\tilde{\sigma}f)_\Sigma,$$

which proves the proposition. \square

4.3. Invariance under symmetries. Let us assume that (\tilde{M}, \tilde{g}) is Riemannian, that \tilde{V} equals $\mathbb{C} \otimes_{\mathbb{S}}^k T^*\tilde{M}$ for $k = 1, 2$ and that $\tilde{D} = \tilde{D}_{k,L} - 2\Lambda$ for some $\Lambda \in \mathbb{R}$, where $\tilde{D}_{k,L}$ are the Lichnerowicz operators for \tilde{g} (see Subsect. 2.4).

We assume that \tilde{X} is a Killing vector field of (\tilde{M}, \tilde{g}) and we denote by $\tilde{\phi}(\cdot)$ the associated flow on $C^\infty(\tilde{M}; \tilde{V})$. We define its generator \tilde{L} by

$$\tilde{\phi}(\sigma) =: e^{i\sigma\tilde{L}}, \quad \sigma \in \mathbb{R}.$$

We have $[\tilde{D}, \tilde{L}] = 0$ and \tilde{L} is a first order differential operator, formally selfadjoint for $(\cdot|\cdot)_{\tilde{V}(\tilde{M})}$. If $u \in \overline{C^\infty}(\Omega^\pm; \tilde{V})$, then $\tilde{\varrho}^\pm \tilde{L}u$ depends only on $\tilde{\varrho}^\pm u$ and $\tilde{D}u|_\Sigma$. We set

$$\tilde{\varrho}^\pm \tilde{L}u =: \tilde{L}_\Sigma \tilde{\varrho}^\pm u + Z_\Sigma \tilde{D}u|_\Sigma.$$

Lemma 4.8. *Assume that \tilde{D} is invertible. Then $[\tilde{L}_\Sigma, \tilde{c}^\pm] = 0$.*

Proof. Let $u \in \overline{C^\infty}(\Omega^\pm; \tilde{V})$ with $\tilde{D}u = 0$ in Ω^\pm and $f = \tilde{\varrho}^\pm u$. Then $\tilde{D}\tilde{L}u = \tilde{L}\tilde{D}u = 0$ in Ω^\pm , hence $\tilde{c}^\pm \tilde{L}_\Sigma f = \tilde{L}_\Sigma f = \tilde{L}_\Sigma \tilde{c}^\pm f$ since $f = \tilde{c}^\pm f$. Therefore $\tilde{c}^\mp \tilde{L}_\Sigma \tilde{c}^\pm = 0$ which proves the lemma. \square

4.4. Wick rotation and Hadamard property. We recall here some results from [16, 14] in a simple situation, which will be sufficient for the de Sitter spacetime considered in the sequel.

Let (M, g) be an *analytic* globally hyperbolic spacetime with a *compact* Cauchy surface Σ . We set $D = D_k$, $V = V_k$, $k = 0, 1, 2$, where D_k, V_k are introduced in 2.5.2.

Using Gaussian normal coordinates to Σ , there exists a neighborhood U of Σ in M and an interval $I \ni 0$ such that (U, g) is isometric to $(I \times \Sigma, -dt^2 + h(t, x))$.

Clearly h is real analytic in t and by Wick rotation in t , we obtain a product $\tilde{I} \times \Sigma$, $\tilde{I} =] - \delta, \delta[$, a complex metric $\tilde{g} = ds^2 + \tilde{h}(s, \mathbf{x})$ on $\tilde{I} \times \Sigma$ and a Wick rotated operator \tilde{D} acting on sections of \tilde{V} . As usual we identify Σ with $\{0\} \times \Sigma$.

We will assume that \tilde{g} is *Riemannian* over $\tilde{I} \times \Sigma$ and that \tilde{g} and \tilde{D} extend to a Riemannian metric \tilde{g} and differential operator \tilde{D} on a compact analytic manifold \tilde{M} with $\tilde{V} = \tilde{V}_k = \mathbb{C} \otimes_s^k T^* \tilde{M}$. \tilde{D} has principal symbol $\xi \cdot \tilde{g}^{-1} \xi \mathbb{1}_{\tilde{V}}$ hence is elliptic.

Since g is real-valued we obtain that

$$\tilde{g}(s, \mathbf{x}) = \tilde{g}(s, \mathbf{x})^* = \tilde{g}(-s, \mathbf{x}), \quad s \in I.$$

By analyticity the reflection $s \mapsto -s$ on $\tilde{I} \times \Sigma$ extends to an involution χ on \tilde{M} , which is isometric for \tilde{g} . We denote by κ the unique lift of χ to the bundle \tilde{V} .

Since $D = D^*$ for the scalar product $(\cdot | \cdot)_{V(M)}$, we obtain that

$$\tilde{D}^* = \kappa \tilde{D} \kappa,$$

where \tilde{D}^* is the adjoint of \tilde{D} for the scalar product $(\cdot | \cdot)_{\tilde{V}(\tilde{M})}$, i.e. the hypotheses in 4.2.2 are satisfied.

4.4.1. Lorentzian charge. We define the Lorentzian traces ϱ_k as in 2.5.3, using the future directed unit normal ∂_t . We systematically identify Hermitian forms with linear operators using the *Hilbertian* scalar product $(\cdot | \cdot)_{\tilde{V}_k(\Sigma) \otimes \mathbb{C}^2}$.

Then the Lorentzian charges \mathbf{q}_k , see 2.5.4 are given by

$$\bar{f} \cdot \mathbf{q}_k f = (f | q_k f)_{\tilde{V}_k(\Sigma) \otimes \mathbb{C}^2}, \quad q_k = \begin{pmatrix} 0 & \kappa_{k\Sigma} \\ \kappa_{k\Sigma} & 0 \end{pmatrix},$$

for

$$\kappa_{0\Sigma} = 1, \quad \kappa_{1\Sigma} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \kappa_{2\Sigma} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

and we use the usual decomposition of $(0, k)$ -tensors.

4.4.2. Euclidean operators. We define the Euclidean traces as in 4.1.3, using the normal vector field $-\partial_s$. Then we have

$$\tilde{\sigma}_k = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad \tilde{q}_k = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad k = 0, 1, 2.$$

As claimed in 4.2.2 we have $q_k = \tilde{\sigma}_k \circ \begin{pmatrix} \kappa_{k\Sigma} & 0 \\ 0 & -\kappa_{k\Sigma} \end{pmatrix}$.

4.4.3. Calderón projectors. Let us finally assume that $\tilde{D} : H^2(\tilde{M}; \tilde{V}) \rightarrow L^2(\tilde{M}; \tilde{V})$ is invertible and let \tilde{c}^\pm be the associated Calderón projectors. We use the coordinates (t, \mathbf{x}, τ, k) on $T^*(I \times \Sigma)$.

We recall that the operators $\mathcal{F}_{k\Sigma}$ identifying Cauchy data for D_k and traces for \tilde{D}_k are defined in Subject. 3.2.

Theorem 4.9. *Let us set*

$$c_k^\pm := \mathcal{F}_{k\Sigma}^{-1} \circ \tilde{c}_k^\pm \circ \mathcal{F}_{k\Sigma} : C^\infty(\Sigma; V_k \otimes \mathbb{C}^2) \rightarrow C^\infty(\Sigma; V_k \otimes \mathbb{C}^2).$$

Then

$$\text{WF}(U_k \circ c_k^\pm)' \subset (\mathcal{N}^\pm \cup \mathcal{F}) \times T^*\Sigma,$$

where $\mathcal{F} = \{k = 0\} \subset T^*M$.

Proof. The theorem is proved in [14, Sect. 5]. In [14] a less natural convention is used for the Wick rotation of tensors and differential operators, which consists in replacing t by is and dt by ds , (except for the background metric g which is transformed into \tilde{g}). Changing from the convention for operators in [14] to the present one is done by conjugation by \mathcal{F}_k . Therefore the theorem follows from [14, Prop 5.4, Prop. 5.21]. \square .

5. WICK ROTATION OF DE SITTER SPACE

5.1. Introduction. The de Sitter space is $dS^4 = \mathbb{R}_t \times \mathbb{S}^3$, equipped with the metric

$$(5.1) \quad g = -dt^2 + \cosh^2(t)h,$$

where h is the canonical metric on $\mathbb{S}^3 = \Sigma$.

If we perform the Wick rotation $t \mapsto is$ we obtain

$$(5.2) \quad \tilde{g} = ds^2 + \cos^2(s)h.$$

The manifold $]-\pi/2, \pi/2[_s \times \mathbb{S}^3$, equipped with the metric \tilde{g} is the Euclidean sphere $\mathbb{S}^4 = \{(x^0, \dots, x^4) : \sum_{i=0}^4 (x^i)^2 = 1\}$, with the two poles $x^0 = \pm 1$ removed. This can easily be seen using the coordinates $(s, \omega) \in [-\pi/2, \pi/2] \times \mathbb{S}^3$ on \mathbb{S}^4 given by

$$x^0 = \sin s, \quad (x^1, \dots, x^4) = \cos s \omega, \quad s \in [-\pi/2, \pi/2], \quad \omega \in \mathbb{S}^3.$$

Therefore the hypotheses in Subsect. 4.4 are satisfied by the de Sitter spacetime dS^4 .

5.2. Lichnerowicz Laplacians. As in Subsect. 3.1 we set

$$\tilde{V}_k = \mathbb{C} \otimes_s^k T^*\mathbb{S}^4, \quad k = 0, 1, 2.$$

Let $\tilde{D}_{k,L}$ be the Lichnerowicz Laplacian on $(\mathbb{S}^4, \tilde{g})$ defined as in Subsect. 2.4 and let \tilde{D}_k the Wick rotation of D_k .

We will denote by $\vec{D}_{k,L}, \vec{d}, \vec{\delta}$ the analogous Lichnerowicz Laplacians, differential and co-differential on the standard sphere (\mathbb{S}^3, h) , which is Einstein with $\Lambda = 2$. We denote by $(\tilde{g}|$ and $|\tilde{g})$ the operators

$$(\tilde{g}| : u_2 \mapsto (\tilde{g}|u_2)_{\tilde{v}_2}, \quad |\tilde{g}) : u_0 \mapsto u_0 \tilde{g}$$

and by $(h|, |h)$ their analogs on \mathbb{S}^3 . The identities recalled in Subsect. 2.4 are independent on the signature of the metric. Therefore we have:

$$(5.3) \quad \begin{aligned} \tilde{D}_{k+1,L} \circ d &= d \circ \tilde{D}_{k,L}, & \delta \circ \tilde{D}_{k+1,L} &= \tilde{D}_{k,L} \circ \delta, \\ (\tilde{g}| \circ \tilde{D}_{2,L} &= \tilde{D}_{0,L} \circ (\tilde{g}|, & |\tilde{g}) \circ \tilde{D}_{0,L} &= \tilde{D}_{2,L} \circ |\tilde{g}), \end{aligned}$$

and the analogous identities for $\vec{D}_{k,L}, \vec{d}, \vec{\delta}$.

Since $(\mathbb{S}^4, \tilde{g})$ is Einstein with $\Lambda = 3$ we have:

$$\tilde{D}_k = \tilde{D}_{k,L} - 6.$$

Since on \mathbb{S}^4 $\tilde{\mathbf{R}}_{abcd} = \tilde{g}_{bd}\tilde{g}_{ac} - \tilde{g}_{ad}\tilde{g}_{bc}$, we have $\widetilde{\text{Riem}} = \mathbb{1} - \frac{1}{2}|\tilde{g}|(\tilde{g})$ hence

$$(5.4) \quad \tilde{D}_2 = -\Delta_2 - |\tilde{g}|(\tilde{g}) + 2\mathbb{1}.$$

5.2.1. *Spectrum of $\tilde{D}_{k,L}$.* The spectral decompositions of $\tilde{D}_{k,L}$ on \mathbb{S}^n for $k = 0, 1, 2$ have been completely described in [4]. We summarize now the results from [4] that we will need.

We embed \mathbb{S}^3 into \mathbb{R}^4 so that $\mathbb{S}^3 = \{(y_1, \dots, y_4) \in \mathbb{R}^4 : \sum_{i=1}^4 y_i^2 = 1\}$. We denote by ψ_i , $1 \leq i \leq 4$ the function equal to the restriction of y_i to \mathbb{S}^3 and by φ_{jk} for $1 \leq j < k \leq 4$ the 1-form equal to the restriction of $y_j dy_k - y_k dy_j$ to \mathbb{S}^3 .

Using the coordinates (s, ω) on \mathbb{S}^4 we set:

$$(5.5) \quad \begin{aligned} \varphi_i &= \psi_i ds - \sin s \cos s d\psi_i, \quad 1 \leq i \leq 4, \\ \varphi_{jk} &= \cos^2 s \psi_{jk}, \quad 1 \leq j < k \leq 4. \end{aligned}$$

As is well known φ_i, φ_{jk} span the 10 dimensional space of Killing 1-forms on \mathbb{S}^4 .

Proposition 5.1. (1) \tilde{D}_2 is invertible and $\tilde{D}_2 \geq 2$ on $\text{Ker}(\tilde{g}|)$.

(2) $\text{Ker } \tilde{D}_1 = \text{Ker } d = \text{Ker } d \cap \text{Ker } \delta = \text{Vect}\{\varphi_i, \varphi_{jk} : 1 \leq i \leq 4, 1 \leq j < k \leq 4\}$.

Proof. We have the direct sum decomposition

$$C^\infty(\mathbb{S}^4; \tilde{V}_2) = C^\infty(\mathbb{S}^4; \tilde{V}_0)\tilde{g} \oplus^\perp \text{Ker}(\tilde{g}|),$$

which is preserved by \tilde{D}_2 . By (5.4) $\tilde{D}_2 \geq 2$ on $\text{Ker}(\tilde{g}|)$ and $\tilde{D}_2 \tilde{g} = -6\tilde{g}$, which proves (1). Let us now prove (2). Since $\text{Ker } \tilde{D}_{1,L} = \{0\}$ we have the direct sum decomposition

$$\tilde{V}_1 = \text{Ker } \delta_a \oplus^\perp \text{Ran } d_a,$$

where we recall that d_a, δ_a are the anti-symmetric differential and co-differential. By (2.11) this decomposition is preserved by $\tilde{D}_{1,L}$ hence by \tilde{D}_1 . From the results in [4, Subsect. 3.6] on the spectrum of $\tilde{D}_{1,L}$ on \mathbb{S}^n we obtain that

$$\begin{aligned} \sigma(\tilde{D}_1|_{\text{Ran } d_a}) &= \{k(k+3) - 6 : k \in \mathbb{N}, k \geq 1\}, \\ \sigma(\tilde{D}_1|_{\text{Ker } \delta_a}) &= \{k(k+3) - 4 : k \in \mathbb{N}, k \geq 1\}. \end{aligned}$$

The eigenspace of \tilde{D}_1 for the eigenvalue 0 is included in $\text{Ker } \delta_a$ and obtained for $k = 1$. Using that $\delta_a = \delta_s$ on \tilde{V}_1 and (2.17) we obtain that it is also included in $\text{Ker } d_s$. Moreover following [4, Subsect. 3.6] $\text{Ker } \tilde{D}_1$ consists of restrictions to \mathbb{S}^4 of 1-forms on \mathbb{R}^5 with linear coefficients. A routine computation then shows that $\text{Ker } \tilde{D}_1 = \text{Vect}\{\varphi_i, \varphi_{jk}\}$. \square

Proposition 5.2. *The Lichnerowicz Laplacians $\vec{D}_{i,L}$ on the standard sphere (\mathbb{S}^3, h) satisfy:*

$$(5.6) \quad \begin{aligned} i) \quad & \sigma(\vec{D}_{0,L}) = \{k(k+2) : k \in \mathbb{N}\}, \\ ii) \quad & \text{Ker}(\vec{D}_{0,L} - 3) = \text{Vect}\{\psi_i : 1 \leq i \leq 4\}, \\ iii) \quad & \sigma(\vec{D}_{1,L}) = \{k(k+2) : k \in \mathbb{N}\} \cup \{k(k+2) + 1 : k \in \mathbb{N}, k \geq 1\}, \\ iv) \quad & \text{Ker}(\vec{D}_{1,L} - 3) = \text{Vect}\{\vec{d}\psi_i : 1 \leq i \leq 4\}, \\ v) \quad & \text{Ker}(\vec{D}_{1,L} - 4) = \text{Vect}\{\psi_{j,k} : 1 \leq j < k \leq 4\} = \text{Ker } \vec{\delta} \cap \text{Ker } \vec{d}, \\ vi) \quad & \vec{d} \circ \vec{d}\psi_i + h\psi_i = 0, \quad 1 \leq i \leq 4. \end{aligned}$$

Proof. *i), ii), iii)* and *iv)* are shown in [4, Subsect. 3.5, 3.6], as is the first statement of *v)*. It is also shown that $\text{Ker}(\vec{D}_{1,L} - 4) \subset \text{Ker } \vec{\delta}$. The identities recalled in Subsect. 2.4 show that $\vec{D}_{1,L} - 4 = \vec{\delta} \circ \vec{d} - \vec{d} \circ \vec{\delta}$ which proves the second statement of *v)*. Finally *vi)* follows from [4, Prop. 3.3]. \square

5.3. Spatial decomposition of \tilde{D}_i . We now explain a useful decomposition of \tilde{D}_i . We use the identifications in 2.1.4, 2.1.5 for $(0, i)$ -tensors.

A lengthy computation using A.2.1 gives for $a(s) = \cos^2(s)$:

$$\begin{aligned} (\tilde{D}_1 w)_s &= -\partial_s^2 w_s + a^{-1} \vec{D}_{0,L} w_s - \dot{a} a^{-2} \vec{\delta} w_\Sigma + (\tfrac{3}{4} \dot{a}^2 a^{-2} - 3) w_s, \\ (\tilde{D}_1 w)_\Sigma &= -(\partial_s - \tfrac{1}{2} \dot{a} a^{-1})^2 w_\Sigma + a^{-1} \vec{D}_{1,L} w_\Sigma - \dot{a} a^{-1} \vec{d} w_s \\ &\quad + (\tfrac{1}{4} \dot{a}^2 a^{-2} - 2a^{-1} - 3) w_\Sigma, \\ (\tilde{D}_2 u)_{ss} &= -\partial_s^2 u_{ss} + a^{-1} \vec{D}_{0,L} u_{ss} - 2\dot{a} a^{-3} \vec{\delta} u_{s\Sigma} \\ &\quad + \tfrac{3}{2} \dot{a}^2 a^{-2} u_{ss} - (\tfrac{1}{4} \dot{a}^2 a^{-3} + 1) (h|u_{\Sigma\Sigma}), \\ (\tilde{D}_2 u)_{s\Sigma} &= -(\partial_s - \tfrac{1}{2} \dot{a} a^{-1})^2 u_{s\Sigma} + a^{-1} \vec{D}_{1,L} u_{s\Sigma} - \tfrac{1}{2} \dot{a} a^{-2} \vec{\delta} u_{\Sigma\Sigma} \\ &\quad - \dot{a} a^{-1} \vec{d} u_{ss} + (\tfrac{3}{2} \dot{a}^2 a^{-2} + 2 - 2a^{-1}) u_{s\Sigma}, \\ (\tilde{D}_2 u)_{\Sigma\Sigma} &= -(\partial_s - \dot{a} a^{-1})^2 u_{\Sigma\Sigma} + a^{-1} \vec{D}_{2,L} u_{\Sigma\Sigma} \\ &\quad - 2a^{-1} \dot{a} \vec{d} u_{s\Sigma} + (2 + \tfrac{1}{2} a^{-2} \dot{a}^2 - 6a^{-1}) u_{\Sigma\Sigma} \\ &\quad + (a^{-1} - 1) h(h|u_{\Sigma\Sigma}) - (2 + \tfrac{1}{2} a^{-1} \dot{a}^2) h u_{ss}. \end{aligned}$$

We set as operators on $C^\infty(\mathbb{S}^4; \tilde{V}_i)$, $0 \leq i \leq 2$ with the above identifications (i.e. w identified with $w_s \oplus w_\Sigma$, etc.):

$$\begin{aligned} \hat{D}_0 &= \vec{D}_{0,L}, \\ \hat{D}_1 &= \vec{D}_{0,L} \oplus \vec{D}_{1,L}, \\ \hat{D}_2 &= \vec{D}_{0,L} \oplus \vec{D}_{1,L} \oplus \vec{D}_{2,L}. \end{aligned}$$

5.3.1. Commutation properties. \hat{D}_i are independent on s and act only in the spatial variables. From the above identities we obtain that

$$[\tilde{D}_i, \hat{D}_i] = 0, \quad \text{hence } [\tilde{D}_i, \mathbb{1}_{\{\lambda\}}(\hat{D}_i)] = 0, \quad \lambda \in \mathbb{R},$$

i.e. \tilde{D}_i commute with the spectral projections of \hat{D}_i .

Moreover from the expressions (A.3), (A.4) in the Appendix, we obtain that

$$(5.7) \quad \hat{D}_i \tilde{K}_{ij} = \tilde{K}_{ij} \hat{D}_j, \quad \hat{D}_j \tilde{K}_{ij}^* = \tilde{K}_{ij}^* \hat{D}_i, \quad \hat{D}_2 \tilde{I} = \tilde{I} \hat{D}_2.$$

5.3.2. *Spectrum of \hat{D}_i .* Clearly $\sigma(\hat{D}_i) = \bigcup_{0 \leq j \leq i} \sigma(\vec{D}_{j,L})$. The eigenvalues and eigenspaces of $\vec{D}_{j,L}$ have been completely described in [4]. In particular one obtains that

$$(5.8) \quad \begin{aligned} \text{Ker}(\hat{D}_2 - 3) &= \text{Vect}\{\psi_i \oplus 0 \oplus 0, 0 \oplus \vec{d}\psi_j \oplus 0, 0 \oplus 0 \oplus \psi_k h : 0 \leq i, j, k \leq 4\}, \\ \text{Ker}(\hat{D}_2 - 4) &= \text{Vect}\{0 \oplus \psi_{jk} \oplus 0 : 0 \leq j < k \leq 4\}. \end{aligned}$$

5.4. **The operators $\tilde{K}_{ij\Sigma}$ and $\tilde{K}_{ij\Sigma}^\dagger$.** We recall that $\Sigma := \mathbb{S}^4 \cap \{s = 0\} \sim \mathbb{S}^3$, and we set

$$\Omega^\pm := \mathbb{S}^4 \cap \{\pm s > 0\}.$$

Let $u_1 \in \overline{C^\infty}(\Omega^\pm; \tilde{V}_1)$ and

$$u_2 = \tilde{K}_{21} u_1 = du_1 - \frac{1}{4} \tilde{g}(\tilde{g}|du_1) \tilde{v}_2.$$

The trace $\tilde{\varrho}_2^\pm u_2$ on Σ depends only on $\tilde{\varrho}_1^\pm u_1$ and on $\tilde{D}_1 u_1|_\Sigma$. One sets then

$$(5.9) \quad \tilde{\varrho}_2^\pm \tilde{K}_{21} u_1 =: \tilde{K}_{21\Sigma} \tilde{\varrho}_1^\pm u_1 + R_\Sigma \tilde{D}_1 u_1|_\Sigma.$$

Similarly if $u_2 \in \overline{C^\infty}(\Omega^\pm; \tilde{V}_2)$ and $u_1 = \delta u_2 = \tilde{K}_{21}^* u_2$, then

$$\tilde{\varrho}_1^\pm \tilde{K}_{21}^* u_2 =: \tilde{K}_{21\Sigma}^\dagger \tilde{\varrho}_2^\pm u_2 + S_\Sigma \tilde{D}_2 u_2|_\Sigma.$$

Note that the notation for $\tilde{K}_{21\Sigma}$, $\tilde{K}_{21\Sigma}^\dagger$ is coherent with the one in (3.2), i.e. $\tilde{K}_{21\Sigma}$, $\tilde{K}_{21\Sigma}^\dagger$ are the Wick rotations of $K_{21\Sigma}$ and $K_{21\Sigma}^\dagger$. In this subsection we will compute the operators $\tilde{K}_{21\Sigma}$, $\tilde{K}_{21\Sigma}^\dagger$ and R_Σ (the operator S_Σ will not be needed). The proofs are given in Appendix A.2.

Lemma 5.3. *If $f = \tilde{K}_{21\Sigma}^\dagger g$ with $f = \begin{pmatrix} f_0 \\ f_1 \end{pmatrix} \in C^\infty(\Sigma; \tilde{V}_1 \otimes \mathbb{C}^2)$ and $g = \begin{pmatrix} g_0 \\ g_1 \end{pmatrix} \in C^\infty(\Sigma; \tilde{V}_2 \otimes \mathbb{C}^2)$, then:*

$$\begin{cases} f_{0s} = 2g_{1ss} + 2\vec{\delta}g_{0s\Sigma}, \\ f_{0\Sigma} = 2g_{1s\Sigma} + \vec{\delta}g_{0\Sigma\Sigma}, \\ f_{1s} = 2(\vec{D}_{0,L} - 3)g_{0ss} + 2\vec{\delta}g_{1s\Sigma} - (h|g_{0\Sigma\Sigma}), \\ f_{1\Sigma} = 2(\vec{D}_{1,L} - 4)g_{0s\Sigma} + \vec{\delta}g_{1\Sigma\Sigma}. \end{cases}$$

Lemma 5.4. *If $g = \tilde{K}_{21\Sigma}f$ with $f = \begin{pmatrix} f_0 \\ f_1 \end{pmatrix} \in C^\infty(\Sigma; \tilde{V}_1 \otimes \mathbb{C}^2)$ and $g = \begin{pmatrix} g_0 \\ g_1 \end{pmatrix} \in C^\infty(\Sigma; \tilde{V}_2 \otimes \mathbb{C}^2)$, then:*

$$\begin{cases} g_{0ss} = \frac{1}{2}(-f_{1s} + \vec{\delta}f_{0\Sigma}), \\ g_{0s\Sigma} = \frac{1}{2}(-f_{1\Sigma} + \vec{d}f_{0s}), \\ g_{0\Sigma\Sigma} = \vec{d}f_{0\Sigma} + \frac{1}{2}(f_{1s} + \vec{\delta}f_{0\Sigma})h, \\ g_{1ss} = \frac{1}{2}(-\vec{D}_{0,L}f_{0s} + \vec{\delta}f_{1\Sigma}), \\ g_{1s\Sigma} = \frac{1}{2}(-(\vec{D}_{1,L} - 4)f_{0\Sigma} + \vec{d}f_{1s}), \\ g_{1\Sigma\Sigma} = \vec{d}f_{1\Sigma} + \frac{1}{2}\vec{\delta}f_{1\Sigma}h + \frac{1}{2}(\vec{D}_{0,L} - 4)f_{0s}h. \end{cases}$$

If $v \in C^\infty(\Sigma; \tilde{V}_1)$ then:

$$\begin{cases} (R_\Sigma v)_{1ss} = \frac{1}{2}v_s, \\ (R_\Sigma v)_{1s\Sigma} = \frac{1}{2}v_\Sigma, \\ (R_\Sigma v)_{1\Sigma\Sigma} = -\frac{1}{2}v_s h, \end{cases}$$

all other entries of $R_\Sigma v$ being equal to 0.

We have similar results for the operators \tilde{K}_{20} and \tilde{K}_{20}^* defined by

$$\begin{aligned} \tilde{K}_{20}u_0 &= \tilde{g}u_0, \\ \tilde{K}_{20}^*u_2 &= -(\tilde{g}|u_2)\tilde{v}_2(\mathbb{S}^4), \quad u_k \in C^\infty(\mathbb{S}^4; \tilde{V}_k). \end{aligned}$$

Note that \tilde{K}_{20} , \tilde{K}_{20}^* are the Wick rotations of K_{20} , K_{20}^* defined in Subsect. 2.8. Moreover if $u_2 = \tilde{K}_{20}u_0$ then $\tilde{\varrho}_2^\pm u_2$ depends only on $\tilde{\varrho}_0^\pm u_0$ and similarly for \tilde{K}_{20}^* . We obtain the following expression for $\tilde{K}_{20\Sigma}^\dagger$.

Lemma 5.5. *If $f = \tilde{K}_{20\Sigma}^\dagger g$ then:*

$$\begin{cases} f_0 = -2g_{0ss} - (h|g_{0\Sigma\Sigma}), \\ f_1 = -2g_{1ss} - (h|g_{1\Sigma\Sigma}). \end{cases}$$

5.4.1. *Commutation properties with \hat{D}_i .*

Lemma 5.6. *We have*

$$(\mathbb{1}_{\{\lambda\}}(\hat{D}_i) \otimes \mathbb{1}_{\mathbb{C}^2}) \circ \tilde{K}_{ij\Sigma} = \tilde{K}_{ij\Sigma} \circ (\mathbb{1}_{\{\lambda\}}(\hat{D}_j) \otimes \mathbb{1}_{\mathbb{C}^2})$$

for $(i, j) = (2, 1)$ or $(i, j) = (2, 0)$.

Proof. We prove the lemma for $(i, j) = (2, 1)$, the proof for $(i, j) = (2, 0)$ being simpler and left to the reader. From the expression of R_Σ in Lemma 5.4 we obtain that

$$(5.10) \quad (\mathbb{1}_{\{\lambda\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2})R_\Sigma = R_\Sigma \mathbb{1}_{\{\lambda\}}(\hat{D}_1).$$

Using (5.9) and the fact that $\hat{D}_2 \tilde{K}_{21} = \tilde{K}_{21} \hat{D}_1$ we have:

$$\begin{aligned} \tilde{\varrho}_2 \tilde{K}_{21} \mathbb{1}_{\{\lambda\}}(\hat{D}_1) u_1 &= \tilde{\varrho}_2 \mathbb{1}_{\{\lambda\}}(\hat{D}_2) \tilde{K}_{21} u_1 = (\mathbb{1}_{\{\lambda\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2}) \tilde{\varrho}_2 \tilde{K}_{21} u_1 \\ &= (\mathbb{1}_{\{\lambda\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2}) \tilde{K}_{21\Sigma} \tilde{\varrho}_1 u_1 + (\mathbb{1}_{\{\lambda\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2}) R_\Sigma \tilde{D}_1 u_1 \upharpoonright_\Sigma. \end{aligned}$$

From (5.9) and (5.10) we have

$$\begin{aligned} \tilde{\varrho}_2 \tilde{K}_{21} \mathbb{1}_{\{\lambda\}}(\hat{D}_1) u_1 &= \tilde{K}_{21\Sigma} \tilde{\varrho}_1 \mathbb{1}_{\{\lambda\}}(\hat{D}_1) u_1 + R_\Sigma \tilde{D}_1 \mathbb{1}_{\{\lambda\}}(\hat{D}_1) u_1 \upharpoonright_\Sigma \\ &= \tilde{K}_{21\Sigma} (\mathbb{1}_{\{\lambda\}}(\hat{D}_1) \otimes \mathbb{1}_{\mathbb{C}^2}) \tilde{\varrho}_1 u_1 + (\mathbb{1}_{\{\lambda\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2}) R_\Sigma \tilde{D}_1 u_1 \upharpoonright_\Sigma, \end{aligned}$$

which proves the lemma. \square

5.5. The operators $K_{ij\Sigma}$ and $K_{ij\Sigma}^\dagger$. We obtain the Lorentzian operators $K_{ij\Sigma}$ and $K_{ij\Sigma}^\dagger$ by conjugation with $\mathcal{F}_{i\Sigma}$, $\mathcal{F}_{j\Sigma}$ as indicated in Subsect. 3.2, i.e.:

$$K_{ij\Sigma}^\dagger = \mathcal{F}_{i\Sigma}^{-1} \circ \tilde{K}_{ij\Sigma} \circ \mathcal{F}_{j\Sigma}, \quad K_{ij\Sigma}^\dagger = \mathcal{F}_{j\Sigma}^{-1} \circ \tilde{K}_{ij\Sigma}^\dagger \circ \mathcal{F}_{i\Sigma}.$$

The explicit expressions of these operators is not important. Since $\mathbb{1}_{\{\lambda\}}(\hat{D}_i)$ commutes with \mathcal{F}_i we obtain from Lemma 5.6 that

$$(\mathbb{1}_{\{\lambda\}}(\hat{D}_i) \otimes \mathbb{1}_{\mathbb{C}^2}) \circ K_{ij\Sigma} = K_{ij\Sigma} \circ (\mathbb{1}_{\{\lambda\}}(\hat{D}_j) \otimes \mathbb{1}_{\mathbb{C}^2}),$$

for $(i, j) = (2, 1)$ or $(i, j) = (2, 0)$.

6. PROPERTIES OF THE TT GAUGE PHYSICAL SPACE

Before studying the Euclidean “vacuum” ω_{eucl} for linearized gravity on dS^4 it is necessary to investigate the TT gauge physical phase space \mathcal{E}_{TT} , see Def. 2.15. Our main result is that $\mathbf{q}_{1,2}$ is *degenerate* on \mathcal{E}_{TT} with an explicit 10 dimensional kernel.

6.1. The kernel of the Lorentzian charge. As a first step we describe a convenient orthogonal decomposition of its Euclidean version

$$\tilde{\mathcal{E}}_{\text{TT}} = \text{Ker } \tilde{K}_{21\Sigma}^\dagger \cap \text{Ker } \tilde{K}_{20\Sigma}^\dagger = \mathcal{F}_{2\Sigma} \mathcal{E}_{\text{TT}}.$$

The proof of the next lemma is given in Appendix A.2.4.

Lemma 6.1. $g = \begin{pmatrix} g_0 \\ g_1 \end{pmatrix} \in \tilde{\mathcal{E}}_{\text{TT}}$ if and only if:

$$\begin{aligned} 1) \quad g_{0ss} &= \frac{3}{4} (\vec{D}_{0,L} - 3)^{-1} \vec{\delta} \vec{\delta} \vec{g}_{0\Sigma\Sigma} - 3\beta_0, \\ 2) \quad g_{1ss} &= \frac{3}{4} (\vec{D}_{0,L} - 6)^{-1} \vec{\delta} \vec{\delta} \vec{g}_{1\Sigma\Sigma}, \\ 3) \quad g_{0s\Sigma} &= -\frac{1}{2} (\vec{D}_{1,L} - 4)^{-1} (1 + \frac{1}{2} \vec{d} (\vec{D}_{0,L} - 6)^{-1} \vec{\delta}) \vec{\delta} \vec{g}_{1\Sigma\Sigma} + \beta_1, \\ 4) \quad g_{1s\Sigma} &= -\frac{1}{2} (1 + \frac{1}{2} \vec{d} (\vec{D}_{0,L} - 3)^{-1} \vec{\delta}) \vec{\delta} \vec{g}_{0\Sigma\Sigma} + \vec{d} \beta_0, \\ 5) \quad g_{0\Sigma\Sigma} &= -\frac{1}{4} (\vec{D}_{0,L} - 3)^{-1} \vec{\delta} \vec{\delta} \vec{g}_{0\Sigma\Sigma} h + \vec{g}_{0\Sigma\Sigma} + \beta_0 h, \\ 6) \quad g_{1\Sigma\Sigma} &= -\frac{1}{4} (\vec{D}_{0,L} - 6)^{-1} \vec{\delta} \vec{\delta} \vec{g}_{1\Sigma\Sigma} h + \vec{g}_{1\Sigma\Sigma}, \end{aligned} \tag{6.1}$$

for $\beta_0 \in \text{Ker}(\vec{D}_{0,L} - 3) = \text{Vect}\{\psi_i\}$, $\beta_1 \in \text{Ker}(\vec{D}_{1,L} - 4) = \text{Vect}\{\psi_{jk}\}$ and $\vec{g}_{i\Sigma\Sigma} \in C^\infty(\Sigma; \tilde{V}_{2\Sigma})$ with $(h|\vec{g}_{i\Sigma\Sigma}) = 0$.

In particular $\tilde{\mathcal{E}}_{\text{TT}}$ and hence \mathcal{E}_{TT} is isomorphic to the space

$$(\text{Ker}(h| \otimes \mathbb{C}^2) \oplus \text{Ker}(\vec{D}_{0,L} - 3) \oplus \text{Ker}(\vec{D}_{1,L} - 4)).$$

Proposition 6.2. (1) *The spectral projections $\mathbb{1}_{\{\lambda\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2}$ preserve \mathcal{E}_{TT} .*
(2) *The kernel of the restriction of \mathbf{q}_2 to \mathcal{E}_{TT} equals $(\mathbb{1}_{\{3,4\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2})\mathcal{E}_{\text{TT}}$.*

The proof is given in Appendix A.2.5.

The space $(\mathbb{1}_{\{3,4\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2})\mathcal{E}_{\text{TT}} = \text{Ker } \mathbf{q}_{I,2}$ is therefore canonically defined.

The space $\mathcal{E}_{\text{TT,reg}} = (\mathbb{1}_{\mathbb{R} \setminus \{3,4\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2})\mathcal{E}_{\text{TT}}$, which will be useful later on, is just a convenient supplementary space.

6.2. Decomposition of \mathcal{E}_{TT} .

6.2.1. *Killing 1-forms.* We denote by $\mathcal{K} \subset \text{Ker}_{\text{sc}} D_1$ the space of *Killing 1-forms* on dS^4 . We have $\mathcal{K} = \text{Ker } d \cap \text{Ker } \delta$ and \mathcal{K} is spanned by the 1-forms $-\psi_i dt + \sinh t \cosh t \vec{d}\psi_i$ and $\cosh^2 t \psi_{jk}$ for $1 \leq i \leq 4$, $1 \leq j < k \leq 4$. Clearly,

$$\mathcal{K} = \mathcal{F}_1^{-1} \text{Ker } \tilde{D}_1,$$

since $\text{Ker } \tilde{D}_1$ is spanned by the Killing 1-forms on \mathbb{S}^4 .

Definition 6.3. *We set*

$$\mathcal{K}_\Sigma := \varrho_1 \mathcal{K},$$

and denote by $\mathcal{K}_\Sigma^{\mathbf{q}_1}$ its orthogonal for the charge \mathbf{q}_1 .

In the next lemma (Lemma 6.4 below) we describe the space $\mathcal{K}_\Sigma^{\mathbf{q}_1}$ explicitly. To this end we set:

$$(6.2) \quad C_{1\Sigma} : C^\infty(\Sigma; V_1 \otimes \mathbb{C}^2) \rightarrow C^\infty(\Sigma, V_1)$$

$$f = \begin{pmatrix} f_0 \\ f_1 \end{pmatrix} \mapsto \begin{pmatrix} \mathbb{1}_{\{3\}}(\vec{D}_{0,L})(f_{1s} + i\vec{\delta}f_{0\Sigma}) \\ \mathbb{1}_{\{4\}}(\vec{D}_{1,L})f_{1\Sigma} \end{pmatrix}.$$

We will need also a corresponding operator acting on Cauchy data for D_2 . We set:

$$(6.3) \quad C_{2\Sigma} : C^\infty(\Sigma; V_2 \otimes \mathbb{C}^2) \rightarrow C^\infty(\Sigma, V_1)$$

$$f = \begin{pmatrix} f_0 \\ f_1 \end{pmatrix} \mapsto \begin{pmatrix} i\mathbb{1}_{\{3\}}(\vec{D}_{0,L})f_{0ss} \\ i\mathbb{1}_{\{4\}}(\vec{D}_{1,L})f_{0s\Sigma} \end{pmatrix}.$$

The Euclidean versions of $C_{k\Sigma}$ defined as the appropriate compositions by $\mathcal{F}_k, \mathcal{F}_{k\Sigma}^{-1}$ are:

$$\tilde{C}_{1\Sigma} : C^\infty(\Sigma; \tilde{V}_1 \otimes \mathbb{C}^2) \rightarrow C^\infty(\Sigma, \tilde{V}_1)$$

$$f = \begin{pmatrix} f_0 \\ f_1 \end{pmatrix} \mapsto \begin{pmatrix} \mathbb{1}_{\{3\}}(\vec{D}_{0,L})(f_{1s} - \vec{\delta}f_{0\Sigma}) \\ \mathbb{1}_{\{4\}}(\vec{D}_{1,L})f_{1\Sigma} \end{pmatrix},$$

$$\tilde{C}_{2\Sigma} : C^\infty(\Sigma; \tilde{V}_2 \otimes \mathbb{C}^2) \rightarrow C^\infty(\Sigma, \tilde{V}_1)$$

$$f = \begin{pmatrix} f_0 \\ f_1 \end{pmatrix} \mapsto \begin{pmatrix} \mathbb{1}_{\{3\}}(\vec{D}_{0,L})f_{0ss} \\ \mathbb{1}_{\{4\}}(\vec{D}_{1,L})f_{0s\Sigma} \end{pmatrix}.$$

Lemma 6.4. *We have*

$$(6.4) \quad \begin{aligned} i) \quad & \text{Ker } C_{1\Sigma} = \mathcal{K}_\Sigma^{\mathbf{q}_1}, \\ ii) \quad & C_{2\Sigma} \circ K_{21\Sigma} = -\frac{1}{2}C_{1\Sigma}. \end{aligned}$$

Proof. *ii)* is a simple computation. To prove *i)* we can work in the Euclidean framework. We have:

$$\tilde{\varrho}_1 \varphi_i = \begin{pmatrix} \psi_i \\ 0 \\ 0 \\ d\psi_i \end{pmatrix}, \quad \tilde{\varrho}_1 \varphi_{jk} = \begin{pmatrix} 0 \\ \psi_{jk} \\ 0 \\ 0 \end{pmatrix},$$

and $q_1 = \begin{pmatrix} 0 & \kappa_1 \\ \kappa_1 & 0 \end{pmatrix}$, for $\kappa_1 = \begin{pmatrix} -1 & 0 \\ 0 & \mathbb{1} \end{pmatrix}$. Therefore

$$\begin{aligned} (\tilde{\varrho}_1 \varphi_i | q_1 f_1)_{\tilde{V}_1(\Sigma) \otimes \mathbb{C}^2} &= -(\psi_i | f_{1s}) + (d\psi_i | f_{0\Sigma}), \\ (\tilde{\varrho}_1 \varphi_{jk} | q_1 f_1)_{\tilde{V}_1(\Sigma) \otimes \mathbb{C}^2} &= (\psi_{jk} | f_{1\Sigma}), \end{aligned}$$

which proves *i)*. \square

From Lemma 6.4 we get in particular

$$K_{21\Sigma} \mathcal{K}_\Sigma^{\mathbf{q}_1} \subset \text{Ker } C_{2\Sigma}.$$

6.2.2. *Decomposition of \mathcal{E}_{TT} .* We set

$$\begin{aligned} \mathcal{E}_{\text{TT,reg}} &= (\mathbb{1}_{\mathbb{R} \setminus \{3,4\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2}) \mathcal{E}_{\text{TT}}, \\ \mathcal{E}_{\text{TT},\lambda} &= (\mathbb{1}_{\{\lambda\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2}) \mathcal{E}_{\text{TT}}, \quad \lambda = 3, 4, \end{aligned}$$

so that

$$(6.5) \quad \mathcal{E}_{\text{TT}} = \mathcal{E}_{\text{TT,reg}} \oplus \mathcal{E}_{\text{TT},3} \oplus \mathcal{E}_{\text{TT},4},$$

and by Prop. 6.2, $\text{Ker } \mathbf{q}_2|_{\mathcal{E}_{\text{TT}}} = \mathcal{E}_{\text{TT},3} \oplus \mathcal{E}_{\text{TT},4}$. We first give an alternative description of $\mathcal{E}_{\text{TT,reg}}$.

Lemma 6.5. *We have*

$$\mathcal{E}_{\text{TT,reg}} = \mathcal{E}_{\text{TT}} \cap \text{Ker } C_{2\Sigma}.$$

Proof. It suffices to prove the Euclidean version of the lemma, i.e. with appropriate compositions by $\mathcal{F}_{k\Sigma}$. We use the description of $\tilde{\mathcal{E}}_{\text{TT}}$ given in Lemma 6.1. It is proved in A.2.5, see (A.7), that $g = \begin{pmatrix} g_0 \\ g_1 \end{pmatrix} \in \tilde{\mathcal{E}}_{\text{TT,reg}}$ iff $\beta_0 = \beta_1 = 0$, where β_0, β_1 appear in (6.1). Now $g \in \text{Ker } \tilde{C}_{2\Sigma}$ iff $\mathbb{1}_{\{3\}}(\vec{D}_{0,L})g_{0ss} = \mathbb{1}_{\{4\}}(\vec{D}_{1,L})g_{0s\Sigma} = 0$, which is equivalent to $\beta_0 = \beta_1 = 0$. \square

Proposition 6.6. *Let*

$$\pi : \mathcal{E}_{\text{TT}} \rightarrow \mathcal{E}_{\text{TT,reg}}$$

be the projection on $\mathcal{E}_{\text{TT,reg}}$ along $\mathcal{E}_{\text{TT},3} \oplus \mathcal{E}_{\text{TT},4}$. Then:

- (1) $\pi = \mathbb{1} + K_{21\Sigma} \circ R$, where R is smoothing and of finite rank,
- (2) $\text{Ran } \pi K_{21\Sigma} \subset K_{21\Sigma} \mathcal{K}_\Sigma^{\mathbf{q}_1}$.

Proof. We work in the Euclidean setting, ie we replace $K_{ij\Sigma}^\dagger, C_{k\Sigma}$ by $\tilde{K}_{ij\Sigma}^\dagger, \tilde{C}_{k\Sigma}$. We start by finding an explicit expression for π . Let $g \in \tilde{\mathcal{E}}_{\text{TT}}$. We look for f such that

$$g + \tilde{K}_{21\Sigma} f \in \tilde{\mathcal{E}}_{\text{TT}} \cap \text{Ker } \tilde{C}_{2\Sigma},$$

or equivalently, using Lemmas 5.4 and 5.5:

$$\begin{cases} f_{1s} + \vec{\delta} f_{0\Sigma} = 0, \\ (\vec{D}_{0,L} - 6)f_{0s} + \vec{\delta} f_{1\Sigma} = 0, \\ \mathbb{1}_{\{3\}}(\vec{D}_{0,L})(f_{1s} - \vec{\delta} f_{0\Sigma}) = 2\mathbb{1}_{\{3\}}(\vec{D}_{0,L})g_{0ss}, \\ \mathbb{1}_{\{4\}}(\vec{D}_{1,L})f_{1\Sigma} = 2\mathbb{1}_{\{4\}}(\vec{D}_{1,L})g_{0s\Sigma}. \end{cases}$$

This is equivalent to

$$\begin{cases} f_{1s} = -\vec{\delta} f_{0\Sigma}, \\ f_{0s} = -(\vec{D}_{0,L} - 6)^{-1} \vec{\delta} f_{1\Sigma}, \\ \mathbb{1}_{\{3\}}(\vec{D}_{0,L})\vec{\delta} f_{0\Sigma} = -\mathbb{1}_{\{3\}}(\vec{D}_{0,L})g_{0ss}, \\ \mathbb{1}_{\{4\}}(\vec{D}_{1,L})f_{1\Sigma} = 2\mathbb{1}_{\{4\}}(\vec{D}_{1,L})g_{0s\Sigma}, \end{cases}$$

which we solve by

$$(6.6) \quad \begin{cases} f = \tilde{R}g, \text{ with} \\ f_{0s} = 0, f_{1s} = \mathbb{1}_{\{3\}}(\vec{D}_{0,L})g_{0ss}, \\ f_{0\Sigma} = -\frac{1}{3}\vec{d}\mathbb{1}_{\{3\}}(\vec{D}_{0,L})g_{0ss}, f_{1\Sigma} = 2\mathbb{1}_{\{4\}}(\vec{D}_{1,L})g_{0s\Sigma}. \end{cases}$$

We set $R = \mathcal{F}_{1\Sigma}^{-1} \tilde{R} \mathcal{F}_{2\Sigma}$ and $\pi g = g + K_{21\Sigma} Rg$. Clearly R is smoothing and of finite rank.

If $g = K_{21\Sigma} f'$ then $\pi g = K_{21\Sigma}(f' + f)$ and $0 = C_{2\Sigma} \pi g = -\frac{1}{2} C_{1\Sigma}(f + f')$ by Lemma 6.4 *ii*), hence $f + f' \in \text{Ker } C_{1\Sigma} = \mathcal{K}_\Sigma^{\mathbf{q}_1}$.

It remains to prove that π is the projection on $\mathcal{E}_{\text{TT,reg}}$ along $\mathcal{E}_{\text{TT},3} \oplus \mathcal{E}_{\text{TT},4}$. By Lemma 6.5 we obtain that $Rg = 0$ if $g \in \mathcal{E}_{\text{TT,reg}}$ hence $\pi = \mathbb{1}$ on $\mathcal{E}_{\text{TT,reg}}$. If $f \in \mathcal{E}_{\text{TT},\lambda}$, $\lambda = 3, 4$, then using the decomposition in Lemma 6.1 and an easy computation we obtain that $K_{21\Sigma} Rf = -f$ hence $\pi f = 0$. This completes the proof of the proposition. \square

Remark 6.7. From Prop. 6.6 we deduce that $\text{Ker } \mathbf{q}_2|_E = \mathcal{E}_{\text{TT},3} \oplus \mathcal{E}_{\text{TT},4} \subset \text{Ran } K_{21\Sigma}$. Therefore $[\mathbf{q}_2]$ is non-degenerate on the physical phase space $\frac{\mathcal{E}_{\text{reg}}}{\mathcal{F}_{\text{reg}}}$.

6.3. Action of symmetries. We now investigate the action of the isometry group $O(1, 4)$ of dS^4 on \mathcal{E}_{TT} .

It is easier in practice, though completely equivalent, to consider the solution space version of \mathcal{E}_{TT} equal to $\text{Ker}_{\text{sc}} D_2 \cap \text{Ker}_{\text{sc}} K_{21} \cap \text{Ker}_{\text{sc}} K_{20}$.

We will also consider the subgroup $O^\uparrow(1, 4)$ of isometries preserving the time orientation, the connected component of identity $SO^\uparrow(1, 4)$, and the subgroup $O(4)$ of isometries preserving the hypersurfaces $t = cst$.

An element $\alpha \in O^\uparrow(1, 4)$ acts \mathbb{C} -linearly on tensor fields by

$$u_k \mapsto \alpha^* u_k, \quad u_k \in C^\infty(dS^4; V_k).$$

Clearly α^* is unitary for $\mathbf{q}_{I,2}$.

6.3.1. *Time reversal.* The time reversal $\tau : dS^4 \in (t, \omega) \mapsto (-t, \omega) \in dS^4$ acts \mathbb{C} anti-linearly on tensor fields by

$$Zu_k := \overline{\tau^* u_k}, \quad u_k \in C^\infty(dS^4; V_k).$$

The map Z is often called the *Wigner time reversal*. Since Z reverses the time orientation we have $ZG_k = -G_k Z$ hence $\overline{Zu_k} \cdot \mathbf{Q}_k Zv_k = \bar{v}_k \cdot \mathbf{Q}_k u_k$ so Z is anti-unitary for \mathbf{Q}_k .

Clearly the action of $O(1,4)$ on tensor fields defined as above preserves $\text{Ran } K_{21}$, $\text{Ker } K_{21}^*$ and $\text{Ker } K_{20}^*$, see Subsect. 2.8 for notation. Therefore this action induces a well-defined action on the TT gauge subspace $\text{Ker}_{\text{sc}} D_2 \cap \text{Ker}_{\text{sc}} K_{21} \cap \text{Ker}_{\text{sc}} K_{20}$ and also on the TT gauge quotient space

$$\frac{\text{Ker}_{\text{sc}} D_2 \cap \text{Ker}_{\text{sc}} K_{21}^* \cap \text{Ker}_{\text{sc}} K_{20}^*}{K_{21} \text{Ker}_{\text{sc}} D_1 \cap \text{Ker}_{\text{sc}} K_{20}^*}.$$

7. THE EUCLIDEAN VACUUM ON DE SITTER SPACE

In this section we define the Euclidean ‘‘vacuum’’ for linearized gravity on dS^4 and prove some of its properties. Our physical phase space is the TT gauge subspace \mathcal{E}_{TT} studied in Sect. 6.

We will see that the Euclidean vacuum is *not a state* on the physical phase space, because its covariances are not positive. Therefore we will use the words ‘*pseudo covariances*’ and ‘*pseudo state*’. Perhaps more importantly, we will see that the Euclidean vacuum is *not* strongly gauge invariant.

We recall that we identify hermitian forms on $C_0^\infty(\Sigma; V_2 \otimes \mathbb{C}^2)$ with linear operators using the Hilbertian scalar product $(\cdot | \cdot)_{\tilde{v}_2(\Sigma) \otimes \mathbb{C}^2}$, obtained from the Riemannian metric \tilde{g} , see 4.4.1.

7.1. **Definition of ω_{eucl} and basic properties.** We recall that $\Sigma = \{s = 0\} \subset \mathbb{S}^4$, $\Omega^\pm = \{\pm s > 0\}$. Since \tilde{D}_2 is invertible, see Prop. 5.1, we can define the Calderón projectors \tilde{c}_2^\pm associated to Ω^\pm and \tilde{D}_2 .

Definition 7.1. Let $c_2^\pm := \mathcal{F}_{2\Sigma}^{-1} \circ \tilde{c}_2^\pm \circ \mathcal{F}_{2\Sigma}$ and

$$\bar{f} \cdot \lambda_{2\Sigma}^\pm f := \pm \bar{f} \cdot \mathbf{q}_{I,2} c_2^\pm f = (f | q_2 \circ I_\Sigma \circ c_2^\pm f)_{\tilde{v}_2(\Sigma) \otimes \mathbb{C}^2}$$

for $f \in C_0^\infty(\Sigma; V_2 \otimes \mathbb{C}^2)$. Then

$$\lambda_{2\Sigma}^\pm = \lambda_{2\Sigma}^{\pm*}, \quad \lambda_{2\Sigma}^+ - \lambda_{2\Sigma}^- = \mathbf{q}_{I,2}.$$

The associated pseudo state ω_{eucl} is called the Euclidean vacuum.

Proof. The fact that $\lambda_{2\Sigma}^\pm$ are Hermitian is proved in Prop. 4.4 in a more general framework. The second statement follows from Prop. 4.3 i). \square

7.1.1. *Hadamard property.* Another property of ω_{eucl} which is relatively easy to prove is the *Hadamard property*. In fact from Thm. 4.9 and the results recalled in 2.7.3 we obtain that if we define as usual the space-time pseudo covariances Λ_2^\pm as in (2.26) and the two-point functions Λ_2^\pm as in 2.24 then

$$\text{WF}(\Lambda_2^\pm)' \subset \mathcal{N}^+ \times \mathcal{N}^+,$$

i.e. ω_{eucl} is a Hadamard pseudo state.

7.1.2. *Commutation properties.* From the fact that $[\tilde{D}_2, \hat{D}_2] = 0$ and that \hat{D}_2 acts only in the spatial variables, we obtain immediately that

$$(7.1) \quad [\mathbb{1}_{\{\lambda\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2}, c_2^\pm] = 0,$$

the same property being also true for \hat{c}_2^\pm .

7.2. **Gauge invariance.** In the next proposition we prove that as a consequence of the existence of Killing 1-forms, c_2^\pm do *not* satisfy the strong TT gauge invariance property in the sense that

$$c_2^\pm : \mathcal{F}_{\text{TT}} \rightarrow \text{Ran } K_{21\Sigma} + \text{Ran } K_{20\Sigma}.$$

However, c_2^\pm satisfy the weak TT gauge invariance explained in 2.8.5.

More precisely, let us recall that \mathcal{K}_Σ is the space of Cauchy data of Killing 1-forms on dS^4 , and $\mathcal{K}_\Sigma^{\mathfrak{q}_1}$ is its orthogonal for \mathfrak{q}_1 . We show that c_2^\pm is only invariant under gauge transformations by elements of $\mathcal{K}_\Sigma^{\mathfrak{q}_1}$. However, this turns out to be sufficient for the weak TT gauge invariance thanks to the observation that

$$\text{Ran}_c K_{21\Sigma}^\dagger \subset \mathcal{K}_\Sigma^{\mathfrak{q}_1}.$$

In fact, for all $K_{21\Sigma}^\dagger f_2 \in \text{Ran}_c K_{21\Sigma}^\dagger$ and all $h_1 \in \mathcal{K}_\Sigma$,

$$\overline{h_1 \cdot \mathfrak{q}_1} K_{21\Sigma}^\dagger f_1 = \overline{K_{21\Sigma} h_1 \cdot \mathfrak{q}_1} f_1 = 0,$$

so $K_{21\Sigma}^\dagger f_2 \in \mathcal{K}_\Sigma^{\mathfrak{q}_1}$.

We set in analogy to Def. 7.1:

$$c_1^\pm := \mathcal{F}_{1\Sigma}^{-1} \circ \hat{c}_1^\pm \circ \mathcal{F}_{1\Sigma},$$

where the Calderón projectors c_1^\pm for \tilde{D}_1 are defined as in 4.2.3, i.e. c_1^\pm are defined *only on* $\mathcal{K}_\Sigma^{\mathfrak{q}_1}$.

Proposition 7.2. *One has*

$$c_2^\pm K_{21\Sigma} f_1 \in \text{Ran } K_{21\Sigma} + \text{Ran } K_{20\Sigma} \text{ iff } f_1 \in \mathcal{K}_\Sigma^{\mathfrak{q}_1},$$

and then

$$c_2^\pm K_{21\Sigma} f_1 = K_{21\Sigma} c_1^\pm f_1.$$

Moreover

$$K_{21\Sigma}^\dagger f_2 = 0 \Rightarrow K_{21\Sigma}^\dagger c_2^\pm f_2 \in \mathcal{K}_\Sigma.$$

Proof. We work in the Euclidean framework. Let $f \in C^\infty(\Sigma; \tilde{V}_1(\Sigma) \otimes \mathbb{C}^2)$ and

$$u_1 = (\tilde{D}_1 + \pi_1)^{-1} \tilde{\varrho}_1^* \tilde{\sigma}_1 f, \quad u_2 = \tilde{K}_{21} u_1,$$

where $\pi_1 = \mathbb{1}_{\{0\}}(\tilde{D}_1)$ is the orthogonal projection on $\text{Ker } \tilde{D}_1 = \text{Ker } d$. By Subsect. 5.4 we have

$$\begin{aligned} \tilde{\varrho}_2^\pm u_2 &= \tilde{K}_{21\Sigma} \tilde{\varrho}_1^\pm u_1 + R_\Sigma \tilde{D}_1 u_1|_\Sigma \\ &= \tilde{K}_{21\Sigma} \tilde{\varrho}_1^\pm u_1 - R_\Sigma \pi_1 u_1|_\Sigma \\ &= \mp \tilde{K}_{21\Sigma} \hat{c}_1^\pm f - R_\Sigma (\pi_1 \tilde{\varrho}_1^* \tilde{\sigma}_1 f)|_\Sigma, \end{aligned}$$

where \hat{c}_1^\pm are defined as in 4.2.5. Let us set

$$Tf := (\pi_1 \tilde{\varrho}_1^* \tilde{\sigma}_1 f)|_\Sigma.$$

Since

$$\tilde{D}_2 u_2 = \tilde{K}_{21} \tilde{D}_1 u_1 = \tilde{K}_{21} (\tilde{D}_1 + \pi_1) u_1 = \tilde{K}_{21} \tilde{\partial}_1^* \sigma_1 f = 0 \text{ in } \Omega^\pm,$$

we have $\tilde{c}_2^\mp \tilde{\partial}_2^\pm u_2 = 0$, hence

$$(7.2) \quad \begin{aligned} \tilde{c}_2^- \tilde{K}_{21\Sigma} \hat{c}_1^+ f + \tilde{c}_2^- R_\Sigma T f &= 0, \\ \tilde{c}_2^+ \tilde{K}_{21\Sigma} \hat{c}_1^- f - \tilde{c}_2^+ R_\Sigma T f &= 0. \end{aligned}$$

Since $\hat{c}_1^+ + \hat{c}_1^- = \mathbb{1}$ and $\tilde{c}_2^+ + \tilde{c}_2^- = \mathbb{1}$ we obtain by (7.2):

$$(7.3) \quad \begin{aligned} \tilde{c}_2^\pm \tilde{K}_{21\Sigma} f &= \tilde{c}_2^\pm \tilde{K}_{21\Sigma} \hat{c}_1^\pm f + \tilde{c}_2^\pm \tilde{K}_{21\Sigma} \hat{c}_1^\mp f \\ &= \tilde{K}_{21\Sigma} \hat{c}_1^\pm f - \tilde{c}_2^\mp \tilde{K}_{21\Sigma} \hat{c}_1^\pm f + \tilde{c}_2^\pm \tilde{K}_{21\Sigma} \hat{c}_1^\mp f \\ &= \tilde{K}_{21\Sigma} \hat{c}_1^\pm f \pm R_\Sigma T f. \end{aligned}$$

Let us now compute Tf . A routine computation shows that

$$\|\varphi_i\|_{\tilde{M}}^2 = \frac{32}{15} \|\psi_i\|_\Sigma^2, \quad \|\varphi_{jk}\|_{\tilde{M}}^2 = \frac{16}{15} \|\psi_{jk}\|_\Sigma^2.$$

Then

$$\tilde{\partial}_1^* \tilde{\sigma}_1 f = -\delta(s) \otimes f_1 + \delta'(s) \otimes f_0, \quad f = \begin{pmatrix} f_0 \\ f_1 \end{pmatrix},$$

and

$$\begin{aligned} \pi_1 \tilde{\partial}_1^* \tilde{\sigma}_1 f &= \sum_{1 \leq i \leq 4} \frac{1}{\|\varphi_i\|^2} \varphi_i (-(\varphi_i(0)|f_1)_\Sigma - (\partial_s \varphi_i(0)|f_0)_\Sigma) \\ &\quad + \sum_{1 \leq j < k \leq 4} \frac{1}{\|\varphi_{jk}\|^2} \varphi_{jk} (-(\varphi_{jk}(0)|f_1)_\Sigma - (\partial_s \varphi_{jk}(0)|f_0)_\Sigma). \end{aligned}$$

We have

$$\begin{aligned} -(\varphi_i(0)|f_1)_\Sigma - (\partial_s \varphi_i(0)|f_0)_\Sigma &= -(\psi_i|f_{1s})_\Sigma + (\vec{d}\psi_i|f_{0\Sigma})_\Sigma, \\ -(\varphi_{jk}(0)|f_1)_\Sigma - (\partial_s \varphi_{jk}(0)|f_0)_\Sigma &= -(\psi_{jk}|f_{1\Sigma})_\Sigma, \end{aligned}$$

hence

$$(7.4) \quad Tf = \begin{pmatrix} \frac{15}{32} \mathbb{1}_{\{3\}} (\vec{D}_{0,L}) (\vec{\delta} f_{0\Sigma} - f_{1s}) \\ -\frac{15}{16} \mathbb{1}_{\{4\}} (\vec{D}_{1,L}) f_{1\Sigma} \end{pmatrix}.$$

We can now complete the proof. Let us see when $R_\Sigma v \in \text{Ran } \tilde{K}_{21\Sigma} + \text{Ran } \tilde{K}_{20\Sigma}$. Recall that $\tilde{K}_{21\Sigma} = \tilde{I}_\Sigma \circ \hat{K}_{21\Sigma}$ where $\hat{K}_{21\Sigma}$ is associated to d and \tilde{I}_Σ to $\tilde{I} = \mathbb{1} - \frac{1}{4} K_{20} K_{20}^*$. Therefore $\text{Ran } \tilde{K}_{21\Sigma} + \text{Ran } \tilde{K}_{20\Sigma} = \text{Ran } \hat{K}_{21\Sigma} + \text{Ran } \tilde{K}_{20\Sigma}$, and $\hat{K}_{21\Sigma}$ is computed in (A.5).

We obtain that $\hat{K}_{21\Sigma} w + \tilde{K}_{20\Sigma} u = R_\Sigma v$ iff

- (1) $-w_{1s} + u_0 = 0$,
- (2) $-w_{1\Sigma} + \vec{d}w_{0s} = 0$,
- (3) $\vec{d}w_{0\Sigma} + u_0 h = 0$,
- (4) $-(\vec{D}_{0,L} - 3)w_{0s} + u_1 = \frac{1}{2}v_s$,
- (5) $\frac{1}{2}(-(\vec{D}_{1,L} - 4)w_{0\Sigma} + \vec{d}w_{1s}) = \frac{1}{2}v_\Sigma$,
- (6) $\vec{d}w_{1\Sigma} + w_{0s} h + u_1 h = -\frac{1}{2}v_s h$.

From (2) and (6) we obtain that

$$\vec{d}^2 w_{0s} + w_{0s} h + u_1 h = -\frac{1}{2} v_s h,$$

which after contracting with h gives

$$-(\vec{D}_{0,L} - 3)w_{0s} = -\frac{3}{2}v_s - 3u_1.$$

This implies that $\mathbb{1}_{\{3\}}(\vec{D}_{0,L})(v_s + 2u_1) = 0$. On the other hand applying $\mathbb{1}_{\{3\}}(\vec{D}_{0,L})$ to (4) we obtain that $\mathbb{1}_{\{3\}}(\vec{D}_{0,L})(v_s - 2u_1) = 0$, hence $\mathbb{1}_{\{3\}}(\vec{D}_{0,L})v_s = 0$.

Next, applying $\mathbb{1}_{\{4\}}(\vec{D}_{1,L})$ to (5), using that $\mathbb{1}_{\{4\}}(\vec{D}_{1,L}) \circ \vec{d} = 0$ we obtain that $\mathbb{1}_{\{4\}}(\vec{D}_{1,L})v_\Sigma = 0$.

Therefore the conditions

$$\begin{cases} \mathbb{1}_{\{3\}}(\vec{D}_{0,L})v_s = 0, \\ \mathbb{1}_{\{4\}}(\vec{D}_{1,L})v_\Sigma = 0, \end{cases}$$

are necessary for $R_\Sigma v \in \text{Ran } \tilde{K}_{21\Sigma} + \text{Ran } \tilde{K}_{20\Sigma}$.

Since $v = Tf$ for T defined in (7.4), we obtain that $f \in \text{Ker } \tilde{C}_{1\Sigma}$ is a necessary condition for $\tilde{c}_2^\pm \tilde{K}_{21\Sigma} f \in \text{Ran } \tilde{K}_{21\Sigma} + \text{Ran } \tilde{K}_{20\Sigma}$.

It is also a sufficient condition since if $f \in \text{Ker } \tilde{C}_{1\Sigma}$ then $v = Tf = 0$, so $R_\Sigma v \in \text{Ran } \tilde{K}_{21\Sigma} + \text{Ran } \tilde{K}_{20\Sigma}$.

Moreover if $Tf = 0$ then $\tilde{\varrho}_1^* \tilde{\sigma}_1 f \in \text{Ker } \tilde{D}_1^\perp$ hence $\hat{c}_1^\pm f = \tilde{c}_1^\pm f$ so $\tilde{c}_2^\pm \tilde{K}_{21\Sigma} f = \tilde{K}_{21\Sigma} \tilde{c}_1^\pm f$ by (7.3). This proves the first statement of the proposition.

Let us now prove the second statement of the proposition. Since $\text{Ker } \tilde{D}_1$ is finite dimensional we need to show that

$$(\tilde{K}_{21\Sigma}^\dagger \tilde{c}_2^\pm f_2 | q_1 f_1) = 0 \quad \forall f_1 \in (\tilde{\varrho}_1 \text{Ker } \tilde{D}_1)^{\mathfrak{q}_1}.$$

We have

$$(\tilde{K}_{21\Sigma}^\dagger \tilde{c}_2^\pm f_2 | q_1 f_1) = (f_2 | q_2 \tilde{c}_2^\pm K_{21\Sigma} f_1) = 0$$

if $f_1 \in (\tilde{\varrho}_1 \text{Ker } \tilde{D}_1)^{\mathfrak{q}_1} = \text{Ker } \tilde{C}_{1\Sigma}$ using the first statement of the proposition and the fact that $\tilde{K}_{21\Sigma}^\dagger f_2 = 0$. \square

Corollary 7.3. *The Cauchy surface covariances $\lambda_{2\Sigma}^\pm$ in Def. 7.1 do not satisfy the strong TT gauge invariance property but satisfy the weak TT gauge invariance property.*

7.3. Positivity. In this subsection we investigate the positivity of the Hermitian forms $\lambda_{2\Sigma}^\pm$ on \mathcal{E}_{TT} by determining their inertia indices. There is a slight inconvenience because $\lambda_{2\Sigma}^\pm = 0$ on $\text{Ran } c_2^\mp$, so strictly speaking we will determine the inertia indices of $\lambda_{2\Sigma}^\pm$ on $\text{Ran } c_2^\pm$.

Equivalently we could also look at the inertia indices of $\lambda_{2\Sigma}^+ + \lambda_{2\Sigma}^-$ on the whole of \mathcal{E}_{TT} .

We use the decomposition (6.5) of \mathcal{E}_{TT} given in Subsect. 6.2.

Since c_2^\pm commute with $\mathbb{1}_{\{\lambda\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2}$ and $\mathbb{1}_{\{\lambda\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2}^* q_2 = q_2 \mathbb{1}_{\{\lambda\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2}$, the three spaces above are orthogonal for $\lambda_{2\Sigma}^\pm$.

7.3.1. *Positivity on $\mathcal{E}_{\text{TT,reg}}$.* We first prove the positivity of $\lambda_{2\Sigma}^\pm$ on $\mathcal{E}_{\text{TT,reg}}$, recalling that $\mathcal{E}_{\text{TT,reg}} = \mathcal{E}_{\text{TT}} \cap \text{Ker } C_{2\Sigma}$. This is the most delicate point of this subsection.

Proposition 7.4. *We have*

$$\lambda_{2\Sigma}^\pm \geq 0 \text{ on } \mathcal{E}_{\text{TT,reg}}.$$

Proof. We prove only the ‘+’ case. It suffices to prove the Euclidean version, i.e.

$$(7.5) \quad (f_2 | q_2 \tilde{I}_\Sigma \tilde{c}_2^+ f_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} \geq 0, \quad f_2 \in \tilde{\mathcal{E}}_{\text{TT}} \cap \text{Ker } \tilde{C}_{2\Sigma}.$$

We will use the restricted gauge invariance stated in Prop. 7.2 to prove (7.5).

Step 1: let $f_2 \in \tilde{\mathcal{E}}_{\text{TT,reg}} = \tilde{\mathcal{E}}_{\text{TT}} \cap \text{Ker } \tilde{C}_{2\Sigma}$ and $u_2 = -(\tilde{D}_2^{-1} \tilde{\varrho}_2^* \tilde{\sigma}_2 f_2)|_{\Omega^+}$, and let

$$(7.6) \quad \tilde{u}_2 = u_2 - \tilde{K}_{21} u_1,$$

for some $u_1 \in \overline{C^\infty}(\Omega^+; \tilde{V}_1)$ with $\tilde{D}_1 u_1 = 0$ in Ω^+ to be determined later. We set $\tilde{f}_2 = \tilde{\varrho}_2^+ \tilde{u}_2$, $f_1 = \tilde{\varrho}_1^+ u_1$. Then

$$\tilde{c}_2^+ f_2 = \tilde{\varrho}_2^+ u_2 = \tilde{f}_2 + \tilde{K}_{21\Sigma} f_1,$$

since $\tilde{D}_1 u_1 = 0$ in Ω^+ . We have

$$\begin{aligned} (f_2 | q_2 \tilde{I}_\Sigma \tilde{c}_2^+ f_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} &= (f_2 | q_2 \tilde{c}_2^+ f_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} \\ &= (f_2 | q_2 \tilde{f}_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} + (f_2 | q_2 \tilde{K}_{21\Sigma} f_1)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} = (f_2 | q_2 \tilde{f}_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2}, \end{aligned}$$

where in the first line we use that $\tilde{I}_\Sigma f_2 = f_2$ since $f_2 \in \text{Ker } \tilde{K}_{20\Sigma}^\dagger$, and in the second line that $f_2 \in \text{Ker } \tilde{K}_{21\Sigma}^\dagger$.

Next, since $\tilde{f}_2 = \tilde{\varrho}_2^+ \tilde{u}_2$ with $\tilde{D}_2 \tilde{u}_2 = 0$ in Ω^+ , we have $\tilde{f}_2 = \tilde{c}_2^+ \tilde{f}_2$, see Prop. 4.3, hence

$$\begin{aligned} (f_2 | q_2 \tilde{f}_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} &= (f_2 | q_2 \tilde{c}_2^+ \tilde{f}_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} = (\tilde{c}_2^+ f_2 | q_2 \tilde{f}_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} \\ &= (\tilde{f}_2 | q_2 \tilde{f}_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} + (\tilde{K}_{21\Sigma} f_1 | q_2 \tilde{f}_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} \\ &= (\tilde{f}_2 | q_2 \tilde{f}_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} + (\tilde{K}_{21\Sigma} f_1 | q_2 \tilde{c}_2^+ f_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} \\ &\quad - (\tilde{K}_{21\Sigma} f_1 | q_2 \tilde{K}_{21\Sigma} f_1)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} \\ &= (\tilde{f}_2 | q_2 \tilde{f}_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} + (\tilde{c}_2^+ \tilde{K}_{21\Sigma} f_1 | q_2 f_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} \\ &\quad - (\tilde{K}_{21\Sigma}^\dagger \tilde{K}_{21\Sigma} f_1 | q_1 f_1)_{\tilde{V}_1(\Sigma) \otimes \mathbb{C}^2} \\ &= (\tilde{f}_2 | q_2 \tilde{f}_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} + (\tilde{c}_2^+ \tilde{K}_{21\Sigma} f_1 | q_2 f_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2}, \end{aligned}$$

since $\tilde{K}_{21\Sigma}^\dagger \tilde{K}_{21\Sigma} = 0$. By Prop. 7.2, if $f_1 \in \text{Ker } \tilde{C}_{1\Sigma}$ then $\tilde{c}_2^+ \tilde{K}_{21\Sigma} f_1 \in \text{Ran } \tilde{K}_{21\Sigma} + \text{Ran } \tilde{K}_{20\Sigma}$ hence

$$(f_2 | q_2 \tilde{c}_2^+ \tilde{K}_{21\Sigma} f_1)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} = 0.$$

In conclusion if $f_1 = \tilde{\varrho}_1^+ u_1 \in \text{Ker } \tilde{C}_{1\Sigma}$, then

$$(f_2 | \tilde{I}_\Sigma q_2 \tilde{c}_2^+ f_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} = (\tilde{f}_2 | q_2 \tilde{f}_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2}.$$

Step 2: since $\tilde{f}_2 = \tilde{\partial}_2^+ \tilde{u}_2$ with $\tilde{D}_2 \tilde{u}_2 = 0$ in Ω^+ we deduce from Lemma 4.1 2) that:

$$(7.7) \quad 2Q(\tilde{u}_2, \tilde{u}_2)_{\Omega^+} = (\tilde{f}_2 | \tilde{q}_2 \tilde{f}_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2}.$$

From the identity (5.4) we obtain that the sesquilinear form Q associated to \tilde{D}_2 is

$$(7.8) \quad Q(\tilde{u}_2, \tilde{u}_2)_{\tilde{M}} = \int_{\tilde{M}} (\tilde{\nabla}_a \tilde{u}_2 \tilde{g}^{ab} \tilde{\nabla}_b \tilde{u}_2 + 2\|\tilde{u}_2\|^2 - |(\tilde{g}|\tilde{u}_2)_{\tilde{V}_2}|^2) d\text{vol}_{\tilde{g}}.$$

Let $v_0 = \tilde{K}_{20}^* \tilde{u}_2 = (\tilde{g}|\tilde{u}_2)_{\tilde{V}_2}$. Then $\tilde{D}_0 v_0 = \tilde{K}_{20}^* \tilde{D}_2 \tilde{u}_2 = 0$ in Ω^+ . If moreover $v_0|_{\Sigma} = 0$, then the extension of v_0 to \mathbb{S}^4 by parity in s is in the kernel of \tilde{D}_0 hence is a constant. Therefore if $(\tilde{g}|\tilde{u}_2)_{\tilde{V}_2}|_{\Sigma} = 0$ we have $(\tilde{g}|\tilde{u}_2)_{\tilde{V}_2} = 0$ in Ω^+ .

It follows then from (7.7), (7.8) that if $(g|\tilde{u}_2)_{\tilde{V}_2}|_{\Sigma} = 0$, one has

$$(\tilde{f}_2 | \tilde{q}_2 \tilde{f}_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} \geq 0.$$

If moreover $\partial_s \tilde{u}_{2s\Sigma}|_{\Sigma} = 0$ then, see (4.4.1) and 4.4.2:

$$(\tilde{f}_2 | \tilde{q}_2 \tilde{f}_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} = (\tilde{f}_2 | q_2 \tilde{f}_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2}.$$

In conclusion, recalling that \tilde{u}_2 is defined in (7.6), if we can find $u_1 \in \overline{C^\infty}(\Omega^+; \tilde{V}_1)$ such that

$$(7.9) \quad \begin{cases} \tilde{D}_1 u_1 = 0 \text{ in } \Omega^+, \\ i) \quad (\tilde{g}|u_2 - \tilde{K}_{21} u_1)|_{\Sigma} = 0, \\ ii) \quad \partial_s (u_2 - \tilde{K}_{21} u_1)_{s\Sigma}|_{\Sigma} = 0, \\ iii) \quad C_{1\Sigma} \tilde{\partial}_1^+ u_1 = 0, \end{cases}$$

then

$$(f_2 | \tilde{I}_\Sigma q_2 \tilde{c}_2^+ f_2)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} \geq 0.$$

Step 3: We now examine the boundary value problem (7.9), leaving aside condition *iii*) for the moment. First we have $\tilde{K}_{20\Sigma}^\dagger \tilde{c}_2^+ f_2 = \tilde{c}_0^+ \tilde{K}_{20\Sigma}^\dagger f_2 = 0$ hence $(\tilde{g}|u_2)|_{\Sigma} = 0$. We obtain that

$$(\tilde{g}|u_2 - \tilde{K}_{21} u_1)|_{\Sigma} = -\delta u_1|_{\Sigma} = \partial_s u_{1s}|_{\Sigma} - \vec{\delta} u_{1\Sigma}|_{\Sigma}.$$

On the other hand we have

$$\partial_s (\tilde{K}_{21} u_1)_{s\Sigma}|_{\Sigma} = \frac{1}{2} (\partial_s^2 u_{1\Sigma} + \vec{d} \partial_s u_{1s} + 2u_{1\Sigma})|_{\Sigma},$$

and since $\tilde{D}_1 u_1 = 0$ in Ω^+ we have (see the computations in A.2.3) $\partial_s^2 u_{1\Sigma}|_{\Sigma} = (\vec{D}_{1,L} - 6)u_{1\Sigma}|_{\Sigma}$.

Therefore, the boundary value problem (7.9) (with condition *iii*) removed) can be written as

$$\begin{cases} \tilde{D}_1 u_1 = 0 \text{ in } \Omega^+, \\ \partial_s u_{1s}|_{\Sigma} - \vec{\delta} u_{1\Sigma}|_{\Sigma} = 0, \\ (\vec{D}_{1,L} - 4)u_{1\Sigma}|_{\Sigma} + \vec{d} \partial_s u_{1s}|_{\Sigma} = 2\partial_s u_{2s\Sigma}|_{\Sigma}, \end{cases}$$

or equivalently

$$(7.10) \quad \begin{cases} \tilde{D}_1 u_1 = 0 \text{ in } \Omega^+, \\ \partial_s u_{1s} \upharpoonright_\Sigma - \vec{\delta} u_{1\Sigma} \upharpoonright_\Sigma = 0, \\ (\vec{D}_{1,L} + \vec{d}\vec{\delta} - 4) u_{1\Sigma} \upharpoonright_\Sigma = 2 \partial_s u_{2s\Sigma} \upharpoonright_\Sigma. \end{cases}$$

Note that $[\vec{D}_{1,L}, \vec{d}\vec{\delta}] = 0$ since $\vec{D}_{1,L} = \vec{d}_a \vec{\delta}_a + \vec{\delta}_a \vec{d}_a$ and $\vec{d}\vec{\delta} = \vec{d}_a \vec{\delta}_a$ on 1-forms. From the results in [4, Sect. 3.6], we obtain that $\text{Ker}(\vec{D}_{1,L} + \vec{d}\vec{\delta} - 4) = \text{Vect}\{\psi_{jk} : 1 \leq j < k \leq 4\}$.

Therefore a necessary condition to solve (7.10) is

$$(7.11) \quad \mathbb{1}_{\{4\}}(\vec{D}_{1,L}) \partial_s u_{2s\Sigma} \upharpoonright_\Sigma = 0.$$

If (7.11) is satisfied, then (7.10) is equivalent to:

$$(7.12) \quad \begin{cases} \tilde{D}_1 u_1 = 0 \text{ in } \Omega^+, \\ \partial_s u_{1s} \upharpoonright_\Sigma = 2 \vec{\delta} (\vec{D}_{1,L} + \vec{d}\vec{\delta} - 4)^{-1} \partial_s u_{2s\Sigma} \upharpoonright_\Sigma, \\ u_{1\Sigma} \upharpoonright_\Sigma = 2 (\vec{D}_{1,L} + \vec{d}\vec{\delta} - 4)^{-1} \partial_s u_{2s\Sigma} \upharpoonright_\Sigma + v_\Sigma, \end{cases}$$

for $v_\Sigma \in \text{Vect}\{\psi_{jk} : 1 \leq j < k \leq 4\}$ arbitrary, where $P^{-1} : \text{Ker } P^\perp \rightarrow \text{Ker } P^\perp$ is the pseudo-inverse of P .

The boundary value problem (7.12) is the Dirichlet realization of \tilde{D}_1 in Ω^+ , studied in Lemma A.1. We find that (7.12) is solvable iff

$$\mathbb{1}_{\{3\}}(\vec{D}_{1,L}) \partial_s u_{2s\Sigma} \upharpoonright_\Sigma = 0.$$

To sum up, we have shown that (7.10) is solvable iff

$$(7.13) \quad \begin{cases} i) \quad \mathbb{1}_{\{4\}}(\vec{D}_{1,L}) \partial_s u_{2s\Sigma} \upharpoonright_\Sigma = 0, \\ ii) \quad \mathbb{1}_{\{3\}}(\vec{D}_{1,L}) \partial_s u_{2s\Sigma} \upharpoonright_\Sigma = 0. \end{cases}$$

The solution is unique modulo $\text{Vect}\{\varphi_i, \varphi_{jk}\}$, since φ_i resp. φ_{jk} solves (7.12) for $\partial_s u_{2s\Sigma} \upharpoonright_\Sigma = 0$ and $v_\Sigma = 0$ resp. $v_\Sigma = \psi_{jk}$.

Since $f_2 \in \tilde{\mathcal{E}}_{\text{TT,reg}}$ we have $f_2 \in \text{Ker } \mathbb{1}_{\{\lambda\}}(\hat{D}_2) \otimes \mathbb{C}^2$ for $\lambda = 3, 4$ by Lemma 6.5. We have $\tilde{\varrho}_2^+ u_2 = \tilde{c}_2^+ f_2$ and by (7.1) this implies that $\tilde{\varrho}_2^+ u_2 \in \text{Ker } \mathbb{1}_{\{\lambda\}}(\hat{D}_2) \otimes \mathbb{C}^2$ for $\lambda = 3, 4$, which implies (7.13).

Therefore for $f_2 \in \tilde{\mathcal{E}}_{\text{TT,reg}}$ the boundary value problem (7.10) is solvable. We choose for u_1 the solution of (7.12) with $v_\Sigma = 0$.

Step 4: to complete the discussion we have to analyze the extra condition *iii*) in (7.9). The condition

$$\mathbb{1}_{\{3\}}(\vec{D}_{0,L})(\partial_s u_{1s} \upharpoonright_\Sigma + \vec{\delta} u_{1\Sigma} \upharpoonright_\Sigma) = 0$$

is satisfied if (7.13) holds since

$$\partial_s u_{1s} \upharpoonright_\Sigma + \vec{\delta} u_{1\Sigma} \upharpoonright_\Sigma = 4 \vec{\delta} (\vec{D}_{1,L} + \vec{d}\vec{\delta} - 4)^{-1} \partial_s u_{2s\Sigma} \upharpoonright_\Sigma$$

and

$$\begin{aligned} \mathbb{1}_{\{3\}}(\vec{D}_{0,L}) \vec{\delta} (\vec{D}_{1,L} + \vec{d}\vec{\delta} - 4)^{-1} &= \vec{\delta} \mathbb{1}_{\{3\}}(\vec{D}_{1,L}) (\vec{D}_{1,L} + \vec{d}\vec{\delta} - 4)^{-1} \\ &= \frac{1}{2} \vec{\delta} \mathbb{1}_{\{3\}}(\vec{D}_{1,L}), \end{aligned}$$

and (7.13) is satisfied.

It remains to discuss the condition:

$$\mathbb{1}_{\{4\}}(\vec{D}_{1,L})\partial_s u_{1\Sigma}|_\Sigma = 0.$$

Let us set $v = \mathbb{1}_{\{4\}}(\hat{D}_1)u_1$, where \hat{D}_1 is defined in Subsect. 5.3. Since \hat{D}_1 acts only in variables on Σ and commutes with \tilde{D}_1 , we have

$$\begin{cases} \tilde{D}_1 v = 0 \text{ in } \Omega^+, \\ v|_\Sigma = \mathbb{1}_{\{4\}}(\hat{D}_1)u_1|_\Sigma, \\ \partial_s v|_\Sigma = \mathbb{1}_{\{4\}}(\hat{D}_1)\partial_s u_1|_\Sigma. \end{cases}$$

Using (7.12) and (5.8) we obtain that:

$$\begin{cases} \tilde{D}_1 v = 0 \text{ in } \Omega^+, \\ \partial_s v|_\Sigma = 0, \\ v|_\Sigma = 0, \end{cases}$$

which by Lemma A.1 implies that $v \in \text{Vect}\{\varphi_i : 1 \leq i \leq 4\}$. Since $\mathbb{1}_{\{4\}}(\hat{D}_1)\varphi_i = 0$ this implies that $v = 0$. Therefore the condition

$$\mathbb{1}_{\{4\}}(\vec{D}_{1,L})\partial_s u_{1\Sigma}|_\Sigma = 0$$

is satisfied and $f_1 = \tilde{\varrho}_1^+ u_1$ belongs to $\text{Ker } C_{1\Sigma}$. \square

7.3.2. Positivity on $\mathcal{E}_{\text{TT},3}$ and $\mathcal{E}_{\text{TT},4}$. It remains to investigate $\lambda_{2\Sigma}^\pm$ on $\mathcal{E}_{\text{TT},3}$ and $\mathcal{E}_{\text{TT},4}$.

Proposition 7.5. *The covariances $\lambda_{2\Sigma}^\pm$ are negative on $\mathcal{E}_{\text{TT},4}$, and $\bar{f}\lambda_{2\Sigma}^\pm f = 0$ implies that $c_2^\pm f = 0$ for $f \in \mathcal{E}_{\text{TT},4}$. In particular $\lambda_{2\Sigma}^+ + \lambda_{2\Sigma}^-$ is negative definite on $\mathcal{E}_{\text{TT},4}$.*

Proof. Let $f \in \mathcal{E}_{\text{TT},4}$. Then

$$\bar{f}\lambda_{2\Sigma}^\pm f = (c_2^\pm f|q_2 c_2^\pm f)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} = -(c_2^\pm f|\tilde{q}_2 c_2^\pm f)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2}.$$

In fact the first equality follows from Prop. 4.4 and the second follows from the fact that $q_2 = -\tilde{q}_2$ on $\mathcal{E}_{\text{TT},4}$. Let us set $u = \tilde{D}_2^{-1}\tilde{\varrho}_2^* \tilde{\sigma} f$ so that $\tilde{c}_2^\pm f = \mp \tilde{\varrho}_2^+ f$. By Lemma 4.1 (2) and the expression in (7.8) for $Q(u, u)$ we obtain using that $(\tilde{g}|u)_{\tilde{V}_2} = 0$ that

$$(\tilde{c}_2^\pm f|\tilde{q}_2 c_2^\pm f)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} = \pm 2 \int_{\Omega^+} (\tilde{\nabla}_a \bar{u} \tilde{g}^{ab} \tilde{\nabla}_b u + 2\|u\|^2) d\text{vol}_{\tilde{g}}.$$

This implies that $(\tilde{c}_2^\pm f|\tilde{q}_2 c_2^\pm f)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} \geq 0$ and if $(\tilde{c}_2^\pm f|\tilde{q}_2 c_2^\pm f)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} = 0$ then $u = 0$ in Ω^\pm , hence $\tilde{c}_2^\pm f = 0$ as claimed. \square

Proposition 7.6. *The covariances $\lambda_{2\Sigma}^\pm$ are positive on $\mathcal{E}_{\text{TT},3}$ and $\bar{f}\lambda_{2\Sigma}^\pm f = 0$ implies that $c_2^\pm f = 0$ for $f \in \mathcal{E}_{\text{TT},3}$. In particular $\lambda_{2\Sigma}^+ + \lambda_{2\Sigma}^-$ is positive definite on $\mathcal{E}_{\text{TT},3}$.*

Proof. Let $f \in \mathcal{E}_{\text{TT},3}$ and $g = c_2^\pm f$. We have $g \in \text{Ran } \mathbb{1}_{\{3\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2}$ and $g \in \text{Ker } \tilde{K}_{20\Sigma}^\dagger$. Using (5.8) this implies that

$$(7.14) \quad g_{iss} = -3u_i, \quad g_{is\Sigma} = \vec{d}v_i, \quad g_{i\Sigma\Sigma} = u_i h, \quad u_i, v_i \in \text{Ker}(\vec{D}_{0,L} - 3),$$

for $i = 0, 1$. Moreover since $f \in \text{Ker } \tilde{K}_{21\Sigma}^\dagger$ we have $\tilde{K}_{21\Sigma}^\dagger g \in \tilde{\varrho}_1 \text{Ker } \tilde{D}_1$, see Prop. 7.2. We use Lemma 5.3 and the expression of $\tilde{\varrho}_1 \text{Ker } \tilde{D}_1$ given in the proof of Lemma 6.4. We obtain using also (7.14) that

$$\begin{cases} 1) & -6u_1 + 6v_0 = \alpha, \\ 2) & 2\vec{d}v_1 - 2\vec{d}u_0 = \beta, \\ 3) & 6v_1 - 6u_0 = 0, \\ 4) & -2\vec{d}v_0 - 2\vec{d}u_1 = \vec{d}\alpha, \end{cases}$$

for $\alpha \in \text{Vect}\{\psi_i\}$ and $\beta \in \text{Vect}\{\psi_{jk}\}$. This is equivalent to $v_1 = u_0$, $\vec{d}v_0 = \frac{1}{2}\vec{d}u_1$ and hence

$$g_{iss} = -3u_i, \quad g_{i\Sigma\Sigma} = u_i h, \quad g_{0s\Sigma} = \frac{1}{2}\vec{d}u_1, \quad g_{1s\Sigma} = \vec{d}u_0.$$

We then compute:

$$\begin{aligned} \bar{f} \cdot \lambda_{2\Sigma}^\pm f &= \pm(f|q_2 c_2^\pm f)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} = (g|q_2 g)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} \\ &= 2(3u_1|3u_0)_{\tilde{V}_0(\Sigma)} + 2(3u_0|3u_1)_{\tilde{V}_0(\Sigma)} - 4(\vec{d}u_0|\frac{1}{2}\vec{d}u_1)_{\tilde{V}_1(\Sigma)} \\ &\quad - 4(\frac{1}{2}\vec{d}u_1|\vec{d}u_0)_{\tilde{V}_1(\Sigma)} + (u_1 h|u_0 h)_{\tilde{V}_2(\Sigma)} + (u_0 h|u_1 h)_{\tilde{V}_2(\Sigma)} \\ &= 18(u_1|u_0)_{\tilde{V}_0(\Sigma)} + 18(u_0|u_1)_{\tilde{V}_0(\Sigma)}, \end{aligned}$$

using that $\vec{d}\vec{u}_i = 3u_i$. Similarly we compute $(g|\tilde{q}_2 g)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2}$ and obtain

$$(g|\tilde{q}_2 g)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} = 30(u_1|u_0)_{\tilde{V}_0(\Sigma)} + 30(u_0|u_1)_{\tilde{V}_0(\Sigma)},$$

hence

$$\bar{f} \cdot \lambda_{2\Sigma}^\pm f = \frac{5}{3}(g|\tilde{q}_2 g)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2}.$$

The same argument as in the proof of Prop. 7.5 shows that $(g|\tilde{q}_2 g)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} \geq 0$ and that $(g|\tilde{q}_2 g)_{\tilde{V}_2(\Sigma) \otimes \mathbb{C}^2} = 0$ implies $g = 0$. This completes the proof of the proposition. \square

7.4. Invariance under symmetries. We now discuss the invariance of ω_{eucl} under the isometry group $O(1, 4)$.

Proposition 7.7. *The pseudo-state ω_{eucl} is invariant under the action of $SO^\uparrow(1, 4)$ and of $O(4)$.*

Proof. Let X be a generator of $so(1, 4)$ viewed as a Killing vector field of dS^4 . We denote by $\phi(\cdot)$ its associated flow on $C^\infty(dS^4; V_2)$ and define its generator L by $\phi(\sigma) =: e^{i\sigma L}$. Note that L commutes with D_2 and I . In particular we can define its Cauchy surface version L_Σ by

$$L_\Sigma = \varrho_2 L U_2.$$

Clearly $[L_\Sigma, I_\Sigma] = 0$ and $L_\Sigma^* q_\Sigma = q_\Sigma L_\Sigma$. The Wick rotation of X denoted by \tilde{X} is a Killing vector field of \mathbb{S}^4 . We use the notation introduced in Subsect. 4.3. By Lemma 4.8, \tilde{L}_Σ commutes with the Calderón projectors \tilde{c}_2^\pm . By analytic continuation, this implies that L_Σ commutes with the Hadamard projectors c_2^\pm , hence that

$$L_\Sigma^* \lambda_{2\Sigma}^\pm = \lambda_{2\Sigma}^\pm L_\Sigma,$$

i.e. $\lambda_{2\Sigma}^\pm$ and the associated pseudo-state ω_{eucl} are invariant under the action of $SO^\uparrow(1, 4)$.

The action of $O(4)$ on tensors commutes with Wick rotation, since it preserves the variable t . It is immediate that the Calderón projectors \tilde{c}_2^\pm are invariant under the action of $O(4)$ on Cauchy data, which implies as above that $\lambda_{2\Sigma}^\pm$ and ω_{eucl} are invariant under the action of $O(4)$. \square

7.4.1. Invariance under time reversal. We recall that time reversal Z is defined in 6.3.1. We denote by Z_Σ its Cauchy surface version, i.e. $Z_\Sigma = \varrho_2 Z U_2$.

Proposition 7.8. *The Euclidean vacuum pseudo state ω_{eucl} is invariant under time reversal, i.e.*

$$\overline{Z_\Sigma f} \cdot \lambda_{2\Sigma}^\pm Z_\Sigma g = \overline{g} \cdot \lambda_{2\Sigma}^\pm f, \quad f, g \in C_0^\infty(\Sigma; V_2 \otimes \mathbb{C}^2).$$

Proof. We have

$$(Z_\Sigma f)_{itt} = \overline{f_{itt}}, \quad (Z_\Sigma f)_{it\Sigma} = -\overline{f_{it\Sigma}}, \quad (Z_\Sigma f)_{i\Sigma\Sigma} = \overline{f_{i\Sigma\Sigma}}, \quad i = 0, 1.$$

We claim that $Z_\Sigma^{-1} c_2^\pm Z_\Sigma = c_2^\pm$, which is equivalent to $\tilde{Z}_\Sigma^{-1} \tilde{c}_2^\pm \tilde{Z}_\Sigma = \tilde{c}_2^\pm$ for $\tilde{Z}_\Sigma = \mathcal{F}_\Sigma Z_\Sigma \mathcal{F}_\Sigma^{-1}$. Using the expression of \mathcal{F}_2 in (3.1) we obtain that

$$(7.15) \quad \tilde{Z}_\Sigma f = -\overline{f}.$$

Since the Wick rotated operator \tilde{D}_2 is real we obtain that $\tilde{Z}_\Sigma^{-1} \tilde{c}_2^\pm \tilde{Z}_\Sigma = \tilde{c}_2^\pm$. Using then that Z_Σ is anti-unitary for $\mathbf{q}_{I,2}$, this implies the proposition. \square

Remark 7.9. *Let us explain our choice for the time reversal, in connection with real fields. We follow the exposition given in [12, Sect. 4.7]. Let (\mathcal{X}, σ) be a real symplectic space. A quasi-free state ω on $\text{CCR}(\mathcal{X}, \sigma)$ is determined by its covariance $\eta \in L_s(\mathcal{X}, \mathcal{X}')$ defined by $\omega(\phi(x_1)\phi(x_2)) = x_1 \eta x_2 + \frac{1}{2} x_1 \sigma x_2$.*

Then $(\mathbb{C}\mathcal{X}, \mathbf{q})$ is a Hermitian space, where $\mathbf{q} = i\sigma_{\mathbb{C}}$ and $\sigma_{\mathbb{C}}$ is the sesquilinear extension of σ . The extension $\tilde{\omega}$ of ω to $\text{CCR}(\mathbb{C}\mathcal{X}, i\sigma_{\mathbb{C}})$ has covariances $\lambda^\pm = \eta_{\mathbb{C}} \pm \frac{1}{2}\mathbf{q}$.

We apply this to $\mathcal{X} = C^\infty(\Sigma; V_{2,\mathbb{R}} \otimes \mathbb{R}^2)$, the space of Cauchy data of real tensors. We obtain that

$$\overline{Z_\Sigma f} \cdot \eta_{\mathbb{C}} Z_\Sigma g = \overline{f \cdot \eta_{\mathbb{C}} g},$$

hence if f, g are real tensors we have $Z_\Sigma f \cdot \eta Z_\Sigma g = f \cdot \eta g$, which is the usual meaning of invariance under time reversal.

7.5. α -vacua. We now define α -vacua, obtained from ω_{eucl} by a Bogoliubov transformation, see [1] for the case of scalar fields. Usually α -vacua are defined via a mode expansion. It is very easy to define them directly, as we will do in this subsection. We refer the reader to [12, Sect. 6.4] for a discussion of the mode expansion construction, in connection with the approach used in this paper.

We denote by $S : C_0^\infty(dS^4; V_k)$ the *linear* time reversal

$$S u_k = \tau^* u_k, \quad u_k \in C_0^\infty(dS^4; V_k).$$

It is often called the *Racah time reversal*.

Again S commutes with D_k, K_{ij}, K_{ij}^* . Denoting by S_Σ is Cauchy surface version, we see that S_Σ is well defined on \mathcal{E}_{TT} and on the quotient space $\mathcal{E}_{\text{TT}}/\mathcal{F}_{\text{TT}}$.

We also see that $S_\Sigma^* \mathbf{q}_{I,2} S_\Sigma = -\mathbf{q}_{I,2}$, which implies that $U_\alpha = e^{\alpha S_\Sigma} = \cosh \alpha \mathbb{1} + \sinh \alpha S_\Sigma$ is unitary for $\mathbf{q}_{I,2}$ for $\alpha \in \mathbb{R}$.

Definition 7.10. *The α -vacuum pseudo-state $\omega_{\alpha, \text{eucl}}$ for $\alpha \in \mathbb{R}$ is defined by the covariances*

$$\lambda_{\alpha, 2\Sigma}^\pm := U_\alpha^* \lambda_{2\Sigma}^\pm U_\alpha.$$

We now discuss the properties of $\lambda_{\alpha, 2\Sigma}^\pm$.

Proposition 7.11. *$\lambda_{\alpha, 2\Sigma}^\pm$ satisfy the same properties as $\lambda_{2\Sigma}^\pm$ expect for the Hadamard condition.*

Proof. The fact that $\lambda_{\alpha, 2\Sigma}^+ - \lambda_{\alpha, 2\Sigma}^- = \mathbf{q}_{I,2}$ follows from the fact that U_α is unitary for $\mathbf{q}_{I,2}$.

U_α commutes with the spectral projections $\mathbb{1}_{\{\lambda\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2}$, hence preserves the subspaces $\mathcal{E}_{\text{TT,reg}}$, $\mathcal{E}_{\text{TT},3}$ and $\mathcal{E}_{\text{TT},4}$, so the positivity properties of $\lambda_{\alpha, 2\Sigma}^\pm$ are unchanged.

The invariance properties under the action of $O(1,4)$ are also unchanged. Finally the Hadamard condition in Prop. 2.10 *iv*) is not satisfied, because S exchanges \mathcal{N}^+ with \mathcal{N}^- . \square

8. THE MODIFIED EUCLIDEAN VACUUM

In this section we show how to modify the Euclidean pseudo state ω_{eucl} to obtain a true, positive and gauge invariant state ω_{mod} on \mathcal{E}_{TT} . One can view this modification as imposing an additional gauge fixing condition. The price to pay is that ω_{mod} is no more invariant under the full isometry group $O(1,4)$, but only under the subgroup $O(4)$.

We recall that the projection $\pi : \mathcal{E}_{\text{TT}} \rightarrow \mathcal{E}_{\text{TT,reg}}$ was defined in Prop. 6.6.

Definition 8.1. *We set*

$$\bar{f} \cdot \lambda_{2\Sigma, \text{mod}}^\pm f := \overline{\pi f} \cdot \lambda_{2\Sigma}^\pm \pi f, \quad f \in \mathcal{E}_{\text{TT}}.$$

Theorem 8.2. *We have:*

- (1) $\lambda_{2\Sigma, \text{mod}}^\pm = \lambda_{2\Sigma, \text{mod}}^{\pm*}$, $\lambda_{2\Sigma, \text{mod}}^+ - \lambda_{2\Sigma, \text{mod}}^- = \mathbf{q}_{I,2}$ on $\text{Ker } K_{21\Sigma}^\dagger$.
- (2) $\lambda_{2\Sigma, \text{mod}}^\pm \geq 0$ on \mathcal{E}_{TT} .

Therefore $\lambda_{2\Sigma, \text{mod}}^\pm$ are the covariances of a state ω_{mod} on \mathcal{E}_{TT} . Moreover,

- (3) ω_{mod} is a Hadamard state.
- (4) ω_{mod} is fully gauge invariant, i.e.

$$\bar{f} \cdot \lambda_{2\Sigma, \text{mod}}^\pm K_{21\Sigma} g = 0 \quad \forall f \in \mathcal{E}_{\text{TT}}, \quad g \in C^\infty(\Sigma; V_1 \otimes \mathbb{C}^2).$$

- (5) ω_{mod} is invariant under the action of $O(4)$.

Proof. The fact that $\lambda_{2\Sigma, \text{mod}}^\pm$ are Hermitian is obvious. We have

$$\lambda_{2\Sigma, \text{mod}}^+ - \lambda_{2\Sigma, \text{mod}}^- = \pi^* \mathbf{q}_{I,2} \pi,$$

and since $\pi f = f + K_{21\Sigma} Rf$, we have $\pi^* \mathbf{q}_{I,2} \pi = \mathbf{q}_{I,2}$ on $\text{Ker } K_{21\Sigma}^\dagger$. This completes the proof of (1).

Property (2) follows from the fact that $\pi f \in \mathcal{E}_{\text{TT,reg}}$ and $\lambda_{2\Sigma}^\pm \geq 0$ on $\mathcal{E}_{\text{TT,reg}}$ by Prop. 7.4.

Property (3) follows from the fact that the spacetime two-point functions $\Lambda_{2\text{eucl}}^\pm$ and $\Lambda_{2\Sigma, \text{mod}}^\pm$ associated to ω_{eucl} and ω_{mod} differ by a smoothing error by Prop. 6.6.

To prove (4) we apply (2) of Prop. 6.6. We obtain that $\pi K_{21\Sigma}g \in K_{21\Sigma}\mathcal{K}_\Sigma^{\text{q1}}$, hence $c_2^+ \pi K_{21\Sigma}g \in \text{Ran } K_{21\Sigma}$ by Prop. 7.2. Since $\pi f \in \text{Ker } K_{21\Sigma}^\dagger$ this implies that $\tilde{f} \cdot \lambda_{2\Sigma, \text{mod}}^\pm K_{21\Sigma}g = 0$ as claimed.

Finally let α be the (linear) action of an element of $O(4)$ on tensors and α_Σ its Cauchy surface version. Since the action of $O(4)$ preserves $t = cst$, we have $\alpha_\Sigma = \begin{pmatrix} \alpha & 0 \\ 0 & \alpha \end{pmatrix}$, which implies that α_Σ commutes with the operator π in Prop. 6.6.

By Prop. 7.7, $\alpha_\Sigma^* \lambda_{2\Sigma}^\pm \alpha_\Sigma = \lambda_{2\Sigma}^\pm$, hence also $\alpha_\Sigma^* \lambda_{2\Sigma, \text{mod}}^\pm \alpha_\Sigma = \lambda_{2\Sigma, \text{mod}}^\pm$, which proves the invariance of ω_{mod} under $O(4)$. \square

Remark 8.3. *One can wonder if ω_{mod} is invariant under the action of $SO^\uparrow(1,4)$. To check this, one chooses a Killing vector field X on dS^4 which is not tangent to Σ . The Wick rotation \tilde{X} of X is then of the form $\tilde{X} = \tilde{g}^{-1} \varphi_i$, where φ_i is one of the Killing 1-forms on \mathbb{S}^4 . To compute the associated operator L_Σ defined in the proof of Prop. 7.7, one can as well compute its Euclidean version $\tilde{L}_\Sigma = \mathcal{F}_{2\Sigma} \circ L_\Sigma \circ \mathcal{F}_\Sigma^{-1}$ associated to the Killing vector field \tilde{X} on \mathbb{S}^4 . Using the expression of \tilde{L}_Σ one can then check that $\mathcal{E}_{\text{TT,reg}}$ is not invariant under the action of L_Σ , so the projection π does not commute with the action of e^{sL_Σ} . Therefore ω_{mod} is not invariant under the action of e^{sL_Σ} .*

APPENDIX A. AUXILIARY PROOFS

A.1. An elliptic boundary value problem. In what follows we work in the setting of Sects. 2–6.

Lemma A.1. *Let $\mathcal{P} : \overline{C^\infty}(\Omega^+; \tilde{V}_1) \rightarrow \overline{C^\infty}(\Omega^+; \tilde{V}_1) \oplus C^\infty(\Sigma; \tilde{V}_1)$ given by*

$$\mathcal{P}u = \begin{pmatrix} \tilde{D}_1 u \\ \partial_s u_s|_\Sigma \\ u_\Sigma|_\Sigma \end{pmatrix}.$$

Then $\text{Ker } \mathcal{P} = \text{Vect}\{\varphi_i|_{\Omega^+} : 1 \leq i \leq 4\}$, and

$$\begin{pmatrix} 0 \\ v \end{pmatrix} \in \text{Ran } \mathcal{P} \text{ iff } \mathbb{1}_{\{3\}}(\vec{D}_{0,L})(v_s + \vec{\delta}v_\Sigma) = 0.$$

Proof. \mathcal{P} corresponds to an elliptic boundary value problem, (see e.g. [5, Chap. 5]) so \mathcal{P} is Fredholm.

If we replace \tilde{D}_1 by $\tilde{D}_1 + \lambda$ the corresponding operator $\mathcal{P}_\lambda : H^2(\Omega^+; \tilde{V}_1) \rightarrow L^2(\Omega^+; \tilde{V}_1) \oplus (H^{\frac{1}{2}}(\Sigma; \mathbb{C}) \oplus H^{3/2}(\Sigma; \tilde{V}_1))$ is invertible for $\lambda \gg 1$. In fact this operator is associated to the quadratic form

$$Q(u, u) = \int_{\Omega^+} ((\nabla u | \nabla u) + (\lambda - 4)(u|u)) d\text{vol}_{\tilde{g}}$$

with domain $\{u \in H^1(\Omega^+; \tilde{V}_1) : u_\Sigma|_\Sigma = 0\}$, which is positive definite for $\lambda \gg 1$. Therefore \mathcal{P} is of index 0.

It is immediate to find $\text{Ker } \mathcal{P}$ by the methods of images. Namely if $\kappa : (s, \omega) \mapsto (-s, \omega)$ is the reflection in Σ , then if $\tilde{D}_1 w = 0$ in Ω^+ with $\partial_s w_s|_{\Sigma} = w_{\Sigma}|_{\Sigma} = 0$ then extending w by $\kappa^* w$ in Ω^- gives a solution \tilde{w} of $\tilde{D}_1 \tilde{w} = 0$ in \mathbb{S}^4 i.e. $\tilde{w} \in \text{Vect}\{\varphi_i, \varphi_{jk}\}$. The boundary condition on Σ implies that $\tilde{w} \in \text{Vect}\{\varphi_i\}$ as claimed.

Finally $\text{Ran } \mathcal{P} = (\text{Ker } \mathcal{P}^*)^\perp$ and using Green's formula we see that $\begin{pmatrix} v \\ f \end{pmatrix} \in \text{Ker } \mathcal{P}^*$ iff $\tilde{D}_1 v = 0$ in Ω^+ and $f_s = -v_s|_{\Sigma}$, $f_{\Sigma} = \partial_s v_{\Sigma}|_{\Sigma}$. For $v = \varphi_i$ we have $v_s|_{\Sigma} = \psi_i$ and $-\partial_s v_{\Sigma}|_{\Sigma} = \vec{d}\psi_i$, which proves the second statement of the lemma. \square

A.2. Various proofs.

A.2.1. *Covariant derivatives on \mathbb{S}^4 .* Recall that we write $w \in C^\infty(\mathbb{S}^4; \tilde{V}_1)$ as $w_s ds + w_{\Sigma}$ and identify w with (w_s, w_{Σ}) . Similarly we write $u \in C^\infty(\mathbb{S}^4; \tilde{V}_2)$ as $u_{ss} ds^2 + ds \otimes u_{s\Sigma} + u_{\Sigma\Sigma}$ and identify u with $(u_{ss}, u_{s\Sigma}, u_{\Sigma\Sigma})$.

We denote by ∇ , resp. $\vec{\nabla}$, the covariant derivative on $(\mathbb{S}^4, \tilde{g})$, resp. (\mathbb{S}^3, h) , and set $\nabla_0 = \nabla_{\partial_s}$. Then a straightforward computation shows that if $X \in T\Sigma$:

$$(A.1) \quad \begin{aligned} (\nabla_0 w)_s &= \partial_s w_s, \\ (\nabla_0 w)_{\Sigma} &= \partial_s w_{\Sigma} - \frac{1}{2} a^{-1} \dot{a} w_{\Sigma}, \\ (\nabla_X w)_s &= \vec{\nabla}_X w_s - \frac{1}{2} a^{-1} \dot{a} w_{\Sigma} X, \\ (\nabla_X w)_{\Sigma} &= \vec{\nabla}_X w_{\Sigma} + \frac{1}{2} \dot{a} h X w_s, \end{aligned}$$

(where $a(s) = \cos^2 s$), and

$$(A.2) \quad \begin{aligned} (\nabla_0 u)_{ss} &= \partial_s u_{ss}, \\ (\nabla_0 u)_{s\Sigma} &= \partial_s u_{s\Sigma} - \frac{1}{2} a^{-1} \dot{a} u_{s\Sigma}, \\ (\nabla_0 u)_{\Sigma\Sigma} &= \partial_s u_{\Sigma\Sigma} - a^{-1} \dot{a} u_{\Sigma\Sigma}, \\ (\nabla_X u)_{ss} &= \vec{\nabla}_X u_{ss} - a^{-1} \dot{a} u_{s\Sigma} X, \\ (\nabla_X u)_{s\Sigma} &= \vec{\nabla}_X u_{s\Sigma} - \frac{1}{2} a^{-1} \dot{a} u_{\Sigma\Sigma} X + \frac{1}{2} \dot{a} h X u_{ss}, \\ (\nabla_X u)_{\Sigma\Sigma} &= \vec{\nabla}_X u_{\Sigma\Sigma} + \frac{1}{2} \dot{a} h X \otimes u_{s\Sigma} + \frac{1}{2} \dot{a} u_{s\Sigma} \otimes h X. \end{aligned}$$

A.2.2. *Proof of Lemma 5.3.* We deduce from (A.2) that

$$(A.3) \quad \begin{aligned} (\delta u)_s &= -2\partial_s u_{ss} + 2a^{-1} \vec{\delta} u_{s\Sigma} + \frac{1}{2} \dot{a} a^{-2} (h|u_{\Sigma\Sigma}) - 3\dot{a} a^{-1} u_{ss}, \\ (\delta u)_{\Sigma} &= -2\partial_s u_{s\Sigma} + \vec{\delta} u_{\Sigma\Sigma} - 3\dot{a} a^{-1} u_{s\Sigma}, \end{aligned}$$

for $a(s) = \cos^2(s)$ hence:

$$\begin{aligned} (\delta u)_s|_{\Sigma} &= (-2\partial_s u_{ss} + 2\vec{\delta} u_{s\Sigma})|_{\Sigma}, \\ (\delta u)_{\Sigma}|_{\Sigma} &= (-2\partial_s u_{s\Sigma} + \vec{\delta} u_{\Sigma\Sigma})|_{\Sigma}, \end{aligned}$$

and

$$\begin{aligned} -\partial_s (\delta u)_s|_{\Sigma} &= (2\partial_s^2 u_{ss} - 2\vec{\delta} \partial_s u_{s\Sigma} - 6u_{ss} + (h|u_{\Sigma\Sigma}))|_{\Sigma}, \\ -\partial_s (\delta u)_{\Sigma}|_{\Sigma} &= (2\partial_s^2 u_{s\Sigma} - \vec{\delta} \partial_s u_{\Sigma\Sigma} - 6u_{s\Sigma})|_{\Sigma}. \end{aligned}$$

If $\tilde{D}_2 u|_\Sigma = 0$ then

$$\begin{aligned} -\partial_s^2 u_{ss}|_\Sigma &= -\vec{D}_{0,L} u_{ss}|_\Sigma + (h|u_{\Sigma\Sigma})|_\Sigma, \\ -\partial_s^2 u_{s\Sigma}|_\Sigma &= -(\vec{D}_{1,L} - 1)u_{s\Sigma}|_\Sigma, \\ -\partial_s^2 u_{\Sigma\Sigma}|_\Sigma &= -(\vec{D}_{2,L} - 6)u_{\Sigma\Sigma} - 2hu_{ss}|_\Sigma, \end{aligned}$$

which proves the lemma. \square

A.2.3. *Proof of Lemma 5.4.* Again, we deduce from (A.1) that

$$\begin{aligned} (dw)_{ss} &= \partial_s w_s, \\ (A.4) \quad (dw)_{s\Sigma} &= \frac{1}{2}(\partial_s w_\Sigma - \dot{a}a^{-1}w_\Sigma + \vec{d}w_s), \\ (dw)_{\Sigma\Sigma} &= \vec{d}w_\Sigma + \frac{1}{2}\dot{a}hw_s, \end{aligned}$$

hence

$$\begin{aligned} (dw)_{ss}|_\Sigma &= \partial_s w_s|_\Sigma, \\ (dw)_{s\Sigma}|_\Sigma &= \frac{1}{2}(\partial_s w_\Sigma + \vec{d}w_s)|_\Sigma, \\ (dw)_{\Sigma\Sigma}|_\Sigma &= \vec{d}w_\Sigma|_\Sigma, \end{aligned}$$

and

$$\begin{aligned} -(\partial_s dw)_{ss}|_\Sigma &= -\partial_s^2 w_s|_\Sigma, \\ -(\partial_s dw)_{s\Sigma}|_\Sigma &= \frac{1}{2}(-\partial_s^2 w_\Sigma - 2w_\Sigma - \vec{d}\partial_s w_s)|_\Sigma, \\ -(\partial_s dw)_{\Sigma\Sigma}|_\Sigma &= -\vec{d}\partial_s w_\Sigma + hw_s. \end{aligned}$$

If $\tilde{D}_1 w|_\Sigma = v$ then

$$\begin{aligned} -\partial_s^2 w_s|_\Sigma &= -(\vec{D}_{0,L} - 3)w_s|_\Sigma + v_s|_\Sigma, \\ -\partial_s^2 w_\Sigma|_\Sigma &= -(\vec{D}_{1,L} - 6)w_\Sigma|_\Sigma + v_\Sigma|_\Sigma. \end{aligned}$$

It follows that if $g = \tilde{\varrho}_2 dw$ and $f = \tilde{\varrho}_1 w$ we have

$$\begin{aligned} (A.5) \quad g_{0ss} &= -f_{1s}, \\ g_{0s\Sigma} &= \frac{1}{2}(-f_{1\Sigma} + \vec{d}f_{0s}), \\ g_{0\Sigma\Sigma} &= \vec{d}f_{0\Sigma}, \\ g_{1ss} &= -(\vec{D}_{0,L} - 3)f_{0s} + v_s, \\ g_{1s\Sigma} &= \frac{1}{2}(-(\vec{D}_{1,L} - 4)f_{0\Sigma} + \vec{d}f_{1s}) + \frac{1}{2}v_\Sigma, \\ g_{1\Sigma\Sigma} &= \vec{d}f_{1\Sigma} + hf_{0s}. \end{aligned}$$

To complete the computation of $\tilde{K}_{21\Sigma}$ we need to compute $\tilde{\varrho}_2 \tilde{I}dw$, where $\tilde{I}u = u - \frac{1}{4}(\tilde{g}|u)_{\tilde{V}_2} \tilde{g}$, i.e.

$$\begin{aligned} (\tilde{I}u)_{ss} &= \frac{1}{2}u_{ss} - \frac{1}{4}(h|u_{\Sigma\Sigma})_{\tilde{V}_{2\Sigma}}, \\ (\tilde{I}u)_{s\Sigma} &= u_{s\Sigma}, \\ (\tilde{I}u)_{\Sigma\Sigma} &= u_{\Sigma\Sigma} - \frac{1}{4}(2u_{ss} + (h|u_{\Sigma\Sigma})_{\tilde{V}_{2\Sigma}})h. \end{aligned}$$

If $g = \tilde{\varrho}_2 \tilde{I} dw$ and $f = \tilde{\varrho}_1 w$, we obtain:

$$\begin{aligned}
g_{0ss} &= \frac{1}{2}(-f_{1s} + \vec{\delta} f_{0\Sigma}), \\
g_{0s\Sigma} &= \frac{1}{2}(-f_{1\Sigma} + \vec{d} f_{0s}), \\
g_{0\Sigma\Sigma} &= \vec{d} f_{0\Sigma} + \frac{1}{2}(f_{1s} + \vec{\delta} f_{0\Sigma})h, \\
g_{1ss} &= \frac{1}{2}(-\vec{D}_{0,L} f_{0s} + \vec{\delta} f_{1\Sigma}) + \frac{1}{2}v_s \upharpoonright_\Sigma, \\
g_{1s\Sigma} &= \frac{1}{2}(-(\vec{D}_{1,L} - 4)f_{0\Sigma} + \vec{d} f_{1s}) + \frac{1}{2}v_\Sigma \upharpoonright_\Sigma, \\
g_{1\Sigma\Sigma} &= \vec{d} f_{1\Sigma} + \frac{1}{2}\vec{\delta} f_{1\Sigma} + \frac{1}{2}(\vec{D}_{0,L} - 4)f_{0s}h - \frac{1}{2}v_s \upharpoonright_\Sigma h,
\end{aligned}
\tag{A.6}$$

which proves the lemma. \square

A.2.4. *Proof of Lemma 6.1.* If P is selfadjoint with non-trivial kernel we denote by $P^{-1} : \text{Ran } P \rightarrow (\text{Ker } P)^\perp$ the generalized inverse of P . From Lemma 5.3 we obtain that $g = \begin{pmatrix} g_0 \\ g_1 \end{pmatrix} \in \text{Ker } \tilde{K}_{21\Sigma}^\dagger \cap \text{Ker } \tilde{K}_{20\Sigma}^\dagger$ iff

- 1) $g_{0ss} + \frac{1}{2}(h|g_{0\Sigma\Sigma}) = 0$,
- 2) $g_{1ss} + \frac{1}{2}(h|g_{1\Sigma\Sigma}) = 0$,
- 3) $g_{1s\Sigma} + \vec{\delta} g_{0s\Sigma} = 0$,
- 4) $2g_{1s\Sigma} + \vec{\delta} g_{0\Sigma\Sigma} = 0$,
- 5) $2(\vec{D}_{0,L} - 3)g_{0ss} + 2\vec{\delta} g_{1s\Sigma} - (h|g_{0\Sigma\Sigma}) = 0$,
- 6) $2(\vec{D}_{1,L} - 4)g_{0s\Sigma} + \vec{\delta} g_{1\Sigma\Sigma} = 0$.

An easy computation shows that this is equivalent to

- 1') $g_{0ss} = -\frac{1}{2}(h|g_{0\Sigma\Sigma})$,
- 2') $g_{1ss} = -\frac{1}{2}(h|g_{1\Sigma\Sigma})$,
- 3') $g_{0s\Sigma} = -\frac{1}{2}(\vec{D}_{1,L} - 4)^{-1}\vec{\delta} g_{1\Sigma\Sigma} + \beta_1$,
- 4') $g_{1s\Sigma} = -\frac{1}{2}\vec{\delta} g_{0\Sigma\Sigma}$,
- 5') $(\vec{D}_{0,L} - 2)(h|g_{0\Sigma\Sigma}) = -\vec{\delta}\vec{\delta} g_{0\Sigma\Sigma}$,
- 6') $(\vec{D}_{0,L} - 4)(h|g_{1\Sigma\Sigma}) = -\vec{\delta}\vec{\delta} g_{1\Sigma\Sigma}$,

for $\beta_1 \in \text{Vect}\{\psi_{jk}\}$. For $u \in C^\infty(\Sigma; V_{2\Sigma})$ we set $u = \alpha h + \bar{u}$ for $(h|\bar{u}) = 0$ and $(h|u) = 6\alpha$. We have then

$$\vec{\delta}\alpha h = -2\vec{d}\alpha, \quad \vec{\delta}\vec{\delta}\alpha h = -2\vec{D}_{0,L}\alpha.$$

Therefore setting $g_{i\Sigma\Sigma} = \alpha_i h + \bar{g}_{i\Sigma\Sigma}$ we obtain from 5') and 6)' that

$$\begin{aligned}
(\vec{D}_{0,L} - 3)\alpha_0 &= -\frac{1}{4}\vec{\delta}\vec{\delta}\bar{g}_{0\Sigma\Sigma}, \\
(\vec{D}_{1,L} - 6)\alpha_1 &= -\frac{1}{4}\vec{\delta}\vec{\delta}\bar{g}_{1\Sigma\Sigma},
\end{aligned}$$

i.e.

$$\begin{aligned}
\alpha_0 &= -\frac{1}{4}(\vec{D}_{0,L} - 3)^{-1}\vec{\delta}\vec{\delta}\bar{g}_{0\Sigma\Sigma} + \beta_0, \\
\alpha_1 &= -\frac{1}{4}(\vec{D}_{1,L} - 6)^{-1}\vec{\delta}\vec{\delta}\bar{g}_{1\Sigma\Sigma},
\end{aligned}$$

for $\beta_0 \in \text{Vect}\{\psi_i\}$. From 3'), 4') we obtain then

$$\begin{aligned}\vec{\delta}g_{0\Sigma\Sigma} &= (1 + \frac{1}{2}\vec{d}(\vec{D}_{0,L} - 3))^{-1}\vec{\delta}\vec{\delta}\vec{g}_{0\Sigma\Sigma} - 2\vec{d}\beta_0, \\ \vec{\delta}g_{1\Sigma\Sigma} &= (1 + \frac{1}{2}\vec{d}(\vec{D}_{0,L} - 6))^{-1}\vec{\delta}\vec{\delta}\vec{g}_{1\Sigma\Sigma},\end{aligned}$$

and finally

$$\begin{aligned}1) \quad g_{0ss} &= \frac{3}{4}(\vec{D}_{0,L} - 3)^{-1}\vec{\delta}\vec{\delta}\vec{g}_{0\Sigma\Sigma} - 3\beta_0, \\ 2) \quad g_{1ss} &= \frac{3}{4}(\vec{D}_{0,L} - 6)^{-1}\vec{\delta}\vec{\delta}\vec{g}_{1\Sigma\Sigma}, \\ 3) \quad g_{0s\Sigma} &= -\frac{1}{2}(\vec{D}_{1,L} - 4)^{-1}(1 + \frac{1}{2}\vec{d}(\vec{D}_{0,L} - 6))^{-1}\vec{\delta}\vec{\delta}\vec{g}_{1\Sigma\Sigma} + \beta_1, \\ 4) \quad g_{1s\Sigma} &= -\frac{1}{2}(1 + \frac{1}{2}\vec{d}(\vec{D}_{0,L} - 3))^{-1}\vec{\delta}\vec{\delta}\vec{g}_{0\Sigma\Sigma} + \vec{d}\beta_0, \\ 5) \quad g_{0\Sigma\Sigma} &= -\frac{1}{4}(\vec{D}_{0,L} - 3)^{-1}\vec{\delta}\vec{\delta}\vec{g}_{0\Sigma\Sigma}h + \vec{g}_{0\Sigma\Sigma} + \beta_0h, \\ 6) \quad g_{1\Sigma\Sigma} &= -\frac{1}{4}(\vec{D}_{0,L} - 6)^{-1}\vec{\delta}\vec{\delta}\vec{g}_{1\Sigma\Sigma}h + \vec{g}_{1\Sigma\Sigma}.\end{aligned}$$

This completes the proof of the lemma. \square

A.2.5. *Proof of Prop. 6.2.* Since \hat{D}_i commutes with \mathcal{F}_i and $\mathcal{F}_{i\Sigma}$ is unitary for q_2 , it suffices to prove the proposition with $\text{Ker } K_{21\Sigma}^\dagger \cap \text{Ker } K_{20\Sigma}^\dagger$ replaced by $\text{Ker } \tilde{K}_{21\Sigma}^\dagger \cap \text{Ker } \tilde{K}_{20\Sigma}^\dagger$.

(1) follows from Lemma 5.6.

Let us now prove (2). Note first that since \mathbf{q}_2 commutes with $\mathbb{1}_{\{\lambda\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2}$ the spaces $(\mathbb{1}_{\mathbb{R}\setminus\{3,4\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2})(\text{Ker } \tilde{K}_{21\Sigma}^\dagger \cap \text{Ker } \tilde{K}_{20\Sigma}^\dagger)$ and $(\mathbb{1}_{\{3,4\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2})(\text{Ker } \tilde{K}_{21\Sigma}^\dagger \cap \text{Ker } \tilde{K}_{20\Sigma}^\dagger)$ are orthogonal for \mathbf{q}_2 .

From (6.1) and the fact that $\vec{d}\vec{d}\psi_i = -\psi_i h$, $\vec{d}\psi_{jk} = \vec{\delta}\psi_{jk} = 0$ we obtain after an easy computation that

$$(A.7) \quad (\mathbb{1}_{\{3,4\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2})g = \begin{pmatrix} -3\beta_0 \\ 0 \\ \beta_1 \\ \vec{d}\beta_0 \\ \beta_0 h \\ 0 \end{pmatrix}.$$

This implies that \mathbf{q}_2 vanishes on $(\mathbb{1}_{\{3,4\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2})(\text{Ker } \tilde{K}_{21\Sigma}^\dagger \cap \text{Ker } \tilde{K}_{20\Sigma}^\dagger)$. Therefore it remains to prove that \mathbf{q}_2 is non degenerate on $(\mathbb{1}_{\mathbb{R}\setminus\{3,4\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2})(\text{Ker } \tilde{K}_{21\Sigma}^\dagger \cap \text{Ker } \tilde{K}_{20\Sigma}^\dagger)$.

To this end we will use an additional decomposition of $(0, 2)$ -tensors on Σ . By e.g. [4, Subsect. 3.7], we can write $u_2 \in C^\infty(\Sigma; V_{2\Sigma})$ as an orthogonal sum

$$u_2 = v_2 + (v_0 h + \vec{d}v_1), \quad v_i \in C^\infty(\Sigma; V_{i\Sigma}), \quad \vec{\delta}v_2 = (h|v_2) = 0.$$

If moreover $(h|u_2) = 0$, we obtain that $v_0 = \frac{1}{3}\vec{\delta}v_1$.

Therefore we can write $\vec{g}_{i\Sigma\Sigma}$ as the orthogonal sum

$$(A.8) \quad \vec{g}_{i\Sigma\Sigma} = u_i + (\vec{d}w_i + \frac{1}{3}\vec{\delta}w_i h), \quad \text{with } \vec{\delta}u_i = (h|u_i) = 0.$$

Note that $\vec{d}w + \frac{1}{3}\vec{\delta}wh = 0$ iff w is a conformal Killing 1-form on \mathbb{S}^3 , i.e. $w \in \text{Vect}\{\vec{d}\psi_i, \psi_{jk}\} = \text{Ran } \mathbb{1}_{\{3,4\}}(\vec{D}_{1,L})$. So there are unique couples (u_i, w_i) with $u_i \in \text{Ker } \vec{\delta} \cap \text{Ker}(h)$, $w_i \in \text{Ker } \mathbb{1}_{\{3,4\}}(\vec{D}_{1,L})$ such that (A.8) is satisfied.

Let now $g \in (\mathbb{1}_{\mathbb{R}\setminus\{3,4\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2})(\text{Ker } \tilde{K}_{21\Sigma}^\dagger \cap \text{Ker } \tilde{K}_{20\Sigma}^\dagger)$, i.e. g is given by (6.1) with $\beta_0 = \beta_1 = 0$. We have

$$\begin{aligned}\vec{\delta}(\vec{d}w_i + \frac{1}{3}\vec{\delta}w_i h) &= (\vec{D}_{1,L} - 4 + \frac{1}{3}\vec{d}\vec{\delta})w_i, \\ \vec{\delta}\vec{\delta}(\vec{d}w_i + \frac{1}{3}\vec{\delta}w_i h) &= \frac{4}{3}(\vec{D}_{0,L} - 3)\vec{\delta}w_i,\end{aligned}$$

if $\mathbb{1}_{\{3,4\}}(\vec{D}_{1,L})w_i = 0$, which using that $\vec{D}_{1,L} = \vec{\delta}\vec{d} - \vec{d}\vec{\delta} + 4$ gives:

- 1) $g_{0ss} = \vec{\delta}w_0$,
- 2) $g_{1ss} = (1 + 3(\vec{D}_{0,L} - 6)^{-1})\vec{\delta}w_1$,
- 3) $g_{0s\Sigma} = -\frac{1}{2}(1 + \vec{d}(\vec{D}_{0,L} - 6)^{-1}\vec{\delta})w_1$,
- 4) $g_{1s\Sigma} = -\frac{1}{2}\vec{\delta}\vec{d}w_0$,
- 5) $g_{0\Sigma\Sigma} = u_0 + \vec{d}w_0$,
- 6) $g_{1\Sigma\Sigma} = u_1 + \vec{d}w_1 - (\vec{D}_{0,L} - 6)^{-1}\vec{\delta}w_1 h$.

For $g, g' \in (\mathbb{1}_{\mathbb{R}\setminus\{3,4\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2})(\text{Ker } \tilde{K}_{21\Sigma}^\dagger \cap \text{Ker } \tilde{K}_{20\Sigma}^\dagger)$, we have

$$\begin{aligned}\vec{g}' \cdot \mathbf{q}_{2\Sigma} g &= ((1 + 3(\vec{D}_{0,L} - 6)^{-1})\vec{\delta}w'_1 | \vec{\delta}w_0) + (\vec{\delta}w'_0 | (1 + 3(\vec{D}_{0,L} - 6)^{-1})\vec{\delta}w_1) \\ &\quad - \frac{1}{2}(\vec{\delta}\vec{d}w'_0 | (1 + \vec{d}(\vec{D}_{0,L} - 6)^{-1}\vec{\delta})w_1) \\ &\quad - \frac{1}{2}((1 + \vec{d}(\vec{D}_{0,L} - 6)^{-1}\vec{\delta})w'_1 | \vec{\delta}\vec{d}w_0) \\ &\quad + (\vec{d}w'_1 - (\vec{D}_{0,L} - 6)^{-1}\vec{\delta}w'_1 h | \vec{d}w_0) \\ &\quad + (\vec{d}w'_0 | \vec{d}w_1 - (\vec{D}_{0,L} - 6)^{-1}\vec{\delta}w_1 h) \\ &\quad + (u'_1 | u_0) + (u'_0 | u_1).\end{aligned}$$

Therefore $\vec{g}' \cdot \mathbf{q}_{2\Sigma} g = 0$ for all $g' g \in (\mathbb{1}_{\mathbb{R}\setminus\{3,4\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2})(\text{Ker } \tilde{K}_{21\Sigma}^\dagger \cap \text{Ker } \tilde{K}_{20\Sigma}^\dagger)$ iff

$$A^*w_0 = Aw_1 = 0,$$

$$u_0 = u_1 = 0.$$

for

$$\begin{aligned}A &= \vec{d}(1 + 3(\vec{D}_{0,L} - 6)^{-1})\vec{\delta} - \frac{1}{2}\vec{\delta}\vec{d}(1 + \vec{d}(\vec{D}_{0,L} - 6)^{-1}\vec{\delta}) \\ &\quad + \vec{\delta}\vec{d} + \vec{d}(\vec{D}_{0,L} - 6)^{-1}\vec{\delta} \\ &= \vec{d}(1 + 4(\vec{D}_{0,L} - 6)^{-1})\vec{\delta} + \frac{1}{2}\vec{\delta}\vec{d}(1 - \vec{d}(\vec{D}_{0,L} - 6)^{-1}\vec{\delta}).\end{aligned}$$

We note that A commutes with $\vec{D}_{1,L}$. Let w an eigenvector of $\vec{D}_{1,L}$ with eigenvalue λ and $\lambda \neq 3, 4$. From [4, Subsect. 3.6] we know that either $w = \vec{d}\alpha$ for $\vec{D}_{0,L}\alpha = \lambda\alpha$, $\lambda = k(k+2)$, $k \in \mathbb{N}, k \geq 2$ or $\vec{\delta}w = 0$, $\lambda = k(k+2) + 1$, $k \in \mathbb{N}, k \geq 2$. Using that $\vec{D}_{1,L} = \vec{\delta}\vec{d} - \vec{d}\vec{\delta} + 4$ we have $\vec{\delta}\vec{d}w = (2\lambda - 4)w$ in the first case and $\vec{\delta}\vec{d}w = (\lambda - 4)w$ in the second case.

In the first case we obtain that $Aw = A^*w = (\lambda - 2)w$ and in the second case that $Aw = A^*w = \frac{1}{2}(\lambda - 4)w$. Since $\lambda \neq 2, 4$ this shows that A

and A^* are injective. Therefore \mathbf{q}_2 is non-degenerate on $(\mathbb{1}_{\mathbb{R} \setminus \{3,4\}}(\hat{D}_2) \otimes \mathbb{1}_{\mathbb{C}^2})(\text{Ker } \tilde{K}_{21\Sigma}^\dagger \cap \text{Ker } \tilde{K}_{20\Sigma}^\dagger)$ as claimed. \square

APPENDIX B. MAXWELL FIELDS

B.1. Maxwell fields on de Sitter space. We briefly treat the analogous case of Maxwell fields on dS^4 . In this case the bundles V_2, V_1 are replaced by V_1, V_0 respectively, D_2 is replaced by $D_{1,L} = \delta_a \circ d_a + d_a \circ \delta_a$, D_0 by $D_{0,L} = -\square_0$, K_{21} by $K_{10} = d_a$. The physical charge $\mathbf{q}_{2,I}$ is replaced by \mathbf{q}_1 . We set

$$\mathcal{E} := \text{Ker } K_{10\Sigma}^\dagger, \quad \mathcal{F} := \text{Ran } K_{10\Sigma},$$

the physical phase space then being \mathcal{E}/\mathcal{F} .

As before we denote with tildes the Wick rotated objects.

We obtain that

$$\tilde{D}_0 u = -\partial_s^2 u + a^{-1} \tilde{D}_{0,L} u + \frac{3}{2} \dot{a} a^{-2} \partial_s u,$$

while the expression for $\tilde{D}_1 w$ is obtained from the one at the beginning of Subsect. 5.3 by adding the constant term $2\Lambda = 6$.

From [4, Thm. 3.2, Prop. 3.9] we know that $\tilde{D}_1 \geq 4$, hence \tilde{D}_1 is invertible, while $\text{Ker } \tilde{D}_0 = \text{Ker } d = \mathbb{C}$, the space of constant functions on \mathbb{S}^4 . Therefore the Calderón projectors \tilde{c}_1^\pm for \tilde{D}_1 are well defined, while those for \tilde{D}_0 exist only on $(\tilde{\varrho}_0 \text{Ker } \tilde{D}_0)^{\mathfrak{q}_0}$.

As before we obtain that \tilde{D}_1 commutes with the spatial operator \hat{D}_1 , hence $[\tilde{c}_1^\pm, \mathbb{1}_{\{\lambda\}}(\hat{D}_1) \otimes \mathbb{1}_{\mathbb{C}^2}] = 0$, and also that $(\mathbb{1}_{\{\lambda\}}(\hat{D}_1) \otimes \mathbb{1}_{\mathbb{C}^2}) K_{10\Sigma} = K_{10\Sigma} (\mathbb{1}_{\{\lambda\}}(\hat{D}_0) \otimes \mathbb{1}_{\mathbb{C}^2})$. We have

$$(B.9) \quad \text{Ker } \hat{D}_1 = \mathbb{C} \oplus 0,$$

where we use the decomposition $w = w_s \oplus w_\Sigma$ of 1-forms on \mathbb{S}^4 .

Defining $\tilde{K}_{10\Sigma}$ and R_Σ as in Subsect. 5.4 we obtain that if $g = \tilde{K}_{10\Sigma} f$ for $f = \begin{pmatrix} f_0 \\ f_1 \end{pmatrix} \in C^\infty(\Sigma; \tilde{V}_0(\Sigma) \otimes \mathbb{C}^2)$ then

$$(B.10) \quad \begin{cases} g_{0s} = -f_1, \\ g_{0\Sigma} = \vec{d}f_0, \\ g_{1s} = -\vec{D}_{0,L} f_0, \\ g_{1\Sigma} = \vec{d}f_1, \end{cases}$$

and if $v \in C^\infty(\Sigma; \tilde{V}_0(\Sigma))$ then $(R_\Sigma v)_{1s} = v$, all other entries being equal to 0.

Similarly we find that if $f = \tilde{K}_{10\Sigma}^\dagger g$, then

$$(B.11) \quad \begin{cases} g_0 = f_{1s} + \vec{\delta} f_{0\Sigma}, \\ g_1 = \vec{D}_{0,L} f_{0s} + \vec{\delta} f_{1\Sigma}. \end{cases}$$

The Lorentzian versions of $K_{10\Sigma}, K_{10\Sigma}^\dagger$ are obtained by conjugation with the $\mathcal{F}_{i\Sigma}$, as explained in Subsect. 3.2.

B.1.1. *Properties of \mathcal{E} .* We split \mathcal{E} as $\mathcal{E}_{\text{reg}} \oplus^\perp \mathcal{E}_{\{0\}}$, with

$$\mathcal{E}_{\{0\}} = (\mathbb{1}_{\{0\}}(\hat{D}_1) \otimes \mathbb{1}_{\mathbb{C}^2})\mathcal{E}.$$

Note that $\mathbb{1}_{\{0\}}(\hat{D}_1) \otimes \mathbb{1}_{\mathbb{C}^2}$ preserves $\mathcal{E} = \text{Ker } K_{10\Sigma}^\dagger$. This decomposition is preserved by \mathbf{q}_1 and \tilde{c}_1^\pm .

Lemma B.1. *The kernel of \mathbf{q}_1 restricted to \mathcal{E} equals \mathcal{F} .*

This implies of course that \mathbf{q}_1 is non-degenerate on \mathcal{E}/\mathcal{F} . We remark that as noticed in [33] (cf. [34] for the analogous statement in the BRST approach), this also follows by a Poincaré duality argument and the fact that the Cauchy surface is compact.

Proof. Clearly $\text{Ran } K_{10\Sigma} \subset \text{Ker } \mathbf{q}_1|_{\mathcal{E}}$. Using (B.9) we see that $\mathcal{E}_{\{0\}} \subset \text{Ran } K_{10\Sigma}$. Moreover \mathcal{E}_{reg} and $\mathcal{E}_{\{0\}}$ are orthogonal for \mathbf{q}_1 . Therefore to complete the proof it suffices to show that $\text{Ker } \mathbf{q}_1|_{\mathcal{E}_{\text{reg}}} = \text{Ran } K_{10\Sigma} \cap \mathcal{E}_{\text{reg}}$. If $f, f' \in \mathcal{E}_{\text{reg}}$ we obtain that

$$\begin{aligned} \bar{f} \cdot \mathbf{q}_1 f' &= -(\vec{\delta} f_{0\Sigma} | \vec{D}_{0,L}^{-1} \vec{\delta} f'_{1\Sigma})_{V_0(\Sigma)} - (\vec{D}_{0,L}^{-1} \vec{\delta} f_{1\Sigma} | \vec{\delta} f'_{0\Sigma})_{V_0(\Sigma)} \\ &\quad + (f_{1\Sigma} | f'_{0\Sigma})_{V_1(\Sigma)} + (f_{0\Sigma} | f'_{1\Sigma})_{V_1(\Sigma)}, \end{aligned}$$

where $\vec{D}_{0,L}^{-1} = (1 - \pi_0)(\vec{D}_{0,L} + \pi_0)^{-1}$ is the pseudo-inverse of $\vec{D}_{0,L}$, $\pi_0 = \mathbb{1}_{\{0\}}(\vec{D}_{0,L})$ and in the sequel $\vec{d}, \vec{\delta}$ are the anti-symmetric differential and codifferential on \mathbb{S}^3 . Using that $\vec{d}\vec{D}_{0,L} = \vec{D}_{1,L}\vec{d}$, $\vec{d}\pi_0 = 0$ and $\vec{D}_{1,L} = \vec{\delta}\vec{d} + \vec{d}\vec{\delta}$, we obtain that

$$\begin{aligned} \bar{f} \cdot \mathbf{q}_1 f' &= -(f_{0\Sigma} | \vec{D}_{1,L}^{-1} \vec{d}\vec{\delta} f'_{1\Sigma})_{V_1(\Sigma)} - (\vec{D}_{1,L}^{-1} \vec{d}\vec{\delta} f_{1\Sigma} | \vec{\delta} f'_{0\Sigma})_{V_1(\Sigma)} \\ &\quad + (f_{1\Sigma} | f'_{0\Sigma})_{V_1(\Sigma)} + (f_{0\Sigma} | f'_{1\Sigma})_{V_1(\Sigma)} \\ &= (f_{1\Sigma} | \vec{D}_{1,L}^{-1} \vec{\delta}\vec{d} f'_{0\Sigma})_{V_1(\Sigma)} + (f_{0\Sigma} | \vec{D}_{1,L}^{-1} \vec{\delta}\vec{d} f'_{1\Sigma})_{V_1(\Sigma)}. \end{aligned}$$

Therefore $f \in \text{Ker } \mathbf{q}_1|_{\mathcal{E}_{\text{reg}}}$ iff $\vec{d}f_{i\Sigma} = 0$, i.e. $f_{i\Sigma} = \vec{d}u_i$ for $u_i \perp \text{Ker } \vec{D}_{0,L}$. Using (B.11) this gives also $f_{1s} = -\vec{D}_{0,L}u_0$ and $\vec{D}_{0,L}f_{0s} = -\vec{D}_{0,L}u_1$. Since $f \in \mathcal{E}_{\text{reg}}$ we have $f_{0,s} \perp \text{Ker } \vec{D}_{0,L}$, which gives $f_{0s} = -u_1$. By (B.10) this means that $f = K_{10\Sigma} \begin{pmatrix} u_0 \\ u_1 \end{pmatrix}$, i.e. $f \in \text{Ran } K_{10\Sigma}$. \square

B.1.2. *The Euclidean vacuum.* As before we define covariances $\lambda_{1\Sigma}^\pm := \pm \mathbf{q}_1 \circ c_1^\pm$, where $c_1^\pm = \mathcal{F}_{1\Sigma}^{-1} \circ \tilde{c}_1^\pm \circ \mathcal{F}_{1\Sigma}$. The associated state ω_{eucl} is again the *Euclidean vacuum*. As in the case of linear gravity, it is not positive on the whole of \mathcal{E} .

We set $\mathcal{K}_\Sigma := \mathcal{F}_{0\Sigma}^{-1} \tilde{\varrho}_0 \text{Ker } \vec{D}_0 = \mathbb{C} \oplus \{0\}$.

B.1.3. *Gauge invariance.*

Lemma B.2. *One has*

$$c_1^\pm K_{10\Sigma} f \in \text{Ran } K_{10\Sigma} \text{ iff } f \in \mathcal{K}_\Sigma^{\mathbf{q}_0}.$$

Moreover if $f \in \text{Ker } K_{10\Sigma}^\dagger$, then $K_{10\Sigma}^\dagger c_1^\pm f \in \mathcal{K}_\Sigma$.

Proof. We mimick the proof of Prop. 7.2. We fix $f \in C^\infty(\Sigma; \tilde{V}_0(\Sigma) \otimes \mathbb{C}^2)$ and set

$$u_0 = (\tilde{D}_0 + \pi_0)^{-1} \tilde{\varrho}_0^* \tilde{\sigma}_0 f, \quad u_1 = \tilde{K}_{10} u_0,$$

where $\pi_0 = \mathbb{1}_{\{0\}}(\tilde{D}_0)$ is the orthogonal projection on $\text{Ker } \tilde{D}_0 = \text{Ker } d$. We obtain that

$$\tilde{\varrho}_1^\pm u_1 = \mp \tilde{K}_{10\Sigma} \hat{c}_0^\pm f - R_\Sigma(\pi_0 \tilde{\varrho}_0^* \tilde{\sigma}_0 f),$$

where \hat{c}_0^\pm are defined as in 4.2.5. We obtain that $\pi_0 \tilde{\varrho}_0^* \tilde{\sigma}_0 f = -\frac{3}{4} \mathbb{1}_{\{0\}}(\tilde{D}_{0,L}) f_1$.

From (B.10) we get that $R_\Sigma(\pi_0 \tilde{\varrho}_0^* \tilde{\sigma}_0 f) = \tilde{K}_{10\Sigma} f'$ iff

$$\begin{cases} -f'_1 = \tilde{d}f'_0 = \tilde{d}f'_1 = 0, \\ \tilde{D}_{0,L} f'_0 = \frac{3}{4} \mathbb{1}_{\{0\}}(\tilde{D}_{0,L}) f_1, \end{cases}$$

ie $\mathbb{1}_{\{0\}}(\tilde{D}_{0,L}) f_1 = 0$. This is equivalent to $f \in (\tilde{\varrho}_0 \text{Ker } \tilde{D}_0)^{\mathfrak{q}_0}$. The second statement is proved as in Prop. 7.2. \square

B.1.4. Positivity.

Proposition B.3. *The covariances $\lambda_{1\Sigma}^\pm$ are positive on \mathcal{E}_{reg} and negative on $\mathcal{E}_{\{0\}}$.*

Proof. Without loss of generality we can work in the Euclidean framework and set $\mathcal{E} = \text{Ker } \tilde{K}_{10\Sigma}^\dagger$ etc.

i) positivity on $\tilde{\mathcal{E}}_{\text{reg}}$:

Let first $f_1 \in \tilde{\mathcal{E}}_{\text{reg}}$ and $u_1 = -(\tilde{D}_1 \tilde{\varrho}_1^* \tilde{\sigma}_1 f_1)|_{\Omega^+}$, $\tilde{u}_1 = u_1 - \tilde{K}_{10} u_0$ for some $u_0 \in \overline{C^\infty}(\Omega^+; \tilde{V}_0)$ with $\tilde{D}_0 u_0 = 0$ to be determined later. We set $\tilde{f}_1 = \tilde{\varrho}_1^+ \tilde{u}_1$, $f_0 = \tilde{\varrho}^+ u_0$. Then

$$\tilde{c}_1^+ f_1 = \tilde{\varrho}_1^+ u_1 = \tilde{f}_1 + \tilde{K}_{10\Sigma} f_0,$$

since $\tilde{D}_0 u_0 = 0$ in Ω^+ . The same computation as in the proof of Prop. 7.4 gives that if $f_0 \in \text{Ker } \tilde{C}_{0\Sigma}$ then $(f_1 | q_1 \tilde{c}_1^+ f_1)_{\tilde{V}_1(\Sigma) \otimes \mathbb{C}^2} = (\tilde{f}_1 | q_1 \tilde{f}_1)_{\tilde{V}_1(\Sigma) \otimes \mathbb{C}^2}$.

Using that $\tilde{D}_1 \tilde{u}_1 = 0$ in Ω^+ and $\tilde{D}_1 \geq 4$ we find as before that

$$(\tilde{f}_1 | \tilde{q}_1 \tilde{f}_1)_{\tilde{V}_1(\Sigma) \otimes \mathbb{C}^2} \geq 0.$$

Moreover if $\tilde{u}_{1s}|_\Sigma = 0$ then $(f_1 | q_1 \tilde{f}_1)_{\tilde{V}_1(\Sigma) \otimes \mathbb{C}^2} = (\tilde{f}_1 | \tilde{q}_1 \tilde{f}_1)_{\tilde{V}_1(\Sigma) \otimes \mathbb{C}^2}$, hence we need to find u_0 such that

$$(B.12) \quad \begin{cases} \tilde{D}_0 u_0 = 0 \text{ in } \Omega^+, \\ i) \quad (\partial_s u_0 + u_{1s})|_\Sigma = 0, \\ ii) \quad \mathbb{1}_{\{0\}}(\tilde{D}_{0,L}) \partial_s u_0|_\Sigma = 0. \end{cases}$$

Arguing as in the proof of Lemma A.1, we find that (B.12) has a solution iff $\mathbb{1}_{\{0\}}(\tilde{D}_{0,L}) u_1|_\Sigma = 0$. Since $\tilde{\varrho}_1^+ u_1 = \tilde{c}_1^+ f_1$ and $[\tilde{c}_1^\pm, \mathbb{1}_{\{\lambda\}}(\hat{D}_1) \otimes \mathbb{1}_{\mathbb{C}^2}] = 0$, this condition is satisfied if $\mathbb{1}_{\{0\}}(\hat{D}_1) \otimes \mathbb{1}_{\mathbb{C}^2} f_1 = 0$. Therefore $\lambda_{1\Sigma}^\pm \geq 0$ on $\tilde{\mathcal{E}}_{\text{reg}}$.

ii) negativity on $\tilde{\mathcal{E}}_{\{0\}}$:

Let $f \in \tilde{\mathcal{E}}_{\{0\}}$. We have $\overline{f} \cdot \lambda_{1\Sigma}^+ f = \overline{f} \cdot \mathfrak{q}_1 c_1^+ f = \overline{c_1^\pm f} \cdot \mathfrak{q}_1 c_1^+ f$. Let $\tilde{f} = c_1^+ f$. Since c_1^\pm commutes with \hat{D}_1 we have $\tilde{f}_0, \tilde{f}_1 \in \text{Ker } \hat{D}_1$ hence $\tilde{f}_0|_\Sigma = \tilde{f}_1|_\Sigma = 0$. Therefore $\overline{f} \cdot \mathfrak{q}_1 \tilde{f} = -\overline{\tilde{f}} \cdot \tilde{\mathfrak{q}}_1 \tilde{f}$. As in the first part of the proof the last quantity is positive. Therefore $\lambda_{1\Sigma}^+$ is negative on $\tilde{\mathcal{E}}_{\{0\}}$. \square

B.1.5. *The modified Euclidean vacuum.* Let $\pi = \mathbb{1}_{\mathbb{R} \setminus \{0\}}(\hat{D}_1) \otimes \mathbb{C}^2$ be the orthogonal projection $\mathcal{E} \rightarrow \mathcal{E}_{\text{reg}}$. For $f \in \mathcal{E}$ we have $((1-\pi)f)_{0s} = \mathbb{1}_{\{0\}}(\vec{D}_{0,L})f_{0s}$, the other components being equal to 0.

We define the modified Euclidean vacuum similarly as before by the covariances

$$\lambda_{1\Sigma, \text{mod}}^\pm := \overline{\pi f} \cdot \lambda_{1\Sigma}^\pm \pi f.$$

As in Prop. 6.6 we have $\text{Ran}(\mathbb{1} - \pi) \subset \text{Ran} K_{10\Sigma}$, $\mathbb{1} - \pi$ is smoothing of rank one, and $\text{Ran} \pi K_{10\Sigma} \subset K_{10\Sigma} \mathcal{K}_\Sigma^{\mathfrak{q}_0}$.

The same arguments as in the proof of Thm. 8.2 show that the associated state ω_{mod} is a Hadamard, fully gauge invariant state on \mathcal{E} . It is only invariant under the action of $O(4)$, not under the full $O(1,4)$ symmetry group.

Acknowledgments. The authors would like to thank Vincent Moncrief and Simone Murro for useful discussions. Support from the grant ANR-20-CE40-0018 is gratefully acknowledged.

REFERENCES

- [1] B. Allen. Graviton propagator in de Sitter space. *Phys. Rev. D*, 34(12):3670–3675, 1986.
- [2] B. Allen, A. Folacci, and A. C. Ottewill. Renormalized graviton stress-energy tensor in curved vacuum space-times. *Phys. Rev. D*, 38(4):1069–1082, 1988.
- [3] M. Benini, C. Dappiaggi, and S. Murro. Radiative observables for linearized gravity on asymptotically flat spacetimes and their boundary induced states. *J. Math. Phys.*, 55(8):082301, 2014.
- [4] M. Boucetta. Spectre des Laplaciens de Lichnerowicz sur les sphères et les projectifs réels. *Publicacions Matemàtiques*, 43:451–483, 1999.
- [5] J. Chazarain and A. Piriou. *Introduction to the Theory of Linear Partial Differential Equations*, volume 14 of *Studies in Mathematics and Its Applications*. Elsevier, 1982.
- [6] C. Dappiaggi and D. Siemssen. Hadamard states for the vector potential on asymptotically flat spacetimes. *Rev. Math. Phys.*, 25(01):1350002, 2013.
- [7] C. J. Fewster and D. S. Hunt. Quantization of linearized gravity in cosmological vacuum spacetimes. *Rev. Math. Phys.*, 25(02):1330003, 2013.
- [8] C. J. Fewster and M. J. Pfenning. A quantum weak energy inequality for spin-one fields in curved space–time. *J. Math. Phys.*, 44(10):4480, 2003.
- [9] F. Finster and A. Strohmaier. Gupta–Bleuler quantization of the Maxwell field in globally hyperbolic space-times. *Ann. Henri Poincaré*, 16(8):1837–1868, 2015.
- [10] E. P. Furlani. Quantization of the electromagnetic field on static space–times. *J. Math. Phys.*, 36(3):1063–1079, 1995.
- [11] J.-P. Gazeau and H. Pejhan. Covariant quantization of the partially massless graviton field in de Sitter spacetime. *Phys. Rev. D*, 108(6):065012, 2023.
- [12] C. Gérard. *Microlocal Analysis of Quantum Fields on Curved Spacetimes*. European Mathematical Society, arXiv:1901.10175, Zürich, 2019.
- [13] C. Gérard. Hadamard states for linearized gravity on spacetimes with compact Cauchy surfaces. *arXiv:2311.11043*, 2023.
- [14] C. Gérard, S. Murro, and M. Wrochna. Wick rotation of linearized gravity in Gaussian time and Calderón projectors. *arXiv:2204.01094*, 2023.
- [15] C. Gérard and M. Wrochna. Hadamard states for the linearized Yang–Mills equation on curved spacetime. *Commun. Math. Phys.*, 337(1):253–320, 2015.
- [16] C. Gérard and M. Wrochna. Analytic Hadamard states, Calderón projectors and Wick rotation near analytic Cauchy surfaces. *Commun. Math. Phys.*, 366(1):29–65, 2019.

- [17] D. Glavan and T. Prokopec. Even the photon propagator must break de Sitter symmetry. *Phys. Lett. B*, 841:137928, 2023.
- [18] T.-P. Hack and A. Schenkel. Linear bosonic and fermionic quantum gauge theories on curved spacetimes. *Gen. Relativ. Gravit.*, 45(5):877–910, 2013.
- [19] A. Higuchi. Quantum linearization instabilities of de Sitter spacetime. I. *Class. Quantum Gravity*, 8(11):1961–1981, 1991.
- [20] A. Higuchi. Quantum linearization instabilities of de Sitter spacetime. II. *Class. Quantum Gravity*, 8(11):1983–2004, 1991.
- [21] A. Higuchi and S. S. Kouris. The covariant graviton propagator in de Sitter spacetime. *Class. Quantum Gravity*, 18(20):4317–4327, 2001.
- [22] A. Higuchi, D. Marolf, and I. A. Morrison. de Sitter invariance of the dS graviton vacuum. *Class. Quantum Gravity*, 28(24):245012, 2011.
- [23] S. Hollands. Renormalized quantum Yang-Mills fields in curved spacetime. *Rev. Math. Phys.*, 20(09):1033–1172, 2008.
- [24] C. Iuliano and J. Zahn. Canonical quantization of Teukolsky fields on Kerr backgrounds. *Phys. Rev. D*, 108(12):125017, 2023.
- [25] A. Lichnerowicz. Propagateurs et commutateurs en relativité générale. *Publ. Mathématiques l’IHÉS*, 10(1):5–56, 1961.
- [26] S. P. Miao, P. J. Mora, N. C. Tsamis, and R. P. Woodard. Perils of analytic continuation. *Phys. Rev. D*, 89(10):104004, 2014.
- [27] S. P. Miao, N. C. Tsamis, and R. P. Woodard. de Sitter breaking through infrared divergences. *J. Math. Phys.*, 51(7), 2010.
- [28] S. P. Miao, N. C. Tsamis, and R. P. Woodard. The graviton propagator in de Donder gauge on de Sitter background. *J. Math. Phys.*, 52(12), 2011.
- [29] V. Moncrief. Decompositions of gravitational perturbations. *J. Math. Phys.*, 16(8):1556–1560, 1975.
- [30] V. Moretti, S. Murro, and D. Volpe. The quantization of Proca fields on globally hyperbolic spacetimes: Hadamard states and Møller operators. *Ann. Henri Poincaré*, 24(9):3055–3111, 2023.
- [31] H. Ringström. *The Cauchy Problem in General Relativity*. European Mathematical Society Publishing House, Zürich, 2009.
- [32] H. Sahlmann and R. Verch. Microlocal spectrum condition and Hadamard form for vector-valued quantum fields in curved spacetime. *Rev. Math. Phys.*, 13(10):1203–1246, 2001.
- [33] K. Sanders, C. Dappiaggi, and T.-P. Hack. Electromagnetism, local covariance, the Aharonov–Bohm effect and Gauss’ law. *Commun. Math. Phys.*, 328(2):625–667, 2014.
- [34] M. Wrochna and J. Zahn. Classical phase space and Hadamard states in the BRST formalism for gauge field theories on curved spacetime. *Rev. Math. Phys.*, 29(04):1750014, 2017.

LABORATOIRE DE MATHÉMATIQUES D’ORSAY, UNIVERSITÉ PARIS-SACLAY, FRANCE
Email address: christian.gerard@math.u-psud.fr

MATHEMATICAL INSTITUTE, UTRECHT UNIVERSITY, THE NETHERLANDS
Email address: m.wrochna@uu.nl