Statistical Inference on the Cumulative Distribution Function using Judgment Post Stratification

Mina Azizi Kouhanestani, * Ehsan Zamanzade[†], and Sareh Goli[‡]

May 1, 2024

Abstract

In this work, we discuss a general class of the estimators for the cumulative distribution function (CDF) based on judgment post stratification (JPS) sampling scheme which includes both empirical and kernel distribution functions. Specifically, we obtain the expectation of the estimators in this class and show that they are asymptotically more efficient than their competitors in simple random sampling (SRS), as long as the rankings are better than random guessing. We find a mild condition that is necessary and sufficient for them to be asymptotically unbiased. We also prove that given the same condition, the estimators in this class are strongly uniformly consistent estimators of the true CDF, and converge in distribution to a normal distribution when the sample size goes to infinity. We then focus on the kernel distribution function (KDF) in the JPS design and obtain the optimal bandwidth. We next carry out a comprehensive Monte Carlo simulation to compare the performance of the KDF in the JPS design for different choices of sample size, set size, ranking quality, parent distribution, kernel function as well as both perfect and imperfect rankings set-ups with its counterpart in SRS design. It is found that the JPS estimator dramatically improves the efficiency of the KDF as compared to its SRS competitor for a wide range of the settings. Finally, we apply the described procedure on a real dataset from medical context to show their usefulness and applicability in practice.

^{*}Department of Mathematical Sciences, Isfahan University of Technology, Isfahan, 84105, Iran. m.azizi@math.iut.ac.ir

[†]Department of Statistics, Faculty of Mathematics and Statistics, University of Isfahan, Isfahan 81746-73441, Iran. Corresponding author: e.zamanzade@sci.ui.ac.ir; ehsanzamanzadeh@yahoo.com

[‡]Department of Mathematical Sciences, Isfahan University of Technology, Isfahan, 84105, Iran. s.goli@iut.ac.ir

Keywords: Nonparametric estimation; Monte Carlo simulation; Judgment ranking; Statistical inference

Mathematics Subject Classifications 2020: 62D05; 62G05

1 Introduction

In many practical studies including ecological, medical, and environmental researches, there are cases in which accurate quantification of characteristic of interest (X) for all units in a sample is difficult or costly but ranking them can be done easily or inexpensively. In such situations, two alternatives sampling designs to simple random sampling (SRS) are ranked set sampling (RSS) and judgment post stratification (JPS) sampling scheme.

RSS was firstly suggested by McIntyre (1952) in his effort for efficient estimation of pasture mean and forage yields. He noticed that although measuring a yield is costly and tedious since it requires harvesting the crops, an agri-expert can produce a good ranking of the yields by eye inspection. To obtain an RSS sample of size n, one first determines the value of H and a vector $\mathbf{N} = (N_1, \ldots, N_H)$, where H is called the set size and N is called the vector of post strata sample sizes such that N_r represents the count of units with rank r that need to be chosen for accurate quantification, ensuring that the sum of all N_r values from r = 1 to H is equal to n. He/She then draws an SRS sample of size $n \times H$ from the population of interest and randomly divides it into n samples of size H. In the next step, each SRS sample of size H is ranked in an increasing magnitude without actual measurement. Lastly, from the N_r ranked samples of size H, units with judgment rank r are marked for accurate quantification (r = 1, ..., H). If $N_1 = ... = N_H$, sampling scheme is called balanced RSS. The term judgment rank is employed to emphasize that the ranking process relies on personal judgment, visual assessment, or an auxiliary variable that strongly correlates with the variable of interest. Hence, it may not be precise and has some errors, which is called imperfect ranking. The best situation is perfect ranking, in which there is no error in the mechanism of ranking and thus judgment ranks coincide with the true ones.

After the introduction of RSS, it was studied in many researches. Takahasi and Wakimoto (1968) showed that the RSS mean estimator is not biased and has no larger variance than the SRS mean estimator of comparable size. Estimation of variance for RSS was considered by Stokes (1980), MacEachern et al. (2002) and Frey and Feeman (2013). Stokes and Sager (1988) and Dümbgen and Zamanzade (2020) studied that the piece-wise linear estimation of the cumulative distribution function (CDF) in RSS. Gulati (2004) and Eftekharian and Razmkhah (2017) developed kernel-type estimators of the CDF and investigated their asymptotic properties. Also RSS method has been utilized to address almost all standard statistical topics such as the two sample problems (Mahdizadeh and Zamanzade, 2017, 2018, 2021; Frey and Zhang, 2019b; Moon et al.,

2020; Zamanzade et al., 2020), the prevalence estimation (Alvandi and Hatefi, 2021; Frey and Zhang, 2019a, 2021) and estimation in a parametric family (Qian et al., 2021; He et al., 2020, 2021; Chen et al., 2018, 2021).

MacEachern et al. (2004) proposed JPS sampling plan as a more flexible and applicable version of RSS. To obtain a JPS sample of size n using the set size H, an SRS sample of size n is drawn and X values for all units are quantified. Then, for each quantified unit, another SRS sample of size H - 1 is drawn to create n sets of size H in total. In the next step, each set of size H is ranked in an increasing magnitude without actual measurement. Finally, in each set, the rank of the quantified unit is noted. The JPS data, therefore, includes n independent and identically distributed (i.i.d) pairs $\{(X_i, R_i), i = 1, \ldots, n\}$, in which X_i denotes the quantified value for the *i*th sampled unit and R_i is its judgment rank among H units in the set. Thus, a JPS is a sampling plan in which an SRS sample is supported with judgment ranks.

While RSS and JPS sampling share a common underlying concept, there are significant distinctions between the two methods. The first one is about when the mechanism of ranking is carried out. In RSS, the ranking process is done **prior** to selecting the units for their exact measurements, so the ranks are strongly connected to the observations and cannot be ignored. Thus, it is not possible to utilize SRS statistical methods for RSS samples. In fact the RSS samples can only be analyzed by appropriate techniques that are specifically developed for them. But in JPS sampling scheme, the ranks are obtained **after** quantifying sample units and therefore they are loosely related to them. So, JPS sampling is more flexible than RSS from practical point of view, since the existing SRS techniques can be readily used for JPS samples by ignoring the ranks information. It is useful when it is believed that the ranking quality is not good or suitable statistical techniques to analyze the JPS sample are not available. Another difference is about the vector of post strata sample sizes: while the vector **N** is determined before the RSS sample is obtained, it is a random vector in JPS sampling. This induces an extra variability to the JPS sample, so the statistical inference using JPS is anticipated to be slightly less efficient than RSS.

Recently, many studies have been conducted in JPS sampling. For instance, Wang et al. (2008) and Frey and Feeman (2012) studied estimation of the population mean in JPS. Ozturk (2012) provided an alternative JPS sampling plan to combine the judgment ranks of different rankers. Wang et al. (2012) developed isotonized estimators for the CDF. The problem of variance estimation has been considered by Frey and Feeman (2013) and Zamanzade (2016). Ozturk (2015) developed sign test and quantile estimators for a JPS sample. Dastbaravarde et al. (2016) discussed

some properties of nonparametric estimation in the JPS setting. Estimation of population proportion for a JPS sample was investigated by Zamanzade and Wang (2017). Zamanzade and Vock (2018) developed perfect judgment ranking for JPS sampling scheme. Omidvar et al. (2018) combined the judgment ranks and obtained the maximum likelihood estimation of model parameters. Ozturk and Kravchuk (2021) presented some alternative JPS estimators to combine rank information from different sources. Dizicheh et al. (2021) studied odds estimation in the JPS and Alvandi and Hatefi (2023) discussed estimation of categorical ordinal populations for a JPS sample

One deficiency concerns with the empirical distribution function (EDF) is that it provides a discrete estimate for a continuous smooth CDF F. Thus, it is not possible to use its derivative to draw statistical inference about any functional of the probability density function (PDF). Moreover, it also fails to properly estimate the CDF beyond the extreme observations. To address these concerns, many studies have been done based on SRS for smooth estimation of the CDF, including Nadaraya (1964), Watson and Leadbetter (1964), Winter (1973) and Yamato (1973).

To the best of our knowledge, the problem of smooth estimation of the CDF has not been addressed in the JPS setting, yet. In Section 2, we introduce a class of estimators for the CDF in the JPS setting in a general form, which contains both empirical and smooth ones, and we study some of their finite sample size properties. We also prove that their asymptotic variances are no larger than their SRS competitors of the same size regardless of ranking quality. In Section 3, we obtain some asymptotic results for our proposed estimators. Specifically, we find the condition for them to be unbiased and strongly uniformly consistent estimators of the true CDF. We also establish their asymptotic normality under the same condition. In Section 4, we obtain some bias corrected results for the smooth CDF estimators and discuss optimal bandwidth selection with respect to minimizing the mean square error (MSE) of the estimators. Section 5, contains an extensive simulation results to compare the performance of the JPS and SRS based CDF estimators. The application of the proposed procedure is illustrated in Section 6. The discussion section, labeled as Section 7, includes some final remarks and outlines potential avenues for future research.

2 Introduction of the CDF Estimator

Let X_1, \ldots, X_n be an SRS sample of size *n* from a population with an unknown CDF *F*. A general class of the CDF estimators of *F* is given by

$$F_{n;srs}(t) = \frac{1}{n} \sum_{i=1}^{n} K_n(t - X_i), \qquad (1)$$

where $K_n(.)$ is a given CDF.

Clearly, $F_{n;srs}(t)$ is a convex combination of the CDFs, and thus it itself is a CDF, as well. In some practical situations, the CDF F(t) is a continuous function in t, so it is reasonable to utilize a continuous CDF as $K_n(.)$. Specially, if it is assumed that the CDF F(t) is an absolutely continuous function in t, then a CDF estimator, which enjoys the said property, can be obtained by taking an absolutely continuous CDF for $K_n(.)$. Thus an estimator for the population PDF fcan be obtained as

$$f_{n;srs}(t) = \frac{1}{n} \sum_{i=1}^{n} k_n (t - X_i),$$

where $k_{n}(.)$ is the first derivative of $K_{n}(.)$.

Obviously, the estimator $F_{n;srs}(t)$ coincides with the EDF $F_{n;srs}^*(t) = \frac{1}{n} \sum_{i=1}^n \mathbb{I}(X_i \le t)$ if $K_n(t) = e(t)$, where $e(t) = \mathbb{I}(t \ge 0)$, and $\mathbb{I}(.)$ is the standard indicator function.

Statistical properties of $F_{n;srs}(t)$ have been discussed by Yamato (1973). He found the necessary and sufficient condition for the estimators in this class to be asymptotically unbiased, and proved their consistencies and normalities under the same condition.

Let $\{(X_i, R_i), i = 1, ..., n\}$ is a JPS sample of size n from a population with an unknown CDF F. As mentioned before, R_i , for i = 1, ..., n is judgment rank of X_i among H units in its set, so, we set $\mathbf{R} = (R_1, ..., R_n)$ as the ranks vector. The variable I_{ir} is defined as follows: If observation X_i has a judgment rank of r (i.e., $R_i = r$), then $I_{ir} = 1$; otherwise, $I_{ir} = 0$. This definition applies to every i ranging from 1 to n and every r ranging from 1 to H. With this definition, if N_r denotes the number of observations X_i in the rth post strata, then we have $N_r = \sum_{i=1}^n I_{ir}$, and the vector of post strata sample sizes $\mathbf{N} = (N_1, \ldots, N_H)$ follows a multinomial distribution with mass parameter n and probability vector $(\frac{1}{H}, \ldots, \frac{1}{H})$. Likewise, if we define $J_r = 1/N_r$ if $N_r > 0$, otherwise $J_r = 0$, for $r = 1, \ldots, H$, then $d_n = \sum_{r=1}^H \mathbb{I}(N_r > 0)$ denotes the number of nonempty post strata which formed during JPS sampling procedure. Furthermore, note that the conditional distribution of each X_i given its rank $R_i = r$, denoted as $F_{[r]}$, is equivalent to the distribution of

the *r*th order statistic $(X_{[r]})$ from a sample of size *H*. Finally, it is worth mentioning that if the same ranking process is applied for all sets of size *H* during the course of JPS, then the ranking mechanism is called **consistent**. According to Presnell and Bohn (1999), if the ranking process is consistent, then for all $t \in \mathbb{R}$, we have

$$F(t) = \frac{1}{H} \sum_{r=1}^{H} F_{[r]}(t).$$

The above equality, which is known as the fundamental equality, plays an important role in establishing our results.

Using above notation, a general class of the CDF estimators in the JPS setting is proposed as

$$F_{n;jps}(t) = \sum_{r=1}^{H} W_r F_{n;[r]}(t),$$
(2)

where $W_r = \frac{\mathbb{I}(N_r > 0)}{d_n}$, $F_{n;[r]}(t) = \frac{1}{N_r} \sum_{i=1}^n K_n (t - X_i) I_{ir}$ if $N_r > 0$, and zero otherwise, where K_n (.) is a given CDF. Similar to SRS, an absolutely continuous estimator of the CDF in the JPS setting can be obtained by an appropriate choice of K_n (.). Furthermore, if we take $K_n(t) = e(t)$, then the EDF in the JPS setting is obtained, which has the following form

$$F_{n;jps}^{*}(t) = \sum_{r=1}^{H} W_r F_{n;[r]}^{*}(t),$$
(3)

where $F_{n;[r]}^{*}(t) = \begin{cases} \frac{1}{N_r} \sum_{i=1}^n \mathbb{I}(X_i \leq t) I_{ir} & \text{if } N_r > 0, \\ 0 & \text{otherwise.} \end{cases}$

To study the properties of $F_{n;jps}(t)$, we first note that both $F_{n;jps}(t)$ and $F_{n;srs}(t)$ have a

common amount of bias. To see this, note that

$$\mathbb{E}(F_{n;jps}(t)) = \mathbb{E}\left(\sum_{r=1}^{H} W_r F_{n;[r]}(t)\right)$$

$$= \mathbb{E}\left(\mathbb{E}\left(\sum_{r=1}^{H} W_r F_{n;[r]}(t) \mid \mathbf{R}\right)\right)$$

$$= \mathbb{E}\left(\sum_{r=1}^{H} W_r \mathbb{E}\left(K_n(t - X_{[r]})\right)\right)$$

$$= \mathbb{E}(W_1) \sum_{r=1}^{H} \mathbb{E}\left(K_n(t - X_{[r]})\right)$$

$$= \frac{1}{H} \sum_{r=1}^{H} \mathbb{E}\left(K_n(t - X_{[r]})\right)$$

$$= \mathbb{E}(F_{n;srs}(t)), \qquad (4)$$

in which the second last equality follows from the fact that W_1, \ldots, W_H are identically distributed with $\mathbb{E}(W_1) = \frac{1}{H}$, and the last equality is an immediate result of the fundamental equality.

To obtain the variance of $F_{n;jps}(t)$, by following lines of Dastbaravarde et al. (2016), and relying on the conditional variance formula, one can obtain

$$\mathbb{V}(F_{n;jps}(t)) = \mathbb{E}(W_1^2 J_1) \sum_{r=1}^H \mathbb{V}\left(K_n(t - X_{[r]})\right) + \frac{H}{H-1} \mathbb{V}(W_1) \left(\sum_{r=1}^H \left(\mathbb{E}\left(K_n(t - X_{[r]})\right) - \mathbb{E}\left(K_n(t - X)\right)\right)^2\right)$$
(5)

Using the fundamental equality, one can show that $\mathbb{V}(K_n(t-X)) = \frac{1}{H} \sum_{r=1}^{H} \mathbb{V}(K_n(t-X_{[r]})) + \frac{1}{H} \sum_{r=1}^{H} (\mathbb{E}(K_n(t-X_{[r]})) - \mathbb{E}(K_n(t-X)))^2)$, and therefore, the variance of $F_{n;jps}(t)$ can be also re-written as

$$\mathbb{V}(F_{n;jps}(t)) = H\mathbb{E}(W_1^2 J_1)\mathbb{V}\left(K_n(t-X)\right) \\ - \left[\mathbb{E}(W_1^2 J_1) - \frac{H}{H-1}\mathbb{V}(W_1)\right] \left(\sum_{r=1}^H \left(\mathbb{E}\left(K_n(t-X_{[r]})\right) - \mathbb{E}\left(K_n(t-X)\right)\right)^2\right).$$
(6)

The properties of the vector (W_1, \ldots, W_H) have been investigated by Dastbaravarde et al. (2016). Specifically, they showed that for $r = 1, \ldots, H$

$$\mathbb{V}(W_r) = \frac{1}{H^2} \sum_{l=1}^{H-1} \left(\frac{l}{H}\right)^{n-1},\tag{7}$$

$$\mathbb{E}(W_r^2 J_r) = \frac{1}{H^n} \left[\frac{1}{n} + \sum_{d_n=2}^H \sum_{j=1}^{d_n-1} \sum_{n_1=1}^{n-d_n+1} \frac{(-1)^{j-1}}{d_n^2 n_1} \binom{H-1}{d_n-1} \binom{d_n-1}{j-1} \binom{n}{n_1} (d_n-j)^{n-n_1} \right].$$
(8)

Also for every fixed H, they have established a proof that as the sample size n approaches infinity, then

$$n\mathbb{V}(W_r) \to 0,$$
(9)

$$nH\mathbb{E}(W_r^2 J_r) \to 1,\tag{10}$$

$$\sqrt{n}\left(W_r - \frac{1}{H}\right) \xrightarrow{\text{a.s.}} 0,$$
(11)

where $\xrightarrow{a.s}$ means the almost sure convergence.

According to (9) and (10), we can write

$$\lim_{n \to \infty} \mathbb{V}(F_{n;jps}(t)) = \mathbb{V}(F_{n;srs}(t))(1-\delta),$$

where $\delta = \frac{1}{H^{\mathbb{V}(K_n(t-X))}} \sum_{r=1}^{H} \left(\mathbb{E} \left(K_n(t-X_{[r]}) \right) - \mathbb{E} \left(K_n(t-X) \right) \right)^2$ is less than one. Thus, we can conclude that asymptotic variance of the CDF estimator in JPS setting is no larger than its SRS competitor of size n, even when the ranking is not perfect but still better than random guessing.

3 Some Asymptotic Results

In this section, we study the asymptotic properties of the CDF estimators in the proposed class, when the set size H is fixed and the sample size n tends to infinity. We prove Glivenko–Cantelli type convergence for the estimators and establish the asymptotic distribution of them. To do so, we rely on the following definition for convergence of a sequence of CDFs.

Definition 1. Let $\{G_n\}$ be a sequence of CDFs and G be a CDF. Then we define the sequence $\{G_n\}$ weakly converges to G and write $G_n \xrightarrow{w} G$, if

$$G_n(t) \to G(t), \quad as \quad n \to \infty,$$

for all $t \in C(G)$, where C(G) denotes the set of all continuity points of G.

The first theorem concerns with asymptotic mean of the JPS estimators of the CDF. Specifically, it finds necessary and sufficient condition for asymptotic unbiasedness of $F_{n;jps}(t)$. Theorem 1 can be simply proven using Lemma 1 in Yamato (1973) and equation (4), and thus its proof is omitted.

Theorem 1. (Asymptotic unbiasedness) Let $\{(X_i, R_i), i = 1, ..., n\}$ be a JPS sample from a population with an unknown CDF F. Then if the mechanism of ranking is consistent, and for a fixed value of H, every estimator of the form $F_{n;jps}(t)$ in (2) is asymptotically unbiased if and only if $K_n \xrightarrow{w} e$.

The next result concerns with Glivenko-Cantelli type convergence property of the estimator $F_{n;jps}(t)$.

Theorem 2. (Strongly uniformly convergence) Let $\{(X_i, R_i), i = 1, ..., n\}$ be a JPS sample from a population with an unknown CDF F. If the mechanism of ranking is consistent and $K_n \xrightarrow{w} e$, then for a fixed value of H, we have

$$\sup_{t \in \mathbb{R}} \left| F_{n;jps}(t) - F(t) \right| \xrightarrow{a.s} 0.$$

Proof. From fundamental equality, we can write

$$\sup_{t \in \mathbb{R}} \left| F_{n;jps}(t) - F(t) \right| = \sup_{t \in \mathbb{R}} \left| \sum_{r=1}^{H} W_r F_{n;[r]}(t) - \frac{1}{H} \sum_{r=1}^{H} F_{[r]}(t) \right|$$
$$\leq \sum_{r=1}^{H} \sup_{t \in \mathbb{R}} \left| W_r F_{n;[r]}(t) - \frac{1}{H} F_{[r]}(t) \right|, \tag{12}$$

where the last inequality follows from triangle inequality. Besides, for every $r = 1, \ldots, H$, we have

$$\begin{split} \sup_{t \in \mathbb{R}} |W_r F_{n;[r]}(t) - \frac{1}{H} F_{[r]}(t)| &\leq \sup_{t \in \mathbb{R}} |W_r F_{n;[r]}(t) - \frac{1}{H} F_{n;[r]}(t)| + \sup_{t \in \mathbb{R}} \left| \frac{1}{H} F_{n;[r]}(t) - \frac{1}{H} F_{[r]}(t) \right| \\ &= \sup_{t \in \mathbb{R}} \left(|W_r - \frac{1}{H}| F_{n;[r]}(t) \right) + \sup_{t \in \mathbb{R}} \left(\frac{1}{H} |F_{n;[r]}(t) - F_{[r]}(t)| \right) \\ &= |W_r - \frac{1}{H}| \sup_{t \in \mathbb{R}} F_{n;[r]}(t) + \frac{1}{H} \sup_{t \in \mathbb{R}} |F_{n;[r]}(t) - F_{[r]}(t)|. \end{split}$$

It is clear from equation (11) we have $(W_r - \frac{1}{H}) \xrightarrow{\text{a.s.}} 0$. Furthermore, from Theorem 3 in Yamato (1973), one can conclude that

$$\sup_{t\in\mathbb{R}}\left|F_{n;[r]}(t)-F_{[r]}(t)\right|\xrightarrow{\mathrm{a.s.}}0,$$

for $r = 1, \ldots, H$, and this completes the proof.

We now establish the asymptotic normality of the JPS estimator.

Theorem 3. (Asymptotic normality) Let $\{(X_i, R_i), i = 1, ..., n\}$ be a JPS sample from a population with an unknown CDF F. If the mechanism of ranking is consistent and $K_n \xrightarrow{w} e$, then for a fixed value of H, we have

$$\sqrt{n} (F_{n;jps}(t) - F(t)) \xrightarrow{d} N \left(0, \frac{1}{H} \sum_{r=1}^{H} F_{[r]}(t) [1 - F_{[r]}(t)] \right),$$

for all $t \in C(F)$ with $F(t) \neq 0$ or 1, where \xrightarrow{d} means convergence in distribution.

Proof. We can write

$$\sqrt{n} \left(F_{n;jps}(t) - F(t) \right) = \sqrt{n} \left(\sum_{r=1}^{H} W_r F_{n;[r]}(t) - F(t) \right)$$
$$= \sqrt{n} \left(\sum_{r=1}^{H} W_r \left(F_{n;[r]}(t) - F_{[r]}(t) \right) \right) + \sqrt{n} \left(\sum_{r=1}^{H} F_{[r]}(t) \left(W_r - \frac{1}{H} \right) \right).$$

From (11), it is clear that $\sqrt{n} \left(\sum_{r=1}^{H} F_{[r]}(t) \left(W_r - \frac{1}{H} \right) \right) \xrightarrow{\text{a.s.}} 0$, thus it is sufficient to obtain asymptotic distribution of $\sqrt{n} \left(\sum_{r=1}^{H} W_r \left(F_{n;[r]}(t) - F_{[r]}(t) \right) \right)$. For this purpose, we define

$$\sqrt{n}\left(\sum_{r=1}^{H} W_r\left(F_{n;[r]}(t) - F_{[r]}(t)\right)\right) = \mathbf{A_n T_n},$$

where $\mathbf{A_n} = \sqrt{n} \left(\sqrt{J_1} W_1, \dots, \sqrt{J_H} W_H \right)$ and

$$\mathbf{T_n} = \left(\sqrt{N_1} \left(F_{n;[1]}(t) - F_{[1]}(t)\right), \dots, \sqrt{N_H} \left(F_{n;[H]}(t) - F_{[H]}(t)\right)\right)^T,$$

where T denotes the transpose operator. Also from (11), we have $(W_r - \frac{1}{H}) \xrightarrow{\text{a.s.}} 0$ and using law of large numbers, we obtain $nJ_r \xrightarrow{\text{a.s.}} H$. Thus $\mathbf{A_n} \xrightarrow{\text{a.s.}} \left(\sqrt{\frac{1}{H}}, \dots, \sqrt{\frac{1}{H}}\right)$.

On the other hand, since $F_{n;[r]}(t)$ s, given the vector of ranks **R**, are conditional independent random variables, we can write

$$P(\bigcap_{r=1}^{H} \sqrt{N_r} \left(F_{n;[r]}(t) - F_{[r]}(t) \right) \le t_r) = E\left[P(\bigcap_{r=1}^{H} \sqrt{N_r} \left(F_{n;[r]}(t) - F_{[r]}(t) \right) \le t_r \mid \mathbf{R}) \right]$$
$$= E\left(\prod_{r=1}^{H} P\left(\sqrt{N_r} (F_{n;[r]}(t) - F_{[r]}(t)) \le t_r \mid \mathbf{R} \right) \right)$$
$$\to E\left(\prod_{r=1}^{H} P\left(T_r \le t_r \right) \right)$$
$$= P\left(\bigcap_{r=1}^{H} T_r \le t_r \right).$$

From Theorem 1 in Yamato (1973), one can conclude that if $K_n \xrightarrow{w} e$, then $T_r = \sqrt{n_r} \left(F_{n;[r]}(t) - F_{[r]}(t) \right)$ follows a mean-zero normal distribution with variance $F_{[r]}(t) \left[1 - F_{[r]}(t) \right]$, for all $t \in C(F_{[r]})$ with $F_{[r]}(t) \neq 0$ or 1.

Thus, we can conclude that the vector $\mathbf{T}_{\mathbf{n}}$ converges in distribution to an H-dimensional normal distribution with zero mean vector and variance covariance matrix \sum , where \sum is a diagonal matrix that its *r*th diagonal element is given by $F_{[r]}(t) [1 - F_{[r]}(t)]$ for $r = 1, \ldots, H$. So, using Slutsky's theorem, we have

$$\sqrt{n}\left(\sum_{r=1}^{H} W_r\left(F_{n;[r]}(t) - F_{[r]}(t)\right)\right) = \mathbf{A_nT_n} \xrightarrow{\mathrm{d}} N\left(0, \frac{1}{H}\sum_{r=1}^{H} F_{[r]}(t)\left[1 - F_{[r]}(t)\right]\right).$$

4 CDF Estimation using Kernel Function

Up to now, we assume that the function $K_n(.)$ is an arbitrary CDF and all asymptotic results obtained under the mild condition that $K_n \xrightarrow{w} e$. However, in the statistical literature, many researchers prefer to impose some more constraints on $K_n(.)$, such that the resulting CDF estimator enjoys some nice properties like smoothness and having less bias. In this section, we impose such constraints on proposed CDF estimator in the JPS setting and then we obtain the optimal estimator in this case. In so doing, we assume that the CDF $K_n(t)$ depends on the sample size n via the parameter h_n , and therefore, it can be written as $K_n(t) = K\left(\frac{t}{h_n}\right)$, where K(.) is a given CDF and the parameter h_n is an smoothing parameter and is called bandwidth. We assume that the given CDF function K(.) can be written as

$$K(t) = \begin{cases} 0 & t < -a, \\ \int_{-a}^{t} k(x) dx & |t| \le a, \\ 1 & t > a, \end{cases}$$
(13)

such that k(.) is a symmetric PDF with a bounded support on [-a, a], and thus it satisfies the following constraints

$$\int_{-a}^{a} k(x)dx = 1, \quad \int_{-a}^{a} xk(x)dx = 0, \quad \int_{-a}^{a} x^{2}k(x)dx \neq 0.$$
(14)

It is also assumed that the bandwidth h_n follows the conditions

$$\lim_{n \to \infty} h_n = 0, \quad \lim_{n \to \infty} nh_n = \infty,$$

indicating that it converges to zero but at a rate slower than n^{-1} . To make the notations less cumbersome, hereafter, we omit the subscript n in h_n . In the statistical literature, the function k (.) satisfying (14) is usually known as kernel function and the method for estimating CDF (PDF) based on the function k (.) is called kernel CDF (PDF) estimation. Therefore, the kernel distribution function (KDF) based on an SRS sample is obtained by replacing (13) in equation (1) as

$$F_{n;srs}^{k}(t) = \frac{1}{n} \sum_{i=1}^{n} K\left(\frac{t - X_{i}}{h}\right).$$
(15)

Azzalini (1981) obtained the optimal value for bandwidth h based on an SRS sample of size n by minimizing large sample MSE of $F_{n;srs}^{k}(t)$ as

$$h_{srs} = \left(\frac{f(t)\left(a - \int_{-a}^{a} K^{2}(x)dx\right)}{n\left(f'(t)\int_{-a}^{a} x^{2}k(x)dx\right)^{2}}\right)^{\frac{1}{3}}$$

Let $\{(X_i, R_i), i = 1, ..., n\}$ is a JPS sample of size *n* from a population with an unknown CDF *F*. The kernel-type estimator of CDF based on the JPS sample can be obtained by substituting (13) in equation (2), and is given by

$$F_{n;jps}^{k}(t) = \sum_{r=1}^{H} W_{r} F_{n;[r]}^{k}(t),$$

where $F_{n;[r]}^k(t) = \frac{1}{N_r} \sum_{i=1}^n K\left(\frac{t-X_i}{h}\right) I_{ir}$ if $N_r > 0$, and zero otherwise. It is worth mentioning that the asymptotic results established in Section 3 hold for kernel-type

It is worth mentioning that the asymptotic results established in Section 3 hold for kernel-type estimators of the CDF, since $K \xrightarrow{w} e$ as the sample size n goes to infinity.

To obtain the mean of a KDF in the JPS setting, we first note that according to (4), one can write

$$\mathbb{E}\left(F_{n;jps}^{k}(t)\right) = \mathbb{E}\left(K\left(\frac{t-X}{h}\right)\right)$$
$$= \int_{-\infty}^{\infty} K\left(\frac{t-x}{h}\right) f(x)dx$$
$$= \int_{-\infty}^{t-ah} f(x)dx + \int_{t-ah}^{t+ah} K\left(\frac{t-x}{h}\right) f(x)dx$$

Note that since k(.) is symmetric around 0, we can write $K(x) = \frac{1}{2} + g(x)$; $|x| \le a$, where g(.) is

an odd function, and therefore we have

$$\mathbb{E}\left(F_{n;jps}^{k}(t)\right) = \int_{-\infty}^{t-ah} f(x)dx + \int_{t-ah}^{t+ah} \left(\frac{1}{2} + g\left(\frac{t-x}{h}\right)\right) f(x)dx$$

= $\frac{1}{2}\left(F(t+ah) + F(t-ah)\right) + \int_{t-ah}^{t+ah} g\left(\frac{t-x}{h}\right) f(x)dx$
= $\frac{1}{2}\left(F(t+ah) + F(t-ah)\right) + h \int_{-a}^{a} g(u) f(t-hu)du.$

Using Taylor series of F(t + ah), F(t - ah) and f(t - hu) around t, we have

$$\begin{split} \mathbb{E}\left(F_{n;jps}^{k}(t)\right) &= F(t) + \frac{a^{2}h^{2}}{2}f'(t) + hf(t)\int_{-a}^{a}g(u)du - h^{2}f'(t)\int_{-a}^{a}ug(u)du + O(h^{3})\\ &= F(t) + \frac{a^{2}h^{2}}{2}f'(t) - h^{2}f'(t)\int_{-a}^{a}ug(u)du + O(h^{3})\\ &= F(t) + h^{2}f'(t)\left(\frac{a^{2}}{2} - \int_{-a}^{a}ug(u)du\right) + O(h^{3}), \end{split}$$

where the second last equality follows from the fact that g(.) is an odd function.

By using integration by parts, we can write

$$\int_{-a}^{a} ug(u)du = \frac{a^2}{2} - \frac{1}{2} \int_{-a}^{a} u^2 k(u)du.$$

Therefore, we have

$$\mathbb{E}\left(F_{n;jps}^{k}(t)\right) = F(t) + \frac{h^{2}}{2}f'(t)\int_{-a}^{a}u^{2}k(u)du + O(h^{3})$$
(16)

$$=F(t) + O(h^2).$$
 (17)

To obtain the variance of $F_{n;jps}^k(t)$, we first require to calculate $\mathbb{V}\left(K\left(\frac{t-X}{h}\right)\right)$. To do so, note that

$$\mathbb{E}\left(K^{2}\left(\frac{t-X}{h}\right)\right) = \int_{-\infty}^{\infty} K^{2}\left(\frac{t-x}{h}\right) f(x)dx$$

$$= \int_{-\infty}^{t-ah} f(x)dx + \int_{t-ah}^{t+ah} K^{2}\left(\frac{t-x}{h}\right) f(x)dx$$

$$= \int_{-\infty}^{t-ah} f(x)dx + \int_{t-ah}^{t+ah} \left(\frac{1}{4} + g\left(\frac{t-x}{h}\right) + g^{2}\left(\frac{t-x}{h}\right)\right) f(x)dx$$

$$= \frac{3}{4}F(t-ah) + \frac{1}{4}F(t+ah) + h\int_{-a}^{a} g(u) f(t-hu)du + h\int_{-a}^{a} g^{2}(u) f(t-hu)du$$

$$= F(t) - \frac{ah}{2}f(t) + hf(t) \int_{-a}^{a} g^{2}(u) du - h^{2}f'(t) \int_{-a}^{a} ug^{2}(u) du + O(h^{2})$$

$$= F(t) - \frac{ah}{2}f(t) + hf(t) \int_{-a}^{a} g^{2}(u) du + O(h^{2})$$

$$= F(t) - hf(t) \left(\frac{a}{2} - \int_{-a}^{a} g^{2}(u) du\right) + O(h^{2}). \tag{18}$$

So, using (17) and (18), one can write

$$\begin{split} \mathbb{V}\left(K\left(\frac{t-X}{h}\right)\right) &= \mathbb{E}\left(K^2\left(\frac{t-X}{h}\right)\right) - \mathbb{E}^2\left(K\left(\frac{t-X}{h}\right)\right) \\ &= F(t)\left[1-F(t)\right] - hf(t)\left(\frac{a}{2} - \int_{-a}^{a} g^2(u)du\right) + O(h^2) \\ &= F(t)\left[1-F(t)\right] - hf(t)\left(a - \int_{-a}^{a} K^2(u)du\right) + O(h^2) \end{split}$$

where the last equality holds owing to $\int_{-a}^{a} g^{2}(u) du = \int_{-a}^{a} K^{2}(u) du - \frac{a}{2}$. Thus, using the relation (6), the MSE of $F_{n;jps}^{k}(t)$ is obtained as

$$MSE\left(F_{n;jps}^{k}(t)\right) = \mathbb{V}(F_{n;jps}^{k}(t)) + \text{bias}^{2}(F_{n;jps}^{k}(t))$$

$$\approx H\mathbb{E}(W_{1}^{2}J_{1})\left(F(t)\left[1 - F(t)\right] - hf(t)\left(a - \int_{-a}^{a} K^{2}(u)du\right)\right) - r(t) + \frac{h^{4}}{4}\left(f'(t)\int_{-a}^{a} u^{2}k(u)du\right)^{2},$$

where r(t) denotes the remaining terms which do not depend on the bandwidth h. Therefore, the optimal bandwidth h in the JPS setting can be obtained by minimizing the MSE of $F_{n;jps}^k(t)$ with respect to h as

$$h_{jps} = \left(\frac{nH\mathbb{E}(W_1^2J_1)f(t)\left(a - \int_{-a}^{a} K^2(x)dx\right)}{n\left(f'(t)\int_{-a}^{a} x^2k(x)dx\right)^2}\right)^{\frac{1}{3}}$$
$$= \left(nH\mathbb{E}(W_1^2J_1)\right)^{\frac{1}{3}}h_{srs}.$$

Remark 1. It is clear from equation (10) that $nH\mathbb{E}(W_1^2J_1) \to 1$, as $n \to \infty$, and therefore, the optimal bandwidth h in the JPS setting is almost the same as the SRS setting for large values of n.

Remark 2. We observe that the optimal bandwidth h depends on quantities f(t) and f'(t) which are often unknown in practice. We estimate these quantities using reference underlying distribution for these quantities (Silverman, 2018). To do so, we take reference distribution as exponential distribution with mean \bar{X} for the situations in which the variable of interest has a positive support such as lifetime data. If the variable of interest has support on real numbers \mathbb{R} , then normal distribution with mean \bar{X} and variance $S^2 = \frac{1}{n} \sum_{i=1}^{n} (X_i - \bar{X})^2$ is considered as the reference distribution.

It is noticeable that we can obtain kernel-type estimator of PDF f, from smooth estimator

 $F_{n;ips}^k(t)$ as follows

$$f_{n;jps}^{k}(t) = \frac{d}{dt} F_{n;jps}^{k}(t) = \sum_{r=1}^{H} W_{r} f_{n;[r]}^{k}(t),$$

where $f_{n;[r]}^k(t) = \frac{1}{N_r} \sum_{i=1}^n k\left(\frac{t-X_i}{h}\right) I_{ir}$ if $N_r > 0$, and zero otherwise.

5 Monte Carlo Simulation

In this section, we conduct a comprehensive Monte Carlo simulation to compare the performance of the KDF estimator based on SRS with its JPS sampling counterpart introduced in section 4, for both perfect and imperfect ranking set-ups. To this end, we have considered six different distributions as the parent distribution: standard normal distribution (N(0,1)), student's tdistribution with 5 degrees of freedom (t_5) , standard laplace distribution (La(0,1)), standard exponential distribution (E(1)), gamma distribution with scale parameter 1 and shape parameter 0.5 (G(0.5,1)), and gamma distribution with scale parameter 1 and shape parameter 2 (G(2,1)). Therefore, both distributions with support on the real numbers \mathbb{R} and lifetime distributions with support on $(0, +\infty)$ are considered in our study.

In this simulation study, we set $n \in \{10, 50, 300\}$, and $H \in \{3, 5, 10\}$, therefore, we can compare the performance of the estimators for small, moderate and large values of sample/set size. We can also observe the effect of increasing sample size (set size) on the performance of the estimators while the set size (sample size) is being fixed.

To perform ranking in the JPS samples, we use the method such as one utilized in linear ranking error model by Dell and Clutter (1972). According to this method, the ranking process is done by an auxiliary variable (Y) correlated with variable of interest (X) as

$$Y = \rho \frac{(X - \mu)}{\sigma} + \sqrt{1 - \rho^2} Z,$$

where μ and σ^2 are mean and variance of X, respectively, and the random variable Z is characterized by being independent of X, which itself follows a standard normal distribution. Also, the user controls quality of ranking by choosing the parameter $\rho \in [0, 1]$, as the correlation coefficient between X and Y. In this simulation study, we select $\rho = 1$ for perfect ranking, $\rho = 0.9$ for good ranking, $\rho = 0.75$ for moderate ranking and $\rho = 0.5$ for weak ranking. The kernel functions utilized in the KDF estimators are epanechnikov kernel function of the form $k(x) = \frac{3}{4}(1-x^2); |x| \le 1$, triangular kernel function of the form $k(x) = 1 - |x|; |x| \le 1$, cosine kernel function of the form $k(x) = \frac{\pi}{4}cos(\frac{\pi}{2}x); |x| \le 1$ and truncated gaussian kernel function of the form $k(x) = \frac{\phi(x)}{\Phi(4) - \Phi(-4)}; |x| \le 4$, where the functions $\phi(.)$ and $\Phi(.)$ represent the PDF and CDF of standard normal distribution, respectively.

The relative efficiency (RE) of $F_{n;srs}^{k}(t)$ with respect to $F_{n;jps}^{k}(t)$ is defined as the ratio of their MSEs, which can be expressed as follows

$$RE(p) = \frac{MSE(F_{n;srs}^k(Q_p))}{MSE(F_{n;ips}^k(Q_p))},$$

where Q_p is the *p*th quantile of the parent distribution. It is clear that an *RE* value larger than 1 indicates an advantage of using JPS estimator instead of SRS one at the *p*th quantile of the parent distribution.

For each (n, H, ρ) , we have generated 100,000 random samples from JPS and SRS designs and obtained RE(p) for $p \in \{0.01, 0.02, \dots, 0.99\}$. Here, we only present the results in Figures 1-6 for epanechnikov kernel function, because we have observed that choice of kernel function does not have much effect on the RE of the estimators.

Figure 1 presents the simulation results for sample size n = 10, and three distributions with support on \mathbb{R} . It is clear from this figure that the patterns of the performance of the CDF estimators are almost the same across different distributions. The RE curves are symmetric and reach their maximum values around their symmetrical point, p = 0.5. Furthermore, the minimum values of the RE curves are observed at the points close to zero/one. The first row panels in Figure 1 display the outcomes for the perfect ranking scenario. It is clear from these panels that the RE curves for H = 3 and H = 5 are almost identical and are slightly higher than that of for H = 10. This can be justified by the fact that empty strata are frequently observed for (n, H) = (10, 10), so the post stratification with set size H = 10 does not contribute much in improving estimation precision. Furthermore, the REs are higher than one except for two narrow intervals at the boundaries of the parent distribution. The simulation results for good ranking case ($\rho = 0.9$) are depicted in the second row of Figure 1. From these panels, it can be observed that the patterns of the RE curves closely resemble those of the perfect ranking case ($\rho = 1$), with a clear difference that the REs are lower in this case and the span of intervals in which the REs are below one become wider. Simulation results for $\rho = 0.75$, and $\rho = 0.5$ are demonstrated in the third and the bottom rows of the Figure 1, respectively. It is evident that although the patterns of RE curves are the same as those of $\rho = 0.9$, RE values decrease as the value of ρ decreases. Specifically, for $\rho = 0.5$, the REs

are below one except for a narrow interval around p = 0.5.

The simulation results for sample size n = 50, and three distributions with support on \mathbb{R} are shown in Figure 2. It is obvious that the RE curves have symmetric forms and are almost the same across different distributions. The three top panels of Figure 2 show the results for $\rho = 1$. It is evident from these panels that the efficiencies of the JPS estimators are increasing function in set size H, and thus RE curve for H = 10 (H = 5) is uniformly higher than that for H = 5(H = 3). Each RE curve reaches its maximum for p = 0.5 and minimum for p = 0.01 and p = 0.99. Furthermore, it has two local minimum points around p = 0.4 and p = 0.6, and falls below one for a very narrow interval close to zero/one. The panels in the second top row of Figure 2 demonstrate the results for $\rho = 0.9$. The patterns of the performances of JPS estimators are almost the same as those for $\rho = 1$, with a obvious difference that the REs are lower in this case. We can observe the performances of the JPS estimators for $\rho = 0.75$ in the panels of the third row of Figure 2. There are two main differences among the RE curves in this case and those for $\rho = 0.9$. First, the RE curves are lower and the spans of the intervals in the REs are below of one are wider for $\rho = 0.75$ than $\rho = 0.9$. Second, the RE curves are not increasing function in H, any more. These differences are also observed when we compare the results for $\rho = 0.5$ in the panels in the bottom row of Figure 2 with those for $\rho = 0.75$. It is of interest to note that the JPS estimator with H = 10 has the lowest efficiency in this case.

The simulation results for sample size n = 300, and three distributions with support on \mathbb{R} are shown in Figure 3. Thus, from this figure, we can observe the asymptotic performance of the estimators. It is clear that REs increase (decrease) when the set size H (ranking quality ρ) increases (decreases) while the other parameters are kept fixed. It is also worth mentioning that the REs never fall below one for n = 300.

Figures 4-6 present simulation results for lifetime distributions for sample sizes n = 10, 50 and 300, respectively. Figure 4 shows the simulation results for lifetime distributions for n = 10. Based on the presented figure, it is evident that the RE curves of the JPS estimators resemble different shapes for different distributions. From the panels in the first row of Figure 4, we observe the JPS estimator for H = 5 (H = 10) usually has the best (worst) performance. The RE curves are not symmetric, but they reach to their maximum around the center of the distribution and (local minimum) minimum around the (lower) upper tail of the parent distribution which is below one. From the panels in the second row of Figure 4, we observe the patterns of the performances of the JPS estimators for $\rho = 0.9$ are almost the same as those for $\rho = 1$, but they are lower. The

simulation results are not in favor of the JPS estimators for $\rho = 0.75$ and $\rho = 0.5$, which are shown in the third and bottom rows of Figure 4, respectively, as their REs are below one for most values of p.

Simulation results for lifetime distributions when the sample size is n = 50 are depicted in Figure 5. Based on the panels displayed in the first row of the provided figure, we can observe that in cases where the ranking process is perfect ($\rho = 1$), the JPS estimator with H = 10 performs better than the others in most considered cases, which is followed by the JPS estimators with H = 5, and H = 3, respectively. However, this pattern does not hold for $\rho = 0.9$, as we see it in the second row of Figure 5, and the results for H = 5 and H = 10 are quite competitive. The simulation results for $\rho = 0.75$ are not in favor of H = 10 as shown in the third row of Figure 5. In fact, the JPS estimator with H = 5 has the best performance for most values of p, which is followed by the estimator with H = 3. The three bottom panels of Figure 5 show the results for $\rho = 0.5$. It is clear from these panels that the REs are close or below one, and the JPS estimator with a smaller set size usually has a better performance in this case.

Simulation results for lifetime distributions and n = 300 are shown in Figure 6. From the provided figure, it can be observed that the REs of the estimators increase as the set size H becomes larger, and the JPS estimator with set size H = 10 has the best performance in most considered case, which is followed by the estimators with set size H = 5, and H = 3, respectively. It is also worth mentioning that the efficiency of the JPS estimators decrease as the ranking quality decreases. Furthermore, the RE curves are mostly higher than one, which indicate superiority of the JPS estimators to their SRS counterpart.

By overall examinations of Figures 1-6, we find out that the REs are increasing function of sample size n. The efficiency of the JPS estimators are usually higher around the center of the parent distribution and lower at its boundaries. Furthermore, the REs decrease with the ranking quality ρ . The optimal value of the set size H, which leads to a higher efficiency depends on both ranking quality (ρ) and sample size (n). Specifically, a larger value of set size H leads to a higher efficiency provided that the sample size is large enough and the ranking quality is sufficiently good. Therefore, small values for set sizes (H = 3, 5) are recommended to be used in practice if the sample size is small or there is any doubt about ranking quality.



Figure 1: The estimated relative efficiency of $F_{n;srs}^k(.)$ to $F_{n;jps}^k(.)$ as a function of p, for n = 10, when H = 3 (represented by red and solid line), H = 5 (represented by blue and dashed line), H = 10 (represented by black and dotted line) and $\rho \in \{1, 0.9, 0.75, 0.5\}$ under N(0, 1), t5 and La(0, 1) distributions.



Figure 2: The estimated relative efficiency of $F_{n;srs}^k(.)$ to $F_{n;jps}^k(.)$ as a function of p, for n = 50, when H = 3 (represented by red and solid line), H = 5 (represented by blue and dashed line), H = 10 (represented by black and dotted line) and $\rho \in \{1, 0.9, 0.75, 0.5\}$ under N(0, 1), t5 and La(0, 1) distributions.



Figure 3: The estimated relative efficiency of $F_{n;srs}^k(.)$ to $F_{n;jps}^k(.)$ as a function of p, for n = 300, when H = 3 (represented by red and solid line), H = 5 (represented by blue and dashed line), H = 10 (represented by black and dotted line) and $\rho \in \{1, 0.9, 0.75, 0.5\}$ under N(0, 1), t5 and La(0, 1) distributions.



Figure 4: The estimated relative efficiency of $F_{n;srs}^k(.)$ to $F_{n;jps}^k(.)$ as a function of p, for n = 10, when H = 3 (represented by red and solid line), H = 5 (represented by blue and dashed line), H = 10 (represented by black and dotted line) and $\rho \in \{1, 0.9, 0.75, 0.5\}$ under E(1), G(0.5, 1) and G(2, 1) distributions.



Figure 5: The estimated relative efficiency of $F_{n;srs}^k(.)$ to $F_{n;jps}^k(.)$ as a function of p, for n = 50, when H = 3 (represented by red and solid line), H = 5 (represented by blue and dashed line), H = 10 (represented by black and dotted line) and $\rho \in \{1, 0.9, 0.75, 0.5\}$ under E(1), G(0.5, 1) and G(2, 1) distributions.



Figure 6: The estimated relative efficiency of $F_{n;srs}^k(.)$ to $F_{n;jps}^k(.)$ as a function of p, for n = 300, when H = 3 (represented by red and solid line), H = 5 (represented by blue and dashed line), H = 10 (represented by black and dotted line) and $\rho \in \{1, 0.9, 0.75, 0.5\}$ under E(1), G(0.5, 1) and G(2, 1) distributions.

6 An Empirical Study: Body Fat Estimation

Our purpose in this section is to demonstrate the applicability and efficiency of the introduced procedure in practice using a real dataset. The dataset is utilized to evaluate the performance of the CDF estimators in JPS and SRS settings where the ranking process is carried out using actual auxiliary variables, rather than ones generated by a model.

It is very important for organizations in health sector such as Food and Drug Administration (FDA), or the National Cancer Institute (NCI) to have an accurate estimation of the distribution of body fat of people in a given population. This estimate is very useful for these organizations, since it can provide insights into the population's risk for some non-communicable illness such as diabetes, cardiovascular diseases, and some certain types of cancers, which usually impose a huge financial burden on the health care system of the society. However, the exact quantifying a person's body fat is a very challenging job and requires some advanced technologies. This is mainly because the body fat has not distributed evenly throughout the body. The exact measurement of a person's body fat is often obtained using some advanced imaging techniques such as Dual-Energy X-ray Absorptiometry (DEXA), Magnetic Resonance Imaging (MRI), or Computerized Tomography (CT) scans. Note that although these methods are very precise, they are very expensive as well, and need some special equipment and expertise which limit their applicability for large-scale population studies or even a standard clinical practice. The body fat percentage variable is correlated with some easily available concomitant variables such as abdomen circumference, chest circumference and weight.

The dataset utilized in this section is known as **bodyfat** dataset, and is accessible online at https://lib.stat.cmu.edu/datasets/bodyfat¹. This dataset contains body fat measurements determined using underwater weighting technique for 252 men along with measurements of some of their body circumferences. In what follows, we assume that this dataset represents our hypothetical population, and it is utilized to conduct a comparison of the performance exhibited by various CDF estimators. Assume that our objective is to estimate the true CDF of body fat among individuals, which is given by

$$F(t) = \frac{1}{252} \sum_{i=1}^{252} \mathbb{I}(X_i \le t),$$

where X_i is the body fat measurement of the *i*th subject in the hypothetical population.

¹Access date: 6 April 2024.

To compare the CDF estimators in the JPS and SRS settings, we set $n \in \{10, 20, 30, 40, 50\}$, $H \in \{3, 5, 10\}$, and for each combination of (n, H), we have drawn 100,000 samples from both of JPS and SRS designs, where all samplings are considered with replacement and therefore the independence assumption is not violated.

To obtain a JPS sample, we assume that ranking process is carried out using one of the concomitant variables, abdomen circumference, chest circumference, and weight. The correlation coefficient between the variable of interest (body fat) and the concomitant variables are $\rho = 0.81$, $\rho = 0.70$, and $\rho = 0.61$, respectively.

To include the perfect ranking case ($\rho = 1$) in our study, we also use the variable of interest for ranking itself. We have utilized epanechnikov function for the kernel function, and compute the RE of the JPS estimator to its SRS counterpart as it is defined in Section 5 for $p \in \{0.1, 0.25, 0.5, 0.75, 0.9\}$. The results are given in Table 1.

The first part of the Table 1 gives the results when the variable of the interest is ranked by itself, so the perfect ranking assumption holds. From the information presented in these rows, it is evident that one of the estimators derived from the JPS sampling outperforms the others and can be considered the superior estimator. except for the cases that $n \leq 20$ and p = 0.1, 0.9. The best estimator in each case depends on the sample size n, and the value of p. However, we observe that the performance of the JPS estimator improves as the value of p approaches to p = 0.5, or sample size n increases. For p = 0.1 or p = 0.9, the best JPS estimator is mostly observed for H = 3. But, for $p \in \{0.25, 0.5, 0.75\}$, the JPS estimator with either H = 5 or H = 10 has the best performance depending on the magnitude of the sample size n.

The other parts of the Table 1 show the results for imperfect ranking case. We observe that although the general patterns of REs remain almost the same, the REs decrease as the value of ρ decreases. Furthermore, the number of cases in which the JPS estimator cannot beat the SRS one increases as the value of ρ reduces. All of these observations align with the results presented in Section 5.

Concomitant		p = 0.1		p = 0.25			p = 0	p = 0.5			p = 0.75			p = 0.9		
Variable	n	H = 3	H = 5	H = 10	H = 3	H = 5	H = 10	H = 3	H = 5	H = 10	H = 3	H = 5	H = 10	H = 3	H = 5	H = 10
	10	0.89	0.89	0.94	1.09	1.11	1.12	1.62	1.71	1.48	1.00	1.00	1.01	0.90	0.90	0.93
Body Fat	20	0.98	0.92	0.91	1.32	1.35	1.32	1.84	2.34	2.21	1.09	1.05	1.03	0.96	0.89	0.88
$(\rho = 1)$	30	1.04	1.00	0.94	1.47	1.65	1.64	1.90	2.65	3.01	1.15	1.15	1.07	1.00	0.95	0.86
	40	1.07	1.07	0.98	1.55	1.89	1.99	1.93	2.75	3.63	1.19	1.22	1.14	1.02	0.98	0.88
	50	1.09	1.11	1.03	1.61	2.04	2.38	1.93	2.76	4.15	1.21	1.26	1.21	1.03	1.01	0.91
	10	0.85	0.84	0.90	0.97	0.97	1.00	1.28	1.32	1.24	0.95	0.95	0.97	0.88	0.88	0.91
Abdomen	20	0.94	0.86	0.84	1.13	1.08	1.05	1.39	1.52	1.46	1.02	0.96	0.95	0.94	0.86	0.85
Circumference	30	0.99	0.92	0.86	1.22	1.23	1.16	1.43	1.61	1.62	1.07	1.03	0.95	0.98	0.91	0.82
$(\rho = 0.81)$	40	1.02	0.98	0.87	1.26	1.35	1.27	1.43	1.65	1.74	1.09	1.09	1.00	1.00	0.95	0.84
	50	1.04	1.02	0.91	1.31	1.42	1.39	1.45	1.67	1.81	1.11	1.12	1.04	1.01	0.98	0.86
	10	0.83	0.82	0.88	0.90	0.91	0.94	1.13	1.15	1.12	0.92	0.92	0.95	0.87	0.87	0.91
Chest	20	0.92	0.82	0.81	1.03	0.95	0.93	1.21	1.23	1.21	0.98	0.92	0.90	0.92	0.85	0.83
Circumference	30	0.96	0.89	0.80	1.10	1.06	0.96	1.23	1.28	1.25	1.03	0.97	0.89	0.96	0.89	0.81
$(\rho = 0.7)$	40	0.99	0.95	0.82	1.14	1.14	1.03	1.24	1.32	1.29	1.05	1.02	0.92	0.98	0.93	0.81
	50	1.01	0.97	0.86	1.16	1.18	1.10	1.24	1.33	1.32	1.05	1.06	0.95	0.99	0.95	0.83
	10	0.82	0.82	0.87	0.89	0.88	0.93	1.08	1.09	1.08	0.91	0.90	0.94	0.87	0.86	0.90
Weight	20	0.90	0.81	0.80	0.99	0.91	0.90	1.14	1.14	1.12	0.97	0.90	0.88	0.92	0.84	0.82
$(\rho=0.61)$	30	0.96	0.88	0.79	1.07	1.02	0.93	1.17	1.19	1.16	1.01	0.95	0.87	0.96	0.88	0.80
	40	0.98	0.93	0.82	1.10	1.09	0.99	1.18	1.22	1.19	1.03	0.99	0.89	0.97	0.92	0.80
	50	0.99	0.96	0.84	1.12	1.14	1.04	1.19	1.25	1.22	1.04	1.02	0.93	0.98	0.94	0.83

Table 1: The estimated relative efficiency of $F_{n;srs}^k(.)$ to $F_{n;jps}^k(.)$ using the bodyfat dataset. In each scenario, the winner is indicated by bold font.

7 Discussion

Judgment post stratification (JPS) sampling plan is beneficial sampling plan for situations in which allocating a judgment rank to a sample unit in a set is far easier/cheaper than precisely quantifying it. It is often considered as a more flexible and practical variation of ranked set sampling (RSS). Despite the extensive utilization of this sampling approach in various fields, the literature lacks discussions on the issue of effectively smooth estimating the cumulative distribution function (CDF).

In this paper, we discussed a general class of the estimators for the population CDF in the JPS setting which includes both empirical and kernel-based ones. We showed that they are more efficient than their competitors in simple random sampling (SRS), as long as the ranking quality is better than random guessing. We found the necessary and sufficient condition for the estimators to be asymptotically unbiased. Also, we studied the Glivenko-Cantelli type convergence and asymptotic normality for the estimators in this class, assuming the same condition holds. We next focused on kernel distribution function (KDF) in the class and proposed optimal bandwidth. We conducted a comparative analysis between the performance of the KDF in JPS sampling and its competitor in SRS using Monte Carlo simulation. This analysis involved investigating various combinations of sample size, set size, ranking quality, parent distribution, and kernel function.

It was found that the JPS estimator outperforms its SRS competitor in the most considered cases. Finally, we showed the applicability and efficiency of our introduced procedure in practice using a real dataset in which real concomitant variables were used for the ranking process.

To the best of our knowledge, this work was the first attempt for estimating the CDF using kernel function based on JPS sampling scheme. So, There is still plenty of room for more research. As an example, consider a parameter of interest denoted by θ , which can be expressed as $\theta = g(F)$, where g(.) represents a known function. One can think of estimating the parameter θ by replacing the CDF F by an appropriate estimator from the discussed class, and establishing its statistical properties. Moreover, it was shown that the true CDF of the post strata often follow the constraint: $F_{[1]}(t) \ge \ldots \ge F_{[H]}(t)$, for all $t \in \mathbb{R}$. However, this constraint may not hold for sample estimates. Wang et al. (2012) improves the EDF in the JPS setting by imposing this limitation onto estimation process. Therefore, another interesting topic for future research can be how to utilize this limitation to improve the performance of the KDF in the JPS setting. These topics can be discussed in future works.

Data Availability Statement

The data supporting the findings of this study can be accessed online at https://lib.stat.cmu.edu/datasets/bodyfat (Access date: 6 April 2024).

References

- Amirhossein Alvandi and Armin Hatefi. Estimation of ordinal population with multi-observer ranked set samples using ties information. *Statistical Methods in Medical Research*, 30(8):1960– 1975, 2021.
- Amirhossein Alvandi and Armin Hatefi. Analysis of ordinal populations from judgment poststratification. *Stats*, 6(3):812–838, 2023.
- Adelchi Azzalini. A note on the estimation of a distribution function and quantiles by a kernel method. *Biometrika*, 68(1):326–328, 1981.
- Haiying Chen, Elizabeth A Stasny, and Douglas A Wolfe. Improved procedures for estimation of disease prevalence using ranked set sampling. *Biometrical Journal: Journal of Mathematical Methods in Biosciences*, 49(4):530–538, 2007.
- Wangxue Chen, Yi Tian, and Minyu Xie. The global minimum variance unbiased estimator of the parameter for a truncated parameter family under the optimal ranked set sampling. *Journal of Statistical Computation and Simulation*, 88(17):3399–3414, 2018.
- Wangxue Chen, Rui Yang, Dongsen Yao, and Chunxian Long. Pareto parameters estimation using moving extremes ranked set sampling. *Statistical Papers*, 62(3):1195–1211, 2021.
- Ali Dastbaravarde, Nasser Reza Arghami, and Majid Sarmad. Some theoretical results concerning non parametric estimation by using a judgment poststratification sample. *Communications in Statistics-Theory and Methods*, 45(8):2181–2203, 2016.
- TR Dell and JL Clutter. Ranked set sampling theory with order statistics background. *Biometrics*, pages 545–555, 1972.
- Mozhgan Alirezaei Dizicheh, Ehsan Zamanzade, and Nasrollah Iranpanah. Efficient estimation of the odds using judgment post stratification. *Brazilian Journal of Probability and Statistics*, 35 (2):375–391, 2021.
- Lutz Dümbgen and Ehsan Zamanzade. Inference on a distribution function from ranked set samples. Annals of the Institute of Statistical Mathematics, 72(1):157–185, 2020.

- Abbas Eftekharian and Mostafa Razmkhah. On estimating the distribution function and odds using ranked set sampling. *Statistics & Probability Letters*, 122:1–10, 2017.
- Jesse Frey. Bootstrap confidence bands for the cdf using ranked-set sampling. Journal of the Korean Statistical Society, 43(3):453–461, 2014.
- Jesse Frey and Timothy G Feeman. An improved mean estimator for judgment post-stratification. Computational Statistics & Data Analysis, 56(2):418–426, 2012.
- Jesse Frey and Timothy G Feeman. Variance estimation using judgment post-stratification. Annals of the Institute of Statistical Mathematics, 65:551–569, 2013.
- Jesse Frey and Yimin Zhang. Improved exact confidence intervals for a proportion using ranked-set sampling. *Journal of the Korean Statistical Society*, 48:493–501, 2019a.
- Jesse Frey and Yimin Zhang. An omnibus two-sample test for ranked-set sampling data. *Journal* of the Korean Statistical Society, 48:106–116, 2019b.
- Jesse Frey and Yimin Zhang. Robust confidence intervals for a proportion using ranked-set sampling. Journal of the Korean Statistical Society, 50:1009–1028, 2021.
- Sneh Gulati. Smooth non-parametric estimation of the distribution function from balanced ranked set samples. *Environmetrics*, 15(5):529–539, 2004.
- Lowell K Halls and Tommy R Dell. Trial of ranked-set sampling for forage yields. *Forest Science*, 12(1):22–26, 1966.
- Xiaofang He, Wangxue Chen, and Wenshu Qian. Maximum likelihood estimators of the parameters of the log-logistic distribution. *Statistical papers*, 61(5):1875–1892, 2020.
- Xiaofang He, Wangxue Chen, and Rui Yang. Modified best linear unbiased estimator of the shape parameter of log-logistic distribution. *Journal of Statistical Computation and Simulation*, 91(2): 383–395, 2021.
- Ralph W Howard, Susan C Jones, Joe K Mauldin, and Raymond H Beal. Abundance, distribution, and colony size estimates for reticulitermes spp.(isoptera: Rhinotermitidae) in southern mississippi. *Environmental Entomology*, 11(6):1290–1293, 1982.

- Paul H Kvam. Ranked set sampling based on binary water quality data with covariates. *Journal of Agricultural, Biological, and Environmental Statistics*, 8:271–279, 2003.
- Steven N MacEachern, Ömer Öztürk, Douglas A Wolfe, and Gregory V Stark. A new ranked set sample estimator of variance. Journal of the Royal Statistical Society: Series B (Statistical Methodology), 64(2):177–188, 2002.
- Steven N MacEachern, Elizabeth A Stasny, and Douglas A Wolfe. Judgement post-stratification with imprecise rankings. *Biometrics*, 60(1):207–215, 2004.
- M Mahdizadeh and Ehsan Zamanzade. Reliability estimation in multistage ranked set sampling. *REVSTAT-Statistical Journal*, 15(4):565–581, 2017.
- M Mahdizadeh and Ehsan Zamanzade. Smooth estimation of the area under the roc curve in multistage ranked set sampling. *Statistical Papers*, 62(4):1753–1776, 2021.
- Mahdi Mahdizadeh and Ehsan Zamanzade. Interval estimation of p (x; y) in ranked set sampling. Computational Statistics, 33(3):1325–1348, 2018.
- GA McIntyre. A method for unbiased selective sampling, using ranked sets. Australian journal of agricultural research, 3(4):385–390, 1952.
- Chul Moon, Xinlei Wang, and Johan Lim. Empirical likelihood inference for area under the roc curve using ranked set samples. *arXiv preprint arXiv:2010.12185*, 2020.
- Elizbar A Nadaraya. Some new estimates for distribution functions. Theory of Probability & Its Applications, 9(3):497–500, 1964.
- Barry D Nussbaum and BK Sinha. Cost effective gasoline sampling using ranked set sampling. In Proceedings of the Section on Statistics and the Environment, pages 83–87. American Statistical Association, 1997.
- Sedigheh Omidvar, Mohammad Jafari Jozani, and Nader Nematollahi. Judgment poststratification in finite mixture modeling: An example in estimating the prevalence of osteoporosis. *Statistics in medicine*, 37(30):4823–4836, 2018.
- Omer Ozturk. Combining ranking information in judgment post stratified and ranked set sampling designs. *Environmental and ecological statistics*, 19(1):73–93, 2012.

- Omer Ozturk. Statistical inference for population quantiles and variance in judgment post-stratified samples. *Quality control and applied statistics*, 60(3):217–218, 2015.
- Omer Ozturk and Olena Kravchuk. Judgment post-stratified assessment combining ranking information from multiple sources, with a field phenotyping example. *Journal of Agricultural*, *Biological and Environmental Statistics*, 26(3):329–348, 2021.
- Omer Ozturk, Omer C Bilgin, and Douglas A Wolfe. Estimation of population mean and variance in flock management: a ranked set sampling approach in a finite population setting. *Journal of Statistical Computation and Simulation*, 75(11):905–919, 2005.
- Brett Presnell and Lora L Bohn. U-statistics and imperfect ranking in ranked set sampling. *Journal* of Nonparametric Statistics, 10(2):111–126, 1999.
- Wenshu Qian, Wangxue Chen, and Xiaofang He. Parameter estimation for the pareto distribution based on ranked set sampling. *Statistical papers*, 62(1):395–417, 2021.
- Hani M Samawi and Omar AM Al-Sagheer. On the estimation of the distribution function using extreme and median ranked set sampling. *Biometrical Journal: Journal of Mathematical Methods* in *Biosciences*, 43(3):357–373, 2001.
- Bernard W Silverman. Density estimation for statistics and data analysis. Routledge, 2018.
- S Lynne Stokes. Estimation of variance using judgment ordered ranked set samples. *Biometrics*, pages 35–42, 1980.
- S Lynne Stokes and Thomas W Sager. Characterization of a ranked-set sample with application to estimating distribution functions. *Journal of the American Statistical Association*, 83(402): 374–381, 1988.
- Koiti Takahasi and Kazumasa Wakimoto. On unbiased estimates of the population mean based on the sample stratified by means of ordering. Annals of the institute of statistical mathematics, 20(1):1–31, 1968.
- Xinlei Wang, Johan Lim, and Lynne Stokes. A nonparametric mean estimator for judgment poststratified data. *Biometrics*, 64(2):355–363, 2008.
- Xinlei Wang, Ke Wang, and Johan Lim. Isotonized cdf estimation from judgment poststratification data with empty strata. *Biometrics*, 68(1):194–202, 2012.

- Xinlei Wang, Johan Lim, and Lynne Stokes. Using ranked set sampling with cluster randomized designs for improved inference on treatment effects. *Journal of the American Statistical* Association, 111(516):1576–1590, 2016.
- Geoffrey S Watson and MR Leadbetter. Hazard analysis ii. Sankhyā: The Indian Journal of Statistics, Series A, pages 101–116, 1964.
- BB Winter. Strong uniform consistency of integrals of density estimators. *Canadian Journal of Statistics*, 1(1-2):247–253, 1973.
- Hajime Yamato. Uniform convergence of an estimator of a distribution function. 1973.
- Ehsan Zamanzade. An isotonized variance estimator for judgment post stratified data. *Journal of the Korean Statistical Society*, 45(1):25–32, 2016.
- Ehsan Zamanzade and Michael Vock. Some nonparametric tests of perfect judgment ranking for judgment post stratification. *Statistical Papers*, 59(3):1085–1100, 2018.
- Ehsan Zamanzade and Xinlei Wang. Estimation of population proportion for judgment poststratification. *Computational Statistics & Data Analysis*, 112:257–269, 2017.
- Ehsan Zamanzade, M Mahdizadeh, and Hani M Samawi. Efficient estimation of cumulative distribution function using moving extreme ranked set sampling with application to reliability. AStA Advances in Statistical Analysis, 104(3):485–502, 2020.