Thomas A. Courtade[∗] Max Fathi[†]

May 2nd 2024

Abstract

We resolve a question of Carrapatoso et al. [\[7\]](#page-11-0) on Gaussian optimality for the sharp constant in Poincaré-Korn inequalities, under a moment constraint. We also prove stability, showing that measures with near-optimal constant are quantitatively close to standard Gaussian.

1 Introduction and Main Result

Let μ be a centered Borel probability measure on \mathbb{R}^n , $n \geq 2$. Let A denote the set of antisymmetric linear maps from \mathbb{R}^n to itself. That is,

$$
\mathcal{A} := \{x \mapsto Ax \; ; A \in M_{n \times n}(\mathbb{R}), A = -A^T\}.
$$

Further define the linear space of vector-valued functions

$$
\mathcal{C}:=\{u:\mathbb{R}^n\to\mathbb{R}^n\ ;\ u\ \text{differentiable},\ \int ud\mu=0,\ \text{and}\ \|\nabla_s u\|_{L^2(\mu)}^2<\infty\},
$$

where $\nabla_s u := \frac{1}{2}((\nabla u) - (\nabla u)^T)$ is the symmetrized gradient of the vector-valued function $u : \mathbb{R}^n \to \mathbb{R}^n$. Since μ is centered^{[1](#page-0-0)}, A is a closed linear subspace of C.

Definition 1. A centered Borel probability measure μ on \mathbb{R}^n satisfies a **Poincaré–Korn in**equality with constant C if

$$
\inf_{a \in \mathcal{A}} \|u - a\|_{L^2(\mu)}^2 \le 2C \|\nabla_s u\|_{L^2(\mu)}^2, \text{ for all } u \in \mathcal{C}.
$$
 (1)

The Poincaré–Korn constant associated to μ , denoted $C_{PK}(\mu)$, is the smallest constant C such that (1) holds.

This type of inequality was introduced in $[7]$. It is inspired by Poincaré inequalities, which control variances of scalar valued functions by the L^2 norm of their gradient, and the Korn inequality from continuous mechanics, which controls the L^2 norm of the gradient of a vector field satisfying some boundary condition by its symmetric part. Both inequalities have found many applications in analysis. They were both originally introduced for uniform measures on domains, but can be extended to general probability densities. In this form, Poincaré inequalities are

$$
\int f^2 d\mu - \left(\int f d\mu\right)^2 \le C_P(\mu) \int |\nabla f|^2 d\mu, \quad \forall f : \mathbb{R}^d \longrightarrow \mathbb{R}^d,
$$
\n(2)

[∗]University of California, Berkeley, United States, Department of Electrical Engineering and Computer Sciences. courtade@berkeley.edu

[†]Université Paris Cité and Sorbonne Université, CNRS, Laboratoire Jacques-Louis Lions and Laboratoire de Probabilités, Statistique et Modélisation, F-75013 Paris, France; and DMA, École normale supérieure, Université PSL, CNRS, 75005 Paris, France; and Institut Universitaire de France. mfathi@lpsm.paris

¹The persistent centering assumption comes without any loss of generality, and is only made for convenience.

where $C_P(\mu)$ is the Poincaré constant of μ , and the right-hand side is to be understood as $+\infty$ if it is not well-defined for the function f . Korn inequalities are of the form

$$
\inf_{a \in \mathcal{A}} \|\nabla(u-a)\|_{L^2(\mu)}^2 \le C_K \|\nabla_s u\|_{L^2(\mu)}^2, \text{ for all } u \in \mathcal{C}.
$$
 (3)

Poincaré inequalities are now a very classical tool in probability and functional analysis, and applications include concentration of measure inequalities and rates of convergence to equilibrium for Markov processes. We refer to the monograph [\[2\]](#page-11-1) for background and many developments. On the other hand, classical Korn inequalities are a tool in kinetic theory and fluid mechanics, going back to [\[19\]](#page-11-2). We shall make no attempt to survey the vast literature, and refer to [\[18\]](#page-11-3) for some background. Best constants were investigated for example in [\[20\]](#page-11-4). Weighted Korn inequalities were recently introduced in [\[7\]](#page-11-0), motivated by hypocoercivity problems in kinetic theory.

Our definition of the Poincaré–Korn constant differs from that in $[7]$ by a factor of 2. This is done to give unit normalization with respect to the standard Gaussian measure γ , defined by

$$
d\gamma(x) := \frac{1}{(2\pi)^{n/2}} e^{-|x|^2/2} dx, \quad x \in \mathbb{R}^n.
$$

Probability measures with sufficiently regular potentials admit a finite Poincaré–Korn constant if they satisfy a Poincaré inequality with finite constant.

Proposition 1. [\[7,](#page-11-0) Theorem 1] We have $C_{PK}(\gamma) = 1$ in any dimension $n \geq 2$. Moreover, if a centered probability measure with density $d\mu = e^{-\phi} dx$ of class C^2 satisfies

 $\forall \epsilon > 0, \exists C_{\epsilon} > 0: \quad \|\nabla^2 \phi(x)\|^2 \leq \epsilon |\nabla \phi(x)|^2 + C_{\epsilon}, \quad \forall x \in \mathbb{R}^n,$

then $C_P(\mu) < \infty \Rightarrow C_{PK}(\mu) < \infty$.

In [\[7\]](#page-11-0), the following conjecture is proposed regarding the rigidity of the Poincaré–Korn inequality:

Conjecture 1. If $d\mu = e^{-\phi}dx$ is centered and isotropic (i.e., $\int x d\mu = 0$ and $\int x x^T d\mu = Id$), and satisfies $\nabla^2 \phi \geq \text{Id}$, then $C_{PK}(\mu) \geq C_{PK}(\gamma)$, with equality only if $\mu = \gamma$.

It turns out that the conjectured statement is indeed true, but the hypotheses are too strong to capture salient rigidity properties of the Poincaré–Korn constant. Namely, the following can be derived as a consequence of known stability results for the Bakry–Émery theorem on \mathbb{R}^n [\[10\]](#page-11-5), or using Caffarelli's contraction theorem (see Appendix A). We make no claim of originality for this statement, which was known in some communities.

Proposition 2. If $d\mu = e^{-\phi}dx$ is centered, isotropic, and satisfies $\nabla^2 \phi \geq \text{Id}$, then $\mu = \gamma$.

Therefore, to study rigidity of the Poincaré–Korn inequality, the assumption of a uniformly convex potential in Conjecture [1](#page-1-0) should be replaced by something else. A natural choice is a moment assumption, which we now define.

Definition 2 (Moment Assumption). We say that μ satisfies the moment assumption if, for all $1 \leq i, j, k \leq n$,

$$
\int x_i d\mu = \int x_i d\gamma = 0;
$$

$$
\int x_i x_j d\mu = \int x_i x_j d\gamma = \delta_{ij};
$$

$$
\int x_i x_j x_k d\mu = \int x_i x_j x_k d\gamma = 0;
$$

and, when $i \neq j$,

$$
\int (x_i^2 + x_j^2)x_j^2 d\mu = \int (x_i^2 + x_j^2)x_j^2 d\gamma = 4.
$$

Remark 1. The first two lines of the moment assumption correspond to μ being centered and isotropic. The third and fourth lines in the moment assumption ensure that μ and γ share mixed third moments and select mixed fourth moments, respectively.

There are many interesting probability measures that satisfy the moment assumption. For example, any product measure whose individual factors share moments up to order 4 with the standard normal will satisfy the moment assumption (and so will mixtures of these measures, and so forth...). Thus, a nontrivial reformulation of Conjecture [1](#page-1-0) is as follows:

Question 1. If μ satisfies the moment assumption, is the lower bound $C_{PK}(\mu) \geq C_{PK}(\gamma)$ true, with equality only if $\mu = \gamma$?

The moment assumption is motivated by the form of extremal functions in the Gaussian Poincaré-Korn inequality. In particular, by considering these as test functions in the Poincaré– Korn inequality for μ , the inequality $C_{PK}(\mu) \geq C_{PK}(\gamma)$ is a consequence of the moment as-sumption (see Proposition [3](#page-5-0) in the sequel). So, it is the rigidity phenomenon that is interesting. We remark that the work of Serres [\[22\]](#page-11-6) already highlights that given a reference measure satisfying a Poincaré inequality with known sharp constant and extremal function, it is possible to study stability of functional inequalities within classes of measures for which the moments of the extremal function match with those under the reference measure.

In this note, we resolve Question [1](#page-2-0) in the affirmative, and further establish quantitative stability of the Poincaré–Korn constant. Such stability results on functional inequalities have been the subject of some recent attention in analysis. For example, there have been many results on stability for sharp functions in classical functional inequalities, including Sobolev inequalities and isoperimetric inequalities, see [\[16,](#page-11-7) [17\]](#page-11-8) for surveys. Stability results under moment constraints have been studied for Poincaré inequalities $[24, 5, 12, 22]$ $[24, 5, 12, 22]$ $[24, 5, 12, 22]$ $[24, 5, 12, 22]$, eigenvalues of diffusion operators $[23]$, as well as fractional [\[1\]](#page-11-11) and free [\[9\]](#page-11-12) Poincaré inequalities. These have been obtained using the combination of Stein's method and variational arguments that we shall use here. In another direction, there have been stability results for sharp constants under convexity or curvature assumptions [\[8,](#page-11-13) [11,](#page-11-14) [15,](#page-11-15) [21\]](#page-11-16).

Our stability result is with respect to the Zolotarev distance of order 2, which controls the same topology as the more familiar W_2 Kantorovich–Wasserstein distance [\[3\]](#page-11-17).

Definition 3. For two Borel probability measures μ, ν on \mathbb{R}^n , the Zolotarev distance of order 2 is defined as

$$
d_{Zol,2}(\mu,\nu) := \sup_{\sup_x \|\nabla^2 f(x)\|_2 \le 1} \int f d\mu - \int f d\nu,
$$

where the supremum is over all $f : \mathbb{R}^n \to \mathbb{R}$ in C^2 with $\sup_{x \in \mathbb{R}^n} ||\nabla^2 f(x)||_2 \leq 1$.

Our main result is as follows.

Theorem 1 (Stability). If μ satisfies the moment assumption, then $C_{PK}(\mu) \geq C_{PK}(\gamma) = 1$, and

$$
d_{Zol,2}(\mu,\gamma) \le c n^2 \sqrt{C_{PK}(\mu)(C_{PK}(\mu) - C_{PK}(\gamma))},
$$

content

where c is a universal constant.

The following is now immediate, and affirmatively answers Question [1.](#page-2-0)

Corollary 1 (Rigidity). If μ satisfies the moment assumption, then $C_{PK}(\mu) \geq C_{PK}(\gamma)$, with equality only if $\mu = \gamma$.

Acknowledgments. We thank Jean Dolbeault for telling us about this problem. TC acknowledges NSF-CCF 1750430, the hospitality of the Laboratoire de Probabilités, Statistique et Modélisation (LPSM) at the Université Paris Cité, and the Invited Professor program of the Fondation Sciences Mathématiques de Paris (FSMP). MF was supported by the Agence Nationale de la Recherche (ANR) Grant ANR-23-CE40-0003 (Project CONVIVIALITY). This work has also received support under the program "Investissement d'Avenir" launched by the French Government and implemented by ANR, with the reference ANR-18-IdEx-0001 as part of its program Emergence.

2 Proof of Main Result

2.1 Notation

For a vector-valued function $u : \mathbb{R}^n \to \mathbb{R}^n$, we define

$$
||u||^2_{L^2(\mu)} := \int |u|^2 d\mu,
$$

where |x| denotes the Euclidean length of $x \in \mathbb{R}^n$. Likewise, for a matrix-valued function $U: \mathbb{R}^n \to M_{n \times n}(\mathbb{R})$, we define

$$
||U||_{L^2(\mu)}^2 := \int U \cdot U d\mu,
$$

where \cdot' will denote the scalar product between matrices (i.e., $A \cdot B := \sum_{ij} [A]_{ij} [B]_{ij}$, for identically dimensioned matrices $A, B \in M_{m \times n}(\mathbb{R})$. The identity matrix is denoted by Id. All vectors are represented in matrix form as column vectors.

In what follows, for $1 \leq i, j \leq n$, we let δ_{ij} denote the usual Kronecker delta function:

$$
\delta_{ij} := \begin{cases} 1 & \text{if } i = j \\ 0 & \text{otherwise.} \end{cases}
$$

2.2 Remarks on the general approach

In [\[11\]](#page-11-14), the authors introduced a general approach for establishing stability of functional inequalities based on approximate integration-by-parts identities and Stein's method; the ideas can also be found in the one-dimensional results of [\[24,](#page-12-0) [5\]](#page-11-9). Interested readers are referred to [\[11\]](#page-11-14) for an overview of the method in abstract settings. We follow this general approach here. Namely, the first step of the proof (Subsection [2.3\)](#page-3-0) is dedicated to proving an approximate Stein identity, which mimics the integration by parts formula for the Gaussian measure on Hermite polynomials of degree 2. The second step (Subsection [2.4\)](#page-8-0) describes our implementation of Stein's method that yields the main stability result.

2.3 An approximate Stein identity

For a differentiable matrix-valued function $V: \mathbb{R}^n \to M_{n \times n}(\mathbb{R})$ and $x \in \mathbb{R}^n$, we write $(x\nabla^T) \cdot V$ to denote the scalar product of the operator $(x\nabla^T)$ and the function V; that is,

$$
(x\nabla^T)\cdot V = \sum_{i,j=1}^n [x\nabla^T]_{ij}[V]_{ij} \cdot := \sum_{i,j=1}^n x_i \partial_j [V]_{ij}.
$$

The following lemma is the main result of this section.

Lemma 1 (Approximate Stein Identity). Let μ satisfy the moment assumption. If $V : \mathbb{R}^n \to$ $M_{n\times n}(\mathbb{R})$ is a matrix-valued function such that each coordinate $[V]_{ij} : \mathbb{R}^n \to \mathbb{R}$ is integrable and differentiable, with $\|\nabla[V]_{ij}\|_{L^2(\mu)} < \infty$, then

$$
\left| \int (xx^T - \mathrm{Id}) \cdot V d\mu - \int (x \nabla^T) \cdot V d\mu \right| \leq K \sqrt{C_{PK}(C_{PK} - 1)} \sum_{i,j=1}^n \|\nabla [V]_{ij}\|_{L^2(\mu)},
$$

for $K := (1 + 10/\sqrt{3}) < 7$.

To provide some perspective, we remark that the classical Stein identity

$$
\int x \cdot \phi d\gamma = \int \mathrm{Id} \cdot \nabla \phi d\gamma, \quad \phi : \mathbb{R}^n \to \mathbb{R}^n
$$

applied to the test function $\phi(x) = V^T(x)x$ gives

$$
\int (xx^T - \mathrm{Id}) \cdot V d\gamma = \int (x \nabla^T) \cdot V d\gamma.
$$

Under certain moment assumptions, this consequence of the classical Stein identity also characterizes the Gaussian measure, so may therefore be regarded as one of many "Stein identities". It is for this reason that we refer to Lemma [1](#page-3-1) as an "approximate Stein identity", which becomes more faithful as $C_{PK}(\mu)$ approaches 1.

The proof of the approximate Stein identity rests on an approximate integration by parts formula enjoyed by near-extremizers of the Poincaré–Korn inequality. To develop it, we assume henceforth that μ has finite second moments. For a function $u \in \mathcal{C}$, define

$$
a_u := \arg\min_{a \in \mathcal{A}} \|u - a\|_{L^2(\mu)}.
$$

Since A is a closed linear subspace of $L^2(\mu)$ and $\mathcal{C} \subset L^2(\mu)$, the function a_u exists and is unique. Moreover, as a projection onto A, the map $u \mapsto a_u$ is linear and equal to identity on A.

The following is an approximate integration by parts formula satisfied by near-extremizers of the Poincar´e–Korn inequality; it does not require the moment assumption, and may therefore be of independent interest.

Lemma 2. Let μ be a centered Borel probability measure on \mathbb{R}^n with finite second moments, and $C_{PK}(\mu) < \infty$. If $\epsilon \geq 0$ and $u \in \mathcal{C}$ satisfy

$$
(2 - (\epsilon/2)^2)C_{PK}(\mu) \|\nabla_s u\|_{L^2(\mu)}^2 \le \inf_{a \in \mathcal{A}} \|u - a\|_{L^2(\mu)}^2,
$$
\n(4)

then for every $v \in \mathcal{C}$, we have

$$
\left| \int (u - a_u) \cdot (v - a_v) d\mu - 2C_{PK}(\mu) \int (\nabla_s u) \cdot (\nabla_s v) d\mu \right| \le \epsilon C_{PK}(\mu) \|\nabla_s u\|_{L^2(\mu)} \|\nabla_s v\|_{L^2(\mu)}.
$$

We briefly remark that if $u \in \mathcal{C}$ is an extremizer in the Poincaré–Korn inequality, then we have the following (exact) integration by parts formula:

$$
\int (u - a_u) \cdot (v - a_v) d\mu = 2C_{PK}(\mu) \int (\nabla_s u) \cdot (\nabla_s v) d\mu, \quad \forall v \in \mathcal{C}.
$$

However, we see no reason to expect that nontrivial extremizers exist in the Poincaré–Korn inequality for general μ , which motivates the approximation in Lemma [2.](#page-4-0) Note that there are examples of measures for which the classical Poincaré inequality has no non-trivial extremal function, including for example the exponential measure, see [\[2,](#page-11-1) Section 4.4.1].

Proof. We'll abbreviate $C_{PK} := C_{PK}(\mu)$ for convenience. Begin by defining the quotient space $\mathcal{Q} := \mathcal{C}/\mathcal{A}$. Note that $\mathcal{A} \subset \text{ker}(\nabla_s)$, so we may define a linear operator D on Q via

$$
D[u] := \nabla_s u, \quad u \in \mathcal{C},
$$

where $[u] \in \mathcal{Q}$ denotes the coset of u. Observe that

$$
\langle [u],[v]\rangle := \int D[u] \cdot D[v] d\mu, \quad [u],[v] \in \mathcal{Q},
$$

defines an inner product on $\mathcal{Q} \times \mathcal{Q}$. Bilinearity and symmetry are self-evident, and positivedefiniteness follows from the Poincaré–Korn inequality, which states

$$
\| [u] \|^2 := \langle [u], [u] \rangle \ge \frac{1}{2C_{PK}} \inf_{a \in \mathcal{A}} \| u - a \|^2_{L^2(\mu)}.
$$

Since the quantity on the right is the quotient norm, positive-definiteness follows. Hence, we are justified in defining a Hilbert space $\mathcal H$ as the completion of $\mathcal Q$ in the norm $\|\cdot\|$, and extending the inner product $\langle \cdot, \cdot \rangle$ to H.

Now, define the operator $T : q \in \mathcal{Q} \mapsto T q$ by

$$
T[u] := u - a_u, \quad u \in \mathcal{C}.
$$

The operator T is linear, and well-defined on $\mathcal Q$ since $u = a_u$ for $u \in \mathcal A$. Now, fix $u \in \mathcal C$. The operator

$$
[v] \in \mathcal{Q} \mapsto \int T[u] \cdot T[v] d\mu
$$

is a bounded linear operator on \mathcal{Q} . Indeed, boundedness follows by the Cauchy–Schwarz and Poincaré–Korn inequalities as

$$
\int T[u] \cdot T[v] d\mu \le 2C_{PK} ||[u] ||[v]||, \text{ for all } v \in \mathcal{C}.
$$

Linearity now follows by linearity of T and the integral. By density of Q in H and the Riesz representation theorem, there is $h \in \mathcal{H}$ with $||h|| \leq ||u||$ such that

$$
\int T[u] \cdot T[v] d\mu = 2C_{PK} \langle h, [v] \rangle, \text{ for all } v \in \mathcal{C}.
$$

Hence, for any $v \in \mathcal{C}$, we have

$$
\int T[u] \cdot T[v] d\mu - 2C_{PK}(\mu) \langle [u], [v] \rangle = 2C_{PK}(\mu) \langle h - [u], [v] \rangle
$$

$$
\leq 2C_{PK}(\mu) ||h - [u]|| ||[v]||.
$$

Opening the square, we have

$$
C_{PK} ||h - [u]||^2 = C_{PK} ||h||^2 - 2C_{PK} \langle h, [u] \rangle + C_{PK} ||[u]||^2
$$

\n
$$
\leq 2C_{PK} ||[u]||^2 - \int T[u] \cdot T[u] d\mu
$$

\n
$$
= 2C_{PK} ||\nabla_s u||^2_{L^2(\mu)} - \inf_{a \in \mathcal{A}} ||u - a||^2_{L^2(\mu)}.
$$

The claim follows.

Proposition 3. Let μ satisfy the moment assumption.

i) For each $u \in \mathcal{C}$, we have $a_u(x) = A_u x$, with

$$
A_u := \frac{1}{2} \int \left(u x^T - x u^T \right) d\mu.
$$

ii) Fix $i, j \in \{1, ..., n\}$ with $i \neq j$. The function $u = (u_1, ..., u_n) \in \mathcal{C}$ defined by

$$
u_k(x) = \delta_{ik}(1 - x_j^2) + \delta_{jk}x_ix_j, \quad 1 \le k \le n
$$
\n⁽⁵⁾

 \Box

satisfies $a_u = 0$ and

$$
[\nabla_s u]_{k\ell} = (\delta_{j\ell}\delta_{jk})x_i - (\delta_{ik}\delta_{j\ell} + \delta_{i\ell}\delta_{jk})\frac{x_j}{2}, \quad 1 \le k, \ell \le n.
$$

iii) We have $C_{PK}(\mu) \geq 1$.

Proof. i) Let $A = -A^T$. By the cyclic property of trace and the isotropic condition in the moment assumption, we may compute

$$
\int (A_u x) \cdot (Ax) d\mu = \text{Tr}(A_u^T A) = \frac{1}{2} \int \text{Tr} (x u^T A - u x^T A) d\mu = \int u \cdot (Ax) d\mu,
$$

where we used antisymmetry of A in the last step. It follows that

$$
\int (u - A_u x) \cdot ad\mu = 0, \ \ \forall a \in \mathcal{A}.
$$

An application of the Hilbert projection theorem proves i).

ii) Fix $i, j \in \{1, \ldots, n\}$ with $i \neq j$. For u given by [\(5\)](#page-5-1), we use i) to evaluate

$$
2[A_u]_{k\ell} = \int (u_k x_\ell - x_k u_\ell) d\mu
$$

=
$$
\int (\delta_{ik} (1 - x_j^2) x_\ell + \delta_{jk} x_i x_j x_\ell - \delta_{i\ell} (1 - x_j^2) x_k - \delta_{j\ell} x_i x_j x_k) d\mu,
$$

which vanishes by the moment assumption^{[2](#page-6-0)}. Next, let ∂_k denote partial derivative with respect to x_k , and observe

$$
2[\nabla_s u]_{k\ell} = \partial_k u_\ell + \partial_\ell u_k
$$

= $\delta_{i\ell} \partial_k (1 - x_j^2) + \delta_{j\ell} \partial_k x_i x_j + \delta_{ik} \partial_\ell (1 - x_j^2) + \delta_{jk} \partial_\ell x_i x_j$
= $\delta_{i\ell} \delta_{jk} (-2x_j) + \delta_{j\ell} (\delta_{ik} x_j + \delta_{jk} x_i) + \delta_{ik} \delta_{j\ell} (-2x_j) + \delta_{jk} (\delta_{i\ell} x_j + \delta_{j\ell} x_i)$
= $2\delta_{j\ell} \delta_{jk} x_i - (\delta_{ik} \delta_{j\ell} + \delta_{i\ell} \delta_{jk}) x_j$.

iii) For the choice of u given by (5) , we use ii) and the moment assumption to evaluate

$$
\inf_{a \in \mathcal{A}} \|u - a\|_{L^2(\mu)}^2 = \|u\|_{L^2(\mu)}^2 = \int x_i^2 x_j^2 d\mu + \int (1 - x_j^2)^2 d\mu = 3
$$

and

$$
2\|\nabla_s u\|_{L^2(\mu)}^2 = 2\int x_i^2 d\mu + 4\int (x_j/2)^2 d\mu = 3.
$$

It now follows by definitions that $C_{PK}(\mu) \geq 1$.

With the necessary ingredients established, we turn our attention to the proof of Lemma [1.](#page-3-1)

Proof of Lemma [1.](#page-3-1) Abbreviate $C_{PK} := C_{PK}(\mu)$. We can assume $C_{PK} < \infty$, else the claim is trivial. Also, the statement is invariant to adding constants to V , so we assume without loss of generality that $\int V d\mu = 0$.

To start, fix $i \neq j$, and let u be given by [\(5\)](#page-5-1). By the moment assumption, this choice of u satisfies [\(4\)](#page-4-1) with

$$
\epsilon = 2\sqrt{2}\sqrt{1 - \frac{1}{C_{PK}}}.\tag{6}
$$

Next, let $\psi : \mathbb{R}^n \to \mathbb{R}$ be integrable and differentiable, satisfying $\int \psi d\mu = 0$ and $\|\nabla \psi\|_{L^2(\mu)} <$ ∞ . Fix $m \in \{1, ..., n\}$ and define $v := (\delta_{1m}, ..., \delta_{nm})\psi$. Since $\int \psi d\mu = 0$ and $\|\nabla_s v\|_{L^2(\mu)} \le$

²Some simple casework shows that, under the assumption that μ is centered and isotropic, $a_u = 0$ for every choice of i, j if and only if all (mixed) third moments vanish.

 $\|\nabla \psi\|_{L^2(\mu)} < \infty$, it follows that $v \in \mathcal{C}$. For A_v defined as in Proposition [3,](#page-5-0) we may compute

$$
\int (u - a_u) \cdot (v - a_v) d\mu = \int u \cdot (v - a_v) d\mu
$$

=
$$
\int u_m \psi d\mu - \sum_{k,\ell} \int u_k [A_v]_{k\ell} x_{\ell} d\mu
$$

=
$$
\int (\delta_{im} (1 - x_j^2) \psi + \delta_{jm} x_i x_m \psi) d\mu - \sum_{\ell} \int ([A_v]_{i\ell} (1 - x_j^2) + [A_v]_{j\ell} x_i x_j x_{\ell}) d\mu
$$

=
$$
\int (\delta_{im} (1 - x_j^2) \psi + \delta_{jm} x_i x_m \psi) d\mu,
$$

where the last line follows from the moment assumption. Next, note that

$$
\int (\nabla_s u) \cdot (\nabla_s v) d\mu = \int (x_i [\nabla_s v]_{jj} - x_j [\nabla_s v]_{ij}) d\mu
$$

=
$$
\int \left(x_i \delta_{jm} \partial_j \psi - \frac{1}{2} x_j [\delta_{jm} \partial_i \psi + \delta_{im} \partial_j \psi] \right) d\mu.
$$

Define $\mathcal{E}(\psi) := 2\sqrt{3}\sqrt{C_{PK}(C_{PK}-1)}\|\nabla\psi\|_{L^2(\mu)}$ $\mathcal{E}(\psi) := 2\sqrt{3}\sqrt{C_{PK}(C_{PK}-1)}\|\nabla\psi\|_{L^2(\mu)}$ $\mathcal{E}(\psi) := 2\sqrt{3}\sqrt{C_{PK}(C_{PK}-1)}\|\nabla\psi\|_{L^2(\mu)}$ for convenience. An application of Lemma 2 with ϵ given in [\(6\)](#page-6-1) yields

$$
\left| \int (\delta_{im}(1-x_j^2)\psi + \delta_{jm} x_i x_m \psi) d\mu - 2C_{PK} \int \left(x_i \delta_{jm} \partial_j \psi - \frac{1}{2} x_j [\delta_{jm} \partial_i \psi + \delta_{im} \partial_j \psi] \right) d\mu \right| \le \mathcal{E}(\psi)
$$

for all i, j, m with $i \neq j$. Taking $m = i$ and $\psi = [V]_{jj}$ gives

$$
\left| \int (x_j^2 - 1)[V]_{jj} d\mu - C_{PK} \int x_j \partial_j [V]_{jj} d\mu \right| \leq \mathcal{E}([V]_{jj}).
$$

On the other hand, taking $m = j$ and $\psi = [V]_{ij}$ gives

$$
\left| \int x_i x_j [V]_{ij} d\mu - 2C_{PK} \int \left(x_i \partial_j [V]_{ij} - \frac{1}{2} x_j \partial_i [V]_{ij} \right) d\mu \right| \leq \mathcal{E}([V]_{ij}).
$$

These can evidently be combined into the single matrix inequality

$$
\left| \int (x_i x_j - \delta_{ij}) [V]_{ij} d\mu - 2C_{PK} \int \left(x_i \partial_j [V]_{ij} - \frac{1}{2} x_j \partial_i [V]_{ij} \right) d\mu \right| \leq \mathcal{E}([V]_{ij}),
$$

holding for all $1 \le i, j \le n$. Summing over all $1 \le i, j \le n$ and applying the triangle inequality gives

$$
\left| \int (xx^T - \mathrm{Id}) \cdot V d\mu - 2C_{PK} \int \left((x \nabla^T) \cdot V - \frac{1}{2} (x \nabla^T) \cdot V^T \right) d\mu \right| \le \mathcal{E}(V),
$$

where $\mathcal{E}(V) := \sum_{i,j=1}^n \mathcal{E}([V]_{ij})$. The same is true when V is replaced by V^T . However, the matrix $(xx^T - Id)$ is symmetric, so we have

$$
\left| \int (xx^T - \mathrm{Id}) \cdot V d\mu - 2C_{PK} \int \left((x \nabla^T) \cdot V^T - \frac{1}{2} (x \nabla^T) \cdot V \right) d\mu \right| \le \mathcal{E}(V),
$$

An application of the triangle inequality gives

$$
\left| C_{PK} \int (x \nabla^T) \cdot V d\mu - C_{PK} \int (x \nabla^T) \cdot V^T d\mu \right| \leq \frac{2}{3} \mathcal{E}(V),
$$

and another gives

$$
\left| \int (xx^T - \mathrm{Id}) \cdot V d\mu - C_{PK} \int (x \nabla^T) \cdot V d\mu \right| \le \frac{5}{3} \mathcal{E}(V).
$$

Two final applications of the triangle inequality followed by Cauchy–Schwarz gives the desired conclusion

$$
\left| \int (xx^T - \text{Id}) \cdot V d\mu - \int (x\nabla^T) \cdot V d\mu \right| \le \frac{5}{3} \mathcal{E}(V) + (C_{PK} - 1) \left| \int (x\nabla^T) \cdot V d\mu \right|
$$

$$
\le \frac{5}{3} \mathcal{E}(V) + (C_{PK} - 1) \sum_{i,j=1}^n \int |x_i \partial_j [V]_{ij} | d\mu
$$

$$
\le \frac{5}{3} \mathcal{E}(V) + (C_{PK} - 1) \sum_{i,j=1}^n ||\partial_j [V]_{ij} ||_{L^2(\mu)}
$$

$$
\le K \sqrt{C_{PK} (C_{PK} - 1)} \sum_{i,j=1}^n ||\nabla [V]_{ij} ||_{L^2(\mu)}.
$$

2.4 Implementation of Stein's method

For a sufficiently smooth function $f : \mathbb{R}^n \to \mathbb{R}^m$, let $D^k f$ denote the tensor of k-th order derivatives. The tensor $D^k f(x)$ can be regarded as a vector in a space of dimension $m \times n^k$, which we equip with its natural Euclidean norm $\|\cdot\|_2$.

We'll need the following Lemma. It combines Barbour's solution to the classical Stein equation in terms of the Ornstein–Uhlenbeck semigroup $(P_t)_{t\geq0}$, defined by

$$
P_t f(x) := \int_{\mathbb{R}^n} f(e^{-t}x + (1 - e^{-2t})^{1/2}z) d\gamma(z), \quad f \in L^1(\gamma),
$$

and the higher-order regularity estimate, that was for example derived in [\[14\]](#page-11-18).

Lemma 3. For $f : \mathbb{R}^n \to \mathbb{R}$ with $\int f d\gamma < \infty$, the function $\varphi_f : \mathbb{R}^n \to \mathbb{R}^n$ defined by

$$
\varphi_f(x) := \nabla \int_0^\infty P_t f(x) dt \tag{7}
$$

 \Box

solves the Poisson equation

$$
f - \int f d\gamma = x \cdot \varphi_f - \text{Tr}(\nabla \varphi_f),\tag{8}
$$

and satisfies

$$
\sup_{x} \|D^{k}\varphi_{f}(x)\|_{2} \le \sup_{x} \|D^{k}f(x)\|_{2}, \quad k \ge 1.
$$
\n(9)

We are now ready to implement Stein's method to prove our main result. In particular, for a given test function $f : \mathbb{R}^n \to \mathbb{R}$ with uniformly bounded second derivatives, we'll bootstrap the solution φ_f to the Stein equation [\(8\)](#page-8-1) to construct a solution $V : \mathbb{R}^n \to M_{n \times n}(\mathbb{R})$ to the integrated second-order Stein equation

$$
\int f d\mu - \int f d\gamma = \int ((xx^T - \mathrm{Id}) \cdot V - (x\nabla^T) \cdot V) d\mu.
$$

The main stability result will then follow from the approximate Stein identity of Lemma [1,](#page-3-1) regularity estimates on V , and definition of the Zolotarev distance.

Proof of Theorem [1.](#page-2-1) Fix any $f : \mathbb{R}^n \to \mathbb{R}$ satisfying $\sup_x ||\nabla^2 f(x)||_2 \leq 1$. Using the classical identity $\nabla(P_t f) = e^{-t} P_t(\nabla f)$ for P_t , we have

$$
\nabla \varphi_f(x) = \int_0^\infty \nabla^2 P_t f(x) dt = \int_0^\infty e^{-2t} P_t(\nabla^2 f)(x) dt.
$$

By the triangle and Jensen inequalities, [\(9\)](#page-8-2), and boundedness of $\nabla^2 f$, [\(7\)](#page-8-3) implies

$$
\sup_{x} \|\nabla \varphi_f(x)\|_2 \le \frac{1}{2} \sup_{x} \|\nabla^2 \varphi_f(x)\|_2 \le \frac{1}{2} \sup_{x} \|\nabla^2 f(x)\|_2 \le \frac{1}{2}.
$$
 (10)

In particular, φ_f is 1/2-Lipschitz and $\nabla \varphi_f$ is 1-Lipschitz. Now, define $a \in \mathbb{R}^n$ and $Q \in M_{n \times n}(\mathbb{R})$ by

$$
a := \varphi_f(0), \quad [Q]_{ij} := (1 + \delta_{ij})[\nabla \varphi_f(0)]_{ij},
$$

and put

$$
g(x) := f(x) - a^T x - \frac{1}{2} x^T Q x.
$$

Using the fact that Hermite polynomials are eigenfunctions of the Ornstein–Uhlenbeck semigroup, we can check that the solution $\varphi_g := \nabla \int_0^\infty P_t g dt$ to the Poisson equation

$$
g - \int g d\gamma = x \cdot \varphi_g - \text{Tr}(\nabla \varphi_g), \tag{11}
$$

is equal to

$$
\varphi_g(x) = \varphi_f(x) - \varphi_f(0) - \nabla \varphi_f(0)x, \quad x \in \mathbb{R}^n.
$$

In particular, φ_g satisfies

$$
\varphi_g(0) = 0
$$
, $\nabla \varphi_g(0) = 0$, and $\nabla^2 \varphi_g(x) = \nabla^2 \varphi_f(x)$. (12)

We now establish some basic regularity properties of φ_g . Combining [\(12\)](#page-9-0) with the Lipschitz estimates established for φ_f , we have

$$
|\varphi_g(x)| \le \frac{1}{2}|x|
$$
, and $\|\nabla \varphi_g(x)\|_2 \le |x|$.

Additionally, by a Taylor expansion around $x = 0$, the properties [\(12\)](#page-9-0) together with boundedness of second-derivatives of φ_f imply the quadratic growth estimate

$$
|\varphi_g(x)| \le \frac{1}{2} \|\nabla^2 \varphi_f(0)\|_2 |x|^2 \le \frac{1}{2}|x|^2.
$$

Next, we define a matrix-valued function $V : \mathbb{R}^n \to M_{n \times n}(\mathbb{R})$ by

$$
V(x) := \begin{cases} \frac{1}{|x|^2} x \varphi_g^T(x) & \text{if } x \neq 0\\ 0 & \text{if } x = 0. \end{cases}
$$

By definition of V and the fact that $\varphi_g(0) = 0$, we have

$$
V^T(x)x = \varphi_g(x), \quad x \in \mathbb{R}^n.
$$

Now, we check the regularity of V. Since φ_g inherits continuity properties from φ_f , it follows that V is continuous on $\mathbb{R}^n \setminus \{0\}$. It is also continuous at $x = 0$, which follows since $|\varphi_g(x)| \leq \frac{1}{2}|x|^2$, and therefore $\lim_{x\to 0} V(x) = 0 = V(0)$.

Evidently, V is differentiable on $\mathbb{R}^n \setminus \{0\}$. For $x \neq 0$, we compute

$$
\nabla([V(x)]_{ij}) = \nabla \frac{x_i[\varphi_g(x)]_j}{|x|^2} = -2x \frac{x_i[\varphi_g(x)]_j}{|x|^4} + \frac{1}{|x|^2} (e_i[\varphi_g(x)]_j + x_i \nabla[\varphi_g(x)]_j).
$$

Using the regularity estimates on φ_g , we obtain

$$
|\nabla([V(x)]_{ij})| \leq |x_i| + \frac{1}{2} + \frac{|x_i|}{|x|} \leq |x_i| + \frac{3}{2}.
$$

Thus, using the moment assumption, we have

$$
\|\nabla([V]_{ij})\|_{L^2(\mu)} = \left(\int |\nabla([V(x)]_{ij})|^2 d\mu(x)\right)^{1/2} \le \left(\int (|x_i| + 3/2)^2 d\mu(x)\right)^{1/2} \le 5/2.
$$
 (13)

Finally, we put everything together to obtain

$$
\int f d\mu - \int f d\gamma = \int g d\mu - \int g d\gamma \tag{14}
$$

$$
= \int (x \cdot \varphi_g - \text{Tr}(\nabla \varphi_g)) d\mu \tag{15}
$$

$$
= \int \left(\left(xx^T - \text{Id} \right) \cdot V - \left(x \nabla^T \right) \cdot V \right) d\mu \tag{16}
$$

$$
\leq K\sqrt{C_{PK}(C_{PK}-1)}\sum_{i,j=1}^{n} \|\nabla[V]_{ij}\|_{L^{2}(\mu)} \tag{17}
$$

$$
< 20n^2\sqrt{C_{PK}(C_{PK}-1)}.\tag{18}
$$

In the above, (14) follows by definition of g and the moment assumption; (15) is (11) , integrated with respect to μ ; [\(16\)](#page-10-2) follows since $V^T(x)x = \varphi_g(x)$; [\(17\)](#page-10-3) follows from the approximate Stein identity of Lemma [1](#page-3-1) applied to (smooth approximations of) V ; [\(18\)](#page-10-4) is the bound [\(13\)](#page-10-5). Taking supremum over f proves the theorem. \Box

A Proof of Proposition [2](#page-1-1)

We first recall Caffarelli's contraction theorem [\[6\]](#page-11-19): if a probability measure $d\mu = e^{-\phi}dx$ satisfies $\nabla^2 \phi \geq \text{Id}$, then there exists a transport map T from γ onto μ that is 1-Lipschitz. This map T is the Brenier map from optimal transport theory.

For T as above, we have $|T(x) - T(y)| \le |x - y|$ for all $x, y \in \mathbb{R}^n$. Moreover,

$$
2n = \int |x - y|^2 d\mu(x) d\mu(y) = \int |T(x) - T(y)|^2 d\gamma(x) d\gamma(y)
$$

$$
\leq \int |x - y|^2 d\gamma(x) d\gamma(y) = 2n.
$$

Hence there is equality throughout, and $|T(x) - T(y)| = |x - y|$, $\gamma^{\otimes 2}$ -a.s. Since T is continuous, the equality holds everywhere. In the case where μ has full support, T is surjective, and therefore T is a surjective isometry. By the Mazur–Ulam theorem, we conclude that T is affine. Since μ is centered and isotropic by assumption, it must be standard Gaussian.

If μ does not have a full support, we can take a convolution with a standard Gaussian, rescaled so that the new measure ν is still isotropic. Since ν has full support and 1-uniform log-concavity is preserved by this operation, we can apply the previous case to deduce that ν is Gaussian. Since a convolution of two measures is Gaussian iff both are Gaussian, it follows that μ is also Gaussian.

References

- [1] B. Arras and C. Houdr´e, On Stein's method for multivariate self-decomposable laws with finite first moment. Electron. J. Probab. 24: 1–33 (2019).
- [2] D. Bakry, I. Gentil and M. Ledoux, Analysis and Geometry of Markov Diffusion Operators. Springer International Publishing, Springer Cham. 2014.
- [3] N. Belili and H. Heinich, Distances de Wasserstein et de Zolotarev. C. R. Acad. Sci. Paris, t. 330, Srie I, p. 811–814, 2000.
- [4] A.A. Borovkov and S.A. Utev. On an inequality and a related characterisation of the normal distribution. Theory of Probability and Its Applications, 28:219– 228, 1984.
- [5] T. Cacoullos, V. Papathanasiou and S. A. Utev, Variational Inequalities with Examples and an Application to the Central Limit Theorem. Ann. Probab. 22(3): 1607-1618 (1994).
- [6] L. Caffarelli, Monotonicity Properties of Optimal Transportation and the FKG and Related Inequalities. Comm. Math. Phys. 214, 547–563, (2000).
- [7] K. Carrapatoso, J. Dolbeault, F. H´erau, S. Mischler and C. Mouhot, Weighted Korn and Poincaré-Korn inequalities in the Euclidean space and associated operators. Arch. Rational Mech. Anal. 243 (3), 1565–1596, 2022.
- [8] F. Cavalletti, A. Mondino and D. Semola, Quantitative Obata's theorem, Anal. PDE 16 (2023), no. 6, 1389-1431.
- [9] G. Cébron, M. Fathi and T. Mai, A note on existence of free Stein kernels, *Proc. Amer.* Math. Soc. 148 (2020), no. 4, 1583-1594.
- [10] X. Cheng and D. Zhou, Eigenvalues of the drifted Laplacian on complete metric measure spaces. Commun. Contemp. Math. Vol. 19, No. 01, 1650001 (2017).
- [11] T. Courtade and M. Fathi, Stability of the Bakry-Emery theorem on $Rⁿ$. J. Funct. Anal. 279 (2020), no. 2, 108523, 28 pp.
- [12] T. Courtade, M. Fathi and A. Pananjady, Existence of Stein Kernels under a Spectral Gap, and Discrepancy Bounds, Ann. IHP: Probab. Stat., 55, 2, 2019.
- [13] G. De Philippis and A. Figalli, Rigidity and stability of Caffarelli's log-concave perturbation theorem. Nonlinear Anal. 154 (2017), 59–70.
- [14] M. Fathi, Higher-order Stein kernels for Gaussian approximation. Studia Math. 256 (2021), no. 3, 241-258.
- [15] M. Fathi, I. Gentil and J. Serres, Stability estimates for the sharp spectral gap bound under a curvature-dimension condition. To appear in Ann. Institut Fourier
- [16] A. Figalli, Stability in geometric and functional inequalities European Congress of Mathematics, 585-599, Eur. Math. Soc., Zurich, 2013.
- [17] A. Figalli, A short review on improvements and stability for some interpolation inequalities Proceedings of ICIAM 2023.
- [18] Horgan, C. O., Korn's Inequalities and Their Applications in Continuum Mechanics. SIAM Review, vol. 37, no. 4, 1995, pp. 491–511.
- [19] Korn, A. Die Eigenschwingungen eines elastischen Körpers mit ruhender Oberfläche. Akad. der Wissensch., Munich, Math. phys. KI. 36 (1906), 351.
- [20] Lewicka, M., and Müller, S. On the optimal constants in Korn's and geometric rigidity estimates, in bounded and unbounded domains, under Neumann boundary conditions. Indiana Univ. Math. J. 65, 2 (2016), 377–397.
- [21] C.H. Mai and S. Ohta, Quantitative estimates for the Bakry-Ledoux isoperimetric inequality, Comment. Math. Helv. 96 (2021), 693-739
- [22] J. Serres, Stability of the Poincaré constant, *Bernoulli* 29 (2) 1297 - 1320, May 2023.
- [23] J. Serres, Stability for higher order eigenvalues in dimension one. Stoch. Proc. Appl., 155, 2023, 459-484.
- [24] S. A. Utev. Probabilistic problems connected with an integro-differential inequality. Sibirsk. Mat. Zh., 30(3):182–186, 220, 1989