The Role of Acetylene in the Chemical Evolution of Carbon Complexity

Evgeniy O. Pentsak¹, Maria S. Murga², Valentine P. Ananikov^{1,*}

¹ Zelinsky Institute of Organic Chemistry, Russian Academy of Sciences, Moscow, 119991, Russia.

 ² Institute of Astronomy, Russian Academy of Sciences, Pyatnitskaya str. 48, Moscow 119017, Russia.

KEYWORDS. acetylene, benzene, PAH, carbon, catalysis, astrochemistry, prebiotic chemistry, chemical evolution

ABSTRACT

Acetylene, among the multitude of organic molecules discovered in space, plays a distinct role in the genesis of organic matter. Characterized by its unique balance of stability and reactivity, acetylene is the simplest unsaturated organic molecule known to have a triple bond. In addition to its inherent chemical properties, acetylene is one of the most prevalent organic molecules found across the Universe, spanning from the icy surfaces of planets and satellites and the cold interstellar medium with low temperatures to hot circumstellar envelopes where temperatures surge to several thousand kelvins. These factors collectively position acetylene as a crucial building block in the molecular diversification of organic molecules and solids present in space. This review comprehensively discusses the formation and expansion of carbon skeletons involving acetylene, ranging from the formation of simple molecules to the origination of the first aromatic ring and ultimately to the formation of nanosized carbon particles. Mechanisms pertinent to both hot environments, such as circumstellar envelopes, and cold environments, including molecular clouds and planetary atmospheres, are explored. In addition, this review contemplates the role of acetylene in the synthesis of prebiotic molecules. A distinct focus is accorded to the recent advancements and future prospects of research into catalytic processes involving acetylene molecules, which is a significant instrument in driving the evolution of carbon complexity in the Universe. The insights garnered from this review underscore the significance of acetylene in astrochemistry and potentially contribute to our understanding of the chemical evolution of the Universe.



Contents

1. Introduction	5
2. Acetylene in Universe. Origin and occurrence	9
2.1. The Jovian Planets	9
2.2. Satellites and Pluto	
2.3. Comets	
2.4. Exoplanets	
2.5. Carbon-rich stars	14
2.6. Cold interstellar medium	14
2.7 Photo-dissociation regions and diffuse interstellar medium	
2.8 Young stellar objects	
3. Acetylene in the network of chemical reactions in space	
3.1. Acetylene as a precursor of small hydrocarbons	
3.2. Acetylene as a precursor of polyynes	
3.3. Acetylene as a precursor of complex organic molecules	24
4. Benzene formation as the first step in PAH evolution	
4.1. Circumstellar envelopes and flame burning models	
4.2. Benzene formation in protoplanetary nebulae	
4.3. Benzene formation in cold interstellar medium	
4.4. Benzene formation in planets and moons	
5. Mechanisms of Formation of Polyaromatic Compounds in Space	40
5.1. PAHs in space and their role in carbon evolution	40
5.2. Mechanisms of PAH formation in hot environments	43
5.2.1. HACA mechanism	
5.2.2. Mechanism of formation of three- and four-ring PAHs in the circumstellar media	
5.2.3. Mechanism of formation of PAHs with five-membered rings	
5.2.4. PAH formation via polyyne polymerization	
5.3. Mechanisms of PAH formation in cold interstellar medium	
5.4. Mechanisms of PAH formation for planets and satellites	
6. Formation of carbon nanoparticles and grains	
6.1. Mechanism of formation of fullerenes and nanotubes	
6.2. Soot formation	
7. Catalytic reactions in space with acetylene	
7.1. Catalytic reactions of acetylene in the laboratory	

7.2. Catalytic reactions involving acetylene in space	60
7.3. PAHs as a catalyst	63
8. Acetylene in prebiotic chemistry evolution	66
8.1. Acetylene in reactions producing prebiotic molecules	66
8.2. From acetylene to nitrogen-bearing PAHs and nucleobases	71
8.3. Role of acetylene in monocarboxylic acid formation	75
9. Conclusions	78

1. Introduction

The past decade has witnessed substantial advancements in the chemistry of acetylene (C_2H_2), sparking novel insights into its role in the development of organic compounds and carbon structures associated with various celestial bodies. Rapid progress in areas such as carbon materials and carbocatalysis has led to the accumulation of a wealth of knowledge concerning acetylene chemistry.¹ This new understanding could inspire fresh breakthroughs in astrochemistry, particularly in relation to catalytic processes.

On Earth, acetylene is both a well-recognized and thoroughly studied molecule. Since its discovery by Davy in 1832² and later by Berthelot in 1860³, acetylene has proven to be both interesting and valuable. Its primary source is industrial production through the pyrolysis of hydrocarbon raw materials.⁴ Acetylene serves as a precursor for many industrial products, including vinyl chloride, vinyl acetate, acrylic acid, 1,4-butynediol, and carbon black. The production and industrial use of acetylene are among the global market indicators^{4,5}, and new potential applications of acetylene have recently been discovered.^{6,7,8,9} For example, new methods for producing carbon materials such as carbon nanotubes and graphene are of particular interest.^{10,11,12,13,14,15,16} The development of cycloaddition reactions involving acetylene has experienced a resurgence in recent years and is another exciting area.¹⁷

In the context of space, the situation around acetylene becomes even more intricate and fascinating. Chemical evolution from monatomic particles to complex carbon systems is a central issue in astrochemistry. The creation of complex aromatic systems is a thermodynamically favored process that pervades the Universe. The existence of numerous polyaromatic compounds, carbon structures such as fullerenes, and macroscopic carbon particles in space attests to this fact.^{18,19,20,21,22,23,24,25} Acetylene is widely considered a primary element in the formation of these structures.

Acetylene is among the simplest organic molecules found abundantly in outer space. It is detectable and readily formed in many different environments: circumstellar envelopes — the hot and dense layers that are lost by asymptotic giant branch (AGB) stars and moving away from them — the atmospheres of planets within and likely beyond the Solar System — in the cold interstellar medium (ISM) at extremely low temperatures and others. Recent studies underscore its significance in the chemistry of exoplanets.^{26,27}

The majority of related research has focused on the mechanisms of chemical reactions triggered by ultraviolet (UV) radiation and cosmic rays in the upper layers of planetary atmospheres, as well as in circumstellar and interstellar media. However, the evolution of organic matter on Earth-like planets or satellites may exhibit unique features. Recent discoveries in the chemistry of atmospheric and surface hydrocarbons on Titan and Enceladus confirm the crucial role of acetylene in the evolution of complex organic structures.^{28,29,30,31,32}

Acetylene molecules serve as excellent precursors for polyaromatics^{33,34} and carbon dust^{35,36} (Figure 1). Although the hydrogen removal/acetylene addition (HACA) paradigm remains the prevailing model for explaining the formation of simple polycyclic aromatic hydrocarbons (PAHs) in space, alternative mechanisms warrant consideration.^{37,38,39} Recent discoveries in chemistry have revealed a catalytic pathway for the formation of aromatic compounds from acetylene molecules. The trimerization of acetylene catalyzed by carbon molecules could unlock a wide variety of organic molecules (Figure 1).

Acetylene is hypothesized to play a significant role in prebiotic systems.^{40,41} Recent studies have indicated its involvement in the metabolic processes of primitive life forms and suggested that it is a probable precursor for numerous prebiotic molecules.⁴² According to the PAH world hypothesis, polyaromatic compounds contribute to the primary assembly of self-replicating informational polymers and likely have other essential roles in prebiotic environments.^{43,44,45} Unraveling the mechanisms of self-sustaining the growth of polyaromatic molecules is a key element in resolving these questions.

Consequently, a wealth of data suggest that acetylene may form bedrock for numerous chemical processes in outer space, leading to the evolution of complex organic molecules and carbon macroparticles from simpler carbon particles (Figure 1). In essence, acetylene could be viewed as a chemical platform for catalyzing the emergence of more intricate carbon species. Our fundamental understanding of its "cosmic" significance is only now beginning to unfold.

Generally, acetylene participates in almost all types of chemical reactions that lead to molecular growth (bottom-up reactions): neutral-neutral reactions, ion-molecular reactions, surface reactions, and nucleation. Neutral-neutral reactions mostly have an activation barrier and dominate in hot and dense environments, such as stellar envelopes and photodissociation regions (PDRs); however, neutral-neutral reactions are not excluded at low temperatures in molecular clouds, as some reactions are barrierless or because the reactants are radicals. Ion-molecular reactions more easily proceed in diffuse ISM (DISM) and in molecular clouds and PDRs due to dipole interactions. The ionization of atoms and molecules is achieved by cosmic rays and UV radiation. Surface chemistry becomes relevant in molecular clouds and photodissociation regions, especially for denser and darker objects and prestellar cores, where many gaseous species are adsorbed on the dust grain surface. Some faraway objects of the Solar System, such as Kuiper's belt objects or comets, can undergo similar chemical reactions because the temperature is also low there. At low temperatures the dust grains are covered by an icy mantle, and interactions between the ice and surface species also occur. Moreover, chemical reactions occur inside the icy mantle. The major trigger of many reactions at low temperatures is cosmic rays (CRs), which, unlike UV photons, penetrate dense cloud cores. However, not only does surface chemistry take place in cold objects, but it is also relevant for radically different conditions typical of hot stellar envelopes where aromatic molecules can be formed on or from the dust surface. Under the same conditions, the nucleation process plays a crucial role in promoting the growth of macroparticles from molecules. Regardless of the type of reaction, acetylene is a starting point for many pathways, leading to cosmic complexity and variety.

Acetylene is also a key reactant of the opposite 'top-down' chemistry, where it is a fragment of dissociation of macromolecules or macroparticles induced by factors such as UV radiation, bombardment by CRs, and shock waves. Thus, acetylene is in some sense a connecting element of the carbon lifecycle. We schematically illustrate the role of acetylene in different types of chemical reactions in Figure 1.

To guide the reader throughout the paper, we provide a summary table (Table 1), which includes chemical reaction types and corresponding sections where these reactions are considered, astronomical objects and their typical physical conditions (number density n_H, temperature T, ionizing source, UV or CR radiation) where these reactions occur. We consider most space objects where acetylene is observed and/or plays an important role in chemical processing of the medium. We note that one type of reactions (not necessarily the same reaction) can correspond to objects with very distinct conditions, and vice versa; many types of reactions can proceed under the same conditions. Further subsections contain more specific tables where individual reactions or chains of reactions under corresponding conditions are presented, although some crossing between reactions may occur because the conditions are similar; thus, it is not always possible to strictly indicate the objects where specific reactions proceed. Some chains are short and fully presented, while long chains are shortened. If rate constants are provided in original papers or alternatively in chemical databases, they are presented in the tables of subsections. Rates are often dependent on temperature, pressure or other parameters; therefore, in these cases, we calculated the rates at fixed values of the parameters, which are averages for the objects listed in the 3rd column. In some cases, quantum yields or other values, instead of rate constants, are given in the original papers; thus, we do not present the kinetics. Finally, there are reactions whose rates are undetermined or not provided.

 Table 1. Summary of acetylene chemistry in different astronomical media.

Type of C chemistry	Astronomical objects	n _H , cm ⁻³ (if gas	Т, К	G ₀ */CR	Subsection
UV photon-initiated	Atmospheres of Jovian	$10^{10} - 10^{16}$	50-300	up to 10^2	21 22 27
chemistry	planets and their satellites ^[46,47]				3.2, 4.2, 4.4, 5.4, 8.1, 8.2
	Kuiper belt objects, comets ^[48]		50	~1	
	PDRs ^[49]	$10^2 - 10^6$	20-1000	up to 10 ⁶	
	Protoplanetary nebulae	107	>200	up to 10 ⁶	
CR/UV photon initiated chemistry	Molecular clouds, prestellar cores	10 ² -10 ⁷	10	any G ₀ / Galactic CR	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$
in ice	Atmospheres of planets and their satellites	10 ¹⁰ -10 ¹⁶	50-300	field ^[51,52,53]	5.4, 8.2
	Kuiper belt objects, comets		50		
Neutral-neutral chemistry	Molecular and dust formation zone of carbon stellar envelopes ^[54]	10 ⁸ -10 ¹⁵	500-2500	not important	2.5, 3.1, 3.2, 4.1, 4.3, 5.2.1, 5.2.2, 5.2.3,
	Molecular clouds, prestellar cores	10 ² -10 ⁷	10		5.2.4, 5.3, 5.4, 6.1, 7.2, 8.2
	Hot cores ^[55]	≥ 10 ⁷	100-300		
Ion-molecular chemistry	PDRs	10 ² -10 ⁶	100-2000	up to 10 ⁶	2.7, 3.1, 3.2, 4.2, 4.4, 8.2
	Protoplanetary nebulae;	107	> 200	up to 10 ⁶	
	Molecular clouds prestellar cores	$10^2 - 10^7 \text{ cm}^{-3}$	10	very low	
	Atmospheres of Jovian planets and their satellites	10 ¹⁰ -10 ¹⁶	50-300	up to 10 ²	
	DISM ^[49]	50-100	100	0.1-100	
Nucleation	Dust formation zone of carbon stellar envelopes.	108-1012	500-2000	not important	6.2
Catalytic chemistry	Molecular clouds, prestellar cores	10 ² -10 ⁷	10	Galactic CR field	7.2, 7.3, 8.2
	Dust formation zone of carbon stellar envelopes	10 ⁸ -10 ¹²	500-2000	not important	
	Planet's satellites	1010-1016	50-300	up to 10 ² / Galactic CR field	
	PDRs, DISM	$10^2 - 10^6$	100-2000	up to 10 ⁶	

* - G_0 is a factor generally used to describe the level of the UV radiation field in space objects. The value of the factor is the ratio between integral intensity of the radiation field in the wavelength interval from 6 to 13.6 eV in an object and the same integral intensity in the solar neighborhood. The spectrum of the radiation field intensity in the Solar neighborhood was adopted from the work of Habing.⁵⁶ Strictly speaking, using this factor for estimation of the radiation field for the planets is incorrect due to the planets being within the HII region of the Sun while the factor is used for characterizing the conditions out of HII regions. However, we are not determined to give any precise estimations of the radiation field intensity, our goal is to show its approximate level to compare to other space objects in common units.

This review elucidates the most pertinent issues related to chemical processes involving acetylene molecules and their derivatives in outer space. The first section presents a brief introduction. The second

section offers a concise overview of the mechanisms of acetylene formation and its prevalence in space. The third section discusses various chemical reactions not associated with aromatic compounds, where acetylene plays a pivotal role. The fourth and fifth sections delve into the formation of an aromatic ring and the expansion of polyaromatic hydrocarbons, respectively. The sixth section examines the creation of nanoparticles, including fullerenes and nanotubes. The seventh section is dedicated to the role of catalytic processes involving acetylene in the spatiotemporal evolution of organic compounds. The eighth section speculates on the possible role of acetylene in prebiotic processes. Finally, the ninth section concludes the review with some closing remarks.



Figure 1. Evolution of carbon species in space with the participation of acetylene.

2. Acetylene in the Universe. Origin and occurrence

Carbon is mainly formed in thermonuclear fusion reaction at the stars of intermediate masses (from ~ $0.8M_{\odot}$ to $8M_{\odot}$).⁵⁷ Furthermore, carbon spreads into the ISM, driven by the stellar winds of these stars. Some fraction of carbon can be produced in massive stars and thrown into the ISM during supernova explosions.⁵⁸ Simple carbonaceous molecules, including acetylene and others (e.g., CO, CO₂, and CN), are formed from this carbon directly inside the parent stellar atmosphere⁵⁹, and the formation of even more molecules occurs further in the ISM. As a result, carbonaceous molecules are widespread throughout the Universe.^{60,61}

Acetylene does not have a permanent dipole moment and therefore does not have the rotational spectrum observed in the radio wave range. Instead, acetylene can be detected by rovibrational transitions that fall into the mid-infrared (mid-IR) range. The rovibrational spectrum of acetylene was both theoretically calculated and obtained in the laboratory.^{62,63} Some of the fundamental modes from this spectrum are observed in many space objects, namely, v^3 at approximately 3 µm, $v^4 + v^5$ at 7.5 µm and v^5 at 13.7 µm. Additionally, several weaker bands corresponding to these modes have been observed via high-resolution spectroscopy.⁵⁴ Acetylene also exhibits many absorption bands in the far-UV range between wavelengths of 0.15 and 0.19 µm, and these bands are also used for acetylene detection.^{47,64} Over the past half century, acetylene has been discovered in a fairly large number of space objects, both inside and outside the Solar system, in the atmospheres of giant planets and their satellites, comets, and circumstellar and interstellar environments. We summarize the observational evidence of the presence of acetylene in space objects in Table S1 in the Supporting Information. Below, we provide detailed information on the detection of acetylene along with the proposed pathways of its formation.

2.1. The Jovian Planets

For the first time, acetylene in space was detected in the atmosphere of Jupiter in 1974 based on the analysis of IR spectra obtained at the Kitt Peak National Observatory.^{65,66} The fundamental mode v^5 at nearly 13.7 µm was found. As a measure of abundance, the term 'mixing ratio' is used in studies of planetary atmospheres. This ratio represents the ratio between the specific gas and the dominant gas in the atmosphere. In the Jupiter atmosphere, the dominant gas is H₂ (similar to that in other Jovian planets), and the mixing ratio of acetylene was estimated to be $8 \cdot 10^{-5}$ relative to H₂. A little later, in 1980, acetylene was discovered on Saturn using the NASA Infrared Telescope Facility (IRTF) at Mauna Kea.^{67,68,69} The acetylene mixing ratio was found to be approximately $2 \cdot 10^{-7,70}$ The launch of Voyager 1 and Voyager 2⁷¹ in 1977 made it possible to significantly extend the available data on the abundance of acetylene in the atmosphere of the planet in the Solar system. In 1981, the IR spectra obtained during the passage of Voyager 1 through the Saturn system were published and showed the presence of acetylene in atmospheres of Saturn and Titan.^{70,72} In 1986, the mixing ratios for Jupiter and Saturn were estimated and found to be $1 \cdot 10^{-7}$ and $3 \cdot 10^{-7}$, respectively.⁷³ Acetylene was determined to be the third most abundant hydrocarbon in the atmosphere of these planets after CH₄ and C₂H₆, while the most abundant molecule was H₂.

In late 2000 and early 2001, the Cassini spacecraft passed near Jupiter, performing a gravityassisted maneuver on its way to Saturn. The Cassini orbiter was equipped with an ultraviolet imaging spectroscope (UVIS) connected to a camera (ISS) and two infrared spectrometers: a composite infrared spectrometer (CIRS) and a visible and infrared mapping spectrometer (VIMS).⁷⁴ CIRS provides the highest resolution map of the acetylene distribution on Jupiter (Figure 2).⁷⁵ A greater intensity of emissions in the region of the poles can be associated with both higher temperature and a greater abundance of gas.⁷⁶ In addition, recent observations of polar regions have revealed local enhancements in the abundances of acetylene and its derivatives that correlate with the location of diffuse auroras.^{77,78} The abundance of acetylene in the upper stratosphere derived from UVIS data is IR thermal emission from the middle stratosphere. In 2020, Melin et al. produced a vertical acetylene abundance profile above 1 mbar, which ensured the consistency of the data obtained by CIRS and UVIS. According to the new model, the abundance of acetylene was $1.21 \pm 0.07 \cdot 10^{-6}$ at 0.1 mbar.⁶⁴



Figure 2. Distribution of acetylene at the north and south poles of Jupiter based on data obtained by the CIRS instrument during the Cassini-Huygens mission. Dec. 2000 – Jan. 2001. Reproduced with permission from ref 75. Copyright 2010 NASA/JPL/GSFC.

Acetylene was detected on other planets, the Uranus and Neptune. In 1986, Broadfoot et al.⁴⁷ measured the acetylene mixing ratio in Uranus as $2 \cdot 10^{-7}$ using data from a Voyager Ultraviolet Spectrometer. This discovery confirmed the findings of Encrenaz et al.⁴⁶ in 1986 based on International Ultraviolet Explorer (IUE) satellite observations. C₂H₂ is the dominant constituent above the 1 mbar level.⁷⁹ Moreover, in 1987, Orton et al.⁸⁰ reported C₂H₂ IR emission at 13.7 µm obtained with NASA IRTF, which is consistent with the maximum mixing ratio of C₂H₂ being equal to 9·10⁻⁹. In the same work, Orton et al. reported recording a Neptune spectrum at approximately 13.7 µm, which is consistent with the emission of stratospheric acetylene with a maximum mixing ratio of 9 · 10⁻⁷. Caldwell et al. in 1988, based on IUE data, independently reported estimates of the acetylene mixing ratio, equal to 4·10⁻⁹, which is approximately 2 orders less than the estimates from the IR spectra.⁸¹ In 1989, Voyager 2 reached the Neptune system, and its atmosphere was examined using an infrared spectrometer (IRIS).⁸² Acetylene was detected with a mixing ratio of 10⁻⁷ to 10⁻⁶ in the atmosphere (approximately 1.5 mbar). For Neptune, Saturn and Jupiter, acetylene is the third most abundant hydrocarbon in the atmosphere after CH₄ and C₂H₆.⁸³

Since the first detections, the contributions of UV radiation and thunderstorms to the origin of acetylene in the atmosphere of gas giants have been actively discussed.⁸⁴ To date, the formation of acetylene has been associated with the photodissociation of methane in the upper stratosphere by UV radiation and the production of radicals that recombine in hydrocarbons with two or more carbon atoms.^{85,86,87} The photolysis of methane with the formation of acetylene occurs in the wavelength range of 120-190 nm.⁸⁸ According to Bar-Nun⁸⁴, incident solar radiation up to 140 nm is absorbed by methane in the atmosphere down to $\sim 5 \cdot 10^{-4}$ bar; at 145 nm, approximately 40% of the radiation is absorbed at 10^{-2} bar.

It should be noted that the degree of branching of CH₄ during UV photodissociation is still debatable. Thus, Romani et al.⁸⁹, based on the results of numerical modeling of methane photochemistry in the Neptune stratosphere, showed the photolytic formation of excited methylene ¹CH₂, ground-state methylene ³CH₂, and CH radicals but not the formation of methyl radical CH₃ as a result of CH₄ photolysis. In Romani's scheme, the formation of acetylene occurs through the addition of a second methane molecule to the CH radical (the product of photolysis of CH₄) to form an ethylene molecule and subsequent photolysis. Another pathway to acetylene is the photolysis of ethane. In turn, ethane is formed as a result of the recombination of methyl radicals formed from methylene or methylidene radicals (Scheme 1A).⁹⁰

Moreover, experimental studies have shown the possibility of the formation of a methyl radical as a result of the photolysis of methane when the sample is irradiated at a vacuum UV (VUV) wavelength of approximately 120 nm.^{91,92} Mordaunt et al.⁹³ also pointed to the formation of CH₃ (and CH) fragments rather than methylene radicals from the photodissociation of CH₄ at 121,6 nm and the importance of this pathway for the chemistry of the atmospheres of the outer planets. As a result, the formation of methyl groups occurs due to the photolysis of methane, subsequent recombination of methyl radicals to form ethane or a methyl radical with a methylene radical to form ethylene, and subsequent photolysis of ethane or ethylene may constitute alternative pathways for the formation of acetylene (Scheme 1B).⁹⁴



Scheme 1. Acetylene formation reactions during methane photolysis. Pathways based on Romani model adapted from the work 89 (A). Pathways through methyl radical based on the work 94 (B).

Thunderstorms in the clouds of Jupiter and Saturn are assumed to be an additional source of acetylene.^{84,95,96} Thunderstorms create extreme conditions by releasing energy from lightning strikes. The electric discharge of lightning causes intense heating of the gas along the current flow up to 30000 K,⁹⁷ which causes thermal dissociation of atmospheric species. In addition, the shock wave resulting from the sudden thermal expansion of the gas has an additional effect. These factors promote the dissociation of molecules into radicals. In the case of Jovian atmospheres, CH_x radicals can combine to form the corresponding compounds. Bar-Nun's calculation⁸⁴ showed that the lightning energy flux on Jupiter is 7.5 erg·cm⁻²·sec⁻¹, and the efficiency of converting heat into lightning energy on Jupiter is 5·10⁻⁴. Furthermore, based on experimental inventions⁹⁸, Bar-Nun noted that a thunderstorm column 5 km in length and 10 cm in diameter at a pressure of 7 bars converted 2.5 moles of methane (34%) to acetylene. As a result, Bar-Nun concluded that 245 strokes per km² per year will result in 7.8 · 10⁻¹⁶ mol·cm⁻²·sec⁻¹ acetylene, which is comparable in scale to the photolytic acetylene formation pathway. However, based on high-spectral-resolution Cassini/CIRS data, Hurley et al.⁹⁹ reported that there was no correlation between the content of upper tropospheric acetylene and thunderstorms 2000 km in size on Saturn. The main reactions described in this section are summarized in Table 2.

Reactions	Kinetics	Objects & conditions
1. a. $CH_4 + hv \rightarrow CH + H + H_2$ b. $CH + CH_4 \rightarrow C_2H_4 + H$ c. $C_2H_4 + hv \rightarrow C_2H_2 + 2H^{[89]}$	$k_{1a} = 2.4 \cdot 10^{-10} \text{ s}^{-1}$ (G ₀ =1, A _V =0) (KIDA ^[100]) $k_{1b} = 2.3 \cdot 10^{-10} \text{ s}^{-1} (\text{T} = 100 \text{ K})$ $k_{1c} = 3 \cdot 10^{-9} \text{ s}^{-1} (\text{G}_0 = 1, \text{A}_{\text{V}}=0)$	Atmospheres of Jovian planets. Conditions: $n_H = 10^{10}-10^{16} \text{ cm}^{-3}$ T = 50-300 K $G_0 = \text{up to } 10^2$
2. a. $CH_4 + hv \rightarrow CH_3 + H^{[86]}$ b. $CH_3 + CH_3 \rightarrow C_2H_6$ c. $C_2H_6 + hv \rightarrow C_2H_4 + H_2$ d. $C_2H_4 + hv \rightarrow C_2H_2 + 2H$	$k_{2a} = 2 \cdot 10^{-8} \text{ s}^{-1} \text{ (p} = 10^{-8} \text{ mbar}) - 3.6 \cdot 10^{-11} \text{ s}^{-1} \text{ (p}$ = 10 ⁻³ mbar) ^[86] $k_{2b} \approx 10^{-24} \text{ cm}^6 \text{ s}^{-1} \text{ (T} = 150 \text{ K}, \text{ n}_{\text{H}} = 10^{12} \text{ cm}^{-3}\text{)}$ $k_{2c} = 2.5 \cdot 10^{-9} \text{ s}^{-1} \text{ (p} = 10^{-8} \text{ mbar}) - 1.1 \cdot 10^{-10} \text{ s}^{-1}$ (p = 10 ⁻³ mbar) $k_{2d} = 1.1 \cdot 10^{-7} \text{ s}^{-1} \text{ (p} = 10^{-8} \text{ mbar}) - 6.6 \cdot 10^{-8} \text{ s}^{-1}$ (p = 10 ⁻³ mbar)	

Table 2. Reactions of acetylene formation in Jovian planets, their kinetics and conditions where the reactions are relevant.

2.2. Satellites and Pluto

In addition to giant planets, acetylene has also been found on other objects in the Solar system. Saturn's moon Titan has been particularly well studied. Unlike giant planets, Titan's atmosphere consists mostly of nitrogen, but hydrocarbons are also present. Voyager 1 flew by Titan in 1980 and probed its atmosphere with IRIS, registering the v^5 band at 13.7 µm.^{101,102} The acetylene mixing ratio was estimated to be $2 \cdot 10^{-6}$ relative to N₂ (for Titan's atmosphere) at an altitude of 130 km. Titan's atmosphere was later explored by the Cassini spacecraft.¹⁰³ Using CIRS mid-IR spectra, the mixing ratio in the stratosphere was found to be $3-4 \cdot 10^{-6}$.^{104,105} The Ion and Neutral Mass Spectrometer (INMS) data of the Cassini spacecraft showed that the acetylene mixing ratio in the thermosphere was $3.1\pm1.1 \cdot 10^{-4}$.^{105,106,107} Based on these studies, acetylene was found to be the third most abundant hydrocarbon in atmosphere of Titan (after CH₄ and C₂H₆).¹⁰⁸ In addition, acetylene is the most abundant unsaturated hydrocarbon in Titan's atmosphere. Similarly, as in the case of the Jovian planets, acetylene is formed as a result of the photolysis of methane in the case of Titan.¹⁰⁹

Using VIMS Cassini, acetylene was also found in solid form in three equatorial regions of Titan: Tui Regio, eastern Shangri-La, and Fensal–Aztlan/Quivira.¹¹⁰ Additionally, with the use of a gas chromatography mass spectrometer (GCMS) of Huygens probe, a trace amount of C_2H_2 gas sublimated/evaporated from the surface was detected.¹¹¹ Another moon on which acetylene was detected in Saturn is Enceladus. Cassini's INMS recorded the presence of acetylene while passing at an altitude 168 km from the surface of the satellite in 2005.³²

Finally, the recent New Horizons¹¹² mission in 2015 provided information on the composition of Pluto's atmosphere and allowed modeling of photochemical processes.¹¹³ The atmosphere of Pluto is very rarefied and consists of gases, mainly nitrogen with an admixture of methane (approximately 0.25%). Moreover, the data from New Horizons's UV spectrograph Alice indicate the absorption signatures of the products of the photochemical transformations of methane to ethylene and acetylene.¹¹⁴ Photochemical models predict that most of the precipitation on Pluto is formed by acetylene.¹¹⁵

Notably, there are near-Earth objects where acetylene was not detected, despite the presence and diversity of organic matter. For example, the Hayabusa2 mission examined the asteroid 162173 Ryugu and returned samples from its surface to Earth in 2020.^{116,117} Analysis of soluble organics revealed the presence

of PAHs, heterocyclic compounds, carboxylic acids, amino acids, amines, etc.¹¹⁸ In addition to the absence of acetylene, the abundance of methane and ethane in the gas samples was low.¹¹⁹ The reactions of acetylene formation in satellites are similar to those for the Jovian planets given in Table 2.

2.3. Comets

Comets are another important class of objects in the Solar system in which acetylene was discovered. The first comet in which acetylene was detected was comet C/1996 B2 (Hyakutake). The detection was performed in 1996 with the NASA IRTF.¹²⁰ The abundance of acetylene was estimated to be 0.3–0.9% relative to water, the dominant molecule in cometary ice.¹²¹ Subsequently, acetylene has been detected on many other comets using a cryogenic echelle spectrometer (CSHELL) at NASA IRTF and via IR spectroscopy with NIRSPEC at the Keck Observatory on Mauna Kea¹²² in amounts up to 0.5% relative to water. In addition, the presence of acetylene in the comets Halley and Churyumov-Gerasimenko was confirmed by the data received by the space missions Giotto¹²³ in 1999 and Rosetta¹²⁴ in 2015, respectively.

The Rosetta mission revealed a high acetylene abundance compared to that of other Jupiter family comets. Moreover, according to data from the Rosetta Orbiter Sensor for Ion and Neutral Analysis (ROSINA), which analyzed the chemical composition of cometary coma, the abundance of C_2H_2 in the summer hemisphere was 0.045% relative to that in water, while in the winter hemisphere, it reached 0.55%. Léna Le Roy et al.¹²⁴ noted that the abundance of ethane is also much greater than that of any other comet. One of the possible pathways for the formation of ethane in comets may involve the passage of acetylene, which is produced in the gas phase, condenses into icy grains, and hydrogenates into dense interstellar clouds.¹²⁵ The amount of acetylene in comets is comparable to its amount in interstellar ices; therefore, this acetylene has an interstellar origin. It might be formed through gas phase ion-molecular or neutral-neutral reactions and then adsorbs onto the icy mantle of dust grains.^{125,120} We refer readers to sections 2.6 and 2.7, where such reactions are discussed.

2.4. Exoplanets

In the last decade, research on the chemical composition of discovered exoplanets has been actively conducted. Acetylene is considered an essential building block for prebiotic systems and may indicate the possibility of abiogenetic processes occurring on planets.^{29,42} Therefore, acetylene is one of the molecules identified in the spectra of exoplanet atmospheres.¹²⁶ Tsiaras et al. (2016) suggested that acetylene may be present in the atmosphere of super-Earth 55 Cancri.¹²⁷ The prediction of the presence of acetylene in the atmosphere of exoplanets was based on data obtained by modeling and data from IR telescopes (see SI Table S1). However, the lack of spectral data over a broad range and at high temperature did not allow us to firmly establish the presence of acetylene. Despite the difficulties in detecting acetylene on exoplanets, the probability of the presence of acetylene molecules on super-Earth exoplanets is estimated to be high.¹²⁶ Tennyson and Yurchenko noted²⁶ that acetylene is probably the most important polyatomic molecule for exoplanet and cool star research, for which there is still no comprehensive hot line list. For a polyatomic molecule, the size of the acetylene and the transitions between rovibrational levels at exoplanet and cool star temperatures give rise to spectra typically spanning the IR region. The problem with acetylene is due to its linear geometry in the equilibrium structure. The rovibrational spectra of linear acetylene requires special consideration when calculating the rovibrational energies. Nevertheless, computational processing is feasible: an extensive effective Hamiltonian fit is available, and the calculations essentially require solving the Schrödinger equation of nuclear motion with some approximations.¹²⁸

To facilitate the detection of important molecular species, such as acetylene, in atmospheres of exoplanets and cool stars, the ExoMol spectroscopic database was created.^{26,27} Molecules suggested to be important for exoplanet spectroscopy are included in the ExoMol molecular line list. Spectral lines of molecules can indicate the type of exoplanet. Spectroscopy of hot gas giants is very different from that of hot rocky objects and cooler objects. The search for acetylene signals on exoplanets continues.^{127,129,130,131,132} There is no information on formation routes; for some exoplanets, routes can be the same as for Solar system planets. Although there is no information on general formation routes for exoplanets, in some cases, the routes can be the same as those for Solar system planets.

2.5. Carbon-rich stars

Acetylene was first discovered outside the Solar system in 1976 in the IR spectrum of IRC +10216 with the 4-m Mayall telescope at the Kitt Peak observatory.¹³³ The detection was based on the band at $\approx 2.4 \,\mu\text{m}$, which is a combination of ν^1 and ν^5 modes. The object IRC +10216 is a well-studied carbon AGB star that is embedded in a thick dust envelope. The detection of acetylene was important, as acetylene is thought to be a precursor of dust. This star was observed in the IR range repeatedly afterwards. For example, Fonfria et al.⁵⁴ performed a detailed analysis of the high-resolution spectra at 11-14 μm and identified many acetylene bands corresponding to the modes ν^2 , ν^4 , ν^5 and their combinations, although many modes are not observed properly. The abundance of C₂H₂ in the envelope of IRC +10216 obtained from the observations is $3 \cdot 10^{-8} - 1 \cdot 10^{-7}$ relative to that of H₂.^{134,135} Subsequently, with a great contribution from space IR missions, the Infrared Space Observatory (ISO)¹³⁶ and Spitzer Space Telescope¹³⁷, the presence of acetylene was also established by the IR spectra in other carbon stars^{138,139} both in the Milky Way and other galaxies^{140,141,142} and in the protoplanetary nebula CRL 618¹⁴³.

Acetylene is one of the most abundant molecules after CO in envelopes around C-rich stars and has a carbon-to-oxygen ratio (C/O) greater than $1.^{140}$ Below (Scheme C and Table 3), the chain of reactions leading to the formation of C₂H₂ is presented, although other chains of neutral-neutral reactions are also possible.^{144,145} First, C₂ is formed in the reaction between atomic C and radical CH (Scheme 2A). Then, an ethynyl radical (C₂H) is formed from the C₂ chain and a hydrogen molecule just after the shock wave front (Scheme 2B). Finally, acetylene appears as a result of an exchange reaction with its radical (Scheme 2C).

Α	С	+	СН.	>	C ₂	+	Н۰
В	C ₂	+	H_2	>	C₂H ∙	+	н۰
С	C₂H⁺	+	H_2		C_2H_2	+	н۰

Scheme 2. Reactions of acetylene formation.

Table 3. Chain of reactions leading to acetylene formation in carbon-rich stellar envelopes.

Reactions	Kinetics	Objects & conditions
a. $C + CH \rightarrow C_2 + H$ b. $C_2 + H_2 \rightarrow C_2H + H$ c. $C_2H + H_2 \rightarrow C_2H_2 + H$ ·[144]	$k_{a} = 2.4 \cdot 10^{-10} \text{ cm}^{3} \text{ s}^{-1} \text{ (KIDA)}$ $k_{b} = 8 \cdot 10^{-11} \text{ cm}^{3} \text{ s}^{-1} \text{ (T} = 2000 \text{ K)}$ (KIDA) $k_{c} = 10^{-10} \text{ cm}^{3} \text{ s}^{-1} \text{ (UMIST)}^{[146]}$	Stellar envelopes $n_{\rm H} = 10^8 - 10^{15} \text{ cm}^{-3}$ T = 500 - 2500 K

2.6. Cold interstellar medium

In cold environments, acetylene can be locked in icy mantles on dust grain surfaces. Astrochemical models predict that the abundance of acetylene is approximately 10^{-3} relative to that of water in ice.^{147,148} The detection of acetylene in solid form is difficult because acetylene has weak features that are blended with much stronger features of H₂O.¹⁴⁹ At the same time, acetylene was indicated in the gas phase by its vibrational modes in a number of star-forming regions and young stellar objects.^{150,151} The observed abundances are relatively high, e.g., a few of 10^{-8} relative to H₂ in the star-forming region Cepheus A East in the work of Sonnentrucker et al.¹⁵⁰ or ~ 10^{-7} relative to H₂ in young stellar objects in the work of Lahius et al.¹⁵¹ Gas-phase chemical reactions provide lower abundances of acetylene – from 10^{-10} to 10^{-8} relative to H₂.¹⁵² Therefore, it is assumed that the observed gas-phase acetylene is a result of sublimation of interstellar ice. Sublimation can occur via thermal or nonthermal desorption. In the case of young stellar

objects where the temperature exceeds 300 K¹⁵¹, thermal desorption is efficient, while in the case of colder objects, desorption can be a result of sputtering caused by shocks.¹⁵⁰

In the gas phase, acetylene can be formed through neutral-neutral and ion-molecular reactions. For example, acetylene can be generated through electron recombination of $C_2H_3^+$, which in turn results in a reaction of methane and C⁺.¹⁵³ Acetylene cations can be generated directly through the same reaction between methane and C⁺.¹⁵³ Alternative pathways of C₂H₂ formation occur on dust ice surfaces. Based on experiments with hydrocarbons in water ice, Hudson & Moore noted that there is no evidence for C₂H₂ formation in ice and suggested that C₂H₂ is part of the natal ice composition.¹⁵⁴ However, the observed abundance is often greater than that predicted by reaction network models.^{155,156} In 2023 in experimental work using the Reaction Apparatus for Surface Characteristic Analysis at Low Temperature (RASCAL) accompanied by temperature-programmed desorption (TPD) and a combination of photostimulated desorption and resonance-enhanced multiphoton ionization (PSD-REMPI) methods. Tsuge et. al. 157 reported that C atoms are weakly bound to the surface of ice and can diffuse on the surface of amorphous solid water, which is the major component of the icy mantle of cosmic dust in cold regions, at temperatures above approximately 30 K. It was determined that the diffusion reaction of C atoms is activated at approximately 22 K on interstellar ice. The authors were able to demonstrate the formation of C_2 molecules. TPD measurements showed an increase in the C_2H_6 signal (m/z 28) in the temperature range from 70 to 90 K for the water ice sample with carbon atoms deposited at 10 K followed by exposure to H atoms. The formation of stable methane (CH_4) or ethane (C_2H_6) molecules through successive hydrogenation reactions of species C and C_2 can be detected by the TPD method. Importantly, the reactions involving the sequential addition of hydrogen atoms to C_2 and C_2H species to form acetylene C_2H_2 are barrier-free. It was also previously shown that the hydrogenation of acetylene to ethylene and ethane can successfully occur in water ice at 10 and 20 K even in the environment of a dark molecular cloud.¹⁵⁸ Moreover, the authors¹⁵⁷ noted that the contribution of the interaction of two CH3 species to the formation of C2H6 should be insignificant in the TPD process. Under the experimental conditions during hydrogenation, CH₃ radicals are immediately converted to methane CH4 due to the barrier-free reaction of CH3 and H; thus, they do not remain on the surface of water ice.



Figure 3. Chemical processes in the gas phase and on the grains. At 20 K < T < 30 K physisorbed C atoms can diffuse, promoting insertion reactions with other C atoms with formation C₂ species. Reproduced with permission from ref 157. Copyright 2023 Springer Nature.

Laboratory studies^{159,160} of the irradiation of methane ice at 10 K energetic electrons as an imitation of processes caused by cosmic rays have shown that the recombination of the resulting methyl radicals, as well as radiolysis with the elimination of atomic hydrogen, can lead to the formation of acetylene. This process can explain the synthesis of acetylene in interstellar ice and comets and in aerosol particles and organic haze layers of hydrocarbon-rich atmospheres of planets and their satellites, such as Titan, which was mentioned above. This pathway is illustrated in Figure 4.



Figure 4. Reaction pathway for the CR- and photoinitiated formation of acetylene from methane.

In addition, Feldman et al. showed¹⁶¹ that the radiolysis of ethylene and ethane at 5 K in noble gas matrices under the influence of X-ray radiation at 20 keV leads to the formation of C_2H_2 molecules as a result of progressive dehydrogenation. The widespread occurrence of acetylene in harsh space environments with high levels of radiation is due not only to the ease of its formation but also to the high stability of the acetylene molecule and its radiation resistance in the series of C_2 hydrocarbons (ethane < ethylene < acetylene). We summarize the reactions considered in this section in Table 4.

Reactions	Kinetics	Objects & conditions
1. Ion-molecular gas phase reactions ^[162] a. $CH_4 + C^+ \rightarrow C_2H_3^+ + H$ b. $C_2H_3^+ + e^- \rightarrow C_2H_2 + H$ 2. $CH_4 + C^+ \rightarrow C_2H_2^+ + H_2$	$k_{1a} = 10^{-9} \text{ cm}^{3} \text{ s}^{-1} \text{ (UMIST)}$ $k_{1b} = 3 \cdot 10^{-7} \text{ cm}^{3} \text{ s}^{-1} \text{ (T} = 100 \text{ K)} \text{ (UMIST)}$ $k_{2} = 3.89 \cdot 10^{-10} \text{ cm}^{3} \text{ s}^{-1} \text{ (UMIST)}$	DISM, molecular clouds. Conditions: T = 10 - 500 K $n_H = 10^{-1} - 10^7 \text{ cm}^{-3}$ $G_0 = \text{from very low to } 10^2$ CR field: Galactic
3. Reactions in methane ice ^[163] a. $2CH_4 \rightarrow C_2H_6 + 2H/H_2$ b. $C_2H_6 \rightarrow C_2H_4 + 2H/H_2$ c. $C_2H_4 \rightarrow C_2H_2 + 2H/H_2$	$k_{3a} = 6.14 \cdot 10^{-5} \text{ s}^{-1}$ $k_{3b} = 3.18 \cdot 10^{-3} \text{ s}^{-1}$ $k_{3c} = 1.22 \cdot 10^{-3} \text{ s}^{-1}$	Ice on dust grains in molecular clouds, in comets, aerosol particles and organic haze layers of planets and their satellites. Conditions: T = 5-100 K
4. Diffuse chemistry on ice surface ^[157] a. $C + C \rightarrow C_2$ b. $C_2 + H \rightarrow C_2H$ c. $C_2H + H \rightarrow C_2H_2$	n.d.	CR field: Galactic G_0 from very low to ~10 ²

Table 4. The reactions of acetylene formation in cold interstellar medium.

2.7 Photo-dissociation regions and diffuse interstellar medium

Acetylene is not detected in PDRs or DISM; moreover, it is highly sensitive to UV radiation.³⁸ Nevertheless, some amount of acetylene may be present in the PDRs and participate in the reactions leading to the different small hydrocarbons C_mH_n , which were detected.^{195,193} Several pathways for the formation of acetylene and acetylene cations at low temperatures were shown above in Section 2.6. As shown above, acetylene can be formed through ion-molecular reactions, namely, through electron recombination of $C_2H_3^+$. At temperatures less than ~300 K, $C_2H_3^+$ can be generated as a result of the reactions 1,2 shown in Table 4.¹⁵³ At higher temperatures, the reactions with excited H₂ can be activated as shown in Scheme 3^{145,164} and Table 5.

$$CH \xrightarrow{C^{+}} C_{2}^{+} \xrightarrow{H_{2}} C_{2}H^{+} \xrightarrow{H_{2}} -H$$

$$\xrightarrow{H_{2}} C_{2}H_{2}^{+} \xrightarrow{H_{2}} C_{2}H_{3}^{+} \xrightarrow{e^{-}} C_{2}H_{2}$$

Scheme 3. Possible routes of acetylene formation in PDRs and DISM.

On the other hand, the acetylene molecule can be obtained from a top-down mechanism. This molecule is the main carbonaceous dissociation channel for PAHs^{165,166,167}. This channel becomes even more important if PAHs are hydrogenated^{168,169} while hydrogenated PAHs are likely present in the ISM.¹⁷⁰ In PDRs, PAHs absorb UV photons with energies up to 13.6 eV. This energy efficiently converts to PAH internal energy,¹⁷¹ which becomes higher than the dissociation energy of approximately 4-5~eV.¹⁷² Although the dissociation process competes with ionization and IR relaxation, its probability was estimated to be high enough that small PAHs would be destroyed in the ISM.¹⁶⁵ The minimum size of PAHs that survive in the ISM varies from 15 to 50 depending on the calculation method and parameters.^{165,173,174} This size is greater under harsh UV radiation from PDRs; hereby, small PAHs are destroyed much quicker than in the diffuse ISM, i.e., this origin of acetylene is time limited. ¹⁷⁵ Larger PAHs can live longer than smaller PAHs, but their dissociation probability is lower; hence, large PAHs should not be considered significant contributors to acetylene. The destruction of PAH clusters likely compensates for the abundance of small PAHs, as they seem to be destroyed in PDRs¹⁷⁶. However, calculations of the impact of PAH cluster destruction have not yet been performed.

The grains from hydrogenated amorphous carbon (HAC) are larger than those from PAHs, and they are not planar. However, irradiation may also lead to the release of acetylene, as shown by Alata et al. (2015). Moreover, they lose other small hydrocarbons (C_2H_4 , C_3H_4 , C_3H_6 , C_4H_4 , etc.^{177,178}), whole PAHs and even fullerenes.¹⁷⁹ However, these experimental results should be used with caution because the hydrogenation level of HAC grains in the ISM should be much lower than that in laboratory samples.^{180,181} Enrichment of the ISM by small hydrocarbons ceases quickly without reverse hydrogenation because hydrogen loss is a much more favorable dissociation channel, while dehydrogenated HAC grains become stable during UV irradiation.¹⁸² Table 5 summarizes the reactions of acetylene formation considered in this section.

Reactions	Kinetics	Objects & conditions
1. Ion-molecular reactions with excited $H_2^{[145, 183]}$ a. $C^+ + CH \rightarrow C_2^+ + H$ b. $C_2^+ + H_2 \rightarrow C_2H^+ + H$ c. $C_2H^+ + H_2 \rightarrow C_2H_2^+ + H$ d. $H_2 + C_2H_2^+ \rightarrow C_2H_3^+ + H$ e. $C_2H_3^+ + e^- \rightarrow C_2H_2 + H$	$k_{Ia} = 3.8 \cdot 10^{-10} \text{ cm}^3 \text{ s}^{-1} \text{ (UMIST)}$ $k_{Ib} = 1.1 \cdot 10^{-9} \text{ cm}^3 \text{ s}^{-1} \text{ (UMIST)}$ $k_{Ic} = 1.1 \cdot 10^{-9} \text{ cm}^3 \text{ s}^{-1} \text{ (UMIST)}$ $k_{Id} = 1.1 \cdot 10^{-11} \text{ cm}^3 \text{ s}^{-1} \text{ (UMIST)}$ $k_{Ie} = 3 \cdot 10^{-7} \text{ cm}^3 \text{ s}^{-1} \text{ (T=100 K)}$ (UMIST)	The reactions may take place in many objects (PDRs, DISM, molecular clouds, protoplanetary disks) where ions can be produced by UV radiation or by CRs. Conditions: T = 10 - 500 K
2. Ion-molecular reactions a. $C^+ + H_2 \rightarrow CH_2 + h\nu$ b. $CH_2^+ + H_2 \rightarrow CH_3^+ + H$ c. $CH_3^+ + e \rightarrow CH_2 + H$ d. $CH_2 + C^+ \rightarrow C_2H^+ + H$ e. $C_2H^+ + H \rightarrow C_2H_2^+ + H^{[184]}$	$k_{2a} = 3 \cdot 10^{-16} \text{ cm}^3 \text{ s}^{-1} (\text{T} = 100 \text{ K})$ (KIDA) $k_{2b} = 1.2 \cdot 10^{-9} \text{ cm}^3 \text{ s}^{-1} (\text{KIDA})$ $k_{2c} = 5 \cdot 10^{-7} \text{ cm}^3 \text{ s}^{-1} (\text{KIDA})$ $k_{2d} = 10^{-9} \text{ cm}^3 \text{ s}^{-1} (100 \text{ K}) (\text{KIDA})$ $k_{2e} = 1.1 \cdot 10^{-9} \text{ cm}^3 \text{ s}^{-1} (\text{KIDA})$	$n_{\rm H} = 10^{-1} - 10^7 \text{ cm}^{-3}$ $G_0 = \text{from very low to } 10^5$ CR field: Galactic
3. Top-down chemistry: PAH/HAC \rightarrow n(C ₂ H ₂)	Rates are sensitive to a parent particle (type, size, charge, C/H) and G_0 , n_H , $T^{[175,185]}$ For a PAH ⁰ with $N_C = 50$ and $G_0 = 1$: $k = 10^{-15} \text{ s}^{-1}$	

Table 5. The reactions of the acetylene formation in PDRs and DISM.

2.8 Young stellar objects

 C_2H_2 was also detected in young stellar objects and on protoplanetary disks via rovibrational modes.^{151, 186,187} The abundance of C_2H_2 was found to be 10⁻⁸-10⁻⁶ relative to H_2 , and this value increases as the temperature increases, which is consistent with the hypothesis that at low temperature, C_2H_2 is locked in ice and appears in gas due to ice sublimation when a young star heats the surrounding gas.¹⁵¹

The gas phase abundance is normally consistent with the estimates of its abundance in ice.¹⁵¹ Recently, C_2H_2 in protoplanetary disks was observed with the James Webb Telescope.¹⁸⁸ While the ratio between the abundances of C_2H_2 and CO_2 is on the order of unity in three objects, it is 10⁴ in disk J 160532.¹⁸⁹ This high abundance cannot be explained by evaporation from ice. Van Dischoek et al.¹⁹⁰ suggested that the enhanced abundance is a result of thermal and/or photodestruction of carbonaceous grains, whether they are PAHs or HACs, similar to what occurs in PDRs (Section 2.7).

The above examples demonstrate the presence of acetylene in a variety of conditions throughout the Universe, both in objects inside the Solar System, in the atmosphere and on the surface of planets and satellites, and beyond, in stellar and interstellar objects. Acetylene molecules can be found in different states and forms, from the cold ice of comets to the hot gases in the circumstellar space. The variety of states and conditions determines the variety of chemical processes. The possibility of acetylene formation along the bottom-up and top-down pathways provides greater dynamism of organic chemical environments and the possibility of the formation of new structures when conditions change. The wide distribution of acetylene makes it a universal molecule, the activity of which is capable of providing a wide network of chemical transformations of both aliphatic organic molecules and large aromatic particles, "pure" hydrocarbon structures and compounds with oxygen, nitrogen, etc. Thus, acetylene can determine and influence the chemistry of various space environments, promoting chemical evolution in space.

3. Acetylene in the network of chemical reactions in space

Acetylene plays an important role in the chemistry of the ISM and planetary systems, including the Solar system. It is a key reactant in the formation of a great number of molecules. The widely used astrochemistry network of chemical reactions, KIDA (Kinetic Database for Astrochemistry),¹⁹¹ includes approximately 9 thousand reactions, approximately 5 hundreds of which are reactions with acetylene. This molecule participates in many different reactions, including general reactions such as charge exchange, absorption of photons or cosmic rays, and specific reactions involving the direct formation of organic molecules from small hydrocarbons to complex large molecules as a parent molecule or a transitional product. While some products of the reactions with acetylene are observed, e.g., small hydrocarbons, and consequently, chemical models can be checked, the detection of complex organic molecules (COMs) is more complicated, and the progression of the corresponding reactions remains questionable. In this section, we consider the reactions with acetylene resulting in nonaromatic products, and the next sections will focus on the network of reactions related to aromatic molecules and dust.

3.1. Acetylene as a precursor of small hydrocarbons

Small hydrocarbons (C₂H, C₄H, c-C₃H₂, l-C₃H⁺, etc.) are observed in various space objects.^{192,193,194} Several pathways lead to their appearance, but the pathways mainly start from acetylene. Acetylene can react with CH via a neutral-neutral reaction leading to C_3H_2 (we omit c- and l-) (Scheme 4A) or participate in a series of ion-molecular reactions that end in $C_{3}H_{2}$ (Scheme 4B). Other examples, namely, the formation of $C_4H_3^+$, $C_4H_2^+$, etc., are given in Table 6.

However, the presented chemical routes do not provide enough hydrocarbons, which are found in some PDRs.^{193,195,196} To explain the high abundance of these hydrocarbons, reactions with excited hydrogen molecules have been suggested.¹⁸³ Hydrogen molecules can be excited if the temperature exceeds a few hundred Kelvin. In particular, a chain of such reactions starting from acetylene can be described as shown in Scheme 4.

Scheme 4. Formation of C₃H₂ molecule from acetylene.

С

It is worth mentioning that these reactions take place only in hot PDRs, such as the Orion Bar PDR, but not in all PDRs with the increased abundance of small hydrocarbons. For instance, the PDR Horsehead nebula is relatively cold, and these reactions do not proceed there. Dissociative reactions induced by CRs and photons can introduce some C_3H_2 but still cannot explain the observed abundance of C_3H_2 and other small hydrocarbons. This underestimation is mainly due to the small abundance of the parent acetylene. As discussed above in Section 2.7, acetylene can be a result of PAH dissociation or other carbonaceous solid particles that are present in such PDRs^{178,193}; however, Murga et al.^{175,182} showed that this channel also cannot help achieve the observed abundances. Thus, alternative mechanisms that can provide additional acetylene or reactions that convert acetylene or related molecules to small hydrocarbons are needed.

Table 6. The reactions that start from acetylene and lead to formation of small hydrocarbons.

Reactions	Kinetics	Objects & conditions
Neutral-neutral reactions: 1. $C_2H_2 + CH \rightarrow C_2H_3 + H$ 2. $C_2H + C_2H_2 \rightarrow C_4H_2 + H$	$k_1 = 3.39 \cdot 10^{-10} \text{ cm}^3 \text{ s}^{-1}$ (UMIST) $k_2 = 1.2 \cdot 10^{-10} \text{ cm}^3 \text{ s}^{-1}$ (T = 100 K) (UMIST)	The reactions may take place in many objects (PDR's, diffuse ISM, molecular clouds, protoplanetary disks) where ions can be produced
Ion-molecular reactions ^[145,183] 3. a. $C_2H_2 + C^+ \rightarrow C_3H^+ + H$ b. $C_3H^+ + H_2 \rightarrow 1/c - C_3H_3^+ + hv$ c. $c - C_3H_3^+ + e^- \rightarrow c - C_3H_2 + H$	$k_{3a} = 1.78 \cdot 10^{-9} \text{ cm}^3 \text{ s}^{-1}$ (UMIST) $k_{3b} = 5.5 \cdot 10^{-12} \text{ cm}^3 \text{ s}^{-1}$ (T=100 K) (KIDA) $k_{3c} = 7 \cdot 10^{-7} \text{ cm}^3 \text{ s}^{-1} \text{ (T} = 100 \text{ K)} \text{ (KIDA)}$	by UV radiation or by CRs. Conditions: T = 10 - 500 K $n_H = 10^{-1} - 10^7 \text{ cm}^{-3}$ $G_0 = \text{from very low to } 10^5$ CR field: Galactic
4. a. $C_2H_2 + C^+ \rightarrow C_3H^+ + H$ b. $C_3H^+ + e^- \rightarrow C_3 + H$ c. $C_3 + H_2 \rightarrow 1/c - C_3H + H$ d. $1/c - C_3H + H_2 \rightarrow 1/c - C_3H_2 + H$	$k_{4a} = 1.78 \cdot 10^{-9} \text{ cm}^3 \text{ s}^{-1}$ (UMIST) $k_{4b} = 2.5 \cdot 10^{-7} \text{ cm}^3 \text{ s}^{-1} (\text{T} = 100 \text{ K}) (\text{UMIST})$ $k_{4c} = 1.2 \cdot 10^{-5} \text{ cm}^3 \text{ s}^{-1} (\text{T} = 100 \text{ K}) (\text{KIDA})$ $k_{4d} = 1.5 \cdot 10^{-7} \text{ cm}^3 \text{ s}^{-1} (\text{T} = 100 \text{ K}) (\text{KIDA})$	
5. $C_2H_2 + C_2H_2^+ \rightarrow C_4H_2^+ + H_2$ 6. $C_2H_2 + C_2H_2^+ \rightarrow C_4H_3^+ + H^{[197]}$	$k_5 = 5.2 \cdot 10^{-10} \text{ cm}^3 \text{ s}^{-1}$ (KIDA) $k_6 = 8.8 \cdot 10^{-10} \text{ cm}^3 \text{ s}^{-1}$ (KIDA)	

3.2. Acetylene as a precursor of polyynes

Acetylene is a parent molecule for different polyynes, cyanopolyynes and methyl-substituted cyanopolyynes. The study of circumstellar envelopes, molecular clouds, hot cores and other objects indicates the presence of polyyne (HC_nH) and cyanopolyyne (HC_nN) chains.^{198,199,200,201,202}

Cernicharo proposed a model²⁰³ in which polyynes, cyanopolyynes, and methyl-polyynes can be formed in PDRs through a network of reactions between $C_n + C_mH_2$, $C_nH + C_mH$, $C_nN + H_2$, $C_nN + C_mH_2$ and others, many of which include acetylene or its derivatives. These reactions could reproduce the observed abundance of polyynes, although Cernicaro emphasized the necessity of both theoretically and experimental checking the parameters of the reactions. The author also suggested that in protoplanetary nebulae such as CRL 618, the large abundance of small cumulenes (H₂C_nH₂) indicates that the growth of small carbon grains with C/H \gg 1 could be dominated by reactions between polyynes and cumulenes.

Reactions such as reactions between the cyan radical CN and acetylene C_2H_2 and its derivative HC_nH have been widely used in astrochemical models of molecular clouds and hot cores to explain HC_nN .^{204,205} It is assumed that the rate constants increase as the temperature decreases. The experiments carried out by Balucani et al. confirmed that some of these reactions (e.g., $C_2H_2 + CN$) may proceed under conditions of circumstellar envelopes in molecular clouds and hot cores.²⁰⁶

In turn, Agundez et al.²⁰⁰ presented a model of the formation of acetylenic chains for the stellar outflow of the C-rich star IRC +10216. This model also includes neutral-neutral reactions between cyan radicals and unsaturated hydrocarbons but is slightly different. According to this model, C_2H_2 and prussic acid (HCN) are formed inside a stellar envelope, while the formation of acetylenic chains occurs further outside the envelope but from the parent molecules C_2H_2 and HCN. First, C_2H_2 and HCN are dissociated by UV photons from the outer radiation field and leave C_2H and CN radicals. The reactions between C_2H and C_2 , C_2H , and C_4H result in C_4H , HC_4H , and HC_6H , respectively. The subsequent photoinduced H loss eventually gives C_4H and C_6H . The reactions between C_2H_2 and its derivatives HC_4H , HC_6H and CN lead to HC₃N, HC₅N and HC₇N, while the subsequent photoinduced H loss from HC₃N and HC₅N provides C_3N and C_5N . The scheme is illustrated in Figure 5.



Figure 5. Pathways from acetylene to polyynes and cyanopolyynes.

Santoro et al.²⁰⁷ experimentally reproduced the conditions of the circumstellar envelopes and studied the chemical evolution of the C_2/C_2H_2 gas mixture. Among the products, polyynes were found, confirming the pathways described above. They emphasized the enormous role of acetylene in the evolution of chemical composition.

In addition to reactions such as $CN + C_2H_2$, which can proceed at low temperature according to Balucani et al.²⁰⁶, there are other barrierless reactions. Gu et al.²⁰⁸ reported that the reaction between ethynyl radical and diacetylene occurs without a barrier, forming triacetylene. Other experiments at low temperature showed that methyltriacetylene ($CH_3(C \equiv C)_3H$) can be generated by a barrierless reaction between diacetylene (HCCCCH) and a 1-propynyl radical (CH_3CC).²⁰⁹ This molecule has been detected at low temperature in the molecular cloud TMC-1²¹⁰; therefore, these reactions most likely take place in such objects and may explain the presence of different polyynes.

Cuylle et al. studied²¹¹ the chemistry of pure C_2H_2 ice under vacuum ultraviolet (VUV) irradiation. Using a combination of infrared and ultraviolet-visible (UV-VIS) spectroscopy, the authors showed the predominance of polymerization and polyyne formation. In this case, the growth of the polyyne chains reached that of $HC_{20}H$ and larger polyyne-like molecules.

The formation of polyyne chains in acetylene ice upon irradiation was also observed in the experiment by G. Compagnini et al.²¹² A total of 2 μ m-thick acetylene C₂H₂ was deposited at 16 K onto a KBr support. Ion irradiation was carried out with an H⁺ beam with an energy of 200 keV. Irradiation of the sample caused the formation of linear carbon chains, the signals of which appeared at 2100, 2170 and 2190 cm⁻¹. The small line widths of these emergent structures indicated that the chains had a distinct length and that they were likely isolated from the rest of the ice matrix. The authors suggested that the chains in question probably contained 8–12 carbon atoms, citing the position of the peak. The results obtained

support the hypothesis that polyynes could be synthesized in comets under the influence of irradiation by CRs and released in gas evaporating from the comet nuclei.

In addition to the neutral carbon chains mentioned above, anions of chains such as C_4H^- , C_6H^- , and C_8H^- were detected in circumstellar envelopes, dark clouds and other objects.^{213,214,215} As electron attachment to C_nH is estimated to be inefficient²¹⁵, it was suggested²¹⁶ that such anionic carbon chains are formed through ion-molecular reactions with acetylene (Scheme 5). The corresponding experiments were carried out, and it was concluded that such reactions could be important for the formation of carbon chain anions.

$$C_{2n}^{-} \xrightarrow{C_2H_2} C_{2n+2}H_m^{-}$$
 (m = 0,1)
 $C_{2n}H^{-} \xrightarrow{C_2H_2} C_{2n+2}H_m^{-}$ (m = 1,2)

Scheme 5. Formation of $C_{2n+2}H_m^-$ anions through ion-molecular reactions with acetylene.

Polyynes and cyanopolyynes can be important components of Titan's haze (see Section 5.4). Kaiser and Mebel, in studies with crossed molecular beams, showed that isoelectronic ethynyl ($C_2H(X_2\Sigma^+)$) and cyano ($CN(X_2\Sigma^+)$) radicals can form on Titan as a result of the photolysis of acetylene (C_2H_2) and hydrogen cyanide (HCN) from solar UV photons.²¹⁷ The data obtained showed that these species, when reacting with unsaturated hydrocarbons (such as acetylene, methylacetylene, and diacetylene), can contribute to the complexation of Titan's haze layers, forming molecules of polyynes and cyanopolyynes.²¹⁷ As shown below in Sections 5.2.4, 5.4 and 8.1, polyynes can participate in PAH formation and in the formation of organonitrogen compounds, which can play an important role in the formation of the topography of Titan and other objects of the Solar system and could play a role in the development of prebiotic chemistry. In Table 7, we summarize the reactions considered in this section.

Reactions	Kinetics	Objects & conditions
Neutral-neutral reactions: 1. $C_2H_2 + CN \rightarrow HC_3N + H$	$k_{I} = 2.38 \cdot 10^{-10} \text{ cm}^{3} \text{ s}^{-1}$ (T=10 K) (KIDA) $k_{I} = 1.42 \cdot 10^{-10} \text{ cm}^{3} \text{ s}^{-1}$ (T = 1000 K) (KIDA)	$\begin{array}{l} \mbox{Stellar envelopes. Conditions:} \\ n_{\rm H} = 10^8 10^{15} \mbox{ cm}^{-3} \\ T = 500 \mbox{-}2500 \mbox{ K} \\ \mbox{Molecular clouds, prestellar} \\ \mbox{cores. Conditions:} \\ n_{\rm H} = 10^2 \mbox{-}10^7 \mbox{ cm}^{-3} \\ T = 10 \mbox{ K} \\ \mbox{G}_0 = \mbox{very low} \\ \mbox{Galactic CR field} \end{array}$

Table 7. The reactions that start from acetylene and lead to formation of polyynes and cyanopolyynes.

- 14			
	2. a. $C_2H_2 + h\nu \rightarrow C_2H + H$ b. $C_2H + C_2H_2 \rightarrow C_4H_2 + H$ c. $C_4H_2 + h\nu \rightarrow C_4H + H$ d. $C_4H + C_2H_2 \rightarrow C_6H_2 + H$ e. $C_6H_2 + h\nu \rightarrow C_6H + H^{[200]}$	$k_{2a} = 4.3 \cdot 10^{-9} \text{ cm}^3 \text{ s}^{-1}$ $k_{2b} = 1.3 \cdot 10^{-10} \text{ cm}^3 \text{ s}^{-1}$ $k_{2c} = 1.1 \cdot 10^{-8} \text{ cm}^3 \text{ s}^{-1}$ $k_{2d} = 3 \cdot 10^{-10} \text{ cm}^3 \text{ s}^{-1}$ $k_{2e} = 1.7 \cdot 10^{-8} \text{ cm}^3 \text{ s}^{-1}$ (all rates are given for T = 100 K, $G_0 = 1)^{[200]}$	PDRs, DISM, molecular clouds, protoplanetary disks. Conditions: T = 10 - 500 K $n_{\text{H}} = 10^{-1} - 10^7 \text{ cm}^{-3}$ $G_0 = \text{from very low to } 10^5$
	3. a. HCN + hv \rightarrow CN + H b. CN + C ₂ H ₂ \rightarrow HC ₃ N + H c. HC ₃ N + hv \rightarrow C ₃ N + H d. C ₃ N + C ₂ H ₂ \rightarrow HC ₅ N + H e. HC ₅ N + hv \rightarrow C ₅ N + H	$k_{3a} = 1.9 \cdot 10^{-9} \text{ cm}^{3} \text{ s}^{-1}$ $k_{3b} = 4.1 \cdot 10^{-10} \text{ cm}^{3} \text{ s}^{-1}$ $k_{3c} = 4.4 \cdot 10^{-9} \text{ cm}^{3} \text{ s}^{-1}$ $k_{3d} = 3.7 \cdot 10^{-10} \text{ cm}^{3} \text{ s}^{-1}$ $k_{3e} = 8.7 \cdot 10^{-9} \text{ cm}^{3} \text{ s}^{-1}$ (all rates are given for T = 100 K, G ₀ = 1) ^[200]	Galactic CR field Titan atmosphere, early Earth ²¹⁷ Conditions: $n_H = 10^{10} \cdot 10^{16} \text{ cm}^{-3}$ $T = 50 \cdot 300 \text{ K}$ G_0 : up to 10 Galactic CR field
	4. $H(C=C)_{2}H$ + $CH_{3}CC \rightarrow$ $CH_{3}(CC)_{3}H^{[209]}$	n.d.	Molecular clouds, prestellar cores. Conditions: $n_{\rm H} = 10^2 \cdot 10^7 \text{ cm}^{-3}$ T = 10 K G_0 is very low Galactic CR field
	5. $C_{2n} + C_2 H_2 \rightarrow C_{2n+2} H_m^- (m = 0,1)$ $C_{2n} H^- + C_2 H_2 \rightarrow C_{2n+2} H_m^- (m = 1,2)^{[216]}$	n.d.	$\begin{array}{l} \mbox{Stellar envelopes. Conditions:} \\ n_{H} = 10^{8} \text{-} 10^{15} \mbox{ cm}^{-3} \\ T = 500\text{-} 2500 \mbox{ K} \\ \mbox{Molecular clouds, prestellar} \\ \mbox{cores.} \\ \mbox{Conditions:} \\ n_{H} = 10^{2} \text{-} 10^{7} \mbox{ cm}^{-3} \\ T = 10 \mbox{ K} \\ \mbox{G}_{0} \mbox{ is very low} \\ \mbox{Galactic CR field} \end{array}$

3.3. Acetylene as a precursor of complex organic molecules

Many COMs are observed in molecular clouds, prestellar cores and protostellar objects.²¹⁸ Among the detected COMs, there are many ethyl- and vinyl-bearing molecules, such as C₂H₅OH, C₂H₃OH, C₂H₅CN, C₂H₃CCCH, and C₂H₃C₃N.¹⁹⁴ Some and perhaps most COMs can form at low temperatures on the ice surface of dust grains or even inside the icy mantle²¹⁹. The interstellar ice mainly consists of H₂O, CO, CO₂, NH₃, CH₄, and CH₃OH.^{220,221} Many different species, including molecules, can be adsorbed on dust grains; these species may lie on or wander the surface and occasionally react with each other or with ice species.^{222,223,224} Energy sources (UV photons, CRs or exothermic reactions) stimulate these reactions. The formed COMs may desorb to the gas phase if enough energy is available, and then, they are observed in radio waves.

Acetylene can be present among species on or inside ice. It can be generated from CH₄, as indicated above in Section 2.6, or trapped from the gas phase. The chemistry of acetylene in interstellar and cometary ices has become the subject of intensive experimental research in recent decades. Experimental studies¹⁶⁰ have shown that acetylene under cold ISM conditions can be the ancestor of a whole family of derivatives, such as diacetylene, vinylacetylene, methylacetylene, and propene, as shown in Figure 6. Specifically, in the experiments, the reactions of acetylene conversion to derivatives were initiated by highly energetic electrons that imitate secondary electrons produced by primary CRs. Furthermore, these derivatives can be the basis for the formation of more complex molecules and carbon nanoparticles (Sections 4, 5, 6).



Figure 6. Experimentally determined¹⁶⁰ dominant reaction pathways for the formation of C_2 -hydrocarbons as well as C_3H_4 , C_3H_6 and C_4H_4 isomers from methane ices.

The transformations of acetylene in ice of various compositions under cryogenic conditions under the influence of radiation were studied by Feldman. In one of his works¹⁶¹, he showed that the decomposition of C_2H_2 in a xenon ice matrix leads to the formation of C_2 species. The clear induction period during which C_2 accumulated indicates that the process proceeded through the decomposition of stabilized C_2H radicals. Conjugated hydrocarbons were detected in acetylene ice in this work, from which the authors concluded that condensation predominates over decomposition during radiation treatment of acetylene ice. This finding is consistent with the findings of other works mentioned above and below.²¹¹

Reactions between acetylene and other species on/in ice may lead to the formation of COMs. One of the COMs, ketene (CH₂CO), was experimentally produced from acetylene-containing ice in a number of works. Specifically, ketene was formed as a result of photolysis and radiolysis of ice mixtures.^{224,,225} Several low-temperature routes involving acetylene and oxygen were investigated experimentally.^{226,227} Notably, these experiments showed that ketene is formed in an ice mixture of C_2H_2 and CO_2 at a temperature of 10 K. First, CO₂ dissociates to CO and O, and then O reacts with C_2H_2 to form ketene. These reactions can explain the observed presence of ketenes in molecular clouds.²²⁸

The interaction of acetylene with a water molecule in an ice matrix at a temperature of 5 K under radiation conditions was experimentally studied by Feldman et al.²²⁹ The authors showed that the formation of a complex with water facilitates the radiation decay of the acetylene molecule. FTIR was used to detect ketene, which is a complex with hydrogen (CH₂CO–H₂), ketenyl radicals (HCCO), methane (CH₄) and carbon monoxide (CO). Such processes can occur in interstellar and cometary ices.

Irradiation of H_2O/CO_2 ices with acetylene also leads to the formation of other (semi)saturated hydrocarbons, such as C_2H_4 , C_2H_6 and other COMs (vinyl alcohol [CH₂CHOH], acetaldehyde [CH₃CHO], ketene [CH₂CO], and ethanol [CH₃CH₂OH]).^{154,230,231,232} Bergner et al.^{225,226,233} suggested that ethylene and ethane are formed by subsequent hydrogenation of acetylene (Scheme 6). The hydrogenation of acetylene to ethane was proven to be efficient under conditions of amorphous solid water in the environment of a dark molecular cloud.¹⁵⁸

Α	C_2H_2	+	н		C ₂ H ₃
В	C_2H_3	+	н	\longrightarrow	C_2H_4
С	C_2H_4	+	н	\longrightarrow	C ₂ H ₅
D	C_2H_5	+	н	>	C_2H_6

Scheme 6. Ethylene, ethane formation by subsequent hydrogenation of acetylene in the ices.

Charnley²³⁴ demonstrated how the combination of hydrogenation and oxidation in H_2O/C_2H_2 ice leads to the formation of vinyl alcohol, acetaldehyde, and ethanol (Scheme 7).



Scheme 7. Formation of formation of vinyl alcohol, acetaldehyde, and ethanol by combination of hydrogenation and oxidation of acetylene in H_2O/C_2H_2 ice.

Hudson et al. suggested that ethanol is formed in an ice mixture of C_2H_2 and H_2O via the following sequence: A) radiolysis of water; B-D) hydrogenation of C_2H_2 to C_2H_5 ; and E) addition of an OH radical to C_2H_5 with C_2H_5OH formation (Scheme 8).

 \rightarrow Hⁱ + OHⁱ by p⁺ radiolysis Α H₂O HC≡CH + - H₂C=CH Β н \rightarrow H₂C=CH₂ С $H_2C=CH' + H'$ → H₃C-CH₂ $H_2C=CH_2 + H$ D Ε H_3C-CH_2 + οн C₂H₅OH

Scheme 8. Ethanol formation in the ices.

According to Chuang et al. 2021, the produced COMs (acetaldehyde, ketene, ethanol) can be explained by reactions that start from the formation of vinyl alcohol in the reactions between C_2H_2 and H and/or OH radicals. Further transformation to acetaldehyde, ketene, or ethanol occurs either by

hydrogenation or energy absorption. These COMs were also produced via a nonenergetic route in the experiment where C_2H_2 , O_2 and H were codeposited on the surface and were not irradiated.²³⁵

Using DFT calculations, Perrerro et al.²³⁶ showed that ethanol can be formed by the reaction of C_2H (an acetylene derivative) with one H_2O molecule. It was assumed in the model that the reaction proceeded in clusters of water ice. Such a process can be barrierless due to the possibility of proton transfer on the boundary molecules of ice water.

Zhou et al. studied²³⁷ the transformation of the ice mixture CO/C₂H₂ at 10 K under high-energy electron irradiation in an ultrahigh vacuum. As a result, two isomers of the C₃H₂O molecule could be detected. Propynal (HCCCHO) was synthesized from a carbon monoxide-acetylene complex via the [HCO... CCH] radical pair. In turn, the addition of triplet carbon monoxide to acetylene in the ground state (or vice versa) occurred with the formation of a cyclic structure (c-C₃H₂O). The authors noted that the heterogeneous formation of both C₃H₂O isomers was orders of magnitude greater than that of gaseous chemistry. The production rates were 5 O 10⁶ molecules cm⁻²s⁻¹ for propynal species (3 O 10⁴ molecules cm⁻²s⁻¹ in gaseous chemistry) and 5 O 10⁶ molecules cm⁻²s⁻¹ for c-C₃H₂O. Thus, the process of formation of C₃H₂O isomers from acetylene can be initiated by energetic electrons produced by CRs penetrating interstellar ice grains in cold molecular clouds.

Wu et al. studied²³⁸ acetylene in an environment of solid nitrogen (N₂) at a temperature of 10 K. Ice was irradiated by VUV photons and electrons. In the first case, the formation of C₂H, CN radicals and C₂N₂ isomers was observed. In the second case, the formation of N₃ and C₂H radicals, as well as various nitriles, was observed. Photolysis of C₂H₂/N₂ ice samples produced 15 times more C₂H radicals than did radiolysis.

The transformation of acetylene in ammonia ice was also studied experimentally. Namely, ice mixtures of NH₃:C₂H₆, NH₃:C₂H₄ and NH₃:C₂H₂ were irradiated by VUV photons at temperatures ranging from 10–50 K. As a result, the formation of imine (CH₃CH=NH) and acetonitrile (CH₃CN) was detected. The highest yields of imine (35%) and acetonitrile (5.2%) were achieved by irradiating the mixtures with C₂H₂ at 10 K.²³⁹ An important discovery was made by Zhang et al.²⁴⁰, who experimentally demonstrated the formation of vinylamine (C₂H₃NH₂) in the ice mixture NH₃:C₂H₂, which was exposed to energetic electrons. Quantum chemical calculations have shown that vinylamine (C₂H₃NH₂) can be formed in two different ways: through barrierless radical-radical recombination of amino (NH₂) and vinyl (C₂H₃) radicals and through a one-step concerted pathway.



Figure 7. Proposed reaction network for the irradiation of C₂H₆, C₂H₄, C₂H₂ and ammonia ice mixtures.

Recently, Molpeceres and Rivilla²⁴¹ computationally (DFT) studied the surface reactions between acetylene, ethylene and ethane and radicals (H, OH, NH₂, CH₃). The reactions with the radicals H and OH were found to be the most important for COM formation, namely, for C_2H_3 , C_2H_5 , C_2H_2OH and C_2H_4OH , while the reactions with NH₂ and CH₃ had a minor role. These formed radicals can further participate in reactions leading to more complex molecules, including those detected in space, although certain pathways have still not been established.

Another example of the likely interaction of acetylene with ammonia is Jupiter's Great Red Spot (GRS).²⁴² In laboratory simulations,²⁴³ a material was obtained by reacting acetylene and ammonia under VUV irradiation, corresponding to solar radiation on Jupiter within and above the upper troposphere. HR-MS and IR spectroscopy confirmed the presence of aliphatic azine, azo, and diazo compounds in the resulting chromophore material. The spectral properties of the resulting chromophore compound were in excellent agreement with the 0.35–1.05 μ m spectra of the core of the GRS obtained by VIMS during the Cassini–Huygens mission. In the high troposphere, upwelling NH₃ molecules can photodissociate and react with downwelling C₂H₂ molecules. The resulting compounds can coat ammonia grains at the top of the GRS clouds. Tholins, which may be products of the reaction of ammonia and acetylene, may also be responsible for the orange-red color of objects on a number of planets and moons in the Solar system (see Section 5.4). More details about the nature of the tholins and the chemical mechanism of action of acetylene with ammonia in the formation of important nitrogen-containing compounds are discussed in Section 8.1. In Table 8, we present the main reactions discussed in the section, although kinetics information is not yet available.

Reactions	Objects & conditions
CO ₂ /C ₂ H ₂ ice 1. a CO ₂ \rightarrow CO +O b. C ₂ H ₂ + O \rightarrow CH ₂ CO [[] 226·227·234 []]	Molecular clouds, prestellar cores. Conditions: T = 10 K G ₀ = very low Galactic CR field
$\begin{aligned} \mathbf{H_2O/C_2H_2 ice} \\ 2. a. H_2O \rightarrow H \cdot + OH \cdot \\ b. C_2H_2 + H \cdot \rightarrow C_2H_3 \\ c. C_2H_3 + H \cdot \rightarrow C_2H_4 \\ d. C_2H_4 + H \cdot \rightarrow C_2H_5 \\ e. C_2H_5 + H \cdot \rightarrow C_2H_6 \\ e. C_2H_5 + OH \cdot \rightarrow C_2H_5OH^{[230]} \end{aligned}$	
3. a. $C_2H_2 + CR$ (or hv) $C_2H + H$ b. $C_2H + H_2O \rightarrow HCCHOH$ c. $HCCHOH + H \rightarrow CH_3CHOH$ d. $CH_3CHOH + H \rightarrow C_2H_5OH^{[236]}$	
4. a. $C_2H_2 + OH$ (from H_2O) CH_2CHOH (vinyl alcohol) b. $CH_2CHOH \rightarrow CH_3CHO$ (acetaldehyde) c. $CH_2CHOH + CR/hv \rightarrow CH_2CO$ (ketene) d. $CH_2CHOH + 2H \rightarrow CH_3CH_2OH$ (ethanol) ^[225]	
5. a. $C_2H_2 + H \cdot \rightarrow CH_2CH$ b. $CH_2CH + O \rightarrow CH_2COH$ c. $CH_2COH + H \cdot \rightarrow CH_2CHOH$ d. $CH_2CHOH + H \cdot \rightarrow CH_3CHOH$ e. $CH_3CHOH + H \cdot \rightarrow C_2H_5OH^{[234]}$	
6. a. $C_2H_2 + H \cdot \rightarrow CH_2CH$ b. $CH_2CH + O \rightarrow CH_2CHO$ c. $CH_2CHO + H \cdot \rightarrow CH_3CHO$ d. $CH_3CHO + H \cdot \rightarrow CH_3CHOH$ e. $CH_3CHOH + H \cdot \rightarrow C_2H_5OH^{[234]}$	
7. a $C_2H_2 + H \cdot \rightarrow C_2H_3$ b. $C_2H_2 + OH \cdot \rightarrow C_2H_2OH$ c. $C_2H_4 + H \cdot \rightarrow C_2H_5$ d. $C_2H_4 + OH \cdot \rightarrow C_2H_4OH^{[241]}$	

Table 8. The reactions of formation of COMs with acetylene as a key reactant.

4. Benzene formation as the first step in PAH evolution

The formation of an aromatic ring is a turning point in chemical evolution from acetylene. This is the first step toward complex PAH molecules. The initial molecules are benzene C_6H_6 and the phenyl radical C_6H_5 , which further act as elementary building blocks for all other high-molecular-weight polycyclic aromatic systems and carbon nanoparticles as well as soot particles and carbon dust. Unlike C_2H_2 , benzene was found only in a few objects, in the protoplanetary nebulae CRL 618¹⁴³ and SMP LMC 11²⁴⁴ and in the protoplanetary disk J160532.¹⁸⁸ Additionally, benzonitrile was detected in the TMC-1 molecular cloud.²⁴⁵ Nevertheless, benzene is present in space objects and plays an important role in chemical reaction networks. The formation of the initial aromatic ring is considered a limiting step that determines the rate of formation of larger polycyclic aromatic systems and particles.^{20,246} This process has even been named a "bottleneck" in macromolecular growth and dust nucleation.³⁸ Moreover, it is important to note that the transition of hydrocarbons from an aliphatic structure to a closed aromatic structure ensures that the system achieves the minimum energy. That is, the process of formation of the benzene ring is thermodynamically determined. Despite its importance, the mechanism of formation of the first aromatic system from aliphatic precursors in various regions of space is still the subject of discussion.

Traditionally, acetylene was considered to be the main precursor of aromatic molecules. Chemical mechanisms for the formation of an aromatic system in outer space were proposed in the 1980s, when the mid-IR interstellar emission bands were interpreted as related to PAH molecules.²⁴⁷ The conditions differ significantly for different space objects; therefore, there are a variety of mechanisms that are determined by the specific physical conditions and chemical composition. Nevertheless, the possible formation of aromatic molecules can be related to two distinct environments: a cold and dense environment, which includes molecular clouds, planets and moons; and a hot/warm environment, which includes stellar envelopes and their remnants, protoplanetary and planetary nebulae. Furthermore, the main pathways are presented. The mechanisms proposed in astrophysics traditionally do not imply the participation of transition metals and almost do not consider catalytic processes. These mechanisms will be discussed in a separate section.

4.1. Circumstellar envelopes and flame burning models

As one of the main sources of cosmic carbon, C-rich AGB stars have attracted the attention of researchers as the basis for models of aromatic ring formation. Several mechanisms for the formation of the benzene ring have been proposed for astrochemical models from the chemistry of flame combustion.^{248,249,250,251} In 1984, Cole et al. suggested that the main mechanism for the formation of aromatic compounds in a flame is the reaction of 1,3-butadienyl with acetylene (Scheme 9A).²⁵² Recently, Frenklach et al. described the main cyclization pathways in a low-pressure acetylene flame soot simulation study. According to their mechanism, the formation of a phenyl radical occurs upon the interaction of 1-buten-3-ynyl with an acetylene molecule through the cyclization of linear C₆H₅ (Scheme 9B).^{253,254} In 1992, Miller and Melius first proposed a mechanism in which ring closure occurs after the reactions of two propargyl radicals as the main pathway for the formation of the benzene ring (Scheme 9C).²⁵⁵



Scheme 9. Various schemes for the formation of an aromatic ring at hot temperature.

Subsequently, Cherchneff noted that the propargyl cyclization pathway is dominant, while Cole's and Frenklach's mechanisms are minor; moreover, these pathways will be realized if the reactants are present in the circumstellar gas.²⁴⁶ The formation of benzene as a result of barrierless self-recombination of the propargyl radical under conditions simulating combustion and in a circumstellar environment using a high-temperature chemical reactor was confirmed in a study by Zhao et al. 2021.²⁵⁶

Regardless of the underlying mechanism, acetylene is the main precursor that initiates this reaction. The formation of the propargyl radical occurs as a result of the reaction of acetylene (C_2H_2) and methylene (CH_2). Other agents originate from vinylacetylene (C_4H_4) by the addition of atomic hydrogen to form 1,3-butadienyl (C_4H_5) or its elimination to form 1-buten-3-ynyl (C_4H_3). In this case, the precursor of vinylacetylene (C_4H_4) is also acetylene.

Other possible mechanisms for the formation of the first aromatic ring have been described.²⁵⁷ For example, in addition to the recombination of propargyl radicals, the reaction between vinylacetylene (C_4H_4) and vinyl radical (C_2H_3) promoted the formation of benzene.^{258,259} Another interesting route is the formation of benzene via cyclopentadienyl (c- C_5H_5), which is generated during the reaction of acetylene with propargyl.²⁶⁰ Lin et al. showed that the reaction of methyl and cyclopentadienyl radicals is a potential source of benzene at high temperatures.^{261,262} This pathway combines the advantages of a highly stable propargyl radical and acetylene as a widely available starting building block.

The main reactions of benzene formation are given in Table 9. It should be noted that the mechanisms presented above, based on the analysis of combustion, control aromatic chemistry at

temperatures of approximately 1000 K and are not applicable to cold interstellar environments (~ 10 K)²⁶³ or to atmospheres of planets and satellites (~ 90 K temperature of the Titan surface, for example²⁶⁴). In addition, these reaction systems do not consider the effects of UV radiation or ion-molecular chemistry.

Reactions	Kinetics	Objects & conditions
1. $C_4H_5 + C_2H_2 \rightarrow C_6H_6 + H^{[252]}$	$k_1 = 5.3 \cdot 10^{-13} \text{ cm}^3 \text{ s}^{-1} (\text{T} = 1000 \text{ K})^{[252]}$	Dust formation zone in stellar envelopes.
2. a. $C_2H_2 + M \rightarrow H_2C_2$ b. $H_2C_2 + C_2H_2 \rightarrow C_4H_4$ c. $C_4H_4 + H \rightarrow C_4H_3 + H_2$ d. $C_4H_3 + C_2H_2 \rightarrow C_6H_6^{[254]}$	$k_{2a} = 2.6 \cdot 10^{-16} \text{ cm}^3 \text{ s}^{-1}$ $k_{2b} = 1.9 \cdot 10^{-12} \text{ cm}^3 \text{ s}^{-1}$ $k_{2c} = 1.4 \cdot 10^{-12} \text{ cm}^3 \text{ s}^{-1}$ $k_{2d} = 2.5 \cdot 10^{-14} \text{ cm}^3 \text{ s}^{-1}$ (all rates are given for T = 1000 K) ^[254]	Conditions: $n_{\rm H} = 10^8 \cdot 10^{12} {\rm cm}^{-3}$ $T = 500 \cdot 2000 {\rm K}$
$\begin{array}{c} 3.\\ C_3H_3 \cdot + C_3H_3 \cdot \rightarrow C_6H_5 + H \cdot \end{array}$	$k_3 = 5.65 \cdot 10^{-11} \text{ cm}^3 \text{ s}^{-1} \text{ [265]}$	

Table 9. The reactions of benzene formation which start from acetylene or its derivatives.

4.2. Benzene formation in protoplanetary nebulae

The presence of benzene as well as di- and triacetylene chains was recorded in protoplanetary nebulae²⁶⁶; therefore, it was proposed that these chains can be formed in such environments. These objects represent the next and short stage after an AGB star's death. In the literature, they are also called post-AGB stars. The fusion process in these stars stops, but the surrounding shell still has a high temperature (several hundred kelvins) and is relatively dense (~ 10^7 cm⁻³). Additionally, the UV radiation intensity from a star gradually increases and may exceed the mean interstellar intensity G₀ by 5 orders of magnitude. This factor can play an important role in the activation of ion-molecular chemical reactions. To the best of the authors' knowledge, to date, a chain of ion-molecular reactions leading to benzene formation has been suggested only in the work of Woods et al.⁵⁰ They presented a gas-phase chemical model of the protoplanetary nebula CRL 618 and showed that the efficient production of benzene is associated with a high flux of ionizing radiation. Woods et al.⁵⁰ reported that HCO⁺ species can transfer a proton to the C₂H₂ parent molecule, triggering a cascade of ionic-molecular transformations. The successive addition of acetylene molecules to cations leads first to a phenyl cation and then to a benzene molecule through the addition of hydrogen and a dissociative reaction (Figure 8). However, this chain of reactions has not been studied either experimentally or theoretically and therefore requires additional approval.



Figure 8. Proposed synthesis of benzene in protoplanetary nebula CRL 618.⁵⁰ Gray color is C atoms, white is hydrogen, and red is oxygen.

Reactions	Kinetics	Objects & conditions
a. $HCO^+ + C_2H_2 \rightarrow C_2H_3^+ + CO$ b. $C_2H_3^+ + C_2H_2 \rightarrow C_4H_3^+ + H_2$ c. $C_4H_3^+ + C_2H_2 \rightarrow cC_6H_5^+$ d. $cC_6H_5^+ + H_2 \rightarrow cC_6H_7^+$ e. $cC_6H_7^+ + e \rightarrow C_6H_6 + H^{[50]}$	$k_{a} = 1.4 \cdot 10^{-9} \text{ cm}^{3} \text{ s}^{-1} \text{ (KIDA)}$ $k_{b} = 7.2 \cdot 10^{-10} \text{ cm}^{3} \text{ s}^{-1} \text{ (KIDA)}$ $k_{c} = 1.4 \cdot 10^{-9} \text{ cm}^{3} \text{ s}^{-1} \text{ (T=250 K) (KIDA)}$ $k_{d} = 6.0 \cdot 10^{-11} \text{ cm}^{3} \text{ s}^{-1} \text{ (T=250 K) (KIDA)}$ $k_{e} = 8.7 \cdot 10^{-7} \text{ cm}^{3} \text{ s}^{-1} \text{ (KIDA)}$	Protoplanetary nebulae. Conditions: T = 250 K $n_H = 10^7 \text{ cm}^{-3}$ $G_0 = 2 \cdot 10^5$, but suitable for similar PDR conditions

Table 10. The chain of ion-molecular reactions which lead to formation of benzene.

4.3. Benzene formation in cold interstellar medium

Jones et al. analyzed the above-described mechanisms based on neutral–neutral reactions from combustion chemistry and ion–molecular mechanisms suitable for AGB and post-AGB stars, respectively, and concluded that these reactions cannot lead to the formation of benzene under conditions of low temperatures of 10 K and pressure in cold molecular clouds and prestellar cores, where the density of matter is 10^{2} - 10^{7} cm⁻³.^{21,267} As a result, Jones et al., based on experiments with a crossed molecular beam in combination with statistical calculations and DFT calculations of the electronic structure, concluded that benzene can be synthesized through a barrierless exoergic reaction of an ethynyl radical and 1,3-butadiene under single collision conditions (Scheme 10).²¹ The authors proposed the following sequence of reactions: 1) ethane is formed as a result of the addition of a hydrogen atom to acetylene or reactions between fragments of the methyl radical (CH₃); 2) the fast reaction of methylidyne radicals (CH) with ethane (C₂H₆) leads to the formation of propene (C₃H₆) and atomic hydrogen; 3) then, propene reacts rapidly with another methylidyne radical, forming 1,3-butadiene (C₄H₆); and 4) in the last step, the ethynyl radical and 1,3-butadiene to form an acyclic isomer.



Scheme 10. Proposed mechanism of benzene formation through a barrierless exoergic reaction of an ethynyl radical and 1,3-butadiene.

Ethane is unlikely to form under gas chemistry conditions. Most likely, its synthesis occurs on dust particles inside the icy mantle from acetylene. Such reactions involving the formation of ethane were described earlier in experiments simulating the conditions of interstellar ice: acetylene hydrogenation of methyl radical recombination.^{159,268} Additionally, high ethane abundance has been detected for the Oort cloud comets by IR observations.¹²¹ When formed in ice^{159,268}, ethane can be released into the gas phase as a result of the thermal desorption of the icy mantle. However, this process occurs only at temperatures higher than ~60 K¹⁵⁹, and it is relevant for hot core objects or comets approaching perihelion. In the case of cold clouds, an icy mantle can be sublimated via nonthermal desorption due to ion bombardment and grain–grain collisions during the propagation of shock waves.^{269,270}

Extensive experiments on barrierless reactions involving the formation of benzene or related molecules have been conducted. Independent experimental evidence for a one-step radiation-induced formation of benzene from isolated trimeric acetylene was presented in a recent study by Lukianova and Feldman.²⁷¹ Using Fourier IR spectroscopy, the authors showed that the decomposition of the acetylene trimer and the accumulation of benzene occur in a frozen inert medium at a very low temperature (6 K). Thus, the $(C_2H_2)_3$ trimer is an ideal precursor for the formation of benzene molecules due to the geometric correspondence between the trimer structure and the shape of the benzene molecule. It is assumed that the formation of benzene in cryogenic films may be related to the cationic mechanism known in the gas phase. This mechanism implies ionization of the trimer followed by almost barrier-free transformation into a stable benzene radical cation. The described mechanism of radiation-induced benzene production without any significant contribution from molecular diffusion could be considered an interesting route for the formation of aromatic rings under cold space conditions. However, the likelihood of the C_2H_2 trimer existing in space conditions is relatively low.

In addition to benzene, benzyl radicals and *ortho*-benzynes can form.^{272,273} These compounds may further form benzene in subsequent reactions. These reactions have also been investigated with quantum mechanical calculations.^{274,275,276} Thus, benzene has been established to be formed not only by fusion but also at very low temperatures.

Reactions	Kinetics	Objects & conditions
1. a. $C_2H_2 + 4H \rightarrow C_2H_6$ (surface reaction) b. $C_2H_6 + CH \rightarrow C_3H_6$ c. $C_3H_6 + CH \rightarrow C_4H_6$ d. $C_4H_6 + C_2H \rightarrow C_6H_6^{21}$	$k_{a} = -$ $k_{b} = 5.8 \cdot 10^{-13} \text{ cm}^{3} \text{ s}^{-1}$ (KIDA) $k_{c} = 3.6 \cdot 10^{-10} \text{ cm}^{3} \text{ s}^{-1}$ (KIDA) $k_{d} = 3.0 \cdot 10^{-10} \text{ cm}^{3} \text{ s}^{-1} \text{ [21]}$	Molecular clouds, prestellar cores. Conditions: T = 10 K $G_0 = very low$ Galactic CR field
2. Trimerization $3C_2H_2 + CR \rightarrow C_6H_6^{271}$	n.d.	

Table 11. The reactions of benzene formation in cold interstellar medium.

4.4. Benzene formation in planets and moons

Benzene has been observed in the atmospheres of Jupiter, Saturn and Titan by ISO and interplanetary spacecrafts such as Voyager and Cassini.²⁷⁷ The paths from small hydrocarbons to benzene in the atmospheres of the giant planets and their satellites are assumed to be similar to the paths considered. Wong et al. proposed a model for predicting the formation of benzene and PAHs in the atmosphere of Jupiter.²⁷⁸ Wong et al. concluded that ion–molecular reactions play the most important role, so the model considers a system of ion–molecular processes and neutral chemistry (Scheme 11). The model included 288 reactions of ion-neutral exchange and 79 reactions of electron-ion recombination. The system of transformations begins with the destruction of methane by energetic particles, in addition to the reactions of acetylene. Neutral chemical reactions contribute only approximately a few percent of the total reactions. Among them are the most important C_3H_3 recombinations and the addition of C_2H_2 to C_4H_3 . The described chemical processes, as the authors suggest, may also be relevant for other giant planets, such as Saturn, and extrasolar giant revolves close to their main orbit and receive high doses of far-UV radiation.



Scheme 11. The most important reaction pathways for the formation of benzene in Jupiter according to the model of Wong et al.²⁷⁸ One of the main chains of reactions designated by red arrows.

Wilson et al. (2003)²⁷⁹, Lebonnois (2005)²⁸⁰, and Vardavas (2008)^{109,281} explored possible ways for benzene to form in Titan's atmosphere (Figure 9). These pathways include 1) recombination of two propargyl radicals, 2) reaction between acetylene and butadiene radical and 3) subsequent reactions of diacetylene with hydrogen and acetylene. Thus, in these models for the conditions of planetary atmospheres and their satellites, the same reactions play a key role as in the models of benzene formation at temperatures of 1000 K and above.

The photochemical model proposed by Vardavas et al.¹⁰⁹ suggested that three-body reactions are the main route for benzene production. However, the predictions of this model give more than a twofold underestimation of the benzene abundance in the thermosphere of Titan. In addition, given the planet's rarefied thermosphere, three-body reactions are unlikely. In addition, the abundance of benzene in the thermosphere of Titan is greater than the abundance of benzene in the stratosphere.^{282,283}


Figure 9. Schematic diagram of the mechanisms of formation of benzene according to the model of Wilson et al.²⁷⁹ M is a third neutral molecule required for energy balance in the reactions.

Based on the analysis of altitude data, Vuitton et al.²⁸⁴ concluded that benzene is synthesized precisely in the thermosphere and is not the result of diffusion processes from the stratosphere, where the mole fraction of benzene is lower (Scheme 12).¹⁰⁴ Vuitton et al. noted that the rate of neutral-neutral reactions is not enough to form a sufficient amount of benzene; therefore, a model based on the chemistry of ion-molecular reactions was proposed for the upper atmosphere (thermosphere and ionosphere) of Titan. Acetylene plays an important role in this model and is responsible for the formation of benzene precursors such as $C_4H_5^+$ and $C_6H_5^+$.²⁸⁵



Scheme 12. The main ion-chemical reactions of the formation of benzene in the atmosphere of Titan.²⁸⁴

Among other substances, benzene can be formed directly from acetylene on the surface of Titan. Experiments on the irradiation of acetylene ices at low temperature with energetic electrons have shown that CR-mediated production of benzene directly from three acetylene molecules on the surface of Titan may be the dominant source of benzene.²⁸⁶

The two singlet ground-state acetylene molecules do not react with each other since the entrance barrier to the reaction is too large and equals to 140 kJ·mol⁻¹. However, the calculations showed that the acetylene molecule in the first excited triplet state C_2H_2 (a^3B_2) can indeed easily (the barrier value is only 4 kJ mol⁻¹) react with the acetylene molecule in the ground state to form the trans- C_4H_4 structure [1] (Figure 10). The resulting molecule can undergo trans-cis isomerization and form [2]. In turn, the triplet structure [2] can react with the second acetylene molecule in the ground state to form a structure [3]. The reaction of cis-CHCHCHCH [2] and C_2H_2 ($X^1\Sigma_g^+$) can be possible even at low temperatures if [2] retains enough energy gained from the initial interaction of two acetylene molecules. Then, the intermediate triplet can easily rearrange into a structure [4], followed by ring closure to triplet benzene [5].



Figure 10. Part of the potential energy surface of the C_6H_6 , triplet involved in the formation of the benzene triplet through reactions with excited acetylene. Reproduced with permission from ref 284. Copyright 2010 The American Astronomical Society.

In their experimental study, Kleimeier et al.²⁸⁷ showed that the photolysis of solid acetylene by low-energy UV photons can lead to the formation of benzene. The authors suppose that this mechanism is the main mechanism for Pluto's atmosphere and other outer planets where acetylene is very abundant according to New Horizons's observations.

The reactions discussed in this section are given in Table 12. We emphasize that over the past few decades, ideas about the mechanisms of synthesis of the first aromatic ring have expanded from models borrowed from combustion chemistry for high-temperature environments to experiments with crossed molecular beams for barrierless reactions under single-collision conditions in cold interstellar environments. Moreover, for almost all the proposed mechanisms, acetylene, as a key building block, takes part, if not directly, in the mechanism of formation of the aromatic ring and then as a precursor of the initial reactants. In this regard, acetylene molecules are exemplary due to their structure, properties, and wide distribution in various cosmic environments, which facilitates their participation in a wide variety of reactions.

Reactions	Kinetics	Objects & conditions
Neutral-neutral reactions 1. $C_3H_3 + C_3H_3 \rightarrow C_6H_5 + H$	$k_1 = 4.98 \cdot 10^{-12} \text{ cm}^3 \text{ s}^{-1} \text{ [278]}$	Planetary atmospheres. Conditions: $n = 1000 \pm 1006 \text{ am}^3$
$2. C_3H_3 + C_3H_3 + M \rightarrow C_6H_6$	$k_2 = 2.1 \cdot 10^{-23} \text{ cm}^6 \text{ s}^{-1}$ (T = 150 K, n _H = 10 ¹² cm ⁻³) ^[280]	T = 50-300 K $G_0 = \text{up to } 10^2$ Galactic CP field
$3. C_2H_2 + C_4H_3 + M \rightarrow C_6H_5$	$k_3 = 2.1 \cdot 10^{-29} \mathrm{cm}^6 \mathrm{s}^{-1} (150 \mathrm{K})^{[278]}$	Galactic CK netu
4. a. $C_2H_2 + h_v \rightarrow C_2H + H$ b. $C_2H_2 + C_2H \rightarrow C_4H_2 + H$ c. $C_4H_2 + H + M \rightarrow C_4H_3$ d. $C_4H_3 + C_2H2 \rightarrow C_6H_4 + H$ e. $C_6H_4 + H + M \rightarrow C_6H_5$ f. $C_6H_5 + H + M \rightarrow C_6H_6$	$k_{4d} = 6.3 \cdot 10^{-20} \text{ cm}^3 \text{ s}^{-1} (150 \text{ K})$ $k_{4f} = 1.7 \cdot 10^{-22} \text{ cm}^6 \text{ s}^{-1} (150 \text{ K})^{[336]}$	
5. $C_4H_5 + C_2H_2 \rightarrow C_6H_6 + H$ (Titan)	$k_5 = 2.9 \cdot 10^{-19} \text{ cm}^3 \text{ s}^{-1} (150 \text{ K})^{[278]}$	
Ion-molecular reactions 6. a. $C_2H_3^+ + C_2H_2 \rightarrow C_4H_3^+ + H_2$ b. $C_4H_3^+ + C_2H_2 \rightarrow cC_6H_5^+$ c. $cC_6H_5^+ + H_2 \rightarrow cC_6H_7^+$ d. $cC_6H_7^+ + e \rightarrow C_6H_6 + H$	$k_{6a} = 2.16 \cdot 10^{-10} \text{ cm}^3 \text{ s}^{-1}$ $k_{6b} = 2.2 \cdot 10^{-10} \text{ cm}^3 \text{ s}^{-1}$ $k_{6c} = 6.0 \cdot 10^{-11} \text{ cm}^3 \text{ s}^{-1}$ $k_{6d} = 7.1 \cdot 10^{-7} \text{ cm}^3 \text{ s}^{-1} \text{ [278]}$	
7. a. $C_2H_4^+ + C_2H_2 \rightarrow C_4H_5^+ + H$ b. $C_4H_5^+ + C_2H_4 \rightarrow C_6H_7^+$ c. $C_6H_7^+ + e \rightarrow C_6H_6 + H^{[284]}$	$k_{7a} = 1.9 \cdot 10^{-10} \text{ cm}^3 \text{ s}^{-1}$ $k_{7b} = 7.4 \cdot 10^{-11} \text{ cm}^3 \text{ s}^{-1}$ $k_{7c} = 5.9 \cdot 10^{-6} \text{ cm}^3 \text{ s}^{-1} (150 \text{ K})^{[284]}$	
Trimerization: 8. $3C_2H_2 + CR \rightarrow C_6H_6$	$k_8 = 3-5 \cdot 10^{-3} \text{ mol eV}^{-1} [286]$	Titan's surface. Conditions: T~100 K, Galactic CR field

Table 12. Summary of considered reactions, kinetics (if available) and conditions where the reactions are relevant

5. Mechanisms for the Formation of Polyaromatic Compounds in Space

5.1. PAHs in space and their role in carbon evolution

The formation of the first aromatic ring is the first step in the growth of a large number of polycyclic aromatic systems. PAHs are responsible for unidentified infrared bands (UIB) in the range from 3 to $20 \ \mu m.^{288,289,290}$ The main bands are positioned at 3.3, 6.2, 7.7, 8.6, and 11.2 μm . These bands appear in most galactic and extragalactic sources²⁹¹. The peak wavelengths and relative strengths of the bands can vary from object to object; nevertheless, the spectra are similar and indicate the ubiquitous presence of PAHs in the ISM. PAHs, their hydrogenated, dehydrogenated²⁹², protonated²⁹³, alkylated^{294,295}, ionized²⁹⁶ derivatives, fullerenes and other related carbonaceous molecules are considered promising candidates for so-called diffuse interstellar bands, which are discrete absorption features superimposed on the interstellar extinction curve in the range from the blue region of visible light (400 nm) to near IR (1200 nm).^{297,298} The

extinction curve in the UV range has a prominent bump at 2175 Å, which is most likely related to electronic transitions in PAHs or related molecules. Moreover, the significant extinction in the UV range and its increase at wavelengths down to 0.1 µm indicate that a large number of small species are comparable to PAH sizes. Other observational features, such as extended red emission and anomalous microwave emission, can also be mentioned as signatures of PAH presence in the ISM, although decoding of these signatures is complicated.²⁹⁹ Despite a great number of observations, specific interstellar PAHs have been scarcely identified. The most precise identification of interstellar PAHs to date involved the discovery of cyanonaphtalene³⁰⁰ as well as indene³⁰¹. PAHs have been identified in several dozen carbonaceous chondrites, such as Murchison, Allende, and Orgueil^{18,302,303,304,305,306,307,308}, and their supposed structures are illustrated in Figure 11.



Figure 11. Possible structures of aromatic molecules found in carbonaceous meteorites.

PAHs and carbon dust are found throughout the cosmos: in circumstellar media; in planetary and reflection nebulae³⁰⁹; in molecular clouds; in interstellar, interplanetary and intergalactic media³¹⁰; and finally in the atmosphere of planets and in meteorites. According to different estimates, up to 20% of carbon can be locked in the form of PAHs, the size of which can exceed 50 carbon atoms.^{60,291,311,312,313,314} These molecules play an important role in the ISM, determining its thermal and ionization balance.¹⁷⁰ PAHs are intermediates between low-molecular-weight hydrocarbons and carbon particles and provide nucleation sites for the formation of carbon dust particles.³¹⁵ Tielens noted⁴⁹ that PAH molecules influence key processes in the evolution and characteristics of interstellar media and galaxies through interactions with gases, photons and energetic ions; provide active surfaces for chemical processes; participate in the regulation of opacity; and influence the heating and cooling of neutral atomic and molecular gases.

PAHs can participate in complex networks of transformations that occur in stellar ejecta and even play the role of catalysts, thereby affecting the overall chemical composition of star formation and planet formation regions.^{291,316} The most impressive example of using PAHs as catalysts is the formation of hydrogen molecules.³¹⁷ In the atmospheres of planets and satellites, PAHs are responsible for the formation of atmospheric organic haze.^{318,319} In addition, the PAH world hypothesis suggests that PAHs could play a role in astrobiological evolution as energy transduction elements, container elements and templating genetic components.^{44,320,321} Therefore, the mechanisms of PAH formation are of fundamental importance for understanding molecular evolution in the universe, the origin of carbon chemistry and prebiotic systems.

In addition to the case of benzene, first, astrochemical models of PAH formation were built from models of combustion flames.³⁸ Considering the temperature in the circumstellar envelopes (~500-2500 K), such borrowing can be considered justified. However, current theoretical studies point out inconsistencies between the stellar yield of PAHs and the estimated abundance of interstellar PAHs. PAHs can be rapidly destroyed by photolysis, interstellar shocks and cosmic ray bombardment.^{165,322,323} The destruction of PAHs occurs through either the loss of the C_2H_2/C_2 -fragment from the periphery of the molecule with the preservation of the aromatic structure or direct ejection of atoms from a skeleton in the case of elastic collisions.^{324,325} This destruction is not compensated by the rate of formation by the mechanisms proposed

for AGB stars. For the lifetime of PAHs in the ISM, there are estimated to be 10⁸ years, which is noticeably shorter than the $2 \cdot 10^9$ years required for the injection of PAHs into the ISM from AGB stars.^{322,323,324,326} Therefore, PAHs should not exist either in the ISM or in meteorites if we consider injection from C-rich AGB stars only. Therefore, the ubiquitous presence of PAHs in the ISM represents a key paradox.^{290,39} On the one hand, this paradox motivates the study of mechanisms in hot stellar envelopes in more detail. On the other hand, it provides reasons to search for alternative mechanisms of PAH formation. Among these alternative mechanisms, a "top-down" scenario can be mentioned. Hirashita proposed a scenario in which PAHs are produced as a result of fragmentation of graphite or amorphous grains in grain-grain collisions at high velocities induced by interstellar shocks.³²⁷ The fingerprint of dust in supernova remnants also hints at the possible formation of PAHs in such highly turbulent media.³²⁸ In turn, Sandstrom et al.³²⁹ showed that the abundance of PAHs is strongly correlated with CO emission, i.e., cold dense clouds, which indicates that PAHs can be generated in such environments. Moreover, the detection of cyanonaphthalene in molecular cloud TMC-1 indicates that the formation must occur in a straightforward manner in this cloud, as such small PAHs are unlikely to survive during a trip from parent stellar envelopes to molecular clouds.^{165,325} To date, several mechanisms that can proceed at low temperatures have been suggested, and C-rich AGB stars are far from being the only source of PAHs.

Thus, the understanding of the formation of PAHs and the role of acetylene in the growth of hydrocarbon molecules has progressed substantially over the past couple of decades from high-temperature mechanisms that can only work in circumstellar envelopes to barrierless mechanisms that work in cold media and cosmic-ray-initiated reactions in acetylene ices on the surface of satellites and planets. Such significant advances in understanding the mechanisms of PAH synthesis are due to experiments involving cross-molecular beams under single-collision conditions, pyrolytic microreactor experiments, and high-level electronic structure calculations.

In a 2021 review, Kaiser and Hansen³³ mentioned five elementary stages that are responsible for the formation of PAHs in a wide range of extreme space conditions, from cold molecular clouds (10 K) and hydrocarbon-rich atmospheres of planets and their moons (50–300 K) to high-temperature circumstellar envelopes of C-rich AGB stars (~1000 K): 1) Hydrogen Abstraction–Acetylene Addition (HACA); 2) Hydrogen Abstraction–Vinylacetylene Addition (HAVA); 3) Phenyl Addition–Dehydrocyclization (PAC); 4) Radical-Radical Reactions (RRR); and 5) Methylidyne Addition-Cyclization-Aromatization (MACA). Recently, He et al. ³³⁰ supplemented the list with the Propargyl Addition–BenzAnnulation (PABA) mechanism. In fact, several alternative mechanisms have been proposed in the literature. For instance, PAHs are suggested to be formed through polyyne polymerization (Section 5.2.4.). In addition to bottom-up mechanisms, PAHs are thought to be generated as a result of the fragmentation of carbon grains exposed to interstellar shocks, as noted above.^{323,327,331} The mechanisms of PAH formation discussed in this review are listed in Table S2 in the Supporting Information.

This variety of pathways makes PAH systems quite complex.³³² Some of the suggested mechanisms can work simultaneously, compete with each other and provide aromatic systems of different sizes and structures, which leads to complicated and varying observational signatures. The acetylene molecule plays an important role in most of the discussed mechanisms, either directly in the formation of complex aromatic systems or as a precursor of key reagents.

In this section, we aim to discuss the mechanisms of PAH formation related to acetylene chemistry, while we omit discussions on other possible PAH formation pathways here. We refer interested readers to a number of specialized reviews in which the full spectrum of mechanisms is described. ^{33,34,35,36,291,326} As in the previous chapter, we divided the mechanisms according to the conditions under which they work; namely, we considered the mechanisms for hot environments applicable to stellar envelopes and protoplanetary nebulae, for cold dense clouds and for Solar system objects. Additionally, we provide information on the reactions leading to the formation of PAHs with five-membered rings.

5.2. Mechanisms of PAH formation in hot environments

5.2.1. HACA mechanism

The most popular mechanism for the growth of an aromatic system larger than benzene is the HACA mechanism, which was explicitly proposed by Bockhorn et al.³³³ Independently, in 1989, Frenklach & Feigelson investigated the mechanism of PAH formation using a chemokinetic approach.²⁵⁴ They specified a similar sequence of chemical reactions for the formation of soot in a hydrocarbon flame and gave the mechanism the name "Hydrogen Abstraction–Acetylene (C_2H_2) Addition" (HACA), which reflects that the sequence includes repeating abstraction of hydrogen followed by the addition of acetylene.³³⁴ On this basis, Frenklach & Feigelson created a model for astrophysical conditions, taking into account the lower pressure and density.

Frenklach noted ³³⁵ that the relationship between "thermodynamic resistance" and reaction reversibility and the kinetic driving force is a defining feature of the "HACA model". In a recent review, Frenklach and Mebel emphasized³⁵ that the kinetic-thermodynamic interaction indeed determines the HACA mechanism but not necessarily the nature of the carbon growth forms. The value of HACA can be extended to cover other forms of carbon growth as long as the kinetic-thermodynamic basis is maintained. In addition, aromatic site activation (the first step in the HACA mechanism) can proceed not only through abstraction but also through hydrogen addition. Thus, Frenklach and Mebel propose considering HACA more generally as an abbreviation for H-Activated-Carbon-Addition, characterized by an underlying kinetic-thermodynamic relationship, where "H-Activated" implies "H-Abstraction", "H-Addition" or even "H-Migration".

However, researchers usually consider the HACA mechanism to be a specific case of aromatic ring formation. In the simplest case, the HACA mechanism starts from benzene (C_6H_6) and leads first to the phenyl radical (C_6H_5) (Scheme 13A). Then, an acetylene molecule is added to form a styrenyl radical (C_8H_7) (Scheme 13B). In the next stage, acetylene is added to the C_8H_7 species (Scheme 13C). In the last stage, cyclization occurs via the elimination of hydrogen to form naphthalene ($C_{10}H_8$), the simplest PAH molecule (Scheme 13D). This process occurs at elevated temperatures up to 1000 K.^{30,336}



Scheme 13. HACA mechanism.

Notably, the reaction described in Scheme 13 can proceed in two different ways.^{33,337,338} The acetylene molecule can directly react with the styrenyl radical (C_8H_7) and form the 4-phenylbuta-1,3-dien-1-yl radical ($C_{10}H_9$). In this case, mutual transformations of the corresponding geometric isomers can occur. As a result, cyclization of the 4-phenyl-(Z)-buta-1,3-dien-1-yl radical ($C_{10}H_9$) occurs with the formation of the 1,8-dihydronaphthalen-1-yl radical ($C_{10}H_9$), from which hydrogen abstraction leads to the formation of naphthalene. This pathway is designated as the Bittner–Howard reaction sequence (Scheme 14).

The second pathway is associated with the isomerization of the styrenyl radical (C_8H_7) to the *ortho*vinylphenyl radical (C_8H_7). Acetylene is attached to this radical, which results in the formation of *ortho*vinylstyrenyl radicals ($C_{10}H_9$). Its cyclization leads to the formation of a 1,2-dihydronaphthalen-1-yl radical ($C_{10}H_9$). Similarly, the elimination of hydrogen leads to the formation of a molecule of naphthalene ($C_{10}H_8$). This pathway is designated as the Frenklach reaction sequence (Scheme 14). Other options for the Frenklach route have also been proposed, involving the formation of an *ortho*-vinylphenyl radical (C_8H_7) via the direct reaction of a phenyl radical (C_6H_5) and C_2H_2 , isomerization of a styrenyl radical (C_8H_7), or a pathway involving H-addition styrene formation.³³⁷





Initially, the mechanism was only hypothesized, but in 2014, it was experimentally confirmed by Parker et al.³⁷ who provided strong evidence that the naphthalene molecule together with phenylacetylene can be formed in a pyrolytic reactor from a phenyl radical and acetylene precursors via the HACA pathway. Photoionization mass spectrometry detection of naphthalene in combination with phenylacetylene successfully demonstrated reaction pathways consistent with the HACA mechanism. In 2016, Kaiser et al.³³⁸ gave additional experimental proof of the easy formation of naphthalene in a simulated combustion environment by the HACA mechanism involving styrenyl C₈H₇ and *ortho*-vinylphenyl C₈H₇ radicals, which are key transitional species of the HACA mechanism.

Frenklach and Feigelson²⁵⁴, who adapted the HACA mechanism to stellar envelopes, concluded that PAH formation should occur within the temperature range between 900 and 1100 K and that large densities and slower stellar winds are required to provide the observed PAH abundance. Their original chains were slightly different from those in Scheme 13 and are presented in Table 13 including the reaction rates. Cherchneff³³⁹ developed the dust condensation model by adding shock-induced chemistry, and it was concluded that in this case, the observed carbonaceous dust mass can be explained by the HACA mechanism only, although subsequent dust destruction in the ISM has not been taken into account.

Another mechanism, the RRR, is supposed to work at high temperatures. For example, this process involves the recombination of two cyclopentadienyl radicals.³⁴⁰ However, this pathway has not been studied in detail as much as HACA has. Therefore, the study of possible pathways is still ongoing.

Reactions	Kinetics	Objects & conditions
a. $C_6H_6 + H \cdot \rightarrow C_6H_5 \cdot + H_2$ b. $C_6H_5 \cdot + C_2H_2 \rightarrow C_8H_7 \cdot$ c. $C_8H_7 \cdot + C_2H_2 \rightarrow C_{10}H_8 + H$	$k_a = 1.5 \cdot 10^{-12} \text{ cm}^3 \text{ s}^{-1} (\text{T} = 1000 \text{ K})$ $k_b = 1.7 \cdot 10^{-11} \text{ cm}^3 \text{ s}^{-1}$ $k_c = 5 \cdot 10^{-10} \text{ cm}^3 \text{ s}^{-1[254]}$	Stellar envelopes (outer edge). Conditions: $n_H = 10^8 \cdot 10^{12} \text{ cm}^{-3}$ T = 500-1500 K

Table 13. The chain of reactions from benzene to naphthalene by the HACA mechanism.

5.2.2. Mechanism of formation of three- and four-ring PAHs in the circumstellar media

A number of experiments have shown that the formation of PAHs with more than two rings via the HACA mechanism is problematic. Kislov et al. performed DFT calculations followed by kinetic analysis of various reaction pathways and showed³⁴¹ that the HACA mechanism essentially ends with the formation of naphthalene ($C_{12}H_8$) and acenaphthalene ($C_{12}H_8$). More recently, Parker et al.³⁴² in 2015 confirmed these

conclusions experimentally. Thus, once naphthalene is formed via the HACA mechanism, further acetylene addition leads to the formation of a five-membered ring instead of a six-membered ring, and acenaphthylene is formed instead of the expected anthracene or phenanthrene.³⁴³ Therefore, the original HACA mechanism does not lead to the growth of six-membered planar PAHs.

Along with the HACA mechanism, it was suggested that aryl-aryl combination reactions can occur during PAH growth.³⁴⁴ In 1994, Sarofim et.al noted that these reactions are underestimated in PAH and soot formation theories³⁴⁵. Biaryls themselves are highly reactive intermediates. They can undergo intramolecular cyclodehydrogenation, and then, upon the addition of acetylene, they turn into highly condensed PAHs.³⁴⁶ Boehm et.al.³⁴⁷ in 1998 investigated the reaction mechanism of PAH growth and showed that aromatic ring–ring condensation is more efficient than the HACA route for the production of high molecular weight aromatic compounds with a short reaction time, while the contribution of HACA increased over longer periods of time. In 2002, Dong and Hüttinger³⁴⁸ reviewed PAH growth reactions and reported that aryl-aryl combination and acetylene addition are the most favorable reactions for PAH growth. The authors also named the model the "particle-filler model", where aromatic hydrocarbons act as molecular species and acetylene is a molecular filler. A general scheme of PAH growth according to this model is illustrated in Figure 12. Depending on the side where acetylene is attached, the final structure of the formed PAHs can be different. For example, Dong and Hüttinger demonstrated a pathway in which naphthalene (C₁₀H₈) and naphthalenyl radical (C₁₀H₇) combine to form binaphthalenyl (C₂₀H₁₄), after which the sequence of dehydrogenation and acetylene addition leads to the formation of PAH C₃₀H₁₄ (Figure 12).



Figure 12. Examples of the «particle-filler model» pathway through aryl condensations and acetylene additions at 4-C bays: for formation of phenanthrene, pyrene (A) and the fully condensed planar PAH (B).³⁴⁸

Later, experiments with crossed molecular beams showed that biphenyl can be formed via a single collision event during the reaction of phenyl radicals with benzene.³⁴⁹ In 2017, Yang et al.³⁹ presented experimental evidence for the formation of phenanthrene (a PAH with three aromatic rings) through the interaction of an *o*-biphenylyl radical and acetylene, as confirmed by DFT calculations. A diagram of the potential energy corresponding to this route is presented in Figure 13. The proposed pathway is initiated by abstraction of a hydrogen atom from the ortho position of the biphenyl molecule, which in turn can be formed through the elemental reaction of benzene with a phenyl radical.³⁴⁹ Then, the radical center of obiphenylyl was added to the acetylene molecule (the reaction barrier was only 10 kJ mol⁻¹). The ring closure of the $C_{14}H_{11}$ collision complex i1 occurs through the passage of a transition state located 20 kJ mol⁻¹ upstream of the complex, resulting in the formation of i2. Notably, the entry barrier can be easily overcome under high-temperature circumstellar conditions. The tricyclic intermediate $C_{14}H_{11}$ (i2) eventually removes atomic hydrogen and aromatizes through a tight transition state to form phenanthrene (p1) and atomic hydrogen. The authors noted that phenanthrene is the preferred product in the temperature range of 500-2500 K and is the main reaction product of o-biphenylyl with acetylene under typical conditions of circumstellar envelopes such as the C-rich star IRC +10216, where temperatures up to 2000 K are maintained near the central star. Generally, this mechanism can be considered a variation of the HACA mechanism, as it includes the same two steps: hydrogen abstraction and acetylene addition.



Figure 13. Proposed reaction mechanism based on stationary points on the $C_{14}H_{10}$ potential energy surface for *o*-biphenynyl reaction with acetylene. Red line indicates the favorable pathway leading to phenanthrene. Reproduced with permission from ref 39. Copyright 2017 John Wiley and Sons.

In 2018, Mebel et al. published a study¹⁸ demonstrating the easy formation of the four-ring PAH pyrene ($C_{16}H_{10}$) in the reaction of the 4-phenanthrenyl radical ($C_{14}H_9$) with acetylene under prevailing conditions in C-rich circumstellar envelopes. According to the mechanism discussed in a previous study, phenanthrene ($C_{14}H_{10}$) molecules can undergo abstraction of a hydrogen atom from the 4-carbon position with the formation of a 4-phenanthrenyl radical.

The proposed route to pyrene through the bimolecular reaction of the 4-phenanthrenyl radical with acetylene occurs in many respects similar to the mechanism of phenanthrene formation discussed above. This process is illustrated in Figure 14. The reaction is initiated by the addition of a 4-phenanthrenyl radical species to an acetylene molecule through an entry barrier of 20 kJ mol⁻¹ to form intermediate i1. Then, the $C_{16}H_{11}$ collision complex ring closes, which leads to intermediate i2 through the entrance barrier at 26 kJ·mol⁻¹. In the last step, intermediate i2 undergoes removal of a hydrogen atom and aromatization to form pyrene as the end product p1. The authors confirmed their conclusions with convincing experimental evidence of the synthesis of a molecule with the molecular formula $C_{16}H_{10}$ in the 4-phenanthrenyl/acetylene system in a high-temperature chemical reactor.



Figure 14. Proposed reaction mechanism based on stationary points on the $C_{16}H_{10}$ potential energy surface for 4-phenanthrenyl radical reaction with acetylene. Red line indicates the favorable pathway leading to pyrene. Reproduced with permission from ref 18. Copyright 2018 Springer Nature.

The pyrene molecule is a key intermediate in the mass growth of PAHs, ultimately leading to the formation of two-dimensional carbon nanostructures such as graphene. The formation of pyrene dimers is the basis for the formation of soot particles.³⁵⁰ However, further experimental measurements of the kinetics of pyrene dimerization coupled with theoretical results showed that a mechanism involving further chemical growth of PAHs to sizes well beyond those of pyrene is necessary.³⁵¹ Mebel et al. proposed a model for the growth of PAHs.¹⁴ The authors noted that the systematic expansion of the ring leading to more complex PAHs is the result of the cooperation of two mechanisms: HACA and HAVA. The HACA mechanism alone cannot explain the formation of PAHs such as benzo[e] pyrene (C₂₀H₁₂). The abstraction of hydrogen and the addition of acetylene to the pyrene molecule can lead to the formation of the $C_{18}H_{11}$ radical, which is expected to undergo cyclization with subsequent loss of atomic hydrogen to cyclopent[cd]pyrene (C₁₈H₁₀). The addition of a second acetylene molecule followed by cyclization and loss of hydrogen to C₂₀H₁₂ is unlikely according to electronic structure calculations^{34,341} and experimental data for the naphthyl/acetylene system.³⁴² Therefore, alternative mechanisms to HACA should complement the formation of extensive polyaromatic systems. Starting with pyrene, the HAVA mechanism provides radial third-order perimeter growth, while the HACA mechanism responds to closing to bays to form fivemembered rings. Sequential implementation of the HACA and HAVA mechanisms allows the synthesis of nonplanar carbon nanostructures containing corannulene subunits (see Section 5.2.3).

In recent studies, Mebel et al. showed that, along with the HACA and HAVA mechanisms, species such as allene (C_3H_4) and methylacetylene (C_3H_4) can also play a role in ring closure during the growth of condensed polyaromatic molecules.³⁵² Experiments in a high-temperature chemical microreactor combined with electronic structure calculations have shown that naphthalene-based isomers, such as 2-(prop-2-yn-1-yl)naphthalene ($C_{13}H_{10}$), 2-(prop-1-yn-1-yl)naphthalene ($C_{13}H_{10}$), and 2-(propa-1,2-dien-1-yl)naphthalene ($C_{13}H_{10}$), can be obtained from the aforementioned $C_{3}H_{4}$ species with the 2-naphthyl isomer. Moreover, it

was possible to record the formation of 3H-benz[*e*]indene ($C_{13}H_{10}$) and 1H-benz[*f*]indene ($C_{13}H_{10}$), which each contain a five-membered ring.³⁵² Later, it was shown that the formation of 1H-phenalene can occur by a similar mechanism. The authors noted that the processes of graphene nanoflakes can start from 1Hphenalene via peripheral expansion of the 1H-phenalene species to the triangular condensed molecules 1,5dihydrodibenzo[*cd*,*mn*]pyrene ($C_{22}H_{14}$) and then 6,11-dihydro-1H-tribenzo[*bc*,*hi*,*no*]coronene ($C_{33}H_{18}$).³⁵³ Despite the fact that acetylene molecules are not directly involved in this mechanism, the molecules allene ($C_{3}H_{4}$) and methylacetylene ($C_{3}H_{4}$) are derivatives of acetylene, and as shown above, their formation can proceed by the interaction of acetylene with a methylene species.¹⁶⁰ This highlights the role of acetylene as a versatile building block for a variety of carbon molecules.

One additional mechanism, the PAC, should also be considered. In the case of PAC, a phenyl radical attaches to biphenyl followed by H abstraction. After subsequent dehydrogenation, cyclization and aromatization, the biphenyl-phenyl system is converted to triphenylene, i.e., four-ring PAHs. Zhao et al. demonstrated the formation of triphenylene through the PAC mechanism. This mechanism may occur in high-temperature and high-pressure regions of space, such as the circumstellar envelopes of C-rich AGB stars, e.g., IRC+10216.³⁵⁴

Notably, in 2021, Green et al. studied the reaction of naphthalene radicals with acetylene (C_2H_2). Using VUV photoionization time-of-flight mass spectrometry at 15 to 50 torr and 500 to 800 K, they recorded the formation of $C_{14}H_{10}$ species, which are probably phenanthrene and anthracene. This study provides experimental evidence that the HACA mechanism is a pathway for the formation of PAHs larger than naphthalene.³⁵⁵

Reactions	Kinetics	Objects conditions:	&
1. a. $C_6H_5 + C_6H_6 \rightarrow C_{12}H_{10} + H$ b. $C_{12}H_{10}$ (biphenyl) + $C_2H_2 \rightarrow$ $C_{14}H_{10}$ (phenanthrene) + 2H c. $C_{14}H_{10} + C_2H_2 \rightarrow C_{16}H_{10}$ (pyrene) + 2H	$k_{1a} = 3.1 \cdot 10^{-13} \text{ cm}^3 \text{ s}^{-1} (\text{T} = 1000 \text{ K})^{[347]}$ $k_{1b} = 4.0 \cdot 10^{-13} - 2.6 \cdot 10^{-12} \text{ cm}^3 \text{ s}^{-1} (\text{p} = 0.04\text{-}100 \text{ atm.}, \text{T} = 1000 \text{ K})^{[39]}$ $k_{1c} = 1.0 \cdot 10^{-12} \text{ cm}^3 \text{ s}^{-1[18]}$	Stellar envelopes (outer edge). Conditions: $n_H = 10^8 \cdot 10^{12} \text{ cm}^{-3}$ T = 500 \cdot 1500 K	
2. $C_{10}H_{7} + C_{3}H_{4} \rightarrow C_{13}H_{10} + H$ $C_{10}H_{7} + C_{3}H_{4} \rightarrow C_{12}H_{8} + CH_{3}$. ^[353]	n.d.		
$3. C_{12}H_{10} + C_6H_5 \rightarrow C_{18}H_{12}$ (triphenylene) + H + H ₂ ^[354]	n.d.		

Table 14. Summary of considered reactions, kinetics (if available) and conditions where the reactions are relevant

5.2.3. Mechanism of formation of PAHs with five-membered rings

The formation of PAHs with five-membered rings is an important step in the formation of detected in space nonplanar carbon molecules, such as fullerene. Several alternative mechanisms for the formation of PAHs with five-membered rings have also been presented in the last decade. Parker et al.³⁵⁶ provided experimental evidence that indene can be formed by the bimolecular gas-phase reaction of a benzyl radical (C_7H_7) with acetylene at 600 K. This mechanism was theoretically predicted as early as 1981 by Bittner and Howard.³⁵⁷ This process includes the addition of C_2H_2 to the CH₂ moiety of the benzyl radical to form a doublet C_9H_9 intermediate. Then, the doublet C_9H_9 intermediate isomerizes through facile ring closure to the *ortho*-carbon of the phenyl ring to form bicyclic intermediate i2, which can be converted to an indene after H loss from the *ortho*-carbon of the phenyl ring (Figure 15).



Figure 15. Proposed mechanism of indene formation via the reaction of the benzyl radical with acetylene. Energies are given in kJ mol⁻¹. Reproduced with permission from ref 354. Copyright 2015 John Wiley and Sons.

The growth of the aromatic system can further proceed through the mechanisms described above involving acetylene or its derivatives. Thus, 5- and 6-indenyl radicals (C_9H_7) can interact with vinylacetylene (C_4H_4). The products of this interaction are tricyclic PAHs, 3H-cyclopenta[*a*]naphthalene ($C_{13}H_{10}$), 1H-cyclopenta[*b*]naphthalene ($C_{13}H_{10}$), and 1H-cyclopenta[*a*]naphthalene ($C_{13}H_{10}$), 1H-cyclopenta[*b*]naphthalene ($C_{13}H_{10}$), and 1H-cyclopenta[*a*]naphthalene ($C_{13}H_{10}$), 360 (Figure 16). In addition, it has recently been shown that the reaction of a benzyl radical with a phenyl (C_6H_5) radical is a pathway for the formation of another important PAH containing a five-membered ring, 9H-fluorene.³⁵⁸ Li et al. showed that the interaction of acetylene with methylene-substituted aromatic compounds carrying a hydrogen atom at the *ortho*-position of the ring can provide a universal pathway for accessing five-membered aromatic compounds ($C_{13}H_{10}$, as indicated above) at elevated temperatures.³⁵⁹ These reactions were carried out in a high-temperature chemical microreactor at 1300 K. In addition to the experiments, corresponding calculations of the energies and molecular parameters of various intermediates and transition states were performed. These calculations have shown that tricyclic PAHs can be formed via a barrierless HAVA mechanism in cold molecular clouds at temperatures of approximately 10 K.³⁶⁰



Figure 16. Proposed mechanism formation of 3H-cyclopenta[*a*]naphthalene, 1H-cyclopenta[*b*]naphthalene, and 1H-cyclopenta[*a*]naphthalene (C₁₃H₁₀) by reaction of 5-/6-indenyl radicals (C₉H₇) with vinylacetylene (C₄H₄). Reproduced with permission from ref 358. Copyright 2020 Royal Society of Chemistry.

An interesting experimental study of the formation of PAHs with five-membered rings under hightemperature conditions was carried out by Necula and Scott.³⁶¹ Using flash vacuum pyrolysis, the authors demonstrated the formation of large PAH molecules at a temperature of approximately 830 K and pressures in the range of 10^{-5} - 10^{-4} atm through the C₂-accretion pathway. The key C–C bond-forming step involved the trapping of aryl radicals by acetylene (C₂H₂). The main reported product of the reaction of acetylene and 1-naphthyl radical was acenaphthylene (C₁₂H₈). Minor amounts of naphthalene (C₁₀H₈) and 2ethynylnaphthalene ($C_{12}H_8$), and traces of pyracylene ($C_{14}H_8$) were also detected. A similar set of products was obtained when 2-naphthyl radicals were used. The proposed mechanism is presented in Scheme 15.



Scheme 15. Proposed reaction mechanism acenaphthylene formation by trapping aryl radicals with acetylene. ³⁵⁹

One of the important molecules with a five-membered ring is corannulene ($C_{20}H_{10}$), which is a building block in the pathway of fullerene formation. A combination of molecular beam experiments and theoretical calculations demonstrated that corannulene can be synthesized in the gas phase through reactions of 7-fluoranthenyl ($C_{16}H_9$) and benzo[*ghi*]fluoranthenen-5-yl ($C_{18}H_9$) radicals with acetylene (Figure 17). This process is initiated by H abstraction from the 7- and 5-positions of fluoranthene ($C_{16}H_{10}$) and benzo[*ghi*]fluoranthenene ($C_{16}H_{10}$) and benzo[*ghi*]fluoranthenene ($C_{16}H_{10}$) and has barriers that can be easily overcome in high-temperature environments of C-rich circumstellar envelopes of AGB stars or under harsh UV radiation fields intrinsic to planetary nebulae.³⁶² Notably, fluoranthene can also be formed via the PAC mechanism when the phenyl radical reacts with naphthalene, similar to the reaction between the phenyl radical and biphenyl described above.



50

Figure 17. Scheme of the formation of corannulene through the following sequence: formation of fluoranthene from the phenyl radical involving steps of HACA (red), HAVA (blue), and PAC (green) mechanisms; the formation of benzo[*ghi*]fluoranthene and corannulene by the HACA mechanism through the reaction of the 7-fluoranthenyl radical with acetylene.³⁶²

Reactions	Kinetics	Objects & conditions
1. $C_7H_7 + C_2H_2 \rightarrow C_9H_8 + H^{[356]}$	n.d.	Stellar envelopes (outer edge). Conditions: $n_H = 10^8-10^{12} \text{ cm}^{-3}$ T = 500-1500 K
2. $C_9H_7 + C_4H_4 \rightarrow C_{13}H_{10} + H^{[360]}$	n.d.	Molecular clouds, prestellar cores. Conditions: $n_H = 10^2 - 10^7 \text{ cm}^{-3}$ T = 10 K $G_0 = \text{very low}$ Galactic CR field
$\begin{array}{c} 3. \ C_7H_7 + C_6H_5 \rightarrow C_{13}H_{10} + \\ H_2^{[330]} \end{array}$	$k_3 = 3 \cdot 10^{-11} \text{ cm}^3 \text{ s}^{-1}$ (T = 1500 K, p = 0.01 atm) ^[330]	Stellar envelopes (outer edge)
4. a. $C_{16}H_9 + C_2H_2 \rightarrow C_{18}H_{10} + H_1$ b. $C_{18}H_{10} \rightarrow C_{18}H_9 + H_1$ c. $C_{18}H_9 + C_2H_2 \rightarrow C_{20}H_{10} + H_1^{[362]}$	$k_{4a} \approx 1.0 \cdot 10^{-11} \text{ cm}^3 \text{ s}^{-1}$ $k_{4c} \approx 3.0 \cdot 10^{-12} \text{ cm}^3 \text{ s}^{-1}$ (both at 1500 K, p = 0.04-100 atm.) ^[362]	Stellar envelopes (outer edge)

Table 15. The reactions which lead to formation of PAHs with five-membered rings.

5.2.4. PAH formation via polyme polymerization

In the 1990s, Krestinin et al. proposed an alternative mechanism for PAH and soot formation.^{363,364,365,366} They considered polyyne polymerization as an alternative route for generating high-molecular-weight carbon species from acetylene. This hypothesis is consistent with the observations of small polyynes along with PAHs in the flame soot formation zone.²⁵² In addition, polyynes and small (transient) cyclic compounds can be formed in discharge plasma sources and are used to create a chemically reactive environment that simulates astrochemical conditions.³⁶⁷ As mentioned above (Section 3.2), carbon-chain polyynic radicals (C_nH), polyynes (HC_nH), cyanopolyynes and methylpolyynes are quite abundant in the C-rich star IRC +10216 and protoplanetary nebula CRL 618 and are formed from acetylene.^{143,266,368}

Bonn, Homann and Wagner suggested³⁶⁹ that polyynes play a key role in the formation of soot. Proponents of the polyyne hypothesis note that the thermodynamic stability of acetylene (and the entire family of $HC_{2n}H$ polyynes, n = 2, 3,...) increases at higher temperatures, while the stability of all other hydrocarbons decreases.³⁷⁰ Krestinin et al. also indicated that polyynes grow rapidly by a one-step ethynyl radical addition mechanism, as opposed to aromatic ring formation. As a result, the concentration of polyynes will be quite high (Scheme 16).

 $HC_{2n}H + C_2H \rightarrow HC_{2n+2}H + H$

Scheme 16. Polyynes growth via one-step ethynyl radical addition mechanism.

These polyynes can form complexes at the surface radical sites of the seed particles, followed by branching during chain polymerization (Scheme 17). As a result of cyclization occurring in such a complex, two new additional radical centers are created. This "propagation" of radical sites during polyyne

polymerization is a key assumption that introduces nucleation into the model. This finding is also consistent with the data of Homann and Wagner³⁶⁹, who noted that the electron spin resonance signal from nascent soot particles under combustion conditions exceeds that from mature soot by more than 100 times, indicating the generation of large amounts of radicals during nucleation. Cherchneff et al. noted that such a mechanism of PAH growth may be relevant in environments such as the wind of the hydrogen-rich R Coronae Borealis star V854 Cen.²⁴⁶



Scheme 17. Formation of a polyyne complex at the surface radical site and its transformation into a more stable structure with subsequent "propagation" of radical centers.³⁷⁰

Reactions	Kinetics	Objects & conditions
$C_{2n}H_2 + C_2H \rightarrow C_{2n+2}H_2 + H^{[363]}$	$k = 3.3 \cdot 10^{-11} \mathrm{cm}^3 \mathrm{s}^{-1}$	$\begin{array}{l} \mbox{Stellar envelopes (outer edge).} \\ \mbox{Conditions:} \\ n_{\rm H} = 10^8 \text{-} 10^{12} \mbox{ cm}^{-3} \\ T = 500 \text{-} 1500 \mbox{ K} \\ \end{array}$ $\begin{array}{l} \mbox{Protoplanetary nebulae.} \\ \mbox{Conditions:} \\ T = 250 \mbox{ K} \\ n_{\rm H} = 10^7 \mbox{ cm}^{-3} \\ \mbox{G}_0 = 2 \cdot 10^5 \end{array}$

Table 16. The reaction of polyyne growth via ethynyl radical addition mechanism.

5.3. Mechanisms of PAH formation in cold interstellar medium

As mentioned above, the observations and theoretically estimated PAH lifetimes require a path of PAH formation in the cold ISM at a temperature of approximately 10 K. Due to such low temperatures and pressures, a vital criterion for potential reaction paths will be their barrierless nature and occurrence under single collision conditions. Reactions by the HACA mechanism involving acetylene have significant entry barriers ranging from 10 to 30 kJ mol⁻¹ and can occur only at high temperatures up to several thousand Kelvin.³⁷¹ However, in the last decade, it has been found that many reactions involving aromatic radicals leading to the formation of PAHs are essentially barrierless.³⁵⁶ Such mechanisms (a single barrierless bimolecular collision) can play a key role in the formation of PAHs in extremely low-temperature environments, such as dense molecular clouds and atmospheres of planets and their satellites in the outer Solar system.

Mebel et al. proposed a number of barrierless mechanisms for the formation of PAHs in cold environments³⁷². Among these growth mechanisms of the aromatic system, the most important are the ethynyl addition mechanism (EAM), HAVA, and the reaction between methylidyne and styrene, leading to the formation of an indene (MACA).²⁹⁰ In all the mechanisms, acetylene can be a precursor of the chain initiator. We provide these reactions in Table 17. Apart from gas-phase reactions, PAHs can likely be formed in astrophysical ices, as will be shown below, together with mechanisms of formation in planetary and satellite atmospheres.

To determine the significance of these barrierless reactions, one needs to include them in astrochemical modeling. Such work was started by McGuire et al. 2022.³⁰⁰ The authors considered the

reactions between phenyl radicals and vinylacetylene and between phenyl radicals and 1,3-butadiene to form naphthalene, which further reacts with cyanide to form cyanonaphtalene. The authors concluded that the considered reactions do not provide the observed abundance of cyanonaphtalene; therefore, other pathways should be taken into account in future modeling.

Reactions	Kinetics	Objects & conditions
1. a. $C_6H_6 + C_2H \rightarrow C_6H_5C_2H + H$ b. $C_6H_5C_2H + C_2H \rightarrow C_{10}H_7$ or c. $C_6H_5C_2H + C_2H \rightarrow C_6H_5(C_2H)_2$ d. $C_6H_5(C_2H)_2 + C_2H \rightarrow C_{12}H_8^{[372]}$	$k_{1a,b,c,d} \sim 10^{-10} \mathrm{cm}^3 \mathrm{s}^{-1} ^{[372]}$	Molecular clouds, prestellar cores. Conditions: $n_H = 10^2 - 10^7 \text{ cm}^{-3}$ T = 10 K $G_0 = \text{very low}$ Galactic CR field + Some reactions suit planetary atmospheres and satellites.
2. $C_6H_5C_2H_3 + CH \rightarrow C_9H_8^{[290]}$	$k_2 \sim 10^{-10} \text{ cm}^3 \text{ s}^{-1[290]}$	The same as reactions #1.
3. $C_6H_5 + C_4H_4 \rightarrow C_{10}H_8 + H^{[34,373]}$	$k_3 \sim 10^{-14} \text{ cm}^3 \text{ s}^{-1} [34]$	Molecular clouds (not planets)

Table 17. The reactions of PAH growth at low temperatures.

5.4. Mechanisms of PAH formation for planets and satellites

In addition to circumstellar envelopes and the ISM, the atmospheres of planets and satellites are no less important for the production of PAHs. In particular, researchers have paid special attention to the chemistry of the atmosphere of Titan. PAHs are considered to be key components of Titan's haze and potential constituents of organic material on Titan's surface.^{374,375} The atmosphere of Titan is a relatively low-temperature environment (surface temperature of 94 K)³⁷⁶; therefore, high-temperature mechanisms are difficult to use to explain the formation of PAHs under such conditions. However, the main explanations for the formation of PAHs in Titan's atmosphere have come from classical high-temperature mechanisms such as HACA.³⁷⁷

Due to the low temperature of Titan, any entry barrier effectively blocks the reaction. Recently, Mebel et al. proposed a new model for the growth of PAHs in the atmosphere of Titan based on the barrierless HAVA mechanism. The authors experimentally proved that tricyclic PAHs, such as anthracene $(C_{14}H_{10})$ and phenanthrene $(C_{14}H_{10})$, can be formed in the atmosphere through bimolecular gas-phase reactions of 1- and 2-naphthyl radicals $(C_{10}H_7)$ with vinylacetylene $(CH_2=CH-C=CH)$.³⁷⁸

In addition to the atmosphere of planets, important chemical evolution processes can occur on the surface of planets. This is how the Cassini-Huygens mission discovered a remarkably rich world on Titan's surface, with complex landscapes and topography.³⁷⁹ One of the amazing features of Titan's landscape is the large longitudinal dunes in the equatorial deserts.^{380,381} Data from Cassini's VIMS indicate the presence of a significant amount of dark organics of undetermined chemical composition and origin.³⁸² It was assumed that the main components of these formations are aromatic hydrocarbons formed as a result of molecular weight growth initiated by solar photons through gas phase ion-molecular and neutral-neutral reactions and surface reactions on the satellite.^{106,293,383},³⁸⁴,³⁸⁵ However, the discrepancy between the particle sizes of the material from the dark dunes at Shangri-La³⁸⁶ (one of the three largest equatorial Titan's sand seas) and the particle size of atmospheric aerosols points to a special way of their formation.³⁸⁰

A comparison of data from the Cassini Synthetic Aperture Radar (SAR) on the location of Titan's dark dune and VIMS data on the location of acetylene ice allowed us to make an assumption about the link between acetylene ices and the formation of dune material.^{28,110,110,387,388} At the same time, most of the energy sources are absorbed by Titan's thick atmosphere. However, energetic CRs are able to penetrate Titan's atmosphere and reach the surface.^{389,390} The transfer of a part of the kinetic energy of the secondary electrons formed by CRs to the acetylene molecule can lead to electronically excited acetylene. As

discussed in Section 4.5, the formation of benzene in ice can occur by the reaction of electronically excited acetylene with a neighboring singlet acetylene molecule. In this case, the barrier can be easily overcome with the help of vibrational energy in one of the reagents. Similar processes of aromatic ring growth with the participation of electronically excited species can lead to naphthalene and phenanthrene on the surface of Titan.

Recently, Kaiser et al. conducted an experiment in which ices of acetylene and deuterated acetylene (C_2D_2) irradiated energetic electrons at 5 K and pressures of a few 10^{-11} torr.²⁸ As a result, the authors recorded the formation of aromatic hydrocarbons, such as naphthalene and phenanthrene. An increase in the dose by a factor of approximately 10 to 27 ± 2 eV per molecule promotes the synthesis of even more complex aromatic compounds, such as chrysene ($C_{18}H_{12}$), perylene ($C_{20}H_{12}$), pentacene ($C_{22}H_{14}$) and coronene ($C_{24}H_{12}$), which bear up to six benzene rings.³⁹¹

X-ray-induced formation of naphthalene from an isolated styrene-acetylene complex in a solid krypton matrix at 6 K was demonstrated by Lukianova and Feldman in 2022.³⁹² The authors used SF₆ as an electron scavenger to determine the conversion mechanism. Using Fourier transform IR spectroscopy, they determined that the formation of a naphthalene radical cation occurred. Thus, the implementation of the cationic pathway is assumed: the cycle is closed due to the formation of an ionized complex, followed by the elimination of a hydrogen molecule (Scheme 18). Thus, it was shown that PAHs can form as a result of the impact of CRs on low-temperature acetylene ice on the surface of Titan.



Scheme 18. Proposed mechanism of naphthalene formation from the styrene-acetylene complex under X-ray irradiation through radical cation.³⁹²

Similar processes, including the formation of aromatic molecules from acetylene at low temperatures, can occur not only on Titan but also on other objects of the Solar system, such as the dwarf planets Pluto³⁹³, Makemake, Ceres^{394,395}, Orcus³⁹⁶, Salacia³⁹⁷, Saturn's satellites Hyperion, Iapetus and Phoebe^{398,399,400}, Jupiter's moons Ganymede and Callisto³⁸⁸, satellites Pluto, Charon, Nikta, Hydra^{401,402} and others, where regions with dark organic matter have been found. Recently, Kaiser et al. showed that the photochemistry of acetylene ice can lead to the formation of PAHs, including five ring molecules, such as dipropylperylene (C₂₆H₂₄).^{403,287} In another experiment simulating the irradiation of methane and acetylene by galactic cosmic rays in the Kuiper belt in ultrahigh vacuum, Kaiser et al.⁴⁰⁴ reported the initial formation of acetylene from methane. In this work, approximately 30 resulting organic molecules were identified, including mono- and tricyclic aromatic compounds (substituted benzenes, naphthalenes and biphenylenes, as well as phenanthrene, phenalene, acenaphthylene and 9H-fluorene). Methyl (CH₃), ethyl (C₂H₅), vinyl (C_2H_3) and allylic (C_3H_5) groups were determinated as substituents. The authors noted that these groups may be potential linkers that connect individual aromatic molecules to complex tholin-like structures. The authors also emphasized a stepwise process of molecular weight growth both in the series of mono-, biand tricyclic molecules through ring annulation and from pure aromatic molecules to mono- and dialkylated PAHs.

What makes this mechanism even more universal is that methane ices have been found on many of the outer bodies of the Solar system. ⁴⁰⁵ Methane, in turn, can be easily converted by ionizing radiation to acetylene.^{159,404} Thus, acetylene becomes a bridge to a variety of complex organic compounds on the hydrocarbon-rich surfaces of planets and their satellites.

The reactions discussed in this section are given in Table 18. Important pathways of PAH formation based on experiments with crossed molecular beams and pyrolytic reactors combined with quantum chemical calculations are listed in Table S3 in the Supporting Information.

Reactions	Kinetics	Objects & conditions
$\begin{array}{l} 1. \ C_{10}H_{7^{*}} + C_{4}H_{4} \rightarrow C_{14}H_{10} \\ + H \end{array}$	$k_{I} \sim 10^{-17} \text{ cm}^{3} \text{ s}^{-1}$ (p = 10 ⁸ atm, T = 70- 180 K) ^[378]	Planetary atmospheres and satellites. Conditions: $n_{\rm H} = 10^{10} \cdot 10^{16} \text{ cm}^{-3}$ T = 50-300 K $G_0 = \text{up to } 10^2$ Galactic CR field.
Ice reactions 2. a. $3C_2H_2 + CR \rightarrow C_6H_6$ b. $C_6H_6 + 2C_2H_2 + CR \rightarrow C_{10}H_8 + H_2$ c. $C_{10}H_8 + 2C_2H_2 + CR \rightarrow C_{14}H_{10} + H_2^{[28,404]}$	$k_{2a} \sim 0.6/0.19$ amu eV ⁻¹ (10/40 K) ^[404] $k_{2b} \sim 0.055/0.08$ amu eV ⁻¹ (10/40 K) ^[404]	Specifically, Titan, Kuiper's belt objects. Conditions: Galactic CR field. T = 50-100 K G ₀ : up to 10 Galactic CR field.
3. $C_6H_5C_2H_3 + C_2H_2 + CR$ $\rightarrow C_{10}H_8^{[392]}$	n.d.	Molecular clouds, prestellar cores. Conditions: $n_{\rm H} = 10^2 \cdot 10^7 \text{ cm}^{-3}$ T: 10 K $G_0 = \text{very low}$ Galactic CR field. + Planetary atmospheres and satellites.

Table 18. The reactions of the PAH growth in planetary atmospheres and their satellites.

6. Formation of carbon nanoparticles and grains

6.1. Mechanism of formation of fullerenes and nanotubes

The formation of other important carbon macromolecules, such as fullerenes and nanotubes, has been studied. Fullerenes are detected in protoplanetary and planetary nebulae; therefore, they must be formed in these objects or, earlier, in stellar envelopes. These particles may form simultaneously or separately from PAHs. Traditionally, fullerenes are produced from carbon vapor in arc charge or, e.g., via laser ablation^{406,407} but these methods scarcely work in stellar envelopes. Several mechanisms suitable for astrophysical conditions have been suggested, including a 'bottom-up' scenario⁴⁰⁸ — closed network growth from C and C₂ gas — and a 'top-down' scenario^{409,410,411,412,413} — formation due to photodissociation of PAHs or HACs.⁴⁰⁹ Despite their diversity, the proposed formation scenarios are of interest and can be taken into account in astrochemical modeling.

Recently, Tuli et al.⁴¹⁴ suggested a new 'bottom-up' pathway to form a building block of fullerenes — a nanobowl $C_{40}H_{10}$. The nanobowl can be obtained through a chain of reactions starting from the corannulene molecule, whose formation was described above in Section 5.2.3. The chain of reactions includes 1) the reaction of the corannulene radical ($C_{20}H_9$) with vynilacetylene (C_4H_4), which results in the formation of benzocorannulene ($C_{24}H_{12}$); 2) benzoannulation of benzocorannulene up to the formation of pentabenzocorranulene ($C_{40}H_{20}$); and 3) ring closure between five attached benzenes with H abstraction and formation of the nanobowl $C_{40}H_{10}$. Curvatures are caused by five-membered rings. The formed nanobowl serves as a building block for buckminsterfullerene C_{60} .

It is likely that along with fullerenes, nanotubes are present in space, although they have not yet been detected. Chen and Li studied the possibility of carbon nanotube formation in space via the HACA mechanism using DFT.²³ The model they proposed included, at the first stage, the formation of biphenyl by the addition of phenyl to benzene with the removal of hydrogen. The formation of triphenyls occurs via a process similar to that of the addition of a second phenyl ring, in addition to the removal of hydrogen. In the next stage, triphenyl isomerizes into a closed three-dimensional structure with the loss of hydrogen.

The next three stages represent the growth of a nanotube with the participation of acetylene according to the HACA mechanism (Figure 18). The authors studied the key transition states involved in forming a nanotube via a similar path. The most interesting point was the possibility of bending the triphenyl molecule to form a ring structure. The displacement vectors tended to close the structure. The authors also examined the internal reaction coordinates of the transition state to confirm the calculations. The calculated barrier for molecular bending was 4.59 eV, which is achievable in the circumstellar envelopes of AGB stars and planetary nebulae.



Figure 18. Scheme of molecular growth of a three-dimensional carbon nanostructure by the NACA mechanism. Reproduced with permission from ref . Copyright 2019 The European Southern Observatory (ESO).

Reactions	Kinetics	Objects & conditions
1. a. $C_{20}H_{10} + H \rightarrow C_{20}H_{9} + H_{2}$ b. $C_{20}H_{9} + C_{4}H_{4} \rightarrow C_{24}H_{12} + H$ c. 4 iterations of a and b: $C_{24}H_{12} + 4x[+H] + 4x[C_{4}H_{4}] \rightarrow$ $C_{40}H_{20} + 4x[H_{2}] + 4x[-H]$ d. $C_{40}H_{20} \rightarrow C_{40}H_{10}$ (nanobowl) + $5H_{2}^{[414]}$	$k_{lb} \sim 3.4 \cdot 10^{-17} \text{ cm}^3 \text{ s}^{-1}$ (T = 1200 K, p = 0.03 atm) ^[414] $k_{ld} \sim 10^7 \text{ s}^{-1}$ (T = 1200 K, p = 0.03 atm) ^[414]	Stellar envelopes (outer edge). Conditions: $n_H = 10^8 \cdot 10^{12} \text{ cm}^{-3}$ T = 500-1500 K
2. a. $C_6H_5 + C_6H_6 \rightarrow C_{12}H_{10} + H$ b. $C_{12}H_{10} + C_6H_5 \rightarrow C_{18}H_{14} + H$ c. $C_{18}H_{14} \rightarrow [\text{isomerization}] \rightarrow$ $C_{18}H_{12} (\text{hnt } 3\text{-}3\text{-}1) + 2H$ d. $C_{18}H_{12} + 3C_2H_2 \rightarrow C_{24}H_{12} (\text{hnt } 3\text{-}3\text{-}2) (\text{nanotube}) + 6H^{[23]}$	n.d.	

Table 19. The reactions which lead to formation of a nanobowl and a nanotube.

6.2. Soot formation

The formation of PAHs is only the first step in the formation of carbon dust grains. The study of dust formation has roots in the study of the practically significant process of carbon black formation.⁴¹⁵ Currently, it is generally accepted that the process of carbon black or soot formation during the pyrolysis of hydrocarbon feedstock includes dehydrogenation, pyrolysis, cracking, and aromatization. An important role is played by the intermediate formation of acetylene, which occurs in flames by pyrolysis of organic fuels at sufficiently high temperatures.^{332,416,417} These processes eventually lead to the formation of a growth

nucleus from PAHs, which are nanosized (1-2 nm) clusters of interconnected carbon atoms. All the underlying mechanisms are currently based on the key role of PAHs in this process (Figure 19).⁴¹⁸

In recent work³⁵, Frenklach and Mebel critically reviewed soot nucleation mechanisms. They quantitatively estimated (by performing kinetic calculations coupled with thermodynamics and DFT calculations) a number of reactions of the dimerization of PAHs. The authors concluded that the most stable PAH dimer is the one in which two PAHs are connected by a doubly bonded bridge, which is termed the E-bridge. The connected rings on the edges are five-membered, and the bridges are aliphatic bonds. The authors emphasize that the dimerization mechanism is related to the HACA mechanism.



Figure 19. The main stages of the process of formation of soot particles with division into low-temperature and high-temperature regions with a conditional boundary of 1500 °C. Reproduced with permission from ref 416. Copyright 2022 Russian Chemical Reviews.

Polyacetylenic chains (polyynes) formed during combustion (see Sections 3.2 and 5.2.4) may play an important role in the formation of large carbon dust particles. Experiments using a special cosmic simulation chamber (COSmIC) facility showed that polyacetylenic chain radicals, as well as H and C_2H_2 , can be formed during the opening of PAH cycles.²⁰ Molecular dynamics studies of ReaxFF by Mao et al.⁴¹⁹ showed that such radicals promote the growth of soot particles. Polyacetylene radicals can link large PAHs such as coronene, ovalene, and circumcoronene to form 'carbon bridges' on the edge (Figure 20, Table 20). As a result, soot particles with stacked structures are formed. In addition, fullerene-like soot particles are formed from polyacetylenic chains at 2500 K.



Figure 20. Key stages of PAH fragmentation and soot formation from circumrooten monomers at 2500 K. Reproduced with permission from ref 417. Copyright 2017 Elsevier.

Using the Stardust machine,⁴²⁰ the final gas-phase molecules and solid nanoparticles obtained from the C_2/C_2H_2 gas mixture were investigated by Santoro et al.²⁰⁷. Among the gas-phase products, polyynes; different aromatic molecules, such as benzene, toluene, styrene, naphthalene, acenaphtalene, and biphenyl; and hydrogenated aliphatic C₅-, C₉-, and C₁₁-clusters were found. Compared with the results of previous experiments carried out in the absence of $C_2H_2^{421}$, the composition of the products was significantly different; therefore, they emphasized the key role of acetylene in the formation of dust analogs. The produced dust analogs were agglomerates with a diameter of several nanometers. The particles represent complex hydrocarbons consisting of sp, sp² and sp³ bonds, i.e., they are a mix of aromatics connected by aliphatic or olefinic bonds. Therefore, all the above-mentioned molecules originated from acetylene, which includes small hydrocarbons, polyacetylenic chains, benzene and PAHs, and participate in the formation of carbon dust.

Table 20. Summary of considered reactions, kinetics (if available) and conditions where the reactions are relevant

Reactions	Kinetics	Objects & conditions
$\begin{array}{rcl} PAH + PAH + C_2H_2 \rightarrow [PAH]_2 \\ (cluster)^{[419]} \end{array}$	n.d.	Stellar envelopes (outer edge). Conditions: $n_H = 10^8 - 10^{12} \text{ cm}^{-3}$ T = 500-1500 K

7. Catalytic reactions in space with acetylene

The abundance of molecules in cold ISM cannot be explained by considering only gas-phase reactions. The ISM, which is mostly gas, also contains approximately 1% dust particles ranging in size from a few tenths of nanometers to several microns.⁶⁰ These dust particles can catalyze many reactions because low-molecular-weight species can agglomerate on their surface. The interaction of gas species with grains can lead to the formation of COMs. Surface catalysis on solid interstellar particles can enable chemical pathways that cannot occur in the gas phase due to reaction barriers.^{422,423} The importance of heterogeneous catalysis in astrochemistry has rapidly grown in recent years. In particular, rich grain-surface chemistry has been described in reviews by Tielens et al.⁶¹ and Cuppen et al.⁴²⁴ One exciting example of a catalytic reaction on the surface of a cosmic dust grain analog is the formation of cyclic prebiotic molecules such as pyrimidinone, uracil, cytosine, purine, urea, and dihydrouracil.^{425,426} Simple compounds of elements such as carbon^{427,428}, silicon⁴²⁹, titanium⁴³⁰, iron⁴³¹ and many other transition metals⁴³², whose high catalytic activity has been well studied in organic synthesis reactions in recent decades, can act as catalytic centers of cosmic particles. Certainly, fascinating discoveries are coming in this astrochemical field in the future.

7.1. Catalytic reactions of acetylene in the laboratory

In our opinion, catalytic pathways for the formation of PAHs from acetylene might have a great potential. In the laboratory, the process of obtaining benzene and aromatic systems from acetylene and alkynes has become a convenient tool in organic chemistry since Reppe proposed the use of transition metals as cycloaddition catalysts in 1948.⁴³³ These reactions make it possible to rapidly construct molecular scaffolds with various functional groups in a single step. For a long time, metal-catalyzed synthesis was limited to the trimerization of acetylene to benzene or led to complex mixtures of isomers in the case of substituted alkynes. For practical applications, product selectivity plays a key role. The creation of substituted benzenes is problematic due to the difficulty in controlling the chemoselectivity during metallacycle formation and regioselectivity upon the introduction of a third alkyne. Since then, the variety of proposed virtuous approaches for the synthesis of substituted benzene molecules using transition metalcatalyzed [2+2+2] cycloadditions has increased significantly.^{434,435} Half a century later, selective trimerization of three different alkyne components was achieved with catalysts based on zirconium⁴³⁶, titanium⁴³⁷ and ruthenium⁴³⁸. Catalyst systems based on cobalt⁴³⁹, rhodium⁴⁴⁰, ruthenium⁴⁴¹, titanium⁴⁴² and nickel⁴⁴³ help solve the problem of regioselectivity and produce 1,2,4-trisubstituted benzene rings. The use of a manganese catalyst made it possible to carry out the [2+2+2] cycloaddition of 1,3-dicarbonyls to terminal acetylenes.⁴⁴⁴ The replacement of manganese with a rhenium catalyst led to the formation of pyrone adducts instead of benzenes.445

In addition to benzene derivatives, transition-metal-catalyzed cyclocotrimerizations are powerful tools for the synthesis of pyridines and their derivatives. Cyclo(co)trimerization of acetylene with nitriles is a convenient path for accessing a wide range of 2-substituted pyridines. These transformations occur using cobalt complexes as catalysts.⁴⁴⁶

The formation of bicyclic product as a result of [2 + 2 + 2]-cycloaddition reaction between acetylene and diynes was studied by More et.al. ⁴⁴⁷ This transformation is catalyzed by cyclopentadienylcobalt dicarbonyl. Notably, metal carbonyls may be widely available in space, where CO is one of the most abundant molecules. Moreover, metal carbonyls were found in interstellar dust clouds⁴⁴⁸ and Jiange H5 chondrite⁴⁴⁹.

Another important catalytic process is the preparation of triazoles in the click reactions of acetylene and azides in the presence of Cu(I).⁴⁵⁰ In the presence of a Au(I) catalyst, it was recently discovered that acetylene can react as a dicarbene with alkenes, giving 1,1'-bi(cyclopropane).⁴⁵¹ Thus, acetylene is an excellent tool for the formation of a wide variety of cycles. These transformations were discussed in detail in a recent review devoted to cycloaddition reactions involving acetylene.¹⁷ We summarize the catalytic reactions considered in this section in Table 21.

Reactions	Catalytic metal	Objects & conditions
1. 2R ₃ -C ₂ H (1,3-dicarbonyl) + C ₂ HOH-R ₁ R ₂ (terminal acetylene) + Cat (Mn complex) \rightarrow C ₆ H ₃ OH-R ₁ R ₂ (R ₃) ₂ \rightarrow C ₆ H ₂ -R ₁ R ₂ (R ₃) ₂ (substituted benzene) + H ₂ O HO HO R ₁ H R ₃ H R ₃ H R R R H R A H R A A A A A A A A A A A A A A A A A A	Mn	not adapted to space objects
2. $2C_2H_2 + C_5N(R)_6 + Cat (Co complex) \rightarrow C_9N(R)_6 (substituted pyridine) + H_2$ $\parallel H_2$ $\downarrow H_2$ $\downarrow H_2$ $\downarrow H_2$ $\downarrow R_R$	Со	not adapted to space objects
3. diyne + C ₂ H ₂ + Cat (CpCo(CO) ₂) \rightarrow bicyclic product Cp - cyclopentadienyl HO HO HO HO HO HO HO HO HO HO HO HO HO	Со	not adapted to space objects
4. azide + C ₂ H ₂ + Cat (CuI) \rightarrow triazole $\downarrow \downarrow \downarrow \downarrow \downarrow \downarrow \downarrow$ N = $\downarrow \downarrow \downarrow \downarrow \downarrow \downarrow$ 4,4'-dicy anodiphenylazidomethane $\downarrow \downarrow \downarrow \downarrow \downarrow \downarrow$ $\downarrow \downarrow \downarrow \downarrow \downarrow \downarrow \downarrow \downarrow$ $\downarrow \downarrow \downarrow \downarrow \downarrow \downarrow \downarrow \downarrow$ $\downarrow \downarrow \downarrow \downarrow \downarrow \downarrow \downarrow \downarrow \downarrow$ $\downarrow \downarrow $	Cu	not adapted to space objects
5. alkene + C ₂ H ₂ + Cat (AuL ⁺) \rightarrow 1,1'-bi(cyclopropane) $R_1 \longrightarrow R_2$ $R_1 \longrightarrow R_2$ + $R_1 \longrightarrow R_2$ + or $H \longrightarrow R_1$	Au	not adapted to space objects

Table 21. Catalytic reactions known in industry but not adapted to astrochemical modeling.

7.2. Catalytic reactions involving acetylene in space

The presence of catalytic species can provide a variety of pathways for the formation of aromatic systems. Metal complexes that are used in laboratories and industry are rare in outer space, although ironnickel particles are present in meteorites and likely in cosmic dust.⁴⁵² The dust particles are either carbonaceous (HACs, graphites, and PAHs) or silicate. Silicate grains consist mostly of pyroxene (MgSiO₃) and olivine (Mg₂SiO₄).¹⁹⁷ Inclusions of aluminum, titanium and other metals were not excluded. Some carbides are also present in the ISM. Specifically, silicate carbide (SiC) is observed in many C-rich stellar envelopes⁴⁵³, and titanium carbide is speculated to also be present in these envelopes.^{454,455} Certainly, catalytic metals are present in the ISM in the gas phase as single atoms or ions, which are traced through their absorption/emission lines.⁴⁵⁶ However, it is doubtful that these materials can act as catalysts in the gas phase, as three-body collisions are assumed to occur, which are quite rare under interstellar conditions.

In 2015, Zhao et al. demonstrated the catalytic conversion of acetylene to PAHs over SiC grains⁴⁵⁷. They observed the formation of aromatic molecules containing up to 20 carbon atoms and even a surface graphene-like coating in their experiments. The authors reported that acetylene activation occurs with the participation of dangling bonds on SiC surfaces, which are potential active sites for chemisorption and activation (Scheme 19). In the first step, acetylene adsorbs onto a silicon dangling bond to generate a silylacetenyl radical. Then, another acetylene molecule attaches to the formed radical to generate a 1-buten-3-ynyl radical, and the next acetylene molecule attaches to the diene. Finally, in the next step, the carbon radical attacks a neighboring Si dangling bond to form an octagonal ring. This ring has an unstable resonance structure with a positive and a negative charge at each end of the triene. The octatomic ring can undergo a rearrangement process to form a benzene ring to obtain the most stable structure. This process likely occurs in the dust formation zone because SiC grains are present there. Additionally, SiC was also found in presolar meteorites and comet materials returned to Earth by the Stardust mission⁴⁵⁸; therefore, the process may take place in other environments in addition to stellar envelopes.



Scheme 19. The proposed mechanism for the catalytic formation of benzene from acetylene on a SiC particle. Reproduced with permission from ref 455. Copyright 2016 Royal Society of Chemistry.

The source of SiC can also be acetylene. Previously, Parker et al. experimentally showed that the silacyclopropenylidene (c-SiC₂H₂) molecule can be synthesized in the gas phase under single-collision conditions via the reaction of a silylidine radical (SiH) with acetylene. Later, Yang et al.⁴⁵⁹ also investigated the reactions between silicon atoms and allene (H₂CCCH₂) and methylacetylene (CH₃CCH), and they obtained SiC₃H₂. In the above works, the authors noted that c-SiC₂H₂ and c-SiC₃H₂ can be photolyzed to carbon-silicon clusters such as silicon c-SiC₃ and c-SiC₂, respectively; thus, they can be precursors of SiC dust grains in the outflows of C-rich AGB stars.⁴⁶⁰

A recent study³⁰ showed that acetylene can be adsorbed on cosmic dust analogs in the temperature range found in Titan's atmosphere below 600 km. Compounds such as Mg₂SiO₄, MgFeSiO₄ and Fe₂SiO₄ can act as catalysts for the cyclotrimerization of acetylene to benzene. The authors showed that C₆H₆ production via C₂H₂ uptake on cosmic dust followed by cyclotrimerization and desorption is likely to be a competitive pathway between 80 and 120 km with gas-phase C₆H₆ production. Other recent studies have also shown that siliceous rocks containing siloxyl radicals can act as catalysts for complex organic matter synthesis.⁴⁶¹ The assumption about the formation of PAHs in interstellar clouds from acetylene due to a metal-catalyzed process was first put forward in 1992. The authors assumed that the formation of PAH organometallic compounds with Fe, Co, Ni, and other metals could be responsible for the observed metal transition depletion.⁴⁶²

Murakami et al. conducted a DFT study simulating acetylene trimerization in the ISM in water ice. The potential energy profiles of this reaction are given in Figure 21. The authors found that in the presence of acetylene and the $Fe^+(H_2O)_n$ cluster as a catalyst, benzene can be formed in one step, including one

transition state structure of three acetylene molecules. Moreover, in the case of single-atom catalysis by Fe⁺, benzene was formed in two stages, including the formation of one C–C σ -bond and two subsequent formations of C–C σ -bonds.⁴⁶³ Subsequently, the authors expanded this study to include other transition metals: Sc, Ti, Mn, Co and Ni.⁴⁶⁴ The number of electrons in the 3d orbitals of the transition metal cation significantly contributed to the catalytic efficiency of the acetylene cyclotrimerization process. The systems with Mn⁺, Co⁺ and Ni⁺ had stable structures in which three acetylene molecules were coordinated to the cationic center of the transition metal. In this case, the energy levels of the transition complexes were higher than the energy levels of the initial reagents. In the case of systems with Sc⁺ and Ti⁺ cations, there is no significant barrier to the reaction path. That is, isolated acetylene and (C₂H₂)₂, together with Sc⁺ and Ti⁺ cations associated with the water cluster (H₂O)₈ and NC⁻, can form a benzene-metal-cation complex without an entry barrier. Thus, according to calculations, Sc⁺ and Ti⁺ cations act as effective catalysts for the cyclotrimerization of acetylene. In Table 22, we summarize the catalytic reactions that were considered in astrochemistry.



Figure 21. Potential energy profiles for the benzene formation by acetylene cyclotrimerization reactions catalyzed by Fe^+ and cluster $Fe^+(H_2O)_8$ obtained at the B3LYP(D3BJ)/def2-SVPP DFT level. Reproduced with permission from ref 461. Copyright 2022 MDPI.

Reactions	Kinetics	Objects & conditions
1. SiC + $3C_2H_2 \rightarrow SiC + C_6H_6^{[457]}$	n.d.	Stellar envelopes. Conditions: $n_H = 10^8 \cdot 10^{15} \text{ cm}^{-3}$ $T = 500 \cdot 2500 \text{ K}$

Table 22. The catalytic reactions which were considered in astrochemistry.

2. a. SiH + C ₂ H ₂ \rightarrow c-SiC ₂ H ₂ b. c-SiC ₂ H ₂ + energy (UV, shocks) \rightarrow c-SiC ₂ + H ₂ ^[460]	n.d.	a) Stellar envelopes. Conditions: $n_{\rm H} = 10^8 \cdot 10^{15} \text{ cm}^{-3}$ T = 500-2500 K
		b) Stellar envelopes (outer edge). Conditions: $n_H = 10^8 \cdot 10^{12} \text{ cm}^{-3}$ $T = 500 \cdot 1500 \text{ K.}$ + PDRs, DISM. Conditions: T = 10 - 500 K $n_H = 10^{-1} \cdot 10^5 \text{ cm}^{-3}$ $G_0 = 1 \text{ to } 10^5$ Galactic CR field.
3. a. Si + C ₃ H ₄ \rightarrow c-SiC ₃ H ₂ + H ₂ b. c-SiC ₃ H ₂ + hv \rightarrow c-SiC ₃ + H ₂ ^[459]	provided many pathways in the work of Yang ^[459]	The same as reactions #2
4. $3C_2H_2 + M(Mg_2SiO_4, MgFeSiO_4 \text{ or}$ Fe ₂ SiO ₄) $\rightarrow C_6H_6^{[30]}$	$k_4 = 1.2 \cdot 10^{-5} \text{ cm}^3 \text{ s}^{-1}$ (T = 150 K) ^[30]	Titan's atmosphere. Conditions: $n_{\rm H}$: 10^{10} - 10^{16} cm ⁻³ T= 100-200 K G ₀ : up to 10^2 Galactic CR field.
5. Ice reactions. $3C_2H_2 + M(Fe^+(H_2O)_n) \rightarrow C_6H_6 + M(Fe^+(H_2O)_n)^{[463]}$	n.d.	Molecular clouds, prestellar cores, comets. Conditions: $n_H = 10^2 \cdot 10^7 \text{ cm}^{-3}$ T = 10 K $G_0 = \text{very low}$ Galactic CR field.

7.3. PAHs as a catalyst

The active development of carbocatalysis in the last decade has shown that important catalytic processes can proceed even without the participation of metals. Several decades ago, PAHs were noted to be appropriate for H₂ formation in PDRs.^{465,466} Functionalized PAHs with oxygen-bearing groups can be a source of small oxygen-bearing molecules under conditions of UV irradiation in the ISM.^{467,468} Recently, it has been shown that PAHs can also be catalysts for many other hydrogenation reactions in the ISM, resulting in the synthesis of molecules such as H₂O and HCO.⁴⁶⁹ Similarly, under certain conditions, PAHs can play a role in the synthesis of benzene from simple precursors. Recently, published DFT calculations and experimental data have confirmed the formation of benzene by trimerization of three acetylene molecules catalyzed by PAH carbene sites at ~900 K.⁴⁷⁰ The authors considered the carbene sites on the zigzag edge of PAHs to be catalytic sites. The considered PAHs of various sizes and configurations from C_6H_6 to $C_{361}H_{51}$.

Quantumochemical calculations of the reaction mechanism made it possible to reconstruct the free energy profiles and visualize the spatial distributions of the spin density for all stages of benzene formation from three acetylene molecules on the carbene centers of PAH molecules (Figure 22). The stepwise addition of acetylene molecules at the carbene center was promoted by continuous spin migration onto the β -carbon of the added alkyne moiety. It has been shown that the trimerization process will dominate linear oligomerization after the addition of three acetylene molecules due to the free energy driving force. Due to the high strength of the C–C bond in the intermediate complex, the removal of benzene and regeneration of the carbocatalyst can be kinetically difficult. The authors considered two possible mechanisms for the

elimination of benzene from a PAH. In the associative pathway, in which the benzene molecule is competitively replaced by an acetylene molecule, a higher activation energy was observed. The associative mechanism is characterized by the predominant localization of unpaired electrons at the carbene center. The dissociative pathway has a lower activation energy and is accompanied by the delocalization of unpaired electrons in a polyaromatic system. The authors also noted that the removal of a benzene molecule by the dissociative mechanism for monoradicals is characterized by a very low activation energy $(1.0-3.0 \text{ kcal mol}^{-1})$ compared to that of carbenes (~26.0 kcal mol⁻¹). Therefore, polyaromatic monoradicals can be promising candidates for carbocatalysts in ISM, although thorough adaptation of this reaction to astrophysical conditions is needed. Interestingly, the catalytic cycle was controlled by reversible fluctuations in the spin density. These vibrations were responsible for the formation of benzene and the reactivation of active sites. This reaction is shown in Table 23, although the kinetics of this reaction, which can be used in astrochemistry, have yet to be determined.



Figure 22. Free energy profile for the acetylene cyclotrimerization reaction catalyzed by $C_{37}H_{14}$. The associative pathway of the product elimination is shown in red; the dissociative pathway is shown in blue. The initial form of the $C_{37}H_{14}$ catalyst, location of the chemically reactive center, and designation of this $C_{37}H_{14}$ structure in the catalytic cycle representations are shown in the magenta frame. Reproduced with permission from ref 468. Copyright 2020 American Chemical Society.

Notably, carbenes are important intermediates in processes leading to carbon nanostructures in the ISM. Previously, carbene species were identified in the interstellar space.⁴⁷¹ Recently, Mebel et al., using crossed molecular beam data with electronic structure calculations and quasiclassical trajectory simulations, reported the fundamental reaction mechanisms leading to the formation of highly reactive carbene species under single-collision conditions relevant to extreme astrochemical conditions.⁴⁷² As a

model, the authors used the reaction of methylidine (CH) with diacetylene (HCCCCH) under single collision conditions to form triplet pentadiynylidene (HCCCCCH) and singlet ethynylcyclopropenylidene ($c-C_5H_2$) carbenes. Considering the prevalence of carbenes in various cosmic media, their role in promoting the evolution of high-molecular-weight organic compounds can be underestimated. The mechanism of the catalytic trimerization of acetylene on the active sites of PAHs and carbon dust particles could be one of the possible ways to synthesize hydrocarbons on aerosol particles and organic haze layers in hydrocarbon-rich atmospheres on planets and their moons. The role of catalytic processes in the evolution of hydrocarbon molecules on the surface of interstellar dust particles, ice, or atmospheres of planets and satellites has yet to be assessed.

Reactions	Kinetics	Objects & conditions
$3C_2H_2 + PAHs \rightarrow C_6H_6 + PAHs^{[470]}$	n.d.	Stellar envelopes (outer edge). Conditions: $n_H = 10^8 \cdot 10^{12} \text{ cm}^{-3}$ T = 500-1500 K

Table 23. The reaction of acetylene trimerization on PAHs.

8. Acetylene in prebiotic chemistry evolution

8.1. Acetylene in reactions producing prebiotic molecules

One important aspect that determines the significance of acetylene chemistry research in space is its possible role in prebiotic chemistry evolution. Acetylene could play an important role in the origin of life, and the presence of acetylene in planetary atmospheres may indicate the possible existence of an extraterrestrial anaerobic ecosystem.⁴² Acetylene may serve as a signature for young Earth-like exoplanets with reducing atmospheres undergoing heavy bombardment.¹²⁶ Such conditions are considered to be similar to those that were intrinsic to the primordial Earth during the Hadean Eon (>4 Gyr ago), when chemical evolution and prebiotic accumulation took place. Theoretical models of the planetary atmosphere and surface that consider photochemical reactions predict steady-state surface C_2H_2 quantities of 0.4% for Earth-like young planets.¹²⁶

HCN is considered one of the key precursors of prebiotic molecules. In turn, an important source of HCN is the photolysis of a mixture of C_2H_2 and NH_3 in planetary atmospheres at temperatures of 130-298 K.⁴⁷³ Recent studies of the photolysis mechanisms of NH_3/C_2H_2 mixtures have shown that this process can be a source of many organonitrogen compounds that are important for prebiological organic chemistry.⁴¹ Under similar conditions, the formation of so-called tholins⁴⁷⁴, which are nitrogen-containing polymeric compounds that are formed on the surface of many planets and icy bodies of the Solar system, primarily Titan, can occur. ^{283,375} The possible role of tholins in the evolution of the prebiotic chemistry of planets, including Pluto⁴⁷⁵, has been noted.⁴⁷⁶ It is likely that tholin formation proceeds through acetylene chemistry analogous to that of cyanopolyynes (see Section 3.2) based on the hypothesized structures of the tholins. However, this pathway has yet to be determined.

In a number of recent experimental and theoretical works, Cable et al. demonstrated the formation of acetylene co-crystals with various organic molecules on the surface of Titan and their influence on the landscape of the satellite. It was experimentally shown that cocrystal C_2H_2 -NH₃ can be formed on Titan at 90 K.⁴⁷⁷ In addition, DFT calculations predicted the existence of three-component co-crystals with the composition 2c-C₆H₆:C₂H₂:HCN under the surface conditions of Titan.⁴⁷⁸ Cable et al.⁴⁷⁹ investigated the formation of co-crystals of acetylene (C₂H₂) and acetonitrile (CH₃CN) on the surface of Titan. The authors noted that similar molecular minerals can be widely distributed in the labyrinth terrains of Saturn's moon. Finally, the same authors⁴⁸⁰ demonstrated the stability of a cocrystal of acetylene and pyridine in the temperature range from 90 K to 180 K. The authors suggested that such acetylene cocrystals could provide concentration, storage and transport of prebiotic and high-energy molecules for the functioning of hypothetical biological systems.

A number of papers have been devoted to formulating speculative hypotheses about the presence of life on Titan using acetylene as an energy source, citing the deficiency of C_2H_2 and H_2 on the satellite's surface as a possible biosignature.^{487,481,482,483,484,485} The proposed models of acetylene metabolism on Titan included the following exothermic reactions (Scheme 20).

A $C_2H_2 + 2H_2 \longrightarrow C_2H_6 \Delta G_0 = -242 \text{ kJ}$

B C_2H_2 + $3H_2$ \longrightarrow $2CH_4$ $\Delta G_0 = -375 \text{ kJ}$

Scheme 20. Proposed models of acetylene metabolism on Titan.

Such metabolism could be stimulated by a strong thermodynamic benefit. However, a number of factors make the existence of life under such conditions unlikely. One of the factors is the low solubility of large molecules in liquid hydrocarbons, which creates problems for the functioning of biopolymers. This factor becomes even more critical given the extremely low temperatures.⁴⁸⁶ In addition, calculations of the biomass value predicted for life on Titan based on similar biochemistry on Earth have shown that the existence of bacteria using acetylene as an energy source on Titan is unlikely.⁴⁸³

Finally, Abbas and Schulze-Makuch proposed three interesting, albeit obviously flawed, pathways for the formation of amino acids as essential prebiogenic components involving acetylene in hydrothermal vents or subsurface Titan.⁴⁸⁷ Catalytic reactions play a key role in the first model, which is illustrated in Scheme 21. The first stage was the trimerization of acetylene with the participation of metal catalysts or clay minerals and zeolites. The formation of substituted benzene could occur due to the participation in the reaction of recombination products of acetylenyl radicals, for example, with methyl or ethynyl radicals. The subsequent formation of nitro-substituted alkylbenzene, according to the authors, could occur as a result of interactions with highly active NO₂⁺ cations that appeared as a result of ionizing radiation. During the next step, the nitro group with hydrogen is converted to an amino group in the presence of a metal catalyst, and an alkyl aminobenzene is subsequently formed. Finally, oxidation of the alkyl radical to a carboxyl group led to the formation of the corresponding amino acid.



Scheme 21. Formation of amino acids from acetylene according to model I by Abbas et al.

In the second model, acetaldehyde is formed from acetylene in the presence of peroxide and hindered borane, which is illustrated in Scheme 22. The authors admit that the presence of hydrogen peroxide on the surface of Titan in significant quantities is unlikely. However, these findings indicate the possibility of peroxide formation from water under the influence of UV radiation. The aldehyde that is formed from acetylene or its derivatives must further react with ammonia and hydrogen cyanide to form the α -amino nitrile. Acid hydrolysis of this compound under hot spring conditions can lead to the formation of an amino acid such as alanine.



Scheme 22. Formation of alanine from acetylene according to the model II by Abbas et al.

The third model (shown in Scheme 23) is associated with the formation of an alkynyl radical and a halogen radical under the influence of UV radiation. These interactions will lead to the appearance of alkynyl halides. In turn, the reaction of an alkynyl halide as a result of a substitution reaction can give the corresponding nitrile. The corresponding carboxylic acid will result from hydrolysis of the nitrile in an acidic environment. The addition of ammonia to a double bond results in the corresponding amino acid. The authors suggested that the first route of amino acid synthesis is the most relevant to the conditions of Titan due to the lack of liquid water.



Scheme 23. Formation of alanine from acetylene according to the model III by Abbas et al.

Considering the role of acetylene in the formation of key prebiotic organics on premordial Earth, acetylene could be expected to be effective in reducing the CO₂- and H₂-rich atmosphere of primitive Earth. In this case, UV photolysis and volcanic lightning produce CH₄ and other light hydrocarbons, which, in turn, can form HCN and other N-species via reactions with dissociated atmospheric N₂. These simple organic molecules can be converted into prebiotic species (nucleobases, fatty acids, amino acids, alcohols, etc.) via aqueous synthesis. In a reducing atmosphere, acetylene could appear in significant quantities. For example, Segura et al. ⁴⁸⁸ modeled the chemical processes that occur under the influence of volcanic lightning under conditions of explosive volcanism. Volcanic lightning was simulated by focusing a high-energy infrared laser beam on a volcanic gas mixture consisting of 64% CH₄, 24% H₂, 10% H₂O and 2% N₂. The authors relied on an accretion model and measured the nitrogen content in Martian meteorites. Among the products, 11 hydrocarbons were identified, and the most common compound was acetylene. Rimmer et al.¹²⁶ studied potential processes that could occur in the atmosphere of a young Earth-like exoplanet under conditions of protoplanetary disk collapse. A mixture of methane (CH₄), carbon monoxide (CO), nitrogen (N₂) and water was used as a model atmosphere. In the simulation of the high-speed impact

of an extraterrestrial body on the early atmosphere of a planet, significant quantities of substances such as acetylene (8%), hydrogen cyanide (5%), and cyanoacetylene (5%) form inside the reactor as does ammonia (1%). The authors concluded that acetylene can be considered one of the signatures of the impacts of bodies from protoplanetary disk on planets with reducing the atmospheres.

Similar chemical processes could have occurred on the early Earth. However, according to modern geophysical studies, after the solidification of the magma ocean after the Moon-forming impact, the atmosphere of Hadean Earth should have been rather neutral, CO₂-rich and H₂-poor.^{489,490,491} Under these conditions, organic syntheses from feedstock atmospheric gases became much less effective⁴⁹², and the delivery of organics via giant impacts or localized synthesis during volcanic eruptions likely became a key process.

Acetylene and other key feedstock molecules for prebiotic chemistry can be produced in hydrothermal and volcanic springs. Currently, hydrocarbon gases can be found in the fumaroles of geothermal areas.⁴⁹³ Acetylene has also been detected in Hawaiian volcanic glasses.⁴⁹⁴ Moreover, modern volcanic activity is very different from the volcanic activity of the primordial Earth. Early experimental simulations of volcanic processes on primordial Earth revealed the formation of unsaturated compounds such as acetylene, propylene, vinyl acetylene, methyl acetylene, and butadiene.⁴⁹⁵ Rimmer et al.⁴⁹⁶ calculated the chemical composition of volcanic gases feeding shallow and superficial vents over a range of pressures and C/N/O ratios in basalt. In the case of an ultrareduced and carbon- and nitrogen-rich gas phase of magma, interactions with groundwater in volcanic vents lead to 10^{-3} –1 mol l⁻¹ concentrations of diacetylene (C₄H₂), acetylene (C₂H₂), cyanoacetylene (HC₃N), hydrogen cyanide (HCN), bisulfite (HSO³⁻), hydrogen sulfide (SH⁻) and soluble iron in wastewater (Figure 23).



Figure 23. Chemistry of a surface hydrothermal vent fed by carbon- and nitrogen-rich magma. The yield of diacetylene (C_4H_2), acetylene (C_2H_2), cyanoacetylene (HC_3N), hydrogen cyanide (HCN), hydrogen (H_2) and methane (CH_4) from the pool is shown in ratios determined by the depth at which the gas is located. The interaction of species under the influence of UV leads to the formation of a small amount of cyanamide (NH_2CN). Reproduced with permission from ref 494. Copyright 2019 MDPI.

Acetylene could have been an energy source for early microorganisms on Earth and could also be an energy source for hypothetical life forms on other planets. *Pelobacter acetylenicus* was discovered in 1985 using acetylene as the sole source of carbon and energy.^{497,498} *Pelobacter acetylenicus* can only grow by acetylene. The key enzyme of these bacteria is acetylene hydratase.⁴⁹⁹ Acetylene hydratase contains transition metals, one [4Fe-4S] cluster and one tungsten atom, coordinated by two molybdopteringuanosine dinucleotide ligands.⁵⁰⁰ During the initial stage of acetylene fermentation, this enzyme catalyzes the hydration of acetylene to acetaldehyde through an enol intermediate. Acetaldehyde is transformed into acetyl-coenzyme CoA. Acetyl-CoA is subsequently used to construct metabolites with C₂-units. At subsequent stages, the formation of ethanol, acetate and hydrogen occurs (Scheme 24). Therefore, acetylene has the potential to fuel complete metabolism on its own.

acetylene hydrating $C_2H_2 + H_2O \xrightarrow{enzyme system} CH_3CHO$

 $2CH_3CHO + H_2O \longrightarrow C_2H_5OH + CH_3COO^- + H^+ + 1/2H_2$ Scheme 24. Acetate formation from acetylene by bacterial metabolism.

The biochemistry of such organisms is much simpler than that of methanogens, which may indicate that acetylene may be the dominant food for primitive organisms early in evolution.⁵⁰¹ Considering these findings, Oremland and Voytek called acetylene a 'fast food' for early life and suggested that such microorganisms that consume acetylene could play a key role in the evolution of the Earth's early biosphere.⁴² In Table 24, we provide the results of prebiotic chemistry discussed in this section.

Reactions	Kinetics	Objects & conditions
1. Mixture $C_2H_2/NH_3 + h\nu \rightarrow [a]$ chain of reactions] $\rightarrow HCN + others^{[41]}$	Rate constants can be found or calculated for the most of reactions in the chain	Atmospheres of Jovian planets and comets. Conditions: $n_H = 10^{10}-10^{16} \text{ cm}^{-3}$ T = 50-300 K $G_0 = \text{up to } 10^2$ Galactic CR field
2. $C_2H_2 + NH_3 \rightarrow \text{co-crystal}$ $C_2H_2:NH_3^{[477]}$ 3. $2c-C_6H_6 + C_2H_2 + HCN \rightarrow \text{co-crystal}$ $2c-C_6H_6:C_2H_2:HCN^{[478]}$ 4. $C_2H_2 + CH_3CN \rightarrow \text{co-crystal}$ $C_2H_2:CH_3CN^{[479]}$	n.d.	Titan's atmosphere. Conditions: $n_H = 10^{10}-10^{16} \text{ cm}^{-3}$ T = 60-300 K $G_0 = up \text{ to } 10$ Galactic CR field
$5. C_{2}H_{2} + 2H_{2} \rightarrow C_{2}H_{6}$ $C_{2}H_{2} + 3H_{2} \rightarrow 2CH_{4}^{[485]}$	n.d.	Titan's atmosphere, planetary atmospheres, probably early Earth.

Table 24. Summary of considered reactions, kinetics (if available) and conditions where the reactions are relevant.

Table 24. Summary of considered reactions, kinetics (if available) and conditions where the reactions are relevant.

Reactions	Kinetics	Objects & conditions
6. $C_2H_2 + H_2O$ [acetylene hydrating enzyme system] $\rightarrow CH_3CHO$ $2CH_3CHO + H_2O \rightarrow C_2H_5OH + CH_3COO^- + H^+ + 1/2H_2^{[499]}$		Conditions: $n_{\rm H} = 10^{10} \cdot 10^{16} \text{ cm}^{-3}$ $T = 50 \cdot 300 \text{ K}$ $G_0 = \text{up to } 10^2$ Galactic CR field.

8.2. From acetylene to nitrogen-bearing PAHs and nucleobases

Previously, we described the mechanisms of PAH formation from acetylene. In this section, specific PAHs that are connected with life precursors are considered. Many COMs, including prebiotic molecules, are formed in astrophysical ices; consequently, the ices and chemical reactions inside them are the subject of interest. Most PAHs are depleted in ices in dense and cold molecular clouds. A fraction of PAHs can attach different functional groups (O, CN, NH₂, CH₃, COOH, etc.) instead of peripheral H atoms.^{502,503,504} Taking into account that the ices mostly consist of water and that there are also other oxygen-bearing molecules such as methanol, oxygen-bearing functional groups, -OH and =OH, are highly expected. Consequently, quinones, ethers and aromatic alcohols are also expected.⁵⁰⁵ For example, the formation of quionone (1,4-napthaquinone) in water ice was traced experimentally⁵⁰³, and this quinone performs important biochemical functions, such as electron transport and oxidative phosphorylation.

Apart from functional groups, carbon atoms in PAHs can be replaced by nitrogen or oxygen. Nitrogen (N-) heterocycles, pyrimidine ($C_4H_4N_2$) and purine ($C_5H_4N_4$), are the main components of DNA and RNA nucleobases. N-heterocycles, associated nucleobases and other biorelevant molecules have been found in meteorites.^{506,507,508,509,510,511,512,513} The detection of ¹⁵N/¹⁴N isotope enrichment⁵¹⁴ as well as the terrestrially rare nucleobases 2,6-diaminopurine and 6,8-diaminopurine in Murchison^{508,509} points to an extraterrestrial origin. N-heterocycles have not yet been detected in the ISM, although some observations can be attributed to their presence. Specifically, Hudgins et al. showed that the commonly observed PAH band at 6.2 µm can be associated with traces nitrogen-substituted PAHs.⁵¹⁵

Previously, it was shown that nitrogenous bases are unlikely to be formed and survive in interstellar and circumstellar environments, as these compounds are highly sensitive to UV radiation. It was estimated that gas phase N-heterocycles are destroyed within 10–100 years in diffuse ISM⁵¹⁶. Their lifetime does not exceed a few hours at a distance of 1 AU from the Sun either. The presence of N-heterocycles at least in meteorites means that there must be mechanisms and conditions that make possible their formation and survival outside the Earth.

Research by Parker et al. showed that the formation of (iso)quinoline (C₉H₇N) probably proceeds with the participation of acetylene molecules in the circumstellar envelopes of evolved C-rich stars. Experiments in a pyrolytic reactor in combination with DFT calculations showed that the formation of (iso)quinoline can occur as a result of the reaction of the *m*-pyridyl radical in series with two molecules of acetylene.⁵¹⁷ In turn, as shown by Kaiser et al., pyridine can be obtained through barrierless elemental reactions: 1) by the reaction of vinyl cyanide (C₂H₃CN) with cyanovinyl (HC=CH–CN)⁵¹⁸ and 2) by the reaction of cyano radical (CN) with 1,3-butadiene (C₄H₆)^{519,520}.

Under low-temperature conditions in molecular clouds (for example, TMC-1 and OMC-1), planetary atmospheres and their satellites (for example, Titan and Pluto), quinoline and isoquinoline (C_9H_7N) can be synthesized via fast, de facto barrierless reactions involving *o*-, *m*-, and *p*-pyridinyl radicals (C_5H_4N) with vinylacetylene (C_4H_4), similar to the HAVA mechanism of naphthalene formation. Molecular beam experiments combined with electronic structure calculations of the reaction (Figure 24) conducted by Mebel et al. point to this mechanism.⁵²¹



Figure 24. Computed potential energy surface for the reaction of p-pyridinyl (C_5H_4N) radical with vinylacetylene (C_4H_4); the energies calculated at the G3(MP2,CC)//B3LYP/6-311G(d,p)+ZPE level of theory. Reproduced with permission from ref 519. Copyright 2021 Royal Society of Chemistry.

Ricca et al. ⁵²² carried out a computational study of possible mechanisms for the formation of PAHs containing nitrogen atoms in Titan's stratosphere. According to their model, the synthesis of pyridine (C_5H_5N) is driven by radical reactions of HCN with C_2H_2 . The authors noted that mechanisms involving the addition of HCN and C_2H_2 to a pure aromatic ring to form an additional ring have barriers lower than 15 kcal/mol. Such barriers are probably too high for Titan, where the temperature is approximately 70-200 K⁵²³, to allow reactions to occur. However, aromatic molecules are strong UV absorbers and can efficiently convert this energy into vibrational energy.⁵²⁴ Therefore, the authors suggested that the reactions are possible even at 170 K due to the vibrational energy. The reducing nature of Titan's atmosphere could help reconstruct abiogenetic processes on the early Earth, although chemical evolution on Titan is limited by surface temperatures that are too low, which hinders the development of biochemical systems.

Rap et al.⁵²⁵ in 2022 demonstrated another possible low-temperature pathway for the formation of nitrogen-containing PAHs, highlighting the underestimation of ion–molecule reactions (Scheme 25). A study of the ion–molecule reaction of pyridinium radical ($C_6H_5N^+$) with acetylene showed branched pathways for the formation of ionic species such as quinolizinium and H-quinolinium. The intermediate structures of the 2-ethynyl-N-pyridine cation, 2-vinyl- β -distone-N-pyridine cation, quinolysinium cation and H-quinoline cation were identified by mass spectrometry.


Scheme 25. Mechanisms of reaction pathways to N-PAHs via ion–molecule reaction of pyridinium radical and acetylene. The different reaction types are designated: radiative association (RA), bimolecular reaction (BI) and termolecular association (TER). Reproduced with permission from ref 523. Copyright 2022 Springer Nature.

Menor-Salvá et al. ⁵²⁶ showed that UV irradiation (in the range of 185–254 nm) of a urea solution subjected to freeze-thaw cycles (from -21 °C to +5 °C) in C₂H₂ leads to the formation of purines, pyrimidines and hydantoin. Guanine, cytosine, and uracil were detected using GC-MS. The authors explained the absence of adenine by the possible derivatization of adenine and guanine with the formation of 8-hydroxyadenine, the peak of which was detected. Moreover, Jeilani et al.²⁹ investigated possible pathways for the formation of pyrimidine bases such as cytosine and uracil under cold conditions relevant to Titan chemistry. The mechanisms for the formation of pyrimidine bases from urea, acetylene and ethynyl radicals were studied by DFT. The authors showed that the precursor of uracil and cytosine may be the biuret molecule. In addition, the pathways for the formation of melamine, ammeline, ammelide, cyanuric acid, uric acid and 8-oxoguanine are shown in Figure 25.



Figure 25. Formation of pyrimidine bases and triazines from urea and acetylene. Red ovals indicate atoms from acetylene.²⁹

The synthesis of cytosine from urea and cyanoacetaldehyde was also examined.⁵²⁷ Cyanoaldehyde may be a product of the hydrolysis of cyanoacetylene, which, in turn, is a derivative of acetylene^{528,529} (cyanoacetylene can be produced by spark discharge in a CH_4/N_2 mixture).^{530,531} Moreover, purine bases can be the product of the reaction of cyanoacetylene with cyanamide through barrierless cyclization through a small number of stages.⁵³²

Another important class of nitrogenous organic compounds is pyrolles. Pyrol structures underlie important biological molecules, such as heme, chlorophyll, cobalamin and others. In his recent work, Seitz et al.⁵³³ demonstrated that pyrrole and dimethylpyrrole can be formed under the simulated volcanic hydrothermal conditions of the Early Earth. As was shown, the starting materials for such important molecules could be acetylene, propine and ammonium salts, the transformations of which were achieved by NiS or CoS as catalysts (Scheme 26).



Scheme 26. Proposed mechanism for the formation of pyrrole from acetylene in the presence of Ni. Begins with the binding of ammonia to the Ni ion (A), begins with the coordination of acetylene on nickel (B). Reproduced with permission from ref 531. Copyright 2021 MDPI.

The important prebiotic role of acetylene and nitrile interactions was noted by Kaiser.²¹⁷ In addition, the mechanisms of the formation of polyaromatic compounds discussed in this review also indicate the important role of acetylene in biochemical evolution from the standpoint of the PAH-world hypothesis. Platts⁵³⁴ and Chen et al. suggested⁴³ that certain types of PAHs can specifically interact with each other. Ionic and π -stacking interactions, as well as hydrogen bonds, can promote the formation of highly ordered structures. Thus, a matrix block polymer with a certain sequence of PAH fragments can be formed. In turn, various neighboring complementary blocks of PAHs, such as polynucleotide sequences, can provide blockspecific replication. The distance between self-ordering PAH species and between nucleotides of RNA and DNA in both cases is approximately 0.34 nm. UV irradiation of astrophysical ice analogs (H₂O, NH₃, and CH₄) with pyrimidine or purine led to the formation of the main nucleases, such as uracil, cytosine, thymine, adenine, guanine and their isomers.^{535,536,537,538,539} In addition, pyrimidine and purine-based nucleases can be formed in a simple astrophysical ice analog consisting of a mixture of H₂O, CO, NH₃, and CH₃OH, i.e., initially without pyridine or purine⁵⁴⁰; thus, the formation of pyrimidines and purines and nucleobases containing these compounds seems to be inevitable. Finally, according to astrochemical modeling, PAHs substantially increase the abundance of several COMs and sulfur-bearing molecules⁵⁴¹, while the latter are considered to be important for several amino acids.^{542,543} We summarize the main prebiotic reactions discussed in this section in Table 25.

Reactions	Kinetics	Objects & conditions	
Ice reactions 1. $C_{10}H_8 + H_2O$ ice $\rightarrow C_{10}H_6O_2$ (1,4- naphthoquinone) ^[503]	n.d.	Molecular clouds, prestellar cores. Conditions: T = 10 K $G_0 = \text{very low}$ Galactic CR field. + Comets, meteorites.	
2. $C_5H_4N + 2C_2H_2 \rightarrow C_9H_7N$ (isoquinoline) ^[517]	n.d.	Stellar envelopes (outer edge). Conditions: $n_H = 10^8 \cdot 10^{12} \text{ cm}^{-3}$ $T = 500 \cdot 1500 \text{ K}$	
3. $C_2H_2CN + C_2H_3CN \rightarrow C_5H_5N$ (pyridine) + $CN^{[518]}$	n.d.	Stellar envelopes (outer edge). Conditions: $n_H = 10^8 \cdot 10^{12} \text{ cm}^{-3}$ $T = 500 \cdot 1500 \text{ K}$ Molecular clouds, prestellar cores. Conditions: $n_H = 10^2 \cdot 10^7 \text{ cm}^{-3}$ T = 10 K $G_0 = \text{very low}$ Galactic CR field.	
4. $CN + C_4H_6 \rightarrow C_5H_5N + H \text{ (pyridine)}^{[520]}$	$k_4 = 10 - 10^4 \text{ s}^{-1}$ ~10 - 10 ⁴ s ⁻¹ (depending on collision energy)	Molecular clouds, prestellar cores. Conditions: T = 10 K $G_0 = very low$ Galactic CR field. + Comets, meteorites.	
5. $C_5H_4N_{\cdot} + C_4H_4 \rightarrow C_9H_7N + H$ (isoquinoline) ^[521]	n.d.		
6. HCN + 2 C ₂ H ₂ \rightarrow C ₅ H ₅ N (pyridine) ^[522]	n.d.	Titan's atmosphere. Conditions: $n_{\rm H}$: 10 ¹⁰ -10 ¹⁶ cm ⁻³ T = 50-300 K G_0 = up to 10 Galactic CR field.	
7. Ice reactions $H_2O/CH_4N_2O(\text{urea})/C_2H_2 + hv \rightarrow \text{guanine},$ adenine, xanthine, uric acid, uracil, cytosine ^[526]	n.d.	Planetary atmospheres, comets, meteorites Conditions: $n_H=10^{10}-10^{16}$ cm ⁻³ T = 50-300 K $G_0 = up$ to 10^2 Galactic CR field.	

Table 25. The reactions of prebiotic chemistry with PAHs.

8.3. Role of acetylene in monocarboxylic acid formation

In addition to amino acids and nucleobases, fatty acids are considered important components in the formation of the first protocellular structures.^{544,545} A number of studies^{546,547,548} suggest that the first membranes encapsulating information polymers consist of fatty acids. Like amino acids and nitrogenous bases, fatty acids are found in meteorites. Moreover, fatty acids are the most common water-soluble organic compounds in carbonaceous chondrites. ⁵⁴⁹ Fatty acids with chain lengths of 2–12 carbon atoms have been found in meteorites. In water, these molecules spontaneously assemble into bilayers that form vesicles. ⁵⁵⁰

There are many possible noncatalytic and catalytic mechanisms for the natural formation of fatty acids in meteorites and under primordial Earth conditions. One such mechanism is based on the reaction of acetylene (C_2H_2) and carbon monoxide (CO) upon contact with nickel sulfide (NiS) in a hot aqueous environment.^{544,545,551} An experimental study showed that this reaction can lead to the formation of C3,5,7,9-monocarboxylic acids. Scheidler et al. ⁵⁵² suggested that such reactions could occur in Hadean volcanic-hydrothermal vents saturated with C_2H_2 and CO. These currents could bring gases into contact with NiS under conditions conducive to the formation of monocarboxylic acids. In this case, a large length of the alkyl radical could be achieved due to the high pressure.

Scheidler et al.⁵⁵² have made some interesting observations linking the proposed pathway with abiogenetic processes. These pathways are illustrated in Scheme 27. For example, for higher C7–9 monocarboxylic acids, the process of formation of the saturated form is not complete. The resulting products are a mixture of unsaturated monocarboxylic acids with one or more double bonds. Double bonds are predominantly in the form of trans isomers. In the case of several double bonds, a conjugated system is generally formed. The authors also noted that the modern organisms Bacteria and Eukarya contain unsaturated fatty acyl lipids within the membranes of saturated lipids. Moreover, the arrangement of double bonds in the middle of the hydrophobic tail is a common feature of natural unsaturated monocarboxylic acids and unsaturated lipids formed during the interaction of C_2H_2 and CO.

The evolution of a complex prebiotic mixture containing C_2H_2 , CO and NiS was also shown in recent works by Sobotta et al.⁵⁵³ and Diederich et al.⁵⁵⁴ A suspension of precipitated NiS was shown to convert acetylene and carbon monoxide into a set of C2–4 products. Moreover, the authors demonstrated the formation of a homologous series of thioacids upon repeated addition of C_2 to the resulting compounds via acetylene. Over 7 days, molecules with alkyl radical sizes ranging from 15 to 27 carbon atoms and a mass range of 230–500 m/z were formed. In this case, only up to four carbon tags from ¹³CO were present. Thus, we note the role of acetylene as the main driver of mass growth and as a building material for the synthesis of high-molecular weight compounds. The general reactions that are part of this synthesis are provided in Table 26.



Scheme 27. Evolution of monocarboxylic acid biosynthesis. Formation of carboxylic acids from acetylene on a nickel catalyst under prebiotic conditions (A); Conversion of acetylene to acetyl-thioester (B); Extant biosynthesis based on acetyl-thioester condensation (C).⁵⁵⁰

The described processes may be relevant for carbonaceous chondrites. Chondrites contain carbon grains stuck together, with water ice in the layers between them. Under the influence of solar radiation, the surface can be heated, and water ice can melt, initiating the synthesis of nitrogen bases, amino acids and fatty acids in the resulting solution.

We conclude that acetylene is a universal, stable, and flexible building block and could be the basis for the formation of a variety of precursors of biochemical molecules. Studying the chemistry of acetylene in space environments can provide important insights into markers indicating various stages of biochemical evolution, from nitrogenous bases to putative PAH templates for the first informational molecules. Moreover, the presence of metal and carbon catalysts can significantly diversify synthetic pathways and increase molecular complexity. Currently, we are still at the origin of knowledge about the role of catalytic processes in the formation of prebiotic molecular systems. Moreover, acetylene has high complementarity with respect to its ability to participate in catalytic cycles involving metals, sulfos, and carbon species, suggesting that this area of research is promising.

Table 26. The reactions of formation monocarboxylic acids from acetylene and other reactants.

Reactions	Kinetics	Objects & conditions
1. $C_2H_2 + CO + NiS + H_2O \rightarrow C3,5,7,9$ -Monocarboxylic Acids ^[552] 2. $C_2H_2 + CO + NiS + H_2O \rightarrow products + C_2H_2 \rightarrow C_5H_5OS$ (thio acids) ^[554]	n.d.	early Earth

9. Conclusions

Acetylene consists of two sp-hybridized carbon atoms connected by one σ and two π bonds. This structure predefines acetylene with a high exothermicity for addition reactions at the triple bond. Additionally, a pronounced shift in the electron density from a proton to a carbon atom results in the high acidity of the acetylene molecules. Due to its structure, the acetylene molecule is a reactive yet sufficiently robust and stable C₂ species. This position makes it an almost ideal building block for a myriad of organic compounds. Acetylene serves as a versatile chemical currency, with the potential to link molecules into complex structures possessing infinite carbon skeletons.

The significant role of acetylene in chemical evolution correlates with the widespread distribution of this molecule across various cosmic environments, spanning from hot circumstellar spaces to cold molecular clouds, planetary atmospheres and icy comets. Acetylene molecules are synthesized under very different conditions in these media. While acetylene is merely one component of the chemical cocktail of the universe, the array of processes in which it participates commands special attention. Despite not being the most prevalent carbon molecule in space, acetylene acts as a critical bridge to the chemistry of complex macromolecular compounds.

Most mechanisms of benzene synthesis across diverse space environments involve acetylene, thereby paving the way to a world of aromatic compounds. The primary mechanism for PAH synthesis, HACA, is rooted in the chemistry surrounding the acetylene molecule. Our understanding of the formation of high-molecular-weight aromatic compounds has undergone a seismic shift over the past decade. The emergence of new mechanisms based on barrierless reactions has radically altered our perception of space-carbon chemistry, shifting focus from high-temperature combustion-based reactions to low-temperature neutral-neutral reactions. This revolutionary concept suggests that PAHs can be formed not only at high temperatures but also at extremely low temperatures, leading to the creation of complex macromolecular compounds in cold the ISM. This includes not only "pure" carbon PAHs but also nitrogen-containing compounds. These mechanisms provide fresh perspectives on the chemistry of celestial bodies such as Titan. Further studies may aid in pinpointing the conditions conducive to determining the origin of PAHs based on their structural characteristics.

Despite advancements in the field, many factors remain elusive. Traditionally, studies of the underlying mechanisms involved the formation of bulk organic matter in rarefied interstellar or circumstellar media. However, critical chemical processes might transpire on the surface of solids at both low and high temperatures. Such processes could be catalytic in nature. For instance, catalytic pathways for the formation of aromatic compounds in space environments are likely underexplored. Catalytic processes, encompassing crucial events such as the formation of aromatic compounds, could occur on the surfaces of comets, asteroids, planets, and interstellar dust grains, as corroborated by recent findings. We can anticipate more exciting discoveries in this area in the coming years.

Our understanding of cosmic chemical processes will continue to broaden in the future. There is no doubt that missions such as the James Webb Space Telescope^{36, 555}, the forthcoming Dragonfly spacecraft^{556, 557} and the Jupiter Icy Moons Explorer⁵⁵⁸ will revolutionize our comprehension of the evolution of organic compounds in space. This approach will facilitate the precise quantification of how physical conditions influence organic transformations, thereby allowing us to track the evolution of acetylene and other organic molecules across various space environments.

ASSOCIATED CONTENT

Supporting Information

Table S1 contains data on the occurrence of acetylene in different space regions and objects; data on the main mechanisms of PAH formation and the role of acetylene in this pathway; Table S2 contains data on several important PAHs and their formation pathways based on crossed molecular beam experiments and pyrolytic reactors combined with quantum chemical calculations (PDF).

AUTHOR INFORMATION

Corresponding Author

* Valentine P. Ananikov – Zelinsky Institute of Organic Chemistry, Russian Academy of Sciences, Moscow 119991, Russia; orcid.org/0000-0002-6447-557X; Email: val@ioc.ac.ru

Authors

Evgeniy O. Pentsak - Zelinsky Institute of Organic Chemistry, Russian Academy of Sciences, Moscow 119991, Russia; orcid.org/0000-0001-9040-0013

Maria S. Murga - Institute of Astronomy, Russian Academy of Sciences, Pyatnitskaya str. 48, Moscow 119017, Russia; orcid.org/0000-0002-7910-6662

Author Contributions

The manuscript was written through contributions of all the authors. All the authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interest.

REFERENCES

¹ Primo, A.; Neatu, F.; Florea, M.; Parvulescu, V.; Garcia, H., Graphenes in the absence of metals as carbocatalysts for selective acetylene hydrogenation and alkene hydrogenation. *Nat. Commun.* **2014**, *5*, 5291. [DOI: 10.1038/ncomms6291]

² Davy E. Notice of a new gaseous bicarburet of hydrogen, *Report of the Sixth Meeting of the British Association for the Advancement of Science* **1837**, 6, 62–63. [https://play.google.com/store/books/details?id=nVFJAAAAcAAJ&rdid=book-nVFJAAAAcAAJ&rdot=1]

³ Berthelot M., Sur une nouvelle série de composés organiques, le quadricarbure d'hydrogène et ses dérivés (Note on a new series of organic compounds, tetra-carbon hydride and its derivatives), *Comptes rendus* **1860**, 3, 50, 805–808. [https://gallica.bnf.fr/ark:/12148/bpt6k3007r/f817.image]

⁴ Pässler, P; Hefner, W; Buckl, K; Meinass, H; Meiswinkel, A; Wernicke, H.-J.; Ebersberg, G; Müller, R; Bässler, J; Behringer, H; Mayer, D. Acetylene, in Ullmann's Encyclopedia of Industrial Chemistry, Wiley-VCH, Weinheim, **2011** [DOI: 10.1002/14356007.a01_097.pub4]

⁵ Schobert, H., Production of acetylene and acetylene-based chemicals from coal. *Chem. Rev.* **2014**, *114* (3), 1743-60. [DOI: 10.1021/cr400276u]

⁶ Trotuş, I.-T.; Zimmermann, T.; Schüth, F. Catalytic reactions of acetylene: A feedstock for the chemical industry revisited. *Chem. Rev.* **2014**, *114*, 1761–1782. [DOI: 10.1021/cr400357r]

⁷ Voronin, V. V.; Ledovskaya, M. S.; Bogachenkov, A. S.; Rodygin, K. S.; Ananikov, V. P., Acetylene in Organic Synthesis: Recent Progress and New Uses. *Molecules* **2018**, *23* (10). [DOI: 10.3390/molecules23102442]

⁸ Rodygin, K. S.; Ananikov, V. P., An efficient metal-free pathway to vinyl thioesters with calcium carbide as the acetylene source. *Green Chem.* **2016**, *18* (2), 482-486. [DOI: 10.1039/C5GC01552A]

⁹ Rodygin, K. S.; Vikenteva, Y. A.; Ananikov, V. P., Calcium-Based Sustainable Chemical Technologies for Total Carbon Recycling. *ChemSusChem* **2019**, *12* (8), 1483-1516. [DOI: 10.1002/cssc.201802412]

¹⁰ Tracy, J.; Zietz, O.; Olson, S.; Jiao, J., Plasma-Enhanced Chemical Vapor Deposition of Acetylene on Codeposited Bimetal Catalysts Increasing Graphene Sheet Continuity Under Low-Temperature Growth Conditions. *Nanoscale Res. Lett.* **2019**, *14* (1), 335. [DOI: 10.1186/s11671-019-3156-y]

¹¹ Osikoya, A. O.; Parlak, O.; Murugan, N. A.; Dikio, E. D.; Moloto, H.; Uzun, L.; Turner, A. P.; Tiwari, A., Acetylene-sourced CVD-synthesised catalytically active graphene for electrochemical biosensing. *Biosens. Bioelectron.* **2017**, *89* (Pt 1), 496-504. [DOI: 10.1016/j.bios.2016.03.063]

¹² Bulyarskiy, S. V.; Gorelik, V. S.; Ryazanov, R. M., Nitrogen Doping of Carbon Nanotubes Synthesized in Flowing Acetylene and Ammonia. *Inorganic Materials* **2020**, *56* (10), 1006-1010. [DOI: 10.1134/S0020168520100027]

¹³ Afrin, R.; Abbas, S. M.; Shah, N. A.; Mustafa, M. F.; Ali, Z.; Ahmad, N., Effect of Varying Inert Gas and Acetylene Concentration on the Synthesis of Carbon Nanotubes. *J. Nanosci. Nanotechnol.* **2016**, *16* (3), 2956-9. [DOI: 10.1166/jnn.2016.12469]

¹⁴ Magrez, A.; Seo, J. W.; Kuznetsov, V. L.; Forro, L., Evidence of an equimolar C2H2-CO2 reaction in the synthesis of carbon nanotubes. *Angew. Chem. Int. Ed. Engl.* **2007**, *46* (3), 441-4. [DOI: 10.1002/anie.200603764]

¹⁵ Zhong, G.; Hofmann, S.; Yan, F.; Telg, H.; Warner, J. H.; Eder, D.; Thomsen, C.; Milne, W. I.; Robertson, J., Acetylene: A Key Growth Precursor for Single-Walled Carbon Nanotube Forests. *J. Phys. Chem. C.* **2009**, *113* (40), 17321-17325. [DOI: 10.1021/jp905134b]

¹⁶ Wang, H.; Zhou, Y.; Qi, M.; Liu, C.; Xu, X.; Ren, Z., Direct Growth of Graphene on Fused Quartz by Atmospheric Pressure Chemical Vapor Deposition with Acetylene. *J. Phys. Chem. C.* **2018**, *123* (4), 2370-2377. [DOI: 10.1021/acs.jpcc.8b08806]

¹⁷ Ananikov, V. P.; Ledovskaya, M. S.; Voronin, V. V.; Rodygin, K. S., Acetylene and Ethylene: Universal C₂ Molecular Units in Cycloaddition Reactions. *Synthesis* **2021**. [DOI: 10.1055/a-1654-2318]

¹⁸ Zhao, L.; Kaiser, R. I.; Xu, B.; Ablikim, U.; Ahmed, M.; Joshi, D.; Veber, G.; Fischer, F. R.; Mebel, A. M., Pyrene synthesis in circumstellar envelopes and its role in the formation of 2D nanostructures. *Nat. Astron.* **2018**, *2* (5), 413-419. [DOI: 10.1038/s41550-018-0399-y]

¹⁹ Zhao, L.; Kaiser, R. I.; Lu, W.; Xu, B. et al. Molecular mass growth through ring expansion in polycyclic aromatic hydrocarbons via radical-radical reactions. *Nat. Commun.* **2019**, *10*, 3689 [DOI: 10.1038/s41467-019-11652-5]

²⁰ Contreras, C. S.; Salama, F. Laboratory Investigations of Polycyclic Aromatic Hydrocarbon Formation and Destruction in the Circumstellar Outflows of Carbon Stars. *Astrophys. J., Suppl. Ser.* **2013**, *208*, 6. [DOI: 10.1088/0067-0049/208/1/6]

²¹ Jones, B. M.; Zhang, F.; Kaiser, R. I.; Jamal, A. et al. Formation of benzene in the interstellar medium. *Proc. Natl. Acad. Sci. U. S. A.* **2011**, *108*, 452. [DOI: 10.1073/pnas.1012468108]

²² Chiar, J. E.; Tielens, A. G. G. M.; Adamson, A. J.; Ricca, A. The Structure, Origin, and Evolution of Interstellar Hydrocarbon Grains. *Astrophys. J.* **2013**, *770*, 78. [DOI: 10.1088/0004-637X/770/1/78]

²³ Chen, T.; Li, A. Synthesizing carbon nanotubes in space. *Astron. Astrophys.* **2019**, *631*, A54. [DOI: 10.1051/0004-6361/201935789]

²⁴ García-Hernández, D. A.; Villaver, E.; García-Lario, P.; Acosta-Pulido, J. A. et al. Infrared Study of Fullerene Planetary Nebulae. *Astrophys. J.* **2012**, *760*, 107. [DOI: 10.1088/0004-637X/760/2/107]

²⁵ García-Hernández, D. A.; Manchado, A.; García-Lario, P.; Stanghellini, L. et al. Formation of Fullerenes in H-Containing Planetary Nebulae. *Astrophys. J.* **2010**, *724*, L39. [DOI: 10.1088/2041-8205/724/1/L39]

²⁶ Tennyson, J.; Yurchenko, S. N. Laboratory spectra of hot molecules: Data needs for hot super-Earth exoplanets. *Mol. Astrophys.* **2017**, 8, 1. [DOI: 10.1016/j.molap.2017.05.002]

²⁷ Chubb, K. L.; Tennyson, J.; Yurchenko, S. N. ExoMol molecular line lists – XXXVII. Spectra of acetylene. *Mon. Not. R. Astron. Soc.* **2020**, *493*, 1531. [DOI: 10.1093/mnras/staa229]

²⁸ Abplanalp, M. J.; Frigge, R.; Kaiser, R. I. Low-temperature synthesis of polycyclic aromatic hydrocarbons in Titan's surface ices and on airless bodies. *Sci. Adv.* **2019**, *5*, eaaw5841. [DOI: 10.1126/sciadv.aaw5841]

²⁹ Jeilani, Y. A.; Fearce, C.; Nguyen, M. T. Acetylene as an essential building block for prebiotic formation of pyrimidine bases on Titan. *Phys. Chem. Chem. Phys.* **2015**, *17*, 24294. [DOI: 10.1039/c5cp03247d]

³⁰ Frankland, V. L.; James, A. D.; Sánchez, J. D. C.; Mangan, T. P. et al. Uptake of acetylene on cosmic dust and production of benzene in Titan's atmosphere. *Icarus* **2016**, *278*, 88. [DOI: 10.1016/j.icarus.2016.06.007]

³¹ Postberg, F.; Khawaja, N.; Abel, B.; Choblet, G. et al. Macromolecular organic compounds from the depths of Enceladus. *Nature* **2018**, *558*, 564. [DOI: 10.1038/s41586-018-0246-4]

³² Waite, J. H., Jr.; Combi, M. R.; Ip, W. H.; Cravens, T. E. et al. Cassini ion and neutral mass spectrometer: Enceladus plume composition and structure. *Science* **2006**, *311*, 1419. [DOI: 10.1126/science.1121290]

³³ Kaiser, R. I.; Hansen, N., An Aromatic Universe-A Physical Chemistry Perspective. J. Phys. Chem. A. **2021**, *125* (18), 3826-3840. [DOI: 10.1021/acs.jpca.1c00606]

³⁴ Mebel, A. M.; Landera, A.; Kaiser, R. I. Formation Mechanisms of Naphthalene and Indene: From the Interstellar Medium to Combustion Flames. *J. Phys. Chem. A* **2017**, *121*, 901. [DOI: 10.1021/acs.jpca.6b09735]

³⁵ Frenklach, M.; Mebel, A. M. On the mechanism of soot nucleation. *Phys. Chem. Chem. Phys.* **2020**, *22*, 5314. [DOI: 10.1039/D0CP00116C]

³⁶ Peeters, E.; Mackie, C.; Candian, A.; Tielens, A., A Spectroscopic View on Cosmic PAH Emission. *Acc. Chem. Res.* **2021**, *54* (8), 1921-1933. [DOI: 10.1021/acs.accounts.0c00747]

³⁷ Parker, D. S.; Kaiser, R. I.; Troy, T. P.; Ahmed, M. Hydrogen abstraction/acetylene addition revealed. *Angew. Chem. Int. Ed. Engl.* **2014**, 53, 7740. [DOI: 10.1002/anie.201404537]

³⁸ Dhanoa, H.; Rawlings, J. M. C. Is acetylene essential for carbon dust formation? *Mon. Not. R. Astron. Soc.* **2014**, *440*, 1786. [DOI: 10.1093/mnras/stu401]

³⁹ Yang, T.; Kaiser, R. I.; Troy, T. P.; Xu, B. et al. HACA's Heritage: A Free-Radical Pathway to Phenanthrene in Circumstellar Envelopes of Asymptotic Giant Branch Stars. *Angew. Chem. Int. Ed. Engl.* **2017**, 56, 4515. [DOI: 10.1002/anie.201701259]

⁴⁰ Ten Brink, F. Living on acetylene. A primordial energy source. *Met. Ions Life Sci.* **2014**, *14*, 15. [DOI: 10.1007/978-94-017-9269-1_2]

⁴¹ Keane, T. C., Mechanism for the Coupled Photochemistry of Ammonia and Acetylene: Implications for Giant Planets, Comets and Interstellar Organic Synthesis. *Origins Life Evol. Biospheres* **2017**, *47* (3), 223-248. [DOI: 10.1007/s11084-017-9545-2]

⁴² Oremland, R. S.; Voytek, M. A. Acetylene as fast food: implications for development of life on anoxic primordial Earth and in the outer solar system. *Astrobiology* **2008**, *8*, 45 [DOI: 10.1089/ast.2007.0183]

⁴³ Ehrenfreund, P.; Rasmussen, S.; Cleaves, J.; Chen, L. Experimentally tracing the key steps in the origin of life: The aromatic world. *Astrobiology* **2006**, *6*, 490. [DOI: 10.1089/ast.2006.6.490]

⁴⁴ Groen, J.; Deamer, D. W.; Kros, A.; Ehrenfreund, P. Polycyclic aromatic hydrocarbons as plausible prebiotic membrane components. *Origins Life Evol. Biospheres* **2012**, *42*, 295. [DOI: 10.1007/s11084-012-9292-3]

⁴⁵ d'Ischia, M.; Manini, P.; Moracci, M.; Saladino, R. et al. Astrochemistry and Astrobiology: Materials Sciencein Wonderland? *Int. J. Mol. Sci.* **2019**, 20. [DOI: 10.3390/ijms20174079]

⁴⁶ Encrenaz, T.; Combes, M.; Atreya, S. K.; Romani, P. N.; Fricke, K.; Moore, V.; Hunt, G.; Wagener, R.; Caldwell, J.; Owen, T.; Butterworth, P., A Study of the Upper-Atmosphere of Uranus Using the Iue. *Astron. Astrophys.* **1986**, *162* (1-2), 317-322. [http://adsabs.harvard.edu/full/1986A%26A...162...317E]

⁴⁷ Broadfoot, A. L.; Herbert, F.; Holberg, J. B.; Hunten, D. M.; Kumar, S.; Sandel, B. R.; Shemansky, D. E.; Smith, G. R.; Yelle, R. V.; Strobel, D. F.; Moos, H. W.; Donahue, T. M.; Atreya, S. K.; Bertaux, J. L.; Blamont, J. E.; McConnell, J. C.; Dessler, A. J.; Linick, S.; Springer, R., Ultraviolet spectrometer observations of uranus. *Science* **1986**, *233* (4759), 74-9. [DOI: 10.1126/science.233.4759.74]

⁴⁸ Jewitt, D. C.; Luu, J., Crystalline water ice on the Kuiper belt object (50000) Quaoar. *Nature* **2004**, *432* (7018), 731-733. [DOI: 10.1038/nature03111].

⁴⁹ Tielens, A. Interstellar PAHs and Dust. In *Planets, Stars and Stellar Systems*; Oswalt, T.D., Gilmore, G., Eds.; Springer, Dordrecht, 2013. [DOI: 10.1007/978-94-007-5612-0_10]

⁵⁰ Woods, P. M.; Millar, T. J.; Zijlstra, A. A.; Herbst, E. The Synthesis of Benzene in the Protoplanetary Nebula CRL 618. *Astrophys. J.* **2002**, *574*, L167. [DOI: 10.1086/342503]

⁵¹ Wiebel-Sooth, Barbara; Biermann, Peter L.; Meyer, Hinrich, Cosmic rays. VII. Individual element spectra: prediction and data. *Astron. Astrophys.* **1998**, *330*, 389-398. [DOI: 10.48550/arXiv.astro-ph/9709253]

⁵² Bringa, E. M.; Kucheyev, S. O.; Loeffler, M. J.; Baragiola, R. A.; Tielens, A. G. G. M.; Dai, Z. R.; Graham, G.; Bajt, S.; Bradley, J. P.; Dukes, C. A.; Felter, T. E.; Torres, D. F.; Breugel, W. v.,

Energetic Processing of Interstellar Silicate Grains by Cosmic Rays. *Astrophys. J.* **2007**, *662* (1), 372-378. [DOI: 10.1086/517865]

⁵³ Padovani, M.; Ivlev, A. V.; Galli, D.; Caselli, P., Cosmic-ray ionisation in circumstellar discs. *Astron. Astrophys.* **2018**, *614*. [DOI: 10.1051/0004-6361/201732202]

⁵⁴ Fonfria, J. P.; Cernicharo, J.; Richter, M. J.; Lacy, J. H., A Detailed Analysis of the Dust Formation Zone of IRC +10216 Derived from Mid-Infrared Bands of C_2H_2 and HCN. *Astrophys. J.* **2008**, *673* (1), 445-469. [DOI: 10.1086/523882]

⁵⁵ Kaufman, M. J.; Hollenbach, D. J.; Tielens, A. G. G. M., High-Temperature Molecular Cores near Massive Stars and Application to the Orion Hot Core. *Astrophys. J.* **1998**, *497* (1), 276-287. [DOI: 10.1086/305444]

⁵⁶ Habing, H. J. The interstellar radiation density between 912 A and 2400 A. *Bulletin of the Astronomical Institutes of the Netherlands* **1968**, *19*, 421. [https://ui.adsabs.harvard.edu/abs/1968BAN....19..421H/abstract]

⁵⁷ Karakas, A. I., Updated stellar yields from asymptotic giant branch models. *Mon. Not. R. Astron. Soc.* **2010**, *403* (3), 1413-1425. [DOI: 10.1111/j.1365-2966.2009.16198.x]

⁵⁸ Farmer, R.; Laplace, E.; de Mink, S. E.; Justham, S., The Cosmic Carbon Footprint of Massive Stars Stripped in Binary Systems. *Astrophys. J.* **2021**, *923* (2). [DOI: 10.3847/1538-4357/ac2f44]

⁵⁹ McCarthy, M. C.; Gottlieb, C. A.; Cernicharo, J., Building blocks of dust: A coordinated laboratory and astronomical study of the archtype AGB carbon star IRC +10216. *J. Mol. Spectrosc.* **2019**, *356*, 7-20. [DOI: 10.1016/j.jms.2018.11.018]

⁶⁰ Ehrenfreund, P.; Spaans, M.; Holm, N. G., The evolution of organic matter in space. *Philos. Trans. A Math. Phys. Eng. Sci.* **2011**, *369* (1936), 538-54. [DOI: 10.1098/rsta.2010.0231]

⁶¹ Tielens, A. G. G. M., The molecular universe. *Rev. Mod. Phys.* **2013**, *85* (3), 1021-1081. [DOI: 10.1103/RevModPhys.85.1021]

 62 Urru, A.; Kozin, I. N.; Mulas, G.; Braams, B. J.; Tennyson, J., Ro-vibrational spectra of C₂H₂based on variational nuclear motion calculations. *Mol. Phys.* **2010**, *108* (15), 1973-1990. [DOI: 10.1080/00268976.2010.499858]

⁶³ Chubb, K. L.; Joseph, M.; Franklin, J.; Choudhury, N.; Furtenbacher, T.; Császár, A. G.; Gaspard, G.; Oguoko, P.; Kelly, A.; Yurchenko, S. N.; Tennyson, J.; Sousa-Silva, C., MARVEL analysis of the measured high-resolution rovibrational spectra of C₂H₂. *J. Quant. Spectrosc. Radiat. Transfer* **2018**, *204*, 42-55. [DOI: 10.1016/j.jqsrt.2017.08.018]

⁶⁴ Melin, H.; Fletcher, L. N.; Irwin, P. G. J.; Edgington, S. G., Jupiter in the Ultraviolet: Acetylene and Ethane Abundances in the Stratosphere of Jupiter from Cassini Observations between 0.15 and 0.19 μm. *Astron. J.* **2020**, *159* (6). [DOI: 10.3847/1538-3881/ab91a6]

⁶⁵ Ridgway, S. T., Jupiter: Identification of ethane and acetylene. *Astrophys. J.* **1974**, *187*. [DOI: 10.1086/181388]

⁶⁶ Kitt Peak National Observatory <u>https://kpno.noirlab.edu/</u> (accessed 2024-03-04).

⁶⁷ NASA Infrared Telescope Facility (IRTF) <u>http://irtfweb.ifa.hawaii.edu/</u> (accessed 2024-03-04).

⁶⁸ Encrenaz, T., Remote sensing of the atmospheres of Jupiter, Saturn and Titan. *Rep. Prog. Phys.* **1990,** *53* (6), 793-836. [DOI: 10.1088/0034-4885/53/6/003]

⁶⁹ Tokunaga, A.T.; Beck, S. C.; Geballe, T.R.; Lacy, J. H. The detection of C2H2 on Saturn and Titan. *Bull. Am. Astron. Soc.* **1980**, *12*, 669 [http://adsabs.harvard.edu/pdf/1980BAAS...12..669T]

⁷⁰ Courtin, R.; Gautier, D.; Marten, A.; Bezard, B.; Hanel, R., The composition of Saturn's atmosphere at northern temperate latitudes from Voyager IRIS spectra - NH₃, PH₃, C₂H₂, C₂H₆, CH₃D, CH₄, and the Saturnian D/H isotopic ratio. *Astrophys. J.* **1984**, 287. [DOI: 10.1086/162748]

⁷¹ Voyager. Mission Overview <u>https://voyager.jpl.nasa.gov/mission/</u> (accessed 2024-03-04).

⁷² Hanel, R.; Conrath, B.; Flasar, F. M.; Kunde, V.; Maguire, W.; Pearl, J.; Pirraglia, J.; Samuelson, R.; Herath, L.; Allison, M.; Cruikshank, D.; Gautier, D.; Gierasch, P.; Horn, L.; Koppany, R.; Ponnamperuma, C., Infrared observations of the saturnian system from voyager 1. *Science* **1981**, *212* (4491), 192-200. [DOI: 10.1126/science.212.4491.192]

⁷³ Noll, K. S.; Knacke, R. F.; Tokunaga, A. T.; Lacy, J. H.; Beck, S.; Serabyn, E., The abundances of ethane and acetylene in the atmospheres of Jupiter and Saturn. *Icarus* **1986**, *65* (2-3), 257-263. [DOI: 10.1016/0019-1035(86)90138-7]

⁷⁴ Esposito, L. W.; Barth, C. A.; Colwell, J. E.; Lawrence, G. M.; McClintock, W. E.; Stewart, A. I. F.; Keller, H. U.; Korth, A.; Lauche, H.; Festou, M. C.; Lane, A. L.; Hansen, C. J.; Maki, J. N.; West, R. A.; Jahn, H.; Reulke, R.; Warlich, K.; Shemansky, D. E.; Yung, Y. L., The Cassini Ultraviolet Imaging Spectrograph Investigation. *Space Sci. Rev.* **2004**, *115* (*1-4*), 299-361. [DOI: 10.1007/s11214-004-1455-8]

⁷⁵ PIA13699: Acetylene around Jupiter's Poles <u>https://photojournal.jpl.nasa.gov/catalog/PIA13699</u> (accessed 2024-03-01).

⁷⁶ Jennings, D. E.; Sada, P. V.; Romani, P. N.; Bjoraker, G. L.; Simon-Miller, A. A., Brightening of Acetylene and Ethane in the South Pole of Saturn from Ground-Based and CIRS Observations. American Astronomical Society, DPS meeting #37, id.60.09; Bulletin of the American Astronomical Society, **2005**, *37*, 757. [https://ui.adsabs.harvard.edu/abs/2005DPS....37.6009J/abstract]

⁷⁷ Fletcher, L. N.; Cavalie, T.; Grassi, D.; Hueso, R.; Lara, L. M.; Kaspi, Y.; Galanti, E.; Greathouse, T. K.; Molyneux, P. M.; Galand, M.; Vallat, C.; Witasse, O.; Lorente, R.; Hartogh, P.; Poulet, F.; Langevin, Y.; Palumbo, P.; Gladstone, G. R.; Retherford, K. D.; Dougherty, M. K.; Wahlund, J. E.; Barabash, S.; Iess, L.; Bruzzone, L.; Hussmann, H.; Gurvits, L. I.; Santolik, O.; Kolmasova, I.; Fischer, G.; Muller-Wodarg, I.; Piccioni, G.; Fouchet, T.; Gerard, J. C.; Sanchez-Lavega, A.; Irwin, P. G. J.; Grodent, D.; Altieri, F.; Mura, A.; Drossart, P.; Kammer, J.; Giles, R.; Cazaux, S.; Jones, G.; Smirnova, M.; Lellouch, E.; Medvedev, A. S.; Moreno, R.; Rezac, L.; Coustenis, A.; Costa, M., Jupiter Science Enabled by ESA's Jupiter Icy Moons Explorer. *Space Sci. Rev.* **2023**, *219* (7), 53. [DOI: 10.1007/s11214-023-00996-6]

⁷⁸ Sinclair, J. A.; Moses, J. I.; Hue, V.; Greathouse, T. K.; Orton, G. S.; Fletcher, L. N.; Irwin, P. G. J., Jupiter's auroral-related stratospheric heating and chemistry III: Abundances of C₂H₄, CH₃C₂H, C₄H₂ and C₆H₆ from Voyager-IRIS and Cassini-CIRS. *Icarus* **2019**, *328*, 176-193. [DOI: 10.1016/j.icarus.2019.03.012]

⁷⁹ Summers, M. E.; Strobel, D. F., Photochemistry of the atmosphere of Uranus. *Astrophys. J.* **1989**, *346*. [DOI: 10.1086/168031]

⁸⁰ Orton, G. S.; Aitken, D. K.; Smith, C.; Roche, P. F.; Caldwell, J.; Snyder, R., The spectra of Uranus and Neptune at 8–14 and 17–23 μm. *Icarus* **1987**, *70* (1), 1-12. [DOI: 10.1016/0019-1035(87)90070-4]

⁸¹ Caldwell, J.; Wagener, R.; Fricke, K.-H., Observations of Neptune and Uranus below 2000 Å with the IUE. *Icarus* **1988**, *74* (1), 133-140. [DOI: 10.1016/0019-1035(88)90035-8]

⁸² Stone, E. C.; Miner, E. D., The voyager 2 encounter with the neptunian system. *Science* **1989**, 246 (4936), 1417-21. [DOI: 10.1126/science.246.4936.1417]

⁸³ Moses, J. I.; Poppe, A. R., Dust Ablation on the Giant Planets: Consequences for Stratospheric Photochemistry. *Icarus* **2017**, *297*, 33-58. [DOI: 10.1016/j.icarus.2017.06.002]

⁸⁴ Bar-nun, A., Acetylene formation on Jupiter: Photolysis or thunderstorms? *Icarus* **1979**, *38* (2), 180-191. [DOI: 10.1016/0019-1035(79)90177-5]

⁸⁵ Moses, J. I.; Cavalie, T.; Fletcher, L. N.; Roman, M. T., Atmospheric chemistry on Uranus and Neptune. *Philos. Trans. A. Math. Phys. Eng. Sci.* **2020**, *378* (2187), 20190477. [DOI: 10.1098/rsta.2019.0477]

⁸⁶ Moses, J., Photochemistry of Saturn's Atmosphere I. Hydrocarbon Chemistry and Comparisons with ISO Observations. *Icarus* **2000**, *143* (2), 244-298. [DOI: 10.1006/icar.1999.6270]

⁸⁷ Burgdorf, M.; Orton, G.; Vancleve, J.; Meadows, V.; Houck, J., Detection of new hydrocarbons in Uranus' atmosphere by infrared spectroscopy. *Icarus* **2006**, *184* (2), 634-637. [DOI: 10.1016/j.icarus.2006.06.006]

⁸⁸ Lo, J.-I.; Lin, M.-Y.; Peng, Y.-C.; Chou, S.-L.; Lu, H.-C.; Cheng, B.-M.; Ogilvie, J. F., Farultraviolet photolysis of solid methane. *Mon. Not. R. Astron. Soc.* **2015**, *451* (1), 159-166. [DOI: 10.1093/mnras/stv935]

⁸⁹ Romani, P. N.; Atreya, S. K., Methane photochemistry and haze production on Neptune. *Icarus* **1988**, *74* (3), 424-445. [DOI: 10.1016/0019-1035(88)90113-3]

⁹⁰. Atreya, S. K. and Romani P. N. Photochemistry and clouds of Jupiter, Saturn, and Uranus. In Recent Advances in Planetary Meterology (G. E. Hint, Ed.), pp. 17-68. Cambridge University Press, Cambridge. **1985**. [https://ui.adsabs.harvard.edu/abs/1985rapm.book...17A/abstract]

⁹¹ Gans, B.; Boye-Peronne, S.; Broquier, M.; Delsaut, M.; Douin, S.; Fellows, C. E.; Halvick, P.; Loison, J. C.; Lucchese, R. R.; Gauyacq, D., Photolysis of methane revisited at 121.6 nm and at 118.2 nm: quantum yields of the primary products, measured by mass spectrometry. *Phys. Chem. Chem. Phys.* **2011**, *13* (18), 8140-52. [DOI: 10.1039/C0CP02627A]

⁹² Blitz, M. A.; Seakins, P. W., Laboratory studies of photochemistry and gas phase radical reaction kinetics relevant to planetary atmospheres. *Chem. Soc. Rev.* **2012**, *41* (19), 6318-47. [DOI: 10.1039/C2CS35204D]

⁹³ Mordaunt, D. H.; Lambert, I. R.; Morley, G. P.; Ashfold, M. N. R.; Dixon, R. N.; Western, C. M.; Schnieder, L.; Welge, K. H., Primary product channels in the photodissociation of methane at 121.6 nm. *J. Chem. Phys.* **1993**, *98* (3), 2054-2065. [DOI: 10.1063/1.464237]

⁹⁴ Bishop, J.; Romani, P. N.; Atreya, S. K., Voyager 2 ultraviolet spectrometer solar occultations at neptune: photochemical modeling of the 125–;165 nm lightcurves. *Planet. Space Sci.* **1998**, *46* (1), 1-20. [DOI: 10.1016/S0032-0633(97)00126-8]

⁹⁵ Gibbard, S.; Levy, E. H.; Lunine, J. I., Generation of lightning in Jupiter's water cloud. *Nature* **1995**, *378* (6557), 592-5. [DOI: 10.1038/378592a0]

⁹⁶ Podolak, M.; Bar-Nun, A., Moist convection and the abundances of lightning-produced CO, C₂H₂, and HCN on Jupiter. *Icarus* **1988**, *75* (3), 566-570. [DOI: 10.1016/0019-1035(88)90165-0]

⁹⁷ Orville, R. E., Spectrum of the lightning stepped leader. *J. Geophys. Res.* **1968**, *73* (22), 6999-7008. [DOI: 10.1029/JB073i022p06999]

⁹⁸ Bar-Nun, A., Thunderstorms on Jupiter. *Icarus* **1975**, *24* (1), 86-94. [DOI: 10.1016/0019-1035(75)90162-1]

⁹⁹ Hurley, J.; Irwin, P. G. J.; Fletcher, L. N.; Moses, J. I.; Hesman, B.; Sinclair, J.; Merlet, C., Observations of upper tropospheric acetylene on Saturn: No apparent correlation with 2000km-sized thunderstorms. *Planet. Space Sci.* **2012**, *65* (1), 21-37. [DOI: 10.1016/j.pss.2011.12.026]

¹⁰⁰ Kinetic Database for Astrochemistry <u>https://kida.astrochem-tools.org</u>.

¹⁰¹ Stone, E. C.; Miner, E. D., Voyager 1 encounter with the saturnian system. *Science* **1981**, *212* (4491), 159-63. [DOI: 10.1126/science.212.4491.159]

 102 Kunde, V. G.; Aikin, A. C.; Hanel, R. A.; Jennings, D. E.; Maguire, W. C.; Samuelson, R. E., C₄H₂, HC₃N and C₂N₂ in Titan's atmosphere. *Nature* **1981**, *292* (5825), 686-688. [DOI: 10.1038/292686a0]

¹⁰³ Cassini-Huygens <u>https://solarsystem.nasa.gov/missions/cassini/overview/</u>.

¹⁰⁴ Coustenis, A.; Achterberg, R. K.; Conrath, B. J.; Jennings, D. E.; Marten, A.; Gautier, D.; Nixon, C. A.; Flasar, F. M.; Teanby, N. A.; Bézard, B.; Samuelson, R. E.; Carlson, R. C.; Lellouch, E.; Bjoraker, G. L.; Romani, P. N.; Taylor, F. W.; Irwin, P. G. J.; Fouchet, T.; Hubert, A.; Orton, G. S.; Kunde, V. G.; Vinatier, S.; Mondellini, J.; Abbas, M. M.; Courtin, R., The composition of Titan's stratosphere from Cassini/CIRS mid-infrared spectra. *Icarus* **2007**, *189* (1), 35-62. [DOI: 10.1016/j.icarus.2006.12.022]

¹⁰⁵ Clark, R. N.; Curchin, J. M.; Barnes, J. W.; Jaumann, R.; Soderblom, L.; Cruikshank, D. P.; Brown, R. H.; Rodriguez, S.; Lunine, J.; Stephan, K.; Hoefen, T. M.; Le Mouélic, S.; Sotin, C.; Baines, K. H.; Buratti, B. J.; Nicholson, P. D., Detection and mapping of hydrocarbon deposits on Titan. *J. Geophys. Res.* **2010**, *115* (E10). [DOI: 10.1029/2009JE003369]

¹⁰⁶ Hörst, S. M., Titan's atmosphere and climate. *J. Geophys. Res.: Planets* **2017**, *122* (3), 432-482. [DOI: 10.1002/2016JE005240]

¹⁰⁷ Cui, J.; Yelle, R. V.; Vuitton, V.; Waite, J. H.; Kasprzak, W. T.; Gell, D. A.; Niemann, H. B.; Müller-Wodarg, I. C. F.; Borggren, N.; Fletcher, G. G.; Patrick, E. L.; Raaen, E.; Magee, B. A., Analysis of Titan's neutral upper atmosphere from Cassini Ion Neutral Mass Spectrometer measurements. *Icarus* **2009**, *200* (2), 581-615. [DOI: 10.1016/j.icarus.2008.12.005]

¹⁰⁸ Fleury, B.; Gudipati, M. S.; Couturier-Tamburelli, I.; Carrasco, N., Photoreactivity of condensed acetylene on Titan aerosols analogues. *Icarus* **2019**, *321*, 358-366. [DOI: 10.1016/j.icarus.2018.11.030]

¹⁰⁹ Lavvas, P. P.; Coustenis, A.; Vardavas, I. M., Coupling photochemistry with haze formation in Titan's atmosphere, Part I: Model description. *Planet. Space Sci.* **2008**, *56* (1), 27-66. [DOI: 10.1016/j.pss.2007.05.026]

¹¹⁰ Singh, S.; McCord, T. B.; Combe, J. P.; Rodriguez, S.; Cornet, T.; Mouélic, S. L.; Clark, R. N.; Maltagliati, L.; Chevrier, V. F., Acetylene on Titan's Surface. *Astrophys. J.* **2016**, *828* (1). [DOI: 10.3847/0004-637X/828/1/55]

¹¹¹ Niemann, H. B.; Atreya, S. K.; Demick, J. E.; Gautier, D.; Haberman, J. A.; Harpold, D. N.; Kasprzak, W. T.; Lunine, J. I.; Owen, T. C.; Raulin, F., Composition of Titan's lower atmosphere and simple surface volatiles as measured by the Cassini-Huygens probe gas chromatograph mass spectrometer experiment. *J. Geophys. Res.* **2010**, *115* (E12). [DOI: 10.1029/2010JE003659]

¹¹² New Horizons <u>https://science.nasa.gov/mission/new-horizons/</u>.

¹¹³ Stern, S. A.; Bagenal, F.; Ennico, K.; Gladstone, G. R.; Grundy, W. M.; McKinnon, W. B.; Moore, J. M.; Olkin, C. B.; Spencer, J. R.; Weaver, H. A.; Young, L. A.; Andert, T.; Andrews, J.; Banks, M.; Bauer, B.; Bauman, J.; Barnouin, O. S.; Bedini, P.; Beisser, K.; Beyer, R. A.; Bhaskaran, S.; Binzel, R. P.; Birath, E.; Bird, M.; Bogan, D. J.; Bowman, A.; Bray, V. J.; Brozovic, M.; Bryan, C.; Buckley, M. R.; Buie, M. W.; Buratti, B. J.; Bushman, S. S.; Calloway, A.; Carcich, B.; Cheng, A. F.; Conard, S.; Conrad, C. A.; Cook, J. C.; Cruikshank, D. P.; Custodio, O. S.; Dalle Ore, C. M.; Deboy, C.; Dischner, Z. J.; Dumont, P.; Earle, A. M.; Elliott, H. A.; Ercol, J.; Ernst, C. M.; Finley, T.; Flanigan, S. H.; Fountain, G.; Freeze, M. J.; Greathouse, T.; Green, J. L.; Guo, Y.; Hahn, M.; Hamilton, D. P.; Hamilton, S. A.; Hanley, J.; Harch, A.; Hart, H. M.; Hersman, C. B.; Hill, A.; Hill, M. E.; Hinson, D. P.; Holdridge, M. E.; Horanyi, M.; Howard, A. D.; Howett, C. J.; Jackman, C.; Jacobson, R. A.; Jennings, D. E.; Kammer, J. A.; Kang, H. K.; Kaufmann, D. E.; Kollmann, P.; Krimigis, S. M.; Kusnierkiewicz, D.; Lauer, T. R.; Lee, J. E.; Lindstrom, K. L.; Linscott, I. R.; Lisse, C. M.; Lunsford, A. W.; Mallder, V. A.; Martin, N.; McComas, D. J.; McNutt, R. L., Jr.; Mehoke, D.; Mehoke, T.; Melin, E. D.; Mutchler, M.; Nelson, D.; Nimmo, F.; Nunez, J. I.; Ocampo, A.; Owen, W. M.; Paetzold, M.; Page, B.; Parker, A. H.; Parker, J. W.; Pelletier, F.; Peterson, J.; Pinkine, N.; Piquette, M.; Porter, S. B.; Protopapa, S.; Redfern, J.; Reitsema, H. J.; Reuter, D. C.; Roberts, J. H.; Robbins, S. J.; Rogers, G.; Rose, D.; Runyon, K.; Retherford, K. D.; Ryschkewitsch, M. G.; Schenk, P.; Schindhelm, E.; Sepan, B.; Showalter, M. R.; Singer, K. N.; Soluri, M.; Taylor, A.; Taylor, H.; Throop, H. B.; Tsang, C. C.; Tyler, G. L.; Umurhan, O. M.; Verbiscer, A. J.; Versteeg, M. H.; Vincent, M.; Webbert, R.; Weidner, S.; Weigle, G. E., 2nd; White, O. L.; Whittenburg, K.; Williams, B. G.; Williams, K.; Williams, S.; Woods, W. W.; Zangari, A. M.; Zirnstein, E., The Pluto system: Initial results from its exploration by New Horizons. *Science* **2015**, 350 (6258), aad1815. [DOI: 10.1126/science.aad1815]

¹¹⁴ Gladstone, G. R.; Stern, S. A.; Ennico, K.; Olkin, C. B.; Weaver, H. A.; Young, L. A.; Summers, M. E.; Strobel, D. F.; Hinson, D. P.; Kammer, J. A.; Parker, A. H.; Steffl, A. J.; Linscott, I. R.; Parker, J. W.; Cheng, A. F.; Slater, D. C.; Versteeg, M. H.; Greathouse, T. K.; Retherford, K. D.; Throop, H.; Cunningham, N. J.; Woods, W. W.; Singer, K. N.; Tsang, C. C.; Schindhelm, R.; Lisse, C. M.; Wong, M. L.; Yung, Y. L.; Zhu, X.; Curdt, W.; Lavvas, P.; Young, E. F.; Tyler, G. L.; New Horizons Science, T., The atmosphere of Pluto as observed by New Horizons. *Science* **2016**, 351 (6279), aad8866. [DOI: 10.1126/science.aad8866]

¹¹⁵ Wong, M. L.; Fan, S.; Gao, P.; Liang, M.-C.; Shia, R.-L.; Yung, Y. L.; Kammer, J. A.; Summers, M. E.; Gladstone, G. R.; Young, L. A.; Olkin, C. B.; Ennico, K.; Weaver, H. A.; Stern, S. A., The photochemistry of Pluto's atmosphere as illuminated by New Horizons. *Icarus* **2017**, 287, 110-115. [DOI: 10.1016/j.icarus.2016.09.028]

¹¹⁶ Yabuta, H.; Cody, G. D.; Engrand, C.; Kebukawa, Y.; De Gregorio, B.; Bonal, L.; Remusat, L.; Stroud, R.; Quirico, E.; Nittler, L.; Hashiguchi, M.; Komatsu, M.; Okumura, T.; Mathurin, J.; Dartois, E.; Duprat, J.; Takahashi, Y.; Takeichi, Y.; Kilcoyne, D.; Yamashita, S.; Dazzi, A.; Deniset-Besseau, A.; Sandford, S.; Martins, Z.; Tamenori, Y.; Ohigashi, T.; Suga, H.; Wakabayashi, D.; Verdier-Paoletti, M.; Mostefaoui, S.; Montagnac, G.; Barosch, J.; Kamide, K.; Shigenaka, M.; Bejach, L.; Matsumoto, M.; Enokido, Y.; Noguchi, T.; Yurimoto, H.; Nakamura, T.; Okazaki, R.; Naraoka, H.; Sakamoto, K.; Connolly, H. C., Jr.; Lauretta, D. S.; Abe, M.; Okada, T.; Yada, T.; Nishimura, M.; Yogata, K.; Nakato, A.; Yoshitake, M.; Iwamae, A.; Furuya, S.; Hatakeda, K.; Miyazaki, A.; Soejima, H.; Hitomi, Y.; Kumagai, K.; Usui, T.; Hayashi, T.; Yamamoto, D.; Fukai, R.; Sugita, S.; Kitazato, K.; Hirata, N.; Honda, R.; Morota, T.; Tatsumi, E.; Sakatani, N.; Namiki, N.; Matsumoto, K.; Noguchi, R.; Wada, K.; Senshu, H.; Ogawa, K.; Yokota, Y.; Ishihara, Y.; Shimaki, Y.; Yamada, M.; Honda, C.; Michikami, T.; Matsuoka, M.; Hirata, N.; Arakawa, M.; Okamoto, C.; Ishiguro, M.; Jaumann, R.; Bibring, J. P.; Grott, M.; Schroder, S.; Otto, K.; Pilorget, C.; Schmitz, N.; Biele, J.; Ho, T. M.; Moussi-Soffys, A.; Miura, A.; Noda, H.; Yamada, T.; Yoshihara, K.; Kawahara, K.; Ikeda, H.; Yamamoto, Y.; Shirai, K.; Kikuchi, S.; Ogawa, N.; Takeuchi, H.; Ono, G.; Mimasu, Y.; Yoshikawa, K.; Takei, Y.; Fujii, A.; Iijima, Y. I.; Nakazawa, S.; Hosoda, S.; Iwata, T.; Hayakawa, M.; Sawada, H.; Yano, H.; Tsukizaki, R.; Ozaki, M.; Terui, F.; Tanaka, S.; Fujimoto, M.; Yoshikawa, M.; Saiki, T.; Tachibana, S.; Watanabe, S. I.; Tsuda, Y., Macromolecular organic matter in samples of the asteroid (162173) Ryugu. Science **2023**, *379* (6634), eabn9057. [DOI: 10.1126/science.abn9057]

¹¹⁷ Potiszil, C.; Tanaka, R.; Kobayashi, K.; Kunihiro, T.; Nakamura, E., The Albedo of Ryugu: Evidence for a High Organic Abundance, as Inferred from the Hayabusa2 Touchdown Maneuver. *Astrobiology* **2020**, *20* (7), 916-921. [DOI 10.1089/ast.2019.2198]

¹¹⁸ Naraoka, H.; Takano, Y.; Dworkin, J. P.; Oba, Y.; Hamase, K.; Furusho, A.; Ogawa, N. O.; Hashiguchi, M.; Fukushima, K.; Aoki, D.; Schmitt-Kopplin, P.; Aponte, J. C.; Parker, E. T.; Glavin, D. P.; McLain, H. L.; Elsila, J. E.; Graham, H. V.; Eiler, J. M.; Orthous-Daunay, F. R.; Wolters, C.; Isa, J.; Vuitton, V.; Thissen, R.; Sakai, S.; Yoshimura, T.; Koga, T.; Ohkouchi, N.; Chikaraishi, Y.; Sugahara, H.; Mita, H.; Furukawa, Y.; Hertkorn, N.; Ruf, A.; Yurimoto, H.; Nakamura, T.; Noguchi, T.; Okazaki, R.; Yabuta, H.; Sakamoto, K.; Tachibana, S.; Connolly, H. C., Jr.; Lauretta, D. S.; Abe, M.; Yada, T.; Nishimura, M.; Yogata, K.; Nakato, A.; Yoshitake, M.; Suzuki, A.; Miyazaki, A.; Furuya, S.; Hatakeda, K.; Soejima, H.; Hitomi, Y.; Kumagai, K.; Usui, T.; Hayashi, T.; Yamamoto, D.; Fukai, R.; Kitazato, K.; Sugita, S.; Namiki, N.; Arakawa, M.; Ikeda, H.; Ishiguro, M.; Hirata, N.; Wada, K.; Ishihara, Y.; Noguchi, R.; Morota, T.; Sakatani, N.; Matsumoto, K.; Senshu, H.; Honda, R.; Tatsumi, E.; Yokota, Y.; Honda, C.; Michikami, T.; Matsuoka, M.; Miura, A.; Noda, H.; Yamada, T.; Yoshihara, K.; Kawahara, K.; Ozaki, M.; Iijima, Y. I.; Yano, H.; Hayakawa, M.; Iwata, T.; Tsukizaki, R.; Sawada, H.; Hosoda, S.; Ogawa, K.; Okamoto, C.; Hirata, N.; Shirai, K.; Shimaki, Y.; Yamada, M.; Okada, T.; Yamamoto, Y.; Takeuchi, H.; Fujii, A.; Takei, Y.; Yoshikawa, K.; Mimasu, Y.; Ono, G.; Ogawa, N.; Kikuchi, S.; Nakazawa, S.; Terui, F.; Tanaka, S.; Saiki, T.; Yoshikawa, M.; Watanabe, S. I.; Tsuda, Y., Soluble organic molecules in samples of the carbonaceous asteroid (162173) Ryugu. Science 2023, 379 (6634), eabn9033. [DOI: 10.1126/science.abn9033]

¹¹⁹ Okazaki, R.; Miura, Y. N.; Takano, Y.; Sawada, H.; Sakamoto, K.; Yada, T.; Yamada, K.; Kawagucci, S.; Matsui, Y.; Hashizume, K.; Ishida, A.; Broadley, M. W.; Marty, B.; Byrne, D.; Furi, E.; Meshik, A.; Pravdivtseva, O.; Busemann, H.; Riebe, M. E. I.; Gilmour, J.; Park, J.; Bajo, K. I.; Righter, K.; Sakai, S.; Sekimoto, S.; Kitajima, F.; Crowther, S. A.; Iwata, N.; Shirai, N.; Ebihara, M.; Yokochi, R.; Nishiizumi, K.; Nagao, K.; Lee, J. I.; Clay, P.; Kano, A.; Caffee, M. W.; Uemura, R.; Inagaki, M.; Krietsch, D.; Maden, C.; Yamamoto, M.; Fawcett, L.; Lawton, T.; Nakamura, T.; Naraoka, H.; Noguchi, T.; Yabuta, H.; Yurimoto, H.; Tsuda, Y.; Watanabe, S. I.; Abe, M.; Arakawa, M.; Fujii, A.; Hayakawa, M.; Hirata, N.; Hirata, N.; Honda, R.; Honda, C.; Hosoda, S.; Iijima, Y. I.; Ikeda, H.; Ishiguro, M.; Ishihara, Y.; Iwata, T.; Kawahara, K.; Kikuchi, S.; Kitazato, K.; Matsumoto, K.; Matsuoka, M.; Michikami, T.; Mimasu, Y.; Miura, A.; Morota, T.; Nakazawa, S.; Namiki, N.; Noda, H.; Noguchi, R.; Ogawa, N.; Ogawa, K.; Okada, T.; Okamoto, C.; Ono, G.; Ozaki, M.; Saiki, T.; Sakatani, N.; Senshu, H.; Shimaki, Y.; Shirai, K.; Sugita, S.; Takei, Y.; Takeuchi, H.; Tanaka, S.; Tatsumi, E.; Terui, F.; Tsukizaki, R.; Wada, K.; Yamada, M.; Yamada, T.; Yamamoto, Y.; Yano, H.; Yokota, Y.; Yoshihara, K.; Yoshikawa, M.; Yoshikawa, K.; Furuya, S.; Hatakeda, K.; Hayashi, T.; Hitomi, Y.; Kumagai, K.; Miyazaki, A.; Nakato, A.; Nishimura, M.; Soejima, H.; Iwamae, A.; Yamamoto, D.; Yogata, K.; Yoshitake, M.; Fukai, R.; Usui, T.; Ireland, T.; Connolly, H. C., Jr.; Lauretta, D. S.; Tachibana, S., First asteroid gas sample delivered by the Hayabusa2 mission: A treasure box from Ryugu. Sci Adv 2022, 8 (46), eabo7239. [DOI: 10.1126/sciadv.abo7239]

¹²⁰ Brooke, T. Y.; Tokunaga, A. T.; Weaver, H. A.; Crovisier, J.; Bockelee-Morvan, D.; Crisp, D., Detection of acetylene in the infrared spectrum of comet Hyakutake. *Nature* **1996**, *383* (6601), 606-8. [DOI: 10.1038/383606a0]

¹²¹ Mumma, M. J.; DiSanti, M. A.; Dello Russo, N.; Magee-Sauer, K.; Gibb, E.; Novak, R., Remote infrared observations of parent volatiles in comets: A window on the early solar system. *Adv. Space Res.* **2003**, *31* (12), 2563-2575. [DOI: 10.1016/S0273-1177(03)00578-7]

¹²² W.M. Keck Observatory <u>https://www.keckobservatory.org/</u>.

¹²³ Eberhardt, P. Comet Halley's Gas Composition and Extended Sources: Results from the Neutral Mass Spectrometer on Giotto. *Space Sci. Rev.* **1999,** *90* (1/2), 45-52. [DOI: 10.1023/A:1005221309219]

¹²⁴ Le Roy, L.; Altwegg, K.; Balsiger, H.; Berthelier, J.-J.; Bieler, A.; Briois, C.; Calmonte, U.; Combi, M. R.; De Keyser, J.; Dhooghe, F.; Fiethe, B.; Fuselier, S. A.; Gasc, S.; Gombosi, T. I.; Hässig, M.; Jäckel, A.; Rubin, M.; Tzou, C.-Y., Inventory of the volatiles on comet

67P/Churyumov-Gerasimenko from Rosetta/ROSINA. *Astron. Astrophys.* **2015**, *583*. [DOI: 10.1051/0004-6361/201526450]

¹²⁵ Mumma, M. J.; DiSanti, M. A.; Dello Russo, N.; Magee-Sauer, K.; Rettig, T. W., Detection of CO and Ethane in Comet 21P/Giacobini-Zinner: Evidence for Variable Chemistry in the Outer Solar Nebula. *Astrophys. J.* **2000**, *531* (2), L155-L159. [DOI: 10.1086/312530]

¹²⁶ Rimmer, P. B.; Ferus, M.; Waldmann, I. P.; Knížek, A.; Kalvaitis, D.; Ivanek, O.; Kubelík, P.; Yurchenko, S. N.; Burian, T.; Dostál, J.; Juha, L.; Dudžák, R.; Krůs, M.; Tennyson, J.; Civiš, S.; Archibald, A. T.; Granville-Willett, A., Identifiable Acetylene Features Predicted for Young Earth-like Exoplanets with Reducing Atmospheres Undergoing Heavy Bombardment. *Astrophys. J.* **2019**, 888 (1). [DOI: 10.3847/1538-4357/ab55e8]

¹²⁷ Tsiaras, A.; Rocchetto, M.; Waldmann, I. P.; Venot, O. et al. Detection of an Atmosphere around the Super-Earth 55 Cancri E. *Astrophys. J.* **2016**, *820*, 99. [DOI: 10.3847/0004-637X/820/2/99]

¹²⁸ Chubb, K. L.; Yachmenev, A.; Tennyson, J.; Yurchenko, S. N., Treating linear molecule HCCH in calculations of rotation-vibration spectra. *J. Chem. Phys.* **2018**, *149* (1), 014101. [DOI: 10.1063/1.5031844]

¹²⁹ Oppenheimer, B. R.; Baranec, C.; Beichman, C.; Brenner, D. at al. Reconnaissance of the Hr 8799 Exosolar System. I. Near-Infrared Spectroscopy. *Astrophys. J.* **2013**, *768*, 24. [DOI: 10.1088/0004-637X/768/1/24]

¹³⁰ de Kok, R. J.; Birkby, J.; Brogi, M.; Schwarz, H. et al. Identifying new opportunities for exoplanet characterisation at high spectral resolution. *Astron. Astrophys.* **2014**, *561*, A150. [DOI: 10.1051/0004-6361/201322947]

¹³¹ Shabram, M.; Fortney, J. J.; Greene, T. P.; Freedman, R. S. Transmission spectra of transiting planet atmospheres model validation and simulations of the hot Neptune GJ 436b for the James Webb Space Telescope. *Astrophys. J.* **2011**, *727*, 65. [DOI: 10.1088/0004-637X/727/2/65]

¹³² Hu, R.; Seager, S. Photochemistry in Terrestrial Exoplanet Atmospheres. Iii. Photochemistry and Thermochemistry in Thick Atmospheres on Super Earths and Mini Neptunes. *Astrophys. J.* **2014**, 784, 63. [DOI: 10.1088/0004-637X/784/1/63]

¹³³ Ridgway, S. T.; Hall, D. N. B.; Kleinmann, S. G.; Weinberger, D. A.; Wojslaw, R. S., Circumstellar acetylene in the infrared spectrum of IRC +10216. *Nature* **1976**, *264* (5584), 345-346. [DOI: 10.1038/264345a0]

¹³⁴ Lacy, J. H.; Evans, N. J., II; Achtermann, J. M.; Bruce, D. E.; Arens, J. F.; Carr, J. S., Discovery of interstellar acetylene. *Astrophys. J.* **1989**, *342*. [DOI: 10.1086/185480]

¹³⁵ Crosas, M.; Menten, K. M.; Young, K.; Phillips, T. G. Radiative Transfer in a Turbulent Expanding Molecular Envelope: Application to Mira. *Astrophys. Space Sci.* **1997**, *251* (1/2), 189-192. [DOI: 10.1023/A:1000740823858]

¹³⁶ The Infrared Space Observatory (ISO) <u>https://www.cosmos.esa.int/web/iso</u> (accessed 2024-03-04).

¹³⁷ Spitzer Space Telescope <u>https://www.spitzer.caltech.edu/</u> (accessed 2024-03-04).

¹³⁸ Jorgensen, U. G.; Hron, J.; Loidl, R., ISO-SWS spectra of the carbon stars TX Psc, V460 Cyg, and TT Cyg. *Astron. Astrophys.* **2000,** 356 (1), 253-266. [https://ui.adsabs.harvard.edu/abs/2000A&A...356..253J/abstract]

¹³⁹ Lagadec, E.; Sloan, G. C.; Zijlstra, A. A.; Mauron, N.; Houck, J. R., Dust and gas in carbon stars towards the Galactic halo. *Mon. Not. R. Astron. Soc.* **2012**, *427* (3), 2588-2596. [DOI: 10.1111/j.1365-2966.2012.21874.x]

¹⁴⁰ Matsuura, M.; Wood, P. R.; Sloan, G. C.; Zijlstra, A. A.; Van Loon, J. T.; Groenewegen, M. A. T.; Blommaert, J. A. D. L.; Cioni, M. R. L.; Feast, M. W.; Habing, H. J.; Hony, S.; Lagadec, E.; Loup, C.; Menzies, J. W.; Waters, L. B. F. M.; Whitelock, P. A., Spitzer observations of acetylene bands in carbon-rich asymptotic giant branch stars in the Large Magellanic Cloud. *Mon. Not. R. Astron. Soc.* **2006**, *371* (1), 415-420. [DOI: 10.1111/j.1365-2966.2006.10664.x]

¹⁴¹ Sloan, G. C.; Kraemer, K. E.; Matsuura, M.; Wood, P. R.; Price, S. D.; Egan, M. P., Mid-Infrared Spectroscopy of Carbon Stars in the Small Magellanic Cloud. *Astrophys. J.* **2006**, *645* (2), 1118-1130. [DOI: 10.1086/504516]

¹⁴² Sloan, G. C.; Matsuura, M.; Lagadec, E.; van Loon, J. T.; Kraemer, K. E.; McDonald, I.; Groenewegen, M. A. T.; Wood, P. R.; Bernard-Salas, J.; Zijlstra, A. A., Carbon-Rich Dust Production in Metal-Poor Galaxies in the Local Group. *Astrophys. J.* **2012**, *752* (2). [DOI: 10.1088/0004-637X/752/2/140]

¹⁴³ Cernicharo, J.; Heras, A. M.; Tielens, A. G. G. M.; Pardo, J. R.; Herpin, F.; Guélin, M.; Waters, L. B. F. M., Infrared Space Observatory's Discovery of C_4H_2 , C_6H_2 , and Benzene in CRL 618. *Astrophys. J.* **2001**, *546* (2), L123-L126. [DOI: 10.1086/318871]

¹⁴⁴ Cherchneff, I., A chemical study of the inner winds of asymptotic giant branch stars. *Astron. Astrophys.* **2006**, *456* (3), 1001-1012. [DOI: 10.1051/0004-6361:20064827]

¹⁴⁵ Agúndez, M.; Cernicharo, J.; Goicoechea, J. R., Formation of simple organic molecules in inner T Tauri disks. *Astron. Astrophys.* **2008**, *483* (3), 831-837. [DOI: 10.1051/0004-6361:20077927]

¹⁴⁶ The UMIST Database for Astrochemistry <u>http://udfa.ajmarkwick.net/index.php</u> (accessed 2024-03-04).

¹⁴⁷ Ruffle, D. P.; Herbst, E., New models of interstellar gas-grain chemistry - I. Surface diffusion rates. *Mon. Not. R. Astron. Soc.* **2002**, *319* (3), 837-850. [DOI: 10.1046/j.1365-8711.2000.03911.x]

¹⁴⁸ Hasegawa, T. I.; Herbst, E., New gas–grain chemical models of quiescent dense interstellar clouds: the effects of H tunnelling reactions and cosmic ray induced desorption. *Mon. Not. R. Astron. Soc.* **1993**, *261* (1), 83-102. [DOI: 10.1093/mnras/261.1.83]

¹⁴⁹ Du, F.; Parise, B.; Bergman, P., Production of interstellar hydrogen peroxide (H₂O₂) on the surface of dust grains. *Astron. Astrophys.* **2012**, *538*. [DOI: 10.1051/0004-6361/201118013]

¹⁵⁰ Sonnentrucker, P.; González-Alfonso, E.; Neufeld, D. A., Detection of Acetylene toward Cepheus A East with Spitzer. *Astrophys. J.* **2007**, *671* (1), L37-L40. [DOI: 10.1086/524705]

¹⁵¹ Lahuis, F.; van Dishoeck, E. F., ISO-SWS spectroscopy of gas-phase C_2H_2 and HCN toward massive young stellar objects. *Astron. Astrophys.* **2000**, *355*, 699-712. [https://adsabs.harvard.edu/full/2000A%26A...355..699L]

¹⁵² Lee, H. H.; Bettens, R. P. A.; Herbst, E., Fractional abundances of molecules in dense interstellar clouds: A compendium of recent model results. *Astron. Astrophys., Suppl. Ser.* **2017**, *119* (1), 111-114. [DOI: 10.1051/aas:1996232]

¹⁵³ Sakai, N.; Yamamoto, S., Warm carbon-chain chemistry. *Chem. Rev.* **2013**, *113* (12), 8981-9015. [DOI: 10.1021/cr4001308]

¹⁵⁴ Moore, M. H.; Hudson, R. L., Infrared Study of Ion-Irradiated Water-Ice Mixtures with Hydrocarbons Relevant to Comets. *Icarus* **1998**, *135* (2), 518-527. [DOI: 10.1006/icar.1998.5996]

¹⁵⁵ Bergin, E. A.; Du, F.; Cleeves, L. I.; Blake, G. A.; Schwarz, K.; Visser, R.; Zhang, K., Hydrocarbon Emission Rings in Protoplanetary Disks Induced by Dust Evolution. *Astrophys. J.* **2016**, *831* (1). [DOI: 10.3847/0004-637X/831/1/101]

¹⁵⁶ Le Gal, R.; Herbst, E.; Dufour, G.; Gratier, P.; Ruaud, M.; Vidal, T. H. G.; Wakelam, V., A new study of the chemical structure of the Horsehead nebula: the influence of grain-surface chemistry. *Astron. Astrophys.* **2017**, 605. [DOI: 10.1051/0004-6361/201730980]

¹⁵⁷ Tsuge, M.; Molpeceres, G.; Aikawa, Y.; Watanabe, N., Surface diffusion of carbon atoms as a driver of interstellar organic chemistry. *Nat. Astron.* **2023**, *7* (11), 1351-1358. [DOI: 10.1038/s41550-023-02071-0]

¹⁵⁸ Kobayashi, H.; Hidaka, H.; Lamberts, T.; Hama, T.; Kawakita, H.; Kästner, J.; Watanabe, N., Hydrogenation and Deuteration of C_2H_2 and C_2H_4 on Cold Grains: A Clue to the Formation Mechanism of C_2H_6 with Astronomical Interest. *Astrophys. J.* **2017**, *837* (2). [DOI: 10.3847/1538-4357/837/2/155]

¹⁵⁹ Bennett, C. J.; Jamieson, C. S.; Osamura, Y.; Kaiser, R. I., Laboratory Studies on the Irradiation of Methane in Interstellar, Cometary, and Solar System Ices. *Astrophys. J.* **2006**, *653* (1), 792-811. [DOI: 10.1086/508561]

¹⁶⁰ Abplanalp, M. J.; Gobi, S.; Kaiser, R. I., On the formation and the isomer specific detection of methylacetylene (CH₃CCH), propene (CH₃CHCH₂), cyclopropane (c-C₃H₆), vinylacetylene (CH₂CHCCH), and 1,3-butadiene (CH₂CHCHCH₂) from interstellar methane ice analogues. *Phys. Chem. Chem. Phys.* **2019**, *21* (10), 5378-5393. [DOI: 10.1039/C8CP03921F]

¹⁶¹ Ryazantsev, S. V.; Zasimov, P. V.; Feldman, V. I., X-ray radiolysis of C2 hydrocarbons in cryogenic media. *Radiat. Phys. Chem.* **2018**, *151*, 253-260. [DOI: 10.1016/j.radphyschem.2018.06.005]

162

Sakai, N.; Yamamoto, S., Warm carbon-chain chemistry. *Chem. Rev.* **2013**, *113* (12), 8981-9015. [DOI: 10.1021/cr4001308] 163

Bennett, C. J.; Jamieson, C. S.; Osamura, Y.; Kaiser, R. I., Laboratory Studies on the Irradiation of Methane in Interstellar, Cometary, and Solar System Ices. *Astrophys. J.* **2006**, *653* (1), 792-811. [DOI: 10.1086/508561]

¹⁶⁴ Kamp, I.; Thi, W.-F.; Woitke, P.; Ilee, J. D., Chemical networks. *EPJ Web Conf.* **2015**, *102*. [DOI: 10.1051/epjconf/201510200014]

¹⁶⁵ Allain, T.; Leach, S.; Sedlmayr, E. Photodestruction of PAHs in the interstellar medium. I. Photodissociation rates for the loss of an acetylenic group. *Astron. Astrophys.* **1996**, *305*, 602. [https://ui.adsabs.harvard.edu/abs/1996A&A...305..602A/abstract]

¹⁶⁶ Zhen, J.; Paardekooper, D. M.; Candian, A.; Linnartz, H.; Tielens, A. G. G. M., Quadrupole ion trap/time-of-flight photo-fragmentation spectrometry of the hexa-peri-hexabenzocoronene (HBC) cation. *Chem. Phys. Lett.* **2014**, *592*, 211-216. [DOI: 10.1016/j.cplett.2013.12.005]

¹⁶⁷ West, B.; Rodriguez Castillo, S.; Sit, A.; Mohamad, S.; Lowe, B.; Joblin, C.; Bodi, A.; Mayer, P. M., Unimolecular reaction energies for polycyclic aromatic hydrocarbon ions. *Phys. Chem. Chem. Phys.* **2018**, *20* (10), 7195-7205. [DOI: 10.1039/c7cp07369k]

¹⁶⁸ Rapacioli, M.; Cazaux, S.; Foley, N.; Simon, A.; Hoekstra, R.; Schlatholter, T., Atomic hydrogen interactions with gas-phase coronene cations: hydrogenation versus fragmentation. *Phys. Chem. Chem. Phys.* **2018**, *20* (35), 22427-22438. [DOI: 10.1039/c8cp03024c]

¹⁶⁹ Quitián-Lara, H. M.; Fantuzzi, F.; Nascimento, M. A. C.; Wolff, W.; Boechat-Roberty, H. M., Hydrogenated Benzene in Circumstellar Environments: Insights into the Photostability of Superhydrogenated PAHs. *Astrophys. J.* **2018**, *854* (1). [DOI: 10.3847/1538-4357/aaa977]

¹⁷⁰ Klærke, B.; Toker, Y.; Rahbek, D. B.; Hornekær, L.; Andersen, L. H., Formation and stability of hydrogenated PAHs in the gas phase. *Astron. Astrophys.* **2012**, *549*. [DOI: 10.1051/0004-6361/201219952]

¹⁷¹ Draine, B. T.; Li, A., Infrared Emission from Interstellar Dust. I. Stochastic Heating of Small Grains. *Astrophys. J.* **2001**, *551* (2), 807-824. [DOI: 10.1086/320227]

¹⁷² Ling, Y.; Lifshitz, C., Time-Dependent Mass Spectra and Breakdown Graphs. 21. C₁₄H₁₀ Isomers. *The J. Phys. Chem. A* **1998**, *102* (4), 708-716. [DOI: 10.1021/jp973167w]

¹⁷³ Le Page, V.; Snow, T. P.; Bierbaum, V. M., Hydrogenation and Charge States of Polycyclic Aromatic Hydrocarbons in Diffuse Clouds. II. Results. *Astrophys. J.* **2003**, *584* (1), 316-330. [DOI: 10.1086/345595]

¹⁷⁴ Murga, M. S.; Wiebe, D. S., Uncertainty in Estimates of Dissociation Rates of Polycyclic Aromatic Hydrocarbons in the Interstellar Medium. *Bull. Lebedev Phys. Inst.* **2023**, *49* (12), 416-421. [DOI: 10.3103/s1068335622120077]

¹⁷⁵ Murga, M. S.; Kirsanova, M. S.; Vasyunin, A. I.; Pavlyuchenkov, Y. N., Impact of PAH photodissociation on the formation of small hydrocarbons in the Orion Bar and the horsehead PDRs. *Mon. Not. R. Astron. Soc.* **2020**, *497* (2), 2327-2339. [DOI: 10.1093/mnras/staa2026]

¹⁷⁶ Pilleri, P.; Montillaud, J.; Berné, O.; Joblin, C., Evaporating very small grains as tracers of the UV radiation field in photo-dissociation regions. *Astron. Astrophys.* **2012**, *542*. [DOI: 10.1051/0004-6361/201015915]

¹⁷⁷ Alata, I.; Cruz-Diaz, G. A.; Muñoz Caro, G. M.; Dartois, E., Vacuum ultraviolet photolysis of hydrogenated amorphous carbons. *Astron. Astrophys.* **2014**, *569*, A119. [DOI: 10.1051/0004-6361/201323118]

¹⁷⁸ Alata, I.; Jallat, A.; Gavilan, L.; Chabot, M.; Cruz-Diaz, G. A.; Munoz Caro, G. M.; Béroff, K.; Dartois, E., Vacuum ultraviolet of hydrogenated amorphous carbons. *Astron. Astrophys.* **2015**, *584*. [DOI: 10.1051/0004-6361/201526368]

¹⁷⁹ Scott, A.; Duley, W. W.; Pinho, G. P., Polycyclic Aromatic Hydrocarbons and Fullerenes as Decomposition Products of Hydrogenated Amorphous Carbon. *Astrophys. J.* **1997**, *489* (2), L193-L195. [DOI: 10.1086/316789]

¹⁸⁰ Li, A.; Draine, B. T., The Carriers of the Interstellar Unidentified Infrared Emission Features: Aromatic or Aliphatic? *Astrophys. J.* **2012**, *760* (2). [DOI: 10.1088/2041-8205/760/2/L35]

¹⁸¹ Jones, A. P.; Fanciullo, L.; Köhler, M.; Verstraete, L.; Guillet, V.; Bocchio, M.; Ysard, N., The evolution of amorphous hydrocarbons in the ISM: dust modelling from a new vantage point. *Astron. Astrophys.* **2013**, *558*. [DOI: 10.1051/0004-6361/201321686]

¹⁸² Murga, M. S.; Vasyunin, A. I.; Kirsanova, M. S., Impact of HAC evolution on the formation of small hydrocarbons in the Orion Bar and the Horsehead PDRs. *Mon. Not. R. Astron. Soc.* **2023**, *519* (2), 2466-2474. [DOI: 10.1093/mnras/stac3656]

¹⁸³ Cuadrado, S.; Goicoechea, J. R.; Pilleri, P.; Cernicharo, J.; Fuente, A.; Joblin, C., The chemistry and spatial distribution of small hydrocarbons in UV-irradiated molecular clouds: the Orion Bar PDR. *Astron. Astrophys.* **2015**, *575*. [DOI: 10.1051/0004-6361/201424568]

hv ¹⁸⁴ Godard, B.; Falgarone, E.; Pineau des Forêts, G., Models of turbulent dissipation regions in the diffuse interstellar medium. *Astron. Astrophys.* **2009**, *495* (3), 847-867. [DOI: 10.1051/0004-6361:200810803]

¹⁸⁵ Murga, M. S.; Kirsanova, M. S.; Wiebe, D. S.; Boley, P. A., Orion Bar as a window to the evolution of PAHs. *Mon. Not. R. Astron. Soc.* **2022,** *509* (1), 800-817. [DOI: 10.1093/mnras/stab3061]

¹⁸⁶ Carr, J. S.; Najita, J. R., Organic molecules and water in the planet formation region of young circumstellar disks. *Science* **2008**, *319* (5869), 1504-6. [DOI: 10.1126/science.1153807]

¹⁸⁷ Bast, J. E.; Lahuis, F.; van Dishoeck, E. F.; Tielens, A. G. G. M., Exploring organic chemistry in planet-forming zones. *Astron. Astrophys.* **2013**, *551*. [DOI: 10.1051/0004-6361/201219908]

¹⁸⁸ van Dishoeck, E. F.; Grant, S.; Tabone, B.; van Gelder, M.; Francis, L.; Tychoniec, L.; Bettoni, G.; Arabhavi, A. M.; Gasman, D.; Nazari, P.; Vlasblom, M.; Kavanagh, P.; Christiaens, V.; Klaassen, P.; Beuther, H.; Henning, T.; Kamp, I., The diverse chemistry of protoplanetary disks as revealed by JWST. *Faraday Discuss.* **2023**, *245* (0), 52-79. [DOI: 10.1039/d3fd00010a]

¹⁸⁹ Tabone, B.; Bettoni, G.; van Dishoeck, E. F.; Arabhavi, A. M.; Grant, S.; Gasman, D.; Henning, T.; Kamp, I.; Güdel, M.; Lagage, P. O.; Ray, T.; Vandenbussche, B.; Abergel, A.; Absil, O.; Argyriou, I.; Barrado, D.; Boccaletti, A.; Bouwman, J.; Caratti o Garatti, A.; Geers, V.; Glauser, A. M.; Justannont, K.; Lahuis, F.; Mueller, M.; Nehmé, C.; Olofsson, G.; Pantin, E.; Scheithauer, S.; Waelkens, C.; Waters, L. B. F. M.; Black, J. H.; Christiaens, V.; Guadarrama, R.; Morales-Calderón, M.; Jang, H.; Kanwar, J.; Pawellek, N.; Perotti, G.; Perrin, A.; Rodgers-Lee, D.; Samland, M.; Schreiber, J.; Schwarz, K.; Colina, L.; Östlin, G.; Wright, G., A rich hydrocarbon chemistry and high C to O ratio in the inner disk around a very low-mass star. *Nat. Astron.* **2023**, 7 (7), 805-814. [DOI: 10.1038/s41550-023-01965-3]

¹⁹⁰ Grant, S. L.; van Dishoeck, E. F.; Tabone, B.; Gasman, D.; Henning, T.; Kamp, I.; Güdel, M.; Lagage, P.-O.; Bettoni, G.; Perotti, G.; Christiaens, V.; Samland, M.; Arabhavi, A. M.; Argyriou, I.; Abergel, A.; Absil, O.; Barrado, D.; Boccaletti, A.; Bouwman, J.; o Garatti, A. C.; Geers, V.; Glauser, A. M.; Guadarrama, R.; Jang, H.; Kanwar, J.; Lahuis, F.; Morales-Calderón, M.; Mueller, M.; Nehmé, C.; Olofsson, G.; Pantin, E.; Pawellek, N.; Ray, T. P.; Rodgers-Lee, D.; Scheithauer, S.; Schreiber, J.; Schwarz, K.; Temmink, M.; Vandenbussche, B.; Vlasblom, M.; Waters, L. B. F. M.; Wright, G.; Colina, L.; Greve, T. R.; Justannont, K.; Östlin, G., MINDS. The Detection of ¹³CO₂ with JWST-MIRI Indicates Abundant CO₂ in a Protoplanetary Disk. *Astrophys. J. Lett.* **2023**, *947* (1). [DOI: 10.3847/2041-8213/acc44b]

¹⁹¹ Wakelam, V.; Herbst, E.; Loison, J. C.; Smith, I. W. M.; Chandrasekaran, V.; Pavone, B.; Adams, N. G.; Bacchus-Montabonel, M. C.; Bergeat, A.; Béroff, K.; Bierbaum, V. M.; Chabot, M.; Dalgarno, A.; van Dishoeck, E. F.; Faure, A.; Geppert, W. D.; Gerlich, D.; Galli, D.; Hébrard, E.; Hersant, F.; Hickson, K. M.; Honvault, P.; Klippenstein, S. J.; Le Picard, S.; Nyman, G.; Pernot, P.; Schlemmer, S.; Selsis, F.; Sims, I. R.; Talbi, D.; Tennyson, J.; Troe, J.; Wester, R.; Wiesenfeld, L., A Kinetic Database for Astrochemistry (Kida). *Astrophys. J., Suppl. Ser.* **2012**, *199* (1). [DOI: 10.1088/0067-0049/199/1/21]

¹⁹² Lucas, R. & Liszt, H. S. Comparative chemistry of diffuse clouds. I. C₂H and C₃H₂. *Astron. Astrophys.* **2000**, *358*, 1069-1076. [https://adsabs.harvard.edu/full/2000A%26A...358.1069L]

¹⁹³ Pety, J.; Teyssier, D.; Fossé, D.; Gerin, M.; Roueff, E.; Abergel, A.; Habart, E.; Cernicharo, J., Are PAHs precursors of small hydrocarbons in photo-dissociation regions? The Horsehead case. *Astron. Astrophys.* **2005**, *435* (3), 885-899. [DOI: 10.1051/0004-6361:20041170]

¹⁹⁴ Pety, J.; Gratier, P.; Guzmán, V.; Roueff, E.; Gerin, M.; Goicoechea, J. R.; Bardeau, S.; Sievers, A.; Le Petit, F.; Le Bourlot, J.; Belloche, A.; Talbi, D., The IRAM-30 m line survey of the Horsehead PDR. *Astron. Astrophys.* **2012**, *548*. [DOI: 10.1051/0004-6361/201220062]

¹⁹⁵ Teyssier, D.; Fossé, D.; Gerin, M.; Pety, J.; Abergel, A.; Roueff, E., Carbon budget and carbon chemistry in Photon Dominated Regions. *Astron. Astrophys.* **2004**, *417* (1), 135-149. [DOI: 10.1051/0004-6361:20034534]

¹⁹⁶ Fuente, A.; Rodriguez-Franco, A.; Garcia-Burillo, S.; Martin-Pintado, J.; Black, J. H., Observational study of reactive ions and radicals in PDRs. *Astron. Astrophys.* **2003**, *406* (3), 899-913. [DOI: 10.1051/0004-6361:20030712]

¹⁹⁷ Tielens, A. G. G. M., *The Physics and Chemistry of the Interstellar Medium*. Publisher: Cambridge University Press. 2012. [DOI: 10.1017/CBO9780511819056]

¹⁹⁸ Morris, M.; Turner, B. E.; Palmer, P.; Zuckerman, B., Cyanoacetylene in dense interstellar clouds. *Astrophys. J.* **1976**, *205*. [DOI: 10.1086/154252]

¹⁹⁹ Gupta, H.; Gottlieb, C. A.; McCarthy, M. C.; Thaddeus, P., A Survey of C4h, C6h, and C6h– with the Green Bank Telescope. *Astrophys. J.* **2009**, *691* (2), 1494-1500. [DOI: 10.1088/0004-637X/691/2/1494]

²⁰⁰ Agundez, M.; Cernicharo, J.; Quintana-Lacaci, G.; Castro-Carrizo, A.; Velilla Prieto, L.; Marcelino, N.; Guelin, M.; Joblin, C.; Martin-Gago, J. A.; Gottlieb, C. A.; Patel, N. A.; McCarthy, M. C., The growth of carbon chains in IRC +10216 mapped with ALMA. *Astron. Astrophys.* **2017**, *601*. [DOI: 10.1051/0004-6361/201630274]

²⁰¹ Pardo, J. R.; Cernicharo, J.; Goicoechea, J. R.; Guelin, M.; Asensio Ramos, A., Molecular Line Survey of CRL 618 from 80 to 276 GHz and Complete Model. *Astrophys. J.* **2007**, *661* (1), 250-261. [DOI: 10.1086/513734]

²⁰² Winnewisser, G.; Walmsley, C. M., Carbon chain molecules in interstellar clouds. *Astrophys. Space Sci.* **1979**, *65* (1), 83-93. [DOI: 10.1007/BF00643491]

²⁰³ Cernicharo, J. The Polymerization of Acetylene, Hydrogen Cyanide, and Carbon Chains in the Neutral Layers of Carbon-rich Protoplanetary Nebulae. *Astrophys. J.* **2004**, *608*, L41. [DOI: 10.1086/422170]

²⁰⁴ Fukuzawa, K.; Osamura, Y.; Schaefer Iii, H. F., Are Neutral-Neutral Reactions Effective for the Carbon-Chain Growth of Cyanopolyynes and Polyacetylenes in Interstellar Space? *Astrophys. J.* **1998**, *505* (1), 278-285. [DOI: 10.1086/306168]

²⁰⁵ Chapman, J. F.; Millar, T. J.; Wardle, M.; Burton, M. G.; Walsh, A. J., Cyanopolyynes in hot cores: modelling G305.2+0.2. *Mon. Not. R. Astron. Soc.* **2009**, *394* (1), 221-230. [DOI: 10.1111/j.1365-2966.2008.14144.x]

²⁰⁶ Balucani, N.; Asvany, O.; Huang, L. C. L.; Lee, Y. T.; Kaiser, R. I.; Osamura, Y.; Bettinger, H. F., Formation of Nitriles in the Interstellar Medium via Reactions of Cyano Radicals, $CN(X 2\Sigma+)$, with Unsaturated Hydrocarbons. *Astrophys. J.* **2000**, *545* (2), 892-906. [DOI: 10.1086/317848]

²⁰⁷ Santoro, G.; Martinez, L.; Lauwaet, K.; Accolla, M.; Tajuelo-Castilla, G.; Merino, P.; Sobrado, J. M.; Pelaez, R. J.; Herrero, V. J.; Tanarro, I.; Mayoral, A. L.; Agundez, M.; Sabbah, H.; Joblin, C.; Cernicharo, J.; Martin-Gago, J. A., The Chemistry of Cosmic Dust Analogues from C, C(2), and C(2)H(2) in C-Rich Circumstellar Envelopes. *Astrophys J* **2020**, *895* (2). [DOI: 10.3847/1538-4357/ab9086]

²⁰⁸ Gu, X.; Kim, Y. S.; Kaiser, R. I.; Mebel, A. M.; Liang, M. C.; Yung, Y. L., Chemical dynamics of triacetylene formation and implications to the synthesis of polyynes in Titan's atmosphere. *Proc Natl Acad Sci U S A* **2009**, *106* (38), 16078-83. [DOI: 10.1073/pnas.0900525106]

²⁰⁹ He, C.; Thomas, A. M.; Galimova, G. R.; Mebel, A. M.; Kaiser, R. I., Gas Phase Formation of the Interstellar Molecule Methyltriacetylene. *Chem. Phys. Chem.* **2019**, *20* (15), 1912-1917. [DOI: 10.1002/cphc.201900305]

²¹⁰ Remijan, A. J.; Hollis, J. M.; Snyder, L. E.; Jewell, P. R.; Lovas, F. J., Methyltriacetylene (CH₃C₆H) toward TMC-1: The Largest Detected Symmetric Top. *Astrophys. J.* **2006**, *643* (1), L37-L40. [DOI: 10.1086/504918]

²¹¹ Cuylle, S. H.; Zhao, D.; Strazzulla, G.; Linnartz, H., Vacuum ultraviolet photochemistry of solid acetylene: a multispectral approach. *Astron. Astrophys.* **2014**, *570*. [DOI: 10.1051/0004-6361/201424379]

²¹² Compagnini, G.; D'Urso, L.; Puglisi, O.; Baratta, G. A.; Strazzulla, G., The irradiation of solid hydrocarbons and the formation of linear carbon chains. *Carbon* **2009**, *47* (6), 1605-1607. [DOI: 10.1016/j.carbon.2009.02.009]

²¹³ Cernicharo, J.; Guélin, M.; Agúndez, M.; Kawaguchi, K.; McCarthy, M.; Thaddeus, P., Astronomical detection of C₄H⁻, the second interstellar anion. *Astron. Astrophys.* **2007**, *4*67 (2), L37-L40. [DOI: 10.1051/0004-6361:20077415]

²¹⁴ Remijan, A. J.; Hollis, J. M.; Lovas, F. J.; Cordiner, M. A.; Millar, T. J.; Markwick-Kemper, A. J.; Jewell, P. R., Detection of C_8H^- and Comparison with C_8H toward IRC +10216. *Astrophys. J.* **2007**, *664* (1), L47-L50. [DOI: 10.1086/520704]

²¹⁵ Millar, T. J.; Walsh, C.; Field, T. A., Negative Ions in Space. *Chem. Rev.* **2017**, *117* (3), 1765-1795. [DOI: 10.1021/acs.chemrev.6b00480]

²¹⁶ Bastian, B.; Michaelsen, T.; Meyer, J.; Wester, R., Anionic Carbon Chain Growth in Reactions of C_2^- , C_4^- , C_6^- , C_2H^- , C_4H^- , and C_6H^- with C_2H_2 . *Astrophys. J.* **2019**, 878 (2). [ref: 10.3847/1538-4357/ab2042]

²¹⁷ Kaiser, R. I.; Mebel, A. M. On the formation of polyacetylenes and cyanopolyacetylenes in Titan's atmosphere and their role in astrobiology. *Chem. Soc. Rev.* **2012**, *41* (16), 5490-501. [DOI: 10.1039/c2cs35068h]

²¹⁸ McGuire, B. A., 2021 Census of Interstellar, Circumstellar, Extragalactic, Protoplanetary Disk, and Exoplanetary Molecules. *Astrophys. J., Suppl. Ser.* **2022**, *259* (2). [DOI: 10.3847/1538-4365/ac2a48]

²¹⁹ Herbst, E.; Garrod, R. T., Synthetic Approaches to Complex Organic Molecules in the Cold Interstellar Medium. *Front. Astron. Space Sci.* **2022**, *8*. [DOI: 10.3389/fspas.2021.789428]

²²⁰ Burke, D. J.; Brown, W. A., Ice in space: surface science investigations of the thermal desorption of model interstellar ices on dust grain analogue surfaces. *Phys. Chem. Chem. Phys.* **2010**, *12* (23), 5947-69. [DOI: 10.1039/B917005G]

²²¹ Oberg, K. I., Photochemistry and Astrochemistry: Photochemical Pathways to Interstellar Complex Organic Molecules. *Chem. Rev.* **2016**, *116* (17), 9631-63. [DOI: 10.1021/acs.chemrev.5b00694]

²²² Wakelam, V.; Loison, J. C.; Mereau, R.; Ruaud, M., Binding energies: New values and impact on the efficiency of chemical desorption. *Mol. Astrophys.* **2017**, *6*, 22-35. [DOI: 10.1016/j.molap.2017.01.002]

²²³ Garrod, R. T.; Weaver, S. L. W.; Herbst, E., Complex Chemistry in Star-forming Regions: An Expanded Gas-Grain Warm-up Chemical Model. *Astrophys. J.* **2008**, *682* (1), 283-302. [DOI: 10.1086/588035]

²²⁴ Jin, M.; Garrod, R. T., Formation of Complex Organic Molecules in Cold Interstellar Environments through Nondiffusive Grain-surface and Ice-mantle Chemistry. *Astrophys. J., Suppl. Ser.* **2020**, *249* (2). [DOI: 10.3847/1538-4365/ab9ec8]

²²⁵ Chuang, K. J.; Fedoseev, G.; Scirè, C.; Baratta, G. A.; Jäger, C.; Henning, T.; Linnartz, H.; Palumbo, M. E., Formation of complex organic molecules in molecular clouds: acetaldehyde, vinyl alcohol, ketene, and ethanol via the "energetic" processing of C_2H_2 ice. *Astron. Astrophys.* **2021**, *650*. [DOI: 10.1051/0004-6361/202140780]

²²⁶ Hudson, R. L.; Loeffler, M. J., Ketene Formation in Interstellar Ices: A Laboratory Study. Astrophys. J. 2013, 773 (2). [DOI: 10.1088/0004-637x/773/2/109]

²²⁷ Bergner, J. B.; Öberg, K. I.; Rajappan, M., Oxygen Atom Reactions with C₂H₆, C₂H₄, and C₂H₂ in Ices. *Astrophys. J.* **2019**, *874* (2). [DOI: 10.3847/1538-4357/ab07b2]

²²⁸ Bergner, J. B.; Öberg, K. I.; Garrod, R. T.; Graninger, D. M., Complex Organic Molecules toward Embedded Low-mass Protostars. *Astrophys. J.* **2017**, *841* (2). [DOI: 10.3847/1538-4357/aa72f6]

²²⁹ Zasimov, P. V.; Ryazantsev, S. V.; Tyurin, D. A.; Feldman, V. I., Radiation-induced chemistry in the C₂H₂–H₂O system at cryogenic temperatures: a matrix isolation study. *Mon. Not. R. Astron. Soc.* **2019**. [DOI: 10.1093/mnras/stz3228]

²³⁰ Hudson, R.; Moore, M., Hydrocarbon Radiation Chemistry in Ices of Cometary Relevance \bigstar . *Icarus* **1997**, *126* (1), 233-235. [DOI: 10.1006/icar.1997.5678]

²³¹ Robert Wu, C., Extreme Ultraviolet Photon-Induced Chemical Reactions in the C2H2–H2O Mixed Ices at 10 K. *Icarus* **2002**, *156* (2), 456-473. [DOI: 10.1006/icar.2001.6801]

²³² Hudson, R. L.; Moore, M. H., Solid-Phase Formation of Interstellar Vinyl Alcohol. *Astrophys. J.* **2003**, *586* (1), L107-L110. [DOI: 10.1086/374580]

²³³ Tielens, A. G.G.M. Grain Surface Chemistry. In *Chemistry and Spectroscopy of Interstellar Molecules*. Bohme, K.K., Ed.; Tokyo: University of Tokyo Press, p. 237, 1992. [https://ui.adsabs.harvard.edu/abs/1992csim.conf..237T/abstract]

²³⁴ Charnley, S. B., Acetaldehyde in star-forming regions. *Adv. Space Res.* **2004**, *33* (1), 23-30. [DOI: 10.1016/j.asr.2003.08.005]

²³⁵ Chuang, K. J.; Fedoseev, G.; Qasim, D.; Ioppolo, S.; Jäger, C.; Henning, T.; Palumbo, M. E.; van Dishoeck, E. F.; Linnartz, H., Formation of complex molecules in translucent clouds: acetaldehyde, vinyl alcohol, ketene, and ethanol via "nonenergetic" processing of C₂H₂ice. *Astron. Astrophys.* **2020**, *635*. [DOI: 10.1051/0004-6361/201937302]

²³⁶ Perrero, J.; Enrique-Romero, J.; Martinez-Bachs, B.; Ceccarelli, C.; Balucani, N.; Ugliengo, P.; Rimola, A., Non-energetic Formation of Ethanol via CCH Reaction with Interstellar H(2)O Ices. A Computational Chemistry Study. *ACS Earth Space Chem.* **2022**, *6* (3), 496-511. [DOI: 10.1021/acsearthspacechem.1c00369]

²³⁷ Zhou, L.; Kaiser, R. I.; Gao, L. G.; Chang, A. H. H.; Liang, M. C.; Yung, Y. L., Pathways to Oxygen-Bearing Molecules in the Interstellar Medium and in Planetary Atmospheres: Cyclopropenone (c-C₃H₂O) and Propynal (HCCCHO). *Astrophys. J.* **2008**, *686* (2), 1493-1502. [DOI: 10.1086/591072]

²³⁸ Wu, Y.-J.; Chuang, S.-J.; Chen, S.-C.; Huang, T.-P., Infrared Spectra of Acetylene Diluted in Solid Nitrogen Upon Irradiation with Vacuum Ultraviolet Light and Electrons. *Astrophys. J.*, *Suppl. Ser.* **2014**, *212* (1). [DOI: 10.1088/0067-0049/212/1/7]

²³⁹ Canta, A.; Öberg, K. I.; Rajappan, M., The Formation of Imines and Nitriles during VUV Photoirradiation of NH₃:C₂H x Ice Mixtures. *Astrophys. J.* **2023**, *953* (1). [DOI: 10.3847/1538-4357/acda99]

²⁴⁰ Zhang, C.; Wang, J.; Turner, A. M.; Marks, J. H.; Chandra, S.; Fortenberry, R. C.; Kaiser, R. I., On the Formation of Vinylamine (C₂H₃NH₂) in Interstellar Ice Analogs. *Astrophys. J.* **2023**, *952* (2). [DOI: 10.3847/1538-4357/acd451]

²⁴¹ Molpeceres, G.; Rivilla, V. M., Radical addition and H abstraction reactions in C_2H_2 , C_2H_4 , and C_2H_6 : A gateway for ethyl- and vinyl-bearing molecules in the interstellar medium. *Astron. Astrophys.* **2022**, *665*. [DOI: 10.1051/0004-6361/202243892]

²⁴² Baines, K. H.; Sromovsky, L. A.; Carlson, R. W.; Momary, T. W.; Fry, P. M., The visual spectrum of Jupiter's Great Red Spot accurately modeled with aerosols produced by photolyzed ammonia reacting with acetylene. *Icarus* **2019**, *330*, 217-229. [DOI: 10.1016/j.icarus.2019.04.008]

²⁴³ Carlson, R. W.; Baines, K. H.; Anderson, M. S.; Filacchione, G.; Simon, A. A., Chromophores from photolyzed ammonia reacting with acetylene: Application to Jupiter's Great Red Spot. *Icarus* **2016**, *274*, 106-115. [DOI: 10.1016/j.icarus.2016.03.008]

²⁴⁴ Bernard-Salas, J.; Peeters, E.; Sloan, G. C.; Cami, J.; Guiles, S.; Houck, J. R., The Spitzer IRS spectrum of SMP LMC 11. *Astrophys. J.* **2006**, *652* (1), L29-L32. [DOI: 10.1086/509777]

²⁴⁵ McGuire, B. A.; Burkhardt, A. M.; Kalenskii, S.; Shingledecker, C. N.; Remijan, A. J.; Herbst, E.; McCarthy, M. C., Detection of the aromatic molecule benzonitrile (c-C(6)H(5)CN) in the interstellar medium. *Science* **2018**, *359* (6372), 202-205. [DOI: 10.1126/science.aao4890]

²⁴⁶ Cherchneff, I. The formation of Polycyclic Aromatic Hydrocarbons in evolved circumstellar environments. *EAS Publ. Ser.* **2011**, *46*, 177. [DOI: 10.1051/eas/1146019]

²⁴⁷ Allamandola L.J., Tielens A.G.G.M., Barker J.R. Infrared Emission from Interstellar PAHs. In: Morfill G.E., Scholer M. (eds) Physical Processes in Interstellar Clouds. NATO ASI Series (Series C: Mathematical and Physical Sciences), vol 210. Springer, Dordrecht, 1987, pp. 305-331. [DOI: 10.1007/978-94-009-3945-5_20]

²⁴⁸ Ehrenfreund, P.; Cami, J., Cosmic carbon chemistry: from the interstellar medium to the early Earth. *Cold Spring Harbor Perspect. Biol.* **2010,** *2* (12), a002097. [DOI: 10.1101/cshperspect.a002097]

²⁴⁹ Kwok, S., The synthesis of organic and inorganic compounds in evolved stars. *Nature* **2004**, *430* (7003), 985-91. [DOI: 10.1038/nature02862]

²⁵⁰ Kwok, S., Delivery of Complex Organic Compounds from Planetary Nebulae to the Solar System. *Int. J. Astrobiol.* **2009**, 8 (3), 161-167. [DOI: 10.1017/S1473550409004492]

²⁵¹ Cherchneff, I.; Barker, J. R.; Tielens, A. G. G. M. Polycyclic aromatic hydrocarbon formation in carbon-rich stellar envelopes. *Astrophys. J.* **1992**, *401*, 269. [DOI: 10.1086/172059]

²⁵² Cole, J. A.; Bittner, J. D.; Longwell, J. P.; Howard, J. B. Formation mechanisms of aromatic compounds in aliphatic flames. *Combust. Flame* **1984**, *56*, 51. [DOI: 10.1016/0010-2180(84)90005-1]

²⁵³ Frenklach, M.; Warnatz, J. Detailed Modeling of PAH Profiles in a Sooting Low-Pressure Acetylene Flame. *Combust. Sci. Technol.* **1987**, *51*, 265. [DOI: 10.1080/00102208708960325]

²⁵⁴ Frenklach, M.; Feigelson, E. D. Formation of polycyclic aromatic hydrocarbons in circumstellar envelopes. *Astrophys. J.* **1989**, *341*, 372. [DOI: 10.1086/167501]

²⁵⁵ Miller, J. A.; Melius, C. F. Kinetic and thermodynamic issues in the formation of aromatic compounds in flames of aliphatic fuels. *Combust. Flame* **1992**, *91*, 21. [DOI: 10.1016/0010-2180(92)90124-8]

²⁵⁶ Zhao, L.; Lu, W.; Ahmed, M.; Zagidullin, M. V.; Azyazov, V. N.; Morozov, A. N.; Mebel, A. M.; Kaiser, R. I., Gas-phase synthesis of benzene via the propargyl radical self-reaction. *Sci. Adv.* **2021**, 7 (21). [DOI: 10.1126/sciadv.abf0360]

²⁵⁷ Richter, H.; Howard, J. B. Formation of polycyclic aromatic hydrocarbons and their growth to soot—a review of chemical reaction pathways. *Prog. Energy Combust. Sci.* **2000**, *26*, 565. [DOI: 10.1016/S0360-1285(00)00009-5]

²⁵⁸ Richter, H.; Howard, J. B. Formation and consumption of single-ring aromatic hydrocarbons and their precursors in premixed acetylene, ethylene and benzene flames. *Phys. Chem. Chem. Phys.* **2002**, *4*, 2038. [DOI: 10.1039/B110089K]

²⁵⁹ Lindstedt, R. P.; Skevis, G., Benzene formation chemistry in premixed, 1,3-butadiene flames. *Twenty-Sixth Symposium (International) on Combustion, Vols 1 and 2* **1996**, 703-709. [DOI: 10.1016/S0082-0784(96)80278-5]

²⁶⁰ Ikeda, E.; Tranter, R. S.; Kiefer, J. H.; Kern, R. D.; Singh, H. J.; Zhang, Q. The pyrolysis of methylcyclopentadiene: Isomerization and formation of aromatics. *Proc. Combust. Inst.* **2000**, *28*, 1725. [DOI: 10.1016/S0082-0784(00)80573-1]

²⁶¹ Moskaleva, L. V.; Mebel, A. M.; Lin, M. C. The CH3+C5H5 reaction: A potential source of benene at high temperatures. *Symposium (International) on Combustion* **1996**, *26*, 521. [DOI: 10.1016/S0082-0784(96)80255-4]

²⁶² Kaiser, R. I.; Zhao, L.; Lu, W.; Ahmed, M.; Zagidullin, M. V.; Azyazov, V. N.; Mebel, A. M., Formation of Benzene and Naphthalene through Cyclopentadienyl-Mediated Radical-Radical Reactions. *J. Phys. Chem. Lett.* **2022**, 13 (1), 208-213. [DOI: 10.1021/acs.jpclett.1c03733]

²⁶³ Kalvāns, J., Temperature Spectra of Interstellar Dust Grains Heated by Cosmic Rays. II. Dark Cloud Cores. *Astrophys. J., Suppl. Ser.* **2018**, *239* (1). [DOI: 10.3847/1538-4365/aae527]

²⁶⁴ Mitchell, J. L.; Lora, J. M., The Climate of Titan. *Annu. Rev. Earth Planet. Sci.* **2016**, *44* (1), 353-380. [DOI: 10.1146/annurev-earth-060115-012428]

²⁶⁵ Alkemade, U.; Homann, K. H., Formation of C₆H₆ Isomers by Recombination of Propynyl in the System Sodium Vapour/Propynylhalide. *Z. Phys. Chem.* **1989**, *161* (1-2), 19-34. [DOI: 10.1524/zpch.1989.161.Part_1_2.019]

²⁶⁶ Cernicharo, J.; Heras, A. M.; Pardo, J. R.; Tielens, A. G. G. M.; Guélin, M.; Dartois, E.; Neri, R.; Waters, L. B. F. M. Methylpolyynes and Small Hydrocarbons in CRL 618. *Astrophys. J.* 2001, 546, L127. [DOI: 10.1086/318872]

²⁶⁷ Kaiser, R. I. Experimental investigation on the formation of carbon-bearing molecules in the interstellar medium via neutral-neutral reactions. *Chem. Rev.* **2002**, *102*, 1309. [DOI: 10.1021/cr970004v]

²⁶⁸ Hiraoka, K.; Takayama, T.; Euchi, A.; Handa, H.; Sato, T. Study of the Reactions of H and D Atoms with Solid C_2H_2 , C_2H_4 , and C_2H_6 at Cryogenic Temperatures. *Astrophys. J.* **2000**, *532*, 1029. [DOI: 10.1086/308612]

²⁶⁹ Flower, D. R.; Forets, G. P. d., Grain-mantle erosion in magnetohydrodynamic shocks. *Mon. Not. R. Astron. Soc.* **1994**, *268* (3), 724-732. [DOI: 10.1093/mnras/268.3.724]

²⁷⁰ Markwick, A. J.; Millar, T. J.; Charnley, S. B., On the Abundance Gradients of Organic Molecules along the TMC-1 Ridge. *Astrophys. J.* **2000**, *535* (1), 256-265. [DOI: 10.1086/308814]

²⁷¹ Lukianova, M. A.; Feldman, V. I., Direct evidence for a single-step radiation-induced assembling of benzene ring from acetylene trimer at cryogenic temperatures. *Radiat. Phys. Chem.* **2021**, *183*. [DOI: 10.1016/j.radphyschem.2021.109417]

²⁷² Dangi, B. B.; Parker, D. S. N.; Yang, T.; Kaiser, R. I.; Mebel, A. M., Gas-Phase Synthesis of the Benzyl Radical ($C_6H_5CH_2$). *Angew. Chem.* **2014**, *126* (18), 4696-4701. [DOI: 10.1002/ange.201310612]

²⁷³ Zhang, F.; Parker, D.; Kim, Y. S.; Kaiser, R. I.; Mebel, A. M., On the formation of orthobenzyne (o- C_6H_4) under single collision conditions and its role in interstellar chemistry. *Astrophys. J.* **2011**, 728 (2). [DOI: 10.1088/0004-637X/728/2/141]

²⁷⁴ Bera, P. P.; Peverati, R.; Head-Gordon, M.; Lee, T. J., Hydrocarbon growth via ion-molecule reactions: computational studies of the isomers of C_4H_2+ , C_6H_2+ and C_6H_4+ and their formation paths from acetylene and its fragments. *Phys Chem Chem Phys* **2015**, *17* (3), 1859-69. [DOI: 10.1039/C4CP04480K]

²⁷⁵ Peverati, R.; Bera, P. P.; Lee, T. J.; Head-Gordon, M., Formation and stability of C_6H_3 + isomers. *J Phys Chem A* **2014**, *118* (43), 10109-16. [DOI: 10.1021/jp5081862]

²⁷⁶ Peverati, R.; Bera, P. P.; Lee, T. J.; Head-Gordon, M., Insights into hydrocarbon chain and aromatic ring formation in the interstellar medium: computational study of the isomers of C_4H_{3+} , C_6H_{3+} and $C6H_{5+}$ and their formation pathways. *Astrophys. J.* **2016**, *830* (2). [DOI: 10.3847/0004-637X/830/2/128]

²⁷⁷ Coustenis, A.; Salama, A.; Schulz, B.; Ott, S.; Lellouch, E.; Encrenaz, T. h.; Gautier, D.; Feuchtgruber, H. Titan's atmosphere from ISO mid-infrared spectroscopy. *Icarus* **2003**, *161*, 383. [DOI: 10.1016/S0019-1035(02)00028-3]

²⁷⁸ Wong, A.-S.; Yung, Y. L.; Friedson, A. J. Benzene and Haze Formation in the Polar Atmosphere of Jupiter. *Geophys. Res. Lett.* **2003**, *30* [DOI: 10.1029/2002GL016661]

²⁷⁹ Wilson, E. H.; Atreya, S. K.; Coustenis, A. Mechanisms for the formation of benzene in the atmosphere of Titan. *J. Geophys. Res.: Planets* **2003**, *108*, n/a. [DOI: 10.1029/2002JE001896]

²⁸⁰ Lebonnois, S. Benzene and aerosol production in Titan and Jupiter's atmospheres: a sensitivity study. *Planet. Space Sci.* **2005**, *53*, 486. [DOI: 10.1016/j.pss.2004.11.004]

²⁸¹ Lavvas, P. P.; Coustenis, A.; Vardavas, I. M. Coupling photochemistry with haze formation in Titan's atmosphere, Part II: Results and validation with Cassini/Huygens data. *Planet. Space Sci.* **2008**, *56*, 67. [DOI: 10.1016/j.pss.2007.05.027]

²⁸² Waite, J. H., Jr.; Niemann, H.; Yelle, R. V.; Kasprzak, W. T.; Cravens, T. E.; Luhmann, J. G.; McNutt, R. L.; Ip, W. H.; Gell, D.; De La Haye, V.; Muller-Wordag, I.; Magee, B.; Borggren, N.; Ledvina, S.; Fletcher, G.; Walter, E.; Miller, R.; Scherer, S.; Thorpe, R.; Xu, J.; Block, B.; Arnett, K., Ion neutral mass spectrometer results from the first flyby of Titan. *Science* **2005**, *308* (5724), 982-6. [DOI: 10.1126/science.1110652]

²⁸³ Waite, J. H., Jr.; Young, D. T.; Cravens, T. E.; Coates, A. J.; Crary, F. J.; Magee, B.; Westlake, J., The process of tholin formation in Titan's upper atmosphere. *Science* **2007**, *316* (5826), 870-5. [DOI: 10.1126/science.1139727]

²⁸⁴ Vuitton, V.; Yelle, R. V.; Cui, J. Formation and distribution of benzene on Titan. J. Geophys. *Res.* **2008**, *113*. [DOI: 10.1029/2007JE002997]

²⁸⁵ Anicich, V. G.; McEwan, M. J. Ion-molecule chemistry in Titan's ionosphere. *Planet. Space Sci.***1997**, *45*, 897. [DOI: 10.1016/S0032-0633(97)00053-6]

²⁸⁶ Zhou, L.; Zheng, W.; Kaiser, R. I.; Landera, A.; Mebel, A. M.; Liang, M.-C.; Yung, Y. L. Cosmic-Ray-Mediated Formation of Benzene on the Surface of Saturn's Moon Titan. *Astrophys. J.* **2010**, *718*, 1243. [DOI: 10.1088/0004-637X/718/2/1243]

²⁸⁷ Kleimeier, N. F.; Liu, Y.; Turner, A. M.; Young, L. A.; Chin, C. H.; Yang, T.; He, X.; Lo, J. I.; Cheng, B. M.; Kaiser, R. I., Excited state photochemically driven surface formation of benzene from acetylene ices on Pluto and in the outer solar system. *Phys. Chem. Chem. Phys.* **2022**, 24 (3), 1424-1436. [DOI: 10.1039/d1cp04959c]

²⁸⁸ Tsuge, M.; Bahou, M.; Wu, Y.-J.; Allamandola, L.; Lee, Y.-P. The Infrared Spectrum of Protonated Ovalene in Solid Para-Hydrogen and Its Possible Contribution to Interstellar Unidentified Infrared Emission. *Astrophys. J.* **2016**, *825*, 96. [DOI: 10.3847/0004-637X/825/2/96]

²⁸⁹ Kim, H. S.; Wagner, D. R.; Saykally, R. J. Single photon infrared emission spectroscopy of the gas phase pyrene cation: support for a polycyclic aromatic hydrocarbon origin of the unidentified infrared emission bands. *Phys. Rev. Lett.* **2001**, *86*, 5691. [DOI: 10.1103/PhysRevLett.86.5691]

²⁹⁰ Doddipatla, S.; Galimova, G. R.; Wei, H.; Thomas, A. M.; He, C.; Yang, Z.; Morozov, A. N.; Shingledecker, C. N.; Mebel, A. M.; Kaiser, R. I. Low-temperature gas-phase formation of indene in the interstellar medium. *Sci. Adv.* **2021**, *7*. [DOI: 10.1126/sciadv.abd4044]

²⁹¹ Tielens, A. G. G. M. Interstellar Polycyclic Aromatic Hydrocarbon Molecules. *Annu. Rev. Astron. Astrophys.* **2008**, *46*, 289. [DOI: 10.1146/annurev.astro.46.060407.145211]

²⁹² Rawlings, M. G.; Adamson, A. J.; Marshall, C. C. M.; Sarre, P. J. Near-infrared diffuse interstellar bands towards Her 36. *Mon. Not. R. Astron. Soc.* **2019**, *485*, 3398. [DOI: 10.1093/mnras/stz663]

²⁹³ Cox, N. L. J.; Cami, J.; Farhang, A.; Smoker, J.; Monreal-Ibero, A.; Lallement, R.; Sarre, P. J.; Marshall, C. C. M.; Smith, K. T.; Evans, C. J.; Royer, P.; Linnartz, H.; Cordiner, M. A.; Joblin, C.; van Loon, J. T.; Foing, B. H.; Bhatt, N. H.; Bron, E.; Elyajouri, M.; de Koter, A.; Ehrenfreund, P.; Javadi, A.; Kaper, L.; Khosroshadi, H. G.; Laverick, M.; Le Petit, F.; Mulas, G.; Roueff, E.; Salama, F.; Spaans, M. The ESO Diffuse Interstellar Bands Large Exploration Survey: EDIBLES I. Project description, survey sample and quality assessment. *Astron. Astrophys.* **2017**, *606*. [DOI: 10.1051/0004-6361/201730912]

²⁹⁴ Ruiterkamp, R.; Halasinski, T.; Salama, F.; Foing, B. H.; Allamandola, L. J.; Schmidt, W.; Ehrenfreund, P. Spectroscopy of large PAHs. *Astron. Astrophys.* **2002**, *390*, 1153. [DOI: 10.1051/0004-6361:20020478]

²⁹⁵ Tsuge, M.; Tseng, C. Y.; Lee, Y. P. Spectroscopy of prospective interstellar ions and radicals isolated in para-hydrogen matrices. *Phys. Chem. Chem. Phys.* **2018**, *20*, 5344. [DOI: 10.1039/C7CP05680J]

²⁹⁶ Ruiterkamp, R.; Cox, N. L. J.; Spaans, M.; Kaper, L.; Foing, B. H.; Salama, F.; Ehrenfreund, P. PAH charge state distribution and DIB carriers: Implications from the line of sight toward HD 147889. *Astron. Astrophys.* **2005**, *432*, 515. [DOI: 10.1051/0004-6361:20040331]

²⁹⁷ Salama, F.; Galazutdinov, G. A.; Krełowski, J.; Biennier, L.; Beletsky, Y.; Song, I.-O., Polycyclic Aromatic Hydrocarbons and the Diffuse Interstellar Bands: A Survey. *Astrophys. J.* **2011**, 728 (2). [DOI: 10.1088/0004-637X/728/2/154]

²⁹⁸ Omont, A.; Bettinger, H. F.; Tönshoff, C., Polyacenes and diffuse interstellar bands. *Astron. Astrophys.* **2019**, *625*. [DOI: 10.1051/0004-6361/201834953]

²⁹⁹ Hensley, B. S.; Draine, B. T., The Astrodust+PAH Model: A Unified Description of the Extinction, Emission, and Polarization from Dust in the Diffuse Interstellar Medium. *Astrophys. J.* **2023**, *948* (1). [DOI: 10.3847/1538-4357/acc4c2]

³⁰⁰ McGuire, B. A.; Loomis, R. A.; Burkhardt, A. M.; Lee, K. L. K.; Shingledecker, C. N.; Charnley, S. B.; Cooke, I. R.; Cordiner, M. A.; Herbst, E.; Kalenskii, S.; Siebert, M. A.; Willis, E. R.; Xue, C.; Remijan, A. J.; McCarthy, M. C., Detection of two interstellar polycyclic aromatic hydrocarbons via spectral matched filtering. *Science* **2021**, *371* (6535), 1265-1269. [DOI: 10.1126/science.abb7535]

³⁰¹ Cernicharo, J.; Agundez, M.; Cabezas, C.; Tercero, B.; Marcelino, N.; Pardo, J. R.; de Vicente, P., Pure hydrocarbon cycles in TMC-1: Discovery of ethynyl cyclopropenylidene, cyclopentadiene and indene. *Astron. Astrophys.* **2021**, *649*. [DOI: 10.1051/0004-6361/202141156]

³⁰² Zenobi, R.; Philippoz, J. M.; Zare, R. N.; Buseck, P. R. Spatially resolved organic analysis of the allende meteorite. *Science* **1989**, *246*, 1026. [DOI: 10.1126/science.246.4933.1026]

³⁰³ Wang, Y.; Huang, Y.; Alexander, C. M. O. D.; Fogel, M.; Cody, G. Molecular and compoundspecific hydrogen isotope analyses of insoluble organic matter from different carbonaceous chondrite groups. *Geochim. Cosmochim. Acta* **2005**, *69*, 3711. [DOI: 10.1016/j.gca.2005.03.008]

³⁰⁴ Messenger, S.; Amari, S.; Gao, X.; Walker, R. M.; Clemett, S. J.; Chillier, X. D. F.; Zare, R. N.; Lewis, R. S. Indigenous Polycyclic Aromatic Hydrocarbons in Circumstellar Graphite Grains from Primitive Meteorites. *Astrophys. J.* **1998**, *502*, 284. [DOI: 10.1086/305874]

³⁰⁵ Hahn, J. H.; Zenobi, R.; Bada, J. L.; Zare, R. N. Application of two-step laser mass spectrometry to cosmogeochemistry: direct analysis of meteorites. *Science* **1988**, *239*, 1523. [DOI: 10.1126/science.239.4847.1523]

³⁰⁶ Plows, F. L.; Elsila, J. E.; Zare, R. N.; Buseck, P. R. Evidence that polycyclic aromatic hydrocarbons in two carbonaceous chondrites predate parent-body formation. *Geochim. Cosmochim. Acta* **2003**, *67*, 1429. [DOI: 10.1016/S0016-7037(02)01277-2]

³⁰⁷ Elsila, J. E.; de Leon, N. P.; Buseck, P. R.; Zare, R. N. Alkylation of polycyclic aromatic hydrocarbons in carbonaceous chondrites. *Geochim. Cosmochim. Acta* **2005**, *69*, 1349. [DOI: 10.1016/j.gca.2004.09.009]

³⁰⁸ Kaiser, K.; Schulz, F.; Maillard, J. F.; Hermann, F.; Pozo, I.; Peña, D.; Cleaves, H. J.; Burton, A. S.; Danger, G.; Afonso, C.; Sandford, S.; Gross, L., Visualization and identification of single meteoritic organic molecules by atomic force microscopy. *Meteorit. Planet. Sci.* **2022**. [DOI: 10.1111/maps.13784]

³⁰⁹ Boersma, C.; Bregman, J.; Allamandola, L. J. The Charge State of Polycyclic Aromatic Hydrocarbons across a Reflection Nebula, an H ii Region, and a Planetary Nebula. *Astrophys. J.* **2018**, *858*, 67. [DOI: 10.3847/1538-4357/aabcbe]

³¹⁰ Ehrenfreund, P., Irvine, W.M., Owen, T., Becker, L., Blank, J., Brucato, J.R., Colangeli, L., Derenne, S., Dutrey, A., Despois, D., Lazcano, A., Robert, F. (Eds.) Astrobiology. Future Perspectives, Springer Netherlands, **2004**. [DOI: 10.1007/1-4020-2305-7]

³¹¹ Duley, W. W. Polycyclic aromatic hydrocarbons, carbon nanoparticles and the diffuse interstellar bands. *Faraday Discuss.* **2006**, *133*, 415. [DOI: 10.1039/B516323D]

³¹² d'Hendecourt, L.; Ehrenfreund, P. Spectroscopic properties of polycyclic aromatic hydrocarbons (PAHs) and astrophysical implications. *Adv. Space Res.* **1997**, *19*, 1023. [DOI: 10.1016/S0273-1177(97)00349-9]

³¹³ Rhee, Y. M.; Lee, T. J.; Gudipati, M. S.; Allamandola, L. J.; Head-Gordon, M. Charged polycyclic aromatic hydrocarbon clusters and the galactic extended red emission. *Proc. Natl. Acad. Sci. U. S. A.* **2007**, *104*, 5274. [DOI: 10.1073/pnas.0609396104]

³¹⁴ Lemmens, A. K.; Rap, D. B.; Thunnissen, J. M. M.; Willemsen, B.; Rijs, A. M. Polycyclic aromatic hydrocarbon formation chemistry in a plasma jet revealed by IR-UV action spectroscopy. *Nat. Commun.* **2020**, *11*, 269. [DOI: 10.1038/s41467-019-14092-3]

³¹⁵ Cook, D. J.; Schlemmer, S.; Balucani, N.; Wagner, D. R.; Steiner, B.; Saykally, R. J. Infrared emission spectra of candidate interstellar aromatic molecules. *Nature* **1996**, *380*, 227. [DOI: 10.1038/380227a0]

³¹⁶ Herbst, E. Astrochemistry and Star Formation: Successes and Challenges, WILEY-VCH, Weinheim, 2006.

³¹⁷ Habart, E.; Boulanger, F.; Verstraete, L.; Walmsley, C. M.; Forêts, G. P. d., Some empirical estimates of the H2formation rate in photon-dominated regions. *Astron. Astrophys.* **2004**, *414* (2), 531-544. [DOI: 10.1051/0004-6361:20031659]

³¹⁸ Zhang, F.; Jones, B.; Maksyutenko, P.; Kaiser, R. I.; Chin, C.; Kislov, V. V.; Mebel, A. M., Formation of the phenyl radical $[C_6H_5(X(2)A1)]$ under single collision conditions: a crossed molecular beam and ab initio study. *J. Am. Chem. Soc.* **2010**, *132* (8), 2672-83. [DOI: 10.1021/ja908559v]

³¹⁹ Sagan, C.; Khare, B. N.; Thompson, W. R.; McDonald, G. D.; Wing, M. R.; Bada, J. L.; Vo-Dinh, T.; Arakawa, E. T. Polycyclic aromatic hydrocarbons in the atmospheres of Titan and Jupiter. *Astrophys. J.* **1993**, *414*, 399. [DOI: 10.1086/173086] ³²⁰ Kahl, G. PAH world hypothesis (polycyclic aromatic hydrocarbons world). In *The Dictionary* of Genomics, Transcriptomics and Proteomics, Kahl, G., Ed.; 2015. [DOI: 10.1002/9783527678679.dg09094]

³²¹ d'Ischia, M.; Manini, P.; Martins, Z.; Remusat, L.; Alexander, C. M. O'D.; Puzzarini, C.; Barone, V.; Saladino, R., Insoluble organic matter in chondrites: Archetypal melanin-like PAH-based multifunctionality at the origin of life? *Phys. Life Rev.* **2021**, *37*, 65-93. [DOI: 10.1016/j.plrev.2021.03.002]

³²² Micelotta, E. R.; Jones, A. P.; Tielens, A. G. G. M. Polycyclic aromatic hydrocarbon processing by cosmic rays. *Astron. Astrophys.* **2010**, *526*, A52. [DOI: 10.1051/0004-6361/201015741]

³²³ Micelotta, E. R.; Jones, A. P.; Tielens, A. G. G. M., Polycyclic aromatic hydrocarbon processing in interstellar shocks. *Astron. Astrophys.* **2010**, *510*, A36. [DOI: 10.1051/0004-6361/200911682]

³²⁴ Micelotta, E. R.; Jones, A. P.; Tielens, A. G. G. M. Polycyclic aromatic hydrocarbon processing in a hot gas. *Astron. Astrophys.* **2010**, *510*, A37. [DOI: 10.1051/0004-6361/200911683]

³²⁵ Murga, M. S.; Wiebe, D. S.; Sivkova, E. E.; Akimkin, V. V. SHIVA: a dust destruction model. *Mon. Not. R. Astron. Soc.* **2019**, *488* (1), 965-977. [DOI: 10.1093/mnras/stz1724]

³²⁶ Parker, D. S.; Kaiser, R. I. On the formation of nitrogen-substituted polycyclic aromatic hydrocarbons (NPAHs) in circumstellar and interstellar environments. *Chem. Soc. Rev.* **2017**, *46*, 452. [DOI: 10.1039/C6CS00714G]

³²⁷ Hirashita, H., Shattering by turbulence as a production source of very small grains. *Mon. Not. R. Astron. Soc.* **2010**, *407* (1), L49-L53. [DOI: 10.1111/j.1745-3933.2010.00902.x]

³²⁸ Matsuura, M.; De Buizer, J. M.; Arendt, R. G.; Dwek, E.; Barlow, M. J.; Bevan, A.; Cigan, P.; Gomez, H. L.; Rho, J.; Wesson, R.; Bouchet, P.; Danziger, J.; Meixner, M., SOFIA mid-infrared observations of Supernova 1987A in 2016 – forward shocks and possible dust re-formation in the post-shocked region. *Mon. Not. R. Astron. Soc.* **2019**, *482* (2), 1715-1723. [DOI: 10.1093/mnras/sty2734]

³²⁹ Sandstrom, K. M.; Bolatto, A. D.; Draine, B. T.; Bot, C.; Stanimirović, S., Thespitzersurvey of the Small Magellanic Cloud (S3mc): Insights into the Life Cycle of Polycyclic Aromatic Hydrocarbons. *Astrophys. J.* **2010**, *715* (2), 701-723. [DOI: 10.1088/0004-637X/715/2/701]

³³⁰ He, C.; Kaiser, R. I.; Lu, W.; Ahmed, M.; Krasnoukhov, V. S.; Pivovarov, P. S.; Zagidullin, M. V.; Azyazov, V. N.; Morozov, A. N.; Mebel, A. M., Unconventional gas-phase preparation of the prototype polycyclic aromatic hydrocarbon naphthalene (C(10)H(8)) via the reaction of benzyl (C(7)H(7)) and propargyl (C(3)H(3)) radicals coupled with hydrogen-atom assisted isomerization. *Chem. Sci.* **2023**, *14* (20), 5369-5378. [DOI: 10.1039/D3SC00911D]

³³¹ Hirashita, H.; Murga, M. S., Self-consistent modelling of aromatic dust species and extinction curves in galaxy evolution. *Mon. Not. R. Astron. Soc.* **2020**, *492* (3), 3779-3793. [DOI: 10.1093/mnras/stz3640]

³³² Lemmens, A. K.; Rap, D. B.; Brunken, S.; Buma, W. J.; Rijs, A. M., Polycyclic aromatic hydrocarbon growth in a benzene discharge explored by IR-UV action spectroscopy. *Phys. Chem. Chem. Phys.* **2022**, 24 (24), 14816-14824. [DOI: 10.1039/d2cp01631a]

³³³ Bockhorn, H.; Fetting, F.; Wenz, H. W. *Berichte der Bunsengesellschaft für physikalische Chemie* **1983**, 87, 1067. [DOI: 10.1002/bbpc.19830871121]

³³⁴ Frenklach, M.; Wang, H. Detailed modeling of soot particle nucleation and growth. *Symposium* (*International*) on Combustion **1991**, *23*, 1559. [DOI: 10.1016/S0082-0784(06)80426-1]

³³⁵ Frenklach, M., Reaction mechanism of soot formation in flames. *Phys. Chem. Chem. Phys.* **2002**, *4* (11), 2028-2037. [DOI: 10.1039/B110045A]

³³⁶ Wang, H.; Frenklach, M. A detailed kinetic modeling study of aromatics formation in laminar premixed acetylene and ethylene flames. *Combust. Flame* **1997**, *110*, 173. [DOI: 10.1016/S0010-2180(97)00068-0]

³³⁷ Chu, T. C.; Buras, Z. J.; Smith, M. C.; Uwagwu, A. B.; Green, W. H., From benzene to naphthalene: direct measurement of reactions and intermediates of phenyl radicals and acetylene. *Phys. Chem. Chem. Phys.* **2019**, *21* (40), 22248-22258. [DOI: 10.1039/C9CP04554F]

³³⁸ Yang, T.; Troy, T. P.; Xu, B.; Kostko, O.; Ahmed, M.; Mebel, A. M.; Kaiser, R. I. Hydrogen-Abstraction/Acetylene-Addition Exposed. *Angew. Chem. Int. Ed. Engl.* **2016**, *55*, 14983. [DOI: 10.1002/anie.201607509]

³³⁹ Cherchneff, I., The inner wind of IRC +10216 revisited: new exotic chemistry and diagnostic for dust condensation in carbon stars. *Astron. Astrophys.* **2012**, *545*. [DOI: 10.1051/0004-6361/201118542]

³⁴⁰ Mebel, A. M.; Kislov, V. V., Can the C(5)H(5) + C(5)H(5) --> C(10)H(10) --> C(10)H(9) + H/C(10)H(8) + H(2) reaction produce naphthalene? An Ab initio/RRKM study. *J. Phys. Chem. A* **2009**, *113* (36), 9825-33. [DOI: 10.1021/jp905931j]

³⁴¹ Kislov, V. V.; Sadovnikov, A. I.; Mebel, A. M. Formation mechanism of polycyclic aromatic hydrocarbons beyond the second aromatic ring. *J. Phys. Chem. A* **2013**, *117*, 4794. [DOI: 10.1021/jp402481y]

³⁴² Parker, D. S.; Kaiser, R. I.; Bandyopadhyay, B.; Kostko, O.; Troy, T. P.; Ahmed, M. Unexpected chemistry from the reaction of naphthyl and acetylene at combustion-like temperatures. *Angew. Chem. Int. Ed. Engl.* **2015**, *54*, 5421. [DOI: 10.1002/anie.201411987]

³⁴³ Kislov, V. V.; Islamova, N. I.; Kolker, A. M.; Lin, S. H.; Mebel, A. M., Hydrogen Abstraction Acetylene Addition and Diels-Alder Mechanisms of PAH Formation: A Detailed Study Using First Principles Calculations. *J. Chem. Theory Comput.* **2005**, *1* (5), 908-24. [DOI: 10.1021/ct0500491]

³⁴⁴ Badger, G. M.; Donnelly, J. K.; Spotswood, T. M., The formation of aromatic hydrocarbons at high temperatures. XXIV. The pyrolysis of some tobacco constituents. *Aust. J. Chem.* **1965**, *18* (8). [DOI: 10.1071/CH9651249]

³⁴⁵ Sarofim A.F., Longwell J.P., Wornat M.J., Mukherjee J. The Role of Biaryl Reactions in PAH and Soot Formation. In: Bockhorn H. (eds) Soot Formation in Combustion. Springer Series in Chemical Physics, vol 59. Springer, Berlin, Heidelberg, 1994. [DOI: 10.1007/978-3-642-85167-4_27]

³⁴⁶ Griesheimer, J.; Homann, K. H. Large molecules, radicals ions, and small soot particles in fuelrich hydrocarbon flames. *Symposium (International) on Combustion* **1998**, *27*, 1753. [DOI: 10.1016/S0082-0784(98)80016-7]

³⁴⁷ Böhm, H.; Jander, H.; Tanke, D. PAH growth and soot formation in the pyrolysis of acetylene and benzene at high temperatures and pressures: Modeling and experiment. *Symp. (Int.) Combust.* **1998**, *27 (1)*, 1605-1612. [DOI: 10.1016/S0082-0784(98)80570-5]

³⁴⁸ Dong, G. L.; Hüttinger, K. J. Consideration of reaction mechanisms leading to pyrolytic carbon of different textures. *Carbon* **2002**, *40*, 2515. [DOI: 10.1016/S0008-6223(02)00174-4]

³⁴⁹ Zhang, F.; Gu, X.; Kaiser, R. I. Formation of the diphenyl molecule in the crossed beam reaction of phenyl radicals with benzene. *J. Chem. Phys.* **2008**, *128*, 084315. [DOI: 10.1063/1.2834221]

³⁵⁰ Schuetz, C. A.; Frenklach, M. Nucleation of soot: Molecular dynamics simulations of pyrene dimerization. *Proc. Combust. Inst.* **2002**, *29*, 2307. [DOI: 10.1016/S1540-7489(02)80281-4]

³⁵¹ Sabbah, H.; Biennier, L.; Klippenstein, S. J.; Sims, I. R.; Rowe, B. R. Exploring the Role of PAHs in the Formation of Soot: Pyrene Dimerization. *J. Phys. Chem. Lett.* **2010**, *1*, 2962. [DOI: 10.1021/jz101033t]

³⁵² He, C.; Fujioka, K.; Nikolayev, A. A.; Zhao, L.; Doddipatla, S.; Azyazov, V. N.; Mebel, A. M.; Sun, R.; Kaiser, R. I., A chemical dynamics study of the reaction of the methylidyne radical (CH, X(2)Pi) with dimethylacetylene (CH(3)CCCH(3), X(1)A(1g)). *Phys. Chem. Chem. Phys.* **2021**, 24 (1), 578-593. [DOI: 10.1039/D1CP04443E]

³⁵³ Zhao, L.; Kaiser, R. I.; Lu, W.; Ahmed, M.; Oleinikov, A. D.; Azyazov, V. N.; Mebel, A. M.; Howlader, A. H.; Wnuk, S. F., Gas phase formation of phenalene via 10pi-aromatic, resonantly stabilized free radical intermediates. *Phys. Chem. Chem. Phys.* **2020**, 22 (27), 15381-15388. [DOI: 10.1039/D0CP02216K]

³⁵⁴ Zhao, L.; Prendergast, M. B.; Kaiser, R. I.; Xu, B.; Ablikim, U.; Ahmed, M.; Sun, B. J.; Chen, Y. L.; Chang, A. H. H.; Mohamed, R. K.; Fischer, F. R., Synthesis of Polycyclic Aromatic Hydrocarbons by Phenyl Addition-Dehydrocyclization: The Third Way. *Angew. Chem. Int. Ed. Engl.* **2019**, *58* (48), 17442-17450. [DOI: 10.1002/anie.201909876]

³⁵⁵ Yang, J.; Smith, M. C.; Prendergast, M. B.; Chu, T. C.; Green, W. H., C(14)H(10) polycyclic aromatic hydrocarbon formation by acetylene addition to naphthalenyl radicals observed. *Phys. Chem. Chem. Phys.* **2021**, 23 (26), 14325-14339. [DOI: 10.1039/d1cp01565f]

³⁵⁶ Parker, D. S.; Kaiser, R. I.; Kostko, O.; Ahmed, M. Selective Formation of Indene through the Reaction of Benzyl Radicals with Acetylene. *ChemPhysChem* **2015**, *16*, 2091. [DOI: 10.1002/cphc.201500313]

³⁵⁷ Bittner, J. D.; Howard, J. B. Composition profiles and reaction mechanisms in a near-sooting premixed benzene/oxygen/argon flame. *Symposium (International) on Combustion* **1981**, *18*, 1105. [DOI: 10.1016/S0082-0784(81)80115-4]

³⁵⁸ He, C.; Kaiser, R. I.; Lu, W.; Ahmed, M.; Pivovarov, P. S.; Kuznetsov, O. V.; Zagidullin, M. V.; Mebel, A. M., Unconventional Pathway in the Gas-Phase Synthesis of 9H-Fluorene (C₁₃H₁₀) via the Radical–Radical Reaction of Benzyl (C₇H₇) with Phenyl (C₆H₅). *Angew. Chem.* **2023**, 135 (6). [DOI: 10.1002/ange.202216972]

³⁵⁹ Li, W.; Zhao, L.; Kaiser, R. I., A unified reaction network on the formation of five-membered ringed polycyclic aromatic hydrocarbons (PAHs) and their role in ring expansion processes through radical-radical reactions. *Phys. Chem. Chem. Phys.* **2023**, 25 (5), 4141-4150. [DOI: 10.1039/D2CP05305E]

³⁶⁰ Zhao, L.; Kaiser, R. I.; Lu, W.; Kostko, O.; Ahmed, M.; Evseev, M. M.; Bashkirov, E. K.; Oleinikov, A. D.; Azyazov, V. N.; Mebel, A. M.; Howlader, A. H.; Wnuk, S. F., Gas phase formation of cyclopentanaphthalene (benzindene) isomers via reactions of 5- and 6-indenyl radicals with vinylacetylene. *Phys. Chem. Chem. Phys.* **2020**, 22 (39), 22493-22500. [DOI: 10.1039/d0cp03846f]

³⁶¹ Necula, A.; Scott, L. T., Trapping Aryl Radicals with Acetylene: Evidence for C2-Accretion as a Mechanism for Polycyclic Aromatic Hydrocarbon Growth1. *J. Am. Chem. Soc.* **2000**, *122* (7), 1548-1549. [DOI: 10.1021/ja992115c]

³⁶² Zhao, L.; Doddipatla, S.; Kaiser, R. I.; Lu, W.; Kostko, O.; Ahmed, M.; Tuli, L. B.; Morozov, A. N.; Howlader, A. H.; Wnuk, S. F.; Mebel, A. M.; Azyazov, V. N.; Mohamed, R. K.; Fischer, F. R., Gas-phase synthesis of corannulene - a molecular building block of fullerenes. *Phys. Chem. Chem. Phys.* 2021, 23 (10), 5740-5749. [DOI: 10.1039/D0CP06537D]

³⁶³ Krestinin, A. V.; Kislov, M. B.; Raevskii, A. V.; Kolesova, O. I.; Stesik, L. N. On the mechanism of soot particle formation. *Kinet. Catal.* **2000**, *41*, 90. [DOI: 10.1007/BF02756146]

³⁶⁴ Krestinin, A. V. Polyyne model of soot formation process. *Symposium (International) on Combustion* **1998**, 27, 1557. [DOI: 10.1016/S0082-0784(98)80564-X]

³⁶⁵ Krestinin, A.V. Sov. J. Chem. Phys., **1990**, 6, 618.

³⁶⁶ Krestinin, A.V. Chem. Phys. Reports, **1994**, 13, 191.

³⁶⁷ McCarthy, M. C.; Chen, W.; Travers, M. J.; Thaddeus, P. Microwave Spectra of 11 Polyyne Carbon Chains. *Astrophys. J., Suppl. Ser.* **2000**, *129*, 611. [DOI: 10.1086/313428]

³⁶⁸ Cernicharo, J.; Guélin, M.; Kahane, C. A λ 2 mm molecular line survey of the C-star envelope. *Astron. Astrophys., Suppl. Ser.* **2000**, *142*, 181. [DOI: 10.1051/aas:2000147]

³⁶⁹ Bonne, U.; Homann, K. H.; Wagner, H. G. Carbon formation in premixed flames. *Symposium* (*International*) on Combustion **1965**, *10*, 503. [DOI: 10.1016/S0082-0784(65)80197-7]

³⁷⁰ Krestinin, A. V. Detailed modeling of soot formation in hydrocarbon pyrolysis. *Combust. Flame* **2000**, *121*, 513. [DOI: 10.1016/S0010-2180(99)00167-4]

³⁷¹ Zhao, L.; Prendergast, M.; Kaiser, R. I.; Xu, B.; Ablikim, U.; Lu, W.; Ahmed, M.; Oleinikov, A. D.; Azyazov, V. N.; Howlader, A. H.; Wnuk, S. F.; Mebel, A. M., How to add a five-membered ring to polycyclic aromatic hydrocarbons (PAHs) - molecular mass growth of the 2-naphthyl radical (C₁₀H₇) to benzindenes (C₁₃H₁₀) as a case study. *Phys. Chem. Chem. Phys.* **2019**, *21* (30), 16737-16750. [DOI: 10.1039/C9CP02930C]

³⁷² Mebel, A. M.; Kislov, V. V.; Kaiser, R. I. Photoinduced mechanism of formation and growth of polycyclic aromatic hydrocarbons in low-temperature environments via successive ethynyl radical additions. *J. Am. Chem. Soc.* **2008**, *130*, 13618. [DOI: 10.1021/ja804198a]

³⁷³ Kaiser, R. I.; Parker, D. S.; Mebel, A. M., Reaction dynamics in astrochemistry: low-temperature pathways to polycyclic aromatic hydrocarbons in the interstellar medium. *Annu. Rev. Phys. Chem.* **2015**, *66*, 43-67. [DOI: 10.1146/annurev-physchem-040214-121502]

³⁷⁴ Wilson, G. R.; Ober, D. M.; Germany, G. A.; Lund, E. J. Nightside auroral zone and polar cap ion outflow as a function of substorm size and phase. *J. Geophys. Res.: Space Phys.* **2004**, *109*. [DOI: 10.1029/2003JA009835]

³⁷⁵ Imanaka, H.; Khare, B. N.; Elsila, J. E.; Bakes, E. L. O.; McKay, C. P.; Cruikshank, D. P.; Sugita, S.; Matsui, T.; Zare, R. N. Laboratory experiments of Titan tholin formed in cold plasma at various pressures: implications for nitrogen-containing polycyclic aromatic compounds in Titan haze. *Icarus* **2004**, *168*, 344. [DOI: 10.1016/j.icarus.2003.12.014]

³⁷⁶ Cottini, V.; Nixon, C. A.; Jennings, D. E.; de Kok, R.; Teanby, N. A.; Irwin, P. G. J.; Flasar, F. M., Spatial and temporal variations in Titan's surface temperatures from Cassini CIRS observations. *Planet. Space Sci.* **2012**, *60* (1), 62-71. [DOI: 10.1016/j.pss.2011.03.015]

³⁷⁷ Wilson, E. H.; Atreya, S. K. Chemical sources of haze formation in Titan's atmosphere. *Planet. Space Sci.* **2003**, *51*, 1017. [DOI: 10.1016/j.pss.2003.06.003]

³⁷⁸ Zhao, L.; Kaiser, R. I.; Xu, B.; Ablikim, U.; Ahmed, M.; Evseev, M. M.; Bashkirov, E. K.; Azyazov, V. N.; Mebel, A. M. Low-temperature formation of polycyclic aromatic hydrocarbons in Titan's atmosphere. *Nature Astronomy* **2018**, *2*, 973. [DOI: 10.1038/s41550-018-0585-y]

³⁷⁹ Fulchignoni, M.; Ferri, F.; Angrilli, F.; Ball, A. J.; Bar-Nun, A.; Barucci, M. A.; Bettanini, C.; Bianchini, G.; Borucki, W.; Colombatti, G.; Coradini, M.; Coustenis, A.; Debei, S.; Falkner, P.; Fanti, G.; Flamini, E.; Gaborit, V.; Grard, R.; Hamelin, M.; Harri, A. M.; Hathi, B.; Jernej, I.; Leese, M. R.; Lehto, A.; Lion Stoppato, P. F.; Lopez-Moreno, J. J.; Makinen, T.; McDonnell, J. A.; McKay, C. P.; Molina-Cuberos, G.; Neubauer, F. M.; Pirronello, V.; Rodrigo, R.; Saggin, B.;

Schwingenschuh, K.; Seiff, A.; Simoes, F.; Svedhem, H.; Tokano, T.; Towner, M. C.; Trautner, R.; Withers, P.; Zarnecki, J. C. In situ measurements of the physical characteristics of Titan's environment. *Nature* **2005**, *438*, 785. [DOI: 10.1038/nature04314]

³⁸⁰ Lorenz, R. D.; Wall, S.; Radebaugh, J.; Boubin, G.; Reffet, E.; Janssen, M.; Stofan, E.; Lopes, R.; Kirk, R.; Elachi, C.; Lunine, J.; Mitchell, K.; Paganelli, F.; Soderblom, L.; Wood, C.; Wye, L.; Zebker, H.; Anderson, Y.; Ostro, S.; Allison, M.; Boehmer, R.; Callahan, P.; Encrenaz, P.; Ori, G. G.; Francescetti, G.; Gim, Y.; Hamilton, G.; Hensley, S.; Johnson, W.; Kelleher, K.; Muhleman, D.; Picardi, G.; Posa, F.; Roth, L.; Seu, R.; Shaffer, S.; Stiles, B.; Vetrella, S.; Flamini, E.; West, R. The sand seas of Titan: Cassini RADAR observations of longitudinal dunes. *Science* 2006, *312*, 724. [DOI: 10.1126/science.1123257]

³⁸¹ Lorenz, R. D.; Mitchell, K. L.; Kirk, R. L.; Hayes, A. G.; Aharonson, O.; Zebker, H. A.; Paillou, P.; Radebaugh, J.; Lunine, J. I.; Janssen, M. A.; Wall, S. D.; Lopes, R. M.; Stiles, B.; Ostro, S.; Mitri, G.; Stofan, E. R. Titan's inventory of organic surface materials. *Geophys. Res. Lett.* **2008**, *35*. [DOI: 10.1029/2007GL032118]

³⁸² Rodriguez, S.; Garcia, A.; Lucas, A.; Appéré, T.; Le Gall, A.; Reffet, E.; Le Corre, L.; Le Mouélic, S.; Cornet, T.; Courrech du Pont, S.; Narteau, C.; Bourgeois, O.; Radebaugh, J.; Arnold, K.; Barnes, J. W.; Stephan, K.; Jaumann, R.; Sotin, C.; Brown, R. H.; Lorenz, R. D.; Turtle, E. P. Global mapping and characterization of Titan's dune fields with Cassini: Correlation between RADAR and VIMS observations. *Icarus* **2014**, *230*, 168. [DOI: 10.1016/j.icarus.2013.11.017]

³⁸³ Hayes, A. G.; Lorenz, R. D.; Lunine, J. I. A post-Cassini view of Titan's methane-based hydrologic cycle. *Nat. Geosci.* **2018**, *11*, 306. [DOI: 10.1038/s41561-018-0103-y]

³⁸⁴ Lunine, J. I.; Lorenz, R. D. Rivers, Lakes, Dunes, and Rain: Crustal Processes in Titan's Methane Cycle. *Annu. Rev. Earth Planet. Sci.* **2009**, *37*, 299. [DOI: 10.1146/annurev.earth.031208.100142]

³⁸⁵ Vuitton, V.; Yelle, R. V.; Klippenstein, S. J.; Hörst, S. M.; Lavvas, P. *Icarus* **2019**, *324*, 120. [DOI: 10.1016/j.icarus.2018.06.013]

³⁸⁶ Lucas, A.; Rodriguez, S.; Lemonnier, F.; Le Gall, A.; MacKenzie, S.; Ferrari, C.; Paillou, P.; Narteau, C., Texture and Composition of Titan's Equatorial Sand Seas Inferred From Cassini SAR Data: Implications for Aeolian Transport and Dune Morphodynamics. *J. Geophys. Res.: Planets* **2019**, *124* (11), 3140-3163. [DOI: 10.1029/2019JE005965]

³⁸⁷ Lunine, J. I.; Hörst, S. M. Organic chemistry on the surface of Titan. *Rendiconti Lincei* **2011**, 22, 183. [DOI: 10.1007/s12210-011-0130-8]

³⁸⁸ McCord, T. B.; Hansen, G. B.; Clark, R. N.; Martin, P. D.; Hibbitts, C. A.; Fanale, F. P.; Granahan, J. C.; Segura, M.; Matson, D. L.; Johnson, T. V.; Carlson, R. W.; Smythe, W. D.; Danielson, G. E. Non-water-ice constituents in the surface material of the icy Galilean satellites from the Galileo near-infrared mapping spectrometer investigation. *J. Geophys. Res.: Planets* **1998**, *103*, 8603. [DOI: 10.1029/98JE00788]

³⁸⁹ Sagan, C.; Reid Thompson, W. Production and condensation of organic gases in the atmosphere of Titan. *Icarus* **1984**, *59*, 133. [DOI: 10.1016/0019-1035(84)90018-6]

³⁹⁰ Molina-Cuberos, G. J.; López-Moreno, J. J.; Rodrigo, R.; Lara, L. M.; O'Brien, K. Ionization by cosmic rays of the atmosphere of Titan. *Planet. Space Sci.* **1999**, *47*, 1347. [DOI: 10.1016/S0032-0633(99)00056-2]

³⁹¹ Kaiser, R. I.; Roessler, K. Theoretical and Laboratory Studies on the Interaction of Cosmic-Ray Particles with Interstellar Ices. I. Synthesis of Polycyclic Aromatic Hydrocarbons by a Cosmic-Ray–induced Multicenter Mechanism. *Astrophys. J.* **1997**, *475*, 144. [DOI: 10.1086/303506] ³⁹² Lukianova, M. A.; Feldman, V. I., Radiation-induced closure of the second aromatic ring: Possible way to PAH starting from a styrene-acetylene complex. *Radiat. Phys. Chem.* **2022**, *191*. [DOI: 10.1016/j.radphyschem.2021.109847]

³⁹³ Protopapa, S.; Olkin, C. B.; Grundy, W. M.; Li, J.-Y.; Verbiscer, A.; Cruikshank, D. P.; Gautier, T.; Quirico, E.; Cook, J. C.; Reuter, D.; Howett, C. J. A.; Stern, A.; Beyer, R. A.; Porter, S.; Young, L. A.; Weaver, H. A.; Ennico, K.; Dalle Ore, C. M.; Scipioni, F.; Singer, K., Disk-resolved Photometric Properties of Pluto and the Coloring Materials across its Surface. *Astron. J.* **2020**, *159* (2). [DOI: 10.3847/1538-3881/ab5e82]

³⁹⁴ Brown, M. E.; Schaller, E. L.; Blake, G. A. Irradiation Products on Dwarf Planet Makemake. *Astron. J.* **2015**, *149*, 105. [DOI: 10.1088/0004-6256/149/3/105]

³⁹⁵ Combe, J.-P.; Singh, S.; Johnson, K. E.; McCord, T. B.; De Sanctis, M. C.; Ammannito, E.; Carrozzo, F. G.; Ciarniello, M.; Frigeri, A.; Raponi, A.; Tosi, F.; Zambon, F.; Scully, J. E. C.; Raymond, C. A.; Russell, C. T. The surface composition of Ceres' Ezinu quadrangle analyzed by the Dawn mission. *Icarus* **2019**, *318*, 124. [DOI: 10.1016/j.icarus.2017.12.039]

³⁹⁶ Barucci, M. A.; Merlin, F.; Guilbert, A.; de Bergh, C.; Alvarez-Candal, A.; Hainaut, O.; Doressoundiram, A.; Dumas, C.; Owen, T.; Coradini, A., Surface composition and temperature of the TNO Orcus. *Astron. Astrophys.* **2008**, *479* (1), L13-L16. [DOI: 10.1051/0004-6361:20079079]

³⁹⁷ Brown, M. E., The Compositions of Kuiper Belt Objects. *Annu. Rev. Earth Planet. Sci.* **2012**, *40* (1), 467-494. [DOI: 10.1146/annurev-earth-042711-105352]

³⁹⁸ Brown, R. H.; Baines, K. H.; Bellucci, G.; Buratti, B. J.; Capaccioni, F.; Cerroni, P.; Clark, R. N.; Coradini, A.; Cruikshank, D. P.; Drossart, P.; Formisano, V.; Jaumann, R.; Langevin, Y.; Matson, D. L.; McCord, T. B.; Mennella, V.; Nelson, R. M.; Nicholson, P. D.; Sicardy, B.; Sotin, C.; Baugh, N.; Griffith, C. A.; Hansen, G. B.; Hibbitts, C. A.; Momary, T. W.; Showalter, M. R. Observations in the Saturn system during approach and orbital insertion, with Cassini's visual and infrared mapping spectrometer (VIMS). *Astron. Astrophys.* **2006**, *446*, 707. [DOI: 10.1051/0004-6361:20053054]

³⁹⁹ Buratti, B. J.; Soderlund, K.; Bauer, J.; Mosher, J. A.; Hicks, M. D.; Simonelli, D. P.; Jaumann, R.; Clark, R. N.; Brown, R. H.; Cruikshank, D. P.; Momary, T. Infrared (0.83–5.1 μm) photometry of Phoebe from the Cassini Visual Infrared Mapping Spectrometer. *Icarus* **2008**, *193*, 309. [DOI: 10.1016/j.icarus.2007.09.014]

⁴⁰⁰ Cruikshank, D. P.; Dalle Ore, C. M.; Clark, R. N.; Pendleton, Y. J. Aromatic and aliphatic organic materials on Iapetus: Analysis of Cassini VIMS data. *Icarus* **2014**, *233*, 306. [DOI: 10.1016/j.icarus.2014.02.011]

⁴⁰¹ Dalle Ore, C. M.; Protopapa, S.; Cook, J. C.; Grundy, W. M.; Cruikshank, D. P.; Verbiscer, A. J.; Ennico, K.; Olkin, C. B.; Stern, S. A.; Weaver, H. A.; Young, L. A. Ices on Charon: Distribution of H2O and NH3 from New Horizons LEISA observations. *Icarus* **2018**, *300*, 21. [DOI: 10.1016/j.icarus.2017.08.026]

⁴⁰² Protopapa, S.; Grundy, W. M.; Reuter, D. C.; Hamilton, D. P.; Dalle Ore, C. M.; Cook, J. C.; Cruikshank, D. P.; Schmitt, B.; Philippe, S.; Quirico, E.; Binzel, R. P.; Earle, A. M.; Ennico, K.; Howett, C. J. A.; Lunsford, A. W.; Olkin, C. B.; Parker, A.; Singer, K. N.; Stern, A.; Verbiscer, A. J.; Weaver, H. A.; Young, L. A. Pluto's global surface composition through pixel-by-pixel Hapke modeling of New Horizons Ralph/LEISA data. *Icarus* **2017**, *287*, 218. [DOI: 10.1016/j.icarus.2016.11.028]

⁴⁰³ Abplanalp, M. J.; Kaiser, R. I., Implications for Extraterrestrial Hydrocarbon Chemistry: Analysis of Acetylene (C_2H_2) and D2-acetylene (C_2D_2) Ices Exposed to Ionizing Radiation via Ultraviolet–Visible Spectroscopy, Infrared Spectroscopy, and Reflectron Time-of-flight Mass Spectrometry. *Astrophys. J.* **2020**, 889 (1). [DOI: 10.3847/1538-4357/ab616c] ⁴⁰⁴ Zhang, C.; Zhu, C.; Turner, A. M.; Antonov, I. O.; Garcia, A. D.; Meinert, C.; Young, L. A.; Jewitt, D. C.; Kaiser, R. I., Processing of methane and acetylene ices by galactic cosmic rays and implications to the color diversity of Kuiper Belt objects. *Sci. Adv.* **2023**, *9* (22), eadg6936. [DOI: 10.1126/sciadv.adg6936]

⁴⁰⁵ Dalle Ore, C. M.; Barucci, M. A.; Emery, J. P.; Cruikshank, D. P.; Dalle Ore, L. V.; Merlin, F.; Alvarez-Candal, A.; de Bergh, C.; Trilling, D. E.; Perna, D.; Fornasier, S.; Mastrapa, R. M. E.; Dotto, E. *Astron. Astrophys.* **2009**, *501*, 349. [DOI: 10.1051/0004-6361/200911752]

⁴⁰⁶ Krätschmer, W.; Lamb, L. D.; Fostiropoulos, K.; Huffman, D. R., Solid C60: a new form of carbon. *Nature* **1990**, *347* (6291), 354-358. [DOI: 10.1038/347354a0]

⁴⁰⁷ Smalley, R. E., Self-assembly of the fullerenes. *Acc. Chem. Res.* **2002**, *25* (3), 98-105. [DOI: 10.1021/ar00015a001],

⁴⁰⁸ Dunk, P. W.; Kaiser, N. K.; Hendrickson, C. L.; Quinn, J. P.; Ewels, C. P.; Nakanishi, Y.; Sasaki, Y.; Shinohara, H.; Marshall, A. G.; Kroto, H. W., Closed network growth of fullerenes. *Nat Commun* **2012**, *3*, 855. [DOI: 10.1038/ncomms1853]

⁴⁰⁹ Micelotta, E. R.; Jones, A. P.; Cami, J.; Peeters, E.; Bernard-Salas, J.; Fanchini, G., The Formation of Cosmic Fullerenes from Arophatic Clusters. *Astrophys. J.* **2012**, *761* (1). [DOI: 10.1088/0004-637X/761/1/35]

⁴¹⁰ Berne, O.; Tielens, A. G., Formation of buckminsterfullerene (C60) in interstellar space. *Proc Natl Acad Sci U S A* **2012**, *109* (2), 401-6. [DOI: 10.1073/pnas.1114207108]

⁴¹¹ Otero, G.; Biddau, G.; Sanchez-Sanchez, C.; Caillard, R.; Lopez, M. F.; Rogero, C.; Palomares, F. J.; Cabello, N.; Basanta, M. A.; Ortega, J.; Mendez, J.; Echavarren, A. M.; Perez, R.; Gomez-Lor, B.; Martin-Gago, J. A., Fullerenes from aromatic precursors by surface-catalysed cyclodehydrogenation. *Nature* **2008**, *454* (7206), 865-8. [DOI: 10.1038/nature07193]

⁴¹² Chuvilin, A.; Kaiser, U.; Bichoutskaia, E.; Besley, N. A.; Khlobystov, A. N., Direct transformation of graphene to fullerene. *Nat. Chem.* **2010**, *2* (6), 450-3. [DOI: 10.1038/nchem.644]

⁴¹³ Berné, O.; Montillaud, J.; Joblin, C., Top-down formation of fullerenes in the interstellar medium (Corrigendum). *Astron. Astrophys.* **2016**, *588*. [DOI: 10.1051/0004-6361/201425338e]

⁴¹⁴ Tuli, L. B.; Goettl, S. J.; Turner, A. M.; Howlader, A. H.; Hemberger, P.; Wnuk, S. F.; Guo, T.; Mebel, A. M.; Kaiser, R. I., Gas phase synthesis of the C40 nano bowl C(40)H(10). *Nat Commun* **2023**, *14* (1), 1527. [DOI: 10.1038/s41467-023-37058-y]

⁴¹⁵ Shaitanov, A. G.; Surovikin, Y. V.; Rezanov, I. V.; Shtertser, A. A.; Ul'yanitskii, V. Y.; Vasil'ev, A. A.; Likholobov, V. A., Synthesis and Study of Nanodispersed Carbon in the Combustion of Acetylene in a Flow Detonation Tube. *Russ. J. Appl. Chem.* **2019**, *91* (12), 2003-2011. [DOI: 10.1134/S1070427218120121]

⁴¹⁶ Martin, J. W.; Salamanca, M.; Kraft, M., Soot inception: Carbonaceous nanoparticle formation in flames. *Prog. Energy Combust. Sci.* **2022**, 88. [DOI: 10.1016/j.pecs.2021.100956]

⁴¹⁷ Gail HP., Sedlmayr E. Dust Formation in Stellar Winds. In: Morfill G.E., Scholer M. (eds) Physical Processes in Interstellar Clouds. NATO ASI Series (Series C: Mathematical and Physical Sciences), vol 210. Springer, Dordrecht, 1987, pp. 275-303 [DOI: 10.1007/978-94-009-3945-5_19]

⁴¹⁸ Mironenko, R. M.; Likholobov, V. A.; Belskaya, O. B., Nanoglobular carbon and palladium - carbon catalysts for liquid-phase hydrogenation of organic compounds. *Russ. Chem. Rev.* **2022**, *91 (1)*, RCR5017. [DOI: 10.1070/RCR5017]
⁴¹⁹ Mao, Q.; van Duin, A. C. T.; Luo, K. H., Formation of incipient soot particles from polycyclic aromatic hydrocarbons: A ReaxFF molecular dynamics study. *Carbon* **2017**, *121*, 380-388. [DOI: 10.1016/j.carbon.2017.06.009]

⁴²⁰ Martinez, L.; Lauwaet, K.; Santoro, G.; Sobrado, J. M.; Pelaez, R. J.; Herrero, V. J.; Tanarro, I.; Ellis, G. J.; Cernicharo, J.; Joblin, C.; Huttel, Y.; Martin-Gago, J. A., Precisely controlled fabrication, manipulation and in-situ analysis of Cu based nanoparticles. *Sci. Rep.* **2018**, *8* (1), 7250. [DOI: 10.1038/s41598-018-25472-y]

⁴²¹ Martinez, L.; Santoro, G.; Merino, P.; Accolla, M.; Lauwaet, K.; Sobrado, J.; Sabbah, H.; Pelaez, R. J.; Herrero, V. J.; Tanarro, I.; Agundez, M.; Martin-Jimenez, A.; Otero, R.; Ellis, G. J.; Joblin, C.; Cernicharo, J.; Martin-Gago, J. A., Prevalence of non-aromatic carbonaceous molecules in the inner regions of circumstellar envelopes. *Nat Astron* **2020**, *4* (1), 97-105. [DOI: 10.1038/s41550-019-0899-4]

⁴²² Cuppen, H. M.; Herbst, E., Simulation of the Formation and Morphology of Ice Mantles on Interstellar Grains. *Astrophys. J.* **2007**, *668* (1), 294-309. [DOI: 10.1086/521014]

⁴²³ Ehrenfreund, P. & Fraser, H. Ice chemistry in space. In Solid state astrochemistry, NATO ASI series (eds V. Pirronello, J. Krelowski & G. Manicy). Dordrecht, The Netherlands: Kluwer Academic Publishers, 2003, pp. 317–356. [DOI: 10.1007/978-94-010-0062-8_10]

⁴²⁴ Cuppen, H. M.; Walsh, C.; Lamberts, T.; Semenov, D.; Garrod, R. T.; Penteado, E. M.; Ioppolo, S., Grain Surface Models and Data for Astrochemistry. *Space Sci. Rev.* **2017**, *212* (1-2), 1-58. [DOI: 10.1007/s11214-016-0319-3]

⁴²⁵ Saladino, R.; Crestini, C.; Neri, V.; Brucato, J. R.; Colangeli, L.; Ciciriello, F.; Di Mauro, E.; Costanzo, G., Synthesis and degradation of nucleic Acid components by formamide and cosmic dust analogues. *ChemBioChem* **2005**, *6* (8), 1368-74. [DOI: 10.1002/cbic.200500035]

⁴²⁶ Potapov, A.; McCoustra, M., Physics and chemistry on the surface of cosmic dust grains: a laboratory view. *Int. Rev. Phys. Chem.* **2021,** *40* (2), 299-364. [DOI: 10.1080/0144235X.2021.1918498]

⁴²⁷ Testa, C.; Zammataro, A.; Pappalardo, A.; Trusso Sfrazzetto, G., Catalysis with carbon nanoparticles. *RSC Adv.* **2019**, *9* (47), 27659-27664. [DOI: 10.1039/c9ra05689k]

⁴²⁸ Pentsak, E. O.; Gordeev, E. G.; Ananikov, V. P., Carbocatalysis: From Acetylene Trimerization to Modern Organic Synthesis. A Review. *Dokl. Phys. Chem.* 2020, 493 (2), 95-122. [DOI: 10.1134/s0012501620380017]

⁴²⁹ Suhasaria, T.; Mennella, V., Catalytic Role of Refractory Interstellar Grain Analogs on H2 Formation. *Front. Astron. Space Sci.* **2021**, *8*. [DOI: 10.3389/fspas.2021.655883]

⁴³⁰ Saladino, R.; Brucato, J. R.; De Sio, A.; Botta, G.; Pace, E.; Gambicorti, L., Photochemical synthesis of citric acid cycle intermediates based on titanium dioxide. *Astrobiology* **2011**, *11* (8), 815-24. [DOI: 10.1089/ast.2011.0652]

⁴³¹ Pareras, G.; Cabedo, V.; McCoustra, M.; Rimola, A., Single-atom catalysis in space: Computational exploration of Fischer–Tropsch reactions in astrophysical environments. *Astron. Astrophys.* **2023**, *680*. [DOI: 10.1051/0004-6361/202347877]

⁴³² Fioroni, M., Astrochemistry of transition metals? The selected cases of [FeN]⁺, [FeNH]⁺ and [(CO)₂FeN]⁺: pathways toward CH₃NH₂ and HNCO. *Phys. Chem. Chem. Phys.* **2014**, *16* (44), 24312-22. [DOI: 10.1039/c4cp03218g]

⁴³³ Reppe W, Schweckendiek WJ. Cyclisierende Polymerisation von Acetylen. III Benzol, Benzolderivate und hydroaromatische Verbindungen. *Justus Liebigs Ann. Chem.* **1948**, *560* (1), 104-116. [DOI: 10.1002/jlac.19485600104]

⁴³⁴ Galan, B. R.; Rovis, T., Beyond Reppe: building substituted arenes by [2+2+2] cycloadditions of alkynes. *Angew. Chem. Int. Ed. Engl.* **2009**, *48* (16), 2830-4. [DOI: 10.1002/anie.200804651]

⁴³⁵ Roglans, A.; Pla-Quintana, A.; Sola, M., Mechanistic Studies of Transition-Metal-Catalyzed
[2+2+2] Cycloaddition Reactions. *Chem. Rev.* 2021, *121* (3), 1894-1979.
[DOI: 10.1021/acs.chemrev.0c00062]

⁴³⁶ Takahashi, T.; Xi, Z.; Yamazaki, A.; Liu, Y.; Nakajima, K.; Kotora, M., Cycloaddition Reaction of Zirconacyclopentadienes to Alkynes: Highly Selective Formation of Benzene Derivatives from Three Different Alkynes. *J. Am. Chem. Soc.* **1998**, *120* (8), 1672-1680. [DOI: 10.1021/ja970869q]

⁴³⁷ Tanaka, R.; Yuza, A.; Watai, Y.; Suzuki, D.; Takayama, Y.; Sato, F.; Urabe, H., One-pot synthesis of metalated pyridines from two acetylenes, a nitrile, and a titanium(II) alkoxide. *J. Am. Chem. Soc.* **2005**, *127* (21), 7774-80. [DOI: 10.1021/ja050261e]

⁴³⁸ Yamamoto, Y.; Ishii, J.; Nishiyama, H.; Itoh, K., Cp(*)RuCl-catalyzed formal intermolecular cyclotrimerization of three unsymmetrical alkynes through a boron temporary tether: regioselective four-component coupling synthesis of phthalides. *J. Am. Chem. Soc.* **2005**, *127* (26), 9625-31. [DOI: 10.1021/ja051377d]

⁴³⁹ Hilt, G.; Hengst, C.; Hess, W., Solvent-Dependent Regiochemical Cyclotrimerisation of Phenylacetylene with Cobalt Catalysts Containing Disulfide Ligands: A Case Study. *Eur. J. Org. Chem.* **2008**, *2008* (13), 2293-2297. [DOI: 10.1002/ejoc.200800106]

⁴⁴⁰ Tanaka, K.; Shirasaka, K., Highly chemo- and regioselective intermolecular cyclotrimerization of alkynes catalyzed by cationic rhodium(I)/modified BINAP complexes. *Org. Lett.* **2003**, *5* (24), 4697-9. [DOI: 10.1021/ol035963s]

⁴⁴¹ Geetharani, K.; Tussupbayev, S.; Borowka, J.; Holthausen, M. C.; Ghosh, S., A mechanistic study of the utilization of arachno-diruthenaborane [(Cp*RuCO)2B2H6] as an active alkyne-cyclotrimerization catalyst. *Chem. - Eur. J.* **2012**, *18* (27), 8482-9. [DOI: 10.1002/chem.201200291]

⁴⁴² Rassadin, V. A.; Nicolas, E.; Six, Y., Ti(OiPr)4/nBuLi: an attractive reagent system for [2+2+2] cyclotrimerisation reactions. *Chem. Commun. (Camb)* **2014,** *50* (57), 7666-9. [DOI: 10.1039/C4CC02698E]

⁴⁴³ Orsino, A. F.; Gutierrez Del Campo, M.; Lutz, M.; Moret, M. E., Enhanced Catalytic Activity of Nickel Complexes of an Adaptive Diphosphine-Benzophenone Ligand in Alkyne Cyclotrimerization. *ACS Catal.* **2019**, *9* (3), 2458-2481. [DOI: 10.1021/acscatal.8b05025]

⁴⁴⁴ Tsuji, H.; Yamagata, K.; Fujimoto, T.; Nakamura, E., Manganese-catalyzed benzene synthesis by [2+2+2] coupling of 1,3-dicarbonyl compound and terminal acetylene. *J. Am. Chem. Soc.* **2008**, *130* (25), 7792-3. [DOI: 10.1021/ja8015186]

⁴⁴⁵ Kuninobu, Y.; Takata, H.; Kawata, A.; Takai, K., Rhenium-catalyzed synthesis of multisubstituted aromatic compounds via C-C single-bond cleavage. *Org. Lett.* **2008**, *10* (14), 3133-5. [DOI: 10.1021/ol801226t]

⁴⁴⁶ Heller, B.; Sundermann, B.; Buschmann, H.; Drexler, H. J.; You, J.; Holzgrabe, U.; Heller, E.; Oehme, G., Photocatalyzed [2 + 2 + 2]-cycloaddition of nitriles with acetylene: an effective method for the synthesis of 2-pyridines under mild conditions. *J. Org. Chem.* **2002**, *67* (13), 4414-22. [DOI: 10.1021/jo011032n]

⁴⁴⁷ More, A. A.; Ramana, C. V., Total synthesis of the putative structure of xylarinol B. *Chem. Asian. J.* **2014**, *9* (6), 1557-62. [DOI: 10.1002/asia.201301647]

⁴⁴⁸ Tielens, A. G.; Wooden, D. H.; Allamandola, L. J.; Bregman, J.; Witteborn, F. C., The infrared spectrum of the Galactic center and the composition of interstellar dust. *Astrophys. J.* **1996**, *461* (1 Pt 1), 210-22. [DOI: 10.1086/177049]

⁴⁴⁹ Xu, Y., Xiao, X., Sun, S., & Ouyang, Z., IR Spectroscopic Evidence of Metal Carbonyl Clusters in the Jiange H5 Chondrite. *Lunar Planet. Sci.* **1996**, *27*, 1457. [https://ui.adsabs.harvard.edu/abs/1996LPI....27.1457X/abstract]

⁴⁵⁰ Doiron, J.; Soultan, A. H.; Richard, R.; Toure, M. M.; Picot, N.; Richard, R.; Cuperlovic-Culf, M.; Robichaud, G. A.; Touaibia, M., Synthesis and structure-activity relationship of 1- and 2-substituted-1,2,3-triazole letrozole-based analogues as aromatase inhibitors. *Eur. J. Med. Chem.* **2011**, *46* (9), 4010-24. [DOI: 10.1016/j.ejmech.2011.05.074]

⁴⁵¹ Scharnagel, D.; Escofet, I.; Armengol-Relats, H.; de Orbe, M. E.; Korber, J. N.; Echavarren, A. M., Acetylene as a Dicarbene Equivalent for Gold(I) Catalysis: Total Synthesis of Waitziacuminone in One Step. *Angew. Chem. Int. Ed. Engl.* **2020**, *59* (12), 4888-4891. [DOI: 10.1002/anie.201915895]

⁴⁵² Pechersky, D. M., Metallic Iron and Nickel in Cretaceous and Cenozoic Sediments: The Results of Thermomagnetic Analysis. *J. Environ. Prot.* 2010, 01 (02), 143-154.
[DOI: 10.4236/jep.2010.12019]

⁴⁵³ Speck, A. K.; Corman, A. B.; Wakeman, K.; Wheeler, C. H.; Thompson, G., Silicon Carbide Absorption Features: Dust Formation in the Outflows of Extreme Carbon Stars. *Astrophys. J.* **2009**, *691* (2), 1202-1221. [DOI: 10.1088/0004-637X/691/2/1202]

⁴⁵⁴ Fujiyoshi, T.; Wright, C. M.; Moore, T. J. T. Mid-infrared spectroscopy of SVS13: silicates, quartz and SiC in a protoplanetary disc. *Mon. Not. R. Astron. Soc.* **2015**, *451*, 3371. [DOI: 10.1093/mnras/stv1171]

⁴⁵⁵ von Helden, G.; Tielens, A. G.; van Heijnsbergen, D.; Duncan, M. A.; Hony, S.; Waters, L. B.; Meijer, G., Titanium carbide nanocrystals in circumstellar environments. *Science* **2000**, 288 (5464), 313-6. [DOI: 10.1126/science.288.5464.313].

⁴⁵⁶ Savage, B. D.; Sembach, K. R., Interstellar Abundances from Absorption-Line Observations with the Hubble Space Telescope. *Annu. Rev. Astron. Astrophys.* **1996**, *34* (1), 279-329. [DOI: 10.1146/annurev.astro.34.1.279]

⁴⁵⁷ Zhao, T. Q.; Li, Q.; Liu, B. S.; Gover, R. K.; Sarre, P. J.; Cheung, A. S. Laboratory astrochemistry: catalytic conversion of acetylene to polycyclic aromatic hydrocarbons over SiC grains. *Phys. Chem. Chem. Phys.* **2016**, *18*, 3489. [DOI: 10.1039/C5CP06425B]

⁴⁵⁸ Speck, A. K.; Thompson, G. D.; Hofmeister, A. M. The Effect of Stellar Evolution on SiC Dust Grain Sizes. *Astrophys. J.* **2005**, *634*, 426. [DOI: 10.1086/496955]

⁴⁵⁹ Yang, T.; Bertels, L.; Dangi, B. B.; Li, X.; Head-Gordon, M.; Kaiser, R. I., Gas phase formation of c-SiC(3) molecules in the circumstellar envelope of carbon stars. *Proc. Natl. Acad. Sci. USA* **2019**, *116* (29), 14471-14478. [DOI: 10.1073/pnas.1810370116]

⁴⁶⁰ Parker, D. S. N.; Wilson, A. V.; Kaiser, R. I.; Mayhall, N. J.; Head-Gordon, M.; Tielens, A. G. G. M., On the formation of silacyclopropenylidene (c-SiC2H2) and its role in the organosilicon chemistry in the interstellar medium. *Astrophys. J.* **2013**, *770* (1). [DOI: 10.1088/0004-637X/770/1/33]

⁴⁶¹ Fioroni, M.; DeYonker, N. J., Complex Organic Matter Synthesis on Siloxyl Radicals in the Presence of CO. *Front. Chem.* **2020**, *8*, 621898. [DOI: 10.3389/fchem.2020.621898]

⁴⁶² Serra, G. C., B.; Saillard, Y.; Le Beuze, A.; Rabaa, H.; Ristorcelli, I.; Klotz, A., Organometallic chemistry in the interstellar medium. I - Are organometallic reactions efficient processes in

astrochemistry? *Astron. Astrophys.* **1992,** 260 (1-2), 489-493. [https://ui.adsabs.harvard.edu/abs/1992A%26A...260..489S/abstract]

⁴⁶³ Murakami, T.; Takayanagi, T., Interstellar Benzene Formation Mechanisms via Acetylene Cyclotrimerization Catalyzed by Fe(+) Attached to Water Ice Clusters: Quantum Chemistry Calculation Study. *Molecules* **2022**, *27* (22). [DOI: 10.3390/molecules27227767]

⁴⁶⁴ Murakami, T.; Matsumoto, N.; Fujihara, T.; Takayanagi, T., Possible Roles of Transition Metal Cations in the Formation of Interstellar Benzene via Catalytic Acetylene Cyclotrimerization. *Molecules* **2023**, 28 (21). [DOI: 10.3390/molecules28217454]

⁴⁶⁵ Wakelam, V.; Bron, E.; Cazaux, S.; Dulieu, F.; Gry, C.; Guillard, P.; Habart, E.; Hornekær, L.; Morisset, S.; Nyman, G.; Pirronello, V.; Price, S. D.; Valdivia, V.; Vidali, G.; Watanabe, N., H 2 formation on interstellar dust grains: The viewpoints of theory, experiments, models and observations. *Mol. Astrophys.* **2017**, *9*, 1-36. [DOI: 10.1016/j.molap.2017.11.001]

⁴⁶⁶ Habart, E.; Boulanger, F.; Verstraete, L.; Forêts, G. P. d.; Falgarone, E.; Abergel, A., H2infrared line emission across the bright side of theρOphiuchi main cloud. *Astron. Astrophys.* **2002**, *397* (2), 623-634. [DOI: 10.1051/0004-6361:20021489]

⁴⁶⁷ Rasmussen, A. P.; Wenzel, G.; Hornekær, L.; Andersen, L. H., Gas-phase electronic action absorption spectra of protonated oxygen-functionalized polycyclic aromatic hydrocarbons (OPAHs). *Astron. Astrophys.* **2023**, *674*. [DOI: 10.1051/0004-6361/202346003]

⁴⁶⁸ Jaganathan, R.; Simonsen, F. D. S.; Thrower, J. D.; Hornekær, L., Enhanced reactivity of oxygen-functionalised PAHs with atomic hydrogen – A route to the formation of small oxygen-carrying molecules. *Astron. Astrophys.* **2022**, *663*. [DOI: 10.1051/0004-6361/202243312]

⁴⁶⁹ Ferullo, R. M.; Zubieta, C. E.; Belelli, P. G. Hydrogenated polycyclic aromatic hydrocarbons (HnPAHs) as catalysts for hydrogenation reactions in the interstellar medium: a quantum chemical model. *Phys. Chem. Chem. Phys.* **2019**, *21*, 12012. [DOI: 10.1039/C9CP02329A]

⁴⁷⁰ Gordeev, E. G.; Pentsak, E. O.; Ananikov, V. P. Carbocatalytic Acetylene Cyclotrimerization: A Key Role of Unpaired Electron Delocalization. *J. Am. Chem. Soc.* **2020**, *142*, 3784. [DOI: 10.1021/jacs.9b10887]

⁴⁷¹ Nakano, R.; Jazzar, R.; Bertrand, G. A crystalline monosubstituted carbine. *Nat. Chem.* **2018**, *10*, 1196. [DOI: 10.1038/s41557-018-0153-1]

⁴⁷² He, C.; Galimova, G. R.; Luo, Y.; Zhao, L.; Eckhardt, A. K.; Sun, R.; Mebel, A. M.; Kaiser, R. I. A chemical dynamics study on the gas-phase formation of triplet and singlet C5H2 carbenes. *Proc. Natl. Acad. Sci. U. S. A.* **2020**, *117*, 30142. [DOI: 10.1073/pnas.2019257117]

⁴⁷³ Kaye, J. A.; Strobel, D. F., HCN formation on Jupiter: The coupled photochemistry of ammonia and acetylene. *Icarus* **1983**, *54* (3), 417-433. [DOI: 10.1016/0019-1035(83)90238-5]

⁴⁷⁴ Sagan, C.; Khare, B. N., Tholins: organic chemistry of interstellar grains and gas. *Nature* **1979**, 277 (5692), 102-107. [DOI: 10.1038/277102a0]

⁴⁷⁵ Cruikshank, D. P.; Materese, C. K.; Pendleton, Y. J.; Boston, P. J.; Grundy, W. M.; Schmitt, B.; Lisse, C. M.; Runyon, K. D.; Keane, J. T.; Beyer, R. A.; Summers, M. E.; Scipioni, F.; Stern, S. A.; Dalle Ore, C. M.; Olkin, C. B.; Young, L. A.; Ennico, K.; Weaver, H. A.; Bray, V. J., Prebiotic Chemistry of Pluto. *Astrobiology* **2019**, *19* (7), 831-848. [DOI: 10.1089/ast.2018.1927]

⁴⁷⁶ Neish, C. D.; Somogyi, Á.; Lunine, J. I.; Smith, M. A., Low temperature hydrolysis of laboratory tholins in ammonia-water solutions: Implications for prebiotic chemistry on Titan. *Icarus* **2009**, *201* (1), 412-421. [DOI: 10.1016/j.icarus.2009.01.003]

⁴⁷⁷ Cable, M. L.; Vu, T. H.; Maynard-Casely, H. E.; Choukroun, M.; Hodyss, R., The Acetylene-Ammonia Co-crystal on Titan. *ACS Earth Space Chem.* **2018**, *2* (4), 366-375. [DOI: 10.1021/acsearthspacechem.7b00135]

⁴⁷⁸ Ennis, C.; Cable, M. L.; Hodyss, R.; Maynard-Casely, H. E., Mixed Hydrocarbon and Cyanide Ice Compositions for Titan's Atmospheric Aerosols: A Ternary-Phase Co-crystal Predicted by Density Functional Theory. *ACS Earth Space Chem.* **2020**, *4* (7), 1195-1200. [DOI: 10.1021/acsearthspacechem.0c00130]

⁴⁷⁹ Cable, M. L.; Vu, T. H.; Malaska, M. J.; Maynard-Casely, H. E.; Choukroun, M.; Hodyss, R., Properties and Behavior of the Acetonitrile–Acetylene Co-Crystal under Titan Surface Conditions. *ACS Earth Space Chem.* **2020**, *4* (8), 1375-1385.
[DOI: 10.1021/acsearthspacechem.0c00129]

⁴⁸⁰ Czaplinski, E. C.; Vu, T. H.; Cable, M. L.; Choukroun, M.; Malaska, M. J.; Hodyss, R., Experimental Characterization of the Pyridine: Acetylene Co-crystal and Implications for Titan's Surface. *ACS Earth Space Chem.* **2023**, *7* (3), 597-608. [DOI: 10.1021/acsearthspacechem.2c00377]

⁴⁸¹ Norman, L. H., Is there life on ... Titan? *Astron. Geophys.* **2011**, *52* (1), 1.39-1.42. [DOI: 10.1111/j.1468-4004.2011.52139.x]

⁴⁸² Seager, S.; Schrenk, M.; Bains, W., An astrophysical view of Earth-based metabolic biosignature gases. *Astrobiology* **2012**, *12* (1), 61-82. [DOI: 10.1089/ast.2010.0489]

⁴⁸³ Seager, S.; Bains, W.; Hu, R., A Biomass-Based Model to Estimate the Plausibility of Exoplanet Biosignature Gases. *Astrophys. J.* **2013**, 775 (2). [DOI: 10.1088/0004-637X/775/2/104]

⁴⁸⁴ Bains, W., Many chemistries could be used to build living systems. Astrobiology **2004**, *4* (2),

137-67. [DOI: 10.1089/153110704323175124]

⁴⁸⁵ McKay, C. P.; Smith, H. D., Possibilities for methanogenic life in liquid methane on the surface of Titan. *Icarus* **2005**, *178* (1), 274-276. [DOI: 10.1016/j.icarus.2005.05.018]

⁴⁸⁶ Catling D.C. Astrobiology: A Very Short Introduction, OUP Oxford, Oxford, UK, 2013.

⁴⁸⁷ Abbas, O.; Schulze-Makuch, D., Acetylene-based pathways for prebiotic evolution on Titan. *Esa. Sp. Publ.* **2002,** *518,* 345-348. [https://ui.adsabs.harvard.edu/abs/2002ESASP.518..345A/abstract]

⁴⁸⁸ Segura, A.; Navarro-Gonzalez, R., Production of low molecular weight hydrocarbons by volcanic eruptions on early Mars. *Orig. Life Evol. Biosph.* **2005**, *35* (5), 477-87. [DOI: 10.1007/s11084-005-6420-3]

⁴⁸⁹ Kasting, J. F., Earth's early atmosphere. *Science* **1993**, *259* (5097), 920-6. [DOI: 10.1126/science.11536547]

⁴⁹⁰ Sossi, P. A.; Burnham, A. D.; Badro, J.; Lanzirotti, A.; Newville, M.; O'Neill, H. S. C., Redox state of Earth's magma ocean and its Venus-like early atmosphere. *Sci. Adv.* **2020**, *6* (48). [DOI: 10.1126/sciadv.abd1387]

⁴⁹¹ Lupu, R. E.; Zahnle, K.; Marley, M. S.; Schaefer, L.; Fegley, B.; Morley, C.; Cahoy, K.; Freedman, R.; Fortney, J. J., The Atmospheres of Earthlike Planets after Giant Impact Events. *Astrophys. J.* **2014**, *784* (1). [DOI: 10.1088/0004-637X/784/1/27]

⁴⁹² Tian, F.; Kasting, J. F.; Zahnle, K., Revisiting HCN formation in Earth's early atmosphere. *Earth Planet. Sci. Lett.* **2011**, *308* (3-4), 417-423. [DOI: 10.1016/j.epsl.2011.06.011]

⁴⁹³ Capaccioni, B.; Martini, M.; Mangani, F.; Giannini, L.; Nappii, G.; Prati, F., Light hydrocarbons in gas-emissions from volcanic areas and geothermal fields. *Geochem. J.* **1993**, 27 (1), 7-17. [DOI: 10.2343/geochemj.27.7]

⁴⁹⁴ Muenow, D. W., High temperature mass spectrometric gas-release studies of Hawaiian volcanic glass: Pele's tears. *Geochim. Cosmochim. Acta* **1973**, *37* (6), 1551-1561. [DOI: 10.1016/0016-7037(73)90090-2]

⁴⁹⁵ Mukhin, L. M., Volcanic processes and synthesis of simple organic compounds on primitive earth. *Orig. Life* **1976**, *7* (4), 355-68. [DOI: 10.1007/BF00927931]

⁴⁹⁶ Rimmer, P. B.; Shorttle, O., Origin of Life's Building Blocks in Carbon- and Nitrogen-Rich Surface Hydrothermal Vents. *Life (Basel)* **2019**, *9* (1). [DOI: 10.3390/life9010012]

⁴⁹⁷ Schink, B., Fermentation of acetylene by an obligate anaerobe, Pelobacter acetylenicus sp. nov. *Arch. Microbiol.* **1985**, *142* (3), 295-301. [DOI: 10.1007/bf00693407]

⁴⁹⁸ Sutton, J. M.; Baesman, S. M.; Fierst, J. L.; Poret-Peterson, A. T.; Oremland, R. S.; Dunlap, D. S.; Akob, D. M., Complete Genome Sequences of Two Acetylene-Fermenting Pelobacter acetylenicus Strains. *Genome Announc.* **2017**, *5* (6). [DOI: 10.1128/genomeA.01572-16]

⁴⁹⁹ Boll, M.; Einsle, O.; Ermler, U.; Kroneck, P. M.; Ullmann, G. M., Structure and Function of the Unusual Tungsten Enzymes Acetylene Hydratase and Class II Benzoyl-Coenzyme A Reductase. *J. Mol. Microbiol. Biotechnol.* **2016**, *26* (1-3), 119-37. [DOI: 10.1159/000440805]

⁵⁰⁰ Tenbrink, F.; Schink, B.; Kroneck, P. M., Exploring the active site of the tungsten, iron-sulfur enzyme acetylene hydratase. *J. Bacteriol.* **2011**, *193* (5), 1229-36. [DOI: 10.1128/jb.01057-10]

⁵⁰¹ Lovett, R. A., Enceladus named sweetest spot for alien life. *Nature* **2011**. [DOI: 10.1038/news.2011.337]

⁵⁰² Bernstein, M. P.; Moore, M. H.; Elsila, J. E.; Sandford, S. A.; Allamandola, L. J.; Zare, R. N., Side Group Addition to the Polycyclic Aromatic Hydrocarbon Coronene by Proton Irradiation in Cosmic Ice Analogs. *Astrophys. J.* **2003**, *582* (1), L25-L29. [DOI: 10.1086/345941]

⁵⁰³ Bernstein, M. P.; Dworkin, J. P.; Sandford, S. A.; Allamandola, L. J., Ultraviolet irradiation of naphthalene in H2O ice: Implications for meteorites and biogenesis. *Meteorit. Planet. Sci.* **2001**, *36* (3), 351-358. [DOI: 10.1111/j.1945-5100.2001.tb01878.x]

⁵⁰⁴ Ashbourn, S. F. M.; Elsila, J. E.; Dworkin, J. P.; Bernstein, M. P.; Sandford, S. A.; Allamandola, L. J., Ultraviolet photolysis of anthracene in H2O interstellar ice analogs: Potential connection to meteoritic organics. *Meteorit. Planet. Sci.* **2007**, *42* (12), 2035-2041. [DOI: 10.1111/j.1945-5100.2007.tb01005.x]

⁵⁰⁵ Bernstein, M. P.; Sandford, S. A.; Allamandola, L. J.; Gillette, J. S.; Clemett, S. J.; Zare, R. N., UV irradiation of polycyclic aromatic hydrocarbons in ices: production of alcohols, quinones, and ethers. *Science* **1999**, *283* (5405), 1135-8. [DOI: 10.1126/science.283.5405.1135]

⁵⁰⁶ Peeters, Z.; Botta, O.; Charnley, S. B.; Ruiterkamp, R.; Ehrenfreund, P. The Astrobiology of Nucleobases. *Astrophys. J.* **2003**, *593*, L129. [DOI: 10.1086/378346]

⁵⁰⁷ Smith, K. E.; Callahan, M. P.; Gerakines, P. A.; Dworkin, J. P.; House, C. H. Investigation of pyridine carboxylic acids in CM2 carbonaceous chondrites: Potential precursor molecules for ancient coenzymes. *Geochim. Cosmochim. Acta* **2014**, *136*, 1. [DOI: 10.1016/j.gca.2014.04.001]

⁵⁰⁸ Callahan, M. P.; Smith, K. E.; Cleaves, H. J., 2nd; Ruzicka, J.; Stern, J. C.; Glavin, D. P.; House, C. H.; Dworkin, J. P. Carbonaceous meteorites contain a wide range of extraterrestrial nucleobases. *Proc. Natl. Acad. Sci. U. S. A.* **2011**, *108*, 13995. [DOI: 10.1073/pnas.1106493108]

⁵⁰⁹ Martins, Z.; Botta, O.; Fogel, M. L.; Sephton, M. A.; Glavin, D. P.; Watson, J. S.; Dworkin, J. P.; Schwartz, A. W.; Ehrenfreund, P. Extraterrestrial nucleobases in the Murchison meteorite. *Earth Planet. Sci. Lett.* **2008**, *270*, 130. [DOI: 10.1016/j.epsl.2008.03.026]

⁵¹⁰ Folsome, C. E.; Lawless, J.; Romiez, M.; Ponnamperuma, C., Heterocyclic compounds indigenous to the Murchison meteorite. *Nature* **1971**, *232* (5306), 108-9. [DOI: 10.1038/232108a0]

⁵¹¹ Hayatsu, R.; Studier, M. H.; Moore, L. P.; Anders, E., Purines and triazines in the Murchison meteorite. *Geochim. Cosmochim. Acta* **1975**, *39* (4), 471-488. [DOI: 10.1016/0016-7037(75)90101-5]

⁵¹² Stoks, P. G.; Schwartz, A. W., Uracil in carbonaceous meteorites. *Nature* **1979**, 282 (5740), 709-710. [DOI: 10.1038/282709a0]

⁵¹³ Martins, Z., The Nitrogen Heterocycle Content of Meteorites and Their Significance for the Origin of Life. *Life (Basel)* **2018**, *8* (3). [DOI: 10.3390/life8030028]

⁵¹⁴ Pizzarello, S.; Holmes, W. Nitrogen-containing compounds in two CR2 meteorites: 15N composition, molecular distribution and precursor molecules. *Geochim. Cosmochim. Acta* **2009**, 73, 2150. [DOI: 10.1016/j.gca.2009.01.022]

⁵¹⁵ Hudgins, D. M.; Bauschlicher, J. C. W.; Allamandola, L. J., Variations in the Peak Position of the 6.2 μm Interstellar Emission Feature: A Tracer of N in the Interstellar Polycyclic Aromatic Hydrocarbon Population. *Astrophys. J.* **2005**, *632* (1), 316-332. [DOI: 10.1086/432495]

⁵¹⁶ Peeters, Z.; Botta, O.; Charnley, S. B.; Kisiel, Z.; Kuan, Y. J.; Ehrenfreund, P. Formation and photostability of N-heterocycles in space. *Astron. Astrophys.* **2005**, *433*, 583. [DOI: 10.1051/0004-6361:20042443]

⁵¹⁷ Parker, D. S. N.; Kaiser, R. I.; Kostko, O.; Troy, T. P.; Ahmed, M.; Mebel, A. M.; Tielens, A. G. G. M. Gas Phase Synthesis of (Iso)Quinoline and Its Role in the Formation of Nucleobases in the Interstellar Medium. *Astrophys. J.* **2015**, *803*, 53. [DOI: 10.1088/0004-637X/803/2/53]

⁵¹⁸ Parker, D. S.; Kaiser, R. I.; Kostko, O.; Troy, T. P.; Ahmed, M.; Sun, B. J.; Chen, S. H.; Chang, A. H. On the formation of pyridine in the interstellar medium. *Phys. Chem. Chem. Phys.* **2015**, *17*, 32000. [DOI: 10.1039/C5CP02960K]

⁵¹⁹ Morales, S. B.; Bennett, C. J.; Le Picard, S. D.; Canosa, A.; Sims, I. R.; Sun, B. J.; Chen, P. H.; Chang, A. H. H.; Kislov, V. V.; Mebel, A. M.; Gu, X.; Zhang, F.; Maksyutenko, P.; Kaiser, R. I. A Crossed Molecular Beam, Low-Temperature Kinetics, and Theoretical Investigation of the Reaction of the Cyano Radical (Cn) with 1,3-Butadiene (C4h6). A Route to Complex Nitrogen-Bearing Molecules in Low-Temperature Extraterrestrial Environments. *Astrophys. J.* **2011**, *742*, 26. [DOI: 10.1088/0004-637X/742/1/26]

⁵²⁰ Sun, B. J.; Huang, C. H.; Chen, S. Y.; Chen, S. H.; Kaiser, R. I.; Chang, A. H. Theoretical study on reaction mechanism of ground-state cyano radical with 1,3-butadiene: prospect of pyridine formation. *J. Phys. Chem.* **2014**, *118*, 7715. [DOI: 10.1021/jp5056864]

⁵²¹ Zhao, L.; Prendergast, M.; Kaiser, R. I.; Xu, B.; Lu, W.; Ahmed, M.; Hasan Howlader, A.; Wnuk, S. F.; Korotchenko, A. S.; Evseev, M. M.; Bashkirov, E. K.; Azyazov, V. N.; Mebel, A. M., A molecular beam and computational study on the barrierless gas phase formation of (iso)quinoline in low temperature extraterrestrial environments. *Phys. Chem. Chem. Phys.* **2021**, 23 (34), 18495-18505. [DOI: 10.1039/d1cp02169a]

⁵²² Ricca, A., A Computational Study of the Mechanisms for the Incorporation of a Nitrogen Atom into Polycyclic Aromatic Hydrocarbons in the Titan Haze. *Icarus* **2001**, *154* (2), 516-521. [DOI: 10.1006/icar.2001.6694]

⁵²³ Lorenz, R. D.; Lunine, J. I. Titan's surface reviewed: the nature of bright and dark terrain. *Planet. Space Sci.* **1997**, *45*, 981. [DOI: 10.1016/S0032-0633(97)00087-1]

⁵²⁴ Allamandola, L. J.; Tielens, A. G.; Barker, J. R. Interstellar polycyclic aromatic hydrocarbons: the infrared emission bands, the excitation/emission mechanism, and the astrophysical implications. *Astrophys. J., Suppl. Ser.* **1989**, *71*, 733. [DOI: 10.1086/191396]

⁵²⁵ Rap, D. B.; Schrauwen, J. G. M.; Marimuthu, A. N.; Redlich, B.; Brünken, S., Low-temperature nitrogen-bearing polycyclic aromatic hydrocarbon formation routes validated by infrared spectroscopy. *Nat. Astron* **2022**, *6* (9), 1059-1067. [DOI: 10.1038/s41550-022-01713-z]

⁵²⁶ Menor-Salvan, C.; Marin-Yaseli, M. R., A new route for the prebiotic synthesis of nucleobases and hydantoins in water/ice solutions involving the photochemistry of acetylene. *Chem. - Eur. J.* **2013**, *19* (20), 6488-97. [DOI: 10.1002/chem.201204313]

⁵²⁷ Shapiro, R., Prebiotic cytosine synthesis: a critical analysis and implications for the origin of life. *Proc. Natl. Acad. Sci. U S A* **1999**, *96* (8), 4396-401. [DOI: 10.1073/pnas.96.8.4396]

⁵²⁸ Albernaz, A. F.; da Silva, W. B.; Correa, E., Isocyanoacetylene and cyanoacetylene formation study from C_2H_2 + CN reaction. *Int. J. Chem. Kinet.* **2022**, *54* (5), 309-316. [DOI: 10.1002/kin.21560]

⁵²⁹ Hopf, H.; Mlynek, C.; McMahon, R. J.; Menke, J. L.; Lesarri, A.; Rosemeyer, M.; Grabow, J. U., On the trimerization of cyanoacetylene: mechanism of formation of tricyanobenzene isomers and laboratory detection of their radio spectra. *Chem. - Eur. J.* **2010**, *16* (47), 14115-23. [DOI: 10.1002/chem.201001648]

⁵³⁰ Miller, S. L.; Urey, H. C., Organic compound synthesis on the primitive earth. *Science* **1959**, *130* (3370), 245-51. [DOI: 10.1126/science.130.3370.245]

⁵³¹ Sanchez, R. A.; Ferris, J. P.; Orgel, L. E., Cyanoacetylene in prebiotic synthesis. *Science* **1966**, *154* (3750), 784-5. [DOI: 10.1126/science.154.3750.784]

⁵³² Kaur, S.; Ohri, A.; Sharma, P., Could Purines Be Formed from Cyanamide and Cyanoacetylene in a Prebiotic Earth Environment? *ACS Omega* **2019**, *4* (7), 12771-12781. [DOI: 10.1021/acsomega.9b01169]

⁵³³ Seitz, C.; Eisenreich, W.; Huber, C., The Abiotic Formation of Pyrrole under Volcanic, Hydrothermal Conditions-An Initial Step towards Life's First Breath? *Life (Basel)* **2021**, *11* (9). [DOI: 10.3390/life11090980]

⁵³⁴ Platts S. N. The PAH World: Discotic polynuclear aromatic compounds as a mesophase scaffolding at the origin of life, http://www.pahworld.com/ (accessed 2004-11-11).

⁵³⁵ Nuevo, M.; Milam, S. N.; Sandford, S. A.; Elsila, J. E.; Dworkin, J. P., Formation of uracil from the ultraviolet photo-irradiation of pyrimidine in pure H2O ices. *Astrobiology* **2009**, *9* (7), 683-95. [DOI: 10.1089/ast.2008.0324]

⁵³⁶ Nuevo, M.; Milam, S. N.; Sandford, S. A., Nucleobases and prebiotic molecules in organic residues produced from the ultraviolet photo-irradiation of pyrimidine in NH(3) and H(2)O+NH(3) ices. *Astrobiology* **2012**, *12* (4), 295-314. [DOI: 10.1089/ast.2011.0726]

⁵³⁷ Nuevo, M.; Materese, C. K.; Sandford, S. A., The Photochemistry of Pyrimidine in Realistic Astrophysical Ices and the Production of Nucleobases. *Astrophys. J.* **2014**, *793* (2). [DOI: 10.1088/0004-637X/793/2/125]

⁵³⁸ Materese, C. K.; Nuevo, M.; Sandford, S. A., The Formation of Nucleobases from the Ultraviolet Photoirradiation of Purine in Simple Astrophysical Ice Analogues. *Astrobiology* **2017**, *17* (8), 761-770. [DOI: 10.1089/ast.2016.1613]

⁵³⁹ Materese, C. K.; Nuevo, M.; McDowell, B. L.; Buffo, C. E.; Sandford, S. A., The Photochemistry of Purine in Ice Analogs Relevant to Dense Interstellar Clouds. *Astrophys. J.* **2018**, *864* (1). [DOI: 10.3847/1538-4357/aad328]

⁵⁴⁰ Oba, Y.; Takano, Y.; Naraoka, H.; Watanabe, N.; Kouchi, A., Nucleobase synthesis in interstellar ices. *Nat Commun* **2019**, *10* (1), 4413. [DOI: 10.1038/s41467-019-12404-1]

⁵⁴¹ Ge, J.; Mardones, D.; Inostroza, N.; Peng, Y., The roles of polycyclic aromatic hydrocarbons in dark cloud chemistry: new constraints on sulphur-bearing species. *Mon. Not. R. Astron. Soc.* **2020**, *497* (3), 3306-3322. [DOI: 10.1093/mnras/staa2098]

⁵⁴² Brosnan, J. T.; Brosnan, M. E., The sulfur-containing amino acids: an overview. *J. Nutr.* **2006**, *136* (6 Suppl), 1636S-1640S. [DOI: 10.1093/jn/136.6.1636S]

⁵⁴³ Bacchus-Montabonel, M. C., Role of sulfur in proton-induced collisions of RNA prebiotic precursors. *Phys. Chem. Chem. Phys.* **2019**, *21* (3), 1428-1434. [DOI: 10.1039/c8cp07204c]

⁵⁴⁴ Cohen, Z. R.; Todd, Z. R.; Wogan, N.; Black, R. A.; Keller, S. L.; Catling, D. C., Plausible Sources of Membrane-Forming Fatty Acids on the Early Earth: A Review of the Literature and an Estimation of Amounts. *ACS Earth Space Chem.* **2023**, *7* (1), 11-27. [DOI: 10.1021/acsearthspacechem.2c00168]

⁵⁴⁵ Lai, J. C. Y.; Pearce, B. K. D.; Pudritz, R. E.; Lee, D., Meteoritic abundances of fatty acids and potential reaction pathways in planetesimals. *Icarus* **2019**, *319*, 685-700. [DOI: 10.1016/j.icarus.2018.09.028]

⁵⁴⁶ Markvoort, A. J.; Pfleger, N.; Staffhorst, R.; Hilbers, P. A.; van Santen, R. A.; Killian, J. A.; de Kruijff, B., Self-reproduction of fatty acid vesicles: a combined experimental and simulation study. *Biophys. J.* **2010**, *99* (5), 1520-8. [DOI: 10.1016/j.bpj.2010.06.057]

⁵⁴⁷ Damer, B.; Deamer, D., Coupled phases and combinatorial selection in fluctuating hydrothermal pools: a scenario to guide experimental approaches to the origin of cellular life. *Life* (*Basel*) **2015**, *5* (1), 872-87. [DOI: 10.3390/life5010872]

⁵⁴⁸ Schrum, J. P.; Zhu, T. F.; Szostak, J. W., The origins of cellular life. *Cold Spring Harb. Perspect. Biol.* **2010**, *2* (9), a002212. [DOI: 10.1101/cshperspect.a002212]

⁵⁴⁹ Huang, Y.; Wang, Y.; Alexandre, M. R.; Lee, T.; Rose-Petruck, C.; Fuller, M.; Pizzarello, S., Molecular and compound-specific isotopic characterization of monocarboxylic acids in carbonaceous meteorites. *Geochim. Cosmochim. Acta* **2005**, *69* (4), 1073-1084. [DOI: 10.1016/j.gca.2004.07.030]

⁵⁵⁰ Naraoka, H.; Shimoyama, A.; Harada, K., Molecular distribution of monocarboxylic acids in Asuka carbonaceous chondrites from Antarctica. *Orig. Life Evol. Biosph.* **1999**, *29* (2), 187-201. [DOI: 10.1023/A:1006547127028]

⁵⁵¹ Rotelli, L.; Trigo-Rodriguez, J. M.; Moyano-Cambero, C. E.; Carota, E.; Botta, L.; Di Mauro, E.; Saladino, R., The key role of meteorites in the formation of relevant prebiotic molecules in a formamide/water environment. *Sci Rep* **2016**, *6*, 38888. [DOI: 10.1038/srep38888]

⁵⁵² Scheidler, C.; Sobotta, J.; Eisenreich, W.; Wachtershauser, G.; Huber, C., Unsaturated C3,5,7,9-Monocarboxylic Acids by Aqueous, One-Pot Carbon Fixation: Possible Relevance for the Origin of Life. *Sci. Rep.* **2016**, *6*, 27595. [DOI: 10.1038/srep27595]

⁵⁵³ Sobotta, J.; Geisberger, T.; Moosmann, C.; Scheidler, C. M.; Eisenreich, W.; Wachtershauser, G.; Huber, C., A Possible Primordial Acetyleno/Carboxydotrophic Core Metabolism. *Life (Basel)* **2020**, *10* (4). [DOI: 10.3390/life10040035]

⁵⁵⁴ Diederich, P.; Ruf, A.; Geisberger, T.; Weidner, L.; Seitz, C.; Eisenreich, W.; Huber, C.; Schmitt-Kopplin, P., C₂-addition patterns emerging from acetylene and nickel sulfide in simulated

prebiotic hydrothermal conditions. *Commun. Chem.* **2023**, *6* (1), 220. [DOI: 10.1038/s42004-023-01021-1]

⁵⁵⁵ James Webb Space Telescope <u>https://webb.nasa.gov</u> (accessed 2024-03-01).

⁵⁵⁶ Lorenz, R. D.; MacKenzie, S. M.; Neish, C. D.; Gall, A. L.; Turtle, E. P.; Barnes, J. W.; Trainer, M. G.; Werynski, A.; Hedgepeth, J.; Karkoschka, E. Selection and Characteristics of the Dragonfly Landing Site near Selk Crater, Titan. *Planet. Sci. J.* **2021**, *2*, 24. [DOI: 10.3847/PSJ/abd08f]

⁵⁵⁷ Dragonfly News <u>https://dragonfly.jhuapl.edu/</u> (accessed 2024-03-01).

⁵⁵⁸ JUICE <u>https://sci.esa.int/web/juice</u> (accessed 2024-03-01).