DEFORMED QUANTUM VERTEX ALGEBRA MODULES ASSOCIATED WITH BRAIDINGS

LUCIA BAGNOLI AND SLAVEN KOŽIĆ

ABSTRACT. We introduce the notion of deformed quantum vertex algebra module associated with a braiding map. We construct two families of braiding maps over the Etingof–Kazhdan quantum vertex algebras associated with the rational *R*-matrices of classical types. We investigate their properties and demonstrate the applications of the corresponding deformed modules to the (generalized) Yangians and reflection algebras.

1. INTRODUCTION

A fundamental problem in the theory of vertex algebras, which goes back to I. Frenkel and Jing [10], is to associate suitable vertex algebra-like structures, quantum vertex algebras to different classes of quantum groups. It is motivated by the rich interplay between vertex algebras and representations of various infinite dimensional Lie algebras; see, e.g. the books by Kac [21] and Lepowsky and Li [26]. Motivated by the ideas of E. Frenkel and Reshetikhin [11], Etingof and Kazhdan [9] introduced the notion of quantum vertex algebra. Furthermore, they constructed first examples of these structures associated with the rational, trigonometric and elliptic *R*-matrix of type A. Later on, some other important generalizations of the notion of vertex algebra, such as *field algebras* of Bakalov and Kac [2], axiomatic G_1 -vertex algebras, inspired by Borcherds' G-vertex algebras [4], and nonlocal vertex algebras of Li [27, 28] and H_D -quantum vertex algebras of Anguelova and Bergvelt [1], were studied. A major step forward in the development of quantum vertex algebra theory was Li's introduction of ϕ -coordinated modules [30] which, in particular, enabled the association of quantum vertex algebras with quantum affine algebras; see, e.g., the papers by Jing, Kong, Li and Tan [17] and Kong [23]. For a more detailed overview of the evolution of quantum vertex algebra theory and its most recent results see [5, 7, 8, 16, 24] and references therein.

Suppose $(V, Y, \mathbf{1})$ is a vertex algebra and W a V-module. The corresponding module map $Y_W(\cdot, z)$ possesses two key properties (see, e.g., [21, 26]), *locality*: for any $u, v \in V$

$$Y_W(u, z_1) Y_W(v, z_2) \sim Y_W(v, z_2) Y_W(u, z_1), \tag{1.1}$$

and the weak associativity: for any $u, v \in V$ and $w \in W$

$$Y_W(u, z_0 + z_2) Y_W(v, z_2) w \sim Y_W(Y(u, z_0)v, z_2) w.$$
(1.2)

The tilde "~" in (1.1) (resp. (1.2)) indicates that the given expressions coincide when multiplied by $(z_1 - z_2)^r$ (resp. $(z_0 + z_2)^r$) for a sufficiently large positive integer r, which depends on the choice of $u, v \in V$ (resp. $u, v \in V, w \in W$). The analogue of (1.1) for a quantum vertex algebra $(V, Y, \mathbf{1}, S)$ and its module W is *S*-locality: for any $u, v \in V$

$$Y_W(z_1) (1 \otimes Y_W(z_2)) \mathcal{S}(z_1 - z_2) (u \otimes v) \sim_h Y_W(v, z_2) Y_W(u, z_1),$$
(1.3)

⁽L. Bagnoli and S. Kožić) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, UNIVERSITY OF ZAGREB, BIJENIČKA CESTA 30, 10000 ZAGREB, CROATIA

E-mail addresses: lucia.bagnoli@math.hr, kslaven@math.hr.

where S is the braiding of V, a certain unitary map satisfying the quantum Yang–Baxter equation; cf. [9,29]. On the other hand, the analogue of (1.2) for quantum vertex algebra modules, the *weak associativity*, is of the same form: for any $u, v \in V$ and $w \in W$

$$Y_W(u, z_0 + z_2) Y_W(v, z_2) w \sim_h Y_W(Y(u, z_0)v, z_2) w;$$
(1.4)

cf. [9,29]. However, both V and W are now topologically free $\mathbb{C}[[h]]$ -modules and the symbol " \sim_h " indicates that the form of S-locality (1.3) (resp. weak associativity (1.4)) generalizes its classical counterpart (1.1) (resp. (1.2)) using the *h*-adic topology. The products of the given expressions with $(z_1 - z_2)^r$ (resp. $(z_0 + z_2)^r$) are required to coincide modulo h^n only, and the power r now depends on the choice of positive integer n as well.

This paper is motivated in part by the notion of H_D -quantum vertex algebra [1], which presents a generalization of Etingof–Kazhdan's quantum vertex algebras [9]. In particular, it features a more general version of S-locality and a deformation of weak associativity which is governed by a certain braiding map. We study a new concept of (μ, ν) -deformed module, or, more briefly, deformed module for a quantum vertex algebra V, such that its analogues of (1.3) and (1.4) also depend on certain maps

$$\mu, \nu \colon V \otimes V \to V \otimes V \otimes \mathbb{C}[z^{\pm 1}][[x^{-1}, h]].$$
(1.5)

In addition, our notion of deformed module conforms to a slightly uncommon truncation condition: its module map is allowed to possess essential singularities at zero while, on the other hand, it is required to be (topologically) truncated from above. The role of this feature is to accommodate the well-known phenomenon which comes from the theory of Yangians. Namely, the techniques and constructions from this framework can be often extended to other classes of algebras associated with the same R-matrix, such as generalized Yangians, twisted Yangians, reflection algebras etc.; see, e.g., [25, 32, 34, 35]. The truncation condition ensures that the theory of deformed modules fits all these algebras, so that some of their fundamental properties naturally emanate from the corresponding Etingof–Kazhdan quantum affine vertex algebra. We shall demonstrate this occurrence in detail in the setting of generalized Yangians of Krylov and Rybnikov [25] and reflection algebras of Molev and Ragoucy [35].

Recall the maps (1.5). In contrast with (1.3), which depends on the original braiding \mathcal{S} of V, the deformed module map $Y_W(\cdot, z)$ satisfies the μ -commutativity: for any $u, v \in V$

$$Y_W(z_1) \big(1 \otimes Y_W(z_2) \big) \mu(-z_2 + z_1, z_2)(u \otimes v) = Y_W(v, z_2) Y_W(u, z_1).$$
(1.6)

Roughly speaking, the map μ is a unitary solution of the quantum Yang–Baxter equation, which is conjugated with S by ν , i.e. we have

$$\mathcal{S}(z)\nu_{21}(-z,x+z) = \nu(z,x)\mu(z,x).$$

We refer to such a pair (μ, ν) as a *compatible pair*. As one may expect, such a structure does no longer need to possess the weak associativity property (1.4). However, we require that it exhibits the ν -associativity: for any $u, v \in V$ and $w \in W$

$$Y_W(u, z_2 + z_0) Y_W(v, z_2) w = Y_W(Y(z_0) \nu(z_0, z_2)(u \otimes v), z_2) w.$$
(1.7)

Let us describe the organization and main results of this manuscript. In Section 2, we provide some preliminaries on the double Yangians and quantum affine vertex algebras of classical types. In Section 3, we introduce and discuss the notion of (μ, ν) -deformed module. Next, in Section 4, we construct two compatible pairs, (σ, ρ) and (μ, ν) . The construction of (σ, ρ) is carried out in the setting of Etingof–Kazhdan's quantum vertex algebras $\mathcal{V}^c(\mathfrak{g}_N)$ associated with rational *R*-matrices of types A-D at an arbitrary level $c \in \mathbb{C}$. Furthermore, we use the fusion procedure for these *R*-matrices to investigate fixed points of σ at the critical level $c = c_{\text{crit}}$. For \mathfrak{g}_N of type A and B, we show that the restriction of σ on the tensor square of quantum Feigin-Frenkel center, i.e. the center of $\mathcal{V}^{\text{crit}}(\mathfrak{g}_N) = \mathcal{V}^{c_{\text{crit}}}(\mathfrak{g}_N)$, is the identity, while for types C and D, we only obtain some partial results of the same form, due to a lack of explicit formulae for complete sets of Segal-Sugawara vectors in these types; cf. [7,18]. Finally, the construction of pair (μ, ν) is given for the quantum affine vertex algebra $\mathcal{V}^{\text{crit}}(\mathfrak{gl}_N)$ associated with rational R-matrix of type A at the critical level. Moreover, some families of fixed points of μ , which come from the quantum Feigin-Frenkel center, are found in this case as well.

As an application, in Section 5 (resp. Section 6), we use the maps from Section 4 to show that the generalized Yangians (as well as ordinary Yangians) of types A-D (resp. certain reflection algebras) are naturally equipped with structure of (σ, ρ) -deformed (resp. (μ, ν) -deformed) $\mathcal{V}^c(\mathfrak{g}_N)$ -module. Next, we establish an equivalence between certain wide classes of modules associated with such algebras and deformed $\mathcal{V}^c(\mathfrak{g}_N)$ -modules. Finally, we show that commutative subalgebras and families of central elements in these algebras naturally arise from the families of fixed points of σ and μ found in Section 4.

2. Preliminaries

In this section, we recall some preliminary definitions and results on the double Yangians of types A-D and the associated quantum vertex algebras.

2.1. Double Yangians of classical types. Fix an integer $N \ge 2$. Let \mathfrak{gl}_N be the general linear, \mathfrak{o}_N the orthogonal and \mathfrak{sp}_N the symplectic Lie algebra, where N is even in the symplectic case. In order to consider all of them simultaneously, we denote by \mathfrak{g}_N any of these Lie algebras. In the case $\mathfrak{g}_N = \mathfrak{o}_N, \mathfrak{sp}_N$, introduce the scalars $\varepsilon_1 = \ldots = \varepsilon_N = 1$ if $\mathfrak{g}_N = \mathfrak{o}_N$ and $\varepsilon_1 = \ldots = \varepsilon_{N/2} = 1$, $\varepsilon_{(N+2)/2} = \ldots = \varepsilon_N = -1$ if $\mathfrak{g}_N = \mathfrak{sp}_N$. For any $i = 1, \ldots, N$ set i' = N - i + 1. Finally, let $G = (g_{ij}) \in \text{End } \mathbb{C}^N$ be the matrix defined by $g_{ij} = \delta_{ij'}\varepsilon_i$ for all $i, j = 1, \ldots, N$. Clearly, G is symmetric in the orthogonal and skewsymmetric in the symplectic case. For any matrix $A = (a_{ij})$ in $\text{End } \mathbb{C}^N$ let $A' = GA^tG^{-1}$, where A^t denotes the transposed matrix $A^t = (a_{ji})$. We have $A' = (\varepsilon_i \varepsilon_j a_{j'i'})$. Set

$$\varepsilon = \begin{cases} 1, & \text{if } \mathfrak{g}_N = \mathfrak{gl}_N, \mathfrak{o}_N, \\ 2, & \text{if } \mathfrak{g}_N = \mathfrak{sp}_N. \end{cases}$$

Let h be a formal parameter and $\mathbb{C}[[h]]$ a commutative ring of formal power series in h. Define the operators I, P and Q in $\operatorname{End} \mathbb{C}^N \otimes \operatorname{End} \mathbb{C}^N$ by

$$I = \sum_{i,j=1}^{N} e_{ii} \otimes e_{jj}, \quad P = \sum_{i,j=1}^{N} e_{ij} \otimes e_{ji} \quad \text{and} \quad Q = \sum_{i,j=1}^{N} \varepsilon_i \varepsilon_j e_{ij} \otimes e_{i'j'},$$

where e_{ij} are matrix units. For $\mathfrak{g}_N = \mathfrak{gl}_N$, we consider the Yang *R*-matrix R(u)

$$R(u) = I - \frac{h}{u}P \tag{2.1}$$

and for $\mathfrak{g}_N = \mathfrak{o}_N, \mathfrak{sp}_N$, the *R*-matrix, which goes back to [37],

$$R(u) = I - \frac{h}{u}P + \frac{h}{u - h\kappa}Q, \quad \text{where} \quad \kappa = \begin{cases} N/2 - 1, & \text{if } \mathfrak{g}_N = \mathfrak{o}_N, \\ N/2 + 1, & \text{if } \mathfrak{g}_N = \mathfrak{sp}_N. \end{cases}$$
(2.2)

The *R*-matrices in (2.1) and (2.2) satisfy the quantum Yang-Baxter equation

$$R_{12}(u)R_{13}(u+v)R_{23}(v) = R_{23}(v)R_{13}(u+v)R_{12}(u).$$
(2.3)

Both sides of (2.3) are regarded as operators on the tensor product $(\mathbb{C}^N)^{\otimes 3}$ and the subscripts indicate the copies of \mathbb{C}^N on which the *R*-matrices act, e.g., $R_{12}(u) = R(u) \otimes 1$.

Let $\overline{R}(u) = f(u/h)R(u)$ be the normalized *R*-matrix, where for $\mathfrak{g}_N = \mathfrak{gl}_N$ (resp. $\mathfrak{g}_N = \mathfrak{o}_N, \mathfrak{sp}_N$), f(u) is a unique series in $1 + u^{-1}\mathbb{C}[[u^{-1}]]$ (resp. $1 + u^{-2}\mathbb{C}[[u^{-1}]]$) which satisfies

$$f(u) f(u+N)^{-1} = f(u) f(-u) = \left(1 - u^{-2}\right)^{-1}$$

(resp. $f(u) f(u+\kappa) = f(u) f(-u) = \left(1 - u^{-2}\right)^{-1}$)

see [7, Subsect. 1.2] and [18, Subsect. 2.2] for more details on f(u). The normalized *R*-matrices possess the *unitarity* property,

$$\overline{R}(u)\overline{R}(-u) = \overline{R}(-u)\overline{R}(u) = 1, \qquad (2.4)$$

and the *crossing symmetry* property,¹

$$\overline{R}(-u)^{t}\overline{R}(u+hN)^{t} = 1 \quad \text{for } \mathfrak{g}_{N} = \mathfrak{gl}_{N}$$
(2.5)

$$\overline{R}(u)\overline{R}(u+h\kappa)' = 1 \quad \text{for } \mathfrak{g}_N = \mathfrak{o}_N, \mathfrak{sp}_N.$$
(2.6)

Alternatively, the crossing symmetry property can be expressed as

$$\overline{R}(-u)_{\mathrm{LR}}\overline{R}(u+hN) = \overline{R}(-u)_{\mathrm{RL}}\overline{R}(u+hN) = 1 \quad \text{for } \mathfrak{g}_N = \mathfrak{gl}_N, \qquad (2.7)$$

$$\overline{R}(u)'_{\text{LR}}\overline{R}(u+h\kappa) = \overline{R}(u)'_{\text{RL}}\overline{R}(u+h\kappa) = 1 \quad \text{for } \mathfrak{g}_N = \mathfrak{o}_N, \mathfrak{sp}_N,$$
(2.8)

where "." (resp. ".") denotes the standard multiplication in End $\mathbb{C}^N \otimes (\text{End } \mathbb{C}^N)^{\text{op}}$ (resp. $(\text{End } \mathbb{C}^N)^{\text{op}} \otimes \text{End } \mathbb{C}^N$) and A^{op} is the opposite algebra of A. In addition, we observe that for $\mathfrak{g}_N = \mathfrak{o}_N, \mathfrak{sp}_N$, unitarity (2.4) and crossing symmetry (2.8) imply $\overline{R}(-u)' = \overline{R}(u+h\kappa)$. This can be used to express the crossing symmetry alternatively as

$$\overline{R}(-u+2h\kappa)_{\mathrm{LR}} \overline{R}(u) = \overline{R}(-u+2h\kappa)_{\mathrm{RL}} \overline{R}(u) = 1 \quad \text{for } \mathfrak{g}_N = \mathfrak{o}_N, \mathfrak{sp}_N.$$
(2.9)

Following [13], we define the *double Yangian* $DY(\mathfrak{gl}_N)$ as the *h*-adically completed associative algebra over the ring $\mathbb{C}[[h]]$ generated by the central element *C* and the elements $t_{ij}^{(\pm r)}$, where $i, j = 1, \ldots, N$ and $r = 1, 2, \ldots$, subject to the defining relations

$$R(u-v)T_1^{\pm}(u)T_2^{\pm}(v) = T_2^{\pm}(v)T_1^{\pm}(u)R(u-v), \qquad (2.10)$$

$$\overline{R}(u - v + h\varepsilon C/2)T_1^-(u)T_2^+(v) = T_2^+(v)T_1^-(u)\overline{R}(u - v - h\varepsilon C/2).$$
(2.11)

The generator matrices $T^{\pm}(u)$ are defined by

$$T^{-}(u) = \sum_{i,j=1}^{N} e_{ij} \otimes t^{-}_{ij}(u) \quad \text{and} \quad T^{+}(u) = \sum_{i,j=1}^{N} e_{ij} \otimes t^{+}_{ij}(u)$$
(2.12)

and the series $t_{ij}^{\pm}(u)$ are given by

$$t_{ij}^{-}(u) = \delta_{ij} + h \sum_{r=1}^{\infty} t_{ij}^{(r)} u^{-r} \quad \text{and} \quad t_{ij}^{+}(u) = \delta_{ij} - h \sum_{r=1}^{\infty} t_{ij}^{(-r)} u^{r-1}.$$
 (2.13)

Next, following [19], we define the *double Yangian* $DY(\mathfrak{g}_N)$ for $\mathfrak{g}_N = \mathfrak{o}_N, \mathfrak{sp}_N$ analogously, using (2.10)–(2.13), but with two additional families of defining relations:

$$T^{-}(u)T^{-}(u+h\kappa)' = 1$$
 and $T^{+}(u)T^{+}(u+h\kappa)' = 1.$ (2.14)

Let us recall two important subalgebras of the double Yangian. The Yangian $Y(\mathfrak{g}_N)$ (resp. the dual Yangian $Y^+(\mathfrak{g}_N)$) is defined as the *h*-adically completed subalgebra of $DY(\mathfrak{g}_N)$ generated by all elements $t_{ij}^{(r)}$ (resp. $t_{ij}^{(-r)}$), where $i, j = 1, \ldots, N$ and $r = 1, 2, \ldots$. By the Poincaré–Birkhoff–Witt theorem for the double Yangian [18, Thm. 2.2] and [19,

¹Note that we have $\overline{R}(u)^{t_1} = \overline{R}(u)^{t_2}$ and $\overline{R}(u)^{\prime_1} = \overline{R}(u)^{\prime_2}$, so these transposed *R*-matrices in (2.5) and (2.6) are denoted simply by $\overline{R}(u)^t$ and $\overline{R}(u)^\prime$, respectively.

Thm. 3.4], $Y(\mathfrak{g}_N)$ (resp. $Y^+(\mathfrak{g}_N)$) is isomorphic to the *h*-adically completed associative algebra over $\mathbb{C}[[h]]$ generated by the elements $t_{ij}^{(r)}$ (resp. $t_{ij}^{(-r)}$), where $i, j = 1, \ldots, N$ and $r = 1, 2, \ldots$, subject to the defining relation (2.10) for $T^-(u)$ (resp. for $T^+(u)$) and, in the case $\mathfrak{g}_N = \mathfrak{o}_N, \mathfrak{sp}_N$, the first (resp. the second) relation in (2.14).

For any $c \in \mathbb{C}$, define the double Yangian $DY_c(\mathfrak{g}_N)$ at the level c as the quotient of the algebra $DY(\mathfrak{g}_N)$ by its *h*-adically completed two-sided ideal generated by C - c. The vacuum module $\mathcal{V}^c(\mathfrak{g}_N)$ at the level c is defined as the *h*-adically completed quotient $DY_c(\mathfrak{g}_N)/I$, where I denotes the *h*-adically completed left ideal in $DY_c(\mathfrak{g}_N)$ generated by the elements $t_{ij}^{(r)}$ with $i, j = 1, \ldots, N$ and $r = 1, 2, \ldots$. By the Poincaré–Birkhoff–Witt theorem for the double Yangian, $\mathcal{V}^c(\mathfrak{g}_N)$ and $Y^+(\mathfrak{g}_N)$ are isomorphic as $\mathbb{C}[[h]]$ -modules.

2.2. Fusion procedure. Let R(u) be the Yang *R*-matrix (2.1). We shall recall a particular case of the fusion procedure for R(u) [20]; see also [32, Sect. 6.4] for more details and references. Recall that the symmetric group \mathfrak{S}_n acts on $(\mathbb{C}^N)^{\otimes n}$ by permuting the tensor factors. Denote by $\mathcal{E}_{[n]}$ the action of the anti-symmetrizer $\frac{1}{n!} \sum_{p \in \mathfrak{S}_n} \operatorname{sgn} p \cdot p \in \mathbb{C}[\mathfrak{S}_n]$ on $(\mathbb{C}^N)^{\otimes n}$. Consider the product

$$R_{[n]}(u) = R_{[n]}(u_1, \dots, u_n) = \frac{1}{n!} \prod_{1 \le i < j \le n} R_{ij}(u_i - u_j), \qquad (2.15)$$

where the factors are ordered lexicographically on the pairs (i, j). Introduce the *n*-tuple

$$u_{[n]} = (u, u - h, \dots, u - (n - 1)h), \qquad (2.16)$$

where u is a single variable. By [20], the consecutive evaluations $u_1 = 0, u_2 = -h, u_3 = -2h, \ldots, u_n = -(n-1)h$ of the variables $u = (u_1, \ldots, u_n)$ in (2.15) are well-defined and they produce the following version of the *fusion procedure*:

$$R_{[n]}(u_1, \dots, u_n)\big|_{u_1=0}\big|_{u_2=-h} \dots \big|_{u_n=-(n-1)h} = \mathcal{E}_{[n]}.$$
(2.17)

Let R(u) be the *R*-matrix (2.2). Following the exposition in [32], we shall recall a special case of the fusion procedure for the Brauer algebra [6]; cf. [33, Sect. 1.2]. Let ω be an indeterminate. Denote by $\mathcal{B}_n(\omega)$ the *Brauer algebra* over $\mathbb{C}(\omega)$ generated by elements s_1, \ldots, s_{n-1} and $\varepsilon_1, \ldots, \varepsilon_{n-1}$, which are subject to certain defining relations; see [32, Sect. 3] for details. Its complex subalgebra generated by s_1, \ldots, s_{n-1} is isomorphic to the group algebra of symmetric group \mathfrak{S}_n and the elements s_i are identified with the transpositions (i, i+1). Denote by $s_{ij} \in \mathcal{B}_n(\omega)$ with i < j the element corresponding to the transposition (i, j) with respect to that isomorphism. Consider the elements $\varepsilon_{ij} \in \mathcal{B}_n(\omega)$ defined by

$$\varepsilon_{j-1j} = \varepsilon_{j-1}$$
 and $\varepsilon_{ij} = s_{ij-1}\varepsilon_{j-1}s_{ij-1}$ for $i < j-1$.

Let $s^{(n)} \in \mathcal{B}_n(\omega)$ be the idempotent which corresponds to the one-dimensional representation of $\mathcal{B}_n(\omega)$ which maps all s_{ij} to the identity and all ε_{ij} to the zero. In the orthogonal case, denote by $\mathcal{E}_{[n]}$ with $n = 1, \ldots, N$ the action of the idempotent $s^{(n)} \in \mathcal{B}_n(N)$ on $(\mathbb{C}^N)^{\otimes n}$ defined by

$$s_{ij} \mapsto P_{ij}$$
 and $\varepsilon_{ij} \mapsto Q_{ij}$ for $i < j$.

In the symplectic case, denote by $\mathcal{E}_{[n]}$ with $n = 1, \ldots, N/2$ the action of the idempotent $s^{(n)} \in \mathcal{B}_n(-N)$ on $(\mathbb{C}^N)^{\otimes n}$ defined by

$$s_{ij} \mapsto -P_{ij}$$
 and $\varepsilon_{ij} \mapsto -Q_{ij}$ for $i < j$.

Let $R_{[n]}(u) = R_{[n]}(u_1, \ldots, u_n)$ be the *R*-matrix product of the same form as (2.15). For a single variable u, introduce the *n*-tuples

$$u_{[n]} = \begin{cases} (u - (n - 1)h, \dots, u - h, u) & \text{for } \mathfrak{g}_N = \mathfrak{o}_N \text{ and } n = 1, \dots, N, \\ (u, u - h, \dots, u - (n - 1)h) & \text{for } \mathfrak{g}_N = \mathfrak{sp}_N \text{ and } n = 1, \dots, N/2. \end{cases}$$
(2.18)

By a special case of the fusion procedure for $\mathcal{B}_n(\omega)$ [14, 15], we have

$$R(u_{[n]}) = \mathcal{E}_{[n]}.\tag{2.19}$$

2.3. Etingof–Kazhdan's construction and the quantum Feigin–Frenkel center. Suppose V is a $\mathbb{C}[[h]]$ -module. We shall denote by $V((z))_h$ (resp. $V[z^{\pm 1}]_h$) the $\mathbb{C}[[h]]$ -module of all power series $a(z) = \sum_{r \in \mathbb{Z}} a_r z^{-r} \in V[[z, z^{-1}]]$ (resp. $a(z) = \sum_{r \geq 0} a_r z^{\pm r} \in V[[z^{\pm 1}]]$) such that $\lim_{r\to\infty} a_r = 0$, where the limit is taken with respect to the *h*-adic topology. Throughout the paper, we assume that the tensor products of $\mathbb{C}[[h]]$ -modules are *h*-adically completed; see [22, Ch. XVI] for more details on the *h*-adic topology. For readers' convenience, we start by recalling the definition of *h*-adic quantum vertex algebra; see [9, Sect. 1.4.1] and [29, Def. 2.20]. In the definition, as well as in the rest of the paper, we use the usual vertex algebraic expansion convention, where the expressions such as $(z_1 + \ldots + z_n)^r$ with r < 0 are expanded in nonnegative powers of z_2, \ldots, z_n . Hence, in particular, we have $(z_1 + z_2)^r \neq (z_2 + z_1)^r$ for r < 0.

Definition 2.1. Let V be a topologically free $\mathbb{C}[[h]]$ -module equipped with a distinguished element $\mathbf{1} \in V$ (the *vacuum vector*), the $\mathbb{C}[[h]]$ -module map

$$Y(\cdot, z) \colon V \otimes V \to V((z))_h,$$

$$u \otimes v \mapsto Y(z)(u \otimes v) = Y(u, z)v = \sum_{r \in \mathbb{Z}} u_r v z^{-r-1},$$
(2.20)

(the vertex operator map) and the $\mathbb{C}[[h]]$ -module map

 $\mathcal{S}(z)\colon V\otimes V\to V\otimes V\otimes \mathbb{C}((z))[[h]]$

(the *braiding*). A quadruple $(V, Y, \mathbf{1}, S)$ is said to be an *h*-adic quantum vertex algebra if it satisfies the following axioms.

(1) The vertex operator map possesses the *weak associativity* property: for any elements $u, v, w \in V$ and $n \in \mathbb{Z}_{\geq 0}$ there exists $r \in \mathbb{Z}_{\geq 0}$ such that

$$\left((z_0 + z_2)^r Y(u, z_0 + z_2) Y(v, z_2) w - (z_0 + z_2)^r Y(Y(u, z_0) v, z_2) w \right) \in h^n V[[z_0^{\pm 1}, z_2^{\pm 1}]].$$

(2) The vacuum vector satisfies the *vacuum property* and the *creation property*

$$Y(\mathbf{1}, z)v = v, \quad Y(v, z) \mathbf{1} \in V[[z]] \text{ and } \lim_{z \to 0} Y(v, z) \mathbf{1} = v \text{ for all } v \in V.$$

(3) The braiding satisfies the *shift condition*

$$[\mathcal{D} \otimes 1, \mathcal{S}(z)] = -\frac{d}{dz}\mathcal{S}(z) \text{ for } \mathcal{D} \in \text{End } V \text{ defined by } \mathcal{D}v = v_{-2}\mathbf{1},$$

the quantum Yang-Baxter equation

$$\mathcal{S}_{12}(z_1)\mathcal{S}_{13}(z_1+z_2)\mathcal{S}_{23}(z_2) = \mathcal{S}_{23}(z_2)\mathcal{S}_{13}(z_1+z_2)\mathcal{S}_{12}(z_1), \qquad (2.21)$$

the unitarity condition

$$S_{21}(z) = S^{-1}(-z),$$
 (2.22)

the *hexagon identity*

$$\mathcal{S}(z_1) (Y(z_2) \otimes 1) = (Y(z_2) \otimes 1) \mathcal{S}_{23}(z_1) \mathcal{S}_{13}(z_1 + z_2), \qquad (2.23)$$

and the *S*-locality: for any $u, v \in V$ and $n \in \mathbb{Z}_{\geq 0}$ there exists $r \in \mathbb{Z}_{\geq 0}$ such that

$$\left((z_1 - z_2)^r Y(z_1) \left(1 \otimes Y(z_2) \right) \left(\mathcal{S}(z_1 - z_2) (u \otimes v) \otimes w \right) - (z_1 - z_2)^r Y(v, z_2) Y(u, z_1) w \right) \in h^n V[[z_1^{\pm 1}, z_2^{\pm 1}]] \quad \text{for all } w \in V.$$
 (2.24)

Let m and n be positive integers, $u = (u_1, \ldots, u_n), v = (v_1, \ldots, v_m)$ families of variables and z a single variable. Label the tensor factors of $(\operatorname{End} \mathbb{C}^N)^{\otimes (n+m)}$ as follows,

$$\underbrace{\left(\operatorname{End} \mathbb{C}^{N}\right)^{\otimes n}}_{1} \otimes \underbrace{\left(\operatorname{End} \mathbb{C}^{N}\right)^{\otimes m}}_{2}.$$
(2.25)

Introduce the formal power series with coefficients in (2.25) by

$$\overline{R}_{nm}^{12}(u|v|z) = \prod_{i=1,\dots,n} \prod_{j=n+1,\dots,n+m} \overline{R}_{ij}(z+u_i-v_{j-n}), \qquad (2.26)$$

$$\overline{\underline{R}}_{nm}^{12}(u|v|z) = \prod_{i=1,\dots,n} \prod_{j=n+1,\dots,n+m} \overline{R}_{ij}(z+u_i+v_{j-n}).$$
(2.27)

In (2.26) and (2.27), the superscripts 1 and 2 indicate the tensor factors in (2.25) while the arrows indicate the order of the factors. For example, we have

$$\overline{R}_{22}^{12}(u|v|z) = \overline{R}_{14}\overline{R}_{13}\overline{R}_{24}\overline{R}_{23} \quad \text{and} \quad \underline{\overline{R}}_{22}^{12}(u|v|z) = \underline{\overline{R}}_{13}\overline{R}_{14}\overline{R}_{23}\overline{R}_{24}$$

for $\overline{R}_{ij} = \overline{R}_{ij}(z+u_i-v_{j-n})$ and $\overline{R}_{ij} = \overline{R}_{ij}(z+u_i+v_{j-n})$. In order to simplify the notation, we write $\overline{R}_{nm}^{12}(u|v) = \overline{R}_{nm}^{12}(u|v|0)$. Furthermore, we shall use the notation

$$T_{[n]}^{\pm}(u) = T_1^{\pm}(u_1) \dots T_n^{\pm}(u_n)$$
 and $T_{[n]}^{\pm}(u|z) = T_1^{\pm}(z+u_1) \dots T_n^{\pm}(z+u_n),$ (2.28)

where the matrix entries of the coefficients of (2.28) are regarded as operators on the vacuum module $\mathcal{V}^{c}(\mathfrak{g}_{N})$. Hence, the coefficients of the expression $T^{\pm}_{[n]}(u|z)$ belong to $(\operatorname{End} \mathbb{C}^N)^{\otimes n} \otimes \operatorname{End} \mathcal{V}^c(\mathfrak{g}_N)$. Finally, we recall the Etingof-Kazhdan construction [9, Thm. 2.3]; see also [7, Thm. 2.2] and [18, Thm. 4.1].

Theorem 2.2. For any $c \in \mathbb{C}$ there exists a unique structure of h-adic quantum vertex algebra on $\mathcal{V}^{c}(\mathfrak{g}_{N})$ such that the vacuum vector is $\mathbf{1} \in \mathcal{V}^{c}(\mathfrak{g}_{N})$, the vertex operator map $Y(\cdot, z)$ is defined by

$$Y(T^{+}_{[n]}(u)\mathbf{1}, z) = T^{+}_{[n]}(u|z)T^{-}_{[n]}(u|z+h\varepsilon c/2)^{-1}$$
(2.29)

and the braiding \mathcal{S} satisfies the identities on $(\operatorname{End} \mathbb{C}^N)^{\otimes n} \otimes (\operatorname{End} \mathbb{C}^N)^{\otimes m} \otimes \mathcal{V}^c(\mathfrak{g}_N) \otimes \mathcal{V}^c(\mathfrak{g}_N)$,

$$S(z) \left(T_{[m]}^{+24}(v) \overline{R}_{nm}^{12}(u|v|z - h\varepsilon c) T_{[n]}^{+13}(u) \overline{R}_{nm}^{12}(u|v|z)^{-1} (\mathbf{1} \otimes \mathbf{1}) \right)$$

= $\overline{R}_{nm}^{12}(u|v|z) T_{[n]}^{+13}(u) \overline{R}_{nm}^{12}(u|v|z + h\varepsilon c)^{-1} T_{[m]}^{+24}(v) (\mathbf{1} \otimes \mathbf{1}).$ (2.30)

Consider the h-adic quantum affine vertex algebras from Theorem 2.2 at the *critical level* c_{crit} , where $c_{\text{crit}} = -N$ for $\mathfrak{g}_N = \mathfrak{gl}_N$, while for $\mathfrak{g}_N = \mathfrak{o}_N$ (resp. $\mathfrak{g}_N = \mathfrak{sp}_N$) we have $c_{\text{crit}} = -N + 2$ (resp. $c_{\text{crit}} = -\frac{N}{2} - 1$). To simplify the notation, we shall denote $\mathcal{V}^{c_{\text{crit}}}(\mathfrak{g}_N)$ by $\mathcal{V}^{\text{crit}}(\mathfrak{g}_N)$. By [7, Thm. 2.4] and [18, Thm. 4.9], all coefficients of the series

$$\mathbb{T}^{+}_{[n]}(u) = \operatorname{tr}_{1,\dots,n} \mathcal{E}_{[n]} T^{+}_{[n]}(u_{[n]}) \mathbf{1} \in \mathcal{V}^{\operatorname{crit}}(\mathfrak{g}_{N})[[u]], \qquad (2.31)$$

where n = 1, ..., N (resp. n = 1, ..., N/2) for $\mathfrak{g}_N = \mathfrak{gl}_N, \mathfrak{o}_N$ (resp. $\mathfrak{g}_N = \mathfrak{sp}_N$), belong to the *center*

$$\mathfrak{z}(\mathcal{V}^{\operatorname{crit}}(\mathfrak{g}_N)) = \left\{ w \in \mathcal{V}^{\operatorname{crit}}(\mathfrak{g}_N) : Y(v, z) w \in \mathcal{V}^{\operatorname{crit}}(\mathfrak{g}_N)[[z]] \text{ for all } v \in \mathcal{V}^{\operatorname{crit}}(\mathfrak{g}_N) \right\}$$

of $\mathcal{V}^{\text{crit}}(\mathfrak{g}_N)$; see [18, Subsect. 3.2] for more information on the notion of center of *h*-adic quantum vertex algebra. Furthermore, by [7, Thm. 2.8] and [18, Thm. 4.9] we have

Theorem 2.3. The center $\mathfrak{z}(\mathcal{V}^{\operatorname{crit}}(\mathfrak{g}_N))$ for $\mathfrak{g}_N = \mathfrak{gl}_N$ and for $\mathfrak{g}_N = \mathfrak{o}_N$ with N odd (i.e. for \mathfrak{g}_N of type A and B) is a commutative associative algebra with respect to the multiplication $a \cdot b = a_{-1}b$. It is topologically generated by the coefficients of $\mathbb{T}^+_{[n]}(u)$, $n = 1, \ldots, N$.

We remark that for $\mathfrak{g}_N = \mathfrak{o}_N, \mathfrak{sp}_N$ with N even (i.e. for \mathfrak{g}_N of type C, D) the coefficients of $\mathbb{T}^+_{[n]}(u), n = 1, \ldots, N/2$ again topologically generate a commutative associative algebra with respect to the multiplication $a \cdot b = a_{-1}b$. However, in this case, such an algebra does not exhaust the entire center $\mathfrak{z}(\mathcal{V}^{\operatorname{crit}}(\mathfrak{g}_N))$; see [7, Cor 2.10].

3. Deformed modules for h-adic quantum vertex algebras

The goal of this section is to present the notion of deformed module. The next definition is motivated by the properties (2.21) and (2.22) of the braiding. Although it envolves an h-adic quantum vertex algebra, it does not depend on its structure, so, alternatively, it can be given over any $\mathbb{C}[[h]]$ -module.

Definition 3.1. Let $(V, Y, \mathbf{1}, S)$ be an *h*-adic quantum vertex algebra. The map

$$\mu(z,x)\colon V\otimes V\to V\otimes V\otimes \mathbb{C}[z^{\pm 1}][[x^{-1},h]]$$

is said to be a braiding map on V, if it satisfies the quantum Yang-Baxter equation

$$\mu_{12}(z_1, x + z_2)\mu_{13}(z_1 + z_2, x)\mu_{23}(z_2, x) = \mu_{23}(z_2, x)\mu_{13}(z_1 + z_2, x)\mu_{12}(z_1, x + z_2), \quad (3.1)$$

the unitarity condition

$$\mu_{21}(z, x - z) = \mu^{-1}(-z, x) \tag{3.2}$$

and the singularity constraint

$$\mu(-x+z,x)(a\otimes b) \in V \otimes V \otimes \mathbb{C}[z][[x^{-1},h]] \text{ for all } a,b \in V.$$
(3.3)

Remark 3.2. Roughly speaking, the singularity constraint (3.3) ensures that the map $\mu = \mu(z, x)$ does not possess a singularity at z = -x. For example, the series

$$m(z,x) = \frac{1}{x+z} = \sum_{k \ge 0} (-1)^k \frac{z^k}{x^{k+1}} \in \mathbb{C}[z][[x^{-1}]] \subset \mathbb{C}[z^{\pm 1}][[x^{-1},h]]$$

does not satisfy such a requirement as $m(-x+z,x) = z^{-1} \notin \mathbb{C}[z][[x^{-1}]]$. We explain the role of this property after Definition 3.5 below.

Let $(V, Y, \mathbf{1}, S)$ be an *h*-adic quantum vertex algebra and

$$\nu(z,x)\colon V\otimes V\to V\otimes V\otimes \mathbb{C}[z^{\pm 1}][[x^{-1},h]]$$
(3.4)

an arbitrary map. By (2.20) and (3.4), the composition $Y(z)\nu(z,x)$ is well-defined and its image belongs to $V((z))[[x^{-1},h]]$. To simplify the notation, we shall denote this composition by $Y^{\nu}(z,x) = Y^{\nu}(\cdot,z,x)$. In what follows, we want to consider the braiding maps over *h*-adic quantum vertex algebras which differ from the original braiding \mathcal{S} . They will need to satisfy certain compatibility conditions, related to the structure of V, as follows.

Definition 3.3. Let $(V, Y, \mathbf{1}, S)$ be an *h*-adic quantum vertex algebra. A braiding map $\mu = \mu(z, x)$ on V is said to be *compatible* if there exists a map (the *intertwining map*) $\nu(z, x)$ as in (3.4) which satisfies

$$S(z)\nu_{21}(-z, x+z) = \nu(z, x)\mu(z, x), \qquad (3.5)$$

$$Y^{\nu}(a, z, x)b \in V[x^{-1}][[z^{\pm 1}, h]] \text{ for all } a, b \in V.$$
(3.6)

Furthermore, the pair (μ, ν) is then said to be a *compatible pair*.

We discuss the role of the requirement (3.6) after Definition 3.5 below.

Example 3.4. One easily deduces from Definition 3.3 that the original braiding S of the *h*-adic quantum vertex algebra $(V, Y, \mathbf{1}, S)$ is compatible. Indeed, the corresponding intertwining map is the embedding $\iota: v_1 \otimes v_2 \mapsto v_1 \otimes v_2 \otimes 1$. Clearly, the both maps of the compatible pair (S, ι) are constant with respect to the second variable x.

Our next goal is to introduce (μ, ν) -deformed modules. In comparison with the notion of *h*-adic quantum vertex algebra, they employ a slightly more general topological setting. Let *W* be a $\mathbb{C}[[h]]$ -module such that there exists a family of $\mathbb{C}[[h]]$ -submodules

$$W = W_0 \supseteq W_1 \supseteq W_2 \supseteq \dots \qquad \text{satisfying} \qquad h^n W \subseteq W_n \quad \text{for all } n \ge 0. \tag{3.7}$$

We shall consider the linear topology τ on W defined by the basis

$$w + W_n$$
 for all $n \ge 0, w \in W$. (3.8)

Denote by $W((z^{-1}))_{\tau}$ the $\mathbb{C}[[h]]$ -module of all formal power series

$$a(z) = \sum_{r \in \mathbb{Z}} a_r z^r \in W[[z^{\pm 1}]]$$
 such that $\lim_{r \to \infty} a_r = 0$,

where the limit is taken with respect to the topology τ . Such a notation naturally extends to the multiple variable case. For example, $a(z_1, z_2) \in W((z_1^{-1}, z_2^{-1}))_{\tau}$ means that the series $a = a(z_1, z_2)$ belongs to $W[[z_1^{\pm 1}, z_2^{\pm 1}]]$ and that for every $n \in \mathbb{Z}_{\geq 0}$ it decomposes as a = b + c for some $b = b(z_1, z_2) \in W((z_1^{-1}, z_2^{-1}))$ and $c = c(z_1, z_2) \in W_n[[z_1, z_2]]$.

Definition 3.5. Let $(V, Y, \mathbf{1}, \mathcal{S})$ be an *h*-adic quantum vertex algebra equipped with a compatible pair (μ, ν) . Let W be a $\mathbb{C}[[h]]$ -module equipped with a descending family of submodules as in (3.7) such that it is complete with respect to the topology τ defined by the basis (3.8). Let $Y_W(\cdot, z)$ be a $\mathbb{C}[[h]]$ -module map

$$Y_W(\cdot, z) \colon V \otimes W \to W((z^{-1}))_{\tau},$$
$$v \otimes w \mapsto Y(z)(v \otimes w) = Y(v, z)w = \sum_{r \in \mathbb{Z}} v_r w z^{-r-1}$$

A pair (W, Y_W) is said to be a (μ, ν) -deformed V-module if the map $Y_W(\cdot, z)$ satisfies the vacuum property

$$Y_W(\mathbf{1}, z)w = w \quad \text{for all } w \in W, \tag{3.9}$$

the truncation condition: for any $u, v \in V$ and $w \in W$ we have

$$Y_W(u, z_1) Y_W(v, z_2) w \in W((z_1^{-1}, z_2^{-1}))_{\tau},$$
(3.10)

the ν -associativity: for any elements $u, v \in V$ and $w \in W$ we have

$$Y_W(u, z_2 + z_0) Y_W(v, z_2) w = Y_W (Y^{\nu}(u, z_0, z_2) v, z_2) w, \qquad (3.11)$$

and the μ -commutativity: for any $u, v \in V$ and $w \in W$ we have

$$Y_W(z_1) (1 \otimes Y_W(z_2)) (\mu(-z_2 + z_1, z_2)(u \otimes v) \otimes w) = Y_W(v, z_2) Y_W(u, z_1) w.$$
(3.12)

Regarding Definition 3.5, the expression on the left-hand side of ν -associativity (3.11) is well-defined due to the truncation condition (3.10), while the right-hand side is well-defined due to (3.6). Furthermore, the singularity constraint (3.3) and the truncation condition (3.10) ensure that the left-hand side of μ -commutativity (3.12) is well-defined.

Definition 3.6. Let $(V, Y, \mathbf{1}, S)$ be an *h*-adic quantum vertex algebra equipped with compatible pairs (μ, ν) and (σ, ρ) . The pairs (μ, ν) and (σ, ρ) are said to be *equivalent* if there exist maps

$$\gamma^{\pm 1}(z,x) \colon V \otimes V \to V \otimes V \otimes \mathbb{C}[z,z^{-1}][[x^{-1},h]],$$

such that $\gamma(z, x)\gamma^{-1}(z, x) = \gamma^{-1}(z, x)\gamma(z, x) = 1$, which satisfy the *shift condition*

$$\gamma^{\pm 1}(z, x) = \gamma^{\pm 1}(-z, x+z) \tag{3.13}$$

and the equalities

 $\mu(z,x) = \gamma^{-1}(z,x)\sigma(z,x)\gamma_{21}(z,x) \quad \text{and} \quad \nu(z,x) = \rho(z,x)\gamma(z,x). \quad (3.14)$

Remark 3.7. The next two simple observations will be useful later on. The unitarity condition (3.2) for the map σ and the shift condition (3.13) for the map γ imply that the map μ , defined by the first identity in (3.14), also satisfies the unitarity (3.2). Moreover, the property (3.5) for the pair (σ , ρ) and the shift condition (3.13) for the map γ imply that the map ν , defined by the second identity in (3.14), also possesses the property (3.5).

4. Compatible braiding maps over h-adic quantum affine vertex algebras

In this section, we construct two compatible pairs over the h-adic quantum vertex algebras from Theorem 2.2 and investigate their properties.

4.1. Compatible pair (σ, ρ) over $\mathcal{V}^c(\mathfrak{g}_N)$. The goal of this subsection is to construct an example of a compatible braiding map on $\mathcal{V}^c(\mathfrak{g}_N)$ which differs from (2.30). The candidate is the map $\sigma = \sigma(z, x)$ from the following lemma, which is constant with respect to the second variable x; cf. Definition 3.1.

Lemma 4.1. There exists a unique map

$$\sigma(z)\colon \mathcal{V}^c(\mathfrak{g}_N)\otimes \mathcal{V}^c(\mathfrak{g}_N)\to \mathcal{V}^c(\mathfrak{g}_N)\otimes \mathcal{V}^c(\mathfrak{g}_N)\otimes \mathbb{C}[[h]][z^{-1}]_h$$

such that on $(\operatorname{End} \mathbb{C}^N)^{\otimes n} \otimes (\operatorname{End} \mathbb{C}^N)^{\otimes m} \otimes \mathcal{V}^c(\mathfrak{g}_N) \otimes \mathcal{V}^c(\mathfrak{g}_N)$ we have

$$\sigma(z) \left(T_{[n]}^{+13}(u) T_{[m]}^{+24}(v) (\mathbf{1} \otimes \mathbf{1}) \right) = \overline{R}_{nm}^{12}(u|v|z) T_{[n]}^{+13}(u) T_{[m]}^{+24}(v) \overline{R}_{nm}^{12}(u|v|z)^{-1} (\mathbf{1} \otimes \mathbf{1}).$$
(4.1)

Moreover, the map σ is a braiding map on $\mathcal{V}^c(\mathfrak{g}_N)$.

Proof. Lemmas 4.1, 4.2 and Proposition 4.3 can be verified by straightforward calculations. They are carried out in parallel to the proofs of [7, Thm. 2.2], [12, Thm. 2.3.8] and [18, Thm. 4.1] and they employ properties (2.3)–(2.6) of the *R*-matrix and the defining relations (2.10), (2.11) and (2.14). Therefore, we omit the proofs of the lemmas and only sketch the proof of Proposition 4.3.

In order to show that σ is compatible, we need to find its intertwining map.

Lemma 4.2. There exists a unique map

$$\rho(z)\colon \mathcal{V}^c(\mathfrak{g}_N)\otimes\mathcal{V}^c(\mathfrak{g}_N)\to\mathcal{V}^c(\mathfrak{g}_N)\otimes\mathcal{V}^c(\mathfrak{g}_N)\otimes\mathbb{C}[[h]][z^{-1}]_h$$

such that on $(\operatorname{End} \mathbb{C}^N)^{\otimes n} \otimes (\operatorname{End} \mathbb{C}^N)^{\otimes m} \otimes \mathcal{V}^c(\mathfrak{g}_N) \otimes \mathcal{V}^c(\mathfrak{g}_N)$ we have

$$\rho(z) \Big(T_{[n]}^{+13}(u) T_{[m]}^{+24}(v) (\mathbf{1} \otimes \mathbf{1}) \Big) = T_{[n]}^{+13}(u) \overline{R}_{nm}^{12}(u|v|z + h\varepsilon c)^{-1} T_{[m]}^{+24}(v) \overline{R}_{nm}^{12}(u|v|z) (\mathbf{1} \otimes \mathbf{1}).$$

$$(4.2)$$

Finally, it remains to show that the maps σ and ρ satisfy Definition 3.3.

Proposition 4.3. The maps σ and ρ form a compatible pair on $\mathcal{V}^{c}(\mathfrak{g}_{N})$.

Proof. In order to verify (3.5), it is useful to observe that the explicit formula for the action of $\rho_{43}(-z)$ on $(\operatorname{End} \mathbb{C}^N)^{\otimes n} \otimes (\operatorname{End} \mathbb{C}^N)^{\otimes m} \otimes \mathcal{V}^c(\mathfrak{g}_N) \otimes \mathcal{V}^c(\mathfrak{g}_N)$ is given by

$$\rho_{43}(-z) \Big(T^{+13}_{[n]}(u) T^{+24}_{[m]}(v) (\mathbf{1} \otimes \mathbf{1}) \Big) = T^{+24}_{[m]}(v) \overline{R}^{12}_{nm}(u|v|z - h\varepsilon c) T^{+13}_{[n]}(u) \overline{R}^{12}_{nm}(u|v|z)^{-1} (\mathbf{1} \otimes \mathbf{1}).$$

By applying the braiding S(z) on the above equality, one finds that the image of the expression $T_{[n]}^{+13}(u)T_{[m]}^{+24}(v)(\mathbf{1}\otimes\mathbf{1})$ under the composition $S(z)\rho_{43}(-z)$ coincides with the right-hand side of (2.30). On the other hand, by using formulae (4.1) and (4.2), one easily checks that the image of the same expression under the composition $\rho(z)\sigma(z)$ coincides with the right-hand side of (2.30) as well. Thus, we conclude that (3.5) holds. The remaining property (3.6) obviously holds as the map $\rho = \rho(z, x)$ is constant with respect to the second variable x.

The maps σ and ρ satisfy the following *hexagon relations*; cf. (2.23).

Proposition 4.4. We have

$$\rho(z_1)\left(Y^{\rho}(z_2)\otimes 1\right) = \left(Y^{\rho}(z_2)\otimes 1\right)\rho_{13}(z_1+z_2)\rho_{23}(z_1),\tag{4.3}$$

$$\sigma(z_1) \left(Y^{\rho}(z_2) \otimes 1 \right) = \left(Y^{\rho}(z_2) \otimes 1 \right) \sigma_{23}(z_1) \sigma_{13}(z_1 + z_2).$$
(4.4)

Proof. Let us prove (4.3). First, we observe that the map Y^{ρ} satisfies the identity

$$Y^{\rho}(T^{+13}_{[n]}(u), z_2)T^{+24}_{[m]}(v)(\mathbf{1} \otimes \mathbf{1}) = T^{+13}_{[n]}(u|z_2)T^{+24}_{[m]}(v)(\mathbf{1} \otimes \mathbf{1}).$$
(4.5)

Indeed, this is easily verified by the use of defining relation (2.11) for the double Yangian and formula (2.29) for the vertex operator map. Clearly, (4.5) implies

$$Y^{\rho}(T_{[n]}^{+14}(u), z_2)T_{[m]}^{+25}(v)T_{[k]}^{+36}(w)(\mathbf{1}\otimes\mathbf{1}\otimes\mathbf{1}) = T_{[n]}^{+14}(u|z_2)T_{[m]}^{+24}(v)T_{[k]}^{+35}(w)(\mathbf{1}\otimes\mathbf{1}\otimes\mathbf{1}).$$
(4.6)

By applying $\rho(z_1)$ to (4.6) and then using (4.2), we find that $\rho(z_1) (Y^{\rho}(z_2) \otimes 1)$ maps

$$T_{[n]}^{+14}(u)T_{[m]}^{+25}(v)T_{[k]}^{+36}(w)(\mathbf{1}\otimes\mathbf{1}\otimes\mathbf{1})$$
(4.7)

to

$$T_{[n]}^{+14}(u|z_2)T_{[m]}^{+24}(v)\overline{R}_{mk}^{23}(v|w|z_1+h\varepsilon c)^{-1}\overline{R}_{nk}^{13}(u|w|z_1+z_2+h\varepsilon c)^{-1} \times T_{[k]}^{+35}(w)\overline{R}_{nk}^{13}(u|w|z_1+z_2)\overline{R}_{mk}^{23}(v|w|z_1)(\mathbf{1}\otimes\mathbf{1}\otimes\mathbf{1}),$$
(4.8)

where the superscripts 1-6 in (4.7) indicate the tensor factors as follows:

$$\underbrace{\left(\operatorname{End} \mathbb{C}^{N}\right)^{\otimes n}}_{\left(\operatorname{End} \mathbb{C}^{N}\right)^{\otimes m}} \otimes \underbrace{\left(\operatorname{End} \mathbb{C}^{N}\right)^{\otimes k}}_{\left(\operatorname{End} \mathbb{C}^{N}\right)^{\otimes k}} \otimes \underbrace{\mathcal{V}^{c}(\mathfrak{g}_{N})}_{\mathcal{V}^{c}(\mathfrak{g}_{N})} \otimes \underbrace{\mathcal{V}^{c}(\mathfrak{g}_{N})}_{\mathcal{V}^{c}(\mathfrak{g}$$

Finally, by using (4.2) and then (4.6), one shows that the image of (4.7) under the composition $(Y^{\rho}(z_2) \otimes 1) \rho_{13}(z_1+z_2) \rho_{23}(z_1)$ equals (4.8) as well, thus proving the hexagon relation (4.3). The remaining relation (4.4) for σ can be verified by analogous arguments which rely on (4.1) and (4.6).

4.2. Compatible pair (μ, ν) over $\mathcal{V}^{\text{crit}}(\mathfrak{gl}_N)$. The goal of this subsection is to construct another compatible pair (μ, ν) on $\mathcal{V}^{\text{crit}}(\mathfrak{gl}_N)$, which is equivalent to (σ, ρ) . We start by fixing the auxiliary map γ from Definition 3.6.

Lemma 4.5. There exists a unique map

$$\gamma(z,x)\colon \mathcal{V}^{\operatorname{crit}}(\mathfrak{gl}_N) \otimes \mathcal{V}^{\operatorname{crit}}(\mathfrak{gl}_N) \to \mathcal{V}^{\operatorname{crit}}(\mathfrak{gl}_N) \otimes \mathcal{V}^{\operatorname{crit}}(\mathfrak{gl}_N) \otimes \mathbb{C}[z^{\pm 1}][[x^{-1},h]]$$

such that on $(\operatorname{End} \mathbb{C}^N)^{\otimes n} \otimes (\operatorname{End} \mathbb{C}^N)^{\otimes m} \otimes \mathcal{V}^{\operatorname{crit}}(\mathfrak{gl}_N) \otimes \mathcal{V}^{\operatorname{crit}}(\mathfrak{gl}_N)$ we have

$$\gamma(z,x) \left(T_{[n]}^{+13}(u) T_{[m]}^{+24}(v) (\mathbf{1} \otimes \mathbf{1}) \right) = T_{[n]}^{+13}(u) \overline{\underline{R}}_{nm}^{12}(u|v|2x+z-hN)^{-1} T_{[m]}^{+24}(v) (\mathbf{1} \otimes \mathbf{1}).$$
(4.10)

Its inverse is given by

$$\gamma^{-1}(z,x) \Big(T^{+13}_{[n]}(u) T^{+24}_{[m]}(v) (\mathbf{1} \otimes \mathbf{1}) \Big) = T^{+13}_{[n]}(u) \overline{\underline{R}}^{12}_{nm}(u|v|2x+z) T^{+24}_{[m]}(v) (\mathbf{1} \otimes \mathbf{1})$$
(4.11)

and both maps satisfy the shift condition (3.13).

Proof. One can show that the maps γ and γ^{-1} are well-defined by arguing as in [9, Lemma 2.1]. The expression for the inverse (4.11) is easily verified by using the crossing symmetry (2.7). Finally, the last assertion of the lemma is evident from (4.10) and (4.11).

Motivated by the first identity in (3.14), we define the map μ by

$$\mu(z,x) = \gamma^{-1}(z,x)\sigma(z)\gamma_{21}(z,x).$$
(4.12)

Lemmas 4.1 and 4.5 imply that its image belongs to $\mathcal{V}^{\text{crit}}(\mathfrak{gl}_N)^{\otimes 2} \otimes \mathbb{C}[z^{\pm 1}][[x^{-1}, h]]$. In fact, one can derive from these lemmas the explicit expression for the action of μ ,

$$\mu(z,x)\Big(T_{[n]}^{+13}(u)T_{[m]}^{+24}(v)(\mathbf{1}\otimes\mathbf{1})\Big) = \overline{\underline{R}}_{nm}^{12}(u|v|2x+z-hN)^{-1} \underset{\mathrm{LR}}{\cdot} \Big(\overline{R}_{nm}^{12}(u|v|z)T_{[n]}^{+13}(u) \\ \times \overline{\underline{R}}_{nm}^{12}(u|v|2x+z)T_{[m]}^{+24}(v)\overline{R}_{nm}^{12}(u|v|z)^{-1}\Big)(\mathbf{1}\otimes\mathbf{1}), \quad (4.13)$$

where " \cdot " denotes the standard multiplication in $(\operatorname{End} \mathbb{C}^N)^{\otimes n} \otimes ((\operatorname{End} \mathbb{C}^N)^{\operatorname{op}})^{\otimes m}$.

Lemma 4.6. The map μ is a braiding map on $\mathcal{V}^{\operatorname{crit}}(\mathfrak{gl}_N)$.

Proof. Recall Definition 3.1. First, note that, due to Remark 3.7, μ possesses unitarity property (3.2). Next, it is clear from (4.13) that μ satisfies the singularity condition (3.3). Hence, it remains to show that the quantum Yang–Baxter equation (3.1) holds. Its proof employs the alternative expression for the action of μ ,

$$\mu(z,x) \left(T_{[m]}^{+24}(v) \overline{\underline{R}}_{nm}^{12}(u|v|2x+z) T_{[n]}^{+13}(u) \overline{R}_{nm}^{12}(u|v|z) (\mathbf{1} \otimes \mathbf{1}) \right)$$

= $\overline{R}_{nm}^{12}(u|v|z) T_{[n]}^{+13}(u) \overline{\underline{R}}_{nm}^{12}(u|v|2x+z) T_{[m]}^{+24}(v) (\mathbf{1} \otimes \mathbf{1}),$ (4.14)

which is obtained from (4.13) by the use of crossing symmetry (2.7). It is sufficient to check that the images of

$$T_{[k]}^{+36}(w)\overline{\underline{R}}_{mk}^{23}T_{[m]}^{+25}(v)\overline{R}_{mk}^{23}\overline{\underline{R}}_{nm}^{12}\overline{\underline{R}}_{nk}^{13}T_{[n]}^{+14}(u)\overline{R}_{nk}^{13}\overline{R}_{nm}^{12}(\mathbf{1}\otimes\mathbf{1}\otimes\mathbf{1})$$
(4.15)

under both sides of (3.1) coincide. In (4.15), we use the notation

$$\overline{R}_{nm}^{12} = \overline{R}_{nm}^{12}(u|v|z_1), \qquad \overline{\underline{R}}_{nm}^{12} = \underline{\overline{R}}_{nm}^{12}(u|v|2x + 2z_2 + z_1), \qquad \overline{R}_{nk}^{13} = \overline{R}_{nk}^{13}(u|w|z_1 + z_2), \\
\underline{\overline{R}}_{nk}^{13} = \underline{\overline{R}}_{nk}^{13}(u|w|2x + z_1 + z_2), \qquad \overline{R}_{mk}^{23} = \overline{R}_{mk}^{23}(v|w|z_2), \qquad \underline{\overline{R}}_{mk}^{23} = \underline{\overline{R}}_{mk}^{23}(v|w|z_2 + z_2)$$

and the meaning of superscripts 1–6 is analogous to (4.9) (except that the three copies of $\mathcal{V}^{c}(\mathfrak{g}_{N})$ are replaced by $\mathcal{V}^{\text{crit}}(\mathfrak{gl}_{N})$).

Let us consider the action of the left-hand side of (3.1) on (4.15). By applying $\mu_{23}(z_2, x)$ to (4.15) and using the fact that $T_{[k]}^{+36}(w)$ and $\overline{\underline{R}}_{nm}^{12}$ commute, we get

$$\overline{R}_{mk}^{23} T_{[m]}^{+25}(v) \overline{\underline{R}}_{mk}^{23} \overline{\underline{R}}_{nm}^{12} T_{[k]}^{+36}(w) \overline{\underline{R}}_{nk}^{13} T_{[n]}^{+14}(u) \overline{R}_{nk}^{13} \overline{R}_{nm}^{12}(\mathbf{1} \otimes \mathbf{1} \otimes \mathbf{1}).$$
(4.16)

Next, we apply $\mu_{13}(z_1 + z_2, x)$ to (4.16), thus getting

$$\overline{R}_{mk}^{23} T_{[m]}^{+25}(v) \overline{\underline{R}}_{mk}^{23} \overline{\underline{R}}_{nm}^{12} \overline{R}_{nk}^{13} T_{[n]}^{+14}(u) \overline{\underline{R}}_{nk}^{13} T_{[k]}^{+36}(w) \overline{R}_{nm}^{12}(\mathbf{1} \otimes \mathbf{1} \otimes \mathbf{1}).$$
(4.17)

By using the quantum Yang–Baxter equation (2.3) one easily verifies the identities

$$\underline{\overline{R}}_{mk}^{23} \underline{\overline{R}}_{nm}^{12} \overline{\overline{R}}_{nk}^{13} = \overline{\overline{R}}_{nk}^{13} \underline{\overline{R}}_{nm}^{12} \underline{\overline{R}}_{mk}^{23} \text{ and } \underline{\overline{R}}_{mk}^{23} \underline{\overline{R}}_{nk}^{13} \overline{\overline{R}}_{nm}^{12} = \overline{\overline{R}}_{nm}^{12} \underline{\overline{R}}_{nk}^{13} \underline{\overline{R}}_{mk}^{23}.$$

By employing these two equalities, along with commutation relations

$$\begin{split} \overline{\underline{R}}_{mk}^{23} T_{[n]}^{+14}(u) &= T_{[n]}^{+14}(u) \overline{\underline{R}}_{mk}^{23}, \qquad T_{[k]}^{+36}(w) \overline{R}_{nm}^{12} = \overline{R}_{nm}^{12} T_{[k]}^{+36}(w), \\ T_{[m]}^{+25}(v) \overline{R}_{nk}^{13} = \overline{R}_{nk}^{13} T_{[m]}^{+25}(v), \end{split}$$

we rewrite (4.17) as

$$\overline{R}_{mk}^{23} \overline{R}_{nk}^{13} T_{[m]}^{+25}(v) \underline{\overline{R}}_{nm}^{12} T_{[n]}^{+14}(u) \overline{R}_{nm}^{12} \underline{\overline{R}}_{nk}^{13} \underline{\overline{R}}_{mk}^{23} T_{[k]}^{+36}(w) (\mathbf{1} \otimes \mathbf{1} \otimes \mathbf{1}).$$
(4.18)

Finally, applying $\mu_{12}(z_1, x + z_2)$ to (4.18) we obtain

$$\overline{R}_{mk}^{23} \overline{R}_{nk}^{13} \overline{R}_{nm}^{12} T_{[n]}^{+14}(u) \underline{\overline{R}}_{nm}^{12} T_{[m]}^{+25}(v) \underline{\overline{R}}_{nk}^{13} \underline{\overline{R}}_{mk}^{23} T_{[k]}^{+36}(w) (\mathbf{1} \otimes \mathbf{1} \otimes \mathbf{1}).$$
(4.19)

Let us consider the action of the right-hand side of (3.1) on (4.15). First, we observe that the quantum Yang–Baxter equation (2.3) implies

$$\overline{R}_{mk}^{23} \overline{R}_{nm}^{12} \overline{R}_{nk}^{13} = \overline{R}_{nk}^{13} \overline{R}_{nm}^{12} \overline{R}_{mk}^{23} \text{ and } \overline{R}_{mk}^{23} \overline{R}_{nk}^{13} \overline{R}_{nm}^{12} = \overline{R}_{nm}^{12} \overline{R}_{nk}^{13} \overline{R}_{mk}^{23}.$$

By using these equalities and the commutation relations

$$T_{[m]}^{+25}(v)\overline{R}_{nk}^{13} = \overline{R}_{nk}^{13}T_{[m]}^{+25}(v) \quad \text{and} \quad \overline{R}_{mk}^{23}T_{[n]}^{+14}(u) = T_{[n]}^{+14}(u)\overline{R}_{mk}^{23},$$

we rewrite (4.15) as

$$T_{[k]}^{+36}(w)\overline{\underline{R}}_{mk}^{23}\overline{\underline{R}}_{nk}^{13}T_{[m]}^{+25}(v)\overline{\underline{R}}_{nm}^{12}T_{[n]}^{+14}(u)\overline{R}_{nm}^{12}\overline{R}_{nk}^{13}\overline{R}_{mk}^{23}(\mathbf{1}\otimes\mathbf{1}\otimes\mathbf{1}).$$
(4.20)

Next, we apply $\mu_{12}(z_1, x + z_2)$ to (4.18), thus getting

$$T_{[k]}^{+36}(w)\overline{\underline{R}}_{mk}^{23}\overline{\underline{R}}_{nk}^{13}\overline{R}_{nm}^{12}T_{[n]}^{+14}(u)\overline{\underline{R}}_{nm}^{12}T_{[m]}^{+25}(v)\overline{R}_{nk}^{13}\overline{R}_{mk}^{23}(\mathbf{1}\otimes\mathbf{1}\otimes\mathbf{1}).$$
(4.21)

We shall also need the following consequences of the Yang–Baxter equation (2.3):

$$\overline{\underline{R}}_{mk}^{23} \overline{\underline{R}}_{nk}^{13} \overline{\overline{R}}_{nm}^{12} = \overline{R}_{nm}^{12} \overline{\underline{R}}_{nk}^{13} \overline{\underline{R}}_{mk}^{23} \quad \text{and} \quad \overline{\underline{R}}_{mk}^{23} \overline{\underline{R}}_{nm}^{12} \overline{\overline{R}}_{nk}^{13} = \overline{R}_{nk}^{13} \overline{\underline{R}}_{nm}^{12} \overline{\underline{R}}_{mk}^{23}$$

By combining them with relations

$$\overline{\underline{R}}_{mk}^{23} T_{[n]}^{+14}(u) = T_{[n]}^{+14}(u) \overline{\underline{R}}_{mk}^{23}, \qquad T_{[m]}^{+25}(v) \overline{R}_{nk}^{13} = \overline{R}_{nk}^{13} T_{[m]}^{+25}(v),
T_{[k]}^{+36}(w) \overline{R}_{nm}^{12} = \overline{R}_{nm}^{12} T_{[k]}^{+36}(w),$$

we turn the expression in (4.21) to

$$\overline{R}_{nm}^{12} T_{[k]}^{+36}(w) \underline{\overline{R}}_{nk}^{13} T_{[n]}^{+14}(u) \overline{R}_{nk}^{13} \underline{\overline{R}}_{nm}^{12} \underline{\overline{R}}_{mk}^{23} T_{[m]}^{+25}(v) \overline{R}_{mk}^{23} (\mathbf{1} \otimes \mathbf{1} \otimes \mathbf{1}).$$
(4.22)

By applying $\mu_{13}(z_1 + z_2, x)$ to (4.22), we get

$$\overline{R}_{nm}^{12} \overline{R}_{nk}^{13} T_{[n]}^{+14}(u) \underline{\overline{R}}_{nk}^{13} T_{[k]}^{+36}(w) \underline{\overline{R}}_{nm}^{12} \underline{\overline{R}}_{mk}^{23} T_{[m]}^{+25}(v) \overline{R}_{mk}^{23}(\mathbf{1} \otimes \mathbf{1} \otimes \mathbf{1}).$$
(4.23)

As $T_{[k]}^{+36}(w)$ and $\overline{\underline{R}}_{nm}^{12}$ commute, the application of $\mu_{23}(z_2, x)$ to (4.23) gives us

$$\overline{R}_{nm}^{12} \overline{R}_{nk}^{13} T_{[n]}^{+14}(u) \underline{\overline{R}}_{nk}^{13} \underline{\overline{R}}_{nm}^{12} \overline{R}_{mk}^{23} T_{[m]}^{+25}(v) \underline{\overline{R}}_{mk}^{23} T_{[k]}^{+36}(w) (\mathbf{1} \otimes \mathbf{1} \otimes \mathbf{1}).$$
(4.24)

The quantum Yang–Baxter equation (2.3) implies

$$\overline{\underline{R}}_{nk}^{13} \overline{\underline{R}}_{nm}^{12} \overline{R}_{mk}^{23} = \overline{R}_{mk}^{23} \overline{\underline{R}}_{nm}^{12} \overline{\underline{R}}_{nk}^{13} \text{ and } \overline{R}_{nm}^{12} \overline{R}_{nk}^{13} \overline{R}_{mk}^{23} = \overline{R}_{mk}^{23} \overline{R}_{nk}^{13} \overline{R}_{nm}^{12}.$$

Finally, by employing these two equalities and the commutativity relations

$$T_{[n]}^{+14}(u)\overline{R}_{mk}^{23} = \overline{R}_{mk}^{23}T_{[n]}^{+14}(u) \quad \text{and} \quad \underline{\overline{R}}_{nk}^{13}T_{[m]}^{+25}(v) = T_{[m]}^{+25}(v)\underline{\overline{R}}_{nk}^{13}$$

we can rewrite (4.24) so that we get the expression in (4.19), which equals the action of the left-hand side of (3.1) on (4.15), thus concluding the proof.

Following the second identity in (3.14), we define the map ν by

$$\nu(z, x) = \rho(z)\gamma(z, x).$$

Lemmas 4.2 and 4.5 imply that its image belongs to $\mathcal{V}^{\text{crit}}(\mathfrak{gl}_N)^{\otimes 2} \otimes \mathbb{C}[z^{\pm 1}][[x^{-1},h]]$ and one can derive from these lemmas the explicit expression for the action of ν ,

$$\nu(z,x) \Big(T^{+13}_{[n]}(u) T^{+24}_{[m]}(v) (\mathbf{1} \otimes \mathbf{1}) \Big) = \underline{\overline{R}}^{12}_{nm}(u|v|2x+z-hN)^{-1} \cdot \Big(T^{+13}_{[n]}(u)$$

$$\times \overline{R}^{12}_{nm}(u|v|z-hN)^{-1} T^{+24}_{[m]}(v) \overline{R}^{12}_{nm}(u|v|z) \Big) (\mathbf{1} \otimes \mathbf{1}),$$
(4.25)

where " \cdot " denotes the standard multiplication in $((\operatorname{End} \mathbb{C}^N)^{\operatorname{op}})^{\otimes n} \otimes (\operatorname{End} \mathbb{C}^N)^{\otimes m}$.

Proposition 4.7. The maps μ and ν form a compatible pair on $\mathcal{V}^{\text{crit}}(\mathfrak{gl}_N)$. Moreover, the pairs (μ, ν) and (σ, ρ) of $\mathcal{V}^{\text{crit}}(\mathfrak{gl}_N)$ are equivalent.

Proof. First, we observe that ν is an intertwining map, so that μ is compatible; recall Definition 3.3. Indeed, property (3.5) holds due to Remark 3.7, while (3.6) is evident from the expressions (2.29) and (4.25) for the action of Y and ν . In fact, using these formulae one easily derives the explicit formula for the action of Y^{ν} ,

$$Y^{\nu}(z,x) \left(T^{+13}_{[n]}(u) T^{+24}_{[m]}(v) (\mathbf{1} \otimes \mathbf{1}) \right) = \underline{\overline{R}}^{12}_{nm}(u|v|2x+z-hN)^{-1} \cdot \left(T^{+13}_{[n]}(u|z) T^{+24}_{[m]}(v) \right) (\mathbf{1} \otimes \mathbf{1}), \qquad (4.26)$$

where "." has the same meaning as in (4.25). Hence, (μ, ν) is a compatible pair. As the shift condition (3.13) is established in Lemma 4.5 and both identities in (3.14) clearly hold, we conclude that the pairs (μ, ν) and (σ, ρ) are equivalent.

Finally, we observe that the map γ satisfies the following *hexagon relation*.

Proposition 4.8. We have

$$\gamma(z_1, x) \left(Y^{\rho}(z_2) \otimes 1 \right) = \left(Y^{\rho}(z_2) \otimes 1 \right) \gamma_{13}(z_1 + z_2, x) \gamma_{23}(z_1, x).$$
(4.27)

Proof. It is sufficient to check that the images of (4.7) under both sides of (4.27) coincide. Of course, the superscripts 4–6 in (4.7) now represent the corresponding copies of $\mathcal{V}^{\text{crit}}(\mathfrak{gl}_N)$; cf. (4.9). Regarding the left-hand side, by using (4.5) and (4.10), we find that the image of (4.7) under $\gamma(z_1, x)$ ($Y^{\rho}(z_2) \otimes 1$) equals

$$T_{[n]}^{+14}(u|z_2)T_{[m]}^{+24}(v)\overline{\underline{R}}_{mk}^{23}(v|w|2x+z_1-hN)^{-1}\overline{\underline{R}}_{nk}^{13}(u|w|2x+z_1+z_2-hN)^{-1}T_{[k]}^{+35}(w)(\mathbf{1}\otimes\mathbf{1}).$$
(4.28)

On the other hand, by (4.10), the image of (4.7) under $\gamma_{13}(z_1+z_2,x)\gamma_{23}(z_1,x)$ is equal to $T_{[n]}^{+14}(u)T_{[m]}^{+25}(v)\overline{\underline{R}}_{mk}^{23}(v|w|2x+z_1-hN)^{-1}\overline{\underline{R}}_{nk}^{13}(u|w|2x+z_1+z_2-hN)^{-1}T_{[k]}^{+36}(w)(\mathbf{1}\otimes\mathbf{1}\otimes\mathbf{1}).$

Finally, (4.5) implies that
$$Y^{\rho}(\cdot, z_2) \otimes 1$$
 maps this expression to (4.28), as required. \Box

4.3. Action of the braiding maps on the quantum Feigin–Frenkel center. In this subsection, we investigate fixed points of the compatible pairs (σ, ρ) and (μ, ν) . Our key results are Lemmas 4.9 and 4.13, which we utilize in Sections 5 and 6 below. From now on, we shall write bar over the number in the superscript to indicate that the corresponding R-matrix product appears in the reversed order, e.g.,

$$\overline{R}_{nm}^{\overline{1}2}(u|v|z) = \prod_{i=1,\dots,n} \stackrel{\longleftarrow}{\prod} \prod_{j=n+1,\dots,n+m} \overline{R}_{ij}, \qquad \overline{R}_{nm}^{1\overline{2}}(u|v|z) = \prod_{i=1,\dots,n} \stackrel{\longrightarrow}{\prod} \prod_{j=n+1,\dots,n+m} \overline{R}_{ij},$$

$$\overline{R}_{nm}^{\overline{12}}(u|v|z) = \prod_{i=1,\dots,n}^{\longleftarrow} \prod_{j=n+1,\dots,n+m}^{\longrightarrow} \overline{R}_{ij}, \qquad \text{where } \overline{R}_{ij} = \overline{R}_{ij}(z+u_i-v_{j-n});$$

recall (2.26). We apply the same convention to the *R*-matrix products defined by (2.27). Also, we denote the reversed products of the operator matrices from (2.28) by

$$\dot{T}_{[n]}^{\pm 12}(u) = T_n^{\pm}(u_n) \dots T_1^{\pm}(u_1)$$
 and $\dot{T}_{[n]}^{\pm 12}(u|z) = T_n^{\pm}(z+u_n) \dots T_1^{\pm}(z+u_1).$

Finally, for any m and n we shall consider the tensor products of the series (2.31),

$$\mathbb{T}^+_{[n,m]}(u,v) = \mathbb{T}^+_{[n]}(u) \otimes \mathbb{T}^+_{[m]}(v) \in \mathcal{V}^{\operatorname{crit}}(\mathfrak{g}_N) \otimes \mathcal{V}^{\operatorname{crit}}(\mathfrak{g}_N)[[u,v]].$$

Consider the maps σ and ρ introduced in Lemmas 4.1 and 4.2, respectively.

Lemma 4.9. (a) The following identities hold in $\mathcal{V}^{\operatorname{crit}}(\mathfrak{g}_N)$:

$$\sigma(z)\Big(\mathbb{T}^+_{[n,m]}(u,v)\Big) = \rho(z)\Big(\mathbb{T}^+_{[n,m]}(u,v)\Big) = \mathbb{T}^+_{[n,m]}(u,v) \quad \text{for any } m \text{ and } n.$$
(4.29)

(b) The following identities hold in $\mathcal{V}^{\operatorname{crit}}(\mathfrak{gl}_N)$:

$$\sigma(z)\Big(w\otimes\mathbb{T}^+_{[N]}(u)\Big) = \rho(z)\Big(w\otimes\mathbb{T}^+_{[N]}(u)\Big) = w\otimes\mathbb{T}^+_{[N]}(u) \quad \text{for any } w\in\mathcal{V}^{\text{crit}}(\mathfrak{gl}_N).$$
(4.30)

Proof. Before we start with the proof, we introduce some new notation. Let

$$\overline{R}_{nm}^{12} = \overline{R}_{nm}^{12}(u_{[n]}|v_{[m]}|z) \quad \text{and} \quad \mathcal{E}_{[n,m]} = \mathcal{E}_{[n]} \otimes \mathcal{E}_{[m]},$$

where $u_{[n]}$ (resp. $\mathcal{E}_{[n]}$) are given by (2.16) and (2.18) (resp. (2.17) and (2.19)), depending on the type of \mathfrak{g}_N . In the proof, we shall use the following well-known properties of the idempotent map $\mathcal{E}_{[n]}$ (see, e.g., [7, Lemma 2.3] and the proof of [18, Thm. 2.4]):

$$\mathcal{E}_{[n]}^{1}\overline{R}_{nm}^{12} = \overline{R}_{nm}^{\bar{1}2}\mathcal{E}_{[n]}^{1}, \quad \mathcal{E}_{[m]}^{2}\overline{R}_{nm}^{12} = \overline{R}_{nm}^{1\bar{2}}\mathcal{E}_{[m]}^{2}, \quad \mathcal{E}_{[n]}^{1}T_{[n]}^{+12}(u_{[n]}) = \overline{T}_{[n]}^{+12}(u_{[n]})\mathcal{E}_{[n]}^{1}.$$
(4.31)

Note that the above equalities imply

$$\mathcal{E}_{[n,m]}^{12}\overline{R}_{nm}^{12} = \overline{R}_{nm}^{\overline{12}}\mathcal{E}_{[n,m]}^{12}, \qquad (4.32)$$

$$\mathcal{E}_{[n,m]}^{12}T_{[n]}^{+13}(u_{[n]})T_{[m]}^{+24}(v_{[m]}) = \tilde{T}_{[m]}^{+24}(v_{[m]})\tilde{T}_{[n]}^{+13}(u_{[n]})\mathcal{E}_{[n,m]}^{12}.$$
(4.33)

(a) Let us prove that all coefficients of $\mathbb{T}^+_{[n,m]}(u,v)$ are fixed points of σ . By (4.1), the image of $\mathbb{T}^+_{[n,m]}(u,v)$ under the braiding map $\sigma(z)$ equals

tr
$$\mathcal{E}_{[n,m]}^{12} \overline{R}_{nm}^{12} T_{[n]}^{+13}(u_{[n]}) T_{[m]}^{+24}(v_{[m]}) (\overline{R}_{nm}^{12})^{-1} (\mathbf{1} \otimes \mathbf{1}),$$

where the trace is taken over the tensor factors $1, \ldots, n+m$. By employing the identities (4.32) and (4.33), along with $(\mathcal{E}_{[n,m]}^{12})^2 = \mathcal{E}_{[n,m]}^{12}$, we rewrite the given expression as follows:

$$\operatorname{tr} \overline{R}_{nm}^{12} \mathcal{E}_{[n,m]}^{12} T_{[n]}^{+13}(u_{[n]}) T_{[m]}^{+24}(v_{[m]}) (\overline{R}_{nm}^{12})^{-1} (\mathbf{1} \otimes \mathbf{1})$$

=
$$\operatorname{tr} \overline{R}_{nm}^{\overline{12}} \mathcal{E}_{[n,m]}^{12} \mathcal{E}_{[n,m]}^{12} T_{[n]}^{+13}(u_{[n]}) T_{[m]}^{+24}(v_{[m]}) (\overline{R}_{nm}^{12})^{-1} (\mathbf{1} \otimes \mathbf{1})$$

=
$$\operatorname{tr} \mathcal{E}_{[n,m]}^{12} \overline{R}_{nm}^{12} \overline{T}_{[n]}^{+13}(u_{[n]}) \overline{T}_{[m]}^{+24}(v_{[m]}) (\overline{R}_{nm}^{\overline{12}})^{-1} \mathcal{E}_{[n,m]}^{12} (\mathbf{1} \otimes \mathbf{1}).$$

Next, we use the cyclic property of the trace to move the left copy of $\mathcal{E}_{[n,m]}^{12}$ to the right:

$$\operatorname{tr} \overline{R}_{nm}^{12} \overline{T}_{[n]}^{+13}(u_{[n]}) \overline{T}_{[m]}^{+24}(v_{[m]}) (\overline{R}_{nm}^{\overline{12}})^{-1} (\mathcal{E}_{[n,m]}^{12})^2 (\mathbf{1} \otimes \mathbf{1}) = \operatorname{tr} \overline{R}_{nm}^{12} \overline{T}_{[n]}^{+13}(u_{[n]}) \overline{T}_{[m]}^{+24}(v_{[m]}) (\overline{R}_{nm}^{\overline{12}})^{-1} \mathcal{E}_{[n,m]}^{12} (\mathbf{1} \otimes \mathbf{1}).$$

Finally, we employ the equalities (4.32) and (4.33) once again, to move $\mathcal{E}_{[n,m]}^{12}$ to the left:

$$\operatorname{tr} \overline{R}_{nm}^{12} \mathcal{E}_{[n,m]}^{12} T_{[n]}^{+13}(u_{[n]}) T_{[m]}^{+24}(v_{[m]}) (\overline{R}_{nm}^{12})^{-1} (\mathbf{1} \otimes \mathbf{1}).$$

By the cyclic property of the trace, the terms \overline{R}_{nm}^{12} and $(\overline{R}_{nm}^{12})^{-1}$ cancel, so we get

$$\operatorname{tr} \mathcal{E}_{[n,m]}^{12} T_{[n]}^{+13}(u_{[n]}) T_{[m]}^{+24}(v_{[m]}) (\mathbf{1} \otimes \mathbf{1}) = \mathbb{T}_{[n,m]}^+(u,v),$$

as required.

The proof that all coefficients of $\mathbb{T}^+_{[n,m]}(u,v)$ are fixed points of ρ is similar. However, some of its intermediate steps differ, so we present it in detail. Let

$$\overline{R}_{nm}^{12*} = \overline{R}_{nm}^{12} (u_{[n]}|v_{[m]}|z + h\varepsilon c_{\text{crit}})^{-1} = \begin{cases} \overline{R}_{nm}^{12} (u_{[n]}|v_{[m]}|z - hN)^{-1} & \text{for } \mathfrak{g}_N = \mathfrak{gl}_N, \\ \overline{R}_{nm}^{12} (u_{[n]}|v_{[m]}|z - 2h\kappa)^{-1} & \text{for } \mathfrak{g}_N = \mathfrak{o}_N, \mathfrak{sp}_N. \end{cases}$$

First, we remark that the first two formulae in (4.31) hold for \overline{R}_{nm}^{12*} as well. By (4.2), the image of $\mathbb{T}^+_{[n,m]}(u,v)$ under the map $\rho(z)$ equals

tr
$$\mathcal{E}_{[n]}^1 \mathcal{E}_{[m]}^2 T_{[n]}^{+13}(u_{[n]}) \overline{R}_{nm}^{12*} T_{[m]}^{+24}(v_{[m]}) \overline{R}_{nm}^{12}(\mathbf{1} \otimes \mathbf{1}),$$

where the trace is again taken over the tensor factors $1, \ldots, n + m$. We now use the identities in (4.31) along with $(\mathcal{E}_{[n]}^1)^2 = \mathcal{E}_{[n]}^1$ and $(\mathcal{E}_{[m]}^2)^2 = \mathcal{E}_{[m]}^2$ to rewrite the given expression as follows:

$$\operatorname{tr} \tilde{T}_{[n]}^{+13}(u_{[n]}) \mathcal{E}_{[n]}^{1} \overline{R}_{nm}^{1\bar{2}*} \mathcal{E}_{[m]}^{2} T_{[m]}^{+24}(v_{[m]}) \overline{R}_{nm}^{12}(\mathbf{1} \otimes \mathbf{1})$$

=
$$\operatorname{tr} \tilde{T}_{[n]}^{+13}(u_{[n]}) \mathcal{E}_{[n]}^{1} \mathcal{E}_{[n]}^{1} \overline{R}_{nm}^{1\bar{2}*} \mathcal{E}_{[m]}^{2} \mathcal{E}_{[m]}^{2} T_{[m]}^{+24}(v_{[m]}) \overline{R}_{nm}^{12}(\mathbf{1} \otimes \mathbf{1}).$$

Next, we use the same identities to move the left (resp. right) copies of $\mathcal{E}_{[n]}^1$ and $\mathcal{E}_{[m]}^2$ all the way to the left (resp. right), thus getting

$$\operatorname{tr} \mathcal{E}_{[n]}^{1} \mathcal{E}_{[m]}^{2} T_{[n]}^{+13}(u_{[n]}) \overline{R}_{nm}^{\overline{1}2*} \overline{T}_{[m]}^{+24}(v_{[m]}) \overline{R}_{nm}^{\overline{1}\overline{2}} \mathcal{E}_{[n]}^{1} \mathcal{E}_{[m]}^{2}(\mathbf{1} \otimes \mathbf{1}).$$

Using the cyclic property of the trace we move the left copies of $\mathcal{E}_{[n]}^1$ and $\mathcal{E}_{[m]}^2$ to the right:

$$\operatorname{tr} T_{[n]}^{+13}(u_{[n]}) \overline{R}_{nm}^{\overline{1}2*} \overline{T}_{[m]}^{+24}(v_{[m]}) \overline{R}_{nm}^{\overline{1}\overline{2}} \mathcal{E}_{[n]}^{1} \mathcal{E}_{[m]}^{2}(\mathbf{1} \otimes \mathbf{1}).$$

Finally, we use the equalities in (4.31) once again, to move $\mathcal{E}_{[n]}^1$ and $\mathcal{E}_{[m]}^2$ to the left:

tr
$$T_{[n]}^{+13}(u_{[n]}) \mathcal{E}_{[n]}^1 \overline{R}_{nm}^{12*} \mathcal{E}_{[m]}^2 T_{[m]}^{+24}(v_{[m]}) \overline{R}_{nm}^{12}(\mathbf{1} \otimes \mathbf{1}).$$

By the cyclic property of the trace, this can be written equivalently as

$$\operatorname{tr} T_{[n]}^{+13}(u_{[n]}) \mathcal{E}_{[n]}^{1} \mathcal{E}_{[m]}^{2} T_{[m]}^{+24}(v_{[m]}) \left(\overline{R}_{nm}^{12*} \cdot \overline{R}_{nm}^{12}\right) (\mathbf{1} \otimes \mathbf{1}), \qquad (4.34)$$

where the symbol "." indicates the standard multiplication in the tensor product algebra $(\operatorname{End} \mathbb{C}^N)^{\otimes n} \otimes ((\operatorname{End} \mathbb{C}^N)^{\operatorname{op}})^{\otimes m}$. By the crossing symmetry properties (2.7) and (2.9), we have $\overline{R}_{nm}^{12*} \cdot \overline{R}_{nm}^{12} = 1$, so the expression in (4.34) is equal to

$$\operatorname{tr} T_{[n]}^{+13}(u_{[n]}) \mathcal{E}_{[n]}^{1} \mathcal{E}_{[m]}^{2} T_{[m]}^{+24}(v_{[m]}) (\mathbf{1} \otimes \mathbf{1}) = \operatorname{tr} \mathcal{E}_{[n]}^{1} T_{[n]}^{+13}(u_{[n]}) \mathcal{E}_{[m]}^{2} T_{[m]}^{+24}(v_{[m]}) (\mathbf{1} \otimes \mathbf{1}) = \mathbb{T}_{[n,m]}^{+}(u,v),$$

as required.

(b) Both equalities can be verified by arguing as in the proof of the first assertion of lemma. In addition, the analogous arguments can be found in the proof of Lemma 4.13 below, so we omit the proof. $\hfill \Box$

Remark 4.10. The series $\mathbb{T}^+_{[N]}(u) \in \mathcal{V}^{\operatorname{crit}}(\mathfrak{gl}_N)[[u]]$, which appears in (4.30), is the wellknown quantum determinant qdet $T^+(u)$ for the dual Yangian $Y^+(\mathfrak{gl}_N)$. Moreover, the corresponding property of the braiding map, $\sigma(z)(w \otimes \mathbb{T}^+_{[N]}(u)) = w \otimes \mathbb{T}^+_{[N]}(u)$ actually holds in $\mathcal{V}^c(\mathfrak{gl}_N)$ for any $c \in \mathbb{C}$, but we use it later at the critical level only.

The first statement of the next corollary for $\mathfrak{g}_N = \mathfrak{o}_N, \mathfrak{sp}_N$ was established in [7, Thm. 2.5]. Although we obtain it here as an immediate consequence of (3.5) and Lemma 4.9, we believe it is worth to single it out, as the braiding of an arbitrary *h*-adic quantum vertex algebra does not need to exhibit such a property; see [8, Thm. 1.4]. The second statement follows by combining Theorem 2.3 and the first statement.

Corollary 4.11. (a) For any m and n we have

$$\mathcal{S}(z)\Big(\mathbb{T}^+_{[n,m]}(u,v)\Big) = \mathbb{T}^+_{[n,m]}(u,v).$$

(b) For \mathfrak{g}_N of type A, B, the restriction of \mathcal{S} to $\mathfrak{z}(\mathcal{V}^{\operatorname{crit}}(\mathfrak{g}_N))^{\otimes 2}$ is the identity.

Analogously to the previous corollary, one also establishes its counterpart for σ .

Corollary 4.12. For \mathfrak{g}_N of type A, B, the restriction of σ to $\mathfrak{z}(\mathcal{V}^{\operatorname{crit}}(\mathfrak{g}_N))^{\otimes 2}$ is the identity.

Finally, we obtain a partial analogue of Lemma 4.9 for the map μ defined by (4.12).

Lemma 4.13. The braiding map μ satisfies

$$\mu(z,x)\Big(\mathbb{T}^+_{[1,1]}(u,v)\Big) = \mathbb{T}^+_{[1,1]}(u,v), \tag{4.35}$$

$$\mu(z,x)\Big(w\otimes\mathbb{T}^+_{[N]}(u)\Big)=w\otimes\mathbb{T}^+_{[N]}(u) \text{ for all } w\in\mathcal{V}^{\operatorname{crit}}(\mathfrak{gl}_N).$$
(4.36)

Proof. Let us prove (4.35). We shall use the notation

 $\overline{R}_{12} = \overline{R}_{12}(z+u-v), \ \overline{\underline{R}}_{12} = \overline{\underline{R}}_{12}(2x+z+u+v), \ \overline{\underline{R}}_{12}^* = \overline{\underline{R}}_{12}(2x+z+u+v-hN)^{-1}.$ By (4.13), the image of $\mathbb{T}^+_{[1,1]}(u,v)$ under the map μ is equal to

tr
$$\underline{\overline{R}}_{12}^*_{\mathrm{LR}} (\overline{R}_{12} T_{13}^+(u) \underline{\overline{R}}_{12} T_{24}^+(v) (\overline{R}_{12})^{-1}) (\mathbf{1} \otimes \mathbf{1}),$$

where the trace is taken over the tensor factors 1 and 2. Due to the cyclic property of the trace and the commutation relation $(\overline{R}_{12})^{-1}\overline{\underline{R}}_{12}^* = \overline{\underline{R}}_{12}^*(\overline{R}_{12})^{-1}$, this equals

$$\operatorname{tr} \overline{\underline{R}}_{12}^{*}_{\operatorname{LR}} \left((\overline{R}_{12})^{-1} \overline{R}_{12} T_{13}^{+}(u) \overline{\underline{R}}_{12} T_{24}^{+}(v) \right) (\mathbf{1} \otimes \mathbf{1}) = \operatorname{tr} \overline{\underline{R}}_{12}^{*}_{\operatorname{LR}} \left(T_{13}^{+}(u) \overline{\underline{R}}_{12} T_{24}^{+}(v) \right) (\mathbf{1} \otimes \mathbf{1}).$$

We use the cyclic property of the trace once more to rewrite the given expression as

$$\operatorname{tr} \underline{\overline{R}}_{12}^* \underbrace{(T_{13}^+(u) \underline{\overline{R}}_{12} T_{24}^+(v))}_{\operatorname{RL}} (\mathbf{1} \otimes \mathbf{1}) = \operatorname{tr} T_{13}^+(u) \left(\underline{\overline{R}}_{12}^* \underbrace{\overline{R}}_{\operatorname{RL}} \underline{\overline{R}}_{12} \right) T_{24}^+(v) (\mathbf{1} \otimes \mathbf{1}).$$

It remains to observe that the expression above is equal to $\mathbb{T}^+_{[1,1]}(u,v)$ since the unitarity and the crossing symmetry relations (2.4) and (2.7) imply $\underline{\overline{R}^*_{12}}_{\mathrm{RL}} \cdot \underline{\overline{R}}_{12} = 1$.

To prove (4.36), we use the property of the anti-symmetrizer,

$$\mathcal{E}_{[N]}\overline{R}_{0N}(u+(N-1)h)\dots\overline{R}_{02}(u+h)\overline{R}_{01}(u) = \mathcal{E}_{[N]}, \qquad (4.37)$$

which holds in the case $\mathfrak{g}_N = \mathfrak{gl}_N$; see [32, Ch. 1]. Furthermore, we use the notation $\overline{R}_{nN}^{12} = \overline{R}_{nN}^{12}(v|u_{[N]}|z), \ \overline{\underline{R}}_{nN}^{12} = \overline{\underline{R}}_{nN}^{12}(v|u_{[N]}|2x+z), \ \overline{\underline{R}}_{nN}^{12*} = \overline{\underline{R}}_{nN}^{12}(v|u_{[N]}|2x+z-hN)^{-1},$ where the *N*-tuple $u_{[N]}$ is given by (2.16). By employing (4.37), one easily proves

$$\mathcal{E}_{[N]}(\overline{R}_{nN}^{12})^{\pm 1} = \mathcal{E}_{[N]} \underline{\overline{R}}_{nN}^{12} = \mathcal{E}_{[N]} \underline{\overline{R}}_{nN}^{12*} = \mathcal{E}_{[N]}.$$
(4.38)

Due to (4.13), the image of $T_{[n]}^{+13}(v) \otimes \mathbb{T}_{[N]}^+(u)$ under the map μ is equal to

$$\operatorname{tr} \mathcal{E}_{[N]}^{2} \overline{\underline{R}}_{nN}^{12*} \cdot \left(\overline{R}_{nN}^{12} T_{[n]}^{+13}(v) \overline{\underline{R}}_{nN}^{12} T_{[N]}^{+24}(u_{[N]}) (\overline{R}_{nN}^{12})^{-1} \right) (\mathbf{1} \otimes \mathbf{1}), \tag{4.39}$$

where the trace is taken over the tensor factors $n + 1, \ldots, n + N$ and "." denotes the standard multiplication in $(\operatorname{End} \mathbb{C}^N)^{\otimes n} \otimes ((\operatorname{End} \mathbb{C}^N)^{\operatorname{op}})^{\otimes N}$. By employing (4.38) and the third equality in (4.31) with n = N, we find that (4.39) equals

$$\operatorname{tr} \mathcal{E}_{[N]}^{2} \overline{\underline{R}}_{nN}^{12*} \left(\overline{R}_{nN}^{12} T_{[n]}^{+13}(v) \overline{\underline{R}}_{nN}^{12} T_{[N]}^{+24}(u_{[N]}) (\overline{R}_{nN}^{12})^{-1} \right) (\mathbf{1} \otimes \mathbf{1})$$

$$= \operatorname{tr} \overline{\underline{R}}_{nN}^{12*} \left(\mathcal{E}_{[N]}^{2} \overline{R}_{nN}^{12} T_{[n]}^{+13}(v) \overline{\underline{R}}_{nN}^{12} T_{[N]}^{+24}(u_{[N]}) (\overline{R}_{nN}^{12})^{-1} \right) (\mathbf{1} \otimes \mathbf{1})$$

$$= \operatorname{tr} \overline{\underline{R}}_{nN}^{12*} \left(T_{[n]}^{+13}(v) \mathcal{E}_{[N]}^{2} \overline{\underline{R}}_{nN}^{12} T_{[N]}^{+24}(u_{[N]}) (\overline{R}_{nN}^{12})^{-1} \right) (\mathbf{1} \otimes \mathbf{1})$$

$$= \operatorname{tr} \overline{\underline{R}}_{nN}^{12*} \left(T_{[n]}^{+13}(v) \mathcal{T}_{[N]}^{+24}(u_{[N]}) \mathcal{E}_{[N]}^{2} (\overline{R}_{nN}^{12})^{-1} \right) (\mathbf{1} \otimes \mathbf{1})$$

$$= \operatorname{tr} \overline{\underline{R}}_{nN}^{12*} \left(T_{[n]}^{+13}(v) \mathcal{T}_{[N]}^{+24}(u_{[N]}) \mathcal{E}_{[N]}^{2} (\overline{R}_{nN}^{12})^{-1} \right) (\mathbf{1} \otimes \mathbf{1})$$

By the cyclic property of the trace and the last equality in (4.38), this equals

$$\operatorname{tr}\left(\mathcal{E}_{[N]}^{2}\overline{\underline{R}}_{nN}^{12*}\right)_{\operatorname{LR}}\left(T_{[n]}^{+13}(v)\overline{T}_{[N]}^{+24}(u_{[N]})\right)(\mathbf{1}\otimes\mathbf{1}) = \operatorname{tr}T_{[n]}^{+13}(v)\overline{T}_{[N]}^{+24}(u_{[N]})\mathcal{E}_{[N]}^{2}(\mathbf{1}\otimes\mathbf{1}).$$

Finally, we use the third equality in (4.31) to move $\mathcal{E}_{[N]}$ to the left, thus getting

tr
$$T_{[n]}^{+13}(v)\mathcal{E}_{[N]}^2 T_{[N]}^{+24}(u_{[N]})(\mathbf{1}\otimes\mathbf{1}) = T_{[n]}^{+13}(v)\otimes\mathbb{T}_{[N]}^+(u),$$

as required.

5. Generalized Yangians as (σ, ρ) -deformed $\mathcal{V}^{c}(\mathfrak{g}_{N})$ -modules

In this section, we associate (σ, ρ) -deformed modules with the generalized Yangians of types A-D and demonstrate their applications. Recall the *R*-matrix R(u) defined by (2.1) (resp. (2.2)) for $\mathfrak{g}_N = \mathfrak{gl}_N$ (resp. $\mathfrak{g}_N = \mathfrak{o}_N, \mathfrak{sp}_N$). We shall also need the *R*-matrix $\widehat{R}(u)$, which is a polynomial in the variable u,

$$\widehat{R}(u) = \begin{cases} u R(u) & \text{for } \mathfrak{g}_N = \mathfrak{gl}_N, \\ u(u - h\kappa) R(u) & \text{for } \mathfrak{g}_N = \mathfrak{o}_N, \mathfrak{sp}_N. \end{cases}$$
(5.1)

Following Krylov and Rybnikov [25, Sect. 3.1], we define the algebra $\mathcal{Y}(\mathfrak{g}_N)$ as the *h*-adically completed associative algebra over the ring $\mathbb{C}[[h]]$ generated by the elements $\lambda_{ij}^{(r)}$, where $i, j = 1, \ldots, N$ and $r \in \mathbb{Z}$, subject to the defining relations

$$\widehat{R}(u-v)\mathcal{L}_1(u)\mathcal{L}_2(v) = \mathcal{L}_2(v)\mathcal{L}_1(u)\widehat{R}(u-v).$$
(5.2)

The generator matrix $\mathcal{L}(u)$ is defined by

$$\mathcal{L}(u) = \sum_{i,j=1}^{N} e_{ij} \otimes \lambda_{ij}(u), \quad \text{where} \quad \lambda_{ij}(u) = \delta_{ij} - h \sum_{r \in \mathbb{Z}} \lambda_{ij}^{(-r)} u^r.$$
(5.3)

Consider the two-sided ideals I_p in $\mathcal{Y}(\mathfrak{g}_N)$, where $p \ge 1$, generated by the elements

$$\lambda_{ij}^{(-r)}$$
 with $r \ge p$ and $h^p \cdot 1$.

Clearly, the family $(I_p)_{p\geq 0}$, where we set $I_0 = \mathcal{Y}(\mathfrak{g}_N)$, satisfies the constraints in (3.7). Denote by τ the topology on $\mathcal{Y}(\mathfrak{g}_N)$ defined by the basis $a+I_p$ for all $a \in \mathcal{Y}(\mathfrak{gl}_N)$ and $p \geq 0$. Consider the algebra completed with respect to τ , $\lim_{\leftarrow} \mathcal{Y}(\mathfrak{g}_N)/I_p$. For $\mathfrak{g}_N = \mathfrak{gl}_N$ (resp. $\mathfrak{g}_N = \mathfrak{o}_N, \mathfrak{sp}_N$), we denote this completed algebra by $\widetilde{\mathcal{Y}}(\mathfrak{gl}_N)$ (resp. $\widetilde{\mathcal{Y}}(\mathfrak{g}_N)'$). Finally, for

 $\mathfrak{g}_N = \mathfrak{o}_N, \mathfrak{sp}_N$, we define the algebra $\widetilde{\mathcal{Y}}(\mathfrak{g}_N)$ as the quotient of $\widetilde{\mathcal{Y}}(\mathfrak{g}_N)'$ over the τ -completed two-sided ideal generated by the relations

$$\mathcal{L}(u)\mathcal{L}(u+h\kappa)' = 1. \tag{5.4}$$

In contrast with the original definition [25, Sect. 3.1] for $\mathfrak{g}_N = \mathfrak{gl}_N$, which is given over the complex field, we consider algebras over the ring $\mathbb{C}[[h]]$ so that they fit the *h*-adic quantum vertex algebra setting. We refer to the algebras $\widetilde{\mathcal{Y}}(\mathfrak{g}_N)$ as generalized Yangians.

In analogy with (2.28), we introduce the notation

$$\mathcal{L}_{[n]}(u) = \mathcal{L}_1(u_1) \dots \mathcal{L}_n(u_n) \quad \text{and} \quad \mathcal{L}_{[n]}(u|z) = \mathcal{L}_1(z+u_1) \dots \mathcal{L}_n(z+u_n).$$
(5.5)

Theorem 5.1. For any $c \in \mathbb{C}$, there exists a unique structure of (σ, ρ) -deformed $\mathcal{V}^c(\mathfrak{g}_N)$ module over $\widetilde{\mathcal{Y}}(\mathfrak{g}_N)$ such that the module map is defined by

$$Y_{\widetilde{\mathcal{Y}}(\mathfrak{g}_N)}(T^+_{[m]}(u)\mathbf{1}, z) = \mathcal{L}_{[m]}(u|z).$$
(5.6)

Proof. The map $Y_{\widetilde{\mathcal{Y}}(\mathfrak{g}_N)}(\cdot, z)$ is well-defined by (5.6) due to the fact that the defining relations for the dual Yangian, (2.10) and (2.14), both for the matrix $T^+(u)$, are of the same form as the defining relations (5.2) and (5.4) for the algebra $\widetilde{\mathcal{Y}}(\mathfrak{g}_N)$. In particular, note that by replacing the *R*-matrix R(u-v) in (2.10) by $\widehat{R}(u-v)$, one obtains equivalent relations. Thus, to prove the theorem, it is sufficient to check that the requirements imposed by Definition 3.5 hold. It is clear from the definition of the topology τ that the image of $Y_{\widetilde{\mathcal{Y}}(\mathfrak{g}_N)}(\cdot, z)$ belongs to $\widetilde{\mathcal{Y}}(\mathfrak{g}_N)((z^{-1}))_{\tau}$ and, in addition, that this map satisfies the truncation condition (3.10). Moreover, the vacuum property (3.9) obviously holds.

As for the ρ -associativity, i.e. (3.11) with $\nu = \rho$, one easily derives from (4.5) and (5.6) that both sides of (3.11) map the expression $T_{[n]}^{+13}(u)T_{[m]}^{+24}(v)$, with n and m being arbitrary and $u = (u_1, \ldots, u_n), v = (v_1, \ldots, v_m)$, to

$$\mathcal{L}_{1k}(z_2+z_0+u_1)\dots\mathcal{L}_{nk}(z_2+z_0+u_n)\mathcal{L}_{n+1k}(z_2+v_1)\dots\mathcal{L}_{n+mk}(z_2+v_m)$$

for k = n + m + 1, so it is evident that $Y_{\widetilde{\mathcal{Y}}(\mathfrak{g}_N)}(\cdot, z)$ possesses the ρ -associativity property. It remains to show that $Y_{\widetilde{\mathcal{Y}}(\mathfrak{g}_N)}(\cdot, z)$ satisfies the σ -commutativity, i.e. (3.12) with $\mu = \sigma$.

By (4.1) and (5.6), the left-hand side of (3.12) maps the expression $T_{[n]}^{+13}(u)T_{[m]}^{+24}(v)$ to

$$\overline{R}_{nm}^{12}(u|v|-z_2+z_1)\mathcal{L}_{[n]}^{13}(u|z_1)\mathcal{L}_{[m]}^{23}(v|z_2)\overline{R}_{nm}^{12}(u|v|-z_2+z_1)^{-1}.$$

By canceling the normalizing functions, this can be written via the R-matrices (5.1) as

$$\widehat{R}_{nm}^{12}(u|v|-z_2+z_1)\mathcal{L}_{[n]}^{13}(u|z_1)\mathcal{L}_{[m]}^{23}(v|z_2)\widehat{R}_{nm}^{12}(u|v|-z_2+z_1)^{-1},$$
(5.7)

where $\widehat{R}_{nm}^{12}(u|v|z)$ is defined as in (2.26) with the factors $\overline{R}_{ij}(z+u_i-v_{j-n})$ being replaced by $\widehat{R}_{ij}(z+u_i-v_{j-n})$. On the other hand, the image of $T_{[n]}^{+13}(u)T_{[m]}^{+24}(v)$ under the righthand side of (3.12) equals

$$\mathcal{L}_{[m]}^{23}(v|z_2)\mathcal{L}_{[n]}^{13}(u|z_1).$$
(5.8)

Finally, it is clear from the defining relations (5.2) that (5.7) and (5.8) coincide, so we conclude that the map $Y_{\widetilde{\mathcal{Y}}(\mathfrak{g}_N)}(\cdot, z)$ is σ -commutative.

Next, we consider the structure of (σ, ρ) -deformed $\mathcal{V}^{\text{crit}}(\mathfrak{g}_N)$ -module over $\mathcal{Y}(\mathfrak{g}_N)$, i.e. we restrict the setting to the critical level. The next corollary is a direct consequence of the σ -commutativity (3.12) and the properties of the map σ given by (4.29) and (4.30).

Corollary 5.2. (a) The coefficients of all

$$\mathbb{L}_{[n]}(z) \coloneqq Y_{\widetilde{\mathcal{Y}}(\mathfrak{g}_N)} \left(\mathbb{T}^+_{[n]}(0), z \right) \in \widetilde{\mathcal{Y}}(\mathfrak{g}_N)[[z^{\pm 1}]]$$
19

generate a commutative subalgebra in $\mathcal{Y}(\mathfrak{g}_N)$.

(b) The coefficients of

$$\mathbb{L}_{[N]}(z) = Y_{\widetilde{\mathcal{Y}}(\mathfrak{gl}_N)} \left(\mathbb{T}^+_{[N]}(0), z \right) \in \widetilde{\mathcal{Y}}(\mathfrak{gl}_N)[[z^{\pm}]]$$

belong to the center of $\widetilde{\mathcal{Y}}(\mathfrak{gl}_N)$.

Remark 5.3. In the case $\mathfrak{g}_N = \mathfrak{gl}_N$, the fact that the coefficients of $\mathbb{L}_{[m]}(z)$ generate a commutative subalgebra was originally established in [25, Sect. 3]. Moreover, a different vertex-operator theoretic interpretation of the series $\mathbb{L}_{[m]}(z)$ for $\mathfrak{g}_N = \mathfrak{gl}_N$ and a direct proof of the second assertion of Corollary 5.2 was given in [2, Sect. 6, 7].

The constructions from this section can be specialized to fit the (ordinary) h-adically completed Yangian Y(\mathfrak{g}_N). Following (2.13), one redefines the series $\lambda_{ij}(u)$ in (5.3) as $\lambda_{ij}(u) = \delta_{ij} + h \sum_{r \ge 1} \lambda_{ij}^{(r)} u^{-r}$ and sets τ to be the *h*-adic topology. Then, the next analogues of Theorem 5.1 and Corollary 5.2 follow by simply adjusting the above arguments.

Theorem 5.4. For any $c \in \mathbb{C}$, there exists a unique structure of (σ, ρ) -deformed $\mathcal{V}^c(\mathfrak{g}_N)$ module over the h-adically completed Yangian $Y(\mathfrak{g}_N)$ such that the module map satisfies

$$Y_{Y(\mathfrak{g}_N)}(T^+_{[m]}(u)\,\mathbf{1},z) = T^-_1(z+u_1)\dots T^-_m(z+u_m).$$
(5.9)

Remark 5.5. Theorem 5.4 can be reformulated as follows. By [18, Prop. 4.2], the hadically completed dual Yangian $Y^+(\mathfrak{g}_N)$ can be equipped with the structure of h-adic quantum vertex algebra $\mathcal{V}^+(\mathfrak{g}_N)$, such that its braiding, which governs (2.24), is the map σ given by (4.1). Moreover, its vertex operator map is the map Y^{ρ} given by (4.5). Thus, $(\sigma, 1)$ is a compatible pair in $\mathcal{V}^+(\mathfrak{g}_N)$. Hence, Theorem 5.4 asserts that the hadically completed Yangian $Y(\mathfrak{g}_N)$ is equipped with a unique structure of $(\sigma, 1)$ -deformed $\mathcal{V}^+(\mathfrak{g}_N)$ -module, such that the corresponding module map is again given by (5.9).

Corollary 5.6. (a) The coefficients of all

$$\mathbb{T}^{-}_{[n]}(z) \coloneqq Y_{\mathcal{Y}(\mathfrak{g}_N)} \big(\mathbb{T}^{+}_{[n]}(0), z \big) \in \mathcal{Y}(\mathfrak{g}_N)[[z^{-1}]]$$

generate a commutative subalgebra in $Y(\mathfrak{g}_N)$.

(b) The coefficients of

$$\mathbb{T}_{[N]}^{-}(z) = Y_{\mathcal{Y}(\mathfrak{gl}_{N})}\big(\mathbb{T}_{[N]}^{+}(0), z\big) \in \mathcal{Y}(\mathfrak{gl}_{N})[[z^{-1}]]$$

belong to the center of $Y(\mathfrak{gl}_N)$.

Remark 5.7. Both statements of Corollary 5.6 are well-known; see [31, Chap. 1.6, 1.14] and [32, Prop. 5.3]. In particular, the series $\mathbb{T}^{-}_{[N]}(z)$ is the quantum determinant for $Y(\mathfrak{gl}_N)$.

Recall that the *extended Yangian* $X(\mathfrak{g}_N)$ is defined as the *h*-adically completed algebra over $\mathbb{C}[[h]]$ generated by the elements $t_{ij}^{(r)}$, where i, j = 1, ..., N and r = 1, 2, ..., subject to the defining relation

$$R(u-v)T_1^{-}(u)T_2^{-}(v) = T_2^{-}(v)T_1^{-}(u)R(u-v),$$
(5.10)

where $T^{-}(u)$ and $t^{-}_{ij}(u)$ are defined as in (2.12) and (2.13), respectfully. Thus, for $\mathfrak{g}_{N} =$ $\mathfrak{o}_N,\mathfrak{sp}_N$, the Yangian $Y(\mathfrak{g}_N)$ is the quotient of the extended Yangian by the first relation in (2.14), while for $\mathfrak{g}_N = \mathfrak{gl}_N$, the algebras $X(\mathfrak{gl}_N)$ and $Y(\mathfrak{gl}_N)$ coincide.

Corollary 5.8. Let W be a topologically free $\mathbb{C}[[h]]$ -module and $c \in \mathbb{C}$. Suppose W is a $Y(\mathfrak{g}_N)$ -module and denote by $T^-(u)_W$ the action of $T^-(u)$ on W. There exists a unique structure of (σ, ρ) -deformed $\mathcal{V}^{c}(\mathfrak{g}_{N})$ -module over W such that the module map satisfies

$$Y_W(T^+_{[n]}(u) \mathbf{1}, z) = T^-_1(z+u_1)_W \dots T^-_n(z+u_n)_W$$

On the other hand, let W be a (σ, ρ) -deformed $\mathcal{V}^c(\mathfrak{g}_N)$ -module such that

$$Y_W(t_{ij}^{(-1)} \mathbf{1}, z) \in \text{Hom}(W, z^{-1}W[[z^{-1}]]) \text{ for all } i, j = 1, \dots, N.$$
(5.11)

There exists a unique structure of $X(\mathfrak{g}_N)$ -module on W satisfying

$$T^{-}(z)_{W} = Y_{W}(T^{+}(0)\mathbf{1}, z).$$
(5.12)

Proof. The first assertion of the corollary is proved by repeating the corresponding arguments from the proof of Theorem 5.1. The second assertion is verified by applying the σ -commutativity (3.12) to $T_{13}^+(0)T_{24}^+(0) \otimes w$ with $w \in W$ and using the explicit expression (4.1) for the braiding map σ . More specifically, this shows that $Y_W(T^+(0)\mathbf{1},z)$ satisfies the RTT-relation for $T^-(u)$ in (2.10). Thus, with $Y_W(T^+(0)\mathbf{1},z)$ being of the same form as $T^-(z)$ due to (5.11), we conclude that (5.12) defines a unique structure of $X(\mathfrak{g}_N)$ -module on W, as required.

6. Reflection algebras as (μ, ν) -deformed $\mathcal{V}^{\operatorname{crit}}(\mathfrak{gl}_N)$ -modules

We start by following [35, Sect. 2] to define a certain class of reflection algebras. As with the generalized Yangians in the previous section, we again modify the original definition so that it is given over the ring $\mathbb{C}[[h]]$. Consider the matrix

$$B(u) = T(u)T(-u)^{-1} \in \operatorname{End} \mathbb{C}^N \otimes \operatorname{Y}(\mathfrak{gl}_N)[[u^{-1}]].$$

It satisfies the *reflection equation*

$$R(u-v)B_1(u)R(u+v)B_2(v) = B_2(v)R(u+v)B_1(u)R(u-v)$$
(6.1)

and the unitarity condition

$$B(u)B(-u) = 1.$$
 (6.2)

The matrix B(u) can be written as

$$B(u) = \sum_{i,j=1}^{N} e_{ij} \otimes b_{ij}(u), \text{ where } b_{ij}(u) = \delta_{ij} + h \sum_{r=1}^{\infty} b_{ij}^{(r)} u^{-r}$$

for some elements $b_{ij}^{(r)} \in \mathcal{Y}(\mathfrak{gl}_N)$. Denote by $\mathcal{B}(N) = \mathcal{B}(N,0)$ the *h*-adically completed subalgebra of the Yangian generated by the elements $b_{ij}^{(r)}$, where $i, j = 1, \ldots, N$ and $r = 1, 2, \ldots$. These generators, along with the relations in (6.1) and (6.2), form a presentation of the algebra $\mathcal{B}(N)$; see [35, Thm. 3.1]. We refer to $\mathcal{B}(N)$ as reflection algebra.

For any $n \ge 1$ and the family of variables $u = (u_1, \ldots, u_n)$ we write

$$B_{[n]}(u|z) = \prod_{i=1,\dots,n} \left(B_i(z+u_i) \overline{R}_{i\,i+1}(2z+u_i+u_{i+1}) \dots \overline{R}_{in}(2z+u_i+u_n) \right).$$
(6.3)

For example, for $B_i = B_i(z + u_i)$ and $\overline{R}_{ij} = \overline{R}_{ij}(2z + u_i + u_j)$, we have

$$B_{[1]}(u|z) = B_1, \quad B_{[2]}(u|z) = B_1 \overline{R}_{12} B_2, \quad B_{[3]}(u|z) = B_1 \overline{R}_{12} \overline{R}_{13} B_2(z+u_2) \overline{R}_{23} B_3.$$

By using such a notation, one can express the generalized version of (6.1), which is a consequence of the quantum Yang–Baxter equation (2.3) and the reflection equation, as

$$R_{nm}^{12}(u|v|z_1 - z_2) B_{[n]}^{13}(u|z_1) \overline{\underline{R}}_{nm}^{12}(u|v|z_1 + z_2) B_{[m]}^{23}(v|z_2)$$

= $B_{[m]}^{23}(v|z_2) \overline{\underline{R}}_{nm}^{12}(u|v|z_1 + z_2) B_{[n]}^{13}(u|z_1) R_{nm}^{12}(u|v|z_1 - z_2),$ (6.4)

where the *R*-matrix product $R_{nm}^{12}(u|v|z)$ is defined as in (2.26) with the factors $\overline{R}_{ij}(z + u_i - v_{j-n})$ being replaced by $R_{ij}(z + u_i - v_{j-n})$. It is worth noting that, as with the original reflection equation (6.1), the identity in (6.4) holds if we replace all terms $(z_1 \pm z_2)$ by $(\pm z_2 + z_1)$. In other words, by expanding all negative powers of $(z_1 \pm z_2)$ in nonnegative

powers of z_1 instead of z_2 , we obtain an equivalent identity. We shall also need the following property of the product (6.3). For the variables $u = (u_1, \ldots, u_n)$ and $v = (v_1, \ldots, v_m)$, let $w = (u, v) = (u_1, \ldots, u_n, v_1, \ldots, v_m)$. Then $B_{[n+m]}(w|z)$ decomposes as

$$B_{[n+m]}(w|z) = B_{[n]}^{13}(u|z)\overline{\underline{R}}_{nm}^{12}(u|v|2z)B_{[m]}^{23}(v|z).$$
(6.5)

Consider the maps γ , μ and ν , as given by (4.10), (4.13) and (4.25), respectively. By Proposition 4.7, the pair (μ, ν) on $\mathcal{V}^{\text{crit}}(\mathfrak{gl}_N)$ is compatible, so one can consider the notion of (μ, ν) -deformed $\mathcal{V}^{\text{crit}}(\mathfrak{gl}_N)$ -module; see Definition 3.5. Recall that the topology τ from this definition is required to satisfy the constraints imposed by (3.7) and (3.8). In this particular case, we define $\mathcal{B}(N)_n = h^n \mathcal{B}(N)$ for $n \ge 0$ and set τ to be the *h*-adic topology, so that both (3.7) and (3.8) clearly hold.

Theorem 6.1. There exists a unique structure of (μ, ν) -deformed $\mathcal{V}^{\operatorname{crit}}(\mathfrak{gl}_N)$ -module over the algebra $\mathcal{B}(N)$ such that the module map is defined by

$$Y_{\mathcal{B}(N)}(T^+_{[m]}(u)\mathbf{1}, z) = B_{[m]}(u|z).$$
(6.6)

Proof. First, let us show that the module map $Y_{\mathcal{B}(N)}(\cdot, z)$ is well-defined by (6.6). It is sufficient to check that the defining relations for the dual Yangian, (2.10) with $T^+(u)$, belong to its kernel. More precisely, let n > i be two positive integers, $u = (u_1, \ldots, u_n)$ a family of variables and $u' = (u_1, \ldots, u_{i-1}, u_{i+1}, u_i, u_{i+2}, \ldots, u_n)$ the *n*-tuple with u_i and u_{i+1} swapped. It suffices to prove that the expression

$$R_{ii+1}T^{+}_{[n]}(u) \mathbf{1} - P_{ii+1}T^{+}_{[n]}(u') P_{ii+1}R_{ii+1}\mathbf{1}, \quad \text{where} \quad R_{ii+1} = R_{ii+1}(u_i - u_{i+1})$$

is annihilated by $Y_{\mathcal{B}(N)}(\cdot, z)$. By using (6.6), we find that it is mapped to

$$R_{ii+1}B_{[n]}(u|z) - P_{ii+1}B_{[n]}(u'|z)P_{ii+1}R_{ii+1}.$$
(6.7)

However, by examining the expression in (6.7), we see that the *R*-matrix R_{ii+1} in the first summand can be moved all the way to the right of $B_{[n]}(u|z)$. Indeed, this is done by the use of the quantum Yang–Baxter equation (2.3) and, also, the reflection equation (6.1) on the tensor factors *i* and *i* + 1. In particular, this turns

$$R_{ii+1}B_i(z+u_i)\overline{R}_{ii+1}(2z+u_i+u_{i+1})B_{i+1}(z+u_{i+1})$$

to

$$B_{i+1}(z+u_{i+1})\overline{R}_{i+1}(2z+u_i+u_{i+1})B_i(z+u_i)R_{i+1}$$

so that $R_{ii+1}B_{[n]}(u|z)$ becomes $P_{ii+1}B_{[n]}(u'|z)P_{ii+1}R_{ii+1}$. Hence, we conclude that (6.7) is equal to zero, as required.

To finish the proof, it remains to check that the module map $Y_{\mathcal{B}(N)}(\cdot, z)$ possesses the properties from Definition 3.5. The vacuum property (3.9) and the truncation condition (3.10) clearly hold. In particular, the later is evident from the fact that the image of $Y_{\mathcal{B}(N)}(\cdot, z)$ belongs to $\mathcal{B}(N)[[z^{-1}]]$. Regarding the ν -associativity, it is clear from (6.6) that the left-hand side of (3.11) maps the expression $T_{[n]}^{+13}(u)T_{[m]}^{+24}(v)(\mathbf{1}\otimes\mathbf{1})$ to

$$B_{[n]}^{13}(u|z_2+z_0)B_{[m]}^{23}(v|z_2).$$
(6.8)

As for the right-hand side of (3.11), by (4.26), $Y^{\nu}(\cdot, z_0, z_2)$ maps the same expression to

$$\overline{\underline{R}}_{nm}^{12}(u|v|2z_2+z_0-hN)^{-1} \underset{\mathrm{RL}}{\cdot} \left(T_{[n]}^{+13}(u|z_0)T_{[m]}^{+24}(v)\right) (\mathbf{1}\otimes\mathbf{1}).$$

Therefore, applying the module map $Y_{\mathcal{B}(N)}(\cdot, z_2)$, we obtain

$$\overline{\underline{R}}_{nm}^{12}(u|v|2z_2+z_0-hN)^{-1} \cdot \underset{\text{RL}}{\overset{22}{\text{RL}}} B_{[n+m]}(w|z_2),$$

where $w = (z_0 + u_1, \ldots, z_0 + u_n, v_1, \ldots, v_m)$. By using (6.5), we rewrite this as

$$\overline{\underline{R}}_{nm}^{12}(u|v|2z_2 + z_0 - hN)^{-1} \underset{\text{RL}}{\cdot} \left(B_{[n]}^{13}(u|z_2 + z_0) \overline{\underline{R}}_{nm}^{12}(u|v|2z_2 + z_0) B_{[m]}^{23}(v|z_2) \right)$$

= $B_{[n]}^{13}(u|z_2 + z_0) \left(\overline{\underline{R}}_{nm}^{12}(u|v|2z_2 + z_0 - hN)^{-1} \underset{\text{RL}}{\cdot} \overline{\underline{R}}_{nm}^{12}(u|v|2z_2 + z_0) \right) B_{[m]}^{23}(v|z_2).$ (6.9)

Finally, by the crossing symmetry property (2.7), all *R*-matrices in (6.9) cancel. Thus, the resulting expression is equal to the image (6.8) of the left-hand side, as required. Hence, we conclude that the ν -associativity (3.11) holds.

Let us prove that the μ -commutativity (3.12) holds. We start by considering the image of $T_{[n]}^{+13}(u)T_{[m]}^{+24}(v)(\mathbf{1}\otimes\mathbf{1})$ under the left-hand side of (3.12). First, we note that, by (4.13), its image under $\mu(-z_2 + z_1, z_2)$ equals

$$\underline{\overline{R}}_{nm}^{12*} \cdot \left(\overline{R}_{nm}^{12} T_{[n]}^{+13}(u) \underline{\overline{R}}_{nm}^{12} T_{[m]}^{+24}(v) (\overline{R}_{nm}^{12})^{-1} \right) (\mathbf{1} \otimes \mathbf{1}), \tag{6.10}$$

where

 $\overline{R}_{nm}^{12} = \overline{R}_{nm}^{12}(u|v| - z_2 + z_1), \ \overline{\underline{R}}_{nm}^{12} = \overline{\underline{R}}_{nm}^{12}(u|v|z_2 + z_1), \ \overline{\underline{R}}_{nm}^{12*} = \overline{\underline{R}}_{nm}^{12}(u|v|z_2 + z_1 - hN)^{-1}.$ Next, by applying $Y_{\mathcal{B}(N)}(z_1)(1 \otimes Y_{\mathcal{B}(N)}(z_2))$ to (6.10), we get

$$\underline{\overline{R}}_{nm}^{12*} \underbrace{\overline{R}}_{LR}^{12} \left(\overline{R}_{nm}^{12} B_{[n]}^{13}(u|z_1) \underline{\overline{R}}_{nm}^{12} B_{[m]}^{23}(v|z_2) (\overline{R}_{nm}^{12})^{-1} \right).$$

By using the generalized reflection equation (6.4), we rewrite the product of the first four factors inside the brackets in the above expression, thus getting

$$\underline{\overline{R}}_{nm}^{12*} \cdot \left(B_{[m]}^{23}(v|z_2) \underline{\overline{R}}_{nm}^{12} B_{[n]}^{13}(u|z_1) \overline{R}_{nm}^{12} (\overline{R}_{nm}^{12})^{-1} \right) = \underline{\overline{R}}_{nm}^{12*} \cdot \left(B_{[m]}^{23}(v|z_2) \underline{\overline{R}}_{nm}^{12} B_{[n]}^{13}(u|z_1) \right).$$

Furthermore, this simplifies to

$$B_{[m]}^{23}(v|z_2) \left(\underline{\overline{R}}_{nm}^{12*} \cdot \underline{\overline{R}}_{nm}^{12} \right) B_{[n]}^{13}(u|z_1) = B_{[m]}^{23}(v|z_2) B_{[n]}^{13}(u|z_1),$$
(6.11)

where the equality in (6.11) is a consequence of the crossing symmetry property (2.7). Finally, it is clear from (6.6) that the image of $T_{[n]}^{+14}(u)T_{[m]}^{+23}(v)(\mathbf{1}\otimes\mathbf{1})$ under the right-hand side of (3.12) equals the right-hand side of (6.11), so we conclude that the μ -commutativity holds, thus finishing the proof.

By combining the μ -commutativity (3.12) and the identities (4.35) and (4.36), we find

Corollary 6.2. (a) The coefficients of the series

$$\mathbb{B}_{[n]}(z) \coloneqq Y_{\mathcal{B}(N)} \left(\mathbb{T}_{[n]}^+(0), z \right) \in \mathcal{B}(N)[[z^{-1}]] \quad \text{with} \quad n = 1, N$$

generate a commutative subalgebra in $\mathcal{B}(N)$.

(b) The coefficients of

$$\mathbb{B}_{[N]}(z) = Y_{\mathcal{B}(N)} \big(\mathbb{B}_{[N]}(0), z \big) \in \mathcal{B}(N)[[z^{-1}]]$$

belong to the center of the reflection algebra $\mathcal{B}(N)$.

Remark 6.3. The commutative subalgebras of reflection algebras in the case of two dimensions go back to the pioneer work of Sklyanin [36]. The series $\mathbb{B}_{[N]}(z)$ from Corollary 6.2 is a multiple of the well-known *Sklyanin determinant* sdet T(u), whose odd coefficients generate the center of $\mathcal{B}(N)$; cf. [35, Thm. 3.4].

At the end, we establish a connection between modules for reflection algebras and (μ, ν) -deformed $\mathcal{V}^{\text{crit}}(\mathfrak{gl}_N)$ -modules. First, following [35, Sect. 2], we define the *reflection*

algebra $\widetilde{\mathcal{B}}(N) = \widetilde{\mathcal{B}}(N, 0)$ as the *h*-adically completed algebra generated by the elements $\widetilde{b}_{ij}^{(r)}$, where $i, j = 1, \ldots, N$ and $r = 1, 2, \ldots$, subject to the *reflection equation*

$$R(u-v)\tilde{B}_{1}(u)R(u+v)\tilde{B}_{2}(v) = \tilde{B}_{2}(v)R(u+v)\tilde{B}_{1}(u)R(u-v), \qquad (6.12)$$

where

$$\widetilde{B}(u) = \sum_{i,j=1}^{N} e_{ij} \otimes \widetilde{b}_{ij}(u) \quad \text{and} \quad \widetilde{b}_{ij}(u) = \delta_{ij} + h \sum_{r=1}^{\infty} \widetilde{b}_{ij}^{(r)} u^{-r}.$$

Corollary 6.4. Let W be a topologically free $\mathbb{C}[[h]]$ -module. Suppose W is a $\widetilde{\mathcal{B}}(N)$ module and denote by $\widetilde{B}(u)_W$ the action of $\widetilde{B}(u)$ on W. There exists a unique structure
of (μ, ν) -deformed $\mathcal{V}^{\operatorname{crit}}(\mathfrak{gl}_N)$ -module over W such that the module map satisfies

$$Y_W(T^+_{[n]}(u) \mathbf{1}, z) = B_{[n]}(u|z)_W,$$

where $\widetilde{B}_{[n]}(u|z)_W$ stands for the operator series obtained from (6.3) by replacing all $B_i(z+u_i)$ with $i = 1, \ldots, n$ by the actions $\widetilde{B}_i(z+u_i)_W$ on W. Conversely, let W be a (μ, ν) -deformed $\mathcal{V}^{\text{crit}}(\mathfrak{gl}_N)$ -module such that

$$Y_W(t_{ij}^{(-1)} \mathbf{1}, z) \in \operatorname{Hom}(W, z^{-1}W[[z^{-1}]]) \text{ for all } i, j = 1, \dots, N.$$
 (6.13)

There exists a unique structure of $\widetilde{\mathcal{B}}(N)$ -module on W satisfying

$$B(z)_W = Y_W(T^+(0)\mathbf{1}, z).$$
(6.14)

Proof. The first assertion of the corollary can be verified by repeating the arguments from the proof of Theorem 6.1. As for the second assertion, by applying the μ -commutativity (3.12) to $T_{13}^+(0)T_{24}^+(0) \otimes w$ with $w \in W$ and using the explicit expression (4.13) for the braiding map μ , one shows that $Y_W(T^+(0)\mathbf{1}, z)$ satisfies the reflection equation (6.12). Thus, with $Y_W(T^+(0)\mathbf{1}, z)$ being of the same form as $\widetilde{B}(z)$ due to (6.13), we conclude that (6.14) defines a unique structure of $\widetilde{\mathcal{B}}(N)$ -module on W.

Acknowledgment

L.B. is member of Gruppo Nazionale per le Strutture Algebriche, Geometriche e le loro Applicazioni (GNSAGA) of the Istituto Nazionale di Alta Matematica (INdAM). This work has been supported by Croatian Science Foundation under the project UIP-2019-04-8488. Furthermore, this work was supported by the project "Implementation of cutting-edge research and its application as part of the Scientific Center of Excellence for Quantum and Complex Systems, and Representations of Lie Algebras", PK.1.1.02, European Union, European Regional Development Fund.

References

- I. I. Anguelova, M. J. Bergvelt, H_D-Quantum vertex algebras and bicharacters, Commun. Contemp. Math. 11 (2009) 937–991; arXiv:0706.1528 [math.QA].
- [2] L. Bagnoli, S. Kožić, Yangian deformations of S-commutative quantum vertex algebras and Bethe subalgebras, Transform. Groups (2024), https://doi.org/10.1007/s00031-023-09837-w; arXiv:2307.03112 [math.QA].
- [3] B. Bakalov, V. G. Kac, *Field algebras*, Int. Math. Res. Not. IMRN (2003), 123–159; arXiv:math/0204282 [math.QA].
- [4] R. E. Borcherds, Vertex algebras, in "Topological Field Theory, Primitive Forms and Related Topics" (Kyoto, 1996), edited by M. Kashiwara, A. Matsuo, K. Saito and I. Satake, Progress in Math. 160, Birkhäuser, Boston, 1998, 35–77; arXiv:q-alg/9706008.
- [5] C. Boyallian, V. Meinardi, An approach to Quantum Conformal Algebra, J. Math. Phys. 63 (2022), 081705; arXiv:2012.15299 [math.QA].

- [6] R. Brauer, On algebras which are connected with the semisimple continuous groups, Ann. Math. 38 (1937), 854–872.
- [7] M. Butorac, N. Jing, S. Kožić, h-Adic quantum vertex algebras associated with rational R-matrix in types B, C and D, Lett. Math. Phys. 109 (2019), 2439–2471; arXiv:1904.03771 [math.QA].
- [8] A. De Sole, M. Gardini, V. G. Kac, On the structure of quantum vertex algebras, J. Math. Phys. 61 (2020), 011701 (29pp); arXiv:1906.05051 [math.QA].
- P. Etingof, D. Kazhdan, Quantization of Lie bialgebras, V, Selecta Math. (N.S.) 6 (2000), 105–130; arXiv:math/9808121 [math.QA].
- [10] I. B. Frenkel, N. Jing, Vertex representations of quantum affine algebras, Proc. Natl. Acad. Sci. USA, 85 (1988), 9373–9377.
- [11] E. Frenkel, N. Reshetikhin, Towards deformed chiral algebras, Quantum Group Symposium, XXI International Colloquium on Group Theoretical Methods in Physics (Goslar, 1996), Heron Press, Sofia, 1997, pp. 27–42; arXiv:q-alg/9706023.
- [12] M. Gardini, Quantum vertex algebras, Ph.D. thesis, Sapienza University of Rome, 2018.
- [13] K. Iohara, Bosonic representations of Yangian double $DY_{\hbar}(\mathfrak{g})$ with $\mathfrak{g} = \mathfrak{gl}_N, \mathfrak{sl}_N, J$. Phys. A 29 (1996), 4593–4621; arXiv:q-alg/9603033.
- [14] A. P. Isaev, A. I. Molev, Fusion procedure for the Brauer algebra, St. Petersburg Math. J. 22 (2011), 437–446; arXiv:0812.4113 [math.RT].
- [15] A. P. Isaev, A. I. Molev and O. V. Ogievetsky, A new fusion procedure for the Brauer algebra and evaluation homomorphisms, Int. Math. Res. Not. IMRN (2012), 2571–2606; arXiv:1101.1336 [math.RT].
- [16] N. Jing, F. Kong, H. Li, S. Tan, Deforming vertex algebras by vertex bialgebras, Commun. Contemp. Math. 26 (2024), 2250067; arXiv:2101.01464 [math.QA].
- [17] N. Jing, F. Kong, H. Li, S. Tan, Twisted quantum affine algebras and equivariant φ -coordinated modules for quantum vertex algebras, arXiv:2212.01895 [math.QA].
- [18] N. Jing, S. Kožić, A. Molev, F. Yang, Center of the quantum affine vertex algebra in type A, J. Algebra 496 (2018), 138–186; arXiv:1603.00237 [math.QA].
- [19] N. Jing, M. Liu, F. Yang, Double Yangians of classical types and their vertex representations, J. Math. Phys. 61 (2020), 051704 (39 pp); arXiv:1810.06484 [math.QA].
- [20] A. Jucys, On the Young operators of the symmetric group, Lietuvos Fizikos Rinkinys 6 (1966), 163–180.
- [21] V. Kac, Vertex algebras for beginners, University Lecture Series, 10. American Mathematical Society, Providence, RI, 1997.
- [22] C. Kassel, *Quantum Groups*, Graduate texts in mathematics; vol. **155**, Springer-Verlag, 1995.
- [23] F. Kong, Quantum affine vertex algebras associated to untwisted quantum affinization algebras, Comm. Math. Phys. 402 (2023), 2577–2625; arXiv:2212.04888 [math.QA].
- [24] F. Kong, Representations of quantum lattice vertex algebras, arXiv:2404.03552 [math.QA].
- [25] V. Krylov, L. Rybnikov, Bethe subalgebras in antidominantly shifted Yangians, Int. Math. Res. Not. IMRN (2024), https://doi.org/10.1093/imrn/rnad327; arXiv:2205.04700 [math.RT].
- [26] J. Lepowsky, H.-S. Li, Introduction to Vertex Operator Algebras and Their Representations, Progress in Math. Vol. 227, Birkhäuser, Boston, 2003.
- [27] H.-S. Li, Axiomatic G_1 -vertex algebras, Commun. Contemp. Math. 5 (2003), 281–327; arXiv:math/0204308 [math.QA].
- [28] H.-S. Li, Nonlocal vertex algebras generated by formal vertex operators, Selecta Math. (N. S.) 11 (2005), 349–397; arXiv:math/0502244 [math.QA].
- [29] H.-S. Li, *ħ*-adic quantum vertex algebras and their modules, Comm. Math. Phys. **296** (2010), 475– 523; arXiv:0812.3156 [math.QA].
- [30] H.-S. Li, φ-Coordinated Quasi-Modules for Quantum Vertex Algebras, Comm. Math. Phys. 308 (2011), 703-741; arXiv:0906.2710 [math.QA].
- [31] A. Molev, Yangians and classical Lie algebras, Mathematical Surveys and Monographs, 143. American Mathematical Society, Providence, RI, 2007.
- [32] A. I. Molev, Feigin-Frenkel center in types B, C and D, Invent. Math. 191 (2013), 1–34; arXiv:1105.2341 [math.RT].
- [33] A. Molev, Sugawara operators for classical Lie algebras, Mathematical Surveys and Monographs, vol. 229, American Mathematical Society, Providence, RI, 2018.
- [34] A. Molev, M. Nazarov, G. Olshanski, Yangians and classical Lie algebras, Russian Math. Surveys 51 (1996), 205–282; arXiv:hep-th/9409025.
- [35] A. I. Molev, E. Ragoucy, Representations of reflection algebras, Rev. Math. Phys. 14 (2002), 317– 342; arXiv:math/0107213 [math.QA].
- [36] E. K. Sklyanin, Boundary conditions for integrable quantum systems, J. Phys. A 21 (1988), 2375– 2389.
- [37] A. B. Zamolodchikov and Al. B. Zamolodchikov, Factorized S-matrices in two dimensions as the exact solutions of certain relativistic quantum field models, Ann. Phys. **120** (1979), 253–291.