

Stability of slow Hamiltonian dynamics from Lieb-Robinson bounds

Daniele Toniolo ^{1,*} and Sougato Bose ¹

¹*Department of Physics and Astronomy, University College London,
Gower Street, London WC1E 6BT, United Kingdom*

We rigorously show that a local spin system giving rise to a slow Hamiltonian dynamics is stable against generic, even time-dependent, local perturbations. The sum of these perturbations can cover a significant amount of the system's size. The stability of the slow dynamics follows from proving that the Lieb-Robinson bound for the dynamics of the total Hamiltonian is the sum of two contributions: the Lieb-Robinson bound of the unperturbed dynamics and an additional term coming from the Lieb-Robinson bound of the perturbations with respect to the unperturbed Hamiltonian. Our results are particularly relevant in the context of the study of the stability of Many-Body-Localized systems, implying that if a so called ergodic region is present in the system, to spread across a certain distance it takes a time proportional to the exponential of such distance. The non-perturbative nature of our result allows us to develop a dual description of the dynamics of a system. As a consequence we are able to prove that the presence of a region of disorder in a ergodic system implies the slowing down of the dynamics in the vicinity of that region.

I. INTRODUCTION

The Lieb-Robinson (L-R) bounds quantify, in non-relativistic quantum physics, the maximal speed at which an operator, that initially acts locally, spreads its action over the system as a function of time because of the Hamiltonian dynamics. This introduces in this context the concept of the lightcone: the effect of the spread of the support of the operator can be detected at a certain distance only after a certain time interval, before that it is exponentially small. The pioneering work of Lieb and Robinson [1], has been improved, in particular by Hastings at the beginning of the 2000's [2], with applications nowadays ranging from condensed matter theory [3] to quantum information and simulation [4, 5] and quantum chaos. The recent reviews [6, 7] collect some of these works. We also refer to [8] for the mathematical foundations of the L-R bounds.

In this work we evaluate the L-R bounds of a system with total Hamiltonian $H + \sum_j h_j$ starting from the L-R bounds of the local Hamiltonian H when $\sum_j h_j$ is an additional local Hamiltonian, that we call perturbation, but that can cover a portion of the system comparable with its size and can have intensity as large as that of H .

If a one dimensional lattice Hamiltonian gives rise to a slow dynamics, as quantified by equation (1), our theorem 1 implies that the dynamics $H + \sum_j h_j$ continues to be slow. The meaning of our result is very transparent considering the case in which the system hosts a single perturbation (2). The spread of the perturbation h , given by the Heisenberg dynamics of the full Hamiltonian $H + h$, happens with a time scale that is exponentially long with the distance as in the unperturbed case with the Hamiltonian H .

Our result, theorem 1, is *non-perturbative* and takes into account all possible kinds of dynamics arising from H . In fact theorem 1 allows us to develop a dual description of the dynamics of a system: when there are a pair of Hamiltonians H and H' , and perturbations $\sum_j h_j$, $\sum_j h'_j$, such that $H + \sum_j h_j = H' + \sum_j h'_j$, then if H or H' gives rise to a slow dynamics then the dynamics of the full Hamiltonian is, locally, slow. This implies, for example, that a region of disorder in a ergodic system slows down the dynamics in the vicinity of that region.

The motivation of our work arises, in the context of the study of Many-Body-Localized (MBL) phases of matter [9–13], from the the long term debate, spurred by the papers [14, 15], regarding the stability of MBL with respect to rare ergodic regions. Our result (2) states that such ergodic regions, within a phase of slow dynamics, propagate at most exponentially slowly, namely to spread across a certain distance they take a time proportional to the exponential of such a distance. We also discuss how regions of anomalous slow dynamics inside an ergodic system globally slow down the dynamics in the vicinity of such regions.

With slow dynamics we precisely mean the class of systems with a L-R bound giving rise to a logarithmic lightcone, defined as follows. Let us consider the operators A and B with supports on the regions $\text{supp}(A)$ and $\text{supp}(B)$ of the lattice, this means that outside $\text{supp}(A)$, respectively $\text{supp}(B)$, they act like the identity. We assume for simplicity

* d.toniolo@ucl.ac.uk; danielet@alumni.ntnu.no

that these supports are simply connected. H_ω gives rise to a slow dynamics when it holds:

$$\mathbb{E} \left\| [e^{iH_\omega t} A e^{-iH_\omega t}, B] \right\| \leq K \|A\| \|B\| t^\beta e^{-\frac{\text{dist}(\text{supp}(A), \text{supp}(B))}{\xi}} \quad (1)$$

With the symbol $\mathbb{E}_\omega(\cdot)$ we denote averaging with respect to all the possible realization of the Hamiltonian H_ω , that, for disordered systems, correspond to the different realizations of disorder $\omega = \{\omega_j\}_{j \in \Lambda}$. Λ denotes the lattice. The bound (1) could in principle arise for a deterministic Hamiltonian, nevertheless an example of this case is unknown to us. This type of L-R bounds have been proven for many-body-localized systems in the context of the so called Local-Integral-of-Motion model [16–18], with $\beta = 1$, and recently, despite only for low energies, from a rigorous analysis of the disordered XXZ model in [19]. The bound (1) in the remarkable work [19] unfortunately suffers from a K that is system's size dependent. In all the following K is assumed to be system's size independent.

The name ‘‘logarithmic lightcone’’ comes from the equality $C t^\beta e^{-\frac{d}{\xi}} = e^{\log(C t^\beta) - \frac{d}{\xi}}$, where we have defined $d := \text{dist}(\text{supp}(A), \text{supp}(B))$. C is a constant depending on K , such that $C t^\beta$ is dimensionless. This means that up to a time t exponentially large in d , $e^{iH_\omega t} A e^{-iH_\omega t}$ and B commute up to an exponentially small correction. In fact defining $C t_{max}^\beta = e^{\frac{d}{2\xi}}$, at $t = t_{max}$ the upper bound in (1) is proportional to $e^{-\frac{d}{2\xi}} \ll 1$, when $d \gg \xi$. $e^{iH_\omega t} A e^{-iH_\omega t}$ is a propagating operator and if we imagine B as a ‘‘detector’’ of such propagation, according to (1) within the time t_{max} there is virtually no detection.

We now informally state our theorem 1, according to equation (7), in the case of a single time-independent perturbation h acting on a system with the L-R bounds (1). h can be a random matrix, modelling a so called ergodic grain, as considered for example by [14, 15]. $H_\omega + h$ is the total Hamiltonian and $e^{-i(H_\omega + h)t}$ the corresponding dynamics. The support of h spreads into the system giving rise to the following L-R bound:

$$\mathbb{E} \left\| [e^{i(H_\omega + h)t} h e^{-i(H_\omega + h)t}, B] \right\| \leq 2K h_{max} \|B\| e^{-\frac{\text{dist}(h, B)}{\xi}} \left(t^\beta + 8 h_{max} \frac{t^{\beta+1}}{\beta + 1} \right) \quad (2)$$

The interpretation of the upper bound in (2) is straightforward: considering for simplicity that the term proportional to $t^{\beta+1}$ dominates on t^β , within the time $c t_{max}^{\beta+1} = e^{\frac{\text{dist}(h, B)}{2\xi}}$, then the RHS of (2) is exponentially small if $\text{dist}(h, B) \ll \xi$. With $\beta = 1$, the presence of the perturbation h makes t_{max} of the order of the square root of the same time interval in the unperturbed case, but still exponentially long in the distance among the supports of h and B . In this sense we call the slow dynamics with $\beta > 0$ of equation (1) stable: the lightcone remains logarithmic.

In the context of the disordered XY model, that is equivalent, through a Jordan-Wigner transformation, to the Anderson model, the bound (1) has been shown in [20] to hold with $\beta = 0$, meaning that at any t there is no spread at all of the support of A . It should be stressed that for disordered systems it is essential to average over the different realizations of the Hamiltonian corresponding to different disorder configurations. This is crucial already at the level of the single particle Anderson model [21, 22]. When the dynamics of H is of Anderson type, namely with $f(t) = \text{constant}$ in (3), theorem 1 agrees with previous results on the destabilization of the many-body Anderson phase by perturbations, considered in [17, 23, 24].

Our theorem 1 can be generalized, with slightly more restrictive assumptions, to systems in dimensions larger than one, we give a sketch of such generalization after the end of the proof of 1. A hint towards the existence of an MBL phase in dimensions larger than one comes by the experiments [25, 26] and theoretical work like [27]. If such a phase would be characterized by a L-R bound like in (1) then our theorem 1 implies the stability of its dynamics.

We now proceed to state our main result, theorem 1. Next we will provide several comments and consider special cases that illustrates the meaning of it. We will also present the dual picture for the description of the dynamics that arises from theorem 1, as mentioned above. The essence of theorem 1 is that for slow dynamics of the MBL-type, like in equation (1), any local perturbation that leaves free a significant part of the region in between A and B , see figure 2, and other than that is arbitrary, leaves invariant the slow dynamics of the system in between A and B , as quantified by the following Lieb-Robinson bounds (5), (6) and (7). The proof of theorem 1 will be given at the end of the paper.

In the following, with abuse of notation, we will denote $\text{dist}(A, B)$ the distance among the supports of the two operators, namely $\text{dist}(\text{supp}(A), \text{supp}(B))$. We explicitly formulate theorem 1 for Hamiltonians that depend on disorder, ω , or more in general that depend on a parameter. Our result applies also in the deterministic case.

Theorem 1. *A one dimensional spin system, defined on the lattice Λ , local Hilbert space \mathbb{C}^r , and with a local Hamiltonian $H_\omega + \sum_j h_{j,\omega}(t)$, is given. For all j , $h_{j,\omega}(t)$ is supported on simply connected regions, that we assume independent from time and the disorder configuration, and we define $h_{max}(t) := \max_{j,\omega} \|h_{j,\omega}(t)\| < \infty$. Two operators A and B supported on the simply connected regions $\text{supp}(A)$ and $\text{supp}(B)$ are also given. The dynamics of H is assumed to give rise to the L-R bound*

$$\mathbb{E} \left\| [e^{iH_\omega t} A e^{-iH_\omega t}, B] \right\| \leq K \|A\| \|B\| e^{-\frac{\text{dist}(A, B)}{\xi}} f(t) \quad (3)$$

with K depending on $\min\{|\text{supp}(A)|, |\text{supp}(B)|\}$, and f positive, non decreasing. The operator of unitary evolution of the full system $H_\omega + \sum_j h_{j,\omega}(t)$, is given by:

$$V_\omega(t) := T \left[\exp \left(-i \left(H_\omega t + \int_0^t ds \sum_j h_{j,\omega}(s) \right) \right) \right] \quad (4)$$

We assume that in between the supports of A and B there is a region free of perturbations $h_{j,\omega}(t)$ of size $\frac{\text{dist}(A,B)}{n}$, with $n \geq 1$, see figure 2. On the left of A and on the right of B there can be a generic configuration of perturbations $h_{j,\omega}$. Then, it holds:

$$\mathbb{E}_\omega \left\| [V_\omega^*(t) A V_\omega(t), B] \right\| \leq 2K \|A\| \|B\| e^{-(1-\frac{1}{2n}) \frac{\text{dist}(A,B)}{\xi}} f(t) + 16K \|A\| \|B\| \xi e^{-\frac{\text{dist}(A,B)}{2n\xi}} \int_0^t ds f(s) h_{\max}(s) \quad (5)$$

If all the perturbations $h_{j,\omega}$ are such that $\text{dist}(h_{j,\omega}, B) \geq \text{dist}(A, B)$, then denoting $d_{\min} := \min_j \{\text{dist}(h_{j,\omega}, B)\}$, under the assumption (3), it holds:

$$\mathbb{E}_\omega \left\| [V_\omega^*(t) A V_\omega(t), B] \right\| \leq 2K \|A\| \|B\| e^{-\frac{\text{dist}(A,B)}{\xi}} f(t) + 16K \|A\| \|B\| \xi e^{-\frac{d_{\min}}{\xi}} \int_0^t ds f(s) h_{\max}(s) \quad (6)$$

If there is only one perturbation $h(t)$ in the system and we look at how its support spreads, again under the assumption (3), it holds:

$$\mathbb{E}_\omega \left\| [V_\omega^*(t) h(t) V_\omega(t), B] \right\| \leq 2K h_{\max} \|B\| e^{-\frac{\text{dist}(h,B)}{\xi}} f(t) + 16K h_{\max}(t) \|B\| e^{-\frac{\text{dist}(h,B)}{\xi}} \int_0^t ds f(s) h_{\max}(s) \quad (7)$$

Before the proof we provide several remarks about the bounds (5), (6) and (7).

The size of the region F free from perturbations, as in figure 2, is $\frac{\text{dist}(A,B)}{n}$. Despite theorem 1 being mathematically correct for all $n > 1$, a meaningful physical interpretation requires n of order 1, namely a small integer. The smaller the free region, the larger n , the larger will become the influence of the perturbations on the L-R bound affecting the factor $e^{-\frac{\text{dist}(A,B)}{2n}}$ in (5). The reason to ask for such free region from interactions, is that, for a fixed time t the meaning of locality is that increasing the distance among the supports of A and B , the bound (5) must decrease and become very small for large distances. Other than leaving a region F free, there is no limit to the number of perturbations $h_{j,\omega}$ that are present in the system. They can have overlapping supports and be, for example, an actual nearest neighbour Hamiltonian $\sum_k h_{k,k+1}$.

We now take full advantage of the fact that the upper bound in (5) is non perturbative and present a dual picture for the description of the dynamics that we mentioned before. We consider a pair of Hamiltonians H and H' , and perturbations $\sum_j h_j$, $\sum_j h'_j$, such that $H + \sum_j h_j = H' + \sum_j h'_j$. For example, let us fix:

$$H = \sum_{j=-L}^{L-1} (\sigma_j^x \sigma_{j+1}^x + \sigma_j^y \sigma_{j+1}^y + \Delta \sigma_j^z \sigma_{j+1}^z) \quad (8)$$

a regular XXZ spin model, and the perturbation

$$\sum_{j=1}^R h_{j,\omega_j} = \sum_{j=1}^R \omega_j \sigma_j^z \quad (9)$$

providing on site disorder over the region R , with ω_j , for instance, taken at random uniformly from $[-W, W]$. In this case $\omega = \{\omega_j\}_{j \in \{1, \dots, R\}}$. As a reference we consider figure 1. The same total Hamiltonian is obtained, for every single realization of the disorder, namely of the variable ω , with

$$H'_\omega = \sum_{j=-L}^{L-1} (\sigma_j^x \sigma_{j+1}^x + \sigma_j^y \sigma_{j+1}^y + \Delta \sigma_j^z \sigma_{j+1}^z + \omega_j \sigma_j^z) \quad (10)$$

and

$$\sum_{j \in R^c} h'_{j,\omega_j} = - \sum_{j \in R^c} \omega_j \sigma_j^z \quad (11)$$




FIG. 1. A representation of the “dual picture” for the study of dynamics from Lieb-Robinson bounds. The final Hamiltonian, that is a regular XXZ model with a disordered region, corresponds to two different pairs $(H, \sum_j h_j)$ and $(H', \sum_j h'_j)$.

where we have denoted R^c the complement of R in $[-L, L]$. In equation (11) the variables ω_j are the same as in (10), therefore the terms $-\sum_{j \in R^c} \omega_j \sigma_j^z$ “undo” the part of H' depending on disorder everywhere but on R . Let us assume that H' gives rise to a slow dynamics, like in (1), and we apply theorem 1 to the Hamiltonian $H' + \sum_j h'_j$. This implies that the support of A , as in figure 1, spreads slowly. In this way we have established that the region of disorder in proximity of the support of A crucially affects the time scale of the spread of the support of A , that is logarithmic. Considering instead the dynamics arising from H , equation (8), with $f(t)$, like in (3), given by $f(t) = e^{\frac{vt}{\xi}}$, we would have obtained that the time scale for the spread of the support of A was linear. The dual description of the dynamics allows a much tighter bound. A recent rigorous result hinting in this direction is given in section 4.5 of [28]. In the proof of theorem 1 we show that taking h'_j depending on disorder, namely on ω_j , is consistent with taking averages over disorder configurations ω .

In the context of the study of the stability of many-body-localized phases the phenomenon of avalanches have gained importance starting with the work of De Roeck and Huveneers [14, 15], and more recently, for example, in the works [29, 30], where the effects of the presence of regions of anomalous low disorder are crucial in destabilizing the MBL phase. When h_ω in (2) is a so called avalanche, as discussed in the introduction, the consequence of our bound (2) is that the avalanche spreads exponentially slowly. A recent work hinting towards MBL phases being more stable than thought is [31]. More explicitly let us consider a perturbation h_ω that “removes” the Hamiltonian H on the region $[0, n]$, and replaces it, for example, with a random matrix. Considering the nearest neighbour Hamiltonian $H_\omega = \sum_{j=-L}^L H_{j,j+1,\omega}$, this is realized by the perturbation $h_\omega = -\sum_{k=1}^n H_{k,k+1,\omega} + T(n)$. The resulting




FIG. 2. Sketching a possible configuration of supports of the operators A , B and h_j of the lattice Λ . Below a configuration with perturbations that are far away from A and B , as described by (6).

Hamiltonian connects the random matrix $T(n)$, for example in the GOE ensemble, as considered by [14, 15], of size $r^n \times r^n$ on the left with $\sum_{j=-L}^{-1} H_{j,j+1,\omega}$ and on the right with $\sum_{j=n}^L H_{j,j+1,\omega}$. The result of (2) is that the spread of $e^{i(H_\omega+h_\omega)t} h_\omega e^{-i(H_\omega+h_\omega)t}$ in time, starting from the region $[0, n]$, happens according to a logarithmic light cone. In the work [32] a GOE matrix has been modelled as an all to all Hubbard model. The bound (2) is meaningful within the time scale such that the right hand side of (2) is smaller than the trivial bound to the commutator, that is $2h_{max} \|B\|$, therefore the larger the size of the GOE matrix, the larger its norm, the shorter the time scale that gives saturation of the bound.

We now comment upon models with $f(t) = \text{constant}$ in (3). The one dimensional XX or XY disordered models, that can be shown to be equivalent to the single particle Anderson model, have been studied for example by [20]. Let us consider a perturbation h supported at one edge of the system. Our theorem 1 gives predictions that qualitatively agree with the findings of [24]. The author of [24] has numerical evidences that the coupling of a disordered XX model with a XXZ -like boundary term causes the entanglement-entropy dynamics to transition from uniformly bounded in time, like proven in [33, 34], to $\log t$, as it would be in a MBL system, [12, 35, 36]. The fact that the upper bound in (2) with $\beta = 0$ goes like t , would lead to the $\log t$ growth of the entanglement entropy, as shown in [16, 18]. Another way of quantifying the slow dynamics of MBL is employing quantum information related concepts like the Holevo quantity [37]. The authors of [23] and [17] have rigorously shown that in a model of exponential dynamical localization, that is $\beta = 0$ in (1), perturbed by an Hamiltonian with a sparse distribution of interactions the transport is subdiffusive. Our result agrees with their, in fact the second term in the upper bound (5) would grow like t in this case. Nevertheless within our theory we cannot rigorously conclude that exponential dynamical localization is destabilized by interactions, or by a boundary term, in fact for this purpose we would need a lower bound in (5) rather than an upper bound. In the context of the study of Schrödinger operators Damanik and Tcheremchantsev [38] have developed methods to evaluate lower bounds to the dynamics.

A given Lieb-Robinson bound can give rise, in the context of systems with fast dynamics, to different types of transport regimes. The relation among the shape of the light cone as predicted by the Lieb-Robinson bound and the transport properties, both of particles and energy is not in a one to one correspondence. For example the authors of [39] have established that linear increase in time of entanglement entropy (that can be put in relation with a linear light cone [4]) happens in a system with diffusive energy transport. Also in [40] a linear lightcone, as predicted by the so called out-of-time-ordered-correlation function (OTOC), that is equivalent to the Lieb-Robinson bound, coexists with diffusive spreading of conserved charges in a quantum unitary circuit. On the other hand a spin system with a linear lightcone like the XXZ model supports ballistic transport [41]. This suggests that the details of the transport properties cannot be capture by the light cones structure, that nevertheless can distinguish fast dynamics, leading to ballistic or diffusive transport, from slow dynamics leading to no transport of particles, like in MBL.

We consider two further particular cases related to (5) and (7).

The first regards the case of slow dynamics, like in (1), arising from H and a single perturbation $h(t)$ varying with a time scale τ : $h(t) = h(\frac{t}{\tau})$. Assuming that $\|h(t)\|$ and $f(t)$ vary continuously in time, it is a consequence of the mean value theorem the existence of $s' \in [0, \frac{t}{\tau}]$, such that:

$$\int_0^t ds \|h(s)\| f(s) = \|h(s')\| \int_0^t ds f(s) \quad (12)$$

This connects with what stated by the authors of [42]. They have numerically found that a local perturbation, adiabatically varying with a time scale τ , is such that, in the infinite-time limit of the dynamics, $t \rightarrow \infty$, and for an infinite system, it causes effects, like charge redistribution, over a length scale of the order of $\log \tau$, both for the Anderson and the MBL dynamics. According to [42] an arbitrarily slow rate of the adiabatic evolution would cause effects on an arbitrarily long length scale. Our upper bound (7) in this case is such that for every finite time $t \leq \bar{t}$, if $\tau \gg \bar{t}$, with $h(0) = 0$ and $\sup_{s' \in [0, \frac{t}{\tau}]} \|h(s')\| \ll \max\{\|A\|, \|B\|\}$, then the effect of the perturbation $h(t)$ is negligible. In fact despite propagating within a logarithmic light-cone, and therefore reaching a region of size at most of order $\log \bar{t}$, the effect of the perturbation is suppressed by the small value of $\|h(s')\|$ respect to the values of $\|A\|$ and $\|B\|$. If instead the initial value of the perturbation $h(0)$ is of the order of $\max\{\|A\|, \|B\|\}$ then the perturbation $h(t)$ has a stronger effect but does not change the nature of the slow dynamics (1).

The second case concerns perturbations varying periodically, $h(t) = h(t + T)$, with high frequency. An application of the Riemann-Lebesgue lemma implies the decoupling of the integral in (7) as follows:

$$\int_0^t ds \|h(s)\| f(s) = \frac{1}{T} \int_0^T ds \|h(s)\| \int_0^t ds f(s) + O\left(T \max_{s \in [0, t]} \|h(s)\| \max_{s \in [0, t]} f(s)\right) \quad (13)$$

We see that in the limit of high frequency, or equivalently short period T , such that

$$O\left(T \max_{s \in [0, t]} \|h(s)\| \max_{s \in [0, t]} f(s)\right) \ll 1 \quad (14)$$

the integral in (13) decouples in the product of two factors, one is the average of $\|h(s)\|$ over a period, and the other one is the integral of $f(s)$.

The extension of our theorem 1 to systems of dimensionality $d > 1$ is straightforward under slightly more stringent assumptions, that consist in requiring the existence of a corona around the support of B free of perturbations, as in figure 3. This splits the space into an inner and outer regions. Clearly in the one dimensional case this is equivalent to say that not only there is a region free of perturbations on the left of B , see figure 2, as asked by the assumption of theorem 1, but also on its right. The bound corresponding to (3) in $d > 1$ would look substantially unaltered, but with a dependence of K on the dimensionality, see for instance [43], and with the factor $e^{-\frac{\text{dist}(A, B)}{2n\xi}}$ replaced by $e^{-\frac{r}{\xi}}$ where we have denoted $2r$ the thickness of the corona free of perturbations, as in figure 3. We give a sketch of the proof of these statement after the end of the proof of theorem 1.

We also note that in [44, 45] already appeared the idea of combining the interaction picture and L-R bounds (see below the proof of theorem 1) but the context in which this idea was applied, the study of long-range interacting systems, is different from ours.

The proof of theorem 1 follows.

Proof. To lighten the notation we drop the dependence of H and h_j on ω , but we restore it when we want to stress averaging.

The so called interaction picture (or representation) of quantum mechanics is based upon:

$$V(t) := T \left[\exp \left(-i \left(Ht + \int_0^t ds h(s) \right) \right) \right] = e^{-iHt} T \left[\exp \left(-i \int_0^t ds e^{iHs} h(s) e^{-iHs} \right) \right] \quad (15)$$

On the left hand side of (15) there is the operator of time evolution of the full system with Hamiltonian $H + h(t)$. $T(\cdot)$ denotes time ordering, according to the rule: latest time goes on the left, $T^{-1}(\cdot)$ instead is the inverse time ordering: the latest time goes on the right, that arises by Hermitian conjugation. This allows to accordingly write the integrals arising from the exponential. Equation (15) is proven showing that both operators are solutions of a




FIG. 3. A sketch in two dimensions of a configuration of perturbations that allows the generalization of (5) to higher dimensions. The inner and outer regions are divided apart from a region free of perturbations, a circular corona in 2D, a spherical one in 3D.

differential equation with a unique solution (once the initial condition $V(0) = \mathbb{1}$ is set), therefore they coincide.

$$i \frac{d}{dt} T \left[\exp \left(-i \left(Ht + \int_0^t ds h(s) \right) \right) \right] = (H + h(t)) T \left[\exp \left(-i \left(Ht + \int_0^t ds h(s) \right) \right) \right] \quad (16)$$

$$i \frac{d}{dt} \left(e^{-iHt} T \left[\exp \left(-i \int_0^t ds e^{iHs} h(s) e^{-iHs} \right) \right] \right) = \quad (17)$$

$$= H e^{-iHt} T \left[\exp \left(-i \int_0^t ds e^{iHs} h(s) e^{-iHs} \right) \right] + e^{-iHt} e^{iHt} h(t) e^{-iHt} T \left[\exp \left(-i \int_0^t ds e^{iHs} h(s) e^{-iHs} \right) \right] \\ = (H + h(t)) e^{-iHt} T \left[\exp \left(-i \int_0^t ds e^{iHs} h(s) e^{-iHs} \right) \right] \quad (18)$$

We now show that if two operators $h_1(t)$ and $h_2(t)$ are such that $\forall (s_1, s_2), [h_1(s_1), h_2(s_2)] = 0$, then:

$$T \left(\exp \left(-i \int_0^t ds (h_1(s) + h_2(s)) \right) \right) = T \left(\exp \left(-i \int_0^t ds h_1(s) \right) \right) T \left(\exp \left(-i \int_0^t ds h_2(s) \right) \right) \quad (19)$$

The argument is that since the LHS and the RHS of (19) satisfy the same differential equation then they coincide. We have that:

$$i \frac{d}{dt} \left[T \left(\exp \left(-i \int_0^t ds h_1(s) \right) \right) T \left(\exp \left(-i \int_0^t ds h_2(s) \right) \right) \right] = \quad (20)$$

$$h_1(t) T \left(\exp \left(-i \int_0^t ds h_1(s) \right) \right) T \left(\exp \left(-i \int_0^t ds h_2(s) \right) \right) + \\ + T \left(\exp \left(-i \int_0^t ds h_1(s) \right) \right) h_2(t) T \left(\exp \left(-i \int_0^t ds h_2(s) \right) \right) \quad (21)$$

$$= (h_1(t) + h_2(t)) T \left(\exp \left(-i \int_0^t ds h_1(s) \right) \right) T \left(\exp \left(-i \int_0^t ds h_2(s) \right) \right) \quad (22)$$

With $V(t)$ as in (4) it is:

$$\| [V^*(t) A V(t), B] \| = \| [e^{iHt} A e^{-iHt}, T \left(e^{-i \int_0^t ds \sum_j e^{-iHs} h_j(s) e^{iHs}} \right) B T^{-1} \left(e^{i \int_0^t ds \sum_j e^{-iHs} h_j(s) e^{iHs}} \right)] \| \quad (23)$$

Let us define

$$\left(e^{iHt} A e^{-iHt} \right)_{B_a(A)} := \frac{1}{r|B_a(A)^c|} \left(\text{Tr}_{B_a(A)^c} e^{iHt}(t) A e^{-iHt} \right) \otimes \mathbb{1}_{B_a(A)^c} \quad (24)$$

that is the partial trace on the Hilbert space associated to the complement, in Λ , of the ball $B_a(A)$, that identifies the union of $\text{supp}(A)$ with the region around $\text{supp}(A)$ within the distance a . $\frac{1}{r^{|B_a(A)^c|}}$ is just a normalization factor, with $|B_a(A)^c|$ denoting the number of sites (the cardinality) of the region $B_a(A)^c$.

One of the main ideas in the following is to use the fact that the Lieb-Robinson bound (3) is equivalent to the bound:

$$\|e^{iHt} A e^{-iHt} - (e^{iHt} A e^{-iHt})_{B_a(A)}\| \leq K \|A\| f(t) e^{-\frac{t}{\xi}} \quad (25)$$

The fact that (3) implies (25) is shown using Haar-integration for example in equations 11 and 12 of [46]. The fact that (25) implies (3) is easily seen as follows:

$$\| [e^{iHt} A e^{-iHt}, B] \| = \| [e^{iHt} A e^{-iHt} - (e^{iHt} A e^{-iHt})_{B_{\text{dist}(A,B)}(A)} + (e^{iHt} A e^{-iHt})_{B_{\text{dist}(A,B)}(A)}, B] \| \quad (26)$$

$$\leq 2 \|B\| \| [e^{iHt} A e^{-iHt} - (e^{iHt} A e^{-iHt})_{B_{\text{dist}(A,B)}(A)}] \| \leq K \|A\| \|B\| f(t) e^{-\frac{\text{dist}(A,B)}{\xi}} \quad (27)$$

In the following definition of $(\widehat{h}_j)_s$ and $(\overline{h}_j)_s$ to have a simple notation we assume the system's lattice to be $\Lambda = [-L, L] \cap \mathbb{Z}$.

We assume for simplicity that the center of the region F free from perturbations in figure 2 corresponds to $x = 0$. We restrict the supports of $e^{-iHs} h_j(s) e^{iHs}$, with $h_j(s)$ supported on the left of the region F , to $[-L, 0]$ and call them $(\widehat{h}_j)_s$, while the perturbations $h_j(s)$ supported on the right of F are such that $e^{-iHs} h_j(s) e^{iHs}$ are replaced with the corresponding ones with support on the region $[1, L]$. The more distant the support of $h_j(s)$ from F , on the right or on the left, the smallest the error associated with the restriction of the support that we have just described.

$$(\widehat{h}_j)_s := \frac{1}{r^{|[-L, 0]^c|}} (\text{Tr}_{[-L, 0]^c} e^{-iHs} h_j(s) e^{iHs}) \otimes \mathbb{1}_{[-L, 0]^c} \quad \text{with } \text{supp}(h_j(s)) \text{ on the left of } F \quad (28)$$

$$(\overline{h}_j)_s := \frac{1}{r^{|[1, L]^c|}} (\text{Tr}_{[1, L]^c} e^{-iHs} h_j(s) e^{iHs}) \otimes \mathbb{1}_{[1, L]^c} \quad \text{with } \text{supp}(h_j(s)) \text{ on the right of } F \quad (29)$$

Each $(\widehat{h}_j)_s$ commutes with each $(\overline{h}_k)_s$ because their supports are disjoint. This implies that we can make use of equation (19). Defining

$$\widehat{T} := T \left(e^{-i \int_0^t ds \sum_j (\widehat{h}_j)_s} \right) \quad \text{and} \quad \overline{T} := T \left(e^{-i \int_0^t ds \sum_k (\overline{h}_k)_s} \right) \quad (30)$$

it holds:

$$T \left(e^{-i \int_0^t ds (\sum_j (\widehat{h}_j)_s + \sum_k (\overline{h}_k)_s)} \right) = \widehat{T} \overline{T} = \overline{T} \widehat{T} \quad (31)$$

Following Appendix B of [47], we now show how to upper bound in norm the difference among two time-ordered operators. With $G(s)$ and $E(s)$ norm-continuous Hermitian operators, it is:

$$\| T \left[\exp \left(-i \int_0^t ds G(s) \right) \right] - T \left[\exp \left(-i \int_0^t ds E(s) \right) \right] \| \quad (32)$$

$$= \| T^{-1} \left[\exp \left(i \int_0^t ds G(s) \right) \right] T \left[\exp \left(-i \int_0^t ds E(s) \right) \right] - \mathbb{1} \| \quad (33)$$

$$= \left\| \int_0^t dr \frac{d}{dr} \left(T^{-1} \left[\exp \left(i \int_0^r ds G(s) \right) \right] T \left[\exp \left(-i \int_0^r ds E(s) \right) \right] \right) \right\| \quad (34)$$

$$= \left\| \int_0^t dr \left(T^{-1} \left[\exp \left(i \int_0^r ds G(s) \right) \right] i (G(r) - E(r)) T \left[\exp \left(-i \int_0^r ds E(s) \right) \right] \right) \right\| \quad (35)$$

$$\leq \int_0^t dr \| G(r) - E(r) \| \quad (36)$$

It follows from equation (32) that:

$$\|T \left(e^{-i \int_0^t ds \sum_j e^{-iHs} h_j(s) e^{iHs}} \right) - \widehat{T} \overline{T}\| \quad (37)$$

$$= \left\| \int_0^t dr \frac{d}{dr} \left(T^{-1} \left[\exp \left(i \int_0^r ds \sum_j e^{-iHs} h_j(s) e^{iHs} \right) \right] \widehat{T} \overline{T} \right) \right\| \quad (38)$$

$$= \left\| \int_0^t dr T^{-1} \left[\exp \left(i \int_0^r ds \sum_j e^{-iHs} h_j(s) e^{iHs} \right) \right] i \left(\sum_j e^{-iHs} h_j(s) e^{iHs} - \sum_l (\widehat{h}_l)_s - \sum_k (\overline{h}_k)_s \right) \widehat{T} \overline{T} \right\| \quad (39)$$

$$\leq \int_0^t ds \left(\sum_j \|e^{-iHs} h_j(s) e^{iHs} - (\widehat{h}_j)_s\| + \sum_k \|e^{-iHs} h_k(s) e^{iHs} - (\overline{h}_k)_s\| \right) \quad (40)$$

To upper bound (40) we realize that the operator A in the Lieb-Robinson bound (3) can be explicitly time dependent and disorder dependent. Let us take C an operator with simply connected support and with $\text{dist}(C, h) = a$, then:

$$\mathbb{E}_\omega \| [e^{iH_\omega t} h_\omega(t) e^{-iH_\omega t}, C] \| = \mathbb{E}_\omega \| [h_\omega(t), e^{-iH_\omega t} C e^{iH_\omega t}] \| = \quad (41)$$

$$\mathbb{E}_\omega \| [h_\omega(t), e^{-iH_\omega t} C e^{iH_\omega t} - (e^{-iH_\omega t} C e^{iH_\omega t})_{B_a(C)} + (e^{-iH_\omega t} C e^{iH_\omega t})_{B_a(C)}] \| \quad (42)$$

$$\mathbb{E}_\omega 2 \| h_\omega(t) \| \| e^{-iH_\omega t} C e^{iH_\omega t} - (e^{-iH_\omega t} C e^{iH_\omega t})_{B_a(C)} \| + \mathbb{E}_\omega \| [h_\omega(t), (e^{-iH_\omega t} C e^{iH_\omega t})_{B_a(C)}] \| \quad (43)$$

$$\leq 2K h_{max}(t) \|C\| f(t) e^{-\frac{a}{\xi}} \quad (44)$$

Where we have introduced the maximal norm over all the local perturbations $h_{i,\omega}$, $h_{max}(s) := \max_{i,\omega} \{ \|h_{i,\omega}(s)\| \}$. (44) in turn implies, using Haar-integration as mentioned above, that:

$$\mathbb{E}_\omega \| e^{iH_\omega t} h_\omega(t) e^{-iH_\omega t} - (e^{iH_\omega t} h_\omega(t) e^{-iH_\omega t})_{B_a(h)} \| \leq 2K h_{max}(t) f(t) e^{-\frac{a}{\xi}} \quad (45)$$

We can now go back to (40) considering the contribution to the integral in (40) of the h_j supported on the left of the region F free from interactions.

$$\sum_j \| e^{-iHs} h_j(s) e^{iHs} - (\widehat{h}_j)_s \| \leq 2K f(s) \sum_j \|h_j(s)\| e^{-\frac{\text{dist}(\text{supp}(h_j), x=0)}{\xi}} \quad (46)$$

$$\leq 2K f(s) h_{max}(s) \sum_j e^{-\frac{\text{dist}(\text{supp}(h_j), x=0)}{\xi}} \leq 2K f(s) h_{max}(s) \sum_{j=\frac{\text{dist}(A,B)}{2n}+1}^{\infty} e^{-\frac{j}{\xi}} \quad (47)$$

Observing that:

$$\sum_{j=\frac{\text{dist}(A,B)}{2n}+1}^{\infty} e^{-\frac{j}{\xi}} = e^{-\left(\frac{\text{dist}(A,B)}{2n}+1\right)\frac{1}{\xi}} \sum_{j=0}^{\infty} e^{-\frac{j}{\xi}} = e^{-\frac{\text{dist}(A,B)}{2n\xi}} \frac{1}{e^{\frac{1}{\xi}} - 1} \leq \xi e^{-\frac{\text{dist}(A,B)}{2n\xi}} \quad (48)$$

By symmetry the contribution to (40) of the h_j supported on the right of the region F is upper bounded by the same quantity. Overall (37) is upper bounded by:

$$\|T \left(e^{-i \int_0^t ds \sum_j e^{-iHs} h_j(s) e^{iHs}} \right) - \widehat{T} \overline{T}\| \leq 4K \xi e^{-\frac{\text{dist}(A,B)}{2n\xi}} \int_0^t ds f(s) h_{max}(s) \quad (49)$$

To shorten the notation we define:

$$T := T \left(e^{-i \int_0^t ds \sum_j e^{-iHs} h_j(s) e^{iHs}} \right) \quad (50)$$

Going back to (37) we have:

$$\|V(t)^* A V(t), B\| = \| [e^{iHt} A e^{-iHt}, T B T^{-1}] \| \quad (51)$$

$$= \| [e^{iHt} A e^{-iHt}, (T - \hat{T}\bar{T} + \hat{T}\bar{T}) B (T^{-1} - \bar{T}^{-1}\hat{T}^{-1} - \bar{T}^{-1}\hat{T}^{-1})] \| \quad (52)$$

$$= \| [e^{iHt} A e^{-iHt}, (T - \hat{T}\bar{T}) B T^{-1} + \hat{T}\bar{T} B (T^{-1} - \bar{T}^{-1}\hat{T}^{-1}) + \hat{T}\bar{T} B \bar{T}^{-1} \hat{T}^{-1}] \| \quad (53)$$

$$\leq 4 K \|A\| \|B\| \|T - \hat{T}\bar{T}\| + \| [e^{iHt} A e^{-iHt}, \bar{T} B \bar{T}^{-1}] \| \quad (54)$$

In the last step we made use of the fact that \hat{T} commutes with both B and \bar{T} because their supports are disjoint. In the next step, since $\bar{T} B \bar{T}^{-1}$ is supported on $x \geq 0$, see figure 2, we restrict the support of $e^{iHt} A e^{-iHt}$ on the left of $x = 0$, that is at a distance $r := (1 - \frac{1}{2n})\text{dist}(A, B)$ from the support of A . The restriction of $e^{iHt} A e^{-iHt}$ to such region and $\bar{T} B \bar{T}^{-1}$ commute. Restarting from (54) we have:

$$\mathbb{E}_\omega \|V(t)^* A V(t), B\| \quad (55)$$

$$\begin{aligned} &\leq 4 K \|A\| \|B\| \mathbb{E}_\omega \|T - \hat{T}\bar{T}\| + \mathbb{E}_\omega \| [e^{iHt} A e^{-iHt} - (e^{iHt} A e^{-iHt})_{B_r(\text{supp}(A))} + (e^{iHt} A e^{-iHt})_{B_r(\text{supp}(A))}, \bar{T} B \bar{T}^{-1}] \| \\ &\leq 2 K \|A\| \|B\| e^{-(1-\frac{1}{2n})\frac{\text{dist}(A, B)}{\xi}} + 16 K \|A\| \|B\| \xi e^{-\frac{\text{dist}(A, B)}{2n\xi}} \int_0^t ds f(s) h_{max}(s) \end{aligned} \quad (56)$$

This completes the proof of the bound given in (5).

We now sketch the proof of (6), all the technical tools and concepts required for it have been already introduced above. We explicitly present the proof of (6) because the assumption that $\forall j, \text{dist}(h_{j,\omega}, B) \geq \text{dist}(A, B)$ is slightly more restrictive than being the region in between A and B free of perturbations. In fact setting $n = 1$ in (5) we get a factor $\frac{1}{2}$ at the exponent of $e^{-\frac{\text{dist}(A, B)}{2\xi}}$, that is absent in (6) where we get $e^{-\frac{\text{dist}(A, B)}{2\xi}}$. Looking, in this setting, at figure 2, we see that the error introduced restricting the support of $e^{iHt} h_j e^{-iHt}$ on the left of the support of B or on its right is at most proportional to $e^{-\frac{d_{min}}{\xi}}$. The summation of the contribution of all h_j is then performed as in (46) and (47).

We end the proof briefly commenting on (7). If there is only one perturbation $h(t)$, and in the L-R bound it is $A = h(t)$, the support of $h(t)$ is on the left of that of B , in this scenario we can apply the bound (6). But with only one perturbation there is no summation, like in (46), (47), to be performed, therefore no factor ξ appearing together with h_{max} in the second term in the right hand side of (7).

This completes the proof of theorem 1. □

We provide a brief account for the generalization of the bound (5) to higher dimensions. The main idea follows those presented in the proof of theorem 1 given above. Namely to split the time-ordered part of the operator of time evolution into two factors, following the logic of equation (31). Referring to figure 3, all the perturbations h_j with support in the outer region, are such that the support of $e^{iHt} h_j e^{-iHt}$ is restricted to the region outside the green circle (in three dimensions that would be a sphere) with an error of at most (for those perturbations sitting just outside the circular corona) of the order of $e^{-\frac{r}{\xi}}$. The perturbations supported in the inner region instead, get the support of $e^{iHt} h_j e^{-iHt}$ restricted till the green circle, with an error again at most of the order of $e^{-\frac{r}{\xi}}$. These two types of restricted operators give rise to a new pair \hat{T} and \bar{T} that commutes as in (31). The rest of the reasoning follows that of the proof of theorem 1.

II. CONCLUDING DISCUSSION

We have proven that an Hamiltonian giving rise to a slow dynamics, as quantified by a Lieb-Robinson bound with a logarithmic lightcone, is stable in a non perturbative fashion. This means that when we are testing the dynamics locally on a certain region, if the perturbations to the Hamiltonian leave enough of this region free, then the dynamics of local operators in this region continues to be characterized by a logarithmic lightcone. The key to prove this fact is the decoupling of the dynamics associated to the perturbations provided by the interaction picture and the fact that, because decoupled, the perturbations that are far away are not going to influence the dynamics locally. Surprisingly we are also able, reversing this logic, to take advantage of perturbations, in particular disorder, to show that in the

vicinity or a disordered region within an ordered system the dynamics slows down. The only input for our theorem 1 is the Lieb-Robinson bound of the unperturbed Hamiltonian under the assumption of locality. Our result not only contributes to the field of many body localization, where it is among the few rigorous ones, but also concerns the general dynamical behavior of Hamiltonian systems.

Acknowledgments

The authors acknowledge support from UKRI grant EP/R029075/1. They're also glad to acknowledge discussions on the topic of Lieb-Robinson bounds and Many-Body-Localization with Abolfazl Bayat, Igor Lerner and George McArdle.

-
- [1] E. H. Lieb and D. Robinson, The finite group velocity of quantum spin systems, *Commun. Math. Phys.* **28**, 251 (1972).
- [2] M. B. Hastings, Lieb-schultz-mattis in higher dimensions, *Phys. Rev. B* **69**, 104431 (2004), [arXiv:cond-mat/0305505 \[cond-mat.str-el\]](#).
- [3] M. B. Hastings and S. Michalakis, Quantization of hall conductance for interacting electrons on a torus, *Communications in Mathematical Physics* **334**, 10.1007/s00220-014-2167-x (2014), [arXiv:1306.1258 \[quant-ph\]](#).
- [4] S. Bravyi, M. B. Hastings, and F. Verstraete, Lieb-robinson bounds and the generation of correlations and topological quantum order, *Phys. Rev. Lett.* **97**, 050401 (2006), [arXiv:quant-ph/0603121](#).
- [5] T. J. Osborne, Simulating adiabatic evolution of gapped spin systems, *Phys. Rev. A* **75**, 032321 (2007), [arXiv:quant-ph/0601019](#).
- [6] Z. Gong and R. Hamazaki, Bounds in nonequilibrium quantum dynamics, *International Journal of Modern Physics B* **36**, 10.1142/s0217979222300079 (2022), [arXiv:2202.02011 \[quant-ph\]](#).
- [7] C.-F. (Anthony) Chen, A. Lucas, and C. Yin, Speed limits and locality in many-body quantum dynamics, *Reports on Progress in Physics* **86**, 116001 (2023), [arXiv:2303.07386 \[quant-ph\]](#).
- [8] B. Nachtergaele, R. Sims, and A. Young, Quasi-locality bounds for quantum lattice systems. i. lieb-robinson bounds, quasi-local maps, and spectral flow automorphisms, *Journal of Mathematical Physics* **60**, 10.1063/1.5095769 (2019), [arXiv:1810.02428 \[math-ph\]](#).
- [9] D. Basko, I. Aleiner, and B. Altshuler, Metal-insulator transition in a weakly interacting many-electron system with localized single-particle states, *Annals of Physics* **321**, 1126–1205 (2006), [arXiv:cond-mat/0506617 \[math-ph\]](#).
- [10] I. V. Gornyi, A. D. Mirlin, and D. G. Polyakov, Interacting electrons in disordered wires: Anderson localization and low- t transport, *Phys. Rev. Lett.* **95**, 206603 (2005), [arXiv:cond-mat/0506411 \[math-ph\]](#).
- [11] V. Oganesyan and D. A. Huse, Localization of interacting fermions at high temperature, *Phys. Rev. B* **75**, 155111 (2007), [arXiv:cond-mat/0610854 \[cond-mat.str-el\]](#).
- [12] M. Žnidarič, T. Prosen, and P. Prelovšek, Many-body localization in the heisenberg xxz magnet in a random field, *Phys. Rev. B* **77**, 064426 (2008), [arXiv:0706.2539 \[quant-ph\]](#).
- [13] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, Colloquium : Many-body localization, thermalization, and entanglement, *Reviews of Modern Physics* **91**, 10.1103/revmodphys.91.021001 (2019), [arXiv:1804.11065 \[cond-mat.dis-nn\]](#).
- [14] W. De Roeck and F. Huveneers, Stability and instability towards delocalization in many-body localization systems, *Phys. Rev. B* **95**, 155129 (2017), [arXiv:1608.01815 \[cond-mat.dis-nn\]](#).
- [15] D. J. Luitz, F. Huveneers, and W. De Roeck, How a small quantum bath can thermalize long localized chains, *Physical Review Letters* **119**, 10.1103/physrevlett.119.150602 (2017), [arXiv:1705.10807 \[cond-mat.str-el\]](#).
- [16] I. H. Kim, A. Chandran, and D. A. Abanin, Local integrals of motion and the logarithmic lightcone in many-body localized systems (2014), [arXiv:1412.3073 \[cond-mat.dis-nn\]](#).
- [17] B. Nachtergaele and J. Reschke, Slow propagation in some disordered quantum spin chains, *Journal of Statistical Physics* **182**, 10.1007/s10955-020-02681-2 (2021), [arXiv:1906.10167 \[math-ph\]](#).
- [18] Y. Zeng, A. Hamma, Y.-R. Zhang, Q. Liu, R. Li, H. Fan, and W.-M. Liu, Logarithmic light cone, slow entanglement growth and scrambling, and quantum memory (2023), [arXiv:2305.08334 \[quant-ph\]](#).
- [19] A. Elgart and A. Klein, Slow propagation of information on the random xxz quantum spin chain (2023), [arXiv:2311.14188 \[math-ph\]](#).
- [20] E. Hamza, R. Sims, and G. Stolz, Dynamical localization in disordered quantum spin systems, *Commun. Math. Phys.* **315**, 215 (2012), [arXiv:1108.3811 \[math-ph\]](#).
- [21] G. Stolz, An introduction to the mathematics of anderson localization, *Contemp Math* **552**, 71 (2011), [arXiv:1104.2317 \[math-ph\]](#).
- [22] R. del Rio, S. Jitomirskaya, Y. Last, and B. Simon, What is localization?, *Phys. Rev. Lett.* **75**, 117 (1995).
- [23] W. De Roeck, F. Huveneers, and S. Olla, Subdiffusion in one-dimensional hamiltonian chains with sparse interactions, *Journal of Statistical Physics* **180**, 678–698 (2020), [arXiv:1909.07322v2 \[math-ph\]](#).
- [24] Y. Huang, Adding boundary terms to anderson localized hamiltonians leads to unbounded growth of entanglement, *Europhysics Letters* **142**, 10001 (2023), [arXiv:2109.07640 \[cond-mat.dis-nn\]](#).

- [25] P. Bordia, H. Lüschen, S. Scherg, S. Gopalakrishnan, M. Knap, U. Schneider, and I. Bloch, Probing slow relaxation and many-body localization in two-dimensional quasiperiodic systems, *Phys. Rev. X* **7**, 041047 (2017), [arXiv:1704.03063 \[cond-mat.quant-gas\]](#).
- [26] J.-y. Choi, S. Hild, J. Zeiher, P. Schauß, A. Rubio-Abadal, T. Yefsah, V. Khemani, D. A. Huse, I. Bloch, and C. Gross, Exploring the many-body localization transition in two dimensions, *Science* **352**, 1547–1552 (2016), [arXiv:1604.04178 \[cond-mat.quant-gas\]](#).
- [27] T. B. Wahl, A. Pal, and S. H. Simon, Signatures of the many-body localized regime in two dimensions, *Nature Physics* **15**, 164–169 (2018), [arXiv:1711.02678 \[cond-mat.dis-nn\]](#).
- [28] M. Gebert, A. Moon, and B. Nachtergaele, A lieb–robinson bound for quantum spin chains with strong on-site impurities, *Reviews in Mathematical Physics* **34**, 10.1142/s0129055x22500076 (2022), [arXiv:arXiv:2104.00968v1 \[math-ph\]](#).
- [29] A. Morningstar, L. Colmenarez, V. Khemani, D. J. Luitz, and D. A. Huse, Avalanches and many-body resonances in many-body localized systems, *Physical Review B* **105**, 10.1103/physrevb.105.174205 (2022), [arXiv:2107.05642 \[cond-mat.dis-nn\]](#).
- [30] H. Ha, A. Morningstar, and D. A. Huse, Many-body resonances in the avalanche instability of many-body localization, *Physical Review Letters* **130**, 10.1103/physrevlett.130.250405 (2023), [arXiv:2301.04658 \[cond-mat.stat-mech\]](#).
- [31] M. Goihl, J. Eisert, and C. Krumnow, Exploration of the stability of many-body localized systems in the presence of a small bath, *Phys. Rev. B* **99**, 195145 (2019).
- [32] I.-D. Potirniche, S. Banerjee, and E. Altman, Exploration of the stability of many-body localization in d larger than 1, *Phys. Rev. B* **99**, 205149 (2019), [arXiv:1805.01475 \[cond-mat.dis-nn\]](#).
- [33] L. Pastur and V. Slavin, Area law scaling for the entropy of disordered quasifree fermions, *Physical Review Letters* **113**, 10.1103/physrevlett.113.150404 (2014), [arXiv:1408.2570 \[quant-ph\]](#).
- [34] H. Abdul-Rahman and G. Stolz, A uniform area law for the entanglement of eigenstates in the disordered xy chain, *Journal of Mathematical Physics* **56**, 10.1063/1.4938573 (2015), [arXiv:1505.02117 \[math-ph\]](#).
- [35] J. H. Bardarson, F. Pollmann, and J. E. Moore, Unbounded growth of entanglement in models of many-body localization, *Physical Review Letters* **109**, 10.1103/physrevlett.109.017202 (2012), [arXiv:1202.5532 \[cond-mat.str-el\]](#).
- [36] M. Serbyn, Z. Papić, and D. A. Abanin, Universal slow growth of entanglement in interacting strongly disordered systems, *Physical Review Letters* **110**, 10.1103/physrevlett.110.260601 (2013), [arXiv:1304.4605 \[cond-mat.str-el\]](#).
- [37] A. Nico-Katz, A. Bayat, and S. Bose, Information-theoretic memory scaling in the many-body localization transition, *Physical Review B* **105**, 10.1103/physrevb.105.205133 (2022), [arXiv:2009.04470 \[quant-ph\]](#).
- [38] D. Damanik and S. Tcheremchantsev, Quantum dynamics via complex analysis methods: General upper bounds without time-averaging and tight lower bounds for the strongly coupled fibonacci hamiltonian, *Journal of Functional Analysis* **255**, 2872 (2008), [arXiv:0801.3399 \[math.SP\]](#).
- [39] H. Kim and D. A. Huse, Ballistic spreading of entanglement in a diffusive nonintegrable system, *Physical Review Letters* **111**, 10.1103/physrevlett.111.127205 (2013), [arXiv:1306.4306 \[quant-ph\]](#).
- [40] T. Rakovszky, F. Pollmann, and C. von Keyserlingk, Diffusive hydrodynamics of out-of-time-ordered correlators with charge conservation, *Physical Review X* **8**, 10.1103/physrevx.8.031058 (2018), [arXiv:1710.09827 \[cond-mat.stat-mech\]](#).
- [41] T. Prosen, Open xxz spin chain: Nonequilibrium steady state and a strict bound on ballistic transport, *Phys. Rev. Lett.* **106**, 217206 (2011), [arXiv:1103.1350 \[cond-mat.str-el\]](#).
- [42] V. Khemani, R. Nandkishore, and S. L. Sondhi, Nonlocal adiabatic response of a localized system to local manipulations, *Nature Physics* **11**, 560–565 (2015), [arXiv:1411.2616 \[cond-mat.dis-nn\]](#).
- [43] L. Masanes, Area law for the entropy of low-energy states, *Physical Review A* **80**, 10.1103/physreva.80.052104 (2009), [arXiv:0907.4672 \[quant-ph\]](#).
- [44] M. Foss-Feig, Z.-X. Gong, C. W. Clark, and A. V. Gorshkov, Nearly linear light cones in long-range interacting quantum systems, *Phys. Rev. Lett.* **114**, 157201 (2015), [arXiv:1410.3466 \[quant-ph\]](#).
- [45] T. Kuwahara and K. Saito, Strictly linear light cones in long-range interacting systems of arbitrary dimensions, *Phys. Rev. X* **10**, 031010 (2020), [arXiv:1910.14477 \[quant-ph\]](#).
- [46] M. B. Hastings, Locality in quantum systems (2010), [arXiv:1008.5137 \[math-ph\]](#).
- [47] M. C. Tran, A. Y. Guo, Y. Su, J. R. Garrison, Z. Eldredge, M. Foss-Feig, A. M. Childs, and A. V. Gorshkov, Locality and digital quantum simulation of power-law interactions, *Physical Review X* **9**, 10.1103/physrevx.9.031006 (2019), [arXiv:1808.05225 \[quant-ph\]](#).