

CARLESON OPERATORS ON DOUBLING METRIC MEASURE SPACES

LARS BECKER, FLORIS VAN DOORN, ASGAR JAMNESHAN,
RAJULA SRIVASTAVA, AND CHRISTOPH THIELE

ABSTRACT. We prove a new generalization of a theorem of Carleson, namely bounds for a generalized Carleson operator on doubling metric measure spaces. Additionally, we explicitly reduce Carleson's classical result on pointwise convergence of Fourier series to this new theorem. Both proofs are presented in great detail, suitable as a blueprint for computer verification using the current capabilities of the software package Lean. Note that even Carleson's classical result has not yet been computer-verified.

CONTENTS

1. Introduction	2
2. Proof of metric space Carleson (Thm 1.2), overview	5
2.1. Auxiliary lemmas	11
3. Proof of metric space Carleson (Thm 1.2)	13
4. Proof of finitary Carleson (Prop 2.1)	17
4.1. Proof of Lemma 4.1, dyadic structure	18
4.2. Proof of Lemma 4.2, tile structure	24
5. Proof of discrete Carleson (Prop. 2.2)	27
5.1. Organisation of the tiles	27
5.2. Proof of Lemma 5.1, the exceptional sets	30
5.3. Auxiliary lemmas	35
5.4. Proof of Lemma 5.2, the forests	38
5.5. Proof of Lemma 5.3, the antichains	43
6. Proof of Proposition 2.3	45
6.1. The density arguments	46
6.2. Proof of Lemma 6.5, the tile correlation bound	50
6.3. Proof of Lemma 6.6, the geometric estimate	53
7. Proof of Proposition 2.4	57
7.1. The pointwise tree estimate	57
7.2. An auxiliary L^2 tree estimate	62
7.3. The quantitative L^2 tree estimate	67
7.4. Almost orthogonality of separated trees	70
7.5. Proof of Lemma 7.18: Tiles with large separation	72
7.6. Proof of Lemma 7.19: The remaining tiles	81
7.7. Forests	85
8. Proof of Prop. 2.5, the Hölder cancellative condition	89
9. Proof of Prop. 2.6, Vitali covering and Hardy–Littlewood	92

Date: May 13, 2024.

10.	Proof of Theorem 1.1	96
10.1.	The classical Carleson theorem	97
10.2.	Piecewise constant functions.	102
10.3.	Proof of Cotlar's Inequality	105
10.4.	The truncated Hilbert transform	109
10.5.	Calderón-Zygmund Decomposition	114
10.6.	The proof of the van der Corput Lemma 10.8	122
10.7.	Partial sums as orthogonal projections	122
10.8.	The error bound	124
10.9.	Carleson on the real line	126
	References	128

1. INTRODUCTION

This preprint is associated with a distributed Lean project accessible under <https://florisvandoorn.com/carleson/>.

In [Car66], L. Carleson addressed a classical question regarding the convergence of Fourier series of continuous functions by proving their pointwise convergence almost everywhere. Theorem 1.1 represents a version of this result.

Let f be a complex-valued, 2π -periodic bounded Borel measurable function on the real line, and for an integer n , define the Fourier coefficient as

$$\widehat{f}_n := \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{-inx} dx. \quad (1.1)$$

Define the partial Fourier sum for $N \geq 0$ as

$$S_N f(x) := \sum_{n=-N}^N \widehat{f}_n e^{inx}. \quad (1.2)$$

Theorem 1.1 (classical Carleson). *Let f be a 2π -periodic complex-valued uniformly continuous function on \mathbb{R} satisfying the bound $|f(x)| \leq 1$ for all $x \in \mathbb{R}$. For all $0 < \epsilon < 1$, there exists a Borel set $E \subset [0, 2\pi]$ with Lebesgue measure $|E| \leq \epsilon$ and a positive integer N_0 such that for all $x \in [0, 2\pi] \setminus E$ and all integers $N > N_0$, we have*

$$|f(x) - S_N f(x)| \leq \epsilon. \quad (1.3)$$

Note that mere continuity implies uniform continuity in the setting of this theorem. By applying this theorem with a sequence of $\epsilon_n := 2^{-n}\delta$ for $n \geq 1$ and taking the union of corresponding exceptional sets E_n , we see that outside a set of measure δ , the partial Fourier sums converge pointwise for $N \rightarrow \infty$. Applying this with a sequence of δ shrinking to zero and taking the intersection of the corresponding exceptional sets, which has measure zero, we see that the Fourier series converges outside a set of measure zero.

The purpose of this paper is twofold. On the one hand, it prepares computer verification of Theorem 1.1 by presenting a very detailed proof as a blueprint for coding in Lean. We pass through a bound for a generalization of the so-called Carleson operator to doubling metric measure spaces.

This generalization is new, and proving these bounds constitutes the second purpose of this paper. This generalization incorporates several results from the recent literature, most prominently bounds for the polynomial Carleson operator of V. Lie [Lie20] as well as its generalization [Zor21]. A computer verification of our theorem will also entail a computer verification for the bulk of the work in these results.

We proceed to introduce the setup for our general theorem. We carry a multi purpose parameter, a natural number

$$a \geq 4 \tag{1.4}$$

in our notation that as it gets larger will allow more general applications but will worsen the constants in the estimates.

A doubling metric measure space (X, ρ, μ, a) is a complete and locally compact metric space (X, ρ) equipped with a σ -finite non-zero Radon–Borel measure μ that satisfies the doubling condition that for all $x \in X$ and all $R > 0$ we have

$$\mu(B(x, 2R)) \leq 2^a \mu(B(x, R)), \tag{1.5}$$

where we have denoted by $B(x, R)$ the open ball of radius R centred at x :

$$B(x, R) := \{y \in X : \rho(x, y) < R\}. \tag{1.6}$$

A collection Θ of real valued continuous functions on the doubling metric measure space (X, ρ, μ, a) is called compatible, if there is a point $o \in X$ where all the functions are equal to 0, and if there exists for each ball $B \subset X$ a metric d_B on Θ , such that the following five properties (1.7), (1.8), (1.9), (1.10), and (1.11) are satisfied. For every ball $B \subset X$

$$\sup_{x, y \in B} |\vartheta(x) - \vartheta(y) - \theta(x) + \theta(y)| \leq d_B(\vartheta, \theta). \tag{1.7}$$

For any two balls $B_1 = B(x_1, R)$, $B_2 = B(x_2, 2R)$ in X with $x_1 \in B_2$ and any $\vartheta, \theta \in \Theta$,

$$d_{B_2}(\vartheta, \theta) \leq 2^a d_{B_1}(\vartheta, \theta). \tag{1.8}$$

For any two balls B_1, B_2 in X with $B_1 \subset B_2$ and any $\vartheta, \theta \in \Theta$

$$d_{B_1}(\vartheta, \theta) \leq d_{B_2}(\vartheta, \theta) \tag{1.9}$$

and for any two balls $B_1 = B(x_1, R)$, $B_2 = B(x_2, 2^a R)$ with $B_1 \subset B_2$, and $\vartheta, \theta \in \Theta$,

$$2d_{B_1}(\vartheta, \theta) \leq d_{B_2}(\vartheta, \theta). \tag{1.10}$$

For every ball B in X and every d_B -ball \tilde{B} of radius $2R$ in Θ , there is a collection \mathcal{B} of at most 2^a many d_B -balls of radius R covering \tilde{B} , that is,

$$\tilde{B} \subset \bigcup \mathcal{B}. \tag{1.11}$$

Further, a compatible collection Θ is called cancellative, if for any ball B in X of radius R , any Lipschitz function $\varphi : X \rightarrow \mathbb{C}$ supported on B , and any $\vartheta, \theta \in \Theta$ we have

$$\left| \int_B e(\vartheta(x) - \theta(x)) \varphi(x) d\mu(x) \right| \leq 2^a \mu(B) \|\varphi\|_{\text{Lip}(B)} (1 + d_B(\vartheta, \theta))^{-\frac{1}{a}}, \tag{1.12}$$

where $\|\cdot\|_{\text{Lip}(B)}$ denotes the inhomogeneous Lipschitz norm on B :

$$\|\varphi\|_{\text{Lip}(B)} = \sup_{x \in B} |\varphi(x)| + R \sup_{x, y \in B, x \neq y} \frac{|\varphi(x) - \varphi(y)|}{\rho(x, y)}.$$

A one-sided Calderón–Zygmund kernel K on the doubling metric measure space (X, ρ, μ, a) is a measurable function

$$K : X \times X \rightarrow \mathbb{C} \quad (1.13)$$

such that for all $x, y', y \in X$ with $x \neq y$, we have

$$|K(x, y)| \leq \frac{2^{a^3}}{V(x, y)} \quad (1.14)$$

and if $2\rho(y, y') \leq \rho(x, y)$, then

$$|K(x, y) - K(x, y')| \leq \left(\frac{\rho(y, y')}{\rho(x, y)} \right)^{\frac{1}{a}} \frac{2^{a^3}}{V(x, y)}, \quad (1.15)$$

where

$$V(x, y) := \mu(B(x, \rho(x, y))).$$

Define the maximally truncated non-tangential singular integral T_* associated with K by

$$T_* f(x) := \sup_{R_1 < R_2} \sup_{\rho(x, x') < R_1} \left| \int_{R_1 < \rho(x', y) < R_2} K(x', y) f(y) d\mu(y) \right|. \quad (1.16)$$

We define the generalized Carleson operator T by

$$Tf(x) := \sup_{\vartheta \in \Theta} \sup_{0 < R_1 < R_2} \left| \int_{R_1 < \rho(x, y) < R_2} K(x, y) f(y) e(\vartheta(y)) d\mu(y) \right|, \quad (1.17)$$

where $e(r) = e^{ir}$.

Our main result is the following restricted weak type estimate for T in the range $1 < q \leq 2$, which by interpolation techniques recovers L^q estimates for the open range $1 < q < 2$.

Theorem 1.2 (metric space Carleson). *For all integers $a \geq 4$ and real numbers $1 < q \leq 2$ the following holds. Let (X, ρ, μ, a) be a doubling metric measure space. Let Θ be a cancellative compatible collection of functions and let K be a one-sided Calderón–Zygmund kernel on (X, ρ, μ, a) . Assume that for every bounded measurable function g on X supported on a set of finite measure we have*

$$\|T_* g\|_2 \leq 2^{a^3} \|g\|_2, \quad (1.18)$$

where T_* is defined in (1.16). Then for all Borel sets F and G in X and all Borel functions $f : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$, we have, with T defined in (1.17),

$$\left| \int_G T f d\mu \right| \leq \frac{2^{450a^3}}{(q-1)^5} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \quad (1.19)$$

In the one-dimensional Euclidean setting, with K representing the Hilbert kernel:

$$K(x, y) = (x - y)^{-1}$$

and Θ denoting the class of linear functions, the operator (1.17) is the classical Carleson operator, which plays a crucial role in proving the almost

everywhere convergence of Fourier series [Car66], [Fef73], [LT00]. The supremum in R_1 and R_2 is often omitted in classical treatments, but considering the maximal truncations can easily be reduced to the case without these truncations.

By replacing Θ with the class of polynomials vanishing at 0 up to some fixed but arbitrary degree, we obtain the polynomial Carleson operator of Lie [Lie09] (quadratic case) and [Lie20]. The case of the class of polynomials with vanishing linear coefficient is simpler and was estimated in [SW01]. The polynomial Carleson operator was generalized to the high-dimensional Euclidean setting in [Zor21] for K being a Calderón-Zygmund kernel with some Hölder regularity.

Doubling metric measure spaces are instances of spaces of homogeneous type. Indeed, by changing from a quasi-metric to an equivalent metric, every space of homogeneous type can be viewed as a doubling metric measure space (cf. [MS79]). Spaces of homogeneous type were introduced by [CW71] as a natural setting for Calderón-Zygmund theory. We refer to the textbook [Ste93] for an account of these spaces.

Our concept of a compatible collection Θ as a natural class of phase functions on a doubling metric measure space does not appear in [Ste93] but is implicitly anticipated in [Zor21] and subsequent work of [Mna22], who proves a Carleson-type theorem for the Malmquist-Takenaka series, which leads to classes of phases related to Blaschke products. A generalization of (1.17) from the previously mentioned Euclidean setting into the anisotropic setting that was suggested in [Zor21] is included in our theory. The polynomial Carleson operator also plays a role in the study of maximally modulated singular Radon transforms along the parabola, see [Ram21] and [Bec24].

For the proof of Theorem 1.2, we largely follow [Zor21], which in turn was inspired by [Lie20]. We make suitable modifications to adapt to our more general setting and have made a few technical improvements in the proof. In particular, in Section 2, we explicitly divide the main work of the proof into mutually independent sections 3, 4, 5, 6, 7, 8, and 9. Some of these sections follow a similar pattern, starting with a subsection dividing the proof into further mutually independent subsections. This modularization of our proof was strongly endorsed in personal communication by the author of [Zor21]. *Acknowledgement.* L.B., F.v.D., R.S., and C.T. were funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC-2047/1 – 390685813. L.B., R.S., and C.T. were also supported by SFB 1060. A.J. is funded by the TÜBİTAK (Scientific and Technological Research Council of Türkiye) under Grant Number 123F122.

2. PROOF OF METRIC SPACE CARLESON (THM 1.2), OVERVIEW

This section organizes the proof of Theorem 1.2 into sections 3, 4, 5, 6, 7, 8, and 9. These sections are mutually independent except for referring to the statements formulated in the present section. Section 3 proves the main Theorem 1.2, while sections 4, 5, 6, 7, 8, and 9 each prove one proposition that is stated in the present section. The present section also introduces all definitions used across these sections.

Subsection 2.1 proves some auxiliary lemmas that are used in more than one of the sections 3-9.

Let a, q be given as in Theorem 1.2.

Define

$$D := 2^{100a^2}, \quad (2.1)$$

$$\kappa := 2^{-10a}, \quad (2.2)$$

and

$$Z := 2^{12a}. \quad (2.3)$$

Let $\psi : \mathbb{R} \rightarrow \mathbb{R}$ be the unique compactly supported, piece-wise linear, continuous function with corners precisely at $\frac{1}{4D}$, $\frac{1}{2D}$, $\frac{1}{4}$ and $\frac{1}{2}$ which satisfies

$$\sum_{s \in \mathbb{Z}} \psi(D^{-s}x) = 1 \quad (2.4)$$

for all $x > 0$. This function vanishes outside $[\frac{1}{4D}, \frac{1}{2}]$, is constant one on $[\frac{1}{2D}, \frac{1}{4}]$, and is Lipschitz with constant $4D$.

Let a doubling metric measure space (X, ρ, μ, a) be given. Let a cancellative compatible collection Θ of functions on X be given. Let $o \in X$ be a point such that $\vartheta(o) = 0$ for all $\vartheta \in \Theta$.

Let a one-sided Calderón–Zygmund kernel K on X be given so that the operator T_* defined in (1.16) satisfies (1.18). Let T be the corresponding operator as defined in (1.17).

For $s \in \mathbb{Z}$, we define

$$K_s(x, y) := K(x, y)\psi(D^{-s}\rho(x, y)), \quad (2.5)$$

so that for each $x, y \in X$ with $x \neq y$ we have

$$K(x, y) = \sum_{s \in \mathbb{Z}} K_s(x, y).$$

In Section 3, we prove Theorem 1.2 from its finitary version, Proposition 2.1 below. Recall that a function from a measure space to a finite set is measurable if the pre-image of each of the elements in the range is measurable.

Proposition 2.1 (finitary Carleson). *Let $\sigma_1, \sigma_2 : X \rightarrow \mathbb{Z}$ be measurable functions with finite range and $\sigma_1 \leq \sigma_2$. Let $Q : X \rightarrow \Theta$ be a measurable function with finite range. Let F, G be bounded Borel sets in X . Then there is a Borel set G' in X with $2\mu(G') \leq \mu(G)$ such that for all Borel functions $f : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$.*

$$\begin{aligned} \int_{G \setminus G'} \left| \sum_{s=\sigma_1(x)}^{\sigma_2(x)} \int K_s(x, y) f(y) e(Q(x)(y)) d\mu(y) \right| d\mu(x) \\ \leq \frac{2^{440a^3}}{(q-1)^4} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \end{aligned} \quad (2.6)$$

Let measurable functions $\sigma_1 \leq \sigma_2 : X \rightarrow \mathbb{Z}$ with finite range be given. Let a measurable function $Q : X \rightarrow \Theta$ with finite range be given. Let bounded Borel sets F, G in X be given. Let S be the smallest integer such that the ranges of σ_1 and σ_2 are contained in $[-S, S]$ and F and G are contained in the ball $B(o, D^S)$.

In Section 4, we prove Proposition 2.1 using a bound for a dyadic model formulated in Proposition 2.2 below.

A grid structure (\mathcal{D}, c, s) on X consists of a finite collection \mathcal{D} of Borel sets in X called dyadic cubes, a surjective function $s: \mathcal{D} \rightarrow [-S, S]$ called scale function, and a function $c: \mathcal{D} \rightarrow X$ called center function such that the five properties (2.7), (2.8), (2.9), (2.10), and (2.11) hold.

For each dyadic cube I and each $-S \leq k < s(I)$ we have

$$I \subset \bigcup_{J \in \mathcal{D}: s(J)=k} J. \quad (2.7)$$

Any two non-disjoint dyadic cubes I, J with $s(I) \leq s(J)$ satisfy

$$I \subset J. \quad (2.8)$$

For any $x \in B(o, D^S)$, and every $k \in [-S, S]$, there is a dyadic cube I with $s(I) = k$ and

$$x \in I. \quad (2.9)$$

For any dyadic cube I ,

$$c(I) \in B(c(I), \frac{1}{4}D^{s(I)}) \subset I \subset B(c(I), 4D^{s(I)}). \quad (2.10)$$

For any dyadic cube I and any t with $tD^{s(I)} \geq D^{-S}$,

$$\mu(\{x \in I : \rho(x, X \setminus I) \leq tD^{s(I)}\}) \leq 2^{100a^2} t^\kappa \mu(I). \quad (2.11)$$

A tile structure $(\mathfrak{P}, \mathcal{I}, \Omega, \mathcal{Q}, c, s)$ for a given grid structure (\mathcal{D}, c, s) is a finite set \mathfrak{P} of elements called tiles with five maps

$$\begin{aligned} \mathcal{I}: \mathfrak{P} &\rightarrow \mathcal{D} \\ \Omega: \mathfrak{P} &\rightarrow \mathcal{P}(\Theta) \\ \mathcal{Q}: \mathfrak{P} &\rightarrow \Theta \\ c: \mathfrak{P} &\rightarrow X \\ s: \mathfrak{P} &\rightarrow \mathbb{Z} \end{aligned}$$

with \mathcal{I} surjective and $\mathcal{P}(\Theta)$ denoting the power set of Θ such that the five properties (2.13), (2.14), (2.15), (2.18), and (2.19) hold. For each dyadic cube I , the restriction of the map Ω to the set

$$\mathfrak{P}(I) = \{\mathfrak{p} : \mathcal{I}(\mathfrak{p}) = I\} \quad (2.12)$$

is injective and we have the disjoint covering property (we use the union symbol with dot on top to denote a disjoint union)

$$\mathcal{Q}(X) \subset \dot{\bigcup}_{\mathfrak{p} \in \mathfrak{P}(I)} \Omega(\mathfrak{p}). \quad (2.13)$$

For any tiles $\mathfrak{p}, \mathfrak{q}$ with $\mathcal{I}(\mathfrak{p}) \subset \mathcal{I}(\mathfrak{q})$ and $\Omega(\mathfrak{p}) \cap \Omega(\mathfrak{q}) \neq \emptyset$ we have

$$\Omega(\mathfrak{q}) \subset \Omega(\mathfrak{p}). \quad (2.14)$$

For each tile \mathfrak{p} ,

$$\mathcal{Q}(\mathfrak{p}) \in B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), 0.2) \subset \Omega(\mathfrak{p}) \subset B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), 1), \quad (2.15)$$

where

$$B_{\mathfrak{p}}(\vartheta, R) := \{\theta \in \Theta : d_{\mathfrak{p}}(\vartheta, \theta) < R\}, \quad (2.16)$$

and

$$d_{\mathbf{p}} := d_{B(c(\mathbf{p}), \frac{1}{4}D^s(\mathbf{p}))}. \quad (2.17)$$

We have for each tile \mathbf{p}

$$c(\mathbf{p}) = c(\mathcal{I}(\mathbf{p})), \quad (2.18)$$

$$s(\mathbf{p}) = s(\mathcal{I}(\mathbf{p})). \quad (2.19)$$

Proposition 2.2 (discrete Carleson). *Let (\mathcal{D}, c, s) be a grid structure and*

$$(\mathfrak{P}, \mathcal{I}, \Omega, \mathcal{Q}, c, s)$$

a tile structure for this grid structure. Define for $\mathbf{p} \in \mathfrak{P}$

$$E(\mathbf{p}) = \{x \in \mathcal{I}(\mathbf{p}) : Q(x) \in \Omega(\mathbf{p}), \sigma_1(x) \leq s(\mathbf{p}) \leq \sigma_2(x)\} \quad (2.20)$$

and

$$T_{\mathbf{p}}f(x) = \mathbf{1}_{E(\mathbf{p})}(x) \int K_{s(\mathbf{p})}(x, y) f(y) e(Q(x)(y) - Q(x)(x)) d\mu(y). \quad (2.21)$$

Then there exists a Borel set G' with $2\mu(G') \leq \mu(G)$ such that for all $f : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$ we have

$$\int_{G \setminus G'} \left| \sum_{\mathbf{p} \in \mathfrak{P}} T_{\mathbf{p}}f(x) \right| d\mu(x) \leq \frac{2^{440a^3}}{(q-1)^4} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \quad (2.22)$$

The proof of Proposition 2.2 is done in Section 5 by a reduction to two further propositions that we state below.

Fix a grid structure (\mathcal{D}, c, s) and a tile structure $(\mathfrak{P}, \mathcal{I}, \Omega, \mathcal{Q}, c, s)$ for this grid structure.

We define the relation

$$\mathbf{p} \leq \mathbf{p}' \quad (2.23)$$

on $\mathfrak{P} \times \mathfrak{P}$ meaning $\mathcal{I}(\mathbf{p}) \subset \mathcal{I}(\mathbf{p}')$ and $\Omega(\mathbf{p}') \subset \Omega(\mathbf{p})$. We further define for $\lambda, \lambda' > 0$ the relation

$$\lambda \mathbf{p} \lesssim \lambda' \mathbf{p}' \quad (2.24)$$

on $\mathfrak{P} \times \mathfrak{P}$ meaning $\mathcal{I}(\mathbf{p}) \subset \mathcal{I}(\mathbf{p}')$ and

$$B_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), \lambda') \subset B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \lambda). \quad (2.25)$$

Define for a tile \mathbf{p} and $\lambda > 0$

$$E_1(\mathbf{p}) := \{x \in \mathcal{I}(\mathbf{p}) \cap G : Q(x) \in \Omega(\mathbf{p})\}, \quad (2.26)$$

$$E_2(\lambda, \mathbf{p}) := \{x \in \mathcal{I}(\mathbf{p}) \cap G : Q(x) \in B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \lambda)\}. \quad (2.27)$$

Given a subset \mathfrak{P}' of \mathfrak{P} , we define $\mathfrak{P}(\mathfrak{P}')$ to be the set of all $\mathbf{p} \in \mathfrak{P}$ such that there exist $\mathbf{p}' \in \mathfrak{P}'$ with $\mathcal{I}(\mathbf{p}) \subset \mathcal{I}(\mathbf{p}')$. Define the densities

$$\text{dens}_1(\mathfrak{P}') := \sup_{\mathbf{p}' \in \mathfrak{P}'} \sup_{\lambda \geq 2} \lambda^{-a} \sup_{\mathbf{p} \in \mathfrak{P}(\mathfrak{P}'), \lambda \mathbf{p} \lesssim \lambda' \mathbf{p}'} \frac{\mu(E_2(\lambda, \mathbf{p}))}{\mu(\mathcal{I}(\mathbf{p}))}, \quad (2.28)$$

$$\text{dens}_2(\mathfrak{P}') := \sup_{\mathbf{p}' \in \mathfrak{P}'} \sup_{r \geq 4D^s(\mathbf{p}')} \frac{\mu(F \cap B(c(\mathbf{p}'), r))}{\mu(B(c(\mathbf{p}'), r))}. \quad (2.29)$$

An antichain is a subset \mathfrak{A} of \mathfrak{P} such that for any distinct $\mathbf{p}, \mathbf{q} \in \mathfrak{A}$ we do not have $\mathbf{p} \leq \mathbf{q}$.

The following proposition is proved in Section 6.

Proposition 2.3 (antichain operator). *For any antichain \mathfrak{A} and for all $f : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$ and all $g : X \rightarrow \mathbb{C}$ with $|g| \leq \mathbf{1}_G$*

$$|\int \overline{g(x)} \sum_{\mathfrak{p} \in \mathfrak{A}} T_{\mathfrak{p}} f(x) d\mu(x)| \quad (2.30)$$

$$\leq \frac{2^{150a^3}}{q-1} \text{dens}_1(\mathfrak{A})^{\frac{q-1}{8a^4}} \text{dens}_2(\mathfrak{A})^{\frac{1}{q}-\frac{1}{2}} \|f\|_2 \|g\|_2. \quad (2.31)$$

Let $n \geq 0$. An n -forest is a pair $(\mathfrak{U}, \mathfrak{T})$ where \mathfrak{U} is a subset of \mathfrak{P} and \mathfrak{T} is a map assigning to each $\mathfrak{u} \in \mathfrak{U}$ a nonempty set $\mathfrak{T}(\mathfrak{u}) \subset \mathfrak{P}$ called tree such that the following properties (2.32), (2.33), (2.34), (2.35), (2.36), and (2.37) hold.

For each $\mathfrak{u} \in \mathfrak{U}$ and each $\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})$ we have

$$4\mathfrak{p} \lesssim 1\mathfrak{u}. \quad (2.32)$$

For each $\mathfrak{u} \in \mathfrak{U}$ and each $\mathfrak{p}, \mathfrak{p}'' \in \mathfrak{T}(\mathfrak{u})$ and $\mathfrak{p}' \in \mathfrak{P}$ we have

$$\mathfrak{p}, \mathfrak{p}'' \in \mathfrak{T}(\mathfrak{u}), \mathfrak{p} \leq \mathfrak{p}' \leq \mathfrak{p}'' \implies \mathfrak{p}' \in \mathfrak{T}(\mathfrak{u}). \quad (2.33)$$

We have

$$\left\| \sum_{\mathfrak{u} \in \mathfrak{U}} \mathbf{1}_{\mathcal{I}(\mathfrak{u})} \right\|_{\infty} \leq 2^n. \quad (2.34)$$

We have for every $\mathfrak{u} \in \mathfrak{U}$

$$\text{dens}_1(\mathfrak{T}(\mathfrak{u})) \leq 2^{4a+1-n}. \quad (2.35)$$

We have for $\mathfrak{u}, \mathfrak{u}' \in \mathfrak{U}$ with $\mathfrak{u} \neq \mathfrak{u}'$ and $\mathfrak{p} \in \mathfrak{T}(\mathfrak{u}')$ with $\mathcal{I}(\mathfrak{p}) \subset \mathcal{I}(\mathfrak{u})$ that

$$d_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), \mathcal{Q}(\mathfrak{u})) > 2^{Z(n+1)}. \quad (2.36)$$

We have for every $\mathfrak{u} \in \mathfrak{U}$ and $\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})$ that

$$B(c(\mathfrak{p}), 8D^{s(\mathfrak{p})}) \subset \mathcal{I}(\mathfrak{u}). \quad (2.37)$$

The following proposition is proved in Section 7.

Proposition 2.4 (forest operator). *For any $n \geq 0$ and any n -forest $(\mathfrak{U}, \mathfrak{T})$ we have for all $f : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$ and all bounded g with bounded support*

$$\begin{aligned} & |\int \overline{g(x)} \sum_{\mathfrak{u} \in \mathfrak{U}} \sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} T_{\mathfrak{p}} f(x) d\mu(x)| \\ & \leq 2^{432a^3} 2^{-\frac{q-1}{q}n} \text{dens}_2\left(\bigcup_{\mathfrak{u} \in \mathfrak{U}} \mathfrak{T}(\mathfrak{u})\right)^{\frac{1}{q}-\frac{1}{2}} \|f\|_2 \|g\|_2. \end{aligned}$$

Theorem 1.2 is formulated at the level of generality for general kernels satisfying the mere Hölder regularity condition (1.15). On the other hand, the cancellative condition (1.12) is a testing condition against more regular, namely Lipschitz functions. To bridge the gap, we follow [Zor21] to observe a variant of (1.12) that we formulate in the following proposition proved in Section 8.

Define

$$\tau := \frac{1}{a}. \quad (2.38)$$

Define for any open ball B of radius R in X the L^∞ -normalized τ -Hölder norm by

$$\|\varphi\|_{C^\tau(B)} = \sup_{x \in B} |\varphi(x)| + R^\tau \sup_{x, y \in B, x \neq y} \frac{|\varphi(x) - \varphi(y)|}{\rho(x, y)^\tau}. \quad (2.39)$$

Proposition 2.5 (Holder van der Corput). *Let $z \in X$ and $R > 0$ and set $B = B(z, R)$. Let $\varphi : X \rightarrow \mathbb{C}$ be supported on B and satisfy $\|\varphi\|_{C^\tau(B)} < \infty$. Let $\vartheta, \theta \in \Theta$. Then*

$$\left| \int e(\vartheta(x) - \theta(x)) \varphi(x) dx \right| \leq 2^{8a} \mu(B) \|\varphi\|_{C^\tau(B)} (1 + d_B(\vartheta, \theta))^{-\frac{1}{2a^2+a^3}}. \quad (2.40)$$

We further formulate a classical Vitali covering result and maximal function estimate that we need throughout several sections. This following proposition will typically be applied to the absolute value of a complex valued function and be proved in Section 9. By a ball B we mean a set $B(x, r)$ with $x \in X$ and $r > 0$ as defined in (1.6). For a finite collection \mathcal{B} of balls in X and $1 \leq p < \infty$ define the measurable function $M_{\mathcal{B}, p} u$ on X by

$$M_{\mathcal{B}, p} u(x) := \left(\sup_{B \in \mathcal{B}} \frac{\mathbf{1}_B(x)}{\mu(B)} \int_B |u(y)|^p d\mu(y) \right)^{\frac{1}{p}}. \quad (2.41)$$

Define further $M_{\mathcal{B}} := M_{\mathcal{B}, 1}$.

Proposition 2.6 (Hardy–Littlewood). *Let \mathcal{B} be a finite collection of balls in X . If for some $\lambda > 0$ and some measurable function $u : X \rightarrow [0, \infty)$ we have*

$$\int_B u(x) d\mu(x) \geq \lambda \mu(B) \quad (2.42)$$

for each $B \in \mathcal{B}$, then

$$\lambda \mu(\bigcup \mathcal{B}) \leq 2^{2a} \int_X u(x) d\mu(x). \quad (2.43)$$

For every measurable function v and $1 \leq p_1 < p_2$ we have

$$\|M_{\mathcal{B}, p_1} v\|_{p_2} \leq 2^{2a} \frac{p_2}{p_2 - p_1} \|v\|_{p_2}. \quad (2.44)$$

Moreover, given any measurable bounded function $w : X \rightarrow \mathbb{C}$ there exists a measurable function $Mw : X \rightarrow [0, \infty)$ such that the following (2.45) and (2.46) hold. For each ball $B \subset X$ and each $x \in B$

$$\frac{1}{\mu(B)} \int_B |w(y)| d\mu(y) \leq Mw(x) \quad (2.45)$$

and for all $1 < p_1 \leq p_2 \leq \infty$

$$\|M(w^{p_1})^{\frac{1}{p_1}}\|_{p_2} \leq 2^{4a} \frac{p_2}{p_2 - p_1} \|w\|_{p_2}. \quad (2.46)$$

This completes the overview of the proof of Theorem 1.2.

2.1. Auxiliary lemmas. We close this section by recording some auxiliary lemmas about the objects defined in Section 2, which will be used in multiple sections to follow.

First, we record an estimate for the metrical entropy numbers of balls in the space Θ equipped with any of the metrics d_B , following from the doubling property (1.11).

Lemma 2.7 (ball metric entropy). *Let $B' \subset X$ be a ball. Let $r > 0$, $\vartheta \in \Theta$ and $k \in \mathbb{N}$. Suppose that $\mathcal{Z} \subset B_{B'}(\vartheta, r2^k)$ satisfies that for all $z, z' \in \mathcal{Z}$ with $z \neq z'$, we have $d_{B'}(z, z') \geq r$. Then*

$$|\mathcal{Z}| \leq 2^{ka}.$$

Proof. By applying property (1.11) k times, we obtain a collection $\mathcal{Z}' \subset \Theta$ with $|\mathcal{Z}'| = 2^{ka}$ and

$$B_{B'}(\vartheta, r2^k) \subset \bigcup_{z' \in \mathcal{Z}'} B_{B'}(z', \frac{r}{2}).$$

Then each $z \in \mathcal{Z}$ is contained in one of the balls $B(z', \frac{r}{2})$, but by the separation assumption no such ball contains more than one element of \mathcal{Z} . Thus $|\mathcal{Z}| \leq |\mathcal{Z}'| = 2^{ka}$. \square

The next lemma concerns monotonicity of the metrics $d_{B(c(I), \frac{1}{4}D^{s(I)})}$ with respect to inclusion of cubes I in a grid.

Lemma 2.8 (monotone cube metrics). *Let (\mathcal{D}, c, s) be a grid structure. Denote for cubes $I \in \mathcal{D}$*

$$I^\circ := B(c(I), \frac{1}{4}D^{s(I)}).$$

Let $I, J \in \mathcal{D}$ with $I \subset J$. Then for all $\vartheta, \theta \in \Theta$ we have

$$d_{I^\circ}(\vartheta, \theta) \leq d_{J^\circ}(\vartheta, \theta),$$

and if $I \neq J$ then we have

$$d_{I^\circ}(\vartheta, \theta) \leq 2^{-95a} d_{J^\circ}(\vartheta, \theta).$$

Proof. If $s(I) \geq s(J)$ then (2.8) and the assumption $I \subset J$ imply $I = J$. Then the lemma holds by reflexivity.

If $s(J) \geq s(I) + 1$, then using the monotonicity property (1.9), (2.1) and (1.10), we get

$$d_{I^\circ}(\vartheta, \theta) \leq d_{B(c(I), 4D^{s(I)})}(\vartheta, \theta) \leq 2^{-100a} d_{B(c(I), 4D^{s(J)})}(\vartheta, \theta). \quad (2.47)$$

Using (2.10), together with the inclusion $I \subset J$, we obtain

$$c(I) \in I \subset J \subset B(c(J), 4D^{s(J)})$$

and consequently by the triangle inequality

$$B(c(I), 4D^{s(J)}) \subset B(c(J), 8D^{s(J)}).$$

Using this together with the monotonicity property (1.9) and (1.8) in (2.47), we obtain

$$\begin{aligned} d_{I^\circ}(\vartheta, \theta) &\leq 2^{-100a} d_{B(c(J), 8D^s(J))}(\vartheta, \theta) \\ &\leq 2^{-100a+5a} d_{B(c(J), \frac{1}{4}D^s(J))}(\vartheta, \theta) \\ &= 2^{-95a} d_{J^\circ}(\vartheta, \theta). \end{aligned}$$

This proves the second inequality claimed in the Lemma, from which the first follows since $a \geq 4$ and hence $2^{-95a} \leq 1$. \square

We also record the following basic estimates for the kernels K_s .

Lemma 2.9 (kernel summand). *Let $-S \leq s \leq S$ and $x, y, y' \in X$. If $K_s(x, y) \neq 0$, then we have*

$$\frac{1}{4}D^{s-1} \leq \rho(x, y) \leq \frac{1}{2}D^s. \quad (2.48)$$

We have

$$|K_s(x, y)| \leq \frac{2^{102a^3}}{\mu(B(x, D^s))} \quad (2.49)$$

and

$$|K_s(x, y) - K_s(x, y')| \leq \frac{2^{150a^3}}{\mu(B(x, D^s))} \left(\frac{\rho(y, y')}{D^s} \right)^{\frac{1}{a}}. \quad (2.50)$$

Proof. By Definition (2.5), the function K_s is the product of K with a function which is supported in the set of all x, y satisfying (2.48). This proves (2.48).

Using (1.14) and the lower bound in (2.48) we obtain

$$|K_s(x, y)| \leq \frac{2^{a^3}}{\mu(B(x, \frac{1}{4}D^{s-1}))} \quad (2.51)$$

Using $D = 2^{100a^2}$ and the doubling property (1.5) $2 + 100a^2$ times estimates the last display by

$$\leq \frac{2^{2a+101a^3}}{\mu(B(x, D^s))}. \quad (2.52)$$

Using $a \geq 4$ proves (2.49).

If $2\rho(y, y') \leq \rho(x, y)$, we obtain similarly with (1.15) and the lower bound in (2.48)

$$|K_s(x, y) - K_s(x, y')| \leq \frac{2^{a^3}}{\mu(B(x, \frac{1}{4}D^{s-1}))} \left(\frac{\rho(y, y')}{\frac{1}{4}D^{s-1}} \right)^{\frac{1}{a}}. \quad (2.53)$$

As above, this is estimated by

$$\leq \frac{4D2^{2a+101a^3}}{\mu(B(x, D^s))} \left(\frac{\rho(y, y')}{D^s} \right)^{\frac{1}{a}} = \frac{2^{2+2a+100a^2+101a^3}}{\mu(B(x, D^s))} \left(\frac{\rho(y, y')}{D^s} \right)^{\frac{1}{a}}. \quad (2.54)$$

Using $a \geq 4$, this proves (2.50) in the case $2\rho(y, y') \leq \rho(x, y)$. If $2\rho(y, y') > \rho(x, y)$, then by the lower bound in (2.48) $2\rho(y, y') > \frac{1}{8}D^s$. Then (2.50) follows from the triangle inequality, (2.49) and $a \geq 4$. \square

3. PROOF OF METRIC SPACE CARLESON (THM 1.2)

Let Borel sets F, G in X be given. We have that

$$X = \bigcup_{R>0} B(o, R), \quad (3.1)$$

because every point of X has finite distance from o .

Lemma 3.1 (R truncation). *For all integers $R > 0$*

$$\begin{aligned} & \int \mathbf{1}_{G \cap B(o, R)} \sup_{1/R < R_1 < R_2 < R} \sup_{\vartheta \in \Theta} |T_{R_1, R_2, \vartheta} \mathbf{1}_F(x)| \, d\mu(x) \\ & \leq \frac{2^{450a^3}}{(q-1)^5} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}, \end{aligned} \quad (3.2)$$

where

$$T_{R_1, R_2, \vartheta} f(x) = \int_{R_1 < \rho(x, y) < R_2} K(x, y) f(y) e(\vartheta(y)) \, d\mu(y). \quad (3.3)$$

We first show how Lemma 3.1 implies Theorem 1.2. As R tends to ∞ , the integrand of the left-hand side of (3.2) grows monotonically toward the integrand of the left-hand side of (1.19) for all x . By Lebesgue's monotone convergence theorem, the left-hand side of (3.2) converges to the left-hand side of (1.19). This verifies Theorem 1.2.

It remains to prove Lemma 3.1. Fix an integer $R > 0$. By replacing G with $G \cap B(o, R)$ if necessary, it suffices to show (3.2) under the assumption that G is contained in $B(o, R)$. We make this assumption. For every $x \in G$, the domain of integration in (3.3) is contained in $B(o, 2R)$. By replacing F with $F \cap B(o, 2R)$ if necessary, it suffices to show (3.2) under the assumption that F is contained in $B(o, 2R)$. We make this assumption.

Using the definition (2.5) of K_s and the partition of unity (2.4), we express (3.3) as the sum of

$$T_{1, s_1, s_2, \vartheta} f(x) := \sum_{s_1 \leq s \leq s_2} \int K_s(x, y) f(y) e(\vartheta(y)) \, d\mu(y) \quad (3.4)$$

and

$$\sum_{s=s_1-2, s_1-1, s_2+1, s_2+2} \int_{R_1 < \rho(x, y) < R_2} K_s(x, y) f(y) e(\vartheta(y)) \, d\mu(y), \quad (3.5)$$

where s_1 is the smallest integer such that $D^{s_1-2}R_2 > \frac{1}{4D}$ and s_2 is the largest integer such that $D^{s_2+2}R_1 < \frac{1}{2}$. We restrict the summation index s by excluding summands with $s < s_1 - 2$ or $s > s_2 + 2$ because for these summands, the function K_s vanishes on the domain of integration. We also omit the restriction in the integral for the summands in (3.4) because in these summands, the support of K_s is contained in the set described by this restriction.

We apply the triangle inequality and estimate the versions of (3.2) separately with $T_{R_1, R_2, \vartheta}$ replaced by (3.4) and by each summand of (3.5). To handle the case (3.4), we employ the following lemma. Here, we utilize the fact that if $\frac{1}{R} \leq R_1 \leq R_2 \leq R$, then s_1 and s_2 as in (3.4) are in an interval $[-S, S]$ for some sufficiently large S depending on R .

Lemma 3.2 (S truncation). *For all integers $S > 0$*

$$\begin{aligned} \int \mathbf{1}_G(x) \max_{-S < s_1 \leq s_2 < S} \sup_{\vartheta \in \Theta} |T_{1,s_1,s_2,\vartheta} \mathbf{1}_F(x)| d\mu(x) \\ \leq \frac{2^{446a^3}}{(q-1)^5} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}, \end{aligned} \quad (3.6)$$

where $T_{1,s_1,s_2,\vartheta}$ is defined in (3.4).

To reduce Lemma 3.1 to Lemma 3.2, we need estimates for the summands in (3.5). Using Lemma 2.9, we obtain for arbitrary s the inequality

$$\begin{aligned} \left| \int_{R_1 < \rho(x,y) < R_2} K_s(x,y) f(y) e(\vartheta(y)) d\mu(y) \right| \\ \leq \frac{2^{102a^3}}{\mu(B(x, D^s))} \int_{B(x, D^s)} \mathbf{1}_F(y) d\mu(y) \leq 2^{102a^3} M \mathbf{1}_F(x), \end{aligned} \quad (3.7)$$

where $M \mathbf{1}_F$ is as defined in Proposition 2.6. Now, the left-hand side of (3.2), with $T_{R_1,R_2,\vartheta}$ replaced by a summand of (3.5), can be estimated using Hölder's inequality and Proposition 2.6 by

$$2^{102a^3} \int \mathbf{1}_G(x) M \mathbf{1}_F(x) d\mu(x) \leq \frac{2^{102a^3+4a} q}{q-1} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}.$$

Applying the triangle inequality to estimate the left-hand side of (3.2) by contributions from the summands in (3.4) and (3.5), using Lemma 3.2 to control the first term, and the above to estimate the contribution from the four summands in (3.5), combined with $a \geq 4$ and $q < 2$, completes the reduction of Lemma 3.1 to Lemma 3.2.

It remains to prove Lemma 3.2. Fix $S > 0$.

Lemma 3.3 (finitary S truncation). *For all finite sets $\tilde{\Theta} \subset \Theta$*

$$\begin{aligned} \int \mathbf{1}_G(x) \max_{-S < s_1 \leq s_2 < S} \sup_{\vartheta \in \tilde{\Theta}} |T_{1,s_1,s_2,\vartheta} \mathbf{1}_F(x)| d\mu(x) \\ \leq \frac{2^{445a^3}}{(q-1)^5} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \end{aligned} \quad (3.8)$$

We reduce Lemma 3.2 to Lemma 3.3. By the Lebesgue monotone convergence theorem, applied to an increasing sequence of finite sets $\tilde{\Theta}$, inequality (3.8) continues to hold for countable $\tilde{\Theta}$.

Let $\epsilon = \frac{1}{2S+1}$. Pick some $\vartheta_0 \in \Theta$. For $k \geq 0$, let the set $\tilde{\Theta}_k$ be a subset of $B_{B(o,2R)}(\vartheta_0, k)$ of maximal size, such that for all $\vartheta, \theta \in \tilde{\Theta}_k$, it holds that $d_{B(o,2R)}(\vartheta, \theta) \geq \epsilon$. Such a set exists, since by Lemma 2.7 there exists an upper bound for the size of such subsets in $B_{B(o,2R)}(\vartheta_0, k)$. Define

$$\tilde{\Theta} := \bigcup_{k \in \mathbb{N}} \tilde{\Theta}_k.$$

Then the set $\tilde{\Theta}$ is at most countable, and it has the property that for any $\theta \in \Theta$, there exists $\vartheta \in \tilde{\Theta}$ with

$$d_{B(o,2R)}(\theta, \vartheta) < \epsilon.$$

For every $\vartheta \in \Theta$, we have

$$|T_{1,s_1,s_2,\vartheta}\mathbf{1}_F(x)| = \left| \sum_{s_1 \leq s \leq s_2} \int K_s(x,y) f(y) e(\vartheta(y) - \vartheta(x)) d\mu(y) \right| \quad (3.9)$$

Moreover, there is a $\tilde{\vartheta} \in \tilde{\Theta}$ with $d_{B(o,2R)}(\vartheta, \tilde{\vartheta}) \leq \epsilon$. Hence,

$$\begin{aligned} & |T_{1,s_1,s_2,\vartheta}\mathbf{1}_F(x)| - |T_{1,s_1,s_2,\tilde{\vartheta}}\mathbf{1}_F(x)| \\ & \leq \sum_{s_1 \leq s \leq s_2} \int |K_s(x,y)| \mathbf{1}_F(y) |e(\vartheta(y) - \vartheta(x)) - e(\tilde{\vartheta}(y) - \tilde{\vartheta}(x))| d\mu(y) \\ & \leq \sum_{s_1 \leq s \leq s_2} \epsilon \int |K_s(x,y)| \mathbf{1}_F(y) d\mu(y) \end{aligned}$$

Using Lemma 2.9, we can estimate the above expression by

$$\begin{aligned} & \sum_{s_1 \leq s \leq s_2} \frac{2^{102a^3}}{\mu(B(x, D^s))^\epsilon} \int_{B(x, D^s)} \mathbf{1}_F(y) d\mu(y) \\ & \leq (2S+1)\epsilon 2^{102a^3} M \mathbf{1}_F(x) \leq 2^{102a^3} M \mathbf{1}_F(x) \end{aligned}$$

We estimate the left-hand-side of (3.6) by the sum of left-hand-side of (3.8) and

$$\int \mathbf{1}_G(x) \max_{-S < s_1 \leq s_2 < S} \sup_{\vartheta \in \Theta} \inf_{\tilde{\vartheta} \in \tilde{\Theta}} (|T_{1,s_1,s_2,\vartheta}| - |T_{1,s_1,s_2,\tilde{\vartheta}}|) \mathbf{1}_F(x) d\mu(x),$$

which, as we have just shown, is estimated by

$$2^{102a^3} \int \mathbf{1}_G(x) M \mathbf{1}_F(x) d\mu(x).$$

By Hölder's inequality and Proposition 2.6 (more precisely, (2.46) with $p = q$), the above is no greater than

$$\frac{2^{102a^3+4a}q}{q-1} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}.$$

Combining this with Lemma 3.3 and the fact that

$$\frac{2^{102a^3+4a}q}{q-1} \leq \frac{2^{445a^3}}{(q-1)^5}$$

proves Lemma 3.2.

It remains to prove Lemma 3.3. Fix a finite set $\tilde{\Theta}$.

Lemma 3.4 (linearized truncation). *Let $\sigma_1, \sigma_2: X \rightarrow \mathbb{Z}$ be measurable functions with finite range $[-S, S]$ and $\sigma_1 \leq \sigma_2$. Let $Q: X \rightarrow \tilde{\Theta}$ be a measurable function. Then we have*

$$\int \mathbf{1}_G(x) |T_{2,\sigma_1,\sigma_2,Q}\mathbf{1}_F(x)| d\mu(x) \leq \frac{2^{445a^3}}{(q-1)^5} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}, \quad (3.10)$$

with

$$T_{2,\sigma_1,\sigma_2,Q}f(x) = \sum_{\sigma_1(x) \leq s \leq \sigma_2(x)} \int K_s(x,y)f(y)e(Q(x)(y) - Q(x)(x)) \, d\mu(y). \quad (3.11)$$

We reduce Lemma 3.3 to Lemma 3.4. For each x , let $\sigma_1(x)$ be the minimal element $s' \in [-S, S]$ such that

$$\max_{s' \leq s_2 < S} \max_{\vartheta \in \tilde{\Theta}} |T_{1,s',s_2,\vartheta} \mathbf{1}_F(x)| = \max_{-S < s_1 \leq s_2 < S} \max_{\vartheta \in \tilde{\Theta}} |T_{1,s_1,s_2,\vartheta} \mathbf{1}_F(x)| := T_{1,x}.$$

Similarly, let $\sigma_2(x)$ be the minimal element $s'' \in [-S, S]$ such that

$$\max_{\vartheta \in \tilde{\Theta}} |T_{1,\sigma_1(x),s'',\vartheta} \mathbf{1}_F(x)| = T_{1,x}.$$

Finally, choose a total order of the finite set $\tilde{\Theta}$ and let $Q(x)$ be the minimal element ϑ with respect to this order such that

$$|T_{1,\sigma_1(x),\sigma_2(x),\vartheta} \mathbf{1}_F(x)| = T_{1,x}.$$

With these choices, and noting that

$$T_{1,\sigma_1(x),\sigma_2(x),Q(x)} \mathbf{1}_F(x) = T_{2,\sigma_1,\sigma_2,Q} \mathbf{1}_F(x),$$

we conclude that the left-hand side of (3.8) and (3.10) are equal. Thus, Lemma 3.3 follows from Lemma 3.4.

It remains to prove Lemma 3.4. Fix σ_1 , σ_2 , and Q as in the lemma. Applying Proposition 2.1 recursively, we obtain a sequence of sets G_n with $G_0 = G$ and, for each $n \geq 0$, $\mu(G_n) \leq 2^{-n} \mu(G)$ and

$$\begin{aligned} \int \mathbf{1}_{G_n \setminus G_{n+1}}(x) |T_{2,\sigma_1,\sigma_2,Q} \mathbf{1}_F(x)| \, d\mu(x) \\ \leq \frac{2^{440a^3}}{(q-1)^4} \mu(G_n)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}, \end{aligned} \quad (3.12)$$

Adding the first n of these inequalities, we obtain by bounding a geometric series

$$\int \mathbf{1}_{G \setminus G_n}(x) |T_{2,\sigma_1,\sigma_2,Q} \mathbf{1}_F(x)| \, d\mu(x) \leq \frac{2^{445a^3}}{(q-1)^5} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \quad (3.13)$$

As the integrand is bounded by

$$\mathbf{1}_{G \setminus G_n}(x) \sum_{-S < s_1 \leq s_2 < S} \sum_{\vartheta \in \tilde{\Theta}} |T_{1,s_1,s_2,\vartheta} \mathbf{1}_F(x)|, \quad (3.14)$$

which by interchange of summation and integration is seen to be integrable, we obtain by Lebesgue's dominated convergence theorem

$$\int \mathbf{1}_G(x) |T_{2,\sigma_1,\sigma_2,Q} \mathbf{1}_F(x)| \, d\mu(x) \leq \frac{2^{445a^3}}{(q-1)^5} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \quad (3.15)$$

This completes the proof of Lemma 3.4 and thus Theorem 1.2.

4. PROOF OF FINITARY CARLESON (PROP 2.1)

To prove Proposition 2.1, we already fixed in Section 2 measurable functions σ_1, σ_2, Q and Borel sets F, G . We have also defined S to be the smallest integer such that the ranges of σ_1 and σ_2 are contained in $[-S, S]$ and F and G are contained in the ball $B(o, D^S)$.

The proof of the next lemma is done in Subsection 4.1, following the construction of dyadic cubes in [Chr90, §3].

Lemma 4.1 (grid existence). *There exists a grid structure (\mathcal{D}, c, s) .*

The next lemma, which we prove in Subsection 4.2, should be compared with the construction in [Zor21, Lemma 2.12].

Lemma 4.2 (tile structure). *For a given grid structure (\mathcal{D}, c, s) , there exists a tile structure $(\mathfrak{P}, \mathcal{I}, \Omega, \mathcal{Q}, c, s)$.*

Choose a grid structure (\mathcal{D}, c, s) with Lemma 4.1 and a tile structure for this grid structure $(\mathfrak{P}, \mathcal{I}, \Omega, \mathcal{Q}, c, s)$ with Lemma 4.2. Applying Proposition 2.2, we obtain a Borel set G' in X with $2\mu(G') \leq \mu(G)$ such that for all Borel functions $f : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$ we have (2.22).

Lemma 4.3 (tile sum operator). *We have for all $x \in G \setminus G'$*

$$\sum_{\mathfrak{p} \in \mathfrak{P}} T_{\mathfrak{p}} f(x) = \sum_{s=\sigma_1(x)}^{\sigma_2(x)} \int K_s(x, y) f(y) e(Q(x)(y) - Q(x)(x)) d\mu(y). \quad (4.1)$$

Proof. Fix $x \in G \setminus G'$. Sorting the tiles \mathfrak{p} on the left-hand-side of (4.1) by the value $s(\mathfrak{p}) \in [-S, S]$, it suffices to prove for every $-S \leq s \leq S$ that

$$\sum_{\mathfrak{p} \in \mathfrak{P}: s(\mathfrak{p})=s} T_{\mathfrak{p}} f(x) = 0 \quad (4.2)$$

if $s \notin [\sigma_1(x), \sigma_2(x)]$ and

$$\sum_{\mathfrak{p} \in \mathfrak{P}: s(\mathfrak{p})=s} T_{\mathfrak{p}} f(x) = \int K_s(x, y) f(y) e(Q(x)(y) - Q(x)(x)) d\mu(y). \quad (4.3)$$

if $s \in [\sigma_1(x), \sigma_2(x)]$. If $s \notin [\sigma_1(x), \sigma_2(x)]$, then by definition of $E(\mathfrak{p})$ we have $x \notin E(\mathfrak{p})$ for any \mathfrak{p} with $s(\mathfrak{p}) = s$ and thus $T_{\mathfrak{p}} f(x) = 0$. This proves (4.2).

Now assume $s \in [\sigma_1(x), \sigma_2(x)]$. By (2.9) and $G \subset B(o, D^S)$, there is at least one $I \in \mathcal{D}$ with $s(I) = s$ and $x \in I$. By (2.8), this I is unique. By (2.13), there is precisely one $\mathfrak{p} \in \mathfrak{P}(I)$ such that $Q(x) \in \Omega(\mathfrak{p})$. Hence there is precisely one $\mathfrak{p} \in \mathfrak{P}$ with $s(\mathfrak{p}) = s$ such that $x \in E(\mathfrak{p})$. For this \mathfrak{p} , the value $T_{\mathfrak{p}} f(x)$ by its definition in (2.21) equals the right-hand side of (4.3). This proves the lemma. \square

We now estimate with Lemma 4.3 and Proposition 2.2

$$\int_{G \setminus G'} \left| \sum_{s=\sigma_1(x)}^{\sigma_2(x)} \int K_s(x, y) f(y) e(Q(x)(y)) d\mu(y) \right| d\mu(x) \quad (4.4)$$

$$= \int_{G \setminus G'} \left| \sum_{s=\sigma_1(x)}^{\sigma_2(x)} \int K_s(x, y) f(y) e(Q(x)(y) - Q(x)(x)) d\mu(y) \right| d\mu(x) \quad (4.5)$$

$$= \int_{G \setminus G'} \left| \sum_{p \in \mathfrak{P}} T_p f(x) \right| d\mu(x) \leq \frac{2^{440a^3}}{(q-1)^4} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \quad (4.6)$$

This proves (2.6) for the chosen set G' and arbitrary f and thus completes the proof of Proposition 2.1.

4.1. Proof of Lemma 4.1, dyadic structure. We begin with the construction of the centers of the dyadic cubes.

Lemma 4.4 (counting balls). *Let $-S \leq k \leq S$. Consider $Y \subset X$ such that for any $y \in Y$, we have*

$$y \in B(o, 4D^S - D^k), \quad (4.7)$$

furthermore, for any $y' \in Y$ with $y \neq y'$, we have

$$B(y, D^k) \cap B(y', D^k) = \emptyset. \quad (4.8)$$

Then the cardinality of Y is bounded by

$$|Y| \leq 2^{3a+200Sa^3}. \quad (4.9)$$

Proof. Let k and Y be given. By applying the doubling property (1.5) inductively, we have for each integer $j \geq 0$

$$\mu(B(y, 2^j D^k)) \leq 2^{aj} \mu(B(y, D^k)). \quad (4.10)$$

Since X is the union of the balls $B(y, 2^j D^k)$ and μ is not zero, at least one of the balls $B(y, 2^j D^k)$ has positive measure, thus $B(y, D^k)$ has positive measure.

Applying (4.10) for $j' = \ln_2(8D^{2S}) = 3 + 2S \cdot 100a^2$ by (2.1), using $-S \leq k \leq S$, $y \in B(o, 4D^S)$, and the triangle inequality, we have

$$B(o, 4D^S) \subset B(y, 8D^S) \subset B(y, 2^{j'} D^k). \quad (4.11)$$

Using the disjointedness of the balls in (4.8), (4.7), and the triangle inequality for ρ , we obtain

$$|Y| \mu(B(o, 4D^S)) \leq 2^{j'a} \sum_{y \in Y} \mu(B(y, D^k)) \quad (4.12)$$

$$\leq 2^{j'a} \mu\left(\bigcup_{y \in Y} B(y, D^k)\right) \leq 2^{j'a} \mu(o, 4D^S). \quad (4.13)$$

As $\mu(o, 4D^S)$ is not zero, the lemma follows. \square

For each $-S \leq k \leq S$, let Y_k be a set of maximal cardinality in X such that $Y = Y_k$ satisfies the properties (4.7) and (4.8). By the upper bound of Lemma 4.4, such a set exists.

For each $-S \leq k \leq S$, choose an enumeration of the points in the finite set Y_k and thus a total order $<$ on Y_k .

Lemma 4.5 (cover big ball). *For each $-S \leq k \leq S$, the ball $B(o, 4D^S - D^k)$ is contained in the union of the balls $B(y, 2D^k)$ with $y \in Y_k$.*

Proof. Let x be any point of $B(o, 4D^S - D^k)$. By maximality of $|Y_k|$, the ball $B(x, D^k)$ intersects one of the balls $B(y, D^k)$ with $y \in Y_k$. By the triangle inequality, $x \in B(y, 2D^k)$. \square

Define the set

$$\mathcal{C} := \{(y, k) : -S \leq k \leq S, y \in Y_k\} \quad (4.14)$$

We totally order the set \mathcal{C} lexicographically by setting $(y, k) < (y', k')$ if $k < k'$ or both $k = k'$ and $y < y'$. In what follows, we define recursively in the sense of this order a function

$$(I_1, I_2, I_3) : \mathcal{C} \rightarrow \mathcal{P}(X) \times \mathcal{P}(X) \times \mathcal{P}(X). \quad (4.15)$$

Assume the sets $I_j(y', k')$ have already been defined for $j = 1, 2, 3$ if $k' < k$ and if $k = k'$ and $y' < y$.

If $k = -S$, define for $j \in \{1, 2\}$ the set $I_j(y, k)$ to be $B(y, jD^{-S})$. If $-S < k$, define for $j \in \{1, 2\}$ and $y \in Y_k$ the set $I_j(y, k)$ to be

$$\bigcup \{I_3(y', k-1) : y' \in Y_{k-1} \cap B(y, jD^k)\}. \quad (4.16)$$

Define for $-S \leq k \leq S$ and $y \in Y_k$

$$I_3(y, k) := I_1(y, k) \cup \left[I_2(y, k) \setminus \left[X_k \cup \bigcup \{I_3(y', k) : y' \in Y_k, y' < y\} \right] \right] \quad (4.17)$$

with

$$X_k := \bigcup \{I_1(y', k) : y' \in Y_k\}. \quad (4.18)$$

Lemma 4.6 (basic grid structure). *For each $-S \leq k \leq S$ and $1 \leq j \leq 3$ the following holds.*

If $j \neq 2$ and for some $x \in X$ and $y_1, y_2 \in Y_k$ we have

$$x \in I_j(y_1, k) \cap I_j(y_2, k), \quad (4.19)$$

then $y_1 = y_2$.

If $j \neq 1$, then

$$B(o, 4D^S - 2D^k) \subset \bigcup_{y \in Y_k} I_j(y, k). \quad (4.20)$$

We have for each $y \in Y_k$,

$$B(y, \frac{1}{2}D^k) \subset I_3(y, k) \subset B(y, 4D^k). \quad (4.21)$$

Proof. We prove these statements simultaneously by induction on the ordered set of pairs (y, k) . Let $-S \leq k \leq S$.

We first consider (4.19) for $j = 1$. If $k = -S$, disjointedness of the sets $I_1(y, -S)$ follows by definition of I_1 and Y_k . If $k > -S$, assume x is in $I_1(y_m, k)$ for $m = 1, 2$. Then, for $m = 1, 2$, there is $z_m \in Y_{k-1} \cap B(y_m, D^k)$ with $x \in I_3(z_m, k-1)$. Using (4.19) inductively for $j = 3$, we conclude $z_1 = z_2$. This implies that the balls $B(y_1, D^k)$ and $B(y_2, D^k)$ intersect. By construction of Y_k , this implies $y_1 = y_2$. This proves (4.19) for $j = 1$.

We next consider (4.19) for $j = 3$. Assume x is in $I_3(y_m, k)$ for $m = 1, 2$ and $y_m \in Y_k$. If x is in X_k , then by definition (4.17), $x \in I_1(y_m, k)$ for $m = 1, 2$. As we have already shown (4.19) for $j = 1$, we conclude $y_1 = y_2$. This completes the proof in case $x \in X_k$, and we may assume x is not in X_k . By definition (4.17), x is not in $I_3(z, k)$ for any z with $z < y_1$ or $z < y_2$. Hence, neither $y_1 < y_2$ nor $y_2 < y_1$, and by totality of the order of Y_k , we have $y_1 = y_2$. This completes the proof of (4.19) for $j = 3$.

We show (4.20) for $j = 2$. In case $k = -S$, this follows from Lemma 4.5. Assume $k > -S$. Let x be a point of $B(o, 4D^S - 2D^k)$. By induction, there is $y' \in Y_{k-1}$ such that $x \in I_3(y', k-1)$. Using the inductive statement (4.21), we obtain $x \in B(y', 4D^{k-1})$. As $D > 4$, by applying the triangle inequality with the points, o , x , and y' , we obtain that $y' \in B(o, 4D^S - D^k)$. By Lemma 4.5, y' is in $B(y, 2D^k)$ for some $y \in Y_k$. It follows that $x \in I_2(y, k)$. This proves (4.20) for $j = 2$.

We show (4.20) for $j = 3$. Let $x \in B(o, 4D^S - 2D^k)$. In case $x \in X_k$, then by definition of X_k we have $x \in I_1(y, k)$ for some $y \in Y_k$ and thus $x \in I_3(y, k)$. We may thus assume $x \notin X_k$. As we have already seen (4.20) for $j = 2$, there is $y \in Y_k$ such that $x \in I_2(y, k)$. We may assume this y is minimal with respect to the order in Y_k . Then $x \in I_3(y, k)$. This proves (4.20) for $j = 3$.

Next, we show the first inclusion in (4.21). Let $x \in B(y, \frac{1}{2}D^k)$. As $I_1(y, k) \subset I_3(y, k)$, it suffices to show $x \in I_1(y, k)$. If $k = -S$, this follows immediately from the assumption on x and the definition of I_1 . Assume $k > -S$. By the inductive statement (4.20) and $D > 4$, there is a $y' \in Y_{k-1}$ such that $x \in I_3(y', k-1)$. By the inductive statement (4.21), we conclude $x \in B(y', 4D^{k-1})$. By the triangle inequality with points x , y , y' , and $D > 4$, we have $y' \in B(y, D^k)$. It follows by definition (4.16) that $I_3(y', k-1) \subset I_1(y, k)$, and thus $x \in I_3(y, k)$. This proves the first inclusion in (4.21).

We show the second inclusion in (4.21). Let $x \in I_3(y, k)$. As $I_1(y, k) \subset I_2(y, k)$ directly from the definition (4.16), it follows by definition (4.17) that $x \in I_2(y, k)$. By definition (4.16), there is $y' \in Y_{k-1} \cap B(y, 2D^k)$ with $x \in I_3(y', k-1)$. By induction, $x \in B(y', 4D^{k-1})$. By the triangle inequality applied to the points x, y', y and $D > 4$, we conclude $x \in B(y, 4D^k)$. This shows the second inclusion in (4.21) and completes the proof of the lemma. \square

Lemma 4.7 (cover by cubes). *Let $-S \leq l \leq k \leq S$ and $y \in Y_k$. We have*

$$I_3(y, k) \subset \bigcup_{y' \in Y_l} I_3(y', l). \quad (4.22)$$

Proof. Let $-S \leq l \leq k \leq S$ and $y \in Y_k$. If $l = k$, the inclusion (4.22) is true from the definition of set union. We may then assume inductively that $k > l$ and the statement of the lemma is true if k is replaced by $k-1$. Let $x \in I_3(y, k)$. By definition (4.17), $x \in I_j(y, k)$ for some $j \in \{1, 2\}$. By (4.16), $x \in I_3(w, k-1)$ for some $w \in Y_{k-1}$. We conclude (4.22) by induction. \square

Lemma 4.8 (dyadic property). *Let $-S \leq l \leq k \leq S$ and $y \in Y_k$ and $y' \in Y_l$ with $I_3(y', l) \cap I_3(y, k) \neq \emptyset$. Then*

$$I_3(y', l) \subset I_3(y, k). \quad (4.23)$$

Proof. Let l, k, y, y' be as in the lemma. Pick $x \in I_3(y', l) \cap I_3(y, k)$. Assume first $l = k$. By (4.19) of Lemma 4.6, we conclude $y' = y$, and thus (4.23). Now assume $l < k$. By induction, we may assume that the statement of the lemma is proven for $k-1$ in place of k .

By Lemma 4.7, there is a $y'' \in Y_{k-1}$ such that $x \in I_3(y'', k-1)$. By induction, we have $I_3(y', l) \subset I_3(y'', k-1)$. If $l = k-1$, then by the

disjointedness property of Lemma 4.6, we see that $y' = y''$, and hence (4.23) follows. For $l < k - 1$, it remains to prove

$$I_3(y'', k - 1) \subset I_3(y, k). \quad (4.24)$$

We make a case distinction and assume first $x \in X_k$. By Definition (4.17), we have $x \in I_1(y, k)$. By Definition (4.16), there is a $v \in Y_{k-1} \cap B(y, D^k)$ with $x \in I_3(v, k - 1)$. By (4.19) of Lemma 4.6, we have $v = y''$. By Definition (4.16), we then have $I_3(y'', k - 1) \subset I_1(y, k)$. Then (4.24) follows by Definition (4.17) in the given case.

Assume now the case $x \notin X_k$. By (4.17), we have $x \in I_2(y, k)$. Moreover, for any $u < y$ in Y_k , we have $x \notin I_3(u, k)$. Let $u < y$. By transitivity of the order in Y_k , we conclude $x \notin I_2(u, k)$. By (4.16) and the disjointedness property of Lemma 4.6, we have $I_3(y'', k - 1) \cap I_2(u, k) = \emptyset$. Similarly, $I_3(y'', k - 1) \cap I_1(u, k) = \emptyset$. Hence $I_3(y'', k - 1) \cap I_3(u, k) = \emptyset$. As $u < y$ was arbitrary, we conclude with (4.17) the claim in the given case. This completes the proof of (4.24), and thus also (4.23). \square

For $-S \leq k' \leq k \leq S$ and $y' \in Y_{k'}$, $y \in Y_k$ write $(y', k'|y, k)$ if $I_3(y', k') \subset I_3(y, k)$ and

$$\inf_{x \in X \setminus I_3(y, k)} \rho(y', x) < 6D^{k'}. \quad (4.25)$$

Lemma 4.9 (transitive boundary). *Assume $-S \leq k'' < k' < k \leq S$ and $y'' \in Y_{k''}$, $y' \in Y_{k'}$, $y \in Y_k$. Assume there is $x \in X$ such that*

$$x \in I_3(y'', k'') \cap I_3(y', k') \cap I_3(y, k). \quad (4.26)$$

If $(y'', k''|y, k)$, then also $(y'', k''|y', k')$ and $(y', k'|y, k)$

Proof. As $x \in I_3(y'', k'') \cap I_3(y', k')$ and $k'' < k'$, we have by Lemma 4.8 that $I_3(y'', k'') \subset I_3(y', k')$. Similarly, $I_3(y', k') \subset I_3(y, k)$. Pick $x' \in X \setminus I_3(y, k)$ such that

$$\rho(y'', x') < 6D^{k''}, \quad (4.27)$$

which exists as $(y'', k''|y, k)$. As $x' \in X \setminus I_3(y', k')$ as well, we conclude $(y'', k''|y', k')$. By the triangle inequality, we have

$$\rho(y', x') \leq \rho(y', x) + \rho(x, y'') + \rho(y'', x') \quad (4.28)$$

Using the choice of x and (4.21) as well as (4.27), we estimate this by

$$< 4D^{k'} + 4D^{k''} + 6D^{k''} \leq 6D^{k'}, \quad (4.29)$$

where we have used $D > 5$ and $k'' < k'$. We conclude $(y', k'|y, k)$. \square

Lemma 4.10 (small boundary). *Let $K = 2^{4a+1}$. For each $-S + K \leq k \leq S$ and $y \in Y_k$ we have*

$$\sum_{z \in Y_{k-K} : (z, k-K|y, k)} \mu(I_3(z, k-K)) \leq \frac{1}{2} \mu(I_3(y, k)). \quad (4.30)$$

Proof. Let K be as in the lemma. Let $-S + K \leq k \leq S$ and $y \in Y_k$.

Pick k' so that $k - K \leq k' \leq k$. For each $y'' \in Y_{k-K}$ with $(y'', k-K|y, k)$, by Lemma 4.7 and Lemma 4.8, there is a unique $y' \in Y_{k'}$ such that

$$I_3(y'', k-K) \subset I_3(y', k') \subset I_3(y, k). \quad (4.31)$$

Using Lemma 4.9, $(y', k'|y, k)$.

We conclude using the disjointedness property of Lemma 4.6 that

$$\sum_{y'':(y'', k-K|y, k)} \mu(I_3(y'', k-K)) \leq \sum_{y':(y', k'|y, k)} \mu(I_3(y', k')). \quad (4.32)$$

Adding over $k-K < k' \leq k$, and using

$$\mu(I_3(y', k')) \leq 2^{4a} \mu(B(y', \frac{1}{4}D^{k'}))$$

from the doubling property (1.5) and (4.21) gives

$$K \sum_{y'':(y'', k-K|y, k)} \mu(I_3(y'', k-K)) \quad (4.33)$$

$$\leq 2^{4a} \sum_{k-K < k' \leq k} \left[\sum_{y':(y', k'|y, k)} \mu(B(y', \frac{1}{4}D^{k'})) \right] \quad (4.34)$$

Each ball $B(y', \frac{1}{4}D^{k'})$ occurring in (4.34) is contained in $I_3(y', k')$ by (4.21) and in turn contained in $I_3(y, k)$ by (4.31). Assume for the moment all these balls are pairwise disjoint. Then by additivity of the measure,

$$K \sum_{y'':(y'', k-K|y, k)} \mu(I_3(y'', k-K)) \leq 2^{4a} \mu(I_3(y, k)) \quad (4.35)$$

which by $K = 2^{4a+1}$ implies (4.30).

It thus remains to prove that the balls occurring in (4.34) are pairwise disjoint. Let (u, l) and (u', l') be two parameter pairs occurring in the sum of (4.34) and let $B(u, \frac{1}{4}D^l)$ and $B(u', \frac{1}{4}D^{l'})$ be the corresponding balls. If $l = l'$, then the balls are equal or disjoint by (4.21) and (4.19) of Lemma 4.6. Assume then without loss of generality that $l' < l$. Towards a contradiction, assume that

$$B(u, \frac{1}{4}D^l) \cap B(u', \frac{1}{4}D^{l'}) \neq \emptyset \quad (4.36)$$

As $(u', l'|y, k)$, there is a point x in $X \setminus I_3(y, k)$ with $\rho(x, u') < 6D^{l'}$. Using $D > 25$, we conclude from the triangle inequality and (4.36) that $x \in B(u, \frac{1}{2}D^l)$. However, $B(u, \frac{1}{2}D^l) \subset I_3(u, l)$, and $I_3(u, l) \subset I_3(y, k)$, a contradiction to $x \notin I_3(y, k)$. This proves the lemma. \square

Lemma 4.11 (smaller boundary). *Let $K = 2^{4a+1}$ and let $n \geq 0$ be an integer. Then for each $-S + nK \leq k \leq S$ we have*

$$\sum_{y' \in Y_{k-nK}:(y', k-nK|y, k)} \mu(I_3(y', k-nK)) \leq 2^{-n} \mu(I_3(y, k)). \quad (4.37)$$

Proof. We prove this by induction on n . If $n = 0$, both sides of (4.37) are equal to $\mu(I_3(y, k))$ by (4.19). If $n = 1$, this follows from Lemma 4.10.

Assume $n > 1$ and (4.37) has been proven for $n-1$. We write (4.37)

$$\sum_{y'' \in Y_{k-nK}:(y'', k-nK|y, k)} \mu(I_3(y'', k-nK)) \quad (4.38)$$

$$= \sum_{y' \in Y_{k-K}:(y', k-K|y, k)} \left[\sum_{y'' \in Y_{k-nK}:(y'', k-nK|y', k-K)} \mu(I_3(y'', k-nK)) \right] \quad (4.39)$$

Applying the induction hypothesis, this is bounded by

$$= \sum_{y' \in Y_{k-K} : (y', k-K | y, k)} 2^{1-n} \mu(I_3(y', k-K)) \quad (4.40)$$

Applying (4.30) gives (4.37), and proves the lemma. \square

Lemma 4.12 (boundary measure). *For each $-S \leq k \leq S$ and $y \in Y_k$ and $0 < t < 1$ with $tD^k \geq D^{-S}$ we have*

$$\mu(\{x \in I_3(y, k) : \rho(x, X \setminus I_3(y, k)) \leq tD^k\}) \leq 2^{100a^2} t^\kappa \mu(I_3(y, k)). \quad (4.41)$$

Proof. Let $x \in I_3(y, k)$ with $\rho(x, X \setminus I_3(y, k)) \leq tD^k$. Let $K = 2^{4a+1}$ as in Lemma 4.11. Let n be the largest integer such that $D^{nK} \leq \frac{1}{t}$, so that $tD^k \leq D^{k-nK}$ and

$$D^{nK} > \frac{1}{tD^K}. \quad (4.42)$$

Let $k' = k - nK$, by the assumption $tD^k \geq D^{-S}$, we have $k' \geq -S$. By (4.22), there exists $y' \in Y_{k'}$ with $x \in I_3(y', k')$. By the squeezing property (4.21) and the assumption on x , we have

$$\rho(y', X \setminus I_3(y, k)) \leq \rho(x, y') + \rho(x, X \setminus I_3(y, k)) \leq 4D^{k'} + tD^k.$$

By the assumption on n and the definition of k' , this is

$$\leq 4D^{k'} + D^{k-nK} < 6D^{k'}.$$

Together with (4.23) thus $(y', k' | y, k)$. We have shown that

$$\begin{aligned} & \{x \in I_3(y, k) : \rho(x, X \setminus I_3(y, k)) \leq tD^k\} \\ & \subset \bigcup_{y' \in Y_{k-nK} : (y', k-nK | y, k)} I_3(y', k-nK). \end{aligned}$$

Using monotonicity and additivity of the measure and Lemma 4.11, we obtain

$$\mu(\{x \in I_3(y, k) : \rho(x, X \setminus I_3(y, k)) \leq tD^k\}) \leq 2^{-n} \mu(I_3(y, k)).$$

By (4.42) and the definition (2.1) of D , this is bounded by

$$t^{1/(100a^2K)} 2^{100a^2} \mu(I_3(y, k)),$$

which completes the proof by the definition (2.2) of κ . \square

Let \mathcal{D} be the set of all $I_3(y, k)$ with $k \in [-S, S]$ and $y \in Y_k$. Define

$$s(I_3(y, k)) := k \quad (4.43)$$

$$c(I_3(y, k)) := y. \quad (4.44)$$

We show that (\mathcal{D}, c, s) constitutes a grid structure. Property (2.10) follows from (4.21), while (2.11) follows from Lemma 4.12.

Let $x \in B(o, D^S)$. We show properties (2.7), (2.8) and (2.9) for (\mathcal{D}, c, s) and x .

We first show (2.7) by contradiction. Then there is an I violating the conclusion of (2.7). Pick such $I = I_3(y, l)$ such that l is minimal. By assumption, we have $-S \leq k < l$; in particular $-S < l$. By definition, $I_3(y, l)$ is contained in $I_1(y, l) \cup I_2(y, l)$, which is contained in the union of $I_3(y', l-1)$ with $y' \in Y_{l-1}$. By minimality of l , each such $I_3(y', l-1)$ is contained in the union of all $I_3(z, k)$ with $z \in Y_k$. This proves (2.7).

We now show (2.8). Assume to get a contradiction that there are non-disjoint $I, J \in \mathcal{D}$ with $s(I) \leq s(J)$ and $I \not\subset J$. We may assume the existence of such I and J with minimal $s(J) - s(I)$. Let $k = s(I)$. Assume first $s(J) = k$. Let $I = I_3(y_1, k)$ and $J = I_3(y_2, k)$ with $y_1, y_2 \in Y_k$. If $y_1 = y_2$, then $I = J$, a contradiction to $I \not\subset J$. If $y_1 \neq y_2$, then $I \cap J = \emptyset$ by (4.19), a contradiction to the non-disjointedness of I, J . Assume now $s(J) > k$. Choose $y \in I \cap J$. By property (2.7), there is $K \in \mathcal{D}$ with $s(K) = s(J) - 1$ and $y \in K$. By construction of J , and pairwise disjointedness of all $I_3(w, s(J) - 1)$ that we have already seen, we have $K \subset J$. By minimality of $s(J)$, we have $I \subset K$. This proves $I \subset J$ and thus (2.8).

We next establish (2.9). Let $-S \leq k \leq S$. Using (4.20) for $j = 3$, we get

$$x \in B(o, D^S) \subset B(o, 4D^S - 2D^k) \subset \bigcup_{y \in Y_k} I_3(y, k). \quad (4.45)$$

Thus, there exists a dyadic cube $I = I_3(y, k)$ with $s(I) = k$ and $x \in I$. This proves (2.9).

4.2. Proof of Lemma 4.2, tile structure. Choose a grid structure (\mathcal{D}, c, s) with Lemma 4.1. Let $I \in \mathcal{D}$. Suppose that

$$\mathcal{Z} \subset \bigcup_{\vartheta \in Q(X)} B_{I^\circ}(\vartheta, 1) \quad (4.46)$$

is such that for any $\vartheta, \theta \in \mathcal{Z}$ we have

$$B_{I^\circ}(\vartheta, 0.3) \cap B_{I^\circ}(\theta, 0.3) = \emptyset. \quad (4.47)$$

By Lemma 2.7 applied to each of the balls $B_{I^\circ}(\vartheta, 1)$, $\vartheta \in Q(X)$, we have

$$|\mathcal{Z}| \leq 2^{2a} |Q(X)|.$$

In particular, there exists a set \mathcal{Z} satisfying both (4.46) and (4.47) of maximal cardinality among all such sets. We pick for each $I \in \mathcal{D}$ such a set $\mathcal{Z}(I)$.

Lemma 4.13 (frequency ball cover). *For each $I \in \mathcal{D}$, we have*

$$Q(X) \subset \bigcup_{\vartheta \in Q(X)} B_{I^\circ}(\vartheta, 1) \subset \bigcup_{z \in \mathcal{Z}(I)} B_{I^\circ}(z, 0.7). \quad (4.48)$$

Proof. To show (4.48) note that the first inclusion is obvious. For the second inclusion let $\theta \in \bigcup_{\vartheta \in Q(X)} B_{I^\circ}(\vartheta, 1)$. By maximality of $\mathcal{Z}(I)$, there must be a point $z \in \mathcal{Z}(I)$ such that $B_{I^\circ}(z, 0.3) \cap B_{I^\circ}(\theta, 0.3) \neq \emptyset$. Else, $\mathcal{Z}(I) \cup \{\theta\}$ would be a set of larger cardinality than $\mathcal{Z}(I)$ satisfying (4.46) and (4.47). Fix such z , and fix a point $z_1 \in B_{I^\circ}(z, 0.3) \cap B_{I^\circ}(\theta, 0.3)$. By the triangle inequality, we deduce that

$$d_{I^\circ}(z, \theta) \leq d_{I^\circ}(z, z_1) + d_{I^\circ}(\theta, z_1) < 0.3 + 0.3 = 0.6,$$

and hence $\theta \in B_{I^\circ}(z, 0.7)$. \square

We define

$$\begin{aligned} \mathfrak{P} &= \{(I, z) : I \in \mathcal{D}, z \in \mathcal{Z}(I)\}, \\ \mathcal{I}((I, z)) &= I \quad \text{and} \quad \mathcal{Q}((I, z)) = z. \end{aligned}$$

We further set

$$s(\mathfrak{p}) = s(\mathcal{I}(\mathfrak{p})), \quad c(\mathfrak{p}) = c(\mathcal{I}(\mathfrak{p})).$$

Then (2.18), (2.19) hold by definition.

It remains to construct the map Ω , and verify properties (2.13), (2.14) and (2.15). We first construct an auxiliary map Ω_1 . For each $I \in \mathcal{D}$, we pick an enumeration of the finite set $\mathcal{Z}(I)$

$$\mathcal{Z}(I) = \{z_1, \dots, z_M\}.$$

We define $\Omega_1 : \mathfrak{P} \mapsto \mathcal{P}(\Theta)$ as below. Set

$$\Omega_1((I, z_1)) = B_{I^\circ}(z_1, 0.7) \setminus \bigcup_{z \in \mathcal{Z}(I) \setminus \{z_1\}} B_{I^\circ}(z, 0.3)$$

and then define iteratively

$$\Omega_1((I, z_k)) = B_{I^\circ}(z_k, 0.7) \setminus \bigcup_{z \in \mathcal{Z}(I) \setminus \{z_k\}} B_{I^\circ}(z, 0.3) \setminus \bigcup_{i=1}^{k-1} \Omega_1((I, z_i)). \quad (4.49)$$

Lemma 4.14 (disjoint frequency cubes). *For each $I \in \mathcal{D}$, and $\mathfrak{p}_1, \mathfrak{p}_2 \in \mathfrak{P}(I)$, if*

$$\Omega_1(\mathfrak{p}_1) \cap \Omega_1(\mathfrak{p}_2) \neq \emptyset,$$

then $\mathfrak{p}_1 = \mathfrak{p}_2$.

Proof. By the definition of the map \mathcal{I} , we have

$$\mathfrak{P}(I) = \{(I, z) : z \in \mathcal{Z}(I)\}.$$

By (4.49), the set $\Omega_1((I, z_k))$ is disjoint from each $\Omega_1((I, z_i))$ with $i < k$. Thus the sets $\Omega_1(\mathfrak{p})$, $\mathfrak{p} \in \mathfrak{P}(I)$ are pairwise disjoint. \square

Lemma 4.15 (frequency cube cover). *For each $I \in \mathcal{D}$, it holds that*

$$\bigcup_{z \in \mathcal{Z}(I)} B_{I^\circ}(z, 0.7) \subset \bigcup_{\mathfrak{p} \in \mathfrak{P}(I)} \Omega_1(\mathfrak{p}). \quad (4.50)$$

For every $\mathfrak{p} \in \mathfrak{P}$, it holds that

$$B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), 0.3) \subset \Omega_1(\mathfrak{p}) \subset B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), 0.7). \quad (4.51)$$

Proof. For (4.51) let $\mathfrak{p} = (I, z)$. The second inclusion in (4.51) then follows from (4.49) and the equality $B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), 0.7) = B_{I^\circ}(z, 0.7)$, which is true by definition. For the first inclusion in (4.51) let $\vartheta \in B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), 0.3)$. Let k be such that $z = z_k$ in the enumeration we chose above. It follows immediately from (4.49) and (4.47) that $\vartheta \notin \Omega_1((I, z_i))$ for all $i < k$. Thus, again from (4.49), we have $\vartheta \in \Omega_1((I, z_k))$.

To show (4.50) let $\vartheta \in \bigcup_{z \in \mathcal{Z}(I)} B_{I^\circ}(z, 0.7)$. If there exists $z \in \mathcal{Z}(I)$ with $\vartheta \in B_{I^\circ}(z, 0.3)$, then

$$z \in \Omega_1((I, z)) \subset \bigcup_{\mathfrak{p} \in \mathfrak{P}(I)} \Omega_1(\mathfrak{p})$$

by the first inclusion in (4.51).

Now suppose that there exists no $z \in \mathcal{Z}(I)$ with $\vartheta \in B_{I^\circ}(z, 0.3)$. Let k be minimal such that $\vartheta \in B_{I^\circ}(z_k, 0.7)$. Since $\Omega_1((I, z_i)) \subset B_{I^\circ}(z_i, 0.7)$ for each i by (4.49), we have that $\vartheta \notin \Omega_1((I, z_i))$ for all $i < k$. Hence $\vartheta \in \Omega_1((I, z_k))$, again by (4.49). \square

Now we are ready to define the function Ω . For all cubes $I \in \mathcal{D}$ such that there exists no $J \in \mathcal{D}$ with $I \subset J$ and $I \neq J$, we define for all $\mathbf{p} \in \mathfrak{P}(I)$

$$\Omega(\mathbf{p}) = \Omega_1(\mathbf{p}). \quad (4.52)$$

For cubes $I \in \mathcal{D}$ for which there exists $J \in \mathcal{D}$ with $I \subset J$ and $I \neq J$, we define Ω by recursion. We can pick an inclusion minimal $J \in \mathcal{D}$ among the finitely many cubes such that $I \subset J$ and $I \neq J$. This J is unique: Suppose that J' is another inclusion minimal cube with $I \subset J'$ and $I \neq J'$. Without loss of generality, we have that $s(J) \leq s(J')$. By (2.8), it follows that $J \subset J'$. Since J' is minimal with respect to inclusion, it follows that $J = J'$. Then we define

$$\Omega(\mathbf{p}) = \bigcup_{z \in \mathcal{Z}(J) \cap \Omega_1(\mathbf{p})} \Omega((J, z)) \cup B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 0.2). \quad (4.53)$$

Lemma 4.16 (dyadic frequency cubes). *With this definition, (2.13), (2.14) and (2.15) hold.*

Proof. First, we prove (2.15). If $I \in \mathcal{D}$ is maximal in \mathcal{D} with respect to set inclusion, then (2.15) holds for all $\mathbf{p} \in \mathfrak{P}(I)$ by (4.52) and (4.51). Now suppose that I is not maximal in \mathcal{D} with respect to set inclusion. Then we may assume by induction that for all $J \in \mathcal{D}$ with $I \subset J$ and all $\mathbf{p}' \in \mathfrak{P}(J)$, (2.15) holds. Let J be the unique minimal cube in \mathcal{D} with $I \subsetneq J$.

Suppose that $\vartheta \in \Omega(\mathbf{p})$. If $\vartheta \in B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 0.2)$, then since

$$B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 0.2) \subset B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 1),$$

we conclude that $\vartheta \in B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 0.7)$. If not, by (4.53), there exists $z \in \mathcal{Z}(J) \cap \Omega_1(\mathbf{p})$ with $\vartheta \in \Omega(J, z)$. Using the triangle inequality and (4.51), we obtain

$$d_{I^\circ}(\mathcal{Q}(\mathbf{p}), \vartheta) \leq d_{I^\circ}(\mathcal{Q}(\mathbf{p}), z) + d_{I^\circ}(z, \vartheta) \leq 0.7 + d_{I^\circ}(z, \vartheta).$$

By Lemma 2.8 and the induction hypothesis, this is estimated by

$$\leq 0.7 + 2^{-95a} d_{J^\circ}(z, \vartheta) \leq 0.7 + 2^{-95a} \cdot 1 < 1.$$

This shows the second inclusion in (2.15). The first inclusion is immediate from (4.53).

Next, we show (2.13). Let $I \in \mathcal{D}$.

If I is maximal with respect to inclusion, then disjointedness of the sets $\Omega(\mathbf{p})$ for $\mathbf{p} \in \mathfrak{P}(I)$ follows from the definition (4.52) and Lemma 4.14. To obtain the inclusion in (2.13) one combines the inclusions (4.48) and (4.50) of Lemma 4.15 with (4.52).

Now we turn to the case where there exists $J \in \mathcal{D}$ with $I \subset J$ and $I \neq J$. In this case we use induction: It suffices to show (2.13) under the assumption that it holds for all cubes $J \in \mathcal{D}$ with $I \subset J$. As shown before definition (4.53), we may choose the unique inclusion minimal such J . To show disjointedness of the sets $\Omega(\mathbf{p})$, $\mathbf{p} \in \mathfrak{P}(I)$ we pick two tiles $\mathbf{p}, \mathbf{p}' \in \mathfrak{P}(I)$ and $\vartheta \in \Omega(\mathbf{p}) \cap \Omega(\mathbf{p}')$. Then we are by (4.53) in one of the following four cases.

1. There exist $z \in \mathcal{Z}(J) \cap \Omega_1(\mathbf{p})$ such that $\vartheta \in \Omega(J, z)$, and there exists $z' \in \mathcal{Z}(J) \cap \Omega_1(\mathbf{p}')$ such that $\vartheta \in \Omega(J, z')$. By the induction hypothesis, that (2.13) holds for J , we must have $z = z'$. By Lemma 4.14, we must then have $\mathbf{p} = \mathbf{p}'$.

2. There exists $z \in \mathcal{Z}(J) \cap \Omega_1(\mathbf{p})$ such that $\vartheta \in \Omega(J, z)$, and $\vartheta \in B_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), 0.2)$. Using the triangle inequality, Lemma 2.8 and (2.15), we obtain

$$d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), z) \leq d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), \vartheta) + d_{\mathbf{p}'}(z, \vartheta) \leq 0.2 + 2^{-95a} \cdot 1 < 0.3.$$

Thus $z \in \Omega_1(\mathbf{p}')$ by (4.51). By Lemma 4.14, it follows that $\mathbf{p} = \mathbf{p}'$.

3. There exists $z' \in \mathcal{Z}(J) \cap \Omega_1(\mathbf{p}')$ such that $\vartheta \in \Omega(J, z')$, and $\vartheta \in B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 0.2)$. This case is the same as case 2., after swapping \mathbf{p} and \mathbf{p}' .

4. We have $\vartheta \in B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 0.2) \cap B_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), 0.2)$. In this case it follows that $\mathbf{p} = \mathbf{p}'$ since the sets $B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 0.2)$ are pairwise disjoint by the inclusion (4.51) and Lemma 4.14.

To show the inclusion in (2.13), let $\vartheta \in Q(X)$. By the induction hypothesis, there exists $\mathbf{p} \in \mathfrak{P}(J)$ such that $\vartheta \in \Omega(\mathbf{p})$. By definition of the set \mathfrak{P} , we have $\mathbf{p} = (J, z)$ for some $z \in \mathcal{Z}(J)$. By (4.46), there exists $x \in X$ with $d_{J^\circ}(Q(x), z) \leq 1$. By Lemma 2.8, it follows that $d_{I^\circ}(Q(x), z) \leq 1$. Thus, by (4.48), there exists $z' \in \mathcal{Z}(I)$ with $z \in B_{I^\circ}(z', 0.7)$. Then by Lemma (4.15) there exists $\mathbf{p}' \in \mathfrak{P}(I)$ with $z \in \mathcal{Z}(J) \cap \Omega_1(\mathbf{p}')$. Consequently, by (4.53), $\vartheta \in \Omega(\mathbf{p}')$. This completes the proof of (2.13).

Finally, we show (2.14). Let $\mathbf{p}, \mathbf{q} \in \mathfrak{P}$ with $\mathcal{I}(\mathbf{p}) \subset \mathcal{I}(\mathbf{q})$ and $\Omega(\mathbf{p}) \cap \Omega(\mathbf{q}) \neq \emptyset$. If we have $s(\mathbf{p}) \geq s(\mathbf{q})$, then it follows from (2.8) that $I = J$, thus $\mathbf{p}, \mathbf{q} \in \mathfrak{P}(I)$. By (2.13) we have then either $\Omega(\mathbf{p}) \cap \Omega(\mathbf{q}) = \emptyset$ or $\Omega(\mathbf{p}) = \Omega(\mathbf{q})$. By the assumption in (2.14) we have $\Omega(\mathbf{p}) \cap \Omega(\mathbf{q}) \neq \emptyset$, so we must have $\Omega(\mathbf{p}) = \Omega(\mathbf{q})$ and in particular $\Omega(\mathbf{q}) \subset \Omega(\mathbf{p})$.

So it remains to show (2.14) under the additional assumption that $s(\mathbf{q}) > s(\mathbf{p})$. In this case, we argue by induction on $s(\mathbf{q}) - s(\mathbf{p})$. By (2.7), there exists a cube $J \in \mathcal{D}$ with $s(J) = s(\mathbf{q}) - 1$ and $J \cap \mathcal{I}(\mathbf{p}) \neq \emptyset$. We pick one such J . By (2.8), we have $\mathcal{I}(\mathbf{p}) \subset J \subset \mathcal{I}(\mathbf{q})$.

By (4.46), there exists $x \in X$ with $d_{\mathbf{q}}(Q(x), \mathcal{Q}(\mathbf{q})) \leq 1$. By Lemma 2.8, it follows that $d_{J^\circ}(Q(x), \mathcal{Q}(\mathbf{q})) \leq 1$. Thus, by (4.48), there exists $z' \in \mathcal{Z}(J)$ with $\mathcal{Q}(\mathbf{q}) \in B_{J^\circ}(z', 0.7)$. Then by Lemma 4.15 there exists $\mathbf{q}' \in \mathfrak{P}(J)$ with $\mathcal{Q}(\mathbf{q}) \in \Omega_1(\mathbf{q}')$. By (4.53), it follows that $\Omega(\mathbf{q}) \subset \Omega(\mathbf{q}')$. Note that then $\mathcal{I}(\mathbf{p}) \subset \mathcal{I}(\mathbf{q}')$ and $\Omega(\mathbf{p}) \cap \Omega(\mathbf{q}') \neq \emptyset$ and $s(\mathbf{q}') - s(\mathbf{p}) = s(\mathbf{q}) - s(\mathbf{p}) - 1$. Thus, we have by the induction hypothesis that $\Omega(\mathbf{q}') \subset \Omega(\mathbf{p})$. This completes the proof. \square

5. PROOF OF DISCRETE CARLESON (PROP. 2.2)

Let a grid structure (\mathcal{D}, c, s) and a tile structure $(\mathfrak{P}, \mathcal{I}, \Omega, \mathcal{Q})$ for this grid structure be given. In Subsection 5.1, we decompose the set \mathfrak{P} of tiles into subsets. Each subset will be controlled by one of three methods. The guiding principle of the decomposition is to be able to apply the forest estimate of Proposition 2.4 to the final subsets defined in (5.23). This application is done in Subsection 5.4. The miscellaneous subsets along the construction of the forests will either be thrown into exceptional sets, which are defined and controlled in Subsection 5.2, or will be controlled by the antichain estimate of Proposition 2.3, which is done in Subsection 5.5. Subsection 5.3 contains some auxiliary lemmas needed for the proofs in Subsections 5.4-5.5.

5.1. Organisation of the tiles. In the following definitions, k, n , and j will be nonnegative integers. Define $\mathcal{C}(G, k)$ to be the set of $I \in \mathcal{D}$ such that

there exists a $J \in \mathcal{D}$ with $I \subset J$ and

$$\mu(G \cap J) > 2^{-k-1} \mu(J), \quad (5.1)$$

but there does not exist a $J \in \mathcal{D}$ with $I \subset J$ and

$$\mu(G \cap J) > 2^{-k} \mu(J). \quad (5.2)$$

Let

$$\mathfrak{P}(k) = \{\mathfrak{p} \in \mathfrak{P} : \mathcal{I}(\mathfrak{p}) \in \mathcal{C}(G, k)\} \quad (5.3)$$

Define $\mathfrak{M}(k, n)$ to be the set of $\mathfrak{p} \in \mathfrak{P}(k)$ such that

$$\mu(E_1(\mathfrak{p})) > 2^{-n} \mu(\mathcal{I}(\mathfrak{p})) \quad (5.4)$$

and there does not exist $\mathfrak{p}' \in \mathfrak{P}(k)$ with $\mathfrak{p}' \neq \mathfrak{p}$ and $\mathfrak{p} \leq \mathfrak{p}'$ such that

$$\mu(E_1(\mathfrak{p}')) > 2^{-n} \mu(\mathcal{I}(\mathfrak{p}')). \quad (5.5)$$

Define for a collection $\mathfrak{P}' \subset \mathfrak{P}(k)$

$$\text{dens}'_k(\mathfrak{P}') := \sup_{\mathfrak{p}' \in \mathfrak{P}'} \sup_{\lambda \geq 2} \lambda^{-a} \sup_{\mathfrak{p} \in \mathfrak{P}(k) : \lambda \mathfrak{p}' \lesssim \lambda \mathfrak{p}} \frac{\mu(E_2(\lambda, \mathfrak{p}))}{\mu(\mathcal{I}(\mathfrak{p}))}. \quad (5.6)$$

Sorting by density, we define

$$\mathfrak{C}(k, n) := \{\mathfrak{p} \in \mathfrak{P}(k) : 2^{4a} 2^{-n} < \text{dens}'_k(\{\mathfrak{p}\}) \leq 2^{4a} 2^{-n+1}\}. \quad (5.7)$$

Following Fefferman [Fef73], we define for $\mathfrak{p} \in \mathfrak{C}(k, n)$

$$\mathfrak{B}(\mathfrak{p}) := \{\mathfrak{m} \in \mathfrak{M}(k, n) : 100\mathfrak{p} \lesssim \mathfrak{m}\} \quad (5.8)$$

and

$$\mathfrak{C}_1(k, n, j) := \{\mathfrak{p} \in \mathfrak{C}(k, n) : 2^j \leq |\mathfrak{B}(\mathfrak{p})| < 2^{j+1}\}. \quad (5.9)$$

and

$$\mathfrak{L}_0(k, n) := \{\mathfrak{p} \in \mathfrak{C}(k, n) : |\mathfrak{B}(\mathfrak{p})| < 1\}. \quad (5.10)$$

Together with the following removal of minimal layers, the splitting into $\mathfrak{C}_1(k, n, j)$ will lead to a separation of trees. Define recursively for $0 \leq l \leq Z(n+1)$

$$\mathfrak{L}_1(k, n, j, l) \quad (5.11)$$

to be the set of minimal elements with respect to \leq in

$$\mathfrak{C}_1(k, n, j) \setminus \bigcup_{0 \leq l' < l} \mathfrak{L}_1(k, n, j, l'). \quad (5.12)$$

Define

$$\mathfrak{C}_2(k, n, j) := \mathfrak{C}_1(k, n, j) \setminus \bigcup_{0 \leq l' \leq Z(n+1)} \mathfrak{L}_1(k, n, j, l'). \quad (5.13)$$

The remaining tile organization will be relative to prospective tree tops, which we define now. Define

$$\mathfrak{U}_1(k, n, j) \quad (5.14)$$

to be the set of all $\mathfrak{u} \in \mathfrak{C}_1(k, n, j)$ such that for all $\mathfrak{p} \in \mathfrak{C}_1(k, n, j)$ with $\mathcal{I}(\mathfrak{u})$ strictly contained in $\mathcal{I}(\mathfrak{p})$ we have $B_{\mathfrak{u}}(\mathcal{Q}(\mathfrak{u}), 100) \cap B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), 100) = \emptyset$.

We first remove the pairs that are outside the immediate reach of any of the prospective tree tops. Define

$$\mathfrak{L}_2(k, n, j) \quad (5.15)$$

to be the set of all $\mathbf{p} \in \mathfrak{C}_2(k, n, j)$ such that there does not exist $\mathbf{u} \in \mathfrak{U}_1(k, n, j)$ with $\mathcal{I}(\mathbf{p}) \neq \mathcal{I}(\mathbf{u})$ and $2\mathbf{p} \lesssim \mathbf{u}$. Define

$$\mathfrak{C}_3(k, n, j) := \mathfrak{C}_2(k, n, j) \setminus \mathfrak{L}_2(k, n, j). \quad (5.16)$$

We next remove the maximal layers. Define recursively for $0 \leq l \leq Z(n+1)$

$$\mathfrak{L}_3(k, n, j, l) \quad (5.17)$$

to be the set of all maximal elements with respect to \leq in

$$\mathfrak{C}_3(k, n, j) \setminus \bigcup_{0 \leq l' < l} \mathfrak{L}_3(k, n, j, l'). \quad (5.18)$$

Define

$$\mathfrak{C}_4(k, n, j) := \mathfrak{C}_3(k, n, j) \setminus \bigcup_{0 \leq l \leq Z(n+1)} \mathfrak{L}_3(k, n, j, l). \quad (5.19)$$

Finally, we remove the boundary pairs relative to the prospective tree tops. Define

$$\mathcal{L}(\mathbf{u}) \quad (5.20)$$

to be the set of all $I \in \mathcal{D}$ with $I \subset \mathcal{I}(\mathbf{u})$ and $s(I) = s(\mathbf{u}) - Z(n+1) - 1$ and

$$B(c(I), 8D^{s(I)}) \not\subset \mathcal{I}(\mathbf{u}). \quad (5.21)$$

Define

$$\mathfrak{L}_4(k, n, j) \quad (5.22)$$

to be the set of all $\mathbf{p} \in \mathfrak{C}_4(k, n, j)$ such that there exists $\mathbf{u} \in \mathfrak{U}_1(k, n, j)$ with $\mathcal{I}(\mathbf{p}) \subset \bigcup \mathcal{L}(\mathbf{u})$, and define

$$\mathfrak{C}_5(k, n, j) := \mathfrak{C}_4(k, n, j) \setminus \mathfrak{L}_4(k, n, j). \quad (5.23)$$

We define three exceptional sets. The first exceptional set G_1 takes into account the ratio of the measures of F and G . Define $\mathfrak{P}_{F,G}$ to be the set of all $\mathbf{p} \in \mathfrak{P}$ with

$$\text{dens}_2(\{\mathbf{p}\}) \geq 2^{2a+5} \frac{\mu(F)}{\mu(G)}. \quad (5.24)$$

Define

$$G_1 := \bigcup_{\mathbf{p} \in \mathfrak{P}_{F,G}} \mathcal{I}(\mathbf{p}). \quad (5.25)$$

For an integer $\lambda \geq 0$, define $A(\lambda, k, n)$ to be the set of all $x \in X$ such that

$$\sum_{\mathbf{p} \in \mathfrak{M}(k, n)} \mathbf{1}_{\mathcal{I}(\mathbf{p})}(x) > 1 + \lambda 2^{n+1} \quad (5.26)$$

and define

$$G_2 := \bigcup_{k \geq 0} \bigcup_{k < n} A(2n+6, k, n). \quad (5.27)$$

Define

$$G_3 := \bigcup_{k \geq 0} \bigcup_{n \geq k} \bigcup_{0 \leq j \leq 2n+3} \bigcup_{\mathbf{p} \in \mathfrak{L}_4(k, n, j)} \mathcal{I}(\mathbf{p}). \quad (5.28)$$

Define $G' = G_1 \cup G_2 \cup G_3$. The following bound of the measure of G' will be proven in Subsection 5.2.

Lemma 5.1 (exceptional set). *We have*

$$\mu(G') \leq 2^{-2} \mu(G). \quad (5.29)$$

In Subsection 5.4, we identify each set $\mathfrak{C}_5(k, n, j)$ outside G' as forest and use Proposition 2.4 to prove the following lemma.

Lemma 5.2 (forest union). *Let*

$$\mathfrak{P}_1 = \bigcup_{k \geq 0} \bigcup_{n \geq k} \bigcup_{0 \leq j \leq 2n+3} \mathfrak{C}_5(k, n, j) \quad (5.30)$$

For all $f : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$ we have

$$\int_{G \setminus G'} \left| \sum_{\mathfrak{p} \in \mathfrak{P}_1} T_{\mathfrak{p}} f \right| d\mu \leq \frac{2^{435a^3}}{(q-1)^3} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \quad (5.31)$$

In Subsection 5.5, we decompose the complement of the set of tiles in Lemma 5.2 and apply the antichain estimate of Proposition 2.3 to prove the following lemma.

Lemma 5.3 (forest complement). *Let*

$$\mathfrak{P}_2 = \mathfrak{P} \setminus \mathfrak{P}_1. \quad (5.32)$$

For all $f : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$ we have

$$\int_{G \setminus G'} \left| \sum_{\mathfrak{p} \in \mathfrak{P}_2} T_{\mathfrak{p}} f \right| d\mu \leq \frac{2^{210a^3}}{(q-1)^4} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \quad (5.33)$$

Proposition 2.2 follows by applying triangle inequality to (2.22) according to the splitting in Lemma 5.2 and Lemma 5.3 and using both Lemmas as well as the bound on the set G' given by Lemma 5.1.

5.2. Proof of Lemma 5.1, the exceptional sets. We prove separate bounds for G_1 , G_2 , and G_3 in Lemmas 5.4, 5.9, and 5.13. Adding up these bounds proves Lemma 5.1.

The bound for G_1 is follows from the Vitali covering lemma, Proposition 2.6.

Lemma 5.4 (first exception). *We have*

$$\mu(G_1) \leq 2^{-4} \mu(G). \quad (5.34)$$

Proof. Let

$$K = 2^{2a+5} \frac{\mu(F)}{\mu(G)}.$$

For each $\mathfrak{p} \in \mathfrak{P}_{F,G}$ pick a $r(\mathfrak{p}) > 4D^{s(\mathfrak{p})}$ with

$$\mu(F \cap B(c(\mathfrak{p}), r(\mathfrak{p}))) \geq K \mu(B(c(\mathfrak{p}), r(\mathfrak{p}))).$$

This ball exists by definition of $\mathfrak{P}_{F,G}$ and dens_2 . By applying Proposition 2.6 to the collection of balls

$$\mathcal{B} = \{B(c(\mathfrak{p}), r(\mathfrak{p})) : \mathfrak{p} \in \mathfrak{P}_{F,G}\}$$

and the function $u = \mathbf{1}_F$, we obtain

$$\mu\left(\bigcup \mathcal{B}\right) \leq 2^{2a+1} K^{-1} \mu(F).$$

We conclude with (2.10) and $r(\mathfrak{p}) > 4D^s(\mathfrak{p})$

$$\mu(G_1) = \mu\left(\bigcup_{\mathfrak{p} \in \mathfrak{P}_{F,G}} \mathcal{I}(\mathfrak{p})\right) \leq \mu\left(\bigcup \mathcal{B}\right) \leq 2^{2a+1} K^{-1} \mu(F) = 2^{-4} \mu(G).$$

□

We turn to the bound of G_2 , which relies on the Dyadic Covering Lemma 5.5 and the John-Nirenberg Lemma 5.8 below.

Lemma 5.5 (dense cover). *For each $k \geq 0$, the union of all dyadic cubes in $\mathcal{C}(G, k)$ has measure at most $2^{k+1} \mu(G)$.*

Proof. The union of dyadic cubes in $\mathcal{C}(G, k)$ is contained the union of elements of the set $\mathcal{M}(k)$ of all dyadic cubes J with $\mu(G \cap J) > 2^{-k-1} \mu(J)$. The union of elements in the set $\mathcal{M}(k)$ is contained in the union of elements in the set $\mathcal{M}^*(k)$ of maximal elements in $\mathcal{M}(k)$ with respect to set inclusion. Hence

$$\mu\left(\bigcup \mathcal{C}(G, k)\right) \leq \mu\left(\bigcup \mathcal{M}^*(k)\right) \leq \sum_{J \in \mathcal{M}^*(k)} \mu(J) \quad (5.35)$$

Using the definition of $\mathcal{M}(k)$ and then the pairwise disjointedness of elements in $\mathcal{M}^*(k)$, we estimate (5.35) by

$$\leq 2^{k+1} \sum_{J \in \mathcal{M}^*(k)} \mu(J \cap G) \leq 2^{k+1} \mu(G). \quad (5.36)$$

This proves the lemma. □

Lemma 5.6 (pairwise disjoint). *If $\mathfrak{p}, \mathfrak{p}' \in \mathfrak{M}(k, n)$ and*

$$E_1(\mathfrak{p}) \cap E_1(\mathfrak{p}') \neq \emptyset, \quad (5.37)$$

then $\mathfrak{p} = \mathfrak{p}'$.

Proof. Let $\mathfrak{p}, \mathfrak{p}'$ be as in the lemma. By definition of E_1 , we have $E_1(\mathfrak{p}) \subset \mathcal{I}(\mathfrak{p})$ and analogously for \mathfrak{p}' , we conclude from (5.37) that $\mathcal{I}(\mathfrak{p}) \cap \mathcal{I}(\mathfrak{p}') \neq \emptyset$. Let without loss of generality $\mathcal{I}(\mathfrak{p})$ be maximal in $\{\mathcal{I}(\mathfrak{p}), \mathcal{I}(\mathfrak{p}')\}$, then $\mathcal{I}(\mathfrak{p}') \subset \mathcal{I}(\mathfrak{p})$. By (5.37), we conclude by definition of E_1 that $\Omega(\mathfrak{p}) \cap \Omega(\mathfrak{p}') \neq \emptyset$. By (2.14) we conclude $\Omega(\mathfrak{p}) \subset \Omega(\mathfrak{p}')$. It follows that $\mathfrak{p}' \leq \mathfrak{p}$. By maximality (5.5) of \mathfrak{p}' , we have $\mathfrak{p}' = \mathfrak{p}$. This proves the lemma. □

Lemma 5.7 (dyadic union). *For each $x \in A(\lambda, k, n)$, there is a dyadic cube I that contains x and is a subset of $A(\lambda, k, n)$.*

Proof. Fix k, n, λ, x as in the lemma such that $x \in A(\lambda, k, n)$. Let \mathcal{M} be the set of dyadic cubes $\mathcal{I}(\mathfrak{p})$ with \mathfrak{p} in $\mathfrak{M}(k, n)$ and $x \in \mathcal{I}(\mathfrak{p})$. By definition of $A(\lambda, k, n)$, the cardinality of \mathcal{M} is at least $1 + \lambda 2^{n+1}$. Let I be a cube of smallest scale in \mathcal{M} . Then I is contained in all cubes of \mathcal{M} . It follows that $I \subset A(\lambda, k, n)$. □

Lemma 5.8 (John Nirenberg). *For all integers $k, n, \lambda \geq 0$, we have*

$$\mu(A(\lambda, k, n)) \leq 2^{k+1-\lambda} \mu(G). \quad (5.38)$$

Proof. Fix k, n as in the lemma and suppress notation to write $A(\lambda)$ for $A(\lambda, k, n)$. We prove the lemma by induction on λ . For $\lambda = 0$, we use that $A(\lambda)$ by definition of $\mathfrak{M}(k, n)$ is contained in the union of elements in $\mathcal{C}(G, k)$. Lemma 5.5 then completes the base of the induction.

Now assume that the statement of Lemma 5.8 is proven for some integer $\lambda \geq 0$. The set $A(\lambda + 1)$ is contained in the set $A(\lambda)$. Let \mathcal{M} be the set of dyadic cubes which are a subset of $A(\lambda)$. By Lemma 5.7, the union of \mathcal{M} is $A(\lambda)$. Let \mathcal{M}^* be the set of maximal dyadic cubes in \mathcal{M} .

Let $L \in \mathcal{M}^*$. For each $x \in L$, we have

$$\sum_{\mathfrak{p} \in \mathfrak{M}(k, n)} \mathbf{1}_{\mathcal{I}(\mathfrak{p})}(x) = \sum_{\mathfrak{p} \in \mathfrak{M}(k, n): \mathcal{I}(\mathfrak{p}) \subset L} \mathbf{1}_{\mathcal{I}(\mathfrak{p})}(x) + \sum_{\mathfrak{p} \in \mathfrak{M}(k, n): \mathcal{I}(\mathfrak{p}) \not\subset L} \mathbf{1}_{\mathcal{I}(\mathfrak{p})}(x). \quad (5.39)$$

If the second sum on the right-hand-side is not zero, there is an element of \mathcal{D} strictly containing L . Let \hat{L} be such a dyadic cube with minimal $s(L)$. Then \hat{L} is contained in $\mathcal{I}(\mathfrak{p})$ for all \mathfrak{p} contributing to the second sum in (5.39). Hence the second sum in (5.39) is constant on \hat{L} . By maximality of L , the second sum is less than $1 + \lambda 2^{n+1}$ somewhere on \hat{L} , thus on all of \hat{L} and consequently also at x . If x is in addition in $A(\lambda + 1)$, then the left-hand-side of (5.39) is at least $1 + (\lambda + 1)2^{n+1}$, so we have by the triangle inequality for the first sum on the right-hand side

$$\sum_{\mathfrak{p} \in \mathfrak{M}(k, n): \mathcal{I}(\mathfrak{p}) \subset L} \mathbf{1}_{\mathcal{I}(\mathfrak{p})}(x) \geq 2^{n+1}. \quad (5.40)$$

By Lemma 5.6, we have

$$\sum_{\mathfrak{p} \in \mathfrak{M}(k, n): \mathcal{I}(\mathfrak{p}) \subset L} \mu(E_1(\mathfrak{p})) \leq \mu(L). \quad (5.41)$$

Multiplying by 2^n and applying (5.4), we obtain

$$\sum_{\mathfrak{p} \in \mathfrak{M}(k, n): \mathcal{I}(\mathfrak{p}) \subset L} \mu(\mathcal{I}(\mathfrak{p})) \leq 2^n \mu(L). \quad (5.42)$$

We then have with (5.40) and (5.42)

$$2^{n+1} \mu(A(\lambda + 1) \cap L) = \int_{A(\lambda+1) \cap L} 2^{n+1} d\mu \quad (5.43)$$

$$\leq \int \sum_{\mathfrak{p} \in \mathfrak{M}(k, n): \mathcal{I}(\mathfrak{p}) \subset L} \mathbf{1}_{\mathcal{I}(\mathfrak{p})} d\mu \leq 2^n \mu(L). \quad (5.44)$$

Hence

$$2\mu(A(\lambda + 1)) = 2 \sum_{L \in \mathcal{M}^*} \mu(A(\lambda + 1) \cap L) \leq \sum_{L \in \mathcal{M}^*} \mu(L) = \mu(A(\lambda)). \quad (5.45)$$

Using the induction hypothesis, this proves (5.38) for $\lambda + 1$ and completes the proof of the lemma. \square

Lemma 5.9 (second exception). *We have*

$$\mu(G_2) \leq 2^{-4} \mu(G). \quad (5.46)$$

Proof. We use Lemma 5.8 and sum twice a geometric series to obtain

$$\sum_{0 \leq k} \sum_{k < n} \mu(A(2n+6, k, n)) \leq \sum_{0 \leq k} \sum_{k < n} 2^{k-5-2n} \mu(G) \quad (5.47)$$

$$\leq \sum_{0 \leq k} 2^{-k-5} \mu(G) \leq 2^{-4} \mu(G). \quad (5.48)$$

This proves the lemma. \square

We turn to the set G_3 .

Lemma 5.10 (top tiles). *We have*

$$\sum_{\mathbf{m} \in \mathfrak{M}(k, n)} \mu(\mathcal{I}(\mathbf{m})) \leq 2^{n+1} 2^{k+1} \mu(G). \quad (5.49)$$

Proof. We write the left-hand side of (5.49)

$$\int \sum_{\mathbf{m} \in \mathfrak{M}(k, n)} \mathbf{1}_{\mathcal{I}(\mathbf{m})}(x) d\mu(x) \leq 2^{n+1} \sum_{\lambda=1}^{|\mathfrak{M}|} \mu(A(\lambda, k, n)). \quad (5.50)$$

Using Lemma 5.8 and then summing a geometric series, we estimate this by

$$\leq 2^{n+1} \sum_{\lambda=1}^{|\mathfrak{M}|} 2^{k+1-\lambda} \mu(G) \leq 2^{n+1} 2^{k+1} \mu(G). \quad (5.51)$$

This proves the lemma. \square

Lemma 5.11 (tree count). *Let $k, n, j \geq 0$. We have for every $x \in X$*

$$\sum_{\mathbf{u} \in \mathfrak{U}_1(k, n, j)} \mathbf{1}_{\mathcal{I}(\mathbf{u})}(x) \leq 2^{-j} 2^{9a} \sum_{\mathbf{m} \in \mathfrak{M}(k, n)} \mathbf{1}_{\mathcal{I}(\mathbf{m})}(x) \quad (5.52)$$

Proof. Let $x \in X$. For each $\mathbf{u} \in \mathfrak{U}_1(k, n, j)$ with $x \in \mathcal{I}(\mathbf{u})$, as $\mathbf{u} \in \mathfrak{C}_1(k, n, j)$, there are at least 2^j elements $\mathbf{m} \in \mathfrak{M}(k, n)$ with $100\mathbf{u} \lesssim \mathbf{m}$ and in particular $x \in \mathcal{I}(\mathbf{m})$. Hence

$$\mathbf{1}_{\mathcal{I}(\mathbf{u})}(x) \leq 2^{-j} \sum_{\mathbf{m} \in \mathfrak{M}(k, n): 100\mathbf{u} \lesssim \mathbf{m}} \mathbf{1}_{\mathcal{I}(\mathbf{m})}(x). \quad (5.53)$$

Conversely, for each $\mathbf{m} \in \mathfrak{M}(k, n)$ with $x \in \mathcal{I}(\mathbf{m})$, let $\mathfrak{U}(\mathbf{m})$ be the set of $\mathbf{u} \in \mathfrak{U}_1(k, n, j)$ with $x \in \mathcal{I}(\mathbf{u})$ and $100\mathbf{u} \lesssim \mathbf{m}$. Summing (5.53) over \mathbf{u} and counting the pairs (\mathbf{u}, \mathbf{m}) with $100\mathbf{u} \lesssim \mathbf{m}$ differently gives

$$\sum_{\mathbf{u} \in \mathfrak{U}_1(k, n, j)} \mathbf{1}_{\mathcal{I}(\mathbf{u})}(x) \leq 2^{-j} \sum_{\mathbf{m} \in \mathfrak{M}(k, n)} \sum_{\mathbf{u} \in \mathfrak{U}(\mathbf{m})} \mathbf{1}_{\mathcal{I}(\mathbf{m})}(x). \quad (5.54)$$

We estimate the number of elements in $\mathfrak{U}(\mathbf{m})$. Let $\mathbf{u} \in \mathfrak{U}(\mathbf{m})$. Then by definition of $\mathfrak{U}(\mathbf{m})$

$$d_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), \mathcal{Q}(\mathbf{m})) \leq 100. \quad (5.55)$$

If \mathbf{u}' is a further element in $\mathfrak{U}(\mathbf{m})$ with $\mathbf{u} \neq \mathbf{u}'$, then

$$\mathcal{Q}(\mathbf{m}) \in B_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), 100) \cap B_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), 100). \quad (5.56)$$

By the last display and definition of $\mathfrak{U}_1(k, n, j)$, none of $\mathcal{I}(\mathbf{u}), \mathcal{I}(\mathbf{u}')$ is strictly contained in the other. As both contain x , we have $\mathcal{I}(\mathbf{u}) = \mathcal{I}(\mathbf{u}')$. We then have $d_{\mathbf{u}} = d_{\mathbf{u}'}$.

By (2.15), the balls $B_u(\mathcal{Q}(u), 0.2)$ and $B_u(\mathcal{Q}(u'), 0.2)$ are contained respectively in $\Omega(u)$ and $\Omega(u')$ and thus are disjoint by (2.13). By (5.55) and the triangle inequality, both balls are contained in $B_u(\mathcal{Q}(m), 100.2)$.

By (1.11) applied nine times, there is a collection of at most 2^{9a} balls of radius 0.2 with respect to the metric d_u which cover the ball $B_u(\mathcal{Q}(m), 100.2)$. Let B' be a ball in this cover. As the center of B' can be in at most one of the disjoint balls $B_u(\mathcal{Q}(u), 0.2)$ and $B_u(\mathcal{Q}(u'), 0.2)$, the ball B' can contain at most one of the points u, u' .

Hence the set $\mathfrak{U}(m)$ has at most 2^{9a} many elements. Inserting this into (5.54) proves the lemma. \square

Lemma 5.12 (boundary exception). *Let $\mathcal{L}(u)$ be as defined in (5.20). We have for each $u \in \mathfrak{U}_1(k, n, l)$,*

$$\mu\left(\bigcup_{I \in \mathcal{L}(u)} I\right) \leq 2^{100a^2} D^{-\kappa Z(n+1)} \mu(\mathcal{I}(u)). \quad (5.57)$$

Proof. Let $u \in \mathfrak{U}_1(k, n, l)$. Let $I \in \mathcal{L}(u)$. Then we have $s(I) = s(u) - Z(n+1) - 1$ and $I \subset \mathcal{I}(u)$ and $B(c(I), 8D^{s(I)}) \not\subset \mathcal{I}(u)$. By (2.10), the set I is contained in $B(c(I), 4D^{s(I)})$. By the triangle inequality, the set I is contained in

$$X(u) := \{x \in \mathcal{I}(u) : \rho(x, X \setminus \mathcal{I}(u)) \leq 12D^{s(u)-Z(n+1)-1}\}. \quad (5.58)$$

By the small boundary property (2.11), noting that

$$12D^{s(u)-Z(n+1)-1} = 12D^{s(I)} > D^{-S},$$

we have

$$\mu(X(u)) \leq 2^{100a^2} (12D^{-Z(n+1)-1})^\kappa \mu(\mathcal{I}(u)).$$

Using $\kappa < 1$ and $D \geq 12$, this proves the lemma. \square

Lemma 5.13 (third exception). *We have*

$$\mu(G_3) \leq 2^{-4} \mu(G). \quad (5.59)$$

Proof. As each $p \in \mathfrak{L}_4(k, n, j)$ is contained in $\cup \mathcal{L}(u)$ for some $u \in \mathfrak{U}_1(k, n, l)$, we have

$$\mu\left(\bigcup_{p \in \mathfrak{L}_4(k, n, j)} \mathcal{I}(p)\right) \leq \sum_{u \in \mathfrak{U}_1(k, n, j)} \mu\left(\bigcup_{I \in \mathcal{L}(u)} I\right). \quad (5.60)$$

Using Lemma 5.12 and then Lemma 5.11, we estimate this further by

$$\leq \sum_{u \in \mathfrak{U}_1(k, n, j)} 2^{100a^2} D^{-\kappa Z(n+1)} \mu(\mathcal{I}(u)) \quad (5.61)$$

$$\leq 2^{100a^2+9a+1-j} \sum_{m \in \mathfrak{M}(k, n)} D^{-\kappa Z(n+1)} \mu(\mathcal{I}(m)). \quad (5.62)$$

Using Lemma 5.10, we estimate this by

$$\leq 2^{100a^2+9a+1-j} D^{-\kappa Z(n+1)} 2^{n+1} 2^{k+1} \mu(G). \quad (5.63)$$

Now we estimate G_3 defined in (5.28) by

$$\mu(G_3) \leq \sum_{k \geq 0} \sum_{n \geq k} \sum_{0 \leq j \leq 2n+3} \mu\left(\bigcup_{p \in \mathfrak{L}_4(k, n, j)} \mathcal{I}(p)\right) \quad (5.64)$$

$$\leq \sum_{k \geq 0} \sum_{n \geq k} \sum_{0 \leq j \leq 2n+3} 2^{100a^2+9a+3+n+k-j} D^{-\kappa Z(n+1)} \mu(G) \quad (5.65)$$

Summing geometric series, using that $D^{\kappa Z} \geq 8$ by (2.1), (2.2) and (2.3), we estimate this by

$$\leq \sum_{k \geq 0} \sum_{n \geq k} 2^{100a^2+9a+4+n+k} D^{-\kappa Z(n+1)} \mu(G) \quad (5.66)$$

$$= \sum_{k \geq 0} 2^{100a^2+9a+4+2k} D^{-\kappa Z(k+1)} \sum_{n \geq k} 2^{n-k} D^{-\kappa Z(n-k)} \mu(G) \quad (5.67)$$

$$\leq \sum_{k \geq 0} 2^{100a^2+9a+5+2k} D^{-\kappa Z(k+1)} \mu(G) \quad (5.68)$$

$$\leq 2^{100a^2+9a+6} D^{-\kappa Z} \mu(G) \quad (5.69)$$

Using $D = 2^{100a^2}$ and $a \geq 4$ and $\kappa Z \geq 2$ by (2.1) and (2.2) proves the lemma. \square

5.3. Auxiliary lemmas. Before proving Lemma 5.2 and Lemma 5.3, we collect some useful properties of \lesssim .

Lemma 5.14 (wiggle order 1). *If $n\mathfrak{p} \lesssim m\mathfrak{p}'$ and $n' \geq n$ and $m \geq m'$ then $n'\mathfrak{p} \lesssim m'\mathfrak{p}'$.*

Proof. This follows immediately from the definition (2.24) of \lesssim and the two inclusions $B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), n) \subset B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), n')$ and $B_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), m') \subset B_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), m)$. \square

Lemma 5.15 (wiggle order 2).

Let $n, m \geq 1$. If $\mathfrak{p}, \mathfrak{p}' \in \mathfrak{P}$ with $\mathcal{I}(\mathfrak{p}) \neq \mathcal{I}(\mathfrak{p}')$ and

$$n\mathfrak{p} \lesssim \mathfrak{p}' \quad (5.70)$$

then

$$(n + 2^{-95a}m)\mathfrak{p} \lesssim m\mathfrak{p}'. \quad (5.71)$$

Proof. The assumption (5.70) together with the definition (2.24) of \lesssim implies that $\mathcal{I}(\mathfrak{p}) \subsetneq \mathcal{I}(\mathfrak{p}')$. Let $\vartheta \in B_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), m)$. Then we have by the triangle inequality

$$d_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), \vartheta) \leq d_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), \mathcal{Q}(\mathfrak{p}')) + d_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}'), \vartheta)$$

Using (5.70) and (2.24) for the first summand, and Lemma 2.8 for the second summand, this is estimated by

$$n + 2^{-95a}d_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), \vartheta) < n + 2^{-95a}m.$$

Thus $B_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), m) \subset B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), n + 2^{-95a}m)$. Combined with $\mathcal{I}(\mathfrak{p}) \subset \mathcal{I}(\mathfrak{p}')$, this yields (5.71). \square

Lemma 5.16 (wiggle order 3). *The following implications hold for all $\mathfrak{q}, \mathfrak{q}' \in \mathfrak{P}$:*

$$\mathfrak{q} \lesssim \mathfrak{q}' \text{ and } \lambda \geq 1 \implies \lambda\mathfrak{q} \lesssim \lambda\mathfrak{q}', \quad (5.72)$$

$$10\mathfrak{q} \lesssim \mathfrak{q}' \text{ and } \mathfrak{q} \neq \mathfrak{q}' \implies 100\mathfrak{q} \lesssim 100\mathfrak{q}', \quad (5.73)$$

$$2\mathfrak{q} \lesssim \mathfrak{q}' \text{ and } \mathfrak{q} \neq \mathfrak{q}' \implies 4\mathfrak{q} \lesssim 500\mathfrak{q}'. \quad (5.74)$$

Proof. All three implications are easy consequences of Lemma 5.14, Lemma 5.15 and the fact that $a \geq 4$. \square

We call a collection \mathfrak{A} of tiles convex if

$$\mathfrak{p} \leq \mathfrak{p}' \leq \mathfrak{p}'' \text{ and } \mathfrak{p}, \mathfrak{p}'' \in \mathfrak{A} \implies \mathfrak{p}' \in \mathfrak{A}. \quad (5.75)$$

Lemma 5.17 (P convex). *For each k , the collection $\mathfrak{P}(k)$ is convex.*

Proof. Suppose that $\mathfrak{p} \leq \mathfrak{p}' \leq \mathfrak{p}''$ and $\mathfrak{p}, \mathfrak{p}'' \in \mathfrak{P}(k)$. By (5.3) we have $\mathcal{I}(\mathfrak{p}), \mathcal{I}(\mathfrak{p}'') \in \mathcal{C}(G, k)$, so there exists by (5.1) some $J \in \mathcal{D}$ with

$$\mathcal{I}(\mathfrak{p}') \subset \mathcal{I}(\mathfrak{p}'') \subset J$$

and $\mu(G \cap J) > 2^{-k-1}\mu(J)$. Thus (5.1) holds for $\mathcal{I}(\mathfrak{p}')$. On the other hand, by (5.2), there exists no $J \in \mathcal{D}$ with $\mathcal{I}(\mathfrak{p}) \subset J$ and $\mu(G \cap J) > 2^{-k}\mu(J)$. Since $\mathcal{I}(\mathfrak{p}) \subset \mathcal{I}(\mathfrak{p}')$, this implies that (5.2) holds for $\mathcal{I}(\mathfrak{p}')$. Hence $\mathcal{I}(\mathfrak{p}') \in \mathcal{C}(G, k)$, and therefore by (5.3) $\mathfrak{p}' \in \mathfrak{P}(k)$. \square

Lemma 5.18 (C convex). *For each k, n , the collection $\mathfrak{C}(k, n)$ is convex.*

Proof. Let $\mathfrak{p} \leq \mathfrak{p}' \leq \mathfrak{p}''$ with $\mathfrak{p}, \mathfrak{p}'' \in \mathfrak{C}(k, n)$. Then, in particular, $\mathfrak{p}, \mathfrak{p}'' \in \mathfrak{P}(k)$, so, by Lemma 5.17, $\mathfrak{p}' \in \mathfrak{P}(k)$. Next, we show that if $\mathfrak{q} \leq \mathfrak{q}' \in \mathfrak{P}(k)$ then $\text{dens}'_k(\{\mathfrak{q}\}) \geq \text{dens}'_k(\{\mathfrak{q}'\})$. If $\mathfrak{p} \in \mathfrak{P}(k)$ and $\lambda \geq 2$ with $\lambda\mathfrak{q}' \lesssim \lambda\mathfrak{p}$, then it follows from $\mathfrak{q} \leq \mathfrak{q}'$, (5.72) of Lemma 5.16 and transitivity of \lesssim that $\lambda\mathfrak{q} \lesssim \lambda\mathfrak{p}$. Thus the supremum in the definition (5.6) of $\text{dens}'_k(\{\mathfrak{q}\})$ is over a superset of the set the supremum in the definition of $\text{dens}'_k(\{\mathfrak{q}'\})$ is taken over, which shows $\text{dens}'_k(\{\mathfrak{q}\}) \geq \text{dens}'_k(\{\mathfrak{q}'\})$. From $\mathfrak{p}' \leq \mathfrak{p}''$, $\mathfrak{p}'' \in \mathfrak{C}(k, n)$ and (5.7) it then follows that

$$2^{4a}2^{-n} < \text{dens}'_k(\{\mathfrak{p}''\}) \leq \text{dens}'_k(\{\mathfrak{p}'\}).$$

Similarly, it follows from $\mathfrak{p} \leq \mathfrak{p}'$, $\mathfrak{p} \in \mathfrak{C}(k, n)$ and (5.7) that

$$\text{dens}'_k(\{\mathfrak{p}'\}) \leq \text{dens}'_k(\{\mathfrak{p}\}) \leq 2^{4a}2^{-n+1}.$$

Thus $\mathfrak{p}' \in \mathfrak{C}(k, n)$. \square

Lemma 5.19 (C1 convex). *For each k, n, j , the collection $\mathfrak{C}_1(k, n, j)$ is convex.*

Proof. Let $\mathfrak{p} \leq \mathfrak{p}' \leq \mathfrak{p}''$ with $\mathfrak{p}, \mathfrak{p}'' \in \mathfrak{C}_1(k, n, j)$. By Lemma 5.18 and the inclusion $\mathfrak{C}_1(k, n, j) \subset \mathfrak{C}(k, n)$, which holds by definition (5.9), we have $\mathfrak{p}' \in \mathfrak{C}(k, n)$. By (5.72) and transitivity of \lesssim we have that $\mathfrak{q} \leq \mathfrak{q}'$ and $100\mathfrak{q}' \lesssim \mathfrak{m}$ imply $100\mathfrak{q} \lesssim \mathfrak{m}$. So, by (5.8), $\mathfrak{B}(\mathfrak{p}'') \subset \mathfrak{B}(\mathfrak{p}') \subset \mathfrak{B}(\mathfrak{p})$. Consequently, by (5.9)

$$2^j \leq |\mathfrak{B}(\mathfrak{p}'')| \leq |\mathfrak{B}(\mathfrak{p}')| \leq |\mathfrak{B}(\mathfrak{p})| < 2^{j+1},$$

thus $\mathfrak{p}' \in \mathfrak{C}_1(k, n, j)$. \square

Lemma 5.20 (C2 convex). *For each k, n, j , the collection $\mathfrak{C}_2(k, n, j)$ is convex.*

Proof. Let $\mathfrak{p} \leq \mathfrak{p}' \leq \mathfrak{p}''$ with $\mathfrak{p}, \mathfrak{p}'' \in \mathfrak{C}_2(k, n, j)$. By (5.13), we have

$$\mathfrak{C}_2(k, n, j) \subset \mathfrak{C}_1(k, n, j).$$

Combined with Lemma 5.19, it follows that $\mathfrak{p}' \in \mathfrak{C}_1(k, n, j)$. Suppose that $\mathfrak{p}' \notin \mathfrak{C}_2(k, n, j)$. By (5.13), this implies that there exists $0 \leq l' \leq Z(n+1)$ with $\mathfrak{p}' \in \mathfrak{L}_1(k, n, j, l')$. By the definition (5.11) of $\mathfrak{L}_1(k, n, j, l')$, this implies that \mathfrak{p} is minimal with respect to \leq in $\mathfrak{C}_1(k, n, j) \setminus \bigcup_{l < l'} \mathfrak{L}_1(k, n, j, l)$. Since $\mathfrak{p} \in \mathfrak{C}_2(k, n, j)$ we must have $\mathfrak{p} \neq \mathfrak{p}'$. Thus $\mathfrak{p} \leq \mathfrak{p}'$ and $\mathfrak{p} \neq \mathfrak{p}'$. By minimality

of \mathbf{p}' it follows that $\mathbf{p} \notin \mathfrak{C}_1(k, n, j) \setminus \bigcup_{l < l'} \mathfrak{L}_1(k, n, j, l)$. But by (5.13) this implies $\mathbf{p} \notin \mathfrak{C}_2(k, n, j)$, a contradiction. \square

Lemma 5.21 (C3 convex). *For each k, n, j , the collection $\mathfrak{C}_3(k, n, j)$ is convex.*

Proof. Let $\mathbf{p} \leq \mathbf{p}' \leq \mathbf{p}''$ with $\mathbf{p}, \mathbf{p}'' \in \mathfrak{C}_3(k, n, j)$. By (5.16) and Lemma 5.20 it follows that $\mathbf{p}' \in \mathfrak{C}_2(k, n, j)$. Suppose that $\mathbf{p}' \notin \mathfrak{C}_3(k, n, j)$. Then, by (5.16) and (5.15), there exists $\mathbf{u} \in \mathfrak{U}_1(k, n, j)$ with $2\mathbf{p}' \lesssim \mathbf{u}$ and $\mathcal{I}(\mathbf{p}') \neq \mathcal{I}(\mathbf{u})$. Together this gives $\mathcal{I}(\mathbf{p}') \subsetneq \mathcal{I}(\mathbf{u})$. From $\mathbf{p}' \leq \mathbf{p}$, (5.72) and transitivity of \lesssim we then have $2\mathbf{p} \lesssim \mathbf{u}$. Also, $\mathcal{I}(\mathbf{p}) \subset \mathcal{I}(\mathbf{p}') \subsetneq \mathcal{I}(\mathbf{u})$, so $\mathcal{I}(\mathbf{p}) \neq \mathcal{I}(\mathbf{u})$. But then $\mathbf{p} \in \mathfrak{L}_2(k, n, j)$, contradicting by (5.16) the assumption $\mathbf{p} \in \mathfrak{C}_3(k, n, j)$. \square

Lemma 5.22 (C4 convex). *For each k, n, j , the collection $\mathfrak{C}_4(k, n, j)$ is convex.*

Proof. Let $\mathbf{p} \leq \mathbf{p}' \leq \mathbf{p}''$ with $\mathbf{p}, \mathbf{p}'' \in \mathfrak{C}_4(k, n, j)$. As before we obtain from the inclusion $\mathfrak{C}_4(k, n, j) \subset \mathfrak{C}_3(k, n, j)$ and Lemma 5.21 that $\mathbf{p}' \in \mathfrak{C}_3(k, n, j)$. Thus, if $\mathbf{p}' \notin \mathfrak{C}_4(k, n, j)$ then by (5.17) there exists l such that $\mathbf{p}' \in \mathfrak{L}_3(k, n, j, l)$. Thus \mathbf{p}' is maximal with respect to \leq in $\mathfrak{C}_3(k, n, j) \setminus \bigcup_{0 \leq l' < l} \mathfrak{L}_3(k, n, j, l')$. Since $\mathbf{p}'' \in \mathfrak{C}_4(k, n, j)$ we must have $\mathbf{p}' \neq \mathbf{p}''$. Thus $\mathbf{p}' \leq \mathbf{p}''$ and $\mathbf{p}' \neq \mathbf{p}''$. By minimality of \mathbf{p}' it follows that $\mathbf{p}'' \notin \mathfrak{C}_3(k, n, j) \setminus \bigcup_{l < l'} \mathfrak{L}_3(k, n, j, l)$. But by (5.19) this implies $\mathbf{p}'' \notin \mathfrak{C}_4(k, n, j)$, a contradiction. \square

Lemma 5.23 (C5 convex). *For each k, n, j , the collection $\mathfrak{C}_5(k, n, j)$ is convex.*

Proof. Let $\mathbf{p} \leq \mathbf{p}' \leq \mathbf{p}''$ with $\mathbf{p}, \mathbf{p}'' \in \mathfrak{C}_5(k, n, j)$. Then $\mathbf{p}, \mathbf{p}'' \in \mathfrak{C}_4(k, n, j)$ by (5.23), and thus by Lemma 5.22 also $\mathbf{p}' \in \mathfrak{C}_4(k, n, j)$. Suppose that $\mathbf{p}' \notin \mathfrak{C}_5(k, n, j)$. By (5.23), it follows that $\mathbf{p}' \in \mathfrak{L}_4(k, n, j)$. By (5.22), there exists $\mathbf{u} \in \mathfrak{U}_1(k, n, j)$ with $\mathcal{I}(\mathbf{p}') \subset \bigcup \mathcal{L}(\mathbf{u})$. Then also $\mathcal{I}(\mathbf{p}) \subset \bigcup \mathcal{L}(\mathbf{u})$, a contradiction. \square

Lemma 5.24 (dens compare). *We have for every $k \geq 0$ and $\mathfrak{P}' \subset \mathfrak{P}(k)$*

$$\text{dens}_1(\mathfrak{P}') \leq \text{dens}'_k(\mathfrak{P}'). \quad (5.76)$$

Proof. It suffices to show that for all $\mathbf{p}' \in \mathfrak{P}'$ and $\lambda \geq 2$ and $\mathbf{p} \in \mathfrak{P}(\mathfrak{P}')$ with $\lambda\mathbf{p}' \lesssim \lambda\mathbf{p}$ we have

$$\frac{\mu(E_2(\lambda, \mathbf{p}))}{\mu(\mathcal{I}(\mathbf{p}))} \leq \sup_{\mathbf{p}'' \in \mathfrak{P}(k): \lambda\mathbf{p}' \lesssim \lambda\mathbf{p}''} \frac{\mu(E_2(\lambda, \mathbf{p}''))}{\mu(\mathcal{I}(\mathbf{p}''))}. \quad (5.77)$$

Let such \mathbf{p}' , λ , \mathbf{p} be given. It suffices to show that $\mathbf{p} \in \mathfrak{P}(k)$, that is, it satisfies (5.1) and (5.2).

We show (5.1). As $\mathbf{p} \in \mathfrak{P}(\mathfrak{P}')$, there exists $\mathbf{p}'' \in \mathfrak{P}'$ with $\mathcal{I}(\mathbf{p}') \subset \mathcal{I}(\mathbf{p}'')$. By assumption on \mathfrak{P}' , we have $\mathbf{p}'' \in \mathfrak{P}(k)$ and there exists $J \in \mathcal{D}$ with $\mathcal{I}(\mathbf{p}'') \subset J$ and

$$\mu(G \cap J) > 2^{-k-1} \mu(J). \quad (5.78)$$

Then also $\mathcal{I}(\mathbf{p}') \subset J$, which proves (5.1) for \mathbf{p} .

We show (5.2). Assume to get a contradiction that there exists $J \in \mathcal{D}$ with $\mathcal{I}(\mathbf{p}) \subset J$ and

$$\mu(G \cap J) > 2^{-k} \mu(J). \quad (5.79)$$

As $\lambda\mathbf{p}' \lesssim \lambda\mathbf{p}$, we have $\mathcal{I}(\mathbf{p}') \subset \mathcal{I}(\mathbf{p})$, and therefore $\mathcal{I}(\mathbf{p}') \subset J$. This contradicts $\mathbf{p}' \in \mathfrak{P}' \subset \mathfrak{P}(k)$. This proves (5.2) for \mathbf{p} . \square

Lemma 5.25 (C dens1). *For each set $\mathfrak{A} \subset \mathfrak{C}(k, n)$, we have*

$$\text{dens}_1(\mathfrak{A}) \leq 2^{4a} 2^{-n+1}.$$

Proof. We have by Lemma 5.24 that $\text{dens}_1(\mathfrak{A}) \leq \text{dens}'_k(\mathfrak{A})$. Since $\mathfrak{A} \subset \mathfrak{C}(k, n)$, it follows from monotonicity of suprema and the definition (5.6) that $\text{dens}'_k(\mathfrak{A}) \leq \text{dens}'_k(\mathfrak{C}(k, n))$. By (5.6) and (5.7), we have

$$\text{dens}'_k(\mathfrak{C}(k, n)) = \sup_{\mathfrak{p} \in \mathfrak{C}(k, n)} \text{dens}'_k(\{\mathfrak{p}\}) \leq 2^{4a} 2^{-n+1}.$$

□

5.4. Proof of Lemma 5.2, the forests. Fix $k, n, j \geq 0$. Define

$$\mathfrak{C}_6(k, n, j)$$

to be the set of all tiles $\mathfrak{p} \in \mathfrak{C}_5(k, n, j)$ such that $\mathcal{I}(\mathfrak{p}) \not\subset G'$. The following chain of lemmas establishes that the set $\mathfrak{C}_6(k, n, j)$ can be written as a union of a small number of n -forests.

For $\mathfrak{u} \in \mathfrak{U}_1(k, n, j)$, define

$$\mathfrak{T}_1(\mathfrak{u}) := \{\mathfrak{p} \in \mathfrak{C}_1(k, n, j) : \mathcal{I}(\mathfrak{p}) \neq \mathcal{I}(\mathfrak{u}), 2\mathfrak{p} \lesssim \mathfrak{u}\}. \quad (5.80)$$

Define

$$\mathfrak{U}_2(k, n, j) := \{\mathfrak{u} \in \mathfrak{U}_1(k, n, j) : \mathfrak{T}_1(\mathfrak{u}) \cap \mathfrak{C}_6(k, n, j) \neq \emptyset\}. \quad (5.81)$$

Define a relation \sim on $\mathfrak{U}_2(k, n, j)$ by setting $\mathfrak{u} \sim \mathfrak{u}'$ for $\mathfrak{u}, \mathfrak{u}' \in \mathfrak{U}_2(k, n, j)$ if $\mathfrak{u} = \mathfrak{u}'$ or there exists \mathfrak{p} in $\mathfrak{T}_1(\mathfrak{u})$ with $10\mathfrak{p} \lesssim \mathfrak{u}'$.

Lemma 5.26 (relation geometry). *If $\mathfrak{u} \sim \mathfrak{u}'$, then $\mathcal{I}(\mathfrak{u}) = \mathcal{I}(\mathfrak{u}')$ and*

$$B_{\mathfrak{u}}(\mathcal{Q}(\mathfrak{u}), 100) \cap B_{\mathfrak{u}'}(\mathcal{Q}(\mathfrak{u}'), 100) \neq \emptyset.$$

Proof. Let $\mathfrak{u}, \mathfrak{u}' \in \mathfrak{U}_2(k, n, j)$ with $\mathfrak{u} \sim \mathfrak{u}'$. If $\mathfrak{u} = \mathfrak{u}'$ then the conclusion of the Lemma clearly holds. Else, there exists $\mathfrak{p} \in \mathfrak{C}_1(k, n, j)$ such that $\mathcal{I}(\mathfrak{p}) \neq \mathcal{I}(\mathfrak{u})$ and $2\mathfrak{p} \lesssim \mathfrak{u}$ and $10\mathfrak{p} \lesssim \mathfrak{u}'$. Using Lemma 5.14 and (5.73) of Lemma 5.16, we deduce that

$$100\mathfrak{p} \lesssim 100\mathfrak{u}, \quad 100\mathfrak{p} \lesssim 100\mathfrak{u}'. \quad (5.82)$$

Now suppose that $B_{\mathfrak{u}}(\mathcal{Q}(\mathfrak{u}), 100) \cap B_{\mathfrak{u}'}(\mathcal{Q}(\mathfrak{u}'), 100) = \emptyset$. Then we have $\mathfrak{B}(\mathfrak{u}) \cap \mathfrak{B}(\mathfrak{u}') = \emptyset$, by the definition (5.8) of \mathfrak{B} and the definition (2.24) of \lesssim , but also $\mathfrak{B}(\mathfrak{u}) \subset \mathfrak{B}(\mathfrak{p})$ and $\mathfrak{B}(\mathfrak{u}') \subset \mathfrak{B}(\mathfrak{p})$, by (5.8), (2.24) and (5.82). Hence,

$$|\mathfrak{B}(\mathfrak{p})| \geq |\mathfrak{B}(\mathfrak{u})| + |\mathfrak{B}(\mathfrak{u}')| \geq 2^j + 2^j = 2^{j+1},$$

which contradicts $\mathfrak{p} \in \mathfrak{C}_1(k, n, j)$. Therefore we must have

$$B_{\mathfrak{u}}(\mathcal{Q}(\mathfrak{u}), 100) \cap B_{\mathfrak{u}'}(\mathcal{Q}(\mathfrak{u}'), 100) \neq \emptyset.$$

It follows from $2\mathfrak{p} \lesssim \mathfrak{u}$ and $10\mathfrak{p} \lesssim \mathfrak{u}'$ that $\mathcal{I}(\mathfrak{p}) \subset \mathcal{I}(\mathfrak{u})$ and $\mathcal{I}(\mathfrak{p}) \subset \mathcal{I}(\mathfrak{u}')$. By (2.8), it follows that $\mathcal{I}(\mathfrak{u})$ and $\mathcal{I}(\mathfrak{u}')$ are nested. Combining this with the conclusion of the last paragraph and definition (5.14) of $\mathfrak{U}_1(k, n, j)$, we obtain that $\mathcal{I}(\mathfrak{u}) = \mathcal{I}(\mathfrak{u}')$. □

Lemma 5.27 (equivalence relation). *For each k, n, j , the relation \sim on $\mathfrak{U}_2(k, n, j)$ is an equivalence relation.*

Proof. Reflexivity holds by definition. For transitivity, suppose that

$$u, u', u'' \in \mathfrak{U}_1(k, n, j)$$

and $u \sim u'$, $u' \sim u''$. By Lemma 5.26, it follows that $\mathcal{I}(u) = \mathcal{I}(u') = \mathcal{I}(u'')$, that there exists

$$\vartheta \in B_u(\mathcal{Q}(u), 100) \cap B_{u'}(\mathcal{Q}(u'), 100)$$

and that there exists

$$\theta \in B_{u'}(\mathcal{Q}(u'), 100) \cap B_{u''}(\mathcal{Q}(u''), 100).$$

If $u = u'$, then $u \sim u''$ holds by assumption. Else, there exists by the definition of \sim some $p \in \mathfrak{T}_1(u)$ with $10p \lesssim u'$. Then we have $2p \lesssim u$ and $p \neq u$ by definition of $\mathfrak{T}_1(u)$, so $4p \lesssim 500u$ by (5.74). For $q \in B_{u''}(\mathcal{Q}(u''), 1)$ it follows by the triangle inequality that

$$\begin{aligned} d_u(\mathcal{Q}(u), q) &\leq d_u(\mathcal{Q}(u), \vartheta) + d_u(\vartheta, \mathcal{Q}(u')) \\ &\quad + d_u(\mathcal{Q}(u'), \theta) + d_u(\theta, \mathcal{Q}(u'')) + d_u(\mathcal{Q}(u''), q). \end{aligned}$$

Using (2.17) and the fact that $\mathcal{I}(u) = \mathcal{I}(u') = \mathcal{I}(u'')$ this equals

$$\begin{aligned} &d_u(\mathcal{Q}(u), \vartheta) + d_{u'}(\vartheta, \mathcal{Q}(u')) \\ &\quad + d_{u'}(\mathcal{Q}(u'), \theta) + d_{u''}(\theta, \mathcal{Q}(u'')) + d_{u''}(\mathcal{Q}(u''), q) \\ &< 100 + 100 + 100 + 100 + 1 < 500. \end{aligned}$$

Since $4p \lesssim 500u$, it follows that $d_p(\mathcal{Q}(p), q) < 4 < 10$. We have shown that $B_{u''}(\mathcal{Q}(u''), 1) \subset B_p(\mathcal{Q}(p), 10)$, combining this with $\mathcal{I}(u'') = \mathcal{I}(u)$ gives $u \sim u''$.

For symmetry suppose that $u \sim u'$. By Lemma (5.26), it follows that $\mathcal{I}(u) = \mathcal{I}(u')$ and that there exists $\vartheta \in B_u(\mathcal{Q}(u), 100) \cap B_{u'}(\mathcal{Q}(u'), 100)$. Again, for $u = u'$ symmetry is obvious. If $u \neq u'$, then there exists $p \in \mathfrak{T}_1(u')$ with $10p \lesssim u$. By definition of $\mathfrak{T}_1(u')$, Lemma 5.14 and (5.74), it follows that

$$10p \lesssim 4p \lesssim 500u'. \quad (5.83)$$

If $q \in B_u(\mathcal{Q}(u), 1)$ then we have from the triangle inequality and the fact that $\mathcal{I}(u) = \mathcal{I}(u')$:

$$\begin{aligned} d_{u'}(\mathcal{Q}(u'), q) &\leq d_{u'}(\mathcal{Q}(u'), \vartheta) + d_{u'}(\vartheta, \mathcal{Q}(u)) + d_{u'}(\mathcal{Q}(u), q) \\ &= d_{u'}(\mathcal{Q}(u'), \vartheta) + d_u(\vartheta, \mathcal{Q}(u)) + d_u(\mathcal{Q}(u), q) \\ &< 100 + 100 + 1 < 500. \end{aligned}$$

Combining this with (5.83) and (2.24), we get

$$B_u(\mathcal{Q}(u), 1) \subset B_p(\mathcal{Q}(p), 10).$$

Since $2p \lesssim u'$, we have $\mathcal{I}(p) \subset \mathcal{I}(u') = \mathcal{I}(u)$. Thus, $10p \lesssim u$ which completes the proof of $u' \sim u$. \square

Choose a set $\mathfrak{U}_3(k, n, j)$ of representatives for the equivalence classes of \sim in $\mathfrak{U}_2(k, n, j)$. Define for each $u \in \mathfrak{U}_3(k, n, j)$

$$\mathfrak{T}_2(u) := \bigcup_{u \sim u'} \mathfrak{T}_1(u') \cap \mathfrak{C}_6(k, n, j). \quad (5.84)$$

Lemma 5.28 (C6 forest). *We have*

$$\mathfrak{C}_6(k, n, j) = \bigcup_{u \in \mathfrak{U}_3(k, n, j)} \mathfrak{T}_2(u). \quad (5.85)$$

Proof. Let $\mathbf{p} \in \mathfrak{C}_6(k, n, j)$. By (5.19) and (5.23), we have $\mathbf{p} \in \mathfrak{C}_3(k, n, j)$. By (5.15) and (5.16), there exists $\mathbf{u} \in \mathfrak{U}_1(k, n, j)$ with $2\mathbf{p} \lesssim \mathbf{u}$ and $\mathcal{I}(\mathbf{p}) \neq \mathcal{I}(\mathbf{u})$, that is, with $\mathbf{p} \in \mathfrak{T}_1(\mathbf{u})$. Then $\mathfrak{T}_1(\mathbf{u})$ is clearly nonempty, so $\mathbf{u} \in \mathfrak{U}_2(k, n, j)$. By the definition of $\mathfrak{U}_3(k, n, j)$, there exists $\mathbf{u}' \in \mathfrak{U}_3(k, n, j)$ with $\mathbf{u} \sim \mathbf{u}'$. By (5.84), we have $\mathbf{p} \in \mathfrak{T}_2(\mathbf{u}')$. \square

Lemma 5.29 (C6 convex). *Let $\mathbf{u} \in \mathfrak{U}_3(k, n, j)$. If $\mathbf{p} \leq \mathbf{p}' \leq \mathbf{p}''$ and $\mathbf{p}, \mathbf{p}'' \in \mathfrak{T}_2(\mathbf{u})$, then $\mathbf{p}' \in \mathfrak{T}_2(\mathbf{u})$.*

Proof. Suppose that $\mathbf{p}, \mathbf{p}'' \in \mathfrak{T}_2(\mathbf{u})$. Then by Lemma 5.23, we have $\mathbf{p}' \in \mathfrak{C}_5(k, n, j)$. Since $\mathbf{p} \in \mathfrak{C}_6(k, n, j)$ we have $\mathcal{I}(\mathbf{p}) \not\subset G'$, hence $\mathcal{I}(\mathbf{p}') \not\subset G'$. This implies $\mathbf{p}' \in \mathfrak{C}_6(k, n, j)$. Since $\mathbf{p}'' \in \mathfrak{T}_2(\mathbf{u})$, we have $2\mathbf{p}'' \lesssim \mathbf{u}'$ and $\mathcal{I}(\mathbf{p}'') \neq \mathcal{I}(\mathbf{u}')$ for some $\mathbf{u}' \sim \mathbf{p}''$. By (5.72), we have $2\mathbf{p}' \lesssim 2\mathbf{p}''$, so by transitivity of \lesssim we have $2\mathbf{p}' \lesssim \mathbf{u}'$. Finally, $\mathcal{I}(\mathbf{p}') \subset \mathcal{I}(\mathbf{p}'')$ implies $\mathcal{I}(\mathbf{p}') \neq \mathcal{I}(\mathbf{u}')$, thus $\mathbf{p}' \in \mathfrak{T}_1(\mathbf{u}') \subset \mathfrak{T}_2(\mathbf{u})$. \square

Lemma 5.30 (forest geometry). *For each $\mathbf{u} \in \mathfrak{U}_3(k, n, j)$, the set $\mathfrak{T}_2(\mathbf{u})$ satisfies (2.32).*

Proof. Let $\mathbf{p} \in \mathfrak{T}_2(\mathbf{u})$. By (5.84), there exists $\mathbf{u}' \sim \mathbf{u}$ with $\mathbf{p} \in \mathfrak{T}_1(\mathbf{u}')$. Then we have $2\mathbf{p} \lesssim \mathbf{u}'$ and $\mathcal{I}(\mathbf{p}) \neq \mathcal{I}(\mathbf{u}')$, so by (5.74) $4\mathbf{p} \lesssim 500\mathbf{u}'$. Further, by Lemma 5.26, we have that $\mathcal{I}(\mathbf{u}') = \mathcal{I}(\mathbf{u})$ and there exists $\vartheta \in B_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), 100) \cap B_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), 100)$. Let $\theta \in B_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), 1)$. Using the triangle inequality and the fact that $\mathcal{I}(\mathbf{u}') = \mathcal{I}(\mathbf{u})$, we obtain

$$\begin{aligned} d_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), \theta) &\leq d_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), \vartheta) + d_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}), \vartheta) + d_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}), \theta) \\ &= d_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), \vartheta) + d_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), \vartheta) + d_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), \theta) \\ &< 100 + 100 + 1 < 500. \end{aligned}$$

Combining this with $4\mathbf{p} \lesssim 500\mathbf{u}'$, we obtain

$$B_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), 1) \subset B_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), 500) \subset B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 4).$$

Together with $\mathcal{I}(\mathbf{p}) \subset \mathcal{I}(\mathbf{u}') = \mathcal{I}(\mathbf{u})$, this gives $4\mathbf{p} \lesssim 1\mathbf{u}$, which is (2.32). \square

Lemma 5.31 (forest convex). *For each $\mathbf{u} \in \mathfrak{U}_3(k, n, j)$, the set $\mathfrak{T}_2(\mathbf{u})$ satisfies the convexity condition (2.33).*

Proof. Let $\mathbf{p}, \mathbf{p}'' \in \mathfrak{T}_2(\mathbf{u})$ and $\mathbf{p}' \in \mathfrak{P}$ with $\mathbf{p} \leq \mathbf{p}' \leq \mathbf{p}''$. By (5.84) we have $\mathbf{p}, \mathbf{p}'' \in \mathfrak{C}_6(k, n, j) \subset \mathfrak{C}_5(k, n, j)$. By Lemma 5.23, we have $\mathbf{p}' \in \mathfrak{C}_5(k, n, j)$. Since $\mathbf{p} \in \mathfrak{C}_6(k, n, j)$ we have $\mathcal{I}(\mathbf{p}) \not\subset G'$, so $\mathcal{I}(\mathbf{p}') \not\subset G'$ and therefore also $\mathbf{p}' \in \mathfrak{C}_6(k, n, j)$.

By (5.84) there exists $\mathbf{u}' \in \mathfrak{U}_2(k, n, j)$ with $\mathbf{p}'' \in \mathfrak{T}_1(\mathbf{u}')$ and hence $2\mathbf{p}'' \lesssim \mathbf{u}'$ and $\mathcal{I}(\mathbf{p}'') \neq \mathcal{I}(\mathbf{u}')$. Together this implies $\mathcal{I}(\mathbf{p}'') \subsetneq \mathcal{I}(\mathbf{u}')$. With the inclusion $\mathcal{I}(\mathbf{p}') \subset \mathcal{I}(\mathbf{p}'')$ from $\mathbf{p}' \leq \mathbf{p}''$, it follows that $\mathcal{I}(\mathbf{p}') \subsetneq \mathcal{I}(\mathbf{u}')$ and hence $\mathcal{I}(\mathbf{p}') \neq \mathcal{I}(\mathbf{u}')$. By (5.72) and transitivity of \lesssim we further have $2\mathbf{p}' \lesssim \mathbf{u}'$, so $\mathbf{p}' \in \mathfrak{T}_1(\mathbf{u}')$. It follows that $\mathbf{p}' \in \mathfrak{T}_2(\mathbf{u})$, which shows (2.33). \square

Lemma 5.32 (forest separation). *For each $\mathbf{u}, \mathbf{u}' \in \mathfrak{U}_3(k, n, j)$ with $\mathbf{u} \neq \mathbf{u}'$ and each $\mathbf{p} \in \mathfrak{T}_2(\mathbf{u})$ with $\mathcal{I}(\mathbf{p}) \subset \mathcal{I}(\mathbf{u}')$ we have*

$$d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{u}')) > 2^{Z(n+1)}. \quad (5.86)$$

Proof. By the definition (5.13) of $\mathfrak{C}_2(k, n, j)$, there exists a tile $\mathbf{p}' \in \mathfrak{C}_1(k, n, j)$ with $\mathbf{p}' \leq \mathbf{p}$ and $s(\mathbf{p}') = s(\mathbf{p}) - Z(n+1)$. By Lemma 2.8 we have

$$d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{u}')) \geq 2^{95aZ(n+1)} d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{u}')).$$

By (5.72) we have $2\mathbf{p}' \lesssim 2\mathbf{p}$, so by transitivity of \lesssim there exists $\mathbf{v} \sim \mathbf{u}$ with $2\mathbf{p}' \lesssim \mathbf{v}$ and $\mathcal{I}(\mathbf{p}') \neq \mathcal{I}(\mathbf{v})$. Since \mathbf{u}, \mathbf{u}' are not equivalent under \sim , we have $\mathbf{v} \not\sim \mathbf{u}'$, thus $10\mathbf{p}' \not\lesssim \mathbf{u}'$. This implies that there exists $q \in B_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), 1) \setminus B_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), 10)$.

From $\mathbf{p}' \leq \mathbf{p}$, $\mathcal{I}(\mathbf{p}') \subset \mathcal{I}(\mathbf{p}) \subset \mathcal{I}(\mathbf{u}')$ and Lemma 2.8 it then follows that

$$\begin{aligned} & d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{u}')) \\ & \geq -d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{p}')) + d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), q) - d_{\mathbf{p}'}(q, \mathcal{Q}(\mathbf{u}')) \\ & \geq -d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{p}')) + d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), q) - d_{\mathbf{u}'}(q, \mathcal{Q}(\mathbf{u}')) \\ & > -1 + 10 - 1 = 8. \end{aligned}$$

The lemma follows by combining the two displays with the fact that $95a \geq 1$. \square

Lemma 5.33 (forest inner). *For each $\mathbf{u} \in \mathfrak{U}_3(k, n, j)$ and each $\mathbf{p} \in \mathfrak{T}_2(\mathbf{u})$ we have*

$$B(c(\mathbf{p}), 8D^{s(\mathbf{p})}) \subset \mathcal{I}(\mathbf{u}). \quad (5.87)$$

Proof. Let $\mathbf{p} \in \mathfrak{T}_2(\mathbf{u})$. Let

$$\mathbf{q} \in \bigcup_{\mathbf{u}' \sim \mathbf{u}} \mathfrak{T}_1(\mathbf{u}') \cap \mathfrak{C}_3(k, n, j) \quad (5.88)$$

be a maximal element of this set with respect to \leq such that $\mathbf{p} \leq \mathbf{q}$. We show that there is no $\mathbf{q}' \in \mathfrak{C}_3(k, n, j)$ with $\mathbf{q} \leq \mathbf{q}'$ and $\mathbf{q} \neq \mathbf{q}'$. Indeed, suppose \mathbf{q}' was such a tile. By (5.16) there exists $\mathbf{u}'' \in \mathfrak{U}_1(k, n, j)$ with $2\mathbf{q}' \lesssim \mathbf{u}''$. Then we have in particular by Lemma 5.14 that $10\mathbf{p} \lesssim \mathbf{u}''$. Let $\mathbf{u}' \sim \mathbf{u}$ be such that $\mathbf{p} \in \mathfrak{T}_1(\mathbf{u}')$. By definition of \sim , we have $\mathbf{u}' \sim \mathbf{u}''$, hence $\mathbf{u} \sim \mathbf{u}''$. This implies that \mathbf{q}' is in the set in (5.88), contradicting maximality of \mathbf{q} .

Let $\mathbf{u}' \sim \mathbf{u}$ with $\mathbf{q} \in \mathfrak{T}_1(\mathbf{u}')$. By the definition (5.80) of \mathfrak{T}_1 , we have $s(\mathbf{p}) < s(\mathbf{u}')$. By Lemma 5.26, we have $s(\mathbf{u}) = s(\mathbf{u}')$, hence $s(\mathbf{q}) < s(\mathbf{u})$. By definition of $\mathfrak{C}_4(k, n, j)$, \mathbf{p} is not in any of the maximal $Z(n+1)$ layers of tiles in $\mathfrak{C}_3(k, n, j)$, and hence $s(\mathbf{p}) \leq s(\mathbf{q}) - Z(n+1) \leq s(\mathbf{u}) - Z(n+1) - 1$.

Thus, there exists some cube $I \in \mathcal{D}$ with $s(I) = s(\mathbf{u}) - Z(n+1) - 1$ and $I \subset \mathcal{I}(\mathbf{u})$ and $\mathcal{I}(\mathbf{p}) \subset I$. Since $\mathbf{p} \in \mathfrak{C}_5(k, n, j)$, we have that $I \notin \mathcal{L}(\mathbf{u})$, so $B(c(I), 8D^{s(I)}) \subset \mathcal{I}(\mathbf{u})$. By the triangle inequality, (2.1) and $a \geq 4$, the same then holds for the subcube $\mathcal{I}(\mathbf{p}) \subset I$. \square

Lemma 5.34 (forest stacking). *It holds that*

$$\sum_{\mathbf{u} \in \mathfrak{U}_3(k, n, j)} \mathbf{1}_{\mathcal{I}(\mathbf{u})} \leq 1 + (4n+12)2^n. \quad (5.89)$$

Proof. Suppose that a point x is contained in more than $1 + (4n+12)2^n$ cubes $\mathcal{I}(\mathbf{u})$ with $\mathbf{u} \in \mathfrak{U}_3(k, n, j)$. Since $\mathfrak{U}_3(k, n, j) \subset \mathfrak{C}_1(k, n, j)$ for each such \mathbf{u} , there exists $\mathbf{m} \in \mathfrak{M}(k, n)$ such that $100\mathbf{u} \lesssim \mathbf{m}$. We fix such an $\mathbf{m}(\mathbf{u}) := \mathbf{m}$ for each \mathbf{u} , and claim that the map $\mathbf{u} \mapsto \mathbf{m}(\mathbf{u})$ is injective. Indeed, assume for $\mathbf{u} \neq \mathbf{u}'$ there is $\mathbf{m} \in \mathfrak{M}(k, n)$ such that $100\mathbf{u} \lesssim \mathbf{m}$ and $100\mathbf{u}' \lesssim \mathbf{m}$. By (2.8), either $\mathcal{I}(\mathbf{u}) \subset \mathcal{I}(\mathbf{u}')$ or $\mathcal{I}(\mathbf{u}') \subset \mathcal{I}(\mathbf{u})$. By (5.14), $B_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), 100) \cap B_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), 100) =$

\emptyset . This contradicts $\Omega(\mathbf{m})$ being contained in both sets by (2.15). Thus x is contained in more than $1 + (4n + 12)2^n$ cubes $\mathcal{I}(\mathbf{m})$, $\mathbf{m} \in \mathfrak{M}(k, n)$. Consequently, we have by (5.26) that $x \in A(2n + 6, k, n) \subset G_2$. Let $\mathcal{I}(\mathbf{u})$ be an inclusion minimal cube among the $\mathcal{I}(\mathbf{u}')$, $\mathbf{u}' \in \mathfrak{U}_3(k, n, j)$ with $x \in \mathcal{I}(\mathbf{u})$. By the dyadic property (2.8), we have $\mathcal{I}(\mathbf{u}) \subset \mathcal{I}(\mathbf{u}')$ for all cubes $\mathcal{I}(\mathbf{u}')$ containing x . Thus

$$\mathcal{I}(\mathbf{u}) \subset \{y : \sum_{\mathbf{u} \in \mathfrak{U}_3(k, n, j)} \mathbf{1}_{\mathcal{I}(\mathbf{u})}(y) > 1 + (4n + 12)2^n\} \subset G_2.$$

Thus $\mathfrak{T}_1(\mathbf{u}) \cap \mathfrak{C}_6(k, n, j) = \emptyset$. This contradicts $\mathbf{u} \in \mathfrak{U}_2(k, n, j)$. \square

Proof of Lemma 5.2. We first fix k, n, j . By (2.21) and (2.20), we have that $\mathbf{1}_{\mathcal{I}(\mathbf{p})}T_{\mathbf{p}}f(x) = T_{\mathbf{p}}f(x)$ and hence $\mathbf{1}_{G \setminus G'}T_{\mathbf{p}}f(x) = 0$ for all $\mathbf{p} \in \mathfrak{C}_5(k, n, j) \setminus \mathfrak{C}_6(k, n, j)$. Thus it suffices to estimate the contribution of the sets $\mathfrak{C}_6(k, n, j)$. By Lemma 5.34, we can decompose $\mathfrak{U}_3(k, n, j)$ as a disjoint union of at most $4n + 13$ collections $\mathfrak{U}_4(k, n, j, l)$, $1 \leq l \leq 4n + 13$, each satisfying

$$\sum_{\mathbf{u} \in \mathfrak{U}_4(k, n, j, l)} \mathbf{1}_{\mathcal{I}(\mathbf{u})} \leq 2^n.$$

By Lemmas 5.30, 5.31, 5.32, 5.33 and 5.25, the pairs

$$(\mathfrak{U}_4(k, n, j, l), \mathfrak{T}_2|_{\mathfrak{U}_4(k, n, j, l)})$$

are n -forests for each k, n, j, l , and by Lemma 5.28, we have

$$\mathfrak{C}_6(k, n, j) = \bigcup_{l=1}^{4n+13} \bigcup_{\mathbf{u} \in \mathfrak{U}_4(k, n, j, l)} \mathfrak{T}_2(\mathbf{u}).$$

Since $\mathcal{I}(\mathbf{p}) \not\subset G_1$ for all $\mathbf{p} \in \mathfrak{C}_6(k, n, j)$, we have $\mathfrak{C}_6(k, n, j) \cap \mathfrak{P}_{F, G} = \emptyset$ and hence

$$\text{dens}_2\left(\bigcup_{\mathbf{u} \in \mathfrak{U}_4(k, n, j, l)} \mathfrak{T}_2(\mathbf{u})\right) \leq 2^{2a+5} \frac{\mu(F)}{\mu(G)}.$$

Using the triangle inequality according to the splitting by k, n, j and l in (5.31) and applying Proposition 2.4 to each term, we obtain the estimate

$$\sum_{k \geq 0} \sum_{n \geq k} (2n + 3)(4n + 13) 2^{432a^3} 2^{-(1-\frac{1}{q})n} \left(2^{2a+5} \frac{\mu(F)}{\mu(G)}\right)^{\frac{1}{q}-\frac{1}{2}} \|f\|_2 \|\mathbf{1}_{G \setminus G'}\|_2$$

for the left hand side of (5.31). Since $|f| \leq \mathbf{1}_F$, we have $\|f\|_2 \leq \mu(F)^{1/2}$, and we have $\|\mathbf{1}_{G \setminus G'}\|_2 \leq \mu(G)^{1/2}$. Combining this with $a \geq 4$, we estimate by

$$2^{433a^3} \mu(F)^{\frac{1}{q}} \mu(G)^{1-\frac{1}{q}} \sum_{k \geq 0} \sum_{n \geq k} n^2 2^{-(1-\frac{1}{q})n}.$$

Interchanging the order of summation, the sum equals

$$\sum_{n \geq 0} n^2 (n + 1) 2^{-\frac{q-1}{q}n} \leq \frac{2^{2a^3}}{(q-1)^3},$$

which completes the proof of the lemma. \square

5.5. Proof of Lemma 5.3, the antichains. Define $\mathfrak{P}_{X \setminus G'}$ to be the set of all $\mathfrak{p} \in \mathfrak{P}$ such that $\mathcal{I}(\mathfrak{p}) \not\subset G'$.

Lemma 5.35 (antichain decomposition). *We have that*

$$\mathfrak{P}_2 \cap \mathfrak{P}_{X \setminus G'} \quad (5.90)$$

$$= \bigcup_{k \geq 0} \bigcup_{n \geq k} \mathfrak{L}_0(k, n) \cap \mathfrak{P}_{X \setminus G'} \quad (5.91)$$

$$\cup \bigcup_{k \geq 0} \bigcup_{n \geq k} \bigcup_{0 \leq j \leq 2n+3} \mathfrak{L}_2(k, n, j) \cap \mathfrak{P}_{X \setminus G'} \quad (5.92)$$

$$\cup \bigcup_{k \geq 0} \bigcup_{n \geq k} \bigcup_{0 \leq j \leq 2n+3} \bigcup_{0 \leq l \leq Z(n+1)} \mathfrak{L}_1(k, n, j, l) \cap \mathfrak{P}_{X \setminus G'} \quad (5.93)$$

$$\cup \bigcup_{k \geq 0} \bigcup_{n \geq k} \bigcup_{0 \leq j \leq 2n+3} \bigcup_{0 \leq l \leq Z(n+1)} \mathfrak{L}_3(k, n, j, l) \cap \mathfrak{P}_{X \setminus G'}. \quad (5.94)$$

Proof. Let $\mathfrak{p} \in \mathfrak{P}_2 \cap \mathfrak{P}_{X \setminus G'}$. Clearly, for every cube $J \in \mathcal{D}$ there exists some $k \geq 0$ such that (5.1) holds, and for no cube $J \in \mathcal{D}$ and no $k < 0$ does (5.2) hold. Thus $\mathfrak{p} \in \mathfrak{P}(k)$ for some $k \geq 0$.

Next, since $E_2(\lambda, \mathfrak{p}') \subset \mathcal{I}(\mathfrak{p}') \cap G$ for every $\lambda \geq 2$ and every tile $\mathfrak{p}' \in \mathfrak{P}(k)$ with $\lambda \mathfrak{p} \lesssim \lambda \mathfrak{p}'$, it follows from (5.2) that $\mu(E_2(\lambda, \mathfrak{p}')) \leq 2^{-k} \mu(\mathcal{I}(\mathfrak{p}'))$ for every such \mathfrak{p}' , so $\text{dens}'_k(\{\mathfrak{p}\}) \leq 2^{-k}$. Combining this with $a \geq 0$, it follows from (5.7) that there exists $n \geq k$ with $\mathfrak{p} \in \mathfrak{C}(k, n)$.

Since $\mathfrak{p} \in \mathfrak{P}_{X \setminus G'}$, we have in particular $\mathcal{I}(\mathfrak{p}) \not\subset A(2n+6, k, n)$, so there exist at most $1 + (4n+12)2^n < 2^{2n+4}$ tiles $\mathfrak{m} \in \mathfrak{M}(k, n)$ with $\mathfrak{p} \leq \mathfrak{m}$. It follows that $\mathfrak{p} \in \mathfrak{L}_0(k, n)$ or $\mathfrak{p} \in \mathfrak{C}_1(k, n, j)$ for some $1 \leq j \leq 2n+3$. In the former case we are done, in the latter case the inclusion to be shown follows immediately from the definitions of the collections \mathfrak{C}_i and \mathfrak{L}_i . \square

Lemma 5.36 (L0 antichain). *We have that*

$$\mathfrak{L}_0(k, n) = \bigcup_{1 \leq l \leq n} \mathfrak{L}_0(k, n, l),$$

where each $\mathfrak{L}_0(k, n, l)$ is an antichain.

Proof. It suffices to show that $\mathfrak{L}_0(k, n)$ contains no chain of length $n+1$. Suppose that we had such a chain $\mathfrak{p}_0 \leq \mathfrak{p}_1 \leq \dots \leq \mathfrak{p}_n$ with $\mathfrak{p}_i \neq \mathfrak{p}_{i+1}$ for $i = 0, \dots, n-1$. By (5.7), we have that $\text{dens}'_k(\{\mathfrak{p}_n\}) > 2^{-n}$. Thus, by (5.6), there exists $\mathfrak{p}' \in \mathfrak{P}(k)$ and $\lambda \geq 2$ with $\lambda \mathfrak{p}_n \leq \lambda \mathfrak{p}'$ and

$$\frac{\mu(E_2(\lambda, \mathfrak{p}'))}{\mu(\mathcal{I}(\mathfrak{p}'))} > \lambda^a 2^{4a} 2^{-n}. \quad (5.95)$$

Let \mathfrak{D} be the set of all $\mathfrak{p}'' \in \mathfrak{P}(k)$ such that we have $\mathcal{I}(\mathfrak{p}'') = \mathcal{I}(\mathfrak{p}')$ and $B_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), \lambda) \cap \Omega(\mathfrak{p}'') \neq \emptyset$. We now show that

$$|\mathfrak{D}| \leq 2^{4a} \lambda^a. \quad (5.96)$$

The balls $B_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}''), 0.2)$, $\mathfrak{p}'' \in \mathfrak{D}$ are disjoint by (2.15), and by the triangle inequality contained in $B_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), \lambda+1)$. By assumption (1.11) on Θ , this ball can be covered with

$$2^{a(\lceil \log_2(\lambda+1) \rceil + 2)} \leq 2^{a(\log_2(\lambda)+4)} = 2^{4a} \lambda^a$$

many $d_{\mathbf{p}'}$ -balls of radius $1/4$. By the triangle inequality, each such ball contains at most one $\mathcal{Q}(\mathbf{p}'')$, and each $\mathcal{Q}(\mathbf{p}'')$ is contained in one of the balls. Thus we get (5.96).

By (2.26) and (2.27) we have $E_2(\lambda, \mathbf{p}') \subset \bigcup_{\mathbf{p}'' \in \mathfrak{D}} E_1(\mathbf{p}'')$, thus

$$2^{4a} \lambda^a 2^{-n} < \sum_{\mathbf{p}'' \in \mathfrak{D}} \frac{\mu(E_1(\mathbf{p}''))}{\mu(\mathcal{I}(\mathbf{p}''))}.$$

Hence there exists a tile $\mathbf{p}'' \in \mathfrak{D}$ with

$$\mu(E_1(\mathbf{p}'')) \geq 2^{-n} \mu(\mathcal{I}(\mathbf{p}')).$$

By the definition (5.5) of $\mathfrak{M}(k, n)$, there exists a tile $\mathbf{m} \in \mathfrak{M}(k, n)$ with $\mathbf{p}' \leq \mathbf{m}$. From (5.95), the inclusion $E_2(\lambda, \mathbf{p}') \subset \mathcal{I}(\mathbf{p}')$ and $a \geq 1$ we obtain

$$2^n \geq 2^{4a} \lambda^a \geq \lambda.$$

From the triangle inequality, Lemma 2.8 and $a \geq 1$, we now obtain for all $\vartheta \in B_{\mathbf{m}}(\mathcal{Q}(\mathbf{m}), 1)$ that

$$\begin{aligned} & d_{\mathbf{p}_0}(\mathcal{Q}(\mathbf{p}_0), \vartheta) \\ & \leq d_{\mathbf{p}_0}(\mathcal{Q}(\mathbf{p}_0), \mathcal{Q}(\mathbf{p}_n)) + d_{\mathbf{p}_0}(\mathcal{Q}(\mathbf{p}_n), \mathcal{Q}(\mathbf{p}')) + d_{\mathbf{p}_0}(\mathcal{Q}(\mathbf{p}'), \mathcal{Q}(\mathbf{p}'')) \\ & \quad + d_{\mathbf{p}_0}(\mathcal{Q}(\mathbf{p}''), \mathcal{Q}(\mathbf{m})) + d_{\mathbf{p}_0}(\mathcal{Q}(\mathbf{m}), \vartheta) \\ & \leq 1 + 2^{-95an} (d_{\mathbf{p}_n}(\mathcal{Q}(\mathbf{p}_n), \mathcal{Q}(\mathbf{p}')) + d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), \mathcal{Q}(\mathbf{p}'')) \\ & \quad + d_{\mathbf{p}''}(\mathcal{Q}(\mathbf{p}''), \mathcal{Q}(\mathbf{m})) + d_{\mathbf{m}}(\mathcal{Q}(\mathbf{m}), \vartheta)) \\ & \leq 1 + 2^{-95an} (\lambda + (\lambda + 1) + 1 + 1) \leq 100. \end{aligned}$$

Thus, by (2.23), $100\mathbf{p}_0 \lesssim \mathbf{m}$, a contradiction to $\mathbf{p}_0 \notin \mathfrak{C}(k, n)$. \square

Lemma 5.37 (L2 antichain). *Each of the sets $\mathfrak{L}_2(k, n, j)$ is an antichain.*

Proof. Suppose that there are $\mathbf{p}_0, \mathbf{p}_1 \in \mathfrak{L}_2(k, n, j)$ with $\mathbf{p}_0 \neq \mathbf{p}_1$ and $\mathbf{p}_0 \leq \mathbf{p}_1$. By Lemma 5.14 and Lemma 5.15, it follows that $2\mathbf{p}_0 \lesssim 200\mathbf{p}_1$. Since $\mathfrak{L}_2(k, n, j)$ is finite, there exists a maximal $l \geq 1$ such that there exists a chain $2\mathbf{p}_0 \lesssim 200\mathbf{p}_1 \lesssim \dots \lesssim 200\mathbf{p}_l$ with $\mathbf{p}_i \neq \mathbf{p}_{i+1}$ for $i = 0, \dots, l-1$. If we have $\mathbf{p}_l \in \mathfrak{U}_1(k, n, j)$, then it follows from $2\mathbf{p}_0 \lesssim 200\mathbf{p}_l \lesssim \mathbf{p}_l$ and (5.15) that $\mathbf{p}_0 \notin \mathfrak{L}_2(k, n, j)$, a contradiction. Thus, by the definition (5.14) of $\mathfrak{U}_1(k, n, j)$, there exists $\mathbf{p}_{l+1} \in \mathfrak{C}_1(k, n, j)$ with $\mathcal{I}(\mathbf{p}_l) \subsetneq \mathcal{I}(\mathbf{p}_{l+1})$ and $\vartheta \in B_{\mathbf{p}_l}(\mathcal{Q}(\mathbf{p}_l), 100) \cap B_{\mathbf{p}_{l+1}}(\mathcal{Q}(\mathbf{p}_{l+1}), 100)$. Using the triangle inequality and Lemma 2.8, one deduces that $200\mathbf{p}_l \lesssim 200\mathbf{p}_{l+1}$. This contradicts maximality of l . \square

Lemma 5.38 (L1 L3 antichain). *Each of the sets $\mathfrak{L}_1(k, n, j, l)$ and $\mathfrak{L}_3(k, n, j, l)$ is an antichain.*

Proof. By its definition (5.11), each set $\mathfrak{L}_1(k, n, j, l)$ is a set of minimal elements in some set of tiles with respect to \leq . If there were distinct $\mathbf{p}, \mathbf{q} \in \mathfrak{L}_1(k, n, j, l)$ with $\mathbf{p} \leq \mathbf{q}$, then \mathbf{q} would not be minimal. Hence such \mathbf{p}, \mathbf{q} do not exist. Similarly, by (5.17), each set $\mathfrak{L}_3(k, n, j, l)$ is a set of maximal elements in some set of tiles with respect to \leq . If there were distinct $\mathbf{p}, \mathbf{q} \in \mathfrak{L}_3(k, n, j, l)$ with $\mathbf{p} \leq \mathbf{q}$, then \mathbf{p} would not be maximal. \square

Proof of Lemma 5.3. If $\mathfrak{p} \notin \mathfrak{P}_{X \setminus G'}$, then $\mathcal{I}(\mathfrak{p}) \subset G'$. By (2.21) and (2.26), it follows that $\mathbf{1}_{G \setminus G'} T_{\mathfrak{p}} f(x) = 0$. We thus have

$$\mathbf{1}_{G \setminus G'} \sum_{\mathfrak{p} \in \mathfrak{P}_2} T_{\mathfrak{p}} f(x) = \mathbf{1}_{G \setminus G'} \sum_{\mathfrak{p} \in \mathfrak{P}_2 \cap \mathfrak{P}_{X \setminus G'}} T_{\mathfrak{p}} f(x).$$

Let $\mathfrak{L}(k, n)$ denote any of the terms $\mathfrak{L}_i(k, n, j, l) \cap \mathfrak{P}_{X \setminus \mathfrak{P}_2}$ on the right hand side of (5.90), where the indices j, l may be void. Then $\mathfrak{L}(k, n)$ is an antichain, by Lemmas 5.36, 5.37, 5.38. Further, we have

$$\text{dens}_1(\mathfrak{L}(k, n)) \leq 2^{4a+1-n}$$

by Lemma 5.25, and we have

$$\text{dens}_2(\mathfrak{L}(k, n)) \leq 2^{2a+5} \frac{\mu(F)}{\mu(G)},$$

since

$$\mathfrak{L}(k, n) \cap \mathfrak{P}_{F, G} \subset \mathfrak{P}_{X \setminus \mathfrak{P}_2} \cap \mathfrak{P}_{F, G} = \emptyset.$$

Applying now the triangle inequality according to the decomposition in Lemma 5.35, and then applying Proposition 2.3 to each term, we obtain the estimate

$$\begin{aligned} &\leq \sum_{k \geq 0} \sum_{n \geq k} (n + (2n + 4) + 2(2n + 4)Z(n + 1)) \\ &\quad \times 2^{201a^3} (q - 1)^{-1} (2^{4a+1-n})^{\frac{q-1}{8a^4}} (2^{2a+5} \frac{\mu(F)}{\mu(G)})^{\frac{1}{q} - \frac{1}{2}} \|f\|_2 \|\mathbf{1}_{G \setminus G'}\|_2. \end{aligned}$$

Because $|f| \leq \mathbf{1}_F$, we have $\|f\|_2 \leq \mu(F)^{1/2}$, and we have $\|\mathbf{1}_{G \setminus G'}\|_2 \leq \mu(G)^{1/2}$. Using this and (2.3), we bound

$$\leq 2^{202a^3} (q - 1)^{-1} \mu(F)^{\frac{1}{q}} \mu(G)^{\frac{1}{q}} \sum_{k \geq 0} \sum_{n \geq k} n^2 2^{-n \frac{q-1}{8a^4}}.$$

The last sum equals, by changing the order of summation,

$$\sum_{n \geq 0} n^2 (n + 1) 2^{-n \frac{q-1}{8a^4}} \leq \frac{2^{8a^3}}{(q - 1)^3}.$$

This completes the proof. \square

6. PROOF OF PROPOSITION 2.3

Let an antichain \mathfrak{A} and functions f, g as in Proposition 2.3 be given. We prove (2.30) in Subsection 6.1 as the geometric mean of two inequalities, each involving one of the two densities. One of these two inequalities will need a careful estimate formulated in Lemma 6.5 of the TT^* correlation between two tile operators. Lemma 6.5 will be proven in Subsection 6.2. The summation of the contributions of these individual correlations will require a geometric Lemma 6.6 counting the relevant tile pairs. Lemma 6.6 will be proven in Subsection 6.3.

6.1. The density arguments. We begin with the following crucial disjointedness property of the sets $E(\mathfrak{p})$ with $\mathfrak{p} \in \mathfrak{A}$.

Lemma 6.1 (tile disjointness). *Let $\mathfrak{p}, \mathfrak{p}' \in \mathfrak{A}$. If there exists an $x \in X$ with $x \in E(\mathfrak{p}) \cap E(\mathfrak{p}')$, then $\mathfrak{p} = \mathfrak{p}'$.*

Proof. Let $\mathfrak{p}, \mathfrak{p}'$ and x be given. Assume without loss of generality that $s(\mathfrak{p}) \leq s(\mathfrak{p}')$. As we have $x \in E(\mathfrak{p}) \subset \mathcal{I}(\mathfrak{p})$ and $x \in E(\mathfrak{p}') \subset \mathcal{I}(\mathfrak{p}')$ by Definition (2.20), we conclude for $i = 1, 2$ that $Q(x) \in \Omega(\mathfrak{p})$ and $Q(x) \in \Omega(\mathfrak{p}')$. By (2.14) we have $\Omega(\mathfrak{p}') \subset \Omega(\mathfrak{p})$. By Definition (2.23), we conclude $\mathfrak{p} \leq \mathfrak{p}'$. As \mathfrak{A} is an antichain, we conclude $\mathfrak{p} = \mathfrak{p}'$. This proves the lemma. \square

Let \mathcal{B} be the collection of balls

$$B(c(\mathfrak{p}), 8D^{s(\mathfrak{p})}) \quad (6.1)$$

with $\mathfrak{p} \in \mathfrak{A}$ and recall the definition of $M_{\mathcal{B}}$ from Definition 2.41.

Lemma 6.2 (maximal bound antichain). *Let $x \in X$. Then*

$$|\sum_{\mathfrak{p} \in \mathfrak{A}} T_{\mathfrak{p}} f(x)| \leq 2^{107a^3} M_{\mathcal{B}} f(x). \quad (6.2)$$

Proof. Fix $x \in X$. By Lemma 6.1, there is at most one $\mathfrak{p} \in \mathfrak{A}$ such that $T_{\mathfrak{p}} f(x)$ is not zero. If there is no such \mathfrak{p} , the estimate (6.2) follows.

Assume there is such a \mathfrak{p} . By definition of $T_{\mathfrak{p}}$ we have $x \in E(\mathfrak{p}) \subset \mathcal{I}(\mathfrak{p})$ and by the squeezing property (2.10)

$$\rho(x, c(\mathfrak{p})) \leq 4D^{s(\mathfrak{p})}. \quad (6.3)$$

Let $y \in X$ with $K_{s(\mathfrak{p})}(x, y) \neq 0$. By Definition (2.5) of $K_{s(\mathfrak{p})}$ we have

$$\frac{1}{4} D^{s(\mathfrak{p})-1} \leq \rho(x, y) \leq \frac{1}{2} D^{s(\mathfrak{p})}. \quad (6.4)$$

The triangle inequality with (6.3) and (6.4) implies

$$\rho(c(\mathfrak{p}), y) \leq 8D^{s(\mathfrak{p})}. \quad (6.5)$$

Using the kernel bound (1.14) and the lower bound in (2.48) we obtain

$$|K_{s(\mathfrak{p})}(x, y)| \leq \frac{2^{a^3}}{\mu(B(x, \frac{1}{4} D^{s(\mathfrak{p})-1}))}. \quad (6.6)$$

Using $D = 2^{100a^2}$ and the doubling property (1.5) $5 + 100a^2$ times estimates the last display by

$$\leq \frac{2^{5a+101a^3}}{\mu(B(x, 8D^{s(\mathfrak{p})}))}. \quad (6.7)$$

Using that $|e(\vartheta)|$ is bounded by 1 for every $\vartheta \in \Theta$, we estimate with the triangle inequality and the above information

$$|T_{\mathfrak{p}} f(x)| \leq \frac{2^{5a+101a^3}}{\mu(B(x, 8D^{s(\mathfrak{p})}))} \int_{\mu(B(x, 8D^{s(\mathfrak{p})}))} |f(y)| dy \quad (6.8)$$

This together with $a \geq 1$ proves the Lemma. \square

Set

$$\tilde{q} = \frac{2q}{1+q}. \quad (6.9)$$

Since $1 < q \leq 2$, we have $1 < \tilde{q} < q \leq 2$.

Lemma 6.3 (dens2 antichain). *We have that*

$$\left| \int \overline{g(x)} \sum_{p \in \mathfrak{A}} T_p f(x) d\mu(x) \right| \leq 2^{111a^2} (q-1)^{-1} \text{dens}_2(\mathfrak{A})^{\frac{1}{q}-\frac{1}{2}} \|f\|_2 \|g\|_2. \quad (6.10)$$

Proof. We have $f = \mathbf{1}_F f$. Using Hölder's inequality, we obtain for each $x \in B'$ and each $B' \in \mathcal{B}$ using $1 < \tilde{q} \leq 2$

$$\frac{1}{\mu(B')} \int_{B'} |f(y)| d\mu(y) \quad (6.11)$$

$$\leq \left(\frac{1}{\mu(B')} \int_{B'} |f(y)|^{\frac{2\tilde{q}}{3\tilde{q}-2}} d\mu(y) \right)^{\frac{3}{2}-\frac{1}{\tilde{q}}} \left(\frac{1}{\mu(B')} \int_{B'} \mathbf{1}_F(y) d\mu(y) \right)^{\frac{1}{\tilde{q}}-\frac{1}{2}} \quad (6.12)$$

$$\leq \left(M_{\mathcal{B}}(|f|^{\frac{2\tilde{q}}{3\tilde{q}-2}})(x) \right)^{\frac{3}{2}-\frac{1}{\tilde{q}}} \text{dens}_2(\mathfrak{A})^{\frac{1}{q}-\frac{1}{2}}. \quad (6.13)$$

Taking the maximum over all B' containing x , we obtain

$$M_{\mathcal{B}}|f| \leq M_{\mathcal{B}, \frac{2\tilde{q}}{3\tilde{q}-2}}|f| \text{dens}_2(\mathfrak{A})^{\frac{1}{q}-\frac{1}{2}}. \quad (6.14)$$

We have with Proposition 2.6

$$\left\| M_{\mathcal{B}, \frac{2\tilde{q}}{3\tilde{q}-2}} f \right\|_2 \leq 2^{2a} (3\tilde{q}-2)(2\tilde{q}-2)^{-1} \|f\|_2. \quad (6.15)$$

Using $1 < \tilde{q} \leq 2$ estimates the last display by

$$2^{2a+2} (\tilde{q}-1)^{-1} \|f\|_2. \quad (6.16)$$

We obtain with Cauchy-Schwarz and then Lemma 6.2

$$\left| \int \overline{g(x)} \sum_{p \in \mathfrak{A}} T_p f(x) d\mu(x) \right| \quad (6.17)$$

$$\leq \|g\|_2 \left\| \sum_{p \in \mathfrak{A}} T_p f \right\|_2 \quad (6.18)$$

$$\leq 2^{107a^2} \|g\|_2 \|M_{\mathcal{B}} f\|_2 \quad (6.19)$$

With (6.14) and (6.16) we can estimate the last display by

$$\leq 2^{107a^2+2a+2} (\tilde{q}-1)^{-1} \|g\|_2 \|f\|_2 \text{dens}_2(\mathfrak{A})^{\frac{1}{q}-\frac{1}{2}} \quad (6.20)$$

Using $a \geq 4$ and $(\tilde{q}-1)^{-1} = (q+1)/(q-1) \leq 3(q-1)^{-1}$ proves the lemma. \square

Lemma 6.4 (dens1 antichain). *Set $p := 4a^4$. We have*

$$\left| \int \overline{g(x)} \sum_{p \in \mathfrak{A}} T_p f(x) d\mu(x) \right| \leq 2^{150a^3} \text{dens}_1(\mathfrak{A})^{\frac{1}{2p}} \|f\|_2 \|g\|_2. \quad (6.21)$$

Proof. We write for the expression inside the absolute values on the left-hand side of (6.21)

$$\sum_{\mathfrak{p} \in \mathfrak{A}} \iint \overline{g(x)} \mathbf{1}_{E(\mathfrak{p})}(x) K_{s(\mathfrak{p})}(x, y) e(Q(x)(y) - Q(x)(x)) f(y) d\mu(y) d\mu(x) \quad (6.22)$$

$$= \int \sum_{\mathfrak{p} \in \mathfrak{A}} \overline{T_{\mathfrak{p}}^* g(y)} f(y) d\mu(y) \quad (6.23)$$

with the adjoint operator

$$T_{\mathfrak{p}}^* g(y) = \int_{E(\mathfrak{p})} \overline{K_{s(\mathfrak{p})}(x, y)} e(-Q(x)(y) + Q(x)(x)) g(x) d\mu(x). \quad (6.24)$$

We have by expanding the square

$$\int \left| \sum_{\mathfrak{p} \in \mathfrak{A}} T_{\mathfrak{p}}^* g(y) \right|^2 d\mu(y) = \int \left(\sum_{\mathfrak{p} \in \mathfrak{A}} T_{\mathfrak{p}}^* g(y) \right) \left(\sum_{\mathfrak{p}' \in \mathfrak{A}} \overline{T_{\mathfrak{p}'}^* g(y)} \right) d\mu(y) \quad (6.25)$$

$$\leq \sum_{\mathfrak{p} \in \mathfrak{A}} \sum_{\mathfrak{p}' \in \mathfrak{A}} \left| \int T_{\mathfrak{p}}^* g(y) \overline{T_{\mathfrak{p}'}^* g(y)} d\mu(y) \right|. \quad (6.26)$$

We split the sum into the terms with $s(\mathfrak{p}') \leq s(\mathfrak{p})$ and $s(\mathfrak{p}) < s(\mathfrak{p}')$. Using the symmetry of each summand, we may switch \mathfrak{p} and \mathfrak{p}' in the second sum. Using further positivity of each summand to replace the condition $s(\mathfrak{p}') < s(\mathfrak{p})$ by $s(\mathfrak{p}') \leq s(\mathfrak{p})$ in the second sum, we estimate (6.26) by

$$\leq 2 \sum_{\mathfrak{p} \in \mathfrak{A}} \sum_{\mathfrak{p}' \in \mathfrak{A}: s(\mathfrak{p}') \leq s(\mathfrak{p})} \left| \int T_{\mathfrak{p}}^* g(y) \overline{T_{\mathfrak{p}'}^* g(y)} d\mu(y) \right|. \quad (6.27)$$

The following basic TT^* estimate will be proved in Subsection 6.2.

Lemma 6.5 (tile correlation). *Let $\mathfrak{p}, \mathfrak{p}' \in \mathfrak{P}$ with $s(\mathfrak{p}') \leq s(\mathfrak{p})$. Then*

$$\left| \int T_{\mathfrak{p}'}^* g \overline{T_{\mathfrak{p}}^* g} \right| \quad (6.28)$$

$$\leq 2^{255a^3} \frac{(1 + d_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), \mathcal{Q}(\mathfrak{p})))^{-1/(2a^2+a^3)}}{\mu(\mathcal{I}(\mathfrak{p}))} \int_{E(\mathfrak{p}')} |g| \int_{E(\mathfrak{p})} |g|. \quad (6.29)$$

Moreover, the term (6.28) vanishes unless

$$\mathcal{I}(\mathfrak{p}') \subset B(c(\mathfrak{p}), 15D^{s(\mathfrak{p})}). \quad (6.30)$$

Define for $\mathfrak{p} \in \mathfrak{P}$

$$B(\mathfrak{p}) := B(c(\mathfrak{p}), 15D^{s(\mathfrak{p})}) \quad (6.31)$$

and define

$$\mathfrak{A}(\mathfrak{p}) := \{\mathfrak{p}' \in \mathfrak{A} : s(\mathfrak{p}') \leq s(\mathfrak{p}) \wedge \mathcal{I}(\mathfrak{p}') \subset B(\mathfrak{p})\}. \quad (6.32)$$

Note that by the squeezing property (2.10) and the doubling property (1.5) applied 6 times we have

$$\mu(B(\mathfrak{p})) \leq 2^{6a} \mu(\mathcal{I}(\mathfrak{p})). \quad (6.33)$$

Using Lemma 6.5 and (6.33), we estimate (6.27) by

$$\leq 2^{255a^3+6a+1} \sum_{\mathbf{p} \in \mathfrak{A}} \int_{E(\mathbf{p})} |g|(y) h(\mathbf{p}) d\mu(y) \quad (6.34)$$

with $h(\mathbf{p})$ defined as

$$\frac{1}{\mu(B(\mathbf{p}))} \int \sum_{\mathbf{p}' \in \mathfrak{A}(\mathbf{p})} (1 + d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), \mathcal{Q}(\mathbf{p})))^{-1/(2a^2+a^3)} (\mathbf{1}_{E(\mathbf{p}')} |g|)(y') d\mu(y'). \quad (6.35)$$

The following lemma will be proved in Subsection 6.3.

Lemma 6.6 (antichain tile count). *Set $p := 4a^4$ and let p' be the dual exponent of p , that is $1/p + 1/p' = 1$. For every $\vartheta \in \Theta$ and every subset \mathfrak{A}' of \mathfrak{A} we have*

$$\left\| \sum_{\mathbf{p} \in \mathfrak{A}'} (1 + d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \vartheta))^{-1/(2a^2+a^3)} \mathbf{1}_{E(\mathbf{p})} \mathbf{1}_G \right\|_p \quad (6.36)$$

$$\leq 2^{104a} \text{dens}_1(\mathfrak{A})^{\frac{1}{p}} \mu(\cup_{\mathbf{p} \in \mathfrak{A}'} I_{\mathbf{p}})^{\frac{1}{p}}. \quad (6.37)$$

Note that $p \geq 4$ since $a > 4$. We estimate $h(\mathbf{p})$ as defined in (6.35) with Hölder using $|g| \leq \mathbf{1}_G$ and $E(\mathbf{p}') \subset B(\mathbf{p})$ by

$$\frac{\|g \mathbf{1}_{B(\mathbf{p})}\|_{p'}}{\mu(B(\mathbf{p}))} \left\| \sum_{\mathbf{p} \in \mathfrak{A}(\mathbf{p})} (1 + d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \Omega(\mathbf{p}'))^{-1/(2a^2+a^3)} \mathbf{1}_{E(\mathbf{p}')} \mathbf{1}_G \right\|_p. \quad (6.38)$$

Then we apply Lemma 6.6 to estimate this by

$$\leq 2^{104a} \frac{\|g \mathbf{1}_{B(\mathbf{p})}\|_{p'}}{\mu(B(\mathbf{p}))} \text{dens}_1(\mathfrak{A})^{\frac{1}{p}} \mu(B(\mathbf{p}))^{\frac{1}{p}}. \quad (6.39)$$

Let \mathcal{B}' be the collection of all balls $B(\mathbf{p})$ with $\mathbf{p} \in \mathfrak{A}$. Then for each $\mathbf{p} \in \mathfrak{A}$ and $x \in B(\mathbf{p})$ we have by definition (2.41) of $M_{\mathcal{B}', p'}$

$$\|g \mathbf{1}_{B(\mathbf{p})}\|_{p'} \leq \mu(B(\mathbf{p}))^{\frac{1}{p'}} M_{\mathcal{B}', p'} g(x). \quad (6.40)$$

Hence we can estimate (6.39) by

$$\leq 2^{104a} (M_{\mathcal{B}', p'} g(x)) \text{dens}_1(\mathfrak{A})^{\frac{1}{p}}. \quad (6.41)$$

With this estimate of $h(\mathbf{p})$, using $E(\mathbf{p}) \subset B(\mathbf{p})$ by construction of $B(\mathbf{p})$, we estimate (6.34) by

$$\leq 2^{255a^3+110a+1} \text{dens}_1(\mathfrak{A})^{\frac{1}{p}} \sum_{\mathbf{p} \in \mathfrak{A}} \int_{E(\mathbf{p})} |g|(y) M_{\mathcal{B}', p'} g(y) dy. \quad (6.42)$$

Using Lemma 6.1, the last display is observed to be

$$= 2^{255a^3+110a+1} \text{dens}_1(\mathfrak{A})^{\frac{1}{p}} \int |g|(y) (M_{\mathcal{B}', p'} g)(y) dy. \quad (6.43)$$

Applying Cauchy-Schwarz and using Proposition 2.6 estimates the last display by

$$2^{255a^3+110a+1} \text{dens}_1(\mathfrak{A})^{\frac{1}{p}} \|g\|_2 \|M_{\mathcal{B}', p'} g\|_2 \quad (6.44)$$

$$\leq 2^{255a^3+110a+3} \frac{2}{2-p'} \text{dens}_1(\mathfrak{A})^{\frac{1}{p}} \|g\|_2^2. \quad (6.45)$$

Using $p > 4$ and thus $1 < p' < \frac{4}{3}$, we estimate the last display by

$$\leq 2^{255a^3+110a+5} \text{dens}_1(\mathfrak{A})^{\frac{1}{p}} \|g\|_2^2. \quad (6.46)$$

Now Lemma 6.4 follows by applying Cauchy-Schwarz on the left-hand side and using $a \geq 4$. \square

We have

$$\left(\frac{1}{\tilde{q}} - \frac{1}{2}\right)(2-q) = \frac{1}{q} - \frac{1}{2}. \quad (6.47)$$

Multiplying the $(2-q)$ -th power of (6.10) and the $(q-1)$ -th power of (6.21) and estimating gives after simplification of some factors

$$\left| \int \overline{g(x)} \sum_{\mathfrak{p} \in \mathfrak{A}} T_{\mathfrak{p}} f(x) d\mu(x) \right| \quad (6.48)$$

$$\leq 2^{150a^3} (q-1)^{-1} \text{dens}_1(\mathfrak{A})^{\frac{q-1}{2p}} \text{dens}_2(\mathfrak{A})^{\frac{1}{q}-\frac{1}{2}} \|f\|_2 \|g\|_2. \quad (6.49)$$

With the definition of p , this implies Proposition 2.3.

6.2. Proof of Lemma 6.5, the tile correlation bound. The next lemma prepares an application of Proposition 2.5.

Lemma 6.7 (correlation kernel bound). *Let $-S \leq s_1 \leq s_2 \leq S$ and let $x_1, x_2 \in X$. Define*

$$\varphi(y) := \overline{K_{s_1}(x_1, y)} K_{s_2}(x_2, y). \quad (6.50)$$

If $\varphi(y) \neq 0$, then

$$y \in B(x_1, D^{s_1}). \quad (6.51)$$

Moreover, we have with $\tau = 1/a$

$$\|\varphi\|_{C^\tau(B(x_1, D^{s_1}))} \leq \frac{2^{254a^3}}{\mu(B(x_1, D^{s_1}))\mu(B(x_2, D^{s_2}))}. \quad (6.52)$$

Proof. If $\varphi(y)$ is not zero, then $K_{s_1}(x_1, y)$ is not zero and thus (2.48) gives (6.51).

We next have for y with (2.49)

$$|\varphi(y)| \leq \frac{2^{204a^3}}{\mu(B(x_1, D^{s_1}))\mu(B(x_2, D^{s_2}))} \quad (6.53)$$

and for $y' \neq y$ additionally with (2.50)

$$|\varphi(y) - \varphi(y')| \quad (6.54)$$

$$\leq |K_{s_1}(x_1, y) - K_{s_1}(x_1, y')| |K_{s_2}(x_2, y)| \quad (6.55)$$

$$+ |K_{s_1}(x_1, y')| |K_{s_2}(x_2, y) - K_{s_2}(x_2, y')| \quad (6.56)$$

$$\leq \frac{2^{252a^3}}{\mu(B(x_1, D^{s_1}))\mu(B(x_2, D^{s_2}))} \left(\left(\frac{\rho(y, y')}{D^{s_1}} \right)^{1/a} + \left(\frac{\rho(y, y')}{D^{s_2}} \right)^{1/a} \right) \quad (6.57)$$

$$\leq \frac{2^{253a^3}}{\mu(B(x_1, D^{s_1}))\mu(B(x_2, D^{s_2}))} \left(\frac{\rho(y, y')}{D^{s_1}} \right)^{1/a}. \quad (6.58)$$

Adding the estimates (6.53) and (6.58) gives (6.52). This proves the lemma. \square

The next lemma is a geometric estimate for two tiles.

Lemma 6.8. *Let $\mathfrak{p}_1, \mathfrak{p}_2 \in \mathfrak{P}$ with $s(\mathfrak{p}_1) \leq s(\mathfrak{p}_2)$. For each $x_1 \in E(\mathfrak{p}_1)$ and $x_2 \in E(\mathfrak{p}_2)$ we have*

$$1 + d_{\mathfrak{p}_1}(\mathcal{Q}(\mathfrak{p}_1), \mathcal{Q}(\mathfrak{p}_2)) \leq 2^{3a}(1 + d_{B(x_1, D^{s(\mathfrak{p}_1)})}(Q(x_1), Q(x_2))). \quad (6.59)$$

Proof. Let $i \in \{1, 2\}$. By Definition (2.20) of E , we have $Q(x_i) \in \Omega(\mathfrak{p}_i)$. With (2.15) we then conclude

$$d_{\mathfrak{p}_i}(Q(x_i), \mathcal{Q}(\mathfrak{p}_i)) \leq 1. \quad (6.60)$$

We have $\mathcal{I}(\mathfrak{p}_1) \subset \mathcal{I}(\mathfrak{p}_2)$ by (2.8). Using Lemma 2.8 it follows that

$$d_{\mathfrak{p}_1}(Q(x_2), \mathcal{Q}(\mathfrak{p}_2)) \leq 1. \quad (6.61)$$

By the triangle inequality, we obtain from (6.60) and (6.61)

$$1 + d_{\mathfrak{p}_1}(\mathcal{Q}(\mathfrak{p}_1), \mathcal{Q}(\mathfrak{p}_2)) \leq 3 + d_{\mathfrak{p}_1}(Q(x_1), Q(x_2)). \quad (6.62)$$

As $x_1 \in \mathcal{I}(\mathfrak{p}_1)$ by Definition (2.20) of E , we have by the squeezing property (2.10)

$$d(x_1, c(\mathfrak{p}_1)) \leq 4D^{s(\mathfrak{p}_1)} \quad (6.63)$$

and thus by (2.10) again and the triangle inequality

$$\mathcal{I}(\mathfrak{p}_1) \subset B(x_1, 8D^{s(\mathfrak{p}_1)}). \quad (6.64)$$

We thus estimate the right-hand side of (6.62) with monotonicity (1.9) of the metrics d_B by

$$\leq 3 + d_{B(x_1, 8D^{s(\mathfrak{p}_1)})}(Q(x_1), Q(x_2)). \quad (6.65)$$

This is further estimated by applying the doubling property (1.8) three times by

$$\leq 3 + 2^{3a}d_{B_1(x_1, D^{s(\mathfrak{p}_1)})}(Q(x_1), Q(x_2)). \quad (6.66)$$

Now (6.59) follows with $a \geq 1$. \square

Lemma 6.9 (tile range support). *For each $\mathfrak{p} \in \mathfrak{P}$, and each $y \in X$, we have that*

$$T_{\mathfrak{p}}^*g(y) \neq 0 \quad (6.67)$$

implies

$$y \in B(c(\mathfrak{p}), 5D^{s(\mathfrak{p})}). \quad (6.68)$$

Proof. Fix \mathfrak{p} and y with (6.67). Then there exists $x \in E(\mathfrak{p})$ with

$$\overline{K_{s(\mathfrak{p})}(x, y)}e(-Q(x)(y) + Q(x)(x))g(x) \neq 0. \quad (6.69)$$

As $E(\mathfrak{p}) \subset \mathcal{I}(\mathfrak{p})$ and by the squeezing property (2.10), we have

$$\rho(x, c(\mathfrak{p})) \leq 4D^{s(\mathfrak{p})}. \quad (6.70)$$

As $K_{s(\mathfrak{p})}(x, y) \neq 0$, we have by (2.48) that

$$\rho(x, y) \leq \frac{1}{2}D^{s(\mathfrak{p})}. \quad (6.71)$$

Now (6.68) follows by the triangle inequality. \square

We now prove Lemma 6.5. We begin with (6.28).

We expand the left-hand side of (6.28) as

$$\left| \int \int_{E(\mathfrak{p}_1)} \overline{K_{s(\mathfrak{p}_1)}(x_1, y)} e(-Q(x_1)(y) + Q(x_1)(x_1)) g(x_1) d\mu(x_1) \right. \quad (6.72)$$

$$\left. \times \int_{E(\mathfrak{p}_2)} K_{s(\mathfrak{p}_2)}(x_2, y) e(Q(x_2)(y) - Q(x_2)(x_2)) \overline{g(x_2)} d\mu(x_2) d\mu(y) \right|. \quad (6.73)$$

By Fubini and the triangle inequality and the fact $|e(Q(x_i)(x_i))| = 1$ for $i = 1, 2$, we can estimate (6.72) from above by

$$\int_{E(\mathfrak{p}_1)} \int_{E(\mathfrak{p}_2)} \mathbf{I}(x_1, x_2) d\mu(x_1) d\mu(x_2). \quad (6.74)$$

with

$$\mathbf{I}(x_1, x_2) := \left| \int e(-Q(x_1)(y) + Q(x_2)(y)) \varphi_{x_1, x_2}(y) d\mu(y) g(x_1) g(x_2) \right| \quad (6.75)$$

We estimate for fixed $x_1 \in E(\mathfrak{p}_1)$ and $x_2 \in E(\mathfrak{p}_2)$ the inner integral of (6.74) with Proposition 2.5. The function $\varphi := \varphi_{x_1, x_2}$ satisfies the assumptions of Proposition 2.5 with $z = x_1$ and $R = D^{s_1}$ by Lemma 6.7. We obtain with $B' := B(x_1, D^{s(\mathfrak{p}_1)})$,

$$\begin{aligned} \mathbf{I}(x_1, x_2) &\leq 2^{8a} \mu(B') \|\varphi\|_{C^\tau(B')} (1 + d_{B'}(Q(x_1), Q(x_2)))^{-1/(2a^2+a^3)} \\ &\leq \frac{2^{254a^3+8a}}{\mu(B(x_2, D^{s(\mathfrak{p}_2)}))} (1 + d_{B'}(Q(x_1), Q(x_2)))^{-1/(2a^2+a^3)}. \end{aligned} \quad (6.76)$$

Using Lemma 6.8 and $a \geq 1$ estimates (6.76) by

$$\leq \frac{2^{254a^3+8a+1}}{\mu(B(x_2, D^{s(\mathfrak{p}_2)}))} (1 + d_{\mathfrak{p}_1}(\mathcal{Q}(\mathfrak{p}_1), \mathcal{Q}(\mathfrak{p}_2)))^{-1/(2a^2+a^3)}. \quad (6.77)$$

As $x_2 \in \mathcal{I}(\mathfrak{p}_2)$ by Definition (2.20) of E , we have by (2.10)

$$\rho(x_2, c(\mathfrak{p}_2)) \leq 4D^{s(\mathfrak{p}_2)} \quad (6.78)$$

and thus by (2.10) again and the triangle inequality

$$\mathcal{I}(\mathfrak{p}_2) \subset B(x_2, 8D^{s(\mathfrak{p}_2)}). \quad (6.79)$$

Using three iterations of the doubling property (1.5) give

$$\mu(\mathcal{I}(\mathfrak{p}_2)) \leq 2^{3a} \mu(B(x_2, D^{s(\mathfrak{p}_2)})). \quad (6.80)$$

With $a \geq 1$ and (6.77) we conclude (6.28).

Now assume the left-hand side of (6.28) is not zero. There is a $y \in X$ with

$$T_{\mathfrak{p}}^* g(y) \overline{T_{\mathfrak{p}}^* g(y)} \neq 0 \quad (6.81)$$

By the triangle inequality and Lemma 6.9, we conclude

$$\rho(c(\mathfrak{p}), c(\mathfrak{p}')) \leq \rho(c(\mathfrak{p}), y) + \rho(c(\mathfrak{p}'), y) \leq 5D^{s(\mathfrak{p})} + 5D^{s(\mathfrak{p}')} \leq 10D^{s(\mathfrak{p})}. \quad (6.82)$$

By the squeezing property (2.10) and the triangle inequality, we conclude

$$\mathcal{I}(\mathfrak{p}') \subset B(c(\mathfrak{p}), 15D^{s(\mathfrak{p})}). \quad (6.83)$$

This completes the proof of Lemma 6.5.

6.3. Proof of Lemma 6.6, the geometric estimate.

Lemma 6.10 (tile reach). *Let $\vartheta \in \Theta$ and $N \geq 0$ be an integer. Let $\mathbf{p}, \mathbf{p}' \in \mathfrak{P}$ with*

$$d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \vartheta) \leq 2^N \quad (6.84)$$

$$d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), \vartheta) \leq 2^N. \quad (6.85)$$

Assume $\mathcal{I}(\mathbf{p}) \subset \mathcal{I}(\mathbf{p}')$ and $s(\mathbf{p}) < s(\mathbf{p}')$. Then

$$2^{N+2}\mathbf{p} \lesssim 2^{N+2}\mathbf{p}'. \quad (6.86)$$

Proof. By Lemma 2.8, we have

$$d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}'), \vartheta) \leq d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), \vartheta) \leq 2^N. \quad (6.87)$$

Together with (6.84) and the triangle inequality, we obtain

$$d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), \mathcal{Q}(\mathbf{p})) \leq 2^{N+1}. \quad (6.88)$$

Now assume

$$\vartheta' \in B_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), 2^{N+2}). \quad (6.89)$$

By the doubling property (1.8), applied five times, we have

$$d_{B(\mathbf{c}(\mathbf{p}'), 8D^{s(\mathbf{p}')})}(\mathcal{Q}(\mathbf{p}'), \vartheta') < 2^{5a+N+2}. \quad (6.90)$$

We have by the squeezing property (2.10)

$$\mathbf{c}(\mathbf{p}) \in B(\mathbf{c}(\mathbf{p}'), 4D^{s(\mathbf{p}')}). \quad (6.91)$$

Hence by the triangle inequality

$$B(\mathbf{c}(\mathbf{p}), 4D^{s(\mathbf{p}')}) \subset B(\mathbf{c}(\mathbf{p}'), 8D^{s(\mathbf{p}')}). \quad (6.92)$$

Together with (6.90) and monotonicity (1.9) of d

$$d_{B(\mathbf{c}(\mathbf{p}), 4D^{s(\mathbf{p}')})}(\mathcal{Q}(\mathbf{p}'), \vartheta') < 2^{5a+N+2}. \quad (6.93)$$

Using the doubling property (1.10) $5a + 2$ times gives

$$d_{B(\mathbf{c}(\mathbf{p}), 2^{2-5a^2-2a}D^{s(\mathbf{p}')})}(\mathcal{Q}(\mathbf{p}'), \vartheta') < 2^N. \quad (6.94)$$

Using $s(\mathbf{p}) < s(\mathbf{p}')$ and $D = 2^{100a^2}$ and $a \geq 4$ gives

$$d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}'), \vartheta') < 2^N. \quad (6.95)$$

With the triangle inequality and (6.88),

$$d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \vartheta') < 2^{N+2}. \quad (6.96)$$

This shows

$$B_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), 2^{N+2}) \subset B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 2^{N+2}). \quad (6.97)$$

This implies (6.86) and completes the proof of the lemma. \square

For $\vartheta \in \Theta$ and $N \geq 0$ define

$$\mathfrak{A}_{\vartheta, N} := \{\mathbf{p} \in \mathfrak{A} : 2^N \leq 1 + d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \vartheta) \leq 2^{N+1}\}. \quad (6.98)$$

Lemma 6.11 (stack density). *Let $\vartheta \in \Theta$, $N \geq 0$ and $L \in \mathcal{D}$. Then*

$$\sum_{\mathbf{p} \in \mathfrak{A}_{\vartheta, N} : \mathcal{I}(\mathbf{p}) = L} \mu(E(\mathbf{p}) \cap G) \leq 2^{a(N+5)} \text{dens}_1(\mathfrak{A}) \mu(L). \quad (6.99)$$

Proof. Let ϑ, N, L be given and set

$$\mathfrak{A}' := \{\mathfrak{p} \in \mathfrak{A}_{\vartheta, N} : \mathcal{I}(\mathfrak{p}) = L\}. \quad (6.100)$$

Let $\mathfrak{p} \in \mathfrak{A}'$. We have by Definition (2.28) using $\lambda = 2$ and the squeezing property (2.15)

$$\mu(E(\mathfrak{p}) \cap G) \leq \mu(E_2(2, \mathfrak{p})) \leq 2^a \text{dens}_1(\mathfrak{A}') \mu(L). \quad (6.101)$$

By the covering property (1.11), applied $N + 4$ times, there is a collection Θ' of at most $2^{a(N+4)}$ elements such that

$$B_{\mathfrak{p}}(\vartheta, 2^{N+1}) \subset \bigcup_{\vartheta' \in \Theta'} B_{\mathfrak{p}}(\vartheta', 0.2). \quad (6.102)$$

As each $\mathcal{Q}(\mathfrak{p})$ with $\mathfrak{p} \in \mathfrak{A}_{\vartheta, N}$ is contained in the left-hand-side of (6.102) by definition, it is in at least one $B_{\mathfrak{p}}(\vartheta', 0.2)$ with $\vartheta' \in \Theta'$.

For two different $\mathfrak{p}, \mathfrak{p}' \in \mathfrak{A}'$, we have by (2.13) that $\Omega(\mathfrak{p})$ and $\Omega(\mathfrak{p}')$ are disjoint and thus by the squeezing property (2.15) we have for every $\vartheta' \in \Theta'$

$$\vartheta' \notin B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), 0.2) \cap B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}'), 0.2). \quad (6.103)$$

Hence at most one of $\mathcal{Q}(\mathfrak{p})$ and $\mathcal{Q}(\mathfrak{p}')$ is in $B_{\mathfrak{p}}(\vartheta', 0.2)$. It follows that there are at most $2^{a(N+4)}$ elements in \mathfrak{A}' . Adding (6.101) over \mathfrak{A}' proves (6.99). \square

Lemma 6.12 (local antichain density). *Let $\vartheta \in \Theta$ and N be an integer. Let \mathfrak{p}_{ϑ} be a tile with $\vartheta \in \Omega(\mathfrak{p}_{\vartheta})$. Then we have*

$$\sum_{\mathfrak{p} \in \mathfrak{A}_{\vartheta, N} : s(\mathfrak{p}_{\vartheta}) < s(\mathfrak{p})} \mu(E(\mathfrak{p}) \cap G \cap \mathcal{I}(\mathfrak{p}_{\vartheta})) \leq \mu(E_2(2^{N+3}, \mathfrak{p}_{\vartheta})). \quad (6.104)$$

Proof. Let \mathfrak{p} be any tile in $\mathfrak{A}_{\vartheta, N}$ with $s(\mathfrak{p}_{\vartheta}) < s(\mathfrak{p})$. By definition of E , the tile contributes zero to the sum on the left-hand side of (6.104) unless $\mathcal{I}(\mathfrak{p}) \cap \mathcal{I}(\mathfrak{p}_{\vartheta}) \neq \emptyset$, which we may assume. With $s(\mathfrak{p}_{\vartheta}) < s(\mathfrak{p})$ and the dyadic property (2.8) we conclude $\mathcal{I}(\mathfrak{p}_{\vartheta}) \subset \mathcal{I}(\mathfrak{p})$. By the squeezing property (2.15), we conclude from $\vartheta \in \Omega(\mathfrak{p}_{\vartheta})$ that

$$\vartheta \in B(\mathcal{Q}(\mathfrak{p}_{\vartheta}), 1). \quad (6.105)$$

We conclude from $\mathfrak{p} \in \mathfrak{A}_{\vartheta, N}$ that

$$\vartheta \in B(\mathcal{Q}(\mathfrak{p}), 2^{N+1}). \quad (6.106)$$

With Lemma 6.10, we conclude

$$2^{N+3} \mathfrak{p}_{\vartheta} \lesssim 2^{N+3} \mathfrak{p}. \quad (6.107)$$

By Definition (2.27) of E_2 , we conclude

$$E(\mathfrak{p}) \cap G \subset E_2(2^{N+3}, \mathfrak{p}_{\vartheta}). \quad (6.108)$$

Using disjointedness of the various $E(\mathfrak{p})$ with $\mathfrak{p} \in \mathfrak{A}$ by Lemma 6.1, we obtain (6.104). This proves the lemma. \square

Lemma 6.13 (global antichain density). *Let $\vartheta \in \Theta$ and let $N \geq 0$ be an integer. Then we have*

$$\sum_{\mathfrak{p} \in \mathfrak{A}_{\vartheta, N}} \mu(E(\mathfrak{p}) \cap G) \leq 2^{101a^3 + Na} \text{dens}_1(\mathfrak{A}) \mu(\cup_{\mathfrak{p} \in \mathfrak{A}} I_{\mathfrak{p}}). \quad (6.109)$$

Proof. Fix ϑ and N . Let \mathfrak{A}' be the set of $\mathfrak{p} \in \mathfrak{A}_{\vartheta, N}$ such that $\mathcal{I}(\mathfrak{p}) \cap G$ is not empty.

Let \mathcal{L} be the collection of dyadic cubes $I \in \mathcal{D}$ such that $I \subset \mathcal{I}(\mathfrak{p})$ for some $\mathfrak{p} \in \mathfrak{A}'$ and if $\mathcal{I}(\mathfrak{p}) \subset I$ for some $\mathfrak{p} \in \mathfrak{A}'$, then $s(\mathfrak{p}) = -S$. By (2.7), for each $\mathfrak{p} \in \mathfrak{A}'$ and each $x \in \mathcal{I}(\mathfrak{p}) \cap G$, there is $I \in \mathcal{D}$ with $s(I) = -S$ and $x \in I$. By (2.8), we have $I \subset \mathcal{I}(\mathfrak{p})$. Hence

$$\mathcal{I}(\mathfrak{p}) \subset \bigcup \{I \in \mathcal{D} : s(I) = -S, I \subset \mathcal{I}(\mathfrak{p})\} \subset \bigcup \mathcal{L}. \quad (6.110)$$

As each $I \in \mathcal{L}$ satisfies $I \subset \mathcal{I}(\mathfrak{p})$ for some \mathfrak{p} in \mathfrak{A}' , we conclude

$$\bigcup \mathcal{L} = \bigcup_{\mathfrak{p} \in \mathfrak{A}'} \mathcal{I}(\mathfrak{p}). \quad (6.111)$$

Let \mathcal{L}^* be the set of maximal elements in \mathcal{L} with respect to set inclusion. By (2.8), the elements in \mathcal{L}^* are pairwise disjoint and we have

$$\bigcup \mathcal{L}^* = \bigcup_{\mathfrak{p} \in \mathfrak{A}'} \mathcal{I}(\mathfrak{p}). \quad (6.112)$$

Using the partition (6.112) into elements of \mathcal{L} in (6.113), it suffices to show for each $L \in \mathcal{L}^*$

$$\sum_{\mathfrak{p} \in \mathfrak{A}'} \mu(E(\mathfrak{p}) \cap G \cap L) \leq 2^{101a^3 + aN} \text{dens}_1(\mathfrak{A}) \mu(L). \quad (6.113)$$

Fix $L \in \mathcal{L}^*$. By definition of L , there exists an element $\mathfrak{p}' \in \mathfrak{A}'$ such that $L \subset \mathcal{I}(\mathfrak{p}')$. Pick such an element \mathfrak{p}' in \mathfrak{A} with minimal $s(\mathfrak{p}')$. As $\mathcal{I}(\mathfrak{p}') \not\subset L$ or $s(L) = -S$ by definition of L , we have with (2.8) that $s(L) < s(\mathfrak{p}')$ or $s(L) = -S$. In particular $s(L) < S$.

If there exists no cube $J \in \mathcal{D}$ with $L \subsetneq J$, then by the definition of \mathcal{L} we must have $s(L) = -S$. By (2.8), this implies that each tile \mathfrak{p} with $E(\mathfrak{p}) \cap L \neq \emptyset$ must satisfy $\mathcal{I}(\mathfrak{p}) = L$. Thus the left hand side of (6.113) equals

$$\sum_{\mathfrak{p} \in \mathfrak{A}' : \mathcal{I}(\mathfrak{p}) = L} \mu(E(\mathfrak{p}) \cap G \cap L). \quad (6.114)$$

Thus in this case (6.113) is a direct consequence of Lemma 6.11 and $a \geq 4$.

We now assume that there exists a cube $J \in \mathcal{D}$ with $L \subsetneq J$. By (2.7), there is an $L' \in \mathcal{D}$ with $s(L') = s(L) + 1$ and $c(L) \in L'$. By (2.8), we have $L \subset L'$.

We split the left-hand side of (6.113) as

$$\sum_{\mathfrak{p} \in \mathfrak{A}' : \mathcal{I}(\mathfrak{p}) = L'} \mu(E(\mathfrak{p}) \cap G \cap L) \quad (6.115)$$

$$+ \sum_{\mathfrak{p} \in \mathfrak{A}' : \mathcal{I}(\mathfrak{p}) \neq L'} \mu(E(\mathfrak{p}) \cap G \cap L), \quad (6.116)$$

We first estimate (6.115) with Lemma 6.11 by

$$\leq \sum_{\mathfrak{p} \in \mathfrak{A}' : \mathcal{I}(\mathfrak{p}) = L'} \mu(E(\mathfrak{p}) \cap G \cap L') \leq 2^{a(N+5)} \text{dens}_1(\mathfrak{A}) \mu(L'). \quad (6.117)$$

We turn to (6.116). Consider the element $\mathfrak{p}' \in \mathfrak{A}'$ as above with $L \subset \mathcal{I}(\mathfrak{p}')$ and $s(L) < s(\mathfrak{p}')$. As $L \subset L'$ and $s(L') = s(L) + 1$, we conclude with the

dyadic property that $L' \subset \mathcal{I}(\mathbf{p}')$. By maximality of L , we have $L' \notin \mathcal{L}$. This together with the existence of the given $\mathbf{p}' \in \mathfrak{A}$ with $L' \subset \mathcal{I}(\mathbf{p}')$ shows by definition of \mathcal{L} that there exists $\mathbf{p}'' \in \mathfrak{A}'$ with $\mathcal{I}(\mathbf{p}'') \subset L'$.

By the covering property (2.13), there exists a unique \mathbf{p}_ϑ with

$$\mathcal{I}(\mathbf{p}_\vartheta) = L'$$

such that $\vartheta \in \Omega(\mathbf{p}_\vartheta)$. Note that

$$\vartheta \in B(\mathcal{Q}(\mathbf{p}_\vartheta), 1) \quad (6.118)$$

and as $\mathbf{p}'' \in \mathfrak{A}_{\vartheta, N}$ that

$$\vartheta \in B(\mathcal{Q}(\mathbf{p}''), 2^{N+1}). \quad (6.119)$$

By Lemma 6.10, we conclude

$$2^{N+3}\mathbf{p}'' \lesssim 2^{N+3}\mathbf{p}_\vartheta. \quad (6.120)$$

As $\mathbf{p}'' \in \mathfrak{A}'$, we have by Definition (2.28) of dens_1 that

$$\mu(E_2(2^{N+3}, \mathbf{p}_\vartheta)) \leq 2^{Na+3a} \text{dens}_1(\mathfrak{A})\mu(L'). \quad (6.121)$$

Now let \mathbf{p} be any tile in the summation set in (6.116), that is, $\mathbf{p} \in \mathfrak{A}'$ and $\mathcal{I}(\mathbf{p}) \neq L'$. Then $\mathcal{I}(\mathbf{p}) \cap L \neq \emptyset$. It follows by the dyadic property (2.8) and the definition of L that $L \subset \mathcal{I}(\mathbf{p})$ and $L \neq \mathcal{I}(\mathbf{p})$. By the dyadic property (2.8), we have $s(L) < s(\mathbf{p})$ and thus $s(L') \leq s(\mathbf{p})$. By the dyadic property (2.8) again, we have $L' \subset \mathcal{I}(\mathbf{p})$. As $L' \neq \mathcal{I}(\mathbf{p})$, we conclude $s(L) < s(\mathbf{p})$. By Lemma 6.12, we can thus estimate (6.116) by

$$\sum_{\mathbf{p} \in \mathfrak{A}': \mathcal{I}(\mathbf{p}) \neq L'} \mu(E(\mathbf{p}) \cap G \cap L') \leq \mu(E_2(2^{N+3}, \mathbf{p}_\vartheta)). \quad (6.122)$$

Using the decomposition into (6.115) and (6.116) and the estimates (6.117), (6.104), (6.121) we obtain the estimate

$$\sum_{\mathbf{p} \in \mathfrak{A}'} \mu(E(\mathbf{p}) \cap G \cap L) \leq (2^{a(N+5)} + 2^{Na+3a}) \text{dens}_1(\mathfrak{A})\mu(L'). \quad (6.123)$$

Using $s(L') = s(L) + 1$ and $D = 2^{100a^2}$ and the squeezing property (2.10) and the doubling property (1.5) $100a^2 + 4$ times, we obtain

$$\mu(L') \leq 2^{100a^3+4a}\mu(L). \quad (6.124)$$

Inserting in (6.123) and using $a \geq 4$ gives (6.113). This completes the proof of the lemma. \square

We turn to the proof of Lemma 6.6.

Proof. Using that \mathfrak{A} is the union of the $\mathfrak{A}_{\vartheta, N}$ with $N \geq 0$, we estimate the left-hand side (6.36) with the triangle inequality by

$$\leq \sum_{N \geq 0} \left\| \sum_{\mathbf{p} \in \mathfrak{A}_{\vartheta, N}} 2^{-N/(2a^2+a^3)} \mathbf{1}_{E(\mathbf{p})} \mathbf{1}_G \right\|_p \quad (6.125)$$

We consider each individual term in this sum and estimate it's p -th power. Using that for each $x \in X$ by Lemma 6.13 there is at most one $\mathfrak{p} \in \mathfrak{A}$ with $x \in E(\mathfrak{p})$, we have

$$\left\| \sum_{\mathfrak{p} \in \mathfrak{A}_{\vartheta, N}} 2^{-N/(2a^2+a^3)} \mathbf{1}_{E(\mathfrak{p})} \mathbf{1}_G \right\|_p^p \quad (6.126)$$

$$= \int_G \left(\sum_{\mathfrak{p} \in \mathfrak{A}_{\vartheta, N}} 2^{-N/(2a^2+a^3)} \mathbf{1}_{E(\mathfrak{p})}(x) \right)^p d\mu(x) \quad (6.127)$$

$$= \int_G \sum_{\mathfrak{p} \in \mathfrak{A}_{\vartheta, N}} 2^{-pN/(2a^2+a^3)} \mathbf{1}_{E(\mathfrak{p})}(x) d\mu(x) \quad (6.128)$$

$$= 2^{-pN/(2a^2+a^3)} \sum_{\mathfrak{p} \in \mathfrak{A}_{\vartheta, N}} \mu(E(\mathfrak{p}) \cap G) \quad (6.129)$$

Using Lemma 6.13, we estimate the last display by

$$\leq 2^{-pN/(2a^2+a^3)+101a^3+Na} \text{dens}_1(\mathfrak{A}) \mu(\cup_{\mathfrak{p} \in \mathfrak{A}} \mathcal{I}(\mathfrak{p})) \quad (6.130)$$

Using that with $a \geq 4$ and since $p = 4a^4$, we have

$$pN/(2a^2 + a^3) \geq 4a^4 N/(3a^3) \geq Na + N. \quad (6.131)$$

Hence we have for (6.130) the upper bound

$$\leq 2^{101a^3-N} \text{dens}_1(\mathfrak{A}) \mu(\cup_{\mathfrak{p} \in \mathfrak{A}} \mathcal{I}(\mathfrak{p})). \quad (6.132)$$

Taking the p -th root and summing over $N \geq 0$ gives for (6.125) the upper bound

$$\leq \left(\sum_{N \geq 0} 2^{-N/p} \right) 2^{101a^3/p} \text{dens}_1(\mathfrak{A})^{\frac{1}{p}} \mu(\cup_{\mathfrak{p} \in \mathfrak{A}} \mathcal{I}(\mathfrak{p}))^{\frac{1}{p}} \quad (6.133)$$

$$\leq \left(1 - 2^{-1/p} \right)^{-1} 2^{101a^3/p} \text{dens}_1(\mathfrak{A})^{\frac{1}{p}} \mu(\cup_{\mathfrak{p} \in \mathfrak{A}} \mathcal{I}(\mathfrak{p}))^{\frac{1}{p}}. \quad (6.134)$$

Using that $p = 4a^4$ and $a \geq 4$, this proves the lemma. \square

7. PROOF OF PROPOSITION 2.4

7.1. The pointwise tree estimate. Fix a forest $(\mathfrak{U}, \mathfrak{T})$. The main result of this subsection is Lemma 7.3, we begin this section with some definitions necessary to state the lemma.

For $\mathfrak{u} \in \mathfrak{U}$ and $x \in X$, we define

$$\sigma(\mathfrak{u}, x) := \{s(\mathfrak{p}) : \mathfrak{p} \in \mathfrak{T}(\mathfrak{u}), x \in E(\mathfrak{p})\}.$$

This is a subset of $\mathbb{Z} \cap [-S, S]$, so has a minimum and a maximum. We set

$$\overline{\sigma}(\mathfrak{u}, x) := \max \sigma(\mathfrak{T}(\mathfrak{u}), x)$$

$$\underline{\sigma}(\mathfrak{u}, x) := \min \sigma(\mathfrak{T}(\mathfrak{u}), x).$$

Lemma 7.1 (convex scales). *For each $\mathfrak{u} \in \mathfrak{U}$, we have*

$$\sigma(\mathfrak{u}, x) = \mathbb{Z} \cap [\underline{\sigma}(\mathfrak{u}, x), \overline{\sigma}(\mathfrak{u}, x)].$$

Proof. Let $s \in \mathbb{Z}$ with $\underline{\sigma}(\mathbf{u}, x) \leq s \leq \overline{\sigma}(\mathbf{u}, x)$. By definition of σ , there exists $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$ with $s(\mathbf{p}) = \underline{\sigma}(\mathbf{u}, x)$ and $x \in E(\mathbf{p})$, and there exists $\mathbf{p}'' \in \mathfrak{T}(\mathbf{u})$ with $s(\mathbf{p}'') = \overline{\sigma}(\mathbf{u}, x)$ and $x \in E(\mathbf{p}'') \subset \mathcal{I}(\mathbf{p}'')$. By property (2.7) of the dyadic grid, there exists a cube $I \in \mathcal{D}$ of scale s with $x \in I$. By property (2.13), there exists a tile $\mathbf{p}' \in \mathfrak{P}(I)$ with $Q(x) \in \Omega(\mathbf{p}')$. By the dyadic property (2.8) we have $\mathcal{I}(\mathbf{p}) \subset \mathcal{I}(\mathbf{p}') \subset \mathcal{I}(\mathbf{p}'')$, and by (2.14), we have $\Omega(\mathbf{p}'') \subset \Omega(\mathbf{p}') \subset \Omega(\mathbf{p})$. Thus $\mathbf{p} \leq \mathbf{p}' \leq \mathbf{p}''$, which gives with the convexity property (2.33) of $\mathfrak{T}(\mathbf{u})$ that $\mathbf{p}' \in \mathfrak{T}(\mathbf{u})$, so $s \in \sigma(\mathbf{u}, x)$. \square

For a nonempty collection of tiles $\mathfrak{S} \subset \mathfrak{P}$ we define

$$\mathcal{J}_0(\mathfrak{S})$$

to be the collection of all dyadic cubes $J \in \mathcal{D}$ such that $s(J) = -S$ or

$$\mathcal{I}(\mathbf{p}) \not\subset B(c(J), 100D^{s(J)+1})$$

for all $\mathbf{p} \in \mathfrak{S}$. We define $\mathcal{J}(\mathfrak{S})$ to be the collection of inclusion maximal cubes in $\mathcal{J}_0(\mathfrak{S})$.

We further define

$$\mathcal{L}_0(\mathfrak{S})$$

to be the collection of dyadic cubes $L \in \mathcal{D}$ such that $s(L) = -S$, or there exists $\mathbf{p} \in \mathfrak{S}$ with $L \subset \mathcal{I}(\mathbf{p})$ and there exists no $\mathbf{p} \in \mathfrak{S}$ with $\mathcal{I}(\mathbf{p}) \subset L$. We define $\mathcal{L}(\mathfrak{S})$ to be the collection of inclusion maximal cubes in $\mathcal{L}_0(\mathfrak{S})$.

Lemma 7.2 (dyadic partitions). *For each $\mathfrak{S} \subset \mathfrak{P}$, we have*

$$\bigcup_{I \in \mathcal{D}} I = \bigcup_{J \in \mathcal{J}(\mathfrak{S})} J \quad (7.1)$$

and

$$\bigcup_{\mathbf{p} \in \mathfrak{S}} \mathcal{I}(\mathbf{p}) = \bigcup_{L \in \mathcal{L}(\mathfrak{S})} L. \quad (7.2)$$

Proof. Since $\mathcal{J}(\mathfrak{S})$ is the set of inclusion maximal cubes in $\mathcal{J}_0(\mathfrak{S})$, cubes in $\mathcal{J}(\mathfrak{S})$ are pairwise disjoint by (2.8). The same applies to $\mathcal{L}(\mathfrak{S})$.

If $x \in \bigcup_{I \in \mathcal{D}} I$, then there exists by (2.7) a cube $I \in \mathcal{D}$ with $x \in I$ and $s(I) = -S$. Then $I \in \mathcal{J}_0(\mathfrak{S})$. There exists an inclusion maximal cube in $\mathcal{J}_0(\mathfrak{S})$ containing I . This cube contains x and is contained in $\mathcal{J}(\mathfrak{S})$. This shows one inclusion in (7.1), the other one follows from $\mathcal{J}(\mathfrak{S}) \subset \mathcal{D}$.

The proof of the two inclusions in (7.2) is similar. \square

For a finite collection of pairwise disjoint cubes \mathcal{C} , define the projection operator

$$P_{\mathcal{C}}f(x) := \sum_{J \in \mathcal{C}} \mathbf{1}_J(x) \frac{1}{\mu(J)} \int_J f(y) \, d\mu(y).$$

Given a scale $-S \leq s \leq S$ and a point $x \in \bigcup_{I \in \mathcal{D}, s(I)=s} I$, there exists a unique cube in \mathcal{D} of scale s containing x by (2.7). We denote it by $I_s(x)$. Define the nontangential maximal operator

$$T_{\mathcal{N}}f(x) := \sup_{-S \leq s_1 < s_2 \leq S} \sup_{x' \in I_{s_1}(x)} \left| \sum_{s=s_1}^{s_2} \int K_s(x', y) f(y) \, d\mu(y) \right|. \quad (7.3)$$

Define for each $\mathbf{u} \in \mathfrak{U}$ the auxiliary operator

$$S_{1,\mathbf{u}}f(x) := \sum_{I \in \mathcal{D}} \mathbf{1}_I(x) \sum_{\substack{J \in \mathcal{J}(\mathfrak{T}(\mathbf{u})) \\ J \subset B(c(I), 16D^{s(I)})}} \frac{D^{(s(J)-s(I))/a}}{\mu(B(c(I), 16D^{s(I)}))} \int_J |f(y)| d\mu(y). \quad (7.4)$$

Define also the collection of balls

$$\mathcal{B} = \{B(c(I), D^{s+s(I)}) : I \in \mathcal{D}, 0 \leq s \leq S+5\}.$$

The following pointwise estimate for operators associated to sets $\mathfrak{T}(\mathbf{u})$ is the main result of this subsection.

Lemma 7.3 (pointwise tree estimate). *Let $\mathbf{u} \in \mathfrak{U}$ and $L \in \mathcal{L}(\mathfrak{T}(\mathbf{u}))$. Let $x, x' \in L$. Then for all bounded functions f with bounded support*

$$\left| \sum_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathbf{p}}[e(\mathcal{Q}(\mathbf{u}))f](x) \right| \leq 2^{151a^3} (M_{\mathcal{B},1} + S_{1,\mathbf{u}}) P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} |f|(x') + |T_{\mathcal{N}} P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} f(x')|, \quad (7.5)$$

Proof. By (2.21), if $T_{\mathbf{p}}[e(-\mathcal{Q}(\mathbf{u}))f](x) \neq 0$, then $x \in E(\mathbf{p})$. Combining this with $|e(\mathcal{Q}(\mathbf{u})(x) - \mathcal{Q}(x)(x))| = 1$, we obtain

$$\begin{aligned} & \left| \sum_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathbf{p}}[e(\mathcal{Q}(\mathbf{u}))f](x) \right| \\ &= \left| \sum_{s \in \sigma(\mathbf{u}, x)} \int e(-\mathcal{Q}(\mathbf{u})(y) + \mathcal{Q}(x)(y) + \mathcal{Q}(\mathbf{u})(x) - \mathcal{Q}(x)(x)) \times \right. \\ & \quad \left. K_s(x, y) f(y) d\mu(y) \right|. \end{aligned}$$

Using the triangle inequality, we bound this by the sum of three terms:

$$\begin{aligned} & \leq \left| \sum_{s \in \sigma(\mathbf{u}, x)} \int (e(-\mathcal{Q}(\mathbf{u})(y) + \mathcal{Q}(x)(y) + \mathcal{Q}(\mathbf{u})(x) - \mathcal{Q}(x)(x)) - 1) \times \right. \\ & \quad \left. K_s(x, y) f(y) d\mu(y) \right| \quad (7.6) \end{aligned}$$

$$+ \left| \sum_{s \in \sigma(\mathbf{u}, x)} \int K_s(x, y) P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} f(y) d\mu(y) \right| \quad (7.7)$$

$$+ \left| \sum_{s \in \sigma(\mathbf{u}, x)} \int K_s(x, y) (f(y) - P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} f(y)) d\mu(y) \right|. \quad (7.8)$$

The proof is completed using the bounds for these three terms proven in Lemma 7.4, Lemma 7.5 and Lemma 7.6. \square

Lemma 7.4 (first tree pointwise). *For all $u \in \mathfrak{U}$, all $L \in \mathcal{L}(\mathfrak{T}(u))$, all $x, x' \in L$ and all bounded f with bounded support, we have*

$$(7.6) \leq 10 \cdot 2^{105a^3} M_{\mathcal{B},1} P_{\mathcal{J}(\mathfrak{T}(u))} |f|(x').$$

Proof. Let $s \in \sigma(u, x)$. If $x, y \in X$ are such that $K_s(x, y) \neq 0$, then, by (2.48), we have $\rho(x, y) \leq 1/2D^s$. By 1-Lipschitz continuity of the function $t \mapsto \exp(it) = e(t)$ and the property (1.7) of the metrics d_B , it follows that

$$\begin{aligned} & |e(-\mathcal{Q}(u)(y) + Q(x)(y) + \mathcal{Q}(u)(x) - Q(x)(x)) - 1| \\ & \leq d_{B(x, 1/2D^s)}(\mathcal{Q}(u), Q(x)). \end{aligned}$$

Let $\mathfrak{p}_s \in \mathfrak{T}(u)$ be a tile with $s(\mathfrak{p}_s) = s$ and $x \in E(\mathfrak{p}_s)$, and let \mathfrak{p}' be a tile with $s(\mathfrak{p}') = \bar{\sigma}(u, x)$ and $x \in E(\mathfrak{p}')$. Using the doubling property (1.8), the definition of $d_{\mathfrak{p}}$ and Lemma 2.8, we can bound the previous display by

$$2^a d_{\mathfrak{p}_s}(\mathcal{Q}(u), Q(x)) \leq 2^a 2^{s-\bar{\sigma}(u, x)} d_{\mathfrak{p}'}(\mathcal{Q}(u), Q(x)).$$

Since $\mathcal{Q}(u) \in B_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), 4)$ by (2.32) and $Q(x) \in \Omega(\mathfrak{p}') \subset B_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), 1)$ by (2.15), this is estimated by

$$\leq 5 \cdot 2^a 2^{s-\bar{\sigma}(u, x)}.$$

Using (2.49), it follows that

$$(7.6) \leq 5 \cdot 2^{103a^3} \sum_{s \in \sigma(x)} 2^{s-\bar{\sigma}(u, x)} \frac{1}{\mu(B(x, D^s))} \int_{B(x, 0.5D^s)} |f(y)| d\mu(y).$$

By (7.1), the collection \mathcal{J} is a partition of X , so this is estimated by

$$5 \cdot 2^{103a^3} \sum_{s \in \sigma(x)} 2^{s-\bar{\sigma}(u, x)} \frac{1}{\mu(B(x, D^s))} \sum_{\substack{J \in \mathcal{J}(\mathfrak{T}(u)) \\ J \cap B(x, 0.5D^s) \neq \emptyset}} \int_J |f(y)| d\mu(y).$$

This expression does not change if we replace $|f|$ by $P_{\mathcal{J}(\mathfrak{T}(u))}|f|$.

Let $J \in \mathcal{J}(\mathfrak{T}(u))$ with $B(x, 0.5D^s) \cap J \neq \emptyset$. By the triangle inequality and since $x \in E(\mathfrak{p}_s) \subset B(c(\mathfrak{p}_s), 4D^s)$, it follows that $B(c(\mathfrak{p}_s), 4.5D^s) \cap J \neq \emptyset$. If $s(J) \geq s$ and $s(J) > -S$, then it follows from the triangle inequality, (2.10) and (2.1) that $\mathcal{I}(\mathfrak{p}_s) \subset B(c(J), 100D^{s(J)+1})$, contradicting $J \in \mathcal{J}(\mathfrak{T}(u))$. Thus $s(J) \leq s-1$ or $s(J) = -S$. If $s(J) = -S$ and $s(J) > s-1$, then $s = -S$. Thus we always have $s(J) \leq s$. It then follows from the triangle inequality and (2.10) that $J \subset B(c(\mathfrak{p}_s), 16D^s)$.

Thus we can continue our chain of estimates with

$$5 \cdot 2^{103a^3} \sum_{s \in \sigma(x)} 2^{s-\bar{\sigma}(u, x)} \frac{1}{\mu(B(x, D^s))} \int_{B(c(\mathfrak{p}_s), 16D^s)} P_{\mathcal{J}(\mathfrak{T}(u))} |f(y)| d\mu(y).$$

We have $B(c(\mathfrak{p}_s), 16D^s) \subset B(x, 32D^s)$, by (2.10) and the triangle inequality, since $x \in \mathcal{I}(\mathfrak{p})$. Combining this with the doubling property (1.5), we obtain

$$\mu(B(c(\mathfrak{p}_s), 16D^s)) \leq 2^{5a} \mu(B(x, D^s)).$$

Since $a \geq 4$, it follows that (7.6) is bounded by

$$2^{104a^3} \sum_{s \in \sigma(x)} 2^{s-\bar{\sigma}(u, x)} \frac{1}{\mu(B(c(\mathfrak{p}_s), 16D^s))} \int_{B(c(\mathfrak{p}_s), 16D^s)} P_{\mathcal{J}(\mathfrak{T}(u))} |f(y)| d\mu(y).$$

Since $L \in \mathcal{L}(\mathfrak{T}(\mathbf{u}))$, we have $s(L) \leq s(\mathbf{p})$ for all $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$. Since $x \in L \cap \mathcal{I}(\mathbf{p}_s)$, it follows by (2.8) that $L \subset \mathcal{I}(\mathbf{p}_s)$, in particular $x' \in \mathcal{I}(\mathbf{p}_s) \subset B(c(\mathbf{p}_s), 16D^s)$. Thus

$$\begin{aligned} &\leq 2^{104a^3} \sum_{s \in \sigma(x)} 2^{s-\bar{\sigma}(\mathbf{u},x)} M_{\mathcal{B},1} P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} |f|(x') \\ &\leq 2^{105a^3} M_{\mathcal{B},1} P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} |f|(x'). \end{aligned}$$

This completes the estimate for term (7.6). \square

Lemma 7.5 (second tree pointwise). *For all $\mathbf{u} \in \mathfrak{U}$, all $L \in \mathcal{L}(\mathfrak{T}(\mathbf{u}))$, all $x, x' \in L$ and all bounded f with bounded support, we have*

$$\left| \sum_{s \in \sigma(\mathbf{u},x)} \int K_s(x,y) P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} f(y) d\mu(y) \right| \leq T_{\mathcal{N}} P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} f(x').$$

Proof. Let $s = \underline{\sigma}(\mathbf{u},x)$. By definition, there exists a tile $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$ with $s(\mathbf{p}) = s$ and $x \in E(\mathbf{p})$. Then $x \in \mathcal{I}(\mathbf{p}) \cap L$. By (2.8) and the definition of $\mathcal{L}(\mathfrak{T}(\mathbf{u}))$, it follows that $L \subset \mathcal{I}(\mathbf{p})$, in particular $x' \in \mathcal{I}(\mathbf{p})$, so $x \in I_s(x')$. The lemma now follows from the definition of $T_{\mathcal{N}}$. \square

Lemma 7.6 (third tree pointwise). *For all $\mathbf{u} \in \mathfrak{U}$, all $L \in \mathcal{L}(\mathfrak{T}(\mathbf{u}))$, all $x, x' \in L$ and all bounded f with bounded support, we have*

$$\begin{aligned} &\left| \sum_{s \in \sigma(\mathbf{u},x)} \int K_s(x,y) (f(y) - P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} f(y)) d\mu(y) \right| \\ &\leq 2^{151a^3} S_{1,\mathbf{u}} P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} |f|(x'). \end{aligned}$$

Proof. We have for $J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}))$:

$$\begin{aligned} &\int_J K_s(x,y) (1 - P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))}) f(y) d\mu(y) \\ &= \int_J \frac{1}{\mu(J)} \int_J K_s(x,y) - K_s(x,z) d\mu(z) f(y) d\mu(y). \end{aligned} \quad (7.9)$$

By (2.50) and (2.10), we have for $y, z \in J$

$$|K_s(x,y) - K_s(x,z)| \leq \frac{2^{150a^3}}{\mu(B(x, D^s))} \left(\frac{8D^{s(J)}}{D^s} \right)^{1/a}.$$

Suppose that $s \in \sigma(\mathbf{u},x)$. If $K_s(x,y) \neq 0$ for some $y \in J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}))$ then, by (2.48), $y \in B(x, 0.5D^s) \cap J \neq \emptyset$. Let $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$ with $s(\mathbf{p}) = s$ and $x \in E(\mathbf{p})$. Then $B(c(\mathbf{p}_s), 4.5D^s) \cap J \neq \emptyset$ by the triangle inequality. If $s(J) \geq s$ and $s(J) > -S$, then it follows from the triangle inequality, (2.10) and (2.1) that $\mathcal{I}(\mathbf{p}) \subset B(c(J), 100D^{s(J)+1})$, contradicting $J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}))$. Thus $s(J) \leq s-1$ or $s(J) = -S$. If $s(J) = -S$ and $s(J) > s-1$, then $s = -S$. So in both cases, $s(J) \leq s$. It then follows from the triangle inequality and (2.10) that $J \subset B(x, 16D^s)$.

Thus, we can estimate (7.8) by

$$2^{150a^3+3/a} \sum_{\mathbf{p} \in \mathfrak{T}} \frac{\mathbf{1}_{E(\mathbf{p})}(x)}{\mu(B(x, D^{s(\mathbf{p})}))} \sum_{\substack{J \in \mathcal{J}(\mathfrak{T}(\mathbf{u})) \\ J \subset B(x, 16D^{s(\mathbf{p})})}} D^{(s(J)-s(\mathbf{p}))/a} \int_J |f|.$$

$$= 2^{150a^3+3/a} \sum_{I \in \mathcal{D}} \sum_{\substack{\mathfrak{p} \in \mathfrak{T} \\ \mathcal{I}(\mathfrak{p})=I}} \frac{\mathbf{1}_{E(\mathfrak{p})}(x)}{\mu(B(x, D^{s(I)}))} \sum_{\substack{J \in \mathcal{J}(\mathfrak{T}(\mathfrak{u})) \\ J \subset B(x, 16D^{s(\mathfrak{p})})}} D^{(s(J)-s(I))/a} \int_J |f|.$$

By (2.13) and (2.20), the sets $E(\mathfrak{p})$ for tiles \mathfrak{p} with $\mathcal{I}(\mathfrak{p}) = I$ are pairwise disjoint. If $x \in E(\mathfrak{p})$ then in particular $x \in \mathcal{I}(\mathfrak{p})$, so by (2.10) $B(c(I), 16D^{s(I)}) \subset B(x, 32D^{s(I)})$. By the doubling property (1.5)

$$\mu(B(c(I), 16D^{s(I)})) \leq 2^{5a} \mu(B(x, D^{s(I)})).$$

Since $a \geq 4$ we can continue our estimate with

$$\leq 2^{151a^3} \sum_{I \in \mathcal{D}} \frac{\mathbf{1}_I(x)}{\mu(B(c(I), 16D^{s(I)}))} \sum_{\substack{J \in \mathcal{J}(\mathfrak{T}(\mathfrak{u})) \\ J \subset B(x, 16D^{s(\mathfrak{p})})}} D^{(s(J)-s(\mathfrak{p}))/a} \int_J |f|.$$

Finally, it follows from the definition of $\mathcal{L}(\mathfrak{T}(\mathfrak{u}))$ that $x \in \mathcal{I}(\mathfrak{p})$ if and only if $x' \in \mathcal{I}(\mathfrak{p})$, thus this equals

$$2^{151a^3} S_{1,\mathfrak{u}} P_{\mathcal{J}(\mathfrak{T}(\mathfrak{u}))} |f|(x').$$

This completes the proof. \square

7.2. An auxiliary L^2 tree estimate. In this subsection we prove the following estimate on L^2 for operators associated to trees.

Lemma 7.7 (tree projection estimate). *Let $\mathfrak{u} \in \mathfrak{U}$. Then we have for all f, g bounded with bounded support*

$$\begin{aligned} & \left| \int_X \sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} \bar{g}(y) T_{\mathfrak{p}} f(y) d\mu(y) \right| \\ & \leq 2^{104a^3} \|P_{\mathcal{J}(\mathfrak{T}(\mathfrak{u}))} f\|_2 \|P_{\mathcal{L}(\mathfrak{T}(\mathfrak{u}))} g\|_2. \end{aligned} \quad (7.10)$$

Below, we deduce Lemma 7.7 from Lemma 7.3 and the following estimates for the operators in Lemma 7.3.

Lemma 7.8 (nontangential operator bound). *For all bounded f with bounded support*

$$\|T_{\mathcal{N}} f\|_2 \leq 2^{103a^3} \|f\|_2.$$

Lemma 7.9 (boundary operator bound). *For all $\mathfrak{u} \in \mathfrak{U}$ and all bounded functions f with bounded support*

$$\|S_{1,\mathfrak{u}} f\|_2 \leq 2^{12a} \|f\|_2. \quad (7.11)$$

Proof of Lemma 7.7. For each $L \in \mathcal{L}(\mathfrak{T}(\mathfrak{u}))$, choose a point $x'(L) \in L$ such that for all $y \in L$

$$\begin{aligned} & (M_{\mathcal{B},1} + S_{1,\mathfrak{u}}) P_{\mathcal{J}(\mathfrak{T}(\mathfrak{u}))} |f|(x') + |T_{\mathcal{N}} P_{\mathcal{J}(\mathfrak{T}(\mathfrak{u}))} f(x')| \\ & \leq 2((M_{\mathcal{B},1} + S_{1,\mathfrak{u}}) P_{\mathcal{J}(\mathfrak{T}(\mathfrak{u}))} |f|(y) + |T_{\mathcal{N}} P_{\mathcal{J}(\mathfrak{T}(\mathfrak{u}))} f(y)|). \end{aligned} \quad (7.12)$$

This point exists since (7.12) is non-negative for each y . Then we have by Lemma 7.3 for each $L \in \mathcal{L}(\mathfrak{T}(\mathfrak{u}))$

$$\int_L |g(y)| \left| \sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} T_{\mathfrak{p}} f(y) \right| d\mu(y)$$

$$\begin{aligned}
&\leq 2^{151a^3} \int_L |g(y)| ((M_{\mathcal{B},1} + S_{1,u})P_{\mathcal{J}(\mathfrak{T}(u))}|f|(x') + |T_{\mathcal{N}}P_{\mathcal{J}(\mathfrak{T}(u))}f(x')|) d\mu(y) \\
&\leq 2^{151a^3} \int_L |g(y)| d\mu(y) \times \\
&\quad \int_L 2((M_{\mathcal{B},1} + S_{1,u})P_{\mathcal{J}(\mathfrak{T}(u))}|f|(y) + |T_{\mathcal{N}}P_{\mathcal{J}(\mathfrak{T}(u))}f(y)|) d\mu(y) \\
&= 2^{151a^3+1} \int_L P_{\mathcal{L}(\mathfrak{T}(u))}|g|(y) \times \\
&\quad ((M_{\mathcal{B},1} + S_{1,u})P_{\mathcal{J}(\mathfrak{T}(u))}|f|(y) + |T_{\mathcal{N}}P_{\mathcal{J}(\mathfrak{T}(u))}f(y)|) d\mu(y).
\end{aligned}$$

By (2.21), we have $T_{\mathfrak{p}}f = \mathbf{1}_{\mathcal{I}(\mathfrak{p})}T_{\mathfrak{p}}f$ for all $\mathfrak{p} \in \mathfrak{P}$, so

$$\left| \int \bar{g}(y) \sum_{\mathfrak{p} \in \mathfrak{T}(u)} T_{\mathfrak{p}}f(y) d\mu(y) \right| = \left| \int_{\bigcup_{\mathfrak{p} \in \mathfrak{T}(u)} \mathcal{I}(\mathfrak{p})} \bar{g}(y) \sum_{\mathfrak{p} \in \mathfrak{T}(u)} T_{\mathfrak{p}}f(y) d\mu(y) \right|.$$

Since $\mathcal{L}(\mathfrak{T}(u))$ partitions $\bigcup_{\mathfrak{p} \in \mathfrak{T}(u)} \mathcal{I}(\mathfrak{p})$ by Lemma 7.2, we get from the triangle inequality

$$\leq \sum_{L \in \mathcal{L}(\mathfrak{T}(u))} \int_L |g(y)| \left| \sum_{\mathfrak{p} \in \mathfrak{T}(u)} T_{\mathfrak{p}}f(y) \right| d\mu(y)$$

which by the above computation is bounded by

$$\begin{aligned}
&2^{151a^3+1} \sum_{L \in \mathcal{L}(\mathfrak{T}(u))} \int_L P_{\mathcal{L}(\mathfrak{T}(u))}|g|(y) \times \\
&\quad ((M_{\mathcal{B},1} + S_{1,u})P_{\mathcal{J}(\mathfrak{T}(u))}|f|(y) + |T_{\mathcal{N}}P_{\mathcal{J}(\mathfrak{T}(u))}f(y)|) d\mu(y) \\
&= 2^{151a^3+1} \int_X P_{\mathcal{L}(\mathfrak{T}(u))}|g|(y) \times \\
&\quad ((M_{\mathcal{B},1} + S_{1,u})P_{\mathcal{J}(\mathfrak{T}(u))}|f|(y) + |T_{\mathcal{N}}P_{\mathcal{J}(\mathfrak{T}(u))}f(y)|) d\mu(y).
\end{aligned}$$

Applying Cauchy-Schwarz and Minkowski's inequality, this is bounded by

$$\begin{aligned}
&2^{151a^3+1} \|P_{\mathcal{L}(\mathfrak{T}(u))}|g|\|_2 \times \\
&\quad (\|M_{\mathcal{B},1}P_{\mathcal{J}(\mathfrak{T}(u))}|f|\|_2 + \|S_{1,u}P_{\mathcal{J}(\mathfrak{T}(u))}|f|\|_2 + \|T_{\mathcal{N}}P_{\mathcal{J}(\mathfrak{T}(u))}f(y)\|_2).
\end{aligned}$$

By Proposition 2.6, Lemma 7.8 and Lemma 7.9, the second factor is at most

$$(2^{2a+1} + 2^{12a})\|P_{\mathcal{J}(\mathfrak{T}(u))}|f|\|_2 + 2^{103a^3}\|P_{\mathcal{J}(\mathfrak{T}(u))}f\|_2.$$

By the triangle inequality we have for all $x \in X$ that $|P_{\mathcal{J}(\mathfrak{T}(u))}f|(x) \leq P_{\mathcal{J}(\mathfrak{T}(u))}|f|(x)$, thus we can further estimate the above by

$$(2^{2a+1} + 2^{12a} + 2^{103a^3})\|P_{\mathcal{J}(\mathfrak{T}(u))}|f|\|_2.$$

This completes the proof since $a \geq 4$. \square

Now we prove the two auxiliary lemmas. We begin with the nontangential maximal operator $T_{\mathcal{N}}$.

Proof of Lemma 7.8. Fix s_1, s_2 . By (2.4) we have for all $x \in (0, \infty)$

$$\sum_{s=s_1}^{s_2} \psi(D^{-s}x) = 1 - \sum_{s < s_1} \psi(D^{-s}x) - \sum_{s > s_2} \psi(D^{-s}x).$$

Since ψ is supported in $[\frac{1}{4D}, \frac{1}{2}]$, the two sums on the right hand side are zero for all $x \in [\frac{1}{2}D^{s_1-1}, \frac{1}{4}D^{s_2-1}]$, hence

$$x \in [\frac{1}{2}D^{s_1-1}, \frac{1}{4}D^{s_2}] \implies \sum_{s=s_1}^{s_2} \psi(D^{-s}x) = 1.$$

Since ψ is supported in $[\frac{1}{4D}, \frac{1}{2}]$, we further have

$$x \notin [\frac{1}{4}D^{s_1-1}, \frac{1}{2}D^{s_2}] \implies \sum_{s=s_1}^{s_2} \psi(D^{-s}x) = 0.$$

Finally, since $\psi \geq 0$ and $\sum_{s \in \mathbb{Z}} \psi(D^{-s}x) = 1$, we have for all x

$$0 \leq \sum_{s=s_1}^{s_2} \psi(D^{-s}x) \leq 1.$$

Let $x' \in I_{s_1}(x)$. By the triangle inequality and (2.10), it holds that $\rho(x, x') \leq 8D^{s_1}$. We have

$$\begin{aligned} & \left| \sum_{s=s_1}^{s_2} \int K_s(x', y) f(y) d\mu(y) \right| \\ &= \left| \int \sum_{s=s_1}^{s_2} \psi(D^{-s}\rho(x', y)) K(x', y) f(y) d\mu(y) \right| \\ &\leq \left| \int_{8D^{s_1} \leq \rho(x', y) \leq \frac{1}{4}D^{s_2}} K(x', y) f(y) d\mu(y) \right| \end{aligned} \quad (7.13)$$

$$+ \int_{\frac{1}{4}D^{s_1-1} \leq \rho(x', y) \leq 8D^{s_1}} |K(x', y)| |f(y)| d\mu(y) \quad (7.14)$$

$$+ \int_{\frac{1}{4}D^{s_2} \leq \rho(x', y) \leq \frac{1}{2}D^{s_2}} |K(x', y)| |f(y)| d\mu(y). \quad (7.15)$$

The first term (7.13) is at most $T_* f(x)$.

The other two terms will be estimated by the finitary maximal function from Proposition 2.6. For the second term (7.14) we use (1.14) which implies that for all y with $\rho(x', y) \geq \frac{1}{4}D^{s_1-1}$, we have

$$|K(x', y)| \leq \frac{2^{a^3}}{\mu(B(x', \frac{1}{4}D^{s_1-1}))}.$$

Using $D = 2^{100a^2}$ and the doubling property (1.5) $6 + 100a^2$ times estimates the last display by

$$\leq \frac{2^{6a+101a^3}}{\mu(B(x', 16D^{s_1}))}. \quad (7.16)$$

By the triangle inequality and (2.10), we have

$$B(x', 8D^{s_1}) \subset B(c(I_{s_1}(x)), 16D^{s(I_{s_1}(x))}).$$

Combining this with (7.16), we conclude that (7.14) is at most

$$2^{6a+101a^3} M_{\mathcal{B},1} f(x).$$

For (7.15) we argue similarly. We have for all y with $\rho(x', y) \geq \frac{1}{4} D^{s_2}$

$$|K(x', y)| \leq \frac{2^{a^3}}{\mu(B(x', \frac{1}{4} D^{s_2}))}.$$

Using the doubling property (1.5) 6 times estimates the last display by

$$\leq \frac{2^{6a+a^3}}{\mu(B(x', 16D^{s_2}))}. \quad (7.17)$$

Note that by (2.8) we have $I_{s_1}(x) \subset I_{s_2}(x)$, in particular $x' \in I_{s_2}(x)$. By the triangle inequality and (2.10), we have

$$B(x', 8D^{s_2}) \subset B(c(I_{s_2}(x)), 16D^{s(I_{s_2}(x))}).$$

Combining this, (7.15) is at most

$$2^{6a+a^3} M_{\mathcal{B},1} f(x).$$

Using $a \geq 4$, taking a supremum over all $x' \in I_{s_1}(x)$ and then a supremum over all $-S \leq s_1 < s_2 \leq S$, we obtain

$$T_{\mathcal{N}} f(x) \leq T_* f(x) + 2^{102a^3} M_{\mathcal{B},1} f(x).$$

The lemma now follows from assumption (1.18), Proposition 2.6 and $a \geq 4$. \square

We need the following lemma to prepare the L^2 -estimate for the auxiliary operators $S_{1,u}$.

Lemma 7.10 (boundary overlap). *For every cube $I \in \mathcal{D}$, there exist at most 2^{8a} cubes $J \in \mathcal{D}$ with $s(J) = s(I)$ and $B(c(I), 16D^{s(I)}) \cap B(c(J), 16D^{s(J)}) \neq \emptyset$.*

Proof. Suppose that $B(c(I), 16D^{s(I)}) \cap B(c(J), 16D^{s(J)}) \neq \emptyset$ and $s(I) = s(J)$. Then $B(c(I), 32D^{s(I)}) \subset B(c(J), 64D^{s(J)})$. Hence by the doubling property (1.5)

$$2^{8a} \mu(B(c(J), \frac{1}{4} D^{s(J)})) \geq \mu(B(c(I), 32D^{s(I)})),$$

and by the triangle inequality, the ball $B(c(J), \frac{1}{4} D^{s(J)})$ is contained in $B(c(I), 32D^{s(I)})$.

If \mathcal{C} is any finite collection of cubes $J \in \mathcal{D}$ satisfying $s(J) = s(I)$ and

$$B(c(I), 16D^{s(I)}) \cap B(c(J), 16D^{s(J)}) \neq \emptyset,$$

then it follows from (2.10) and pairwise disjointness of cubes of the same scale (2.8) that the balls $B(c(J), \frac{1}{4} D^{s(J)})$ are pairwise disjoint. Hence

$$\begin{aligned} \mu(B(c(I), 32D^{s(I)})) &\geq \sum_{J \in \mathcal{C}} \mu(B(c(J), \frac{1}{4} D^{s(J)})) \\ &\geq |\mathcal{C}| 2^{-8a} \mu(B(c(I), 32D^{s(I)})). \end{aligned}$$

Since μ is doubling and $\mu \neq 0$, we have $\mu(B(c(I), 32D^{s(I)})) > 0$. The lemma follows after dividing by $2^{-8a} \mu(B(c(I), 32D^{s(I)}))$. \square

Now we can bound the operators $S_{1,u}$.

Proof of Lemma 7.9. Note that by definition, $S_{1,u}f$ is a finite sum of indicator functions of cubes $I \in \mathcal{D}$ for each locally integrable f , and hence is bounded, has bounded support and is integrable. Let g be another function with the same three properties. Then $\bar{g}S_{1,u}f$ is integrable, and we have

$$\begin{aligned}
& \left| \int \bar{g}(y) S_{1,u}f(y) \, d\mu(y) \right| \\
&= \left| \sum_{I \in \mathcal{D}} \frac{1}{\mu(B(c(I), 16D^{s(I)}))} \int_I \bar{g}(y) \, d\mu(y) \right. \\
&\quad \times \left. \sum_{J \in \mathcal{J} : J \subseteq B(c(I), 16D^{s(I)})} D^{(s(J)-s(I))/a} \int_J |f(y)| \, d\mu(y) \right| \\
&\leq \sum_{I \in \mathcal{D}} \frac{1}{\mu(B(c(I), 16D^{s(I)}))} \int_{B(c(I), 16D^{s(I)})} |g(y)| \, d\mu(y) \\
&\quad \times \sum_{J \in \mathcal{J} : J \subseteq B(c(I), 16D^{s(I)})} D^{(s(J)-s(I))/a} \int_J |f(y)| \, d\mu(y).
\end{aligned}$$

Changing the order of summation and using $J \subset B(c(I), 16D^{s(I)})$ to bound the first average integral by $M_{\mathcal{B},1}|g|(y)$ for any $y \in J$, we obtain

$$\leq \sum_{J \in \mathcal{J}} \int_J |f(y)| M_{\mathcal{B},1}|g|(y) \, d\mu(y) \sum_{I \in \mathcal{D} : J \subset B(c(I), 16D^{s(I)})} D^{(s(J)-s(I))/a}. \quad (7.18)$$

By (2.10) and (2.1) the condition $J \subset B(c(I), 16D^{s(I)})$ implies $s(I) \geq s(J)$. By Lemma 7.10, there are at most 2^{8a} cubes I at each scale with $J \subset B(c(I), D^{s(I)})$. By convexity of $t \mapsto D^t$ and since $D \geq 2$, we have for all $-1 \leq t \leq 0$

$$D^t \leq 1 + t \left(1 - \frac{1}{D} \right) \leq 1 + \frac{1}{2}t,$$

so $(1 - D^{-1/a})^{-1} \leq 2a \leq 2^a$. Using this estimate for the sum of the geometric series, we conclude that (7.18) is at most

$$2^{9a} \sum_{J \in \mathcal{J}} \int_J |f(y)| M_{\mathcal{B},1}|g|(y) \, d\mu(y).$$

The collection \mathcal{J} is a partition of X , so this equals

$$2^{9a} \int_X |f(y)| M_{\mathcal{B},1}|g|(y) \, d\mu(y).$$

Using Cauchy-Schwarz and Proposition 2.6 we conclude

$$\left| \int \bar{g} S_{1,u}f \, d\mu \right| \leq 2^{11a+1} \|g\|_2 \|f\|_2.$$

The lemma now follows by choosing $g = S_{1,u}f$ and dividing on both sides by the finite $\|S_{1,u}f\|_2$. \square

7.3. The quantitative L^2 tree estimate. The main result of this subsection is the following quantitative bound for operators associated to trees, with decay in the densities dens_1 and dens_2 .

Lemma 7.11 (densities tree bound). *Let $\mathbf{u} \in \mathfrak{U}$. Then for all f, g bounded with bounded support*

$$\left| \int_X \bar{g} \sum_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathbf{p}} f \, d\mu \right| \leq 2^{155a^3} \text{dens}_1(\mathfrak{T}(\mathbf{u}))^{1/2} \|f\|_2 \|g\|_2. \quad (7.19)$$

If $|f| \leq \mathbf{1}_F$, then we have

$$\left| \int_X \bar{g} \sum_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathbf{p}} f \, d\mu \right| \leq 2^{256a^3} \text{dens}_1(\mathfrak{T}(\mathbf{u}))^{1/2} \text{dens}_2(\mathfrak{T}(\mathbf{u}))^{1/2} \|f\|_2 \|g\|_2. \quad (7.20)$$

Below, we deduce this lemma from Lemma 7.7 and the following two estimates controlling the size of support of the operator and its adjoint.

Lemma 7.12 (local dens_1 tree bound). *Let $\mathbf{u} \in \mathfrak{U}$ and $L \in \mathcal{L}(\mathfrak{T}(\mathbf{u}))$. Then*

$$\mu(L \cap \bigcup_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} E(\mathbf{p})) \leq 2^{101a^3} \text{dens}_1(\mathfrak{T}(\mathbf{u})) \mu(L). \quad (7.21)$$

Lemma 7.13 (local dens_2 tree bound). *Let $J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}))$ be such that there exist $\mathbf{q} \in \mathfrak{T}(\mathbf{u})$ with $J \cap \mathcal{I}(\mathbf{q}) \neq \emptyset$. Then*

$$\mu(F \cap J) \leq 2^{200a^3+19} \text{dens}_2(\mathfrak{T}(\mathbf{u})).$$

Proof of Lemma 7.11. Denote

$$\mathcal{E}(\mathbf{u}) = \bigcup_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} E(\mathbf{p}).$$

Then we have

$$\left| \int_X \bar{g} \sum_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathbf{p}} f \, d\mu \right| = \left| \int_X \overline{g \mathbf{1}_{\mathcal{E}(\mathbf{u})}} \sum_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathbf{p}} f \, d\mu \right|.$$

By Lemma 7.7, this is bounded by

$$\leq 2^{104a^3} \|P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} |f|\|_2 \|P_{\mathcal{L}(\mathfrak{T}(\mathbf{u}))} \mathbf{1}_{\mathcal{E}(\mathbf{u})} g\|_2. \quad (7.22)$$

We bound the two factors separately. We have

$$\|P_{\mathcal{L}(\mathfrak{T}(\mathbf{u}))} \mathbf{1}_{\mathcal{E}(\mathbf{u})} g\|_2 = \left(\sum_{L \in \mathcal{L}(\mathfrak{T}(\mathbf{u}))} \frac{1}{\mu(L)} \left(\int_{L \cap \mathcal{E}(\mathbf{u})} |g(y)| \, d\mu(y) \right)^2 \right)^{1/2}.$$

By Cauchy-Schwarz and Lemma 7.12 this is at most

$$\leq \left(\sum_{L \in \mathcal{L}(\mathfrak{T}(\mathbf{u}))} 2^{101a^3} \text{dens}_1(\mathfrak{T}(\mathbf{u})) \int_{L \cap \mathcal{E}(\mathbf{u})} |g(y)|^2 \, d\mu(y) \right)^{1/2}.$$

Since cubes $L \in \mathcal{L}(\mathfrak{T}(\mathbf{u}))$ are pairwise disjoint by Lemma 7.2, this is

$$\leq 2^{51a^3} \text{dens}_1(\mathfrak{T}(\mathbf{u}))^{1/2} \|g\|_2. \quad (7.23)$$

Similarly, we have

$$\|P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))}|f|\|_2 = \left(\sum_{J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}))} \frac{1}{\mu(J)} \left(\int_J |f(y)| d\mu(y) \right)^2 \right)^{1/2}. \quad (7.24)$$

By Cauchy-Schwarz, this is

$$\leq \left(\sum_{J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}))} \int_J |f(y)|^2 d\mu(y) \right)^{1/2}.$$

Since cubes in $\mathcal{J}(\mathfrak{T}(\mathbf{u}))$ are pairwise disjoint by Lemma 7.2, this at most

$$\|f\|_2. \quad (7.25)$$

Combining (7.22), (7.23) and (7.25) and using $a \geq 4$ gives (7.19).

If $f \leq \mathbf{1}_F$ then $f = f\mathbf{1}_F$, so

$$\left(\sum_{J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}))} \int_J |f(y)|^2 d\mu(y) \right)^{1/2} = \left(\sum_{J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}))} \int_{J \cap F} |f(y)|^2 d\mu(y) \right)^{1/2}.$$

We estimate as before, using now Lemma 7.13 and Cauchy-Schwarz, and obtain that this is

$$\leq 2^{100a^3+10} \text{dens}_2(\mathfrak{T}(\mathbf{u}))^{1/2} \|f\|_2.$$

Combining this with (7.22), (7.23) and $a \geq 4$ gives (7.20). \square

Now we prove the two auxiliary estimates.

Proof of Lemma 7.12. If the set on the right hand side is empty, then (7.21) holds. If not, then there exists $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$ with $L \cap \mathcal{I}(\mathbf{p}) \neq \emptyset$.

Suppose first that there exists such \mathbf{p} with $s(\mathbf{p}) \leq s(L)$. Then by (2.8) $\mathcal{I}(\mathbf{p}) \subset L$, which gives by the definition of $\mathcal{L}(\mathfrak{T}(\mathbf{u}))$ that $s(L) = -S$ and hence $L = \mathcal{I}(\mathbf{p})$. Let $\mathbf{q} \in \mathfrak{T}(\mathbf{u})$ with $E(\mathbf{q}) \cap L \neq \emptyset$. Since $s(L) = -S \leq s(\mathbf{q})$ it follows from (2.8) that $\mathcal{I}(\mathbf{p}) = L \subset \mathcal{I}(\mathbf{q})$. We have then by Lemma 2.8

$$\begin{aligned} d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{q})) &\leq d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{u})) + d_{\mathbf{p}}(\mathcal{Q}(\mathbf{q}), \mathcal{Q}(\mathbf{u})) \\ &\leq d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{u})) + d_{\mathbf{q}}(\mathcal{Q}(\mathbf{q}), \mathcal{Q}(\mathbf{u})). \end{aligned}$$

Using that $\mathbf{p}, \mathbf{q} \in \mathfrak{T}(\mathbf{u})$ and (2.32), this is at most 8. Using again the triangle inequality and Lemma 2.8, we obtain that for each $q \in B_{\mathbf{q}}(\mathcal{Q}(\mathbf{q}), 1)$

$$d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), q) \leq d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{q})) + d_{\mathbf{q}}(\mathcal{Q}(\mathbf{q}), q) \leq 9.$$

Thus $L \cap E(\mathbf{q}) \subset E_2(9, \mathbf{p})$. We obtain

$$\mu(L \cap \bigcup_{\mathbf{q} \in \mathfrak{T}(\mathbf{u})} E(\mathbf{q})) \leq \mu(E_2(9, \mathbf{p})).$$

By the definition of dens_1 , this is bounded by

$$9^a \text{dens}_1(\mathfrak{T}(\mathbf{u})) \mu(\mathcal{I}(\mathbf{p})) = 9^a \text{dens}_1(\mathfrak{T}(\mathbf{u})) \mu(L).$$

Since $a \geq 4$, (7.21) follows in this case.

Now suppose that for each $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$ with $L \cap E(\mathbf{p}) \neq \emptyset$, we have $s(\mathbf{p}) > s(L)$. Since there exists at least one such \mathbf{p} , there exists in particular at least one cube $L'' \in \mathcal{D}$ with $L \subset L''$ and $s(L'') > s(L)$. By (2.7), there

exists $L' \in \mathcal{D}$ with $L \subset L'$ and $s(L') = s(L) + 1$. By the definition of $\mathcal{L}(\mathfrak{T}(u))$ there exists a tile $\mathfrak{p}'' \in \mathfrak{T}(u)$ with $\mathcal{I}(\mathfrak{p}'') \subset L'$. Let \mathfrak{p}' be the unique tile such that $\mathcal{I}(\mathfrak{p}') = L'$ and such that $\Omega(u) \cap \Omega(\mathfrak{p}') \neq \emptyset$. Since by (2.32) $s(\mathfrak{p}') = s(L') \leq s(\mathfrak{p}) < s(u)$, we have by (2.8) and (2.14) that $\Omega(u) \subset \Omega(\mathfrak{p}')$. Let $\mathfrak{q} \in \mathfrak{T}(u)$ with $L \cap E(\mathfrak{q}) \neq \emptyset$. As shown above, this implies $s(\mathfrak{q}) \geq s(L')$, so by (2.8) $L' \subset \mathcal{I}(\mathfrak{q})$. If $q \in B_q(\mathcal{Q}(\mathfrak{q}), 1)$, then by a similar calculation as above, using the triangle inequality, Lemma 2.8 and (2.32), we obtain

$$d_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), q) \leq d_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), \mathcal{Q}(\mathfrak{q})) + d_q(\mathcal{Q}(\mathfrak{q}), q) \leq 6.$$

Thus $L \cap E(\mathfrak{q}) \subset E_2(6, \mathfrak{p}')$. Since $\mathcal{I}(\mathfrak{p}'') \subset \mathcal{I}(\mathfrak{p}') \subset \mathcal{I}(\mathfrak{p})$ and $\mathfrak{p}'', \mathfrak{p} \in \mathfrak{T}(u)$, we have $\mathfrak{p}' \in \mathfrak{P}(\mathfrak{T}(u))$. We deduce using the definition (2.28) of dens_1

$$\mu(L \cap \bigcup_{\mathfrak{q} \in \mathfrak{T}(u)} E(\mathfrak{q})) \leq \mu(E_2(6, \mathfrak{p}')) \leq 6^a \text{dens}_1(\mathfrak{T}(u)) \mu(L').$$

Using the doubling property (1.5), (2.10), and $a \geq 4$ this is estimated by

$$6^a 2^{100a^3+5} \text{dens}_1(\mathfrak{T}(u)) \mu(L) \leq 2^{101a^3} \text{dens}_1(\mathfrak{T}(u)) \mu(L).$$

This completes the proof. \square

Proof of Lemma 7.13. Suppose first that there exists a tile $\mathfrak{p} \in \mathfrak{T}(u)$ with $\mathcal{I}(\mathfrak{p}) \subset B(c(J), 100D^{s(J)+1})$. By the definition of $\mathcal{J}(\mathfrak{T}(u))$, this implies that $s(J) = -S$, and in particular $s(\mathfrak{p}) \geq s(J)$. Using the triangle inequality and (2.10) it follows that $J \subset B(c(\mathfrak{p}), 200D^{s(\mathfrak{p})+1})$. From the doubling property (1.5), $D = 2^{100a^2}$ and (2.10), we obtain

$$\mu(\mathcal{I}(\mathfrak{p})) \leq 2^{100a^3+9} \mu(J)$$

and hence

$$\mu(B(c(\mathfrak{p}), 200D^{s(\mathfrak{p})+1})) \leq 2^{200a^3+19} \mu(J).$$

With the definition (2.29) of dens_2 it follows that

$$\begin{aligned} \mu(J \cap F) &\leq \mu(B(c(\mathfrak{p}), 200D^{s(\mathfrak{p})+1}) \cap F) \\ &\leq \text{dens}_2(\mathfrak{T}(u)) \mu(B(c(\mathfrak{p}), 200D^{s(\mathfrak{p})+1})) \leq 2^{200a^3+19} \text{dens}_2(\mathfrak{T}(u)) \mu(J), \end{aligned}$$

completing the proof in this case.

Now suppose that there does not exist a tile $\mathfrak{p} \in \mathfrak{T}(u)$ with $\mathcal{I}(\mathfrak{p}) \subset B(c(J), 100D^{s(J)+1})$. If we had $s(\mathfrak{q}) \leq s(J)$, then by (2.8) and (2.10) $\mathcal{I}(\mathfrak{q}) \subset J \subset B(c(J), 100D^{s(J)+1})$, contradicting our assumption. Thus $s(\mathfrak{q}) > s(J)$. Then, by (2.7) and (2.8), there exists some cube $J' \in \mathcal{D}$ with $s(J') = s(J) + 1$ and $J \subset J'$. By definition of $\mathcal{J}(\mathfrak{T}(u))$ there exists some $\mathfrak{p} \in \mathfrak{T}(u)$ such that $\mathcal{I}(\mathfrak{p}) \subset B(c(J'), 100D^{s(J')+1})$. From the doubling property (1.5), $D = 2^{100a^2}$ and (2.10), we obtain

$$\mu(B(c(\mathfrak{p}), 4D^{s(\mathfrak{p})})) \leq 2^{4a} \mu(\mathcal{I}(\mathfrak{p})) \leq 2^{200a^3+14} \mu(J). \quad (7.26)$$

If $J \subset B(c(\mathfrak{p}), 4D^{s(\mathfrak{p})})$, then we bound

$$\mu(J \cap F) \leq \mu(B(c(\mathfrak{p}), 4D^{s(\mathfrak{p})}) \cap F)$$

and use the definition (2.29)

$$\leq \text{dens}_2(\mathfrak{T}(u)) \mu(B(c(\mathfrak{p}), 4D^{s(\mathfrak{p})})) \leq 2^{200a^3+14} \mu(J).$$

From now on we assume $J \not\subset B(c(\mathfrak{p}), 4D^{s(\mathfrak{p})})$. Since

$$c(\mathfrak{p}) \in \mathcal{I}(\mathfrak{p}) \subset B(c(J'), 100D^{s(J')+1}),$$

we have by (2.10) and the triangle inequality

$$J \subset J' \subset B(c(J'), 4D^{s(J')}) \subset B(c(\mathfrak{p}), 104D^{s(J')+1}).$$

In particular this implies $104D^{s(J')+1} > 4D^{s(\mathfrak{p})}$. By the triangle inequality we also have

$$B(c(\mathfrak{p}), 104D^{s(J')+1}) \subset B(c(J), 204D^{s(J')+1}),$$

so from the doubling property (1.5)

$$\mu(B(c(\mathfrak{p}), 104D^{s(J')+1})) \leq 2^{200a^3+10}\mu(J).$$

From here one completes the proof as in the other cases. \square

7.4. Almost orthogonality of separated trees. The main result of this subsection is the almost orthogonality estimate for operators associated to distinct trees in a forest in Lemma 7.17 below. We will deduce it from Lemmas 7.18 and 7.19, which are proven in Subsections 7.5 and 7.6, respectively. Before stating it, we introduce some relevant notation.

The adjoint of the operator $T_{\mathfrak{p}}$ defined in (2.21) is given by

$$T_{\mathfrak{p}}^*g(x) = \int_{E(\mathfrak{p})} \overline{K_{s(\mathfrak{p})}(y, x)} e(-Q(y)(x) + Q(y)(y))g(y) d\mu(y). \quad (7.27)$$

Lemma 7.14 (adjoint tile support). *For each $\mathfrak{p} \in \mathfrak{P}$, we have*

$$T_{\mathfrak{p}}^*g = \mathbf{1}_{B(c(\mathfrak{p}), 5D^{s(\mathfrak{p})})} T_{\mathfrak{p}}^* \mathbf{1}_{\mathcal{I}(\mathfrak{p})} g.$$

For each $\mathfrak{u} \in \mathfrak{U}$ and each $\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})$, we have

$$T_{\mathfrak{p}}^*g = \mathbf{1}_{\mathcal{I}(\mathfrak{u})} T_{\mathfrak{p}}^* \mathbf{1}_{\mathcal{I}(\mathfrak{u})} g.$$

Proof. By (2.32), $E(\mathfrak{p}) \subset \mathcal{I}(\mathfrak{p}) \subset \mathcal{I}(\mathfrak{u})$. Thus by (7.27)

$$\begin{aligned} T_{\mathfrak{p}}^*g(x) &= T_{\mathfrak{p}}^*(\mathbf{1}_{\mathcal{I}(\mathfrak{p})}g)(x) \\ &= \int_{E(\mathfrak{p})} \overline{K_{s(\mathfrak{p})}(y, x)} e(-Q(y)(x) + Q(y)(y)) \mathbf{1}_{\mathcal{I}(\mathfrak{p})}(y)g(y) d\mu(y). \end{aligned}$$

If this integral is not 0, then there exists $y \in \mathcal{I}(\mathfrak{p})$ such that $K_{s(\mathfrak{p})}(y, x) \neq 0$. By (2.48), (2.10) and the triangle inequality, it follows that

$$x \in B(c(\mathfrak{p}), 5D^{s(\mathfrak{p})}).$$

Thus

$$T_{\mathfrak{p}}^*g(x) = \mathbf{1}_{B(c(\mathfrak{p}), 5D^{s(\mathfrak{p})})}(x) T_{\mathfrak{p}}^*(\mathbf{1}_{\mathcal{I}(\mathfrak{p})}g)(x).$$

The second claimed equation follows now since $\mathcal{I}(\mathfrak{p}) \subset \mathcal{I}(\mathfrak{u})$ and by (2.37) $B(c(\mathfrak{p}), 5D^{s(\mathfrak{p})}) \subset \mathcal{I}(\mathfrak{u})$. \square

Lemma 7.15 (adjoint tree estimate). *For all bounded g with bounded support, we have that*

$$\left\| \sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} T_{\mathfrak{p}}^*g \right\|_2 \leq 2^{155a^3} \text{dens}_1(\mathfrak{T}(\mathfrak{u}))^{1/2} \|g\|_2.$$

Proof. By Cauchy-Schwarz and Lemma 7.11, we have for all bounded f, g with bounded support that

$$\begin{aligned} \left| \int_X \overline{\sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} T_{\mathfrak{p}}^* g f} d\mu \right| &= \left| \int_X g \sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} T_{\mathfrak{p}} f d\mu \right| \\ &\leq 2^{155a^3} \text{dens}_1(\mathfrak{T}(\mathfrak{u}))^{1/2} \|g\|_2 \|f\|_2. \end{aligned} \quad (7.28)$$

Let $f = \sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} T_{\mathfrak{p}}^* g$. If g is bounded and has bounded support, then the same is true for f . In particular $\|f\|_2 < \infty$. Dividing (7.28) by $\|f\|_2$ completes the proof. \square

We define

$$S_{2,\mathfrak{u}}g := \left| \sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} T_{\mathfrak{p}}^* g \right| + M_{\mathcal{B},1}g + |g|.$$

Lemma 7.16 (adjoint tree control). *We have for all bounded g with bounded support*

$$\|S_{2,\mathfrak{u}}g\|_2 \leq 2^{156a^3} \|g\|_2.$$

Proof. This follows immediately from Minkowski's inequality, Proposition 2.6 and Lemma 7.15, using that $a \geq 4$. \square

Now we are ready to state the main result of this subsection.

Lemma 7.17 (correlation separated trees). *For any $\mathfrak{u}_1 \neq \mathfrak{u}_2 \in \mathfrak{U}$ and all bounded g_1, g_2 with bounded support, we have*

$$\left| \int_X \sum_{\mathfrak{p}_1 \in \mathfrak{T}(\mathfrak{u}_1)} \sum_{\mathfrak{p}_2 \in \mathfrak{T}(\mathfrak{u}_2)} T_{\mathfrak{p}_1}^* g_1 \overline{T_{\mathfrak{p}_2}^* g_2} d\mu \right| \quad (7.29)$$

$$\leq 2^{550a^3-3n} \prod_{j=1}^2 \|S_{2,\mathfrak{u}_j} g_j\|_{L^2(\mathcal{I}(\mathfrak{u}_1) \cap \mathcal{I}(\mathfrak{u}_2))}. \quad (7.30)$$

Proof of Lemma 7.17. By Lemma 7.14 and (2.8), the left hand side (7.29) is 0 unless $\mathcal{I}(\mathfrak{u}_1) \subset \mathcal{I}(\mathfrak{u}_2)$ or $\mathcal{I}(\mathfrak{u}_2) \subset \mathcal{I}(\mathfrak{u}_1)$. Without loss of generality we assume that $\mathcal{I}(\mathfrak{u}_1) \subset \mathcal{I}(\mathfrak{u}_2)$.

Define

$$\mathfrak{S} := \{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u}_1) \cup \mathfrak{T}(\mathfrak{u}_2) : d_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{u}_1), \mathcal{Q}(\mathfrak{u}_2)) \geq 2^{Zn/2}\}. \quad (7.31)$$

Lemma 7.17 follows by combining the definition (2.3) of Z with the following two lemmas.

Lemma 7.18 (correlation distant tree parts). *We have for all $\mathfrak{u}_1 \neq \mathfrak{u}_2 \in \mathfrak{U}$ with $\mathcal{I}(\mathfrak{u}_1) \subset \mathcal{I}(\mathfrak{u}_2)$ and all bounded g_1, g_2 with bounded support*

$$\left| \int_X \sum_{\mathfrak{p}_1 \in \mathfrak{T}(\mathfrak{u}_1)} \sum_{\mathfrak{p}_2 \in \mathfrak{T}(\mathfrak{u}_2) \cap \mathfrak{S}} T_{\mathfrak{p}_1}^* g_1 \overline{T_{\mathfrak{p}_2}^* g_2} d\mu \right| \quad (7.32)$$

$$\leq 2^{541a^3} 2^{-Zn/(4a^2+2a^3)} \prod_{j=1}^2 \|S_{2,\mathfrak{u}_j} g_j\|_{L^2(\mathcal{I}(\mathfrak{u}_1))}. \quad (7.33)$$

Lemma 7.19 (correlation near tree parts). *We have for all $u_1 \neq u_2 \in \mathfrak{U}$ with $\mathcal{I}(u_1) \subset \mathcal{I}(u_2)$ and all bounded g_1, g_2 with bounded support*

$$\left| \int_X \sum_{\mathfrak{p}_1 \in \mathfrak{T}(u_1)} \sum_{\mathfrak{p}_2 \in \mathfrak{T}(u_2) \setminus \mathfrak{S}} T_{\mathfrak{p}_1}^* g_1 \overline{T_{\mathfrak{p}_2}^* g_2} d\mu \right| \quad (7.34)$$

$$\leq 2^{222a^3} 2^{-Zn2^{-10a}} \prod_{j=1}^2 \|S_{2,u_j} g_j\|_{L^2(\mathcal{I}(u_1))}. \quad (7.35)$$

□

In the proofs of both lemmas, we will need the following observation.

Lemma 7.20 (overlap implies distance). *Let $u_1 \neq u_2 \in \mathfrak{U}$ with $\mathcal{I}(u_1) \subset \mathcal{I}(u_2)$. If $\mathfrak{p} \in \mathfrak{T}(u_1) \cup \mathfrak{T}(u_2)$ with $\mathcal{I}(\mathfrak{p}) \cap \mathcal{I}(u_1) \neq \emptyset$, then $\mathfrak{p} \in \mathfrak{S}$. In particular, we have $\mathfrak{T}(u_1) \subset \mathfrak{S}$.*

Proof. Suppose first that $\mathfrak{p} \in \mathfrak{T}(u_1)$. Then $\mathcal{I}(\mathfrak{p}) \subset \mathcal{I}(u_1) \subset \mathcal{I}(u_2)$, by (2.32). Thus we have by the separation condition (2.36), (2.15), (2.32) and the triangle inequality

$$\begin{aligned} d_{\mathfrak{p}}(\mathcal{Q}(u_1), \mathcal{Q}(u_2)) &\geq d_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), \mathcal{Q}(u_2)) - d_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), \mathcal{Q}(u_1)) \\ &\geq 2^{Z(n+1)} - 4 \\ &\geq 2^{Zn}, \end{aligned}$$

using that $Z = 2^{12a} \geq 4$. Hence $\mathfrak{p} \in \mathfrak{S}$.

Suppose now that $\mathfrak{p} \in \mathfrak{T}(u_2)$. If $\mathcal{I}(\mathfrak{p}) \subset \mathcal{I}(u_1)$, then the same argument as above with u_1 and u_2 swapped shows $\mathfrak{p} \in \mathfrak{S}$. If $\mathcal{I}(\mathfrak{p}) \not\subset \mathcal{I}(u_1)$ then, by (2.8), $\mathcal{I}(u_1) \subset \mathcal{I}(\mathfrak{p})$. Pick $\mathfrak{p}' \in \mathfrak{T}(u_1)$, we have $\mathcal{I}(\mathfrak{p}') \subset \mathcal{I}(u_1) \subset \mathcal{I}(\mathfrak{p})$. Hence, by Lemma 2.8 and the first paragraph

$$d_{\mathfrak{p}}(\mathcal{Q}(u_1), \mathcal{Q}(u_2)) \geq d_{\mathfrak{p}'}(\mathcal{Q}(u_1), \mathcal{Q}(u_2)) \geq 2^{Zn},$$

so $\mathfrak{p} \in \mathfrak{S}$. □

To simplify the notation, we will write at various places throughout the proof of Lemmas 7.18 and 7.19 for a subset $\mathfrak{C} \subset \mathfrak{P}$

$$T_{\mathfrak{C}} f := \sum_{\mathfrak{p} \in \mathfrak{C}} T_{\mathfrak{p}} f, \quad T_{\mathfrak{C}}^* g := \sum_{\mathfrak{p} \in \mathfrak{C}} T_{\mathfrak{p}}^* g.$$

7.5. Proof of Lemma 7.18: Tiles with large separation. Lemma 7.18 follows from the van der Corput estimate in Proposition 2.5. We apply this proposition in Subsubsection 7.5.3. To prepare this application, we first, in Subsubsection 7.5.1, construct a suitable partition of unity, and show then, in Subsubsection 7.5.2 the Hölder estimates needed to apply Proposition 2.5.

7.5.1. A partition of unity. Define

$$\mathcal{J}' = \{J \in \mathcal{J}(\mathfrak{S}) : J \subset \mathcal{I}(u_1)\}.$$

Lemma 7.21 (dyadic partition 1). *We have that*

$$\mathcal{I}(u_1) = \dot{\bigcup}_{J \in \mathcal{J}'} J.$$

Proof. By Lemma 7.2, it remains only to show that each $J \in \mathcal{J}(\mathfrak{S})$ with $J \cap \mathcal{I}(\mathfrak{u}_1) \neq \emptyset$ is in \mathcal{J}' . But if $J \notin \mathcal{J}'$, then by (2.8) $\mathcal{I}(\mathfrak{u}_1) \subsetneq J$. Pick $\mathfrak{p} \in \mathfrak{T}(\mathfrak{u}_1) \subset \mathfrak{S}$. Then $\mathcal{I}(\mathfrak{p}) \subsetneq J$. This contradicts the definition of $\mathcal{J}(\mathfrak{S})$. \square

For cubes $J \in \mathcal{D}$, denote

$$B(J) := B(c(J), 8D^{s(J)}). \quad (7.36)$$

The main result of this subsection is the following.

Lemma 7.22 (Lipschitz partition unity). *There exists a family of functions χ_J , $J \in \mathcal{J}'$ such that*

$$\mathbf{1}_{\mathcal{I}(\mathfrak{u}_1)} = \sum_{J \in \mathcal{J}'} \chi_J, \quad (7.37)$$

and for all $J \in \mathcal{J}'$ and all $y, y' \in \mathcal{I}(\mathfrak{u}_1)$

$$0 \leq \chi_J(y) \leq \mathbf{1}_{B(J)}(y), \quad (7.38)$$

$$|\chi_J(y) - \chi_J(y')| \leq 2^{226a^3} \frac{\rho(y, y')}{D^{s(J)}}. \quad (7.39)$$

In the proof, we will use the following auxiliary lemma.

Lemma 7.23 (moderate scale change). *If $J, J' \in \mathcal{J}'$ with*

$$B(J) \cap B(J') \neq \emptyset,$$

then $|s(J) - s(J')| \leq 1$.

Proof of Lemma 7.22. For each cube $J \in \mathcal{J}$ let

$$\tilde{\chi}_J(y) = \max\{0, 8 - D^{-s(J)}\rho(y, c(J))\},$$

and set

$$a(y) = \sum_{J \in \mathcal{J}'} \tilde{\chi}_J(y).$$

We define

$$\chi_J(y) := \frac{\tilde{\chi}_J(y)}{a(y)}.$$

Then, due to (2.37) and (7.36), the properties (7.37) and (7.38) are clearly true. Estimate (7.39) follows from (7.38) if $y, y' \notin B(J)$. Thus we can assume that $y \in B(J)$. We have by the triangle inequality

$$|\chi_J(y) - \chi_J(y')| \leq \frac{|\tilde{\chi}_J(y) - \tilde{\chi}_J(y')|}{a(y)} + \frac{\tilde{\chi}_J(y')|a(y) - a(y')|}{a(y)a(y')}$$

Since $\tilde{\chi}_J(z) \geq 4$ for all $z \in B(c(J), 4) \supset J$ and by Lemma 7.21, we have that $a(z) \geq 4$ for all $z \in \mathcal{I}(\mathfrak{u}_1)$. So we can estimate the above further by

$$\leq 2^{-2}(|\tilde{\chi}_J(y) - \tilde{\chi}_J(y')| + \tilde{\chi}_J(y')|a(y) - a(y')|).$$

If $y' \notin B(c(\mathfrak{p}), 8D^{s(\mathfrak{p})})$ then the second summand vanishes. Else, we can estimate the above, using also that $|\tilde{\chi}_J(y')| \leq 8$, by

$$\leq 2^{-2}|\tilde{\chi}_J(y) - \tilde{\chi}_J(y')| + 2 \sum_{\substack{J' \in \mathcal{J}' \\ B(c(J'), 8D^{s(J')}) \cap B(c(J), 8D^{s(J)}) \neq \emptyset}} |\tilde{\chi}_{J'}(y) - \tilde{\chi}_{J'}(y')|.$$

By the triangle inequality, we have for all dyadic cubes $I \in \mathcal{J}'$

$$|\tilde{\chi}_I(y) - \tilde{\chi}_I(y')| \leq \rho(y, y')D^{-s(I)}.$$

Using this above, we obtain

$$|\chi_J(y) - \chi_J(y')| \leq \rho(y, y') \left(\frac{1}{4} D^{-s(J)} + 2 \sum_{\substack{J' \in \mathcal{J}' \\ B(J') \cap B(J) \neq \emptyset}} D^{-s(J')} \right).$$

By Lemma 7.23, this is at most

$$\frac{\rho(y, y')}{D^{s(J)}} \left(\frac{1}{4} + 2D |\{J' \in \mathcal{J}' : B(J') \cap B(J) \neq \emptyset\}| \right).$$

By (2.10) and Lemma 7.21, the balls $B(c(J'), \frac{1}{4}D^{s(J')})$ are pairwise disjoint, so by Lemma 7.23 the balls $B(c(J'), \frac{1}{4}D^{s(J)-1})$ are also disjoint. By the triangle inequality and Lemma 7.23, each such ball for J' in the set of the last display is contained in

$$B(c(J), 9D^{s(J)+1}).$$

By the doubling property (1.5), we further have

$$\mu\left(B(c(J'), \frac{1}{4}D^{s(J')})\right) \geq 2^{-200a^3-6} \mu(B(c(J), 9D^{s(J)+1}))$$

for each such ball. Thus

$$|\{J' \in \mathcal{J}' : B(J') \cap B(J) \neq \emptyset\}| \leq 2^{200a^3+6}.$$

Recalling that $D = 2^{100a^2}$, we obtain

$$\frac{1}{4} + 2D |\{J' \in \mathcal{J}' : B(J') \cap B(J) \neq \emptyset\}| \leq 2^{200a^3+100a^2+8}.$$

Since $a \geq 4$, (7.39) follows. \square

Proof of Lemma 7.23. Suppose that $s(J') < s(J) - 1$. Then $s(J) > -S$. Thus, by the definition of \mathcal{J}' there exists no $\mathbf{p} \in \mathfrak{S}$ with

$$\mathcal{I}(\mathbf{p}) \subset B(c(J), 100D^{s(J)+1}). \quad (7.40)$$

Since $s(J') < s(J)$ and $J', J \subset \mathcal{I}(\mathbf{u}_1)$, we have $J' \subsetneq \mathcal{I}(\mathbf{u}_1)$. By (2.7), (2.8) there exists a cube $J'' \in \mathcal{D}$ with $J \subset J''$ and $s(J'') = s(J') + 1$. By the definition of \mathcal{J}' , there exists a tile $\mathbf{p} \in \mathfrak{S}$ with

$$\mathcal{I}(\mathbf{p}) \subset B(c(J''), 100D^{s(J')+2}). \quad (7.41)$$

But by the triangle inequality and (2.1), we have

$$B(c(J''), 100D^{s(J')+2}) \subset B(c(J), 100D^{s(J)+1}),$$

which contradicts (7.40) and (7.41). \square

7.5.2. Hölder estimates for adjoint tree operators. Let $g_1, g_2 : X \rightarrow \mathbb{C}$ be bounded with bounded support. Define for $J \in \mathcal{J}'$

$$h_J(y) := \chi_J(y) \cdot (e(\mathcal{Q}(\mathbf{u}_1)(y))T_{\mathfrak{T}(\mathbf{u}_1)}^* g_1(y)) \cdot \overline{(e(\mathcal{Q}(\mathbf{u}_2)(y))T_{\mathfrak{T}(\mathbf{u}_2) \cap \mathfrak{S}}^* g_2(y))}. \quad (7.42)$$

The main result of this subsection is the following τ -Hölder estimate for h_J , where $\tau = 1/a$.

Lemma 7.24 (Holder correlation tree). *We have for all $J \in \mathcal{J}'$ that*

$$\|h_J\|_{C^\tau(B(c(J), 8D^s(J)))} \leq 2^{535a^3} \prod_{j=1,2} \left(\inf_{B(c(J), \frac{1}{8}D^s(J))} |T_{\mathfrak{T}(\mathbf{u}_j)}^* g_j| + \inf_J M_{\mathcal{B},1} |g_j| \right). \quad (7.43)$$

We will prove this lemma at the end of this section, after establishing several auxiliary results.

We begin with the following Hölder continuity estimate for adjoints of operators associated to tiles.

Lemma 7.25 (Holder correlation tile). *Let $\mathbf{u} \in \mathfrak{U}$ and $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$. Then for all $y, y' \in X$ and all bounded g with bounded support, we have*

$$\begin{aligned} & |e(\mathcal{Q}(\mathbf{u})(y))T_{\mathbf{p}}^* g(y) - e(\mathcal{Q}(\mathbf{u})(y'))T_{\mathbf{p}}^* g(y')| \\ & \leq \frac{2^{151a^3}}{\mu(B(c(\mathbf{p}), 4D^s(\mathbf{p})))} \left(\frac{\rho(y, y')}{D^s(\mathbf{p})} \right)^{1/a} \int_{E(\mathbf{p})} |g(x)| d\mu(x). \end{aligned} \quad (7.44)$$

Proof. By (7.27), we have

$$\begin{aligned} & |e(\mathcal{Q}(\mathbf{u})(y))T_{\mathbf{p}}^* g(y) - e(\mathcal{Q}(\mathbf{u})(y'))T_{\mathbf{p}}^* g(y')| \\ & = \left| \int_{E(\mathbf{p})} e(Q(x)(y) - Q(x)(y') + \mathcal{Q}(\mathbf{u})(y) - \mathcal{Q}(\mathbf{u})(y')) \overline{K_{s(\mathbf{p})}(x, y)} g(x) \right. \\ & \quad \left. - e(Q(x)(y) - Q(x)(y') + \mathcal{Q}(\mathbf{u})(y') - \mathcal{Q}(\mathbf{u})(y)) \overline{K_{s(\mathbf{p})}(x, y')} g(x) d\mu(x) \right| \\ & \leq \int_{E(\mathbf{p})} |g(x)| |e(Q(x)(y) - Q(x)(y') - \mathcal{Q}(\mathbf{u})(y) + \mathcal{Q}(\mathbf{u})(y')) \overline{K_{s(\mathbf{p})}(x, y)} \\ & \quad - \overline{K_{s(\mathbf{p})}(x, y')}| d\mu(x) \\ & \leq \int_{E(\mathbf{p})} |g(x)| |e(-Q(x)(y) + Q(x)(y') + \mathcal{Q}(\mathbf{u})(y) - \mathcal{Q}(\mathbf{u})(y')) - 1| \\ & \quad \times |\overline{K_{s(\mathbf{p})}(x, y)}| d\mu(x) \end{aligned} \quad (7.45)$$

$$+ \int_{E(\mathbf{p})} |g(x)| |\overline{K_{s(\mathbf{p})}(x, y)} - \overline{K_{s(\mathbf{p})}(x, y')}| d\mu(x). \quad (7.46)$$

By the oscillation estimate (1.7), we have

$$\begin{aligned} & |-Q(x)(y) + Q(x)(y') + \mathcal{Q}(\mathbf{u})(y) - \mathcal{Q}(\mathbf{u})(y')| \\ & \leq d_{B(y, \rho(y, y'))}(Q(x), \mathcal{Q}(\mathbf{u})). \end{aligned} \quad (7.47)$$

Suppose that $y, y' \in B(c(\mathbf{p}), 5D^s(\mathbf{p}))$, so that $\rho(y, y') \leq 10D^s(\mathbf{p})$. Let $k \in \mathbb{Z}$ be such that $2^{ak}\rho(y, y') \leq 10D^s(\mathbf{p})$ but $2^{a(k+1)}\rho(y, y') > 10D^s(\mathbf{p})$. In particular, $k \geq 0$. Then, using (1.8), we can bound (7.47) from above by

$$2^{-k} d_{B(c(\mathbf{p}), 10D^s(\mathbf{p}))}(Q(x), \mathcal{Q}(\mathbf{u})) \leq 2^{6a-k} d_{\mathbf{p}}(Q(x), \mathcal{Q}(\mathbf{u})).$$

Since $x \in E(\mathbf{p})$ we have $Q(x) \in \Omega(\mathbf{p}) \subset B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 1)$, and since $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$ we have $\mathcal{Q}(\mathbf{u}) \in B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 4)$, so this is estimated by

$$\leq 5 \cdot 2^{6a-k}.$$

By definition of k , we have

$$k \leq \frac{1}{a} \log_2 \left(\frac{10D^{s(\mathfrak{p})}}{\rho(y, y')} \right),$$

which gives

$$|-Q(x)(y) + Q(x)(y') + \mathcal{Q}(\mathfrak{u})(y) - \mathcal{Q}(\mathfrak{u})(y')| \leq 5 \cdot 2^{6a} \left(\frac{\rho(y, y')}{10D^{s(\mathfrak{p})}} \right)^{1/a}. \quad (7.48)$$

For all $x \in \mathcal{I}(\mathfrak{p})$, we have by (1.5) that

$$\mu(B(x, D^{s(\mathfrak{p})})) \geq 2^{-3a} \mu(B(\mathfrak{c}(\mathfrak{p}), 4D^{s(\mathfrak{p})})).$$

Combining the above with (2.49), (2.50) and (7.48), we obtain

$$(7.45) + (7.46) \leq \frac{2^{3a}}{\mu(B(\mathfrak{c}(\mathfrak{p}), 4D^{s(\mathfrak{p})}))} \int_{E(\mathfrak{p})} |g(x)| d\mu(x) \times \\ (2^{102a^3} \cdot 5 \cdot 2^{6a} \left(\frac{\rho(y, y')}{D^{s(\mathfrak{p})}} \right)^{1/a} + 2^{150a^3} \left(\frac{\rho(y, y')}{D^{s(\mathfrak{p})}} \right)^{1/a})$$

Since $\rho(y, y') \leq 10D^{s(\mathfrak{p})}$, we conclude

$$(7.45) + (7.46) \leq \frac{2^{151a^3}}{\mu(B(\mathfrak{c}(\mathfrak{p}), 4D^{s(\mathfrak{p})}))} \left(\frac{\rho(y, y')}{D^{s(\mathfrak{p})}} \right)^{1/a} \int_{E(\mathfrak{p})} |g(x)| d\mu(x).$$

Next, if $y, y' \notin B(\mathfrak{c}(\mathfrak{p}), 5D^{s(\mathfrak{p})})$, then $T_{\mathfrak{p}}^* g(y) = T_{\mathfrak{p}}^* g(y') = 0$, by Lemma 7.14. Then (7.44) holds.

Finally, if $y \in B(\mathfrak{c}(\mathfrak{p}), 5D^{s(\mathfrak{p})})$ and $y' \notin B(\mathfrak{c}(\mathfrak{p}), 5D^{s(\mathfrak{p})})$, then

$$|e(\mathcal{Q}(\mathfrak{u})(y))T_{\mathfrak{p}}^* g(y) - e(\mathcal{Q}(\mathfrak{u})(y'))T_{\mathfrak{p}}^* g(y')| = |T_{\mathfrak{p}}^* g(y)| \\ \leq \int_{E(\mathfrak{p})} |K_{s(\mathfrak{p})}(x, y)| |g(x)| d\mu(x).$$

By the same argument used to prove (2.52), this is bounded by

$$\leq 2^{102a^3} \int_{E(\mathfrak{p})} \frac{1}{\mu(B(x, D^s))} \psi(D^{-s}\rho(x, y)) |g(x)| d\mu(x). \quad (7.49)$$

It follows from the definition of ψ that

$$\psi(x) \leq \max\{0, (2 - 4x)^{1/a}\}.$$

Now for all $x \in E(\mathfrak{p})$, it follows by the triangle inequality and (2.10) that

$$2 - 4D^{-s(\mathfrak{p})}\rho(x, y) \leq 2 - 4D^{-s(\mathfrak{p})}\rho(y, \mathfrak{c}(\mathfrak{p})) + 4D^{-s(\mathfrak{p})}\rho(x, \mathfrak{c}(\mathfrak{p})) \\ \leq 18 - 4D^{-s(\mathfrak{p})}\rho(y, \mathfrak{c}(\mathfrak{p})) \leq 4D^{-s(\mathfrak{p})}\rho(y, y') - 2.$$

Combining the above with the previous estimate on ψ , we get

$$\psi(D^{-s(\mathfrak{p})}\rho(x, y)) \leq 4(D^{-s(\mathfrak{p})}\rho(y, y'))^{1/a}.$$

Further, we obtain from the doubling property (1.5) and (2.10) that

$$\mu(B(x, D^s)) \geq 2^{-2a} \mu(\mathfrak{c}(\mathfrak{p}), 4D^s).$$

Plugging this into (7.49) and using $a \geq 4$, we get

$$|T_{\mathfrak{p}}^* g(y)| \leq \frac{2^{103a^3}}{\mu(B(\mathfrak{c}(\mathfrak{p}), 4D^{s(\mathfrak{p})}))} \left(\frac{\rho(y, y')}{D^{s(\mathfrak{p})}} \right)^{1/a} \int_{E(\mathfrak{p})} |g(x)| d\mu(y),$$

which completes the proof of the lemma. \square

Recall that

$$B(J) := B(c(J), 8D^{s(J)}).$$

We also denote

$$B^\circ(J) := B(c(J), \frac{1}{8}D^{s(J)}).$$

Lemma 7.26 (limited scale impact). *Let $\mathfrak{p} \in \mathfrak{T}_2 \setminus \mathfrak{S}$, $J \in \mathcal{J}'$ and suppose that*

$$B(\mathcal{I}(\mathfrak{p})) \cap B^\circ(J) \neq \emptyset.$$

Then

$$s(J) \leq s(\mathfrak{p}) \leq s(J) + 10a^2 + 2.$$

Proof. For the first estimate, assume that $s(\mathfrak{p}) < s(J)$, then in particular $s(\mathfrak{p}) \leq s(\mathfrak{u}_1)$. Since $\mathfrak{p} \notin \mathfrak{S}$, we have by Lemma 7.20 that $\mathcal{I}(\mathfrak{p}) \cap \mathcal{I}(\mathfrak{u}_1) = \emptyset$. Since $B(c(J), \frac{1}{4}D^{s(J)}) \subset \mathcal{I}(J) \subset \mathcal{I}(\mathfrak{u}_1)$, this implies

$$\rho(c(J), c(\mathfrak{p})) \geq \frac{1}{4}D^{s(J)}.$$

On the other hand

$$\rho(c(J), c(\mathfrak{p})) \leq \frac{1}{8}D^{s(J)} + 8D^{s(\mathfrak{p})},$$

by our assumption. Thus $D^{s(\mathfrak{p})} \geq 64D^{s(J)}$, which contradicts (2.1) and $a \geq 4$.

For the second estimate, assume that $s(\mathfrak{p}) > s(J) + 10a^2 + 2$. Since $J \in \mathcal{J}'$, we have $J \subsetneq \mathcal{I}(\mathfrak{u}_1)$. Thus there exists $J' \in \mathcal{D}$ with $J \subset J'$ and $s(J') = s(J) + 1$, by (2.7) and (2.8). By definition of \mathcal{J}' , there exists some $\mathfrak{p}' \in \mathfrak{S}$ such that $\mathcal{I}(\mathfrak{p}') \subset B(c(J'), 100D^{s(J')+2})$. On the other hand, since $B(\mathcal{I}(\mathfrak{p})) \cap B^\circ(J) \neq \emptyset$, by the triangle inequality it holds that

$$B(c(J'), 100D^{s(J)+10a^2+2}) \subset B(c(\mathfrak{p}), 10D^{s(\mathfrak{p})}).$$

Using the definition of \mathfrak{S} , we have

$$2^{Zn/2} \leq d_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{u}_1), \mathcal{Q}(\mathfrak{u}_2)) \leq d_{B(c(J'), 100D^{s(J)+2})}(\mathcal{Q}(\mathfrak{u}_1), \mathcal{Q}(\mathfrak{u}_2)).$$

By (1.10), this is

$$\begin{aligned} &\leq 2^{-10a} d_{B(c(J'), 100D^{s(J)+10a^2+2})}(\mathcal{Q}(\mathfrak{u}_1), \mathcal{Q}(\mathfrak{u}_2)) \\ &\leq 2^{-10a} d_{B(c(\mathfrak{p}), 10D^{s(\mathfrak{p})})}(\mathcal{Q}(\mathfrak{u}_1), \mathcal{Q}(\mathfrak{u}_2)), \end{aligned}$$

and by (1.8) and the definition of \mathfrak{S}

$$\leq 2^{-4a} d_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{u}_1), \mathcal{Q}(\mathfrak{u}_2)) \leq 2^{-4a} 2^{Zn/2}.$$

This is a contradiction, the second estimate follows. \square

Lemma 7.27 (local tree control). *For all $J \in \mathcal{J}'$ and all bounded g with bounded support*

$$\sup_{B^\circ(J)} |T_{\mathfrak{T}(\mathfrak{u}_2) \setminus \mathfrak{S}}^* g| \leq 2^{104a^3} \inf_J M_{\mathcal{B},1} |g|$$

Proof. By the triangle inequality and since $T_{\mathfrak{p}}^*g = \mathbf{1}_{B(c(\mathfrak{p}), 5D^{s(\mathfrak{p})})}T_{\mathfrak{p}}^*g$, we have

$$\sup_{B^\circ(J)} |T_{\mathfrak{I}(\mathfrak{u}_2) \setminus \mathfrak{S}}^*g| \leq \sup_{B^\circ(J)} \sum_{\substack{\mathfrak{p} \in \mathfrak{I}(\mathfrak{u}_2) \setminus \mathfrak{S} \\ B(\mathcal{I}(\mathfrak{p})) \cap B^\circ(J) \neq \emptyset}} |T_{\mathfrak{p}}^*g|.$$

By Lemma 7.26, this is at most

$$\sum_{s=s(J)}^{s(J)+10a^2+2} \sum_{\substack{\mathfrak{p} \in \mathfrak{P}, s(\mathfrak{p})=s \\ B(\mathcal{I}(\mathfrak{p})) \cap B^\circ(J) \neq \emptyset}} \sup_{B^\circ(J)} |T_{\mathfrak{p}}^*g|. \quad (7.50)$$

If $x \in E(\mathfrak{p})$ and $B(\mathcal{I}(\mathfrak{p})) \cap B^\circ(J) \neq \emptyset$, then

$$B(c(J), 16D^{s(\mathfrak{p})}) \subset B(x, 32D^{s(\mathfrak{p})}),$$

by (2.10) and the triangle inequality. Using the doubling property (1.5), it follows that

$$\mu(B(x, D^{s(\mathfrak{p})})) \geq 2^{-5a} \mu(B(c(J), 16D^{s(\mathfrak{p})})).$$

Using (7.27), (2.49) and that $a \geq 4$, we bound (7.50) by

$$2^{103a^3} \sum_{s=s(J)}^{s(J)+10a^2+2} \sum_{\substack{\mathfrak{p} \in \mathfrak{P}, s(\mathfrak{p})=s \\ B(\mathcal{I}(\mathfrak{p})) \cap B^\circ(J) \neq \emptyset}} \frac{1}{\mu(B(c(J), 16D^s))} \int_{E(\mathfrak{p})} |g| d\mu.$$

For each $I \in \mathcal{D}$, the sets $E(\mathfrak{p})$ for $\mathfrak{p} \in \mathfrak{P}$ with $\mathcal{I}(\mathfrak{p}) = I$ are pairwise disjoint by (2.20) and (2.13). Further, if $B(\mathcal{I}(\mathfrak{p})) \cap B^\circ(J) \neq \emptyset$ and $s(\mathfrak{p}) \geq s(J)$, then $E(\mathfrak{p}) \subset B(c(J), 32D^{s(\mathfrak{p})})$. Thus the last display is bounded by

$$\begin{aligned} & 2^{103a^3} \sum_{s=s(J)}^{s(J)+10a^2+2} \frac{1}{\mu(B(c(J), 32D^s))} \int_{B(c(J), 16D^s)} |g| d\mu \\ & \leq \inf_{x' \in J} 2^{103a^3} (10a^2 + 3) M_{\mathcal{B},1} |g|. \end{aligned}$$

The lemma follows since for $a \geq 4$ it holds that $10a^2 + 3 \leq 2^{a^3}$. \square

Lemma 7.28 (scales impacting interval). *Let $\mathfrak{C} = \mathfrak{I}(\mathfrak{u}_1)$ or $\mathfrak{C} = \mathfrak{I}(\mathfrak{u}_2) \cap \mathfrak{S}$. Then for each $J \in \mathcal{J}'$ and $\mathfrak{p} \in \mathfrak{C}$ with $B(\mathcal{I}(\mathfrak{p})) \cap B(J) \neq \emptyset$, we have $s(\mathfrak{p}) \geq s(J)$.*

Proof. By Lemma 7.20, we have that in both cases, $\mathfrak{C} \subset \mathfrak{S}$. If $\mathfrak{p} \in \mathfrak{C}$ with $B(\mathcal{I}(\mathfrak{p})) \cap B(J) \neq \emptyset$ and $s(\mathfrak{p}) < s(J)$, then $\mathcal{I}(\mathfrak{p}) \subset B(c(J), 100D^{s(J)+1})$. Since $\mathfrak{p} \in \mathfrak{S}$, it follows from the definition of \mathcal{J}' that $s(J) = -S$, which contradicts $s(\mathfrak{p}) < s(J)$. \square

Lemma 7.29 (global tree control 1). *Let $\mathfrak{C} = \mathfrak{I}(\mathfrak{u}_1)$ or $\mathfrak{C} = \mathfrak{I}(\mathfrak{u}_2) \cap \mathfrak{S}$. Then for each $J \in \mathcal{J}'$ and all bounded g with bounded support, we have*

$$\sup_{B(J)} |T_{\mathfrak{C}}^*g| \leq \inf_{B^\circ(J)} |T_{\mathfrak{C}}^*g| + 2^{154a^3} \inf_J M_{\mathcal{B},1} |g| \quad (7.51)$$

and for all $y, y' \in B(J)$

$$\begin{aligned} & |e(\mathcal{Q}(\mathfrak{u})(y))T_{\mathfrak{C}}^*g(y) - e(\mathcal{Q}(\mathfrak{u})(y'))T_{\mathfrak{C}}^*g(y')| \\ & \leq 2^{153a^3} \left(\frac{\rho(y, y')}{D^{s(J)}} \right)^{1/a} \inf_J M_{\mathcal{B},1} |g|. \end{aligned} \quad (7.52)$$

Proof. Note that (7.51) follows from (7.52), since for $y' \in B^\circ(J)$, by the triangle inequality,

$$\left(\frac{\rho(y, y')}{D^{s(J)}}\right)^{1/a} \leq \left(8 + \frac{1}{8}\right)^{1/a} \leq 2^{a^3}.$$

By the triangle inequality, Lemma 7.14 and Lemma 7.25, we have for all $y, y' \in B(J)$

$$\begin{aligned} & |e(\mathcal{Q}(\mathbf{u})(y))T_{\mathfrak{C}}^*g(y) - e(\mathcal{Q}(\mathbf{u})(y'))T_{\mathfrak{C}}^*g(y')| \\ & \leq \sum_{\substack{\mathbf{p} \in \mathfrak{C} \\ B(\mathcal{I}(\mathbf{p})) \cap B(J) \neq \emptyset}} |e(\mathcal{Q}(\mathbf{u})(y))T_{\mathbf{p}}^*g(y) - e(\mathcal{Q}(\mathbf{u})(y'))T_{\mathbf{p}}^*g(y')| \\ & \leq 2^{151a^3} \rho(y, y')^{1/a} \sum_{\substack{\mathbf{p} \in \mathfrak{C} \\ B(\mathcal{I}(\mathbf{p})) \cap B(J) \neq \emptyset}} \frac{D^{-s(\mathbf{p})/a}}{\mu(B(c(\mathbf{p}), 4D^{s(\mathbf{p})}))} \int_{E(\mathbf{p})} |g| d\mu. \end{aligned} \quad (7.53)$$

By Lemma 7.28, we have $s(\mathbf{p}) \geq s(J)$ for all \mathbf{p} occurring in the sum. Further, for each $s \geq s(J)$, the sets $E(\mathbf{p})$ for $\mathbf{p} \in \mathfrak{P}$ with $s(\mathbf{p}) = s$ are pairwise disjoint by (2.20) and (2.13), and contained in $B(c(J), 32D^s)$ by (2.10) and the triangle inequality. Using also the doubling estimate (1.5), we obtain that the expression in the last display can be estimated by

$$\begin{aligned} & 2^{151a^3} \rho(y, y')^{1/a} \sum_{S \geq s \geq s(J)} D^{-s/a} \frac{2^{3a}}{\mu(B(c(J), 32D^s))} \int_{B(c(J), 32D^s)} |g| d\mu \\ & \leq 2^{152a^3} \left(\frac{\rho(y, y')}{D^s}\right)^{1/a} \sum_{S \geq s \geq s(J)} D^{(s(J)-s)/a} \inf_J M_{\mathcal{B},1} |g|. \end{aligned}$$

By convexity of $t \mapsto D^t$ and since $D \geq 2$, we have for all $-1 \leq t \leq 0$

$$D^t \leq 1 + t(1 - \frac{1}{D}) \leq 1 + \frac{1}{2}t.$$

Since $-1 \leq -1/a < 0$, it follows that

$$\sum_{S \geq s \geq s(J)} D^{(s(J)-s)/a} \leq \frac{1}{1 - D^{-1/a}} \leq 2a \leq 2^a.$$

Estimate (7.52), and therefore the lemma, follow. \square

Lemma 7.30 (global tree control 2). *We have for all $J \in \mathcal{J}'$ and all bounded g with bounded support*

$$\sup_{B(J)} |T_{\mathfrak{I}_2 \cap \mathfrak{S}}^*g| \leq \inf_{B^\circ(J)} |T_{\mathfrak{I}_2}^*g| + 2^{155a^3} \inf_J M_{\mathcal{B},1} |g|.$$

Proof. By Lemma 7.29

$$\begin{aligned} & \sup_{B(J)} |T_{\mathfrak{I}(u_2) \cap \mathfrak{S}}^*g| \leq \inf_{B^\circ(J)} |T_{\mathfrak{I}(u_2)}^*g| + 2^{154a^3} \inf_J M_{\mathcal{B},1} |g| \\ & \leq \inf_{B^\circ(J)} |T_{\mathfrak{I}(u_2)}^*g| + \sup_{B^\circ(J)} |T_{\mathfrak{I}(u_2) \setminus \mathfrak{S}}^*g| + 2^{154a^3} \inf_J M_{\mathcal{B},1} |g|, \end{aligned}$$

and by Lemma 7.27

$$\leq \inf_{B^\circ(J)} |T_{\mathfrak{I}(u_2)}^*g| + (2^{104a^3} + 2^{154a^3}) \inf_J M_{\mathcal{B},1} |g|.$$

This completes the proof. \square

Proof of Lemma 7.24. Let P be the product on the right hand side of (7.43), and h_J as defined in (7.42).

By (7.38) and Lemma 7.14, the function h_J is supported in $B(J) \cap \mathcal{I}(\mathbf{u}_1)$. By (7.38) and Lemma 7.29, we have for all $y \in B(J)$:

$$|h_J(y)| \leq 2^{308a^3} P.$$

We have by the triangle inequality

$$\begin{aligned} & |h_J(y) - h_J(y')| \\ & \leq |\chi_J(y) - \chi_J(y')| |T_{\mathfrak{T}(\mathbf{u}_1)}^* g_1(y)| |T_{\mathfrak{T}(\mathbf{u}_2) \cap \mathfrak{S}}^* g_2(y)| \end{aligned} \quad (7.54)$$

$$+ |\chi_J(y')| |T_{\mathfrak{T}(\mathbf{u}_1)}^* g_1(y) - T_{\mathfrak{T}(\mathbf{u}_1)}^* g_1(y')| |T_{\mathfrak{T}(\mathbf{u}_2) \cap \mathfrak{S}}^* g_2(y)| \quad (7.55)$$

$$+ |\chi_J(y')| |T_{\mathfrak{T}(\mathbf{u}_1)}^* g_1(y')| |T_{\mathfrak{T}(\mathbf{u}_2) \cap \mathfrak{S}}^* g_2(y) - T_{\mathfrak{T}(\mathbf{u}_2) \cap \mathfrak{S}}^* g_2(y')|. \quad (7.56)$$

As h_J is supported in $\mathcal{I}(\mathbf{u}_1)$, we can assume without loss of generality that $y' \in \mathcal{I}(\mathbf{u}_1)$. If $y \notin \mathcal{I}(\mathbf{u}_1)$, then (7.54) vanishes. If $y \in \mathcal{I}(\mathbf{u}_1)$ then we have by (7.39), Lemma 7.29 and Lemma 7.30

$$(7.54) \leq 2^{534a^3} \frac{\rho(y, y')}{D^{s(J)}} P,$$

where P denotes the product on the right hand side of (7.43).

By (7.38), Lemma 7.29 and Lemma 7.30, we have

$$(7.55) \leq 2^{310a^3} P.$$

By (7.38), and twice Lemma 7.29, we have

$$(7.56) \leq 2^{308a^3} P.$$

Using that $\rho(y, y') \leq 16D^{s(J)}$ and $a \geq 4$, the lemma follows. \square

7.5.3. The van der Corput estimate.

Lemma 7.31 (lower oscillation bound). *For all $J \in \mathcal{J}'$, we have that*

$$d_{B(J)}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)) \geq 2^{-201a^3} 2^{Zn/2}.$$

Proof. Since $\emptyset \neq \mathfrak{T}(\mathbf{u}_1) \subset \mathfrak{S}$ by Lemma 7.20, there exists at least one tile $\mathbf{p} \in \mathcal{S}$ with $\mathcal{I}(\mathbf{p}) \subsetneq \mathcal{I}(\mathbf{u}_1)$. Thus $\mathcal{I}(\mathbf{u}_1) \notin \mathcal{J}'$, so $J \subsetneq \mathcal{I}(\mathbf{u}_1)$. Thus there exists a cube $J' \in \mathcal{D}$ with $J \subset J'$ and $s(J') = s(J) + 1$, by (2.7) and (2.8). By definition of \mathcal{J}' and the triangle inequality, there exists $\mathbf{p} \in \mathfrak{S}$ such that

$$\mathcal{I}(\mathbf{p}) \subset B(c(J'), 100D^{s(J')+1}) \subset B(c(J), 128D^{s(J)+2}).$$

Thus, by definition of \mathfrak{S} :

$$2^{Zn/2} \leq d_{\mathbf{p}}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)) \leq d_{B(c(J), 128D^{s(J)+2})}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)).$$

By the doubling property (1.8), this is

$$\leq 2^{200a^3+4a} d_{B(J)}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)),$$

which gives the lemma using $a \geq 4$. \square

Now we are ready to prove Lemma 7.18.

Proof of Lemma 7.18. We have

$$(7.32) = \left| \int_X T_{\mathfrak{T}(\mathbf{u}_1)}^* g_1 \overline{T_{\mathfrak{T}(\mathbf{u}_2) \cap \mathfrak{S}}^* g_2} \right|.$$

By Lemma 7.14, the right hand side is supported in $\mathcal{I}(\mathbf{u}_1)$. Using (7.37) of Lemma 7.22 and the definition (7.42) of h_J , we thus have

$$\leq \sum_{J \in \mathcal{J}'} \left| \int_{B(J)} e(\mathcal{Q}(\mathbf{u}_2)(y) - \mathcal{Q}(\mathbf{u}_1)(y)) h_J(y) d\mu(y) \right|.$$

Using Proposition 2.5 with the ball $B(J)$, we bound this by

$$\leq 2^{8a} \sum_{J \in \mathcal{J}'} \mu(B(J)) \|h_J\|_{C^\tau(B(J))} (1 + d_{B(J)}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_1)))^{-1/(2a^2+a^3)}.$$

Using Lemma 7.24, Lemma 7.31 and $a \geq 4$, we have that the above is bounded from above by

$$\begin{aligned} &\leq 2^{540a^3} 2^{-Zn/(4a^2+2a^3)} \sum_{J \in \mathcal{J}'} \mu(B(J)) \\ &\quad \times \prod_{j=1}^2 \left(\inf_{B^\circ(J)} |T_{\mathfrak{T}(\mathbf{u}_j)}^* g_j| + \inf_J M_{\mathcal{B},1} g_j \right). \end{aligned} \quad (7.57)$$

By the doubling property (1.5)

$$\mu(B(J)) \leq 2^{6a} \mu(B^\circ(J)),$$

thus

$$\begin{aligned} &\mu(B(J)) \prod_{j=1}^2 \left(\inf_{B^\circ(J)} |T_{\mathfrak{T}(\mathbf{u}_j)}^* g_j| + \inf_J M_{\mathcal{B},1} g_j \right) \\ &\leq 2^{6a} \int_{B^\circ(J)} \prod_{j=1}^2 (|T_{\mathfrak{T}(\mathbf{u}_j)}^* g_j|(x) + M_{\mathcal{B},1} g_j(x)) d\mu(x) \\ &\leq 2^{6a} \int_J \prod_{j=1}^2 (|T_{\mathfrak{T}(\mathbf{u}_j)}^* g_j|(x) + M_{\mathcal{B},1} g_j(x)) d\mu(x). \end{aligned}$$

Summing over $J \in \mathcal{J}'$, we obtain

$$(7.57) \leq 2^{541a^3} 2^{-Zn/(4a^2+2a^3)} \int_X \prod_{j=1}^2 (|T_{\mathfrak{T}(\mathbf{u}_j)}^* g_j|(x) + M_{\mathcal{B},1} g_j(x)) d\mu(x).$$

Applying the Cauchy-Schwarz inequality, Lemma 7.18 follows. \square

7.6. Proof of Lemma 7.19: The remaining tiles. We define

$$\mathcal{J}' = \{J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}_1)) : J \subset \mathcal{I}(\mathbf{u}_1)\},$$

note that this is different from the \mathcal{J}' defined in the previous subsection.

Lemma 7.32 (dyadic partition 2). *We have*

$$\mathcal{I}(\mathbf{u}_1) = \dot{\bigcup}_{J \in \mathcal{J}'} J.$$

Proof. By Lemma 7.2, it remains only to show that each $J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}_1))$ with $J \cap \mathcal{I}(\mathbf{u}_1) \neq \emptyset$ is in \mathcal{J}' . But if $J \notin \mathcal{J}'$, then by (2.8) $\mathcal{I}(\mathbf{u}_1) \subsetneq J$. Pick $\mathbf{p} \in \mathfrak{T}(\mathbf{u}_1)$. Then $\mathcal{I}(\mathbf{p}) \subsetneq J$. This contradicts the definition of $\mathcal{J}(\mathfrak{T}(\mathbf{u}_1))$. \square

Lemma 7.19 follows from the following key estimate.

Lemma 7.33 (bound for tree projection). *We have*

$$\|P_{\mathcal{J}'}|T_{\mathfrak{T}_2 \setminus \mathfrak{S}}^* g_2|\|_2 \leq 2^{118a^3} 2^{-\frac{100}{202a} Zn\kappa} \|\mathbf{1}_{\mathcal{I}(\mathbf{u}_1)} M_{B,1} |g_2|\|_2$$

We prove this lemma below. First, we deduce Lemma 7.19.

Proof of Lemma 7.19. By Lemma 7.7 and Lemma 7.14, we have

$$(7.34) \leq 2^{104a^3} \|P_{\mathcal{L}(\mathfrak{T}(\mathbf{u}_1))} |g_1 \mathbf{1}_{\mathcal{I}(\mathbf{u}_1)}|\|_2 \|\mathbf{1}_{\mathcal{I}(\mathbf{u}_1)} P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}_1))} |T_{\mathfrak{T}(\mathbf{u}_2) \setminus \mathfrak{S}}^* g_2|\|_2.$$

It follows from the definition of the projection operator P and Jensen's inequality that

$$\|P_{\mathcal{L}(\mathfrak{T}(\mathbf{u}_1))} |g_1 \mathbf{1}_{\mathcal{I}(\mathbf{u}_1)}|\|_2 \leq \|g_1 \mathbf{1}_{\mathcal{I}(\mathbf{u}_1)}\|_2.$$

Since cubes in \mathcal{J}' are pairwise disjoint and by Lemma 7.32, a cube $J \in \mathcal{J}'$ intersect $\mathcal{I}(\mathbf{u}_1)$ if and only if $J \in \mathcal{J}'$. Thus

$$\mathbf{1}_{\mathcal{I}(\mathbf{u}_1)} P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}_1))} |T_{\mathfrak{T}(\mathbf{u}_2) \setminus \mathfrak{S}}^* g_2| = P_{\mathcal{J}'} |T_{\mathfrak{T}(\mathbf{u}_2) \setminus \mathfrak{S}}^* g_2|.$$

Combining this with Lemma 7.33, the definition (2.2) and $a \geq 4$ proves the lemma. \square

We need two more auxiliary lemmas before we prove Lemma 7.33.

Lemma 7.34 (thin scale impact). *If $\mathbf{p} \in \mathfrak{T}_2 \setminus \mathfrak{S}$ and $J \in \mathcal{J}'$ with $B(\mathcal{I}(\mathbf{p})) \cap B(J) \neq \emptyset$, then*

$$s(\mathbf{p}) \leq s(J) + 2 - \frac{Zn}{202a^3}.$$

Proof. Suppose that $s(\mathbf{p}) > s(J) + 2 - \frac{Zn}{202a^3} =: s(J) - s_1$. Then, we have

$$\rho(c(\mathbf{p}), c(J)) \leq 8D^{s(J)} + 8D^{s(\mathbf{p})} \leq 16D^{s(\mathbf{p})+s_1}.$$

There exists a tile $\mathbf{q} \in \mathfrak{T}(\mathbf{u}_1)$. By (2.32), it satisfies $\mathcal{I}(\mathbf{q}) \subsetneq \mathcal{I}(\mathbf{u}_1)$. Thus $\mathcal{I}(\mathbf{u}_1) \notin \mathcal{J}'$. It follows that $J \subsetneq \mathcal{I}(\mathbf{u}_1)$. By (2.7) and (2.8), there exists a cube $J' \in \mathcal{D}$ with $J \subset J'$ and $s(J') = s(J) + 1$. By definition of \mathcal{J}' , there exists a tile $\mathbf{p}' \in \mathfrak{T}(\mathbf{u}_1)$ with

$$\mathcal{I}(\mathbf{p}') \subset B(c(J'), 100D^{s(J')+1}).$$

By the triangle inequality, the definition (2.1) and $a \geq 4$, we have

$$B(c(J'), 100D^{s(J')+1}) \subset B(c(\mathbf{p}), 128D^{s(\mathbf{p})+s_1+1}).$$

Since $\mathbf{p}' \in \mathfrak{T}(\mathbf{u}_1)$ and $\mathcal{I}(\mathbf{u}_1) \subset \mathcal{I}(\mathbf{u}_2)$, we have by (2.36)

$$d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), \mathcal{Q}(\mathbf{u}_2)) > 2^{Z(n+1)}.$$

Hence, by (2.32), the triangle inequality and using that by (2.3) $Z \geq 2$

$$d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)) > 2^{Z(n+1)} - 4 \geq 2^{Zn}.$$

It follows that

$$2^{Zn} \leq d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)) \leq d_{B(c(\mathbf{p}), 128D^{s(\mathbf{p})+s_1+1})}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)).$$

Using (1.8), we obtain

$$\leq 2^{9a+100a^3(s_1+1)} d_{\mathfrak{p}}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)).$$

Since $\mathfrak{p}' \notin \mathfrak{S}$ this is bounded by

$$\leq 2^{9a+100a^3(s_1+1)} 2^{Zn/2}.$$

Thus

$$Zn/2 \leq 9a + 100a^3(s_1 + 1),$$

contradicting the definition of s_1 . \square

Lemma 7.35 (square function count). *For each $J \in \mathcal{J}'$, we have*

$$\frac{1}{\mu(J)} \int_J \left(\sum_{\substack{I \in \mathcal{D}, s(I)=s(J)-s \\ I \cap \mathcal{I}(\mathbf{u}_1)=\emptyset \\ J \cap B(I) \neq \emptyset}} \mathbf{1}_{B(I)} \right)^2 d\mu \leq 2^{104a^2} (8D^{-s})^\kappa.$$

Proof of Lemma 7.35. Since $J \in \mathcal{J}'$ we have $J \subset \mathcal{I}(\mathbf{u}_1)$. Thus, if $B(I) \cap J \neq \emptyset$ then

$$B(I) \cap J \subset \{x \in J : \rho(x, X \setminus J) \leq 8D^{s(I)}\}. \quad (7.58)$$

Furthermore, for each s the balls $B(I)$ with $s(I) = s$ have bounded overlap: Consider the collection $\mathcal{D}_{s,x}$ of all $I \in \mathcal{D}$ with $x \in B(I)$ and $s(I) = s$. By (2.10) and (2.8), the balls $B(c(I), \frac{1}{4}D^{s(I)})$, $I \in \mathcal{D}_{s,x}$ are disjoint, and by the triangle inequality, they are contained in $B(x, 9D^s)$. By the doubling property (1.5), we have

$$\mu(B(x, 9D^s)) \leq 2^{8a} \mu(B(c(I), \frac{1}{4}D^{s(I)}))$$

for each $I \in \mathcal{D}_{s,x}$. Thus

$$\mu(B(x, 9D^s)) \geq \sum_{I \in \mathcal{D}_{s,x}} \mu(B(c(I), \frac{1}{4}D^{s(I)})) \geq 2^{-8a} |\mathcal{D}_{s,x}| \mu(B(x, 9D^s)).$$

Dividing by the positive $\mu(B(x, 9D^s))$, we obtain that for each x

$$\left(\sum_{\substack{I \in \mathcal{D}, s(I)=s(J)-s \\ I \cap \mathcal{I}(\mathbf{u}_1)=\emptyset \\ J \cap B(I) \neq \emptyset}} \mathbf{1}_{B(I)}(x) \right)^2 = |\mathcal{D}_{s(J)-s,x}|^2 \leq 2^{16a}. \quad (7.59)$$

Combining (7.58), (7.59) and the small boundary property (2.11), noting that $8D^{s(I)} = 8D^{-s} D^{s(J)}$, the lemma follows. \square

Proof of Lemma 7.33. Expanding the definition of $P_{\mathcal{J}'}$, we have

$$\begin{aligned} & \|P_{\mathcal{J}'} |T_{\mathfrak{T}_2 \setminus \mathfrak{S}}^* g_2| \|_2 \\ &= \left(\sum_{J \in \mathcal{J}'} \frac{1}{\mu(J)} \left| \int_J \sum_{\mathfrak{p} \in \mathfrak{T}(\mathbf{u}_2) \setminus \mathfrak{S}} T_{\mathfrak{p}}^* g_2 d\mu(y) \right|^2 \right)^{1/2}. \end{aligned}$$

We split the innermost sum according to the scale of the tile \mathbf{p} , and then apply the triangle inequality and Minkowski's inequality:

$$\leq \sum_{s=-S}^S \left(\sum_{J \in \mathcal{J}'} \frac{1}{\mu(J)} \left| \int_J \sum_{\substack{\mathbf{p} \in \mathfrak{T}(\mathbf{u}_2) \setminus \mathfrak{S} \\ s(\mathbf{p})=s}} T_{\mathbf{p}}^* g_2 \, d\mu(y) \right|^2 \right)^{1/2}.$$

By Lemma 7.14, the integral in the last display is 0 if $J \cap B(\mathcal{I}(\mathbf{p})) = \emptyset$. By Lemma 7.34, it follows with $s_1 := \frac{Z_n}{202a^3} - 2$:

$$= \sum_{s=s_1}^{s_1+2S} \left(\sum_{J \in \mathcal{J}'} \frac{1}{\mu(J)} \left| \int_J \sum_{\substack{\mathbf{p} \in \mathfrak{T}(\mathbf{u}_2) \setminus \mathfrak{S} \\ s(\mathbf{p})=s(J)-s \\ J \cap B(\mathcal{I}(\mathbf{p})) \neq \emptyset}} T_{\mathbf{p}}^* g_2 \, d\mu(y) \right|^2 \right)^{1/2}. \quad (7.60)$$

We have by Lemma 7.14 and (2.49)

$$\begin{aligned} & \int |T_{\mathbf{p}}^* g_2|(y) \, d\mu(y) \\ & \leq 2^{102a^3} \int_{B(\mathcal{I}(\mathbf{p}))} \int \frac{1}{\mu(B(x, D^{s(\mathbf{p})}))} \mathbf{1}_{E(\mathbf{p})}(x) |g_2|(x) \, d\mu(x) \, d\mu(y). \end{aligned}$$

If $x \in E(\mathbf{p}) \subset \mathcal{I}(\mathbf{p})$, then we have by (2.10) that

$$B(c(\mathbf{p}), 4D^{s(\mathbf{p})}) \subset B(\mathcal{I}(\mathbf{p})).$$

Using the doubling property (1.5), it follows that

$$\mu(B(c(\mathbf{p}), 4D^{s(\mathbf{p})})) \leq 2^{3a} \mu(B(x, D^s)).$$

Thus, using also $a \geq 4$

$$\begin{aligned} & \int |T_{\mathbf{p}}^* g_2|(y) \, d\mu(y) \\ & \leq 2^{103a^3} \int_{B(\mathcal{I}(\mathbf{p}))} \int \frac{1}{\mu(B(c(\mathbf{p}), 4D^{s(\mathbf{p})}))} \mathbf{1}_{E(\mathbf{p})}(x) |g_2|(x) \, d\mu(x) \, d\mu(y). \end{aligned}$$

Since for each $I \in \mathcal{D}$ the sets $E(\mathbf{p}), \mathbf{p} \in \mathfrak{P}(I)$ are disjoint, it follows that

$$\begin{aligned} & \left| \int_J \sum_{\substack{\mathbf{p} \in \mathfrak{T}(\mathbf{u}_2) \setminus \mathfrak{S} \\ \mathcal{I}(\mathbf{p})=I \\ J \cap B(\mathcal{I}(\mathbf{p})) \neq \emptyset}} T_{\mathbf{p}}^* g_2 \, d\mu \right| \\ & \leq 2^{103a^3} \int_J \mathbf{1}_{B(I)} \frac{1}{\mu(B(c(\mathbf{p}), 4D^{s(\mathbf{p})}))} \int_{B(c(\mathbf{p}), 4D^{s(\mathbf{p})})} |g_2|(x) \, d\mu(x) \\ & \leq 2^{103a^3} \int_J M_{\mathcal{B},1} |g_2|(y) \mathbf{1}_{B(I)}(y) \, d\mu(y). \end{aligned}$$

By Lemma 7.20, we have $\mathcal{I}(\mathbf{p}) \cap \mathcal{I}(\mathbf{u}_1) = \emptyset$ for all $\mathbf{p} \in \mathfrak{T}(\mathbf{u}_2) \setminus \mathfrak{S}$. Thus we can estimate (7.60) by

$$2^{103a^3} \sum_{s=s_1}^{s_1+2S} \left(\sum_{J \in \mathcal{J}'} \frac{1}{\mu(J)} \left| \int_J \sum_{\substack{I \in \mathcal{D}, s(I)=s(J)-s \\ I \cap \mathcal{I}(\mathbf{u}_1)=\emptyset \\ J \cap B(I) \neq \emptyset}} M_{\mathcal{B},1} |g_2| \mathbf{1}_{B(I)} \, d\mu \right|^2 \right)^{\frac{1}{2}},$$

which is by Cauchy-Schwarz at most

$$2^{103a^3} \sum_{s=s_1}^{s_1+2S} \left(\sum_{J \in \mathcal{J}'} \int_J (M_{\mathcal{B},1}|g_2|)^2 \frac{1}{\mu(J)} \int_J \left(\sum_{\substack{I \in \mathcal{D}, s(I)=s(J)-s \\ I \cap \mathcal{I}(\mathbf{u}_1)=\emptyset \\ J \cap B(I) \neq \emptyset}} \mathbf{1}_{B(I)} \right)^2 d\mu \right)^{\frac{1}{2}}. \quad (7.61)$$

Using Lemma 7.35, we bound (7.61) by

$$2^{103a^3} \sum_{s=s_1}^{s_1+2S} \left(\sum_{J \in \mathcal{J}'} \int_J (M_{\mathcal{B},1}|g_2|)^2 2^{104a^2} (8D^{-s})^\kappa \right)^{\frac{1}{2}},$$

and, since dyadic cubes in \mathcal{J}' form a partition of $\mathcal{I}(\mathbf{u}_1)$ by Lemma 7.32, $\kappa \leq 1$ by (2.2), and $a \geq 4$

$$\begin{aligned} &\leq 2^{116a^3} \sum_{s=s_1}^{s_1+2S} D^{-s\kappa/2} \|\mathbf{1}_{\mathcal{I}(\mathbf{u}_1)} M_{\mathcal{B},1}|g_2|\|_2 \\ &\leq 2^{116a^3} D^{-s_1\kappa/2} \frac{1}{1 - D^{-\kappa/2}} \|\mathbf{1}_{\mathcal{I}(\mathbf{u}_1)} M_{\mathcal{B},1}|g_2|\|_2. \end{aligned}$$

By convexity of $t \mapsto D^t$ and since $D \geq 2$, we have for all $-1 \leq t \leq 0$

$$D^t \leq 1 + t(1 - \frac{1}{D}) \leq 1 + \frac{1}{2}t.$$

Using this for $t = -\kappa/2$ and using that $s_1 = \frac{Zn}{202a^3} - 2$ and the definitions (2.1) and (2.2) of κ and D

$$\begin{aligned} &\leq 2^{116a^3} 2^{-100a^2(\frac{Zn}{202a^3}-2)\frac{\kappa}{2}} \frac{2}{\kappa} \|\mathbf{1}_{\mathcal{I}(\mathbf{u}_1)} M_{\mathcal{B},1}|g_2|\|_2 \\ &\leq \frac{2^{117a^3+1}}{\kappa} 2^{-\frac{100}{202a}Zn\kappa} \|\mathbf{1}_{\mathcal{I}(\mathbf{u}_1)} M_{\mathcal{B},1}|g_2|\|_2. \end{aligned}$$

Using the definition (2.2) of κ and $a \geq 4$, the lemma follows. \square

7.7. Forests. In this subsection, we complete the proof of Proposition 2.4 from the results of the previous subsections.

Define an n -row to be an n -forest $(\mathfrak{U}, \mathfrak{T})$, i.e. satisfying conditions (2.32) - (2.37), such that in addition the sets $\mathcal{I}(\mathbf{u}), \mathbf{u} \in \mathfrak{U}$ are pairwise disjoint.

Lemma 7.36 (forest row decomposition). *Let $(\mathfrak{U}, \mathfrak{T})$ be an n -forest. Then there exists a decomposition*

$$\mathfrak{U} = \dot{\bigcup}_{1 \leq j \leq 2^n} \mathfrak{U}_j$$

such that for all $j = 1, \dots, 2^n$ the pair $(\mathfrak{U}_j, \mathfrak{T}|_{\mathfrak{U}_j})$ is an n -row.

Proof. Define recursively \mathfrak{U}_j to be a maximal disjoint set of tiles \mathbf{u} in

$$\mathfrak{U} \setminus \bigcup_{j' < j} \mathfrak{U}_{j'}$$

with inclusion maximal $\mathcal{I}(\mathbf{u})$. Properties (2.32), -(2.37) for $(\mathfrak{U}_j, \mathfrak{T}|_{\mathfrak{U}_j})$ follow immediately from the corresponding properties for $(\mathfrak{U}, \mathfrak{T})$, and the cubes

$\mathcal{I}(\mathbf{u}), \mathbf{u} \in \mathfrak{U}_j$ are disjoint by definition. The collections \mathfrak{U}_j are also disjoint by definition.

Now we show by induction on j that each point is contained in at most $2^n - j$ cubes $\mathcal{I}(\mathbf{u})$ with $\mathbf{u} \in \mathfrak{U} \setminus \bigcup_{j' \leq j} \mathfrak{U}_{j'}$. This implies that $\bigcup_{j=1}^{2^n} \mathfrak{U}_j = \mathfrak{U}$, which completes the proof of the Lemma. For $j = 0$ each point is contained in at most 2^n cubes by (2.34). For larger j , if x is contained in any cube $\mathcal{I}(\mathbf{u})$ with $\mathbf{u} \in \mathfrak{U} \setminus \bigcup_{j' < j} \mathfrak{U}_{j'}$, then it is contained in a maximal such cube. Thus it is contained in a cube in $\mathcal{I}(\mathbf{u})$ with $\mathbf{u} \in \mathfrak{U}_j$. Thus the number $\mathbf{u} \in \mathfrak{U} \setminus \bigcup_{j' \leq j} \mathfrak{U}_{j'}$ with $x \in \mathcal{I}(\mathbf{u})$ is zero, or is less than the number of $\mathbf{u} \in \mathfrak{U} \setminus \bigcup_{j' \leq j-1} \mathfrak{U}_{j'}$ with $x \in \mathcal{I}(\mathbf{u})$ by at least one. \square

We pick a decomposition of the forest $(\mathfrak{U}, \mathfrak{T})$ into 2^n n -rows

$$(\mathfrak{U}_j, \mathfrak{T}_j) := (\mathfrak{U}_j, \mathfrak{T}|_{\mathfrak{U}_j})$$

as in Lemma 7.36.

Lemma 7.37 (row bound). *For each $1 \leq j \leq 2^n$ and each bounded g with bounded support, we have*

$$\left\| \sum_{\mathbf{u} \in \mathfrak{U}_j} \sum_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathbf{p}}^* g \right\|_2 \leq 2^{156a^3} 2^{-n/2} \|g\|_2 \quad (7.62)$$

and

$$\left\| \sum_{\mathbf{u} \in \mathfrak{U}_j} \sum_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} \mathbf{1}_F T_{\mathbf{p}}^* g \right\|_2 \leq 2^{257a^3} 2^{-n/2} \text{dens}_2\left(\bigcup_{\mathbf{u} \in \mathfrak{U}} \mathfrak{T}(\mathbf{u})\right)^{1/2} \|g\|_2. \quad (7.63)$$

Proof. By Lemma 7.11 and the density assumption (2.35), we have for each $\mathbf{u} \in \mathfrak{U}$ and all bounded f of bounded support that

$$\left\| \sum_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathbf{p}} f \right\|_2 \leq 2^{155a^3} 2^{(4a+1-n)/2} \|f\|_2 \quad (7.64)$$

and

$$\left\| \sum_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathbf{p}} \mathbf{1}_F f \right\|_2 \leq 2^{256a^3} 2^{(4a+1-n)/2} \text{dens}_2(\mathfrak{T}(\mathbf{u}))^{1/2} \|f\|_2. \quad (7.65)$$

Since for each j the top cubes $\mathcal{I}(\mathbf{u}), \mathbf{u} \in \mathfrak{U}_j$ are disjoint, we further have for all bounded g of bounded support by Lemma 7.14

$$\begin{aligned} & \left\| \mathbf{1}_F \sum_{\mathbf{u} \in \mathfrak{U}_j} \sum_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathbf{p}}^* g \right\|_2^2 = \left\| \mathbf{1}_F \sum_{\mathbf{u} \in \mathfrak{U}_j} \sum_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} \mathbf{1}_{\mathcal{I}(\mathbf{u})} T_{\mathbf{p}}^* \mathbf{1}_{\mathcal{I}(\mathbf{u})} g \right\|_2^2 \\ &= \sum_{\mathbf{u} \in \mathfrak{U}_j} \int_{\mathcal{I}(\mathbf{u})} \left| \mathbf{1}_F \sum_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathbf{p}}^* \mathbf{1}_{\mathcal{I}(\mathbf{u})} g \right|^2 d\mu \leq \sum_{\mathbf{u} \in \mathfrak{U}_j} \left\| \sum_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} \mathbf{1}_F T_{\mathbf{p}}^* \mathbf{1}_{\mathcal{I}(\mathbf{u})} g \right\|_2^2. \end{aligned}$$

Applying the estimate for the adjoint operator following from equation (7.65), we obtain

$$\leq 2^{256a^3} 2^{(4a+1-n)/2} \max_{u \in \mathfrak{U}_j} \text{dens}_2(\mathfrak{T}(u))^{1/2} \sum_{u \in \mathfrak{U}_j} \|\mathbf{1}_{\mathcal{I}(u)} g\|_2^2.$$

Again by disjointedness of the cubes $\mathcal{I}(u)$, this is estimated by

$$2^{256a^3} 2^{(4a+1-n)/2} \max_{u \in \mathfrak{U}_j} \text{dens}_2(\mathfrak{T}(u))^{1/2} \|g\|_2^2.$$

Thus, (7.63) follows, since $a \geq 4$. The proof of (7.62) from (7.64) is the same up to replacing F by X . \square

Lemma 7.38 (row correlation). *For all $1 \leq j < j' \leq 2^n$ and for all bounded g_1, g_2 with bounded support, it holds that*

$$\left| \int \sum_{u \in \mathfrak{U}_j} \sum_{u' \in \mathfrak{U}_{j'}} \sum_{p \in \mathfrak{T}_j(u)} \sum_{p' \in \mathfrak{T}_{j'}(u')} T_p^* g_1 \overline{T_{p'}^* g_2} d\mu \right| \leq 2^{862a^3-3n} \|g_1\|_2 \|g_2\|_2.$$

Proof. To save some space we will write for subsets $\mathfrak{C} \subset \mathfrak{P}$

$$T_{\mathfrak{C}}^* = \sum_{p \in \mathfrak{C}} T_p^*.$$

We have by Lemma 7.14 and the triangle inequality that

$$\begin{aligned} & \left| \int \sum_{u \in \mathfrak{U}_j} \sum_{u' \in \mathfrak{U}_{j'}} \sum_{p \in \mathfrak{T}_j(u)} \sum_{p' \in \mathfrak{T}_{j'}(u')} T_p^* g_1 \overline{T_{p'}^* g_2} d\mu \right| \\ & \leq \sum_{u \in \mathfrak{U}_j} \sum_{u' \in \mathfrak{U}_{j'}} \left| \int T_{\mathfrak{T}_j(u)}^* (\mathbf{1}_{\mathcal{I}(u)} g_1) \overline{T_{\mathfrak{T}_{j'}(u')}^* (\mathbf{1}_{\mathcal{I}(u')} g_2)} d\mu \right|. \end{aligned}$$

By Lemma 7.17, this is bounded by

$$2^{550a^3-3n} \sum_{u \in \mathfrak{U}_j} \sum_{u' \in \mathfrak{U}_{j'}} \|S_{2,u} g_1\|_{L^2(\mathcal{I}(u') \cap \mathcal{I}(u))} \|S_{2,u'} g_2\|_{L^2(\mathcal{I}(u') \cap \mathcal{I}(u))}. \quad (7.66)$$

We apply the Cauchy-Schwarz inequality in the form

$$\sum_{i \in M} a_i b_i \leq \left(\sum_{i \in M} a_i^2 \right)^{1/2} \left(\sum_{i \in M} b_i^2 \right)^{1/2}$$

to the outer two sums:

$$\begin{aligned} & \leq 2^{550a^3-3n} \left(\sum_{u \in \mathfrak{U}_j} \sum_{u' \in \mathfrak{U}_{j'}} \|S_{2,u} g_1\|_{L^2(\mathcal{I}(u') \cap \mathcal{I}(u))}^2 \right)^{1/2} \\ & \quad \left(\sum_{u \in \mathfrak{U}_j} \sum_{u' \in \mathfrak{U}_{j'}} \|S_{2,u'} g_2\|_{L^2(\mathcal{I}(u') \cap \mathcal{I}(u))}^2 \right)^{1/2}. \end{aligned}$$

By pairwise disjointedness of the sets $\mathcal{I}(u)$ for $u \in \mathfrak{U}_j$ and of the sets $\mathcal{I}(u')$ for $u' \in \mathfrak{U}_{j'}$, we have

$$\sum_{u \in \mathfrak{U}_j} \sum_{u' \in \mathfrak{U}_{j'}} \|S_{2,u} g_1\|_{L^2(\mathcal{I}(u') \cap \mathcal{I}(u))}^2 = \sum_{u \in \mathfrak{U}_j} \sum_{u' \in \mathfrak{U}_{j'}} \int_{\mathcal{I}(u) \cap \mathcal{I}(u')} |S_{2,u} g_1(y)|^2 d\mu(y)$$

$$\leq \int_X |S_{2,u}g_1(y)|^2 d\mu(y) = \|S_{2,u}g_1\|_2^2.$$

Arguing similar for g_2 , we can estimate (7.66) to be

$$\leq 2^{550a^3-3n} \|S_{2,u}g_1\|_2 \|S_{2,u'}g_2\|_2.$$

The lemma now follows from Lemma 7.16. \square

Define for $1 \leq j \leq 2^n$

$$E_j := \bigcup_{u \in \mathcal{U}_j} \bigcup_{p \in \mathfrak{T}(u)} E(p).$$

Lemma 7.39 (disjoint row support). *The sets E_j , $1 \leq j \leq 2^n$ are pairwise disjoint.*

Proof. Suppose that $p \in \mathfrak{T}(u)$ and $p' \in \mathfrak{T}(u')$ with $u \neq u'$ and $x \in E(p) \cap E(p')$. Suppose without loss of generality that $s(p) \leq s(p')$. Then $x \in \mathcal{I}(p) \cap \mathcal{I}(p') \subset \mathcal{I}(u')$. By (2.8) it follows that $\mathcal{I}(p) \subset \mathcal{I}(u')$. By (2.36), it follows that

$$d_p(\mathcal{Q}(p), \mathcal{Q}(u')) > 2^{Z(n+1)}.$$

By the triangle inequality. Lemma 2.8 and (2.32) it follows that

$$\begin{aligned} d_p(\mathcal{Q}(p), \mathcal{Q}(p')) &\geq d_p(\mathcal{Q}(p), \mathcal{Q}(u')) - d_p(\mathcal{Q}(p'), \mathcal{Q}(u')) \\ &> 2^{Z(n+1)} - d_{p'}(\mathcal{Q}(p'), \mathcal{Q}(u')) \\ &\geq 2^{Z(n+1)} - 4. \end{aligned}$$

Since $Z \geq 3$ by (2.3), it follows that $\mathcal{Q}(p') \notin B_p(\mathcal{Q}(p), 1)$, so $\Omega(p') \not\subset \Omega(p)$ by (2.15). Hence, by (2.14), $\Omega(p) \cap \Omega(p') = \emptyset$. But if $x \in E(p) \cap E(p')$ then $\mathcal{Q}(x) \in \Omega(p) \cap \Omega(p')$. This is a contradiction, and the lemma follows. \square

Now we prove Proposition 2.4.

Proof of Proposition 2.4. To save some space, we will write

$$T_{\mathfrak{R}_j}^* = \sum_{u \in \mathcal{U}_j} \sum_{p \in \mathfrak{T}(u)} T_p^*.$$

By (7.27), we have for each j

$$T_{\mathfrak{R}_j}^* g = \sum_{u \in \mathcal{U}_j} \sum_{p \in \mathfrak{T}(u)} T_p^* g = \sum_{u \in \mathcal{U}_j} \sum_{p \in \mathfrak{T}(u)} T_p^* \mathbf{1}_{E_j} g = T_{\mathfrak{R}_j}^* \mathbf{1}_{E_j} g.$$

Hence, by Lemma 7.36,

$$\begin{aligned} \left\| \sum_{u \in \mathcal{U}} \sum_{p \in \mathfrak{T}(u)} T_p^* g \right\|_2^2 &= \left\| \sum_{j=1}^{2^n} T_{\mathfrak{R}_j}^* g \right\|_2^2 = \left\| \sum_{j=1}^{2^n} T_{\mathfrak{R}_j}^* \mathbf{1}_{E_j} g \right\|_2^2 \\ &= \int_X \left| \sum_{j=1}^{2^n} T_{\mathfrak{R}_j}^* \mathbf{1}_{E_j} g \right|^2 d\mu \\ &= \sum_{j=1}^{2^n} \int_X |T_{\mathfrak{R}_j}^* \mathbf{1}_{E_j} g|^2 + \sum_{j=1}^{2^n} \sum_{\substack{j'=1 \\ j' \neq j}}^{2^n} \int_X \overline{T_{\mathfrak{R}_j}^* \mathbf{1}_{E_j} g} T_{\mathfrak{R}_{j'}}^* \mathbf{1}_{E_{j'}} g d\mu. \end{aligned}$$

We use Lemma 7.37 to estimate each term in the first sum, and Lemma 7.38 to bound each term in the second sum:

$$\leq 2^{312a^3-n} \sum_{j=1}^{2^n} \|\mathbf{1}_{E_j} g\|_2^2 + 2^{862a^3-3n} \sum_{j=1}^{2^n} \sum_{j'=1}^{2^n} \|\mathbf{1}_{E_j} g\|_2 \|\mathbf{1}_{E_{j'}} g\|_2.$$

By Cauchy-Schwarz in the second two sums, this is at most

$$2^{862a^3} (2^{-n} + 2^n 2^{-3n}) \sum_{j=1}^n \|\mathbf{1}_{E_j} g\|_2^2,$$

and by disjointedness of the sets E_j , this is at most

$$2^{863a^3-n} \|g\|_2^2.$$

Taking adjoints and square roots, it follows that for all f

$$\left\| \sum_{u \in \mathfrak{U}} \sum_{p \in \mathfrak{T}(u)} T_p f \right\|_2 \leq 2^{432a^3 - \frac{n}{2}} \|f\|_2. \quad (7.67)$$

On the other hand, we have by disjointedness of the sets E_j from Lemma 7.39

$$\left\| \sum_{u \in \mathfrak{U}} \sum_{p \in \mathfrak{T}(u)} T_p f \right\|_2^2 = \left\| \sum_{j=1}^{2^n} \mathbf{1}_{E_j} T_{\mathfrak{H}_j} f \right\|_2^2 = \sum_{j=1}^{2^n} \|\mathbf{1}_{E_j} T_{\mathfrak{H}_j} f\|_2^2.$$

If $|f| \leq \mathbf{1}_F$ then we obtain from Lemma 7.37 and taking square roots that

$$\begin{aligned} &\leq 2^{257a^3} \text{dens}_2\left(\bigcup_{u \in \mathfrak{U}} \mathfrak{T}(u)\right)^{\frac{1}{2}} 2^{-\frac{n}{2}} \left(\sum_{j=1}^{2^n} \|f\|_2^2\right)^{\frac{1}{2}} \\ &= 2^{257a^3} \text{dens}_2\left(\bigcup_{u \in \mathfrak{U}} \mathfrak{T}(u)\right)^{\frac{1}{2}} \|f\|_2. \end{aligned} \quad (7.68)$$

Proposition 2.4 follows by taking the product of the $(2 - \frac{2}{q})$ -th power of (7.67) and the $(\frac{2}{q} - 1)$ -st power of (7.68). \square

8. PROOF OF PROP. 2.5, THE HÖLDER CANCELLATIVE CONDITION

We need the following auxiliary lemma. Recall that $\tau = 1/a$.

Lemma 8.1 (Lipschitz Holder approximation). *Let $z \in X$ and $R > 0$. Let $\varphi : X \rightarrow \mathbb{C}$ be a function supported in the ball $B := B(z, R)$ with finite norm $\|\varphi\|_{C^\tau(B)}$. Let $0 < t \leq 1$. There exists a function $\tilde{\varphi} : X \rightarrow \mathbb{C}$, supported in $B(z, 2R)$, such that for every $x \in X$*

$$|\varphi(x) - \tilde{\varphi}(x)| \leq t^\tau \|\varphi\|_{C^\tau(B)} \quad (8.1)$$

and

$$\|\tilde{\varphi}\|_{\text{Lip}(B(z, 2R))} \leq 2^{4a} t^{-1-a} \|\varphi\|_{C^\tau(B)}. \quad (8.2)$$

Proof. Define for $x, y \in X$ the Lipschitz and thus measurable function

$$L(x, y) := \max\{0, 1 - \frac{\rho(x, y)}{tR}\}. \quad (8.3)$$

We have that $L(x, y) \neq 0$ implies

$$y \in B(x, tR). \quad (8.4)$$

We have for $y \in B(x, 2^{-1}tR)$ that

$$|L(x, y)| \geq 2^{-1}. \quad (8.5)$$

Hence

$$\int L(x, y) d\mu(y) \geq 2^{-1} \mu(B(x, 2^{-1}tR)). \quad (8.6)$$

Let n be the smallest integer so that

$$2^n t \geq 1. \quad (8.7)$$

Iterating $n + 2$ times the doubling condition (1.5), we obtain

$$\int L(x, y) d\mu(y) \geq 2^{-1-a(n+2)} \mu(B(x, 2R)). \quad (8.8)$$

Now define

$$\tilde{\varphi}(x) := \left(\int L(x, y) d\mu(y) \right)^{-1} \int L(x, y) \varphi(y) d\mu(y).$$

Using that φ is supported in $B(z, R)$ and (8.4), we have that $\tilde{\varphi}$ is supported in $B(z, 2R)$.

We prove (8.1). For any $x \in X$, using that L is nonnegative,

$$\left(\int L(x, y) d\mu(y) \right) |\varphi(x) - \tilde{\varphi}(x)| \quad (8.9)$$

$$= \left| \int L(x, y) (\varphi(x) - \varphi(y)) d\mu(y) \right|. \quad (8.10)$$

Using (8.4), we estimate the last display by

$$\leq \int_{B(x, tR)} L(x, y) |\varphi(x) - \varphi(y)| d\mu(y). \quad (8.11)$$

Using the definition of $\|\varphi\|_{C^\tau(B)}$, we estimate the last display further by

$$\leq \left(\int_{B(x, tR)} L(x, y) \rho(x, y)^\tau d\mu(y) \right) \|\varphi\|_{C^\tau(B)} R^{-\tau}. \quad (8.12)$$

Using the condition on the domain of integration to estimate $\rho(x, y)$ by tR and then expanding the domain by positivity of the integrand, we estimate this further by

$$\leq \left(\int L(x, y) d\mu(y) \right) \|\varphi\|_{C^\tau(B)} t^\tau. \quad (8.13)$$

Dividing the string of inequalities from (8.9) to (8.13) by the positive integral of L proves (8.1).

We turn to (8.2). For every $x \in X$, we have

$$\left| \int L(x, y) d\mu(y) \right| |\tilde{\varphi}(x)| = \left| \int L(x, y) \varphi(y) d\mu(y) \right| \quad (8.14)$$

$$\leq \left| \int L(x, y) d\mu(y) \right| \sup_{x' \in X} |\varphi(x')|. \quad (8.15)$$

As φ is supported on B , dividing by the integral of L , we obtain

$$|\tilde{\varphi}(x)| \leq \sup_{x' \in B} |\varphi(x')| \leq \|\varphi\|_{C^\tau(B)}. \quad (8.16)$$

If $\rho(x, x') \geq R$, then we have by the triangle inequality

$$R \frac{|\tilde{\varphi}(x') - \tilde{\varphi}(x)|}{\rho(x, x')} \leq 2 \sup_{x'' \in X} |\tilde{\varphi}(x'')| \leq 2\|\varphi\|_{C^\tau(B)}. \quad (8.17)$$

Now assume $\rho(x, x') < R$. For $y \in X$ we have by the triangle inequality and a two fold case distinction for the maximum in the definition of L ,

$$|L(x, y) - L(x', y)| \leq \frac{\rho(x, x')}{tR}. \quad (8.18)$$

We compute with (8.18), first adding and subtracting a term in the integral,

$$\left(\int L(x, y) d\mu(y) \right) |\tilde{\varphi}(x') - \tilde{\varphi}(x)| = \quad (8.19)$$

$$\left| \int L(x, y) \tilde{\varphi}(x') - L(x, y) \tilde{\varphi}(x) + L(x', y) \tilde{\varphi}(x') - L(x', y) \tilde{\varphi}(x) d\mu(y) \right|. \quad (8.20)$$

Grouping the second and third and the first and fourth term, we obtain using the definition of $\tilde{\varphi}$ and Fubini,

$$\leq \left| \int (L(x', y) - L(x, y)) \varphi(y) d\mu(y) \right| \quad (8.21)$$

$$+ \left| \int L(x, y) d\mu(y) - \int L(x', y) d\mu(y) \right| |\tilde{\varphi}(x')| \quad (8.22)$$

$$\leq 2 \int |L(x, y) - L(x', y)| d\mu(y) \|\varphi\|_{C^\tau(B)}, \quad (8.23)$$

where in the last inequality we have used (8.16). Using further (8.18) and the support of L , we estimate the last display by

$$\leq 2 \frac{\rho(x, x')}{tR} \mu(B(x, tR) \cup B(x', tR)) \|\varphi\|_{C^\tau(B)}. \quad (8.24)$$

Using $\rho(x, x') < R$ and the triangle inequality, we estimate the last display by

$$\leq 2 \frac{\rho(x, x')}{tR} \mu(B(x, 2R)) \|\varphi\|_{C^\tau(B)}. \quad (8.25)$$

Dividing by the integral over L and using (8.8) and (8.7), we obtain

$$\frac{R|\tilde{\varphi}(x') - \tilde{\varphi}(x)|}{\rho(x, x')} \leq 2^{2+a(n+2)} t^{-1} \|\varphi\|_{C^\tau(B)} \leq 2^{2+3a} t^{-1-a} \|\varphi\|_{C^\tau(B)}. \quad (8.26)$$

Combining (8.17) and (8.26) using $a \geq 4$ and $t \leq 1$ and adding (8.16) proves (8.2) and completes the proof of Lemma 8.1. \square

We turn to the proof of Proposition 2.5. Let $z \in X$ and $R > 0$ and set $B = B(z, R)$. Let φ be given as in Proposition 2.5. Set

$$t := (1 + d_B(\vartheta, \theta))^{-\frac{\tau}{2+a}} \quad (8.27)$$

and define $\tilde{\varphi}$ as in Lemma 8.1. Let ϑ and θ be in Θ . Then

$$\left| \int e(\vartheta(x) - \theta(x)) \varphi(x) d\mu(x) \right| \quad (8.28)$$

$$\leq \left| \int e(\vartheta(x) - \theta(x)) \tilde{\varphi}(x) d\mu(x) \right| \quad (8.29)$$

$$+ \left| \int e(\vartheta(x) - \theta(x)) (\varphi(x) - \tilde{\varphi}(x)) d\mu(x) \right| \quad (8.30)$$

Using the cancellative condition (1.12) of Θ on the ball $B(z, 2R)$, the term (8.29) is bounded above by

$$2^a \mu(B(z, 2R)) \|\tilde{\varphi}\|_{\text{Lip}(B(z, 2R))} (1 + d_{B(z, 2R)}(\vartheta, \theta))^{-\tau}. \quad (8.31)$$

Using the doubling condition (1.5), the inequality (8.2), and the estimate $d_B \leq d_{B(z, 2R)}$ from the definition, we estimate (8.31) from above by

$$2^{6a} t^{-1-a} \mu(B) \|\varphi\|_{C^\tau(B)} (1 + d_B(\vartheta, \theta))^{-\tau}. \quad (8.32)$$

The term (8.30) we estimate using (8.1) and that ϑ and θ are real and thus $e(\vartheta)$ and $e(\theta)$ bounded in absolute value by 1. We obtain for (8.30) with (1.5) the upper bound

$$\mu(B(z, 2R)) t^\tau \|\varphi\|_{C^\tau(B)} \leq 2^a \mu(B) t^\tau \|\varphi\|_{C^\tau(B)}. \quad (8.33)$$

Using the definition (8.27) of t and adding (8.32) and (8.33) estimates (8.28) from above by

$$2^{6a} \mu(B) \|\varphi\|_{C^\tau(B)} (1 + d_B(\vartheta, \theta))^{-\frac{\tau}{2+a}} \quad (8.34)$$

$$+ 2^a \mu(B) \|\varphi\|_{C^\tau(B)} (1 + d_B(\vartheta, \theta))^{-\frac{\tau^2}{2+a}}. \quad (8.35)$$

$$\leq 2^{1+6a} \mu(B) \|\varphi\|_{C^\tau(B)} (1 + d_B(\vartheta, \theta))^{-\frac{\tau^2}{2+a}}, \quad (8.36)$$

where we used $\tau \leq 1$. This completes the proof of Proposition 2.5.

9. PROOF OF PROP. 2.6, VITALI COVERING AND HARDY–LITTLEWOOD

We begin with a classical representation of the Lebesgue norm.

Lemma 9.1 (layer cake representation). *Let $1 \leq p < \infty$. Then for any measurable function $u : X \rightarrow [0, \infty)$ on the measure space X relative to the measure μ we have*

$$\|u\|_p^p = p \int_0^\infty \lambda^{p-1} \mu(\{x : u(x) \geq \lambda\}) d\lambda. \quad (9.1)$$

Proof. The left-hand side of (9.1) is by definition

$$\int_X u(x)^p d\mu(x). \quad (9.2)$$

Writing $u(x)$ as an elementary integral in λ and then using Fubini, we write for the last display

$$= \int_X \int_0^{u(x)} p \lambda^{p-1} d\lambda d\mu(x) \quad (9.3)$$

$$= p \int_0^\infty \lambda^{p-1} \mu(\{x : u(x) \geq \lambda\}) d\lambda. \quad (9.4)$$

This proves the lemma. \square

We turn to the proof of Proposition 2.6. Let the collection \mathcal{B} be given. We first show (2.43).

We recursively choose a finite sequence $B_i \in \mathcal{B}$ for $i \geq 0$ as follows. Assume $B_{i'}$ is already chosen for $0 \leq i' < i$. If there exists a ball $B_i \in \mathcal{B}$ so that B_i is disjoint from all $B_{i'}$ with $0 \leq i' < i$, then choose such a ball $B_i = B(x_i, r_i)$ with maximal r_i .

If there is no such ball, stop the selection and set $i'' := i$.

By disjointedness of the chosen balls and since $0 \leq u$, we have

$$\sum_{0 \leq i < i''} \int_{B_i} u(x) d\mu(x) \leq \int_X u(x) d\mu(x). \quad (9.5)$$

By (2.42), we conclude

$$\lambda \sum_{0 \leq i < i''} \mu(B_i) \leq \int_X u(x) d\mu(x). \quad (9.6)$$

Let $x \in \bigcup \mathcal{B}$. Choose a ball $B' = B(x', r') \in \mathcal{B}$ such that $x \in B'$. If B' is one of the selected balls, then

$$x \in \bigcup_{0 \leq i < i''} B_i \subset \bigcup_{0 \leq i < i''} B(x_i, 3r_i). \quad (9.7)$$

If B' is not one of the selected balls, then as it is not selected at time i'' , there is a selected ball B_i with $B' \cap B_i \neq \emptyset$. Choose such B_i with minimal index i . As B' is therefore disjoint from all balls $B_{i'}$ with $i' < i$ and as it was not selected in place of B_i , we have $r_i \geq r'$.

Using a point y in the intersection of B_i and B' , we conclude by the triangle inequality

$$\rho(x_i, x') \leq \rho(x_i, y) + \rho(x', y) \leq r_i + r' \leq 2r_i. \quad (9.8)$$

By the triangle inequality again, we further conclude

$$\rho(x_i, x) \leq \rho(x_i, x') + \rho(x', x) \leq 2r_i + r' \leq 3r_i. \quad (9.9)$$

It follows that

$$x \in \bigcup_{0 \leq i < i''} B(x_i, 3r_i). \quad (9.10)$$

With (9.7) and (9.10), we conclude

$$\bigcup \mathcal{B} \subset \bigcup_{0 \leq i < i''} B(x_i, 3r_i). \quad (9.11)$$

With the doubling property (1.5) applied twice, we conclude

$$\mu\left(\bigcup \mathcal{B}\right) \leq \sum_{0 \leq i < i''} \mu(B(x_i, 3r_i)) \leq 2^{2a} \sum_{0 \leq i < i''} \mu(B_i). \quad (9.12)$$

With (9.6) and (9.12) we conclude (2.43).

We turn to the proof of (2.44). We first consider the case $p_1 = 1$ and recall $M_{\mathcal{B}} = M_{\mathcal{B},1}$. We write for the p_2 -th power of left-hand side of (2.44) with Lemma 9.1 and a change of variables

$$\|M_{\mathcal{B}}u(x)\|_{p_2}^{p_2} = p_2 \int_0^\infty \lambda^{p_2-1} \mu(\{x : M_{\mathcal{B}}u(x) \geq \lambda\}) d\lambda \quad (9.13)$$

$$= 2^{p_2} p_2 \int_0^\infty \lambda^{p_2-1} \mu(\{x : M_{\mathcal{B}} u(x) \geq 2\lambda\}) d\lambda. \quad (9.14)$$

Fix $\lambda \geq 0$ and let $x \in X$ satisfy $M_{\mathcal{B}} u(x) \geq 2\lambda$. By definition of $M_{\mathcal{B}}$, there is a ball $B' \in \mathcal{B}$ such that $x \in B'$ and

$$\int_{B'} u(y) d\mu(y) \geq 2\lambda \mu(B'). \quad (9.15)$$

Define $u_\lambda(y) := 0$ if $|u(y)| < \lambda$ and $u_\lambda(y) := u(y)$ if $|u(y)| \geq \lambda$. Then with (9.15)

$$\int_{B'} u_\lambda(y) d\mu(y) = \int_{B'} u(y) d\mu(y) - \int_{B'} (u - u_\lambda)(y) d\mu(y) \quad (9.16)$$

$$\geq 2\lambda \mu(B') - \int_{B'} (u - u_\lambda)(y) d\mu(y). \quad (9.17)$$

As $(u - u_\lambda)(y) \leq \lambda$ by definition, we can estimate the last display by

$$\geq 2\lambda \mu(B') - \int_{B'} \lambda d\mu(y) = \lambda \mu(B'). \quad (9.18)$$

Hence x is contained in $\bigcup(\mathcal{B}_\lambda)$, where \mathcal{B}_λ is the collection of balls B'' in \mathcal{B} such that

$$\int_{B''} u_\lambda(y) d\mu(y) \geq \lambda \mu(B''). \quad (9.19)$$

We have thus seen

$$\{x : M_{\mathcal{B}} u(x) \geq 2\lambda\} \subset \bigcup \mathcal{B}_\lambda. \quad (9.20)$$

Applying (2.43) to the collection \mathcal{B}_λ gives

$$\lambda \mu(\{x : M_{\mathcal{B}} u(x) \geq 2\lambda\}) \leq 2^{2a} \int u_\lambda(x) dx. \quad (9.21)$$

With Lemma 9.1,

$$\lambda \mu(\{x : M_{\mathcal{B}} u(x) \geq 2\lambda\}) \leq 2^{2a} \int_0^\infty \mu(\{x : |u_\lambda(x)| \geq \lambda'\}) d\lambda'. \quad (9.22)$$

By definition of h_λ , making a case distinction between $\lambda \geq \lambda'$ and $\lambda < \lambda'$, we see that

$$\mu(\{x : |u_\lambda(x)| \geq \lambda'\}) \leq \mu(\{x : |u(x)| \geq \max(\lambda, \lambda')\}). \quad (9.23)$$

We obtain with (9.14), (9.22), and (9.23)

$$\|M_{\mathcal{B}} u(x)\|_{p_2}^{p_2} \quad (9.24)$$

$$\leq 2^{p_2+2a} p_2 \int_0^\infty \lambda^{p_2-2} \int_0^\infty \mu(\{x : |u(x)| \geq \max(\lambda, \lambda')\}) d\lambda' d\lambda. \quad (9.25)$$

We split the integral into $\lambda \geq \lambda'$ and $\lambda < \lambda'$ and resolve the maximum correspondingly. We have for $\lambda \geq \lambda'$ with Lemma 9.1

$$\int_0^\infty \lambda^{p_2-2} \int_0^\lambda \mu(\{x : |u(x)| \geq \lambda\}) d\lambda' d\lambda \quad (9.26)$$

$$= \int_0^\infty \lambda^{p_2-1} \mu(\{x : |u(x)| \geq \lambda\}) d\lambda. \quad (9.27)$$

$$= p_2^{-1} \|u\|_{p_2}^{p_2}. \quad (9.28)$$

We have for $\lambda < \lambda'$ with Fubini and Lemma 9.1

$$\int_0^\infty \lambda^{p_2-2} \int_\lambda^\infty \mu(\{x : |u(x)| \geq \lambda'\}) d\lambda' d\lambda. \quad (9.29)$$

$$= \int_0^\infty \int_0^{\lambda'} \lambda^{p_2-2} \mu(\{x : |u(x)| \geq \lambda'\}) d\lambda d\lambda'. \quad (9.30)$$

$$= (p_2 - 1)^{-1} \int_0^\infty (\lambda')^{p_2-1} \mu(\{x : |u(x)| \geq \lambda'\}) d\lambda'. \quad (9.31)$$

$$= (p_2 - 1)^{-1} p_2^{-1} \|u\|_{p_2}^{p_2}. \quad (9.32)$$

Adding the two estimates (9.28) and (9.32) gives

$$\|M_{\mathcal{B}}u(x)\|_{p_2}^{p_2} \leq 2^{p_2+2a} (1 + (p_2 - 1)^{-1}) \|u\|_{p_2}^{p_2} = 2^{p_2+2a} p_2 (p_2 - 1)^{-1} \|u\|_{p_2}^{p_2}. \quad (9.33)$$

With $a \geq 1$ and $p_2 > 1$, taking the p_2 -th root, we obtain (2.44). We turn to the case of general $1 \leq p_1 < p_2$. We have

$$M_{\mathcal{B},p_1}u = (M_{\mathcal{B}}(|u|^{p_1}))^{\frac{1}{p_1}}. \quad (9.34)$$

Applying the special case of (2.44) for $M_{\mathcal{B}}$ gives

$$\|M_{\mathcal{B},p_1}u\|_{p_2} = \|M_{\mathcal{B}}(|u|^{p_1})\|_{p_2/p_1}^{\frac{1}{p_1}} \quad (9.35)$$

$$\leq 2^{2a} (p_2/p_1) (p_2/p_1 - 1)^{-1} \|(|u|^{p_1})\|_{p_2/p_1}^{\frac{1}{p_1}} = 2^{2a} p_2 (p_2 - p_1)^{-1} \|u\|_{p_2}. \quad (9.36)$$

This proves (2.44) in general.

Now we construct the operator M satisfying (2.45) and (2.46).

Lemma 9.2 (covering separable space). *For each $r > 0$, there exists a countable collection $C(r) \subset X$ of points such that*

$$X \subset \bigcup_{c \in C(r)} B(c, r).$$

Proof. It clearly suffices to construct finite collections $C(r, k)$ such that

$$B(o, r2^k) \subset \bigcup_{c \in C(r, k)} B(c, r),$$

since then the collection $C(r) = \bigcup_{k \in \mathbb{N}} C(r, k)$ has the desired property.

Suppose that $Y \subset B(o, r2^k)$ is a collection of points such that for all $y, y' \in Y$ with $y \neq y'$, we have $\rho(y, y') \geq r$. Then the balls $B(y, r/2)$ are pairwise disjoint and contained in $B(o, r2^{k+1})$. If $y \in B(o, r)$, then $B(o, r2^{k+1}) \subset B(y, r2^{k+2})$. Thus, by the doubling property (1.5),

$$\mu(B(y, \frac{r}{2})) \geq 2^{-(k+2)a} \mu(B(o, r2^{k+1})).$$

Thus, we have

$$\mu(B(o, r2^{k+1})) \geq \sum_{y \in Y} \mu(B(y, \frac{r}{2})) \geq |Y| 2^{-(k+2)a} \mu(B(o, r2^{k+1})).$$

We conclude that $|Y| \leq 2^{(k+2)a}$. In particular, there exists a set Y of maximal cardinality. Define $C(r, k)$ to be such a set.

If $x \in B(o, r2^k)$ and $x \notin C(r, k)$, then there must exist $y \in C(r, k)$ with $\rho(x, y) < r$. Thus $C(r, k)$ has the desired property. \square

For each $k \in \mathbb{N}$ we choose a countable set $C(2^k)$ as in the lemma. Define

$$\mathcal{B}_\infty = \{B(c, 2^k) : c \in C(2^k), k \in \mathbb{N}\}.$$

By Lemma 9.2, this is a countable collection of balls. We choose an enumeration $\mathcal{B}_\infty = \{B_1, \dots\}$ and define

$$\mathcal{B}_n = \{B_1, \dots, B_n\}.$$

We define

$$Mw := 2^{2a} \sup_{n \in \mathbb{N}} M_{\mathcal{B}_n} w.$$

This function is measurable for each measurable w , since it is a countable supremum of measurable functions. Estimate (2.46) follows immediately from (2.44) and the monotone convergence theorem.

It remains to show (2.45). Let $B = B(x, r) \subset X$. Let k be the smallest integer such that $2^k \geq r$, in particular we have $2^k < 2r$. By definition of $C(2^k)$, there exists $c \in C(2^k)$ with $x \in B(c, 2^k)$. By the triangle inequality, we have $B(c, 2^k) \subset B(x, 4r)$, and hence by the doubling property (1.5)

$$\mu(B(c, 2^k)) \leq 2^{2a} \mu(B(x, r)).$$

It follows that for each $z \in B(x, r)$

$$\begin{aligned} \frac{1}{\mu(B(x, r))} \int_{B(x, r)} |w(y)| d\mu(y) &\leq \frac{2^{2a}}{\mu(B(c, 2^k))} \int_{B(c, 2^k)} |w(y)| d\mu(y) \\ &\leq Mw(z). \end{aligned}$$

This completes the proof.

10. PROOF OF THEOREM 1.1

The convergence of partial Fourier sums is proved in Subsection 10.1 in two steps. In the first step, we establish convergence on a suitable dense subclass of functions. We choose piece-wise constant functions as subclass, the convergence is stated in Lemma 10.2 and proved in Subsection 10.2. In the second step, one controls the relevant error of approximating a general function by a function in the subclass. This is stated in Lemma 10.3 and proved in Subsection 10.8. The proof relies on a bound on the real Carleson maximal operator stated in Lemma 10.4 and proved in Subsection 10.9. This latter proof refers to the main Carleson Theorem 1.2. Two assumptions in Theorem 1.2 require more work. The boundedness of the nontangential maximal operator T^* defined in (1.16) is established in Lemma 10.5 using L^2 and weak L^1 bounds for the Hilbert transform, Lemmas 10.6 and 10.7. These lemmas are proved in Subsections 10.3, 10.4 and 10.5. The cancellative property is verified by Lemma 10.8, which is proved in Subsection 10.6. Several further auxiliary lemmas are stated and proved in Subsection 10.1, the proof of one of these auxiliary lemmas, Lemma 10.12, is done in Subsection 10.7.

All subsections past Subsection 10.1 are mutually independent.

Subsections 10.3 uses bounds for the Hardy–Littlewood maximal function on the real line. There may be a better path through Lean than we choose here. We refer to Proposition 2.6, the assumptions of it, namely that the real line fits into the setting of Section 2, is done in Subsection 10.9.

10.1. The classical Carleson theorem. Let a uniformly continuous 2π -periodic function $f : \mathbb{R} \rightarrow \mathbb{C}$, bounded in absolute value by 1, be given. Let $0 < \epsilon < 1$ be given.

By uniform continuity of f , there is a $0 < \delta < \pi$ such that for all $x, x' \in \mathbb{R}$ with $|x - x'| \leq \delta$ we have

$$|f(x) - f(x')| \leq 2^{-2^{50}} \epsilon^2. \quad (10.1)$$

Let K be the Gaussian bracket of $\frac{2\pi}{\delta} + 1$, that is the unique integer with

$$K \leq \frac{2\pi}{\delta} + 1 < K + 1, \quad (10.2)$$

and note that $K > 2$ by assumption on δ . For each $x \in \mathbb{R}$, let $k(x)$ be the Gaussian bracket of $Kx/2\pi$, that is the unique integer such that

$$k(x) \leq \frac{Kx}{2\pi} < k(x) + 1. \quad (10.3)$$

Define

$$f_0(x) := f\left(\frac{2\pi k(x)}{K}\right). \quad (10.4)$$

Lemma 10.1 (piecewise constant approximation). *The function f_0 is measurable. The function f_0 is 2π -periodic. The function f_0 satisfies for all $x \in \mathbb{R}$:*

$$|f(x) - f_0(x)| \leq 2^{-2^{50}} \epsilon^2, \quad (10.5)$$

$$|f_0(x)| \leq 1. \quad (10.6)$$

Proof. Let F be any set in \mathbb{C} . We show that $f_0^{-1}(F)$ is measurable. As $Z := \{2\pi k/K, k \in \mathbb{Z}\}$ is countable, it suffices to show that $f_0^{-1}(F) \setminus Z$ is measurable. It then suffices to show that $f_0^{-1}(F) \setminus Z$ is open. Let $x \in f_0^{-1}(F) \setminus Z$. There is a k such that $x \in [2\pi k/K, 2\pi(k+1)/K)$. As $x \notin Z$, we have that $x \in (2\pi k/K, 2\pi(k+1)/K)$. But f_0 is constant on $(2\pi k/K, 2\pi(k+1)/K)$, hence $(2\pi k/K, 2\pi(k+1)/K)$ is a subset of $f_0^{-1}(F) \setminus Z$. This proves that $f_0^{-1}(F) \setminus Z$ is open and completes the proof of measurability of f_0 .

To see that f_0 is 2π -periodic, we observe by applying (10.3) to $x + 2\pi$ and subtracting K that

$$k(x + 2\pi) - K \leq \frac{Kx}{2\pi} < k(x + 2\pi) - K + 1 \quad (10.7)$$

and hence

$$k(x + 2\pi) - K = k(x) \quad (10.8)$$

and hence by Definition (10.4)

$$f_0(x + 2\pi) = f\left(\frac{2\pi k(x + 2\pi)}{K}\right) = f\left(\frac{2\pi(k(x) + K)}{K}\right) = f\left(\frac{2\pi k(x)}{K} + 2\pi\right), \quad (10.9)$$

which by 2π -periodicity of f is equal to

$$= f\left(\frac{2\pi k(x)}{K}\right) = f_0(x). \quad (10.10)$$

This proves that f_0 is 2π -periodic

To see (10.5), observe that by (10.3) used twice and then by definition of K we have

$$0 \leq x - \frac{2\pi k(x)}{K} \leq \frac{2\pi}{K} \leq \delta. \quad (10.11)$$

Hence, by choice of δ , for all $x \in \mathbb{R}$, we obtain (10.5).

Finally (10.5) follows because $|f|$ by assumption is bounded by 1. This completes the proof of the lemma. \square

Define the set E_1 to be

$$E_1 = \bigcup_{k=0}^K \left[\frac{2\pi}{K} \left(k - \frac{\epsilon}{16\pi} \right), \frac{2\pi}{K} \left(k + \frac{\epsilon}{16\pi} \right) \right]. \quad (10.12)$$

Then we have for the Lebesgue measure of E_1 , using $K > 1$,

$$|E_1| \leq \sum_{k=0}^K \frac{\epsilon}{4K} = \frac{\epsilon}{4} \left(1 + \frac{1}{K} \right) \leq \frac{\epsilon}{2}. \quad (10.13)$$

We prove in Subsection 10.2:

Lemma 10.2 (convergence for piecewise constant). *For all $N > \frac{2^{25}K^2}{\epsilon^3}$ and*

$$x \in [0, 2\pi) \setminus E_1 \quad (10.14)$$

we have

$$|S_N f_0(x) - f_0(x)| \leq \frac{\epsilon}{4}. \quad (10.15)$$

We prove in Subsection 10.8:

Lemma 10.3 (control approximation effect). *There is a set $E_2 \subset \mathbb{R}$ with Lebesgue measure $|E_2| \leq \frac{\epsilon}{2}$ such that for all*

$$x \in [0, 2\pi) \setminus E_2 \quad (10.16)$$

we have

$$\sup_{N \geq 0} |S_N f(x) - S_N f_0(x)| \leq \frac{\epsilon}{4}. \quad (10.17)$$

Define

$$E := E_1 \cup E_2. \quad (10.18)$$

Then

$$|E| \leq |E_1| + |E_2| \leq \frac{\epsilon}{2} + \frac{\epsilon}{2} \leq \epsilon. \quad (10.19)$$

Let N_0 be the unique integer such that

$$N_0 - 1 \leq 2^{25} K^2 \epsilon^{-3} < N_0. \quad (10.20)$$

For every

$$x \in [0, 1) \setminus E, \quad (10.21)$$

and every $N > N_0$ we have by the triangle inequality

$$\begin{aligned} & |f(x) - S_N f(x)| \\ & \leq |f(x) - f_0(x)| + |f_0(x) - S_N f_0(x)| + |S_N f_0(x) - S_N f(x)|. \end{aligned} \quad (10.22)$$

Using (10.1) and Lemmas 10.2 and 10.3, we estimate (10.22) by

$$\leq 2^{-250} \epsilon^2 + \frac{\epsilon}{4} + \frac{\epsilon}{4} \leq \epsilon. \quad (10.23)$$

This shows (1.3) for the given E and N_0 and completes the proof of Theorem 1.1.

Let $\kappa : \mathbb{R} \rightarrow \mathbb{R}$ be the function defined by $\kappa(0) = 0$ and for $0 < |x| < 1$

$$\kappa(x) = \frac{1 - |x|}{1 - e^{ix}} \quad (10.24)$$

and for $|x| \geq 1$

$$\kappa(x) = 0. \quad (10.25)$$

Note that this function is continuous at every point x with $|x| > 0$.

The proof of Lemma 10.3 will use the following Lemma 10.4, which itself is proven in Subsection 10.9 as an application of Theorem 1.2.

Lemma 10.4 (real Carleson). *Let F, G be Borel subsets of \mathbb{R} with finite measure. Let f be a bounded measurable function on \mathbb{R} with $|f| \leq \mathbf{1}_F$. Then*

$$\left| \int_G T f(x) dx \right| \leq 2^{240} |F|^{\frac{1}{2}} |G|^{\frac{1}{2}}, \quad (10.26)$$

where

$$T f(x) = \sup_{n \in \mathbb{Z}} \sup_{r > 0} \left| \int_{r < |x-y| < 1} f(y) \kappa(x-y) e^{iny} dy \right|. \quad (10.27)$$

One of the main assumption of Theorem 1.2, concerning the operator T_* defined in (1.16), is verified by the following lemma, which is proved in Subsection 10.3.

Lemma 10.5 (nontangential Hilbert). *For every bounded measurable function g with bounded support we have*

$$\|T_* g\|_2 \leq 2^{43} \|g\|_2, \quad (10.28)$$

where

$$T_* g(x) := \sup_{0 < r_1 < r_2 < 1} \sup_{|x-x'| < r_1} \frac{1}{2\pi} \left| \int_{r_1 < |x'-y| < r_2} g(y) \kappa(x'-y) dy \right|. \quad (10.29)$$

The proof of Lemma 10.5 relies on the next two auxiliary Lemmas.

For $r \in (0, 1)$, $x \in \mathbb{R}$ and a bounded, measurable function g on \mathbb{R} with bounded support, we define

$$H_r g(x) := \int_{r < |x-y| < 1} g(y) \kappa(x-y) dy. \quad (10.30)$$

The following Lemma is proved in Section 10.4

Lemma 10.6 (Hilbert strong 2 2). *Let $0 < r < 1$. Let f be a bounded, measurable function on \mathbb{R} with bounded support. Then*

$$\|H_r f\|_2 \leq 2^{13} \|f\|_2. \quad (10.31)$$

The following Lemma is proved in Subsection 10.5

Lemma 10.7 (Hilbert weak 1 1). *Let f be a bounded measurable function on \mathbb{R} with bounded support. Let $\alpha > 0$. Then for all $r \in (0, 1)$, we have*

$$\mu(\{x \in \mathbb{R} : |H_r f(x)| > \alpha\}) \leq \frac{2^{19}}{\alpha} \int |f(y)| dy. \quad (10.32)$$

The next lemma will be used to verify that the collection Θ of modulation functions in our application of Theorem 1.2 satisfies the condition (1.12). It is proved in Subsection 10.6.

Lemma 10.8 (van der Corput). *Let $\alpha < \beta$ be real numbers. Let $g : \mathbb{R} \rightarrow \mathbb{C}$ be a measurable function and assume*

$$\|g\|_{Lip(\alpha, \beta)} := \sup_{\alpha \leq x \leq \beta} |g(x)| + |\beta - \alpha| \sup_{\alpha \leq x < y \leq \beta} \frac{|g(y) - g(x)|}{|y - x|} < \infty. \quad (10.33)$$

Then for any $0 \leq \alpha < \beta \leq 2\pi$ we have

$$\int_{\alpha}^{\beta} g(x) e^{-inx} dx \leq 2\pi |\beta - \alpha| \|g\|_{Lip(\alpha, \beta)} (1 + |n| |\beta - \alpha|)^{-1}. \quad (10.34)$$

We close this section with six lemmas that are used across the following subsections.

Lemma 10.9 (mean zero oscillation). *Let $n \in \mathbb{Z}$ with $n \neq 0$, then*

$$\int_0^{2\pi} e^{inx} dx = 0. \quad (10.35)$$

Proof. We have

$$\int_0^{2\pi} e^{inx} dx = \left[\frac{1}{in} e^{inx} \right]_0^{2\pi} = \frac{1}{in} (e^{2\pi in} - e^{2\pi i 0}) = \frac{1}{in} (1 - 1) = 0. \quad \square$$

Lemma 10.10 (Dirichlet kernel). *We have for every 2π -periodic bounded measurable f and every $N \geq 0$*

$$S_N f(x) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(y) K_N(x - y) dy \quad (10.36)$$

where K_N is the 2π -periodic continuous function of \mathbb{R} given by

$$\sum_{n=-N}^N e^{inx'}. \quad (10.37)$$

We have for $e^{ix'} \neq 1$ that

$$K_N(x') = \frac{e^{iNx'}}{1 - e^{-ix'}} + \frac{e^{-iNx'}}{1 - e^{ix'}}. \quad (10.38)$$

Proof. We have by definitions and interchanging sum and integral

$$\begin{aligned} S_N f(x) &= \sum_{n=-N}^N \hat{f}_n e^{inx} \\ &= \sum_{n=-N}^N \frac{1}{2\pi} \int_{-\pi}^{\pi} f(y) e^{in(x-y)} dy \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(y) \sum_{n=-N}^N e^{in(x-y)} dy. \end{aligned} \quad (10.39)$$

This proves the first statement of the lemma. By a telescoping sum, we have for every $x' \in \mathbb{R}$

$$\left(e^{\frac{1}{2}ix'} - e^{-\frac{1}{2}ix'}\right) \sum_{n=-N}^N e^{inx'} = e^{(N+\frac{1}{2})ix'} - e^{-(N+\frac{1}{2})ix'}. \quad (10.40)$$

If $e^{ix'} \neq 1$, the first factor on the left-hand side is not 0 and we may divide by this factor to obtain

$$\sum_{n=-N}^N e^{inx'} = \frac{e^{i(N+\frac{1}{2})x'}}{e^{\frac{1}{2}ix'} - e^{-\frac{1}{2}ix'}} - \frac{e^{-i(N+\frac{1}{2})x'}}{e^{\frac{1}{2}ix'} - e^{-\frac{1}{2}ix'}} = \frac{e^{iNx'}}{1 - e^{-ix'}} + \frac{e^{-iNx'}}{1 - e^{ix'}}. \quad (10.41)$$

This proves the second part of the lemma. \square

Lemma 10.11 (lower secant bound). *Let $\eta > 0$ and $-2\pi + \eta \leq x \leq 2\pi - \eta$ with $|x| \geq \eta$. Then*

$$|1 - e^{ix}| \geq \frac{\eta}{8} \quad (10.42)$$

Proof. We have

$$|1 - e^{ix}| = \sqrt{(1 - \cos(x))^2 + \sin^2(x)} \geq |\sin(x)|.$$

If $0 \leq x \leq \frac{\pi}{2}$, then we have from concavity of \sin on $[0, \pi]$ and $\sin(0) = 0$ and $\sin(\frac{\pi}{2}) = 1$

$$|\sin(x)| \geq \frac{2}{\pi}x \geq \frac{2}{\pi}\eta.$$

When $x \in \frac{m\pi}{2} + [0, \frac{\pi}{2}]$ for $m \in \{-4, -3, -2, -1, 1, 2, 3\}$ one can argue similarly. \square

The following lemma will be proved in Subsection 10.7.

Lemma 10.12 (spectral projection bound). *Let f be a bounded 2π -periodic measurable function. Then, for all $N \geq 0$*

$$\|S_N f\|_{L^2[-\pi, \pi]} \leq \|f\|_{L^2[-\pi, \pi]}. \quad (10.43)$$

Lemma 10.13 (Hilbert kernel bound). *For $x, y \in \mathbb{R}$ with $x \neq y$ we have*

$$|\kappa(x - y)| \leq 2^4(2|x - y|)^{-1}. \quad (10.44)$$

Proof. Fix $x \neq y$. If $\kappa(x - y)$ is zero, then (10.44) is evident. Assume $\kappa(x - y)$ is not zero, then $0 < |x - y| < 1$. We have

$$|\kappa(x - y)| = \left| \frac{1 - |x - y|}{1 - e^{i(x-y)}} \right|. \quad (10.45)$$

We estimate with Lemma 10.11

$$|\kappa(x - y)| \leq \frac{1}{|1 - e^{i(x-y)}|} \leq \frac{8}{|x - y|}. \quad (10.46)$$

This proves (10.44) in the given case and completes the proof of the lemma. \square

Lemma 10.14 (Hilbert kernel regularity). *For $x, y, y' \in \mathbb{R}$ with $x \neq y, y'$ and*

$$2|y - y'| \leq |x - y|, \quad (10.47)$$

we have

$$|\kappa(x - y) - \kappa(x - y')| \leq 2^{10} \frac{1}{|x - y|} \frac{|y - y'|}{|x - y|}. \quad (10.48)$$

Proof. Upon replacing y by $y - x$ and y' by $y' - x$ on the left-hand side of (10.47), we can assume that $x = 0$. Then the assumption (10.47) implies that y and y' have the same sign. Since $\kappa(y) = \bar{\kappa}(-y)$ we can assume that they are both positive. Then it follows from (10.47) that

$$\frac{y}{2} \leq y'.$$

We distinguish four cases. If $y, y' \leq 1$, then we have

$$|\kappa(-y) - \kappa(-y')| = \left| \frac{1 - y}{1 - e^{-iy}} - \frac{1 - y'}{1 - e^{-iy'}} \right|$$

and by the fundamental theorem of calculus

$$= \left| \int_{y'}^y \frac{-1 + e^{-it} + i(1 - t)e^{it}}{(1 - e^{-it})^2} dt \right|.$$

Using $y' \geq \frac{y}{2}$ and Lemma 10.11, we bound this by

$$\leq |y - y'| \sup_{\frac{y}{2} \leq t \leq 1} \frac{3}{|1 - e^{-it}|^2} \leq 3|y - y'| \left(8\frac{2}{y}\right)^2 \leq 2^{10} \frac{|y - y'|}{|y|^2}.$$

If $y \leq 1$ and $y' > 1$, then $\kappa(-y') = 0$ and we have from the first case

$$|\kappa(-y) - \kappa(-y')| = |\kappa(-y) - \kappa(-1)| \leq 2^{10} \frac{|y - 1|}{|y|^2} \leq 2^{10} \frac{|y - y'|}{|y|^2}.$$

Similarly, if $y > 1$ and $y' \leq 1$, then $\kappa(-y) = 0$ and we have by the same computation as for the first case

$$|\kappa(-y) - \kappa(-y')| = |\kappa(-y') - \kappa(-1)| \leq 2^{10} \frac{|y - 1|}{|y|^2} \leq 2^{10} \frac{|y - y'|}{|y|^2}.$$

Finally, if $y, y' > 1$ then

$$|\kappa(-y) - \kappa(-y')| = 0 \leq 2^{10} \frac{|y - y'|}{|y|^2}.$$

□

10.2. Piecewise constant functions. We first compute the partial Fourier sums for constant functions.

Lemma 10.15 (constant function). *If h satisfies $h(x) = h(0)$ for all $x \in \mathbb{R}$, then for all $N \geq 0$ we have $S_N h = h$.*

Proof. We compute with Lemma 10.9 for $n \in \mathbb{Z}$ with $n \neq 0$,

$$\hat{h}_n = \frac{1}{2\pi} \int_0^{2\pi} h(y) e^{-iny} dy = \frac{h(0)}{2\pi} \int_0^{2\pi} e^{-iny} dy = 0. \quad (10.49)$$

hence for $N \geq 0$

$$S_n h(x) = \sum_{n=-N}^N \widehat{h}_n e^{inx} = \widehat{h}_0 = \frac{1}{2\pi} \int_0^{2\pi} h(y) dy = \frac{h(0)}{2\pi} \int_0^{2\pi} dy = h(x). \quad (10.50)$$

This proves the lemma. \square

Lemma 10.16 (Dirichlet kernel bound). *Let $\eta > 0$. Let*

$$-2\pi + \eta \leq \alpha < \beta \leq 2\pi - \eta \quad (10.51)$$

and assume

$$[\alpha, \beta] \cap [-\eta, \eta] = \emptyset. \quad (10.52)$$

Then

$$\left| \int_{\alpha}^{\beta} \frac{e^{iNx}}{1 - e^{-ix}} dx \right| \leq \frac{64}{N} \left(\frac{|\beta - \alpha|}{\eta^2} + \frac{1}{\eta} \right). \quad (10.53)$$

Proof. Note first that the integral in (10.53) is well defined as the denominator of the integrand is bounded away from zero by Lemma 10.11 and thus the integrand is continuous. We have with partial integration

$$\begin{aligned} & \int_{\alpha}^{\beta} \frac{e^{-iNx}}{1 - e^{ix}} dx \\ &= \frac{1}{-iN} \int_{\alpha}^{\beta} e^{-iNx} \frac{ie^{ix}}{(1 - e^{ix})^2} dx + \left[\frac{1}{-iN} \frac{e^{-iNx}}{1 - e^{ix}} \right]_{\alpha}^{\beta}. \end{aligned} \quad (10.54)$$

We estimate the two summands in (10.54) separately. We have for the first summand in (10.54) with Lemma 10.11,

$$\begin{aligned} & \left| \frac{1}{-iN} \int_{\alpha}^{\beta} e^{-iNx} \frac{ie^{ix}}{(1 - e^{ix})^2} dx \right| \\ & \leq \frac{1}{N} \int_{\alpha}^{\beta} |1 - e^{-ix}|^{-2} dx \leq \frac{|\beta - \alpha|}{N} \sup_{x \in [\alpha, \beta]} |1 - e^{-ix}|^{-2} \end{aligned} \quad (10.55)$$

$$\leq \frac{64|\beta - \alpha|}{\eta^2 N} \quad (10.56)$$

We have for the second summand in (10.54) with Lemma 10.11 again

$$\begin{aligned} & \left| \left[\frac{1}{-iN} \frac{e^{-iNx}}{1 - e^{ix}} \right]_{\alpha}^{\beta} \right| \\ & \leq \frac{1}{N} (|1 - e^{i\alpha}|^{-1} + |1 - e^{i\beta}|^{-1}) \leq \frac{16}{\eta N}. \end{aligned} \quad (10.57)$$

Using the triangle inequality in (10.54) and applying the estimates (10.55) and (10.57) proves the lemma. \square

Let \mathcal{G} be the class of 2π -periodic measurable functions g which satisfy for all $x \in \mathbb{R}$

$$g\left(\frac{2\pi k(x)}{K}\right) = g(x) \quad (10.58)$$

and

$$|g(x)| \leq 2. \quad (10.59)$$

Lemma 10.17 (flat interval partial sum). *Let $g \in \mathcal{G}$. Assume $x \in [0, 2\pi) \setminus E_1$ and $g(x) = 0$. Then for all $N > \frac{2^{25}K^2}{\epsilon^3}$*

$$|S_N g(x)| \leq \frac{\epsilon}{4}, \quad (10.60)$$

Proof. With Lemma 10.10, breaking up the domain of integration into a partition of subintervals,

$$\begin{aligned} |S_N(g)| &= \frac{1}{2\pi} \left| \int_{-\pi}^{\pi} g(y) K_N(x-y) dy \right| \\ &= \frac{1}{2\pi} \left| \sum_{k=0}^{K-1} \int_{\frac{2\pi k}{K}}^{\frac{2\pi(k+1)}{K}} g(y) K_N(x-y) dy \right|. \end{aligned} \quad (10.61)$$

Using $g \in \mathcal{G}$ and that $k(y) = k$ for each

$$y \in \left[\frac{2\pi k}{K}, \frac{2\pi(k+1)}{K} \right), \quad (10.62)$$

and then applying the triangle inequality with the upper bound on $|g|$ and the identity $g(x) = g(2\pi k(x)/K) = 0$, we estimate (10.61) by

$$\begin{aligned} &= \frac{1}{2\pi} \left| \sum_{k=0}^{K-1} g\left(\frac{2\pi k}{K}\right) \int_{\frac{2\pi k}{K}}^{\frac{2\pi(k+1)}{K}} K_N(x-y) dy \right| \\ &\leq \frac{1}{\pi} \sum_{0 \leq k < K, k \neq k(x)} \left| \int_{\frac{2\pi k}{K}}^{\frac{2\pi(k+1)}{K}} K_N(x-y) dy \right|. \end{aligned} \quad (10.63)$$

Doing a variable substitution, we obtain for (10.63)

$$\frac{1}{\pi} \sum_{0 \leq k < K, k \neq k(x)} \left| \int_{x-2\pi(k+1)/K}^{x-2\pi k/K} K_N(y) dy \right|. \quad (10.64)$$

Fix $0 \leq k < K$ with $k \neq k(x)$ and set

$$\alpha = x - 2\pi(k+1)/K, \quad (10.65)$$

$$\beta = x - 2\pi k/K. \quad (10.66)$$

As $x \in [0, 2\pi) \setminus E_1$, we have with $\eta = \epsilon/(8K)$

$$\alpha \geq x - 2\pi \geq -2\pi + \eta, \quad (10.67)$$

$$\beta \leq x \leq 2\pi - \eta. \quad (10.68)$$

If $\beta < 0$, then $2\pi k/K > x$ and as $x \notin E_1$ also $2\pi k - \eta \geq x$. Hence $\beta \leq -\eta$. If $\beta \geq 0$, then $2\pi k/K \leq x$. As $k \neq k(x)$, we also have $2\pi(k+1)/K \leq x$. As $x \notin E_1$, we have $2\pi(k+1)/K + \eta \leq x$. It follows that $0 < \alpha$. In both cases, we have seen (10.52).

With Lemma 10.10 and the triangle inequality, it follows that

$$\left| \int_{x-2\pi(k+1)/K}^{x-2\pi k/K} K_N(y) dy \right| \leq \left| \int_{\alpha}^{\beta} \frac{e^{iNy}}{1 - e^{-iy}} dy \right| + \left| \int_{\alpha}^{\beta} \frac{e^{-iNy}}{1 - e^{iy}} dy \right| \quad (10.69)$$

Using that the two integrals on the right-hand side are complex conjugates of each other, and using Lemma 10.16 and $\epsilon < 2\pi$, this is bounded by

$$\leq 2 \left| \int_{\alpha}^{\beta} \frac{e^{iNy}}{1 - e^{-iy}} dy \right| \leq \frac{128}{N} \left(\frac{|\beta - \alpha|}{\eta^2} + \frac{1}{\eta} \right) \leq \frac{2^{20}K}{N\epsilon^2} \leq 2^{-3}\epsilon K^{-1}. \quad (10.70)$$

Inserting this in (10.63) and adding up proves the lemma. \square

We now prove Lemma 10.2.

Let $x \in [0, 2\pi) \setminus E_1$ and $N > \frac{2^{25}K^2}{\epsilon^3}$. Let h be the function which is constant equal to $f_0(x)$ and let $g = f_0 - h$. By the bound on f and the triangle inequality, $|g|$ is bounded by 2. Hence g is in the class \mathcal{G} . We also have $g(x) = 0$. Using Lemma 10.15, we obtain

$$S_N f_0(x) - f_0(x) = S_N(g + h)(x) - S_N h(x) = S_N g(x). \quad (10.71)$$

Lemma 10.2 now follows by Lemma 10.17.

10.3. Proof of Cotlar's Inequality.

Lemma 10.18 (estimate x shift). *Let $0 < r < 1$ and $x \in \mathbb{R}$. Let g be a bounded measurable function with bounded support on \mathbb{R} . Let Mg be the Hardy–Littlewood function defined in Proposition 2.6. Then for all x' with $|x - x'| < r$.*

$$\left| \int_{r < |x-y| < 1} g(y) \kappa(x-y) dy - \int_{r < |x'-y| < 1} g(y) \kappa(x'-y) dy \right| \leq 2^{13} Mg(x). \quad (10.72)$$

Proof. First note that the conditions $|x - y| < 1$ and $|x' - y| < 1$ may be removed as the factor containing κ vanishes without these conditions.

We split the first integral in (10.72) into the domains $r < |x - y| \leq 2r$ and $2r < |x - y|$. The integral over the first domain we estimate by (10.73) below. For the second domain, we observe with $|x - x'| < r$ and the triangle inequality that $r < |x' - y|$. We therefore combine on this domain with the corresponding part of the second integral in (10.72) and estimate that by (10.74) below. The remaining part of the second integral in (10.72) we estimate by (10.75). Overall, we have estimated (10.72) by

$$\int_{r < |x-y| \leq 2r} |g(y)| |\kappa(x-y)| dy \quad (10.73)$$

$$+ \left| \int_{2r < |x-y|} g(y) (\kappa(x'-y) - \kappa(x-y)) dy \right| \quad (10.74)$$

$$+ \left| \int_{r < |x'-y|, r < |x-y| < 2r} |g(y) \kappa(x'-y)| dy \right| \quad (10.75)$$

Using the bound on κ in Lemma 10.13, we estimate (10.73) by

$$\frac{8}{r} \int_{|x-y| < 2r} |g(y)| dy. \quad (10.76)$$

Using the definition of Mg , we estimate (10.76) by

$$\leq 32Mg(x). \quad (10.77)$$

Similarly, in the domain of (10.75) we note by the triangle inequality and assumption on x' that $|x' - y| \leq 3r$ and thus we estimate (10.75) by

$$\frac{8}{r} \int_{|x'-y| < 3r} |g(y)| dy \leq 48Mg(x) \quad (10.78)$$

We turn to the term (10.74). Let ν be the smallest integer such that $2^\nu > 2/r$. Using that the kernel vanishes unless $|x - y| < 2$, we decompose and estimate (10.74) with the triangle inequality by

$$\sum_{j=1}^{\nu} \int_{2^j r < |x-y| \leq 2^{j+1} r} |g(y)| |\kappa(x' - y) - \kappa(x - y)| dy. \quad (10.79)$$

Using Lemma 10.14, we estimate (10.79) by

$$\begin{aligned} & 2^{10} \sum_{j=1}^{\nu} \int_{2^j r < |x-y| \leq 2^{j+1} r} |g(y)| \frac{|x - x'|}{|x - y|^2} dy \\ & \leq 2^{10} \sum_{j=1}^{\nu} \frac{1}{2^{2j} r} \int_{2^j r < |x-y| \leq 2^{j+1} r} |g(y)| dy \\ & \leq 2^{10} \sum_{j=1}^{\nu} 2^{1-j} M g(x). \end{aligned} \quad (10.80)$$

Using a geometric series, we estimate (10.80) by

$$\leq 2^{11} M g(x). \quad (10.81)$$

Summing the estimates for (10.73), (10.74), and (10.75) proves the lemma. \square

Recall that

$$H_r g(x) = \int_{r < |x-y| < 1} g(y) \kappa(x - y) dy.$$

Lemma 10.19 (Cotlar control). *Let $0 < r < r_1 < 1$ and $x \in \mathbb{R}$. Let g be a bounded measurable function with bounded support on \mathbb{R} . Let Mg be the Hardy–Littlewood function defined in Proposition 2.6. Then for all $x' \in \mathbb{R}$ with $|x' - x| < \frac{r_1}{4}$ we have*

$$|H_{r_1} g(x)| \leq |H_r(g - g\mathbf{1}_{[x-\frac{r_1}{2}, x+\frac{r_1}{2}]}) (x')| + 2^{15} M g(x). \quad (10.82)$$

Proof. Let x and x' be given with $|x' - x| < \frac{r_1}{4}$. By an application of Lemma 10.18, we estimate the left-hand-side of (10.82) by

$$|H_{r_1}(g)(x')| + 2^{13} M g(x). \quad (10.83)$$

We have

$$H_{r_1}(g)(x') = \int_{r_1 < |x'-y| < 1} g(y) \kappa(x' - y) dy. \quad (10.84)$$

On the domain $r_1 < |x' - y|$, we have $\frac{r_1}{2} < |x - y|$. Hence we may write for (10.84)

$$\begin{aligned} H_{r_1}(g)(x') &= \int_{r_1 < |x'-y| < 1} (g - g\mathbf{1}_{[x-\frac{r_1}{2}, x+\frac{r_1}{2}]}) (y) \kappa(x' - y) dy \\ &= H_{r_1}(g - g\mathbf{1}_{[x-\frac{r_1}{2}, x+\frac{r_1}{2}]}) (x'). \end{aligned} \quad (10.85)$$

Combining the estimate (10.83) with the identification (10.85), we obtain

$$|H_{r_1} g(x)| \leq |H_{r_1}(g - g\mathbf{1}_{[x-\frac{r_1}{2}, x+\frac{r_1}{2}]}) (x')| + 2^{13} M g(x). \quad (10.86)$$

We have

$$\begin{aligned}
& (H_r - H_{r_1})(g - g\mathbf{1}_{[x-\frac{r_1}{2}, x+\frac{r_1}{2}]})(x') \\
&= \int_{r < |x'-y| < r_1} (g - g\mathbf{1}_{[x-\frac{r_1}{2}, x+\frac{r_1}{2}]})(y)k(x'-y) dy. \tag{10.87}
\end{aligned}$$

Assume first $r \geq \frac{r_1}{4}$. Then we estimate (10.87) with Lemma 10.13 by

$$\begin{aligned}
& \int_{\frac{r_1}{4} < |x'-y| < r_1} |g(y)k(x'-y)| dy \\
& \leq \frac{32}{r_1} \int_{|x'-y| < r_1} |g(y)| dy \leq 64Mg(x'). \tag{10.88}
\end{aligned}$$

Assume now $r \leq \frac{r_1}{4}$. As $|x'-y| \leq \frac{r_1}{4}$ implies $|x-y| \leq \frac{r_1}{2}$, we see that (10.87) equals

$$\int_{\frac{r_1}{4} < |x'-y| < r_1} (g - g\mathbf{1}_{[x-\frac{r_1}{2}, x+\frac{r_1}{2}]})(y)k(x'-y) dy,$$

which we again estimate as above by (10.88). In both cases, (10.82) follows by the triangle inequality from (10.86) and the estimate for (10.87). \square

Lemma 10.20 (Cotlar sets). *Let $0 < r < r_1 < 1$ and $x \in \mathbb{R}$. Let g be a bounded measurable function with bounded support on \mathbb{R} . Let Mg be the Hardy–Littlewood maximal function defined in Proposition 2.6. Let $x \in \mathbb{R}$. Then the measure $|F_1|$ of the set F_1 of all $x' \in [x - \frac{r_1}{4}, x + \frac{r_1}{4}]$ such that*

$$|H_r(g)(x')| > 4M(H_rg)(x) \tag{10.89}$$

is less than or equal to $r_1/8$. Moreover, the measure $|F_2|$ of the set F_2 of all $x' \in [x - \frac{r_1}{4}, x + \frac{r_1}{4}]$ such that

$$|H_r(g\mathbf{1}_{[x-\frac{r_1}{2}, x+\frac{r_1}{2}]})| > 2^{22}Mg(x) \tag{10.90}$$

is less than $r_1/8$.

Proof. Let r, r_1, x and g be given. If $M(H_rg)(x) = 0$, then H_rg is constant zero and F_1 is empty and the estimate on F_1 trivial. Assume $M(H_rg)(x) > 0$. We have with (10.89)

$$M(H_rg)(x) \geq \frac{2}{r_1} \int_{|x'-x| < \frac{r_1}{4}} |H_rg(x')| dx' \tag{10.91}$$

$$\geq \frac{2}{r_1} \int_{F_1} 4M(H_rg)(x) dx' \tag{10.92}$$

Dividing by $M(H_rg)(x)$ gives

$$1 \geq \frac{8}{r_1} |F_1|. \tag{10.93}$$

This gives the desired bound for the measure of F_1 .

We turn to the set F_2 . Similarly as above we may assume $Mg(x) > 0$. The set F_2 is then estimated with Lemma 10.7 by

$$\frac{2^{19}}{2^{22}Mg(x)} \int |g\mathbf{1}_{[x-\frac{r_1}{2}, x+\frac{r_1}{2}]}|(y) dy \tag{10.94}$$

$$\leq \frac{2^{19}}{2^{22}Mg(x)} r_1 Mg(x) = \frac{r_1}{8}. \tag{10.95}$$

This gives the desired bound for the measure of F_2 . \square

Lemma 10.21 (Cotlar estimate). *Let $0 < r < r_1 < 1$ and $x \in \mathbb{R}$. Let g be a bounded measurable function with bounded support on \mathbb{R} . Let Mg and $M(H_r g)$ be the respective Hardy–Littlewood maximal functions defined in Proposition 2.6. Then for all $x \in \mathbb{R}$*

$$\left| \int_{r_1 < |x-y| < 1} g(y) \kappa(x-y) dy \right| \quad (10.96)$$

$$\leq 2^2 M(H_r g)(x) + 2^{23} Mg(x). \quad (10.97)$$

Proof. By Lemma 10.19, the measure of the set of all $x' \in [x - \frac{r_1}{4}, x + \frac{r_1}{4}]$ such that at least one of the conditions (10.89) and (10.90) is satisfied is at most $r_1/4$ and hence not all of $x' \in [x - \frac{r_1}{4}, x + \frac{r_1}{4}]$. Pick an $x' \in [x - \frac{r_1}{4}, x + \frac{r_1}{4}]$ such that both conditions are not satisfied. Applying Lemma 10.19 for this x' and using the triangle inequality estimates the left-hand side of (10.96) by

$$4M(H_r g)(x) + 2^{22} Mg(x) + 2^{15} Mg(x). \quad (10.98)$$

This proves the lemma. \square

Lemma 10.22 (simple nontangential Hilbert). *For every $0 < r < 1$ and every bounded measurable function g with bounded support we have*

$$\|T_r g\|_2 \leq 2^{42} \|g\|_2, \quad (10.99)$$

where

$$T_r g(x) := \sup_{r < r_1 < 1} \sup_{|x-x'| < r_1} \frac{1}{2\pi} \left| \int_{r_1 < |x'-y| < 1} g(y) \kappa(x'-y) dy \right|. \quad (10.100)$$

Proof. With Lemma 10.18 and the triangle inequality, we estimate for every $x \in \mathbb{R}$

$$T_r g(x) \leq 2^{13} Mg(x) + \sup_{r < r_1 < 1} \frac{1}{2\pi} \left| \int_{r_1 < |x-y| < 1} g(y) \kappa(x-y) dy \right|. \quad (10.101)$$

Using further Lemma 10.21, we estimate

$$T_r g(x) \leq 2^{13} Mg(x) + 2^{23} Mg(x) + 2^2 M(H_r g)(x). \quad (10.102)$$

Taking the L^2 norm and using Proposition 2.6 with $a = 4$ and $p_2 = 2$ and $p_1 = 1$, we obtain

$$\|T_r g\|_2 \leq 2^{24} \|Mg\|_2 + 2^2 \|M(H_r g)\|_2 \quad (10.103)$$

$$\leq 2^{41} \|g\|_2 + 2^{19} \|H_r(g)\|_2. \quad (10.104)$$

Applying Lemma 10.6, gives

$$\|T_r g\|_2 \leq 2^{41} \|g\|_2 + 2^{32} \|g\|_2. \quad (10.105)$$

This shows (10.99) and completes the proof of the lemma. \square

Proof of Lemma 10.5. Fix g as in the Lemma. Applying Lemma 10.22 with a sequence of r tending to 0 and using Lebesgue monotone convergence shows

$$\|T_0 g\|_2 \leq 2^{42} \|g\|_2, \quad (10.106)$$

where

$$T_0 g(x) := \sup_{0 < r_1 < 1} \sup_{|x-x'| < r_1} \frac{1}{2\pi} \left| \int_{r_1 < |x'-y| < 1} g(y) \kappa(x' - y) dy \right|. \quad (10.107)$$

We now write by the triangle inequality

$$\begin{aligned} T_* g(x) &\leq \sup_{0 < r_1 < r_2 < 1} \sup_{|x-x'| < r_1} \frac{1}{2\pi} \left| \int_{r_1 < |x'-y| < 1} g(y) \kappa(x' - y) dy \right| \\ &\quad + \sup_{0 < r_1 < r_2 < 1} \sup_{|x-x'| < r_1} \frac{1}{2\pi} \left| \int_{r_2 < |x'-y| < 1} g(y) \kappa(x' - y) dy \right|. \end{aligned} \quad (10.108)$$

Noting that the first integral does not depend on r_2 and estimating the second integral by the larger supremum over all $|x - x'| < r_2$, at which time the integral does not depend on r_1 , we estimate (10.108) by

$$\begin{aligned} &\sup_{0 < r_1 < 1} \sup_{|x-x'| < r_1} \frac{1}{2\pi} \left| \int_{r_1 < |x'-y| < 1} g(y) \kappa(x' - y) dy \right| \\ &\quad + \sup_{0 < r_2 < 1} \sup_{|x-x'| < r_2} \frac{1}{2\pi} \left| \int_{r_2 < |x'-y| < 1} g(y) \kappa(x' - y) dy \right|. \end{aligned} \quad (10.109)$$

Applying the triangle inequality on the left-hand side of (10.28) and applying (10.106) twice proves (10.28). This completes the proof of Lemma 10.5. \square

10.4. The truncated Hilbert transform. Let M_n be the modulation operator acting on measurable 2π -periodic functions defined by

$$M_n g(x) = g(x) e^{inx}. \quad (10.110)$$

Define the approximate Hilbert transform by

$$L_N g = \frac{1}{N} \sum_{n=0}^{N-1} M_{-n-N} S_{N+n} M_{N+n} g. \quad (10.111)$$

Lemma 10.23 (modulated averaged projection). *We have for every bounded measurable 2π -periodic function g*

$$\|L_N g\|_{L^2[-\pi, \pi]} \leq \|g\|_{L^2[-\pi, \pi]}. \quad (10.112)$$

Proof. We have

$$\|M_n g\|_{L^2[-\pi, \pi]}^2 = \int_{-\pi}^{\pi} |e^{inx} g(x)|^2 dx = \int_{-\pi}^{\pi} |g(x)|^2 dx = \|g\|_{L^2[-\pi, \pi]}^2. \quad (10.113)$$

We have by the triangle inequality, the square root of the identity in (10.113), and Lemma 10.12

$$\begin{aligned} \|L_N g\|_{L^2[-\pi, \pi]} &= \left\| \frac{1}{N} \sum_{n=0}^{N-1} M_{-n-N} S_{N+n} M_{N+n} g \right\|_{L^2[-\pi, \pi]} \\ &\leq \frac{1}{N} \sum_{n=0}^{N-1} \|M_{-n-N} S_{N+n} M_{N+n} g\|_{L^2[-\pi, \pi]} = \frac{1}{N} \sum_{n=0}^{N-1} \|S_{N+n} M_{N+n} g\|_{L^2[-\pi, \pi]} \end{aligned}$$

$$\leq \frac{1}{N} \sum_{n=0}^{N-1} \|M_{N+n}g\|_{L^2[-\pi,\pi]} = \frac{1}{N} \sum_{n=0}^{N-1} \|g\|_{L^2[-\pi,\pi]} = \|g\|_{L^2[-\pi,\pi]}. \quad (10.114)$$

This proves (10.113) and completes the proof of the lemma. \square

Lemma 10.24 (periodic domain shift). *Let f be a bounded 2π -periodic function. We have for any $0 \leq x \leq 2\pi$ that*

$$\int_0^{2\pi} f(y) dy = \int_{-x}^{2\pi-x} f(y) dy = \int_{-\pi}^{\pi} f(y-x) dy. \quad (10.115)$$

Proof. We have by periodicity and change of variables

$$\int_{-x}^0 f(y) dy = \int_{-x}^0 f(y+2\pi) dy = \int_{2\pi-x}^{2\pi} f(y) dy. \quad (10.116)$$

We then have by breaking up the domain of integration and using (10.116)

$$\begin{aligned} \int_0^{2\pi} f(y) dy &= \int_0^{2\pi-x} f(y) dy + \int_{2\pi-x}^{2\pi} f(y) dy \\ &= \int_0^{2\pi-x} f(y) dy + \int_{-x}^0 f(y) dy = \int_{-x}^{2\pi-x} f(y) dy. \end{aligned} \quad (10.117)$$

This proves the first identity of the lemma. The second identity follows by substitution of y by $y-x$. \square

Lemma 10.25 (Young convolution). *Let f and g be two bounded non-negative measurable 2π -periodic functions on \mathbb{R} . Then*

$$\left(\int_{-\pi}^{\pi} \left(\int_{-\pi}^{\pi} f(y)g(x-y) dy \right)^2 dx \right)^{\frac{1}{2}} \leq \|f\|_{L^2[-\pi,\pi]} \|g\|_{L^1[-\pi,\pi]}. \quad (10.118)$$

Proof. Using Fubini and Lemma 10.24, we observe

$$\begin{aligned} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} f(y)^2 g(x-y) dy dx &= \int_{-\pi}^{\pi} f(y)^2 \int_{-\pi}^{\pi} g(x-y) dx dy \\ &= \int_{-\pi}^{\pi} f(y)^2 \int_{-\pi}^{\pi} g(x) dx dy = \|f\|_{L^2[-\pi,\pi]}^2 \|g\|_{L^1[-\pi,\pi]}. \end{aligned} \quad (10.119)$$

Let h be the nonnegative square root of g , then h is bounded and 2π -periodic with $h^2 = g$. We estimate the square of the left-hand side of (10.118) with Cauchy-Schwarz and then with (10.119) by

$$\begin{aligned} &\int_{-\pi}^{\pi} \left(\int_{-\pi}^{\pi} f(y)h(x-y)h(x-y) dy \right)^2 dx \\ &\leq \int_{-\pi}^{\pi} \left(\int_{-\pi}^{\pi} f(y)^2 g(x-y) dy \right) \left(\int_{-\pi}^{\pi} g(x-y) dy \right) dx \\ &= \|f\|_{L^2[-\pi,\pi]}^2 \|g\|_{L^1[-\pi,\pi]}. \end{aligned}$$

Taking square roots, this proves the lemma. \square

For $0 < r < 1$, Define the kernel k_r to be the 2π -periodic function

$$|k_r(x)| := \min \left(r^{-1}, 1 + \frac{r}{|1 - e^{ix}|^2} \right), \quad (10.120)$$

where the minimum is understood to be r^{-1} in case $1 = e^{ix}$.

Lemma 10.26 (integrable bump convolution). *Let g, f be bounded measurable 2π -periodic functions. Let $0 < r < \pi$. Assume we have for all $0 \leq x \leq 2\pi$*

$$|g(x)| \leq k_r(x). \quad (10.121)$$

Let

$$h(x) = \int_{-\pi}^{\pi} f(y)g(x-y) dy. \quad (10.122)$$

Then

$$\|h\|_{L^2[-\pi, \pi]} \leq 2^5 \|f\|_{L^2[-\pi, \pi]}. \quad (10.123)$$

Proof. From monotonicity of the integral and (10.121),

$$\|g\|_{L^1[-\pi, \pi]} \leq \int_{-\pi}^{\pi} k_r(x) dx. \quad (10.124)$$

Using the symmetry $k_r(x) = k_r(-x)$, the assumption, and Lemma 10.11, the last display is equal to

$$\begin{aligned} &= 2 \int_0^{\pi} \min\left(\frac{1}{r}, 1 + \frac{r}{|1 - e^{ix}|^2}\right) dx \\ &\leq 2 \int_0^r \frac{1}{r} dx + 2 \int_r^{\pi} 1 + \frac{64r}{x^2} dx \\ &\leq 2 + 2\pi + 2 \left(\frac{64r}{r} - \frac{64r}{\pi} \right) \leq 2^5. \end{aligned} \quad (10.125)$$

Together with Lemma 10.25, this proves the lemma. \square

Lemma 10.27 (dirichlet approximation). *Let $0 < r < 1$. Let N be the smallest integer larger than $\frac{1}{r}$. There is a 2π -periodic continuous function L' on \mathbb{R} that satisfies for all $-\pi \leq x \leq \pi$ and all 2π -periodic bounded measurable functions f on \mathbb{R}*

$$L_N f(x) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(y) L'(x-y) dy \quad (10.126)$$

and

$$|L'(x) - \mathbf{1}_{\{y: r < |y| < 1\}} \kappa(x)| \leq 2^5 k_r(x). \quad (10.127)$$

Proof. We have by definition and Lemma 10.10

$$L_N g(x) = \frac{1}{N} \sum_{n=0}^{N-1} \int_{-\pi}^{\pi} e^{-i(N+n)x} K_{N+n}(x-y) e^{i(N+n)y} g(y) dy. \quad (10.128)$$

This is of the form (10.126) with the continuous function

$$L'(x) = \frac{1}{N} \sum_{n=0}^{N-1} K_{N+n}(x) e^{-i(N+n)x}. \quad (10.129)$$

With (10.37) of Lemma 10.10 we have $|K_N(x)| \leq N$ for every x and thus

$$|L'(x)| \leq \frac{1}{N} \sum_{n=0}^{N-1} (N+n) \leq 2N \leq 2^2 r^{-1}. \quad (10.130)$$

Therefore, for $|x| \in [0, r) \cup (1, \pi]$, we have

$$|L'(x) - \mathbf{1}_{\{y: r < |y| < 1\}}(x) \kappa(x)| = |L'(x)| \leq 2^2 r^{-1}. \quad (10.131)$$

This proves (10.127) for $|x| \in [0, r)$ since $k_r(x) = r^{-1}$ in this case.

For $e^{ix'} \neq 1$ and may use the expression (10.38) for K_N in Lemma 10.10 to obtain

$$\begin{aligned} L'(x) &= \frac{1}{N} \sum_{n=0}^{N-1} \left(\frac{e^{i(N+n)x}}{1 - e^{-ix}} + \frac{e^{-i(N+n)x}}{1 - e^{ix}} \right) e^{-i(N+n)x} \\ &= \frac{1}{N} \sum_{n=0}^{N-1} \left(\frac{1}{1 - e^{-ix}} + \frac{e^{-i2(N+n)x}}{1 - e^{ix}} \right) \\ &= \frac{1}{1 - e^{-ix}} + \frac{1}{N} \frac{e^{-i2Nx}}{1 - e^{ix}} \sum_{n=0}^{N-1} e^{-i2nx} \end{aligned} \quad (10.132)$$

and thus

$$L'(x) - \mathbf{1}_{\{y: r < |y| < 1\}} \kappa(x) = L''(x) + \frac{1 - \mathbf{1}_{\{y: r < |y| < 1\}}(x)(1 - |x|)}{1 - e^{-ix}}, \quad (10.133)$$

where

$$L''(x) := \frac{1}{N} \frac{e^{-i2Nx}}{1 - e^{ix}} \sum_{n=0}^{N-1} e^{-i2nx}.$$

For $x \in [-\pi, r] \cup [r, \pi]$, we have using Lemma 10.11 that

$$\begin{aligned} \left| \frac{1 - \mathbf{1}_{\{y: r < |y| < 1\}}(x)(1 - |x|)}{1 - e^{-ix}} \right| &= \left| \frac{\min(|x|, 1)}{1 - e^{-ix}} \right| \leq \frac{8 \min(|x|, 1)}{|x|} \\ &\leq 2^3 \cdot 1 \leq 2^3 k_r(x). \end{aligned} \quad (10.134)$$

Next, we need to estimate $L''(x)$. If the real part of e^{ix} is negative, we have

$$1 \leq |1 - e^{ix}| \leq 2. \quad (10.135)$$

and hence

$$|L''(x)| \leq \frac{1}{N} \sum_{n=0}^{N-1} 1 = 1 \leq 1 + \frac{r}{|1 - e^{ix}|^2}. \quad (10.136)$$

If the real part of e^{ix} is positive and in particular while still $e^{ix} \neq \pm 1$, then we have by telescoping

$$(1 - e^{-2ix}) \sum_{n=0}^{N-1} e^{-i2nx} = 1 - e^{-i2Nx}. \quad (10.137)$$

As $e^{-2ix} \neq 1$, we may divide by $1 - e^{-2ix}$ and insert this into (10.132) to obtain

$$L''(x) = \frac{1}{N} \frac{e^{-i2Nx}}{1 - e^{ix}} \frac{1 - e^{-i2Nx}}{1 - e^{-2ix}}. \quad (10.138)$$

Hence, with Lemma 10.11 and nonnegativity of the real part of e^{ix}

$$\begin{aligned} |L''(x)| &\leq \frac{2}{N} \frac{1}{|1 - e^{ix}|} \frac{1}{|1 - e^{-2ix}|} \\ &= \frac{2}{N} \frac{1}{|1 - e^{ix}|^2} \frac{1}{|1 + e^{ix}|} \leq \frac{4r}{|1 - e^{ix}|^2} \leq 2^2 \left(1 + \frac{r}{|1 - e^{ix}|^2} \right) \end{aligned} \quad (10.139)$$

Inequalities (10.130), (10.131), (10.133), (10.134), (10.136), and (10.139) prove (10.127). This completes the proof of the lemma. \square

We now prove Lemma 10.6.

Proof of Lemma 10.6. We first show that if f is supported in $[-3/2, 3/2]$, then

$$\|H_r f\|_{L^2(\mathbb{R})} \leq 2^{16} \|f\|_{L^2(\mathbb{R})}. \quad (10.140)$$

Let \tilde{f} be the 2π -periodic extension of f to \mathbb{R} . Let N be the smallest integer larger than $\frac{1}{r}$. Then, by Lemma 10.27 and the triangle inequality, for $x \in [-\pi, \pi]$ we have

$$|H_r \tilde{f}(x)| \leq 2\pi |L_N \tilde{f}(x)| + 2^5 \left| \int_{-\pi}^{\pi} k_r(x-y) \tilde{f}(y) dy \right|.$$

Taking L^2 norm over the interval $[-\pi, \pi]$ and using its sub-additivity, we get

$$\begin{aligned} & \|H_r \tilde{f}\|_{L^2([-\pi, \pi])} \\ & \leq 2\pi \|L_N \tilde{f}\|_{L^2([-\pi, \pi])} + 2^5 \left(\int_{-\pi}^{\pi} \left| \int_{-\pi}^{\pi} k_r(x-y) \tilde{f}(y) dy \right|^2 dx \right)^{\frac{1}{2}}. \end{aligned}$$

Since k_r is supported in $[-1, 1]$, we have that $H_r f$ is supported in $[-5/2, 5/2]$ and agrees there with $H_r \tilde{f}(x)$. Using Lemma 10.23 and Lemma 10.26, we conclude

$$\|H_r f\|_{L^2(\mathbb{R})} \leq \|H_r \tilde{f}\|_{L^2([-\pi, \pi])} \leq 2\pi \|f\|_{L^2(\mathbb{R})} + 2^{10} \|f\|_{L^2(\mathbb{R})}, \quad (10.141)$$

which gives (10.140).

Suppose now that f is supported in $[c, c+3]$ for some $c \in \mathbb{R}$. Then the function $g(x) = f(c + \frac{3}{2} + x)$ is supported in $[-3/2, 3/2]$. By a change of variables in (10.30), we have $H_r g(x) = H_r f(c + 3/2 + x)$. Thus, by (10.140)

$$\|H_r g\|_2 = \|H_r f\|_2 \leq 2^{11} \|f\|_2 = \|g\|_2. \quad (10.142)$$

Let now f be arbitrary. Since $\kappa(x) = 0$ for $|x| > 1$, we have for all $x \in [c+1, c+2]$

$$H_r f(x) = H_r(f \mathbf{1}_{[c, c+3]})(x).$$

Thus

$$\int_{c+1}^{c+2} |H_r f(x)|^2 dx \leq \int_{\mathbb{R}} |H_r(f \mathbf{1}_{[c, c+3]})(x)|^2 dx.$$

Applying the bound (10.142), this is

$$\leq 2^{11} \int_c^{c+3} |f(x)|^2 dx.$$

Summing over all $c \in \mathbb{Z}$, we obtain

$$\int_{\mathbb{R}} |H_r f(x)|^2 dx \leq 3 \cdot 2^{11} \int_{\mathbb{R}} |f(x)|^2 dx.$$

This completes the proof. \square

10.5. Calderón-Zygmund Decomposition. For $I = [s, t) \subset \mathbb{R}$, we define

$$\text{bi}(I) := \left\{ \left[s, \frac{s+t}{2} \right), \left[\frac{s+t}{2}, t \right) \right\}. \quad (10.143)$$

In what follows, we write μ for the Lebesgue measure on \mathbb{R} .

Lemma 10.28 (interval bisection). *Let $I = [s, t) \subset \mathbb{R}$ be a bounded, right-open interval. Let $I_1, I_2 \in \text{bi}(I)$ with $I_1 \neq I_2$. Then*

$$\mu(I_1) = \mu(I_2) = \frac{\mu(I)}{2}. \quad (10.144)$$

Further,

$$I = I_1 \cup I_2, \quad (10.145)$$

and the intervals I_1 and I_2 are disjoint.

Proof. This in some form can be taken from the Lean library. \square

For a bounded interval $I = [a, b) \subset \mathbb{R}$ and a non-negative integer n , we inductively define

$$\text{ch}_0(I) := \{I\}, \quad \text{ch}_n(I) = \cup_{J \in \text{ch}_{n-1}(I)} \text{bi}(J). \quad (10.146)$$

Lemma 10.29 (bisection children). *Let $I = [s, t)$ and let n be a non-negative integer. Then the intervals in $\text{ch}_n(I)$ are pairwise disjoint. For any $J \in \text{ch}_n(I)$,*

$$\mu(J) = \frac{\mu(I)}{2^n} = \frac{t-s}{2^n}. \quad (10.147)$$

Further,

$$I = \cup_{J \in \text{ch}_n(I)} J \quad (10.148)$$

and

$$|\text{ch}_n(I)| = 2^{n+1}. \quad (10.149)$$

Proof. This follows by induction on n , using Lemma 10.28. \square

Lemma 10.30 (Lebesgue differentiation). *Let f be a bounded measurable function with bounded support. Then for μ almost every x , we have*

$$\frac{1}{\mu(I_n)} \int_{I_n} f(y) dy = f(x),$$

where $\{I_n\}_{n \geq 1}$ is a sequence of intervals such that $x \in I_n$ for each $n \geq 1$ and

$$\lim_{n \rightarrow \infty} \mu(I_n) = 0.$$

Proof. This follows from the Lebesgue differentiation theorem, which is already formalized in Lean. \square

Lemma 10.31 (stopping time). *Let f be a bounded, measurable function with bounded support on \mathbb{R} . Let $\alpha > 0$. Then there exists $A \subset \mathbb{R}$ such that the following properties (10.150), (10.151), (10.152), (10.153), and (10.154) are satisfied. For all $x \in A$*

$$|f(x)| \leq \alpha. \quad (10.150)$$

The set $\mathbb{R} \setminus A$ can be decomposed into a countable union of intervals

$$\mathbb{R} \setminus A = \cup_j I_j = \cup_j [s_j, t_j), \quad (10.151)$$

such that

$$[s_j, t_j) \cap [s_{j'}, t_{j'}) = \emptyset \text{ for } j \neq j'. \quad (10.152)$$

For each j ,

$$\frac{\alpha}{2} \leq \frac{1}{t_j - s_j} \int_{s_j}^{t_j} |f(y)| dy \leq \alpha. \quad (10.153)$$

Further,

$$\sum_j (t_j - s_j) \leq \frac{2}{\alpha} \int |f(y)| dy. \quad (10.154)$$

Proof. Since f is bounded with bounded support, there exists a non-negative integer ℓ such that

$$f(x) = 0 \text{ for } x \notin [-2^{\ell-1}, 2^{\ell-1}), \quad (10.155)$$

and

$$2^{-\ell} \int_{-2^{\ell-1}}^{2^{\ell-1}} |f(y)| dy < \alpha.$$

Let

$$I_0 := [-2^\ell, 2^\ell), \quad \mathcal{Q}_0 := \{I_0\},$$

and

$$\tilde{\mathcal{Q}}_1 := \text{bi}(I_0).$$

For $n \geq 1$, we inductively define

$$\mathcal{Q}_n := \left\{ I \in \tilde{\mathcal{Q}}_n : \frac{1}{\mu(I)} \int_I |f(y)| dy > \frac{\alpha}{2} \right\},$$

and

$$\tilde{\mathcal{Q}}_{n+1} := \cup_{I \in \tilde{\mathcal{Q}}_n \setminus \mathcal{Q}_n} \text{bi}(I).$$

Finally, let

$$\mathcal{Q} := \cup_{n \geq 1} \mathcal{Q}_n.$$

For each n we have

$$\mathcal{Q}_n \subseteq \text{ch}_n(I_0).$$

Therefore, by Lemma 10.29,

$$|\mathcal{Q}_n| \leq \frac{\mu(I_0)}{2^n} = 2^{\ell+1-n}.$$

It follows that \mathcal{Q} is a countable union of finite sets and hence, is countable itself. Let $\{I_j\}_{j \geq 1}$ be an enumeration of this set, with

$$I_j := [s_j, t_j),$$

for $s_j, t_j \in I_0$ and $s_j < t_j$. We set

$$A := \mathbb{R} \setminus \cup_{j \geq 1} [s_j, t_j).$$

Then (10.151) holds by definition. We next show (10.152). Let $I_j, I_{j'} \in \mathcal{Q}$. Then, there exist $1 \leq n, n'$ such that $I_j \in \mathcal{Q}_n$ and $I_{j'} \in \mathcal{Q}_{n'}$. If $n = n'$, (10.152) follows from Lemma 10.29 since $\mathcal{Q}_n \subseteq \text{ch}_n(I_0)$. Otherwise, assume without loss of generality that $n < n'$. Then by construction, there exists $J \in \tilde{\mathcal{Q}}_n \setminus \mathcal{Q}_n$ such that $I_{j'} \subset J$. Since $I_j \in \mathcal{Q}_n$, it follows that $I_j \neq J$. Since $I_j, J \in \tilde{\mathcal{Q}}_n \subset \text{ch}_n(I_0)$, by Lemma 10.29, we deduce that they are disjoint. Since $I_{j'} \subset J$, we conclude that I_j and $I_{j'}$ are disjoint as well. This proves (10.152).

To see (10.153), let $I_j = (s_j, t_j) \in \mathcal{Q}$. Then $I_j \in \mathcal{Q}_n$ for some $n \geq 1$. By definition of \mathcal{Q}_n , we have

$$\frac{\alpha}{2} < \frac{1}{\mu(I)} \int_I |f(y)| dy = \frac{1}{t_j - s_j} \int_{s_j}^{t_j} |f(y)| dy.$$

As $I_j \in \mathcal{Q}_n \subseteq \tilde{\mathcal{Q}}_n$, by definition of the latter set, there exists $J \in \tilde{\mathcal{Q}}_{n-1} \setminus \mathcal{Q}_{n-1}$ with $I \in \text{bi}(J)$. Since $J \notin \mathcal{Q}_{n-1}$, we conclude that

$$\frac{1}{\mu(J)} \int_J |f(y)| dy \leq \frac{\alpha}{2}.$$

Using Lemma 10.28 and the above, we get

$$\frac{1}{t_j - s_j} \int_{s_j}^{t_j} |f(y)| dy = \frac{1}{\mu(I)} \int_I |f(y)| dy \leq \frac{2}{\mu(J)} \int_J |f(y)| dy \leq \alpha.$$

This establishes (10.153). Using this, we see that for each j , we have

$$t_j - s_j \leq \frac{2}{\alpha} \int_{s_j}^{t_j} |f(y)| dy.$$

Summing up in j and using the disjointness property (10.152), we get

$$\sum_j (t_j - s_j) \leq \frac{2}{\alpha} \int_{s_j}^{t_j} |f(y)| dy = \frac{2}{\alpha} \int |f(y)| dy.$$

Finally, we show (10.150). Let $x \in A$. If $x \notin [-2^\ell, 2^\ell]$, then by (10.155) that $f(x) = 0$. Thus (10.150) is true in this case. Alternately, let $x \in I_0 \cap A$. Then for each n , there exists $I(n) \in \tilde{\mathcal{Q}}_n$ such that $x \in I(n)$ and

$$\frac{1}{\mu(I(n))} \int_{I(n)} |f(y)| dy \leq \alpha. \quad (10.156)$$

Since $\mu(I(n)) = 2^{\ell+1-n}$, we have

$$\lim_{n \rightarrow \infty} \mu(I(n)) = 0.$$

By Lemma 10.30, we also have

$$\lim_{n \rightarrow \infty} \frac{1}{\mu(I(n))} \int_{I(n)} |f(y)| dy = |f(x)|$$

for almost every $x \in A$. Combining the above with (10.156), we conclude that

$$|f(x)| \leq \alpha.$$

This finishes the proof of (10.150), and hence the lemma. \square

Calderón-Zygmund decomposition is a tool to extend L^2 bounds to L^p bounds with $p < 2$ or to the so-called weak $(1,1)$ type endpoint bound. It is classical and can be found in [Ste93].

Lemma 10.32 (Calderon Zygmund decomposition). *Let f be a bounded, measurable function with bounded support. Let $\alpha > 0$ and $\gamma \in (0, 1)$. Then there exists a measurable functions g , a countable family of disjoint intervals*

$I_j = [s_j, t_j)$, and a countable family of measurable functions $\{g_j\}_{j \geq 1}$ such that for almost every $x \in \mathbb{R}$

$$f(x) = g(x) + \sum_{j \geq 1} b_j(x) \quad (10.157)$$

and such that the following holds. For almost every $x \in \mathbb{R}$,

$$|g(x)| \leq \gamma\alpha. \quad (10.158)$$

We have

$$\int |g(y)| dy \leq \int |f(y)| dy. \quad (10.159)$$

For every j

$$\text{supp } b_j \subset I_j. \quad (10.160)$$

For every j

$$\int_{I_j} b_j(x) dx = 0, \quad (10.161)$$

and

$$\int_{I_j} |b_j(x)| dx \leq 2\gamma\alpha. \quad (10.162)$$

We have

$$\sum_j (t_j - s_j) \leq \frac{2}{\gamma\alpha} \int |f(y)| dy \quad (10.163)$$

and

$$\sum_j \int_{I_j} |b_j(y)| dy \leq 2 \int |f(y)| dy. \quad (10.164)$$

Proof. Applying Lemma 10.31 to f and $\gamma\alpha$, we obtain a collection I_j of intervals such that the conditions (10.150)-(10.154) are satisfied. We set $A = \mathbb{R} \setminus \cup_j I_j$ and

$$g(x) := \begin{cases} f(x), & x \in A, \\ \frac{1}{\mu(I_j)} \int_{I_j} f(y) dy, & x \in (s_j, t_j), \\ 0, & x \in [s_j, t_j] \setminus (s_j, t_j), \end{cases} \quad (10.165)$$

and, for each j ,

$$b_j(x) := \begin{cases} f(x) - \frac{1}{\mu(I_j)} \int_{I_j} f(y) dy, & x \in (s_j, t_j), \\ 0, & x \notin (s_j, t_j). \end{cases} \quad (10.166)$$

Then (10.160) and (10.163) are true by construction and Lemma 10.31. Further, let $b(x) = \sum_j b_j(x)$. Then

$$f(x) = g(x) + b(x) = g(x) + \sum_j b_j(x),$$

for all x not in the measure zero set $\cup_j \{s_j, t_j\}$.

For almost every $x \in A$, we get using (10.150)

$$|g(x)| \leq \gamma\alpha.$$

In the case when neither of the above is true, we have $g(x) = 0$ by definition. Thus, we obtain (10.158).

To prove (10.159), we estimate

$$\int |g(y)| dy \leq \int_A |f(y)| dy + \sum_j \int_{I_j} \frac{1}{\mu(I_j)} \int_{I_j} |f(y)| dy.$$

Since the intervals I_j are disjoint, and $A = \mathbb{R} \setminus \cup_j I_j$, we conclude

$$\int |g(y)| dy \leq \int_A |f(y)| dy + \sum_j \int_{I_j} |f(y)| dy = \int |f(y)| dy.$$

This establishes (10.159). If $x \in I_j$ for some j , (10.153) yields

$$|g(x)| \leq \frac{1}{\mu(I_j)} \int_{I_j} |f(y)| dy \leq \gamma \alpha.$$

Further, for each j , it follows from the definition of t_j that

$$\begin{aligned} \int_{I_j} b_j(x) dx &= \int_{I_j} f(x) dx - \int_{I_j} \frac{1}{\mu(I_j)} \int_{I_j} f(y) dy dx \\ &= \int_{I_j} f(x) dx - \int_{I_j} f(y) dy = 0. \end{aligned}$$

This establishes (10.161).

Using the triangle inequality, we have that

$$\begin{aligned} \int_{I_j} |b_j(y)| dy &\leq \int_{I_j} |f(y)| dy + \int_{I_j} \frac{1}{\mu(I_j)} \int_{I_j} |f(x)| dx dy \\ &= 2 \int_{I_j} |f(y)| dy. \end{aligned} \tag{10.167}$$

Dividing both sides by $\mu(I_j)$ and using (10.153), we obtain (10.162).

Further, summing up (10.167) in j yields

$$\sum_j \int_{I_j} |b_j(y)| dy \leq \sum_j \int_{I_j} |f(y)| dy + \sum_j \int_{I_j} \frac{1}{\mu(I_j)} \int_{I_j} |f(x)| dx dy.$$

Using the disjointedness of (s_j, t_j) , we get

$$\sum_j \int_{I_j} |b_j(y)| dy \leq 2 \int |f(y)| dy.$$

This proves (10.164), and completes the proof. \square

Proof of Lemma 10.7. Using Lemma 10.32 for f and $2^{-10}\alpha$, we obtain the decomposition

$$f = g + b = g + \sum_j b_j$$

such that the properties (10.157)-(10.164) are satisfied with $\gamma = 2^{-10}$. For each j , let

$$I_j = [s_j, t_j), \quad c_j := \frac{s_j + t_j}{2}, \tag{10.168}$$

and

$$I_j^* = [s_j - 2(t_j - s_j), t_j + 2(t_j - s_j)]. \tag{10.169}$$

Then I_j^* is an interval with the same center as I_j but with

$$\mu(I_j^*) = 5(t_j - s_j) = 5\mu(I_j). \tag{10.170}$$

Let

$$\Omega := \cup_j I_j^*. \quad (10.171)$$

By definition, for each $x \in \mathbb{R} \setminus \Omega$ and $y \in I_j$,

$$|x - c_j| > \frac{5(t_j - s_j)}{2} \geq 5|y - c_j|, \quad (10.172)$$

and

$$|x - y| > 2(t_j - s_j). \quad (10.173)$$

It follows by the triangle inequality and subadditivity of μ that

$$\begin{aligned} & \mu(\{x \in \mathbb{R} : |H_r f(x)| > \alpha\}) \\ & \leq \mu(\{x \in \mathbb{R} : |H_r g(x)| > \alpha/2\}) + \mu(\{x \in \mathbb{R} : |H_r b(x)| > \alpha/2\}). \end{aligned} \quad (10.174)$$

We estimate using monotonicity of the integral

$$\mu(\{x \in \mathbb{R} : |H_r g(x)| > \alpha/2\}) \leq \frac{4}{\alpha^2} \int |H_r g(y)|^2 dy.$$

Using Lemma 10.6 followed by (10.158) and (10.159), we estimate the right hand side above by

$$\leq 2^{26} \frac{4}{\alpha^2} \int |g(y)|^2 dy \leq \frac{2^{18}}{\alpha} \int |g(y)| dy \leq \frac{2^{18}}{\alpha} \int |f(y)| dy.$$

Thus, we conclude

$$\mu(\{x \in \mathbb{R} : |H_r g(x)| > \alpha/2\}) \leq \frac{2^{18}}{\alpha} \int |f(y)| dy. \quad (10.175)$$

Next, we estimate

$$\begin{aligned} & \mu(\{x \in \mathbb{R} : |H_r b(x)| > \alpha/2\}) \\ & \leq \mu(\Omega) + \mu(\{x \in \mathbb{R} \setminus \Omega : |H_r b(x)| > \alpha/2\}). \end{aligned}$$

Using (10.170) and (10.163), we conclude that

$$\mu(\Omega) \leq \sum_j \mu(I_j^*) = 5 \sum_j (t_j - s_j) \leq \frac{2^{13}}{\alpha} \int |f(y)| dy. \quad (10.176)$$

We now focus on estimating the remaining term

$$\mu(\{x \in \mathbb{R} \setminus \Omega : |H_r b(x)| > \alpha/2\}).$$

For $x \in \mathbb{R} \setminus \Omega$, define

$$\begin{aligned} \mathcal{J}_1(x) &:= \{j : [s_j, t_j] \cap [x - r, x + r] = \emptyset\}, \\ \mathcal{J}_2(x) &:= \{j : |y - x| = r \text{ for some } y \in [s_j, t_j]\}, \\ \mathcal{J}_3(x) &:= \{j : [s_j, t_j] \subset [x - r, x + r]\}. \end{aligned}$$

Since $H_r b_j(x) = 0$ for all $j \in \mathcal{J}_3(x)$, using the triangle inequality and the decomposition above, we get

$$|H_r b(x)| \leq \sum_{j \in \mathcal{J}_1(x)} |H_r b_j(x)| + \sum_{j \in \mathcal{J}_2(x)} |H_r b_j(x)|. \quad (10.177)$$

Further, for $j \in \mathcal{J}_1(x)$, we have

$$H_r b_j(x) = \int_{I_j} \kappa(x - y) \mathbf{1}_{\{z: r < |z| < 1\}}(x - y) b_j(y) dy = \int_{I_j} \kappa(x - y) b_j(y) dy.$$

Using (10.161), the above is equal to

$$\int_{I_j} (\kappa(x-y) - \kappa(x-c_j)) b_j(y) dy.$$

Thus, using the triangle inequality, (10.172), (10.173) and Lemma 10.14, we can estimate

$$\begin{aligned} \sum_{j \in \mathcal{J}_1(x)} |H_r b_j(x)| &\leq \sum_{j \in \mathcal{J}_1(x)} 2^{10} \int_{I_j} \frac{|y-c_j|}{|x-c_j|^2} |b_j(y)| dy \\ &\leq 2^{10} \sum_j \frac{(t_j - s_j)}{|x-c_j|^2} \int_{I_j} |b_j(y)| dy := F_1(x). \end{aligned} \quad (10.178)$$

Next, we estimate the second sum in (10.177). For each $j \in \mathcal{J}_2(x)$, set

$$d_j := \frac{1}{t_j - s_j} \int_{I_j} \mathbf{1}_{\{z: r < |z| < 1\}} (x-y) b_j(y) dy.$$

Then by (10.162)

$$|d_j| \leq 2 \cdot 2^{-10} \alpha = 2^{-9} \alpha. \quad (10.179)$$

For each $j \in \mathcal{J}_2(x)$, we have

$$\begin{aligned} H_r b_j(x) &= \int_{I_j} \kappa(x-y) (\mathbf{1}_{\{z: r < |z| < 1\}} (x-y) b_j(y) - d_j) dy \\ &\quad + \int_{I_j} d_j \kappa(x-y) dy. \\ &= \int_{I_j} (\kappa(x-y) - \kappa(x-c_j)) (\mathbf{1}_{\{z: r < |z| < 1\}} (x-y) b_j(y) - d_j) dy \\ &\quad + \int_{I_j} d_j \kappa(x-y) dy. \end{aligned}$$

Thus, using the triangle inequality, the estimate above and (10.179), we obtain

$$\begin{aligned} |H_r b_j(x)| &\leq \int_{I_j} |(\kappa(x-y) - \kappa(x-c_j))| (|b_j(y)| + 2^{-9} \alpha) dy + 2^{-9} \alpha \int_{I_j} |\kappa(x-y)| dy. \end{aligned} \quad (10.180)$$

Using (10.172), (10.173) and Lemma 10.14, and arguing as in (10.178), we get the the first term above can be estimated by

$$2^{10} \sum_j \frac{(t_j - s_j)}{|x-c_j|^2} \left(\int_{I_j} |b_j(y)| + 2^{-9} \alpha (t_j - s_j) \right) = F_1(x) + F_2(x), \quad (10.181)$$

with F_1 as in (10.178) and

$$F_2(x) := 2 \sum_j \frac{(t_j - s_j)}{|x-c_j|^2} \alpha (t_j - s_j). \quad (10.182)$$

For each $j \in \mathcal{J}_2(x)$, let $y_j \in [s_j, t_j]$ be such that

$$|x - y_j| = r.$$

Using (10.173), we have

$$r = |x - y_j| > 2(t_j - s_j).$$

Further, using the triangle inequality, for each $y \in I_j$, we obtain

$$|x - y| \leq |x - y_j| + |y - y_j| \leq r + (t_j - s_j) < \frac{3r}{2},$$

and

$$|x - y| \geq |x - y_j| - |y - y_j| \geq r - (t_j - s_j) > \frac{r}{2}.$$

We conclude that

$$[s_j, t_j) \subset \left\{ y : \frac{r}{2} < |x - y| < \frac{3r}{2} \right\}.$$

Using this and Lemma 10.13, we get

$$\int_{I_j} |\kappa(x - y)| dy \leq \int_{\{y: \frac{r}{2} < |x - y| < \frac{3r}{2}\}} \frac{8}{|x - y|} dy \leq 32. \quad (10.183)$$

Combining (10.177), (10.178), (10.180), (10.181) and (10.183), we get

$$|H_r b(x)| \leq 2F_1(x) + F_2(x) + 2^{-4}\alpha.$$

Using the triangle inequality, we deduce that

$$\begin{aligned} & \mu(\{x \in \Omega : |H_r b(x)| > \alpha/4\}) \leq \\ & \mu(\{x \in \Omega : |F_1(x)| > 2^{-4}\alpha\}) + \mu(\{x \in \Omega : |F_2(x)| > 2^{-4}\alpha\}). \end{aligned} \quad (10.184)$$

We estimate

$$\mu(\{x \in \Omega : |F_1(x)| > 2^{-4}\alpha\}) \leq \frac{2^{14}}{\alpha} \sum_j \int_{I_j} |b_j(y)| dy \int_{\Omega} \frac{(t_j - s_j)}{|x - c_j|^2} dx. \quad (10.185)$$

Using (10.172), we can bound

$$\int_{\Omega} \frac{(t_j - s_j)}{|x - c_j|^2} dx \leq 2(t_j - s_j) \int_{5|t_j - s_j|}^{\infty} \frac{1}{t^2} dt = \frac{2}{5}. \quad (10.186)$$

Plugging (10.186) into (10.185) and using (10.164), we conclude that

$$\mu(\{x \in \Omega : |F_1(x)| > 2^{-4}\alpha\}) \leq \frac{2^{14}}{\alpha} \int_{\Omega} |f(y)| dy. \quad (10.187)$$

Similarly, we estimate

$$\mu(\{x \in \Omega : |F_2(x)| > 2^{-4}\alpha\}) \leq \frac{2^5}{\alpha} \sum_j \alpha(t_j - s_j) \int_{\Omega} \frac{(t_j - s_j)}{|x - c_j|^2} dx.$$

Using (10.186) and (10.163), this is bounded by

$$2^4 \sum_j (t_j - s_j) \leq \frac{2^{15}}{\alpha} \int_{\Omega} |f(y)| dy. \quad (10.188)$$

Combining estimates (10.174), (10.175), (10.176), (10.184), (10.187) and (10.188) yields (10.32). \square

10.6. The proof of the van der Corput Lemma 10.8. Let g be a Lipschitz continuous function as in the lemma. We have

$$e^{in(x+\pi/n)} = -e^{inx}.$$

Using this, we write

$$\int_{\alpha}^{\beta} g(x)e^{-inx} dx = \frac{1}{2} \int_{\alpha}^{\beta} g(x)e^{inx} dx - \frac{1}{2} \int_{\alpha}^{\beta} g(x)e^{in(x+\pi/n)} dx.$$

We split the first integral at $\alpha + \frac{\pi}{n}$ and the second one at $\beta - \frac{\pi}{n}$, and make a change of variables in the second part of the first integral to obtain

$$\begin{aligned} &= \frac{1}{2} \int_{\alpha}^{\alpha+\frac{\pi}{n}} g(x)e^{inx} dx - \frac{1}{2} \int_{\beta-\frac{\pi}{n}}^{\beta} g(x)e^{in(x+\pi/n)} dx \\ &\quad + \frac{1}{2} \int_{\alpha+\frac{\pi}{n}}^{\beta} (g(x) - g(x - \frac{\pi}{n}))e^{inx} dx. \end{aligned}$$

The sum of the first two terms is by the triangle inequality bounded by

$$\frac{\pi}{n} \sup_{x \in [\alpha, \beta]} |g(x)|.$$

The third term is by the triangle inequality at most

$$\begin{aligned} &\frac{1}{2} \int_{\alpha+\frac{\pi}{n}}^{\beta} |g(x) - g(x - \frac{\pi}{n})| dx \\ &\leq \frac{\pi}{2n} \sup_{\alpha \leq x < y \leq \beta} \frac{|g(x) - g(y)|}{|x - y|} |\beta - \alpha|. \end{aligned}$$

Adding the two terms, we obtain

$$\left| \int_{\alpha}^{\beta} g(x)e^{-inx} dx \right| \leq \frac{\pi}{n} \|g\|_{\text{Lip}(\alpha, \beta)}.$$

By the triangle inequality, we also have

$$\left| \int_{\alpha}^{\beta} g(x)e^{-inx} dx \right| \leq |\beta - \alpha| \sup_{x \in [\alpha, \beta]} |g(x)| \leq |\beta - \alpha| \|g\|_{\text{Lip}(\alpha, \beta)}.$$

This completes the proof of the lemma, using that

$$\min\{|\beta - \alpha|, \frac{\pi}{n}\} \leq 2\pi|\beta - \alpha|(1 + n|\beta - \alpha|)^{-1}.$$

10.7. Partial sums as orthogonal projections. This subsection proves Lemma 10.12

Lemma 10.33 (partial sum projection). *Let f be a bounded 2π -periodic measurable function. Then, for all $N \geq 0$*

$$S_N(S_N f) = S_N f. \quad (10.189)$$

Proof. Let $N > 0$ be given. With K_N as in Lemma 10.10,

$$\begin{aligned} S_N(S_N f)(x) &= \frac{1}{2\pi} \int_0^{2\pi} S_N f(y) K_N(x - y) dy \\ &= \frac{1}{(2\pi)^2} \int_0^{2\pi} \int_0^{2\pi} f(y') K_N(y - y') K_N(x - y) dy' dy. \end{aligned} \quad (10.190)$$

We have by Lemma 10.10

$$\begin{aligned}
& \frac{1}{2\pi} \int_0^{2\pi} K_N(y - y') K_N(x - y) dy \\
&= \frac{1}{2\pi} \sum_{n=-N}^N \sum_{n'=-N}^N \int_0^{2\pi} e^{in(y-y')} e^{in'(x-y)} dy \\
&= \frac{1}{2\pi} \sum_{n=-N}^N \sum_{n'=-N}^N e^{i(n'x - ny')} \int_0^{2\pi} e^{i(n-n')y} dy. \tag{10.191}
\end{aligned}$$

By Lemma 10.9, the summands for $n \neq n'$ vanish. We obtain for (10.191)

$$= \frac{1}{2\pi} \sum_{n=-N}^N e^{in(x-y')} \int_0^{2\pi} dy = K_N(x - y'). \tag{10.192}$$

Applying Fubini in (10.190) and using (10.192) gives

$$S_N(S_N f)(x) = \frac{1}{2\pi} \int_0^{2\pi} f(y') K(x - y') dy' = S_N f(x) \tag{10.193}$$

This proves the lemma. \square

Lemma 10.34 (partial sum selfadjoint). *We have for any 2π -periodic bounded measurable g, f that*

$$\int_0^{2\pi} \overline{S_N f(x)} g(x) dx = \int_0^{2\pi} \overline{f(x)} S_N g(x) dx. \tag{10.194}$$

Proof. We have with K_N as in Lemma 10.10 for every x

$$\overline{K_N(x)} = \sum_{n=-N}^N \overline{e^{inx}} = \sum_{n=-N}^N e^{-inx} = K_N(-x). \tag{10.195}$$

Further, with Lemma 10.10 and Fubini

$$\begin{aligned}
& \int_0^{2\pi} \overline{S_N f(x)} g(x) dx = \frac{1}{2\pi} \int_0^{2\pi} \int_{-\pi}^{\pi} \overline{f(y) K_N(x - y)} g(x) dy dx \\
&= \frac{1}{2\pi} \int_0^{2\pi} \int_{-\pi}^{\pi} \overline{f(y)} K_N(y - x) g(x) dx dy = \int_0^{2\pi} \overline{f(x)} S_N g(x) dx. \tag{10.196}
\end{aligned}$$

This proves the lemma. \square

We turn to the proof of Lemma 10.12.

We have with Lemma 10.34, then Lemma 10.33 and the Lemma 10.34 again

$$\begin{aligned}
& \int_0^{2\pi} S_N f(x) \overline{S_N f(x)} dx \int_0^{2\pi} f(x) \overline{S_N(S_N f)(x)} dx \\
&= \int_0^{2\pi} f(x) \overline{S_N f(x)} dx = \int_0^{2\pi} S_N f(x) \overline{f(x)} dx. \tag{10.197}
\end{aligned}$$

We have by the distributive law

$$\int_0^{2\pi} (f(x) - S_N f(x)) \overline{(f(x) - S_N f(x))} dx = \tag{10.198}$$

$$\int_0^{2\pi} f(x)\overline{f(x)} - S_N f(x)\overline{f(x)} - f(x)\overline{S_N f(x)} + S_N f(x)\overline{S_N f(x)} dx$$

Using the various identities expressed in (10.197), this becomes

$$= \int_0^{2\pi} f(x)\overline{f(x)} dx - \int_0^{2\pi} S_N f(x)\overline{S_N f(x)} dx. \quad (10.199)$$

As (10.198) has nonnegative integrand and is thus nonnegative, we conclude

$$\int_0^{2\pi} S_N f(x)\overline{S_N f(x)} dx \leq \int_0^{2\pi} f(x)\overline{f(x)} dx. \quad (10.200)$$

As both sides are positive, we may take the square root of this inequality. This completes the proof of the lemma.

10.8. The error bound. We prove Lemma 10.3. Define

$$E_2 := \{x \in [0, 2\pi) : \sup_{N>0} |S_N f(x) - S_N f_0(x)| > \frac{\epsilon}{4}\}.$$

Then (10.17) clearly holds, and it remains to show that $|E_2| \leq \frac{\epsilon}{2}$. This will follow from Lemma 10.4.

Let $x \in E_2$. Then there exists $N > 0$ with

$$|S_N f(x) - S_N f_0(x)| > \frac{\epsilon}{4},$$

pick such N . We have with Lemma 10.10

$$\frac{\epsilon}{4} < |S_N f(x) - S_N f_0(x)| = \frac{1}{2\pi} \left| \int_0^{2\pi} (f(y) - f_0(y)) K_N(x - y) dy \right|.$$

We make a change of variables, replacing y by $x - y$. Then we use 2π -periodicity of f , f_0 and K_N to shift the domain of integration to obtain

$$= \frac{1}{2\pi} \left| \int_{-\pi}^{\pi} (f(x - y) - f_0(x - y)) K_N(y) dy \right|.$$

Using the triangle inequality, we split this as

$$\begin{aligned} &\leq \frac{1}{2\pi} \left| \int_{-\pi}^{\pi} (f(x - y) - f_0(x - y)) \max(1 - |y|, 0) K_N(y) dy \right| \\ &\quad + \frac{1}{2\pi} \left| \int_{-\pi}^{\pi} (f(x - y) - f_0(x - y)) \min(|y|, 1) K_N(y) dy \right|. \end{aligned}$$

Note that all integrals are well defined, since K_N is by (10.37) bounded by $2N + 1$. Using (10.38) and the definition (10.24) of κ , we rewrite the two terms and obtain

$$\begin{aligned} \frac{\epsilon}{4} &< \frac{1}{2\pi} \left| \int_{-\pi}^{\pi} (f(x - y) - f_0(x - y)) (e^{iNy} \overline{\kappa}(y) + e^{-iNy} \kappa(y)) dy \right| \quad (10.201) \\ &+ \frac{1}{2\pi} \left| \int_{-\pi}^{\pi} (f(x - y) - f_0(x - y)) (e^{iNy} \frac{\min\{|y|, 1\}}{1 - e^{-iy}} + e^{-iNy} \frac{\min\{|y|, 1\}}{1 - e^{iy}}) dy \right|. \end{aligned} \quad (10.202)$$

By Lemma 10.11 with $\eta = |y|$, we have for $-1 \leq y \leq 1$

$$|e^{iNy} \frac{\min\{|y|, 1\}}{1 - e^{-iy}}| = \frac{|y|}{|1 - e^{iy}|} \leq 8.$$

By Lemma 10.11 with $\eta = 1$, we have for $1 \leq |y| \leq \pi$

$$|e^{iNy} \frac{\min\{|y|, 1\}}{1 - e^{-iy}}| = \frac{1}{|1 - e^{iy}|} \leq 8.$$

Thus we obtain using the triangle inequality and (10.5)

$$(10.202) \leq \frac{16}{2\pi} \int_{-\pi}^{\pi} |f(x-y) - f_0(x-y)| dy \leq 2^{4-2^{50}} \epsilon^2.$$

Consequently, we have that

$$\frac{\epsilon}{8} \leq \frac{1}{2\pi} \left| \int_{-\pi}^{\pi} (f(x-y) - f_0(x-y))(e^{iNy}\overline{\kappa}(y) + e^{-iNy}\kappa(y)) dy \right|.$$

By dominated convergence and since $\kappa(y) = 0$ for $|y| > 1$, this equals

$$= \frac{1}{2\pi} \lim_{r \rightarrow 0^+} \left| \int_{r < |y| < 1} (f(x-y) - f_0(x-y))(e^{iNy}\overline{\kappa}(y) + e^{-iNy}\kappa(y)) dy \right|.$$

Let $h = (f - f_0)\mathbf{1}_{[-\pi, 3\pi]}$. Since $x \in [0, 2\pi]$, the above integral does not change if we replace $(f - f_0)$ by h . We do that, apply the triangle inequality and bound the limits by suprema

$$\begin{aligned} &\leq \frac{1}{2\pi} \sup_{r > 0} \left| \int_{r < |y| < 1} h(x-y) e^{-iNy}\kappa(y) dy \right| \\ &\quad + \frac{1}{2\pi} \sup_{r > 0} \left| \int_{r < |y| < 1} \bar{h}(x-y) e^{-iNy}\kappa(y) dy \right|. \end{aligned}$$

By the definition (10.27) of T , this is

$$\leq \frac{1}{2\pi} (Th(x) + T\bar{h}(x)).$$

Thus for each $x \in E_2$, at least one of $Th(x)$ and $T\bar{h}(x)$ is larger than $\frac{\epsilon}{16}$. Thus at least one of Th and $T\bar{h}$ is $\geq \frac{\epsilon}{16}$ on a subset E'_2 of E_2 with $2|E'_2| \geq |E_2|$. Without loss of generality this is Th . By assumption (10.5), we have $|2^{2^{50}} \epsilon^{-2} h| \leq \mathbf{1}_{[-\pi, 3\pi]}$. Applying Lemma 10.4 with $F = [-\pi, 3\pi]$ and $G = E'_2$, it follows that

$$\begin{aligned} \frac{\epsilon}{16} |E'_2| &\leq \int_{E'_2} Th(x) dx = 2^{-2^{50}} \epsilon^2 \int_{E'_2} T(2^{2^{50}} \epsilon^{-2} h)(x) dx \\ &\leq 2^{-2^{50}} \epsilon^2 \cdot 2^{2^{40}} |F|^{\frac{1}{2}} |E'_2|^{\frac{1}{2}} \leq 2^{-2^{40}} \frac{\epsilon^2}{16} |E'_2|^{\frac{1}{2}}. \end{aligned}$$

Rearranging, we obtain

$$|E'_2| \leq 2^{-2^{41}} \epsilon^2 \leq \frac{\epsilon}{4}.$$

This completes the proof using $|E_2| \leq 2|E'_2|$.

10.9. Carleson on the real line. We prove Lemma 10.4.

Consider the standard distance function

$$\rho(x, y) = |x - y| \quad (10.203)$$

on the real line \mathbb{R} .

Lemma 10.35 (real line metric). *The space (\mathbb{R}, ρ) is a complete locally compact metric space.*

Proof. This is part of the Lean library. \square

Lemma 10.36 (real line ball). *For $x \in \mathbb{R}$ and $R > 0$, the ball $B(x, R)$ is the interval $(x - R, x + R)$*

Proof. Let $x' \in B(x, R)$. By definition of the ball, $|x' - x| < R$. It follows that $x' - x < R$ and $x - x' < R$. It follows $x' < x + R$ and $x' > x - R$. This implies $x' \in (x - R, x + R)$. Conversely, let $x' \in (x - R, x + R)$. Then $x' < x + R$ and $x' > x - R$. It follows that $x' - x < R$ and $x - x' < R$. It follows that $|x' - x| < R$, hence $x' \in B(x, R)$. This proves the lemma. \square

We consider the Lebesgue measure μ on \mathbb{R} .

Lemma 10.37 (real line measure). *The measure μ is a sigma-finite non-zero Radon-Borel measure on \mathbb{R} .*

Proof. This is part of the Lean library. \square

Lemma 10.38 (real line ball measure). *We have for every $x \in \mathbb{R}$ and $R > 0$*

$$\mu(B(x, R)) = 2R. \quad (10.204)$$

Proof. We have with Lemma 10.36

$$\mu(B(x, R)) = \mu((x - R, x + R)) = 2R. \quad (10.205)$$

\square

Lemma 10.39 (real line doubling). *We have for every $x \in \mathbb{R}$ and $R > 0$*

$$\mu(B(x, 2R)) = 2\mu(B(x, R)). \quad (10.206)$$

Proof. We have with Lemma 10.38

$$\mu(B(x, 2R)) = 4R = 2\mu(B(x, R)). \quad (10.207)$$

This proves the lemma. \square

The preceding four lemmas show that $(\mathbb{R}, \rho, \mu, 4)$ is a doubling metric measure space. Indeed, we even show that $(\mathbb{R}, \rho, \mu, 1)$ is a doubling metric measure space, but we may relax the estimate in Lemma 10.39 to conclude that $(\mathbb{R}, \rho, \mu, 4)$ is a doubling metric measure space.

For each $n \in \mathbb{Z}$ define $\vartheta_n : \mathbb{R} \rightarrow \mathbb{R}$ by

$$\vartheta_n(x) = nx. \quad (10.208)$$

Let Θ be the collection $\{\vartheta_n, n \in \mathbb{Z}\}$. Note that for every $n \in \mathbb{Z}$ we have $\vartheta_n(0) = 0$. Define

$$d_{B(x, R)}(\vartheta_n, \vartheta_m) := 2R|n - m|. \quad (10.209)$$

Lemma 10.40 (frequency metric). *For every $R > 0$ and $x \in X$, the function $d_{B(x, R)}$ is a metric on Θ .*

Proof. This follows immediately from the fact that the standard metric on \mathbb{Z} is a metric. \square

Lemma 10.41 (oscillation control). *For every $R > 0$ and $x \in X$, and for all $n, m \in \mathbb{Z}$, we have*

$$\sup_{y, y' \in B(x, R)} |ny - ny' - my + my'| \leq 2|n - m|R. \quad (10.210)$$

Proof. The right hand side of (10.210) equals

$$\sup_{y, y' \in B(x, R)} |(n - m)(y - x) - (n - m)(y' - x)|.$$

The lemma then follows from the triangle inequality. \square

Lemma 10.42 (frequency monotone). *For any $x, x' \in X$ and $R, R' > 0$ with $B(x, R) \subset B(x', R')$, and for any $n, m \in \mathbb{Z}$*

$$d_{B(x, R)}(\vartheta_n, \vartheta_m) \leq d_{B(x', R')}(\vartheta_n, \vartheta_m).$$

Proof. This follows immediately from the definition (10.209) and $R \leq R'$. \square

Lemma 10.43 (frequency ball doubling). *For any $x, x' \in \mathbb{R}$ and $R > 0$ with $x \in B(x', 2R)$ and any $n, m \in \mathbb{Z}$, we have*

$$d_{B(x', 2R)}(\vartheta_n, \vartheta_m) \leq 2d_{B(x, R)}(\vartheta_n, \vartheta_m). \quad (10.211)$$

Proof. With (10.209), both sides of (10.211) are equal to $4R|n - m|$. This proves the lemma. \square

Lemma 10.44 (frequency ball growth). *For any $x, x' \in \mathbb{R}$ and $R > 0$ with $B(x, R) \subset B(x', 2R)$ and any $n, m \in \mathbb{Z}$, we have*

$$2d_{B(x, R)}(\vartheta_n, \vartheta_m) \leq d_{B(x', 2R)}(\vartheta_n, \vartheta_m). \quad (10.212)$$

Proof. With (10.209), both sides of (10.211) are equal to $4R|n - m|$. This proves the lemma. \square

Lemma 10.45 (integer ball cover). *For every $x \in \mathbb{R}$ and $R > 0$ and every $n \in \mathbb{Z}$ and $R' > 0$, there exist $m_1, m_2, m_3 \in \mathbb{Z}$ such that*

$$B' \subset B_1 \cup B_2 \cup B_3, \quad (10.213)$$

where

$$B' = \{\vartheta \in \Theta : d_{B(x, R)}(\vartheta, \vartheta_n) < 2R'\} \quad (10.214)$$

and for $j = 1, 2, 3$

$$B_j = \{\vartheta \in \Theta : d_{B(x, R)}(\vartheta, \vartheta_{m_j}) < R'\}. \quad (10.215)$$

Proof. Let m_1 be the largest integer smaller than or equal to $n - R'/2$. Let $m_2 = n$. Let m_3 be the smallest integer larger than or equal to $n + R'/2$.

Let $\vartheta_{n'} \in B'$, then with (10.209), we have

$$2R|n - n'| < 2R'. \quad (10.216)$$

Assume first $n' \leq n - R'/2$. By definition of m_1 , we have $n' \leq m_1$. With (10.216) we have

$$R|m_1 - n'| = R(m_1 - n') = R(m_1 - n) + R(n - n')$$

$$< -\frac{R'}{2} + R' = -\frac{R'}{2}. \quad (10.217)$$

We conclude $\vartheta_{n'} \in B_1$.

Assume next $n - R'/2 < n' < n + R'/2$. Then $\vartheta_{n'} \in B_2$.

Assume finally that $n + R'/2 \leq n'$. By definition of m_3 , we have $m_3 \leq n'$. With (10.216) we have

$$\begin{aligned} R|m_3 - n'| &= R(n' - m_3) = R(n' - n) + R(n - m_3) \\ &< R' - \frac{R'}{2} = -\frac{R'}{2}. \end{aligned} \quad (10.218)$$

We conclude $\vartheta_{n'} \in B_1$. This completes the proof of the lemma. \square

Lemma 10.46 (real van der Corput). *For any $x \in \mathbb{R}$ and $R > 0$ and any function $\varphi : X \rightarrow \mathbb{C}$ supported on $B' = B(x, R)$ such that*

$$\|\varphi\|_{\text{Lip}(B')} = \sup_{x \in B'} |\varphi(x)| + R \sup_{x, y \in B', x \neq y} \frac{|\varphi(x) - \varphi(y)|}{\rho(x, y)} \quad (10.219)$$

is finite and for any $n, m \in \mathbb{Z}$, we have

$$\left| \int_{B'} e(\vartheta_n(x) - \vartheta_m(x)) \varphi(x) d\mu(x) \right| \leq 2\pi\mu(B') \frac{\|\varphi\|_{\text{Lip}(B')}}{1 + d_{B'}(\vartheta_n, \vartheta_m)}. \quad (10.220)$$

Proof. Set $n' = n - m$. Then we have to prove

$$\left| \int_{x-R}^{x+R} e^{in'y} \varphi(y) dy \right| \leq 4\pi R \|\varphi\|_{\text{Lip}(B')} (1 + 2R|n'|)^{-1}. \quad (10.221)$$

This follows from Lemma 10.8 with $\alpha = x - R$ and $\beta = x + R$. \square

The preceding chain of lemmas establish that Θ is a cancellative, compatible collection of functions on $(\mathbb{R}, \rho, \mu, 4)$. Again, some of the statements in these lemmas are stronger than what is needed for $a = 4$, but can be relaxed to give the desired conclusion for $a = 4$.

With κ as near (10.24), define the function $K : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{C}$ as in Theorem 1.2 by

$$K(x, y) := \kappa(x - y). \quad (10.222)$$

The function K is continuous outside the diagonal $x = y$ and vanishes on the diagonal. Hence it is measurable.

By the Lemmas 10.13 and 10.14, it follows that K is a one-sided Calderón–Zygmund kernel on $(\mathbb{R}, \rho, \mu, 4)$.

The operator T^* defined in (1.16) coincides in our setting with the operator T^* defined in (10.29). By Lemma 10.5, this operator satisfies the bound (1.18).

Thus the assumptions of Theorem 1.2 are all satisfied. Applying the Theorem, Lemma 10.4 follows.

REFERENCES

- [Bec24] L. Becker. Maximal polynomial modulations of singular Radon transforms. *J. Funct. Anal.*, 286(6):Paper No. 110299, 46, 2024 (cited on page 5).
- [Car66] L. Carleson. On convergence and growth of partial sums of Fourier series. *Acta Math.*, 116:135–157, 1966 (cited on pages 2, 5).

- [Chr90] M. Christ. A $t(b)$ theorem with remarks on analytic capacity and the Cauchy integral. *Colloquium Mathematicum*, 2(60-61):601–628, 1990 (cited on page 17).
- [CW71] R. R. Coifman and G. Weiss. *Analyse harmonique non-commutative sur certains espaces homogènes*. Lecture Notes in Mathematics, Vol. 242. Springer-Verlag, Berlin-New York, 1971. Étude de certaines intégrales singulières (cited on page 5).
- [Fef73] C. Fefferman. Pointwise convergence of Fourier series. *Ann. of Math. (2)*, 98:551–571, 1973 (cited on pages 5, 28).
- [Lie09] V. Lie. The (weak- L^2) boundedness of the quadratic Carleson operator. *Geom. Funct. Anal.*, 19(2):457–497, 2009 (cited on page 5).
- [Lie20] V. Lie. The polynomial Carleson operator. *Ann. of Math. (2)*, 192(1):47–163, 2020 (cited on pages 3, 5).
- [LT00] M. Lacey and C. Thiele. A proof of boundedness of the Carleson operator. *Math. Res. Lett.*, 7(4):361–370, 2000 (cited on page 5).
- [Mna22] G. Mnatsakanyan. On almost-everywhere convergence of Malmquist Takenaka series. *J. Funct. Anal.*, 282(12):Paper No. 109461, 33, 2022 (cited on page 5).
- [MS79] R. A. Macías and C. Segovia. Lipschitz functions on spaces of homogeneous type. *Adv. in Math.*, 33(3):257–270, 1979 (cited on page 5).
- [Ram21] J. P. G. Ramos. The Hilbert transform along the parabola, the polynomial Carleson theorem and oscillatory singular integrals. *Math. Ann.*, 379(1-2):159–185, 2021 (cited on page 5).
- [Ste93] E. M. Stein. *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, volume 43 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1993, pages xiv+695. With the assistance of Timothy S. Murphy, *Mono-graphs in Harmonic Analysis, III* (cited on pages 5, 116).
- [SW01] E. M. Stein and S. Wainger. Oscillatory integrals related to Carleson’s theorem. *Math. Res. Lett.*, 8(5-6):789–800, 2001 (cited on page 5).
- [Zor21] P. Zorin-Kranich. Maximal polynomial modulations of singular integrals. *Adv. Math.*, 386:Paper No. 107832, 40, 2021 (cited on pages 3, 5, 9, 17).

MATHEMATICAL INSTITUTE, UNIVERSITY OF BONN, ENDENICHER ALLEE 60, 53115,
BONN, GERMANY

Email address: `becker@math.uni-bonn.de`

MATHEMATICAL INSTITUTE, UNIVERSITY OF BONN, ENDENICHER ALLEE 60, 53115,
BONN, GERMANY

Email address: `vdoorn@math.uni-bonn.de`

COLLEGE OF SCIENCES, KOÇ UNIVERSITY, RUMELİ FENERİ YOLU, 34450 SARIYER,
İSTANBUL, TÜRKİYE

Email address: `ajamneshan@ku.edu.tr`

MATHEMATICAL INSTITUTE, UNIVERSITY OF BONN, ENDENICHER ALLEE 60, 53115,
BONN, GERMANY, AND

MAX PLANCK INSTITUTE FOR MATHEMATICS, VIVATSGASSE 7, 53111, BONN, GERMANY.

Email address: `rajulas@math.uni-bonn.de`

MATHEMATICAL INSTITUTE, UNIVERSITY OF BONN, ENDENICHER ALLEE 60, 53115,
BONN, GERMANY

Email address: `thiele@math.uni-bonn.de`