

# Tuning parameter selection for the adaptive nuclear norm regularized trace regression

Pan Shang, Lingchen Kong, Yiting Ma \*

May 14, 2024

**Abstract:** Regularized models have been applied in lots of areas, with high-dimensional data sets being popular. Because tuning parameter decides the theoretical performance and computational efficiency of the regularized models, tuning parameter selection is a basic and important issue. We consider the tuning parameter selection for adaptive nuclear norm regularized trace regression, which achieves by the Bayesian information criterion (BIC). The proposed BIC is established with the help of an unbiased estimator of degrees of freedom. Under some regularized conditions, this BIC is proved to achieve the rank consistency of the tuning parameter selection. That is the model solution under selected tuning parameter converges to the true solution and has the same rank with that of the true solution in probability. Some numerical results are presented to evaluate the performance of the proposed BIC on tuning parameter selection.

*Keywords:* Tuning parameter selection, Adaptive nuclear norm regularized trace regression, Bayesian information criterion, Degrees of freedom

*MSCcodes(2020):* 62J12, 62H12, 90C25, 15A24

## 1 Introduction

To deal with high-dimensional data sets with special structures, there are plenty of works that focus on regularized models. These models can be unified as a minimization problem, whose objective function is composed with a loss function and some regularizer. So, existing works consider different loss functions and regularizer under different data settings. For instance, the famous Lasso[34] and lots of similar models[35, 56, 48, 55, 57, 11, 49] inducing sparsity, regularized matrix regression[52] and some other models inducing low-rank solution[47, 30, 24, 22, 46, 53, 43, 54, 26, 52, 51, 9, 10, 14] and so on. Undoubtedly, these regularized models have been applied to lots of areas, see e.g., bioinformatics[50], engineering[28], signal process[7], system identification[23] and so on. Among these models, there is

---

\*Address: Pan Shang is with Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China; Lingchen Kong and Yiting Ma are with the School of Mathematics and Statistics, Beijing Jiaotong University, Beijing, 100044, China.

E-mail: pshang@amss.ac.cn

an interesting one called the adaptive nuclear norm regularized trace regression, which is proposed by Bach[3] and achieves the rank consistency. Therefore, we focus on this model in the paper.

For regularized models, their loss function and regularizer are summed together by the so-called tuning parameter. It seems that tuning parameter is insignificant, while the selection of this parameter decides the theory performance and computational effect of these models. Therefore, there are some tools can be used to select this parameter[44], such as Akaike information criterion (AIC,[2]), Bayesian information criterion (BIC,[31, 38, 39, 40, 41, 19, 12, 15, 33, 45, 1]), cross-validation ([16, 17, 20, 6, 5]), and so on. Here, we review some related works about existing tuning parameter selection methods for regularized models, which include regularized liner models with the unknown variable being in vector form and regularized matrix models with matrix as variable, respectively. The following results mainly focus on tuning parameter selection for regularized linear models. Wang et al. [40] established a BIC for SCAD under the case that the sample size  $n$  and feature size  $p$  satisfying  $n > p$  and  $p$  fixed. Wang et al. [39] considered the BIC for adaptive Lasso under  $n > p$  and  $p$  diverged. Wang and Zhu [41] builded up a high dimensional BIC (HBIC) for adaptive elastic net under  $n < p$  and  $p$  diverged. Kim et al. [19] builded a generalized information criterion for SCAD and MCP under different data settings, including  $n < p$  with  $p$  fixed and diverged. Hirose et al. [15] considered the extended regularized information criterion for adaptive Lasso. Li et al. [45] established the regularized information criterion for adaptive group regularized generalized linear models. These proposed information criteria all were proved to have the consistency property, that is the selected tuning parameter will lead to a solution with the same active index as the true one in probability, when the sample size increases. So, these regularized linear models all have an important property, which is the selection consistency. In addition, there are a few tuning parameter selection results for regularized matrix regression. Zhou and Li [52] proposed the AIC and BIC for regularized matrix regression under the orthogonal assumption of prediction matrix. Yuan [46] calculated the degree of freedom of regularized matrix regression, which is the basic for some tuning parameter selection methods. However, these results of regularized matrix regression have no theoretical guarantees of consistency.

In this paper, we tends to select the tuning parameter for adaptive nuclear norm regularized trace regression proposed by [3]. The core reason we consider this model is its rank consistency, i.e., the model solution achieved the true rank in probability with the sample size increasing. In addition, this model also includes the regularized matrix regression models in [52] and [48] as special cases. We first establish some basic results, including the subdifferential and conjugate function of the adaptive nuclear norm. These results help to analyze the model and provide tools to establish its degrees of freedom, which is an essential element to set up the tuning parameter selection criterion. Based on the definition in [8], we calculate the unbiased estimation of degrees of freedom, according to the model optimality and basic results of the model. Although the degrees of freedom seems like a complex form, we prove that this result includes the existing result in [52] as a special case. With the help of the estimator of the degrees of freedom, we propose a BIC to select the tuning parameter for the adaptive nuclear norm regularized trace regression. The proposed BIC is proved to achieve the rank selection consistency in probability,

under some regularized conditions. That is, the tuning parameter selected by BIC achieves selection consistency and rank selection consistency simultaneously. On numerical experiments, we compare the proposed BIC with AIC, AICc and cross validation for tuning parameter selection on some simulation data and real data sets, which illustrates the efficiency of our criterion.

This paper is organized as follows. We review the adaptive nuclear norm regularized trace regression and present some basic results in Section 2. In Section 3, we establish the degrees of freedom and BIC with theoretical guarantees. Some numerical experiments and conclusion are showed in Section 4 and Section 5, respectively.

*Notations:* Let  $M \in \mathbb{R}^{p_1 \times p_2}$  be a matrix.  $\text{vec}(M)$  denotes the vector in  $\mathbb{R}^{p_1 p_2}$  obtained by stacking its columns into a single vector. For any  $j \in \{1, 2, \dots, p_1\}$  and  $k \in \{1, 2, \dots, p_2\}$ ,  $M_{:,j}$  means the  $j$ th column and  $M_{k,:}$  means the  $k$ th row of  $M$ . Suppose  $M$  has a singular value decomposition with nonincreasing singular values  $\sigma_1(M) \geq \dots \geq \sigma_r(M) > 0$  and  $r \leq \min\{p_1, p_2\}$  is the rank of  $M$ . There are some related norms with singular values of  $M$ . The Frobenius norm  $\|\cdot\|_F$  is defined as  $\|M\|_F = \sqrt{\sum_{i=1}^{p_1} \sum_{j=1}^{p_2} M_{i,j}^2} = \sqrt{\sigma_1^2(M) + \dots + \sigma_r^2(M)}$ . The nuclear norm  $\|\cdot\|_*$  is the sum of non-zero singular values, i.e.,  $\|M\|_* = \sum_{i=1}^r \sigma_i(M)$ . The spectral norm  $\|\cdot\|_2$  is the largest singular value, i.e.,  $\|M\|_2 = \sigma_1(M)$ . A symmetric matrix  $M \in \mathbb{R}^{p \times p}$  is called positive semidefinite (positive definite), denoted as  $M \geq 0$  ( $M > 0$ ), if  $\mathbf{x}^\top M \mathbf{x} \geq 0$  ( $\mathbf{x}^\top M \mathbf{x} > 0$ ) holds for any  $0 \neq \mathbf{x} \in \mathbb{R}^p$ . Any  $M \in \mathbb{O}^p$  means that  $M \in \mathbb{R}^{p \times p}$  and  $M^\top M = I_p$ . For any vector  $\mathbf{x} \in \mathbb{R}^p$ , the norm  $\|\cdot\|$  is defined as  $\|\mathbf{x}\| = \sqrt{\sum_{i=1}^p x_i^2}$ . The notation  $\mathbf{x} \in \mathbb{R}_{++}^p$  means all elements of  $\mathbf{x}$  are positive. The indicator function of a set  $\mathcal{A}$  is denoted as  $\delta_{\mathcal{A}}(\cdot)$ , which means the value of this function is zero when variable in the set  $\mathcal{A}$  and is  $\infty$  otherwise. For any index set  $I$ , its cardinal number is denoted as  $|I|$  that counts the number in the index set  $I$ , and its complementary set is denoted as  $I^c$ . In this paper, the notation 0 may represent scalar, vector and matrix, which can be inferred from the context.

## 2 Preliminaries

This section presents the adaptive nuclear norm regularized trace regression with its analysis and show some optimal results of this model, which are foundations of the tuning parameter selection.

### 2.1 Model Analysis

The statistical model of the trace regression is

$$y = \langle X, B^* \rangle + \epsilon,$$

where  $X \in \mathbb{R}^{p_1 \times p_2}$  is the predictor,  $y \in \mathbb{R}$  is the responsor,  $\epsilon \in \mathbb{R}$  is a random error and  $B^* \in \mathbb{R}^{p_1 \times p_2}$  is the true coefficient. By sampling  $n$  times, we get

$$y_i = \langle X_i, B^* \rangle + \epsilon_i, \quad i = 1, 2, \dots, n.$$

Let  $\mathcal{X} = (\text{vec}(X_1), \dots, \text{vec}(X_n))^\top \in \mathbb{R}^{n \times p_1 p_2}$ ,  $\mathbf{y} = (y_1, \dots, y_n)^\top \in \mathbb{R}^n$  and  $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n)^\top \in \mathbb{R}^n$ . These sample models can be written as

$$\mathbf{y} = \mathcal{X} \text{vec}(B^*) + \varepsilon.$$

To estimate the unknown matrix  $B^*$ , there are some literatures focus on the nuclear norm regularized trace regression, such as Koltchinskii et al. [36], Negahban and Wainwright [27], Zhou and Li [52] and so on, which is

$$\min_{B \in \mathbb{R}^{p_1 \times p_2}} \left\{ \frac{1}{2n} \sum_{i=1}^n (y_i - \langle X_i, B \rangle)^2 + \lambda \|B\|_* \right\}.$$

Here,  $\lambda > 0$  is the tuning parameter. However, this model can hardly achieve the rank consistent solution in high-dimensional case as in Bach [3]. So, an adaptive version was proposed as

$$\min_{B \in \mathbb{R}^{p_1 \times p_2}} \left\{ \frac{1}{2n} \sum_{i=1}^n (y_i - \langle X_i, B \rangle)^2 + \lambda \|W_1 B W_2\|_* \right\}, \quad (1)$$

where  $W_1$  and  $W_2$  are given weight matrixes based on the solution of least square trace regression. The detailed results of these matrixes are showed as follows. Let

$$\hat{B}_{LS} = \underset{B \in \mathbb{R}^{p_1 \times p_2}}{\text{argmin}} \left\{ \frac{1}{2n} \sum_{i=1}^n (y_i - \langle X_i, B \rangle)^2 \right\}.$$

Suppose  $\hat{B}_{LS}$  has full rank. Let the singular value decomposition of  $\hat{B}_{LS}$  be  $\hat{B}_{LS} = U_{LS} \text{Diag}(s^{LS}) V_{LS}^\top$ , where  $U_{LS} \in \mathbb{O}^{p_1}$ ,  $V_{LS} \in \mathbb{O}^{p_2}$ ,  $s^{LS} \in \mathbb{R}^{\min\{p_1, p_2\}}$  and  $\text{Diag}(s^{LS})$  is a matrix whose diagonal vector is the singular vector  $s^{LS}$ .  $s^{LS}$  can be completed by  $n^{-1/2}$  to reach dimensions  $p_1$  or  $p_2$ . Then,

$$W_1 = U_{LS} \text{Diag}(s^{LS})^{-\gamma} U_{LS}^\top, \quad W_2 = V_{LS} \text{Diag}(s^{LS})^{-\gamma} V_{LS}^\top, \quad \gamma \in (0, 1].$$

Based on expressions of  $W_1$  and  $W_2$ , they are obvious symmetric matrixes, i.e.,  $W_1^\top = W_1$  and  $W_2^\top = W_2$ . In addition,  $W_1^{-1} = U_{LS} \text{Diag}(s^{LS})^\gamma U_{LS}^\top$  and  $W_2^{-1} = V_{LS} \text{Diag}(s^{LS})^\gamma V_{LS}^\top$ .

## 2.2 Basic Results

In this section, some related results are proposed and reviewed, which provide theoretical guarantees for the rest of this paper. The following definitions are from Rockafellar [29].

**Definition 2.1.** Let  $f : \mathbb{R}^{p_1 \times p_2} \rightarrow (-\infty, +\infty]$  be a proper closed convex function and let  $M \in \mathbb{R}^{p_1 \times p_2}$ . The subdifferential of  $f$  at  $M$  is denoted by  $\partial f(M)$  and

$$\partial f(M) = \{G \in \mathbb{R}^{p_1 \times p_2} : f(N) \geq f(M) + \langle G, N - M \rangle, \forall N \in \mathbb{R}^{p_1 \times p_2}\}.$$

According to this definition, we compute the subdifferential of  $f(M) = \|W_1 M W_2\|_*$  as follows.

**Proposition 2.1.** For any  $M \in \mathbb{R}^{p_1 \times p_2}$ , let  $r$  be the rank of  $W_1 M W_2$ . The subdifferential of  $\|W_1 M W_2\|_*$  is

$$\partial \|W_1 M W_2\|_* = \{W_1 (U_r V_r^\top + N) W_2 : W_1 M W_2 = U_r \text{Diag}(\sigma_r) V_r^\top, \|N\|_2 \leq 1, U_r^\top N = 0, N V_r = 0\},$$

where  $U_r \in \mathbb{R}^{p_1 \times r}$ ,  $V_r \in \mathbb{R}^{p_2 \times r}$ ,  $\sigma_r \in \mathbb{R}_{++}^r$  and  $N \in \mathbb{R}^{p_1 \times p_2}$ .

*Proof.* It is clear that  $f(M) = h(g(M))$  with  $g(M) = W_1 M W_2$  and  $h(M) = \lambda \|M\|_*$ . Because  $g$  is a linear transformation, according to [4, Theorem 3.43],

$$\partial f(M) = g^\top (\partial_{g(M)} h(g(M)))$$

with  $g^\top$  being the adjoint operator of  $g$ .

According to the definition of the adjoint operator,  $\langle g(M), N \rangle = \langle M, g^\top(N) \rangle$ , which leads to

$$g^\top(N) = W_1^\top N W_2^\top = W_1 N W_2.$$

Based on the singular value decomposition of  $W_1 M W_2$  and the subdifferential of the nuclear norm in Watson [42], it holds that

$$\partial_{g(M)} h(g(M)) = \{U_r V_r^\top + N : \|N\|_2 \leq 1, U_r^\top N = 0, N V_r = 0\}.$$

Combing these results, the desired result can be obtained. □

There are some basic calculation rule in [18, 25] that are foundations of our result, which we review them in the next proposition.

**Proposition 2.2.** Let  $A, B, C, D$  be any matrix in compatible dimensions, the following claims hold.

i)  $\text{vec}(ABC) = (C^\top \otimes A) \text{vec}(B)$ , where  $\otimes$  means the Kronecker product.

ii)  $\text{vec}(A^\top) = K \text{vec}(A)$ , where  $K$  is the permutation matrix.

iii)  $(A \otimes B)(C \otimes D) = AC \otimes BD$ .

iv)  $\frac{\partial \text{vec}(ABC)}{\partial \text{vec}^\top(B)} = C^\top \otimes A$ .

### 3 Bayesian Information Criterion

In this section, we tend to build up a Bayesian Information Criterion (BIC) and prove its consistency to select the appropriate tuning parameter for (1).

According to the definition of traditional BIC, a key basis is degrees of freedom of the model. So, we calculate degrees of freedom of (1) in the next theorem.

**Theorem 3.1.** For any  $\lambda \geq 0$ , let  $r$  be the rank of  $W_1 \hat{B}(\lambda) W_2$ . Suppose the singular value decomposition of  $W_1 \hat{B}(\lambda) W_2$  is  $W_1 \hat{B}(\lambda) W_2 = U_r \text{Diag}(b_r) V_r^\top$  with  $U_r \in \mathbb{R}^{p_1 \times r}$ ,  $V_r \in \mathbb{R}^{p_2 \times r}$  and  $b_r \in \mathbb{R}_{++}^r$ . An unbiased estimation of degrees of freedom of (1) under  $\lambda$  satisfies that

$$\hat{d}f_\lambda = \frac{1}{n} \sum_{k=1}^n \text{vec}(X_k)^\top M_r^+ \text{vec}(X_k).$$

Here,  $M_r = \frac{1}{n} \sum_{k=1}^n \text{vec}(X_k) \text{vec}(X_k)^\top + \lambda M_r^{(1)} + M_r^{(2)}$  is assumed full rank or symmetric, with  $M_r^{(1)}$  and  $M_r^{(2)}$  defined in Lemma 3.1 and Lemma 3.2, respectively.

*Proof.* As the result in Stein [32], an unbiased estimation of degrees of freedom of (1) under  $\lambda$  is

$$\hat{d}f_\lambda = \sum_{k=1}^n \frac{\partial y_k}{\partial y_k} = \sum_{k=1}^n \left\langle X_k, \frac{\partial \hat{B}(\lambda)}{\partial y_k} \right\rangle = \sum_{k=1}^n \left\langle \text{vec}(X_k), \frac{\partial \text{vec}(\hat{B}(\lambda))}{\partial y_k} \right\rangle.$$

According to the optimal condition of (1) and the subdifferential of  $\|W_1 B W_2\|_*$  in Proposition 2.1, there is a matrix  $N$  such that  $\|N\|_2 \leq 1$ ,  $U_r^\top N = 0$ ,  $N V_r = 0$  and

$$-\frac{1}{n} \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i + \lambda W_1 G(\hat{B}(\lambda)) W_2 + \lambda W_1 N W_2 = 0, \quad (2)$$

where  $G(\hat{B}(\lambda)) = U_r V_r^\top$ . Vectorizing this equality and using  $i$  in Proposition 2.2, we get that

$$\frac{1}{n} \sum_{i=1}^n y_i \text{vec}(X_i) = \frac{1}{n} \sum_{i=1}^n \text{vec}(X_i) \text{vec}(X_i)^\top \text{vec}(\hat{B}(\lambda)) + \lambda (W_2 \otimes W_1) \text{vec}(G(\hat{B}(\lambda))) + \lambda (W_2 \otimes W_1) \text{vec}(N). \quad (3)$$

The result of the partial differential of the equation (3) with respect of  $y_k$  is

$$\begin{aligned} \frac{1}{n} \text{vec}(X_k) &= \left[ \frac{1}{n} \sum_{i=1}^n \text{vec}(X_i) \text{vec}(X_i)^\top + \lambda (W_2 \otimes W_1) \frac{\partial \text{vec}(G(\hat{B}(\lambda)))}{\partial \text{vec}^\top(\hat{B}(\lambda))} + \lambda (W_2 \otimes W_1) \frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(\hat{B}(\lambda))} \right] \frac{\partial \text{vec}(\hat{B}(\lambda))}{\partial y_k} \\ &:= \left[ \frac{1}{n} \sum_{i=1}^n \text{vec}(X_i) \text{vec}(X_i)^\top + \lambda M_r^{(1)} + M_r^{(2)} \right] \frac{\partial \text{vec}(\hat{B}(\lambda))}{\partial y_k}, \end{aligned} \quad (4)$$

where

$$M_r^{(1)} = (W_2 \otimes W_1) \frac{\partial \text{vec}(G(\hat{B}(\lambda)))}{\partial \text{vec}^\top(\hat{B}(\lambda))}, \quad M_r^{(2)} = \lambda (W_2 \otimes W_1) \frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(\hat{B}(\lambda))}.$$

Denote  $M_r = \frac{1}{n} \sum_{i=1}^n \text{vec}(X_i) \text{vec}(X_i)^\top + \lambda M_r^{(1)} + M_r^{(2)}$ . Then, the equation (4) is transformed as

$$\frac{1}{n} \text{vec}(X_k) = M_r \frac{\partial \text{vec}(\hat{B}(\lambda))}{\partial y_k}. \quad (5)$$

To get the result of  $\hat{d}f_\lambda$ , we assume that  $M_r$  has full rank or it is a symmetric matrix. The results are illustrated as follows.

If  $M_r$  has full rank, it is obvious that  $\frac{\partial \text{vec}(\hat{B}(\lambda))}{\partial y_k} = \frac{1}{n} M_r^{-1} \text{vec}(X_k)$  and  $\hat{d}f_\lambda = \frac{1}{n} \sum_{k=1}^n \text{vec}(X_k)^\top M_r^{-1} \text{vec}(X_k)$ .

If  $M_r$  is symmetric, suppose the eigenvalue decomposition of  $M_r$  is

$$M_r = V \begin{pmatrix} \text{Diag}(\sigma(M_r)), & 0 \\ 0, & 0 \end{pmatrix} V^\top,$$

where  $V = (\mathbf{v}_1, \dots, \mathbf{v}_{r_M}, \mathbf{v}_{r_M+1}, \dots, \mathbf{v}_{p_1 p_2}) = (V_{r_M}, V_{r_M}^\perp) \in \mathbb{O}^{p_1 p_2}$ ,  $V_{r_M}^\top V_{r_M} = 0$  and  $\sigma(M_r) \in \mathbb{R}^{r_M}$  with  $r_M$  being the rank of  $M_r$ . Then, the pseudo inverse matrix of  $M_r$  is

$$M_r^+ = V \begin{pmatrix} \text{Diag}(\sigma(M_r))^{-1}, & 0 \\ 0, & 0 \end{pmatrix} V^\top.$$

Based on the equation (5), we have

$$\begin{aligned} \frac{1}{n} V_{r_M}^\top \text{vec}(X_k) &= V_{r_M}^\top M_r \frac{\partial \text{vec}(\hat{B}(\lambda))}{\partial y_k} = V_{r_M}^\top V \begin{pmatrix} \text{Diag}(\sigma(M_r)), & 0 \\ 0, & 0 \end{pmatrix} V^\top \frac{\partial \text{vec}(\hat{B}(\lambda))}{\partial y_k} \\ &= V_{r_M}^\top V \begin{pmatrix} \text{Diag}(\sigma(M_r)) V_{r_M}^\top \frac{\partial \text{vec}(\hat{B}(\lambda))}{\partial y_k} \\ 0 \end{pmatrix} = \left( V_{r_M}^\top V_{r_M}, V_{r_M}^\top V_{r_M}^\perp \right) \begin{pmatrix} \text{Diag}(\sigma(M_r)) V_{r_M}^\top \frac{\partial \text{vec}(\hat{B}(\lambda))}{\partial y_k} \\ 0 \end{pmatrix} \\ &= \left( 0, I_{p_1 p_2 - r_M} \right) \begin{pmatrix} \text{Diag}(\sigma(M_r)) V_{r_M}^\top \frac{\partial \text{vec}(\hat{B}(\lambda))}{\partial y_k} \\ 0 \end{pmatrix} = 0. \end{aligned} \quad (6)$$

Combining the fact that  $M_r^+ M_r = V_{r_M} V_{r_M}^\top$  and (6), we have

$$\begin{aligned} \hat{\text{d}}f_\lambda &= \sum_{k=1}^n \left\langle \text{vec}(X_k), \frac{\partial \text{vec}(\hat{B}(\lambda))}{\partial y_k} \right\rangle \\ &= \sum_{k=1}^n \text{vec}(X_k)^\top V V^\top \frac{\partial \text{vec}(\hat{B}(\lambda))}{\partial y_k} = \sum_{k=1}^n \text{vec}(X_k)^\top M_r^+ M_r \frac{\partial \text{vec}(\hat{B}(\lambda))}{\partial y_k} + \sum_{k=1}^n \text{vec}(X_k)^\top V_{r_M}^\perp V_{r_M}^\top \frac{\partial \text{vec}(\hat{B}(\lambda))}{\partial y_k} \\ &= \frac{1}{n} \sum_{k=1}^n \text{vec}(X_k)^\top M_r^+ \text{vec}(X_k) + \sum_{k=1}^n \text{vec}(X_k)^\top V_{r_M}^\perp V_{r_M}^\top \frac{\partial \text{vec}(\hat{B}(\lambda))}{\partial y_k} \\ &\stackrel{(5)}{=} \frac{1}{n} \sum_{k=1}^n \text{vec}(X_k)^\top M_r^+ \text{vec}(X_k) + \sum_{k=1}^n \left\langle V_{r_M}^\perp \text{vec}(X_k), V_{r_M}^\top \frac{\partial \text{vec}(\hat{B}(\lambda))}{\partial y_k} \right\rangle \\ &= \frac{1}{n} \sum_{k=1}^n \text{vec}(X_k)^\top M_r^+ \text{vec}(X_k). \end{aligned} \quad (6)$$

These two results leads to Theorem 3.1. □

Next, we give detailed results of  $M_r^{(1)}$  and  $M_r^{(2)}$  in following lemmas.

**Lemma 3.1.** *For any  $\lambda \geq 0$ , let  $r$  be the rank of  $W_1 \hat{B}(\lambda) W_2$ . Suppose the singular value decomposition of  $W_1 \hat{B}(\lambda) W_2$  is  $W_1 \hat{B}(\lambda) W_2 = U_r \text{Diag}(b_r) V_r^\top$  with  $U_r \in \mathbb{R}^{p_1 \times r}$ ,  $V_r \in \mathbb{R}^{p_2 \times r}$  and  $b_r \in \mathbb{R}_{++}^r$ . Then,*

$$M_r^{(1)} = (W_2 \otimes W_1) \frac{\partial \text{vec}(G(\hat{B}(\lambda)))}{\partial \text{vec}^\top(\hat{B}(\lambda))} = [W_2 V_r \text{Diag}(b_r)^{-1} V_r^\top W_2] \otimes W_1^2 + W_2^2 \otimes [W_1 U_r \text{Diag}(b_r)^{-1} U_r^\top W_1]. \quad (7)$$

*Proof.* Note that  $M_r^{(1)} = (W_2 \otimes W_1) \frac{\partial \text{vec}(G(\hat{B}(\lambda)))}{\partial \text{vec}^\top(\hat{B}(\lambda))}$ . It is obvious that

$$\frac{\partial \text{vec}(G(\hat{B}(\lambda)))}{\partial \text{vec}^\top(\hat{B}(\lambda))} = \frac{\partial \text{vec}(G(\hat{B}(\lambda)))}{\partial \text{vec}^\top(U_r)} \cdot \frac{\partial \text{vec}(U_r)}{\partial \text{vec}^\top(\hat{B}(\lambda))} + \frac{\partial \text{vec}(G(\hat{B}(\lambda)))}{\partial \text{vec}^\top(V_r^\top)} \cdot \frac{\partial \text{vec}(V_r^\top)}{\partial \text{vec}^\top(\hat{B}(\lambda))}. \quad (8)$$

(i). According to the fact that  $i$  in Proposition 2.2, the following results hold.

$$\begin{aligned} \frac{\partial \text{vec}(G(\hat{B}(\lambda)))}{\partial \text{vec}^\top(U_r)} &= \frac{\partial \text{vec}(I_{p_1} U_r V_r^\top)}{\partial \text{vec}^\top(U_r)} = \frac{\partial [(V_r \otimes I_{p_1}) \text{vec}(U_r)]}{\partial \text{vec}^\top(U_r)} = V_r \otimes I_{p_1}. \\ \frac{\partial \text{vec}(G(\hat{B}(\lambda)))}{\partial \text{vec}^\top(V_r^\top)} &= \frac{\partial \text{vec}(U_r V_r^\top I_{p_2})}{\partial \text{vec}^\top(V_r^\top)} = \frac{\partial [(I_{p_2} \otimes U_r) \text{vec}(V_r^\top)]}{\partial \text{vec}^\top(V_r^\top)} = I_{p_2} \otimes U_r. \end{aligned}$$

(ii). According to the fact that  $U_r$  and  $V_r$  have full column rank, there are  $U_r^\top U_r = I_r$  and  $V_r^\top V_r = I_r$ . Combining these facts with  $W_1 \hat{B}(\lambda) W_2 = U_r \text{Diag}(b_r) V_r^\top$ , we have

$$\begin{aligned} U_r &= W_1 \hat{B}(\lambda) W_2 V_r \text{Diag}(b_r)^{-1}, \\ V_r^\top &= \text{Diag}(b_r)^{-1} U_r^\top W_1 \hat{B}(\lambda) W_2, \end{aligned}$$

which leads to

$$\begin{aligned} \text{vec}(U_r) &= [(\text{Diag}(b_r)^{-1} V_r^\top W_2) \otimes W_1] \text{vec}(\hat{B}(\lambda)), \\ \text{vec}(V_r^\top) &= [W_2 \otimes (\text{Diag}(b_r)^{-1} U_r^\top W_1)] \text{vec}(\hat{B}(\lambda)). \end{aligned}$$

Thus,

$$\frac{\partial \text{vec}(U_r)}{\partial \text{vec}^\top(\hat{B}(\lambda))} = [\text{Diag}(b_r)^{-1} V_r^\top W_2] \otimes W_1, \quad \frac{\partial \text{vec}(V_r^\top)}{\partial \text{vec}^\top(\hat{B}(\lambda))} = W_2 \otimes [\text{Diag}(b_r)^{-1} U_r^\top W_1].$$

Combining results in (i) and (ii), we know that

$$M_r^{(1)} = (W_2 \otimes W_1) \frac{\partial \text{vec}(G(\hat{B}(\lambda)))}{\partial \text{vec}^\top(\hat{B}(\lambda))} = [W_2 V_r \text{Diag}(b_r)^{-1} V_r^\top W_2] \otimes W_1^2 + W_2^2 \otimes [W_1 U_r \text{Diag}(b_r)^{-1} U_r^\top W_1].$$

□

It is obvious that  $W_2 V_r \text{Diag}(b_r)^{-1} V_r^\top W_2 \geq 0$ ,  $W_1^2 > 0$ ,  $W_2^2 > 0$  and  $W_1 U_r \text{Diag}(b_r)^{-1} U_r^\top W_1 \geq 0$ . Due to [25, Corollary 2.1], we have  $(W_2 V_r \text{Diag}(b_r)^{-1} V_r^\top W_2) \otimes W_1^2 \geq 0$  and  $W_2^2 \otimes (W_1 U_r \text{Diag}(b_r)^{-1} U_r^\top W_1) \geq 0$ , which leads to the matrix  $M_r^{(1)}$  is positive semidefinite.

Now, we calculate the result of  $M_r^{(2)}$ , which is decided by  $\frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(\hat{B}(\lambda))}$ .

**Lemma 3.2.** *For any  $\lambda \geq 0$ , let  $r$  be the rank of  $W_1 \hat{B}(\lambda) W_2$ . Suppose the singular value decomposition of  $W_1 \hat{B}(\lambda) W_2$  is  $W_1 \hat{B}(\lambda) W_2 = U_r \text{Diag}(b_r) V_r^\top$  with  $U_r \in \mathbb{R}^{p_1 \times r}$ ,  $V_r \in \mathbb{R}^{p_2 \times r}$  and  $b_r \in \mathbb{R}_{++}^r$ . Then,*

$$\begin{aligned} M_r^{(2)} &= - [I_{p_2} \otimes (W_1 (I_{p_1} - U_r U_r^\top) W_1^{-1})] \cdot \frac{\mathcal{X}^\top \mathcal{X}}{n} \cdot [(W_2^{-1} V_r V_r^\top W_2) \otimes I_{p_1}] \\ &\quad - [(W_2 (I_{p_2} - V_r V_r^\top) W_2^{-1}) \otimes I_{p_1}] \cdot \frac{\mathcal{X}^\top \mathcal{X}}{n} \cdot [I_{p_2} \otimes (W_1^{-1} U_r U_r^\top W_1)] \\ &\quad - [(W_2 (I_{p_2} - V_r V_r^\top) W_2^{-1}) \otimes I_{p_1}] \cdot \frac{\mathcal{X}^\top \mathcal{X}}{n} \cdot [(W_2^{-1} V_r V_r^\top W_2) \otimes (W_1^{-1} U_r U_r^\top W_1)] \\ &\quad - \lambda [(E_\lambda^\top W_1^{-1}) \otimes W_1] (I_{p_1}^2 + K_{p_1}^2) \left[ ((W_1 \hat{B}(\lambda) W_2)^+ )^\top W_2 \right] \otimes W_1 \\ &\quad - \lambda [W_2 \otimes (E_\lambda W_2^{-1})] (I_{p_2}^2 + K_{p_2}^2) \left[ W_2 \otimes ((W_1 \hat{B}(\lambda) W_2)^+ W_1) \right], \end{aligned} \quad (9)$$



where  $E_\lambda = \frac{1}{n} \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i$  and  $(W_1 \hat{B}(\lambda) W_2)^+ = V_r \text{Diag}(b_r)^{-1} U_r^\top$ .

*Proof.* Note that  $M_r^{(2)} = \lambda (W_2 \otimes W_1) \frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(\hat{B}(\lambda))}$ . Based on the optimality condition (2), the matrix  $N$  can be expressed as  $N = \frac{1}{n\lambda} W_1^{-1} \left( \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i \right) W_2^{-1} - U_r V_r^\top$ . According to  $U_r^\top N = 0$  and  $N V_r = 0$ , the following results hold.

$$\frac{1}{n\lambda} U_r^\top W_1^{-1} \left( \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i \right) W_2^{-1} = V_r^\top \text{ and } \frac{1}{n\lambda} W_1^{-1} \left( \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i \right) W_2^{-1} V_r = U_r.$$

Replacing these results into  $N$ , then  $N$  can be expressed as

$$\begin{aligned} N &= \frac{1}{n\lambda} W_1^{-1} \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i W_2^{-1} \\ &\quad - \frac{1}{n^2 \lambda^2} W_1^{-1} \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i W_2^{-1} V_r U_r^\top W_1^{-1} \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i W_2^{-1} \\ &= \frac{1}{n\lambda} W_1^{-1} \left( \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i \right) W_2^{-1} \left[ I_{p_2} - \frac{1}{n\lambda} V_r U_r^\top W_1^{-1} \left( \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i \right) W_2^{-1} \right] \\ &= \left[ I_{p_1} - \frac{1}{n\lambda} W_1^{-1} \left( \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i \right) W_2^{-1} V_r U_r^\top \right] \frac{1}{n\lambda} W_1^{-1} \left( \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i \right) W_2^{-1}. \end{aligned}$$

Let  $\tilde{N} = \frac{1}{n\lambda} W_1^{-1} \left( \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i \right) W_2^{-1}$ . Then,

$$\begin{aligned} N &= \frac{1}{n\lambda} W_1^{-1} \left( \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i \right) W_2^{-1} [I_{p_2} - V_r V_r^\top] = \tilde{N} [I_{p_2} - V_r V_r^\top] \\ &= \frac{1}{n\lambda} [I_{p_1} - U_r U_r^\top] W_1^{-1} \left( \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i \right) W_2^{-1} = [I_{p_1} - U_r U_r^\top] \tilde{N}. \end{aligned} \tag{10}$$

To verify this  $N$  satisfies the required conditions, there is only one left part needed to be proved, i.e.,  $\|N\|_2 \leq 1$ . According to the equation (2),

$$\tilde{N} = \frac{1}{n\lambda} W_1^{-1} \left( \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i \right) W_2^{-1} = \partial \|\hat{B}(\lambda)\|_*,$$

which leads to  $\|\tilde{N}\|_2 \leq 1$ . Based on the singular inequalities in [18],

$$\|N\|_2 = \|\tilde{N} [I_{p_2} - V_r V_r^\top]\|_2 \leq \|\tilde{N}\|_2 \cdot \|I_{p_2} - V_r V_r^\top\|_2 \leq 1.$$

So, the matrix  $N$  in (2) can be expressed as in (10).

Now, we calculate the result of  $\frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(\hat{B}(\lambda))}$ . It is sure that

$$\frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(\hat{B}(\lambda))} = \frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(U_r)} \cdot \frac{\partial \text{vec}(U_r)}{\partial \text{vec}^\top(\hat{B}(\lambda))} + \frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(V_r^\top)} \cdot \frac{\partial \text{vec}(V_r^\top)}{\partial \text{vec}^\top(\hat{B}(\lambda))} + \frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(\text{Diag}(b_r))} \cdot \frac{\partial \text{vec}(\text{Diag}(b_r))}{\partial \text{vec}^\top(\hat{B}(\lambda))}.$$

(a). Based on the expression of  $N$ ,  $\frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(U_r)} = \frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(\tilde{N})} \cdot \frac{\partial \text{vec}(\tilde{N})}{\partial \text{vec}^\top(U_r)} + \frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(I_{p_1} - U_r U_r^\top)} \cdot \frac{\partial \text{vec}(I_{p_1} - U_r U_r^\top)}{\partial \text{vec}^\top(U_r)}$ .

$$\frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(\tilde{N})} = \frac{\partial \text{vec}((I_{p_2} \otimes (I_{p_1} - U_r U_r^\top)) \text{vec}(\tilde{N}))}{\partial \text{vec}^\top(\tilde{N})} = I_{p_2} \otimes (I_{p_1} - U_r U_r^\top).$$

Due to the fact that

$$\begin{aligned} \text{vec}(\tilde{N}) &= \frac{1}{n\lambda} (W_2^{-1} \otimes W_1^{-1}) \text{vec} \left( \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i \right) \\ &= \frac{1}{n\lambda} (W_2^{-1} \otimes W_1^{-1}) [\mathcal{X}^\top \mathbf{y} - \mathcal{X}^\top \mathcal{X} \text{vec}(\hat{B}(\lambda))] \\ &= \frac{1}{n\lambda} (W_2^{-1} \otimes W_1^{-1}) [\mathcal{X}^\top \mathbf{y} - \mathcal{X}^\top \mathcal{X} ((W_2^{-1} V_r \text{Diag}(b_r)) \otimes W_1^{-1}) \text{vec}(U_r)], \end{aligned}$$

the result of  $\frac{\partial \text{vec}(\tilde{N})}{\partial \text{vec}^\top(U_r)}$  is

$$\frac{\partial \text{vec}(\tilde{N})}{\partial \text{vec}^\top(U_r)} = -\frac{1}{n\lambda} (W_2^{-1} \otimes W_1^{-1}) \mathcal{X}^\top \mathcal{X} [(W_2^{-1} V_r \text{Diag}(b_r)) \otimes W_1^{-1}].$$

It is easy to get that

$$\frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(I_{p_1} - U_r U_r^\top)} = \frac{\partial \left( (\tilde{N}^\top \otimes I_{p_1}) \text{vec}(I_{p_1} - U_r U_r^\top) \right)}{\partial \text{vec}^\top(I_{p_1} - U_r U_r^\top)} = \tilde{N}^\top \otimes I_{p_1}.$$

The claim *iii* in Proposition 2.2 leads to  $\text{vec}(U_r^\top) = K_{r p_1} \text{vec}(U_r)$  and

$$\begin{aligned} \frac{\partial \text{vec}(I_{p_1} - U_r U_r^\top)}{\partial \text{vec}^\top(U_r)} &= -\frac{\partial \text{vec}(U_r U_r^\top)}{\partial \text{vec}^\top(U_r)} - \frac{\partial \text{vec}(U_r U_r^\top)}{\partial \text{vec}^\top(U_r^\top)} \cdot \frac{\partial \text{vec}(U_r^\top)}{\partial \text{vec}^\top(U_r)} \\ &= -U_r \otimes I_{p_1} - (I_{p_1} \otimes U_r) K_{r p_1} \\ &= -U_r \otimes I_{p_1} - K_{p_1^2} (U_r \otimes I_{p_1}) = -(I_{p_1^2} + K_{p_1^2}) (U_r \otimes I_{p_1}). \end{aligned}$$

Hence,

$$\begin{aligned} \frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(U_r)} &= -\frac{1}{n\lambda} [W_2^{-1} \otimes (I_{p_1} - U_r U_r^\top) W_1^{-1}] \mathcal{X}^\top \mathcal{X} [(W_2^{-1} V_r \text{Diag}(b_r)) \otimes W_1^{-1}] \\ &\quad - (\tilde{N}^\top \otimes I_{p_1}) (I_{p_1^2} + K_{p_1^2}) (U_r \otimes I_{p_1}). \end{aligned}$$

(b). Based on the expression of  $N$ ,  $\frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(V_r^\top)} = \frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(\tilde{N})} \cdot \frac{\partial \text{vec}(\tilde{N})}{\partial \text{vec}^\top(V_r^\top)} + \frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(I_{p_2} - V_r V_r^\top)} \cdot \frac{\partial \text{vec}(I_{p_2} - V_r V_r^\top)}{\partial \text{vec}^\top(V_r^\top)}$ .

$$\frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(\tilde{N})} = \frac{\partial \text{vec}(((I_{p_2} - V_r V_r^\top) \otimes I_{p_1}) \text{vec}(\tilde{N}))}{\partial \text{vec}^\top(\tilde{N})} = (I_{p_2} - V_r V_r^\top) \otimes I_{p_1}.$$

Due to the fact that

$$\begin{aligned} \text{vec}(\tilde{N}) &= \frac{1}{n\lambda} (W_2^{-1} \otimes W_1^{-1}) \text{vec} \left( \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i \right) \\ &= \frac{1}{n\lambda} (W_2^{-1} \otimes W_1^{-1}) [\mathcal{X}^\top \mathbf{y} - \mathcal{X}^\top \mathcal{X} \text{vec}(\hat{B}(\lambda))] \\ &= \frac{1}{n\lambda} (W_2^{-1} \otimes W_1^{-1}) [\mathcal{X}^\top \mathbf{y} - \mathcal{X}^\top \mathcal{X} (W_2^{-1} \otimes (W_1^{-1} U_r \text{Diag}(b_r))) \text{vec}(V_r^\top)], \end{aligned}$$

the result of  $\frac{\partial \text{vec}(\tilde{N})}{\partial \text{vec}^\top(V_r^\top)}$  is

$$\frac{\partial \text{vec}(\tilde{N})}{\partial \text{vec}^\top(V_r^\top)} = -\frac{1}{n\lambda} (W_2^{-1} \otimes W_1^{-1}) \mathcal{X}^\top \mathcal{X} [W_2^{-1} \otimes (W_1^{-1} U_r \text{Diag}(b_r))].$$

It is easy to get that

$$\frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(I_{p_2} - V_r V_r^\top)} = \frac{\partial \text{vec}((I_{p_2} \otimes \tilde{N}) \text{vec}(I_{p_2} - V_r V_r^\top))}{\partial \text{vec}^\top(I_{p_2} - V_r V_r^\top)} = I_{p_2} \otimes \tilde{N}.$$

From the result that  $\text{vec}(V^\top) = K_{rp_2} \text{vec}(V_r)$ , we have

$$\begin{aligned} \frac{\partial \text{vec}(I_{p_2} - V_r V_r^\top)}{\partial \text{vec}^\top(V_r^\top)} &= -\frac{\partial \text{vec}(V_r V_r^\top)}{\partial \text{vec}^\top(V_r^\top)} - \frac{\partial \text{vec}(V_r V_r^\top)}{\partial \text{vec}^\top(V_r)} \cdot \frac{\partial \text{vec}(V_r)}{\partial \text{vec}^\top(V_r^\top)} \\ &= -I_{p_2} \otimes V_r - (V_r \otimes I_{p_1}) K_{rp_2} \\ &= -I_{p_2} \otimes V_r - K_{p_2^2} (I_{p_2} \otimes V_r) = -(I_{p_2^2} + K_{p_2^2}) (I_{p_2} \otimes V_r). \end{aligned}$$

Hence,

$$\begin{aligned} \frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(V_r^\top)} &= -\frac{1}{n\lambda} [((I_{p_2} - V_r V_r^\top) W_2^{-1}) \otimes W_1^{-1}] \mathcal{X}^\top \mathcal{X} [W_2^{-1} \otimes (W_1^{-1} U_r \text{Diag}(b_r))] \\ &\quad - (I_{p_2} \otimes \tilde{N}) (I_{p_2^2} + K_{p_2^2}) (I_{p_2} \otimes V_r). \end{aligned}$$

(c). In the similar way, the result of  $\frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(\text{Diag}(b_r))}$  is showed as follows.

$$\begin{aligned} \frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(\text{Diag}(b_r))} &= \frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(\tilde{N})} \cdot \frac{\partial \text{vec}(\tilde{N})}{\partial \text{vec}^\top(\text{Diag}(b_r))} \\ &= -\frac{1}{n\lambda} [(I_{p_2} - V_r V_r^\top) \otimes I_{p_1}] \cdot (W_2^{-1} \otimes W_1^{-1}) \mathcal{X}^\top \mathcal{X} [(W_2^{-1} V_r) \otimes (W_1^{-1} U_r)] \\ &= -\frac{1}{n\lambda} [((I_{p_2} - V_r V_r^\top) W_2^{-1}) \otimes W_1^{-1}] \mathcal{X}^\top \mathcal{X} [(W_2^{-1} V_r) \otimes (W_1^{-1} U_r)]. \end{aligned}$$

(d). Because  $W_1 \hat{B}(\lambda) W_2 = U_r \text{Diag}(b_r) V_r^\top$ , then  $\text{Diag}(b_r) = U_r^\top W_1 \hat{B}(\lambda) W_2 V_r$  and

$$\frac{\partial \text{vec}(\text{Diag}(b_r))}{\partial \text{vec}^\top(\hat{B}(\lambda))} = (V_r^\top W_2) \otimes (U_r^\top W_1).$$

In addition, results of  $\frac{\partial \text{vec}(U_r)}{\partial \text{vec}^\top(\hat{B}(\lambda))}$  and  $\frac{\partial \text{vec}(V_r^\top)}{\partial \text{vec}^\top(\hat{B}(\lambda))}$  have been computed in Lemma 3.1.

From (a)-(d), we know that

$$\begin{aligned} M_r^{(2)} &= \lambda (W_2 \otimes W_1) \cdot \frac{\partial \text{vec}(N)}{\partial \text{vec}^\top(\hat{B}(\lambda))} \\ &= -[I_{p_2} \otimes (W_1 (I_{p_1} - U_r U_r^\top) W_1^{-1})] \cdot \frac{\mathcal{X}^\top \mathcal{X}}{n} \cdot [(W_2^{-1} V_r V_r^\top W_2) \otimes I_{p_1}] \\ &\quad - [(W_2 (I_{p_2} - V_r V_r^\top) W_2^{-1}) \otimes I_{p_1}] \cdot \frac{\mathcal{X}^\top \mathcal{X}}{n} \cdot [I_{p_2} \otimes (W_1^{-1} U_r U_r^\top W_1)] \\ &\quad - [(W_2 (I_{p_2} - V_r V_r^\top) W_2^{-1}) \otimes I_{p_1}] \cdot \frac{\mathcal{X}^\top \mathcal{X}}{n} \cdot [(W_2^{-1} V_r V_r^\top W_2) \otimes (W_1^{-1} U_r U_r^\top W_1)] \\ &\quad - \lambda [(W_2 \tilde{N}^\top) \otimes W_1] (I_{p_1^2} + K_{p_1^2}) [(U_r \text{Diag}(b_r)^{-1} V_r^\top W_2) \otimes W_1] \\ &\quad - \lambda [W_2 \otimes (W_1 \tilde{N})] (I_{p_2^2} + K_{p_2^2}) [W_2 \otimes (V_r \text{Diag}(b_r)^{-1} U_r^\top W_1)]. \end{aligned}$$

By simple computation, we know that  $(W_1 \hat{B}(\lambda) W_2)^+ = V_r \text{Diag}(b_r)^{-1} U_r^\top$ . Replace the expression of  $\tilde{N}$  and  $E_\lambda = \frac{1}{n} \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle) X_i$  into  $M_r^{(2)}$ . Then, we have

$$\begin{aligned} M_r^{(2)} = & - [I_{p_2} \otimes (W_1 (I_{p_1} - U_r U_r^\top) W_1^{-1})] \cdot \frac{\mathcal{X}^\top \mathcal{X}}{n} \cdot [(W_2^{-1} V_r V_r^\top W_2) \otimes I_{p_1}] \\ & - [(W_2 (I_{p_2} - V_r V_r^\top) W_2^{-1}) \otimes I_{p_1}] \cdot \frac{\mathcal{X}^\top \mathcal{X}}{n} \cdot [I_{p_2} \otimes (W_1^{-1} U_r U_r^\top W_1)] \\ & - [(W_2 (I_{p_2} - V_r V_r^\top) W_2^{-1}) \otimes I_{p_1}] \cdot \frac{\mathcal{X}^\top \mathcal{X}}{n} \cdot [(W_2^{-1} V_r V_r^\top W_2) \otimes (W_1^{-1} U_r U_r^\top W_1)] \\ & - \lambda [(E_\lambda^\top W_1^{-1}) \otimes W_1] (I_{p_1^2} + K_{p_1^2}) \left[ \left( (W_1 \hat{B}(\lambda) W_2)^+ \right)^\top W_2 \right] \otimes W_1 \\ & - \lambda [W_2 \otimes (E_\lambda W_2^{-1})] (I_{p_2^2} + K_{p_2^2}) \left[ W_2 \otimes \left( (W_1 \hat{B}(\lambda) W_2)^+ W_1 \right) \right]. \end{aligned}$$

□

**Remark 3.1.** *There are some facts about the result of this degrees of freedom.*

In (1), if  $\lambda = 0$  and  $B_{1S}$  has full rank, we have  $M_r^{(2)} = 0$  and  $M_r = \frac{1}{n} \sum_{i=1}^n \text{vec}(X_i) \text{vec}(X_i)^\top$ , which leads to

$$\hat{\text{df}}_\lambda = \frac{1}{n} \sum_{k=1}^n \text{vec}(X_k)^\top \left[ \frac{1}{n} \sum_{i=1}^n \text{vec}(X_i) \text{vec}(X_i)^\top \right]^+ \text{vec}(X_k) = \text{rank} \left( \frac{1}{n} \sum_{i=1}^n \text{vec}(X_i) \text{vec}(X_i)^\top \right).$$

When  $\lambda$  is sufficiently large such that  $\hat{B}(\lambda) = 0$ , we have  $M_r^{(1)} = 0$  and  $M_r^{(2)} = 0$ , which also leads to

$$\hat{\text{df}}_\lambda = \text{rank} \left( \frac{1}{n} \sum_{i=1}^n \text{vec}(X_i) \text{vec}(X_i)^\top \right).$$

In Zhou and Li [52], they considered the special case of (1), with the weight matrixes  $W_1 = I_{p_1}$  and  $W_2 = I_{p_2}$ . In [52], they presented the value of degrees of freedom of their model, under the assumption that  $\sum_{i=1}^n \text{vec}(X_i) \text{vec}(X_i)^\top = I_{p_1 p_2}$ . Under this condition and  $\lambda = 0$ , our degrees of freedom is  $\hat{\text{df}}_0 = p_1 p_2$ , which equals to the degrees of freedom in their paper.

For any  $\lambda \geq 0$ , we define a Bayesian information criterion (BIC) as

$$\text{BIC}_\lambda = \log \left( \frac{1}{n} \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle)^2 \right) + \hat{\text{df}}_\lambda \cdot \frac{\log(n)}{n}, \quad (11)$$

where  $\hat{B}(\lambda)$  is the solution of (1) and  $\hat{\text{df}}_\lambda$  is given in Theorem 3.1. Then, the optimal tuning parameter is

$$\lambda^* = \underset{\lambda \geq 0}{\text{argmin}} \text{BIC}_\lambda = \underset{\lambda \geq 0}{\text{argmin}} \left\{ \log \left( \frac{1}{n} \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle)^2 \right) + \hat{\text{df}}_\lambda \cdot \frac{\log(n)}{n} \right\}.$$

To prove the selection result of this new defined BIC, we introduce some technique conditions and new notations as follows.

- (C1)  $\{X_i\}_{i=1}^n$  are sampled i.i.d. from  $X$  and  $\{y_i\}_{i=1}^n$  are sampled i.i.d. from  $y$ . The predictor  $X$  is standardized with  $\|X\|_F = 1$ .  $X$  and  $y$  have finite fourth order moments, i.e.,

$$\max_{j,k} \left\{ \mathbb{E} \left( X_{j,k}^4 \right) \right\} < \infty \text{ and } \mathbb{E}(y^4) < \infty.$$

- (C2)  $\exists B^*$  such that  $B^* \neq 0$ ,  $\text{rank}(B^*) = r^* < \min\{p_1, p_2\}$  and

$$\mathbb{E}(y_i | X_1, X_2, \dots, X_n) = \langle X_i, B^* \rangle, \quad i = 1, 2, \dots, n.$$

$$\text{Var}(y_i | X_1, X_2, \dots, X_n) = \sigma^2, \quad i = 1, 2, \dots, n.$$

- (C3)  $\Sigma = \mathbb{E}(\text{vec}(X) \text{vec}(X)^\top)$  is positive definite, which means there is a  $\kappa > 0$  such that the smallest eigenvalue of  $\Sigma$  is larger than  $\kappa$ .
- (C4)  $X$  and  $\epsilon$  are independent with  $\epsilon \sim \mathcal{N}(0, \sigma^2)$ .

**Remark 3.2.** Condition (C2) states that there exists a low rank matrix  $B^*$  such that the relationship of  $X$  and  $y$  is linear, which is a common assumption and ensures that the true solution has low rank. Conditions (C1) and (C3) are nature sampling assumptions. Condition (C4) guarantees that the random error is independent with the predictor.

Under conditions (C1)-(C4), Bach [3, Theorem 15] proved the consistency and rank consistency of the solution of (1). That is, if  $\lambda_n n^{1/2+\gamma/2} \rightarrow \infty$  and  $\lambda_n n^{1/2} \rightarrow 0$  with  $n \rightarrow \infty$ ,  $\lim_{n \rightarrow \infty} P(\hat{B}(\lambda_n) = B^*) = 1$  and  $\lim_{n \rightarrow \infty} P(r_{\lambda_n} = r^*) = 1$ .

Let  $\Omega_-$  denote the underfitted case, which means  $\Omega_- = \{\lambda : 0 < r_\lambda < r^*\}$ . Let  $\Omega_+$  denote the overfitted case, which means  $\Omega_+ = \{\lambda : r_\lambda > r^*\}$ . Next, we prove the rank selection consistency of  $\text{BIC}_\lambda$ .

**Theorem 3.2.** Assume technical conditions (C1)-(C4). Let  $\lambda_n n^{1/2+\gamma/2} \rightarrow \infty$  and  $\lambda_n n^{1/2} \rightarrow 0$  with  $n \rightarrow \infty$ . For any  $\lambda \in \Omega_+ \cup \Omega_-$ ,

$$\lim_{n \rightarrow \infty} P(\text{BIC}_\lambda > \text{BIC}_{\lambda_n}) = 1,$$

*Proof.* For any  $\lambda \in \Omega_+ \cup \Omega_-$ ,

$$\text{BIC}_\lambda - \text{BIC}_{\lambda_n} = \underbrace{\log \left( \frac{1}{n} \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda) \rangle)^2 \right) - \log \left( \frac{1}{n} \sum_{i=1}^n (y_i - \langle X_i, \hat{B}(\lambda_n) \rangle)^2 \right)}_a + \underbrace{\hat{\text{df}}_\lambda \cdot \frac{\log(n)}{n} - \hat{\text{df}}_{\lambda_n} \cdot \frac{\log(n)}{n}}_b.$$

For a, we know that

$$\begin{aligned} a &= \log \left( \frac{1}{n} \|\mathbf{y} - \mathcal{X} \text{vec}(\hat{B}(\lambda))\|^2 \right) - \log \left( \frac{1}{n} \|\mathbf{y} - \mathcal{X} \text{vec}(\hat{B}(\lambda_n))\|^2 \right) \\ &= \log \left( \frac{\|\mathbf{y} - \mathcal{X} \text{vec}(\hat{B}(\lambda))\|^2 / n}{\|\mathbf{y} - \mathcal{X} \text{vec}(\hat{B}(\lambda_n))\|^2 / n} \right) = \log \left( \frac{\|\mathbf{y} - \mathcal{X} \text{vec}(\hat{B}(\lambda_n)) + \mathcal{X} \text{vec}(\hat{B}(\lambda_n) - \hat{B}(\lambda))\|^2 / n}{\|\mathbf{y} - \mathcal{X} \text{vec}(\hat{B}(\lambda_n))\|^2 / n} \right) \\ &= \log \left( 1 + \underbrace{\frac{\text{vec}(\hat{B}(\lambda_n) - \hat{B}(\lambda))^\top (\mathcal{X}^\top \mathcal{X} / n) \text{vec}(\hat{B}(\lambda_n) - \hat{B}(\lambda))}{\|\mathbf{y} - \mathcal{X} \text{vec}(\hat{B}(\lambda_n))\|^2 / n}}_{a1} + \underbrace{\frac{2(\mathbf{y} - \mathcal{X} \text{vec}(\hat{B}(\lambda_n)))^\top \mathcal{X} \text{vec}(\hat{B}(\lambda_n) - \hat{B}(\lambda)) / n}{\|\mathbf{y} - \mathcal{X} \text{vec}(\hat{B}(\lambda_n))\|^2 / n}}_{a2} \right) \end{aligned}$$

For any  $x$ , we have  $\log(1+x) \geq \min\{\log(2), 0.5x\}$ . So,  $a \geq \min\{\log(2), 0.5(a1+a2)\}$ . Next, we compute these values. Remark 3.2 illustrates that  $P(\hat{B}(\lambda_n) = B^*) \rightarrow 1$  when  $n \rightarrow \infty$ , which leads to

$$\text{vec}(\hat{B}(\lambda_n) - \hat{B}(\lambda))^\top \frac{\mathcal{X}^\top \mathcal{X}}{n} \text{vec}(\hat{B}(\lambda_n) - \hat{B}(\lambda)) \rightarrow \text{vec}(B^* - \hat{B}(\lambda))^\top E(\text{vec}(X)\text{vec}(X)^\top) \text{vec}(B^* - \hat{B}(\lambda)).$$

Based on the fact that  $\mathbf{y} = \mathcal{X} \text{vec}(B^*) + \varepsilon$  and the condition (C4),

$$\frac{\|\mathbf{y} - \mathcal{X} \text{vec}(\hat{B}(\lambda_n))\|^2}{n} = \frac{\|\mathcal{X} \text{vec}(B^* - \hat{B}(\lambda))\|^2}{n} + \frac{2}{n} \varepsilon^\top \mathcal{X} \text{vec}(B^* - \hat{B}(\lambda)) + \frac{\varepsilon^\top \varepsilon}{n} \rightarrow \sigma^2.$$

Combing these facts and (C3), we have  $P(a1 > \kappa \|B^* - \hat{B}(\lambda)\|_F^2 / \sigma^2) \rightarrow 1$  and  $P(a2 = 0) \rightarrow 1$  when  $n \rightarrow \infty$ , which leads to

$$P\left(a \geq \min\left\{\log(2), \frac{\kappa}{2\sigma^2} \|B^* - \hat{B}(\lambda)\|_F^2\right\}\right) \rightarrow 1 \text{ with } n \rightarrow \infty.$$

For  $b$ , we consider the bound of  $\hat{\text{df}}_\lambda$ . By Cauchy-Schwartz inequality, it is sure that

$$\begin{aligned} \hat{\text{df}}_\lambda &= \frac{1}{n} \sum_{k=1}^n \text{vec}(X_k)^\top M_r^+ \text{vec}(X_k) = \left\langle \frac{1}{n} \sum_{k=1}^n \text{vec}(X_k) \text{vec}(X_k)^\top, M_r^+ \right\rangle \\ &\leq \left\| \frac{1}{n} \sum_{k=1}^n \text{vec}(X_k) \text{vec}(X_k)^\top \right\|_F \cdot \|M_r^+\|_2 \leq \frac{1}{n} \sum_{k=1}^n \|X_k\|_F^2 \cdot \|M_r^+\|_2. \end{aligned}$$

When  $n \rightarrow \infty$ ,  $\frac{1}{n} \sum_{k=1}^n \|X_k\|_F^2 \rightarrow E(\|X\|_F^2)$ , which is bounded because of (C1). In addition,  $\|M_r^+\|_2 = \frac{1}{\sigma_r(M_r)}$  with  $\tilde{r}$  being the rank of  $M_r$ , which is bounded when  $n \rightarrow \infty$ . Combining the fact that  $\frac{\log(n)}{n} \rightarrow 0$ , we have  $\log(2) + b > 0$  and  $\frac{\kappa}{2\sigma^2} \|B^* - \hat{B}(\lambda)\|_F^2 + b > 0$  with probability tending to one. Therefore, the result about  $P(\text{BIC}_\lambda - \text{BIC}_{\lambda_n} > 0) \rightarrow 1$  holds, when  $n \rightarrow \infty$ .  $\square$

According to this theorem, the information criterion  $\text{BIC}_\lambda$  will not select the underfitted or overfitted models when the sample size increasing, which means  $\text{BIC}_\lambda$  can select the true model consistently.

## 4 Numerical Experiments

This section will show some numerical results of our proposed Bayesian information criterion (BIC) for tuning parameter selection. To compare our method with the popular tuning parameter selection methods, we extend the corresponding expressions for the model (1) without further proofs, which includes Akaike Information Criterion (AIC), Akaike Information Criterion corrected (AICc) and cross validation. The difference between BIC, AIC and AICc is on the second term of their expressions, which are showed as follows.

AIC is defined as

$$\text{AIC}_\lambda = \log\left(\sum_{i=1}^n \frac{(y_i - \langle X_i, \hat{B}(\lambda) \rangle)^2}{n}\right) + \hat{\text{df}}_\lambda \cdot \frac{2}{n}.$$

AICc is defined as

$$\text{AICc}_\lambda = \log\left(\sum_{i=1}^n \frac{(y_i - \langle X_i, \hat{B}(\lambda) \rangle)^2}{n}\right) + \hat{\text{df}}_\lambda \cdot \frac{2}{n} + \frac{2\hat{\text{df}}_\lambda \cdot (\hat{\text{df}}_\lambda + 1)}{n - \hat{\text{df}}_\lambda - 1}.$$

$K$ -fold cross-validation is a popular method used to evaluate and compare the generalization ability to predict new data. This approach works by splitting the dataset into  $K$  smaller subsets (folds). One of these folds is used as a test set to validate the performance of the model, while the other  $K - 1$  subsets are used as training sets for the model. This process is repeated  $K$  times, and each time selects a different fold as the test set and the rest as the training set. Finally, the  $K$  test results are averaged or otherwise integrated to obtain an overall model performance evaluation. The usual choice of  $K$  is 5 and 10, which are 5-fold cross validation and 10-fold cross validation, respectively.

In the following numerical experiments, we compare our BIC with AIC, AICc, 5-fold cross validation and 10-fold cross validation. The comparison indexes include the selected best tuning parameter  $\lambda^*$ , the mean square error (MSE) on the predictor  $y$ , the rank of the selected solution  $r$  and the computational time. Here, the MSE is defined as  $\|\hat{y} - y\|^2$ , where  $\hat{y}$  is the predicted response variable under the selected tuning parameter. Our numerical experiments are all implemented on MATLAB R2023b with AMD Ryzen 5 5600H with Radeon Graphics 3.30 GHz and 16G RAM.

sample size	method	$\lambda^*$	MSE	r	time (s)
$n = 10^3$	BIC	22.860	7.2380	<b>2</b>	<b>101.69</b>
	AIC	0.0600	0.0808	6	<b>101.69</b>
	AIC <sub>c</sub>	0.0003	150.23	15	<b>101.69</b>
	5-fold CV	0.1361	0.0259	<b>2</b>	178.12
	10-fold CV	0.1361	0.0259	<b>2</b>	672.54
$n = 10^2$	BIC	0.8335	0.0154	<b>4</b>	<b>121.22</b>
	AIC	0.8335	0.0154	<b>4</b>	<b>121.22</b>
	AICc	0.0004	0.0002	15	<b>121.22</b>
	5-fold CV	0.0006	$1.22 \times 10^{-8}$	14	344.23
	10-fold CV	0.0006	$1.22 \times 10^{-8}$	14	5112.1
$n = 50$	BIC	6.8011	0.0154	<b>4</b>	<b>110.38</b>
	AIC	6.8011	0.0154	<b>4</b>	<b>110.38</b>
	AICc	0.0906	0.0002	<b>4</b>	<b>110.38</b>
	5-fold CV	0.0026	$1.22 \times 10^{-8}$	10	344.38
	10-fold CV	0.0026	$1.22 \times 10^{-8}$	10	484.55

Table 1: Simulation results on prediction matrix dimension as  $p_1 = 15$  and  $p_2 = 45$ . The sample size includes  $10^3$ ,  $10^2$  and 50, respectively.

## 4.1 Simulation Data

As in Bach [3], we simulate a lots of data sets with different values  $p_1$ ,  $p_2$  and  $n$ , where the prediction matrixes follow the standard normal distribution/Gaussian distribution. Here, we select  $p_1 \in \{15, 25, 35\}$ ,  $p_2 \in \{30, 45\}$  and  $n \in \{10^3, 10^2, 50\}$ . We generate random i.i.d. data matrixes  $P_i \in \mathbb{R}^{p_1}$  and  $Q_i \in \mathbb{R}^{p_2}$ , and we set a true solution  $B^* \in \mathbb{R}^{p \times q}$  with rank 2. All elements of these matrix distribute with the standard normal distribution. Then,  $y_i = \langle P_i Q_i^\top, B^* \rangle + \epsilon_i, i = 1, 2, \dots, n$ , where  $\epsilon_i$  have i.i.d components with normal distributions with zero mean and 0.1 standard variance. To select the best tuning parameter, same as the setting in Bach [3], we set the tuning parameter sequence as  $\lambda_k = e^{\log(\lambda_{\max}) + (k-1) \times \frac{\log(\lambda_{\min}) - \log(\lambda_{\max})}{100}}$  with  $k = 1, 2, \dots, 100$ . The numerical results are reported in Tables 1 to 3.

sample size	method	$\lambda^*$	MSE	r	time (s)
$n = 10^3$	BIC	0.1092	0.1210	<b>3</b>	<b>56.763</b>
	AIC	0.0218	0.0487	7	<b>56.763</b>
	AICc	0.0489	0.0569	4	<b>56.763</b>
	5-fold CV	0.0218	0.0487	7	128.60
	10-fold CV	0.0218	0.0487	7	266.08
$n = 10^2$	BIC	0.0176	0.0001	14	<b>89.548</b>
	AIC	0.0176	0.0001	14	<b>89.548</b>
	AICc	0.0596	0.0003	<b>10</b>	<b>89.548</b>
	5-fold CV	0.0176	0.0001	14	301.95
	10-fold CV	0.0176	0.0001	14	802.78
$n = 50$	BIC	0.6608	0.0629	<b>6</b>	<b>112.48</b>
	AIC	0.6608	0.0629	<b>6</b>	<b>112.48</b>
	AICc	0.0277	0.3611	12	<b>112.48</b>
	5-fold CV	0.0910	0.0292	7	259.31
	10-fold CV	0.0612	0.0279	9	498.99

Table 2: Simulation results on prediction matrix dimension as  $p_1 = 25$  and  $p_2 = 30$ . The sample size includes  $10^3$ ,  $10^2$  and 50, respectively.

Based on these results in Tables 1 to 3, there are some conclusions. (i) Comparing to the other tuning parameter methods, our proposed BIC will select the best tuning parameter with the smallest solution rank. Because the true rank of the solution is 2, BIC will tend to select the best tuning parameter with the same true solution rank. (ii) The computational time of BIC is same with AIC and AICc, and obviously smaller than that of 5-fold CV and 10-fold CV. Because BIC, AIC and AICc are different only on the second term, their computational time are same. Comparing to cross validation, our BIC surely reduce the computational time, because that our method does not need to solve the model many times as cross validation. (iii) BIC generally performs same with the other methods on MSE and the MSE under BIC is larger than cross validation in some special cases, which is a trade off of the computational effect and



sample size	method	$\lambda^*$	MSE	r	time (s)
$n = 10^3$	BIC	0.0089	0.0134	<b>13</b>	<b>190.15</b>
	AIC	0.0026	0.0050	22	<b>190.15</b>
	AICc	0.0059	0.0088	16	<b>190.15</b>
	5-fold CV	0.0007	0.0025	30	566.70
	10-fold CV	0.0007	0.0025	30	1113.4
$n = 10^2$	BIC	0.4744	0.2561	<b>10</b>	<b>178.11</b>
	AIC	0.4744	0.2561	<b>10</b>	<b>178.11</b>
	AICc	0.3883	0.3430	<b>10</b>	<b>178.11</b>
	5-fold CV	0.2602	0.1944	11	655.69
	10-fold CV	0.2602	0.1944	11	1125.49
$n = 50$	BIC	1.2896	66.480	<b>7</b>	<b>158.60</b>
	AIC	1.2896	66.480	<b>7</b>	<b>158.60</b>
	AICc	0.0009	158.11	25	<b>158.60</b>
	5-fold CV	1.0588	58.339	<b>7</b>	448.32
	10-fold CV	1.0588	58.338	<b>7</b>	1397.4

Table 3: Simulation results on prediction matrix dimension as  $p_1 = 35$  and  $p_2 = 30$ . The sample size includes  $10^3$ ,  $10^2$  and 50, respectively.

computational cost.

## 4.2 Real Data

This section applies the tuning parameter selection process of the model (1) on the COVID-19 data set and Bike data set, and compares BIC with AIC, AICc, 5-fold CV and 10-fold CV in Table 4 and Table 5. For every data set, we standardize the data set by columns, i.e., every number is subtracted by the column mean and divided by the column standard deviation. To select the best tuning parameter, we set the tuning parameter sequence as  $\lambda_k = 0.618^k \lambda_{\max}$  with  $k = 1, 2, \dots, 20$  and  $\lambda_{\max} = \left\| W_1^\top \cdots \left( \sum_{i=1}^n y_i X_i \right) W_2^\top \right\|_2$ . Here,  $\lambda_{\max}$  is the smallest tuning parameter such that the solution of (1) is zero.

The COVID-19 dataset ([21, 37]) consists of daily measurements related to COVID-19 for 138 countries around the world. This data set records the newly confirmed case in the period June 13, 2020 to July 12, 2020. In addition, this data also includes the 41 COVID-19 related government policies in each day, i.e., school-closing, restrictions on gathering, stay-at-home requirement, income support and so on. Each of these policies may have several levels, for example, school closing includes no closing, recommend closing, require some closing (e.g. just high school) or require all closing, which varied during the 30-day period. Therefore, for every sample, its prediction matrix is  $X \in \mathbb{R}^{41 \times 30}$  and response is the newly confirmed case. The sample size is  $n = 138$ .

Method	$\lambda^*$	MSE	rank	time
BIC	0.1346	0.0231	<b>1</b>	<b>19.088</b>
AIC	0.1346	0.0231	<b>1</b>	<b>19.088</b>
AICc	0.1346	0.0231	<b>1</b>	<b>19.088</b>
5-fold CV	0.1346	0.0231	<b>1</b>	37.855
10-fold CV	0.1346	0.0231	<b>1</b>	74.117

Table 4: COVID data set results with  $p_1 = 41$ ,  $p_2 = 30$  and  $n = 138$ .

The Bike sharing data set [13] includes matrixes data from January 1st, 2011 to December 31th, 2012. For every data during this period, the recorded data includes 6 weather conditions (including sunny, mist or others, temperature, apparent temperature, humidity and wind speed) every hour, which makes the prediction matrix dimension as  $X \in \mathbb{R}^{24 \times 6}$  and the sample size  $n = 731$ . The response variable is the daily aggregated count of rented bikes.

Method	$\lambda^*$	MSE	rank	time
BIC	1.5074	0.5173	<b>1</b>	<b>1.6880</b>
AIC	0.3557	0.4911	2	<b>1.6880</b>
AICc	0.0011	0.4029	6	<b>1.6880</b>
5-fold CV	0.0011	0.4029	6	7.1090
10-fold CV	0.0011	0.4029	6	9.2680

Table 5: Bike sharing data set results with  $p_1 = 24$ ,  $p_2 = 6$  and  $n = 731$ .

On these real data sets, our proposed BIC performs better than AIC and AICc because that the rank under  $\lambda^*$  with BIC is the smallest and their MSE values are similar. In addition, BIC outperforms 5-fold CV and 10-fold CV on computational time.

## 5 Conclusion

To select the best tuning parameter of the adaptive nuclear norm regularized trace regression model, we propose a Bayesian information criterion (BIC) in this paper. We first compute the unbiased estimator of the degrees of the freedom of the model, which is the basic element of the BIC. Under this estimator, we build up the BIC and prove its rank selection consistency, i.e., BIC will select the tuning parameter achieving the true solution and true solution rank in probability with the sample size increasing. To evaluate the performance of BIC, we compare it with AIC, AICc, 5-fold CV and 10-fold CV on some simulation data and real data sets. The numerical results show that BIC performs better than the other methods.

## Acknowledgements

This work was supported by the National Natural Science Foundation of China (12371322).

## References

- [1] Antonino Abbruzzo, Ivan Vujačić, Angelo M Mineo, and Ernst C Wit. Selecting the tuning parameter in penalized gaussian graphical models. *Statistics and Computing*, 29(3):559–569, 2019.
- [2] Hirotugu Akaike. Statistical predictor identification. *Annals of the Institute of Statistical Mathematics*, 22(1):203–217, 1970.
- [3] Francis R Bach. Consistency of trace norm minimization. *Journal of Machine Learning Research*, 9:1019–1048, 2008.
- [4] Amir Beck. *First-Order Methods in Optimization*. SIAM, 2017.
- [5] Denis Chetverikov, Zhipeng Liao, and Victor Chernozhukov. On cross-validated lasso in high dimensions. *The Annals of Statistics*, 49(3):1300–1317, 2021.
- [6] Abhirup Datta and Hui Zou. A note on cross-validation for lasso under measurement errors. *Technometrics*, 62(4):549–556, 2020.
- [7] Mark A Davenport and Justin Romberg. An overview of low-rank matrix recovery from incomplete observations. *IEEE Journal of Selected Topics in Signal Processing*, 10(4):608–622, 2016.
- [8] Bradley Efron. The estimation of prediction error: covariance penalties and cross-validation. *Journal of the American Statistical Association*, 99(467):619–632, 2004.
- [9] Andreas Elsener and Sara van de Geer. Robust low-rank matrix estimation. *The Annals of Statistics*, 46(6):3481–3509, 2018.
- [10] Jianqing Fan, Wenyan Gong, and Ziwei Zhu. Generalized high-dimensional trace regression via nuclear norm regularization. *Journal of Econometrics*, 212(1):177–202, 2019.
- [11] Jianqing Fan and Runze Li. Variable selection via nonconcave penalized likelihood and its oracle properties. *Journal of the American Statistical Association*, 96(456):1348–1360, 2001.
- [12] Yingying Fan and Cheng Yong Tang. Tuning parameter selection in high dimensional penalized likelihood. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 75(3):531–552, 2013.
- [13] Hadi Fanaee-T and Joao Gama. Event labeling combining ensemble detectors and background knowledge. *Progress in Artificial Intelligence*, 2:113–127, 2014.

- [14] Nima Hamidi and Mohsen Bayati. On low-rank trace regression under general sampling distribution. *Journal of Machine Learning Research*, 23(321):1–49, 2022.
- [15] Kei Hirose, Shohei Tateishi, and Sadanori Konishi. Tuning parameter selection in sparse regression modeling. *Computational Statistics and Data Analysis*, 59:28–40, 2013.
- [16] Darren Homrighausen and Daniel J McDonald. The lasso, persistence, and cross-validation. In *the International Conference on Machine Learning*, pages 1031–1039. PMLR, 2013.
- [17] Darren Homrighausen and Daniel J McDonald. Risk consistency of cross-validation with lasso-type procedures. *Statistica Sinica*, 27:1017–1036, 2017.
- [18] Roger A Horn and Charles R Johnson. *Matrix analysis*. Cambridge university press, 2012.
- [19] Yongdai Kim, Sunghoon Kwon, and Hosik Choi. Consistent model selection criteria on high dimensions. *Journal of Machine Learning Research*, 13(1):1037–1057, 2012.
- [20] Jing Lei. Cross-validation with confidence. *Journal of the American Statistical Association*, 115(532):1978–1997, 2020.
- [21] Mei Li, Lingchen Kong, and Zhihua Su. Double fused lasso regularized regression with both matrix and vector valued predictors. *Electronic Journal of Statistics*, 15:1909–1950, 2021.
- [22] Han Liu, Lie Wang, and Tuo Zhao. Multivariate regression with calibration. In *Advances in Neural Information Processing Systems*, pages 127–135, 2014.
- [23] Zhang Liu and Lieven Vandenberghe. Interior-point method for nuclear norm approximation with application to system identification. *SIAM Journal on Matrix Analysis and Applications*, 31(3):1235–1256, 2010.
- [24] Zhaosong Lu, Renato DC Monteiro, and Ming Yuan. Convex optimization methods for dimension reduction and coefficient estimation in multivariate linear regression. *Mathematical Programming*, 131(1):163–194, 2012.
- [25] Jan R Magnus and Heinz Neudecker. *Matrix differential calculus with applications in statistics and econometrics*. John Wiley & Sons, 2019.
- [26] Rahul Mazumder, Trevor Hastie, and Robert Tibshirani. Spectral regularization algorithms for learning large incomplete matrices. *Journal of Machine Learning Research*, 11(11):2287–2322, 2010.
- [27] Sahand Negahban and Martin J Wainwright. Estimation of (near) low-rank matrices with noise and high-dimensional scaling. *The Annals of Statistics*, 39(2):1069–1097, 2011.
- [28] Rassoul Noorossana, M Eyvazian, Amirhossein Amiri, and Mahmoud A Mahmoud. Statistical monitoring of multivariate multiple linear regression profiles in phase i with calibration application. *Quality and Reliability Engineering International*, 26(3):291–303, 2010.

- [29] Ralph Tyrell Rockafellar. *Convex Analysis*. Princeton University Press, 2015.
- [30] Adam J Rothman, Elizaveta Levina, and Ji Zhu. Sparse multivariate regression with covariance estimation. *Journal of Computational and Graphical Statistics*, 19(4):947–962, 2010.
- [31] Gideon Schwarz. Estimating the dimension of a model. *The Annals of Statistics*, 6(2):461–464, 1978.
- [32] Charles M Stein. Estimation of the mean of a multivariate normal distribution. *The Annals of Statistics*, pages 1135–1151, 1981.
- [33] Wei Sun, Junhui Wang, and Yixin Fang. Consistent selection of tuning parameters via variable selection stability. *Journal of Machine Learning Research*, 14(1):3419–3440, 2013.
- [34] Robert Tibshirani. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society: Series B (Methodological)*, 58(1):267–288, 1996.
- [35] Robert Tibshirani, Michael Saunders, Saharon Rosset, Ji Zhu, and Keith Knight. Sparsity and smoothness via the fused lasso [j]. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 67(1):91–108, 2005.
- [36] Koltchinskii Vladimir, Lounici Karim, and Alexandre B. Tsybakov. Nuclear-norm penalization and optimal rates for noisy low-rank matrix completion. *The Annals of Statistics*, 39(5):2302–2329, 2011.
- [37] O Wahltinez, K Murphy, M Brenner, et al. Covid-19 open-data: curating a fine-grained, global-scale data repository for sars-cov-2. 2020. *Work in progress*, 2020.
- [38] Hansheng Wang and Chenlei Leng. Unified lasso estimation by least squares approximation. *Journal of the American Statistical Association*, 102(479):1039–1048, 2007.
- [39] Hansheng Wang, Bo Li, and Chenlei Leng. Shrinkage tuning parameter selection with a diverging number of parameters. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 71(3):671–683, 2009.
- [40] Hansheng Wang, Runze Li, and Chih-Ling Tsai. Tuning parameter selectors for the smoothly clipped absolute deviation method. *Biometrika*, 94(3):553–568, 2007.
- [41] Tao Wang and Lixing Zhu. Consistent tuning parameter selection in high dimensional sparse linear regression. *Journal of Multivariate Analysis*, 102(7):1141–1151, 2011.
- [42] G Alistair Watson. Characterization of the subdifferential of some matrix norms. *Linear Algebra and its Applications*, 170(0):33–45, 1992.
- [43] Zhenyu Wei and Thomas CM Lee. High-dimensional multi-task learning using multivariate regression and generalized fiducial inference. *Journal of Computational and Graphical Statistics*, 31:1–15, 2022.

- [44] Yunan Wu and Lan Wang. A survey of tuning parameter selection for high-dimensional regression. *Annual Review of Statistics and Its Application*, 7:209–226, 2020.
- [45] Li Yaguang, Wu Yaohua, and Jin Baisuo. Consistent tuning parameter selection in high-dimensional group-penalized regression. *Science China Mathematics*, 62(4):751–770, 2019.
- [46] Ming Yuan. Degrees of freedom in low rank matrix estimation. *Science China Mathematics*, 59(12):2485–2502, 2016.
- [47] Ming Yuan, Ali Ekici, Zhaosong Lu, and Renato Monteiro. Dimension reduction and coefficient estimation in multivariate linear regression. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 69(3):329–346, 2007.
- [48] Ming Yuan and Yi Lin. Model selection and estimation in regression with grouped variables. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 68(1):49–67, 2006.
- [49] Cun-Hui Zhang. Nearly unbiased variable selection under minimax concave penalty. *The Annals of Statistics*, 38(2):894–942, 2010.
- [50] Bo Zhao, Justin P Haldar, Cornelius Brinegar, and Zhi-Pei Liang. Low rank matrix recovery for real-time cardiac mri. In *2010 IEEE International Symposium on Biomedical Imaging: from Nano to Macro*, pages 996–999. IEEE, 2010.
- [51] Junlong Zhao, Lu Niu, and Shushi Zhan. Trace regression model with simultaneously low rank and row (column) sparse parameter. *Computational Statistics and Data Analysis*, 116:1–18, 2017.
- [52] Hua Zhou and Lexin Li. Regularized matrix regression. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 76(2):463–483, 2014.
- [53] Yunzhang Zhu. A convex optimization formulation for multivariate regression. In *Advances in Neural Information Processing Systems*, pages 17652–17661, 2020.
- [54] Changliang Zou, Yuan Ke, and Wenyang Zhang. Estimation of low rank high-dimensional multivariate linear models for multi-response data. *Journal of the American Statistical Association*, 117(538):693–703, 2022.
- [55] Hui Zou. The adaptive lasso and its oracle properties. *Journal of the American Statistical Association*, 101(476):1418–1429, 2006.
- [56] Hui Zou and Trevor Hastie. Regularization and variable selection via the elastic net. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 67(2):301–320, 2005.
- [57] Hui Zou and Hao Helen Zhang. On the adaptive elastic-net with a diverging number of parameters. *The Annals of Statistics*, 37(4):1733–1751, 2009.