

REMARKS ON b -ENRICHED NONEXPANSIVE MAPPINGS

RIZWAN ANJUM* AND MUJAHID ABBAS

ABSTRACT. In this note, we analyzed the concept of enriched nonexpansive which was proposed in “Approximating fixed points of enriched nonexpansive mappings by Krasnoselskij iteration in Hilbert spaces” (Carpathian J. Math., 35(2019), No. 3, 293-304.) Through our analysis, we conclude that the idea of enriched nonexpansive needs reconsideration, as it coincides with well known concept of nonexpansive. Our findings provide an insights into the existing literature and highlight the need for further investigations and clarifications in the existing literature on a metric-fixed point theory.

1. INTRODUCTION

During the 1960s, there was a notable upsurge in the investigation of nonexpansive mappings, largely propelled by two significant factors. One catalyst was Browder’s exploration of the interplay between monotone operators and nonexpansive mappings [32, 33, 34, 35, 36, 37], which sparked considerable interest in the field. Additionally, Kirk’s seminal paper [24] investigated the crucial role of the geometric properties of the norm in establishing the existence of fixed points for nonexpansive mappings.

Let C be a nonempty closed and convex subset in a real Hilbert space H equipped with a norm $\|\cdot\|$ derived from an inner product $\langle \cdot, \cdot \rangle$. A mapping $T : C \rightarrow C$ is called nonexpansive if it satisfies the contractive condition given by:

$$(1.1) \quad \|Tx - Ty\| \leq \|x - y\|, \quad x, y \in C.$$

The study of nonexpansive mappings is a significant and actively researched topic in nonlinear analysis. The reader interested in fixed point theory of nonexpansive mappings is referred to [25, 26, 27, 28, 29, 30, 31] and references mentioned therein.

Next, we state the Browder-Goede-Kirk fixed point theorem for nonexpansive mappings, originally presented as Theorem 4 in [37] and reproduced as Theorem 3.1 in [25].

Theorem 1. *Let C be a closed bounded convex subset of the Hilbert space H and $T : C \rightarrow C$ a nonexpansive mapping. Then T has at least one fixed point.*

Berinde in [9] introduced and examined a new class of mappings known as the class of enriched nonexpansive mappings, which broadens the scope of nonexpansive mappings.

2000 *Mathematics Subject Classification.* 47H09. 47H10 .

Key words and phrases. fixed point, fixed point theorem, b -enriched nonexpansive
Corresponding author: Rizwan Anjum (rizwananjum1723@gmail.com) .

Definition 1. [9] Let $(X, \|\cdot\|)$ be a normed space. A mapping $T : X \rightarrow X$ is called an enriched nonexpansive or b -enriched nonexpansive if there exists $b \in [0, +\infty)$ such that the following condition holds:

$$(1.2) \quad \|b(x - y) + Tx - Ty\| \leq (b + 1) \|x - y\|, \quad \forall x, y \in X.$$

It was stated in [9] that the class of enriched nonexpansive mappings is larger than the class of nonexpansive mappings. Indeed, if we take $b = 0$ in (1.2), then we obtain (1.1).

In [9], it was demonstrated that every enriched nonexpansive mapping defined on a Hilbert space possesses a fixed point that can be approximated using the Krasnoselskij iterative scheme employing the condition of the additional property of demicompactness of the enriched nonexpansive mapping T . Now, we state the main result in [9].

Theorem 2. [9] Let C be a closed bounded convex subset of the Hilbert space H and $T : C \rightarrow C$ a b -enriched nonexpansive mapping. Then T has at least one fixed point.

By conducting a search on MathSciNet, it is observed that there are 105 papers indexed with the term "enriched nonexpansive" in their titles (see [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 38]). Additionally, there are 2,350 papers that mentioned the term "enriched nonexpansive" anywhere in the text, as of March 16, 2024.

The main objective of this paper is to demonstrate that the class of enriched nonexpansive mappings does not contribute much, as it does not go away from the class of nonexpansive mappings. Therefore, as a conclusion, we suggest the researchers in the field of metric fixed point theory to reconsider Theorem 2 for additional exploration. Our results investigate significant review of the current literature and the need to reconsider the continued investigation in this direction.

REMARKS ON b -ENRICHED NONEXPANSIVE MAPPINGS

In this section, we prove that the fixed point of enriched nonexpansive mappings can be obtained by using the fixed point theory of nonexpansive mappings.

The subsequent remark provides the justification of the above comment.

Remark 1. Any b -enriched nonexpansive (1.2) can be simplified as follows:

$$\begin{aligned} & \|b(x - y) + Tx - Ty\| \leq (b + 1) \|x - y\| \\ & \frac{1}{b + 1} \|b(x - y) + Tx - Ty\| \leq \|x - y\| \\ & \left\| \frac{b(x - y) + Tx - Ty}{b + 1} \right\| \leq \|x - y\| \\ & \left\| \frac{bx + Tx}{b + 1} - \frac{by + Ty}{b + 1} \right\| \leq \|x - y\|. \end{aligned}$$

Therefore, the b -enriched nonexpansive condition (1.2) reduces to the condition:

$$(1.3) \quad \|Sx - Sy\| \leq \|x - y\|, \quad \text{for all } x, y \in X$$

where $S : X \rightarrow X$ is defined as:

$$(1.4) \quad Sx = \frac{bx + Tx}{b+1}, \quad \forall x \in X.$$

It follows from (1.3) that the mapping S defined in (1.4) is a nonexpansive mapping.

In conclusion, the b -enriched nonexpansive condition (1.2) can be rephrased to a nonexpansive mapping contractive condition. We refer readers to the proof of Theorem 2, where it was established that every b -enriched nonexpansive mapping possesses at least one fixed point (see, Theorem 2.2 of [9]). Through Remark 1, we show that the set of fixed points of S is equal to the set of fixed point of T .

Remark 2. From Remark 1, it can be observed that every b -enriched nonexpansive mapping T can be transformed into a nonexpansive mapping S as defined in (1.4). By applying Theorem 1, we know that S possesses at least one fixed point, denote it by $x^* \in C$. This implies:

$$\begin{aligned} Sx^* &= \frac{bx^* + Tx^*}{b+1} = x^* \\ bx^* + Tx^* &= (b+1)x^* \\ bx^* + Tx^* &= bx^* + x^* \\ Tx^* &= x^*. \end{aligned}$$

Therefore, x^* is indeed the fixed point of T , thereby proving Theorem 2.

MODIFICATION OF b -ENRICHED NONEXPANSIVE MAPPINGS

In this section, we will give the modified definition of b -enriched nonexpansive mappings and demonstrate fixed point theorems for such mappings. We now present the following concept.

Definition 2. Let $(X, \|\cdot\|)$ be a normed space. A mapping $T : X \rightarrow X$ is called an modified enriched nonexpansive or b -modified enriched nonexpansive if there exists $b \in [0, +\infty)$ such that the following condition holds:

$$(1.5) \quad \|b(x-y) + Tx - Ty\| \leq \|x-y\|, \quad \forall x, y \in X.$$

The remark below illustrates that the definition of b -modified enriched nonexpansive, as in 2, aligns accurately with the definition of enriched nonexpansive as specified in the Definition 1.

Remark 3. Any b -modified enriched nonexpansive (1.5) can be simplified as follows:

$$(1.6) \quad \|Sx - Sy\| \leq \|x - y\|, \quad \text{for all } x, y \in X$$

where $S : X \rightarrow X$ is defined as:

$$(1.7) \quad Sx = bx + Tx, \quad \forall x \in X.$$

We are interested in the conditions under which a mapping T satisfying (1.5) has at the least one fixed point. It has been observed that the contractive condition satisfied by b -modified enriched nonexpansive can be reformulated as shown in (1.6) to obtain a nonexpansive mappings S (1.7). The Browder-Goede-Kirk fixed-point theorem 1 guarantees the existence of at least one fixed point of a nonexpansive mapping under certain conditions. Note that, if $x^* \in X$ is the fixed point of S , then

$x^*(1-b) = Tx^*$ implies that x^* is not the fixed point of T . This was not the case with the definition of enriched nonexpansive mappings as specified in Definition 1 and mentioned in Remarks 1 and 2.

Clearly, the class of b -modified enriched nonexpansive mappings includes the class of enriched nonexpansive mappings; that is, every mapping satisfying (1.1) reduces to (1.5) for $b = 0$.

Now, we present the examples of mapping which are b -modified enriched nonexpansive mappings but not a nonexpansive mappings.

Example 1. Let $X = \mathbb{R}$ be the set of real numbers equipped with the usual norm, and $T : X \rightarrow X$ be defined by $Tx = 100 - 2x$ for all $x \in \mathbb{R}$. Obviously, T is a 3-modified enriched nonexpansive mapping. Indeed, for all $x, y \in \mathbb{R}$, we have:

$$\begin{aligned} \|3(x-y) + Tx - Ty\| &= \|3(x-y) + (100 - 2x) - (100 - 2y)\| \\ &= \|3x - 3y + 100 - 2x - 100 + 2y\| \\ &= \|x - y\|. \end{aligned}$$

Example 2. Let $X = \mathbb{R}$, and $b \in (0, +\infty)$ is a fixed real number. Let $T_b : X \rightarrow X$ be defined by $T_b x = (1-b)x$ for all $x \in \mathbb{R}$. Then T_b is a b -modified enriched nonexpansive mapping. Indeed, for all $x, y \in \mathbb{R}$, we have:

$$\begin{aligned} \|b(x-y) + T_b x - T_b y\| &= \|b(x-y) + (1-b)x - (1-b)y\| \\ &= \|bx - by + x - bx - y + by\| \\ &= \|x - y\|. \end{aligned}$$

Our first result shows that in order to prove the existence of fixed points of b -modified enriched nonexpansive mappings, we do not require the rich geometric structure of a nonempty closed and convex subset in a real Hilbert space, but this only holds when $b \neq 0$. Furthermore, a unique fixed point exists. We establish these results within the framework of a Banach space.

Before we present our main theorem, we need the following concept from [7].

Remark 4. [7] If T is a self mapping on a normed space space X , then for any $\lambda \in (0, 1)$, an averaged mapping T_λ given by

$$(1.8) \quad T_\lambda x = (1-\lambda)x + \lambda Tx, \quad \forall x \in X.$$

Moreover,

$$(1.9) \quad \text{Fix}(T_\lambda) = \{x \in X : T_\lambda x = x\} = \{x \in X : Tx = x\} = \text{Fix}(T).$$

We start with following result.

Theorem 3. Let $(X, \|\cdot\|)$ be Banach space and $T : X \rightarrow X$ a b -modified enriched nonexpansive mapping with $b \neq 0$. Then

- (1) $\text{Fix}(T) = \{x^*\}$;
- (2) there exists $\lambda \in (0, 1)$ such that the Krasnoselskii iteration associated to T , that is, the sequence $\{x_n\}_{n=0}^\infty$, given by

$$(1.10) \quad x_{n+1} = (1-\lambda)x_n + \lambda Tx_n, \quad n \geq 0,$$

converges to x^* for any initial guess $x_0 \in X$.

Proof. Let us denote $\lambda = \frac{1}{b+1}$. Obviously, $\lambda \in (0, 1)$ and the b -modified enriched nonexpansive (1.5) becomes

$$\|(1 - \lambda)x + Tx - ((1 - \lambda)y + Ty)\| \leq \lambda \|x - y\|, \quad \forall x, y \in X,$$

which can be written in an equivalent form as

$$(1.11) \quad \|T_\lambda x - T_\lambda y\| \leq \lambda \|x - y\|, \quad \forall x, y \in X,$$

where T_λ is the averaged mapping defined in (1.8). Since $\lambda \in (0, 1)$, (1.11) gives that T_λ is a Banach contraction mapping.

In view of (1.8), the Krasnoselskij iterative process $\{x_n\}_{n=0}^\infty$, defined by (1.10) is exactly the Picard iteration associated with T_λ , that is,

$$(1.12) \quad x_{n+1} = T_\lambda x_n, n \geq 0.$$

Since X is a Banach space and T_λ is a Banach contraction mapping on X , it follows from the Banach fixed-point theorem (Theorem 2.1 in [25]) in the setting of Banach spaces that T_λ has a unique fixed point, denoted as $x^* \in X$, and the Picard iteration associated with T_λ as defined in (1.12) converges to x^* .

Also, noting from (1.9), we observe that T has a unique fixed point since T_λ does. This completes the proof. \square

From Theorem 3, we do not derive Theorem 1 as a direct consequence. This is because Theorem 3 is established under the assumption that $b \neq 0$. The rationale behind imposing the condition $b \neq 0$ in Theorem 3 is to establish a minimal requirement on the structure of X to ensure that every b -modified enriched nonexpansive mapping T defined on X possesses at least one fixed point. This condition is met when X is a Banach space. To extend this conclusion to all $b \in [0, +\infty)$, we present our subsequent theorem within the framework of Hilbert spaces.

Theorem 4. *Let C be a closed bounded convex subset of the Hilbert space H and $T : C \rightarrow C$ a b -modified enriched nonexpansive mapping. Then T has at least one fixed point.*

Proof. We divide the proof into the following two cases.

Case 1. $b = 0$. In this case, the b -modified enriched nonexpansive mapping (1.5) becomes

$$(1.13) \quad \|Tx - Ty\| \leq \|x - y\|, \quad \forall x, y \in C,$$

That is, T is an nonexpansive mapping and hence by Theorem 1, T has at least one fixed point.

Case 2. If $b > 0$. Then $\lambda = \frac{1}{b+1} \in (0, 1)$. Following arguments similar to those given in the proof of Theorem 3, we establish that T_λ is a Banach contraction on C . Furthermore, since C is a closed subset of a Hilbert space, it is complete. The result then follows by the Banach fixed-point theorem. This implies that T_λ possesses a unique fixed point, as indicated by (1.9). Therefore, T also possesses a unique fixed point. \square

Remark 5. *We can derive Theorem 1 by setting $b = 0$ in Theorem 4. This results in obtaining 1 as a corollary of our main result.*

COMPETING INTERESTS

The authors declare that they have no competing interests.

REFERENCES

- [1] Panja S., Roy, K., Saha, M.: Wardowski type enriched contractive mappings with their fixed points. *J Anal* (2023). <https://doi.org/10.1007/s41478-023-00626-y>
- [2] Anjum R, Abbas M, Iqbal H.: *Generalized enriched cyclic contractions with application to generalized iterated function system*. *Chaos, Solitons and Fractals*. **154**, 111591 (2022).
- [3] Anjum R, Abbas M, Berinde V.: *Enriched multivalued contractions with applications to differential inclusions and dynamic programming*. *Symmetry*. **13**(8), 1350 (2021).
- [4] Anjum R, Abbas M, Berinde V.: *Equivalence of certain iteration processes obtained by two new classes of operators*. *Mathematics*. **9**(18), 2292 (2021).
- [5] Anjum R, Abbas M.: *Common fixed point theorem for modified Kannan enriched contraction pair in Banach spaces and its applications*. *Filomat*. **35**(8), 2485-95 (2021).
- [6] Babu G.V, Rao C.S, Mounika P.: *A fixed point theorem on enriched (ψ, ϕ, λ) -weakly contractive map in convex metric space*. *Journal of Ramanujan Society of Mathematics and Mathematical Sciences*. **10**(1), (2022).
- [7] Berinde V, Păcurar M.: *Approximating fixed points of enriched contractions in Banach spaces*. *Journal of Fixed Point Theory and Applications*. **22**. 1-10 (2020).
- [8] Berinde V, Păcurar M, Kannan's fixed point approximation for solving split feasibility and variational inequality problems. *Journal of Computational and Applied Mathematics*. 377-427 (2020).
- [9] Berinde V.: *Approximating fixed points of enriched nonexpansive mappings by Krasnoselskii iteration in Hilbert spaces*. *Carpathian Journal of Mathematics*. **35**(3), 293-304 (2019).
- [10] Berinde V, Păcurar M.: *Approximating fixed points of enriched Chatterjea contractions by Krasnoselskii iterative algorithm in Banach spaces*. *Journal of Fixed Point Theory and Applications*. **23**, 1-16 (2021).
- [11] Berinde V, Păcurar M.: *Fixed point theorems for enriched Ćirić-Reich-Rus contractions in Banach spaces and convex metric spaces*. *Carpathian Journal of Mathematics*. **37**, 173-184 (2021).
- [12] Chugh R, Batra C.: *Fixed point theorems of enriched Ćirić type and enriched Hardy-Rogers contractions*. *Numerical Algebra, Control and Optimization*. 0-0 (2023).
- [13] Faraji H, Radenović S.T.: *Some fixed point results of enriched contractions by Krasnoselskii iterative method in partially ordered Banach spaces*. *Trans. A Razmadze Math. Inst.* to appear. (2020).
- [14] García, G., *A generalization of the (b, θ) -enriched contractions based on the degree of non-densifiability*. *Asian-European Journal of Mathematics*. **15**(09), 2250168 (2022).
- [15] Górnicki J, Bisht R.K.: *Around averaged mappings*. *Journal of Fixed Point Theory and Applications*. 1-12 (2021).
- [16] Mondal P, Garai H, Dey L.K.: *On some enriched contractions in Banach spaces*. *arXiv preprint arXiv*. **2006**, 11500 (2020).
- [17] Mondal P, Garai H, Dey L.K.: *On some enriched contractions in Banach spaces and an application*. *Filomat*. **35**(15), 5017-29 (2021).
- [18] Marchiş A.: *Common fixed point theorems for enriched Jungck contractions in Banach spaces*. *Journal of Fixed Point Theory and Applications*. **23**(4), 76 (2021).
- [19] Rawat S, Kukreti S, Dimri R.C.: *Fixed point results for enriched ordered contractions in noncommutative Banach spaces*. *The Journal of Analysis*. **30**(4), 1555-1566 (2022).
- [20] Rawat S, Bartwal A, Dimri R.C.: *Approximation and existence of fixed points via interpolative enriched contractions*. *FILOMAT*. **37**(16), 5455-5467 (2023).
- [21] Simsek E, Yildirim I.: *Krasnoselskii iteration process for approximating fixed points of enriched generalized nonexpansive mappings in Banach spaces*. *Carpathian Mathematical Publications*. **14**(1), 86-94 (2022).
- [22] Simsek E, Yildirim I.: *On new fixed point results for some classes of enriched mappings in N -Banach Spaces*. *Communications in Nonlinear Analysis*. **10**(1) 1-9 (2022).
- [23] Salisu S, Kumam P, Sriwongsa S.: *On fixed points of enriched contractions and enriched nonexpansive mappings*. *Carpathian Journal of Mathematics*. **39**(1), 237-254 (2023).
- [24] Nithiarayaphaks W, Sintunavarat W.: *On approximating fixed points of weak enriched contraction mappings via Kirk's iterative algorithm in Banach spaces*. *Carpathian Journal of Mathematics*. **39**(2), (2023).

- [25] Berinde, V.: Iterative Approximation of Fixed Points, Lecture Notes in Mathematics. Springer, Berlin, (2007)
- [26] Chidume, C.: Geometric properties of Banach spaces and nonlinear iterations. In Lecture Notes in Mathematics; Springer: London, UK, (2009)
- [27] Goebel K., Reich, S.: Uniform Convexity, Hyperbolic Geometry, and Nonexpansive Mappings; Monographs and Textbooks in Pure and Applied Mathematics; Marcel Dekker, Inc.: New York, NY, USA, (1984)
- [28] Reich S., Zaslavski A. J.: Asymptotic behavior of inexact orbits of nonexpansive mappings, Topological Methods in Nonlinear Analysis. 1-11 (2020) DOI: 10.12775/TMNA.2020.043
- [29] Reich S., Zaslavski A.J.: A fixed point result for mean nonexpansive mappings, Optimization. <https://doi.org/10.1080/02331934.2020.1727903>
- [30] Reich S., Salinas Z.: Weak convergence of infinite products of operators in Hadamard spaces, Rendiconti del Circolo Matematico di Palermo Series 2, DOI: 10.1007/s12215-015-0218-6
- [31] Reich, S.: Weak convergence theorems for nonexpansive mappings in Banach spaces, J. Math. Anal. Appl. 67, 274–276 (1979)
- [32] Browder, F. E.: Nonlinear monotone operators and convex sets in Banach spaces, Bull. Amer. Math. Soc. 71 (1965), 780–785
- [33] Browder, F. E.: Nonexpansive nonlinear operators in a Banach space, Proc. Nat. Acad. Sci. 54 (1965), 1041–1044
- [34] Browder, F. E.: Convergence theorems for sequences of nonlinear operators in Banach spaces, Mathematische Zeitschrift 100 (1967), 201–225
- [35] Browder, F. E.: Convergence of approximants to fixed points of non-expansive nonlinear mappings in Banach spaces, Arch. Rational Mech. Anal. 24 (1967), 82–90
- [36] Browder, F. E., Petryshyn, W. V.: The solution by iteration of nonlinear functional equations in Banach spaces, Bull. Amer. Math. Soc. 72 (1966), 571–575.
- [37] Browder, F. E., Petryshyn, W. V.: Construction of fixed points of nonlinear mappings in Hilbert spaces, J. Math. Anal. Appl. 20 (1967), 197–228
- [38] Berinde, V.: A Modified Krasnosel'skii-Mann Iterative Algorithm for Approximating Fixed Points of Enriched Nonexpansive Mappings. Symmetry. 14, 123 (2022)

RIZWAN ANJUM, DEPARTMENT OF MATHEMATICS, DIVISION OF SCIENCE AND TECHNOLOGY, UNIVERSITY OF EDUCATION, LAHORE 54770, PAKISTAN.

Email address: rizwananjum1723@gmail.com

DEPARTMENT OF MECHANICAL ENGINEERING SCIENCES, FACULTY OF ENGINEERING AND THE BUILT ENVIRONMENT, DOORNFONTEIN CAMPUS, UNIVERSITY OF JOHANNESBURG, SOUTH AFRICA

Email address: mujahida@uj.ac.za