

Self-supervised contrastive learning unveils cortical folding pattern linked to prematurity

Julien Laval

JULIEN.LAVAL@CEA.FR

Denis Rivière

DENIS.RIVIERE@CEA.FR

Aymeric Gaudin

AYMERIC.GAUDIN@CEA.FR

Vincent Frouin

VINCENT.FROUIN@CEA.FR

Jessica Dubois

JESSICA.DUBOIS@CEA.FR

Andrea Gondova

ANDREA.GONDOVA@ETU.U-PARIS.FR

Jean-François Mangin

JEAN-FRANCOIS.MANGIN@CEA.FR

Joël Chavas

JOEL.CHAVAS@CEA.FR

NeuroSpin, CEA Saclay, Université Paris-Saclay, France

Editors: Under Review for MIDL 2024

Abstract

Brain folding patterns have been reported to carry clinically relevant information. The brain folds mainly during the last trimester of pregnancy, and the process might be durably disturbed by preterm birth. Yet little is known about preterm-specific patterns. In this work, we train a self-supervised model (SimCLR) on the UKBioBank cohort (21070 adults) to represent the right superior temporal sulcus (STS) region and apply it to sulci images of 374 babies from the dHCP database, containing preterms and full-terms, and acquired at 40 weeks post-menstrual age. We find a lower variability in the preterm embeddings, supported by the identification of a knob pattern, missing in the extremely preterm population. The code can be found on [Github](#).

Keywords: Brain, sulci, preterms, self-supervised learning, contrastive learning

1. Introduction

Brain folding can be described using sulci, which are the grooves separating surface humps, called gyri. Main sulci are common to all individuals, but they differ in shape. Recurrent shape patterns can be identified, and some have been reported to be correlated to pathologies ([Mellerio et al., 2015](#); [Kabat and Król, 2012](#)). Therefore, studying folding patterns can pave the way for better diagnosis and understanding of brain diseases. In particular, finding preterm-specific patterns could lead to a better understanding of their developmental trajectories and their susceptibility to neurodevelopmental diseases ([Hee Chung et al., 2020](#)). In this paper, we apply a self-supervised contrastive learning algorithm based on SimCLR ([Chen et al., 2020](#)) to extract preterm-specific patterns.

2. Methods

The dHCP database: We process a subset of 374 subjects of the Developing Human Connectome Project (dHCP) database ([Edwards et al., 2022](#)) containing subjects born between 23 and 41 post-menstrual weeks and classified according to their degree of prematurity (Table 1). All images were acquired at ~ 40 weeks post-menstrual using the same machine and protocol.

Degree of prematurity	number of subjects
Extremely preterm (< 28 weeks)	19
Very preterm (28 ≤ weeks < 32)	35
Moderate to late preterm (32 ≤ weeks < 37)	43
full-term (≥ 37 weeks)	277

Table 1: Distribution of birth ages in dHCP, according to WHO classification

Inputs: The starting points are structural MR images of the brain. They are processed through the BrainVisa Morphologist pipeline¹ that skeletonizes a negative cast of the brain. It transforms the sulci into surfaces (3D objects of one-voxel width) following the middle of the folding. In practice, we use regional crops so that the algorithm can focus on specific sulci using the deep_folding toolbox (Chavas et al., 2022)². Cortical skeleton images are affinely normalized in the Talairach space with a 2mm voxel size. This paper focuses on the right superior temporal sulcus (STS). It was chosen because its development spans the whole third trimester of pregnancy (Dubois et al., 2008).

Model parameters and evaluation: Gaudin et al. (2024) optimized a SimCLR pipeline applied to the cingular region of the brain, classifying the presence of the paracingular sulcus as a pretext task. They selected the backbone, the augmentations, and hyperparameters. The main augmentation is called "branch clipping". It removes branches of the skeleton until a given percentage of voxels is removed. No further optimization was performed in the present study because we want to evaluate the performance (thus the generalizability) of a model optimized on another brain region³. To assess the model performances, we apply a two-layer linear Support Vector Classifier to the model embeddings of dHCP for each prematurity group (the negative class being the full-terms) and measure the 3-fold cross-validation AUC (with class-based stratification).

3. Experiments and results

We train the model on the STS crops of 21070 adult brains from UkBioBank (Sudlow et al., 2015). Next, we compute the embeddings of the dHCP subjects and perform classification (§2) to assess performances. We reach an AUC of **87.4(1.0)** for the extremely preterm class (Fig. 1a), confirming the ability of the model to capture information related to prematurity. Using UMAP (McInnes et al., 2020), we visualize the embedding space. The more premature the babies, the less variable they seem to be (see Laval (2024) for the UMAP). We quantify it in the 10-dimension latent space and find significant loss of variability for extremely and very preterms compared to full-terms (Fig. 1b). Then, we focus on the extremely preterms to characterize the variability drop visually. We select the top-predicted full-terms because they are more likely to contrast with preterms, and we identify an STS pattern shared by all top-predicted subjects, which is in stark contrast with typical extremely preterm STS: a long sulcus with a deep large knob at its anterior extremity. This pattern is then

1. <https://brainvisa.info>

2. [github: neurospin/deep_folding](https://github.com/neurospin/deep_folding)

3. The main parameters are the following: backbone: ConvNet, input size: (1, 22, 49, 46), embedding size: 10, number of trainable parameters: 452 172, loss temperature: 0.1, drop rate: 0.05, branch clipping percentage: 40%, batch size: 16, lr: 0.0004, weight decay: 5.0e-05, max epochs: 250.

identified in at least 5/19 randomly selected full-terms, which confirms quantitatively that it is missing in the extremely preterm population ($p = 0.008$ using Barnard’s exact test, Fig. 1c).




Figure 1: **a)** AUC between preterms and full-terms for each class of prematurity (averaged over 3 trainings). **b)** Average pairwise Euclidean distance within each group in the latent space ($\pm 2\sigma$ in blue). An F-test is performed to compute a p-value on the decrease of the variance ^a for each preterm category compared to full-term. (, 28): $p = 0.01$, (28, 32): $p = 0.01$, (32, 37): $p = 0.1$. **c)** STS visualization (red) in whole brain context (sulci in blue) of dHCP subjects sampled based on the extremely preterm classification score. In each group, 19 subjects are sampled for the long knob pattern identification (marked in green). Only 4 are shown here, Laval (2024) gives exhaustive visualization.

^a computed as the average squared Euclidean distance to the barycenter of the distribution.

4. Conclusion

In this work, using SimCLR, we identify an STS pattern absent from the extremely preterm data, suggesting a more unique developmental trajectory for preterms than full-terms. We showcase the transferability of the model optimized on adults in the cingular region to baby data in the STS region. Future work will be to repeat such analysis in all brain regions for the three classes of prematurity to uncover other regions altered by preterm birth.

Acknowledgments

This work was funded by the ANR grants FOLDDICO and BABYTOUCH.

References

- Joël Chavas, Louise Guillon, Marco Pascucci, Benoît Dufumier, Denis Rivière, and Jean-François Mangin. Unsupervised Representation Learning of Cingulate Cortical Folding Patterns. In Linwei Wang, Qi Dou, P. Thomas Fletcher, Stefanie Speidel, and Shuo Li, editors, *Medical Image Computing and Computer Assisted Intervention – MICCAI 2022*, Lecture Notes in Computer Science, pages 77–87, Cham, 2022. Springer Nature Switzerland. ISBN 978-3-031-16431-6.
- Ting Chen, Simon Kornblith, Mohammad Norouzi, and Geoffrey Hinton. A Simple Framework for Contrastive Learning of Visual Representations. 2020. doi: 10.48550/ARXIV.2002.05709. URL <https://arxiv.org/abs/2002.05709>. Publisher: arXiv Version Number: 3.
- J Dubois, M Benders, A Cachia, F Lazeyras, R Ha-Vinh Leuchter, S. V. Sizonenko, C Borradori-Tolsa, J. F. Mangin, and P. S. Hüppi. Mapping the Early Cortical Folding Process in the Preterm Newborn Brain. *Cerebral Cortex*, 18(6):1444–1454, June 2008. ISSN 1047-3211. doi: 10.1093/cercor/bhm180. URL <https://doi.org/10.1093/cercor/bhm180>.
- A. David Edwards, Daniel Rueckert, Stephen M. Smith, Samy Abo Seada, Amir Alansary, Jennifer Almalbis, Joanna Allsop, Jesper Andersson, Tomoki Arichi, Sophie Arulkumar, Matteo Bastiani, Dafnis Batalle, Luke Baxter, Jelena Bozek, Eleanor Braithwaite, Jacqueline Brandon, Olivia Carney, Andrew Chew, Daan Christiaens, Raymond Chung, Kathleen Colford, Lucilio Cordero-Grande, Serena J. Counsell, Harriet Cullen, John Cupitt, Charles Curtis, Alice Davidson, Maria Deprez, Louise Dillon, Konstantina Dimitrakopoulou, Ralica Dimitrova, Eugene Duff, Shona Falconer, Seyedeh-Rezvan Farahibozorg, Sean P. Fitzgibbon, Jianliang Gao, Andreia Gaspar, Nicholas Harper, Sam J. Harrison, Emer J. Hughes, Jana Hutter, Mark Jenkinson, Saad Jbabdi, Emily Jones, Vyacheslav Karolis, Vanessa Kyriakopoulou, Gregor Lenz, Antonios Makropoulos, Shaihan Malik, Luke Mason, Filippo Mortari, Chiara Nosarti, Rita G. Nunes, Camilla O’Keeffe, Jonathan O’Muircheartaigh, Hamel Patel, Jonathan Passerat-Palmbach, Maximilian Pietsch, Anthony N. Price, Emma C. Robinson, Mary A. Rutherford, Andreas Schuh, Stamatios Sotiropoulos, Johannes Steinweg, Rui Pedro Azeredo Gomes Teixeira, Tencho Tenev, Jacques-Donald Tournier, Nora Tusor, Alena Uus, Katy Vecchiato, Logan Z. J. Williams, Robert Wright, Julia Wurie, and Joseph V. Hajnal. The Developing Human Connectome Project Neonatal Data Release. *Frontiers in Neuroscience*, 16, 2022. ISSN 1662-453X. URL <https://www.frontiersin.org/articles/10.3389/fnins.2022.886772>.
- Aymeric Gaudin, Louise Guillon, Clara Fischer, Arnaud Cachia, Denis Rivière, Jean-François Mangin, and Joël Chavas. Optimizing contrastive learning for cortical folding pattern detection. *SPIE 2024, in press*, 2024.
- Estefani Hee Chung, Jesse Chou, and Kelly A. Brown. Neurodevelopmental outcomes of preterm infants: a recent literature review. *Translational Pediatrics*, 9(Suppl 1):S3–S8, February 2020. ISSN 2224-4344. doi: 10.21037/tp.2019.09.10.

- Joanna Kabat and Przemysław Król. Ogniskowa dysplazja korowa – stan obecny wiedzy. *Polish Journal of Radiology*, 77(2):35–43, 2012. ISSN 0137-7183. doi: 10.12659/PJR.882968. URL <http://www.polradiol.com/index.php?archives/article/882968>.
- Julien Laval. Self-supervised learning for preterm-specific variability of cortical folding in the superior temporal sulcus region. Technical report, CEA - Commissariat à l'énergie atomique et aux énergies alternatives, April 2024. URL <https://hal.science/hal-04538440>.
- Leland McInnes, John Healy, and James Melville. UMAP: Uniform Manifold Approximation and Projection for Dimension Reduction, September 2020. URL <http://arxiv.org/abs/1802.03426>. arXiv:1802.03426 [cs, stat].
- Charles Mellerio, Pauline Roca, Francine Chassoux, Florian Danière, Arnaud Cachia, Stéphanie Lion, Olivier Naggara, Bertrand Devaux, Jean-François Meder, and Catherine Oppenheim. The Power Button Sign: A Newly Described Central Sulcal Pattern on Surface Rendering MR Images of Type 2 Focal Cortical Dysplasia. *Radiology*, 274(2): 500–507, February 2015. ISSN 0033-8419, 1527-1315. doi: 10.1148/radiol.14140773. URL <http://pubs.rsna.org/doi/10.1148/radiol.14140773>.
- Cathie Sudlow, John Gallacher, Naomi Allen, Valerie Beral, Paul Burton, John Danesh, Paul Downey, Paul Elliott, Jane Green, Martin Landray, Bette Liu, Paul Matthews, Giok Ong, Jill Pell, Alan Silman, Alan Young, Tim Sprosen, Tim Peakman, and Rory Collins. UK Biobank: An Open Access Resource for Identifying the Causes of a Wide Range of Complex Diseases of Middle and Old Age. *PLOS Medicine*, 12(3):e1001779, March 2015. ISSN 1549-1676. doi: 10.1371/journal.pmed.1001779. URL <https://dx.plos.org/10.1371/journal.pmed.1001779>.