

High dimensional test for functional covariates

Huaqing Jin and Fei Jiang *

Department of Epidemiology and Biostatistics,
University of California, San Francisco

May 16, 2024

Abstract

As medical devices become more complex, they routinely collect extensive and complicated data. While classical regressions typically examine the relationship between an outcome and a vector of predictors, it becomes imperative to identify the relationship with predictors possessing functional structures. In this article, we introduce a novel inference procedure for examining the relationship between outcomes and large-scale functional predictors. We target testing the linear hypothesis on the functional parameters under the generalized functional linear regression framework, where the number of the functional parameters grows with the sample size. We develop the estimation procedure for the high dimensional generalized functional linear model incorporating B-spline functional approximation and amenable regularization. Furthermore, we construct a procedure that is able to test the local alternative hypothesis on the linear combinations of the functional parameters. We establish the statistical guarantees in terms of non-asymptotic convergence of the parameter estimation and the oracle property and asymptotic normality of the estimators. Moreover, we derive the asymptotic distribution of the test statistic. We carry out intensive simulations and illustrate with a new dataset from an Alzheimer's disease magnetoencephalography study.

Keywords: Alzheimer's disease, B-spline, Generalized functional linear model, Magnetoencephalography, Neuroimaging data.

*Email: fei.jiang@ucsf.edu

1 Introduction

Functional regression is widely used to model the relationship between a scalar response and functional predictors. Traditional approaches typically only involve low-dimensional functional predictors. Among the studies, Yuan & Cai (2010) and Cai & Yuan (2012) investigate parameter estimation in functional linear regression using a single functional predictor. Lian (2013) introduces a shrinkage functional estimator for functional linear regression in a finite-dimensional setting. Furthermore, Kong et al. (2016) explore a partial functional linear regression with finite-dimensional functional predictors and high-dimensional baseline covariates. In addition, Lei (2014), Hilgert et al. (2013) and Shang & Cheng (2015) have formulated testing procedures to assess hypotheses concerning the finite-dimensional functional predictors in the generalized functional linear models. However, in numerous applications, the number of functional predictors can grow with the sample size. Examples can be found in neuroimaging analyses that focus on the relationship between a disease marker and temporal signals from several brain regions of interest (ROI) recorded by magnetoencephalography (MEG), electroencephalogram (EEG) or functional magnetic resonance imaging (fMRI). Under this context, Gertheiss et al. (2013) adopt a penalized likelihood method for variable selection under the generalized functional linear model framework with an application to a neuroimaging dataset. Fan et al. (2015) propose a functional additive regression model to handle a large number of functional predictors. Zhang, Zhang, Ma & Fang (2022) consider the subgroup analysis problem under the high-dimensional functional regression framework. Zhang, Fang & Zhang (2022) propose to estimate the high dimensional functional parameters in the generalized additive model through a penalized regression method utilizing group minimax concave penalty. Xue et al. (2023) introduce a procedure for estimating high-dimensional functional parameters, designed specifically

to analyze EEG data. Despite the existence of various estimation techniques, options for hypothesis testing on high-dimensional functional parameters in the generalized functional linear model framework remain scarce. To the best of our knowledge, the bootstrap method proposed by Xue & Yao (2021) stands out as the sole known testing approach in this area specifically for the functional linear regression. However, this method does not extend to the nonlinear regression models. Moreover, it lacks the capability to test for local alternative hypotheses, incorporate baseline covariates and does not aim to test linear hypotheses. To address these limitations, we introduce a novel hypothesis testing procedure that applies to high-dimensional functional predictors and goes beyond the functional linear regression. This innovative approach is built on the generalized functional linear regression and extended to test local alternatives and linear hypotheses of the functional parameters of interest.

We formulate a high-dimensional hypothesis testing procedure for functional predictors (HDFP) under the generalized functional linear regression framework. Our work is motivated by the Alzheimer’s disease (AD) study (Jin et al. 2023) that aims to investigate the association between the features from MEG recordings and the cognitive function in 82 AD patients and 61 healthy controls. A critical question to explore is the relationship between the power spectrum densities (PSD) of the MEG data and cognitive function, which can be measured by cognitive scores such as the mini-mental state examination (MMSE) score (Kurlowicz & Wallace 1999). Here, a PSD from one brain region of interest (ROI) is characterized as the square of the Fourier transform of the raw time series across a sequence of frequencies. Understanding the association is crucial as it can lead us to create new treatments aiming at suppressing abnormal waveforms, thereby potentially slowing down the neurodegenerative process (Bosco et al. 2023). To improve interpretability, rather than us-

ing raw time series data as predictors, we map the signals recorded from the scalp onto the brain surface. We then consolidate the voxel-level time series into region-level aggregates, which leads to a 68-dimensional functional predictor based on the Desikan-Killiany (DK) atlas (Desikan et al. 2006). Finally, Fourier transformations are applied independently to each brain signal to generate region-wise PSDs over the whole brain. Conventionally, the average PSDs over all frequencies or a subset of frequencies are used as a predictor of the cognitive function (Wang et al. 2015). This approach ignores the functional pattern of PSD, resulting in the predictors that do not sufficiently explain the variations in the outcome. We apply this approach to study the relationship between the MMSE score and the 68 dimensional average PSDs (over all frequencies) through a linear regression model adjusted for age, gender, and education level. Figure 1A shows that the PSDs from only two ROIs significantly affect the MMSE score with the p-values less than $0.05/68$. This finding deviates from the results in the existing literature (Jagust 2018, Jin et al. 2023). We further calculate the average PSD within the specific frequency bands, namely delta (2HZ–4HZ), theta (4HZ–8HZ), alpha (8HZ–12HZ), and beta (12HZ–35HZ) bands and conduct similar linear regression analysis. We then identify the significant ROIs corresponding to each frequency band. Figure 1B shows that the significant ROI varies across the frequency bands, suggesting that the effect of the PSDs on the MMSE score varies across the frequencies. This preliminary study indicates that incorporating functional parameters that vary with frequency is essential for explaining the relationship between PSD and cognitive function.

To study the functional relationship between the PSDs over a large number of ROIs and the cognitive function or the disease indicator, we pursue a high dimensional generalized functional linear regression. We develop a hypothesis testing procedure to identify the important linear combinations of the function predictors. Specifically, we target the




Figure 1: **A:** Negative log p-values from the significant ROIs selected via fitting MMSE score on each ROI's average PSD, adjusted for age, gender, and education level. The significant ROIs are identified by the p-values less than $0.05/68$. **B:** Negative log p-values from the significant ROIs selected via fitting MMSE score on each ROI's average PSD at delta, theta, alpha, and beta bands, adjusted for age, gender, and education level. The significant ROIs are identified by the p-values less than $0.05/68/4$.

association between a response variable Y and a high-dimensional covariate of interest $\mathbf{X}(s)$ adjusted by the confounder \mathbf{Z} . The association between Y and $\mathbf{X}(s)$ varies with s . In our motivating example, Y is the MMSE score, $\mathbf{X}(s)$ is the PSD at the frequency s , and \mathbf{Z} contains information about a subject's age, gender, and education level. We model the conditional distribution of Y given \mathbf{X} and \mathbf{Z} through $f\left\{Y, \boldsymbol{\alpha}^T \mathbf{Z} + \int_0^1 \boldsymbol{\beta}^T(s) \mathbf{X}(s) ds\right\}$, where $f(\cdot)$ is a conditional density function in the exponential family. When $\mathbf{X}(s)$ is a one dimensional function, we can approximate $\boldsymbol{\beta}(s)$ through a basis expansion $\mathbf{B}(s)^T \boldsymbol{\gamma}$, where $\boldsymbol{\gamma}$ is an unknown parameter. In this case, the regression problem simplifies to a generalized linear model, and the hypothesis on $\boldsymbol{\beta}(s)$ can be transformed to a linear hypothesis on $\boldsymbol{\gamma}$. However when $\mathbf{X}(s)$ is a high-dimensional function, it is unclear whether any of the existing

proposed tests are applicable. The first obstacle in constructing valid hypothesis testing procedures is that the asymptotic distribution of the resulting estimator has not been established. We utilize the B-splines to approximate the functional parameters and derive the asymptotic normality of the functional estimator when the amenable penalties (Loh & Wainwright 2015, 2017) are used for the parameter estimation. Even after establishing the asymptotic normal properties, it is still challenging to construct the testing under the high dimensional setting for the generalized functional linear regression. This is because under the amenable penalties (Loh & Wainwright 2015, 2017), the asymptotic normality of the estimators is built on a minimal signal condition, which requires the norm of the nonzero elements in $\boldsymbol{\beta}(s)$ to be at least of order λ . Here λ is the penalty parameter which goes to zero when the sample size increases. Now consider testing the null hypothesis $\beta_1(s) = 0$ versus the alternative $\beta_1(s) = h_n(s)$, where $\beta_1(s)$ is the first element of $\boldsymbol{\beta}(s)$. The minimal signal condition implies that the test will have no power in testing the local alternative when $(\int h_n(s)^2 ds)^{1/2} \leq \lambda$. To address this issue, we eliminate the penalties on the subset of parameters involved in the test. However, it remains uncertain how quickly the dimension of this parameter subset can increase while still maintaining adequate test power and the number of basis functions required to approximate the function. To clarify this, we have derived the convergence properties of the estimators. These properties specify the rate at which the dimension of the parameter can grow relative to the sample size, while considering an increasing number of B-spline bases. This analysis ensures consistency, asymptotic normality, and adequate testing power.

We briefly summarize our contributions as follows. First, we develop a new estimation procedure for the high-dimensional generalized functional linear model with amenable regularization. Second, we show the variable selection consistency and the consistency of

the resulting estimator. We provide the explicit convergence rate of the estimator. Third, we derive the asymptotic normality of the estimator for the nonzero parameters and the parameters to test. Fourth, we construct a test statistic for linear hypothesis in the form of $\mathbf{C}\boldsymbol{\beta}(s) = \mathbf{t}(s)$, where \mathbf{C} is a known matrix and \mathbf{t} is a known function. Fifth, we derive the asymptotic distribution of the test statistics. These five essential components collectively allow us to perform hypothesis testing for a generalized functional linear model with high-dimensional functional parameters.

Our proposal is related to but also clearly different from existing high-dimensional test procedure. First, our model captures varying effects through functional parameters. In this sense, it is similar in spirit to Xue & Yao (2021)'s bootstrap-based test. Nevertheless, our approach is extended to the generalized functional linear model, accommodating models with baseline covariates. Additionally, our approach is capable of testing linear and local alternatives hypotheses, which is similar to Shi et al. (2019)'s solution. However, we consider functional parameters and primarily focus on a group sparsity type of penalties (Yuan & Lin 2006).

The rest of the paper is organized as follows. In Section 2, we present the model, discuss the estimation procedure and introduce the test statistic. In Section 3 we present the theoretic properties of the estimators and the test statistic. In Section 4, we validate the theoretical properties of the proposed method, and perform the benchmark comparison through simulation studies. Finally, we apply the proposed method to analyze the data from the AD MEG study in Section 5.

Notations: We first introduce some general notations that will be used throughout the text. For a matrix \mathbf{M} , let $\|\mathbf{M}\|_{\max}$ be the matrix maximum norm, $\|\mathbf{M}\|_{\infty}$ be the L_{∞} norm and $\|\mathbf{M}\|_p$ be the L_p norm. For a general vector \mathbf{a} , let $\|\mathbf{a}\|_{\infty}$ be the vector sup-norm,

$\|\mathbf{a}\|_p$ be the vector l_p -norm. Let \mathbf{e}_j be the unit vector with 1 on its j th entry. For a vector $\mathbf{v} = (v_1, \dots, v_p)^\top$, let $\text{supp}(\mathbf{v})$ be the set of indices with $v_i \neq 0$ and $\|\mathbf{v}\|_0 = |\text{supp}(\mathbf{v})|$, where $|\mathcal{U}|$ stands for the cardinality of the set \mathcal{U} . For a vector $\mathbf{v} \in \mathbb{R}^p$ and a subset $S \subseteq (1, \dots, p)$, we use $\mathbf{v}_S \in \mathbb{R}^S$ to denote the vector obtained by restricting \mathbf{v} on the set S . For a matrix \mathbf{M} , denote \mathbf{M}_k and $\mathbf{M}_{\cdot j}$ be the k th row and j th column of matrix \mathbf{M} . Further, we define $\|X\|_{\psi_1} \equiv \sup_{k \geq 1} k^{-1} E(|X|^k)^{1/k}$, and $\|X\|_{\psi_2} \equiv \sup_{k \geq 1} k^{-1/2} E(|X|^k)^{1/k}$. Following Fletcher & Watson (1980), for an arbitrary norm $\|\cdot\|_A$ and its dual norm $\|\cdot\|_D$, we define $\partial\|\mathbf{x}\|_A$ as the set $\{\mathbf{v} : \|\mathbf{x}\|_A = \mathbf{v}^\top \mathbf{x}, \|\mathbf{v}\|_D \leq 1\}$. Thus, for an arbitrary vector $\mathbf{x} = (x_1, \dots, x_p)^\top$, $\partial\|\mathbf{x}\|_1 = \{\mathbf{v} = (v_1, \dots, v_p)^\top : v_j = \text{sign}(x_j) \text{ if } x_j \neq 0, \text{ and } |v_j| \leq 1 \text{ if } x_j = 0\}$, and $\partial\|\mathbf{x}\|_2 = \{\mathbf{v} = (v_1, \dots, v_p)^\top : v_j = x_j / \|\mathbf{x}\|_2\}$. Following the definition in Loh & Wainwright (2012), a random vector \mathbf{V} is sub-Gaussian or sub-exponential if for any given unit vector \mathbf{u} , $\mathbf{u}^\top \mathbf{V}$ is sub-Gaussian or sub-exponential, respectively.

2 Model, Estimation, and Test Statistic

2.1 Problem formulation

Consider a high-dimensional functional covariate $\mathbf{X}(s) = \{X_1(s), \dots, X_d(s)\}^\top \in \mathbb{R}^d, s \in [0, \tau]$, where d is the number of the functional predictors. Without loss of generality, we assume $\tau = 1$, because we can always project any compact intervals to $[0, 1]$. In our application, we consider $X_j(s)$ to be the PSD (in \log_{10} scale) of MEG/EEG recording from j th region of interest. We consider the PSD rather than the original time series because the PSD analysis of the MEG/EEG data is a common practice in the neuroimaging community, due to its ability to provide a more stable and less noisy representation of the brain signals (Ranasinghe et al. 2022, Wang et al. 2015). Note that while we focus on the PSD in this

paper, our method can also be applied to the original time series data.

Let $\mathbf{Z} = (Z_1, \dots, Z_q)^\top$ be a vector of baseline covariates with $q < \infty$. Let $p = d + q$ and Y be the outcome of interest, such as cognitive scores or disease indicator. We consider the conditional distribution of Y_i given $\mathbf{X}_i, \mathbf{Z}_i$ for the independent identically distributed $\{(Y_i, \mathbf{X}_i, \mathbf{Z}_i), i = 1, \dots, n\}$ as

$$f \left\{ Y_i, \boldsymbol{\alpha}_0^\top \mathbf{Z}_i + \int_0^1 \boldsymbol{\beta}^\top(s) \mathbf{X}_i(s) ds \right\} \\ = \exp \left[\frac{Y \left\{ \boldsymbol{\alpha}_0^\top \mathbf{Z}_i + \int_0^1 \boldsymbol{\beta}^\top(s) \mathbf{X}(s) ds \right\} - \psi \left\{ \boldsymbol{\alpha}_0^\top \mathbf{Z}_i + \int_0^1 \boldsymbol{\beta}^\top(s) \mathbf{X}(s) ds \right\}}{c(\sigma)} \right], \quad (1)$$

where $\boldsymbol{\beta}(s) = \{\beta_1(s), \dots, \beta_d(s)\}^\top$ represents the functional effects, $\boldsymbol{\alpha}_0$ is an unknown true effect of the baseline covariates, $c(\sigma)$ is a scale parameter, and f is a known conditional density function of Y given $\{\mathbf{Z}, \mathbf{X}(s)\}$ in the exponential family. Since we are interested in $\boldsymbol{\alpha}_0, \boldsymbol{\beta}(s)$, we rescale the loss function and assume $c(\sigma) = 1$.

We aim to test the hypothesis that

$$\mathbf{H}_0 : \mathbf{C}\boldsymbol{\beta}_{\mathcal{M}}(s) = \mathbf{t}(s) \text{ vs. } \mathbf{H}_1 : \mathbf{C}\boldsymbol{\beta}_{\mathcal{M}}(s) = \mathbf{t}(s) + \mathbf{h}_n(s), \quad (2)$$

where $\mathcal{M} \subset (1, \dots, d)$ with $|\mathcal{M}| = m$ is the subset of indices for the functional coefficients that we are interested in to test; \mathbf{C} is an $r \times m$ matrix and $\mathbf{t}(s)$ and $\mathbf{h}_n(s)$ are the known functions. Here $\mathbf{h}_n(s)$ quantifies the effect size under the alternative hypothesis. For example, if we are interested in testing the difference of the effects from the first two functional predictors, we can set $\mathbf{C} = [1, -1]$, $\mathbf{t}(s) \equiv \mathbf{0}$ and $\mathbf{h}_n(s)$ to be some nonzero function. In summary, by varying $\mathbf{C}, \mathbf{t}, \mathbf{h}_n$, and \mathcal{M} , we can generate different linear hypotheses to test.

2.2 Parameter estimation

Assume $\beta_j(s) \in C^\omega([0, 1]), j = 1, \dots, d$, is a smooth function with ω th order derivatives, we approximate $\beta_j(s)$ by a B-spline function $\mathbf{B}(s)^\top \boldsymbol{\gamma}_{0j}$, where $\mathbf{B}(s)$ is the N dimensional B-

spline basis function and γ_{0j} is the true spline coefficients. Let $\mathcal{S} = \{j : \beta_j(s) \neq 0, j \in \mathcal{M}^c\}$ and $k_0 = |\mathcal{S}|$, and $\mathbf{\Gamma}_0 \equiv (\gamma_{0j}, j = 1, \dots, d)$.

Furthermore, we consider the loss function as the negative of the logarithm of the likelihood function

$$\mathcal{L}(\boldsymbol{\alpha}, \mathbf{\Gamma}) = -n^{-1} \sum_{i=1}^n \log \left[f \left\{ Y_i, \boldsymbol{\alpha}^T \mathbf{Z}_i + \int_0^1 \mathbf{B}(s)^T \mathbf{\Gamma} \mathbf{X}_i(s) ds \right\} \right],$$

where $\mathbf{\Gamma} = (\gamma_j, j = 1, \dots, d)$. Let $\psi'_i(\boldsymbol{\alpha}, \mathbf{\Gamma}), \psi''_i(\boldsymbol{\alpha}, \mathbf{\Gamma})$ be the first and second derivatives of $\psi(t)$ evaluated at $t = \boldsymbol{\alpha}^T \mathbf{Z}_i + \int_0^1 \mathbf{B}(s)^T \mathbf{\Gamma} \mathbf{X}_i(s) ds$. Slightly abusing the notations, we define $\psi_i\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\}, \psi'_i\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\}, \psi''_i\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\}$ be the cumulant function and its first and second derivatives of $\psi_i(t)$ evaluated at $t = \boldsymbol{\alpha}^T \mathbf{Z}_i + \int_0^1 \boldsymbol{\beta}(s)^T \mathbf{X}_i(s) ds$. In a generalized linear model $\psi''_i(\boldsymbol{\alpha}, \mathbf{\Gamma}) > 0$ and we assume that there is a constant C_ψ such that $|\psi''_i(\cdot, \cdot)| \leq C_\psi$ uniformly. These conditions are often used in the generalized linear model to ensure the estimation consistency (Loh & Wainwright 2015).

Suppose we are interested in testing whether $\mathbf{C}\boldsymbol{\beta}_{\mathcal{M}}(s) = \mathbf{t}(s)$ or not, when estimating $\boldsymbol{\alpha}, \mathbf{\Gamma}$, we minimize the regularized loss function

$$\hat{\boldsymbol{\alpha}}, \hat{\mathbf{\Gamma}} = \operatorname{argmin}_{\boldsymbol{\alpha}, \mathbf{\Gamma}} \left\{ \mathcal{L}(\boldsymbol{\alpha}, \mathbf{\Gamma}) + \sum_{j \in \mathcal{M}^c} \rho_\lambda(\|N^{-1/2} \boldsymbol{\gamma}_j\|_2) \right\}, \text{ such that } \|\boldsymbol{\alpha}\|_1 + \sum_{j=1}^d \|N^{-1/2} \boldsymbol{\gamma}_j\|_2 \leq R, \quad (3)$$

where ρ_λ is some regularized function and λ is the tuning parameter, and R is a constant that greater than $2\|\boldsymbol{\alpha}_0\|_1 + 2\sum_{j=1}^d \|N^{-1/2} \boldsymbol{\gamma}_{0j}\|_2$, $\hat{\boldsymbol{\alpha}}$ and $\hat{\mathbf{\Gamma}} = (\hat{\boldsymbol{\gamma}}_j, j = 1, \dots, d)$ are the estimators of $\boldsymbol{\alpha}$ and $\mathbf{\Gamma}$, respectively. The additional constraints $\|\boldsymbol{\alpha}_0\|_1 + 2\sum_{j=1}^d \|N^{-1/2} \boldsymbol{\gamma}_{0j}\|_2$ are only used to facilitate the theoretical derivation to control the magnitude of the estimator and have little empirical effect. Note that in (3), we do not impose the penalization on $\beta_j(s)$ for $j \in \mathcal{M}$, and hence do not shrink the estimators corresponding to the functional coefficients involved in the hypothesis. Since we do not shrink the estimators, the procedure can effectively test the local hypothesis.

2.3 Computational algorithm

For notation simplicity, we define a generic $q + dN$ dimensional vector $\boldsymbol{\theta} \equiv (\boldsymbol{\theta}_1^T, \boldsymbol{\theta}_{2j}^T, j = 1, \dots, d)^T \equiv \{\boldsymbol{\alpha}^T, N^{-1/2}\text{vec}(\boldsymbol{\Gamma})^T\}^T$, $\boldsymbol{\theta}_0 \equiv (\boldsymbol{\theta}_{01}^T, \boldsymbol{\theta}_{02j}^T, j = 1, \dots, d)^T \equiv \{\boldsymbol{\alpha}_0^T, N^{-1/2}\text{vec}(\boldsymbol{\Gamma}_0)^T\}^T$ and $\hat{\boldsymbol{\theta}} = (\hat{\boldsymbol{\theta}}_1^T, \hat{\boldsymbol{\theta}}_{2j}^T, j = 1, \dots, d)^T \equiv \{\hat{\boldsymbol{\alpha}}^T, N^{-1/2}\text{vec}(\hat{\boldsymbol{\Gamma}})^T\}^T$. Without loss of generality, we assume the last $(p - q - m - k_0)N$ element of $\boldsymbol{\theta}_0$ are zeros. We define $\boldsymbol{\theta}_{\mathcal{F}} = \{\boldsymbol{\theta}_1^T, \boldsymbol{\theta}_{2j}^T, j \in \mathcal{F}\}$ where \mathcal{F} is a subset of $\{1, \dots, d\}$ and define $\boldsymbol{\theta}_{0\mathcal{F}}$ and $\hat{\boldsymbol{\theta}}_{\mathcal{F}}$ in the similar way and denote $\mathcal{L}(\boldsymbol{\theta}) \equiv \mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\Gamma})$.

We utilize the contractive Peaceman-Rachford splitting method (CPRSM) (He et al. 2014) to solve for $\boldsymbol{\alpha}$ and $\boldsymbol{\Gamma}$ in (3). More specifically, let $\mathbf{D} = [\mathbf{0}_{dN, q}, \mathbf{I}_{dN \times dN}]$ and let $f_1(\boldsymbol{\theta}) = \mathcal{L}(\boldsymbol{\theta})$, where $\boldsymbol{\theta} = (\boldsymbol{\theta}_1^T, \boldsymbol{\theta}_{2j}^T, j = 1, \dots, d)$, and $f_2(\boldsymbol{\Gamma}) = \sum_{j=1}^{T-1} \rho_\lambda(\|N^{-1/2}\boldsymbol{\gamma}_j\|_2)$. We estimate $\boldsymbol{\alpha}$ and $\boldsymbol{\Gamma}$ by minimizing

$$f_1(\boldsymbol{\theta}) + f_2(\boldsymbol{\Gamma}) \text{ subject to } \mathbf{D}\boldsymbol{\theta} - N^{-1/2}\text{vec}(\boldsymbol{\Gamma}) = \mathbf{0} \text{ and } \|\boldsymbol{\theta}_1\|_1 + \sum_{j=1}^d \|\boldsymbol{\theta}_{2j}\|_2 \leq R. \quad (4)$$

At the k th iteration, CPRSM updates $\boldsymbol{\theta}^{(k)}$ as

$$\begin{aligned} \boldsymbol{\theta}^{k+1/2} &= \underset{\boldsymbol{\theta}}{\text{argmin}} f_1(\boldsymbol{\theta}) - \boldsymbol{\rho}^{(k)\text{T}}\{\mathbf{D}\boldsymbol{\theta} - N^{-1/2}\text{vec}(\boldsymbol{\Gamma}^{(k)})\} \\ &\quad + \frac{\beta}{2} \|\mathbf{D}\boldsymbol{\theta} - N^{-1/2}\text{vec}(\boldsymbol{\Gamma}^{(k)})\|_2^2, \end{aligned} \quad (5)$$

$$\check{\boldsymbol{\theta}}_1, \|\check{\boldsymbol{\theta}}_{2j}\|_2 = \text{Proj}_{\{\boldsymbol{\theta}: \|\boldsymbol{\theta}_1\|_1 + \sum_{j=1}^d \|\boldsymbol{\theta}_{2j}\|_2 \leq R\}}([\boldsymbol{\theta}_1^{(k+1/2)\text{T}}, \|\boldsymbol{\theta}_{2j}^{(k+1/2)}\|_2, j = 1, \dots, d]^T), \quad (6)$$

$$\boldsymbol{\theta}^{(k+1)} = (\check{\boldsymbol{\theta}}_1^T, \boldsymbol{\theta}_{2j}^{(k+1/2)\text{T}} \|\check{\boldsymbol{\theta}}_{2j}\|_2 / \|\boldsymbol{\theta}_{2j}^{(k+1/2)}\|_2, j = 1, \dots, d)^T, \quad (7)$$

$$\boldsymbol{\rho}^{k+1/2} = \boldsymbol{\rho}^{(k)} - \alpha\beta\{\mathbf{D}\boldsymbol{\theta}^{(k+1)} - N^{-1/2}\text{vec}(\boldsymbol{\Gamma}^{(k)})\}, \quad (8)$$

$$\begin{aligned} \boldsymbol{\Gamma}^{(k+1)} &= \underset{\boldsymbol{\Gamma}}{\text{argmin}} f_2(\boldsymbol{\Gamma}) - \boldsymbol{\rho}^{(k+1/2)\text{T}}\{\mathbf{D}\boldsymbol{\theta}^{(k+1)} - N^{-1/2}\text{vec}(\boldsymbol{\Gamma})\} \\ &\quad + \frac{\beta}{2} \|\mathbf{D}\boldsymbol{\theta}^{(k+1)} - N^{-1/2}\text{vec}(\boldsymbol{\Gamma})\|_2^2, \end{aligned} \quad (9)$$

$$\boldsymbol{\rho}^{(k+1)} = \boldsymbol{\rho}^{(k+1/2)} - \alpha\beta\{\mathbf{D}\boldsymbol{\theta}^{(k+1)} - N^{-1/2}\text{vec}(\boldsymbol{\Gamma}^{(k+1)})\}, \quad (10)$$

where (α, β) are the relaxation and penalty parameters in CPRSM and $\text{Proj}_C(\cdot)$ projects a vector to the feasible set C . When $\mathcal{L}(\boldsymbol{\theta})$ is the logarithm of a Gaussian likelihood,

(5) reduces to a quadratic minimization which can be solved analytically. When $\mathcal{L}(\boldsymbol{\theta})$ is the log-likelihood from the other generalized linear models, such as the logistic model, (5) can be solved using Newton-Raphson method iteratively. The projection step (6) can be achieved by adopting the linear programming method (Duchi et al. 2008). (9) is a standard penalized quadratic minimization step. Throughout the paper, we choose $\rho_\lambda(\cdot)$ as the smoothly clipped absolute deviation (SCAD) penalty (Fan & Li 2001), which is an amenable penalty satisfying Conditions (A1)–(A6) in Appendix D of the supplementary material (Loh & Wainwright 2015, 2017). When using the SCAD penalty, the minimization in (9) can be addressed by first calculating the analytical solution of the minimization without the penalty term, and then applying SCAD shrinkage to the resulting solution.

We summarize the CPRSM algorithm in Algorithm 1 in Appendix C of the supplementary material and more details of the computational algorithm are also provided there.

2.4 Test statistic

We propose a test statistic to test the hypothesis (2). Define

$$\mathbf{Q}\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\} \equiv E \left\{ \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)) \left(\left[\mathbf{Z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} \right\}.$$

The covariance matrix of the residuals is

$$\boldsymbol{\Sigma}\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\} \equiv E \left\{ \left([\psi_i'\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\} - Y_i] \left[\mathbf{Z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} \right\}.$$

Further, define

$$\mathbf{A} = (\mathbf{C} \otimes \mathbf{I}_{N \times N}) [\mathbf{0}_{mN \times q}, \mathbf{I}_{mN \times mN}, \mathbf{0}_{mN \times k_0N}],$$

$$\boldsymbol{\Psi}(\boldsymbol{\Sigma}, \mathbf{Q}, \boldsymbol{\beta}) = \mathbf{Q}\{\boldsymbol{\alpha}_0, \boldsymbol{\beta}(\cdot)\}_{\mathcal{M} \cup S, \mathcal{M} \cup S}^{-1} \boldsymbol{\Sigma}_{\mathcal{M} \cup S, \mathcal{M} \cup S} \mathbf{Q}\{\boldsymbol{\alpha}_0, \boldsymbol{\beta}(\cdot)\}_{\mathcal{M} \cup S, \mathcal{M} \cup S}^{-1},$$

where $\mathbf{M}_{\mathcal{M} \cup S, \mathcal{M} \cup S}$ is the top left $\{q + (m + k_0)N\} \times \{q + (m + k_0)N\}$ of an arbitrary matrix \mathbf{M} .

The sample versions of $\mathbf{Q}\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\}$, $\boldsymbol{\Sigma}\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\}$ and $\boldsymbol{\Psi}(\boldsymbol{\Sigma}, \mathbf{Q}, \boldsymbol{\beta})$ based on the estimator $\hat{\boldsymbol{\theta}}$ are denoted by $\hat{\mathbf{Q}}(\hat{\boldsymbol{\theta})}$, $\hat{\boldsymbol{\Sigma}}(\hat{\boldsymbol{\theta})}$ and $\boldsymbol{\Psi}(\hat{\boldsymbol{\Sigma}}, \hat{\mathbf{Q}}, \hat{\boldsymbol{\theta}})$, respectively. Then the test statistic is defined as

$$T = n \left(\mathbf{A}\hat{\boldsymbol{\theta}}_{\mathcal{M} \cup \mathcal{S}} - N^{-1/2} \text{vec} \left[\left\{ \int_0^1 \mathbf{B}(s)\mathbf{B}(s)^\top ds \right\}^{-1} \int_0^1 \mathbf{B}(s)\mathbf{t}^\top(s) ds \right] \right)^\top \mathbf{A}\boldsymbol{\Psi}(\hat{\boldsymbol{\Sigma}}, \hat{\mathbf{Q}}, \hat{\boldsymbol{\theta}})^{-1} \mathbf{A}^\top \left(\mathbf{A}\hat{\boldsymbol{\theta}}_{\mathcal{M} \cup \mathcal{S}} - N^{-1/2} \text{vec} \left[\left\{ \int_0^1 \mathbf{B}(s)\mathbf{B}(s)^\top ds \right\}^{-1} \int_0^1 \mathbf{B}(s)\mathbf{t}^\top(s) ds \right] \right). \quad (11)$$

As we will show in the next section, the test statistic T asymptotically follows central chi-square distribution with degree of freedom rN under the null hypothesis. Therefore, to conduct the hypothesis test (2) with significance level α , we reject the null hypothesis if $T > \chi_{1-\alpha, rN}^2$, where $\chi_{1-\alpha, rN}^2$ is the $(1 - \alpha)$ th quantile of the central chi-square distribution with degree of freedom rN .

3 Theoretical properties

We demonstrate the statistical consistency of $\hat{\boldsymbol{\alpha}}$ and $\hat{\boldsymbol{\Gamma}}$. This is followed by establishing the asymptotic normality of the resulting estimators. Finally, we prove that the test statistic constructed in Section 2.4 follows a chi-squared distribution.

3.1 Conditions

Before presenting the main results, we firstly list a set of necessary conditions.

(C1) Assume $c_\psi < \psi''(\cdot) < C_\psi$ and $0 < |\psi'''(\cdot)| < \infty$.

(C2) Assume for an arbitrary unit vector $\mathbf{u} = (u_1, \dots, u_d)^\top$, $\sum_{j=1}^d u_j \int_0^1 X_{ij}^2(s) ds$ is a sub-exponential distributed random variable with $\|\sum_{j=1}^d u_j \int_0^1 X_{ij}^2(s) ds\|_{\psi_1} \leq A_1$. Furthermore, $\int_0^1 \mathbf{X}_i(s) ds$ is a sub-Gaussian random vector, with $\|\int_0^1 \mathbf{u}^\top \mathbf{X}_i(s) ds\|_{\psi_2} \leq A_2$,

and \mathbf{Z}_i is a sub-Gaussian random vector with $\|\mathbf{v}^T \mathbf{Z}_i\|_{\psi_2} \leq A_3$ for any unit vector \mathbf{v} .

And $Y_i - \psi'_i\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\}$ is a mean zero sub-Gaussian random variable.

- (C3) Define the knots $\tau_{-b+1} = \dots = \tau_0 = 0 < \tau_1 < \dots < \tau_N < 1 = \tau_{N+1} = \dots = \tau_{N+b}$, where K is the number of interior knots and $[0, 1]$ is divided into $K + 1$ subintervals. Assume $\beta_j(s) \in C^\omega([0, 1])$, and there is $N = K + b$ dimensional $\boldsymbol{\gamma}_{0j}$, and a b th order Bspline with $b > \omega$ such that $\sup_{s \in [0, 1]} |\mathbf{B}^T(s) \boldsymbol{\gamma}_{0j} - \beta_j(s)| = O_p(N^{-\omega})$, $j = 1, \dots, d$. The smallest and largest distances between two knots are h_{\min}, h_{\max} , respectively. Assume $h_{\max}/h_{\min} \leq H < \infty$. Furthermore, assume $Nn^{-1/(2\omega+1)} \rightarrow \infty$.

- (C4) Assume $n \geq c_0 \log(p)$, $k_0 \leq c'_0 \log(p)$, $N \leq c'_0 \log(p)$ and $q \leq c'_0 \log(p)$ for constants $c_0, c'_0 > 0$.

- (C5) Define $\alpha_{\min}(\mathbf{M})$ and $\alpha_{\max}(\mathbf{M})$ to be the minimal and maximal eigenvalue of \mathbf{M} , respectively. Assume

$$\begin{aligned} 0 < \alpha_{\min} &\leq \alpha_{\min} \left[E \left\{ \left(\left[\mathbf{Z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} \right\} \right] \\ &\leq \alpha_{\max} \left[E \left\{ \left(\left[\mathbf{Z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} \right\} \right] \leq \alpha_{\max} < \infty. \end{aligned}$$

Condition (C1) and (C5) guarantee the boundedness and the invertibility of the second derivative of the noise free negative log likelihood. Condition (C2) implies that the integration of \mathbf{X}_j should exhibit a sub-Gaussian decay in its tail, and it also suggests that each element in \mathbf{Z}_i is a sub-Gaussian random variable. Similar requirement is also assumed in Loh & Wainwright (2012, 2015, 2017). Condition (C3) is the basic assumption to ensure the accuracy of B-spline approximation. Condition (C4) establishes the relationship between sample size, the number of unknown parameters, and the number of knots in the B-spline necessary to ensure consistent estimation.

3.2 Consistency

We first show $\widehat{\boldsymbol{\theta}}$ converges to $\boldsymbol{\theta}_0$ with high probability in Theorem 1.

Theorem 1. Denote $\widehat{\mathbf{v}} \equiv (\widehat{\mathbf{v}}_1^T, \widehat{\mathbf{v}}_{2j}^T, j = 1, \dots, d)^T \equiv \widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0 \equiv \{\widehat{\boldsymbol{\alpha}}^T - \boldsymbol{\alpha}_0^T, N^{-1/2}\text{vec}(\widehat{\boldsymbol{\Gamma}})^T - N^{-1/2}\text{vec}(\boldsymbol{\Gamma}_0)^T\}^T$, where $\widehat{\boldsymbol{\alpha}}, \widehat{\boldsymbol{\Gamma}}$ are the solution for (3). Assume Conditions (A1)–(A6) in the supplementary material and (C1)–(C5) hold and $c_\psi \alpha_{\min} > 3\mu/2$. Furthermore, assume

$$\lambda \geq \max \left\{ 2\tau_1 R \log\{\max(p, n)\}/n, C_0 \sqrt{\log\{\max(p, n)\}/n} + Ck_0^{1/2} N^{-\omega} \right\} / 4,$$

where C_0, C are positive constants defined in Lemma A.15, we have there are constants $c_1 > 0$ such that $\|\widehat{\mathbf{v}}\|_2 \leq 3\lambda\sqrt{q+m+k_0}/(c_\psi\alpha_{\min} - 3\mu/2)$, with probability greater than $1 - 2 \exp \{-c_1 n \min(c_\psi^2 \alpha_{\min}^2 / 54^2, 1) / 2\} - 20 \max(p, n)^{-1}$. And

$$\|\widehat{\mathbf{v}}_1\|_1 + \sum_{j=1}^d \|\widehat{\mathbf{v}}_{2j}\|_2 \leq 12\lambda(q+m+k_0)/(c_\psi\alpha_{\min} - 3\mu/2),$$

with probability greater than $1 - 2 \exp \{-c_1 n \min(c_\psi^2 \alpha_{\min}^2 / 54^2, 1) / 2\} - 20 \max(p, n)^{-1}$.

When choosing $\lambda = O(\log\{\max(p, n)\}/n + N^{-\omega})$ and q, m, k_0 are finite constants, Theorem 1 shows that the L_1 distance $\|\widehat{\boldsymbol{\alpha}} - \boldsymbol{\alpha}_0\|_1$ and the L_2 distance $\sum_{j=1}^d \|N^{-1/2}(\widehat{\boldsymbol{\gamma}}_j)^T - N^{-1/2}(\boldsymbol{\gamma}_{0j})\|_2$ is vanished in the rate of $O(\log\{\max(p, n)\}/n + N^{-\omega})$ with high probability.

In Theorem 2, we show that that under the condition that $(q+m+k_0)\sqrt{\log\{\max(p, n)\}/n} \rightarrow 0$ and $\lambda = O_p\{[\log\{\max(p, n)\}/n]^{1/4}\}$, minimizing the objective function (3) yields a unique solution which approaches true estimator $\boldsymbol{\theta}_0$ with high probability.

Theorem 2. Assume Conditions (A1)–(A6) in the supplementary material and (C1)–(C5) hold. Further assume that $\mu < c_\psi \alpha_{\min} / 2$ and $24\lambda(q+m+k_0)/(c_\psi\alpha_{\min} - 3\mu/2) \leq R$,

$$\lambda \geq \max \left\{ 2\tau_1 R \log(p)/n, C_0 \sqrt{\log\{\max(p, n)\}/n} + Ck_0^{1/2} N^{-\omega} \right\} / 4,$$

and

$$n \geq \max \left[8\tau_1(\log(p) + N + q) \{ \sqrt{q} + \sqrt{m} + (2/\delta + 2)\sqrt{k_0} \}^2 / (c_\psi \alpha_{\min}/2 - \mu), \right. \\ \left. \tau_1 \{ \log(p) + N + q \} (q + m + k_0) / (c_\psi \alpha_{\min}/2 - \mu) \right].$$

Furthermore, assume $(q + m + k_0)\sqrt{\log\{\max(p, n)\}/n} \rightarrow 0$, $k_0 N^{-2\omega} = o[\log\{\max(p, n)\}/n]$,

and $\lambda = O_p\{[\log\{\max(p, n)\}/n]^{1/4}\}$. Let \mathbf{E}_j be a matrix so that $\mathbf{E}_j^T(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) = \hat{\boldsymbol{\gamma}}_j - \boldsymbol{\Gamma}_{0:j}$,

assume

$$\sup_{j \in S} \sum_{k \in S} \|\mathbf{E}_j^T \{\mathbf{Q}_{\mathcal{M} \cup S, \mathcal{M} \cup S}(\boldsymbol{\theta}^*)\}^{-1} \mathbf{E}_k\|_2 \leq c_\infty.$$

Then we have $\hat{\boldsymbol{\theta}}$ is an unique minimizer for (3) and

$$\hat{\boldsymbol{\theta}}_{\mathcal{M} \cup S} - \boldsymbol{\theta}_{0, \mathcal{M} \cup S} = -\{\mathbf{Q}_{\mathcal{M} \cup S, \mathcal{M} \cup S}\{\boldsymbol{\alpha}_0, \boldsymbol{\beta}(\cdot)\}\}^{-1} \\ \times \left\{ \sum_{i=1}^n \{\psi'_i\{\boldsymbol{\alpha}_0, \boldsymbol{\beta}(s)\} - Y_i\} \left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right\}_{\mathcal{M} \cup S} \{1 + o_p(1)\},$$

and $\hat{\boldsymbol{\theta}}_{(\mathcal{M} \cup S)^c} = \mathbf{0}$ with probability greater than $1 - \exp\{-O(n)\} - O\{\max(p, n)^{-1}\}$.

The proof of the theorem follows the standard prime-dual construction introduced in Wainwright (2009) that is illustrated in Theorem A.1 in the supplementary material. We first show when $\mathcal{M} \cup S$ is given,

$$\mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) + N^{-1/2} \sum_{j \in \mathcal{M}^c} \rho_\lambda(\|\boldsymbol{\gamma}_j\|_2), \\ \text{such that } \|\boldsymbol{\alpha}\|_1 + N^{-1/2} \sum_{j=1}^d \|\boldsymbol{\gamma}_j\|_2 \leq R, \text{ and } \boldsymbol{\gamma}_j = \mathbf{0} \text{ for } j \in (\mathcal{M} \cup S)^c,$$

has an unique local minimizer in the interior of the feasible set. Then we show that all the functional stationary points of (3) must have support in $\mathcal{M} \cup S$. Therefore, the local minimizers of (3) are actually the local minimizers of the above oracle loss function, so are also unique.

Remark 1. Let \mathbf{E}_0 be the matrix so that $\mathbf{E}_0^T(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) = \hat{\boldsymbol{\alpha}} - \boldsymbol{\alpha}_0$ and \mathbf{E}_j be the matrix so that $\mathbf{E}_j^T(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) = N^{-1/2}\hat{\boldsymbol{\gamma}}_j - N^{-1/2}\boldsymbol{\Gamma}_{0j}$. Theorem 2 leads to the asymptotic normality of the estimator $\hat{\boldsymbol{\theta}}_{\mathcal{M} \cup \mathcal{S}}$ that

$$\sqrt{n}(\hat{\boldsymbol{\alpha}} - \boldsymbol{\alpha}_0) \sim N(0, \mathbf{E}_0^T \boldsymbol{\Psi} \mathbf{E}_0). \quad (12)$$

Furthermore, for each s ,

$$N^{-1/2}n^{1/2}\{\mathbf{B}(s)^T \hat{\boldsymbol{\gamma}}_j - \beta_j(s)\} = O_p[N^{-1/2}n^{1/2}\{\mathbf{B}(s)^T \hat{\boldsymbol{\gamma}}_j - \mathbf{B}(s)^T \boldsymbol{\gamma}_{0j}\}].$$

This holds because $N^{-1/2}n^{1/2}\{\mathbf{B}(s)^T \hat{\boldsymbol{\gamma}}_j - \mathbf{B}(s)^T \boldsymbol{\gamma}_{0j}\} = O_p(1)$ similar to the results in Jiang et al. (2015) and $N^{-1/2}n^{1/2}\{\mathbf{B}(s)^T \boldsymbol{\gamma}_{0j} - \beta_j(s)\} = O(N^{-1/2-\omega}n^{1/2}) = o_p(1)$ by Condition (C3). Therefore,

$$N^{-1/2}n^{1/2}\{\mathbf{B}(s)^T \hat{\boldsymbol{\gamma}}_j - \beta_j(s)\} \sim N\{0, \mathbf{B}(s)^T \mathbf{E}_j^T \boldsymbol{\Psi} \mathbf{E}_j \mathbf{B}(s)\}. \quad (13)$$

We then can construct the confidence intervals (CIs) for $\boldsymbol{\alpha}_0$ and the point-wise CI for $\beta_j(s)$ based on (12) and (13)

We now show that the test statistic T in (11) is asymptotically chi-square distributed under both null and alternative hypotheses.

Define

$$\begin{aligned} \boldsymbol{\omega}_n &= -\sqrt{n} \mathbf{A} \mathbf{Q} \{\boldsymbol{\alpha}_0, \boldsymbol{\beta}(\cdot)\}_{\mathcal{M} \cup \mathcal{S}, \mathcal{M} \cup \mathcal{S}}^{-1} \\ &\quad \times \left\{ n^{-1} \sum_{i=1}^n \{\psi'_i \{\boldsymbol{\alpha}_0, \boldsymbol{\beta}(s)\} - Y_i\} \left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right\}_{\mathcal{M} \cup \mathcal{S}}, \end{aligned}$$

and define $\mathbf{c}_n = \text{vec} \left[\sqrt{n} \boldsymbol{\Psi}^{-1/2}(\boldsymbol{\Sigma}, \mathbf{Q}, \boldsymbol{\beta}) N^{-1/2} \left\{ \int_0^1 \mathbf{B}(s) \mathbf{B}(s)^T ds \right\}^{-1} \int_0^1 \mathbf{B}(s) \mathbf{h}_n^T(s) ds \right]$. Let $T_0 = \boldsymbol{\Psi}^{-1/2}(\boldsymbol{\Sigma}, \mathbf{Q}, \boldsymbol{\beta})(\boldsymbol{\omega}_n + \mathbf{c}_n)$.

We first construct the asymptotic distribution of T_0 , Then we show that the test statistic T is asymptotically equivalent to T_0 .

Lemma 1. *Assume Conditions (C1) and (C5) holds. Furthermore, assume $(rN)^{1/4}n^{-1/2} \rightarrow 0$ and*

$$E \left[\left\| \Psi^{-1/2}(\boldsymbol{\Sigma}, \mathbf{Q}, \boldsymbol{\beta}) \mathbf{A} \mathbf{Q}_{\mathcal{M} \cup \mathcal{S}, \mathcal{M} \cup \mathcal{S}}^{-1} \{ \boldsymbol{\alpha}_0, \boldsymbol{\beta}(\cdot) \}_{\mathcal{M} \cup \mathcal{S}, \mathcal{M} \cup \mathcal{S}}^{-1} \right. \right. \\ \left. \left. \times \{ \psi'_i \{ \boldsymbol{\alpha}_0, \boldsymbol{\beta}(s) \} - Y_i \} \left[\mathbf{Z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]_{\mathcal{M} \cup \mathcal{S}}^T \right\|_2^3 \right] = O(1).$$

We have $\lim_{n \rightarrow \infty} | \Pr(T_0 \leq x) - \Pr \{ \chi^2(rN, \mathbf{c}_n^T \boldsymbol{\Psi}^{-1}(\boldsymbol{\Sigma}, \mathbf{Q}, \boldsymbol{\beta}) \mathbf{c}_n) \leq x \} | = 0$, where $\chi^2(r, \gamma)$ is a r degree of freedom non-central chi-square distribution, with non-centrality parameter γ .

In Lemma 1, we show that the test statistic T_0 is asymptotically chi-square distributed under both null and alternative hypotheses with different non-central parameters. Note that $\mathbf{h}_n(s) = 0$ under the null hypothesis, and therefore the non-central parameter satisfies $\mathbf{c}_n^T \boldsymbol{\Psi}^{-1}(\boldsymbol{\Sigma}, \mathbf{Q}, \boldsymbol{\beta}) \mathbf{c}_n = 0$. We then show that the test statistic T is asymptotically equivalent to T_0 in Theorem 3.

Theorem 3. *Assume Conditions (A1)–(A6) in the supplementary material and (C1)–(C5) hold. Furthermore, assume the conditions in Theorem 2 hold, we have $T - T_0 = o(rN)$. In addition, assume $(rN)^{1/4}n^{-1/2} \rightarrow 0$ and*

$$E \left[\left\| \Psi^{-1/2}(\boldsymbol{\Sigma}, \mathbf{Q}, \boldsymbol{\beta}) \mathbf{A} \mathbf{Q}_{\mathcal{M} \cup \mathcal{S}, \mathcal{M} \cup \mathcal{S}}^{-1} \{ \boldsymbol{\alpha}_0, \boldsymbol{\beta}(\cdot) \}_{\mathcal{M} \cup \mathcal{S}, \mathcal{M} \cup \mathcal{S}}^{-1} \right. \right. \\ \left. \left. \times \{ \psi'_i \{ \boldsymbol{\alpha}_0, \boldsymbol{\beta}(s) \} - Y_i \} \left[\mathbf{Z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]_{\mathcal{M} \cup \mathcal{S}}^T \right\|_2^3 \right] = O(1).$$

Then $\lim_{n \rightarrow \infty} | \Pr(T \leq x) - \Pr \{ \chi^2(rN, \mathbf{c}_n^T \boldsymbol{\Psi}^{-1}(\boldsymbol{\Sigma}, \mathbf{Q}, \boldsymbol{\beta}) \mathbf{c}_n) \leq x \} | = 0$, where $\chi^2(r, \gamma)$ is a r degree of freedom non-central chi-square distribution, with non-centrality parameter γ .

Theorem 3 shows that the test statistic T is asymptotically $\chi^2(rN)$ under the null hypothesis. Therefore, if we reject the null hypothesis if $T > \chi_{1-\alpha, rN}^2$, we can preserve the type I error under α nominal level asymptotically.

4 Numerical studies

4.1 Simulation configuration

We carry out comprehensive numerical analyses to assess both the theoretical attributes and the performance of the proposed HDFP method. Data are generated using the model specified in (1), with Gaussian and Bernoulli distributions corresponding to the linear and logistic model for the continuous and binary outcomes, respectively. For the linear model, random errors are drawn from either a standard normal distribution or a t -distribution with three degrees of freedom, and these errors are subsequently standardized to have a variance of one. For the logistic model, we model Y given $\alpha_0^T \mathbf{Z}_i + \int_0^1 \boldsymbol{\beta}^T(s) \mathbf{X}(s) ds$ as following a Bernoulli distribution.

The $\boldsymbol{\beta}(s)$ is constructed using Fourier basis following Xue & Yao (2021). Specifically, we set

$$\beta_j(s) = c_j \sum_{k=1}^{50} \eta_k \phi_k(s), \eta_k = I_{\{k \leq 4\}}(1.2 - 0.2k) + 0.4 I_{\{5 \leq k \leq 50\}}(k - 3)^{-4}, \text{ for } k = 1, \dots, 50,$$

where $I_{\{\cdot\}}$ is an indicator function, $c_j = 0$ for $j = 3, \dots, d-1$ and c_j is chosen differently for each j with $j = 1, 2, d$ under various settings to control the signal-to-noise ratio and $\phi_k(s)$'s are the Fourier basis functions. Specifically, $\phi_1(s) = 1$, $\phi_{2k}(s) = \sqrt{2} \cos \{k\pi(2s - 1)\}$ for $1 \leq k \leq 25$ and $\phi_{2k-1}(s) = \sqrt{2} \sin \{(k - 1)\pi(2s - 1)\}$ for $2 \leq k \leq 25$. We consider two types of functional predictors, one is the uncorrelated functional predictors generated based on the B-spline basis and the other is highly-correlated functional predictors generated with the spectral graph model (SGM) (Verma et al. 2022) to mimic the real PSD data in our motivating example.

The uncorrelated $\mathbf{X}(s)$ is generated as follows,

$$X_j(s) = \sum_{k=1}^{10} \xi_{jk} B_k(s) + 0.5 \epsilon_j(s), \quad (14)$$

where ξ_{jk} is generated from $N(0, 5^2)$, $B_k(s)$ is the k th B-spline basis function and $\epsilon_j(s)$ is the standard Gaussian process. Here $\epsilon_j(s)$ and ξ_{jk} are independent for different j . Therefore, the correlation between different functional predictors is zero.

The highly-correlated $\mathbf{X}(s)$ is generated from the SGM, which simulates the PSD of the human brain activity at the resting state through Verma et al. (2022). After creating the PSDs from the SGM, we centered the PSD at each ROI and added Gaussian noise with a mean of 0 and a variance of 100 to the PSD to mimic real data. The detail of generating the PSDs is presented in Verma et al. (2022). The resulting PSDs have strong spatially correlation with average pair-wise correlation coefficient around 0.6 empirically.

Unless stated otherwise, $\boldsymbol{\alpha}_0 = [\alpha_{00}, -1, 2]$ under both the linear and logistic models. For the linear model, the intercept α_{00} is 5. For the logistic model, α_{00} is adaptively adjusted to ensure the probability of success remains at 0.5. The two baseline covariates in \mathbf{Z} are derived respectively from the standard normal distribution and a Bernoulli distribution with a probability of 0.5.

In the implementation, the number of B-spline bases (N and the penalty parameter λ are optimized using a 10-fold cross-validation (CV) process aimed at minimizing the prediction error. The number of B-spline bases is chosen from the set $[4, 6, 8, 10, 12]$. We select λ from $[0, 1.5]$ with increments of 0.1 under the linear model and from $(0, 0.05, 0.1, 0.2, 0.3, 0.4)$ under the logistic model. The parameter a in the SCAD penalty is set to be 3.7 as recommended by Fan & Song (2010), and $\alpha = 0.9$, $\beta \in [0.5, 10]$ to ensure rapid convergence according to He et al. (2014). We set $R = 2e5$ to guarantee it to be larger than $\|\boldsymbol{\alpha}_0\|_1 + 2 \sum_{j=1}^d \|N^{-1/2} \boldsymbol{\gamma}_{0j}\|_2$. The CPRSM algorithm terminates when the norm of the difference between the parameters in two consecutive iterations is less than $5e-4$ or when the number of iterations reaches 2000. All results in this section are based on 1000 repetitions.




Figure 2: **A:** $\|\hat{\alpha} - \alpha_0\|_2$ for the coefficients of the baseline covariates when the sample size n varies under the linear and logistic scenarios. The shadow area represents the 25% and 75% quantiles of the estimated error over 1000 simulations. **B and C:** Empirical coverage probabilities of the 95% confidence interval of α_{01} and α_{02} , respectively under linear and logistic models.

4.2 Evaluate the asymptotic property of α

We use the asymptotic normality outlined in Remark 1 to construct 95% CIs for the coefficients of the baseline covariates α . We validate the empirical coverage of these theoretical CIs for the coefficients of the baseline covariates α and confirm their consistency. We choose $d = 200$ and vary the sample size n from 100 to 3200. The data are derived from a linear model with standard normal random errors and a logistic model. The uncorrelated functional predictors are generated as specified in (14). Additionally, we set $c_j = 2$ for $j = d$ and $c_j = 0$ for all other values in (14) to generate the functional coefficients. In this scenario, only the last functional covariate is associated with the outcome. Since only α is the parameter of interest, we apply penalties to all the functional coefficients in the estimation.

For each given sample size, we repeat the simulation 1000 times and the results are

shown in Figure 2, where we report both the average estimation error $\|\hat{\boldsymbol{\alpha}} - \boldsymbol{\alpha}_0\|_2$ and the empirical coverage of the 95% CI for $\boldsymbol{\alpha}_0$ over the 1000 simulations. Figure 2A shows that the average estimation error decreases when the sample size increases, which validates the consistency of the estimator. Figures 2B and 2C illustrate that the CI coverage probabilities for α_{01} and α_{02} in the linear model are close to 95% across all evaluated sample sizes. Additionally, the coverage probability approaches 95% more closely as the sample size increases. A similar trend is observed under the logistic model, although the coverage probability only reaches 95% when the sample size exceeds 300. This suggests that the estimation is more efficient under the linear model.

4.3 Evaluate the hypothesis testing procedure

We then evaluate the performance of HDFP in testing linear functional hypotheses. We compare our method with the bootstrap method proposed by Xue & Yao (2021). Note that the bootstrap method is developed under the linear model with no baseline covariate, and therefore, our comparative study is conducted without baseline covariates. All the tuning parameters in the bootstrap method are chosen through the 10-fold CV.

We choose $d = 200$ and sample size $n = 100$. We generate the uncorrelated functional predictor from (14) and correlated predictor from the SGM as described in Section 4.1. We choose $c_d = 1$ and varying c_1 and c_2 in (14) under different simulation settings. Specifically, we aim to test the null hypotheses that $\mathbf{H}_{0,1} : \beta_1(s) = 0$ and $\mathbf{H}_{0,2} : \beta_1(s) = \beta_2(s) = 0$. To assess the performance for testing $\mathbf{H}_{0,1}$, we generate functional parameters according to (14) with $c_1 = c$ and $c_2 = 0$, varying c from 0 to 0.4. To evaluate the performance for testing $\mathbf{H}_{0,2}$, we generate functional parameters from (14) with both $c_1 = c_2 = c$, varying c from 0 to 0.4.

Table 1: The empirical size and power of hypothesis testing for the comparison between HDFP and bootstrap methods under the linear model with normal noise and t noise based on 1000 repetitions. We choose number of functional covariates $d = 200$, sample size $n = 100$ and significance level $\alpha = 0.05$. $c = 0$ indicates the null hypothesis is true. Err. dist. standards for the distribution of the random error. Type I errors are in **bold**.

Type of $\mathbf{X}(s)$	Err. dist.	c	HDFP	Bootstrap	HDFP	Bootstrap
			$\beta_1(s) = 0$		$\beta_1(s) = \beta_2(s) = 0$	
Uncorrelated	$N(0, 1)$	0.0	0.045	0.047	0.050	0.049
		0.1	0.115	0.115	0.167	0.099
		0.2	0.377	0.361	0.637	0.388
		0.4	0.862	0.930	0.975	0.854
	$t(3)$	0.0	0.038	0.048	0.044	0.035
		0.1	0.134	0.118	0.154	0.099
		0.2	0.403	0.370	0.615	0.379
		0.4	0.909	0.933	0.941	0.915
Highly-correlated (PSD)	$N(0, 1)$	0.0	0.038	0.036	0.040	0.024
		0.1	0.114	0.037	0.150	0.035
		0.2	0.187	0.048	0.496	0.063
		0.4	0.624	0.051	0.935	0.055
	$t(3)$	0.0	0.026	0.027	0.038	0.019
		0.1	0.066	0.030	0.130	0.027
		0.2	0.176	0.051	0.304	0.020
		0.4	0.628	0.144	0.944	0.052

We show the size and power of testing $\mathbf{H}_{0,1}$ and $\mathbf{H}_{0,2}$ in Table 1 under various settings. Table 1 shows that when the functional predictors are uncorrelated, both HDFP and the bootstrap method control the type I error ($c = 0$) when testing $\mathbf{H}_{0,1}$, while HDFP and the bootstrap yield similar powers under the alternative hypotheses when $c \neq 0$. Furthermore, when the hypothesis involve two functional parameters as those in $\mathbf{H}_{0,2}$, HDFP yields greater power than the bootstrap method does. When the functional predictors are highly-correlated, both methods control the type I error under the nominal level. However, the

Table 2: The empirical size and power of hypothesis testing for the HDFP under the linear and logistic models based on 1000 repetitions. The set of interested functional coefficients is $\beta_1(s)$. We choose number of functional covariates $d = 68$, sample size $n = 200$ and significance level $\alpha = 0.05$. The type I errors are in **bold**.

\mathbf{H}_0	c	$n = 200$			$n = 500$		
		$N(0, 1)$	$t(3)$	Logistic	$N(0, 1)$	$t(3)$	Logistic
$\mathbf{H}_{0,1}^1$	0.0	0.054	0.051	0.046	0.044	0.048	0.032
	0.1	0.649	0.732	0.149	0.998	0.999	0.180
	0.2	0.997	1.000	0.314	1.000	1.000	0.594
	0.4	1.000	1.000	0.794	1.000	1.000	1.000
$\mathbf{H}_{0,2}^1$	0.0	0.555	0.645	0.116	0.997	0.996	0.197
	0.1	0.054	0.053	0.064	0.041	0.049	0.038
	0.2	0.640	0.730	0.103	0.997	0.999	0.138
	0.4	1.000	1.000	0.486	1.000	1.000	0.884
$\mathbf{H}_{0,3}^1$	0.0	0.996	1.000	0.383	1.000	1.000	0.659
	0.1	0.573	0.638	0.155	0.997	0.996	0.169
	0.2	0.053	0.049	0.054	0.043	0.043	0.059
	0.4	0.998	1.000	0.209	1.000	1.000	0.449

bootstrap method only achieves a power of less than 0.15 in all scenarios. The results indicate that the HDFP method outperforms the bootstrap method, particularly when the functional predictors are highly correlated.

4.4 Simulation evaluates the testing procedure under the real data setting

We assess the performance of the HDFP method in a more realistic setting when the distributions of the functional predictors mimic those in the real data. Therefore, we choose $d = 68$ and consider $n = 200$ and $n = 500$. Furthermore, we generate a continuous and a binary baseline covariate from the standard normal distribution and the Bernoulli

distribution with probability 0.5, respectively.

The baseline effect is $\boldsymbol{\alpha}_0 = [5, -1, 2]$ under the linear model. Whereas, the baseline effect is $\boldsymbol{\alpha}_0 = [\alpha_{00}, -1, 2]$ for logistic model where α_{00} is adaptively chosen in each scenario to ensure that the positive and negative samples remain balanced. The functional predictors are generated by the SGM (Verma et al. 2022), and the functional coefficients $\boldsymbol{\beta}(s)$ are generated from the Fourier basis as described in (14) with $c_d = 2$, $c_j = 0$ for $j \neq 1, 2, d$. We vary (c_1, c_2) under different simulation settings.

We consider two sets of hypothesis testings, where we aim to test

$$\mathbf{H}_{0,1}^1 : \beta_1(s) = 2c \sum_{k=1}^{50} \eta_k \phi_k(s), c = 0; \quad \mathbf{H}_{0,1}^2 : \beta_1(s) - \beta_2(s) = 2c \sum_{k=1}^{50} \eta_k \phi_k(s), c = 0.$$

$$\mathbf{H}_{0,2}^1 : \beta_1(s) = 2c \sum_{k=1}^{50} \eta_k \phi_k(s), c = 0.1; \quad \mathbf{H}_{0,2}^2 : \beta_1(s) - 0.8\beta_2(s) = (0.2c + 0.2) \sum_{k=1}^{50} \eta_k \phi_k(s), c = 0.$$

$$\mathbf{H}_{0,3}^1 : \beta_1(s) = 2c \sum_{k=1}^{50} \eta_k \phi_k(s), c = 0.2; \quad \mathbf{H}_{0,3}^2 : 4\beta_1(s) - 3\beta_2(s) = (8c + 1) \sum_{k=1}^{50} \eta_k \phi_k(s), c = 0.$$

We set the significance level $\alpha = 0.05$ and repeat the simulation 1000 times. Table 2 shows the results of testing hypotheses $\mathbf{H}_{0,1}^1$ to $\mathbf{H}_{0,3}^1$, where only one parameter is involved in the test. HDFP effectively controls the type I errors under the linear model. However, it shows slightly inflated type I error rates when testing $\mathbf{H}_{0,3}^1$ under the logistic model, which is likely a finite sample phenomenon. Furthermore, the power increases as the discrepancy between the true value of c and its value under the null hypothesis increases. Moreover, a larger sample size also results in a higher power. Table 3 shows the results from testing the hypotheses $\mathbf{H}_{0,1}^2$ to $\mathbf{H}_{0,3}^2$ that involve the linear combinations of two parameters. The type I errors are well controlled when the sample size reaches 500 in all scenarios. Furthermore, the power increases when c deviates from 0, and when the sample size increases. In summary, the simulation studies demonstrate that HDFP has satisfactory performance in testing linear hypotheses under both linear and logistic models.

Table 3: The empirical size and power of linear hypothesis testing for the HDFP under the linear and logistic models based on 1000 repetitions. The functional parameters of interested are $\beta_1(s), \beta_2(s)$. We choose number of functional covariates $d = 68$, sample size $n = 200$ and significance level $\alpha = 0.05$. The value of c corresponding to the true null hypothesis is in **bold**.

\mathbf{H}_0	c	$n = 200$			$n = 500$		
		$N(0, 1)$	$t(3)$	Logistic	$N(0, 1)$	$t(3)$	Logistic
$\mathbf{H}_{0,1}^2$	0.0	0.054	0.063	0.041	0.046	0.049	0.051
	0.1	0.248	0.251	0.066	0.833	0.811	0.093
	0.2	0.849	0.858	0.088	1.000	1.000	0.202
	0.4	1.000	1.000	0.227	1.000	1.000	0.608
$\mathbf{H}_{0,2}^2$	0.0	0.052	0.063	0.041	0.048	0.047	0.051
	0.1	0.311	0.323	0.059	0.916	0.893	0.080
	0.2	0.921	0.923	0.062	1.000	1.000	0.180
	0.4	1.000	1.000	0.161	1.000	1.000	0.607
$\mathbf{H}_{0,3}^2$	0.0	0.053	0.062	0.040	0.047	0.049	0.050
	0.1	0.336	0.345	0.057	0.930	0.919	0.075
	0.2	0.929	0.934	0.063	1.000	1.000	0.178
	0.4	1.000	1.000	0.149	1.000	1.000	0.600

5 Real data analysis

5.1 Data preprocessing

We apply HDFP to analyze our AD data, which includes 85 AD patients and 61 healthy subjects. The dataset includes the PSDs evaluated at 74 continuous frequencies of the MEG source time series at 68 ROIs, where the detailed preprocessing procedure is presented in Appendix A of the supplementary material. The outcome is the MMSE score that assesses cognitive impairment. The lower the score, the worse the cognitive function (Arevalo-Rodriguez et al. 2021). In addition, we consider three baseline covariates, gender, age and number of years of education.

5.2 Comparison with other methods

We apply the HDFP method to study the association between the PSDs and the MMSE adjusted for baseline covariates and test the null hypothesis that $\mathbf{H}_0 : \beta_j(s) = 0$ for $j = 1, \dots, 68$. The significant level are set as $\alpha = 0.05/68$ to account for the multiple testing. All implementation details are the same as in the Section 4.1, except that λ is chosen in the set of $\{0.1, 0.3, 0.5, 0.7, 1.1, 1.3\}$ through CV.

We include the bootstrap method proposed by Xue & Yao (2021) as the benchmark. Since the bootstrap method does not account for the baseline covariates, we employ a projection step to eliminate the influence of baseline covariates. Specifically, consider $\tilde{\mathbf{Z}}$ as an $n \times (p+1)$ matrix where the first column consists of ones and the remaining columns represent the baseline covariates. Let $\mathbf{P} = \mathbf{I} - \tilde{\mathbf{Z}}(\tilde{\mathbf{Z}}^T \tilde{\mathbf{Z}})^{-1} \tilde{\mathbf{Z}}^T$ and $\mathbf{Y} = (Y_i, i = 1, \dots, n)^T$. We generate the new outcomes as $\mathbf{P}\mathbf{Y}$ and the new functional predictors as $\mathbf{P}\{\mathbf{X}_1(t), \dots, \mathbf{X}_n(t)\}^T$. Then we implement the bootstrap method considering the new outcomes and the new functional predictors. Here we remove the covariate effect because $\mathbf{P}\tilde{\mathbf{Z}} = \mathbf{0}$.

We demonstrate significant ROIs selected via HDFP with the p-values less than 0.05/68 in Figure 3A. We do not present the results from the bootstrap method as no significant ROIs are identified after applying the Bonferroni correction. Overall, the significant ROIs identified by HDFP exhibit high symmetry between the left and right hemispheres, which is intuitive and consistent with the existing literature (Jagust 2018). Most of the significant ROIs are located in the inferior and posterior temporal cortices and the posterior parietal-occipital cortices, which reflects the similar regional distribution of the glucose hypometabolism and the high tau accumulation in the AD brain (Jagust 2018, Jin et al. 2023).

We further analyze the accuracies of HDFP and the bootstrap method in classifying AD




Figure 3: **A:** Negative log p-values from the significant ROIs selected via HDFP. Only the significant regions with p-value less than $0.05/68$ are shown. **B:** The receiver operating characteristic curves and corresponding AUCs for HDFP and the bootstrap method in distinguishing healthy subjects with AD patients.

and healthy controls based on their chosen significant ROIs that associated with the MMSE score. In evaluating HDFP, we construct the logistic model between the disease indicator and the PSDs from the significant ROIs, adjusting for the baseline covariates. We minimize (3) for the parameter estimation and apply penalties to all the functional predictors. In evaluating the bootstrap method, given that it fails to identify any significant PSDs, we proceed with a standard logistic regression between the disease indicator and the baseline covariates. We compare the area under the receiver operating characteristic (ROC) curve (AUC) of both methods. Figure 3 illustrates that HDFP achieves better classification accuracy, indicating that the PSDs from selected ROIs indeed offer valuable information for distinguishing AD from control samples.

We further conduct subsequent analyses on the HDFP function estimators. We establish point-wise CIs at the $100(1 - 0.05/68)\%$ level for each functional coefficient across the 68 ROIs. If the CI for a specific frequency point does not cover zero, the functional effect




Figure 4: The percentage of the significant ROIs based on the point-wise $100(1-0.05/68)\%$ confidence intervals for each frequency point among the 68 ROIs via HDFP method.

is considered as significantly different from zero at that frequency. We then draw the percentage of the significant ROIs in across frequencies in Figure 4. Figure 4 shows that the majority of significant regions are concentrated in the theta and low-beta ($[12, 20]$ Hz) frequency bands. This suggests the theta and low-beta bands could play a crucial role in cognitive function, consistent with the findings in Ranasinghe et al. (2022).

Furthermore, as shown in Figure 3A, the significant regions are not completely symmetric. This raise a question of whether the functional pattern is asymmetric in affecting the cognitive function. To explore this, we further examine the hypothesis regarding whether the effects from the PSDs derived from the left and right hemispheres are equal. We categorize the significant ROIs into two groups: Group 1 consists of the ROIs that are significant in both the left and right hemispheres, while Group 2 comprises ROIs that are significant in only one hemisphere. We denote \mathcal{G}_1 be the set of indices of the ROIs in Group 1 and \mathcal{G}_2 be the set of indices of the ROIs in Group 2. If $\beta_j(s)$ is the functional coefficient corresponding to the j th ROI in one hemisphere, we denote $\tilde{\beta}_j(s)$ as the functional coefficient corresponding to the j th ROI in the other hemisphere.

We then test the two null hypotheses that $\mathbf{H}_{0,1}^r : \tilde{\beta}_j(s) - \beta_j(s) = 0$ for all $j \in \mathcal{G}_1$ and

$\mathbf{H}_{0,2}^r : \tilde{\beta}_j(s) - \beta_j(s) = 0$ for all $j \in \mathcal{G}_2$. All the implementation details are the same as in the Section 5.2. In $\mathbf{H}_{0,1}^r$ where ROIs are significant in both hemispheres, we obtain test statistics $T = 24.418$ with pvalue 0.418, showing that no significant difference between the left and right hemispheres. In $\mathbf{H}_{0,2}^r$ where ROIs are significant in only one hemisphere, we obtain test statistics $T = 60.248$ with pvalue 3.786×10^{-4} , indicating a significant difference between the left and right hemispheres in at least one ROI. This analysis suggests that there are indeed asymmetry in the PSD effects from a subset of ROIs.

Finally, we summarize the effects from the baseline covariates on the MMSE score in Table 4, which shows that the education level is a strong predictor on the cognitive function, with higher education correlating to higher MMSE scores on average. It indicates that an additional year of education is associated with an increase in the MMSE score, and thus lower probability of suffering from AD, which is consistent with the existing literature (Sando et al. 2008).

Table 4: The estimated coefficients of the baseline covariates and the corresponding 95% confidence intervals after applying the HDFP to the MEG data.

Baseline covariate	Effect	95% confidence interval
Gender	-0.502	[-1.920, 0.916]
Age	0.388	[-0.241, 1.017]
Years of education	0.417	[0.155, 0.679]

6 Conclusion and discussion

We introduce the novel HDFP method for testing linear hypotheses regarding functional parameters under the generalized functional linear model. We develop a regularization technique for estimating functional parameters and validate the asymptotic properties of

the estimators. Moreover, we design a hypothesis testing procedure capable of assessing local alternative hypotheses. We apply the proposed method to analyze an AD MEG study. Our method achieves higher interpretability and the prediction accuracy than the existing method. HDFP is generalizable to study other functional data, such as fMRI data and EEG data. Further research along this line is ongoing in our group.

When presenting a large number of functional predictors, one can reduce the computational complexity by preprocessing the data using a screening method (Fan & Li 2001, Fan & Lv 2008). We have conducted preliminary studies on adopting a screening preprocessing step prior to implementing HDFP. These studies suggest that such preprocessing not only significantly reduces computational complexity but also preserves the error rates in the hypothesis testing.

The programming code for the simulation studies and the real data analysis is included in the supplementary material.

References

- Arevalo-Rodriguez, I., Smailagic, N., Roqué-Figuls, M., Ciapponi, A., Sanchez-Perez, E., Giannakou, A., Pedraza, O. L., Cosp, X. B. & Cullum, S. (2021), ‘Mini-mental state examination (mmse) for the early detection of dementia in people with mild cognitive impairment (mci)’, *Cochrane Database of Systematic Reviews* (7).
- Bosco, F., Guarnieri, L., Rania, V., Palma, E., Citraro, R., Corasaniti, M. T., Leo, A. & De Sarro, G. (2023), ‘Antiseizure medications in Alzheimer’s disease from preclinical to clinical evidence’, *International Journal of Molecular Sciences* **24**(16), 12639.
- Cai, T. T. & Yuan, M. (2012), ‘Minimax and adaptive prediction for functional linear

- regression’, *Journal of the American Statistical Association* **107**(499), 1201–1216.
- Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., Buckner, R. L., Dale, A. M., Maguire, R. P., Hyman, B. T. et al. (2006), ‘An automated labeling system for subdividing the human cerebral cortex on mri scans into gyral based regions of interest’, *NeuroImage* **31**(3), 968–980.
- Duchi, J., Shalev-Shwartz, S., Singer, Y. & Chandra, T. (2008), Efficient projections onto the l_1 -ball for learning in high dimensions, *in* ‘Proceedings of the 25th international conference on Machine learning’, pp. 272–279.
- Fan, J. & Li, R. (2001), ‘Variable selection via nonconcave penalized likelihood and its oracle properties’, *Journal of the American statistical Association* **96**(456), 1348–1360.
- Fan, J. & Lv, J. (2008), ‘Sure independence screening for ultrahigh dimensional feature space’, *Journal of the Royal Statistical Society Series B: Statistical Methodology* **70**(5), 849–911.
- Fan, J. & Song, R. (2010), ‘Sure independence screening in generalized linear models with np-dimensionality’, *The Annals of Statistics* **38**(6), 3567–3604.
- Fan, Y., James, G. M. & Radchenko, P. (2015), ‘Functional additive regression’, *The Annals of Statistics* **43**(5), 2296–2325.
- Fletcher, R. & Watson, G. A. (1980), ‘First and second order conditions for a class of nondifferentiable optimization problems’, *Mathematical Programming* **18**(1), 291–307.
- Gertheiss, J., Maity, A. & Staicu, A.-M. (2013), ‘Variable selection in generalized functional linear models’, *Stat* **2**(1), 86–101.

- He, B., Liu, H., Wang, Z. & Yuan, X. (2014), ‘A strictly contractive peaceman–rachford splitting method for convex programming’, *SIAM Journal on Optimization* **24**(3), 1011–1040.
- Hilgert, N., Mas, A. & Verzelen, N. (2013), ‘Minimax adaptive tests for the functional linear model’, *The Annals of Statistics* pp. 838–869.
- Jagust, W. (2018), ‘Imaging the evolution and pathophysiology of Alzheimer disease’, *Nature Reviews Neuroscience* **19**(11), 687–700.
- Jiang, F., Ma, Y. & Wang, Y. (2015), ‘Fused kernel-spline smoothing for repeatedly measured outcomes in a generalized partially linear model with functional single index’, *Annals of statistics* **43**(5), 1929.
- Jin, H., Ranasinghe, K. G., Prabhu, P., Dale, C., Gao, Y., Kudo, K., Vessel, K., Raj, A., Nagarajan, S. S. & Jiang, F. (2023), ‘Dynamic functional connectivity MEG features of Alzheimer’s disease’, *NeuroImage* **281**, 120358.
- Kong, D., Xue, K., Yao, F. & Zhang, H. H. (2016), ‘Partially functional linear regression in high dimensions’, *Biometrika* **103**(1), 147–159.
- Kurlowicz, L. & Wallace, M. (1999), ‘The mini-mental state examination (MMSE)’, *Journal of Gerontological Nursing* **25**(5), 8–9.
- Lei, J. (2014), ‘Adaptive global testing for functional linear models’, *Journal of the American Statistical Association* **109**(506), 624–634.
- Lian, H. (2013), ‘Shrinkage estimation and selection for multiple functional regression’, *Statistica Sinica* pp. 51–74.

- Loh, P.-L. & Wainwright, M. J. (2012), ‘High-dimensional regression with noisy and missing data: provable guarantees with nonconvexity’, *The Annals of Statistics* **40**(3), 1637–1664.
- Loh, P.-L. & Wainwright, M. J. (2015), ‘Regularized m-estimators with nonconvexity: Statistical and algorithmic theory for local optima’, *Journal of Machine Learning Research* **16**, 559–616.
- Loh, P.-L. & Wainwright, M. J. (2017), ‘Support recovery without incoherence: A case for nonconvex regularization’, *The Annals of Statistics* **45**(6), 2455–2482.
- Ranasinghe, K. G., Verma, P., Cai, C., Xie, X., Kudo, K., Gao, X., Lerner, H., Mizuiri, D., Strom, A., Iaccarino, L. et al. (2022), ‘Altered excitatory and inhibitory neuronal subpopulation parameters are distinctly associated with tau and amyloid in Alzheimer’s disease’, *Elife* **11**, e77850.
- Sando, S. B., Melquist, S., Cannon, A., Hutton, M., Sletvold, O., Saltvedt, I., White, L. R., Lydersen, S. & Aasly, J. (2008), ‘Risk-reducing effect of education in Alzheimer’s disease’, *International Journal of Geriatric Psychiatry* **23**(11), 1156–1162.
- Shang, Z. & Cheng, G. (2015), ‘Nonparametric inference in generalized functional linear models’, *The Annals of Statistics* pp. 1742–1773.
- Shi, C., Song, R., Chen, Z. & Li, R. (2019), ‘Linear hypothesis testing for high dimensional generalized linear models’, *Annals of statistics* **47**(5), 2671.
- Verma, P., Nagarajan, S. & Raj, A. (2022), ‘Spectral graph theory of brain oscillations—revisited and improved’, *NeuroImage* **249**, 118919.
- Wainwright, M. J. (2009), ‘Sharp thresholds for high-dimensional and noisy sparsity recovery

- ery using l1 constrained quadratic programming lasso', *IEEE Transactions on Information Theory* **55**(5), 2183–2202.
- Wang, R., Wang, J., Yu, H., Wei, X., Yang, C. & Deng, B. (2015), 'Power spectral density and coherence analysis of Alzheimer's EEG', *Cognitive Neurodynamics* **9**, 291–304.
- Xue, K., Yang, J. & Yao, F. (2023), 'Optimal linear discriminant analysis for high-dimensional functional data', *Journal of the American Statistical Association* pp. 1–10.
- Xue, K. & Yao, F. (2021), 'Hypothesis testing in large-scale functional linear regression', *Statistica Sinica* **31**(2), 1101–1123.
- Yuan, M. & Cai, T. T. (2010), 'A reproducing kernel hilbert space approach to functional linear regression', *The Annals of Statistics* **38**(6), 3412–3444.
- Yuan, M. & Lin, Y. (2006), 'Model selection and estimation in regression with grouped variables', *Journal of the Royal Statistical Society Series B: Statistical Methodology* **68**(1), 49–67.
- Zhang, X., Fang, K. & Zhang, Q. (2022), 'Multivariate functional generalized additive models', *Journal of Statistical Computation and Simulation* **92**(4), 875–893.
- Zhang, X., Zhang, Q., Ma, S. & Fang, K. (2022), 'Subgroup analysis for high-dimensional functional regression', *Journal of Multivariate Analysis* **192**, 105100.

Supplementary of “High dimensional test for functional covariates”

Appendix A Preprocessing of real data

In this section, we outline the preprocessing of real data in the MEG study (refer to Section 5). This includes our methods for extracting the Power Spectral Density (PSD) from MEG recordings and procedures for outlier removal. For information on acquiring MEG recordings under the DK atlas, please refer to Jin et al. [2023].

To obtain the PSD of the MEG recordings, we firstly filter the MEG data with a band-pass filter of 1 – 40 Hz and then calculate the PSD with multitaper method [Das and Babadi, 2020] with the `pmtm` function in MATLAB. We uniformly sample 100 points within the interval [1, 40] Hz for each ROI. After the PSD is obtained, we further use a convolve function in MATLAB to smooth the PSD. We then select the PSD with [2, 35] Hz as the functional predictors $\mathbf{X}(s)$ and convert the PSD to dB scale. Note that the frequency range [2, 35] Hz includes the delta, theta, alpha and beta bands, which are commonly investigated in the MEG data analysis [Mandal et al., 2018]. After selecting PSD within [2, 35] Hz, the PSD functional predictors contain 74 points for each ROI. The PSD data in the dB scale within the [2, 35] Hz from MEG recordings should be typically greater than 0 [Ranasinghe et al., 2022], so we discard the data whose mean of PSD is less than 0 as outliers and remove them from the analysis. After all the preprocessing, our dataset includes 61 healthy subjects and 82 patients with AD ($n = 143$), with three baseline covariates and the PSD functional predictors includes 74 points for each ROI.

Appendix B Additional notations

Here we define some additional notations that are used in Appendix. Some of them are not very conventional and depends on the context.

Let $\mathbf{Q}_{\mathcal{M} \cup \mathcal{S}, \mathcal{M} \cup \mathcal{S}}(\boldsymbol{\theta})$, $\mathbf{Q}_{\mathcal{M} \cup \mathcal{S}, (\mathcal{M} \cup \mathcal{S})^c}(\boldsymbol{\theta})$, $\mathbf{Q}_{(\mathcal{M} \cup \mathcal{S})^c, \mathcal{M} \cup \mathcal{S}}(\boldsymbol{\theta})$, $\mathbf{Q}_{(\mathcal{M} \cup \mathcal{S})^c, (\mathcal{M} \cup \mathcal{S})^c}(\boldsymbol{\theta})$ be the top left $\{q + (m + k_0)N\} \times \{q + (m + k_0)N\}$, bottom left $\{q + (m + k_0)N\} \times \{(d - m - k_0)N\}$, top right $\{(d - m - k_0)N\} \times \{q + (m + k_0)N\}$, bottom right blocks of $\{(d - m - k_0)N\} \times \{(p - k_0)N\}$. Moreover, we define $\hat{\mathbf{Q}}(\boldsymbol{\theta}) = \partial^2 \mathcal{L}(\boldsymbol{\theta}) / \partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T$ and define $\hat{\mathbf{Q}}_{\mathcal{M} \cup \mathcal{S}, \mathcal{M} \cup \mathcal{S}}(\boldsymbol{\theta})$, $\hat{\mathbf{Q}}_{\mathcal{M} \cup \mathcal{S}, (\mathcal{M} \cup \mathcal{S})^c}(\boldsymbol{\theta})$, $\hat{\mathbf{Q}}_{(\mathcal{M} \cup \mathcal{S})^c, \mathcal{M} \cup \mathcal{S}}(\boldsymbol{\theta})$, $\hat{\mathbf{Q}}_{(\mathcal{M} \cup \mathcal{S})^c, (\mathcal{M} \cup \mathcal{S})^c}(\boldsymbol{\theta})$ similarly. For any vector $\mathbf{a} = (\mathbf{a}_1^T, \mathbf{a}_{2j}^T, j = 1, \dots, d)^T$, we define the subset of \mathbf{a} as $\mathbf{a}_{\mathcal{F}} \equiv (\mathbf{a}_1^T, \mathbf{a}_{2j}^T, j \in \mathcal{F})^T$ for an arbitrary index set \mathcal{F} , and $\mathbf{a}_{2\mathcal{F}} = (\mathbf{a}_{2j}^T, j \in \mathcal{F})^T$.

Appendix C Algorithm and Details of CPRSM

The CPRSM algorithm is in Algorithm 1 and more details of the algorithm are provided in the following.

- (5) is a simple quadratic minimization under the linear case or using Newton-Raphson under the logistic case. Let

$$g(\boldsymbol{\theta}) = f_1(\boldsymbol{\theta}) - \boldsymbol{\rho}^{(k)T} \{\mathbf{D}\boldsymbol{\theta} - N^{-1/2} \text{vec}(\boldsymbol{\Gamma}^{(k)})\} + \frac{\beta}{2} \|\mathbf{D}\boldsymbol{\theta} - N^{-1/2} \text{vec}(\boldsymbol{\Gamma}^{(k)})\|_2^2.$$

We have the following first and second derivatives,

$$\begin{aligned} \frac{\partial g(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} &= \frac{\partial f_1(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} - \mathbf{D}^T \boldsymbol{\rho}^{(k)} + \beta \left\{ \mathbf{D}^T \mathbf{D} \boldsymbol{\theta} - N^{-1/2} \mathbf{D}^T \text{vec}(\boldsymbol{\Gamma}^{(k)}) \right\} \\ \frac{\partial^2 g(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}^T \partial \boldsymbol{\theta}} &= \frac{\partial^2 f_1(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}^T \partial \boldsymbol{\theta}} + \beta \mathbf{D}^T \mathbf{D}. \end{aligned}$$

Under the linear case, we can simply solve

$$\frac{\partial g(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} = \mathbf{0}.$$

Algorithm 1 Contractive Peaceman-Rachford splitting method (CPRSM)

Data: Dataset $\{Y_i, \mathbf{X}_i(s), \mathbf{Z}_i\}_{i=1}^n$

Input: \mathcal{M} : the set of indices for the functional covariates of interest; λ : the penalty parameter; N : the number of B-spline basis functions; ϵ : the stop tolerance; R : the constraint constant; (α, β) : the relaxation and penalty parameters in CPRSM, $\alpha = 0.9$ following He et al. [2014]. $(\mathbf{\Gamma}^{(0)}, \boldsymbol{\rho}^{(0)})$: the initial values of $\mathbf{\Gamma}$ and $\boldsymbol{\rho}$, typically $\mathbf{\Gamma}^{(0)} = \mathbf{0}$ and $\boldsymbol{\rho}^{(0)} = \mathbf{0}$, K : the maximal iteration number.

Output: Estimators $\hat{\boldsymbol{\alpha}}$ and $\hat{\mathbf{\Gamma}}$.

```
1: for k = 0:K do
2:   Update  $\boldsymbol{\theta}^{(k)}$  following (5)–(7) to obtain  $\boldsymbol{\theta}^{(k+1)}$ .
3:   Update  $\boldsymbol{\rho}^{(k)}$  following (8) to obtain  $\boldsymbol{\rho}^{(k+1/2)}$ .
4:   Update  $\mathbf{\Gamma}^{(k)}$  following (9) to obtain  $\mathbf{\Gamma}^{(k+1)}$ .
5:   Update  $\boldsymbol{\rho}^{(k+1/2)}$  following (10) to obtain  $\boldsymbol{\rho}^{(k+1)}$ .
6:    $\Delta_{\boldsymbol{\rho}} = \boldsymbol{\rho}^{(k+1)} - \boldsymbol{\rho}^{(k)}$ ,  $\Delta_{\boldsymbol{\theta}} = \boldsymbol{\theta}^{(k+1)} - \boldsymbol{\theta}^{(k)}$ ,  $\Delta_0 = N^{1/2}\mathbf{D}\boldsymbol{\theta}^{(k+1)} - \text{vec}(\mathbf{\Gamma}^{(k+1)})$ 
7:    $\Delta = \max(\|\Delta_{\boldsymbol{\rho}}\|_2/\|\boldsymbol{\rho}^{(k+1)}\|_2, \|\Delta_{\boldsymbol{\theta}}\|_2/\|\boldsymbol{\theta}^{(k+1)}\|_2, \|\Delta_0\|_2/\|\text{vec}(\mathbf{\Gamma}^{(k+1)})\|_2)$ 
8:   if  $\Delta < \epsilon$  then
9:     BREAK
10:  end if
11: end for
12:  $\hat{\boldsymbol{\alpha}} = \boldsymbol{\theta}_1^{(k+1)}$ ,  $\hat{\mathbf{\Gamma}} = \mathbf{\Gamma}^{(k+1)}$ .
```

Under the logistic case, we can use Newton-Raphson method with the update

$$\boldsymbol{\theta}^{(k)(l+1)} = \boldsymbol{\theta}^{(k)(l)} - \left\{ \frac{\partial^2 g(\boldsymbol{\theta}^{(k)(l)})}{\partial \boldsymbol{\theta}^T \partial \boldsymbol{\theta}} \right\}^{-1} \frac{\partial g(\boldsymbol{\theta}^{(k)(l)})}{\partial \boldsymbol{\theta}}.$$

2. The projection (6) can be achieved by adopting the linear programming method of Duchi et al. [2008].

3. (9) is a quadratic minimization with SCAD penalization. We can write the fifth line as

$$\text{argmin}_{\mathbf{\Gamma}} f_2(\mathbf{\Gamma}) + g_2(\mathbf{\Gamma})$$

where $g_2(\mathbf{\Gamma}) = -\boldsymbol{\rho}^{(k+1/2)T} \{ \mathbf{D}\boldsymbol{\theta}^{(k+1)} - N^{-1/2}\text{vec}(\mathbf{\Gamma}) \} + \frac{\beta}{2} \| \mathbf{D}\boldsymbol{\theta}^{(k+1)} - N^{-1/2}\text{vec}(\mathbf{\Gamma}) \|_2^2$. So

$$\frac{\partial g_2}{\partial N^{-1/2}\text{vec}(\mathbf{\Gamma})} = \boldsymbol{\rho}^{(k+1/2)} + \beta \left\{ N^{-1/2}\text{vec}(\mathbf{\Gamma}) - \mathbf{D}\boldsymbol{\theta}^{(k+1)} \right\} = \mathbf{0}.$$

So

$$N^{-1/2}\text{vec}(\mathbf{\Gamma}^{(k+1/2)}) = \mathbf{D}\boldsymbol{\theta}^{(k+1)} - \boldsymbol{\rho}^{(k+1/2)}/\beta.$$

Then it is equivalent to solve

$$\frac{\beta}{2} \left\| N^{-1/2}\text{vec}(\mathbf{\Gamma}^{(k+1/2)}) - N^{-1/2}\text{vec}(\mathbf{\Gamma}) \right\|_2^2 + \sum_{j=1}^{T-1} \rho_\lambda(\|N^{-1/2}\boldsymbol{\gamma}_j\|_2)$$

We find $N^{-1/2}\boldsymbol{\gamma}_j^{(k+1)}$ as the projection of $N^{-1/2}\boldsymbol{\gamma}_j^{(k+1/2)}$ to the ball

$$\begin{aligned} \mathbb{B}_2\{(\|N^{-1/2}\boldsymbol{\gamma}_j^{(k+1/2)}\|_2 - \lambda/\beta)_+\} & \quad \text{if} \quad \|N^{-1/2}\boldsymbol{\gamma}_j^{(k+1/2)}\|_2 \leq \lambda + \lambda/\beta \\ \mathbb{B}_2[\{(a-1)\beta\|N^{-1/2}\boldsymbol{\gamma}_j^{(k+1/2)}\|_2 - a\lambda\}/(a\beta - \beta - 1)] & \quad \text{if} \quad \lambda + \lambda/\beta < \|N^{-1/2}\boldsymbol{\gamma}_j^{(k+1/2)}\|_2 \leq a\lambda \\ \mathbb{B}_2(\|N^{-1/2}\boldsymbol{\gamma}_j^{(k+1/2)}\|_2) & \quad \text{if} \quad \|N^{-1/2}\boldsymbol{\gamma}_j^{(k+1/2)}\|_2 > a\lambda. \end{aligned}$$

Appendix D Amenable penalty

We first list a set of assumptions on the univariate penalty function ρ_λ which are similar to those in Loh and Wainwright [2015, 2017].

- (A1) The function $\rho_\lambda(t)$ satisfies $\rho_\lambda(0) = 0$ and is symmetric around zero.
- (A2) On the nonnegative real line $t \geq 0$, the function $\rho_\lambda(t)$ is nondecreasing. Furthermore, $\rho_\lambda(t)$ is subadditive, i.e. $\rho_\lambda(t_1 + t_2) \leq \rho_\lambda(t_1) + \rho_\lambda(t_2)$ for all $t_1, t_2 \geq 0$.
- (A3) For $t > 0$, the function $\rho_\lambda(t)/t$ is non-increasing in t .
- (A4) The function $\rho_\lambda(t)$ is differentiable at all $t \neq 0$ and sub-differentiable at $t = 0$, with $\lim_{t \rightarrow 0^+} \rho'_\lambda(t) = \lambda$, where $\rho'(t)$ denotes the derivative of $\rho(t)$. Together with the symmetric Condition in (A1), this leads to $\lim_{t \rightarrow 0^-} \rho'_\lambda(t) = -\lambda$.
- (A5) There exists $\mu > 0$ so that $\rho_\lambda(t) + \mu t^2/2$ is convex.

(A6) There exists a $\gamma \in (0, +\infty)$ such that $\rho'_\lambda(t) = 0$ for all $t \geq \gamma\lambda$.

In Conditions (A1)–(A6), the first three are general requirements for the penalty function [Zhang and Zhang, 2012]. Condition (A4) excludes the regularizers that are not differentiable at zero, for example, the lasso penalty. Condition (A5) is referred to as the weak convexity condition [Chen and Gu, 2014, Vial, 1982] which controls the level of nonconvexity of the penalty function. Condition (A6) is to remove the estimation bias via allow penalty to be zero if the estimator is $\gamma\lambda$ away from zero.

We also need some additional conditions on the model and the data to facilitate the development of the theoretical properties. We discuss these conditions here but defer the detailed conditions to Section Appendix D in the supplementary material. Condition (C1) requires the boundedness of the function $\psi(t)$ in the generalized linear model. Condition (C2) imposes the sub-Gaussian assumption on the functional covariates, baseline covariates as well as the error term. Condition (C3) restricts the smoothness of the functional coefficients $\beta_j(s)$, and it also requires b-spline can approximate the true functional coefficients well enough. Condition (C4) shows the relationship between the sample size n , the number of B-spline basis N and the number of functional covariates d and baseline covariates q . It requires that the sample size should be as least in the order of $\log(p)$ to ensure the consistency of the estimator. Condition (C5) guarantees the boundedness and invertibility of the second order derivative of the log-likelihood function.

Appendix E Some useful lemmas

Define an auxiliary function $q_\lambda(t) = \lambda t - \rho_\lambda(t)$, where $q_\lambda(t) - \mu t^2/2$ is concave and everywhere differentiable in t as shown in Lemma A.5.

Lemma A.1. *There is a constant $D_r > 0$ such that for each spline $\sum_{k=1}^N c_k B_k(t)$, and for each $1 \leq l \leq \infty$*

$$D_r \|\mathbf{c}'\|_l \leq \left[\int_0^1 \left\{ \sum_{k=1}^N c_k B_k(t) \right\}^l dt \right]^{1/l} \leq \|\mathbf{c}'\|_l,$$

where $\mathbf{c}' = \{c_k \{(t_k - t_{k-b})/b\}^{1/l}, k = 1, \dots, N\}^T$.

Proof: This is a direct consequence of Theorem 5.4.2 on page 145 in DeVore and Lorentz [1993]. \square

Lemma A.2. *Conditions (A1)–(A4) imply that ρ_λ is λ -Lipschitz and all sub gradients and derivatives of ρ_λ are bounded by λ in magnitude. Conditions (A1)–(A5) imply*

$$\lambda \|\gamma_j\|_2 \leq \rho_\lambda(\|\gamma_j\|_2) + \mu/2 \|\gamma_j\|_2^2,$$

Proof: This lemma is a direct consequence of Lemma 4 in Loh and Wainwright [2015].

Lemma A.3. *Suppose ρ_λ satisfies Conditions (A1)–(A5). Let $\mathbf{V} \in \mathbb{R}^{N \times d-m}$, let \mathcal{A} be the index set of k largest elements of $\|\mathbf{V}_{\cdot j}\|_2, j = 1, \dots, d-m$ in magnitude, and let \mathcal{A}^c be the index set of the remaining $d-m-k_0$ columns of \mathbf{V} . Suppose $\xi > 0$ and satisfies*

$$\xi \sum_{j \in \mathcal{A}} \rho_\lambda(\|\mathbf{V}_{\cdot j}\|_2) - \sum_{j \in \mathcal{A}^c} \rho_\lambda(\|\mathbf{V}_{\cdot j}\|_2) \geq 0.$$

Then

$$\xi \sum_{j \in \mathcal{A}} \rho_\lambda(\|\mathbf{V}_{\cdot j}\|_2) - \sum_{j \in \mathcal{A}^c} \rho_\lambda(\|\mathbf{V}_{\cdot j}\|_2) \leq \lambda \left(\xi \sum_{j \in \mathcal{A}} \|\mathbf{V}_{\cdot j}\|_2 - \sum_{j \in \mathcal{A}^c} \|\mathbf{V}_{\cdot j}\|_2 \right).$$

Moreover, if $\mathbf{\Gamma}_{0, \mathcal{M}^c} = (\|\gamma_{0j}\|_2, j \in \mathcal{M}^c)^T$ is k -sparse, then suppose there are $\mathbf{\Gamma}_{\mathcal{M}^c} = \{\gamma_j, j \in \mathcal{M}^c\}$ such that $\xi \sum_{j \in \mathcal{M}^c} \{\rho_\lambda(\|\gamma_{0j}\|_2) - \rho_\lambda(\|\gamma_j\|_2)\} > 0$ and $\xi \geq 1$, we have

$$\sum_{j \in \mathcal{M}^c} \xi \rho_\lambda(\|\gamma_{0j}\|_2) - \rho_\lambda(\|\gamma_j\|_2) \leq \lambda \left(\xi \sum_{j \in \mathcal{A}} \|\mathbf{V}_{\cdot j}\|_2 - \sum_{j \in \mathcal{A}^c} \|\mathbf{V}_{\cdot j}\|_2 \right),$$

where $\mathbf{V} = \mathbf{\Gamma}_{\mathcal{M}^c} - \mathbf{\Gamma}_{0, \mathcal{M}^c}$, \mathcal{A} is the index set of the k largest $\|\mathbf{V}_{\cdot j}\|_2$, and \mathcal{A}^c is the index set of the remaining $d - m - k$ columns of \mathbf{V} .

Proof: This lemma is a direct consequence of Lemma 5 in Loh and Wainwright [2015]. \square

Lemma A.4. *Suppose ρ_λ is (μ, γ) -amenable, $\|\hat{\boldsymbol{\gamma}}_j\|_2 \geq \lambda\gamma$ for $j \in \mathcal{M} \cup S$, then $q'_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2) = \lambda$.*

Proof: Because ρ_λ is (μ, γ) -amenable, $\rho'(\|\hat{\boldsymbol{\gamma}}_j\|_2) = 0$ by Condition (A6) and (A1). Hence $q'_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2) = \partial\lambda\|\hat{\boldsymbol{\gamma}}_j\|_2/\partial\|\hat{\boldsymbol{\gamma}}_j\|_2 = \lambda$. This proves the result. \square

Lemma A.5. *Consider a μ -amenable regularizer ρ_λ . Then*

(a) $|\rho'_\lambda(t)| \leq \lambda$ for all $t \neq 0$.

(b) *The function $q_\lambda(\|\mathbf{t}\|_2) - \mu/2\|\mathbf{t}\|_2^2$ is concave in $\|\mathbf{t}\|_2$ and everywhere differentiable in \mathbf{t} , where $q_\lambda(\|\mathbf{t}\|_2) = \lambda\|\mathbf{t}\|_2 - \rho_\lambda(\|\mathbf{t}\|_2)$.*

Proof: This lemma is a direct consequence of Lemma 5 in Loh and Wainwright [2017]. \square

Lemma A.6. *Let $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{p \times p}$ be invertible. For any matrix norm $\|\cdot\|$, we have*

$$\|\mathbf{A}^{-1} - \mathbf{B}^{-1}\| \leq \frac{\|\mathbf{A}^{-1}\|^2 \|\mathbf{A} - \mathbf{B}\|}{1 - \|\mathbf{A}^{-1}\| \|\mathbf{A} - \mathbf{B}\|}.$$

In particular, if $\|\mathbf{A}^{-1}\| \|\mathbf{A} - \mathbf{B}\| \leq 1/2$, then $\|\mathbf{A}^{-1} - \mathbf{B}^{-1}\| = O(\|\mathbf{A}^{-1}\|^2 \|\mathbf{A} - \mathbf{B}\|)$.

Proof: This lemma is Lemma 11 in Loh and Wainwright [2017]. \square

Lemma A.7. *Suppose $\mathbf{X}_1, \dots, \mathbf{X}_n$ are independent m dimensional random vector which satisfies, $E(\mathbf{X}_i) = 0$ and $\sum_{i=1}^n \text{cov}(\mathbf{X}_i) = \mathbf{I}_m$. Let \mathbf{Z} be a m -dimensional standard multivariate normal vector, then*

$$\sup_{\mathcal{C}} |\Pr(\sum_{i=1}^n \mathbf{X}_i \in \mathcal{C}) - \Pr(\mathbf{Z} \in \mathcal{C})| = O(m^{1/4} \sum_{i=1}^n E\|\mathbf{X}_i\|_2^3).$$

Proof: The lemma follows Theorem 1.1 in Bentkus [2005].

Appendix F Main proofs

Write $\boldsymbol{\theta} = \{\boldsymbol{\alpha}^\top, N^{-1/2}\text{vec}(\boldsymbol{\Gamma})^\top\}^\top$. For notation simplicity, we denote $\mathcal{L}(\boldsymbol{\theta}) \equiv \mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\Gamma})$. The derivative of $\mathcal{L}(\boldsymbol{\theta})$ with respect to $\boldsymbol{\theta}$ is

$$\frac{\partial \mathcal{L}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} = n^{-1} \sum_{i=1}^n \{\psi'_i(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) - Y_i\} \left[\mathbf{z}_i^\top, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^\top \right]^\top$$

and the second derivative of $\mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\Gamma})$ with respect to $\boldsymbol{\theta}$ is

$$\frac{\partial^2 \mathcal{L}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^\top} = n^{-1} \sum_{i=1}^n \psi''_i(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^\top, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^\top \right]^\top \right)^{\otimes 2}.$$

Denote $\tilde{\boldsymbol{\theta}} = \{\tilde{\boldsymbol{\alpha}}^\top, \text{vec}(\tilde{\boldsymbol{\Gamma}})^\top\}^\top$ be a stationary points for (3), respectively, which satisfies the first order condition that

$$\partial \mathcal{L}(\tilde{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^\top (\boldsymbol{\theta} - \tilde{\boldsymbol{\theta}}) + N^{-1/2} \sum_{j \in \mathcal{M}^c} \partial \rho_\lambda(\|\tilde{\boldsymbol{\gamma}}_j\|_2) / \partial \tilde{\boldsymbol{\gamma}}_j^\top (\boldsymbol{\gamma}_j - \tilde{\boldsymbol{\gamma}}_j) \geq 0 \quad (1)$$

Lemma A.8. *Assume Conditions (C2) and (C3) hold, we have for any given unit vector $\mathbf{v} = (v_{jk}, j = 1, \dots, d, k = 1, \dots, N)$ with $\|\mathbf{v}\|_2 = 1$, $N^{1/2} \sum_{j=1}^d \sum_{k=1}^N v_{jk} \int_0^1 X_{ij}(s) B_k(s) ds$ is a sub-Gaussian random variable. Therefore,*

$$\left[\mathbf{z}_i^\top, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^\top \right]^\top$$

is a sub-Gaussian random vector.

Proof: For a given unit vector $\mathbf{v} = (v_{jk}, j = 1, \dots, d, k = 1, \dots, N)$ with $\|\mathbf{v}\|_2 = 1$. Denote $\mathbf{v}_{j\cdot} = (v_{j1}, \dots, v_{jN})^\top$.

$$\begin{aligned} & \left\{ \sum_{j=1}^d \sum_{k=1}^N v_{jk} \int_0^1 X_{ij}(s) N^{1/2} B_k(s) ds \right\}^l \\ & \leq \left[\sum_{j=1}^d \sum_{k=1}^N \left\{ v_{jk}^2 \int_0^1 X_{ij}^2(s) ds \int_0^1 N B_k^2(s) ds \right\}^{1/2} \right]^l \end{aligned}$$

$$\begin{aligned}
&\leq \left[\sum_{j=1}^d \sum_{k=1}^N \left\{ v_{jk}^2 \int_0^1 X_{ij}^2(s) ds \int_0^1 NB_k(s) ds \right\} \right]^{1/2}{}^l \\
&\leq \left[\sum_{j=1}^d \sum_{k=1}^N \left\{ v_{jk}^2 \int_0^1 X_{ij}^2(s) ds \int_0^1 NB_k(s) ds \right\} \right]^{l/2} \\
&= \left[\sum_{j=1}^d \int_0^1 \sum_{k=1}^N \left\{ v_{jk}^2 \int_0^1 X_{ij}^2(s) ds NB_k(s) \right\} ds \right]^{l/2} \\
&\leq \left[\sum_{j=1}^d \sum_{k=1}^N v_{jk}^2 \int_0^1 X_{ij}^2(s) ds N(\tau_k - \tau_{k-b})/b \right]^{l/2} \\
&\leq \left[\sum_{j=1}^d \|\mathbf{v}_{j\cdot}\|_2 \int_0^1 X_{ij}^2(s) ds \sum_{k=1}^N \frac{v_{jk}^2}{\|\mathbf{v}_{j\cdot}\|_2} H \right]^{l/2} \\
&\leq \left[H \sum_{j=1}^d \|\mathbf{v}_{j\cdot}\|_2 \int_0^1 X_{ij}^2(s) ds \right]^{l/2},
\end{aligned}$$

where the second inequality holds because $B_k(s) \in [0, 1]$, the sixth line holds by Lemma A.1, the last line holds because by Condition (C3), $\tau_k - \tau_{k-b} \leq h_{\max}b \leq h_{\min}Hb$ and fact that $h_{\min}N \leq 1$. Furthermore $\sum_{k=1}^d v_{jk}^2/\|\mathbf{v}_{j\cdot}\|_2 = \|\mathbf{v}_{j\cdot}\|_2 \leq \|\mathbf{v}\|_2 = 1$. Hence,

$$E \left\{ N^{1/2} \sum_{j=1}^d \sum_{k=1}^N v_{jk} \int_0^1 X_{ij}(s) B_k(s) ds \right\}^l \leq E \left[H \sum_{j=1}^d \|\mathbf{v}_{j\cdot}\|_2 \int_0^1 X_{ij}^2(s) ds \right]^{l/2}. \quad (2)$$

Now because for any unit vector \mathbf{u} , $\sum_{j=1}^N u_j \int_0^1 X_{ij}^2(s) ds$ is a sub-exponential distributed random variable by Condition (C2), and $\{\|\mathbf{v}_{j\cdot}\|_2, j = 1, \dots, d\}^T$ is a unit vector, by the definition of sub-exponential random variable in (5.15) in [Vershynin, 2010], we have there is a constant $C < \infty$ such that

$$\left(E \left[\sum_{j=1}^d \|\mathbf{v}_{j\cdot}\|_2 \int_0^1 X_{ij}^2(s) ds \right]^{l/2} \right)^{2/l} = Cl/2$$

which implies

$$E \left[\sum_{j=1}^d \|\mathbf{v}_{j\cdot}\|_2 \int_0^1 X_{ij}^2(s) ds \right]^{l/2} = (l/2)^{l/2} C^{l/2}.$$

Plug the above result to (2), we have

$$\begin{aligned} & l^{-1/2} \left[E \left\{ N^{1/2} \sum_{j=1}^d \sum_{k=1}^N v_{jk} \int_0^1 X_{ij}(s) B_k(s) ds \right\}^l \right]^{1/l} \\ &= l^{-1/2} \{(Hl/2)^{l/2} C^{l/2}\}^{1/l} = l^{-1/2} (l/2)^{1/2} H^{1/2} C^{1/2} = (HC/2)^{1/2} \end{aligned}$$

Therefore, we have $\|N^{1/2} \sum_{j=1}^d \sum_{k=1}^N v_{jk} \int_0^1 X_{ij}(s) B_k(s) ds\|_{\psi_2} < \infty$. By the definition of the sub-Gaussian random variable as shown in (5.11) in [Vershynin, 2010], we have $N^{1/2} \sum_{j=1}^d \sum_{k=1}^N v_{jk} \int_0^1 X_{ij}(s)$ is a sub-Gaussian random variable. Now because \mathbf{Z}_i is a fixed dimensional sub-Gaussian vector, we have

$$\left[\mathbf{Z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T$$

is a sub-Gaussian random vector. This proves the result. \square

Lemma A.9. *For any constant $s > 1$, we have*

$$\mathbb{B}_1(\sqrt{s}) \cap \mathbb{B}_2(1) \subseteq \text{cl}\{\text{conv}\{\mathbb{B}_0(s) \cap \mathbb{B}_2(3)\}\},$$

where the \mathbb{B}_k balls with radius r , $\mathbb{B}_k(r)$, $k = 0, 1, 2$, are taken in p -dimensional space, and $\text{cl}(\cdot)$ and $\text{conv}(\cdot)$ denote the topological closure and convex hull, respectively.

Proof: From Lemma 11 in Loh and Wainwright [2012], we get

$$\mathbb{B}_1(\sqrt{s}) \cap \mathbb{B}_2(1) \subseteq 3\text{cl}\{\text{conv}\{\mathbb{B}_0(s) \cap \mathbb{B}_2(3)\}\}.$$

Here for a set A , $3A$ is defined as the set that satisfies $\sup_{\boldsymbol{\theta} \in 3A} \langle \boldsymbol{\theta}, \mathbf{z} \rangle = 3 \sup_{\boldsymbol{\theta} \in A} \langle \boldsymbol{\theta}, \mathbf{z} \rangle$ for any \mathbf{z} . Let U be a subset of $\{1, \dots, p\}$ and \mathbf{z}_U be the subvector of \mathbf{z} with only the elements whose indices in U retained. Now when $A = \text{cl}\{\text{conv}\{\mathbb{B}_0(s) \cap \mathbb{B}_2(1)\}\}$, we get $\sup_{\boldsymbol{\theta} \in 3A} \langle \boldsymbol{\theta}, \mathbf{z} \rangle = 3 \max_{|U|=|s|} \sup_{\|\boldsymbol{\theta}_U\|_2 \leq 1} \langle \boldsymbol{\theta}_U, \mathbf{z}_U \rangle = \max_{|U|=|s|} \sup_{\|\boldsymbol{\theta}_U\|_2 \leq 3} \langle \boldsymbol{\theta}_U, \mathbf{z}_U \rangle = 3\|\mathbf{z}_U\|_2$, hence $3\text{cl}\{\text{conv}\{\mathbb{B}_0(s) \cap \mathbb{B}_2(1)\}\} = \text{cl}\{\text{conv}\{\mathbb{B}_0(s) \cap \mathbb{B}_2(3)\}\}$. Thus the results hold.

This proves the Lemma. \square

Lemma A.10. For a fixed matrix $\mathbf{\Omega} \in \mathbb{R}^{(q+Nd) \times (q+Nd)}$, parameter $s \geq 1$, and tolerance $\delta > 0$. For \mathbf{V} be a $N \times d$ dimensional matrix with v_{ij} be its i, j th element. Furthermore, let $\mathbf{W} = (w_1, \dots, w_q)^\top$ be a q dimensional vector. Define

$$\mathbb{K}(t) = \{(\mathbf{W}^\top, \mathbf{V}^\top)^\top \in \mathbf{R}^{q+Nd}, \|\{(\mathbf{W}^\top, \|\mathbf{V}_{\cdot j}\|_2, j = 1, \dots, d)\}^\top\|_0 \leq t, \|\{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top\|_2 \leq 1\}.$$

Suppose the following condition holdes

$$|\{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top \mathbf{\Omega} \{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top| \leq \delta, \quad \forall \{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top \in \mathbb{K}(2s). \quad (3)$$

Then

$$\begin{aligned} & |\{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top \mathbf{\Omega} \{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top| \\ & \leq 27\delta \left[\|\{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top\|_2^2 + s^{-1} \left\{ \|\mathbf{W}\|_1 + \sum_{j=1}^d \|\mathbf{V}_{\cdot j}\|_2 \right\}^2 \right], \forall \mathbf{W} \in \mathbb{R}^q, \mathbf{V} \in \mathbb{R}^{N \times d}. \end{aligned}$$

Proof: Define

$$\mathbb{C}(s) \equiv \left[\{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top : \|\mathbf{W}\|_1 + \sum_{j=1}^d \|\mathbf{V}_{\cdot j}\|_2 \leq \sqrt{s} \|\{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top\|_2 \right].$$

First note that $\|\{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top\|_2 = \|(\mathbf{W}^\top, \|\mathbf{V}_{\cdot j}\|_2, j = 1, \dots, d)^\top\|_2$. Therefore, for $\{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top \in \mathbb{C}(s)$, by Lemma A.9, we have $(\mathbf{W}^\top, \|\mathbf{V}_{\cdot j}\|_2, j = 1, \dots, d)^\top \in 3\text{cl}\{\text{conv}\{\mathbb{B}_0(s) \cap \mathbb{B}_2(3)\}\}$. Consider a weighted linear combination of the form $\{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top = \sum_k a_k \{\mathbf{W}^{k\top}, \text{vec}(\mathbf{V}^k)^\top\}^\top$, with weights $a_k \geq 0$ such that $\sum_k a_k = 1$ and $\|(\mathbf{W}^{k\top}, \|\mathbf{V}_{\cdot j}^k\|_2, j = 1, \dots, d)^\top\|_0 \leq s$ and $\|\{\mathbf{W}^{k\top}, \text{vec}(\mathbf{V}^k)^\top\}^\top\|_2 \leq 3$ for each k . Then we can write

$$\begin{aligned} \{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top \mathbf{\Omega} \{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top &= \sum_k a_k \left\{ \mathbf{W}^{k\top}, \text{vec}(\mathbf{V}^k)^\top \right\} \mathbf{\Omega} \sum_k a_k \left\{ \mathbf{W}^{k\top}, \text{vec}(\mathbf{V}^k)^\top \right\}^\top \\ &= \sum_{k,l} a_k a_l \left\{ \mathbf{W}^{k\top}, \text{vec}(\mathbf{V}^k)^\top \right\} \mathbf{\Omega} \left\{ \mathbf{W}^{l\top}, \text{vec}(\mathbf{V}^l)^\top \right\}^\top. \end{aligned}$$

Now because $1/3 \left\{ \mathbf{W}^{k\top}, \text{vec}(\mathbf{V}^k)^\top \right\}^\top$, $1/3 \left\{ \mathbf{W}^{l\top}, \text{vec}(\mathbf{V}^l)^\top \right\}^\top$ and

$$1/6 \left[\left\{ \mathbf{W}^{k\top}, \text{vec}(\mathbf{V}^k)^\top \right\} + \left\{ \mathbf{W}^{l\top}, \text{vec}(\mathbf{V}^l)^\top \right\} \right]$$

are in $\mathbb{K}(2s)$, follow (3), we can write

$$\begin{aligned}
& \left| \left\{ \mathbf{W}^{k\top}, \text{vec}(\mathbf{V}^k)^\top \right\} \boldsymbol{\Omega} \left\{ \mathbf{W}^{l\top}, \text{vec}(\mathbf{V}^l)^\top \right\}^\top \right| \\
&= \frac{1}{2} \left| \left[\left\{ \mathbf{W}^{k\top}, \text{vec}(\mathbf{V}^k)^\top \right\} + \left\{ \mathbf{W}^{l\top}, \text{vec}(\mathbf{V}^l)^\top \right\} \right] \boldsymbol{\Omega} \left[\left\{ \mathbf{W}^{k\top}, \text{vec}(\mathbf{V}^k)^\top \right\} + \left\{ \mathbf{W}^{l\top}, \text{vec}(\mathbf{V}^l)^\top \right\} \right]^\top \right. \\
&\quad \left. - \left\{ \mathbf{W}^{k\top}, \text{vec}(\mathbf{V}^k)^\top \right\} \boldsymbol{\Omega} \left\{ \mathbf{W}^{k\top}, \text{vec}(\mathbf{V}^k)^\top \right\}^\top - \left\{ \mathbf{W}^{l\top}, \text{vec}(\mathbf{V}^l)^\top \right\} \boldsymbol{\Omega} \left\{ \mathbf{W}^{l\top}, \text{vec}(\mathbf{V}^l)^\top \right\}^\top \right| \\
&\leq \frac{1}{2} (36\delta + 9\delta + 9\delta) = 27\delta,
\end{aligned}$$

for all k, l . Therefore, $|\{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\} \boldsymbol{\Omega} \{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top| \leq \sum_{k,l} a_k a_l (27\delta) \leq 27\delta \|\mathbf{a}\|_2^2 = 27\delta$. Therefore,

$$\{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\} \boldsymbol{\Omega} \{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top \leq 27\delta, \forall \{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top \in \mathbb{C}(s). \quad (4)$$

On the other hand for $\{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top \notin \mathbb{C}(s)$, we have

$$\frac{\{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\} \boldsymbol{\Omega} \{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top}{\left(\|\mathbf{W}\|_1 + \sum_{j=1}^d \|\mathbf{V}_{\cdot j}\|_2 \right)^2} \leq \frac{1}{s} \sup_{\mathbf{u} \in \mathbb{C}(s)} |\mathbf{u}^\top \boldsymbol{\Omega} \mathbf{u}| \leq \frac{27\delta}{s},$$

where $\mathbf{u} \equiv \{\mathbf{W}_1^\top, \text{vec}(\mathbf{V}_1)^\top\}^\top \equiv \sqrt{s} \{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top / (\|\mathbf{W}\|_1 + \sum_{j=1}^d \|\mathbf{V}_{\cdot j}\|_2)$. $\mathbf{u} \in \mathbb{C}(s)$ because $\|\mathbf{W}\|_1 + \sum_{j=1}^d \|\mathbf{V}_{\cdot j}\|_2 \geq \sqrt{s} \|\{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top\|_2$, $\|\mathbf{u}\|_2 \leq 1$ and $\|\mathbf{W}_1\|_1 + \sum_{j=1}^d \|\mathbf{V}_{1 \cdot j}\|_2 = \sqrt{s} (\|\mathbf{W}\|_1 + \sum_{j=1}^d \|\mathbf{V}_{\cdot j}\|_2) / (\|\mathbf{W}\|_1 + \sum_{j=1}^d \|\mathbf{V}_{\cdot j}\|_2) = \sqrt{s}$. Combine with (4), we have

$$\{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\} \boldsymbol{\Omega} \{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top \leq 27\delta \left\{ \|\{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top\|_2^2 + \frac{1}{s} \left(\|\mathbf{W}\|_1 + \sum_{j=1}^d \|\mathbf{V}_{\cdot j}\|_2 \right)^2 \right\}$$

□

Lemma A.11. *Assume Conditions (C1), (C2) and (C3) hold, there is a constant $c > 0$ such that for any given vector \mathbf{v} ,*

$$\begin{aligned}
& \Pr \left[\left| \mathbf{v}^\top \sum_{i=1}^n \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{Z}_i^\top, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^\top \right]^\top \right)^{\otimes 2} \right. \right. \\
& \quad \left. \left. - \mathbf{v}^\top E \left\{ \sum_{i=1}^n \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{Z}_i^\top, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^\top \right]^\top \right)^{\otimes 2} \right\} \mathbf{v} \right| \geq nt \right] \leq 2 \exp \{-cn \min(t^2, t)\}.
\end{aligned}$$

Proof: By Lemma A.8

$$\left[\mathbf{z}_i, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^{\text{T}} \right]^{\text{T}}$$

is a sub-Gaussian random vector. Furthermore $0 < \psi''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) < \infty$ by Condition (C1), we have

$$\psi''(\boldsymbol{\alpha}, \boldsymbol{\Gamma})^{1/2} \left[\mathbf{z}_i^{\text{T}}, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^{\text{T}} \right]^{\text{T}}$$

is sub-Gaussian vector. Hence, follow Lemma 14 (G.1) in [Loh and Wainwright, 2012], we have there is a constant $c > 0$ such that

$$\begin{aligned} & \Pr \left[\left| \mathbf{v}^{\text{T}} \sum_{i=1}^n \psi''_i(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^{\text{T}}, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^{\text{T}} \right]^{\text{T}} \right)^{\otimes 2} \right| \mathbf{v} \right. \\ & \left. - \mathbf{v}^{\text{T}} E \left\{ \sum_{i=1}^n \psi''_i(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^{\text{T}}, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^{\text{T}} \right]^{\text{T}} \right)^{\otimes 2} \right\} \right| \geq nt \right] \leq 2 \exp \{ cn \min(t^2, t) \}. \end{aligned}$$

This proves the result. \square

Lemma A.12. *Assume Conditions (C1), (C2) and (C3) hold, there is a constant $c_1 > 0$ such that*

$$\begin{aligned} & \Pr \left(\sup_{\mathbf{v} \in \mathbb{K}(2s)} \left| \sum_{i=1}^n \mathbf{v}^{\text{T}} \left[\psi''_i(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^{\text{T}}, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^{\text{T}} \right]^{\text{T}} \right)^{\otimes 2} \right. \right. \right. \\ & \left. \left. - E \left\{ \psi''_i(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^{\text{T}}, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^{\text{T}} \right]^{\text{T}} \right)^{\otimes 2} \right\} \right] \right| > nt \right) \\ & \leq 2 \exp \{ -c_1 n \min(t^2, t) + 2s \log(9p) \}. \end{aligned}$$

Proof: For each subset $\mathcal{U} \subset (1, \dots, p)$, we define the set $S_{\mathcal{U}}$ as $S_{\mathcal{U}} = [\{\mathbf{W}^{\text{T}}, \text{vec}(\mathbf{V})^{\text{T}}\}^{\text{T}} \in \mathbb{R}^{p+Nd}, \|\{\mathbf{W}^{\text{T}}, \text{vec}(\mathbf{V})^{\text{T}}\}^{\text{T}}\|_2 \leq 1, \text{supp}\{\{\mathbf{W}^{\text{T}}, \|\mathbf{V}_{\cdot j}\|_2, j = 1, \dots, d\}^{\text{T}} \subseteq \mathcal{U}\}]$, and note that $\mathbb{K}(2s) = \cup_{|\mathcal{U}| \leq 2s} S_{\mathcal{U}}$. We define $\mathcal{A} = \{\mathbf{u}_1, \dots, \mathbf{u}_m\} \subset S_{\mathcal{U}}$ to be a $1/3$ -cover of $S_{\mathcal{U}}$, if for every $\mathbf{v} \in S_{\mathcal{U}}$, there is some $\mathbf{u}_i \in \mathcal{A}$ such that $\|\mathbf{v} - \mathbf{u}_i\|_2 \leq 1/3$. Define $\Delta \mathbf{v} = \mathbf{v} - \mathbf{u}_j$ where

$\mathbf{u}_j = \arg \min_{\mathbf{u}_i} \|\mathbf{v} - \mathbf{u}_i\|_2$. We have $\|\Delta \mathbf{v}\|_2 \leq 1/3$. The same as those shown in Lemma 15 in Loh and Wainwright [2012], by Ledoux and Talagrand [2013], we can construct \mathcal{A} with $|\mathcal{A}| < 9^{2sN+q}$. Define

$$\begin{aligned} \Phi(\mathbf{v}_1, \mathbf{v}_2) &= \mathbf{v}_1^\top n^{-1} \sum_{i=1}^n \left[\psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^\top, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^\top \right]^\top \right)^{\otimes 2} \right. \\ &\quad \left. - E \left\{ \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^\top, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^\top \right]^\top \right)^{\otimes 2} \right\} \right] \mathbf{v}_2. \end{aligned}$$

We have

$$\begin{aligned} &|\Phi(\mathbf{v}, \mathbf{v})| \\ &= |\Phi(\Delta \mathbf{v} + \mathbf{u}_j, \Delta \mathbf{v} + \mathbf{u}_j)| \\ &\leq \max_i |\Phi(\mathbf{u}_i, \mathbf{u}_i)| + \max_i |\Phi(\Delta \mathbf{v}, \mathbf{u}_i)| + \max_i |\Phi(\mathbf{u}_i, \Delta \mathbf{v})| + |\Phi(\Delta \mathbf{v}, \Delta \mathbf{v})| \\ &\leq \max_i |\Phi(\mathbf{u}_i, \mathbf{u}_i)| + 2 \max_i |\Phi(\Delta \mathbf{v}, \mathbf{u}_i)| + |\Phi(\Delta \mathbf{v}, \Delta \mathbf{v})|. \end{aligned}$$

Hence,

$$\sup_{\mathbf{v} \in S_{\mathcal{U}}} |\Phi(\mathbf{v}, \mathbf{v})| \leq \max_i |\Phi(\mathbf{u}_i, \mathbf{u}_i)| + 2 \sup_{\mathbf{v} \in S_{\mathcal{U}}} \max_i |\Phi(\Delta \mathbf{v}, \mathbf{u}_i)| + \sup_{\mathbf{v} \in S_{\mathcal{U}}} |\Phi(\Delta \mathbf{v}, \Delta \mathbf{v})|.$$

Since $\|3\Delta \mathbf{v}\|_2 \leq 1$ and $\text{supp}(3\Delta \mathbf{v}) \subseteq \mathcal{U}$, $3\Delta \mathbf{v} \in S_{\mathcal{U}}$. It follows that

$$\begin{aligned} &\sup_{\mathbf{v} \in S_{\mathcal{U}}} |\Phi(\mathbf{v}, \mathbf{v})| \\ &\leq \max_i |\Phi(\mathbf{u}_i, \mathbf{u}_i)| + 2/3 \sup_{\mathbf{v} \in S_{\mathcal{U}}} \max_i |\Phi(3\Delta \mathbf{v}, \mathbf{u}_i)| + 1/9 \sup_{\mathbf{v} \in S_{\mathcal{U}}} |\Phi(3\Delta \mathbf{v}, 3\Delta \mathbf{v})| \\ &\leq \max_i |\Phi(\mathbf{u}_i, \mathbf{u}_i)| + 2/3 \{ \sup_{\mathbf{v} \in S_{\mathcal{U}}} |\Phi(3\Delta \mathbf{v}, 3\Delta \mathbf{v})| \}^{1/2} \{ \max_i |\Phi(\mathbf{u}_i, \mathbf{u}_i)| \}^{1/2} + 1/9 \sup_{\mathbf{v} \in S_{\mathcal{U}}} |\Phi(\mathbf{v}, \mathbf{v})| \\ &\leq \max_i |\Phi(\mathbf{u}_i, \mathbf{u}_i)| + \sup_{\mathbf{v} \in S_{\mathcal{U}}} \{ 2/3 |\Phi(\mathbf{v}, \mathbf{v})| + 1/9 |\Phi(\mathbf{v}, \mathbf{v})| \}. \end{aligned}$$

Hence, $\sup_{\mathbf{v} \in S_{\mathcal{U}}} |\Phi(\mathbf{v}, \mathbf{v})| \leq 9/2 \max_i |\Phi(\mathbf{u}_i, \mathbf{u}_i)|$. By Lemma A.11 and a union bound, we

have

$$\Pr \left(\sup_{\mathbf{v} \in S_{\mathcal{U}}} \left| \sum_{i=1}^n \mathbf{v}^\top \left[\psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^\top, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^\top \right]^\top \right)^{\otimes 2} \right] \right| \right)$$

$$\begin{aligned}
& -E \left\{ \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} \right\} \left| \mathbf{v} \right| > 9/2nt \Big) \\
\leq & 9^{2sN+q} 2 \exp \{ -cn \min(t^2, t) \}. \tag{5}
\end{aligned}$$

Now replacing t with $2/9t$, we have there is a constant c_1

$$\begin{aligned}
& \Pr \left(\sup_{\mathbf{v} \in S_{\mathcal{U}}} \left| \sum_{i=1}^n \mathbf{v}^T \left[\psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} \right] \right| > 9/2nt \right) \\
& -E \left\{ \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} \right\} \left| \mathbf{v} \right| > nt \Big) \\
\leq & 9^{2sN+q} 2 \exp \{ -c_1 n \min(t^2, t) \}.
\end{aligned}$$

Finally, taking a union bound over the $\binom{p}{2s}$ choices of \mathcal{U} , and noting that $\binom{p}{2s} \leq p^{2s}$, we have

$$\begin{aligned}
& \Pr \left(\sup_{\mathbf{v} \in \mathbb{K}(2s)} \left| \sum_{i=1}^n \mathbf{v}^T \left[\psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} \right] \right| > nt \right) \\
& -E \left\{ \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} \right\} \left| \mathbf{v} \right| > nt \Big) \\
\leq & 2 \exp \{ -c_1 n \min(t^2, t) + (2sN + q) \log(9) + 2s \log(p) \}.
\end{aligned}$$

This proves the result. \square

Lemma A.13. *Assume Conditions (C1)–(C5) hold, $\forall \mathbf{W} \in \mathbb{R}^q, \mathbf{V} \in \mathbb{R}^{N \times d}$, there is constant*

$c_1 > 0$, such that

$$\begin{aligned}
& \{\mathbf{W}^T, \text{vec}(\mathbf{V})^T\} n^{-1} \sum_{i=1}^n \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} \{\mathbf{W}^T, \text{vec}(\mathbf{V})^T\}^T \\
& \geq c_{\psi} \alpha_{\min} \|\{\mathbf{W}^T, \text{vec}(\mathbf{V})^T\}^T\|_2^2 / 2 - \tau(n, p) \left\{ \|\mathbf{W}\|_1 + \sum_{j=1}^d \|\mathbf{V}_{\cdot j}\|_2 \right\}^2,
\end{aligned}$$

and

$$\{\mathbf{W}^T, \text{vec}(\mathbf{V})^T\} n^{-1} \sum_{i=1}^n \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} \{\mathbf{W}^T, \text{vec}(\mathbf{V})^T\}^T$$

$$\leq 3c_\psi\alpha_{\max}\|\{\mathbf{W}^\top, \text{vec}(\mathbf{V})^\top\}^\top\|_2^2/2 + \tau(n, p) \left\{ \|\mathbf{W}\|_1 + \sum_{j=1}^d \|\mathbf{V}_{\cdot j}\|_2 \right\}^2$$

with probability greater than $1 - 2 \exp\{-c_1 n \min(c_\psi^2 \alpha_{\min}^2 / 54^2, 1) / 2\}$, where $\tau(n, p) = \tau_1 \{\log(p) + N + q\} / n$, where $\tau_1 = 5c_\psi \alpha_{\min} / \{c_1 \min(c_\psi^2 \alpha_{\min}^2 / 54^2, 1)\}$.

Proof: Let

$$\begin{aligned} \boldsymbol{\Omega} = & n^{-1} \sum_{i=1}^n \left[\psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^\top, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^\top \right]^\top \right)^{\otimes 2} \right. \\ & \left. - E \left\{ \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^\top, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^\top \right]^\top \right)^{\otimes 2} \right\} \right]. \end{aligned}$$

Let

$$t = c_\psi \alpha_{\min} \left[E \left\{ \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^\top, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^\top \right]^\top \right)^{\otimes 2} \right\} \right] / 54 = c_\psi \alpha_{\min} / 54,$$

by Lemma (A.12), for $p \geq 9$, we have

$$\begin{aligned} & \Pr \left(\sup_{\mathbf{v} \in S_U} \left| \sum_{i=1}^n \mathbf{v}^\top \left[\psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^\top, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^\top \right]^\top \right)^{\otimes 2} \right. \right. \right. \\ & \quad \left. \left. - E \left\{ \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^\top, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^\top \right]^\top \right)^{\otimes 2} \right\} \right] \mathbf{v} \right| > nc_\psi \alpha_{\min} / 54 \right) \\ & \leq \Pr \left(\sup_{\mathbf{v} \in S_U} \left| \sum_{i=1}^n \mathbf{v}^\top \left[\psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^\top, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^\top \right]^\top \right)^{\otimes 2} \right. \right. \right. \\ & \quad \left. \left. - E \left\{ \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^\top, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^\top \right]^\top \right)^{\otimes 2} \right\} \right] \mathbf{v} \right| > n \min(c_\psi \alpha_{\min} / 54, 1) \right) \\ & \leq 2 \exp\{-c_1 n \min(c_\psi^2 \alpha_{\min}^2 / 54^2, 1) + (2sN + q) \log 9 + 2s \log(p)\} \\ & \leq 2 \exp\{-c_1 n \min(c_\psi^2 \alpha_{\min}^2 / 54^2, 1) + 5s(N + q + \log(p))\}. \end{aligned}$$

Let $s = c_1 n \min(\alpha_{\min}^2 / 54^2, 1) / [10\{N + q + \log(p)\}]$, we obtain

$$\Pr \left(\sup_{\mathbf{v} \in S_U} \left| \sum_{i=1}^n \mathbf{v}^\top \left[\psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^\top, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^\top \right]^\top \right)^{\otimes 2} \right. \right. \right. \right.$$

$$\begin{aligned}
& -E \left\{ \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{Z}_i^{\text{T}}, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^{\text{T}} \right]^{\text{T}} \right)^{\otimes 2} \right\} \mathbf{v} \Big| > nc_\psi \alpha_{\min}/54 \Big) \\
& \leq 2 \exp \left\{ -c_1 n \min \left(c_\psi^2 \alpha_{\min}^2 / 54^2, 1 \right) / 2 \right\}.
\end{aligned}$$

Now by Lemma A.10, we have

$$\begin{aligned}
& \left| \{ \mathbf{W}^{\text{T}}, \text{vec}(\mathbf{V})^{\text{T}} \} n^{-1} \sum_{i=1}^n \left[\psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{Z}_i^{\text{T}}, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^{\text{T}} \right]^{\text{T}} \right)^{\otimes 2} \right. \right. \\
& \left. \left. - E \left\{ \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{Z}_i^{\text{T}}, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^{\text{T}} \right]^{\text{T}} \right)^{\otimes 2} \right\} \right] \{ \mathbf{W}^{\text{T}}, \text{vec}(\mathbf{V})^{\text{T}} \}^{\text{T}} \right| \\
& \leq c_\psi \alpha_{\min} / 2 + s^{-1} \left\{ \|\mathbf{W}\|_1 + \sum_{j=1}^d \|\mathbf{V}_{\cdot j}\|_2 \right\}^2, \forall \mathbf{W} \in \mathbb{R}^q, \mathbf{V} \in \mathbb{R}^{N \times d}.
\end{aligned}$$

This implies $\forall \mathbf{W} \in \mathbb{R}^q, \mathbf{V} \in \mathbb{R}^{N \times d}$,

$$\begin{aligned}
& \{ \mathbf{W}^{\text{T}}, \text{vec}(\mathbf{V})^{\text{T}} \} n^{-1} \sum_{i=1}^n \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{Z}_i^{\text{T}}, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^{\text{T}} \right]^{\text{T}} \right)^{\otimes 2} \{ \mathbf{W}^{\text{T}}, \text{vec}(\mathbf{V})^{\text{T}} \}^{\text{T}} \\
& \geq c_\psi \alpha_{\min} \|\{ \mathbf{W}^{\text{T}}, \text{vec}(\mathbf{V})^{\text{T}} \}^{\text{T}}\|_2^2 / 2 - s^{-1} \left\{ \|\mathbf{W}\|_1 + \sum_{j=1}^d \|\mathbf{V}_{\cdot j}\|_2 \right\}^2,
\end{aligned}$$

and

$$\begin{aligned}
& \{ \mathbf{W}^{\text{T}}, \text{vec}(\mathbf{V})^{\text{T}} \} n^{-1} \sum_{i=1}^n \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{Z}_i^{\text{T}}, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^{\text{T}} \right]^{\text{T}} \right)^{\otimes 2} \{ \mathbf{W}^{\text{T}}, \text{vec}(\mathbf{V})^{\text{T}} \}^{\text{T}} \\
& \leq 3c_\psi \alpha_{\max} \|\{ \mathbf{W}^{\text{T}}, \text{vec}(\mathbf{V})^{\text{T}} \}^{\text{T}}\|_2^2 / 2 + s^{-1} \left\{ \|\mathbf{W}\|_1 + \sum_{j=1}^d \|\mathbf{V}_{\cdot j}\|_2 \right\}^2
\end{aligned}$$

with probability greater than $1 - 2 \exp \left\{ -c_1 n \min \left(c_\psi^2 \alpha_{\min}^2 / 54^2, 1 \right) / 2 \right\}$. Recall that we have choose $s = c_1 n \min \left(c_\psi^2 \alpha_{\min}^2 / 54^2, 1 \right) / [10\{N + q + \log(p)\}]$. Now select $\tau(n, p) = c_\psi \alpha_{\min} / (2s)$, we prove the result. \square

Lemma A.14. *Assume Conditions (C1)–(C4) hold, there are constants $C > 0$ such that*

By union bound over $k = 1, \dots, q$ and $j = 1, \dots, d$, we have

$$\max \left(\left\| n^{-1} \sum_{i=1}^n [\psi'(\boldsymbol{\alpha}, \boldsymbol{\Gamma}_0) - \psi'\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\}] \mathbf{Z}_i \right\|_\infty, \right),$$

$$\begin{aligned} & \left\| n^{-1} \sum_{i=1}^n [\psi'(\boldsymbol{\alpha}, \boldsymbol{\Gamma}_0) - \psi'\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\}] N^{1/2} \int_0^1 X_{ij}(s) \mathbf{B}(s) ds \right\|_2, \quad j = 1, \dots, d \\ & \leq C k_0^{1/2} N^{-\omega} \end{aligned}$$

with probability $1 - 8p^{-1}$

Proof: First by Condition (C1) and (C3), we can show that there is a positive constant C_1 such that

$$\begin{aligned} & \left| \sum_{i=1}^n [\psi'(\boldsymbol{\alpha}, \boldsymbol{\Gamma}_0) - \psi'\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\}] Z_{ij} \right| \\ & = \sum_{i=1}^n |\psi''\{\boldsymbol{\alpha}, \boldsymbol{\beta}^*(\cdot)\}| \left\{ \int_0^1 \|\boldsymbol{\Gamma}_0^T \mathbf{B}(s) - \boldsymbol{\beta}(s)\|_2^2 ds \right\}^{1/2} \left(\int_0^1 \mathbf{X}_{i\mathcal{S}}(s)^T \mathbf{X}_{i\mathcal{S}}(s) ds \right)^{1/2} |Z_{ij}| \\ & \leq \sum_{i=1}^n |\psi''\{\boldsymbol{\alpha}, \boldsymbol{\beta}^*(\cdot)\}| \left\{ \int_0^1 \|\boldsymbol{\Gamma}_0^T \mathbf{B}(s) - \boldsymbol{\beta}(s)\|_2^2 ds \right\}^{1/2} \left(\int_0^1 \mathbf{X}_{i\mathcal{S}}^T(s) \mathbf{X}_{i\mathcal{S}}(s) ds \right)^{1/2} |Z_{ij}| \\ & = C_1 k_0^{1/2} 2 N^{-\omega} \sum_{i=1}^n \left(\int_0^1 \mathbf{X}_{i\mathcal{S}}^T(s) \mathbf{X}_{i\mathcal{S}}(s) ds \right)^{1/2} |Z_{ij}| \\ & \leq C_1 k_0^{1/2} N^{-\omega} \sum_{i=1}^n \left\{ \left(\int_0^1 \mathbf{X}_{i\mathcal{S}}^T(s) \mathbf{X}_{i\mathcal{S}}(s) ds \right) + |Z_{ij}|^2 \right\}. \end{aligned} \quad (6)$$

Now by Condition (C2), for any unit vector \mathbf{v} , $\mathbf{v}^T \int_0^1 \mathbf{X}_{i\mathcal{S}} \mathbf{X}_{i\mathcal{S}}(s)^T ds \mathbf{v}$ is a sub-exponential random variable and Z_{ij}^2 is a sub-exponential random variable, by Proposition 5.16 the Bernstein-type inequality in Vershynin [2010] and union bounded as those used in (5) and fact that $|\mathcal{S}| = k_0$, we have

$$\begin{aligned} & \Pr \left[\left| \sum_{i=1}^n \left\{ \left(\int_0^1 \mathbf{X}_{i\mathcal{S}}^T(s) \mathbf{X}_{i\mathcal{S}}(s) ds \right) - E \left(\int_0^1 \mathbf{X}_{i\mathcal{S}}^T(s) \mathbf{X}_{i\mathcal{S}}(s) ds \right) \right\} \right| \geq nt \right] \\ & \leq 2 \exp \left\{ -c'_3 n \min(t^2, t) + k_0 \log(9) \right\} \\ & \leq 2 \exp \left\{ -c'_3 n \min(t^2, t) + c'_0 \log\{\max(p, n)\} \log(9) \right\} \end{aligned}$$

for some constant $c'_3 > 0$, where the last inequality holds by Condition (C4). Therefore, by

selecting $t = B \sqrt{c_0 \log\{\max(p, n)\}/n}$, we have

$$\frac{\sum_{i=1}^n \int_0^1 \mathbf{X}_i^T(s) \mathbf{X}_i(s) ds}{n} \leq E \left(\int_0^1 \mathbf{X}_i^T(s) \mathbf{X}_i(s) ds \right) + B \sqrt{c_0 \log\{\max(p, n)\}/n}$$

with probability greater than $1 - 2 \exp \left[-c'_3 n \min \left\{ B^2 c_0 \log \{ \max(p, n) \} / n, B \sqrt{c_0 \log \{ \max(p, n) \} / n} \right\} \right]$,

where B satisfies $c'_3 c_0 \min(B^2, B) = c'_0 \log(9) + 2$ and c_0 is defined in Condition (C4) so that $c_0 \log \{ \max(p, n) \} / n \leq 1$. Similarly, we can show that

$$\frac{\sum_{i=1}^n Z_{ij}^2}{n} \leq E(Z_{ij}^2) + B_1 \sqrt{c_0 \log \{ \max(p, n) \} / n}$$

with probability greater than $1 - 2 \exp[-2 \log \{ \max(p, n) \}]$ for some constant $c'_3 > 0$ and

$c'_4 c_0 \min(B_1^2, B_1) = 2$. Plug in (6), we have

$$\begin{aligned} & \left| n^{-1} \sum_{i=1}^n [\psi'(\boldsymbol{\alpha}, \boldsymbol{\Gamma}_0) - \psi'\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\}] Z_{ij} \right| \\ & \leq C_1 k_0^{1/2} N^{-\omega} \left\{ E \left(\int_0^1 \mathbf{X}_i^\top(s) \mathbf{X}_i(s) ds \right) + E(Z_{ij}) + (B + B_1) \sqrt{c_0 \log \{ \max(p, n) \} / n} \right\} \end{aligned} \quad (7)$$

with probability greater than $1 - 4 \exp[-2 \log \{ \max(p, n) \}]$.

Similar to (6)

$$\begin{aligned} & \left\| \sum_{i=1}^n [\psi'(\boldsymbol{\alpha}, \boldsymbol{\Gamma}_0) - \psi'\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\}] N^{1/2} \int_0^1 X_{ij}(s) \mathbf{B}(s) ds \right\|_2 \\ & \leq C_2 k_0^{1/2} N^{-\omega} \sum_{i=1}^n \left\{ \left(\int_0^1 \mathbf{X}_i^\top(s) \mathbf{X}_i(s) ds \right) + N \int_0^1 X_{ij}(s) \mathbf{B}(s)^\top ds \int_0^1 X_{ij}(s) \mathbf{B}(s) ds \right\}. \end{aligned}$$

Also because $N \int_0^1 X_{ij}(s) \mathbf{B}(s)^\top ds$ is a sub-Gaussian random vector as shown in Lemma A.8,

similar to (7), there are constants B_3, B_4 such that

$$\begin{aligned} & \left\| n^{-1} \sum_{i=1}^n [\psi'(\boldsymbol{\alpha}, \boldsymbol{\Gamma}_0) - \psi'\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\}] N^{1/2} \int_0^1 X_{ij}(s) \mathbf{B}(s) ds \right\|_2 \\ & \leq C_2 k_0^{1/2} N^{-\omega} \left[E \left(\int_0^1 \mathbf{X}_i^\top(s) \mathbf{X}_i(s) ds \right) \right. \\ & \quad \left. + E \left\{ N \int_0^1 X_{ij}(s) \mathbf{B}(s)^\top ds \int_0^1 X_{ij}(s) \mathbf{B}(s) ds \right\} + (B_3 + B_4) \sqrt{\log \{ \max(p, n) \} / n} \right], \end{aligned}$$

with probability $1 - 4 \exp[-2 \log \{ \max(p, n) \}]$.

Now because for any unit vector \mathbf{v} , $\int_0^1 \mathbf{v}^\top \mathbf{X}_i(s) \mathbf{X}_i(s)^\top \mathbf{v} ds$, $N \int_0^1 X_{ij}(s) \mathbf{B}(s)^\top \mathbf{v} ds \int_0^1 X_{ij}(s) \mathbf{v}^\top \mathbf{B}(s) ds$, and Z_{ij}^2 are all sub-exponential random variable, their expectations are uniformly bounded

by Condition (C2) and by Condition (C4), we have there is a constant $C > 0$, for any given j, k , we have

$$\begin{aligned} & \max \left(\left| n^{-1} \sum_{i=1}^n [\psi'(\boldsymbol{\alpha}, \boldsymbol{\Gamma}_0) - \psi'\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\}] Z_{ik} \right|, \right. \\ & \left. \left\| n^{-1} \sum_{i=1}^n [\psi'(\boldsymbol{\alpha}, \boldsymbol{\Gamma}_0) - \psi'\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\}] N^{1/2} \int_0^1 X_{ij}(s) \mathbf{B}(s) ds \right\|_2 \right) \\ & \leq C k_0^{1/2} N^{-\omega} \end{aligned}$$

with probability $1 - 8 \exp[-2 \log\{\max(p, n)\}]$. By union bound over $k = 1, \dots, q$ and $j = 1, \dots, d$, we have

$$\begin{aligned} & \max \left(\left\| n^{-1} \sum_{i=1}^n [\psi'(\boldsymbol{\alpha}, \boldsymbol{\Gamma}_0) - \psi'\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\}] \mathbf{Z}_i \right\|_\infty, \right. \\ & \left. \left\| n^{-1} \sum_{i=1}^n [\psi'(\boldsymbol{\alpha}, \boldsymbol{\Gamma}_0) - \psi'\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\}] N^{1/2} \int_0^1 X_{ij}(s) \mathbf{B}(s) ds \right\|_2, j = 1, \dots, d \right) \\ & \leq C k_0^{1/2} N^{-\omega} \end{aligned}$$

with probability $1 - 8 \max(p, n)^{-1}$ □

Lemma A.15. *Assume Conditions (C1)–(C4) hold, there are constants $C_0, C > 0$*

$$\begin{aligned} & \max \left\{ \left\| n^{-1} \sum_{i=1}^n \{\psi'_i(\boldsymbol{\alpha}, \boldsymbol{\Gamma}_0) - Y_i\} \mathbf{Z}_i \right\|_\infty, \right. \\ & \left. \left\| n^{-1} \sum_{i=1}^n \{\psi'_i(\boldsymbol{\alpha}, \boldsymbol{\Gamma}_0) - Y_i\} \left[N^{1/2} \left\{ \int_0^1 X_{ij}(s) \otimes \mathbf{B}(s) ds \right\}^\top \right]^\top \right\|_2, j = 1, \dots, d \right\} \\ & \leq C_0 \sqrt{\log\{\max(p, n)\}/n} + C k_0^{1/2} N^{-\omega} \end{aligned}$$

with probability $1 - 20 \max(p, n)^{-1}$.

Proof:

$$n^{-1} \sum_{i=1}^n \{\psi'_i(\boldsymbol{\alpha}, \boldsymbol{\Gamma}_0) - Y_i\} \left[\mathbf{Z}_i^\top, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^\top \right]^\top$$

$$\begin{aligned}
&= n^{-1} \sum_{i=1}^n [\psi'_i\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\} - Y_i] \left[\mathbf{Z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \\
&\quad + n^{-1} \sum_{i=1}^n [\psi'_i(\boldsymbol{\alpha}, \boldsymbol{\Gamma}_0) - \psi'_i\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\}] \left[\mathbf{Z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \quad (8)
\end{aligned}$$

For the first term, follow Lemma 14 in Loh and Wainwright [2012], we have

$$\Pr \left\{ \left\| n^{-1} \sum_{i=1}^n \{\psi'_i\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\} - Y_i\} \mathbf{Z}_i \right\|_{\infty} \geq t \right\} \quad (9)$$

$$\leq 6q \exp\{-c_3 n \min(t^2, t)\} \quad (10)$$

for some constants $c_3 > 0$. Furthermore for any given j and any unit vector \mathbf{v} , let $\mathbf{U} = [\{\psi'_i(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) - Y_i\}, i = 1, \dots, n]$, $\mathbf{V} = \left(\mathbf{v}^T \left[N^{1/2} \left\{ \int_0^1 X_{ij}(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T, i = 1, \dots, n \right)$,

$$\begin{aligned}
& \left| n^{-1} \sum_{i=1}^n \{\psi'_i\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\} - Y_i\} \mathbf{v}^T \left[N^{1/2} \left\{ \int_0^1 X_{ij}(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right| \\
&= n^{-1} |\mathbf{U}^T \mathbf{V}| \\
&= \frac{1}{2} \{\Phi(\mathbf{U} + \mathbf{V}) - \Phi(\mathbf{U}) - \Phi(\mathbf{V})\}
\end{aligned}$$

where $\Phi(\mathbf{U}) \equiv \frac{\|\mathbf{U}\|_2^2}{n} - E\left(\frac{\|\mathbf{U}\|_2^2}{n}\right)$. Since by Condition (C2) and Lemma A.8, \mathbf{U} , \mathbf{V} are both sub-Gaussian, we have

$$\left| n^{-1} \sum_{i=1}^n \{\psi'_i\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\} - Y_i\} \mathbf{v}^T \left[N^{1/2} \left\{ \int_0^1 X_{ij}(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right| \leq 3t/2,$$

with probability greater than $1 - 6 \exp\{c_4 n \min(t^2, t)\}$. Now by the union bound, similar to

(5) for some constant $c_4 > 0$, we have

$$\begin{aligned}
& \Pr \left\{ \max_{j=1, \dots, d} \left\| n^{-1} \sum_{i=1}^n \{\psi'_i\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\} - Y_i\} \left[N^{1/2} \left\{ \int_0^1 X_{ij}(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right\|_2 \geq t \right\} \\
&= \Pr \left\{ \max_{j=1, \dots, d} \sup_{\|\mathbf{v}\|_2 \leq 1} \left| n^{-1} \sum_{i=1}^n \{\psi'_i\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\} - Y_i\} \mathbf{v}^T \left[N^{1/2} \left\{ \int_0^1 X_{ij}(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right| \geq t \right\} \\
&\leq 6p \exp[-c_4 n \min(t^2, t) + N \{\log(9)\}].
\end{aligned}$$

Combine with (9), let we have

$$\begin{aligned} & \max \left\{ \left\| n^{-1} \sum_{i=1}^n \{\psi'_i\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\} - Y_i\} \mathbf{Z}_i \right\|_{\infty}, \right. \\ & \left. \left\| n^{-1} \sum_{i=1}^n \{\psi'_i\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\} - Y_i\} \left[N^{1/2} \left\{ \int_0^1 X_{ij}(s) \otimes \mathbf{B}(s) ds \right\}^{\Gamma} \right]^{\Gamma} \right\|_2, j = 1, \dots, d \right\} \\ & \leq t \end{aligned}$$

with probability greater than

$$1 - 6p \exp[-c_4 n \min(t^2, t) + N \{\log(9)\}] - 6q \exp\{-c_3 n \min(t^2, t)\}$$

which is larger than

$$1 - 12p \exp[-\min(c_3, c_5) n \min(t^2, t) + c'_0 \log(p) \{\log(9)\}],$$

where c'_0 is defined in Condition (C4). Now select $t = C_0 \sqrt{c_0 \log\{\max(p, n)\}/n}$ so that $c_0 \min(c_3, c_5) \min(C_0, C_0^2) = c'_0 \log(9) + 2$, where c_0 is defined in Condition (C4), we have

$$\begin{aligned} & \max \left\{ \left\| n^{-1} \sum_{i=1}^n \{\psi'_i\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\} - Y_i\} \mathbf{Z}_i \right\|_{\infty}, \right. \\ & \left. \left\| n^{-1} \sum_{i=1}^n \{\psi'_i\{\boldsymbol{\alpha}, \boldsymbol{\beta}(\cdot)\} - Y_i\} \left[N^{1/2} \left\{ \int_0^1 X_{ij}(s) \otimes \mathbf{B}(s) ds \right\}^{\Gamma} \right]^{\Gamma} \right\|_2, j = 1, \dots, d \right\} \\ & \leq C_0 \sqrt{\log\{\max(p, n)\}/n} \end{aligned}$$

with probability greater than $1 - 12 \max(p, n)^{-1}$. Now combine with (8) and Lemma A.14,

we have there is a constant $C_0 > 0$

$$\begin{aligned} & \max \left\{ \left\| n^{-1} \sum_{i=1}^n \{\psi'_i(\boldsymbol{\alpha}, \boldsymbol{\Gamma}_0) - Y_i\} \mathbf{Z}_i \right\|_{\infty}, \right. \\ & \left. \left\| n^{-1} \sum_{i=1}^n \{\psi'_i(\boldsymbol{\alpha}, \boldsymbol{\Gamma}_0) - Y_i\} \left[N^{1/2} \left\{ \int_0^1 X_{ij}(s) \otimes \mathbf{B}(s) ds \right\}^{\Gamma} \right]^{\Gamma} \right\|_2, j = 1, \dots, d \right\} \\ & \leq C_0 \sqrt{\log\{\max(p, n)\}/n} + C k_0^{1/2} N^{-\omega} \end{aligned}$$

with probability $1 - 20 \max(p, n)^{-1}$. □

Proof of Theorem 1:

First by Taylor expansion of the first order derivative,

$$\left\{ \partial \mathcal{L}(\hat{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^T - \partial \mathcal{L}(\boldsymbol{\theta}_0) / \partial \boldsymbol{\theta}^T \right\} \hat{\mathbf{v}} = \hat{\mathbf{v}}^T \partial^2 \mathcal{L}(\boldsymbol{\theta}^*) / \partial \boldsymbol{\theta} \boldsymbol{\theta}^T \hat{\mathbf{v}},$$

where $\boldsymbol{\theta}^*$ is a point on the line between $\hat{\boldsymbol{\theta}}$ and $\boldsymbol{\theta}_0$, and hence is in the feasible set. Now by

Lemma A.13, we obtain

$$\left\{ \partial \mathcal{L}(\hat{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^T - \partial \mathcal{L}(\boldsymbol{\theta}_0) / \partial \boldsymbol{\theta}^T \right\} \hat{\mathbf{v}} \geq c_\psi \alpha_{\min} \|\hat{\mathbf{v}}\|_2^2 / 2 - \tau_1 \log(p) / n \left\{ \|\hat{\mathbf{v}}_1\|_1 + \sum_{j=1}^d \|\hat{\mathbf{v}}_{2j}\|_2 \right\}^2. \quad (11)$$

Since function $\rho_\lambda(\|\boldsymbol{\gamma}_j\|_2) + \mu/2\|\boldsymbol{\gamma}_j\|_2^2$ is convex in $\|\boldsymbol{\gamma}_j\|_2$ by Condition (A5), and L_2 norm is convex, we have $\rho_\lambda(\|\boldsymbol{\gamma}_j\|_2) + \mu/2\|\boldsymbol{\gamma}_j\|_2^2$ is a convex function on $\boldsymbol{\gamma}_j$.

Therefore, we have

$$\rho_\lambda(\|\boldsymbol{\gamma}_{0j}\|_2) + \mu/2\|\boldsymbol{\Gamma}_{0 \cdot j}\|_2^2 - \rho_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2) - \mu/2\|\hat{\boldsymbol{\gamma}}_j\|_2^2 \geq \left\{ \partial \rho_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2) / \partial \boldsymbol{\gamma}_j^T + \mu \hat{\boldsymbol{\gamma}}_j^T \right\} (\boldsymbol{\gamma}_{0j} - \hat{\boldsymbol{\gamma}}_j),$$

which implies

$$\begin{aligned} & \rho_\lambda(\|\boldsymbol{\Gamma}_{0 \cdot j}\|_2) - \rho_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2) + \mu/2\|\hat{\boldsymbol{\gamma}}_j - \boldsymbol{\gamma}_{0j}\|_2^2 \\ & \geq \partial \rho_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2) / \partial \boldsymbol{\gamma}_j^T (\boldsymbol{\gamma}_{0j} - \hat{\boldsymbol{\gamma}}_j). \end{aligned}$$

Therefore, combine with (1), we have

$$N^{-1/2} \sum_{j \in \mathcal{M}^c} \left\{ \rho_\lambda(\|\boldsymbol{\Gamma}_{0 \cdot j}\|_2) - \rho_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2) + \mu/2\|\hat{\boldsymbol{\gamma}}_j - \boldsymbol{\Gamma}_{0 \cdot j}\|_2^2 \right\} \geq \partial \mathcal{L}(\hat{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^T (\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0).$$

Combining with (11), we have

$$\begin{aligned} & c_\psi \alpha_{\min} \|\hat{\mathbf{v}}\|_2^2 / 2 - \tau_1 \log(p) / n \left\{ \|\hat{\mathbf{v}}_1\|_1 + \sum_{j=1}^d \|\hat{\mathbf{v}}_{2j}\|_2 \right\}^2 \\ & \leq -\partial \mathcal{L}(\boldsymbol{\theta}_0) / \partial \boldsymbol{\theta}^T \hat{\mathbf{v}} + N^{-1/2} \sum_{j \in \mathcal{M}^c} \left\{ \rho_\lambda(\|\boldsymbol{\Gamma}_{0 \cdot j}\|_2) - \rho_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2) + \mu/2\|\hat{\boldsymbol{\gamma}}_j - \boldsymbol{\Gamma}_{0 \cdot j}\|_2^2 \right\} \end{aligned}$$

$$\leq -\partial\mathcal{L}(\boldsymbol{\theta}_0)/\partial\boldsymbol{\theta}^T\widehat{\mathbf{v}} + N^{-1/2} \sum_{j \in \mathcal{M}^c} \{\rho_\lambda(\|\mathbf{\Gamma}_{0:j}\|_2) - \rho_\lambda(\|\widehat{\boldsymbol{\gamma}}_j\|_2) + \mu/2\|\widehat{\mathbf{v}}\|_2^2\},$$

with probability greater than $1 - 2 \exp\{-c_1 n \min(c_\psi^2 \alpha_{\min}^2/54^2, 1)/2\}$. This implies

$$\begin{aligned} & (c_\psi \alpha_{\min}/2 - \mu/2)\|\widehat{\mathbf{v}}\|_2^2 \\ \leq & \tau_1 \log(p)/n \left\{ \|\widehat{\mathbf{v}}_1\|_1 + \sum_{j=1}^d \|\widehat{\mathbf{v}}_{2j}\|_2 \right\}^2 - \partial\mathcal{L}(\boldsymbol{\theta}_0)/\partial\boldsymbol{\theta}^T\widehat{\mathbf{v}} + N^{-1/2} \sum_{j \in \mathcal{M}^c} \{\rho_\lambda(\|\mathbf{\Gamma}_{0:j}\|_2) - \rho_\lambda(\|\widehat{\boldsymbol{\gamma}}_j\|_2)\} \\ \leq & \tau_1 \log(p)/n \left\{ \|\widehat{\mathbf{v}}_1\|_1 + \sum_{j=1}^d \|\widehat{\mathbf{v}}_{2j}\|_2 \right\}^2 - \partial\mathcal{L}(\boldsymbol{\theta}_0)/\partial\boldsymbol{\alpha}^T\widehat{\mathbf{v}}_1 - \sum_{j=1}^d \partial\mathcal{L}(\boldsymbol{\theta}_0)/\partial\boldsymbol{\gamma}_j^T\widehat{\mathbf{v}}_{2j} \\ & + N^{-1/2} \sum_{j \in \mathcal{M}^c} \{\rho_\lambda(\|\mathbf{\Gamma}_{0:j}\|_2) - \rho_\lambda(\|\widehat{\boldsymbol{\gamma}}_j\|_2)\} \\ \leq & \tau_1 \log(p)/n \left\{ \|\widehat{\mathbf{v}}_1\|_1 + \sum_{j=1}^d \|\widehat{\mathbf{v}}_{2j}\|_2 \right\}^2 + \max_{j=1, \dots, d} \{\|\partial\mathcal{L}(\boldsymbol{\theta}_0)/\partial\boldsymbol{\alpha}\|_\infty, \|\partial\mathcal{L}(\boldsymbol{\theta}_0)/\partial\boldsymbol{\gamma}_j\|_2\} \left\{ \|\widehat{\mathbf{v}}_1\|_1 + \sum_{j=1}^d \|\widehat{\mathbf{v}}_{2j}\|_2 \right\} \\ & + N^{-1/2} \sum_{j \in \mathcal{M}^c} \{\rho_\lambda(\|\mathbf{\Gamma}_{0:j}\|_2) - \rho_\lambda(\|\widehat{\boldsymbol{\gamma}}_j\|_2)\} \\ \leq & \left[2\tau_1 R \log(p)/n + \max_{j=1, \dots, d} \{\|\partial\mathcal{L}(\boldsymbol{\theta}_0)/\partial\boldsymbol{\alpha}\|_\infty, \|\partial\mathcal{L}(\boldsymbol{\theta}_0)/\partial\boldsymbol{\gamma}_j\|_2\} \right] \left\{ \|\widehat{\mathbf{v}}_1\|_1 + \sum_{j=1}^d \|\widehat{\mathbf{v}}_{2j}\|_2 \right\} \\ & + N^{-1/2} \sum_{j \in \mathcal{M}^c} \{\rho_\lambda(\|\mathbf{\Gamma}_{0:j}\|_2) - \rho_\lambda(\|\widehat{\boldsymbol{\gamma}}_j\|_2)\} \\ \leq & \lambda/2 \left\{ \|\widehat{\mathbf{v}}_1\|_1 + \sum_{j=1}^d \|\widehat{\mathbf{v}}_{2j}\|_2 \right\} + N^{-1/2} \sum_{j \in \mathcal{M}^c} \{\rho_\lambda(\|\mathbf{\Gamma}_{0:j}\|_2) - \rho_\lambda(\|\widehat{\boldsymbol{\gamma}}_j\|_2)\} \\ = & \lambda/2 \left\{ \|\widehat{\mathbf{v}}_1\|_1 + \sum_{j \in \mathcal{M}} \|\widehat{\mathbf{v}}_{2j}\|_2 + \sum_{j \in \mathcal{M}^c} \|\widehat{\mathbf{v}}_{2j}\|_2 \right\} + N^{-1/2} \sum_{j \in \mathcal{M}^c} \{\rho_\lambda(\|\mathbf{\Gamma}_{0:j}\|_2) - \rho_\lambda(\|\widehat{\boldsymbol{\gamma}}_j\|_2)\} \\ \leq & \lambda/2 \left\{ \|\widehat{\mathbf{v}}_1\|_1 + \sum_{j \in \mathcal{M}} \|\widehat{\mathbf{v}}_{2j}\|_2 + \sum_{j \in \mathcal{M}^c} \rho_\lambda(\|\widehat{\mathbf{v}}_{2j}\|_2)/\lambda + \mu/(2\lambda)\|\widehat{\mathbf{v}}_{2j}\|_2^2 \right\} + N^{-1/2} \sum_{j \in \mathcal{M}^c} \{\rho_\lambda(\|\mathbf{\Gamma}_{0:j}\|_2) - \rho_\lambda(\|\widehat{\boldsymbol{\gamma}}_j\|_2)\} \\ \leq & \lambda/2 \left(\|\widehat{\mathbf{v}}_1\|_1 + \sum_{j \in \mathcal{M}} \|\widehat{\mathbf{v}}_{2j}\|_2 \right) + \frac{\sum_{j \in \mathcal{M}^c} \{\rho_\lambda(\|\widehat{\boldsymbol{\gamma}}_j\|_2) + \rho_\lambda(\|\mathbf{\Gamma}_{0:j}\|_2)\}}{2} + \mu/4\|\widehat{\mathbf{v}}\|_2^2 \\ & + N^{-1/2} \sum_{j \in \mathcal{M}^c} \{\rho_\lambda(\|\mathbf{\Gamma}_{0:j}\|_2) - \rho_\lambda(\|\widehat{\boldsymbol{\gamma}}_j\|_2)\} \\ = & \lambda/2 \left(\|\widehat{\mathbf{v}}_1\|_1 + \sum_{j \in \mathcal{M}} \|\widehat{\mathbf{v}}_{2j}\|_2 \right) + \mu/4\|\widehat{\mathbf{v}}\|_2^2 + N^{-1/2}/2 \sum_{j \in \mathcal{M}^c} \{3\rho_\lambda(\|\mathbf{\Gamma}_{0:j}\|_2) - \rho_\lambda(\|\widehat{\boldsymbol{\gamma}}_j\|_2)\}, \end{aligned} \tag{12}$$

with probability greater than $1 - 2 \exp\{-c_1 n \min(c_\psi^2 \alpha_{\min}^2/54^2, 1)/2\} - 20 \max(p, n)^{-1}$. Here

the sixth line holds by the assumption that

$$\lambda \geq \max \left\{ 2\tau_1 R \log\{\max(p)\}/n, C_0 \sqrt{\log\{\max(p, n)\}/n} + Ck_0^{1/2} N^{-\omega} \right\} / 4,$$

and the fact that

$$\max_{j=1, \dots, d} \left\{ \|\partial \mathcal{L}(\boldsymbol{\theta}_0) / \partial \boldsymbol{\alpha}\|_\infty, \|\partial \mathcal{L}(\boldsymbol{\theta}_0) / \partial \boldsymbol{\gamma}_j\|_2 \right\} \leq C_0 \sqrt{\log\{\max(p, n)\}/n} + Ck_0^{1/2} N^{-\omega},$$

with probability greater than $1 - 20 \max(p, n)^{-1}$ by Lemma A.15. The eighth line holds by Lemma A.2. Now if

$$\sum_{j \in \mathcal{M}^c} \{3\rho_\lambda(\|\boldsymbol{\Gamma}_{0,j}\|_2) - \rho_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2)\} \leq 0,$$

by (12), we have

$$(c_\psi \alpha_{\min} / 2 - 3\mu / 4) \|\hat{\mathbf{v}}\|_2^2 \leq \lambda / 2 \left(\|\hat{\mathbf{v}}_1\|_1 + \sum_{j \in \mathcal{M}} \|\hat{\mathbf{v}}_{2j}\|_2 \right) \leq \lambda / 2 \sqrt{q + m} \|\hat{\mathbf{v}}\|_2,$$

which implies

$$\|\hat{\mathbf{v}}\|_2 \leq \lambda \sqrt{q + m} / (c_\psi \alpha_{\min} - 3\mu / 2). \quad (13)$$

On the other hand if

$$\sum_{j \in \mathcal{M}^c} \{3\rho_\lambda(\|\boldsymbol{\Gamma}_{0,j}\|_2) - \rho_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2)\} > 0.$$

by Lemma (A.3), we have

$$0 \leq N^{-1/2} \sum_{j \in \mathcal{M}^c} \{3\rho_\lambda(\|\boldsymbol{\Gamma}_{0,j}\|_2) - \rho_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2)\} \leq \lambda \left(3 \sum_{j \in \mathcal{A}} \|\hat{\mathbf{v}}_{2j}\|_2 - \sum_{j \in \mathcal{A}^c} \|\hat{\mathbf{v}}_{2j}\|_2 \right), \quad (14)$$

where \mathcal{A} is the index set of the k_0 largest $\|\hat{\mathbf{v}}_{2j}\|_2$, $j \in \mathcal{M}^c$, and \mathcal{A}^c is the index set of the remaining $d - m - k_0$ of $\hat{\mathbf{v}}_{2j}$, $j \in \mathcal{M}^c$, which implies

$$\sum_{j \in \mathcal{A}^c} \|\hat{\mathbf{v}}_{2j}\|_2 \leq 3 \sum_{j \in \mathcal{A}} \|\hat{\mathbf{v}}_{2j}\|_2$$

and hence

$$\sum_{j \in \mathcal{M}^c} \|\widehat{\mathbf{v}}_{2j}\|_2 \leq 4 \sum_{j \in \mathcal{A}} \|\widehat{\mathbf{v}}_{2j}\|_2.$$

Plug (14) into (12), we have

$$\begin{aligned} (c_\psi \alpha_{\min}/2 - 3\mu/4) \|\widehat{\mathbf{v}}\|_2^2 &\leq \lambda/2 \left(\|\widehat{\mathbf{v}}_1\|_1 + \sum_{j \in \mathcal{M}} \|\widehat{\mathbf{v}}_{2j}\|_2 \right) + \lambda/2 \left(3 \sum_{j \in \mathcal{A}} \|\widehat{\mathbf{v}}_{2j}\|_2 - \sum_{j \in \mathcal{A}^c} \|\widehat{\mathbf{v}}_{2j}\|_2 \right) \\ &\leq \lambda/2 \left(\|\widehat{\mathbf{v}}_1\|_1 + \sum_{j \in \mathcal{M}} \|\widehat{\mathbf{v}}_{2j}\|_2 \right) + \lambda/2 \left(3 \sum_{j \in \mathcal{A}} \|\widehat{\mathbf{v}}_{2j}\|_2 \right) \\ &\leq 3\lambda/2 \left(\|\widehat{\mathbf{v}}_1\|_1 + \sum_{j \in \mathcal{M}} \|\widehat{\mathbf{v}}_{2j}\|_2 + \sum_{j \in \mathcal{A}} \|\widehat{\mathbf{v}}_{2j}\|_2 \right) \\ &\leq 3\lambda/2 \sqrt{q+m+k_0} \left\{ \|\widehat{\mathbf{v}}_1\|_2^2 + \sum_{j \in \mathcal{M} \cup \mathcal{A}} \|\widehat{\mathbf{v}}_{2j}\|_2^2 \right\}^{1/2} \\ &\leq 3\lambda \sqrt{q+m+k_0}/2 \|\widehat{\mathbf{v}}\|_2. \end{aligned}$$

This gives

$$\|\widehat{\mathbf{v}}\|_2 \leq 3\lambda \sqrt{q+m+k_0}/(c_\psi \alpha_{\min} - 3\mu/2). \quad (15)$$

Finally, since

$$\sum_{j \in \mathcal{A}^c} \|\widehat{\mathbf{v}}_{2j}\|_2 \leq 3 \sum_{j \in \mathcal{A}} \|\widehat{\mathbf{v}}_{2j}\|_2$$

we have

$$\begin{aligned} &\|\widehat{\mathbf{v}}_1\|_1 + \sum_{j=1}^d \|\widehat{\mathbf{v}}_{2j}\|_2 \\ &= \|\widehat{\mathbf{v}}_1\|_1 + \sum_{j \in \mathcal{M}} \|\widehat{\mathbf{v}}_{2j}\|_2 + \sum_{j \in \mathcal{A}} \|\widehat{\mathbf{v}}_{2j}\|_2 + \sum_{j \in \mathcal{A}^c} \|\widehat{\mathbf{v}}_{2j}\|_2 \\ &\leq \|\widehat{\mathbf{v}}_1\|_1 + \sum_{j \in \mathcal{M}} \|\widehat{\mathbf{v}}_{2j}\|_2 + 4 \sum_{j \in \mathcal{A}} \|\widehat{\mathbf{v}}_{2j}\|_2 \\ &\leq 4\sqrt{q+m+k_0} \|\widehat{\mathbf{v}}\|_2 \\ &\leq 12\lambda(q+m+k_0)/(c_\psi \alpha_{\min} - 3\mu/2). \end{aligned} \quad (16)$$

Combine (13), (15) and (16), we prove the result. \square

Lemma A.16. Let $\mathbf{x}^* = (\mathbf{x}_1^{*\top}, \mathbf{x}_{2j}^{*\top}, j = 1, \dots, d)^\top$ is feasible for the program

$$\min_{\mathbf{x} \equiv (\mathbf{x}_1^\top, \mathbf{x}_{2j}^\top, j=1, \dots, d)^\top} \left\{ f(\mathbf{x}) - \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_j\|_2) + \lambda \sum_{j \in \mathcal{M}^c} \|\mathbf{x}_j\|_2 \right\} \text{ such that } \|\mathbf{x}_1\|_1 + \sum_{j=1}^d \|\mathbf{x}_{2j}\|_2 \leq R, \quad (17)$$

where $f(\mathbf{x}) \in C^2$, $g(\|\mathbf{x}\|)$ is differentiable in \mathbf{x} , and $g(\|\mathbf{x}\|_2) - \kappa/2\|\mathbf{x}\|_2^2$ is concave in $\|\mathbf{x}_{2j}\|_2$.

Assume there are $\mu_1^* > 0$, $\mathbf{v}_j^* \in \partial\|\mathbf{x}_{2j}^*\|_2$ for $j \in \mathcal{M}^c$ and $\mathbf{v}_j^* = 0$ for $j \in \mathcal{M}$, $\mathbf{v}^* = (\mathbf{0}_q^\top, \mathbf{v}_j^{*\top}, j = 1, \dots, d)^\top$, and $\mathbf{w}^* \in \partial\{\|\mathbf{x}_1^*\|_1 + \sum_{j=1}^d \|\mathbf{x}_{2j}^*\|_2\}$ such that

$$\mu_1^* \left\{ R - \|\mathbf{x}_1\|_1 - \sum_{j=1}^d \|\mathbf{x}_{2j}\|_2 \right\} = 0 \quad (18)$$

$$\frac{\partial f(\mathbf{x}^*)}{\partial \mathbf{x}^*} - \frac{\partial \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^*\|_2)}{\partial \mathbf{x}^*} + \lambda \mathbf{v}^* + \mu_1^* \mathbf{w}^* = 0 \quad (19)$$

$$\mathbf{s}^\top \frac{\partial^2 f(\mathbf{x})}{\partial \mathbf{x} \partial \mathbf{x}^\top} \mathbf{s} > \kappa, \forall \mathbf{s} \in G^*, \quad (20)$$

where

$$G^* \equiv \left\{ \mathbf{s} \equiv (\mathbf{s}_1^\top, \mathbf{s}_{2j}^\top, j = 1, \dots, d)^\top \in \mathbb{R}^{q+Nd} : \|\mathbf{s}\|_2 = 1; \right. \\ \left. \sup_{\mathbf{w} \in \partial\{\|\mathbf{x}_1^*\|_1 + \sum_{j=1}^d \|\mathbf{x}_{2j}^*\|_2\}} \mathbf{s}^\top \mathbf{w} \leq 0 \text{ if } \|\mathbf{x}_1^*\|_1 + \sum_{j=1}^d \|\mathbf{x}_{2j}^*\|_2 \leq R; \right. \\ \left. \sup_{(\mathbf{v}_j \in \partial\|\mathbf{x}_{2j}^*\|_2, j \in \mathcal{M}^c)} \mathbf{s}^\top \left(\frac{\partial f(\mathbf{x}^*)}{\partial \mathbf{x}^*} - \frac{\partial \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^*\|_2)}{\partial \mathbf{x}^*} \right) + \lambda \sum_{j \in \mathcal{M}^c} \mathbf{s}_{2j}^\top \mathbf{v}_j = 0; \right. \\ \left. \mu_1^* \sup_{\mathbf{w} \in \partial\{\|\mathbf{x}_1^*\|_1 + \sum_{j=1}^d \|\mathbf{x}_{2j}^*\|_2\}} \mathbf{s}^\top \mathbf{w} = 0 \right\}.$$

Then \mathbf{x}^* is an isolated local minimum of the program (17)

Proof: Suppose \mathbf{x}^* is not an isolated local minimum. Then there is a sequence $\{\mathbf{x}^k \equiv (\mathbf{x}_1^{k\top}, \mathbf{x}_{2j}^{k\top}, j = 1, \dots, d)^\top\}$, so that $\mathbf{x}^k \rightarrow \mathbf{x}^*$ and

$$\phi(\mathbf{x}^k) \leq \phi(\mathbf{x}^*),$$

where $\phi(\mathbf{x}) = f(\mathbf{x}) - \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_j\|_2) + \lambda \sum_{j \in \mathcal{M}^c} \|\mathbf{x}_j\|_2$. Let $\mathbf{s}^k \equiv (\mathbf{x}^k - \mathbf{x}^*) / \|\mathbf{x}^k - \mathbf{x}^*\|_2$, so (\mathbf{s}^k) is a set of feasible directions. Since $(\mathbf{s}^k) \subset \mathbb{B}_2(1)$, where $\mathbb{B}_2(1)$ is the L_2 ball with radius 1,

the set must possess a point of accumulation $\mathbf{s} \in \mathbb{B}_2(1)$, and we can extract a convergence subsequence such that $(\mathbf{s}^k) \rightarrow \mathbf{s}$. With a slight abuse of notation, we still use (\mathbf{s}^k) to denote the subsequence. We will show that $\mathbf{s} \in G^*$.

First because the feasible region is closed, \mathbf{s} is also in the feasible direction of \mathbf{x}^* . If $\|\mathbf{x}_1^*\|_1 + \sum_{j=1}^d \|\mathbf{x}_{2j}^*\|_2 = R$, by the sub-gradient of convexity function of $\|\cdot\|_1$ and $\|\cdot\|_2$, we have

$$\begin{aligned}
0 &\geq \|\mathbf{x}_1^k\|_1 + \sum_{j=1}^d \|\mathbf{x}_{2j}^k\|_2 - \|\mathbf{x}_1^*\|_1 - \sum_{j=1}^d \|\mathbf{x}_{2j}^*\|_2 \\
&= \|\mathbf{x}_1^*\|_1 + \|\mathbf{x}^k - \mathbf{x}^*\|_2 \mathbf{s}_1^k - \|\mathbf{x}_1^*\|_1 + \sum_{j=1}^d \{ \|\mathbf{x}_{2j}^*\|_2 + \|\mathbf{x}^k - \mathbf{x}^*\|_2 \mathbf{s}_{2j}^k - \|\mathbf{x}_{2j}^*\|_2 \} \\
&\geq \|\mathbf{x}^k - \mathbf{x}^*\|_2 \mathbf{s}_1^{kT} \mathbf{w}_1 + \|\mathbf{x}^k - \mathbf{x}^*\|_2 \sum_{j=1}^d \mathbf{s}_{2j}^{kT} \mathbf{w}_{2j} \\
&= \|\mathbf{x}^k - \mathbf{x}^*\|_2 \mathbf{s}^{kT} \mathbf{w},
\end{aligned}$$

where $\mathbf{w} = (\mathbf{w}_1^T, \mathbf{w}_{2j}^T, j = 1, \dots, d)^T \in \partial(\|\mathbf{x}_1^*\|_1 + \sum_{j=1}^d \|\mathbf{x}_{2j}^*\|_2)$. When $k \rightarrow \infty$, this leads to

$$\sup_{\mathbf{w} \in \partial\{\|\mathbf{x}_1^*\|_1 + \sum_{j=1}^d \|\mathbf{x}_{2j}^*\|_2\}} \mathbf{s}^T \mathbf{w} \leq 0. \tag{21}$$

Furthermore, (18) also implies that if $\|\mathbf{x}_1^*\|_1 + \sum_{j=1}^d \|\mathbf{x}_{2j}^*\|_2 \neq R$, then $\mu_1^* = 0$. Since $\mu_1^* \geq 0$, we have

$$\mu_1^* \sup_{\mathbf{w} \in \partial\{\|\mathbf{x}_1^*\|_1 + \sum_{j=1}^d \|\mathbf{x}_{2j}^*\|_2\}} \mathbf{s}^T \mathbf{w} \leq 0. \tag{22}$$

Moreover, by (19), we have

$$\mathbf{s}^T \frac{\partial f(\mathbf{x}^*)}{\partial \mathbf{x}^*} - \sum_{j \in \mathcal{M}^c} \mathbf{s}_{2j}^T \frac{\partial g(\|\mathbf{x}_{2j}^*\|_2)}{\partial \mathbf{x}_{2j}} + \lambda \sum_{j \in \mathcal{M}^c} \mathbf{s}_{2j}^T \mathbf{v}_j^* + \mu_1^* \mathbf{s}^T \mathbf{w}^* = 0.$$

Together with (22), we have

$$\mathbf{s}^T \frac{\partial f(\mathbf{x}^*)}{\partial \mathbf{x}^*} - \sum_{j \in \mathcal{M}^c} \mathbf{s}_{2j}^T \frac{\partial g(\|\mathbf{x}_{2j}^*\|_2)}{\partial \mathbf{x}_{2j}} + \lambda \sum_{j \in \mathcal{M}^c} \mathbf{s}_{2j}^T \mathbf{v}_j^* = -\mu_1^* \mathbf{s}^T \mathbf{w}^* \geq 0. \tag{23}$$

By the definition of sub-gradient, we have

$$\|\mathbf{x}_{2j}^* + \|\mathbf{x}^k - \mathbf{x}^*\|_2 \mathbf{s}_{2j}^k\|_2 - \|\mathbf{x}_{2j}^*\|_2 \geq \|\mathbf{x}^k - \mathbf{x}^*\|_2 \mathbf{s}_{2j}^{k\top} \mathbf{v}_j$$

for all $\mathbf{v}_j \in \partial\|\mathbf{x}_{2j}^*\|_2, j \in \mathcal{M}^c$ and all k . Furthermore because

$$\|\mathbf{x}_{2j}^* + \|\mathbf{x}^k - \mathbf{x}^*\|_2 \mathbf{s}_{2j}^k\|_2 - \|\mathbf{x}_{2j}^*\|_2 = \|\mathbf{x}_{2j}^k\|_2 - \|\mathbf{x}_{2j}^*\|_2,$$

we have

$$\mathbf{s}_{2j}^{\top} \mathbf{v}_j = \lim_{k \rightarrow \infty} \mathbf{s}_{2j}^{k\top} \mathbf{v}_j \leq \lim_{k \rightarrow \infty} \frac{\|\mathbf{x}_{2j}^k\|_2 - \|\mathbf{x}_{2j}^*\|_2}{\|\mathbf{x}^k - \mathbf{x}^*\|_2}. \quad (24)$$

Furthermore,

$$\begin{aligned} & \mathbf{s}^{\top} \frac{\partial f(\mathbf{x}^*)}{\partial \mathbf{x}^*} - \sum_{j \in \mathcal{M}^c} \mathbf{s}_{2j}^{\top} \frac{\partial g(\|\mathbf{x}_{2j}^*\|_2)}{\partial \mathbf{x}_{2j}^*} \\ &= \lim_{k \rightarrow \infty} \mathbf{s}^{k\top} \frac{\partial f(\mathbf{x}^*)}{\partial \mathbf{x}^*} - \sum_{j \in \mathcal{M}^c} \mathbf{s}_{2j}^{k\top} \frac{\partial g(\|\mathbf{x}_{2j}^*\|_2)}{\partial \mathbf{x}_{2j}^*} \\ &= \lim_{k \rightarrow \infty} \frac{\langle \mathbf{x}^k - \mathbf{x}^*, \frac{\partial f(\mathbf{x}^*)}{\partial \mathbf{x}^*} \rangle - \sum_{j \in \mathcal{M}^c} \langle \mathbf{x}_{2j}^k - \mathbf{x}_{2j}^*, \frac{\partial g(\|\mathbf{x}_{2j}^*\|_2)}{\partial \mathbf{x}_{2j}^*} \rangle}{\|\mathbf{x}^k - \mathbf{x}^*\|_2} \\ &= \lim_{k \rightarrow \infty} \frac{f(\mathbf{x}^k) - \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^k\|_2) - f(\mathbf{x}^*) + \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^*\|_2)}{\|\mathbf{x}^k - \mathbf{x}^*\|_2} \end{aligned}$$

since $\mathbf{x}^k \rightarrow \mathbf{x}^*$ and $f(\mathbf{x}^k) - \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^k\|_2)$ is differentiable in \mathbf{x}^k . Combine with (24), we

have

$$\begin{aligned} & \sup_{(\mathbf{v}_j \in \partial\|\mathbf{x}_{2j}^*\|_2, j \in \mathcal{M}^c)} \mathbf{s}^{\top} \frac{\partial f(\mathbf{x}^*)}{\partial \mathbf{x}^*} - \sum_{j \in \mathcal{M}^c} \mathbf{s}_{2j}^{\top} \frac{\partial g(\|\mathbf{x}_{2j}^*\|_2)}{\partial \mathbf{x}_{2j}^*} + \lambda \sum_{j \in \mathcal{M}^c} \mathbf{s}_{2j}^{\top} \mathbf{v}_j \\ & \leq \lim_{k \rightarrow \infty} \frac{\phi(\mathbf{x}^k) - \phi(\mathbf{x}^*)}{\|\mathbf{x}^k - \mathbf{x}^*\|_2} \leq 0. \end{aligned}$$

Combine with (23), we have

$$\sup_{(\mathbf{v}_j \in \partial\|\mathbf{x}_{2j}^*\|_2, j \in \mathcal{M}^c)} \mathbf{s}^{\top} \frac{\partial f(\mathbf{x}^*)}{\partial \mathbf{x}^*} - \sum_{j \in \mathcal{M}^c} \mathbf{s}_{2j}^{\top} \frac{\partial g(\|\mathbf{x}_{2j}^*\|_2)}{\partial \mathbf{x}_{2j}^*} + \lambda \sum_{j \in \mathcal{M}^c} \mathbf{s}_{2j}^{\top} \mathbf{v}_j = \mathbf{0}. \quad (25)$$

Now by (19), we have

$$\mu_1^* \mathbf{s}^\top \mathbf{w}^* = 0.$$

Together with (22), we have

$$\mu_1^* \sup_{\mathbf{w} \in \partial\{\|\mathbf{x}_1^*\|_1 + \sum_{j=1}^d \|\mathbf{x}_{2j}^*\|_2\}} \mathbf{s}^\top \mathbf{w} = 0.$$

Combine with (21) and (25), we have $\mathbf{s} \in G^*$.

By the convexity of L_1 , L_2 norms, we have

$$\|\mathbf{x}_1^k\|_1 - \|\mathbf{x}_1^*\|_1 + \sum_{j=1}^d \{\|\mathbf{x}_{2j}^k\|_2 - \|\mathbf{x}_{2j}^*\|_2\} \geq \mathbf{x}_1^{k^\top} \mathbf{w}_1 - \|\mathbf{x}_1^*\|_1 + \sum_{j=1}^d (\mathbf{x}_{2j}^{k^\top} \mathbf{w}_{2j} - \|\mathbf{x}_{2j}^*\|_2)$$

for $\mathbf{w} = (\mathbf{w}_1^\top, \mathbf{w}_{2j}^\top, j = 1, \dots, d)^\top \in \partial(\|\mathbf{x}_1^*\|_1 + \sum_{j=1}^d \|\mathbf{x}_{2j}^*\|_2)$. Therefore, $\mathbf{x}_1^{k^\top} \mathbf{w}_1 + \sum_{j=1}^d \mathbf{x}_{2j}^{k^\top} \mathbf{w}_{2j} \leq \|\mathbf{x}_1^k\|_1 + \sum_{j=1}^d \|\mathbf{x}_{2j}^k\|_2 \leq R$. Therefore,

$$\begin{aligned} \phi(\mathbf{x}^k) &= f(\mathbf{x}^k) - \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^k\|_2) + \lambda \sum_{j \in \mathcal{M}^c} \|\mathbf{x}_{2j}^k\|_2 \\ &\geq f(\mathbf{x}^k) - \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^k\|_2) + \lambda \sum_{j \in \mathcal{M}^c} \mathbf{x}_{2j}^{k^\top} \mathbf{v}_j^* + \mu_1^* \left(\mathbf{x}_1^{k^\top} \mathbf{w}_1^* + \sum_{j=1}^d \mathbf{x}_{2j}^{k^\top} \mathbf{w}_{2j}^* - R \right), \end{aligned}$$

for all $\mathbf{v}^* = (\mathbf{0}_q, \mathbf{v}_j^{*\top}, j = 1, \dots, d)$, $\mathbf{v}_j^* \in \partial\|\mathbf{x}_{2j}^*\|_2$ for $j \in \mathcal{M}^c$ and $\mathbf{v}_j^* = 0$ for $j \in \mathcal{M}$. And

$$\begin{aligned} \phi(\mathbf{x}^*) &= f(\mathbf{x}^*) - \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^*\|_2) + \lambda \sum_{j \in \mathcal{M}^c} \|\mathbf{x}_{2j}^*\|_2 \\ &= f(\mathbf{x}^*) - \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^*\|_2) + \lambda \sum_{j \in \mathcal{M}^c} \mathbf{x}_{2j}^{*\top} \mathbf{v}_j^* + \mu_1^* \left(\mathbf{x}_1^{*\top} \mathbf{w}_1^* + \sum_{j=1}^d \mathbf{x}_{2j}^{*\top} \mathbf{w}_{2j}^* - R \right). \end{aligned}$$

The equality holds because $\mu_1^* \left(\mathbf{x}_1^{*\top} \mathbf{w}_1^* + \sum_{j=1}^d \mathbf{x}_{2j}^{*\top} \mathbf{w}_{2j}^* - R \right) = 0$, and $\mathbf{x}_{2j}^{*\top} \mathbf{v}_j^* = \|\mathbf{x}_{2j}^*\|_2$.

Therefore, we can write

$$\begin{aligned} &\lim_{k \rightarrow \infty} \{\phi(\mathbf{x}^k) - \phi(\mathbf{x}^*)\} / \|\mathbf{x}^k - \mathbf{x}^*\|_2^2 \\ &\geq \lim_{k \rightarrow \infty} \left\{ f(\mathbf{x}^k) - f(\mathbf{x}^*) - \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^k\|_2) + \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^*\|_2) + \langle \lambda \mathbf{v}^* + \mu_1^* \mathbf{w}^*, \mathbf{x}^k - \mathbf{x}^* \rangle \right\} \end{aligned}$$

$$\begin{aligned}
& / \|\mathbf{x}^k - \mathbf{x}^*\|_2^2 \\
= & \lim_{k \rightarrow \infty} \left\{ f(\mathbf{x}^k) - f(\mathbf{x}^*) - \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^k\|_2) + \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^*\|_2) \right\} / \|\mathbf{x}^k - \mathbf{x}^*\|_2^2 \\
& - \left\langle \frac{\partial f(\mathbf{x}^*)}{\partial \mathbf{x}^*} - \frac{\partial \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^*\|_2)}{\partial \mathbf{x}^*}, \mathbf{x}^k - \mathbf{x}^* \right\rangle / \|\mathbf{x}^k - \mathbf{x}^*\|_2^2 \\
= & \lim_{k \rightarrow \infty} \left\{ f(\mathbf{x}^k) - f(\mathbf{x}^*) - \left\langle \frac{\partial f(\mathbf{x}^*)}{\partial \mathbf{x}^*}, \mathbf{x}^k - \mathbf{x}^* \right\rangle \right\} / \|\mathbf{x}^k - \mathbf{x}^*\|_2^2 \\
& - \lim_{k \rightarrow \infty} \left\{ \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^k\|_2) - \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^*\|_2) - \left\langle \frac{\partial \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^*\|_2)}{\partial \mathbf{x}^*}, \mathbf{x}^k - \mathbf{x}^* \right\rangle \right\} \\
& / \|\mathbf{x}^k - \mathbf{x}^*\|_2^2 \tag{26}
\end{aligned}$$

Now because $g(\|\mathbf{x}\|_2) - \kappa/2\|\mathbf{x}\|_2^2$ is concave in $\|\mathbf{x}\|_2$, and hence $-g(\|\mathbf{x}\|_2) + \kappa/2\|\mathbf{x}\|_2^2$ is convex in \mathbf{x} . Therefore, we have

$$- \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^k\|_2) + \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^*\|_2) + \left\langle \frac{\partial \sum_{j \in \mathcal{M}^c} g(\|\mathbf{x}_{2j}^*\|_2)}{\partial \mathbf{x}^*}, \mathbf{x}^k - \mathbf{x}^* \right\rangle \geq -k/2\|\mathbf{x}^k - \mathbf{x}^*\|_2^2.$$

Furthermore note that $\phi(\mathbf{x}^k) - \phi(\mathbf{x}^*) \leq 0$, combine with (26), we have

$$\lim_{k \rightarrow \infty} \left\{ f(\mathbf{x}^k) - f(\mathbf{x}^*) - \left\langle \frac{\partial f(\mathbf{x}^*)}{\partial \mathbf{x}^*}, \mathbf{x}^k - \mathbf{x}^* \right\rangle \right\} / \|\mathbf{x}^k - \mathbf{x}^*\|_2^2 - k/2 \leq 0,$$

which by Talyor expansion implies

$$\begin{aligned}
\mathbf{s}^\top \frac{\partial^2 f(\mathbf{x}^*)}{\partial \mathbf{x} \partial \mathbf{x}^\top} \mathbf{s} &= \lim_{k \rightarrow \infty} (\mathbf{x}^k - \mathbf{x}^*)^\top \frac{\partial^2 f(\mathbf{x}^*)}{\partial \mathbf{x} \partial \mathbf{x}^\top} (\mathbf{x}^k - \mathbf{x}^*) / \|\mathbf{x}^k - \mathbf{x}^*\|_2^2 \\
&\leq \lim_{k \rightarrow \infty} \{ \kappa \|\mathbf{x}^k - \mathbf{x}^*\|_2^2 + o(\|\mathbf{x}^k - \mathbf{x}^*\|_2^2) \} / \|\mathbf{x}^k - \mathbf{x}^*\|_2^2 \\
&\leq \kappa,
\end{aligned}$$

which contradicts with (20). Hence \mathbf{x}^* must be an isolated local minimum. This proves the result. \square

Lemma A.17. *Assume Conditions (A1)–(A6) and (C1) – (C5) hold. Assume $\mu < c_\psi \alpha_{\min}/2$, $\delta \in [4R\tau_1 \log(p)/n/\lambda, 1]$, $\max_{j \in (\mathcal{M} \cup \mathcal{S})^c} \|\hat{\mathbf{z}}_j\|_2 \leq 1 - \delta$ for $\hat{\mathbf{z}}_j \in \partial \|N^{-1/2} \hat{\gamma}_j\|_2$. Furthermore, assume that $n \geq 8\tau_1(\log(p) + N + q) \{ \sqrt{q} + \sqrt{m} + (2/\delta + 2)\sqrt{k_0} \}^2 / (c_\psi \alpha_{\min}/2 - \mu)$. Suppose*

$\tilde{\boldsymbol{\theta}} \equiv \{\tilde{\boldsymbol{\alpha}}^T, N^{-1/2} \text{vec}(\tilde{\boldsymbol{\Gamma}})\}^T$ is a stationary point of program 3 and $\hat{\boldsymbol{\theta}} \equiv \{\hat{\boldsymbol{\alpha}}^T, N^{-1/2} \text{vec}(\hat{\boldsymbol{\Gamma}})\}^T$ is the interior local minimizer of 3 and satisfies $\text{supp}\{\|\hat{\boldsymbol{\gamma}}_j\|_2, j = 1, \dots, d\} \subset \mathcal{M} \cup S$. Then there is a constant $c_1 > 0$ such that $\text{supp}\{\|\tilde{\boldsymbol{\gamma}}_j\|_2, j = 1, \dots, d\} \subset \mathcal{M} \cup S$ with probability $1 - 2 \exp\{-c_1 n \min(c_\psi^2 \alpha_{\min}^2 / 54^2, 1) / 2\}$.

Proof: Let $\tilde{\mathbf{v}} \equiv (\tilde{\mathbf{v}}_1^T, \tilde{\mathbf{v}}_{2j}^T, j = 1, \dots, d)^T \equiv \tilde{\boldsymbol{\theta}} - \hat{\boldsymbol{\theta}} = \{\tilde{\boldsymbol{\alpha}}^T - \hat{\boldsymbol{\alpha}}^T, N^{-1/2} \text{vec}(\tilde{\boldsymbol{\Gamma}})^T - N^{-1/2} \text{vec}(\hat{\boldsymbol{\Gamma}})^T\}^T$ by the Taylor expansion of the first order derivative we have

$$\left\{ \partial \mathcal{L}(\tilde{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^T - \partial \mathcal{L}(\hat{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^T \right\} \tilde{\mathbf{v}} = \tilde{\mathbf{v}}^T \partial^2 \mathcal{L}(\boldsymbol{\theta}^*) / \partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T \tilde{\mathbf{v}},$$

where $\boldsymbol{\theta}^*$ is a point on the line connecting $\hat{\boldsymbol{\theta}}$ and $\tilde{\boldsymbol{\theta}}$, and hence in the feasible set. Therefore, by Lemma A.13, we have

$$\left\{ \partial \mathcal{L}(\tilde{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^T - \partial \mathcal{L}(\hat{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^T \right\} \tilde{\mathbf{v}} \geq c_\psi \alpha_{\min} / 2 \|\tilde{\mathbf{v}}\|_2^2 - \tau_1 \{\log(p) + N + q\} / n \left\{ \|\tilde{\mathbf{v}}_1\|_1 + \sum_{j=1, \dots, d} \|\tilde{\mathbf{v}}_{2j}\|_2 \right\}^2,$$

with probability $1 - 2 \exp\{-c_1 n \min(c_\psi^2 \alpha_{\min}^2 / 54^2, 1) / 2\}$. Now note that

$$\begin{aligned} & \mathcal{L}(\boldsymbol{\theta}) - N^{-1/2} \sum_{j \in \mathcal{M}^c} q_\lambda(\|\boldsymbol{\gamma}_j\|_2) \\ = & \mathcal{L}(\boldsymbol{\theta}) - N^{-1/2} \sum_{j \in \mathcal{M}^c} \mu \|\boldsymbol{\gamma}_j\|_2^2 / 2 + N^{-1/2} \sum_{j \in \mathcal{M}^c} \mu \|\boldsymbol{\gamma}_j\|_2^2 / 2 - N^{-1/2} \sum_{j \in \mathcal{M}^c} q_\lambda(\|\boldsymbol{\gamma}_j\|_2), \end{aligned}$$

and $\sum_{j \in \mathcal{M}^c} \mu \|\boldsymbol{\gamma}_j\|_2^2 / 2 - \sum_{j \in \mathcal{M}^c} q_\lambda(\|\boldsymbol{\gamma}_j\|_2)$ is convex by Lemma A.5, and hence for any $\boldsymbol{\theta}$ in the feasible set, we have

$$\begin{aligned} & \tilde{\mathbf{v}}^T \frac{\partial \left[\mathcal{L}(\boldsymbol{\theta}) - N^{-1/2} \sum_{j \in \mathcal{M}^c} q_\lambda(\|\boldsymbol{\gamma}_j\|_2) \right]}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} \tilde{\mathbf{v}} \\ \geq & \tilde{\mathbf{v}}^T \frac{\partial \mathcal{L}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} \tilde{\mathbf{v}} - \mu \sum_{j \in \mathcal{M}^c} \tilde{\mathbf{v}}_{2j}^T \tilde{\mathbf{v}}_{2j} \\ \geq & (c_\psi \alpha_{\min} / 2 - \mu) \|\tilde{\mathbf{v}}\|_2^2 - \tau_1 \{\log(p) + N + q\} / n \left\{ \|\tilde{\mathbf{v}}_1\|_1 + \sum_{j=1, \dots, d} \|\tilde{\mathbf{v}}_{2j}\|_2 \right\}^2. \end{aligned}$$

Furthermore, the first order condition implies

$$\left\{ \partial \mathcal{L}(\tilde{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta} + N^{-1/2} \partial \sum_{j \in \mathcal{M}^c} \rho_\lambda(\|\boldsymbol{\gamma}_j\|_2) / \partial \boldsymbol{\theta} \right\}^T (\hat{\boldsymbol{\theta}} - \tilde{\boldsymbol{\theta}}) \geq 0.$$

Therefore, we have

$$0 \leq \left\{ \frac{\partial \mathcal{L}(\tilde{\boldsymbol{\theta}})}{\partial \boldsymbol{\theta}^T} - N^{-1/2} \frac{\partial \sum_{j \in \mathcal{M}^c} q_\lambda(\|\boldsymbol{\gamma}_j\|_2)}{\partial \boldsymbol{\theta}^T} \right\} (\hat{\boldsymbol{\theta}} - \tilde{\boldsymbol{\theta}}) + \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \tilde{\mathbf{z}}_j^T (\hat{\boldsymbol{\gamma}}_j - \tilde{\boldsymbol{\gamma}}_j),$$

where $\tilde{\mathbf{z}}_j \in \partial \|N^{-1/2} \tilde{\boldsymbol{\gamma}}_j\|_2$ for $j \in \mathcal{M}^c$ and $\tilde{\mathbf{z}}_j = \mathbf{0}$ for $j \in \mathcal{M}$. And define $\tilde{\mathbf{z}} = (\mathbf{0}_q^T, \tilde{\mathbf{z}}_j^T, j = 1, \dots, d)^T$. Furthermore, because $\hat{\boldsymbol{\theta}}$ is the interior local minimum, for $\hat{\mathbf{z}}_j \in \partial \|N^{-1/2} \hat{\boldsymbol{\gamma}}_j\|_2$ for $j \in \mathcal{M}^c$ and $\hat{\mathbf{z}}_j = \mathbf{0}$ for $j \in \mathcal{M}$, and let $\hat{\mathbf{z}} = (\mathbf{0}_q^T, \hat{\mathbf{z}}_j^T, j = 1, \dots, d)^T$, we have

$$\begin{aligned} \mathbf{0} &= \partial \mathcal{L}(\hat{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta} + N^{-1/2} \partial \sum_{j \in \mathcal{M}^c} \rho_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2) / \partial \boldsymbol{\theta} \\ &= \partial \mathcal{L}(\tilde{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta} - N^{-1/2} \partial \sum_{j \in \mathcal{M}^c} q_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2) / \partial \boldsymbol{\theta} + \lambda \hat{\mathbf{z}}, \end{aligned}$$

which leads to

$$\left\{ \partial \mathcal{L}(\hat{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^T + N^{-1/2} \partial \sum_{j \in \mathcal{M}^c} \rho_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2) / \partial \boldsymbol{\theta}^T \right\} \tilde{\mathbf{v}} + \lambda \hat{\mathbf{z}}^T \tilde{\mathbf{v}} = 0.$$

Hence

$$\begin{aligned} 0 &\leq \left\{ \partial \mathcal{L}(\hat{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^T + N^{-1/2} \partial \sum_{j \in \mathcal{M}^c} \rho_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2) / \partial \boldsymbol{\theta}^T \right\} \tilde{\mathbf{v}} - \left\{ \partial \mathcal{L}(\tilde{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^T + N^{-1/2} \partial \sum_{j \in \mathcal{M}^c} \rho_\lambda(\|\tilde{\boldsymbol{\gamma}}_j\|_2) / \partial \boldsymbol{\theta}^T \right\} \tilde{\mathbf{v}} \\ &\quad + \lambda \hat{\mathbf{z}}^T (\tilde{\boldsymbol{\theta}} - \hat{\boldsymbol{\theta}}) - \lambda \tilde{\mathbf{z}}^T (\tilde{\boldsymbol{\theta}} - \hat{\boldsymbol{\theta}}) \\ &= \left\{ \partial \mathcal{L}(\hat{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^T + N^{-1/2} \partial \sum_{j \in \mathcal{M}^c} \rho_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2) / \partial \boldsymbol{\theta}^T \right\} \tilde{\mathbf{v}} - \left\{ \partial \mathcal{L}(\tilde{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^T + N^{-1/2} \partial \sum_{j \in \mathcal{M}^c} \rho_\lambda(\|\tilde{\boldsymbol{\gamma}}_j\|_2) / \partial \boldsymbol{\theta}^T \right\} \tilde{\mathbf{v}} \\ &\quad + \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \hat{\mathbf{z}}_j^T \tilde{\boldsymbol{\Gamma}}_j - \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \|\hat{\boldsymbol{\gamma}}_j\|_2 + \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \tilde{\mathbf{z}}_j^T \hat{\boldsymbol{\gamma}}_j, \end{aligned}$$

which implies

$$\begin{aligned} &\lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \|\tilde{\boldsymbol{\gamma}}_j\|_2 - \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \hat{\mathbf{z}}_j^T \tilde{\boldsymbol{\gamma}}_j \\ &\leq \left\{ \partial \mathcal{L}(\hat{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^T + N^{-1/2} \partial \sum_{j \in \mathcal{M}^c} \rho_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2) / \partial \boldsymbol{\theta}^T \right\} \tilde{\mathbf{v}} - \left\{ \partial \mathcal{L}(\tilde{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^T + N^{-1/2} \partial \sum_{j \in \mathcal{M}^c} \rho_\lambda(\|\tilde{\boldsymbol{\gamma}}_j\|_2) / \partial \boldsymbol{\theta}^T \right\} \tilde{\mathbf{v}} \\ &\quad - \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \|\hat{\boldsymbol{\gamma}}_j\|_2 + \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \tilde{\mathbf{z}}_j^T \hat{\boldsymbol{\gamma}}_j \\ &\leq \left\{ \partial \mathcal{L}(\hat{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^T + N^{-1/2} \partial \sum_{j \in \mathcal{M}^c} \rho_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2) / \partial \boldsymbol{\theta}^T \right\} \tilde{\mathbf{v}} - \left\{ \partial \mathcal{L}(\tilde{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^T + N^{-1/2} \partial \sum_{j \in \mathcal{M}^c} \rho_\lambda(\|\tilde{\boldsymbol{\gamma}}_j\|_2) / \partial \boldsymbol{\theta}^T \right\} \tilde{\mathbf{v}} \end{aligned}$$

$$\leq -(c_\psi \alpha_{\min}/2 - \mu) \|\tilde{\mathbf{v}}\|_2^2 + \tau_1 \{\log(p) + N + q\}/n \left\{ \|\tilde{\mathbf{v}}_1\|_1 + \sum_{j=1, \dots, d} \|\tilde{\mathbf{v}}_{2j}\|_2 \right\}^2. \quad (27)$$

The second inequality holds by the definition of the $\partial \|\cdot\|_2$, and the last line holds by the Taylor expansion. Now suppose the following relation holds,

$$\|\tilde{\mathbf{v}}_1\|_1 + \sum_{j=1}^d \|\tilde{\mathbf{v}}_{2j}\|_2 \leq 2\{\sqrt{q} + \sqrt{m} + (2/\delta + 2)\sqrt{k_0}\} \|\tilde{\mathbf{v}}\|_2.$$

(27) implies

$$\begin{aligned} & \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \|\tilde{\gamma}_j\|_2 - \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \hat{\mathbf{z}}_j^T \tilde{\gamma}_j \\ & \leq \left\{ 4\tau_1 (\log(p) + N + q) \{\sqrt{q} + \sqrt{m} + (2/\delta + 2)\sqrt{k_0}\}^2 n^{-1} - (c_\psi \alpha_{\min}/2 - \mu) \right\} \|\tilde{\mathbf{v}}\|_2^2. \end{aligned}$$

Now because $n \geq 8\tau_1 (\log(p) + N + q) \{\sqrt{q} + \sqrt{m} + (2/\delta + 2)\sqrt{k_0}\}^2 / (c_\psi \alpha_{\min}/2 - \mu)$, we have

$$\begin{aligned} \mathbf{0} & = \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \|\tilde{\gamma}_j\|_2 - \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \|\tilde{\gamma}_j\|_2 \leq \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \|\tilde{\gamma}_j\|_2 - \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \hat{\mathbf{z}}_j^T \tilde{\gamma}_j \\ & \leq -(c_\psi \alpha_{\min}/2 - \mu) \|\tilde{\mathbf{v}}\|_2^2/2. \end{aligned}$$

The first inequality holds by the fact that $\|\tilde{\mathbf{z}}_j\|_2 \leq 1$. Therefore, we have

$$\lambda \sum_{j \in \mathcal{M}^c} \|N^{-1/2} \tilde{\gamma}_j\|_2 = \lambda \sum_{j \in \mathcal{M}^c} \hat{\mathbf{z}}_j^T N^{-1/2} \tilde{\gamma}_j.$$

Now because $\max_{j \in (\mathcal{M} \cup S)^c} \|\hat{\mathbf{z}}_j\|_2 < 1$, we conclude that $\tilde{\gamma}_j = \mathbf{0}$ for $j \in (\mathcal{M} \cup S)^c$. Hence $\text{supp}\{\|\tilde{\gamma}_j\|_2, j = 1, \dots, d\} \subset \mathcal{M} \cup S$. This would prove the claim in the statement.

Thus, we only need to show that

$$\|\tilde{\mathbf{v}}_1\|_1 + \sum_{j=1}^d \|\tilde{\mathbf{v}}_{2j}\|_2 \leq 2\{\sqrt{q} + \sqrt{m} + (2/\delta + 2)\sqrt{k_0}\} \|\tilde{\mathbf{v}}\|_2.$$

From (27), we have

$$(c_\psi \alpha_{\min}/2 - \mu) \|\tilde{\mathbf{v}}\|_2^2 - \tau_1 \{\log(p) + N + q\}/n \left\{ \|\tilde{\mathbf{v}}_1\|_1 + \sum_{j=1, \dots, d} \|\tilde{\mathbf{v}}_{2j}\|_2 \right\}^2$$

$$\begin{aligned}
&\leq \left\{ \partial \mathcal{L}(\tilde{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^T + N^{-1/2} \partial \sum_{j \in \mathcal{M}^c} \rho_\lambda(\|\tilde{\boldsymbol{\gamma}}_j\|_2) / \partial \boldsymbol{\theta}^T \right\} \tilde{\mathbf{v}} - \left\{ \partial \mathcal{L}(\hat{\boldsymbol{\theta}}) / \partial \boldsymbol{\theta}^T + N^{-1/2} \partial \sum_{j \in \mathcal{M}^c} \rho_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2) / \partial \boldsymbol{\theta}^T \right\} \tilde{\mathbf{v}} \\
&\leq \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \hat{\mathbf{z}}_j^T \tilde{\boldsymbol{\gamma}}_j - \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \|\hat{\boldsymbol{\gamma}}_j\|_2 - \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \|\tilde{\boldsymbol{\gamma}}_j\|_2 + \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \hat{\mathbf{z}}_j^T \hat{\boldsymbol{\gamma}}_j \\
&= \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \tilde{\mathbf{z}}_j^T \hat{\boldsymbol{\gamma}}_j - \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \|\tilde{\boldsymbol{\gamma}}_j\|_2 + \lambda \hat{\mathbf{z}}^T \tilde{\mathbf{v}}. \tag{28}
\end{aligned}$$

Now because $\text{supp}(\{\|\hat{\boldsymbol{\gamma}}_{j,\cdot}\|_2, j = 1, \dots, d\}) \subset \mathcal{M} \cup S$, we have

$$\begin{aligned}
&\lambda \sum_{j \in \mathcal{M}^c} \tilde{\mathbf{z}}_j^T \hat{\boldsymbol{\gamma}}_j - \lambda \sum_{j \in \mathcal{M}^c} \|\tilde{\boldsymbol{\gamma}}_j\|_2 \\
&\leq \lambda \sum_{j \in \mathcal{M}^c} \|\hat{\boldsymbol{\gamma}}_j\|_2 - \lambda \sum_{j \in \mathcal{M}^c} \|\tilde{\boldsymbol{\gamma}}_j\|_2 \\
&= \lambda \sum_{j \in S} \|\hat{\boldsymbol{\gamma}}_j\|_2 - \lambda \sum_{j \in S} \|\tilde{\boldsymbol{\gamma}}_j\|_2 - \lambda \sum_{j \in (\mathcal{M} \cup S)^c} \|\tilde{\boldsymbol{\gamma}}_j\|_2 \\
&= \lambda \sum_{j \in S} \|\hat{\boldsymbol{\gamma}}_j\|_2 - \lambda \sum_{j \in S} \|\tilde{\boldsymbol{\gamma}}_j\|_2 - (\lambda \sum_{j \in (\mathcal{M} \cup S)^c} \|\tilde{\boldsymbol{\gamma}}_j\|_2 - \lambda \sum_{j \in (\mathcal{M} \cup S)^c} \|\hat{\boldsymbol{\gamma}}_j\|_2) \\
&= \lambda \sum_{j \in S} \|\tilde{\boldsymbol{\gamma}}_j - \hat{\boldsymbol{\gamma}}_j - \tilde{\boldsymbol{\gamma}}_j\|_2 - \lambda \sum_{j \in S} \|\tilde{\boldsymbol{\gamma}}_j\|_2 - (\lambda \sum_{j \in (\mathcal{M} \cup S)^c} \|\tilde{\boldsymbol{\gamma}}_j\|_2 - \lambda \sum_{j \in (\mathcal{M} \cup S)^c} \|\tilde{\boldsymbol{\gamma}}_j + \hat{\boldsymbol{\gamma}}_j - \tilde{\boldsymbol{\gamma}}_j\|_2) \\
&\leq \lambda N^{1/2} \sum_{j \in S} \|\tilde{\mathbf{v}}_j\|_2 - \lambda N^{1/2} \sum_{j \in (\mathcal{M} \cup S)^c} \|\tilde{\mathbf{v}}_j\|_2. \tag{29}
\end{aligned}$$

In addition,

$$\begin{aligned}
\lambda \hat{\mathbf{z}}^T \tilde{\mathbf{v}} &= \lambda \sum_{j \in S} \hat{\mathbf{z}}_j^T \tilde{\mathbf{v}}_{2j} + \lambda \sum_{j \in (\mathcal{M} \cup S)^c} \hat{\mathbf{z}}_j^T \tilde{\mathbf{v}}_{2j} \\
&\leq \lambda \max_{j \in S} \|\hat{\mathbf{z}}_j\|_2 \sum_{j \in S} \|\tilde{\mathbf{v}}_{2j}\|_2 + \lambda \max_{j \in (\mathcal{M} \cup S)^c} \|\hat{\mathbf{z}}_j\|_2 \sum_{j \in (\mathcal{M} \cup S)^c} \|\tilde{\mathbf{v}}_{2j}\|_2 \\
&\leq \lambda \sum_{j \in S} \|\tilde{\mathbf{v}}_{2j}\|_2 + \lambda(1 - \delta) \sum_{j \in (\mathcal{M} \cup S)^c} \|\tilde{\mathbf{v}}_j\|_2.
\end{aligned}$$

Combine with (28) and (29), we have

$$\begin{aligned}
&-\tau_1 \{\log(p) + N + q\} / n \left\{ \|\tilde{\mathbf{v}}_1\|_1 + \sum_{j=1, \dots, d} \|\tilde{\mathbf{v}}_{2j}\|_2 \right\}^2 \\
&\leq (c_\psi \alpha_{\min} / 2 - \mu) \|\tilde{\mathbf{v}}\|_2^2 - \tau_1 \{\log(p) + N + q\} / n \left\{ \|\tilde{\mathbf{v}}_1\|_1 + \sum_{j=1, \dots, d} \|\tilde{\mathbf{v}}_{2j}\|_2 \right\}^2 \\
&\leq \lambda \sum_{j \in S} \|\tilde{\mathbf{v}}_{2j}\|_2 - \lambda \sum_{j \in (\mathcal{M} \cup S)^c} \|\tilde{\mathbf{v}}_{2j}\|_2 + \lambda \sum_{j \in S} \|\tilde{\mathbf{v}}_{2j}\|_2 + \lambda(1 - \delta) \sum_{j \in (\mathcal{M} \cup S)^c} \|\tilde{\mathbf{v}}_{2j}\|_2 \\
&= 2\lambda \sum_{j \in S} \|\tilde{\mathbf{v}}_{2j}\|_2 - \lambda \delta \sum_{j \in (\mathcal{M} \cup S)^c} \|\tilde{\mathbf{v}}_{2j}\|_2.
\end{aligned}$$

Now because $\delta\lambda/2 \geq 2\tau_1 R\{\log(p) + N + q\}/n \geq \tau_1\{\log(p) + N + q\}/n \left\{ \|\tilde{\mathbf{v}}_1\|_1 + \sum_{j=1}^d \|\tilde{\mathbf{v}}_{2j}\|_2 \right\}$,

the above display implies

$$-2^{-1}\delta\lambda \left\{ \|\tilde{\mathbf{v}}_1\|_1 + \sum_{j=1}^d \|\tilde{\mathbf{v}}_{2j}\|_2 \right\} \leq 2\lambda \sum_{j \in S} \|\tilde{\mathbf{v}}_{2j}\|_2 - \lambda\delta \sum_{j \in (\mathcal{M} \cup S)^c} \|\tilde{\mathbf{v}}_{2j}\|_2,$$

which implies

$$2^{-1}\delta\lambda \sum_{j \in (\mathcal{M} \cup S)^c} \|\tilde{\mathbf{v}}_{2j}\|_2 \leq 2^{-1}\delta\lambda \|\tilde{\mathbf{v}}_1\|_1 + 2^{-1}\delta\lambda \sum_{j \in \mathcal{M}} \|\tilde{\mathbf{v}}_{2j}\|_2 + (2 + 2^{-1}\delta)\lambda \sum_{j \in S} \|\tilde{\mathbf{v}}_{2j}\|_2.$$

Therefore,

$$\begin{aligned} \|\tilde{\mathbf{v}}_1\|_1 + \sum_{j=1}^d \|\tilde{\mathbf{v}}_{2j}\|_2 &\leq \|\tilde{\mathbf{v}}_1\|_1 + \sum_{j \in \mathcal{M}} \|\tilde{\mathbf{v}}_{2j}\|_2 + \sum_{j \in S} \|\tilde{\mathbf{v}}_{2j}\|_2 + \sum_{j \in (\mathcal{M} \cup S)^c} \|\tilde{\mathbf{v}}_{2j}\|_2 \\ &\leq \|\tilde{\mathbf{v}}_1\|_1 + \sum_{j \in \mathcal{M}} \|\tilde{\mathbf{v}}_{2j}\|_2 + \sum_{j \in S} \|\tilde{\mathbf{v}}_{2j}\|_2 + \|\tilde{\mathbf{v}}_1\|_1 + \sum_{j \in \mathcal{M}} \|\tilde{\mathbf{v}}_{2j}\|_2 + (4 + \delta)/\delta \sum_{j \in S} \|\tilde{\mathbf{v}}_{2j}\|_2 \\ &\leq 2\{\sqrt{q} + \sqrt{m} + (2/\delta + 2)\sqrt{k_0}\} \|\tilde{\mathbf{v}}\|_2. \end{aligned}$$

This completes the proof. \square

Lemma A.18. *Assume Conditions (A1)–(A6) and (C1) – (C5) hold, assume $\mu < c_\psi\alpha_{\min}/2$ and $n > \tau_1\{\log(p) + N + q\}(q + m + k_0)/(c_\psi\alpha_{\min}/2 - \mu)$. Then the function $\mathcal{L}(\boldsymbol{\theta}) + N^{-1/2} \sum_{j \in \mathcal{M}^c} q\lambda(\|\boldsymbol{\gamma}_j\|_2)$ and $\mathcal{L}(\boldsymbol{\theta}) - \mu N^{-1/2} \sum_{j \in \mathcal{M}^c} \|\boldsymbol{\gamma}_j\|_2/2$ are strictly convexity over the set $[\boldsymbol{\theta} : \boldsymbol{\theta} = \{\boldsymbol{\alpha}^T, N^{-1/2}\text{vec}(\boldsymbol{\Gamma})^T\}^T; \boldsymbol{\gamma}_j = \mathbf{0} \ \forall j \in (\mathcal{M} \cup S)^c]$.*

Proof: First define a vector $\mathbf{v} = (\mathbf{v}_1^T, \mathbf{v}_{2j}^T, j = 1, \dots, d)^T$ with $\|\mathbf{v}_{2j}\|_2 > 0$ for $j \in \mathcal{M} \cup S$ and $\|\mathbf{v}_{2j}\|_2 = 0$, otherwise. By Lemma A.13, we have for $\boldsymbol{\theta}$ in the feasible set of program (30), we have

$$\mathbf{v}^T \partial \mathcal{L}(\boldsymbol{\theta}) / \partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T \mathbf{v} \geq c_\psi\alpha_{\min} \|\mathbf{v}\|_2^2 - \tau_1\{\log(p) + N + q\}/n \left\{ \|\mathbf{v}_1\|_1 + \sum_{j=1}^d \|\mathbf{v}_{2j}\|_2^2 \right\}^2.$$

Therefore, we have

$$\mathbf{v}^T \partial \mathcal{L}(\boldsymbol{\theta}) / \partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T \mathbf{v} \geq \{c_\psi\alpha_{\min} - \tau_1\{\log(p) + N + q\}(q + m + k_0)/n\} \|\mathbf{v}_{2j}\|_2^2,$$

and

$$\mathbf{v}^T \partial \mathcal{L}(\boldsymbol{\theta}) / \partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T \mathbf{v} - \mu \|\mathbf{v}\|_2^2 \geq \{c_\psi \alpha_{\min} - \mu - \tau_1 \{\log(p) + N + q\} (q + m + k_0) / n\} \|\mathbf{v}_{2j}\|_2^2.$$

Now because $\{c_\psi \alpha_{\min} - \mu - \tau_1 \{\log(p) + N + q\} (q + m + k_0) / n\} > 0$, $\mathcal{L}(\boldsymbol{\theta}) - \mu \sum_{j \in \mathcal{M}^c} \|\boldsymbol{\gamma}_j\|_2^2 / 2$ is strictly convex on the set that $[\boldsymbol{\theta} : \boldsymbol{\theta} = \{\boldsymbol{\alpha}^T, \text{vec}(\boldsymbol{\Gamma})^T\}^T; \boldsymbol{\gamma}_j = \mathbf{0} \ \forall j \in (\mathcal{M} \cup S)^c]$. Finally, because

$$\begin{aligned} & \mathcal{L}(\boldsymbol{\theta}) - N^{-1/2} \sum_{j \in \mathcal{M}^c} q_\lambda(\|\boldsymbol{\gamma}_j\|_2) \\ = & \mathcal{L}(\boldsymbol{\theta}) - \mu N^{-1/2} \sum_{j \in \mathcal{M}^c} \|\boldsymbol{\gamma}_j\|_2^2 / 2 + \left\{ \mu N^{-1/2} \sum_{j \in \mathcal{M}^c} \|\boldsymbol{\gamma}_j\|_2^2 / 2 - N^{-1/2} \sum_{j \in \mathcal{M}^c} q_\lambda(\|\boldsymbol{\gamma}_j\|_2) \right\}, \end{aligned}$$

where the second part is convex by Lemma A.5. Therefore, $\mathcal{L}(\boldsymbol{\theta}) - N^{-1/2} \sum_{j \in \mathcal{M}^c} q_\lambda(\|\boldsymbol{\gamma}_j\|_2)$ and $\mathcal{L}(\boldsymbol{\theta}) + N^{-1/2} \sum_{j \in \mathcal{M}^c} \rho_\lambda(\|\boldsymbol{\gamma}_j\|_2) = \mathcal{L}(\boldsymbol{\theta}) - N^{-1/2} \sum_{j \in \mathcal{M}^c} q_\lambda(\|\boldsymbol{\gamma}_j\|_2) + \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \|\boldsymbol{\gamma}_j\|_2$ are strictly convex over the set $[\boldsymbol{\theta} : \boldsymbol{\theta} = \{\boldsymbol{\alpha}^T, \text{vec}(\boldsymbol{\Gamma})^T\}^T; \boldsymbol{\gamma}_j = \mathbf{0} \ \forall j \in (\mathcal{M} \cup S)^c]$. This proves the result. \square

Suppose the indices of nonzero $\boldsymbol{\beta}_j(\cdot), j \in \mathcal{M}^c$ are given, we minimize

$$\begin{aligned} & \mathcal{L}(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) + N^{-1/2} \sum_{j \in \mathcal{M}^c} \rho_\lambda(\|\boldsymbol{\gamma}_j\|_2), \\ & \text{such that } \|\boldsymbol{\alpha}\|_1 + N^{-1/2} \sum_{j=1}^d \|\boldsymbol{\gamma}_j\|_2 \leq R, \text{ and } \boldsymbol{\gamma}_j = \mathbf{0} \text{ for } j \in (\mathcal{M} \cup S)^c, \end{aligned} \quad (30)$$

with respect to $\boldsymbol{\alpha}, \boldsymbol{\Gamma}$ to obtain the estimator, which named oracle estimator.

In Theorem A.1, we show that the minimizer of (3) is unique under some conditions. Moreover, it approaches the oracle estimator, i.e., the estimator of (30) when the indices of nonzero $\boldsymbol{\beta}_j(\cdot), j \in \mathcal{M}^c$ are given.

Theorem A.1. *Assume Conditions (A1)–(A6) and (C1) – (C5) hold. Further assume that $\mu < c_\psi \alpha_{\min} / 2$ and $24\lambda(q + m + k_0) / (c_\psi \alpha_{\min} - 3\mu/2) \leq R$,*

$$\lambda \geq \max \left\{ 2\tau_1 R \log(p) / n, C_0 \sqrt{\log\{\max(p, n)\}} / n + Ck_0^{1/2} N^{-\omega} \right\} / 4.$$

Let $\hat{\boldsymbol{\theta}}_{\mathcal{M} \cup \mathcal{S}} = \{\hat{\boldsymbol{\alpha}}^T, N^{-1/2} \hat{\boldsymbol{\gamma}}_j^T, j \in \mathcal{M} \cup \mathcal{S}\}$ be the minimizer for (30). Without loss of generality, we define $\hat{\boldsymbol{\theta}} = \{\hat{\boldsymbol{\alpha}}^T, N^{-1/2} \hat{\boldsymbol{\gamma}}_j^T, j \in \mathcal{M} \cup \mathcal{S}, \mathbf{0}_{d-m-k_0}^T\}^T$, where we assume the first $q + (m + k_0)N$ elements are nonzero, and satisfies

$$\frac{\partial \mathcal{L}(\hat{\boldsymbol{\theta}})}{\partial \boldsymbol{\theta}} - N^{-1/2} \frac{\partial \sum_{j \in \mathcal{M}^c} q_\lambda(\|\hat{\boldsymbol{\Gamma}}_{\cdot j}\|_2)}{\partial \boldsymbol{\theta}} + \lambda \hat{\mathbf{z}} = \mathbf{0}, \quad (31)$$

where $\hat{\mathbf{z}} = (\mathbf{0}_q, \hat{\mathbf{z}}_j^T, j = 1, \dots, d)^T$, and $\hat{\mathbf{z}}_j \in \partial \|N^{-1/2} \hat{\boldsymbol{\Gamma}}_{\cdot j}\|_2$. Assume there is a $\delta \in [4R\tau_1 \log(p)/n/\lambda, 1]$, and $\max_{j \in (\mathcal{M} \cup \mathcal{S})^c} \|\hat{\mathbf{z}}_j\|_2 \leq 1 - \delta$ and

$$n \geq \max \left[8\tau_1(\log(p) + N + q) \{ \sqrt{q} + \sqrt{m} + (2/\delta + 2)\sqrt{k_0} \}^2 / (c_\psi \alpha_{\min}/2 - \mu), \right. \\ \left. \tau_1 \{ \log(p) + N + q \} (q + m + k_0) / (c_\psi \alpha_{\min}/2 - \mu) \right].$$

Then (3) has a unique minimizer $\hat{\boldsymbol{\theta}}$.

Proof: Step 1: By Lemma A.18, $\hat{\boldsymbol{\theta}}_{\mathcal{M} \cup \mathcal{S}}$ be the minimizer for (30). It is easy to see that the condition in Theorem 1 holds and $\sum_{j \in (\mathcal{M} \cup \mathcal{S})^c} \|\hat{\boldsymbol{\Gamma}}_{\cdot j} - \boldsymbol{\gamma}_{0j}\|_2 \leq 3 \sum_{j \in \mathcal{M}^c \cap \mathcal{S}} \|\hat{\boldsymbol{\Gamma}}_{\cdot j} - \boldsymbol{\gamma}_{0j}\|_2$ because $\sum_{j \in (\mathcal{M} \cup \mathcal{S})^c} \|\hat{\boldsymbol{\Gamma}}_{\cdot j} - \boldsymbol{\gamma}_{0j}\|_2 = 0$. Therefore, applying Theorem 1, we have

$$\|\hat{\boldsymbol{\alpha}}\|_1 + N^{-1/2} \sum_{j=1}^d \|\hat{\boldsymbol{\Gamma}}_{\cdot j}\|_2 \leq 12\lambda(q + m + k_0) / (c_\psi \alpha_{\min} - 3\mu/2) + \|\boldsymbol{\alpha}_0\|_1 + N^{-1/2} \sum_{j=1}^d \|\boldsymbol{\Gamma}_{0 \cdot j}\|_2 \\ \leq R,$$

and hence $\hat{\boldsymbol{\theta}}_{\mathcal{M} \cup \mathcal{S}}$ is in the feasible region. The second inequality holds by the assumption that $24\lambda(q + m + k_0) / (c_\psi \alpha_{\min} - 3\mu/2) \leq R$ in the theorem statement.

Step 2: We show that $\hat{\boldsymbol{\theta}}$ is a local minimum (3) by verifying the conditions in Lemma A.16 in the supplementary material. Because

$$\mathcal{L}(\boldsymbol{\theta}) + N^{-1/2} \sum_{j \in \mathcal{M}^c} \rho_\lambda \|\boldsymbol{\gamma}_j\|_2 = \mathcal{L}(\boldsymbol{\theta}) - N^{-1/2} \sum_{j \in \mathcal{M}^c} q_\lambda \|\boldsymbol{\gamma}_j\|_2 + \lambda N^{-1/2} \sum_{j \in \mathcal{M}^c} \|\boldsymbol{\gamma}_j\|_2,$$

we can write $f = \mathcal{L}$, $g = q_\lambda$ and $(\mathbf{x}^*, \mathbf{v}^*, \mathbf{w}^*, \mu_1^*) = (\hat{\boldsymbol{\theta}}, \hat{\mathbf{z}}, \hat{\mathbf{z}}_1, 0)$, where $\hat{\mathbf{z}} = (\mathbf{0}_q^T, \hat{\mathbf{z}}_j^T, j = 1, \dots, d)^T$ and $\hat{\mathbf{z}}_j \in \partial \|N^{-1/2} \hat{\boldsymbol{\Gamma}}_{\cdot j}\|_2$ and $\hat{\mathbf{z}}_1 \in \partial \{ \|\hat{\boldsymbol{\alpha}}\|_1 + N^{-1/2} \sum_{j=1}^d \|\hat{\boldsymbol{\gamma}}_j\|_2 \}$. Lemma A.5 in the

supplementary material ensures the concavity and differentiability of $g(\|\mathbf{x}\|_2) - \mu/2\|\mathbf{x}\|_2^2$ in $\|\mathbf{x}\|_2$ and \mathbf{x} , respectively. Furthermore, since $\mu_1^* = 0$, (18) is satisfied. (19) is satisfied by our construction in (31). Therefore, it remains to verify (20). We first show that if $\mathbf{s} \in G^*$, $\mathbf{s}_{2j} = \mathbf{0}$ for $j \in (\mathcal{M} \cup S)^c$, and therefore (19) only need to be satisfied for the vectors in the form of $\mathbf{s} = (\mathbf{s}_1^T, \mathbf{s}_{2j}^T, j \in \mathcal{M} \cup S, \mathbf{0}_{d-m-k_0}^T)^T$. Suppose it does not hold so that there is a $\boldsymbol{\nu} \in G^*$ such that $\boldsymbol{\nu} = (\boldsymbol{\nu}_1^T, \boldsymbol{\nu}_{2j}^T, j \in \mathcal{M} \cup S, \mathbf{0}_{d-m-k_0}^T)^T$. We define $\mathbf{z}' = (\mathbf{0}_q^T, \mathbf{z}'_{2j}, j = 1, \dots, d)$ such that $\mathbf{z}'_{2k} = \hat{\mathbf{z}}_{2k}$ for $k \neq j$, and $\mathbf{z}'_{2k} = \partial\|\boldsymbol{\nu}_{2k}\|_2$ for $k = j$. Clearly $\mathbf{z}'_{2k} \in \partial\|N^{-1/2}\hat{\boldsymbol{\gamma}}_k\|_2$, $k = 1, \dots, d$, $\|\mathbf{z}'_{2j}\| = 1$ because $\boldsymbol{\nu}_{2j} > \mathbf{0}$, and

$$\lambda \boldsymbol{\nu}^T \mathbf{z}' > \lambda \boldsymbol{\nu}^T \hat{\mathbf{z}},$$

because $\max_{j \in (\mathcal{M} \cup S)^c} \|\hat{\mathbf{z}}_{2j}\|_2 < 1$. Therefore,

$$\begin{aligned} & \boldsymbol{\nu}^T \left[\frac{\partial \mathcal{L}(\hat{\boldsymbol{\theta}})}{\partial \boldsymbol{\theta}} - N^{-1/2} \frac{\partial \sum_{j \in \mathcal{M}^c} q_\lambda(\|\boldsymbol{\gamma}_j\|_2)}{\partial \boldsymbol{\theta}} \right] + \lambda \boldsymbol{\nu}^T \mathbf{z}' \\ & > \boldsymbol{\nu}^T \left[\frac{\partial \mathcal{L}(\hat{\boldsymbol{\theta}})}{\partial \boldsymbol{\theta}} - N^{-1/2} \frac{\partial \sum_{j \in \mathcal{M}^c} q_\lambda(\|\boldsymbol{\gamma}_j\|_2)}{\partial \boldsymbol{\theta}} \right] + \lambda \boldsymbol{\nu}^T \hat{\mathbf{z}} \\ & = 0, \end{aligned}$$

which contradicts with the requirement of G^* that

$$\sup_{(\mathbf{v}_j \in \partial\|\hat{\boldsymbol{\gamma}}_j\|_1, j \in \mathcal{M}^c)} \boldsymbol{\nu}^T \left[\frac{\partial \mathcal{L}(\hat{\boldsymbol{\theta}})}{\partial \boldsymbol{\theta}} - N^{-1/2} \frac{\partial \sum_{j \in \mathcal{M}^c} q_\lambda(\|\boldsymbol{\gamma}_j\|_2)}{\partial \boldsymbol{\theta}} \right] + \lambda \sum_{j \in \mathcal{M}^c} \boldsymbol{\nu}_{2j}^T \mathbf{v}_j = 0.$$

Therefore, we conclude that if $\mathbf{s} \in G^*$, $\mathbf{s}_{2j} = \mathbf{0}$ for $j \in (\mathcal{M} \cup S)^c$. Now by Lemma A.18 in the supplementary material and the fact that $\hat{\boldsymbol{\gamma}}_j = \mathbf{0}$ for $j \in (\mathcal{M} \cup S)^c$, we conclude that (20) holds by setting $\kappa = \mu$. Hence all conditions of Lemma A.16 in the supplementary material are satisfied, so we conclude $\hat{\boldsymbol{\theta}}$ is an isolated local minimizer of (3).

Now by Lemma A.17, because $\hat{\boldsymbol{\gamma}}_j = \mathbf{0}$ for $j \in (\mathcal{M} \cup S)^c$ and $\hat{\boldsymbol{\theta}}$ is an interior minimizer, $\tilde{\boldsymbol{\gamma}}_j = \mathbf{0}$ for $j \in (\mathcal{M} \cup S)^c$ for any given stationary point $\tilde{\boldsymbol{\theta}}$ of (3). Hence, we write any

stationary point in the form of $\tilde{\boldsymbol{\theta}} = (\tilde{\boldsymbol{\alpha}}_1^T, N^{-1/2}\tilde{\boldsymbol{\gamma}}_j^T, j \in \mathcal{M} \cup S, \mathbf{0}_{d-m-k_0}^T)^T$, where $\tilde{\boldsymbol{\theta}}_{\mathcal{M} \cup S} = (\tilde{\boldsymbol{\alpha}}_1^T, N^{-1/2}\tilde{\boldsymbol{\gamma}}_j^T, j \in \mathcal{M} \cup S)^T$ is a stationary point for (30). Furthermore, (30) is strictly convex by Lemma A.18, and $\tilde{\boldsymbol{\theta}}_{\mathcal{M} \cup S}$ is unique in the feasible set and $\hat{\boldsymbol{\theta}}_{\mathcal{M} \cup S}$ and $\tilde{\boldsymbol{\theta}}_{\mathcal{M} \cup S}$ are unique. Therefore, $\hat{\boldsymbol{\theta}}$ is the unique local minimum. This proves the result.

Let $\boldsymbol{\theta}^*$ is the point on the line connecting $\hat{\boldsymbol{\theta}}$ and $\boldsymbol{\theta}_0$. We write

$$\hat{\mathbf{Q}}(\boldsymbol{\theta}^*) = \begin{Bmatrix} \hat{\mathbf{Q}}_{(\mathcal{M} \cup S)(\mathcal{M} \cup S)}(\boldsymbol{\theta}^*) & \hat{\mathbf{Q}}_{(\mathcal{M} \cup S)(\mathcal{M} \cup S)^c}(\boldsymbol{\theta}^*) \\ \hat{\mathbf{Q}}_{(\mathcal{M} \cup S)^c(\mathcal{M} \cup S)}(\boldsymbol{\theta}^*) & \hat{\mathbf{Q}}_{(\mathcal{M} \cup S)^c(\mathcal{M} \cup S)^c}(\boldsymbol{\theta}^*) \end{Bmatrix}. \quad (32)$$

Then by the construction in (A.1), let $\hat{\boldsymbol{\theta}} = (\hat{\boldsymbol{\theta}}_{\mathcal{M} \cup S}^T, \mathbf{0}_{(\mathcal{M} \cup S)^c}^T)^T$, $\hat{\boldsymbol{\theta}}_{\mathcal{M} \cup S}$ is the minimizer for (30), then we have

$$\begin{aligned} & \hat{\mathbf{Q}}(\boldsymbol{\theta}^*)(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) + \begin{bmatrix} \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} - \left\{ \frac{\partial \sum_{j \in \mathcal{M}^c} q_\lambda(\|\boldsymbol{\gamma}_{0j}\|_2)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} \\ \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{2(\mathcal{M} \cup S)^c} - \left\{ \frac{\partial \sum_{j \in \mathcal{M}^c} q_\lambda(\|\boldsymbol{\gamma}_{0j}\|_2)}{\partial \boldsymbol{\theta}} \right\}_{2(\mathcal{M} \cup S)^c} \end{bmatrix} \\ & + \lambda \begin{Bmatrix} \hat{\mathbf{z}}_{\mathcal{M} \cup S} \\ \hat{\mathbf{z}}_{2(\mathcal{M} \cup S)^c} \end{Bmatrix} = \mathbf{0}. \end{aligned}$$

Taking the upper $q + m + k_0$ non-zero component, we get

$$\begin{aligned} & \hat{\boldsymbol{\theta}}_{\mathcal{M} \cup S} - \boldsymbol{\theta}_{0\mathcal{M} \cup S} \\ & = \{ \hat{\mathbf{Q}}_{\mathcal{M} \cup S, \mathcal{M} \cup S}(\boldsymbol{\theta}^*) \}^{-1} \left[- \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} + \left\{ \frac{\partial \sum_{j \in \mathcal{M}^c} q_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} - \lambda \hat{\mathbf{z}}_{\mathcal{M} \cup S} \right] \end{aligned} \quad (33)$$

while taking the lower $d - m - k_0$ components, this leads to

$$\begin{aligned} \hat{\mathbf{z}}_{(\mathcal{M} \cup S)^c} & = \lambda^{-1} \left[\left\{ \frac{\partial \sum_{j \in \mathcal{M}^c} q_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2)}{\partial \boldsymbol{\theta}} \right\}_{2(\mathcal{M} \cup S)^c} - \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}_0} \right\}_{2(\mathcal{M} \cup S)^c} \right] \\ & \quad - \lambda^{-1} \hat{\mathbf{Q}}_{(\mathcal{M} \cup S)^c(\mathcal{M} \cup S)}(\boldsymbol{\theta}^*)(\hat{\boldsymbol{\theta}}_{\mathcal{M} \cup S} - \boldsymbol{\theta}_{\mathcal{M} \cup S}) \\ & = \lambda^{-1} \left[\left\{ \frac{\partial \sum_{j \in \mathcal{M}^c} q_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2)}{\partial \boldsymbol{\theta}} \right\}_{2(\mathcal{M} \cup S)^c} - \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{2(\mathcal{M} \cup S)^c} \right] \\ & \quad + \lambda^{-1} \hat{\mathbf{Q}}_{(\mathcal{M} \cup S)^c(\mathcal{M} \cup S)}(\boldsymbol{\theta}^*) \{ \hat{\mathbf{Q}}_{(\mathcal{M} \cup S)(\mathcal{M} \cup S)}(\boldsymbol{\theta}^*) \}^{-1} \\ & \quad \times \left(\left[\left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} - \left\{ \frac{\partial \sum_{j \in \mathcal{M}^c} q_\lambda(\|\hat{\boldsymbol{\gamma}}_j\|_2)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} \right] + \lambda \hat{\mathbf{z}}_{\mathcal{M} \cup S} \right). \end{aligned}$$

Further by Condition (A4), we have

$$\begin{aligned} \left\{ \frac{\partial \sum_{j \in \mathcal{M}^c} q_\lambda(\|\hat{\gamma}_j\|_2)}{\partial \boldsymbol{\theta}} \right\}_{2(\mathcal{M} \cup S)^c} &= \left[\left\{ \frac{\partial \lambda \|\hat{\gamma}_j\|_2}{\partial \boldsymbol{\gamma}_j^T} \right\} - \left\{ \frac{\partial \rho_\lambda(\|\hat{\gamma}_j\|_2)}{\partial \boldsymbol{\gamma}_j^T} \right\}, j \in (\mathcal{M} \cup S)^c \right]^T \\ &= \mathbf{0}. \end{aligned}$$

Therefore, we have

$$\begin{aligned} \hat{\mathbf{z}}_{2(\mathcal{M} \cup S)^c} &= \lambda^{-1} \left[- \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{2(\mathcal{M} \cup S)^c} \right] + \lambda^{-1} \hat{\mathbf{Q}}_{(\mathcal{M} \cup S)^c(\mathcal{M} \cup S)}(\boldsymbol{\theta}^*) \left\{ \hat{\mathbf{Q}}_{(\mathcal{M} \cup S)(\mathcal{M} \cup S)}(\boldsymbol{\theta}^*) \right\}^{-1} \\ &\quad \times \left(\left[\left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} - \left\{ \frac{\partial \sum_{j \in \mathcal{M}^c} q_\lambda(\|\hat{\gamma}_j\|_2)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} \right] + \lambda \hat{\mathbf{z}}_{\mathcal{M} \cup S} \right). \end{aligned}$$

Hence, to use Theorem 1, we must show that $\max_{j \in (\mathcal{M} \cup S)^c} \|\hat{\mathbf{z}}_j\|_2 < 1$. \square

Proof of Theorem 2: First of all $\hat{\boldsymbol{\theta}}_{2(\mathcal{M} \cup S)^c} = \mathbf{0}$ by construction. So for any unit vector

$\mathbf{v} = (\mathbf{v}_1, \mathbf{v}_{2j}^T, j = 1, \dots, d)$, $\mathbf{w} = (\mathbf{w}_1^T, \mathbf{w}_{2j}^T, j = 1, \dots, d)^T$, recall that

$$\mathbf{v} \frac{\partial^2 \mathcal{L}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} \mathbf{w} = \mathbf{v}^T \sum_{i=1}^n \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} \mathbf{w}$$

Denote $\nabla^3 \mathcal{L}(\boldsymbol{\beta})$ to be the third order gradient of \mathcal{L} , we have

$$\begin{aligned} &\mathbf{v} \nabla^3 \mathcal{L}(\boldsymbol{\beta}) \mathbf{w} \\ &= n^{-1} \mathbf{v}^T \sum_{i=1}^n \psi_i''(\boldsymbol{\alpha}, \boldsymbol{\Gamma}) \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} \mathbf{w} \\ &\quad \times \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right). \end{aligned}$$

Here we consider vector \mathbf{v}, \mathbf{w} such that $\|\mathbf{v}_{2j}\|_2 > 0, \|\mathbf{w}_{2j}\|_2 > 0$ for $j \in \mathcal{M} \cup S$ and $\|\mathbf{v}_{2j}\|_2 =$

$0, \|\mathbf{w}_{2j}\|_2 = 0$ otherwise. First by Theorem 1, we have

$$\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0\|_2 \leq 3\lambda \sqrt{q + m + k_0} / (c_\psi \alpha_{\min} - 3\mu/2).$$

Let $\mathbb{K} \equiv \{\mathbf{v} = (\mathbf{v}_1^T, \mathbf{v}_{2j}^T, j = 1, \dots, d)^T, \mathbf{v}_{2j} = \mathbf{0} \text{ for } j \in (\mathcal{M} \cup S)^c, \|\mathbf{v}\|_2 \leq 1\}$, then there are

constants $a_1 > 0, a_2 > 0$ such that

$$\sup_{\mathbf{v}, \mathbf{w} \in \mathbb{K}} \left| \mathbf{v}^T \left\{ \hat{\mathbf{Q}}(\boldsymbol{\theta}^*) - \frac{\partial^2 \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} \right\} \mathbf{w} \right|$$

$$\begin{aligned}
&\leq \sup_{\mathbf{v}, \mathbf{w} \in \mathbb{K}} |n^{-1} \mathbf{v}^T \sum_{i=1}^n \psi_i'''(\boldsymbol{\alpha}^{**}, \boldsymbol{\Gamma}^{**}) \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} \mathbf{w} \\
&\quad \times \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right) (\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) \\
&\leq \{3\lambda \sqrt{q+m+k_0} / (c_\psi \alpha_{\min} - 3\mu/2)\} \\
&\quad \times \sup_{\mathbf{v}, \mathbf{w} \in \mathbb{K}} \left| n^{-1} \mathbf{v}^T \sum_{i=1}^n |\psi_i'''(\boldsymbol{\alpha}^{**}, \boldsymbol{\Gamma}^{**})| \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} \mathbf{w} \right. \\
&\quad \times \left. \left| \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right) (\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) / \{3\lambda \sqrt{q+m+k_0} / (c_\psi \alpha_{\min} - 3\mu/2)\} \right| \right| \\
&\leq (a_1 + \log(p) \sqrt{q+m+k_0}/n) \{3\lambda \sqrt{q+m+k_0} / (c_\psi \alpha_{\min} - 3\mu/2)\} \\
&= O_p \left(\sqrt{q+m+k_0} [\sqrt{\log\{\max(p, n)\}}/n + k_0^{1/2} N^{-\omega}] \right) \tag{34}
\end{aligned}$$

with probability greater than $1 - 2 \exp\{-a_2 n \min(c_\psi^2 \alpha_{\min}^2 / 54^2, 1) / 2\} - 20 \max(p, n)^{-1}$, where

$\boldsymbol{\theta}^{**} \equiv (\boldsymbol{\alpha}^{**T}, \text{vec}(\boldsymbol{\Gamma}^{**})^T)^T$ is on the line connecting $\boldsymbol{\theta}^*$ and $\boldsymbol{\theta}_0$. Here the second to the last line holds by using the same arguments as those lead to Theorem A.1 and the fact that $|\psi_i'''(\boldsymbol{\alpha}, \boldsymbol{\Gamma})| > 0$ by Condition (C1), and $\left| \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right) (\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) / \{3\lambda \sqrt{q+m+k_0} / (c_\psi \alpha_{\min} - 3\mu/2)\} \right| = O_p(1)$. Furthermore, by Lemma A.12, we have

$$\begin{aligned}
&\sup_{\mathbf{v}, \mathbf{w} \in \mathbb{K}} \mathbf{v}^T \left\{ \frac{\partial^2 \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} - \mathbf{Q}(\boldsymbol{\theta}_0) \right\} \mathbf{w} \\
&= \sqrt{\log(\max(p, n)) / n} \\
&= o_p(\sqrt{q+m+k_0} [\sqrt{\log\{\max(p, n)\}}/n + k_0^{1/2} N^{-\omega}]), \tag{35}
\end{aligned}$$

with probability greater than $1 - a_3 \max(p, n)^{-1}$ for some constants $a_3 > 0$. Now define

$\mathbb{K}_1 \equiv \{\mathbf{v} = (\mathbf{v}_1^T, \mathbf{v}_{2j}^T, j = 1, \dots, d)^T, \|\mathbf{v}_{2j}\|_2 > \mathbf{0} \text{ for only one } j, \|\mathbf{v}\|_2 \leq 1\}$, similar to (34)

there are constants $a_4, a_5 > 0$ such that

$$\sup_{\mathbf{v} \in \mathbb{K}_1, \mathbf{w} \in \mathbb{K}} \left| \mathbf{v}^T \left\{ \hat{\mathbf{Q}}(\boldsymbol{\theta}^*) - \frac{\partial^2 \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} \right\} \mathbf{w} \right|$$

$$\begin{aligned}
&\leq \sup_{\mathbf{v} \in \mathbb{K}_1, \mathbf{w} \in \mathbb{K}} |n^{-1} \mathbf{v}_1^T \sum_{i=1}^n \psi_i'''(\boldsymbol{\alpha}^{**}, \boldsymbol{\Gamma}^{**}) \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} \mathbf{w} \\
&\quad \times \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right) (\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) \\
&\leq \{3\lambda\sqrt{q+m+k_0}/(c_\psi\alpha_{\min} - 3\mu/2)\} \\
&\quad \times \sup_{\mathbf{v} \in \mathbb{K}_1, \mathbf{w} \in \mathbb{K}} \left| n^{-1} \mathbf{v}^T \sum_{i=1}^n |\psi_i'''(\boldsymbol{\alpha}^{**}, \boldsymbol{\Gamma}^{**})| \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} \mathbf{w} \right. \\
&\quad \times \left| \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right) (\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) / \{3\lambda\sqrt{q+m+k_0}/(c_\psi\alpha_{\min} - 3\mu/2)\} \right| \\
&\leq \{3\lambda\sqrt{q+m+k_0}/(c_\psi\alpha_{\min} - 3\mu/2)\} / 4 \\
&\quad \times \sup_{\mathbf{v} \in \mathbb{K}_1, \mathbf{w} \in \mathbb{K}} \left| n^{-1} (\mathbf{v} + \mathbf{w})^T \sum_{i=1}^n |\psi_i'''(\boldsymbol{\alpha}^{**}, \boldsymbol{\Gamma}^{**})| \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} (\mathbf{v} + \mathbf{w}) \right. \\
&\quad \times \left| \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right) (\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) / \{3\lambda\sqrt{q+m+k_0}/(c_\psi\alpha_{\min} - 3\mu/2)\} \right| \\
&\quad + \{3\lambda\sqrt{q+m+k_0}/(c_\psi\alpha_{\min} - 3\mu/2)\} / 4 \\
&\quad \times \sup_{\mathbf{v} \in \mathbb{K}_1, \mathbf{w} \in \mathbb{K}} \left| n^{-1} (\mathbf{v} - \mathbf{w})^T \sum_{i=1}^n |\psi_i'''(\boldsymbol{\alpha}^{**}, \boldsymbol{\Gamma}^{**})| \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)^{\otimes 2} (\mathbf{v} - \mathbf{w}) \right. \\
&\quad \times \left| \left(\left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right) (\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) / \{3\lambda\sqrt{q+m+k_0}/(c_\psi\alpha_{\min} - 3\mu/2)\} \right| \\
&\leq [a_4 + \{\log(p) + q + N\}\sqrt{q+m+k_0} + 1/n] \{3\lambda\sqrt{q+m+k_0}/(c_\psi\alpha_{\min} - 3\mu/2)\} \\
&= O_p \left(\sqrt{q+m+k_0} [\sqrt{\log\{\max(p, n)\}}/n + k_0^{1/2} N^{-\omega}] \right), \tag{36}
\end{aligned}$$

with probability $1 - 2 \exp\{-a_5 n \min(c_\psi^2 \alpha_{\min}^2 / 54^2, 1) / 2\} - 20 \max(p, n)^{-1}$. Similar to (35)

$$\begin{aligned}
&\sup_{\mathbf{v} \in \mathbb{K}_1, \mathbf{w} \in \mathbb{K}} \mathbf{v}^T \left\{ \frac{\partial^2 \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} - \mathbf{Q}(\boldsymbol{\theta}_0) \right\} \mathbf{w} \\
&= o_p(\sqrt{q+m+k_0} [\sqrt{\log\{\max(p, n)\}}/n + k_0^{1/2} N^{-\omega}]). \tag{37}
\end{aligned}$$

Combine (34) and (35) and Lemma A.6 in the supplementary material, we have

$$\| \hat{\mathbf{Q}}_{\mathcal{M} \cup \mathcal{S}, \mathcal{M} \cup \mathcal{S}}(\boldsymbol{\theta}^*)^{-1} - \mathbf{Q}_{\mathcal{M} \cup \mathcal{S}, \mathcal{M} \cup \mathcal{S}}(\boldsymbol{\theta}_0)^{-1} \|_2$$

$$\begin{aligned}
&\leq O_p(\widehat{\mathbf{Q}}_{\mathcal{M} \cup S, \mathcal{M} \cup S}(\boldsymbol{\theta}^*)^{-1} - \mathbf{Q}_{\mathcal{M} \cup S, \mathcal{M} \cup S}(\boldsymbol{\theta}_0)^{-1}) \\
&= O_p\left(\sqrt{q+m+k_0}[\sqrt{\log\{\max(p,n)\}/n} + k_0^{1/2}N^{-\omega}]\right), \tag{38}
\end{aligned}$$

with probability $1 - \exp\{-O(n)\} - O\{\max(p,n)^{-1}\}$, and therefore

$$\begin{aligned}
&\sup_{j \in \mathcal{M} \cup S} \sum_{k \in S} \|\mathbf{E}_j^T [\{\widehat{\mathbf{Q}}_{\mathcal{M} \cup S, \mathcal{M} \cup S}(\boldsymbol{\theta}^*)\}^{-1} - \{\mathbf{Q}_{\mathcal{M} \cup S, \mathcal{M} \cup S}(\boldsymbol{\theta}^*)\}^{-1}] \mathbf{E}_k\|_2 \\
&= O_p\left\{(q+m+k_0)[\sqrt{\log\{\max(p,n)\}/n} + k_0^{1/2}N^{-\omega}]\right\} \\
&= o_p(1), \tag{39}
\end{aligned}$$

where \mathbf{E}_j is a matrix so that $\mathbf{E}_j^T(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) = N^{-1/2}\widehat{\boldsymbol{\gamma}}_j - N^{-1/2}\boldsymbol{\Gamma}_{0,j}$.

Furthermore, let $\mathbb{K}_2 \equiv \{\mathbf{v} = (\mathbf{v}_1^T, \mathbf{v}_{2j}^T, j = 1, \dots, d)^T, \|\mathbf{v}_{2j}\|_2 > \mathbf{0} \text{ for only one } j \in \mathcal{M} \cup S, \|\mathbf{v}_{2j}\|_2 = \mathbf{0} \text{ for } j \in (\mathcal{M} \cup S)^c, \|\mathbf{v}\|_2 \leq 1\}$, then

$$\begin{aligned}
&\sup_{\mathbf{v} \in \mathbb{K}_1} \left| \mathbf{v}_{\mathcal{M} \cup S}^T \widehat{\mathbf{Q}}_{(\mathcal{M} \cup S)^c, \mathcal{M} \cup S}(\boldsymbol{\theta}^*) \{\widehat{\mathbf{Q}}_{(\mathcal{M} \cup S), \mathcal{M} \cup S}(\boldsymbol{\theta}^*)\}^{-1} \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} \right| \\
&\leq \sup_{\mathbf{v} \in \mathbb{K}_1} \left| \mathbf{v}_{\mathcal{M} \cup S}^T [\widehat{\mathbf{Q}}_{(\mathcal{M} \cup S)^c, \mathcal{M} \cup S}(\boldsymbol{\theta}^*) - \mathbf{Q}_{(\mathcal{M} \cup S)^c, \mathcal{M} \cup S}(\boldsymbol{\theta}_0)] \{\widehat{\mathbf{Q}}_{(\mathcal{M} \cup S), \mathcal{M} \cup S}(\boldsymbol{\theta}^*)\}^{-1} \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} \right| \\
&\quad + \sup_{\mathbf{v} \in \mathbb{K}_1} \left| \mathbf{v}_{\mathcal{M} \cup S}^T \mathbf{Q}_{(\mathcal{M} \cup S)^c, \mathcal{M} \cup S}(\boldsymbol{\theta}_0) \{\widehat{\mathbf{Q}}_{(\mathcal{M} \cup S), \mathcal{M} \cup S}(\boldsymbol{\theta}^*)\}^{-1} \right. \\
&\quad \left. - \{\mathbf{Q}_{(\mathcal{M} \cup S), \mathcal{M} \cup S}(\boldsymbol{\theta}_0)\}^{-1} \right\} \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} \right| \\
&\quad + \sup_{\mathbf{v} \in \mathbb{K}_1} \left| \mathbf{v}_{\mathcal{M} \cup S}^T \mathbf{Q}_{(\mathcal{M} \cup S)^c, \mathcal{M} \cup S}(\boldsymbol{\theta}_0) \{\mathbf{Q}_{(\mathcal{M} \cup S), \mathcal{M} \cup S}(\boldsymbol{\theta}_0)\}^{-1} \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} \right| \\
&\leq \sup_{\mathbf{v} \in \mathbb{K}_1, \mathbf{w} \in \mathbb{K}} \left| \mathbf{v}_{1(\mathcal{M} \cup S)^c}^T [\widehat{\mathbf{Q}}_{(\mathcal{M} \cup S)^c, \mathcal{M} \cup S}(\boldsymbol{\theta}^*) - \mathbf{Q}_{(\mathcal{M} \cup S)^c, \mathcal{M} \cup S}(\boldsymbol{\theta}_0)] \mathbf{w}_{\mathcal{M} \cup S} \right| \\
&\quad \times \left\| \{\widehat{\mathbf{Q}}_{(\mathcal{M} \cup S), \mathcal{M} \cup S}(\boldsymbol{\theta}^*)\}^{-1} \right\|_2 \left\| \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} \right\|_2 \\
&\quad + \left\| \mathbf{Q}_{(\mathcal{M} \cup S)^c, \mathcal{M} \cup S}(\boldsymbol{\theta}_0) \right\|_2 \left\| \{\widehat{\mathbf{Q}}_{(\mathcal{M} \cup S), \mathcal{M} \cup S}(\boldsymbol{\theta}^*)\}^{-1} - \{\mathbf{Q}_{(\mathcal{M} \cup S), \mathcal{M} \cup S}(\boldsymbol{\theta}_0)\}^{-1} \right\|_2 \left\| \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} \right\|_2 \\
&\quad + \sup_{\mathbf{v} \in \mathbb{K}_1} \left| \mathbf{v}_{\mathcal{M} \cup S}^T \mathbf{Q}_{(\mathcal{M} \cup S)^c, \mathcal{M} \cup S}(\boldsymbol{\theta}_0) \{\mathbf{Q}_{(\mathcal{M} \cup S), \mathcal{M} \cup S}(\boldsymbol{\theta}_0)\}^{-1} \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} \right| \\
&\leq O_p\left((q+m+k_0)[\sqrt{\log\{\max(p,n)\}/n} + k_0^{1/2}N^{-\omega}]^2\right) \tag{40} \\
&\quad + \sup_{\mathbf{v} \in \mathbb{K}_1} \left| \mathbf{v}_{\mathcal{M} \cup S}^T \mathbf{Q}_{(\mathcal{M} \cup S)^c, \mathcal{M} \cup S}(\boldsymbol{\theta}_0) \{\mathbf{Q}_{(\mathcal{M} \cup S), \mathcal{M} \cup S}(\boldsymbol{\theta}_0)\}^{-1} \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} \right|
\end{aligned}$$

$$\begin{aligned}
&= O_p \left((q + m + k_0) [\sqrt{\log\{\max(p, n)\}/n} + k_0^{1/2} N^{-\omega}]^2 \right. \\
&\quad \left. + \sqrt{q + m + k_0} [\sqrt{\log\{\max(p, n)\}/n} + k_0^{1/2} N^{-\omega}] \right) \tag{41}
\end{aligned}$$

$$= O_p \left(\sqrt{q + m + k_0} [\sqrt{\log\{\max(p, n)\}/n} + k_0^{1/2} N^{-\omega}] \right), \tag{42}$$

with probability greater than $1 - \exp\{-O(n)\} - O\{\max(p, n)^{-1}\}$, (40) is obtained by using (36) and (37) and (38) and the fact that

$$\begin{aligned}
&\|\{\widehat{\mathbf{Q}}_{(\mathcal{M} \cup \mathcal{S}), \mathcal{M} \cup \mathcal{S}}(\boldsymbol{\beta}^*)\}^{-1}\|_2 \\
&\leq \|\{\widehat{\mathbf{Q}}_{(\mathcal{M} \cup \mathcal{S}), \mathcal{M} \cup \mathcal{S}}(\boldsymbol{\beta}^*)\}^{-1} - [\mathbf{Q}_{(\mathcal{M} \cup \mathcal{S}), \mathcal{M} \cup \mathcal{S}}(\boldsymbol{\beta})]^{-1}\|_2 + \|\{\mathbf{Q}_{(\mathcal{M} \cup \mathcal{S}), \mathcal{M} \cup \mathcal{S}}(\boldsymbol{\beta})\}^{-1}\|_2 \\
&\leq O_p(1).
\end{aligned}$$

And (41) holds by using the same argument as those lead to Lemma A.15. (42) holds by the assumption that $(q + m + k_0)\sqrt{\log\{\max(p, n)\}/n} \rightarrow 0$ and $k_0 N^{-2\omega} = O[\log\{\max(p, n)\}/n]$. By the assumption that $k_0 N^{-2\omega} = O[\log\{\max(p, n)\}/n]$, $\lambda = O_p([\log\{\max(p, n)\}/n]^{1/4})$, and the assumption that we have

$$\sup_{\mathbf{v} \in \mathbb{K}_1} \left| \mathbf{v}_{\mathcal{M} \cup \mathcal{S}}^T \widehat{\mathbf{Q}}_{(\mathcal{M} \cup \mathcal{S})^c, \mathcal{M} \cup \mathcal{S}}(\boldsymbol{\theta}^*) \{\widehat{\mathbf{Q}}_{(\mathcal{M} \cup \mathcal{S}), \mathcal{M} \cup \mathcal{S}}(\boldsymbol{\theta}^*)\}^{-1} \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup \mathcal{S}} \right| = o_p(\lambda). \tag{43}$$

Therefore,

$$\begin{aligned}
&\max_{j \in (\mathcal{M} \cup \mathcal{S})^c} \|\widehat{\mathbf{z}}_j\|_2 \tag{44} \\
&\leq \lambda^{-1} \sup_{\mathbf{v} \in \mathbb{K}_1} \left| \mathbf{v}_{2(\mathcal{M} \cup \mathcal{S})^c}^T \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{2(\mathcal{M} \cup \mathcal{S})^c} \right| \\
&\quad + \lambda^{-1} \sup_{\mathbf{v} \in \mathbb{K}_1} \left| \mathbf{v}_{2(\mathcal{M} \cup \mathcal{S})^c}^T \widehat{\mathbf{Q}}_{(\mathcal{M} \cup \mathcal{S})^c(\mathcal{M} \cup \mathcal{S})}(\boldsymbol{\theta}^*) \{\widehat{\mathbf{Q}}_{(\mathcal{M} \cup \mathcal{S})(\mathcal{M} \cup \mathcal{S})}(\boldsymbol{\theta}^*)\}^{-1} \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup \mathcal{S}} \right| \\
&\quad - \sup_{\mathbf{v} \in \mathbb{K}_1} \left| \mathbf{v}_{2(\mathcal{M} \cup \mathcal{S})^c}^T \widehat{\mathbf{Q}}_{(\mathcal{M} \cup \mathcal{S})^c(\mathcal{M} \cup \mathcal{S})}(\boldsymbol{\theta}^*) \{\widehat{\mathbf{Q}}_{(\mathcal{M} \cup \mathcal{S})(\mathcal{M} \cup \mathcal{S})}(\boldsymbol{\theta}^*)\}^{-1} \right. \\
&\quad \times \left[\left\{ \frac{\partial \sum_{j \in \mathcal{M}^c} q_\lambda(\|\widehat{\boldsymbol{\gamma}}_j\|_2)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup \mathcal{S}} - \lambda \widehat{\mathbf{z}}_{\mathcal{M} \cup \mathcal{S}} \right] \left. \right| \\
&\leq - \sup_{\mathbf{v} \in \mathbb{K}_1} \left| \mathbf{v}_{2(\mathcal{M} \cup \mathcal{S})^c}^T \widehat{\mathbf{Q}}_{(\mathcal{M} \cup \mathcal{S})^c(\mathcal{M} \cup \mathcal{S})}(\boldsymbol{\theta}^*) \{\widehat{\mathbf{Q}}_{(\mathcal{M} \cup \mathcal{S})(\mathcal{M} \cup \mathcal{S})}(\boldsymbol{\theta}^*)\}^{-1} \right.
\end{aligned}$$

$$\times \left[\left\{ \frac{\partial \sum_{j \in \mathcal{M}^c} q_\lambda(\|\hat{\gamma}_j\|_2)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} - \lambda \hat{\mathbf{z}}_{\mathcal{M} \cup S} \right] + o_p(\lambda)$$

where the last inequality holds by using the same arguments leads to Lemma A.15 and (43). Also by (33) we have for $j \in \mathcal{M} \cup S$, recall that \mathbf{E}_j is a matrix so that $\mathbf{E}_j^T(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) = N^{-1/2}\hat{\boldsymbol{\gamma}}_j - N^{-1/2}\boldsymbol{\Gamma}_{0,j}$

$$\begin{aligned} & \sup_{j \in S} N^{-1/2} \|\hat{\boldsymbol{\gamma}}_j - \boldsymbol{\Gamma}_{0,j}\|_2 \\ & \leq \sup_{j \in S} \|\mathbf{E}_j^T \{\hat{\mathbf{Q}}_{\mathcal{M} \cup S, \mathcal{M} \cup S}(\boldsymbol{\theta}^*)\}^{-1} \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S}\|_2 \\ & \quad + \sup_{j \in S} \|\mathbf{E}_j^T \{\hat{\mathbf{Q}}_{\mathcal{M} \cup S, \mathcal{M} \cup S}(\boldsymbol{\theta}^*)\}^{-1} \left[\left\{ \frac{\partial \sum_{j \in \mathcal{M}^c} q_\lambda(\|\hat{\gamma}_j\|_2)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} - \lambda \hat{\mathbf{z}}_{\mathcal{M} \cup S} \right]\| \\ & = \sup_{j \in S} \|\mathbf{E}_j^T \{\hat{\mathbf{Q}}_{\mathcal{M} \cup S, \mathcal{M} \cup S}(\boldsymbol{\theta}^*)\}^{-1} \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S}\| \\ & \quad + \sup_{j \in \mathcal{M} \cup S} \left\| \sum_{k \in S} \mathbf{E}_j^T \{\hat{\mathbf{Q}}_{\mathcal{M} \cup S, \mathcal{M} \cup S}(\boldsymbol{\theta}^*)\}^{-1} \mathbf{E}_k \left[\frac{\partial q_\lambda(\|\hat{\gamma}_k\|_2)}{\partial \hat{\gamma}_k} - \lambda \hat{\mathbf{z}}_k \right] \right\| \\ & = o_p(\lambda) + \lambda \sup_{j \in S} \sum_{k \in S} \|\mathbf{E}_j^T \{\hat{\mathbf{Q}}_{\mathcal{M} \cup S, \mathcal{M} \cup S}(\boldsymbol{\theta}^*)\}^{-1} \mathbf{E}_k\|_2 \\ & \leq o_p(\lambda) + 2\lambda c_\infty \\ & \leq 3\lambda c_\infty, \end{aligned} \tag{45}$$

where the third to the last line holds by Lemma A.5 that $\sup_{k \in S} \left\| \frac{\partial q_\lambda(\|\hat{\gamma}_k\|_2)}{\partial \hat{\gamma}_k} - \lambda \hat{\mathbf{z}}_k \right\|_2 \leq \lambda$.

The second to the last line holds because

$$\begin{aligned} & \sup_{j \in S} \sum_{k \in S} \|\mathbf{E}_j^T \{\hat{\mathbf{Q}}_{\mathcal{M} \cup S, \mathcal{M} \cup S}(\boldsymbol{\theta}^*)\}^{-1} \mathbf{E}_k\|_2 \\ & \leq \sup_{j \in S} \sum_{k \in S} \|\mathbf{E}_j^T [\{\hat{\mathbf{Q}}_{\mathcal{M} \cup S, \mathcal{M} \cup S}(\boldsymbol{\theta}^*)\}^{-1} - \{\mathbf{Q}_{\mathcal{M} \cup S, \mathcal{M} \cup S}(\boldsymbol{\theta}^*)\}^{-1}] \mathbf{E}_k\|_2 \\ & \quad + \sup_{j \in S} \sum_{k \in S} \|\mathbf{E}_j^T \{\mathbf{Q}_{\mathcal{M} \cup S, \mathcal{M} \cup S}(\boldsymbol{\theta}^*)\}^{-1} \mathbf{E}_k\|_2 \\ & \leq 2c_\infty, \end{aligned}$$

by the assumption that $\sup_{j \in S} \sum_{k \in S} \|\mathbf{E}_j^T \{\mathbf{Q}_{\mathcal{M} \cup S, \mathcal{M} \cup S}(\boldsymbol{\theta}^*)\}^{-1} \mathbf{E}_k\|_2 \leq c_\infty$ and (39). Now be-

cause $\min_{j \in S} (\|\boldsymbol{\gamma}_j\|_2) \geq \lambda(\gamma + 3c_\infty)$, by (45), we have

$$\min_{j \in S} (\|\hat{\boldsymbol{\gamma}}_j\|_2) \geq \lambda\gamma,$$

by Lemma A.4 in the in the supplementary material, we have

$$\sup_{k \in S} \left[\frac{\partial q_\lambda(\|\hat{\boldsymbol{\gamma}}_k\|_2)}{\partial \hat{\boldsymbol{\gamma}}_k} - \lambda \hat{\boldsymbol{z}}_k \right] = \mathbf{0}. \quad (46)$$

Now plug this result in (44), we have

$$\max_{j \in (\mathcal{M} \cup S)^c} \|\hat{\boldsymbol{z}}_j\|_2 = o_p(1),$$

and therefore, $\max_{j \in (\mathcal{M} \cup S)^c} \|\hat{\boldsymbol{z}}_j\|_2 < 1$ with probability greater than $1 - \exp\{-O(n)\} - O\{\max(p, n)^{-1}\}$.

Therefore, Theorem A.1 applies so that $\hat{\boldsymbol{\theta}} = \{\hat{\boldsymbol{\theta}}_{\mathcal{M} \cup S}^T, \mathbf{0}^T\}^T$ is the unique solution for (3) and also plug (46) in to (45)

$$\begin{aligned} & \hat{\boldsymbol{\theta}}_{\mathcal{M} \cup S} - \boldsymbol{\theta}_{0, \mathcal{M} \cup S} \\ &= -\{\mathbf{Q}_{\mathcal{M} \cup S, \mathcal{M} \cup S}(\boldsymbol{\theta}_0)\}^{-1} \left\{ \frac{\partial \mathcal{L}(\boldsymbol{\theta}_0)}{\partial \boldsymbol{\theta}} \right\}_{\mathcal{M} \cup S} + o_p(1) \\ &= -\{\mathbf{Q}_{\mathcal{M} \cup S, \mathcal{M} \cup S}\{\boldsymbol{\alpha}_0, \boldsymbol{\beta}(\cdot)\}\}^{-1} \left\{ n^{-1} \sum_{i=1}^n \{\psi'_i\{\boldsymbol{\alpha}_0, \boldsymbol{\beta}(s)\} - Y_i\} \left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right\}_{\mathcal{M} \cup S} \\ & \quad \times \{1 + o_p(1)\}, \end{aligned}$$

with probability greater than $1 - \exp\{-(n)\} - O\{\max(p, n)^{-1}\}$, where the last inequality holds by using the same arguments as those lead to Lemma A.15, and the fact $\sup_{s \in [0, 1]} |\mathbf{B}^T(s) \boldsymbol{\gamma}_{0j} - \beta_j(s)| = O_p(N^{-\omega})$ in Condition (C3) and the condition the assumption that $k_0 N^{-2\omega} = o[\log\{\max(p, n)\}/n]$. This proves the result. \square

Proof of Lemma 1:

$$\begin{aligned} & \boldsymbol{\Psi}^{-1/2}(\boldsymbol{\Sigma}, \mathbf{Q}, \boldsymbol{\beta}) \boldsymbol{\omega}_n \\ &= -\sqrt{n} \boldsymbol{\Psi}^{-1/2}(\boldsymbol{\Sigma}, \mathbf{Q}, \boldsymbol{\beta}) \mathbf{A} \mathbf{Q}_{\mathcal{M} \cup S, \mathcal{M} \cup S}^{-1} \{\boldsymbol{\alpha}_0, \boldsymbol{\beta}(\cdot)\}_{\mathcal{M} \cup S, \mathcal{M} \cup S}^{-1} \end{aligned}$$

$$\left\{ n^{-1} \sum_{i=1}^n \{ \psi'_i \{ \boldsymbol{\alpha}_0, \boldsymbol{\beta}(s) \} - Y_i \} \left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right\}_{\mathcal{M} \cup \mathcal{S}}.$$

It is easy to see that

$$\begin{aligned} & \sum_{i=1}^n \text{cov} \left(\boldsymbol{\Psi}^{-1/2}(\boldsymbol{\Sigma}, \mathbf{Q}, \boldsymbol{\beta}) \mathbf{A} \mathbf{Q}_{\mathcal{M} \cup \mathcal{S}, \mathcal{M} \cup \mathcal{S}}^{-1} \{ \boldsymbol{\alpha}_0, \boldsymbol{\beta}(\cdot) \}_{\mathcal{M} \cup \mathcal{S}, \mathcal{M} \cup \mathcal{S}}^{-1} \right. \\ & \left. \times n^{-1/2} \{ \psi'_i \{ \boldsymbol{\alpha}_0, \boldsymbol{\beta}(s) \} - Y_i \} \left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right)_{\mathcal{M} \cup \mathcal{S}} = \mathbf{I}_{rN \times rN}. \end{aligned}$$

Also,

$$\begin{aligned} & (rN)^{1/4} \sum_{i=1}^n E \left[\left\| \boldsymbol{\Psi}^{-1/2}(\boldsymbol{\Sigma}, \mathbf{Q}, \boldsymbol{\beta}) \mathbf{A} \mathbf{Q}_{\mathcal{M} \cup \mathcal{S}, \mathcal{M} \cup \mathcal{S}}^{-1} \{ \boldsymbol{\alpha}_0, \boldsymbol{\beta}(\cdot) \}_{\mathcal{M} \cup \mathcal{S}, \mathcal{M} \cup \mathcal{S}}^{-1} \right. \right. \\ & \left. \left. \times n^{-1/2} \{ \psi'_i \{ \boldsymbol{\alpha}_0, \boldsymbol{\beta}(s) \} - Y_i \} \left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right\|_2^3 \right] \\ & = O\{(rN)^{1/4} n^{-1/2}\} = o(1), \end{aligned}$$

where the second to the last equality holds by the assumption that

$$\begin{aligned} & E \left[\left\| \boldsymbol{\Psi}^{-1/2}(\boldsymbol{\Sigma}, \mathbf{Q}, \boldsymbol{\beta}) \mathbf{A} \mathbf{Q}_{\mathcal{M} \cup \mathcal{S}, \mathcal{M} \cup \mathcal{S}}^{-1} \{ \boldsymbol{\alpha}_0, \boldsymbol{\beta}(\cdot) \}_{\mathcal{M} \cup \mathcal{S}, \mathcal{M} \cup \mathcal{S}}^{-1} \right. \right. \\ & \left. \left. \times \{ \psi'_i \{ \boldsymbol{\alpha}_0, \boldsymbol{\beta}(s) \} - Y_i \} \left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right\|_2^3 \right] = O(1), \end{aligned}$$

and the last equality holds by the assumption that $(rN)^{1/4} n^{-1/2} \rightarrow 0$. Using the Lemma A.7 from the supplementary material, let \mathcal{Z} be a standard Gaussian random variable, we have

$$\limsup_{n \rightarrow \infty} \sup_{\mathcal{C}} \left| \Pr \left[\boldsymbol{\Psi}^{-1/2}(\boldsymbol{\Sigma}, \mathbf{Q}, \boldsymbol{\beta}) \boldsymbol{\omega}_n \in \mathcal{C} \right] - \Pr(\mathcal{Z} \in \mathcal{C}) \right| = 0.$$

Now we choose $\mathcal{C}_x = \{ \mathbf{z} : \|\mathbf{z} - \mathbf{c}_n\|_2 \leq x \}$, we have

$$\lim_{n \rightarrow \infty} \left| \Pr \left[\boldsymbol{\Psi}^{-1/2}(\boldsymbol{\Sigma}, \mathbf{Q}, \boldsymbol{\beta}) \boldsymbol{\omega}_n \in \mathcal{C}_x \right] - \Pr(\mathcal{Z} \in \mathcal{C}_x) \right| = 0,$$

which implies

$$\lim_{n \rightarrow \infty} |\Pr(T_0 \leq x) - \Pr\{\chi^2(rN, \mathbf{c}_n^T \Psi^{-1}(\Sigma, \mathbf{Q}, \beta) \mathbf{c}_n) \leq x\}| = 0,$$

where $\chi^2(r, \gamma)$ is a r degree of freedom non-central chi-square distribution, with non-centrality parameter γ . □

Proof of Theorem 3: From Theorem 2, we have

$$\begin{aligned} & \widehat{\boldsymbol{\theta}}_{\mathcal{M} \cup \mathcal{S}} - \boldsymbol{\theta}_{0, \mathcal{M} \cup \mathcal{S}} \\ &= -[\mathbf{Q}_{\mathcal{M} \cup \mathcal{S}, \mathcal{M} \cup \mathcal{S}}\{\boldsymbol{\alpha}_0, \beta(\cdot)\}]^{-1} \left\{ \sum_{i=1}^n \{\psi'_i\{\boldsymbol{\alpha}_0, \beta(s)\} - Y_i\} \left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right\}_{\mathcal{M} \cup \mathcal{S}} \\ & \quad \times \{1 + o_p(1)\} \end{aligned}$$

and $\widehat{\boldsymbol{\theta}}_{(\mathcal{M} \cup \mathcal{S})^c} = \mathbf{0}$ with probability greater than $1 - \exp\{-O(n)\} - O\{\max(p, n)^{-1}\}$. Let

$\widehat{\boldsymbol{\theta}}_{\mathcal{M}} \equiv (\widehat{\boldsymbol{\theta}}_1^T, \widehat{\boldsymbol{\theta}}_{2j}^T, j \in \mathcal{M})^T$, and $\boldsymbol{\theta}_{0, \mathcal{M}} \equiv (\boldsymbol{\theta}_{01}^T, \boldsymbol{\theta}_{02j}^T, j \in \mathcal{M})^T$. Then

$$\begin{aligned} & \sqrt{n} \mathbf{A}(\widehat{\boldsymbol{\theta}}_{\mathcal{M}} - \boldsymbol{\theta}_{0, \mathcal{M}}) \\ &= -\sqrt{n} \mathbf{A}[\mathbf{Q}_{\mathcal{M} \cup \mathcal{S}, \mathcal{M} \cup \mathcal{S}}\{\boldsymbol{\alpha}_0, \beta(\cdot)\}]^{-1} \\ & \quad \times \left\{ \sum_{i=1}^n \{\psi'_i\{\boldsymbol{\alpha}_0, \beta(s)\} - Y_i\} \left[\mathbf{z}_i^T, N^{1/2} \left\{ \int_0^1 \mathbf{X}_i(s) \otimes \mathbf{B}(s) ds \right\}^T \right]^T \right\}_{\mathcal{M} \cup \mathcal{S}} \\ &= \boldsymbol{\omega}_n \{1 + o_p(1)\}. \end{aligned}$$

Furthermore, because when $\mathbf{C}\boldsymbol{\beta}_{\mathcal{M}}(s) = \mathbf{t}(s) + \mathbf{h}_n(s)$, we have

$$\begin{aligned} & \text{vec} \left[\left\{ \int_0^1 \mathbf{B}(s) \mathbf{B}(s)^T ds \right\}^{-1} \int_0^1 \mathbf{B}(s) \boldsymbol{\beta}_{\mathcal{M}}(s)^T ds \mathbf{C}^T \right] \\ &= \text{vec} \left[\left\{ \int_0^1 \mathbf{B}(s) \mathbf{B}(s)^T ds \right\}^{-1} \int_0^1 \mathbf{B}(s) \{\mathbf{t}(s)^T + \mathbf{h}_n^T(s)\} ds \right]. \end{aligned}$$

Hence we have

$$\sqrt{n} \Psi^{-1/2}(\widehat{\Sigma}, \widehat{\mathbf{Q}}, \widehat{\boldsymbol{\theta}}) \left(\mathbf{A} \widehat{\boldsymbol{\theta}}_{\mathcal{M}} - N^{-1/2} \text{vec} \left[\left\{ \int_0^1 \mathbf{B}(s) \mathbf{B}(s)^T ds \right\}^{-1} \int_0^1 \mathbf{B}(s) \mathbf{t}(s)^T ds \right] \right)$$

$$\begin{aligned}
&= \sqrt{n}\Psi^{-1/2}(\widehat{\Sigma}, \widehat{\mathbf{Q}}, \widehat{\boldsymbol{\theta}}) \left(\mathbf{A}\widehat{\boldsymbol{\theta}}_{\mathcal{M}} - \mathbf{A}\widehat{\boldsymbol{\theta}}_{\mathcal{M}} \right) \\
&\quad + \sqrt{n}\Psi^{-1/2}(\widehat{\Sigma}, \widehat{\mathbf{Q}}, \widehat{\boldsymbol{\theta}}) \left(\mathbf{A}\widehat{\boldsymbol{\theta}}_{\mathcal{M}} - N^{-1/2}\text{vec} \left[\left\{ \int_0^1 \mathbf{B}(s)\mathbf{B}(s)^\text{T} ds \right\}^{-1} \int_0^1 \mathbf{B}(s)\{\mathbf{t}(s)^\text{T} + \mathbf{h}_n(s)^\text{T}\} ds \right] \right) \\
&\quad + \sqrt{n}\Psi^{-1/2}(\widehat{\Sigma}, \widehat{\mathbf{Q}}, \widehat{\boldsymbol{\theta}}) \left(\text{vec} \left[N^{-1/2} \left\{ \int_0^1 \mathbf{B}(s)\mathbf{B}(s)^\text{T} ds \right\}^{-1} \int_0^1 \mathbf{B}(s)\mathbf{h}_n(s)^\text{T} ds \right] \right) \\
&= \sqrt{n}\Psi^{-1/2}(\widehat{\Sigma}, \widehat{\mathbf{Q}}, \widehat{\boldsymbol{\theta}})(\boldsymbol{\omega}_n + \mathbf{c}_n)\{1 + o_p(1)\} + O_p\{n^{1/2}(rN)^{1/2}N^{-1/2-\omega}\},
\end{aligned}$$

where the last equality holds by the Condition that $\sup_{s \in [0,1]} |\mathbf{B}(s)^\text{T} N^{-1/2} \boldsymbol{\gamma}_{0j} - N^{-1/2} \beta_j(s)| = N^{-\omega-1/2}$. Therefore,

$$\begin{aligned}
&n \left(\mathbf{A}\widehat{\boldsymbol{\theta}}_{\mathcal{M}} - N^{-1/2}\text{vec} \left[\left\{ \int_0^1 \mathbf{B}(s)\mathbf{B}(s)^\text{T} ds \right\}^{-1} \int_0^1 \mathbf{B}(s)\mathbf{t}(s)^\text{T} ds \right] \right)^\text{T} \\
&\quad \times \Psi^{-1}(\widehat{\Sigma}, \widehat{\mathbf{Q}}, \widehat{\boldsymbol{\theta}}) \left(\mathbf{A}\widehat{\boldsymbol{\theta}}_{\mathcal{M}} - N^{-1/2}\text{vec} \left[\left\{ \int_0^1 \mathbf{B}(s)\mathbf{B}(s)^\text{T} ds \right\}^{-1} \int_0^1 \mathbf{B}(s)\mathbf{t}(s)^\text{T} ds \right] \right) \\
&= n(\boldsymbol{\omega}_n + \mathbf{c}_n)^\text{T} \Psi^{-1/2}(\widehat{\Sigma}, \widehat{\mathbf{Q}}, \widehat{\boldsymbol{\theta}})(\boldsymbol{\omega}_n + \mathbf{c}_n)\{1 + o_p(1)\} + o_p\{rN\} \\
&= n(\boldsymbol{\omega}_n + \mathbf{c}_n)^\text{T} \Psi^{-1/2}(\boldsymbol{\Sigma}, \mathbf{Q}, \boldsymbol{\beta})(\boldsymbol{\omega}_n + \mathbf{c}_n)\{1 + o_p(1)\} + o_p\{rN\}.
\end{aligned}$$

The second to the last equality holds by the assumption that $nN^{-2\omega-1} \rightarrow 0$ and the last equality holds by the consistency of $\widehat{\Sigma}$, $\widehat{\mathbf{Q}}$, and $\widehat{\boldsymbol{\theta}}$. Therefore, we have $T - T_0 = o_p(rN)$.

This proves the result. \square

References

Vidmantas Bentkus. A lyapunov-type bound in r^d . *Theory of Probability & Its Applications*, 49(2):311–323, 2005.

Laming Chen and Yuantao Gu. The convergence guarantees of a non-convex approach for sparse recovery. *IEEE Transactions on Signal Processing*, 62(15):3754–3767, 2014.

Proloy Das and Behtash Babadi. Multitaper spectral analysis of neuronal spiking activity driven by latent stationary processes. *Signal Processing*, 170:107429, 2020.

- Ronald A DeVore and George G Lorentz. *Constructive approximation*, volume 303. Springer Science & Business Media, New York, 1993.
- John Duchi, Shai Shalev-Shwartz, Yoram Singer, and Tushar Chandra. Efficient projections onto the l_1 -ball for learning in high dimensions. In *Proceedings of the 25th international conference on Machine learning*, pages 272–279, 2008.
- Bingsheng He, Han Liu, Zhaoran Wang, and Xiaoming Yuan. A strictly contractive peaceman–rachford splitting method for convex programming. *SIAM Journal on Optimization*, 24(3):1011–1040, 2014.
- Huaqing Jin, Kamalini G Ranasinghe, Pooja Prabhu, Corby Dale, Yijing Gao, Kiwamu Kudo, Keith Vossel, Ashish Raj, Srikantan S Nagarajan, and Fei Jiang. Dynamic functional connectivity MEG features of Alzheimer’s disease. *NeuroImage*, 281:120358, 2023. ISSN 1053-8119. doi: <https://doi.org/10.1016/j.neuroimage.2023.120358>.
- Michel Ledoux and Michel Talagrand. *Probability in Banach Spaces: isoperimetry and processes*. Springer Science & Business Media, 2013.
- Po-Ling Loh and Martin J Wainwright. High-dimensional regression with noisy and missing data: provable guarantees with nonconvexity. *The Annals of Statistics*, 40(3):1637–1664, 2012.
- Po-Ling Loh and Martin J Wainwright. Regularized m-estimators with nonconvexity: Statistical and algorithmic theory for local optima. *Journal of Machine Learning Research*, 16:559–616, 2015.

- Po-Ling Loh and Martin J. Wainwright. Support recovery without incoherence: A case for nonconvex regularization. *The Annals of Statistics*, 45(6):2455–2482, 2017.
- Pravat K Mandal, Anwesha Banerjee, Manjari Tripathi, and Ankita Sharma. A comprehensive review of magnetoencephalography (MEG) studies for brain functionality in healthy aging and Alzheimer’s disease (ad). *Frontiers in computational neuroscience*, 12:60, 2018.
- Kamalini G Ranasinghe, Parul Verma, Chang Cai, Xihe Xie, Kiwamu Kudo, Xiao Gao, Hannah Lerner, Danielle Mizuiri, Amelia Strom, Leonardo Iaccarino, et al. Altered excitatory and inhibitory neuronal subpopulation parameters are distinctly associated with tau and amyloid in Alzheimer’s disease. *Elife*, 11:e77850, 2022.
- Roman Vershynin. Introduction to the non-asymptotic analysis of random matrices. *arXiv preprint arXiv:1011.3027*, 2010.
- Jean-Philippe Vial. Strong convexity of sets and functions. *Journal of Mathematical Economics*, 9(1-2):187–205, 1982.
- Cun-Hui Zhang and Tong Zhang. A general theory of concave regularization for high-dimensional sparse estimation problems. *Statistical Science*, 27(4):576–593, 2012.