

On the effect of derivative interactions in quantum field theory

Karl-Henning Rehren¹

¹Institut für Theoretische Physik, Georg-August-Universität Göttingen, 37077 Göttingen, Germany.

May 16, 2024

Abstract

There exist several good reasons why one may wish to add a total derivative to an interaction in quantum field theory, e.g., in order to improve the perturbative construction. Unlike in classical field theory, adding derivatives in general changes the theory. The analysis whether and how this can be prevented, is presently limited to perturbative orders g^n , $n \leq 3$. We drastically simplify it by an all-orders formula, which also allows to answer some salient structural questions. The method is part of a larger program to (re)derive interactions of particles by quantum consistency conditions, rather than a classical principle of gauge invariance.

1 Introduction

Total derivatives: motivations. Adding a total four-derivative $\partial_\mu V^\mu$ (of sufficiently rapid decay) to the Lagrangian density $L(x)$ of a classical field theory does not affect the Euler-Lagrange equations of motion. The reason is basically that the equations of motion are equivalent to Hamilton’s principle extremalizing the action $\int d^4x L(x)$. The total derivative contributes a boundary term to the action, that vanishes if V^μ has sufficiently fast decay.

The same is not true in quantum field theory. Most notably, the S-matrix

$$S_{L_{\text{int}}} = T e^{i \int d^4x L_{\text{int}}(x)} \quad (1.1)$$

is sensitive to derivative terms because the time-ordering does not commute with the time-derivative.

The problem has been previously studied in the context of BRST theory by the Scharf group, under the label “perturbative gauge invariance (PGI)” [17]: The cubic interaction density $L_{\text{int}}^{\text{BRST}}$ (including gauge-fixing and ghost terms) in the Standard Model (SM) is BRST invariant only up to a total derivative. Therefore, the integral $\int d^4x L_{\text{int}}^{\text{BRST}}$ is BRST invariant, but it is not obvious that the same is true for its time-ordered exponential (1.1). It was discovered that BRST invariance of the S-matrix requires higher-order interactions (like the quartic self-coupling of gluons or the self-coupling of the

*Email: krehren@uni-goettingen.de

Higgs) that can be recursively determined from the cubic interaction. In this way, parts of the SM were re-derived without assuming gauge invariance.

The present work uses techniques from the PGI analysis. It also addresses its recursive structure, but our main motivation comes from a conceptually complementary direction [16] which we call “autonomous” because it is intrinsically quantum, referring neither to canonical quantization nor gauge or BRST invariance.

We considerably extend the scope. There are two variants (called “LQ” and “LV”, see below.) The first is similar to PGI: the algebra of free fields admits a derivation, that we denote by δ . It may be the BRST variation $\delta_{\text{BRST}}(X) = [Q_{\text{BRST}}, X]_{\pm}$ as in PGI. Or it may be the ordinary variation $\delta_c(X(c))$ with respect to some auxiliary parameter or function c on which the interaction density may depend, while the S-matrix must not depend on c . In either case one wants to have

$$\delta S_{L_{\text{int}}} \stackrel{!}{=} 0, \quad \text{with} \quad L_{\text{int}} = gL_1 + \frac{g^2}{2}L_2 + \dots \quad (1.2)$$

At first order in the coupling constant g , this requires $g \int d^4x \delta L_1(x) = 0$, hence δL_1 must be the derivative of some quantity Q_1 of sufficiently rapid decay:

$$\delta L_1 = \partial_\mu Q_1^\mu. \quad (1.3)$$

We call such a structure an “ L - Q pair”. The prototype of an L - Q pair is that of string-localized QED:

$$\delta_c(A_\mu(c)j^\mu) = \partial_\mu(wj^\mu) \quad (1.4)$$

where $A_\mu(c)$ is a string-localized vector potential and w another string-localized field, both constructed on the Fock space of the photon, with the property $\delta_c A_\mu(c) = \partial_\mu w$.

Given such a pair as a necessary condition to begin with, the aim is then to develop the recursive scheme to determine L_n ($n = 2, 3, \dots$) so that (1.2) is fulfilled.

The second variant is more ambitious, by relating an interaction L_{int} to another interaction K_{int} , of which one enjoys one desirable feature which the other one doesn't, and vice versa. These features can be “Hilbert space” (unitarity), “renormalizability”, or “locality”, see [14] for several examples. Then we are asking for conditions that

$$S_{L_{\text{int}}} \stackrel{!}{=} S_{K_{\text{int}}}, \quad \text{with} \quad L_{\text{int}} = gL_1 + \frac{g^2}{2}L_2 + \dots, \quad K_{\text{int}} = gK_1 + \frac{g^2}{2}K_2 + \dots \quad (1.5)$$

so that the same S-matrix enjoys both desirable features.

Because the first perturbative orders $ig \int d^4x K_1(x)$ resp. $ig \int d^4x L_1(x)$ of the S-matrices in (1.5) do not involve time-ordering, K_1 and L_1 can only differ by a total derivative:

$$L_1(x) = K_1(x) + \partial_\mu V_1^\mu(x). \quad (1.6)$$

We call such a structure an “ L - V pair”. Like (1.4) for LQ, the prototype of an L - V pair is that of string-localized QED:

$$A_\mu(c)j^\mu = A_\mu^K j^\mu + \partial_\mu(\phi j^\mu) \quad (1.7)$$

where $A_\mu(c)$ is as in (1.4), embedded into the Fock space with indefinite metric (a Krein space) generated from the vacuum by the gauge potential A_μ^K , where also the string-localized field ϕ can be constructed.

Given such a pair, the aim is to develop the recursive scheme to determine L_n and K_n ($n = 2, 3, \dots$) so that (a strengthened version of) (1.5) is fulfilled.

The remarkable fact is that all interactions of the SM admit L - V pairs. What is more: all L - V pairs with a given field content from the SM involve the familiar cubic interactions of the SM, e.g., the cubic non-abelian YM interactions or minimal interactions of fermions with vector bosons. Recall that local and covariant massless vector gauge potentials are only defined on a Krein space, while local interactions involving massive vector bosons are non-renormalizable. This already names the main instances of the above indicated “desirable features” that may fail with either L_{int} or K_{int} . There are two ways of dealing with massive vector fields: one usually replaces them by massless gauge fields and invokes the “Higgs mechanism to make them massive”; alternatively, one may replace them by massive gauge fields plus a scalar Stückelberg field of the same mass. In both cases one buys renormalizability by indefinite metric and extra degrees of freedom, which then have to be eliminated by BRST.

Our main motivation, however, comes from “string-localized quantum field theory (sQFT)” which allows to describe both photons and massive vector bosons by covariant vector potentials on a Hilbert space, of the same UV dimension as gauge fields, which makes the said interactions renormalizable. These desirable features are bought by a weakening of the localization of the vector potentials: they are localized (in the sense of commutation relations) along a “string” or spacelike cone extending from x to infinity, labelled by a parameter function c . The point is that such potentials can be constructed without unphysical degrees of freedom on the Fock space over the respective unitary Wigner representation of the Poincaré group, and that they naturally give rise to L - Q and L - V pairs.

sQFT is reviewed in more detail in [16]. All that needs to interest us here is that one has a parameter c to vary with so as to get L - Q -pairs with respect to the derivation $\delta = \delta_c$, and one has L - V pairs connecting a renormalizable string-localized interaction on the Hilbert space either with a local gauge interaction on Krein space, or with a local but non-renormalizable interaction (or the latter two with each other) [14, 16].

Obstructions and induced interactions. The common theme of both variants, LQ or LV, is that by imposing the condition (1.2) resp. (1.5), higher-order interactions are recursively determined, starting from an initial L - Q pair or L - V pair. Because Hilbert space, covariance and locality are fundamental principles of QFT, and renormalizability is an essential feature to make reliable predictions, these conditions are indispensable for a reasonable perturbative construction. sQFT can therefore be understood as a scheme to ensure fundamental principles, that predicts interactions without recourse to a gauge principle that violates the said principles. This point is elaborated in [16].

The cubic parts (1.3) or (1.6) are the only input of a model, from which higher-order interactions are “induced” by asking the validity of (1.2) or (1.5). A “free Lagrangian” plays no role, when the free fields are constructed by second quantization of the Wigner representations of the Poincaré group. Where propagators of derivative fields are not unique, the freedom can be helpful to enforce the validity of (1.2) or (1.5).

The method is recursive in the perturbative order; for a first glimpse see the next section. However, as one proceeds to higher orders, the recursion becomes more and more intricate. In the first attempt [14] we were able to go until third order “by hand”, but there was no emergent pattern to proceed to higher orders.

It is the purpose of the present paper to formulate the conditions (1.2) resp. (1.5) at all orders at one stroke. The result is given, for each perturbative order n , as a formula for an “obstruction” (against the validity at order n when only interactions up to order $k < n$ are taken into account), that must be cancelled by interactions of order n .

This requires the obstruction computed by the formula to have a very special form. If this is not the case in a model, then (1.2) resp. (1.5) is impossible to satisfy. On the other hand, if the cancellation is possible, then we call the obstruction “resolvable”, and L_n (and K_n) the “induced” higher-order interactions.

It is, however, *not* the purpose of this paper to decide whether the obstructions are resolvable at each order. As of now, the only available method is a case-by-case investigation, see [15]. Remarkably, it is successful in sQFT for most subtheories of the SM. A few notable exceptions with non-resolvable obstructions follow a “lock-key scenario”: when a suitable other L - Q or L - V pair is added, the non-resolvable obstructions in all cross channels cancel each other. In other words: some inconsistent subtheories become consistent upon inclusion of further interactions. The latter may even require further particles.

E.g., the minimal interactions of QCD are inconsistent without the gluon self-interactions, and self-interactions of massive vector bosons are inconsistent without Higgs couplings. At second order, the known quartic interactions of the SM are induced, and the induction intrinsically terminates at third order, i.e., at the power counting bound for renormalizability.

Moreover, consistency at third order determines some numerical parameters, including chirality of the weak interaction and the precise shape of the Higgs potential, see [9, 15, 16] for more details.

Thus, the whole business in sQFT is about identifying admissible first-order interactions, *and* about determining higher-order from initial first-order interactions. The underlying mechanism is consistency with fundamental principles. Substantial parts of the SM of particle physics can be built up in this way, and the expectation is that this is true for the entire SM.

Out of the scope of the present paper are subtleties of analytic nature, that may arise especially in the actual computation of obstructions in sQFT, based on propagators (time-ordered two-point functions) of linear string-localized fields. Yet in all case studies based on subtheories of the Standard Model, these subtleties can be dealt with at least pragmatically. For a discussion, see [15]. A deeper functional-analytic investigation is certainly needed.

We do also not address loop corrections and UV renormalization, because the method to determine induced interactions proceeds already at tree-level (although we have reason to believe that Prop. 2.4 and Prop. 3.3, relying on the Master Ward Identity (MWI) in Lemma 2.1, also hold at unrenormalized loop level). The higher-order interactions thus found are then the starting point for a full loop analysis, as in all other approaches. The distinction is that sQFT allows to work with power-counting renormalizable Hilbert space interactions, even for couplings of massive vector bosons where local QFT is either non-renormalizable, or has to evade into Krein space.

1.1 A first glimpse at the recursive structure of the problem

LQ setting. Let an L - Q pair (1.3) be given. The perturbative expansion of the S-matrix up to second order is

$$S = 1 + ig \cdot \int dx L_1(x) + \frac{(ig)^2}{2} \cdot \left(\iint dx dx' T[L_1(x)L_1(x')] - i \int dx L_2(x) \right) + O(g^3).$$

When the derivation δ is applied, the first-order integral vanishes by the L - Q pair condition (1.3), while the second-order term becomes

$$\iint dx dx' (T[\partial_\mu Q_1^\mu(x)L_1(x')] + (x \leftrightarrow x')) - i \int dx \delta L_2(x). \quad (1.8)$$

The double integral does not vanish in general because time-ordering does not commute with derivatives. In this case, a suitable interaction L_2 is needed to cancel the double integral.

It is convenient to subtract from the integrand of the double integral in (1.8) the derivative term $\partial_\mu^x T[Q_1^\mu(x)L_1(x')]$. The resulting integrand is called the “second-order obstruction of the S-matrix” $O_{\text{LQ}}^{(2)}(x, x')$. The subtracted term is itself a derivative and does not contribute. Yet, the subtraction is convenient because combinations of the form

$$O_{Y(x)}(X(x')) := T[\partial_\mu^x Y^\mu(x)X(x')] - \partial_\mu^x T[Y^\mu(x)X(x')] \quad (1.9)$$

are, for local fields Y^μ and X , supported at $x = x'$, i.e., they exhibit factors $\delta_{xx'}$ or derivatives thereof. (For string-localized fields, x' may lie on the string or cone emanating from x , or vice versa, see [15, 16].) Expressions of the form (1.9) frequently appear in QFT. E.g., when $Y^\mu = j^\mu$ is the conserved Dirac current, the vanishing of $O_j(X')$ for neutral fields X is well-known as a “Ward identity”, while $O_{j(x)}(\psi(x')) = \psi(x)\delta_{xx'}$.

For Y fixed, we call $X(x') \mapsto O_{Y(x)}(X(x'))$ an “obstruction map” on the algebra of Wick polynomials. We shall often write (1.9) in short-hand notation as

$$O_Y(X') \equiv [T, \partial]YX' \equiv [T, \partial_\mu]Y^\mu(x)X(x').$$

As mentioned earlier, we shall henceforth only consider the tree-level contribution, which we shall denote by the same symbol. By Wick’s theorem, the tree-level contributions have exactly one contraction. This entails that obstruction maps are derivations w.r.t. the Wick product, and that they lower the degree of homogeneity in the free fields by 2. In particular, if Q_1 and L_1 are both cubic, then $O_{Q_1}(L_1)|^{\text{tree}}$ is quartic.

Specifically, one finds (see Sect. 3)

$$O_{\text{LQ}}^{(2)}(x, x') = O_{Q_1(x)}(L_1(x')) + O_{Q_1(x')}(L_1(x)). \quad (1.10)$$

In order for (1.8) to vanish, one must have

$$-iO_{\text{LQ}}^{(2)}(x, x') \stackrel{!}{=} \delta L_2(x) \cdot \delta_{xx'} - \mathfrak{S}_2 \partial_\mu^x Q_2^\mu(x; x'), \quad (1.11)$$

where $\mathfrak{S}_2 f(x, x') = \frac{1}{2}(f(x, x') + f(x', x))$ is the symmetrization in two variables. L_2 and Q_2 , if they exist, are determined by this cancellation condition (“resolution of the obstruction”).

It is important to keep track of the derivative term ∂Q_2 because (1.11) will be used “under the T -product” at higher orders of the recursion, where the derivative term cannot be ignored:

The expansion of (1.2) to third order will contain $T[\delta L_1(x)L_1(x')L_1(x'')]$ and $T[\delta L_1(x)L_2(x')]$, as well as $T[L_1(x)\delta L_2(x')]$. One uses (1.3) and (1.11) to write $\delta L_1(x')$ as $\partial_\mu Q_1^\mu(x')$, and $\delta L_2(x')$ as an integral over $O^{(2)}(x', x'')$ and $\partial_\mu' Q_2^\mu(x'; x'')$. As at second order, one replaces $T\partial \dots$ by $[T, \partial] \dots$ wherever it occurs. The resulting term $T[\partial Q_1 L_1(x)L_1(x')L_1(x'')] - \partial^x T[Q_1 L_1(x)L_1(x')L_1(x'')]$ is evaluated with the help of the MWI, Lemma 2.1. It turns out that the term $T[L_1(x)O^{(2)}(x', x'')]$ drops out, and one is left with the third-order obstruction expressed entirely in terms of obstruction maps:

$$\frac{(ig)^3}{6} O_{\text{LQ}}^{(3)}(x, x', x'') = -\frac{g^3}{2} \mathfrak{S}_3(O_{Q_1(x)}(L_2(x')) \cdot \delta_{x'x''} + O_{Q_2(x, x')}(L_1(x''))), \quad (1.12)$$

where $O_{Q_2(x, x')}$ is defined as in (1.9) with the derivative ∂^x acting only on x . This must be cancelled by

$$-i^2 O_{\text{LQ}}^{(3)}(x, x', x'') \stackrel{!}{=} \delta L_3(x) \cdot \delta_{xx'x''} - \mathfrak{S}_3 \partial_\mu^x Q_3^\mu(x; x', x''). \quad (1.13)$$

LV setting. The combinatorics in the LV setting is much more involved. In fact, we shall require a strengthened version of (1.5) including an arbitrary cutoff function, replacing g by $g\chi(x)$ in every integral. This is necessary because we consider the “local S-matrices” $S[\chi]$ at each order as operator-valued distributions, to be used for the process of causal renormalization as in the Epstein-Glaser framework. The obstructions of the S-matrix at each order are understood as symmetric kernels of operator-valued distributions with argument $\chi^{\otimes n}$, i.e., as the integrands multiplying $\prod_{i=1}^n \chi(x_i)$.

This prescription requires to include terms of the form “ $V \circ \partial$ ” (see (1.20) in Sect. 1.3 and [14]) in one of the two sides in the desired identity (1.5), where the derivative acts on the cutoff function. Such terms obviously vanish in the adiabatic limit $\chi(x) \rightarrow 1$. Nevertheless, we must keep track of them and integrate by parts all derivatives onto the fields, where they act outside the time-ordering.

Like the “subtraction by hand” of derivatives made in the LQ setting, this step ensures that all obstructions of the S-matrix arrange into combinations of obstruction maps. At first order,

$$S_{L_{\text{int}}+V \circ \partial}[\chi] - S_{K_{\text{int}}}[\chi] = ig \int dx \left(\chi(x) (L_1(x) - K_1(x)) + \partial_\mu \chi(x) V_1^\mu(x) \right)$$

which equals $ig \int dx \partial_\mu (\chi(x) V_1^\mu(x)) = 0$ by the L - V pair condition. The expansion of (1.5) at second order gives

$$\frac{(ig)^2}{2} \iint dx dx' \left(T[(L_1 + V_1 \partial^x)(L'_1 + V'_1 \partial^{x'})] - T[K_1 K'_1] - i((L_2 - K_2) \delta_{xx'} + V_2 \partial^x) \right) \chi(x) \chi(x').$$

The derivatives are understood to act on the test functions. Integrating them by parts (so that they act outside the T -product) and using $L_1 = K_1 + \partial V_1$, one gets

$$\begin{aligned} & S_{L_{\text{int}}+V \circ \partial}[\chi] - S_{K_{\text{int}}}[\chi] \\ &= \frac{(ig)^2}{2} \cdot \iint dx dx' \chi(x) \chi(x') \left(O^{(2)}(x, x') - i((L_2(x) - K_2(x)) \delta_{xx'} - \partial_\mu^x V_2^\mu(x; x')) \right) + O(g^3), \end{aligned} \quad (1.14)$$

where

$$O^{(2)}(x, x') = \mathfrak{S}_2(O_{V_1}(L'_1 + K'_1) - \partial_\mu^x O_{V'_1}(V_1^\mu)). \quad (1.15)$$

This second-order obstruction must be cancelled by

$$-iO^{(2)}(x, x') \stackrel{!}{=} (L_2(x) - K_2(x)) \cdot \delta_{xx'} - \mathfrak{S}_2 \partial_\mu^x V_2^\mu(x; x'). \quad (1.16)$$

L_2 , K_2 and V_2 , if they exist with the specified respective features as explained in the motivations, are determined by this resolution of the obstruction.¹

The expression for $O^{(3)}(x, x', x'')$ in terms of obstruction maps is an intricate expression. It was computed in [14] (again with the use of the MWI) under model-specific assumptions on the form of V_2^μ , and found to involve iterated obstruction maps $O_{V_1} \circ O_{V'_1}$ as well as the “nested” obstruction map $O_{O_{V_1}(V'_1)}$.

1.2 Main results

All-order obstruction formulas. It is the main aim of this paper to give expressions for the obstructions of the S-matrix at all orders in terms of obstruction maps, generalizing (1.10), (1.12), (1.15).

¹In concrete cases, the resolution may admit free parameters which are fixed at the next order.

This task is very easy in the L - Q setting, and is drastically simplified in the L - V setting (as compared to the first attempt in [14]) with an advantageous “reparametrization”. To illustrate the gist at second order: the manifest derivative term in the obstruction (1.15) is part of the derivative term in (1.16) that must resolve the obstruction. Equivalently, the obstruction is resolved by finding L_2 , K_2 and U_2 that solve the simpler equation

$$-i\mathfrak{S}_2 O_{V_1}(L'_1 + K'_1) \stackrel{!}{=} (L_2 - K_2) \cdot \delta_{xx'} - \mathfrak{S}_2 \partial_\mu^x U_2^\mu(x; x').$$

Consequently, $V_2^\mu(x; x') = U_2^\mu(x; x') - iO_{V_1}(V_1^\mu)$, and the “nested” obstruction appearing at third order will be absorbed in O_{U_2} . The same will happen recursively at all higher orders.

Apart from this simplification that will make the recursive structure far more transparent in terms of U_n rather than V_n , see Sect. 1.3, the present setup admits derivative terms with fields of the form $Q_n^\mu(x_1; x_2, \dots, x_n)$ and $V_n^\mu(x_1; x_2, \dots, x_n)$ (or U_n) that are supported at $x_i = x$ or (in the sQFT context) at x_i on the string emanating from x . It was observed in models with non-abelian self-interactions of vector bosons that such fields naturally appear. This generalization does not harm because Q_n do not appear in the S-matrix, and the contributions from V_n will disappear in the adiabatic limit.

The dressed field. A main advantage of the LV setting is that it allows to define a “dressed field”, whose role we now explain.

Recall that in the BRST approach, interacting charged fields like the Dirac field are not defined on the BRST Hilbert space. If we consider an L - V pair with the BRST interaction K_{int} on a Krein space and a string-localized interaction L_{int} on the BRST Hilbert space \mathcal{H} (which can be thought of as embedded into \mathcal{K} , see [13]), then the dressed field $\Phi|_{[g]}$ associated with a free field Φ is a power series of free fields on \mathcal{K} such that

$$\Phi|_{L_{\text{int}}}(x) = \Phi|_{K_{\text{int}}}(x) \tag{1.17}$$

holds, where $\cdot|_{L_{\text{int}}}$ stands for the perturbative interacting field obtained by Bogoliubov’s formula

$$\Phi|_{L_{\text{int}}}(x) = (T e^{i \int L_{\text{int}}})^* T[\Phi(x) e^{i \int L_{\text{int}}}] \tag{1.18}$$

The latter is expanded as a power series in retarded integrals over Wick products of fields. (Of course, (1.18) requires renormalization. As we shall not address renormalization in this paper, we claim and prove (1.17) only at tree level.)

The formula (1.17) addresses a critical issue of sQFT: if the interaction L_{int} is not local, standard arguments with Bogoliubov’s formula fail to show that the interacting field is a local field; in fact, all control of commutation relations is lost because of the support of retarded integrals over Wick products of string-localized fields.

On the other hand, because K_{int} is local, the dressed field with interaction K_{int} enjoys the same localization as the dressed field. The point is that the dressed field is a power series in the fields appearing in L_{int} , and therefore inherits its string-localization (commutativity for space-like separated strings). Thus, by virtue of (1.17), also the field with interaction L_{int} is in the worst case string-localized. If it happens to be local, then it qualifies as an observable of the theory. In either case, it is defined on \mathcal{H} . String-localized charged fields were anticipated by inspection of Gauß’ Law in QED [5], and more abstractly by the localization structure of charged superselection sectors in Algebraic QFT [1]. sQFT provides a handle to actually construct them (perturbatively).

The assignment $\Phi \mapsto \Phi_{[g]}$ is in principle recursively determined by the L - V pair. In [14] we had found the first few orders, without a compelling pattern emerging. In Sect. 4, we shall give an all-orders formula for $\Phi_{[g]}$ which is surprisingly simple thanks to the above-mentioned “reparametrization” from V to U .

1.3 Outline

We write L for L_{int} and K for K_{int} from now on.

The time-ordered exponential in (1.1) is a formal expression that needs UV renormalizations and IR regularizations. The Glaser-Epstein method rather starts from “local” S-matrices with smooth space-time cutoffs $\chi(x)$ of the coupling constant. They are regarded as symmetric distributions with tensor products of χ as test functions. Each χ is an IR regularization. On the other hand, the UV-renormalization can be reformulated as a problem of extension of distributions in position space to coinciding points where products of propagators in loop contributions are a-priori non-defined, rather than a subtraction scheme in momentum space.

Because the recursive determination of higher-order interactions is a condition for consistency with fundamental principles, we shall impose the validity of (1.2) and (1.5) only at tree level, as a *necessary* condition. In particular, we do not address UV renormalization in this paper. At tree level, the adiabatic limit $\chi(x) \rightarrow 1$ is unproblematic, and we shall suppress the cutoff function χ in the LQ setting altogether. Also, all obstruction maps are understood at tree level.

With these simplifications, we develop a method to determine $O_{\text{LQ}}^{(n)}$ at all orders (Prop. 2.4) by exploiting the property that δ is a derivation, as well as the MWI, Lemma 2.1.

In the LV setting we are more ambitious by imposing the validity of (1.5) with a cutoff function, as a prerequisite for the UV renormalization at loop level. This makes the task much more difficult. Moreover, there is no derivation δ at our disposal. As for the former complication, we shall demand the equality

$$Te^{iK[\chi]} = Te^{i(L+V \circ \partial)[\chi]}, \quad (1.19)$$

where (in distributional notation)

$$K[\chi] = \sum_{n \geq 1} \frac{g^n}{n!} K_n(\chi^n), \quad L[\chi] = \sum_{n \geq 1} \frac{g^n}{n!} L_n(\chi^n), \quad V^\mu \circ \partial[\chi] = \sum_{n \geq 1} \frac{g^n}{n!} V_n^\mu(\partial_\mu \chi \otimes \chi^{\otimes n-1}). \quad (1.20)$$

The more general form of $V_n(x_1; x_2, \dots, x_n)$ as compared to the ansatz in [14] means a gain in flexibility to satisfy (1.19), that is unavoidable in non-abelian models, and at the same time harmless because V becomes irrelevant in the adiabatic limit.

In sQFT, $K_n(x)$ and $L_n(x, c)$ are Wick polynomials in local resp. string-localized fields.

As for the absence in LV of a derivation δ , that was instrumental in LQ, we find a “substitute” in the form of an interpolation between K_{int} and L_{int} with a parameter t , such that the derivation property of d/dt can be exploited.

We shall then prove (Prop. 3.1) that along with (1.19) there also holds an interpolation

$$Te^{iK[\chi]} = Te^{i(\ell(t)+v^\mu(t) \circ \partial_\mu)[\chi]}, \quad (1.21)$$

with fields $\ell(t)$ and $v^\mu(t)$ interpolating $\ell(0) = K$ with $\ell(1) = L$, and $v^\mu(0) = 0$ with $v^\mu(1) = V^\mu$.

The proof introduces a field U^μ (another power series of Wick polynomials like V^μ in (1.20)). The point is that $v(t) = tU + \dots$ and $V = v(1) = U + \dots$ can be expressed as power series in U (involving iterated obstruction maps O_U as in (1.9)). The latter power series can be inverted for U as a power series in V (involving also “nested” obstruction maps such as $O_{O_V(V)}$). Because V is itself a power series in g , so is U , with $U_1 = V_1$.

The passage from V to U is the “change of variables” mentioned in Sect. 1.2. In particular, it will absorb all “nested” obstruction maps.

Prop. 3.3 then provides a formula (3.9) for L as a power series in the obstruction map O_U , acting on K and on $\partial_\mu U^\mu$. This formula is the main result. In fact, the field V is not needed for this formula. For the validity of (1.19), it suffices to know that V exists as a power series in U . Its precise form is uninteresting because it disappears in the adiabatic limit anyway.

The main formula (3.9) can be rewritten in several ways. Particularly useful is a rewriting in such a way (Cor. 3.4) that the obstructions of the S-matrix at each perturbative order g^n can be directly read off as iterated obstructions $O_{U_{k_i}}$ acting on L_k and K_k (with $k + \sum_i k_i = n$). This surprisingly simple formula at all orders is the main technical improvement over the “mess” in [14] already at $n = 3$.

Finally, in Sect. 4, we establish a formula for the “dressed field” at all orders.

2 The all-orders L - Q formula

Working in perturbation theory means working with formal power series in the coupling constant g . The interactions in (1.2) and (1.20) are themselves power series (and in fact in most cases polynomials) in g with leading order g . Since power series in power series in g are again power series in g , it is legitimate, and will turn out to be most advantageous, to work and display the results in terms of power series in $L_{\text{int}}(g) = \sum_{n=1}^{\infty} \frac{g^n}{n!} L_n$ and $Q^\mu = \sum_{n=1}^{\infty} \frac{g^n}{n!} Q_n$, resp. L_{int} , K_{int} and V in Sect. 3. The order-by-order decomposition into orders g^n is then a trivial step.

Moreover, we may formally include total delta functions δ_{x_1, \dots, x_n} into L_n and K_n , and suppress all arguments, integrals and (in LV) cutoff functions. This is just a device to keep the notation simple, enabling us to concentrate on the recursive structure. The arguments etc. can unequivocally re-installed at the end, with (1.12) and (1.13) as a “blueprint”: Each L_k on the right-hand side comes with an argument x and a total delta function in k arguments, and each Q_k has one distinguished argument (on which the derivative acts) and is symmetric in its remaining $k - 1$ arguments.

Let us assume that (1.2) holds. By the derivation property of δ it follows

$$iT[\delta L \cdot e^{i \int L}] = 0. \quad (2.1)$$

At each order, this amounts to the resolution of an obstruction

$$i^n O_{LQ}^{(n)} + i(\delta L_n - \partial_\mu Q_n^\mu) \stackrel{!}{=} 0, \quad (2.2)$$

where $O_{LQ}^{(n)}$ is the contribution of all L_k and Q_k with $k < n$ to the integrand of an n -fold integral, and at each order a new derivative term Q_n may be admitted, that vanishes when integrated over. The task is to compute $O_{LQ}^{(n)}$ in terms of L_k and Q_k ($k < n$), in order to determine L_n and Q_n .

Defining also

$$O_{LQ} = \sum_{n=2}^{\infty} \frac{i^{n-1} g^n}{n!} O_{LQ}^{(n)},$$

we write the resolution of obstructions (2.2) at all orders as

$$O_{LQ} + \delta L - \partial_\mu Q^\mu = 0. \quad (2.3)$$

We insert (2.3) into (2.1), and subtract a derivative term as in Sect. 1.1:

$$T[-O_{LQ}e^{iL}] + T[\partial_\mu Q^\mu e^{iL}] - \partial_\mu T[Q^\mu e^{iL}] = 0. \quad (2.4)$$

At this point, we need the MWI, which holds identically at tree level by virtue of Wick's theorem. In fact, it also holds at unrenormalized loop level and is often imposed as a renormalization condition [3]:

Lemma 2.1. (*Master Ward Identity, see [3]*). *For Wick polynomials Y and X_i , it holds*

$$\begin{aligned} O_Y(X_1, \dots, X_n) &\equiv T[\partial_\mu Y^\mu(x) \cdot X_1(x_1) \dots X_n(x_n)] - \partial_\mu^x T[Y^\mu(x) \cdot X_1(x_1) \dots X_n(x_n)] \\ &= \sum_{i=1}^n T[X_1(x_1) \dots O_{Y(x)}(X_i(x_i)) \dots X_n(x_n)]. \end{aligned} \quad (2.5)$$

In sQFT, the MWI must be extended *mutatis mutandis* to string-localized fields φ, φ' , and to multi-local fields $Y_m^\mu(y_1, \dots, y_m)$, such that the derivative is on the first argument y_1 . The former extension is unproblematic when the time-ordering is w.r.t. the ‘‘apex’’ x of the string-localized fields $\varphi(x, c)$. There are some subtle issues with the latter extension. For the time being, we assume the MWI to hold, i.e., all subtleties have been properly settled. (In all SM applications of sQFT this is the case, see the discussion in [15], and also [12].)

The ‘‘derivation-like’’ structure of (2.5) implies

Corollary 2.2. *It holds*

$$T[\partial_\mu Y^\mu(x)e^{iL}] - \partial_\mu^x T[Y^\mu(x)e^{iL}] = iT[O_Y(L)e^{iL}]. \quad (2.6)$$

Proof: Expand the exponential as a power series and apply Lemma 2.1 to each term. \square

Cor. 2.2 turns (2.4) into

$$T[-O_{LQ}e^{iL}] + iT[O_Q(L)e^{iL}] = 0. \quad (2.7)$$

Now we invoke an elementary Lemma:

Lemma 2.3. *Let $P(g) = \sum_{n \geq 0} g^n p_n$ and $E(g) = 1 + \sum_{n \geq 1} g^n e_n$ be power series in g with coefficients in some algebra with 1. If $PE = 0$, then $P = 0$.*

Proof by induction: Obviously, $PE = 0$ implies $p_0 = 0$. Assume $p_k = 0$ for all $k < n$. Then the term of PE at order n is p_n . Hence $p_n = 0$. \square

Lemma 2.3 applies to (2.7) with $E = e^{iL}$ and $P = O_{LQ} - iO_Q(L)$. Thus, we have already derived the desired all-orders result:

Proposition 2.4. (*All-orders LQ obstruction*). *If (2.1) and (2.3) hold, then*

$$O_{LQ} = iO_Q(L). \quad (2.8)$$

Expanded in g , this is

$$i^{n-2}O_{LQ}^{(n)} = \sum_{k_1+k_2=n} \frac{n!}{k_1!k_2!} O_{Q_{k_1}}(L_{k_2}), \quad (2.9)$$

where obviously all $k_1, k_2 < n$. Conversely, if (2.2) holds for each n with $O_{LQ}^{(n)}$ computed by (2.9), i.e., if all obstructions are resolved at every order, then (2.1) follows.

Proof: The first statement was just proven. The converse statement follows by doing the same steps backwards. \square

(1.12) and (1.13) and their analogues at all orders are obtained by re-installing the arguments and delta functions in (2.9) and (2.2).

3 The all-orders L - V formula

3.1 The main result

We adopt a similar efficient short-hand notation as in Sect. 2, keeping however track of the cutoff function. We think of $K_n(\chi^n)$ as $(K_n \delta_{\text{tot}})(\chi^{\otimes n})$ (and similarly with L), where δ_{tot} is the total delta function in n arguments x_1, \dots, x_n . We suppress writing δ_{tot} , however. We write $K[\chi]$ and $L[\chi]$ for the power series of distributions evaluated on $\chi^{\otimes n}$.

When V_n appears inside a time-ordered product with test function $\partial\chi \otimes \chi^{\otimes(n-1)}$, one can integrate the derivative by parts so that it acts on the time-ordered product of fields. Then, all test functions are just tensor powers of χ , which means that the corresponding distributional kernels are completely symmetric. In this sense, we also write $V \circ \partial[\chi]$.

Proposition 3.1. (*Interpolation*). *Suppose that (1.19) holds:*

$$Te^{iK[\chi]} = Te^{i(L+V^\mu \circ \partial_\mu)[\chi]} \quad (3.1)$$

with power series (in g) of Wick polynomials L, K, V^μ as in (1.20). Then there exist power series of Wick polynomials $\ell(t)$ and $v^\mu(t)$, such that also (1.21) holds:

$$Te^{iK[\chi]} = Te^{i(\ell(t)+v^\mu(t)\partial_\mu)[\chi]}, \quad (3.2)$$

and $\ell(t)$ and $v^\mu(t)$ interpolate $\ell(0) = K$ with $\ell(1) = L$, and $v^\mu(0) = 0$ with $v^\mu(1) = V^\mu$.

For the proof, we shall need another Lemma, which states the solution of a simple inhomogeneous linear differential equation. It is elementary to prove:

Lemma 3.2. *Let M be a linear map on some vector space S . Then the inhomogeneous initial value problem for a function $s(t) \in S$*

$$\dot{s}(t) - M(s(t)) = a \quad \text{with} \quad s(0) = s_0 \quad (3.3)$$

is solved (as a power series in t) by

$$s(t) = \exp(tM)(s_0) + F(tM)(ta), \quad \text{where} \quad F(u) := \frac{e^u - 1}{u} = \sum_{r=0}^{\infty} \frac{u^r}{(r+1)!}. \quad (3.4)$$

Proof of Prop. 3.1: $Te^{i(\ell(t)+v(t)\partial)[\chi]}$ (with $\ell(t)$ and $v(t)$ still to be determined) is independent of t if

$$T[\dot{\ell}(t)e^{i(\ell(t)+v(t)\partial)[\chi]}] = \partial_\mu T[v^\mu(t)e^{i(\ell(t)+v(t)\partial)[\chi]}]. \quad (3.5)$$

On the other hand, Cor. 2.2 with L replaced by $(\ell(t) + v(t)\partial)[\chi]$ yields (for any U)

$$[T, \partial_\mu]U^\mu e^{i(\ell(t)+v(t)\partial)[\chi]} = iT[O_U(\ell(t))e^{i(\ell(t)+v(t)\partial)[\chi]}] - \partial_\mu T[O_U(v^\mu(t))e^{i(\ell(t)+v(t)\partial)[\chi]}],$$

equivalently

$$T[(\partial_\mu U^\mu - iO_U(\ell(t)))e^{i(\ell(t)+v(t)\partial)[x]}] = \partial_\mu T[(U^\mu - iO_U(v^\mu(t)))e^{i(\ell(t)+v(t)\partial)[x]}].$$

By comparison with (3.5), we therefore need to specify U such that

$$\dot{\ell}(t) = \partial_\mu U^\mu - iO_U(\ell(t)) \quad \text{and} \quad \dot{v}^\mu(t) = U^\mu - iO_U(v^\mu(t)), \quad (3.6)$$

ensuring that the right-hand side of (3.2) is independent of t . Its value at $t = 0$ is the left-hand side provided $\ell(0) = K$ and $v^\mu(0) = 0$.

Lemma 3.2 gives the solutions $\ell(t)$ and $v(t)$ of the initial value problems (3.6) in terms of the (yet unknown) U^μ :

$$\ell(t) = \exp(-itO_U)(K) + F(-itO_U)(t\partial_\mu U^\mu), \quad v^\mu(t) = F(-itO_U)(tU^\mu). \quad (3.7)$$

$F(u) = \frac{e^u - 1}{u}$ is the function given in Lemma 3.2, understood as a power series.

Now, (3.7) determines $v^\mu(1)$ as a power series in obstruction maps O_U acting on U^μ . The desired value $v^\mu(1) = V^\mu$ is obtained if

$$V^\mu = F(-iO_U)(U^\mu) = U^\mu - \frac{i}{2}O_U(U^\mu) - \frac{1}{6}O_U^2(U^\mu) + \dots \quad (3.8)$$

This power series can be inverted:

$$U^\mu = V^\mu + \frac{i}{2}O_V(V^\mu) - \frac{1}{4}O_{O_V(V)}(V^\mu) - \frac{1}{12}O_V^2(V^\mu) + \dots,$$

and therefore determines U in terms of V . Since V is itself a power series in the coupling constant g , so is U . Thus, we have established (3.2), but we still have to show that $\ell(1) = L$.

For this purpose, it suffices to notice that when K and V in (3.1) are given, then L is unique if it exists. Namely, at each perturbative order the right-hand side of (3.1) is a sum of L_n and terms involving only V_k , and L_k with $k < n$, which fixes L_n recursively. Therefore, since (3.7) at $t = 1$ is (3.1) with $\ell(1)$ in the place of L , it follows that $\ell(1) = L$. \square

Proposition 3.3. (“LV identity”). (i) *The identity (3.1) holds if and only if*

$$L = \exp(-iO_U)(K) + F(-iO_U)(\partial U), \quad (3.9)$$

with $U = U(V)$ given by the inverse of the power series $V(U) = F(-iO_U)(U)$ as in (3.8), where $F(u) = \frac{e^u - 1}{u}$ as in Lemma 3.2.

(ii) *It also holds*

$$K = \exp(iO_U)(L) - F(iO_U)(\partial U), \quad (3.10)$$

exhibiting a perfect symmetry under $L \leftrightarrow K$ and $U \leftrightarrow -U$. In particular, (3.1) is also equivalent to

$$Te^{iL[x]} = Te^{i(K - \tilde{V} \circ \partial)[x]}$$

with $\tilde{V} = F(+iO_U)(U)$.

Proof: The first formula (3.9) is just the evaluation of the solution $\ell(t)$ in (3.7) at $t = 1$. Therefore, (3.1) implies (3.9) by Prop. 3.1. Conversely, the proof of Prop. 3.1 shows that (3.2) with $\ell(t)$ and $v(t)$ given by (3.7) is independent of t . Putting $t = 1$, one gets (3.1).

For the second formula (3.10), we write $F(u) = \int_0^1 dt e^{tu}$ in (3.9), apply e^{iO_U} on both sides, and substitute $1 - t \rightarrow t$ in the integral. \square

One can solve (3.9) for ∂U and then rewrite it in a way such that the obstruction maps act only on the interactions L and K :

Corollary 3.4. (All-orders LV obstruction). With $G(u)$ the even power series $G(u) = 1 - \frac{1}{2} \cot(\frac{1}{2}u) = \frac{1}{12}u^2 + \frac{1}{360}u^4 + \dots$, it holds

$$L - K - \partial_\mu U^\mu = G(O_U)(L - K) - \frac{i}{2}O_U(L + K). \quad (3.11)$$

Proof: Apply $\exp(\frac{i}{2}O_U)$ to (3.9), and solve the resulting

$$\exp(\frac{i}{2}O_U)(L) - \exp(-\frac{i}{2}O_U)(K) = iH(\frac{1}{2}O_U)(\partial U), \quad \text{where} \quad H(u) := \frac{\sin(u)}{u}$$

for ∂U by using that the inverse of a power series $1 + \dots$ is the reciprocal power series. The reason why $L - K$ is added on both sides will be clear from Rem. 3.5.(ii) and (iii). \square

Remarks 3.5. (i) Cor. 3.4 manifestly exhibits the symmetry mentioned in Prop. 3.3.

(ii) Cor. 3.4 is used to recursively determine the higher-order interactions L_n and K_n along with U_n^μ . By expanding the right-hand side to order g^n , one obtains the distributional kernel of the n -th order obstruction of the S -matrix, to be evaluated on $\chi^{\otimes n}$. It arises in terms of $O_{U_{k_i}}$ acting on $L_k \pm K_k$, with $k + \sum_{i=1}^r k_i = n$ (in particular only $k < n$ and $k_i < n$ appear). Remarkably, the odd part in $O_{U_{k_i}}$ is in fact linear ($r = 1$). At lowest orders:

$$\begin{aligned} L_2 - K_2 - \partial U_2 &= -i O_{U_1}(L_1 + K_1), \\ L_3 - K_3 - \partial U_3 &= \frac{1}{2} O_{U_1}^2(L_1 - K_1) - \frac{3i}{2} (O_{U_1}(L_2 + K_2) + O_{U_2}(L_1 + K_1)), \\ L_4 - K_4 - \partial U_4 &= O_{U_1}^2(L_2 - K_2) + (O_{U_1}O_{U_2} + O_{U_2}O_{U_1})(L_1 - K_1) \\ &\quad - 2i O_{U_1}(L_3 + K_3) - 3i O_{U_2}(L_2 + K_2) - 2i O_{U_3}(L_1 + K_1). \end{aligned} \quad (3.12)$$

The symmetric re-installment of arguments x_1, \dots, x_n and implicit delta functions that are suppressed in the present simplified notation, is unequivocal.

(iii) The right-hand sides of (3.12) are the n -th order obstructions of the S -matrix remaining after the passage from V to U . As explained in Sect. 1.2, one must compute the obstructions, and resolve them by the quantities on the left-hand sides, in order to determine the induced higher-order interactions L_n and K_n .

(iv) When comparing (3.12) with the second and third-order formulas given in [14], one should observe, apart from the reparametrization from V to U , that in [14] the present $V(x; x')$ was expanded into delta functions and their derivatives (where special feature of the model were exploited). Moreover, a major distinction is that in (3.12), the obstruction maps act only on the interactions L_k and K_k , thanks to the elimination of ∂U in Cor. 3.4. This further simplifies actual computations.

(v) As emphasized in Sect. 1.2: We did not prove that the obstructions on the right-hand sides of (3.12) are of the form that can be resolved at all orders. (3.11) only provides a simple formula for the combinatorics on the right-hand side, needed to compute the obstructions at order g^n , once the obstruction of orders $k < n$ have been resolved. Prop. 3.3 is also true without special features of K_n and L_n (like factoring a total delta function or being local or string-localized), that were only subsidiary conditions (“resolvability”) necessary to interpret $K_n(x)$ and $L_n(x)$ as interaction densities.

4 Interacting quantum fields

Interacting fields as operator-valued distributions are defined perturbatively by Bogoliubov's formula

$$\Phi|_L(f) = \int dx f(x) \Phi|_L(x) := -i \left(T e^{i \int dx \chi(x) L(x)} \right)^* \partial_s \left(T e^{i \int dx (\chi(x) L(x) + f(x) \Phi(x))} \right) \Big|_{s=0} \quad (4.1)$$

in the adiabatic limit $\chi(x) \rightarrow 1$.

The same formula is employed in sQFT, but if L is string-localized, it does not allow to assess the localization properties (in the sense of commutation relations) of the interacting fields $\Phi|_L$ relative to each other. We have discussed in Sect. 1.2 how this can be achieved with the help of the dressed field.

4.1 The dressed field

For a free Wick polynomial $\Phi(x)$, we define the dressed field $\Phi_{[g\chi]}(x)$ such that the “magic formula” holds for the interacting fields with a cutoff function:

$$\Phi|_{L[\chi]}(f) = \Phi_{[g\chi]}|_{K[\chi]}(f). \quad (4.2)$$

The task is to give a formula for $\Phi_{[g\chi]}$ such that (4.2) holds. Setting $\chi = 1$, gives (1.17).

Proposition 4.1. (*Dressed field*). *With a cutoff function χ , the dressed field is a power series in $g\chi$, given by*

$$\Phi_{[g\chi]}(f) = \exp(iO_U[\chi])(\Phi(f)), \quad (4.3)$$

with the obstruction map O_U as in Sect. 3 understood, at each order n , as a distribution in x_1, \dots, x_n .

Proof: Assume that (3.1) holds without the insertion. Then, by (4.1), we have to determine the dressed field such that

$$\partial_s T e^{i(K[\chi] + s\Phi_{[g\chi]}(f))} \Big|_{s=0} = \partial_s T e^{i(L + V^{\mu\circ\partial_\mu}[\chi] + s\Phi(f))} \Big|_{s=0} \quad (4.4)$$

holds. Let $K_s = K + s\Phi_{[g\chi]}(f)$. Then, by Prop. 3.3, (3.1) holds with K and L replaced by K_s and L_s where L_s is computed with (3.9):

$$L_s = \exp(-iO_U)(K_s) + F(-iO_U)(\partial U) = L + s \exp(-iO_U[\chi])(\Phi_{[g\chi]}(f)).$$

This argument is legitimate because no specific properties of the power series K and L were assumed in the proof of the proposition, see Rem. 3.5.(v). Thus, Prop. 3.3 gives (4.4) (without the derivative w.r.t. s), provided $\exp(-iO_U[\chi])(\Phi_{[g\chi]}(f)) = \Phi(f)$. Applying $\exp(iO_U[\chi])$, we get (4.3). \square

In sQFT, due to the obstruction maps involving string-localized fields U , the dressed field $\Phi_{[g\chi]}(x)$ will be a perturbative series in Wick products of $g^n \Phi_{[n]}(x; x_1, \dots, x_n)$ evaluated on $\chi^{\otimes n}$, where $\Phi_{[n]}(x; x_1, \dots, x_n)$ are supported at x_i lying on the string or cone emanating from x or from x_j . In particular, if the cone is convex, then $\Phi_{[n]}$ are string-localized.

In the adiabatic limit $\chi \rightarrow 1$, one has to integrate over all x_i . The dressed field then becomes a power series in g whose coefficients may be string-integrals over Wick products of (string-integrals over Wick products of ...) string-localized free fields.

We illustrate this by the example of dressed Dirac fields. In QED one finds [13]

$$\psi_{[g]}(x) = :e^{ig\phi(x,c)}: \cdot \psi(x), \quad (4.5)$$

where $\phi(x, c)$ is a smeared integral over the gauge potential along a string from x to infinity (i.e., a smeared abelian Wilson operator). In this case, the exponential is just a Wick power series in string-localized free fields², which is itself a string-localized field. (In fact, due to the infrared divergence of the field ϕ , *only* its exponential can be defined and turns $\psi_{[g]}$ into an “infrafield”, creating states on which the mass operator has no point spectrum. This feature reflects the “photon cloud” accompanying a charged particle.)

The dressed quark triplet field in QCD is more complicated [15]. We have computed it until third order, finding that the exponential in (4.5) is replaced by a power series in string-integrals over Wick products of (string-integrals over Wick products of . . .) string-localized Lie-algebra valued free fields. The nested structure of these integrals precisely amounts to path-ordering in the case when the string is just a line from x to ∞ . In the smeared case, the exponential still enjoys properties of a “smeared Wilson operator”³; in particular, the dressed field is invariant when classical gauge transformations that are trivial at infinity act on the free quantum gauge potential and Dirac field it is made of.

4.2 The algebra property

Of prominent interest is the case when $\Phi_{[g]}(x)$ is a power series of Wick products of local fields $\varphi(x)$, i.e., $\Phi_{[g]}$ is local relative to other free fields. This will not be the case in general.

Free fields Φ for which $\Phi_{[g]}$ is a local free field (order by order), are called “seeds” for local interacting fields, see [14, Sect. 3.6]. Namely, because Bogoliubov’s formula with a local interaction preserves the relative localization, the corresponding interacting fields in the adiabatic limit $\Phi|_L = \Phi_{[g]}|_K$ are local relative to each other and to other interacting fields.

The question was raised in [14] whether the seeds form an algebra under the Wick product. Thanks to Prop. 4.1, the answer is affirmative.

Proposition 4.2. (*Algebra property of seeds*). *The seeds of local interacting fields form an algebra under the Wick product.*

Proof: Φ_i ($i = 1, 2$) are seeds if and only if all $\Phi_{i,[n]}$ are local. By Prop. 4.1, this is the case if and only if all $U_U^r(\Phi_i)$ are local. Because obstruction maps O_Y are derivations w.r.t. the Wick product (see Sect. 1.1), it follows

$$O_U^r(: \Phi_1 \Phi_2 :) = \sum_{r_1+r_2=r} \frac{r!}{r_1!r_2!} : O_U^{r_1}(\Phi_1) O_U^{r_2}(\Phi_2) : .$$

It follows that $: \Phi_1 \Phi_2 :$ is a seed if Φ_i are seeds. This proves the proposition. \square

²Quantities of the form (4.5) were previously considered by many authors [2, 10, 11] as *classical* expressions with the motive to quantize only gauge invariant quantities. Steinmann’s idea [18] to formulate perturbation theory as a perturbation of a free *quantum* field (4.5) as a starting point is very close to ours in (4.2), with the distinction that Steinmann *chose* (4.5), while sQFT *derives* it.

³We are not aware of an independent notion of “path-ordering” for exponentials of smeared line integrals.

5 Outlook

We have reported substantial progress on the combinatorial structure of the recursive scheme to control the effect of “adding derivatives to the interaction” (and eliminate it by resolving obstructions).

Our main motivation is string-localized QFT, which allows to reformulate (and in fact predict) Standard Model interactions in an autonomous way (intrinsically quantum, referring neither to canonical quantization nor gauge or BRST invariance) in the “LQ setting” where the string-variation is given by derivatives, and to show equivalence with gauge theory approaches by means of the “LV setting” which allows to compare S-matrices obtained with interactions that differ by a total derivative at first order.

Yet, the results found are independent of the special application to sQFT, and may as well be used to compare, say, different local methods to deal with interactions of massive vector bosons. (An example is given in [14, Sect. 4.5].)

An interesting question that could be addressed with the present methods is what makes the recursive LQ or LV scheme terminate (at least at tree level). Empirically, the sequence $(L_n)_n$ terminates at L_2 whenever L_1 is power-counting renormalizable, and the same is true in the LV setting for $(K_n)_n$. But there exist cases [14] where L_1 is renormalizable while K_1 is not, and then $(L_n)_n$ terminates and $(K_n)_n$ does not. The example of perturbative gravity, treated in PGI and in sQFT in the LQ setting [4, 8], has L_1 non-renormalizable and $(L_n)_n$ does not terminate. Superficially thinking, the latter is not a surprise since also the expansion of the Einstein-Hilbert Lagrangian doesn't terminate. But the Einstein-Hilbert Lagrangian is not an input of the approach; and more generally “renormalizability” never explicitly enters in the present formalism. So what is the reason for these surprising findings?

We also do not know how to pass from LV to LQ. At first order this is trivial because each L - V pair with $\delta K_1 = 0$ gives rise to an L - Q pair by $Q_1^\mu = \delta V_1^\mu$. At higher orders the connection is less clear because δ acts both on L and on V (or U). The expectation remains that “LQ is some kind of infinitesimal version of LV”.

Finally, there is the “Elephant in the Room” question: Why do gauge interactions so swiftly give rise to L - Q pairs and L - V pairs, and, more importantly, why do all such pairs that are found without assuming gauge invariance (especially in the autonomous LQ setting) induce higher-order gauge interactions? Beyond the obvious pattern seen in (1.4) and (1.7), there is presently no deeper understanding for these remarkable facts that could shed light on the “ontological” question: Why is gauge theory so successful, despite its many unphysical features?

References

- [1] D. Buchholz, K. Fredenhagen: Locality and the structure of particle states, *Commun. Math. Phys.* 84 (1982) 1–54.
- [2] P.A.M. Dirac: Gauge-invariant formulation of Quantum Electrodynamics, *Canadian J. Phys.* 33 (1955) 650–660.
- [3] M. Dütsch: *From Classical Field Theory to Perturbative Quantum Field Theory*, Birkhäuser 2019.
- [4] M. Dütsch: Proof of perturbative gauge invariance for tree diagrams to all orders, *Ann. Phys. (Leipzig)* 14 (2005) 438–461.

- [5] R. Ferrari, L.E. Picasso, F. Strocchi: Some remarks on local operators in Quantum Electrodynamics, *Commun. Math. Phys.* 35 (1974) 25–38.
- [6] C. Gass: Renormalization in string-localized field theories: a microlocal analysis, *Ann. H. Poinc.* 23 (2022) 3493–3523.
- [7] C. Gass, J.M. Gracia-Bondía, J. Mund: Revisiting the Okubo–Marshak argument, *Symmetry* 13 (2021) 1645.
- [8] C. Gass, J.M. Gracia-Bondía, K.-H. Rehren: Quantum general covariance. *Class. Qu. Grav.* 40 (2023) 195016.
- [9] J.M. Gracia-Bondía, J. Mund, J.C. Várilly: The chirality theorem. *Ann. Henri Poincaré* 19 (2018) 843–874.
- [10] P. Jordan: Zur Quantenelektrodynamik, *Zeitschr. Phys.* 95 (1935) 202–209.
- [11] S. Mandelstam: Quantum electrodynamics without potentials, *Ann. Phys.* 19 (1962) 1–24.
- [12] N. Münch: Bachelor’s thesis, Göttingen 2024.
- [13] J. Mund, K.-H. Rehren, B. Schroer: Infraparticle quantum fields and the formation of photon clouds, *JHEP* 04 (2022) 083.
- [14] J. Mund, K.-H. Rehren, B. Schroer: How the Higgs potential got its shape, *Nucl. Phys. B* 987 (2023) 116109.
- [15] K.-H. Rehren: Nonabelian models in sQFT. arXiv:2405.nnnnn.
- [16] K.-H. Rehren, L.T. Cardoso, C. Gass, J.M. Gracia-Bondía, B. Schroer, J.C. Várilly: sQFT – an autonomous explanation of the interactions of quantum particles. arXiv:2405.nnnnn.
- [17] G. Scharf: *Quantum Gauge Theories: A true ghost story.* Wiley 2001.
- [18] O. Steinmann: Perturbative QED in terms of gauge invariant fields, *Ann. Phys.* 157 (1984) 232–254.