

Note on the Taub-NUT Instanton Metric

Joon-Hwi Kim¹

¹Walter Burke Institute for Theoretical Physics, California Institute of Technology, Pasadena, CA 91125

It is shown that a complex coordinate transformation maps the Taub-NUT instanton metric to a Kerr-Schild metric. This metric involves a semi-infinite line defect as a gravitational analog of the Dirac string, much like the original metric. Also, it corresponds to self-dual dyon in electromagnetism through three versions of classical double copy, one of them being a novel nonlocal operator form. The relevance to Newman-Janis algorithm is briefly noted.

Introduction.—Recently, an exciting program dubbed *classical double copy* [1–17] has refined analogies between gravity and electromagnetism into concrete mappings between classical solutions while being rooted in the study of scattering amplitudes [17–21]. The prime example is the *Kerr-Schild classical double copy*, established in the seminal work [1]. Namely, it is shown that a stationary Kerr-Schild solution $g = \eta + \Phi \ell \otimes \ell$ in general relativity maps to a stationary solution in Maxwell theory described by the gauge potential $A = \Phi \ell$, by taking a “single copy” of the null one-form ℓ . Conversely, one envisions a gravitational counterpart $g = \eta + \Phi \ell \otimes \ell$ of a stationary solution $A = \Phi \ell$ in electromagnetism, by taking a “double copy” of ℓ . Well-known instances of this correspondence are Schwarzschild to point charge and Kerr to a rotating solution dubbed $\sqrt{\text{Kerr}}$ [1, 22].

From this angle, it is natural to expect that the Taub-NUT solution [23, 24] will correspond to a dyon, given the interpretation of the NUT charge as the gravitomagnetic monopole [16, 25–31]. However, only a “double Kerr-Schild” metric $g = \eta + \Phi_1 \ell_1 \otimes \ell_1 + \Phi_2 \ell_2 \otimes \ell_2$ has been explicitly known for Taub-NUT [32, 33]. Thus, a mapping to electromagnetic dyons had to be achieved by a *variant* of the Kerr-Schild double copy [6, 16], which yet offered an intriguing new perspective.

Remarkably, in the present note we point out that the straightforward application of the *original* Kerr-Schild double copy is in fact possible for the *self-dual* Taub-NUT (SDTN) solution, which is an extremal case of the Taub-NUT solution that is gaining a revived interest [34–38] due to its intriguing physical properties [39, 40]. The crucial fact is that the SDTN solution remarkably admits a Kerr-Schild metric in the ordinary sense: $g = \eta + \Phi \ell \otimes \ell$.

Firstly, a Kerr-Schild metric is obtained by Kerr-Schild double copying the gauge potential of a self-dual dyon (SDD) in electromagnetism. Then we show that it describes the SDTN solution by explicitly constructing a coordinate transformation to the instanton metric of Gibbons and Hawking [41–45]. Curiously, this change of coordinates turns out to be inherently complex-geometrical in Lorentzian signature, which is consistent with the identification of the SDTN solution as a complex saddle.

Secondly, the physical interpretation of the solution as a gravitational dyon is established within the Kerr-Schild description. Most importantly, we confirm the characteristic *Misner string* geometry—the gravitational analog [25, 26] of the Dirac string [46, 47]. Notably, finding its

distributional source [26, 27] becomes a simple problem in Kerr-Schild coordinates, thanks to a linearization [48].

Thirdly, the classical double copy is established in two more ways. We first show that the Weyl double copy relations given by Refs. [2–4], established directly in terms of curvature tensors without referencing an explicit gauge potential nor metric, are concretely reproduced in our explicit Kerr-Schild description. This check clarifies a subtlety in the previous constructions [3, 4] and also illustrates that our Kerr-Schild coordinates are secretly the “Cartesian” coordinates used in the twistor theory context [3, 4, 34, 49–55]. Next, we propose an exotic *new* form of classical double copy employing a *nonlocal* operator, which implements the Dirac strings and traces back to propagators between dyonic sources [56–60].

Lastly, we briefly remark on how we originally arrived at this metric, which was by deciphering the Newman-Janis shift—the magical property of spinning black holes noticed in the curious “Note on the Kerr Spinning Particle Metric” [61].

Derivation of SDTN metric from double copy.—The gauge potential of a dyon carrying electric and magnetic charges $(q, -iq)$ is given by

$$A = \frac{q}{4\pi r} \left(-dt - ir(1 - \cos\theta) d\phi \right), \quad (1)$$

where we have set up spherical coordinates and oriented the Dirac string [46, 47] along the negative z -axis (south pole). This describes a complexified solution where the field strength $F = dA$ is self-dual: $*F = +iF$ [62].

Now adding a total derivative $(q/4\pi) d \log(r+z)$ gauge transforms this to a Kerr-Schild form:

$$A = \frac{q}{4\pi r} \left(-dt + \frac{x+iy}{r+z} dx + \frac{y-ix}{r+z} dy + dz \right), \quad (2)$$

where $x = r \sin\theta \cos\phi$, $y = r \sin\theta \sin\phi$, $z = r \cos\theta$ are the Cartesian coordinates. Importantly, the terms inside the bracket comprise a null one-form.

With this understanding, the Kerr-Schild classical double copy [1] maps Eq. (2) to the following line element, by literally taking “two copies” of the null one-form:

$$ds^2 = -dt^2 + dx^2 + dy^2 + dz^2 + \frac{m}{4\pi r} \left(-dt + \frac{x+iy}{r+z} dx + \frac{y-ix}{r+z} dy + dz \right)^2. \quad (3)$$

Here we have adopted $8\pi G = 1$ units and replaced the charge q with the mass m . Remarkably, Eq. (3) is a metric of the SDTN solution whose mass and NUT charge are m and $-im$, where complexified vacuum Einstein's equations are satisfied off the negative z -axis.

The diffeomorphism.—To show this, we simply construct an explicit coordinate transformation to the known metric of the SDTN solution. It is convenient to employ complex coordinates $\zeta = (x + iy)/(r + z)$, $\tilde{\zeta} = (x - iy)/(r + z)$ that arise from stereographically projecting the two-sphere with respect to the south pole so that Eq. (3) translates to

$$ds^2 = -dt^2 + dr^2 + r^2 \frac{4d\tilde{\zeta}d\zeta}{(1 + \tilde{\zeta}\zeta)^2} + \frac{m}{4\pi r} \left(-dt + dr - r \frac{2\tilde{\zeta}d\zeta}{1 + \tilde{\zeta}\zeta} \right)^2. \quad (4)$$

Amusingly, applying a diffeomorphism given by

$$t \mapsto t, \quad r \mapsto r + \frac{m}{4\pi}, \quad (5)$$

$$\zeta \mapsto \zeta, \quad \tilde{\zeta} \mapsto \frac{1}{1 + (m/4\pi r)(1 + \tilde{\zeta}\zeta)} \tilde{\zeta},$$

we obtain

$$ds^2 = - \left(1 + \frac{m}{4\pi r} \right)^{-1} \left(dt + \frac{m}{4\pi r} \left(dr - r \frac{2\tilde{\zeta}d\zeta}{1 + \tilde{\zeta}\zeta} \right) \right)^2 + \left(1 + \frac{m}{4\pi r} \right) \left(dr^2 + r^2 \frac{4d\tilde{\zeta}d\zeta}{(1 + \tilde{\zeta}\zeta)^2} \right), \quad (6)$$

which is the Gibbons-Hawking metric [41, 42] [63] of the SDTN solution with mass and NUT charge m and $-im$!

This diffeomorphism is surely invertible. In fact, as is elaborated on in the appendix, it turns out that Eq. (5) traces back to the flow along the null geodesic congruence associated with the Kerr-Schild metric. As a result, the inverse map is simply Eq. (5) with m replaced with $-m$.

This completes our proof that the Kerr-Schild and Gibbons-Hawking metrics in Eqs. (3) and (6) are diffeomorphic. In turn, we establish that the SDTN solution admits a construction from the Kerr-Schild classical double copy. Previously, these facts had been only *conjectured* by Ref. [64] in split signature: see the note added.

Meanwhile, the astute reader may have noticed an unusual feature of the diffeomorphism in Eq. (5). Namely, it describes a genuine *complexified diffeomorphism*, since ζ is frozen while $\tilde{\zeta}$ gets scrambled; hence the notation $\tilde{\zeta}$ instead of $\bar{\zeta}$. However crucially, this is a legitimate operation because the Lorentzian signature SDTN solution is inherently a complexified solution as a self-dual solution. In fact, the SDD gauge potential described in Eq. (2) is a solution to complexified Maxwell's equations, due to the imaginary magnetic charge $-iq$. In the precise same sense, the SDTN spacetime is a solution to “complex general relativity” as a holomorphic field theory governing

holomorphic metrics of complex four-manifolds [54, 65–67] (i.e., belongs to the “holomorphic, complexified category” [34]). Especially, the imaginary NUT charge $-im$ essentially makes the metric complex-valued. As a consequence of this interpretation, ζ and $\tilde{\zeta}$ can be transformed independently as *holomorphic* coordinates.

Physical interpretation.—We shall now examine the physical validity of the Kerr-Schild metric in Eq. (3). To begin with, we point out that Eq. (3) *itself* is unusual as a Kerr-Schild metric, in fact. While the Kerr-Schild potential is simply $\Phi = m/4\pi r$, the null one-form ℓ involves a *semi-infinite line defect* along the negative z -axis, which is exactly how the gravitomagnetic charge is encoded:

$$\ell = -dt + \frac{x + iy}{r + z} dx + \frac{y - ix}{r + z} dy + dz. \quad (7)$$

This is intriguing because the null one-form is usually regular wherever the Kerr-Schild potential is regular, which is especially true for the double Kerr-Schild construction of Taub-NUT in Ref. [6].

Crucially, as in the original metric in Eq. (6), the interpretation of this line defect should be the *Misner string* [25, 26]: the gravitational analog of the Dirac string, in-putting a gravitomagnetic flux of $-im$ at the origin. To show this, we recall that there are at least three senses in which the Taub-NUT metric [25, 26] is a gravitational analog of a dyon: (a) the Misner string is invisible as a coordinate artifact if the gravitational analog of the Dirac quantization condition holds [25], (b) the Misner string implements a flux tube of time monodromy [16, 28, 29], and (c) the distributional stress-energy of the string describes an energy-less thin solenoid of mass current [26, 27]. Our goal now is to show that all of these properties hold in the Kerr-Schild description as well.

First, we derive the Dirac quantization condition. As a primer, consider the SDD gauge potential in Eq. (2). Adding a term $(q/2\pi)d \log \zeta$ repositions its Dirac string along the positive z -axis (north pole). This encodes the complexified group-valued transition function $\zeta^{ieq/2\pi}$, where e is the electromagnetic coupling [16]. Demanding its single-valuedness within the overlap of the two patches covering either north or south poles [68] implies $ieq \in 2\pi\mathbb{Z}$, which is precisely the Dirac quantization condition from the imaginary magnetic charge $-iq$.

The gravitational Dirac quantization has been derived in a similar fashion, namely by hiding the string with overlapping coordinate patches [25]. In the Kerr-Schild description, we find that the diffeomorphism

$$t \mapsto t + \frac{m}{2\pi} \log \zeta, \quad \tilde{\zeta} \mapsto \frac{1 - (m/4\pi r)(1 + \tilde{\zeta}\zeta)/\tilde{\zeta}\zeta}{1 + (m/4\pi r)(1 + \tilde{\zeta}\zeta)} \tilde{\zeta} \quad (8)$$

with r and ζ fixed transforms Eq. (4) to

$$ds^2 = -dt^2 + dr^2 + r^2 \frac{4d\tilde{\zeta}d\zeta}{(1 + \tilde{\zeta}\zeta)^2} + \frac{m}{4\pi r} \left(-dt + dr + r \frac{2d\zeta}{\zeta(1 + \tilde{\zeta}\zeta)} \right)^2, \quad (9)$$

which develops a string defect along the positive z -axis (north pole) [69]. For this diffeomorphism to be single-valued within the overlapping region, the time coordinate should be periodic as $t \sim t + im\mathbb{Z}$, given the $2\pi i$ periodicity of $\log \zeta$. As a result, the energy e [70] of a test particle is quantized as $iem \in 2\pi\mathbb{Z}$. This is precisely the gravitational analog of the Dirac quantization condition in the presence of the imaginary NUT charge $-im$ [16, 28, 29].

Second, we study the near-string geometries in an explicit manner. Let $\xi := x + iy$, $\bar{\xi} := x - iy$ be complex coordinates for the (complexified) x - y plane, and suppose a positive infinitesimal ϵ . Applying the replacement $z \mapsto -1/\epsilon + z$ to the gauge potential in Eq. (2) and then taking the limit $\epsilon \rightarrow 0$, we obtain $A = (q/2\pi)d \log \bar{\xi}$. This precisely describes an infinite Dirac string as a thin tube of magnetic flux iq flowing in the $-z$ direction, which should be clear from its field strength $F = dA$:

$$F = \frac{q}{2\pi} dd \log \bar{\xi} = iq \delta^{(2)}(\bar{\xi}, \bar{\xi}) d\bar{\xi} \wedge d\bar{\xi}. \quad (10)$$

Here $\delta^{(2)}(\bar{\xi}, \bar{\xi}) = (1/2\pi i) \partial(1/\bar{\xi})/\partial\bar{\xi}$ describes the Dirac delta distribution peaked at $\bar{\xi} = 0$ [71].

For the line element in Eq. (3), the same limiting procedure leaves out a divergent part, which interestingly can be compensated by a shift $\xi \mapsto \xi + ((m/\pi)(1/\epsilon - z) - 3m^2/4\pi^2)/\xi$. Amusingly, the metric then becomes simply Minkowski, $ds^2 = -dt'^2 + dz'^2 + d\bar{\xi}d\xi$, but with multi-valued time:

$$t' = t + \frac{m}{2\pi} \log \bar{\xi}, \quad z' = z - \frac{m}{\pi} \log \bar{\xi}. \quad (11)$$

In other words, in the vicinity of the string the geometry approaches to the image of empty spacetime upon a periodically multivalued diffeomorphism (large gauge transformation [72, 73]). In this geometry, a Sagnac interferometer measures the “time translation monodromy” $\oint dt' = im$ as the gravitational Aharonov-Bohm phase [29, 30, 74, 75] [76]. Clearly, the flux of this monodromy is localized along $\bar{\xi} = 0$:

$$ddt' = \frac{m}{2\pi} dd \log \bar{\xi} = im \delta^{(2)}(\bar{\xi}, \bar{\xi}) d\bar{\xi} \wedge d\bar{\xi}, \quad (12)$$

which is curiously the “double copy” of Eq. (10) [77]. Therefore, we conclude that the string defect of Eq. (3) indeed describes a thin flux tube of time monodromy.

Finally, we explicitly derive the distributional stress-energy tensor of this string. In general one can take various approaches [78–80], but here we particularly consider the distributional stress-energy that follows from the “harmonic-gauge linearization” of Ref. [48] which the Kerr-Schild description enables as a luxury.

The harmonic-gauge linearization employs a crucial property that the congruence of $\ell^\mu \partial_\mu$ can enjoy: geodesic and shear-free (GSF). Given such assumptions, it follows that $T^{\mu\nu} = u^{(\mu} J^{\nu)}$ can be identified as the source for the gravitational solution [48]. Here J^μ denotes the source for the electromagnetic solution, and $u^\mu = \delta^\mu_0$ is the unit timelike Killing vector. Applying this formula, we immediately reproduce Bonnor’s stress-energy tensor [26, 27],




FIG. 1. The Misner string is a “screw dislocation” in spacetime, inducing time monodromy as an Aharonov-Bohm phase.

describing a point mass m attached to an energy-less [81] half-infinite rotating rod! The GSF property of our null one-form in Eq. (7) can be shown from an identity $\partial_\mu \ell_\nu = (1/2r)(\eta_{\mu\nu} + u_\mu u_\nu - (\ell - u)_\mu (\ell - u)_\nu + i \varepsilon_{\mu\nu\rho\sigma} u^\rho \ell^\sigma)$, while the single copy source J^μ easily follows by acting on the Laplacian $\square := -\eta^{\mu\nu} \partial_\mu \partial_\nu$ to Eq. (1). This shows that the string defect of Eq. (3) indeed describes a thin solenoid of mass current.

To sum up, we have convinced ourselves from multiple angles that the Kerr-Schild metric in Eq. (3) indeed exhibits the geometrical and physical characteristics of a Taub-NUT spacetime arising from the Misner string. This analysis demonstrates that the Kerr-Schild description of the solution is not merely a formal construct but physical. Moreover, the quick derivation of the distributional source demonstrates its usefulness as well.

Weyl double copy in position space.—Finally, let us unlock more classical double copy relations for the SDTN solution. First of all, we consider the *Weyl double copy* and explicate how the Kerr-Schild metric in Eq. (3) resolves a subtlety in the previous constructions [3, 4].

We begin with a brief review. The Weyl double copy directly constructs the Weyl curvatures of algebraically special spacetimes by “squaring” electromagnetic field strengths [2–5], while not necessarily referencing a metric or a gauge potential. For example, take the Coulomb and Schwarzschild solutions. In the spinor language, their field strength and Weyl tensor are given as the following:

$$F_{\dot{\alpha}\dot{\beta}} = \frac{2q}{4\pi r^2} \tilde{o}_{(\dot{\alpha}} \tilde{l}_{\dot{\beta})}, \quad C_{\dot{\alpha}\dot{\beta}\dot{\gamma}\dot{\delta}} = -\frac{6m}{4\pi r^3} \tilde{o}_{(\dot{\alpha}} \tilde{o}_{\dot{\beta}} \tilde{l}_{\dot{\gamma}} \tilde{l}_{\dot{\delta})}, \quad (13)$$

where we have omitted the anti-self-dual counterparts. The Weyl double copy is the squaring relation $C_{\dot{\alpha}\dot{\beta}\dot{\gamma}\dot{\delta}} = F_{(\dot{\alpha}\dot{\beta}} F_{\dot{\gamma}\dot{\delta})} / S$ [2], which holds with $S \propto 1/r$.

The Weyl double copy correspondences between the SDD and SDTN solutions are established in Refs. [3, 4] as the “self-dual parts” of the Coulomb and Schwarzschild solutions. That is, the self-dual Weyl curvature (and the principal spinors) of the SDD/SDTN solution is simply *postulated* to be that of the Coulomb/Schwarzschild solution. This physical anticipation, however, may appear a little nontrivial if one recalls the nonlinearity of gravity [82]. Especially, the spin-2 field $\phi_{\dot{\alpha}\dot{\beta}\dot{\gamma}\dot{\delta}}$ obtained from the

Penrose transform describes only the linearized Weyl tensor in the flat background [50]. To promote it to the fully nonlinear solution one necessarily has to appeal to a linearization, which is famously possible in the Kerr-Schild gauge [48, 83, 84].

Certainly, the explicit Kerr-Schild metric we have established clearly resolves this subtlety and concretely validates the Weyl double copy constructions in Refs. [3, 4]. Through a straightforward computation of spin connection coefficients in a tetrad frame, we explicitly checked that the self-dual Weyl tensor of the *nonlinear* SDTN solution as defined in Eq. (3) indeed is equal to that of the Schwarzschild solution with the same principal spinors. Specifically, the null one-form in Eq. (7) boils down to

$$-\frac{1}{2}\ell_\mu(\sigma^\mu)_{\alpha\dot{\alpha}} = o_\alpha\tilde{o}_{\dot{\alpha}}, \quad \tilde{o}^{\dot{\alpha}} = \begin{pmatrix} 1 \\ \zeta \end{pmatrix}, \quad o^\alpha = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad (14)$$

where $\tilde{o}^{\dot{\alpha}}$ is one of the principal spinors in Eq. (13). Computing the Weyl tensor then reveals the other principal spinor, $\tilde{l}^{\dot{\alpha}} = (-\zeta, 1)/(1 + \zeta\bar{\zeta})$. Crucially, the *real* combinations $[\tilde{o}_{\dot{\alpha}}]^*\tilde{o}_{\dot{\beta}}$, $[\tilde{l}_{\dot{\alpha}}]^*\tilde{l}_{\dot{\beta}}$ of these principal spinors exactly reproduce the null one-forms $-dt \pm dr$ associated with the Kerr-Schild metrics of the Schwarzschild solution [3, 4], hence confirming the statement that SDTN is the “self-dual part” of Schwarzschild.

Note that this relationship between SDTN and Schwarzschild could be traced back to the fact that the condition for a null vector field $\tilde{\pi}^{\dot{\alpha}}(x)\pi^\alpha(x)$ to be GSF is given by a *chiral* equation for each copy of the spinors: in flat background, the equation for the right-handed reads [50, 85]

$$\tilde{\pi}^{\dot{\alpha}}(x)\tilde{\pi}^{\dot{\beta}}(x)(\partial_{\alpha\dot{\alpha}}\tilde{\pi}_{\dot{\beta}}(x)) = 0. \quad (15)$$

If $\tilde{\pi}^{\dot{\alpha}}(x)$ is a solution to Eq. (15), then both the *complexified, self-dual* combination $o_\alpha\tilde{\pi}_{\dot{\beta}}(x)$ (with constant o_α) and the *real, non-self-dual* combination $[\tilde{\pi}_{\dot{\alpha}}(x)]^*\tilde{\pi}_{\dot{\beta}}(x)$ define null GSF one-forms, from which the idea of taking the “self-dual part” is validated [86].

Weyl double copy in twistor space.—Next, let us revisit the Weyl double copy from *twistor space* [3, 4], which eventually illuminates how the Kerr-Schild coordinates have been secretly used in the twistor theory context.

To smoothly motivate the twistor space, we consider a curious theorem due to Kerr [50, 87, 88]: any solution to Eq. (15) arises as a zero locus of a holomorphic function $f(\tilde{Z})$ homogeneous in its four complex arguments $\tilde{Z}^A = (\tilde{\pi}_{\dot{\alpha}}, \tilde{\omega}^\alpha)$ upon plugging in the “incidence relation” $\tilde{\omega}^\alpha = \tilde{\pi}_{\dot{\alpha}}x^{\dot{\alpha}\alpha}$ [87]. This curious complex variable \tilde{Z}^A is called a *twistor* [87] and is known to represent a null ray in complexified Minkowski space [87].

Taking inspirations from Refs. [89–91], Ref. [4] proposed the following function for constructing the GSF congruences associated with the SDTN solution [92]:

$$f(\tilde{Z}) = -\tilde{\omega}^\alpha u_{\alpha\dot{\alpha}}\tilde{\pi}^{\dot{\alpha}}, \quad u_{\alpha\dot{\alpha}} := u_\mu(\sigma^\mu)_{\alpha\dot{\alpha}}. \quad (16)$$

Indeed, $f(\tilde{\pi}, \tilde{\pi}x) = 0$ with the above function exactly yields our principal spinors $\tilde{o}_{\dot{\alpha}}, \tilde{l}_{\dot{\alpha}}$ as $\tilde{\pi}_{\dot{\alpha}}(x) \propto \tilde{o}_{\dot{\alpha}}, \tilde{l}_{\dot{\alpha}}$.

Remarkably, this twistor function turns out to encode not only the congruences but the entire geometry of the (linearized) SDTN spacetime through a contour integral formula known as Penrose transform [93, 94], where the “localization” of $\tilde{\pi}_{\dot{\alpha}}$ to zeroes of $f(\tilde{\pi}, \tilde{\pi}x)$ arises from a residue computation [3, 4, 50]:

$$\phi_{\dot{\alpha}_1 \dots \dot{\alpha}_{2h}}(x) = \oint \frac{\tilde{\pi}_{\dot{\gamma}} d\tilde{\pi}^{\dot{\gamma}}}{2\pi i} \frac{\tilde{\pi}_{\dot{\alpha}_1} \dots \tilde{\pi}_{\dot{\alpha}_{2h}}}{(f(\tilde{\pi}, \tilde{\pi}x))^{h+1}}. \quad (17)$$

Evaluating these contour integrals for $h = 1, 2$ readily reproduces the spin-1 and spin-2 fields in Eq. (13). The calculations are almost immediate if one notes that $f(\tilde{\pi}, \tilde{\pi}x) = -\tilde{\pi}_{\dot{\alpha}}\tilde{\chi}^{\dot{\alpha}\dot{\beta}}\tilde{\pi}^{\dot{\beta}} = -r\sigma$ with $\tilde{\chi}_{\dot{\alpha}\dot{\beta}} = r\tilde{o}_{(\dot{\alpha}}\tilde{l}_{\dot{\beta})}$ when the contour is parameterized as $\tilde{\pi}_{\dot{\alpha}} = \tilde{o}_{\dot{\alpha}} + \sigma\tilde{l}_{\dot{\alpha}}$, where

$$\tilde{\chi}^{\dot{\alpha}\dot{\beta}} = -x^\mu u^\nu (\tilde{\sigma}_{\mu\nu})^{\dot{\alpha}\dot{\beta}} = -\frac{1}{2}(\vec{x} \cdot \vec{\sigma})^{\dot{\alpha}\dot{\beta}} \quad (18)$$

describes the self-dual part of the bivector $2x^{[\mu}u^{\nu]}$.

In this manner, the curvature tensors arise as the Penrose transform of functions $\mathcal{M}_h(\tilde{Z}) = (f(\tilde{Z}))^{-h-1}$ in twistor space, which are in fact scattering amplitudes of a massive object absorbing a negative-helicity force carrier [49] [95]. The twistor-space Weyl double copy is the squaring relation $\mathcal{M}_2(\tilde{Z}) = (\mathcal{M}_1(\tilde{Z}))^2/\mathcal{M}_0(\tilde{Z})$ between these “twistor-space amplitudes.”

Crucially, this rederivation of the twistor-space construction of the SDTN solution is not merely a review. Throughout, the coordinates $x^\mu = (\sigma^\mu)_{\alpha\dot{\alpha}}x^{\dot{\alpha}\alpha}$ have always referred to the Kerr-Schild coordinates defined by the metric in Eq. (3). This explicitly shows that our Kerr-Schild coordinates are exactly the Cartesian coordinates which the twistor-space construction of the SDTN solution employs in terms of Penrose transform. In fact, this may not be surprising because our Kerr-Schild coordinates turn out to be just Plebański’s coordinates [54, 55] leading to the second heavenly equation. To conclude, twistor-theoretic perspectives on the SDTN solution have presumed the Kerr-Schild coordinates, even if the explicit metric may have not been written down as in Eq. (3) or shown to be equivalent to the previously documented form (Gibbons-Hawking [41] or Taub-NUT [23, 24]).

As a side note, we also remark that $\tilde{\chi}^{\dot{\alpha}\dot{\beta}}$ in Eq. (18) is in fact the Killing spinor [50–53] of the SDTN solution, from which a “position vector” \vec{x} is identified [34]. The identity of this \vec{x} should be evident now.

Operator Kerr-Schild double copy.—Lastly, we propose a *nonlocal operator Kerr-Schild double copy* as a new result, which interestingly differs from the self-dual double copy previously established in Refs. [1, 17].

In operator Kerr-Schild double copy, the null one-form ℓ_μ in the Kerr-Schild ansatz is promoted to a differential operator \hat{k}_μ . Especially, it can factorize as $(\sigma^\mu)_{\alpha\dot{\alpha}}\hat{k}_\mu = -2o_\alpha\hat{k}_{\dot{\alpha}}$ so that the fields are manifestly described in the lightcone gauge [1, 7, 8]. Intriguingly, we find that the gauge potential in Eq. (2) and the metric in Eq. (3) precisely arise in this way if the operator $\hat{k}_{\dot{\alpha}}$ is given by

$$\hat{k}_{\dot{\alpha}} = \left(-\frac{\partial_x + i\partial_y}{\partial_z}, 1 \right), \quad (19)$$

which is *nonlocal*. Specifically, direct computation shows that $\hat{k}_{\hat{\alpha}}(1/r) = (1/r)\bar{o}_{\hat{\alpha}}$ and $\hat{k}_{\hat{\alpha}}\hat{k}_{\hat{\beta}}(1/r) = (1/r)\bar{o}_{\hat{\alpha}}\bar{o}_{\hat{\beta}}$, so a classical double copy correspondence between SDD and SDTN is realized in the following form:

$$A_{\alpha\hat{\alpha}} = -2o_{\alpha}\hat{k}_{\hat{\alpha}}\frac{q}{4\pi r}, \quad h_{\alpha\hat{\alpha}\beta\hat{\beta}} = 4o_{\alpha}o_{\beta}\hat{k}_{\hat{\alpha}}\hat{k}_{\hat{\beta}}\frac{m}{4\pi r}, \quad (20)$$

where $h_{\mu\nu} := g_{\mu\nu} - \eta_{\mu\nu}$ denotes the ‘‘graviton field.’’

Interestingly, we find that this exotic new form of classical double copy is demystified from an electric-magnetic dual perturbation theory of dyons [56–60]. Note first that the distributional source of the SDD is explicitly given by

$$J^{\mu} = \partial_{\nu}F^{\mu\nu} = \left(1, -i\frac{\vec{n}\times\vec{\partial}}{\vec{n}\cdot\vec{\partial}}\right)q\delta^{(3)}(\vec{x}), \quad (21)$$

where \vec{n} is the unit vector along the z -axis [96]. This describes an extended object: the anti-derivative $(-\vec{n}\cdot\vec{\partial})^{-1} = (-\partial_z)^{-1}$ stacks magnetic dipoles $(-\vec{n}\times\vec{\partial})(-iq\delta^{(3)}(\vec{x}))$ along the negative z -axis, implementing a thin solenoid. On the other hand, physically we know that this extended object mimics an isolated point dyon $(q, -iq)$ localized at the origin, which is described by the following source:

$$\begin{aligned} J^{\mu} &= \partial_{\nu}F^{\mu\nu} = qu^{\mu}\delta^{(3)}(\vec{x}), \\ J^{*\mu} &= -\partial_{\nu}{}^*F^{\mu\nu} = -iqu^{\mu}\delta^{(3)}(\vec{x}). \end{aligned} \quad (22)$$

Specifically, $F_{\mu\nu}$ sourced by Eq. (21) is identical to $F_{\mu\nu}$ sourced by Eq. (22) at all points outside the solenoid [97]. Yet crucially, Eq. (22) involves a bonafide violation of the Bianchi identity: $\partial_{[\rho}F_{\mu\nu]} \neq 0$. Hence it does not define a source in the Maxwell theory but instead belongs to its *extension* admitting both electric and magnetic sources considered by Dirac [98].

As a dynamical field theory, a Lagrangian formulation of this extended Maxwell theory is given by Zwanziger’s ‘‘two-potential electrodynamics’’ where the dual photon field A_{μ}^* is employed to incorporate the Bianchi violation [56, 99]. This means the gauge potential of the SDD could have been derived from the dyonic source in Eq. (22) by using the propagators of Zwanziger theory!

The Zwanziger propagators can be found in Refs. [56–59]. Here, we note that a single spinor equation succinctly summarizes them [100]:

$$\hat{\Delta}_{\alpha\hat{\alpha}\beta\hat{\beta}}^{+-} = \frac{n_{\alpha\hat{\beta}}\partial_{\beta\hat{\alpha}}}{-n\cdot\partial}\frac{1}{\square}, \quad (23)$$

where $n_{\alpha\hat{\alpha}}$ is an auxiliary four-vector that traces back to the Dirac string. Eq. (23) describes the position-space propagator from A_{μ}^{-} to A_{μ}^{+} where $A_{\mu}^{\pm} := \frac{1}{2}(A_{\mu} \pm iA_{\mu}^*)$, while other propagators between A_{μ}^{\pm} vanish. Meanwhile, the sources coupling to A_{μ}^{\pm} are given by $J_{\pm}^{\mu} = J^{\mu} \mp iJ^{*\mu}$, which are $J_{+}^{\mu} = 0$, $J_{-}^{\mu} = 2qu^{\mu}\delta^{(3)}(\vec{x})$ for Eq. (22). As a result, the photon field $A_{\mu} = A_{\mu}^{+} + A_{\mu}^{-}$ sourced by Eq. (22) is found as

$$A_{\alpha\hat{\alpha}} = A_{\alpha\hat{\alpha}}^{+} = \frac{n_{\alpha\hat{\beta}}u^{\hat{\beta}\beta}\partial_{\beta\hat{\alpha}}}{n\cdot\partial}\frac{q}{4\pi r}. \quad (24)$$

Then a simple calculation shows that Eqs. (19) and (20) are exactly reproduced when the auxiliary vector is taken to be null as $n_{\alpha\hat{\alpha}} = -o_{\alpha}\bar{o}_{\hat{\alpha}}$, yielding $\hat{k}_{\hat{\alpha}} = -\partial_{1\hat{\alpha}}/\partial_{0\hat{0}}$.

Amusingly, the graviton field arises in the same fashion by using the *linearized gravity equivalent of Zwanziger propagator*, which is simply given by double copying Eq. (23) as is advocated in Ref. [60]! In the gravitational case the source dual to $T^{\mu\nu} = u^{(\mu}J^{\nu)}$ is given by $T_{+}^{\mu\nu} = 0$ and $T_{-}^{\mu\nu} = 2mu^{\mu}u^{\nu}\delta^{(3)}(\vec{x})$ [101], describing a static self-dual gravitational dyon. As a result, the linearized graviton field $h_{\mu\nu} = h_{\mu\nu}^{+} + h_{\mu\nu}^{-}$ is found as

$$h_{\alpha\hat{\alpha}\beta\hat{\beta}} = h_{\alpha\hat{\alpha}\beta\hat{\beta}}^{+} = \frac{n_{\alpha\hat{\gamma}}u^{\hat{\gamma}\gamma}\partial_{\gamma\hat{\alpha}}}{n\cdot\partial}\frac{n_{\beta\hat{\delta}}u^{\hat{\delta}\delta}\partial_{\delta\hat{\beta}}}{n\cdot\partial}\frac{m}{4\pi r}, \quad (25)$$

which is in turn identified as the full metric perturbation $g_{\mu\nu} - \eta_{\mu\nu}$ by virtue of the linearization in the Kerr-Schild gauge [48, 83, 84]. This precisely explains the double copy structure proposed in Eq. (20) as we claimed earlier.

A few comments are in order. Firstly, it is instructive to revisit this construction from the on-shell perspective. It is easy to check that the ‘‘numerator factor’’ of the off-shell propagator in Eq. (23) correctly factorizes into the null polarization vectors with $n_{\alpha\hat{\alpha}} = -o_{\alpha}\bar{o}_{\hat{\alpha}}$, from which its double copy can be argued [60]. The double copy of sources from $J_{-}^{\mu} = 2qu^{\mu}\delta^{(3)}(\vec{x})$ to $T_{-}^{\mu\nu} = 2mu^{\mu}u^{\nu}\delta^{(3)}(\vec{x})$ realizes the squaring relation of the massive on-shell amplitudes in terms of the x -factor [60, 102–104]. Further, one may also explicitly connect the off-shell constructed fields in Eqs. (24) and (25) to the Penrose transform in Eq. (17) through a parameterization of off-shell momentum employed in Ref. [49].

Secondly, it will be interesting to examine whether this ‘‘Zwanziger double copy’’ persists for generic Taub-NUT [105]. Its failure would signal the significance of not only linearization by Kerr-Schild gauge but also self-duality.

Lastly, we report a strange fact that the SDTN solution *fails* to be the *self-dual double copy* [1, 7, 17] of SDD. Such a dissonance is disturbing, especially since the self-dual double copy is arguably the most natural option for self-dual solutions and will imply fascinating conclusions such as ‘‘SDTN black hole *is* a large- N dyon’’ by being rooted on Moyal deformation [106–109] of Plebański [54]’s second heavenly equation [110, 111]. Specifically, the *local* operator $\hat{k}_{\hat{\alpha}} = \partial_{0\hat{\alpha}}$ of the self-dual double copy makes the Plebański scalar field different for each helicity: $A_{\alpha\hat{\alpha}} = -2o_{\alpha}\partial_{0\hat{\alpha}}(q\zeta/4\pi)$, $h_{\alpha\hat{\alpha}\beta\hat{\beta}} = 4o_{\alpha}o_{\beta}\partial_{0\hat{\alpha}}\partial_{0\hat{\beta}}(m(2r+z)\zeta^2/24\pi)$ [112]. This puzzle seems to deserve a further investigation, possibly by comparing with the case of Eguchi-Hanson instanton [7].

Conclusions.—In this note, we have shown that a complexified diffeomorphism reveals a Kerr-Schild metric for the SDTN solution. Crucially, the validity of this diffeomorphism and the metric is verified by highlighting their intrinsically complex nature. Especially, we have verified that the Kerr-Schild metric also exhibits the Misner string defect, supporting its physical interpretation as a gravitational dyon. Further explicitly, we have established three concrete senses in which the SDTN solu-

tion is precisely the gravitational counterpart of a SDD in electromagnetism: Kerr-Schild, Weyl, and operator Kerr-Schild classical double copy. Notably, the last one adds to the inventory of classical double copies as a new entry. We have identified its physical origin as the double copy of propagators between dyonic sources.

It is interesting to realize that the Gibbons–Hawking ansatz had already suggested a sense in which self-dual electromagnetic objects correspond to self-dual gravitational objects, far before the advent of the classical double copy program. It is amusing to see how this correspondence has become strengthened for SDTN and SDD from three new perspectives. Especially, the idea that the Misner string is a gravitational analog of Dirac string is concretely confirmed in the first and third double copy constructions. On a related note, we can also frame this work from the angle of “classical double copy constructions of gravitational instantons.” A noteworthy example has been given by Ref. [7], where the Eguchi–Hanson instanton [113–115] is constructed from the self-dual double copy (cf. Ref. [116]). This work adds another instance to the discourse, i.e., the “Taub–NUT instanton.”

The Kerr–Schild metric of the SDTN solution conveys an aesthetic elegance in terms of its minimalist and spinorial nature as manifested in Eq. (14). To investigate what facilitates this remarkable gem, we could examine its relation to the self-dual limit of the *double* Kerr–Schild metric employed in Ref. [6]. Finding a diffeomorphism bridging between the two would shed an insight [117].

Meanwhile, as we have clarified the Kerr–Schild *coordinates* themselves have been already employed in the spinorial/twistorial formalisms. In this light, the metric in Eq. (3) can be thought of as providing an explicit construction of a “potential” for the Weyl curvature which the Penrose transform or Killing spinor has described.

Yet, we believe there is a value in having a more explicit description. In particular, it will be interesting to explore further consequences and applications of our Kerr–Schild metric: asymptotic symmetries [118, 119] [120], revisiting the integrability of test-particle motion [39, 40], black hole perturbation theory [34, 35, 121, 122], or finding exact graviton propagators [123–125] to carry out self-force calculations [126–128] in an idealized set-up, to list a few directions. Especially, explorations in the scattering context are likely to shed insights on Kerr black hole scattering as a certain integrable sector on account of the Newman–Janis shift: the “self-dual part” of the Kerr solution is a SDTN solution displaced along an imaginary direction [1, 22, 34–36, 61, 129–133].

Amusingly, about a year ago we found a vivid physical picture that explains this fact explicitly by means of complex diffeomorphisms. Namely, as depicted in Fig. 2, we realized that a simple complex coordinate transformation maps the Kerr metric in Kerr–Schild coordinates to another Kerr–Schild metric describing two point masses joined by a Misner string. By zooming into the upper tip of the string, we obtained—by construction—a Kerr–Schild metric, which was precisely the metric in Eq. (3).




FIG. 2. The very physical origin of the Newman–Janis shift is the fact that the Kerr black hole is secretly a “dumbbell” of self-dual and anti-self-dual Taub–NUT instantons [134].

And crucially, this had to be describing a Taub–NUT instanton according to the Newman–Janis shift. The complex diffeomorphism in Eq. (5) was precisely a result of explicitly verifying this strong conviction. This is how we originally derived the Kerr–Schild metric reported in this note. We hope to release more details soon [134].

Note added. Shortly after the initial release of this work, the author was informed that a split-signature version of the Kerr–Schild metric in Eq. (3) was previously *conjectured* in Ref. [64]. Then, about a week after this notice, Ref. [135] appeared, providing an *existence proof* for a real diffeomorphism in split signature but “without determining an explicit diffeomorphism carrying one line element to the other” [135]. In contrast, this work directly provides a concrete *constructive proof* in the form of an explicit diffeomorphism, for all signatures by taking the complexified point of view [136]. Especially, it is interesting to observe that our constructive proof in fact resolves the question left by Ref. [135]: taking the real slice $t, x, iy, z \in \mathbb{R}$ makes the two metrics in Eqs. (3) and (6) as well as the diffeomorphism in (5) all real for real mass $m \in \mathbb{R}$, while also exactly reproducing Ref. [64]’s conjectured metric, Eq. (6.16) [137].

Acknowledgements. We would like to thank Clifford Cheung, Jung-Wook Kim, Andres Luna, Donal O’Connell, Nabha Shah, Justin Vines, and Chirs D. White for discussions and insightful comments. We are also grateful to Gabriel Herczeg, Max Pezzelle, and Jash Desai for bringing the author’s attention to the works [64, 135] and the follow-up discussions. This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of High Energy Physics, under Award Number DE-SC0011632. J.-H.K. is also supported by Ilju Academy and Culture Foundation.

Appendix: Kerr–Schild to Gibbons–Hawking by the null geodesic flow

Suppose a vector field ℓ^μ in complexified Minkowski space that defines a geodesic congruence: $\ell^\mu{}_{,\nu}\ell^\nu = 0$. The time- ϵ flow generated by this vector field is simply $x^\mu \mapsto x^\mu + \epsilon\ell^\mu(x)$, as geodesics are straight lines in flat space. The pull-back of the flat metric $\eta = \eta_{\mu\nu} dx^\mu \otimes dx^\nu$ is $\eta + 2\epsilon\ell_{(\mu,\nu)} dx^\mu \otimes dx^\nu + \epsilon^2\eta_{\rho\sigma}\ell^\rho{}_{,\mu}\ell^\sigma{}_{,\nu} dx^\mu \otimes dx^\nu$.

The null vector field associated with the Kerr–Schild metric of the SDTN solution, $\ell^\mu = (1, \zeta, -i\zeta, 1)$, is

stationary, geodesic, and shear-free. With the self-duality condition, they together implies that $2\partial_\nu\ell^\mu = \rho\tilde{m}^\mu m_\nu$, when described with a null tetrad $(\ell^\mu, n^\mu = 2u^\mu - \ell^\mu, m^\mu, \tilde{m}^\mu)$ where $u^\mu := \delta^\mu_0$. The complex expansion computes to $\rho = 1/r$.

As a result, it follows that the flat metric transforms to $\eta + \epsilon\rho\tilde{m}\odot m$ if the geodesic flow $x^\mu \mapsto x^\mu + \epsilon\ell^\mu(x)$ is used as a diffeomorphism, where \odot denotes the symmetrized tensor product. In turn, the Kerr-Schild metric $\eta + \phi\ell^2$ transforms to

$$\eta + \epsilon\rho\tilde{m}\odot m + \phi'\ell^2, \quad (26)$$

where $\phi'(x) := \phi(x + \epsilon\ell(x))$.

Meanwhile, using $\eta = -2u\odot\ell + \ell^2 + \tilde{m}\odot m$, the Gibbons-Hawking metric $-(1+\phi)^{-1}(u - \phi(\ell - u))^2 +$

$(1+\phi)(u^2 + \eta)$ boils down to

$$\eta + \phi\tilde{m}\odot m + (\phi^{-1} + 1)^{-1}\ell^2. \quad (27)$$

Therefore, if we can flow along the geodesic congruence such that $\phi^{-1} + 1 = \phi'^{-1}$ and $\epsilon\rho = \phi$, then the Kerr-Schild metric is mapped to the Gibbons-Hawking metric. Evidently, $\epsilon = m/4\pi$ satisfies both of these conditions. Thus the diffeomorphism from the Kerr-Schild to Gibbons-Hawking metrics is given by $t \mapsto t + m/4\pi$, $x \mapsto x + m\zeta/4\pi$, $y \mapsto y - im\zeta/4\pi$, $z \mapsto z + m/4\pi$, which is equivalent to Eq. (5) up to dropping the constant shift in the time coordinate. It could be interesting if this “trick” also proves to be useful for more general cases such as non-self-dual solutions or multi-centered Gibbons-Hawking metrics.

-
- [1] R. Monteiro, D. O’Connell, and C. D. White, “Black holes and the double copy,” *Journal of High Energy Physics* **2014** no. 12, (2014) 1–23.
- [2] A. Luna, R. Monteiro, I. Nicholson, and D. O’Connell, “Type D spacetimes and the Weyl double copy,” *Classical and Quantum Gravity* **36** no. 6, (2019) 065003.
- [3] E. Chacón, S. Nagy, and C. D. White, “The Weyl double copy from twistor space,” *Journal of High Energy Physics* **2021** no. 5, (2021) 1–40.
- [4] C. D. White, “Twistorial foundation for the classical double copy,” *Physical Review Letters* **126** no. 6, (2021) 061602.
- [5] A. Luna, N. Moynihan, and C. D. White, “Why is the Weyl double copy local in position space?,” *JHEP* **12** (2022) 046, [arXiv:2208.08548 \[hep-th\]](#).
- [6] A. Luna, R. Monteiro, D. O’Connell, and C. D. White, “The classical double copy for Taub-NUT spacetime,” *Phys. Lett. B* **750** (2015) 272–277, [arXiv:1507.01869 \[hep-th\]](#).
- [7] D. S. Berman, E. Chacón, A. Luna, and C. D. White, “The self-dual classical double copy, and the Eguchi-Hanson instanton,” *JHEP* **01** (2019) 107, [arXiv:1809.04063 \[hep-th\]](#).
- [8] G. Elor, K. Farnsworth, M. L. Graesser, and G. Herczeg, “The Newman-Penrose map and the classical double copy,” *Journal of High Energy Physics* **2020** no. 12, (2020) 1–37.
- [9] I. Bah, R. Dempsey, and P. Weck, “Kerr-Schild double copy and complex worldlines,” *Journal of High Energy Physics* **2020** no. 2, (2020) 1–34.
- [10] M. Carrillo González, B. Melcher, K. Ratliff, S. Watson, and C. D. White, “The classical double copy in three spacetime dimensions,” *JHEP* **07** (2019) 167, [arXiv:1904.11001 \[hep-th\]](#).
- [11] M. Carrillo-González, R. Penco, and M. Trodden, “The classical double copy in maximally symmetric spacetimes,” *JHEP* **04** (2018) 028, [arXiv:1711.01296 \[hep-th\]](#).
- [12] R. Monteiro, D. O’Connell, D. Peinador Veiga, and M. Sergola, “Classical solutions and their double copy in split signature,” *JHEP* **05** (2021) 268, [arXiv:2012.11190 \[hep-th\]](#).
- [13] R. Monteiro, S. Nagy, D. O’Connell, D. Peinador Veiga, and M. Sergola, “NS-NS spacetimes from amplitudes,” *JHEP* **06** (2022) 021, [arXiv:2112.08336 \[hep-th\]](#).
- [14] N. Bahjat-Abbas, A. Luna, and C. D. White, “The Kerr-Schild double copy in curved spacetime,” *JHEP* **12** (2017) 004, [arXiv:1710.01953 \[hep-th\]](#).
- [15] A. Banerjee, E. O. Colgáin, J. A. Rosabal, and H. Yavartanoo, “Ehlers as EM duality in the double copy,” *Phys. Rev. D* **102** (2020) 126017, [arXiv:1912.02597 \[hep-th\]](#).
- [16] L. Alfonsi, C. D. White, and S. Wikeley, “Topology and Wilson lines: global aspects of the double copy,” *JHEP* **07** (2020) 091, [arXiv:2004.07181 \[hep-th\]](#).
- [17] R. Monteiro and D. O’Connell, “The kinematic algebra from the self-dual sector,” *Journal of High Energy Physics* **2011** no. 7, (2011) 1–24.
- [18] H. Kawai, D. C. Lewellen, and S. H. H. Tye, “A Relation Between Tree Amplitudes of Closed and Open Strings,” *Nucl. Phys. B* **269** (1986) 1–23.
- [19] Z. Bern, J. J. M. Carrasco, and H. Johansson, “Perturbative Quantum Gravity as a Double Copy of Gauge Theory,” *Phys. Rev. Lett.* **105** (2010) 061602, [arXiv:1004.0476 \[hep-th\]](#).
- [20] Z. Bern, T. Dennen, Y.-t. Huang, and M. Kiermaier, “Gravity as the Square of Gauge Theory,” *Phys. Rev. D* **82** (2010) 065003, [arXiv:1004.0693 \[hep-th\]](#).
- [21] Z. Bern, J. J. Carrasco, M. Chiodaroli, H. Johansson, and R. Roiban, “The duality between color and kinematics and its applications,” [arXiv:1909.01358 \[hep-th\]](#).
- [22] N. Arkani-Hamed, Y.-t. Huang, and D. O’Connell, “Kerr black holes as elementary particles,” *JHEP* **01** (2020) 046, [arXiv:1906.10100 \[hep-th\]](#).
- [23] A. H. Taub, “Empty space-times admitting a three parameter group of motions,” *Annals Math.* **53** (1951) 472–490.
- [24] E. Newman, L. Tamburino, and T. Unti, “Empty space generalization of the Schwarzschild metric,” *J. Math. Phys.* **4** (1963) 915.
- [25] C. W. Misner, “The Flatter regions of Newman, Unti and Tamburino’s generalized Schwarzschild space,” *J. Math. Phys.* **4** (1963) 924–938.
- [26] W. B. Bonnor, “A new interpretation of the NUT metric in general relativity,” *Math. Proc. Cambridge Phil. Soc.* **66** no. 1, (1969) 145–151.
- [27] A. Sackfield, “Physical interpretation of NUT metric,” in *Mathematical Proceedings of the Cambridge Philosophical Society*, vol. 1, pp. 89–94, Cambridge University Press, 1971.
- [28] Y. Cho, “Magnetic theory of gravitation,” in *Primordial Nucleosynthesis and Evolution of Early Universe*, pp. 203–211. Springer, 1991.
- [29] J. Dowker and J. Roche, “The gravitational analogues of magnetic monopoles,” *Proceedings of the Physical Society (1958-1967)* **92** no. 1, (1967) 1.
- [30] J. Samuel and B. Iyer, “A gravitational analog of the Dirac monopole,” *Current Science* (1986) 818–824.
- [31] R. Maartens and B. A. Bassett, “Gravito-electromagnetism,” *Classical and Quantum Gravity* **15** no. 3, (1998) 705.
- [32] J. F. Plebański, “A class of solutions of Einstein-Maxwell equations,” *Annals Phys.* **90** no. 1, (1975) 196–255.
- [33] Z. W. Chong, G. W. Gibbons, H. Lu, and C. N. Pope,

- “Separability and killing tensors in Kerr-Taub-NUT-de Sitter metrics in higher dimensions,” *Phys. Lett. B* **609** (2005) 124–132, [arXiv:hep-th/0405061](#).
- [34] T. Adamo, G. Bogna, L. Mason, and A. Sharma, “Scattering on self-dual Taub-NUT,” [arXiv:2309.03834 \[hep-th\]](#).
- [35] A. Guevara and U. Kol, “Self Dual Black Holes as the Hydrogen Atom,” [arXiv:2311.07933 \[hep-th\]](#).
- [36] E. Crawley, A. Guevara, N. Miller, and A. Strominger, “Black holes in Klein space,” *JHEP* **10** (2022) 135, [arXiv:2112.03954 \[hep-th\]](#).
- [37] E. Crawley, A. Guevara, E. Himwich, and A. Strominger, “Self-dual black holes in celestial holography,” *JHEP* **09** (2023) 109, [arXiv:2302.06661 \[hep-th\]](#).
- [38] T. Adamo, G. Bogna, L. Mason, and A. Sharma, “Gluon scattering on the self-dual dyon,” [arXiv:2406.09165 \[hep-th\]](#).
- [39] G. W. Gibbons and P. J. Ruback, “The Hidden Symmetries of Taub-NUT and Monopole Scattering,” *Phys. Lett. B* **188** (1987) 226.
- [40] G. W. Gibbons and P. J. Ruback, “The Hidden Symmetries of Multicenter Metrics,” *Commun. Math. Phys.* **115** (1988) 267.
- [41] S. W. Hawking, “Gravitational instantons,” *Physics Letters A* **60** no. 2, (1977) 81–83.
- [42] G. W. Gibbons and S. W. Hawking, “Gravitational Multi-Instantons,” *Phys. Lett. B* **78** (1978) 430.
- [43] G. W. Gibbons and S. W. Hawking, “Classification of Gravitational Instanton Symmetries,” *Commun. Math. Phys.* **66** (1979) 291–310.
- [44] T. Eguchi, P. B. Gilkey, and A. J. Hanson, “Gravitation, Gauge Theories and Differential Geometry,” *Phys. Rept.* **66** (1980) 213.
- [45] M. Dunajski, *Solitons, instantons, and twistors*. Oxford University Press, 2024.
- [46] P. A. M. Dirac, “Quantised singularities in the electromagnetic field,,” *Proc. Roy. Soc. Lond. A* **133** no. 821, (1931) 60–72.
- [47] P. A. M. Dirac, “The Theory of magnetic poles,” *Phys. Rev.* **74** (1948) 817–830.
- [48] J. Vines, “Scattering of two spinning black holes in post-Minkowskian gravity, to all orders in spin, and effective-one-body mappings,” *Classical and Quantum Gravity* **35** no. 8, (2018) 084002.
- [49] A. Guevara, “Reconstructing Classical Spacetimes from the S-Matrix in Twistor Space,” [arXiv:2112.05111 \[hep-th\]](#).
- [50] R. Penrose and W. Rindler, “Spinors and space-time: Volume 2, spinor and twistor methods in space-time geometry,.”
- [51] M. Walker and R. Penrose, “On quadratic first integrals of the geodesic equations for type [22] spacetimes,” *Commun. Math. Phys.* **18** (1970) 265–274.
- [52] L. P. Hughston, R. Penrose, P. Sommers, and M. Walker, “On a quadratic first integral for the charged particle orbits in the charged Kerr solution,” *Commun. Math. Phys.* **27** (1972) 303–308.
- [53] L. P. Hughston and P. Sommers, “Spacetimes with killing tensors,” *Communications in Mathematical Physics* **32** (1973) 147–152.
- [54] J. F. Plebanski, “Some solutions of complex Einstein equations,” *Journal of Mathematical Physics* **16** no. 12, (1975) 2395–2402.
- [55] T. Adamo, L. Mason, and A. Sharma, “Twistor sigma models for quaternionic geometry and graviton scattering,” *Adv. Theor. Math. Phys.* **27** (2023) 623–681, [arXiv:2103.16984 \[hep-th\]](#).
- [56] D. Zwanziger, “Local Lagrangian quantum field theory of electric and magnetic charges,” *Phys. Rev. D* **3** (1971) 880.
- [57] F. V. Gubarev, M. I. Polikarpov, and V. I. Zakharov, “Monopole - anti-monopole interaction in Abelian Higgs model,” *Phys. Lett. B* **438** (1998) 147–153, [arXiv:hep-th/9805175](#).
- [58] Y. Shnir, “Magnetic monopoles,” *Phys. Part. Nucl. Lett.* **8** (2011) 749–754.
- [59] J. Terning and C. B. Verhaaren, “Spurious Poles in the Scattering of Electric and Magnetic Charges,” *JHEP* **12** (2020) 153, [arXiv:2010.02232 \[hep-th\]](#).
- [60] N. Moynihan and J. Murugan, “On-shell electric-magnetic duality and the dual graviton,” *Phys. Rev. D* **105** no. 6, (2022) 066025, [arXiv:2002.11085 \[hep-th\]](#).
- [61] E. T. Newman and A. I. Janis, “Note on the Kerr spinning particle metric,” *J. Math. Phys.* **6** (1965) 915–917.
- [62] Technically, this holds off the Dirac string or when the Dirac string is hidden.
- [63] Note that this metric has a residual freedom of gauge transforming t .
- [64] D. A. Easson, G. Herczeg, T. Manton, and M. Pezzelle, “Isometries and the double copy,” *JHEP* **09** (2023) 162, [arXiv:2306.13687 \[gr-qc\]](#).
- [65] J. F. Plebanski, “On the separation of Einsteinian substructures,” *J. Math. Phys.* **18** (1977) 2511–2520.
- [66] E. T. Newman, “The Bondi-Metzner-Sachs group: Its complexification and some related curious consequences,” in *General relativity and gravitation: proceedings of the seventh international conference (GR7), Tel-Aviv University, June 23-28, 1974, Shaviv, Giora and Rosen, Joseph, Relativity and Gravitation, 1975* (1975) .
- [67] E. T. Newman, “Heaven and its properties,” *General Relativity and Gravitation* **7** no. 1, (1976) 107–111.
- [68] T. T. Wu and C. N. Yang, “Concept of Nonintegrable Phase Factors and Global Formulation of Gauge Fields,” *Phys. Rev. D* **12** (1975) 3845–3857.
- [69] Note that repositioning the string commutes with the double copy: the single copy of Eq. (9) exactly describes the SDD with north-pointing Dirac string.
- [70] defined as a constant since the spacetime is stationary
- [71] Note that $\bar{\xi} \neq \xi$.
- [72] D. Tong, “Gauge theory,” *Lecture notes, DAMTP Cambridge* **10** (2018) .
- [73] C. Cheung, M. Derda, J.-H. Kim, V. Nevoa, I. Rothstein, and N. Shah, “Generalized Symmetry in Dynamical Gravity,” [arXiv:2403.01837 \[hep-th\]](#).
- [74] J. Sakurai, “Comments on quantum-mechanical interference due to the earth’s rotation,” *Physical Review D* **21** no. 10, (1980) 2993.
- [75] R. L. Zimmerman and B. Y. Shahir, “Geodesics for the NUT metric and gravitational monopoles,” *General relativity and gravitation* **21** no. 8, (1989) 821–848.
- [76] To understand this quickly, note that the Coriolis force $2m\vec{v} \times \vec{\omega}$ is analogous to the Lorentz force $q\vec{v} \times \vec{B}$ [74].
- [77] In this sense the “field strength” of gravity is torsion: note that Eq. (12) can be interpreted as the torsion two-form generated by the multivalued diffeomorphism ($ddx^\mu = \Gamma^\mu{}_{\nu\rho} dx^\nu \wedge dx^\rho = \frac{1}{2} T^\mu{}_{\rho\sigma} dx^\rho \wedge dx^\sigma$) [138–140]. This might be suggesting a realization of color-kinematics duality in terms of Einstein-Cartan theory or teleparallel gravity.
- [78] W. Israel, “Source of the Kerr metric,” *Physical Review D* **2** no. 4, (1970) 641.
- [79] H. Balasin and H. Nachbagauer, “Distributional energy momentum tensor of the Kerr-Newman space-time family,” *Class. Quant. Grav.* **11** (1994) 1453–1462, [arXiv:gr-qc/9312028](#).
- [80] H. Balasin and H. Nachbagauer, “The energy-momentum tensor of a black hole, or what curves the Schwarzschild geometry?,” *Class. Quant. Grav.* **10** (1993) 2271, [arXiv:gr-qc/9305009](#).
- [81] in the sense that T^{00} vanishes
- [82] In particular, the Newman-Janis shift is essentially the same argument applied to the Kerr solution [1, 34, 61, 129–131], but concretely verifying that SDTN is the “self-dual part”³ of Kerr requires a nontrivial discussion [36, 132–134].
- [83] A. I. Harte and J. Vines, “Generating exact solutions to Einstein’s equation using linearized approximations,” *Phys. Rev. D* **94** no. 8, (2016) 084009, [arXiv:1608.04359 \[gr-qc\]](#).
- [84] B. C. Xanthopoulos, “Exact vacuum solutions of einstein’s equation from linearized solutions,” *Journal of Mathematical Physics* **19** no. 7, (1978) 1607–1609.
- [85] S. A. Huggett and K. P. Tod, *An Introduction to Twistor Theory*. London Mathematical Society Student Texts. Cambridge University Press, 2 ed., 1994.

- [86] It would be interesting to experiment a further mix-and-match of null GSF congruences, such as one copy of SDTN congruence for the right-handed tensored with one copy of Robinson congruence for the left-handed.
- [87] R. Penrose, “Twistor algebra,” *Journal of Mathematical Physics* **8** no. 2, (1967) 345–366.
- [88] E. T. Newman, “Maxwell fields and shear-free null geodesic congruences,” *Classical and Quantum Gravity* **21** no. 13, (2004) 3197.
- [89] R. Penrose and G. A. J. Sparling, “The twistor quadrille,” *Twistor Newsletter* **1** (1976) 10–13.
- [90] G. A. J. Sparling, “The non-linear graviton representing the analogue of schwarzschild or kerr black holes,” *Twistor Newsletter* **1** (1976) 14–17.
- [91] L. P. Hughston and R. S. Ward, *Advances in twistor theory*, vol. 37. Pitman, 1979.
- [92] This function can be written as $i\tilde{Z}^A u_\mu(\gamma^\mu)_A{}^C \bar{I}_{CB} \tilde{Z}^B$, where \bar{I}_{AB} denotes infinity twistor.
- [93] R. Penrose, “Solutions of the zero-rest-mass equations,” *J. Math. Phys.* **10** (1969) 38–39.
- [94] M. Atiyah, M. Dunajski, and L. Mason, “Twistor theory at fifty: from contour integrals to twistor strings,” *Proc. Roy. Soc. Lond. A* **473** no. 2206, (2017) 20170530, [arXiv:1704.07464 \[hep-th\]](#).
- [95] Ref. [49] has half-Fourier transformed only the massless leg to the twistor space. For the possibility of half-Fourier transforming the massive legs as well, see Ref. [141].
- [96] In fact, the orientation of the Dirac string is determined by not only the three-vector \vec{n} here but also the prescription for the anti-derivative $(\vec{n}\cdot\partial)^{-1}$. In particular, it is placed along the negative z -axis as in Eq. (1) if $\vec{n}\cdot\partial = \partial_z$ and $(\partial_z)^{-1}(1/r) = \log(r+z)$.
- [97] This traces back to the well-known fact in electromagnetism that the magnetic field of Gilbertian and Ampèrian dipoles is identical up to the contact term [142]. In fact, the operator $(\vec{n}\times\partial)/(\vec{n}\cdot\partial)$ in \vec{J} given by Eq. (21) connects between the Gilbertian (separated opposite magnetic monopoles) and Ampèrian pictures (electric current loop) of magnetic dipole, so dropping it gives the magnetic charge density J^{*0} given by Eq. (22).
- [98] P. A. M. Dirac, “The theory of magnetic poles,” *Physical Review* **74** no. 7, (1948) 817.
- [99] D. Zwanziger, “Quantum field theory of particles with both electric and magnetic charges,” *Phys. Rev.* **176** (1968) 1489–1495.
- [100] As will be clear from the exposition in Ref. [60], this propagator essentially traces back to the XZC presentation [143, 144] of massless propagator employing an auxiliary massless vector.
- [101] This is the gravitational analog of “Gilbert-Ampère duality” described in footnote [96]. Note that a static gravitational dyon $(m, -im)$ is precisely described by $T^{\mu\nu} = mu^\mu u^\nu \delta^{(3)}(\vec{x})$ and $T^{*\mu\nu} = -imu^\mu u^\nu \delta^{(3)}(\vec{x})$.
- [102] Y.-T. Huang, U. Kol, and D. O’Connell, “Double copy of electric-magnetic duality,” *Phys. Rev. D* **102** no. 4, (2020) 046005, [arXiv:1911.06318 \[hep-th\]](#).
- [103] W. T. Emond, Y.-T. Huang, U. Kol, N. Moynihan, and D. O’Connell, “Amplitudes from Coulomb to Kerr-Taub-NUT,” *JHEP* **05** (2022) 055, [arXiv:2010.07861 \[hep-th\]](#).
- [104] N. Arkani-Hamed, T.-C. Huang, and Y.-t. Huang, “Scattering amplitudes for all masses and spins,” *JHEP* **11** (2021) 070, [arXiv:1709.04891 \[hep-th\]](#).
- [105] Duff construction [145] from the Gilbertian linearized sources might be not straightforward by the failure of electric-magnetic duality due to the inherent nonlinearity of gravity [146]. Meanwhile, switching between the Ampèrian and Gilbertian pictures at the nonlinear level might be realized by the duality observed in Ref. [147].
- [106] J. F. Plebanski and H. Garcia-Compean, “A q deformed version of the heavenly equations,” *Int. J. Mod. Phys. A* **10** (1995) 3371–3379, [arXiv:hep-th/9405073](#).
- [107] Q.-H. Park, “Selfdual Yang-Mills (+ gravity) as a 2-D sigma model,” *Phys. Lett. B* **257** (1991) 105–110.
- [108] Q.-H. Park, “2-D sigma model approach to 4-D instantons,” *Int. J. Mod. Phys. A* **7** (1992) 1415–1448.
- [109] V. Husain, “Selfdual gravity as a two-dimensional theory and conservation laws,” *Class. Quant. Grav.* **11** (1994) 927–938, [arXiv:gr-qc/9310003](#).
- [110] C. Cheung, J. Mangan, J. Parra-Martinez, and N. Shah, “Non-perturbative Double Copy in Flatland,” *Phys. Rev. Lett.* **129** no. 22, (2022) 221602, [arXiv:2204.07130 \[hep-th\]](#).
- [111] K. Armstrong-Williams, C. D. White, and S. Wikeley, “Non-perturbative aspects of the self-dual double copy,” *JHEP* **08** (2022) 160, [arXiv:2205.02136 \[hep-th\]](#).
- [112] Naively, incorporating the nonabelian color structure seems to not solve the puzzle: as a diagnosis one can check whether the color Killing vector equation [148] in the single copy configuration is mapped to the Killing equation for gravity and find failure. Fundamentally, there seems to be a difference in how the mass dimension increases by one when the gauge potential is mapped to the graviton field, between the self-dual double copy construction of the Eguchi-Hanson instanton in Ref. [7] and the double copy constructions of Taub-NUT instanton presented here. The former keeps the “matter coupling” (denoted λ in Ref. [7]) the same while $\hat{k}_{\hat{\alpha}} = \partial_{0\hat{\alpha}}$ imparts a mass dimension. However, the latter imparts the mass dimension by changing the matter coupling from q to m .
- [113] T. Eguchi and A. J. Hanson, “Asymptotically flat selfdual solutions to euclidean gravity,” *Phys. Lett. B* **74** (1978) 249–251.
- [114] T. Eguchi and A. J. Hanson, “Selfdual solutions to euclidean gravity,” *Annals Phys.* **120** (1979) 82.
- [115] T. Eguchi and A. J. Hanson, “Gravitational instantons,” *Gen. Rel. Grav.* **11** (1979) 315–320.
- [116] K. P. Tod, “Self-dual Kerr–Schild metrics and null Maxwell fields,” *J. Math. Phys.* **23** no. 6, (1982) 1147–1148.
- [117] We thank Andres Luna for suggesting this direction.
- [118] H. Godazgar, M. Godazgar, and C. N. Pope, “Dual gravitational charges and soft theorems,” *JHEP* **10** (2019) 123, [arXiv:1908.01164 \[hep-th\]](#).
- [119] H. Godazgar, M. Godazgar, and C. N. Pope, “Taub-NUT from the Dirac monopole,” *Phys. Lett. B* **798** (2019) 134938, [arXiv:1908.05962 \[hep-th\]](#).
- [120] Note also that the Kerr-Schild metric in Eq. (3) nicely falls into the BMS category of asymptotically flat metrics [149–151], with mass aspect m , angular momentum aspects zero, and $C_{\zeta\zeta} = 4M\zeta^2/(1+\zeta\bar{\zeta})^2$, $C_{\zeta\bar{\zeta}} = 0$. The Gibbons-Hawking metric in Eq. (6), in contrast, fails to satisfy the falloff conditions as $g_{\zeta\bar{\zeta}} = 4r^2(1+\zeta\bar{\zeta})^{-2} + \mathcal{O}(r^{-1})$.
- [121] S. A. Teukolsky, “Perturbations of a rotating black hole. 1. Fundamental equations for gravitational electromagnetic and neutrino field perturbations,” *Astrophys. J.* **185** (1973) 635–647.
- [122] W. H. Press and S. A. Teukolsky, “Perturbations of a Rotating Black Hole. II. Dynamical Stability of the Kerr Metric,” *Astrophys. J.* **185** (1973) 649–674.
- [123] M. Sasaki and H. Tagoshi, “Analytic black hole perturbation approach to gravitational radiation,” *Living Rev. Rel.* **6** (2003) 6, [arXiv:gr-qc/0306120](#).
- [124] S. Mano, H. Suzuki, and E. Takasugi, “Analytic solutions of the Teukolsky equation and their low frequency expansions,” *Prog. Theor. Phys.* **95** (1996) 1079–1096, [arXiv:gr-qc/9603020](#).
- [125] G. Bonelli, C. Iossa, D. P. Lichtig, and A. Tanzini, “Exact solution of Kerr black hole perturbations via CFT2 and instanton counting: Greybody factor, quasinormal modes, and Love numbers,” *Phys. Rev. D* **105** no. 4, (2022) 044047, [arXiv:2105.04483 \[hep-th\]](#).
- [126] E. Poisson, A. Pound, and I. Vega, “The Motion of point particles in curved spacetime,” *Living Rev. Rel.* **14** (2011) 7, [arXiv:1102.0529 \[gr-qc\]](#).
- [127] B. S. DeWitt and R. W. Brehme, “Radiation damping in a gravitational field,” *Annals Phys.* **9** (1960) 220–259.
- [128] L. Barack and A. Ori, “Mode sum regularization approach for the selfforce in black hole space-time,” *Phys. Rev. D* **61** (2000) 061502, [arXiv:gr-qc/9912010](#).
- [129] E. T. Newman, “Maxwell’s equations and complex Minkowski space,” *J. Math. Phys.* **14** (1973) 102–103.

- [130] E. T. Newman, “On a classical, geometric origin of magnetic moments, spin angular momentum and the Dirac gyromagnetic ratio,” *Phys. Rev. D* **65** (2002) 104005, [arXiv:gr-qc/0201055](#).
- [131] T. Adamo and E. T. Newman, “The Kerr-Newman metric: a review,” *arXiv preprint arXiv:1410.6626* (2014) .
- [132] D. J. Gross and M. J. Perry, “Magnetic Monopoles in Kaluza-Klein Theories,” *Nucl. Phys. B* **226** (1983) 29–48.
- [133] A. M. Ghezelbash, R. B. Mann, and R. D. Sorkin, “The Disjointed thermodynamics of rotating black holes with a NUT twist,” *Nucl. Phys. B* **775** (2007) 95–119, [arXiv:hep-th/0703030](#).
- [134] J.-H. Kim, “Spinning black holes and the Dirac string.” To appear.
- [135] J. Desai, G. Herczeg, D. McNutt, and M. Pezzelle, “Taub-NUT Instanton as the Self-dual Analog of Kerr,” [arXiv:2405.15946 \[hep-th\]](#).
- [136] Note that this approach of first working with complexified spacetime and fields and then specializing to particular real sections to derive results in any signature is in accordance with the common practice in twistor theory (see [152], for example).
- [137] To be explicit, the line element $-dt^2 + dx^2 - d\tilde{y}^2 + dz^2 + (K/r)(-dt + (x + j\tilde{y})(dx - j d\tilde{y})/(r + z) + dz)^2$ maps to $-(1 + K/r)^{-1} [dt + (K/r)((x + j\tilde{y})(dx - j d\tilde{y})/(r + z) + dz)]^2 + (1 + K/r)(dx^2 - d\tilde{y}^2 + dz^2)$ through $t \mapsto t, z \mapsto z + K, x \mapsto x + K(x + j\tilde{y})/(r + z), \tilde{y} \mapsto \tilde{y} + j^{-1}K(x + j\tilde{y})/(r + z)$, with $r := (x^2 - \tilde{y}^2 + z^2)^{1/2}$ and $j^2 = 1$. The sign choice for $j = \pm 1$ depends on the sign of the NUT charge or the orientation of the volume form, while $j = +1$ reproduces Eq. (6.16) of Ref. [64]. It could be interesting if a connection to split-complex numbers can be made.
- [138] P. Fiziev and H. Kleinert, “New action principle for classical particle trajectories in spaces with torsion,” *EPL* **35** (1996) 241–246, [arXiv:hep-th/9503074](#).
- [139] H. Kleinert and A. Pelster, “Autoparallels From a New Action Principle,” *Gen. Rel. Grav.* **31** (1999) 1439, [arXiv:gr-qc/9605028](#).
- [140] H. Kleinert, *Multivalued fields. In condensed matter, electromagnetism, and gravitation*. World Scientific, 2008.
- [141] J.-H. Kim, “Asymptotic Spinspacetime,” [arXiv:2309.11886 \[hep-th\]](#).
- [142] D. J. Griffiths, *Introduction to electrodynamics; 3rd ed.* Prentice-Hall, Upper Saddle River, NJ, 1999. Chapter 6. Magnetic Fields in Matter.
- [143] Z. Xu, D.-H. Zhang, and L. Chang, “Helicity Amplitudes for Multiple Bremsstrahlung in Massless Nonabelian Gauge Theories,” *Nucl. Phys. B* **291** (1987) 392–428.
- [144] H. K. Dreiner, H. E. Haber, and S. P. Martin, “Two-component spinor techniques and Feynman rules for quantum field theory and supersymmetry,” *Phys. Rept.* **494** (2010) 1–196, [arXiv:0812.1594 \[hep-ph\]](#).
- [145] M. J. Duff, “Quantum Tree Graphs and the Schwarzschild Solution,” *Phys. Rev. D* **7** (1973) 2317–2326.
- [146] R. Monteiro, “No U(1) ‘electric-magnetic’ duality in Einstein gravity,” [arXiv:2312.02351 \[hep-th\]](#).
- [147] U. Kol and S.-T. Yau, “Duality in Gauge Theory, Gravity and String Theory,” [arXiv:2311.07934 \[hep-th\]](#).
- [148] L. F. Abbott and S. Deser, “Charge definition in non-abelian gauge theories,” *Physics Letters B* **116** no. 4, (1982) 259–263.
- [149] H. Bondi, M. G. J. van der Burg, and A. W. K. Metzner, “Gravitational waves in general relativity. 7. Waves from axisymmetric isolated systems,” *Proc. Roy. Soc. Lond. A* **269** (1962) 21–52.
- [150] R. K. Sachs, “Gravitational waves in general relativity. 8. Waves in asymptotically flat space-times,” *Proc. Roy. Soc. Lond. A* **270** (1962) 103–126.
- [151] A. Strominger, “Lectures on the Infrared Structure of Gravity and Gauge Theory.” no note, 3, 2017.
- [152] T. Adamo, “Lectures on twistor theory,” *PoS Modave2017* (2018) 003, [arXiv:1712.02196 \[hep-th\]](#).