# On embeddability of Coxeter groups into the Riordan group

Tian-Xiao He<sup>1</sup> and Nikolai A. Krylov<sup>2</sup>

<sup>1</sup>Department of Mathematics, Illinois Wesleyan University, 1312 Park Street, Bloomington IL 61702, USA email: the@iwu.edu <sup>2</sup>Department of Mathematics, Siena College, 515 Loudon Road, Loudonville NY 12211, USA email: nkrylov@siena.edu

#### Abstract

We prove that a Coxeter group containing an element of finite order, which is generated by two non-commuting involutions, can not be embedded into the Riordan group.

*2020 Mathematics Subject Classification*: 20F05, 20F55, 20H20, 05A05, 05E16 *Keywords*: Coxeter group, Riordan group, Riordan array, Involution.

#### 1 Introduction

The Riordan group consists of infinite lower-triangular matrices with no zeroes on the main diagonal. For example, the alternating Pascal's matrix, where the *n*-th row consists of the signed binomial coefficients (assuming that  $\binom{n}{k}$  ${k \choose k} = 0$ , if  $k > n$ )

<span id="page-0-0"></span>
$$
P_1 := \left\{ (-1)^k \binom{n}{k} \right\}_{n,k \ge 0} = \begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ 1 & -1 & 0 & 0 & \cdots \\ 1 & -2 & 1 & 0 & \cdots \\ 1 & -3 & 3 & -1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}, \quad (1.1)
$$

is a famous element of this group. Each column of  $P_1$  consists of the coefficients of a formal power series (f.p.s.): the zeroth column is given by the geometric series  $1/(1-t)$ , and the k-th column, for an arbitrary  $k \geq 0$ , is given by the coefficients of  $(-t)^k/(1-t)^{k+1}$ . The group operation is the usual row-by-column matrix multiplication, and one quickly notices that the square of  $P_1$  will give us the infinite identity matrix with ones on the main diagonal and zeros everywhere else. Thus,  $P_1$  is an element of order 2, which is called an involution.

The Riordan group was introduced by Shapiro et al. [\[20\]](#page-12-0) in 1991. The elements of this group, called Riordan arrays, proved to be useful in evaluating combinatorial sums in a uniform and constructive way. A Riordan array is defined as a pair of formal power series with coefficients in an arbitrary field  $K$  (e.g.  $\mathbb{C}$ ). When there are no zeroes on the main diagonal, we say that the Riordan array is proper.

More formally, let us consider the set of all formal power series (f.p.s.)  $\mathcal{F} = \mathbb{K}[[t]]$  in indeterminate t with coefficients in K; the *order* of  $f(t) \in \mathcal{F}$ ,  $f(t) = \sum_{k=0}^{\infty} f_k t^k$   $(f_k \in \mathbb{K})$ , is the minimal number  $r \in \mathbb{N}_0 = \{0\} \cup \mathbb{N}$  such that  $f_r \neq 0$ . Denote by  $\mathcal{F}_r$  the set of formal power series of order r. Let  $g(t) \in \mathcal{F}_0$  and  $f(t) \in \mathcal{F}_1$ ; the pair  $(g(t), f(t))$  defines the *(proper) Riordan array*

 $A = (a_{n,k})_{n,k \ge 0} = (g(t), f(t))$ 

having

$$
a_{n,k} = [t^n]g(t)f(t)^k,
$$
\n(1.2)

where  $[t^n]h(t)$  denotes the coefficient of  $t^n$  in the expansion of a f.p.s.  $h(t)$ . The reader will find all the main properties of the functionals  $[t^n]$  in the Monthly's paper by Merlini et al.  $[14]$  from 2007. It is a classical fact now that the set of all such Riordan arrays forms a multiplicative group, called the Riordan group, and denoted by  $\mathcal{R}(\mathbb{K})$  (see, for example, [\[1\]](#page-10-0), [\[4\]](#page-11-1), [\[9\]](#page-11-2), [\[19\]](#page-12-1), [\[20\]](#page-12-0)). The group operation  $*$  in  $\mathcal{R}(\mathbb{K})$  (will be omitted below for brevity) is written in terms of the pairs of the f.p.s. as

$$
(g_1(t), f_1(t)) * (g_2(t), f_2(t)) = (g_1(t)g_2(f_1(t)), f_2(f_1(t))), \quad (1.3)
$$

with the Riordan array  $I = (1, t)$  acting as the group identity. The inverse of the Riordan array  $(g(t), f(t))$  is the pair

$$
(g(t), f(t))^{-1} = \left(\frac{1}{g(\bar{f}(t))}, \bar{f}(t)\right),
$$

where we used the standard notation  $\bar{f}(t)$  for the compositional inverse of  $f(t)$ . Thus,  $\bar{f}(f(t)) = t$  and  $f(\bar{f}(t)) = t$ . Notice, that for  $g(t) \in \mathcal{F}_0$  and

 $f(t) \in \mathcal{F}_1$  to have the multiplicative and compositional inverses respectively, we must require that  $g(0) = [t^0]g(t) \neq 0$  and  $f'(0) = [t^1]f(t) \neq 0$  in the field K. The Riordan group  $\mathcal{R}(\mathbb{K})$ , is the semidirect product of two proper subgroups: the Appell subgroup and the associated (or Lagrange) subgroup ([\[1\]](#page-10-0), [\[19\]](#page-12-1)). The Appell subgroup  $A$  is abelian, normal, and consists of the Riordan arrays  $(g(t), t)$ . The Lagrange subgroup  $\mathcal L$  consists of the Riordan arrays  $(1, f(t))$ .

If G is a multiplicative group with the identity e, an element  $q \in G$  is called an involution if it is of oder 2, i.e., if  $g^2 = e$ . Thus,  $(g, f) \in \mathcal{R}(\mathbb{K})$ is an (Riordan) involution if  $(g, f)^2 = (g \cdot g(f), f(f)) = (1, t)$ . Namely,  $g(f) = g \circ f = 1/g$  and  $f = \overline{f}$ . The alternating Pascal's matrix  $P_1$  is an example of such involution. The classical Pascal's matrix consisting of the non-negative binomial coefficients is a so-called pseudo-involution. An element  $(q, f)$  in the Riordan group is said to be a pseudo-involution (PI) if and only if

 $(g, f)(1, -t) = (g, -f)$  or  $(1, -t)(g, f) = (g(-t), f(-t))$ 

is an involution. Notice that  $(1, -t) \in \mathcal{L}$  is an involution which is not in the center of  $\mathcal{R}(\mathbb{K})$ , while  $(-1, t) \in \mathcal{A}$  is an involution, which commutes with every element of  $\mathcal{R}(\mathbb{K})$ .

Since its introduction, the group  $\mathcal{R}(\mathbb{K})$  has been of great interest in combinatorics and by now, there are numerous results clarifying the algebraic structure of this group. For instance, in Luzón et al.  $[13]$ , the Riordan group over a field of characteristic zero  $\mathcal{R}(\mathbb{K})$ , is described as an inverse limit of a certain inverse sequence of groups  $\mathcal{R}_n(\mathbb{K})$  formed by  $(n+1)\times(n+1)$  invertible lower triangular matrices; i.e.,  $\mathcal{R}_n(\mathbb{K})$  is the *n*th truncation of  $\mathcal{R}(\mathbb{K})$ . Luzón, Morón, and Prieto-Martinez [\[12\]](#page-11-4) proved that any element in the group  $\mathcal{R}(\mathbb{K})$ or  $\mathcal{R}_n(\mathbb{K})$  for  $n \geq 1$  generated by the involutions in the group is the product of at most four of them. The elements of finite order, and the involutions and pseudo-involutions in particular, have been studied by many mathematicians. Here is by no means exhaustive, a list of the corresponding publications: [\[2\]](#page-10-1), [\[3\]](#page-11-5), [\[5\]](#page-11-6), [\[6\]](#page-11-7), [\[7\]](#page-11-8), [\[9\]](#page-11-2), [\[10\]](#page-11-9), [\[12\]](#page-11-4), [\[18\]](#page-12-2).

The alternating Pascal's matrix  $P_1$  in [\(1.1\)](#page-0-0) is easily generalized to produce infinitely many similar involutions. Take any  $b \in \mathbb{K}$ , and consider the generalized Pascal's matrix (see §2 of [\[3\]](#page-11-5)) of the form

<span id="page-2-0"></span>
$$
P_b = \left(\frac{1}{1 - bt}, \frac{-t}{1 - bt}\right). \tag{1.4}
$$

We leave it to the reader to check that  $P<sub>b</sub>$  is indeed an involution, and that  $P_{b_1}$  and  $P_{b_2}$  do not commute, as long as  $b_1 \neq b_2$ .

It is well known that the symmetric group  $S_n$ , consisting of all bijections from the set  $\{1, 2, \ldots, n\}$  to itself, is generated by transpositions (permuta-tions of order 2), i.e. involutions. It is also known (see [\[8\]](#page-11-10), §6.2) that  $S_n$  has the following presentation in terms of generators and relations

<span id="page-3-1"></span>
$$
S_n \cong \langle \sigma_1, \ldots, \sigma_{n-1} | \sigma_i^2 = 1, \sigma_i \sigma_j = \sigma_j \sigma_i \text{ for } |i - j| > 1, (\sigma_i \sigma_{i+1})^3 = 1 \rangle, (1.5)
$$

where  $\sigma_i = (i \ i+1), \ \forall i \in \{1, \ldots, n-1\}$  are the generating involutions. In particular, for  $n = 3$  we have

<span id="page-3-0"></span>
$$
S_3 \cong \langle \sigma_1, \sigma_2 \mid \sigma_1^2 = \sigma_2^2 = (\sigma_1 \sigma_2)^3 = 1 \rangle. \tag{1.6}
$$

Moreover, the elements of  $S_3$  can be naturally embedded, preserving the group operation, into the general linear group of  $3 \times 3$  matrices  $GL(3, \mathbb{R})$  as

$$
e \leftrightarrow \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \sigma_1 \leftrightarrow \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \sigma_2 \leftrightarrow \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix},
$$

$$
\sigma_1 \sigma_2 \leftrightarrow \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad \sigma_2 \sigma_1 \leftrightarrow \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \quad \sigma_1 \sigma_2 \sigma_1 \leftrightarrow \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}.
$$

This is an example of a faithful representation of  $S_3$  by  $3 \times 3$  permutation matrices. Analogous representation of  $\mathcal{S}_n$  exists for every  $n \in \mathbb{N}$  (see, for example, [\[17\]](#page-12-3), §1.3). The Riordan group consists of infinite lower triangular matrices and has plenty of involutions, so a natural question arises: what are faithful representations of  $S_n$  by Riordan arrays? Surprisingly, it turns out that when  $n \geq 3$ , there are none. It will follow from our theorem proven below, that for  $n \geq 3$ ,  $S_n$  can not be embedded into the Riordan group. But first note, that since  $\mathbb R$  has only two roots of unity,  $\pm 1$ , the only realvalued Riordan arrays of finite order, which are different from the identity, are the elements of order 2, and  $\mathcal{R}(\mathbb{R})$  can not contain an element  $r \neq 1$ , such that  $r^3 = 1$ . However, if we consider the Riordan arrays over  $\mathbb{C}$ , then using the complex primitive roots of unity, one easily generalizes the idea of the alternating Pascal's matrix, and produces an element of arbitrary finite order (see  $[6]$ ). Thus, from now on, we consider the Riordan group over the complex numbers, and denote this group simply by  $\mathcal{R}$ .

In the view of symmetry of the products, we are also inspired by the palindromes of (Riordan) involutions and pseudo-involutions to consider the identification of the product of three involutions in a palindromic form. For

instance, taking two involutions  $\sigma_1$  and  $\sigma_2$ , it is immediate that the palindromes  $\sigma_1 \sigma_2 \sigma_1$  and  $\sigma_2 \sigma_1 \sigma_2$  will be also involutions. The same statement holds true for the pseudo-involutions (see section 3 of  $[10]$ ). For involutions  $\sigma_1$  and  $\sigma_2$ , the identities  $(\sigma_1 \sigma_2)^3 = 1$  and  $\sigma_1 \sigma_2 \sigma_1 = \sigma_2 \sigma_1 \sigma_2$  are equivalent. Therefore, if we could find two non-commuting Riordan involutions  $\sigma_1$  and  $\sigma_2$  so that the palindromes  $\sigma_1 \sigma_2 \sigma_1$  and  $\sigma_2 \sigma_1 \sigma_2$  were equal, the presentation  $(1.6)$  would imply that  $S_3$  is embeddable into the Riordan group. However, this case seems unlikely, if we try to find such involutions among the Pascal type Riordan arrays we have mentioned above [\(1.4\)](#page-2-0). Indeed, for the Riordan arrays with  $b_1 \neq b_2$ 

$$
\sigma_1 = (g_1, f_1) = \left(\frac{1}{1 - b_1 t}, \frac{at}{1 - b_1 t}\right)
$$
  

$$
\sigma_2 = (g_2, f_2) = \left(\frac{1}{1 - b_2 t}, \frac{at}{1 - b_2 t}\right),
$$
 (1.7)

we may find that

<span id="page-4-1"></span><span id="page-4-0"></span>
$$
\sigma_1 \sigma_2 \sigma_1 = \sigma_2 \sigma_1 \sigma_2 \tag{1.8}
$$

holds if and only if  $a = \frac{1+\sqrt{3}i}{2}$  $\frac{\sqrt{3}i}{2}$  or  $\frac{1-\sqrt{3}i}{2}$  $\frac{\sqrt{3}i}{2}$ . This implies that  $(1.8)$  does not hold for involutions or pseudo-involutions  $\sigma_1$  and  $\sigma_2$  in the form shown in [\(1.7\)](#page-4-1).

In the next section, we will recall some preliminary knowledge about Coxeter groups, including the symmetric groups, dihedral groups, etc., and show how to embed two particular Coxeter groups (the Klein four-group and the infinite dihedral group) into  $\mathcal{R}$ . In the last section, we use the truncated Riordan arrays and  $\mathcal{R}_n(\mathbb{C})$  defined in [\[13\]](#page-11-3), to prove that the embeddability of non-abelian Coxeter groups into the Riordan group does not hold in general.

### 2 Coxeter groups

A Coxeter group G can be introduced using generators and defining relations as

<span id="page-4-2"></span>
$$
G \cong \langle r_1, \dots, r_k \mid (r_i r_j)^{m_{ij}} = 1 \rangle, \tag{2.1}
$$

where  $m_{ii} = 1$  and  $m_{ij} = m_{ji} \geq 2$  is either an integer or  $\infty$  for  $i \neq j$ . If there is no relation between  $r_i$ , and  $r_j$ ,  $(r_i r_j)^{m_{ij}} = 1$  for any integer  $m_{ij} \geq 2$ , we assume  $m_{ij} = \infty$ . (see §5.1 of [\[11\]](#page-11-11)). In particular, the symmetric group  $S_n$ of order n!, and the dihedral group  $\mathcal{D}_n$  of order  $2n$  are examples of Coxeter groups. Recall that  $S_n$  has a presentation  $(1.5)$ , and  $\mathcal{D}_n$  has a presentation

$$
\mathcal{D}_n \cong \langle r_1, r_2 \mid r_1^2 = r_2^2 = 1, (r_1 r_2)^n = 1 \rangle. \tag{2.2}
$$

The relations  $(r_{ii})^1 = 1$  in [\(2.1\)](#page-4-2), mean that all the generators of G are involutions, and G is commutative if and only if  $m_{ij} \leq 2$ , for all  $i \neq j$ . Thus, if  $m_{ij} > 2$ , the involutions  $r_i$  and  $r_j$  do not commute, while if we have a group G where all  $m_{ij} \leq 2$ , then the group is commutative. For example, the Klein four-group

$$
K_4 = \mathcal{D}_2 = \langle r_1, r_2 | r_1^2 = r_2^2 = 1, (r_1 r_2)^2 = 1 \rangle \cong C_2 \times C_2
$$

is a Coxeter group  $(C_2$  stands for the cyclic group of order two) and can be embedded into the Riordan group  $\mathcal R$  using the Riordan arrays

$$
r_1 := P_0 = (1, -t),
$$
 and  $r_2 := (-1, t).$ 

Another example of a Coxeter group, which can be embedded into  $\mathcal R$  is the infinite dihedral group  $\mathcal{D}_{\infty}$ . This group is isomorphic to the semidirect product  $\mathbb{Z} \rtimes C_2$  (see §2.2 of [\[16\]](#page-12-4)), and has a presentation

$$
\mathcal{D}_{\infty} \cong \langle s, t \mid s^2 = t^2 = 1 \rangle.
$$

We prove a bit more general result. Let  $\mathbb{D} \subset \mathbb{C}$  be an integral domain, denote by  $\mathbb{D}^*$  the multiplicative group of units of  $\mathbb{D}$ , and consider the subgroup  $\Delta(2,\mathbb{D})$  of  $GL(2,\mathbb{D})$  consisting of the matrices

$$
\begin{pmatrix} 1 & 0 \\ b & a \end{pmatrix}, \text{ where } b \in \mathbb{D}, a \in \mathbb{D}^*.
$$

Define a map from  $\Delta(2,\mathbb{D})$  into the Lagrange subgroup  $\mathcal L$  of the Riordan group  $\mathcal R$  by

<span id="page-5-1"></span>
$$
f: \begin{pmatrix} 1 & 0 \\ b & a \end{pmatrix} \longmapsto \begin{pmatrix} 1, \frac{ax}{1+bx} \end{pmatrix} \in \mathcal{L}.
$$
 (2.3)

It is easy to check that this  $f$  is a monomorphism. Next lemma shows that  $\Delta(2,\mathbb{D})$  is isomorphic to a semidirect product of  $\mathbb{D}$  and  $\mathbb{D}^*$ .

**Lemma 1.**  $\Delta(2,\mathbb{D})$  *is isomorphic to the semidirect product of the additive group* D *and multiplicative group of units* D ∗ *, that is*

$$
\Delta(2,\mathbb{D}) \cong \mathbb{D} \rtimes_{\varphi} \mathbb{D}^*,
$$

where  $\varphi : \mathbb{D}^* \to Aut(\mathbb{D})$  *is a homomorphism defined by*  $\varphi(a)(b) = \varphi_a(b) = ab$ .

*Proof.* The binary operation  $\circ$  on  $\mathbb{D} \rtimes_{\varphi} \mathbb{D}^*$  is defined by

<span id="page-5-0"></span>
$$
(b, a) \circ (y, x) = (b + \varphi_a(y), ax) = (b + ay, ax). \tag{2.4}
$$

Since we clearly have

$$
\mathbb{D} \cong \left\{ \begin{pmatrix} 1 & 0 \\ b & 1 \end{pmatrix} \mid b \in \mathbb{D} \right\} \triangleleft \Delta(2, \mathbb{D}), \text{ and } \mathbb{D}^* \cong \left\{ \begin{pmatrix} 1 & 0 \\ 0 & a \end{pmatrix} \mid a \in \mathbb{D}^* \right\},\
$$

it is natural to associate the matrix

$$
\begin{pmatrix} 1 & 0 \\ b & a \end{pmatrix}
$$

with the pair  $(b, a) \in \mathbb{D} \rtimes_{\varphi} \mathbb{D}^*$ . Then the automorphism  $\varphi_a : \mathbb{D} \to \mathbb{D}$ corresponding to  $a \in \mathbb{D}^*$  will be written as

$$
\varphi_a: \begin{pmatrix} 1 & 0 \\ x & 1 \end{pmatrix} \to \begin{pmatrix} 1 & 0 \\ ax & 1 \end{pmatrix},
$$

and we can write the binary operation  $(2.4)$  in terms of the matrices as

$$
\left(\begin{pmatrix} 1 & 0 \\ b & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & a \end{pmatrix}\right) \circ \left(\begin{pmatrix} 1 & 0 \\ y & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & x \end{pmatrix}\right) = \left(\begin{pmatrix} 1 & 0 \\ b + ay & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & ax \end{pmatrix}\right).
$$

Since in  $\Delta(2,\mathbb{D})$  we have

$$
\begin{pmatrix} 1 & 0 \ b & a \end{pmatrix} \cdot \begin{pmatrix} 1 & 0 \ y & x \end{pmatrix} = \begin{pmatrix} 1 & 0 \ b + ay & ax \end{pmatrix},
$$

 $\Box$ 

the proof is finished.

Corollary 2. *There is a faithful representation of the infinite dihedral group* D<sup>∞</sup> *by Riordan arrays.*

*Proof.* Take  $\mathbb{D} = \mathbb{Z}$  and apply the monomorphism f defined in [\(2.3\)](#page-5-1) to the group  $\Delta(2,\mathbb{Z}) \cong \mathcal{D}_{\infty}$ .  $\Box$ 

# 3 The main theorem.

Our next result shows in particular, that finite non-commutative Coxeter groups in general, can not be embedded into  $\mathcal{R}$ .

**Theorem 3.** Let G be a Coxeter group with at least two generators  $r_1$  and  $r_2$ *, such that their product*  $r_1r_2$  *has a finite order*  $m \geq 3$ *, i.e.*  $(r_1r_2)^m = 1$ *. Then there exists no monomorphism*

$$
\mu: G \hookrightarrow \mathcal{R}.
$$

*Proof.* Suppose there exists such a monomorphism  $\mu$ . We let  $\sigma_i := \mu(r_i)$ , so  $\sigma_i \in \mathcal{R}$  are the elements of the Riordan group  $\mathcal{R}$ , which satisfy the relations

$$
\sigma_i^2 = 1
$$
, and  $(\sigma_1 \sigma_2)^m = (1, t)$ ,  $m \ge 3$ . (3.1)

Thus, we assume that there are two Riordan involutions the product of which has order  $m \geq 3$ . Take the f.p.s.  $q_i(t)$  and  $f_i(t)$  such that

$$
\sigma_i = (g_i(t), f_i(t)), \ i \in \{1, 2\}.
$$

Using the group operation in  $\mathcal R$  together with  $\sigma_i^2 = 1$ , we see that  $f_i$  and  $g_i$ satisfy

<span id="page-7-0"></span>
$$
f_i(f_i(t)) = t
$$
, and  $g_i(t)g_i(f_i(t)) = 1$ . (3.2)

Let us write each  $f_i(t)$  as a series expansion

$$
f_1(t) = \sum_{i \ge 1} a_i t^i, a_1 \ne 0
$$
 and  $f_2(t) = \sum_{i \ge 1} b_i t^i, b_1 \ne 0$ .

In order to satisfy the first identity of  $(3.2)$ , we must have  $a_1^2 = b_1^2 = 1$ , and it is a simple math induction exercise to show that if  $a_1 = 1$  then  $f_1(t) = t$ . This is an element of compositional order one, and for  $\sigma_1$  to be a proper (i.e. not the identity) involution, we must have  $\sigma_1 = (-1, t)$ . But  $(-1, t)$ commutes with every element of  $\mathcal{R}$ , and can't be  $\mu(r_1)$ , since  $r_1$  and  $r_2$  do not commute. Therefore we can assume that  $a_1 = b_1 = -1$ . Consider now the Riordan arrays

$$
\hat{\sigma}_1 := (1, f_1(t)) = \begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ 0 & -1 & 0 & 0 & \cdots \\ 0 & a_2 & 1 & 0 & \cdots \\ 0 & a_3 & -2a_2 & -1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},
$$

and  $\hat{\sigma}_2 := (1, f_2(t))$  where all  $a_i$  are replaced by the corresponding  $b_i$ . Since the *m*-th compositional power of  $(f_1 \circ f_2)$  satisfies

$$
(f_1 \circ f_2)^{(m)} = Id,
$$

we must also have

<span id="page-7-1"></span>
$$
\left(\begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ 0 & -1 & 0 & 0 & \cdots \\ 0 & a_2 & 1 & 0 & \cdots \\ 0 & a_3 & -2a_2 & -1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \cdot \begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ 0 & -1 & 0 & 0 & \cdots \\ 0 & b_2 & 1 & 0 & \cdots \\ 0 & b_3 & -2b_2 & -1 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \right)^m = Id. (3.3)
$$

Moreover, since the matrices are lower triangular, the identity [\(3.3\)](#page-7-1) must hold true for every product of the truncated matrices of the same size corresponding to  $\hat{\sigma}_1$  and  $\hat{\sigma}_2$ . In particular,

$$
\begin{pmatrix} 1 & 0 \ 0 & 1 \end{pmatrix} = \left( \begin{pmatrix} -1 & 0 \ a_2 & 1 \end{pmatrix} \cdot \begin{pmatrix} -1 & 0 \ b_2 & 1 \end{pmatrix} \right)^m = \begin{pmatrix} 1 & 0 \ m(b_2 - a_2) & 1 \end{pmatrix}, \quad (3.4)
$$

which gives  $b_2 = a_2$ . Now we use the strong mathematical induction to show that the existence of a monomorphism  $\mu$  must imply  $f_1(t) = f_2(t)$ . Let us assume that  $f_1(t) \neq f_2(t)$ . Then there exists the smallest integer  $n \geq 3$  such that  $a_n \neq b_n$  and  $a_i = b_i, \forall i \in \{1, 2, \ldots, n-1\}$ . Consider the truncated matrices of order  $n+1$  corresponding to the elements  $\hat{\sigma}_i$  and remove the zeroth row and column, so the upper left element in each truncated matrix will be −1. We enote the truncated matrices by  $\tau_i$  respectively. Since  $a_i = b_i, \forall i \in \{1, 2, \ldots, n-1\},$  we can write these matrices as

<span id="page-8-0"></span>
$$
\tau_1 = \begin{pmatrix}\n-1 & 0 & 0 & \cdots & 0 \\
a_2 & 1 & 0 & \cdots & 0 \\
a_3 & -2a_2 & -1 & \cdots & 0 \\
\vdots & \vdots & \vdots & \ddots & \vdots \\
a_n & [t^n]f_1^2(t) & [t^n]f_1^3(t) & \cdots & (-1)^n\n\end{pmatrix}
$$
\n(3.5)

and

<span id="page-8-1"></span>
$$
\tau_2 = \begin{pmatrix}\n-1 & 0 & 0 & \cdots & 0 \\
a_2 & 1 & 0 & \cdots & 0 \\
a_3 & -2a_2 & -1 & \cdots & 0 \\
\vdots & \vdots & \vdots & \ddots & \vdots \\
b_n & [t^n]f_1^2(t) & [t^n]f_1^3(t) & \cdots & (-1)^n\n\end{pmatrix},
$$
\n(3.6)

so  $\tau_1$  and  $\tau_2$  have all the corresponding entries the same, except possibly for the first entries in the last (n-th) rows, which are  $a_n$  in  $\tau_1$  and  $b_n$  in  $\tau_2$ . Since  $\tau_i$  has order 2, and all the entries in  $\tau_1$  and  $\tau_2$  are the same except for  $a_n$  and  $b_n$ , we deduce that the product matrix  $\tau_1\tau_2$  will have 1 on the main diagonal, and zeroes everywhere else except possibly the first element of the n-th row. If we multiply the *n*-th row of  $(3.5)$  by its first column, it follows again from the equality  $\tau_1^2 = Id$  that

<span id="page-8-2"></span>
$$
-a_n + [t^n]f_1^2(t)a_2 + [t^n]f_1^3(t)a_3 + \dots + [t^n]f_1^{n-1}(t)a_{n-1} + (-1)^n a_n = 0.
$$
 (3.7)

On the other hand, if we multiply the *n*-th row of  $(3.5)$  by the first column of  $(3.6)$ , we will get the expression

<span id="page-8-3"></span>
$$
-a_n + [t^n]f_1^2(t)a_2 + [t^n]f_1^3(t)a_3 + \dots + [t^n]f_1^{n-1}(t)a_{n-1} + (-1)^n b_n.
$$
 (3.8)

Comparing  $(3.7)$  and  $(3.8)$ , we see that the product of the last row of  $(3.5)$ by the first column of  $(3.6)$  equals  $(-1)^n(b_n - a_n)$ . Hence the matrix of the product  $\tau_1\tau_2$  equals

$$
\Pi_n = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 0 \\ X & 0 & 0 & \cdots & 0 & 1 \end{pmatrix},
$$

where  $X = (-1)^n (b_n - a_n)$ . The mathematical induction shows that for any  $m \geq 1$  we have

$$
\Pi_n^m = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & 0 \\ mX & 0 & 0 & \cdots & 0 & 1 \end{pmatrix},
$$

and since  $\Pi_n^m = Id$ , we must have  $X = 0$ . In particular,  $b_n = a_n$ , which contradicts the assumption that  $a_n \neq b_n$ . Therefore  $f_1(t) = f_2(t)$  and we can write our Riordan involutions  $\sigma_1$  and  $\sigma_2$  as

$$
\sigma_1 = (g_1(t), f(t))
$$
 and  $\sigma_2 = (g_2(t), f(t)),$ 

where  $f(t) = f_1(t) = f_2(t)$ . The second of  $(3.2)$  identities implies that  $g_i(f(t)) = 1/g_i(t)$ . Therefore  $\sigma_1 \sigma_2$  equals

<span id="page-9-0"></span>
$$
(g_1(t), f(t))(g_2(t), f(t)) = (g_1(t)g_2(f(t)), f(f(t))) = \left(\frac{g_1(t)}{g_2(t)}, t\right). \quad (3.9)
$$

Hence  $\sigma_1 \sigma_2$  is an element of the commutative (Appell) subgroup of  $\mathcal{R}$ . The second of [\(3.2\)](#page-7-0) identities also means that  $g_i^2(0) = 1$ . Since  $(\sigma_1 \sigma_2)^m = (1, t)$ , [\(3.9\)](#page-9-0) implies

$$
(1,t) = \left(\frac{g_1(t)}{g_2(t)}, t\right)^m = \left(\frac{g_1^m(t)}{g_2^m(t)}, t\right),
$$

so  $g_1^m(t) = g_2^m(t)$ , and if m is even,  $g_1^m(0) = 1$ . In this case Theorem 3 of [\[15\]](#page-11-12) says that  $g_1(t) = g_2(t)$ , which contradicts the assumption that  $r_1$  and  $r_2$  do not commute. If m is odd, and  $g_1(0) = 1$  then  $g_2(0) = 1$  and we obtain the same contradiction, so let us assume that  $g_i(0) = -1$  for each  $i \in \{1, 2\}$ . But in this case we can consider two involutions  $\bar{\sigma}_i := (-g_i(t), f(t))$  and apply the previous argument to obtain the equality  $-g_1(t) = -g_2(t)$  and finish the proof by contradiction.  $\Box$  **Corollary 4.** For any  $n \geq 3$ , neither  $S_n$ , no  $\mathcal{D}_n$  can be embedded into R.

Corollary 5. *The projective linear group* PGL(2, Z) *can not be embedded into the Riordan group* R*.*

*Proof.* It follows from Example 2 in §5.1 of [\[11\]](#page-11-11), where it is shown that  $PGL(2, \mathbb{Z})$  is isomorphic to a Coxeter group

$$
\text{PGL}(2,\mathbb{Z}) \cong \langle r_1, r_2, r_3 \mid r_i^2 = (r_1 r_2)^3 = (r_1 r_3)^2 = 1 \rangle.
$$

 $\Box$ 

Corollary 6. *A product of two non-commuting Riordan involutions has the infinite order.*

One may wonder here, if a product of more than two non-commuting Riordan involutions can have a finite order. The generalized Pascal's matrices [\(1.4\)](#page-2-0) give the affirmative answer to this question. Since  $\sum^{2n+1}$  $_{k=1}$  $(-1)^{k-1}k = n + 1,$ the induction shows that the product of  $2n + 1$  Riordan involutions  $P_b$  with  $b \in \{1, 2, \ldots, 2n + 1\},\$ 

$$
P_1 \cdot P_2 \cdot \ldots \cdot P_{2n+1} = P_{n+1}
$$

is again the Riordan involution, so it has the order 2. Finally, the product of four non-commuting involutions can have order 1, since  $P_1 \cdot P_2 \cdot P_5 \cdot P_4 = (1, t)$ .

### Acknowledgement

The authors sincerely thank the organizers of the Special Session on Riordan Arrays of the AMS 2024 Spring Eastern Sectional Meeting for providing an effective platform for academic exchange and cooperation. The authors also thank Lou Shapiro and Alex Burstein for the fruitful discussions about involutions and pseudo-involutions during the conference.

# <span id="page-10-0"></span>References

- <span id="page-10-1"></span>[1] P. Barry, *Riordan Arrays: A Primer*, Logic Press, 2016.
- [2] A. Burstein and L.W. Shapiro, Pseudo-Involutions in the Riordan Group, *Journal of Integer Sequences*, Vol 25 (2022), Article 22.3.6.
- <span id="page-11-5"></span><span id="page-11-1"></span>[3] N.T. Cameron and A. Nkwanta, On some (pseudo) involutions in the Riordan group, *Journal of Integer Sequences*, Vol 8 (2005), Article 05.3.7.
- <span id="page-11-6"></span>[4] N.T. Cameron and A. Nkwanta, Riordan Matrices and Lattice Path Enumeration, *Notices of the AMS*, Vol. 70, no. 2 (2023), 231–243.
- [5] Gi-Sang Cheon, Hana Kim, Simple proofs of open problems about the structure of involutions in the Riordan group, *Linear Algebra Appl.* 428 (2008) 930–940.
- <span id="page-11-7"></span>[6] Gi-Sang Cheon, Hana Kim, The elements of finite order in the Riordan group over the complex field, *Linear Algebra Appl.* 439 (2013) 4032– 4046.
- <span id="page-11-10"></span><span id="page-11-8"></span>[7] M. M. Cohen, Elements of finite order in the Riordan group and their eigenvectors, *Linear Algebra Appl.* 602 (2020), 264–280.
- [8] H.S.M. Coxeter, W.O.J. Moser, *Generators and Relations for Discrete Groups*, 4th Edition, Ergebnisse der Mathematik und ihrer Grenzgebiete 14.
- <span id="page-11-2"></span>[9] D.E. Davenport, S.K. Frankson, L.W. Shapiro, L.C. Woodson, An Invitation to the Riordan Group, *Enumerative Combinatorics and Applications*, ECA 4:3 (2024), Article # S2S1.
- <span id="page-11-9"></span>[10] T.-X. He and L.W. Shapiro, Palindromes and pseudo-involution multiplication, *Linear Algebra Appl.* 593 (2020), 1–17.
- <span id="page-11-11"></span><span id="page-11-4"></span>[11] J. Humphreys *Reflection groups and Coxeter groups*, Cambridge University Press, 1990.
- $[12]$  A. Luzón, M.A. Morón, L.F. Prieto-Martínez, The group generated by Riordan involutions, *Revista Matem´atica Complutense* 35 (2018), 199– 217.
- <span id="page-11-3"></span>[13] A. Luzón, D. Merlini, M.A. Morón, L.F. Prieto-Martínez and R. Sprugnoli, Some inverse limit approaches to the Riordan group, *Linear Algebra Appl.* 491 (2016), 239–262.
- <span id="page-11-12"></span><span id="page-11-0"></span>[14] D. Merlini, R. Sprugnoli, M.C. Verri, The Method of Coefficients, *The American Mathematical Monthly*, 2007, Vol. 114, No. 1, 40–57.
- [15] Ivan Niven, Formal power series, *Am. Math. Mon.* 76 (1969), 871–889.
- <span id="page-12-4"></span><span id="page-12-3"></span>[16] Derek J.S. Robinson, *A Course in the Theory of Groups*, Springer GTM  $#80$ , Second Edition, 1996.
- [17] B. Sagan *The Symmetric Group Representations, Combinatorial Algorithms, and Symmetric Functions*, Springer GTM # 203, 2nd edition, 2001.
- <span id="page-12-2"></span>[18] L.W. Shapiro, Some open questions about random walks, involutions, limiting distributions and generating functions, *Advances in Applied Math.* 27 (2001), 585–596.
- <span id="page-12-1"></span><span id="page-12-0"></span>[19] L.W. Shapiro, R. Sprugnoli, P. Barry, G. Cheon, T.-X. He, D. Merlini, W. Wang, *The Riordan Group and Applications*, Springer, 2022.
- [20] L.W. Shapiro, S. Getu, W.-J. Woan, L. Woodson, The Riordan group, *Discrete Appl. Math.* 34 (1991), 229–239.